Title: Linking Choroid Plexus Enlargement with Plasma Analyte and Structural Phenotypes in Clinical High Risk for Psychosis: A Multisite Neuroimaging Study

Short Title: Choroid Plexus Enlargement in Clinical High Risk for Psychosis

Authors: Deepthi Bannai[1], Martin Reuter[2-4], Rachal Hegde[1], Dung Hoang[1], Iniya Adhan[1], Swetha Gandu[1], Sovannarath Pong[1], Alexandra Zeng[1], Nick Raymond[1], Victor Zeng[1], Yoonho Chung[5], George He[5], Daqiang Sun[6], Theo G.M. van Erp[7], Jean Addington[8], Carrie E. Bearden[6], Kristin Cadenhead[9], Barbara Cornblatt[10], Daniel H. Mathalon[11], Thomas McGlashan[12], Clark Jeffries[13], William Stone[1,14], Ming Tsuang[9], Elaine Walker[5], Scott W. Woods[12], Tyrone D. Cannon[5,12], Diana Perkins[13,15], Matcheri Keshavan[1,14], Paulo Lizano[1,14,16]

[1]Department of Psychiatry, Beth Israel Deaconess Medical Center, Boston, MA, USA
[2]German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany
[3]A.A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Boston, MA, USA
[4]Department of Radiology, Harvard Medical School, Boston, MA, USA
[5]Department of Psychology, Yale University, New Haven, CT, USA
[6]Semel Institute for Neuroscience and Human Behavior and Department of Psychology, UCLA, Los Angeles, CA, USA
[7]Clinical Translational Neuroscience Laboratory, Department of Psychiatry and Human Behavior, UC Irvine, Irvine, CA, USA
[8]Hotchkins Brain Institute, Department of Psychiatry, University of Calgary, Calgary, AB, Canada
[9]Department of Psychiatry, UCSD, San Diego, CA, USA
[10]Department of Psychiatry, Zucker Hillside Hospital, Queens, NY, USA
[11]Department of Psychiatry, UCSF, San Francisco, CA, USA
[12]Department of Psychiatry, Yale University, New Haven, CT, USA
[13]Renaissance Computing Institute, University of North Carolina, Chapel Hill, NC, USA
[14]Department of Psychiatry, Harvard Medical School, Boston, MA, USA
[15]Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA
[16]Division of Translational Neuroscience, Beth Israel Deaconess Medical Center, Boston, MA, USA

Corresponding Author:
Paulo Lizano MD, PhD
75 Fenwood Road Room 612, Boston, MA 02115
Email: plizano@bidmc.harvard.edu
Phone: 617.754.1227

Key words: choroid plexus, lateral ventricle, clinical high risk for psychosis, inflammation

Key Points: 83 words
Abstract: 350 words
Text: 2999 words
Tables: 1
Figures: 3

Abstract

Background: Choroid plexus (ChP) enlargement has been described in first-episode psychosis and psychosis spectrum disorders, but whether ChP enlargement occurs before disease onset is unknown. This study investigated whether ChP volume is enlarged in individuals with clinical high-risk (CHR) for psychosis and whether these changes are related to clinical, cortical, and plasma analyte measures.

Methods: Clinical and neuroimaging data from the North American Prodrome Longitudinal Study 2 (NAPLS2) was used for analysis. A total of 509 participants (169 controls, 340 CHR) were recruited across 8 sites. Conversion status was determined until 2-years of follow-up, with 36 patients developing psychosis. The lateral ventricle ChP was manually segmented from all available baseline brain scans. A subsample of 84 participants (31 controls, 53 CHR) had plasma analyte and neuroimaging data.

Results: Compared to controls, CHR overall ($d=0.22$, $p=0.017$) and converters ($d=0.21$, $p=0.041$) demonstrated higher ChP volumes, but not nonconverters. In CHR, greater ChP volume correlated with lower cortical ($r=-0.22$, $p<0.001$) and subcortical gray matter volume ($r=-0.21$, $p<0.001$), total white matter volume ($r=-0.28$, $p<0.001$), and larger lateral ventricle volume ($r=0.63$, $p<0.001$). In CHR, greater ChP volume, but not lateral ventricle volume, correlated with higher C3 ($r=0.39$, $p=0.005$) and TSH ($r=0.31$, $p=0.026$), and lower MMP7 ($r=-0.30$, $p=0.032$) and UMOD ($r=-0.33$, $p=0.019$).

Conclusions and Relevance: CHR and converters to psychosis demonstrated significantly larger ChP volumes compared to controls. ChP enlargement was associated with cortical volume reduction and increased peripheral inflammatory markers. These findings suggest that the ChP may be a key biomarker in psychosis disease onset and conversion.

Main Text

Introduction:

The choroid plexus (ChP) projects into the four cerebral ventricles and is the principal source of cerebrospinal fluid (CSF). Together, the ChP and CSF play an important role in brain physiology[1,2], buoyancy[3], and homeostasis by providing physical, enzymatic, and immunological barriers to the brain[4]. Neuroimaging studies have observed ChP changes with aging[5–7] and across various neurodevelopmental[8–10] and brain disorders[11–13], suggesting that the ChP may play a key role in brain disorders. Despite growing evidence implicating the ChP in neurodevelopment and brain disorders, it has not been the focus of commonly used neuroimaging tools, which causes it to be poorly segmented, mislabeled, and incorrectly quantified[14]. Therefore, there is a critical need to elucidate the role of the ChP in brain disorders using more accurate segmentation of the ChP.

The earliest observations of ChP morphological changes in schizophrenia and bipolar disorders occurred in the early 20[th] century, where cystic formations, fibrosis, lipid deposits, and hypersecretory phenotypes were found in ChP epithelial cells, as well as endothelial cell degradation[15–17]. We previously reported on ChP enlargement in a large sample of patients with schizophrenia, schizoaffective, and psychotic bipolar disorder compared to healthy controls[18]. Larger ChP volume was associated with worse overall cognition, but no correlations with clinical measures were identified. A link between ChP volume and brain structure was established, with greater ChP volume relating to smaller gray matter and subcortical volume, larger ventricular volume, and lower white matter microstructure. Enlarged ChP volume correlated with higher peripheral levels of interleukin-6, suggesting that inflammation may play a role in ChP structural changes observed in psychosis and inflammation-meditated CSF hypersecretion[19,20]. These findings were replicated in a study in first-episode schizophrenia, where the greater ChP volume observed also correlated with higher allostatic load (indexed by subclinical cardiovascular,

metabolic, neuroendocrine, and immune markers)[21]. Another study reported larger ChP volume in schizophrenia patients with orofacial tardive dyskinesia and identified associations with elevated N-methyl-D-aspartate receptor antibody levels[22]. Taken together, these results suggest that the ChP may be an important structure in mechanisms underlying psychosis pathology. However, no studies to date have investigated ChP volumes in a population at clinical high risk (CHR) for psychosis to determine whether ChP enlargement is associated with future psychosis risk.

The results from these neuroimaging studies are promising; however, the use of FreeSurfer for ChP segmentation limits their interpretation, primarily due to poor ChP segmentation by FreeSurfer[14]. Recently, three tools have been developed to provide better ChP parcellation: Gaussian Mixture Model (GMM; Alzheimer's)[23], an optimized 3D U-Net[24] (aging healthy individuals), and an axial multi-layer perceptron (Axial-MLP 8)[25]. While robust, each technique only provides adequate segmentation performance, reflected in average Dice Coefficients (DC) of 64 for GMM, 71 for Axial-MLP8, and 73 for 3D U-Net. Therefore, in this study, we aim to analyze ChP volume in individuals at CHR for psychosis using manually segmented lateral ventricle ChP from the North American Prodrome Longitudinal Study 2 (NAPLS2) and will compare these manually-segmented values to two techniques previously used to study ChP volume in neuropsychiatric disorders: FreeSurfer 6.0 and GMM. We further aim to understand whether greater ChP volume is related to clinical, neuroimaging, and plasma analyte measures. We hypothesized that the CHR group will demonstrate ChP enlargement compared to healthy controls (HC) and that converters to psychosis will have higher volumes. Finally, we hypothesized that greater ChP volumes in CHR would be associated with worse cognition, global brain volume reduction, and higher plasma analyte levels.

Methods:

Participants

Participants were part of NAPLS2, a prospective multisite study analyzing psychosis predictors in individuals 12-35 years of age meeting Criteria of Psychosis Risk Syndromes[26]. Institutional Review Board approval was obtained from all sites[27]. Participants included 521 individuals (175 HC and 346 CHR) who had completed T1 MRI scans. Each participant provided written consent and parents/guardians provided consent for minors with minors providing assent. Exclusion criteria included current or lifetime Axis I psychotic disorder, IQ<70, past history of a central nervous system disorder, substance dependence within the past 6 months, and prodromal symptoms caused by an Axis I disorders. Non-psychotic DSM-IV disorders were not exclusionary (e.g. substance abuse disorder, anxiety disorders, or Axis II disorders) if they did not account for psychosis-risk symptoms. Exclusionary criteria for HC included meeting psychosis-risk syndrome criteria, any current or past psychotic disorder or Cluster A personality disorder, first-degree family history of any psychotic disorder or any disorder involving psychotic symptoms, and currently using psychotropic medications. Symptoms were assessed using the Scale of Psychosis-risk Symptoms (SOPS)[27,28]. The Measurement and Treatment Research to Improve Cognition in Schizophrenia (MATRICS)[29–31] and Global Assessment of Functioning (GAF)[32] measures were gathered to evaluate cognition and functioning, respectively. Social cognition was tested using The Awareness of Social Inference Test (TASIT)[33]. A questionnaire documenting trauma from before the age of 16 was collected. Conversion status was determined at 12-month-follow-up or at the point of conversion to psychosis for either a psychotic illness or bipolar disorder (bipolar 1/2, cyclothymic, or not-otherwise-specified). Substance use status was categorized by alcohol, amphetamine, cannabis, cocaine, hallucinogen, or polysubstance use. Illness duration categories (short or long) were determined by median split (30.9 weeks). Antipsychotic status and average daily chlorpromazine (CPZ) equivalents were collected[34]. One HC was dropped due to conversion to psychosis. Lastly, 5 participants were dropped due to missing diagnostic information.

MRI data

Subjects had baseline MRI scans that were free of visually detectable artifacts. A detailed

explanation of scanner parameters and site harmonization was detailed previously[35]. Five sites

used Siemens 3T with a 12-channel head coil (Emory, Harvard, Yale, University of California

Los Angeles, and University of North Carolina) and 3 used GE 3T with an 8-channel head coil

(Zucker Hillside, University of California San Diego, and University of Calgary). Images were

acquired in the sagittal plane with a 1x1x1.2mm resolution. Siemens devices used an MPRAGE

sequence with a 256x240x176mm field of view, TR/TE/T1=2300/2.91/900ms, and a 9-degree

flip angle. GE scanners used an IR-SPGR sequence with a 26cm field of view,

TR/TE/TI=7.0/minimum full/400ms, and an 8-degree flip angle.

Choroid plexus segmentation

T_1-MPRAGE images were processed through FreeSurfer v6.0[36]. Upon visual inspection,

FreeSurfer lateral ventricle ChP maps were found to segment ~50% of the structure (Figure

1A,D,G,J). Thus, a Gaussian Mixture Model (GMM), previously developed for ChP analysis in

Alzheimer's disease, was used to segment ChP[23]. However, GMM ChP parcellations were

found to be inaccurate (Figure 1B,E,H,K) and we decided to manually-segment the ChP for all

scans. 3D Slicer v4.11 was used for manual segmentation by 8 trained raters (IA, DB, SG, RH,

SP, NR, AZ) in a first pass. In the second pass, brains were given to a new rater for further

checking and editing, and a final evaluation was done from an expert rater (DB) to ensure

cohesion of manual segmentation across all raters. DB was trained by PL, a senior

neuropsychiatrist with 8 years of clinical and image processing experience. The trigonum

collaterale was used as a reliable starting point for segmentation. Manual segmentation started

in the axial>sagittal>coronal views to ensure the entirety of the ChP was captured and that

voxels from surrounding brain structures (e.g. fornix, hippocampus, corpus collosum, and

thalamus) were excluded. A 3D inspection was performed at the final stage to ensure proper

overall segmentation (Figure 1A-C). Third and fourth ventricle ChP were excluded to better compare our segmentation results with FreeSurfer and GMM, both of which only segment lateral ventricle ChP. ChP volume measures were extracted using 'fslstats' (FMRIB Software Library, v6.0). FreeSurfer was used to extract intracranial volume (ICV), lateral ventricle volume (LVV), total and subcortical gray matter volumes (GMV), and total white matter volume (WMV). Cortical measures greater than 4 standard deviations (SD) were removed (N=8) and those between 3-4 SD (N=21) were winsorized to the 3^{rd} SD. Manual segmentation was considered the gold standard with respect to FreeSurfer 6.0 and GMM segmentations of the lateral ventricle ChP.

Plasma analyte collection

84 participants (33 HC and 51 CHR) had both MRI and plasma analyte data. A comprehensive detail of blood collection/assay has been provided previously[37]. In brief, blood was collected at baseline, processed within 120min and stored at -80°C. 185 analytes were immunoassayed blind to the subject's clinical status. 44 analytes were previously excluded due to medication or substance use confounds[37]. 13 analytes were not detected in >25% of subjects and were excluded (Supplementary Table 1). CHR vs HC comparisons were performed on the remaining 185 analytes and significant analytes were used for subsequent analyses.

Statistical Analysis

Sociodemographic variables were compared between groups via Chi-square or analysis of variance (ANOVA) tests for categorical and continuous data, respectively. ANOVAs were performed to test the moderating effects of age, sex, race, site, ICV, and total GMV on manually-segmented ChP volume. Significant moderators (p<0.05) were used as covariates in ChP analysis and included age, sex, race, and site (Supplementary Table 2). ICV was used as an additional covariate for ChP measures. DC[38] and absolute intraclass correlation[39] (ICC) were

used to measure the accuracy and reliability of ChP segmentation between FreeSurfer, GMM, and manual segmentation. DC is a measure of the number of overlapping voxels segmented as ChP in two masks with values ranging from 0-100, where 0 designates no overlap and 100 reflects total overlap. Absolute ICC is a measure of agreement between two measurements and ranges from 0 to 1 (1=high reliability). General linear models, controlling for confounding variables, were used for group contrasts between CHR and HC. Post hoc comparisons examining conversion (converters vs non-converters), antipsychotic, duration (short vs long), and substance use status were performed. Effect sizes were calculated using Cohen's d. To analyze whether ChP volume results were impacted by LVV or GMV, ChP group contrasts were performed using 1) LVV and 2) LVV and GMV in place of ICV as a covariate. Partial Spearman correlations were used to analyze associations between ChP volume and clinical (n= 10), brain (n= 4), and analyte measures (n=15), and calculated p-values were corrected for multiple comparisons using the false discovery rate (FDR) method (q-values)[40]. All statistical analyses were performed using R statistical software (version 3.5.1).

Results:

Demographics

The final sample included 340 CHR and 169 HC (Table 1). HC and CHR demonstrated no race differences, but differed for age *(p=0.003)*, sex *(p=0.025)*, and site *(p=0.006)*. CHR had higher rates of substance use disorder (*p=0.004*), SOPS (total, positive, negative, disorganization) and trauma scores *(p<0.001)*. CHR had lower GAF, MATRICS, and TASIT scores *(p<0.001)*. 105 (30%) CHR were on a first- or second-generation antipsychotic, with an average daily chlorpromazine dosage of 187.8mg. In CHR with available data on conversion status (n=277), 36 (13%) converted to psychosis.

Accuracy and reliability of ChP segmentation

Compared to manual segmentation, FreeSurfer *(DC=29; left ICC=0.41, p<0.001; right ICC=0.44, p<0.001; bilateral ICC=0.44, p<0.001;* Figure 1N,M) and GMM *(DC=67; left ICC=0.45, p<0.001; right DC=0.51, p<0.001; bilateral ICC=0.53, p<0.001)* demonstrated poor accuracy and reliability. Compared to FreeSurfer, GMM demonstrated significantly higher accuracy *(ΔDC=0.37, p<0.001)*. Thus, manually segmented ChP volumes were used for primary analyses.

ChP differences

CHR demonstrated higher ChP volume compared to HC *(left: d=0.22, p=0.02, q=0.02; right: d=0.22, p=0.02, q=0.02; bilateral: d=0.22, p=0.017, q=0.02;* Figure 2A, Supplementary Table 3). This effect did not hold after controlling for LVV or LVV+GMV (Supplementary Table 3). For group differences using FreeSurfer or GMM ChP measures see Supplementary Table 3. In CHR, post hoc comparisons were performed using conversion (36 converters, 231 non-converters), antipsychotic (105 antipsychotics, 241 no antipsychotics), illness duration (155 short, 156 long), and substance use statuses (24 substance use, 322 no substance use; Supplementary Table 4). Compared to HC, converters had higher ChP volumes *(d=0.15, p=0.021*, Figure 2B; Supplementary Table 5). CHR not on antipsychotics demonstrated larger ChP compared to HC *(d=0.21, p=0.041)*, while CHR on antipsychotics had larger ChP compared to those not on antipsychotics *(d=0.041, p=0.035;* Figure 2C). CHR with longer illness durations had larger ChP compared to HC *(d=0.18, p=0.008;* Figure 2D). Lastly, CHR both with and without a history of substance abuse had significantly higher ChP volumes compared to HC *(no-substance use d=0.20, p=0.035; substance use: d=0.56, p=0.015;* Figure 2E).

Clinical correlations

No significant correlations were observed between ChP volume and SOPS, global functioning, MATRICS, trauma, or social cognitive scores in the overall, CHR or HC groups (Supplementary Table 6). In CHR, no associations were noted between ChP volume and illness duration or CPZ equivalence.

Neuroimaging correlations

In CHR, larger ChP was associated with lower total GMV *(r=-0.22, p<0.001, q<0.001)*, subcortical GMV *(d=-0.21, p<0.001, q<0.001)*, and WMV *(r=-0.28, p<0.001, q<0.001),* as well as with higher LVV *(r=0.63, p<0.001, q<0.001;* Figure 3, Supplementary Table 7). HC demonstrated significant correlations between higher ChP and lower WMV *(r=-0.27, p<0.001, q<0.001)* and higher LVV *(r=0.66, p<0.001, q<0.001).*

Exploratory plasma analyte correlations

There were no sociodemographic differences between HC and CHR in the subset of participants with both MRI and analyte data (Supplementary Table 8). In CHR, greater ChP volume was associated with higher levels of thyroid stimulating hormone (TSH; *r=0.39, p=0.004*) and lower levels of alpha-2 macroglobulin (A2M; *r=-0.29, p=0.037*), lymphocyte secreted protein I-309 (CCL1; *r=-0.42, p=0.002*), myeloperoxidase (MPO; *r=-0.39, p=0.006*), and tissue inhibitor of metalloproteinases-1 (TIMP-1; *r=-0.29, p=0.041;* Figure 4). Non-converters demonstrated negative relationships between ChP volume and peripheral levels of A2M *(r=-0.43, p=0.01)*, CCL1 *(r=-0.43, p=0.01)*, MPO *(r=-0.43, p=0.01)*, TIMP-1 *(r=-0.43, p=0.01)*, and vascular cell adhesion molecule-1 (VCAM-1; *r=-0.43, p=0.01*). Interestingly, converters showed strong correlations between enlarged ChP volume and greater peripheral AXL receptor tyrosine kinase (AXL; *r=-0.43, p=0.01*), creatine kinase-MB (CKM; *r=-0.43, p=0.01*), epidermal growth factor receptor (EGFR; *r=-0.43, p=0.01*), glutathione S-transferase alpha 1 (GSTA1; *r=-0.43, p=0.01*), intracellular adhesion molecule-1 (ICAM-1; *r=-0.43, p=0.01*), myoglobin (MB; *r=-0.43, p=0.01*),

and E-selectin (SELE; *r=-0.43, p=0.01*) levels. In controls, ChP volume was associated with

higher placental growth factor (PGF; *r=0.42, p=0.048*) and lower endothelin-1 (EDN1; *r=-0.55,*

p=0.046), interleukin-3 (IL-3; *r=-0.36, p=0.041*) and interleukin-4 (IL-4; *r=-0.59, p=0.004*).

Discussion:

Our study demonstrated that patients with CHR, who converted to psychosis, with longer illness

durations, or on antipsychotics had larger ChP volume compared to HC. While no clinical

correlations were observed in CHR, enlarged ChP was associated with lower cortical and

subcortical volumes, and larger LVV. Larger ChP was also associated with elevated

complement C3 and TSH levels and lower MMP7 and UMOD levels. FreeSurfer and GMM

demonstrated poor accuracy/reliability for ChP segmentation compared to manual

segmentation. Together, these results suggest that the ChP may be an important risk and

treatment biomarker for CHR individuals.

These results extend upon prior findings of lateral ventricle ChP enlargement in first-episode

schizophrenia[21] and chronic psychosis[18] using a more accurate method for ChP segmentation.

ChP enlargement has been reported in other neuropsychiatric disorders, such as Alzheimer's[12],

multiple sclerosis (MS)[41,42], neuromyelitis optica[41], and major depressive disorder[13]. Normal size

criterion for the ChP does not exist to compare our findings to, but one study in children[5] (0-16

years) and two studies in adults (18-94 years)[6,7] have attempted to do this. However, these

studies have been limited by small sample sizes, measurement types (thickness vs volume),

neuroimaging modality (MRI vs ultrasound), narrow age categories (mostly older adults), and

use of less accurate ChP segmentation algorithms, which make it challenging to understand the

role of ChP morphology in psychotic disorders.

What ChP enlargement might mean for patients with psychosis has yet to be determined, but

recent studies have shed light on this matter. For example, Lizano et al. (2019)[18] observed

relationships between ChP enlargement and larger LVV and lower total GMV in psychotic

disorders. In MS, ChP enlargement was associated with higher LVV, greater WM lesions, lower

brain volume, and normal-appearing WMV[42]. In depression, researchers found that ChP

enlargement correlated with markers of central inflammation, namely greater mitochondrial

translocator protein (TSPO) expression within the anterior cingulate, prefrontal, and insular

cortices[13]. ChP enlargement was also associated with higher neuroinflammation in MS,

reflected by greater [18]F-DPA-714 binding in the thalamus[42]. The ChP of MS patients

demonstrated greater TSPO expression compared to controls, suggesting an increased

presence of immune cells within the ChP epithelium. Similarly, greater gadolinium enhancement

of lateral and 4[th] ventricle ChP in MS was identified suggesting a potential increase in ChP

stromal capillary permeability[41]. These studies suggest that ChP enlargement is associated with

inflammatory and microvascular changes within the ChP itself, as well as other parenchymal

areas.

An additional gap in the literature pertains to the effects of circulating peripheral analytes on

ChP morphology. Previous work has demonstrated associations between ChP enlargement and

higher peripheral IL6[18] in those with chronic psychosis and with greater allostatic load in first-

episode SZ[21]. We could not examine IL6 in our exploratory correlational analysis since it did not

survive our quality control. However, we found that enlarged ChP in CHR correlated with higher

TSH and lower A2M, CCL1, MPO and TIMP-1 levels. Patients with first-episode SZ have been

found to have significantly lower peripheral TSH levels compared to patients with multi-episode

SZ, who demonstrated significantly higher TSH levels albeit with a significant degree of

heterogeneity[43]. In our sample, CHR on antipsychotics had significantly higher TSH levels

(mean TSH=1.95, SD=0.78) compared to CHR not on antipsychotics (mean TSH=1.41, SD=0.7;

p=0.02, d=0.75) which may reflect a potential hypothyroidism caused by antipsychotic usage[44].

Further studies are necessary to better understand the relationship between TSH levels and

antipsychotic usage as well as between elevated TSH levels and enlarged ChP. Reductions in

serum A2M have been observed in individuals at high-risk for SZ as well as in patients with SZ[45,46]. A2M is a broad protease inhibitor as well as a cytokine transporter and can be thought of as able to inhibit inflammatory cytokines and cascades[47], which supports its observed inverse association with ChP volume in CHR. In contrast, CCL1 and MPO are analytes associated with pro-inflammatory processes and are secreted by immune cells and expressed by neutrophil granulocytes, respectively. We expected both to have positive relationships with ChP volume in CHR, linking peripheral and central inflammation. In mice, Ito et al. (2019)[48] linked CCL1, intraventricular not serum, to regulatory T-cell infiltration of the brain after stroke. Elevated serum MPO has been described in multiple sclerosis[49], depression[50], and schizophrenia[51]. Interestingly, Baranyi et al. (2022)[52] described lower MPO levels in patients with acute myocardial infarction (AMI) induced depression compared to AMI patients that did not develop depression. This suggests that there may be an interaction occurring between inflammatory and neuropsychiatric states with regards to MPO serum levels, but further research is needed. Lastly, TIMP-1 acts as an inhibitor to matrix metalloproteinases and is key in maintaining the composition of the extracellular matrix. While elevated serum TIMP-1 and MMP9 have been described in SZ[53], TIMP-1 has also been shown to play a protective role in blood-brain-barrier (BBB) integrity after damage via a mechanism independent of matrix metalloproteinase activity[54]. More research studying the mechanisms of TIMP-1 in SZ are needed to better elucidate the role of TIMP-1 in disease and protective states.

In conclusion, our results suggest that the ChP is a biomarker of particular interest for neurodevelopment and CHR for psychosis. Strengths of this study include the use of manually-segmented ChP volume in the largest sample size of CHR patients. Limitations include 1) a sample size for examining ChP to peripheral analyte relationships, 2) lack of 3rd/4th ventricle ChP segmentation given their proximity to important brain areas (hippocampus and cerebellum), and 3) cross-sectional neuroimaging design. Future studies should focus on these remaining gaps to would allow for a more compressive understanding of ChP morphology in psychosis.

Acknowledgements:

The authors would like to acknowledge Ehsan Tadayon for his help regarding choroid plexus segmentation techniques and for use of his Gaussian Mixture Model and the late Dr. Larry J. Seidman for his contribution to the field of CHR for psychosis research. The authors thank the participants at each recruitment site who took part in this study.

References:

1. Kaiser K, Bryja V. Choroid Plexus: The Orchestrator of Long-Range Signalling Within the CNS. *Int J Mol Sci*. 2020;21(13):4760. doi:10.3390/ijms21134760

2. Damkier HH, Brown PD, Praetorius J. Cerebrospinal Fluid Secretion by the Choroid Plexus. *Physiol Rev*. 2013;93:46.

3. Fame RM, Cortés☐Campos C, Sive HL. Brain Ventricular System and Cerebrospinal Fluid Development and Function: Light at the End of the Tube: A Primer with Latest Insights. *BioEssays*. 2020;42(3):1900186. doi:10.1002/bies.201900186

4. Liddelow SA. Development of the choroid plexus and blood-CSF barrier. *Front Neurosci*. 2015;9. doi:10.3389/fnins.2015.00032

5. Madhukar M, Choudhary AK, Boal DK, Dias MS, Iantosca MR. Choroid plexus: normal size criteria on neuroimaging. *Surg Radiol Anat*. 2012;34(10):887-895. doi:10.1007/s00276-012-0980-5

6. İmamoğlu H, Erkoç MF, Şalk İ, Okur A, Aydemir EI. Evaluation of the normal choroid plexus size in adults with Magnetic Resonance Imaging. *Cumhur Med J*. 2013;35(1):51-54. doi:10.7197/1305-0028.1661

7. Alisch JSR, Kiely M, Triebswetter C, et al. Characterization of Age-Related Differences in the Human Choroid Plexus Volume, Microstructural Integrity, and Blood Perfusion Using Multiparameter Magnetic Resonance Imaging. *Front Aging Neurosci*. 2021;13:734992. doi:10.3389/fnagi.2021.734992

8. Levman J, Vasung L, MacDonald P, et al. Regional volumetric abnormalities in pediatric autism revealed by structural magnetic resonance imaging. *Int J Dev Neurosci*. 2018;71(1):34-45. doi:10.1016/j.ijdevneu.2018.08.001

9. Lin YJ, Chiu NC, Chen HJ, Huang JY, Ho CS. Cranial ultrasonographic screening findings among healthy neonates and their association with neurodevelopmental outcomes. *Pediatr Neonatol*. 2021;62(2):158-164. doi:10.1016/j.pedneo.2020.10.012

10. Chang H, Tsai CM, Hou CY, Tseng SH, Lee JC, Tsai ML. Multiple subependymal pseudocysts in neonates play a role in later attention deficit hyperactivity and autistic spectrum disorder. *J Formos Med Assoc*. 2019;118(3):692-699. doi:10.1016/j.jfma.2018.08.007

11. Egorova N, Gottlieb E, Khlif MS, Spratt NJ, Brodtmann A. Choroid plexus volume after stroke. *Int J Stroke*. 2019;14(9):923-930. doi:10.1177/1747493019851277

12. Tadayon E, Pascual-Leone A, Press D, Santarnecchi E. Choroid plexus volume is associated with levels of CSF proteins: relevance for Alzheimer's and Parkinson's disease. *Neurobiol Aging*. 2020;89:108-117. doi:10.1016/j.neurobiolaging.2020.01.005

13. Althubaity N, Schubert J, Martins D, et al. Choroid plexus enlargement is associated with neuroinflammation and reduction of blood brain barrier permeability in depression. *NeuroImage Clin*. 2022;33:102926. doi:10.1016/j.nicl.2021.102926

14. Bannai D, Lutz O, Lizano P. Neuroimaging considerations when investigating choroid plexus morphology in idiopathic psychosis. *Schizophr Res*. 2020;224:19-21. doi:10.1016/j.schres.2020.07.013

15. Kitabayashi S. The choroid plexuses in organic disease of the brain and in schizophrenia. *J Nerv Ment Disord*. 1922;56:21-26.

16. Morowoka T. The Microscopical Examination of the Choroid Plexus in Generat Paralysis of the Insane, and other forms of Mental Disease. *J Nerv Ment Disord*. Published online 1921:14.

17. Taft AE. A NOTE ON THE PATHOLOGY OF THE CHOROID PLEXUS IN GENERAL PARALYSIS. *Arch Neurol Psychiatry*. 1922;7(2):177. doi:10.1001/archneurpsyc.1922.02190080026003

18. Lizano P, Lutz O, Ling G, et al. Association of Choroid Plexus Enlargement With Cognitive, Inflammatory, and Structural Phenotypes Across the Psychosis Spectrum. *Am J Psychiatry*. 2019;176(7):564-572. doi:10.1176/appi.ajp.2019.18070825

19. Zhang Z, Tan Q, Guo P, Jia Z, Liu X, Feng H. *NLRP3 Inflammasome-Mediated Cerebrospinal Fluid Hypersecretion in Choroid Plexus Contributes to Hydrocephalus after Hemorrhage*.; 2020. doi:10.21203/rs.3.rs-109549/v1

20. Karimy JK, Zhang J, Kurland DB, et al. Inflammation-dependent cerebrospinal fluid hypersecretion by the choroid plexus epithelium in posthemorrhagic hydrocephalus. *Nat Med*. 2017;23(8):997-1003. doi:10.1038/nm.4361

21. Zhou YF, Huang JC, Zhang P, et al. Choroid Plexus Enlargement and Allostatic Load in Schizophrenia. *Schizophr Bull*. 2020;46(3):722-731. doi:10.1093/schbul/sbz100

22. Li N, Huang J, Zhang P, et al. N-methyl-D-aspartate receptor antibody and the choroid plexus in schizophrenia patients with tardive dyskinesia. *J Psychiatr Res*. 2021;142:290-298. doi:10.1016/j.jpsychires.2021.08.010

23. Tadayon E, Moret B, Sprugnoli G, Monti L, Pascual-Leone A, Santarnecchi E. Improving Choroid Plexus Segmentation in the Healthy and Diseased Brain: Relevance for Tau-PET Imaging in Dementia. *J Alzheimers Dis*. 2020;74(4):1057-1068. doi:10.3233/JAD-190706

24. Zhao L, Feng X, Meyer CH, Alsop DC. Choroid Plexus Segmentation Using Optimized 3D U-Net. In: *2020 IEEE 17th International Symposium on Biomedical Imaging (ISBI)*. IEEE; 2020:381-384. doi:10.1109/ISBI45749.2020.9098443

25. Schmidt-Mengin M, Ricigliano VAG, Bodini B, et al. Axial multi-layer perceptron architecture for automatic segmentation of choroid plexus in multiple sclerosis. *ArXiv210903778 Cs Eess Q-Bio*. Published online November 21, 2021. Accessed January 12, 2022. http://arxiv.org/abs/2109.03778

26. Addington J, Cadenhead KS, Cornblatt BA, et al. North American Prodrome Longitudinal Study (NAPLS 2): Overview and recruitment. *Schizophr Res*. 2012;142(1-3):77-82. doi:10.1016/j.schres.2012.09.012

27. Miller TJ, McGlashan TH, Rosen JL, et al. Prospective Diagnosis of the Initial Prodrome for Schizophrenia Based on the Structured Interview for Prodromal Syndromes: Preliminary Evidence of Interrater Reliability and Predictive Validity. *Am J Psychiatry*. 2002;159(5):863-865. doi:10.1176/appi.ajp.159.5.863

28. McGlashan T, Walsh B, Woods S. *The Psychosis-Risk Syndrome: Handbook for Diagnosis and Follow-Up*. Oxford University Press; 2010.

29. Green MF, Nuechterlein KH, Kern RS, et al. Functional Co-Primary Measures for Clinical Trials in Schizophrenia: Results From the MATRICS Psychometric and Standardization Study. *Am J Psychiatry*. 2008;165(2):221-228. doi:10.1176/appi.ajp.2007.07010089

30. Kern RS, Nuechterlein KH, Green MF, et al. The MATRICS Consensus Cognitive Battery, Part 2: Co-Norming and Standardization. *Am J Psychiatry*. 2008;165(2):214-220. doi:10.1176/appi.ajp.2007.07010043

31. Nuechterlein KH, Green MF, Kern RS, et al. The MATRICS Consensus Cognitive Battery, Part 1: Test Selection, Reliability, and Validity. *Am J Psychiatry*. 2008;165(2):203-213. doi:10.1176/appi.ajp.2007.07010042

32. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. American Psychiatric Association; 2000.

33. McDonald S, Flanagan S, Rollins J, Kinch J. TASIT: A New Clinical Tool for Assessing Social Perception After Traumatic Brain Injury. *J Head Trauma Rehabil*. 2003;18(3):219-238. doi:10.1097/00001199-200305000-00001

34. Cao H, Chung Y, McEwen SC, et al. Progressive reconfiguration of resting-state brain networks as psychosis develops: Preliminary results from the North American Prodrome Longitudinal Study (NAPLS) consortium. *Schizophr Res*. 2020;226:30-37. doi:10.1016/j.schres.2019.01.017

35. Cannon TD, Sun F, McEwen SJ, et al. Reliability of neuroanatomical measurements in a multisite longitudinal study of youth at risk for psychosis: Reliability of Multisite, Longitudinal MRI. *Hum Brain Mapp*. 2014;35(5):2424-2434. doi:10.1002/hbm.22338

36. Fischl B. FreeSurfer. *NeuroImage*. 2012;62(2):774-781. doi:10.1016/j.neuroimage.2012.01.021

37. Perkins DO, Jeffries CD, Addington J, et al. Towards a Psychosis Risk Blood Diagnostic for Persons Experiencing High-Risk Symptoms: Preliminary Results From the NAPLS Project. *Schizophr Bull*. 2015;41(2):419-428. doi:10.1093/schbul/sbu099

38. Dice LR. Measures of the Amount of Ecologic Association Between Species. *Ecology*. 1945;26(3):297-302. doi:10.2307/1932409

39. McGraw KO, Wong SP. Forming Inferences About Some Intraclass Correlation Coefficients. *Psychol Methods*. 1996;1(1):30-46.

40. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol*. 1995;57(1):289-300.

41. Kim H, Lim YM, Kim G, et al. Choroid plexus changes on magnetic resonance imaging in multiple sclerosis and neuromyelitis optica spectrum disorder. *J Neurol Sci*. 2020;415:116904. doi:10.1016/j.jns.2020.116904

42. Ricigliano VA, Morena E, Colombi A, et al. Choroid Plexus Enlargement in Inflammatory Multiple Sclerosis: 3.0-T MRI and Translocator Protein PET Evaluation. *Radiology*. 2021;301(1):166-177. doi:10.1148/radiol.2021204426

43. Misiak B, Stramecki F, Stańczykiewicz B, Frydecka D, Lubeiro A. Vascular endothelial growth factor in patients with schizophrenia: A systematic review and meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry*. 2018;86:24-29. doi:10.1016/j.pnpbp.2018.05.005

44. Khalil RB, Richa S. Thyroid Adverse Effects of Psychotropic Drugs: A Review. *Clin Neuropharmacol*. 2011;34(6):8.

45. Heurich M, Föcking M, Mongan D, Cagney G, Cotter DR. Dysregulation of complement and coagulation pathways: emerging mechanisms in the development of psychosis. *Mol Psychiatry*. 2022;27(1):127-140. doi:10.1038/s41380-021-01197-9

46. Hayes LN, Severance EG, Leek JT, et al. Inflammatory Molecular Signature Associated With Infectious Agents in Psychosis. *Schizophr Bull*. 2014;40(5):963-972. doi:10.1093/schbul/sbu052

47. Vandooren J, Itoh Y. Alpha-2-Macroglobulin in Inflammation, Immunity and Infections. *Front Immunol*. 2021;12:803244. doi:10.3389/fimmu.2021.803244

48. Ito M, Komai K, Mise-Omata S, et al. Brain regulatory T cells suppress astrogliosis and potentiate neurological recovery. *Nature*. 2019;565(7738):246-250. doi:10.1038/s41586-018-0824-5

49. Strzepa A, Pritchard KA, Dittel BN. Myeloperoxidase: A new player in autoimmunity. *Cell Immunol*. 2017;317:1-8. doi:10.1016/j.cellimm.2017.05.002

50. Vaccarino V, Brennan ML, Miller AH, et al. Association of Major Depressive Disorder with Serum Myeloperoxidase and Other Markers of Inflammation: A Twin Study. *Biol Psychiatry*. 2008;64(6):476-483. doi:10.1016/j.biopsych.2008.04.023

51. Al-Asmari A, Khan MdW. Inflammation and schizophrenia: Alterations in cytokine levels and perturbation in antioxidative defense systems. *Hum Exp Toxicol*. 2014;33(2):115-122. doi:10.1177/0960327113493305

52. Baranyi A, Enko D, Meinitzer A, et al. Myeloperoxidase as a Potential Biomarker of Acute-Myocardial-Infarction-Induced Depression and Suppression of the Innate Immune System. *Antioxidants*. 2022;11(11):2083. doi:10.3390/antiox11112083

53. Domenici E, Willé DR, Tozzi F, et al. Plasma Protein Biomarkers for Depression and Schizophrenia by Multi Analyte Profiling of Case-Control Collections. Domschke K, ed. *PLoS ONE*. 2010;5(2):e9166. doi:10.1371/journal.pone.0009166

54. Tang J, Kang Y, Huang L, Wu L, Peng Y. TIMP1 preserves the blood–brain barrier through interacting with CD63/integrin 1 complex and regulating downstream FAK/RhoA signaling. *Acta Pharm Sin B*. 2020;10(6):987-1003. doi:10.1016/j.apsb.2020.02.015

55. Chen Y, Zhao Z, Lin F, Wang L, Lin Z, Yue W. Associations Between Genotype and Peripheral Complement Proteins in First-Episode Psychosis: Evidences From C3 and C4. *Front Genet*. 2021;12:647246. doi:10.3389/fgene.2021.647246

56. Laskaris L, Zalesky A, Weickert CS, et al. Investigation of peripheral complement factors across stages of psychosis. *Schizophr Res*. 2019;204:30-37. doi:10.1016/j.schres.2018.11.035

57. Mayilyan KR, Weinberger DR, Sim RB. The complement system in schizophrenia. *Drug News Perspect*. 2008;21(4):200. doi:10.1358/dnp.2008.21.4.1213349

58. Balusu S, Van Wonterghem E, De Rycke R, et al. Identification of a novel mechanism of blood–brain communication during peripheral inflammation via choroid plexus-derived extracellular vesicles. *EMBO Mol Med*. 2016;8(10):1162-1183. doi:10.15252/emmm.201606271

59. Vandendriessche C, Balusu S, Van Cauwenberghe C, et al. Importance of extracellular vesicle secretion at the blood–cerebrospinal fluid interface in the pathogenesis of Alzheimer's disease. *Acta Neuropathol Commun*. 2021;9(1):143. doi:10.1186/s40478-021-01245-z

60. Ogaki A, Ikegaya Y, Koyama R. Extracellular Vesicles Taken up by Astrocytes. *Int J Mol Sci*. 2021;22(19):10553. doi:10.3390/ijms221910553

61. Berretta S. Extracellular matrix abnormalities in schizophrenia. *Neuropharmacology*. 2012;62(3):1584-1597. doi:10.1016/j.neuropharm.2011.08.010

62. Frye MA, Nassan M, Jenkins GD, et al. Feasibility of investigating differential proteomic expression in depression: implications for biomarker development in mood disorders. *Transl Psychiatry*. 2015;5(12):e689-e689. doi:10.1038/tp.2015.185

63. Futtrup J, Margolinsky R, Benros ME, et al. Blood-brain barrier pathology in patients with severe mental disorders: a systematic review and meta-analysis of biomarkers in case-control studies. *Brain Behav Immun - Health*. 2020;6:100102. doi:10.1016/j.bbih.2020.100102

64. Omori W, Hattori K, Kajitani N, et al. Increased Matrix Metalloproteinases in Cerebrospinal Fluids of Patients With Major Depressive Disorder and Schizophrenia. *Int J Neuropsychopharmacol*. 2020;23(11):713-720. doi:10.1093/ijnp/pyaa049

65. Castellazzi M, Ligi D, Contaldi E, et al. Multiplex Matrix Metalloproteinases Analysis in the Cerebrospinal Fluid Reveals Potential Specific Patterns in Multiple Sclerosis Patients. *Front Neurol*. 2018;9:1080. doi:10.3389/fneur.2018.01080

66. Nichols P, Urriola J, Miller S, et al. Blood-brain barrier dysfunction significantly correlates with serum matrix metalloproteinase-7 (MMP-7) following traumatic brain injury. *NeuroImage Clin*. 2021;31:102741. doi:10.1016/j.nicl.2021.102741

Figure 1. Schematic showing differences in choroid plexus segmentation between FreeSurfer 6.0 (Figures 1A, 1D, 1G, and 1J), Gaussian Mixture Model (Figures 1B, 1E, 1H, and 1K), and manual segmentation (Figures 1C, 1F, 1I, and 1L). While letters stand for: 'A', anterior; 'P', posterior; 'R', right; 'L', left. Panel M demonstrates the difference in segmentation accuracy, quantified using Dice coefficients. Panel N enumerates intra-class correlation values across the three segmentation techniques.

	FreeSurfer–Manual		GMM–Manual		FreeSurfer–GMM	
	ICC	p	ICC	p	ICC	p
Left	0.41	<0.001	0.45	<0.001	0.34	<0.001
Right	0.44	<0.001	0.41	<0.001	0.44	<0.001
Bilateral	0.41	<0.001	0.53	<0.001	0.44	<0.001

GMM, Gaussian mixture model; ICC, intra-class correlation. Note that choroid plexus measures from each segmentation technique were used to calculate the ICC. Individually significant between segmentation techniques are denoted in bold.

Figure 2. Panel A denotes diagnostic group differences between controls and CHR. Choroid plexus volume was corrected for age, sex, race, site number, and intracranial volume. Panels B-E demonstrate subgroup differences between controls and (B) non-converters and converters, (C) individuals on and not on antipsychotics, (D) individuals with short and long illness durations, and (E) individuals with and without a history of substance use. Note: ChP measures are covaried for age, sex, race, site number, and total intracranial volume.

Figure 3. Choroid plexus volume correlations with cortical measures.

Figure 4. Choroid plexus correlations with analyte measures within CHR, non-converters, converters, and HC.

Table 1. Clinical and demographics information.

Variable	HC (n=169)	CHR (n=340)	Test Stat (χ^2,F)	p
Age (years)	20.1 (4.8)	18.9 (4.1)	8.92	0.003
Sex (Female/Male)	87/82	138/202	5	0.025
Race (CA/OT)	93/76	192/147	0.062	0.81
Site Number (1-8)	20/16/26/20/23/16/23/25	38/45/23/37/55/33/79/30	19.6	0.006
Antipsychotic Status (Yes/No)	-	241/105	-	-
CPZ Equivalent	-	187.8 (216.6)	-	-
Converter/Non-Converter	-	36/231	-	-
Substance Use Disorder (Yes/No)	1/168	24/332	8.25	0.004
Duration of Illness (weeks)	-	51.4 (75.7)	-	-
SOPS Total Score	4.29 (5.99)	37.6 (13.0)	1010	<0.001
SOPS Positive	0.94 (1.57)	12.0 (4.2)	1110	<0.001
SOPS Negative	1.43 (2.36)	11.6 (6.9)	416	<0.001
SOPS Disorganization	0.62 (1.15)	5.19 (3.3)	307	<0.001
GAF	84.12 (10.51)	48.9 (11.4)	1140	<0.001
MATRICS Composite	432.5 (65.1)	391.2 (71.0)	36.8	<0.001
Documentation of Trauma	5.56 (2.5)	7.41 (2.7)	56.3	<0.001
Social Cognition (TASIT)	54.7 (5.4)	52.9 (6.0)	10.2	0.002

HC, healthy controls; CA, Caucasian; OT, other; CPZ, chlorpromazine; SOPS, Scale of Prodromal Symptoms; GAF, Global Assessment of Functioning; MATRICS, Measurement and Treatment Research to Improve Cognition in Schizophrenia; BACS, Brief Assessment of Cognition in Schizophrenia; TASIT, The Awareness of Social Inference Test. Note: Sites 1-8 are as follows: University of California Los Angeles, Emory, Harvard, Zucker Hillside Hospital, University of North Carolina, University of California San Diego, University of Calgary, and Yale. Statistical significance is denoted in bold.