Synaptic mechanisms for associative learning in the cerebellar nuclei

Robin Broersen[1,2,†], Catarina Albergaria[3,4,†], Daniela Carulli[5,†], Megan R. Carey[3,#,*], Cathrin B. Canto[1,2,#,*]

and Chris I. De Zeeuw[1,2,*]

[1]Department of Cerebellar Coordination and Cognition, Netherlands Institute for Neuroscience, Royal

Netherlands Academy of Arts and Sciences, Amsterdam, The Netherlands

[2]Department of Neuroscience, Erasmus MC, Rotterdam, The Netherlands

[3]Neuroscience Program, Champalimaud Center for the Unknown, Lisbon, Portugal

[4]Current address: University College London, Sainsbury Wellcome Centre, London, United Kingdom

[5]Laboratory for Neuroregeneration, Netherlands Institute for Neuroscience, Royal Netherlands

Academy of Arts and Sciences, Amsterdam, The Netherlands

[†]Shared first authorship

[#]Shared senior authorship

*Correspondence: c.dezeeuw@erasmusmc.nl, c.canto@nin.knaw.nl,

megan.carey@neuro.fchampalimaud.org

Abstract

Associative learning during delay eyeblink conditioning (EBC) depends on an intact cerebellum. However, the relative contribution of changes in the cerebellar nuclei to learning remains a subject of ongoing debate. In particular, little is known about the changes in synaptic inputs to cerebellar nuclei neurons that take place during EBC and how they shape the membrane potential of these neurons. Here, we probed the ability of these inputs to support associative learning in mice, and investigated structural and cell-physiological changes within the cerebellar nuclei during learning. We find that optogenetic stimulation of mossy fiber afferents to the anterior interposed nucleus (AIP) can substitute for a conditioned stimulus and is sufficient to elicit conditioned responses (CRs) that are adaptively well-timed. Further, EBC induces structural changes in mossy fiber and inhibitory inputs, but not of climbing fiber inputs, and it leads to changes in subthreshold processing of AIP neurons that correlate with conditioned eyelid movements. The changes in synaptic and spiking activity that precede the CRs allow for a decoder to distinguish trials with a CR. Our data reveal how structural and physiological modifications of synaptic inputs to cerebellar nuclei neurons can facilitate learning.

Introduction

The cerebellum has a remarkable ability to learn sensorimotor associations. One well-known example of such associative learning is Pavlovian eyeblink conditioning (EBC), where well-timed conditioned eyelid closures (conditioned response; CR) arise after pairing a neutral conditioned stimulus (CS) with an aversive unconditioned stimulus (US) at a fixed interval, with both stimuli co-terminating in time[1,2]. The main inputs of the cerebellar system that convey the CS and US information have been characterized. The current working hypothesis is that the CS is at least partially conveyed to granule cells in the cerebellar cortex by mossy fiber (MF) axons originating in the pontine nuclei (PN)[3,4]. Granule cells give rise to parallel fibers and innervate Purkinje cells (PCs). Climbing fibers, derived from the inferior olive and powerfully innervating PCs, respond strongly to the US[5]. As incoming MFs and climbing fibers innervate both the cerebellar cortex and the cerebellar nuclei (CN)[6–10], both sites are strong candidates that may contribute to the plasticity mechanisms that underlie associative learning.

In the cerebellar cortex, synaptic weight changes at parallel fiber synapses onto PCs[11–13] and molecular layer interneurons[14,15], potentially in combination with intrinsic mechanisms[16] and structural changes of axonal fibers[17,18], lead predominantly to reductions in PC simple spike activity during the CS-US interval[14,19,20]. This in turn leads to disinhibition of CN neurons[21,22] and ultimately increased spiking activity of premotor and motor neurons downstream in the red nucleus and facial nucleus, respectively[23,24], thereby generating a conditioned eyelid movement.

Within the complex of the CN, the anterior interposed nucleus (AIP) contains most EBC encoding neurons[23,24,27]. Manipulating their activity disrupts both the acquisition and expression of CRs[28–30]. Furthermore, CRs remain partly intact after removal of the cerebellar cortex in fully conditioned mice[31], suggesting that part of the memory trace is stored at the level of the nuclei[32]. However, in contrast to the synaptic plasticity mechanisms in the cerebellar cortex that contribute to eyeblink conditioning, those in the CN remain largely obscure.

Here, we investigated the role of MF synaptic inputs to CN neurons in EBC. We show that using optogenetic stimulation of MF terminals in the AIP as a conditioned stimulus results in surprisingly well-timed eyelid movements over the course of conditioning. Moreover, during EBC, the density of inputs from excitatory MFs, as well as inhibitory inputs, is increased. Finally, in vivo whole-cell recordings of AIP neurons in awake head-fixed mice during EBC reveal the dynamics of their membrane potential (V_m) in relation to associative learning. Presentation of the CS induces predominantly V_m depolarizations in AIP neurons, but both the timing and the directionality of the resulting eyelid movements reverse during conditioning. V_m depolarizations precede the CR in conditioned mice, but follow eyelid openings in naive mice. Using linear discrimination analyses, we

70 show that synaptic as well as spike activity can successfully inform a decoder to distinguish trials with

71 and without a CR. Together, our findings raise the intriguing possibility that there is the capacity for

72 sensorimotor learning and temporal coding within the CN themselves, carried by both structural and

73 physiological changes at the level of individual nuclei neurons.

Results

Optogenetic stimulation of mossy fibers in the cerebellar nuclei is sufficient to induce well-timed EBC

Previous work has shown that optogenetic stimulation of MF terminals within the cerebellar cortex can act as a CS for EBC[25]. We asked whether optogenetic stimulation of MF terminals in the AIP could also act as a CS for EBC (Fig. 1a), and, if so, whether the learned responses could be distinguished from those driven by a cortical MF CS (from here on termed MF-CTX CS). To test this, we placed optic fiber implants just over the AIP of Thy1-ChR2/EYFP transgenic mice expressing Channelrhodopsin-2 (ChR2) in cerebellar MFs (from here on termed MF-ChR2 mice). MF afferents to AIP prominently express ChR2-YFP in these mice (Fig. 1b). We used 350 ms low-power (subthreshold for movement) laser stimulation as a CS, paired with a 50 ms air-puff US delivered to the ipsilateral eye. Similar to learning to a MF-CTX CS[25], mice learned to make CRs in response to optogenetic stimulation of MF terminals in the AIP (from here on termed MF-AIP CS, Fig. 1c).

The conditioned responses to a MF-AIP CS increased gradually during learning and were well-timed in that eyelid closures peaked at the end of the CS-US inter-stimulus interval (ISI) of 300 ms (Fig. 1c,d). Surprisingly, unlike learning to either a sensory CS or the learning driven by MFs in the cerebellar cortex[25], learning to a MF-AIP CS was not enhanced by locomotor activity (main-effect of locomotion: $F_{(1,10)}$ = 2.688, P = 0.132, main-effect of session: $F_{(2.168,\ 21.68)}$ = 61.51, P < 0.001, interaction-effect locomotion x session: $F_{(7,\ 70)}$ = 0.619, P = 0.738, 2-way repeated measure ANOVA), and there was no difference in how quickly animals learned on a slow self-paced vs. fast motorized treadmill (AIP: average onset session 3.3 vs. 3.1, $t_{(10)}$ = -0.447, P = 0.66; CTX: average onset session 2.1 vs. 4.6, $t_{(12)}$ = 5.376, P < 0.001, unpaired *t*-test, two-tailed; Fig. 1d,e). The differential effects of locomotion on EBC for MF-AIP vs. MF-CTX CSs parallel the previous finding that locomotor activity preferentially enhanced eyelid closures evoked through the cerebellar cortex and not the nuclei[25] and they suggest that different mechanisms may govern learning through the two pathways.

We further examined the specificity of optogenetic stimulation of MF terminals in the AIP vs. CTX by combining optogenetic MF CS stimulation in AIP with the use of inhibitory Designer Receptors Exclusively Activated by Designer Drugs (DREADDs)[26,27] expressed in cerebellar granule cells, the targets of MFs in the cerebellar cortex (Fig. 1f). These mice expressed ChR2 in MFs as well as the inhibitory DREADD hM4Di in granule cells, as confirmed by immunohistochemistry (Fig. 1g; mice from here on termed MF-ChR2-gc-DREADD). Administration of the DREADD agonist clozapine-N-oxide (CNO) intraperitoneally to MF-ChR2-gc-DREADD mice prior to each training session had no effect on learning to an optogenetic MF-AIP CS, compared to CNO administration in non-DREADD expressing

MF-ChR2 littermate controls (main-effect of DREADD: $F_{(1,10)}$ = 0.762, P = 0.403, main-effect of session: $F_{(2.348,23.48)}$ = 128.4, P < 0.001, interaction-effect DREADD x session: $F_{(5,50)}$ = 2.267, P = 0.062, 2-way repeated measure ANOVA; Fig. 1h). In contrast, CNO impaired learning to either a MF-CTX CS (Fig. 1i) or a visual CS (Fig. 1j) in MF-ChR2-gc-DREADD mice (MF-CTX CS: main-effect of DREADD: $F_{(1,13)}$ = 9.186, P < 0.001, main-effect of session: $F_{(2.770,32.45)}$ = 31.3, P < 0.001, interaction-effect DREADD x session: $F_{(7,82)}$ = 2.665, P = 0.016; visual CS: main-effect of DREADD: $F_{(1,14)}$ = 10.81, P = 0.005, main-effect of session: $F_{(2.394,27.13)}$ = 18.74, P < 0.001, interaction-effect DREADD x session: $F_{(9,102)}$ = 3.133, P = 0.002, linear mixed models). Together, the differential effects of locomotor activity and granule cell inhibition on learning to a MF-AIP vs. MF-CTX CS confirm the selective stimulation of the MF-CN pathway.

In all of the experiments presented so far, the CRs generated by a MF-AIP CS appeared to be properly timed, with the peak amplitude appearing just before the expected air-puff, similar to a cortical MF CS[25] or a visual CS (Supplementary Fig. 1a-d). However, these experiments do not speak to the capacity to learn to a longer ISI, nor to the ability to adjust the timing of CRs adaptively. To test this, we designed an experiment that allowed us to look at the ability of individual mice to adapt their CRs to changing ISIs (Supplementary Fig. 1e). Interestingly, mice subjected to MF-AIP CS (Supplementary Fig. 1f), as well as those subjected to MF-CTX CS (Supplementary Fig. 1g), could learn to a long (500ms) ISI, and adaptively time their CRs to different ISIs (Supplementary Fig. 1h). Moreover, prolonging the ISI, when training to a MF-AIP CS, resulted in smaller average amplitudes of the CR (Z = 10, P = 0.034, N = 4 mice, Wilcoxon, one-tailed; Supplementary Fig. 1f), similar to previous studies on the impact of longer ISI protocols in both wild types and mutants with cerebellar cortical defects[28].

Conditioning induces structural changes in pontine mossy fiber input to the AIP

Our finding that optogenetic activation of MF inputs in the AIP can serve as a reliable CS suggests that plasticity mechanisms in the AIP can directly contribute to learning and storage of the associative memory trace. We therefore asked what structural changes in inputs to the AIP occur during normal associative learning to a sensory CS. Mice were subjected to 10 days of conditioning or pseudo-conditioning ('pseudo', *i.e.,* CS and US at random rather than fixed ISIs, preventing learning; Supplementary Fig. 2a), and we subsequently evaluated the number of excitatory and inhibitory synapses (Fig. 2). The vast majority of excitatory synapses in the CN belong to terminals of MFs and climbing fibers, whereas inhibitory synapses belong mainly to PCs. Presynaptic excitatory terminals can be detected by using antibodies against the vesicular glutamate transporter 1 (VGLUT1) or 2 (VGLUT2). Neurons in the pontine nuclei (PN) sending MFs to the cerebellum predominantly express

VGLUT1 mRNA, whereas neurons in the inferior olive, from which the climbing fibers originate, express

VGLUT2 mRNA[29,30].

We found a significantly increased density of VGLUT1+ puncta in the AIP of conditioned mice

when compared to pseudo-conditioned mice (conditioned vs. pseudo: $t_{(22)}$ = 2.352, P = 0.028, unpaired

t-test, two-tailed, cond: N = 11, pseudo: N = 13 mice; Fig. 2a-c). No significant difference was detected

between the ipsilateral and contralateral side in conditioned mice (ipsi vs. contra: $t_{(10)}$ = 1.876, P =

0.090, unpaired *t*-test, two-tailed, N = 11). Interestingly, we found a positive correlation between

VGLUT1 density in the ipsilateral AIP and the mean CR amplitude of the last conditioning session (r^2 =

0.35, P = 0.056, Pearson's correlation, N = 11), suggesting that the number of VGLUT1+ terminals in

the ipsilateral AIP coincides with a larger conditioned eyelid response (Fig. 2e). No correlation was

found for the contralateral AIP or for both sides pooled (contra AIP: r^2 = 0.074, P = 0.42; bilateral AIP:

r^2 = 0.20, P = 0.17, Pearson's correlation; Fig. 2f) or in pseudo-conditioned mice (contra AIP: r^2 = 0.026,

P = 0.62; ipsi AIP: r^2 = 0.087, P = 0.35; bilateral AIP: r^2 = 0.032, P = 0.58, Pearson's correlation). As for

VGLUT2+ terminals, no significant change was found in the AIP after conditioning (conditioned vs.

pseudo: $t_{(8)}$ = 0.9532, P = 0.37, cond: N = 4, pseudo: N = 6; Fig. 2d) and no significant correlation was

observed between VGLUT2 density and CR amplitude (contra AIP: r^2 = 0.025, P = 0.84; ipsi AIP: r^2 =

0.0090, P = 0.90; bilateral AIP: r^2 = 0.0012, P = 0.97, Pearson's correlation, N = 4).

To confirm that the increases in VGLUT1+ puncta represent changes in pontocerebellar MFs,

we overexpressed yellow fluorescent protein (YFP) in MFs by injecting mice with AAV9-hSyn-

hChR2(H134R)-EYFP bilaterally in the PN. Mice were subsequently conditioned or pseudo-conditioned

with a unilateral US on the left side (Supplementary Fig. 2a) and we evaluated the density of YFP+

axonal varicosities in the AIP (number/MF length; Fig. 2g-i). The vast majority of YFP+ varicosities were

also VGLUT1+ (>95%), indicating that we quantified largely VGLUT1+ MF axons. We found a significant

increase in the amount of varicosities in the AIP in conditioned mice compared to pseudo-conditioned

mice (bilateral AIP in pseudo vs. conditioned mice: $t_{(4)}$ = 4.689, P = 0.0094, unpaired *t*-test, two-tailed,

cond: N = 3, pseudo: N = 3; Fig. 2g-i). Again, no significant difference was detected between the

ipsilateral and contralateral side in conditioned mice ($t_{(3)}$ = 0.9401, P = 0.416, unpaired *t*-test, two-

tailed, N = 3).

We also investigated whether the increase in number of MF synapses in the AIP of conditioned

mice is accompanied by changes in MF terminals in the eyeblink area of the cerebellar cortex. MF

terminals (also known as rosettes) consist of multiple glutamatergic synapses on granule cell

dendrites. Rosettes were visualized thanks to YFP-reactivity in AAV-injected mice. The vast majority

(>95%) of YFP+ rosettes contained VGLUT1 (Fig. 2j). When we quantified the size of YFP+ rosettes in

the ipsilateral and contralateral side of conditioned mice, we again did not detect any difference (D =

0.08, P = 0.30, Kolmogorov-Smirnov test, N = 5). Moreover, no difference was present between

conditioned and pseudo-conditioned mice (D = 0.0016, P > 0.99, Kolmogorov-Smirnov test, cond: n =

613 YFP+ rosettes, N = 5 mice, pseudo: n = 761, N = 4; Fig. 2k).

To investigate whether the inhibitory input to the AIP also changes at the structural level

during conditioning, we quantified the number of gephyrin+ clusters in putative excitatory AIP

neurons (*i.e.,* neurons with a soma size >250 μm^2)[32,33]. Gephyrin is the core scaffolding protein in the

inhibitory post-synaptic density, which anchors glycine receptors and GABAA receptors at

postsynaptic sites[34,35]. The density of gephyrin puncta in the AIP was significantly increased in

conditioned mice ($t_{(8)}$ = 3.042, P = 0.016, unpaired *t*-test, two-tailed, cond: N = 4, pseudo: N = 6; Fig.

2l-o) as compared to pseudo-conditioned mice (data from both sides were pooled as gephyrin puncta

densities were comparable, $t_{(2)}$ = 0.241, P = 0.832, paired *t*-test, two-tailed, N = 4). To determine

whether the increase in gephyrin+ puncta can represent an increased input from PC terminals, we

quantified the percentage of gephyrin+ puncta that were apposed to calbindin-positive boutons (Fig.

2n). Given that calbindin is expressed by PCs as well as by climbing fibers[36,37], but only PC terminals

are inhibitory, gephyrin+ puncta that are apposed to calbindin-positive boutons represent inhibitory

connections from PCs. We found that in both conditioned and pseudo-conditioned mice, around 98%

of gephyrin+ puncta were apposed to PC terminals (conditioned mice: 97.4 ± 0.54%; pseudo-

conditioned mice: 98.15 ± 0.35%; *U* = 480.5, P = 0.551, Mann-Whitney *U* test, two-tailed, cond: N = 3,

pseudo: N = 3), indicating that the increase in gephyrin puncta following conditioning probably largely

reflects a change in PC transmission.

Together, these data indicate that terminals of pontocerebellar MFs to the AIP as well as

inhibitory PC axons increase bilaterally over the course of associative learning, whereas MF inputs to

the main eyeblink region in the cerebellar cortex remain unchanged.

Optogenetic stimulation of the pontine mossy fibers after EBC results predominantly in spike

facilitation in AIP neurons

Following conditioning, the predominant response of AIP neurons is spike facilitation induced by the

CS, with the spike rate increasing during learning[22]. We next set out to investigate whether activity in

MFs from the PN can lead to activity changes in the AIP after conditioning. We made juxtasomal loose-

patch recordings from AIP neurons of awake mice that had been injected with AAVs expressing the

excitatory opsin Chronos in the PN (Supplementary Fig. 3a,b). Our recordings at the end of

conditioning day 10 showed that 75% of the neurons showed changes in spike activity during

optogenetic stimulation of the pons with 465 nm (blue) light (Supplementary Fig. 3c). Blue light

stimulation resulted in spike facilitation with a latency shorter than 10 ms in 62% of the neurons,
suggestive of direct excitatory MF activation.

To confirm that this spike facilitation was due to a direct MF connection and not due to
disinhibition in the MF - granule cell - PC pathway, we subsequently made whole-cell recordings of CN
neurons in anesthetized GABRA6::ChR2EYFP mice that selectively express ChR2 in granule cells
(Supplementary Fig. 4a,b). Most (*i.e.,* 13 out of 15) of the recorded AIP neurons showed inhibition
rather than excitation in response to gc-ChR2 stimulation (Supplementary Fig. 4c,d), indicating that
activation of the MF - granule cell - PC pathway is unlikely to result in spike facilitation in CN neurons
and that the facilitation outcomes highlighted above could indeed result from direct MF to CN neuron
stimulation.

Optogenetic stimulation of the pontine nuclei leads to larger eyelid movements after conditioning

We next tested whether eyelid movements induced by optogenetic stimulation of PN neurons change
over conditioning. Mice were injected with AAVs expressing Chronos and/or eArch3.0 in the PN. Co-
injecting the two opsins allowed us to both activate and inhibit PN neurons using light stimulation of
different wavelengths. Next, we conditioned mice to a sensory CS for 10 days (Supplementary Fig. 2b).
We recorded eyelid movements on day 0 (D0), day 5 (D5) and day 10 (D10) that were generated in
response to optogenetic stimulation at a consistent light intensity across days (Supplementary Fig.
3d). In response to blue light stimulation, we found that conditioned mice made increased short-
latency eyelid closures on D10 compared to D0 (area under the curve 0-150 ms: $F_{(2,6)}$ = 6.883, P = 0.028,
D0 vs. D10: P = 0.023, one-way repeated-measures ANOVA with Tukey's multiple comparisons test, N
= 4 mice), while the peak-time of eyelid closures remained unchanged (86.8 ± 4.52 ms after light onset,
$F_{(1.254,3.761)}$ = 2.308, P = 0.214, one-way repeated-measures ANOVA; Supplementary Fig. 3e). Eyelid
closures were not observed during amber light stimulation inhibiting PN neurons ($F_{(1.594,8.764)}$ = 0.281,
P = 0.714, linear mixed models, N = 9 mice; Supplementary Fig. 3f), ruling out that this was a non-
specific reflexive eyelid closure to onset of the light. Together, these results suggest that increased
VGLUT1+ MFs to the AIP in conditioned mice may support larger eyelid movements after conditioning.

Membrane potential dynamics during CS-evoked eyelid movements

The data presented above raise the question as to how the changes in excitatory MF inputs and
inhibitory inputs shape the membrane potential (V_m) of AIP neurons over the course of EBC. We
therefore recorded from AIP neurons in whole-cell configuration during EBC in awake head-fixed mice

on a treadmill (Fig. 3a). A green LED flash (CS) signaled the arrival of a corneal air-puff (US) into the left eye with an ISI of 250 ms. We subjected naive and fully conditioned mice, the latter of which were conditioned for at least 10 days (Supplementary Fig. 2c), to a single session of *in vivo* whole-cell recordings in current-clamp mode targeting the AIP (Fig. 3a). Correct targeting was validated by inspection of a fluorescent tracer (Fig. 3b), electrical microstimulation and CS-evoked spike response patterns[22,38]. All mice were presented with alternating CS-US paired and CS or US unpaired trials during recordings. As expected, conditioned mice showed significantly more CRs than naive mice and their CRs showed higher amplitudes (%CR: $P < 0.001$, $U = 0$, Mann-Whitney U test, two-tailed; CR amplitude: $P < 0.001$, $U = 3$, Mann-Whitney U test, two-tailed; N = 12 conditioned/11 naive mice Fig. 3c). Interestingly, in 66.7% of the recordings, naive mice showed eyelid openings (EO) in response to the CS, movements directly opposite to classical CRs (%EO: $P = 0.015$, $U = 58$, Mann-Whitney U test, two-tailed; EO amplitude: $P = 0.056$, $U = 69$, Mann-Whitney U test, two-tailed, n = 10 recordings). Thus, naive and conditioned mice show directionally opposite eyelid responses to the CS, providing an opportunity to study the membrane potential of AIP neurons during stimulus-evoked eyelid movements towards different directions.

To investigate the physiological mechanisms underlying these opposite movements, simultaneous recordings of the V_m in current-clamp mode and eyelid movements were acquired (n = 15 neurons per group; Fig. 1a, Table 1). *In vitro* recorded AIP neurons can be classified into multiple cell-types[33,39], but classification based on *in vivo* electrophysiological properties alone appeared problematic[40,41]. AIP neurons showed a diversity of V_m responses to paired and unpaired CS and US stimuli, as well as responses during spontaneous voluntary eyelid movements (Fig. 3d,e, arrowheads). The population response including all neurons showed a transient V_m depolarization in naive mice and a more sustained depolarization in conditioned mice, coinciding with directionally opposite eyelid movements (Fig. 3f). ROC analysis revealed that the percentage of CS-responding neurons was 73.3% in naive mice (n = 11) and 80% in conditioned mice (n = 12; Fig. 3g). Most CS-responders showed V_m depolarizations after CS onset (naive, 53.3% of all neurons; conditioned, 66.7%; Fig. 3d, e). Since these depolarizations could be the result of excitatory input or reductions in tonic inhibitory input from PCs, we classified these neurons as 'CS-activated' (Supplementary Movie 1). A subset of neurons showed V_m hyperpolarizations and were classified as 'CS-suppressed' (naive, 20%; conditioned, 13.3%; Supplementary Fig. 5a-c; Supplementary Movie 2). The remaining neurons did not show any V_m responses evoked by the CS ('CS-no response', naive, 26.7%; conditioned, 20%; Supplementary Fig. 5d).

Overall, the distribution of CS responses was comparable between groups ($\chi^2_{(2)} = 1.083$, $P = 0.582$, chi-square test, n = 15 per group) (Fig. 3g). Since CS-activated AIP neurons were most prevalent,

and eyelid movements are thought to be driven by increased spike activity in excitatory projection neurons to downstream facial and oculomotor nuclei[23], we focused on CS-activated AIP neurons in our subsequent analysis. V_m depolarizations in CS-activated neurons in naive and conditioned neurons were similar in amplitude ($t_{(16)}$ = 0.857, P = 0.404, unpaired t-test, two-tailed, n = 10 conditioned/8 naive), peak time ($t_{(16)}$ = 1.567, P = 0.137, unpaired t-test, two-tailed) and area during the last 200 ms before the US (U = 35, P = 0.697, Mann-Whitney U test, two-tailed). However, the area during the last 50 ms before the US was significantly lower in naive neurons compared to conditioned neurons (U = 17, P = 0.043, Mann-Whitney U test, two-tailed; Fig. 3h,i, gray area). This shows that CS-evoked depolarizations underlying directionally opposite eyelid movements are transient in CS-activated naive neurons, but sustained in conditioned neurons.

V_m depolarizations reflect CRs in a subset of conditioned AIP neurons

To further investigate how conditioned eyelid movements are encoded, we separated V_m responses in trials with a CR (CR trials) and trials with no-eyelid response (NR trials; Fig. 4a). The initial rising phase of depolarizations was comparable between CR and NR trials, but higher sustained depolarizations on CR trials occurred during the last 50 ms before the US (Fig. 4a, gray area). ROC and correlation analysis revealed that only 33.3% of conditioned CS-activated responders showed CR dependency (Fig. 4b), whereas 66.7% of neurons responded similarly on CR and NR trials (Fig. 4c, Supplementary Fig. 6a). CR-dependent neurons showed larger V_m amplitudes (main-effect of eyelid response: $F_{(1,7)}$ = 6.734, P = 0.036, interaction-effect eyelid response vs. CR dependency: $F_{(1,7)}$ = 6.972, P = 0.033, CR vs. NR for CR-dependent group: P = 0.03, 2-way repeated measure ANOVA with Šídák's multiple comparisons test, n = 3 CR-dep/6 CR-indep) and area during the last 50 ms before the US on CR versus NR trials (main-effect of eyelid response: $F_{(1,7)}$ = 7.064, P = 0.033, interaction-effect eyelid response vs. CR dependency: $F_{(1,7)}$ = 9.567, P = 0.018, CR vs. NR for CR-dependent group: P = 0.019, 2-way repeated measure ANOVA with Šídák's multiple comparisons test; Fig. 4d). Furthermore, on a group level, V_m amplitudes and area correlated with eyelid movements (V_m amplitude: r = 0.435, P = 0.04; area: r = 0.545, P = 0.004, Spearman, two-tailed, Bonferroni-corrected, n = 34 trials, N = 3 neurons; Supplementary Fig. 6b). In contrast, for CR-independent neurons, V_m amplitudes and area did not correlate with eyelid movements (V_m amplitude: r = 0.14, P = 0.189; area: r = 0.035, P = 1, Spearman, two-tailed, Bonferroni-corrected, n = 203 trials, N = 6 neurons; Supplementary Fig. 6c). Among CS-suppressed neurons we did not find a clear relationship between V_m hyperpolarizing responses and eyelid movements (V_m amplitude: r = 0.157, P = 0.751; area: r = 0.172, P = 0.6, Spearman, two-tailed, Bonferroni-corrected, n = 72 trials, N = 2 neurons; Supplementary Fig. 5e,f). In

summary, CS processing can be widely observed in the membrane potential of many AIP neurons and part of them also show depolarizations in a CR-dependent fashion.

V_m depolarizations reflect eyelid openings in naive AIP neurons

All naive CS-activated neurons for which we recorded both EO and NR trials showed EO dependency (Fig. 4e,f). V_m amplitude, but not the area during the last 50 ms before the US, was larger on EO trials than on NR trials (V_m amplitude: $t_{(3)}$ = 6.258, P = 0.008; Fig. 4g; area: $t_{(3)}$ = 1.679, P = 0.192, paired *t*-test, two-tailed, n = 4 neurons). On a group level both the V_m amplitude and area correlated significantly with eyelid amplitude (V_m amplitude: *r* = 0.462, P < 0.001; area: *r* = 0.413, P < 0.001, Spearman, two-tailed, Bonferroni-corrected, n = 135 trials, N = 8 neurons; Supplementary Fig. 6d). However, CS-suppressed naive neurons did not show larger V_m hyperpolarizations on EO trials compared to NR trials (V_m amplitude: $t_{(2)}$ = 2.934, P = 0.099; area: $t_{(2)}$ = 1.429, P = 0.289, paired *t*-test, two-tailed, n = 3 neurons). Neither did we find a correlation between V_m hyperpolarizations and eyelid movements (V_m amplitude: *r* = 0.084, P = 0.927; area: *r* = 0.104, P = 0.726, Spearman, two-tailed, Bonferroni-corrected, n = 78 trials, N = 3 neurons; Supplementary Fig. 5g,h). Together, our data suggest that AIP neurons widely show CS-evoked V_m responses reflecting conditioned and unconditioned CS-eyelid movements, but with a varying degree of correlation with the movement.

Conditioning, but not pseudo-conditioning, leads to CS-evoked whole-body movements

One possibility for the varying degree of correlation with eyelid movements is that a subset of our recorded AIP neurons processes body movements other than the eyelid. Indeed, cerebellar rostral anterior interpositus nucleus neurons have previously been shown to drive coordinated whole-body movements[42]. To investigate whether conditioning leads to CS-evoked body movements, we acquired video recordings of mice subjected to conditioning or pseudo-conditioning and calculated the sum of frame-by-frame pixel changes as a measure of macroscopic body movements (movement index; Supplementary Fig. 7a, Supplementary Movie 3). During the first training session (D1), before the emergence of CRs, mice showed a short-latency response to CS onset (Supplementary Fig. 7b, black arrowhead) followed by a moderate increase in movement. After 10 days of training (D10), conditioned mice developed increased CS-evoked movements (Supplementary Fig. 7b, red arrowhead), whereas pseudo mice did not (main-effect of training day: $F_{(1,13)}$ = 0.098, P = 0.76, main-effect of group: $F_{(1,13)}$ = 7.296, P = 0.018, interaction-effect training day vs. group : $F_{(1,13)}$ = 6.355, P = 0.026, D10 cond vs. pseudo: P = 0.002, 2-way repeated measure ANOVA with Šídák's multiple

comparisons test, cond: N = 9 mice, pseudo: N = 6 mice; Supplementary Fig. 7d). The short-latency CS-responses did not change during training and remained comparable between groups (main-effect of training day: $F_{(1,13)}$ = 0.161, P = 0.694, main-effect of group: $F_{(1,13)}$ = 0.876, P = 0.366, interaction-effect training day vs. group : $F_{(1,13)}$ = 0.951, P = 0.347, 2-way repeated measure ANOVA). Both conditioned and pseudo mice showed a higher variability in baseline movements on D10 versus D1, but this effect was comparable between groups (main-effect of training day: $F_{(1,13)}$ = 9.857, P = 0.008, main-effect of group: $F_{(1,13)}$ = 0.302, P = 0.592, interaction-effect training day vs. group: $F_{(1,13)}$ = 0.019, P = 0.891, 2-way repeated measure ANOVA; Supplementary Fig. 7e).

Notably, the magnitude of CS-evoked movements correlated significantly with CR amplitude on a trial-by-trial basis for 55.6% (5 out of 9) of conditioned mice. The US alone also evoked movements, but neither conditioning nor pseudo-conditioning changed the magnitude of these movements (main-effect of training day: $F_{(1,8)}$ = 3.41, P = 0.102, main-effect of group: $F_{(1,8)}$ = 2.016, P = 0.193, interaction-effect training day vs. group: $F_{(1,8)}$ = 0.481, P = 0.508, 2-way repeated measure ANOVA; Supplementary Fig. 7c,f). These results indicate that eyeblink conditioning concurrently leads to generalized body movements, overall correlated with eyelid movements. Therefore, we cannot discount the possibility that at least part of the V_m responses in AIP neurons represent movements of other body parts.

V_m responses precede conditioned eyelid movements, but follow CS-evoked eyelid openings

We reasoned that if AIP neurons play a causative role in generating CRs, V_m responses should precede the onset of CRs. In contrast, unconditioned EOs in naive mice may well be the result of cerebral cortical activity[43,44] and their onset would not necessarily precede the onset of EOs. To test this, we quantified the onset of V_m and eyelid responses blindly on a trial-by-trial basis (Fig. 4h). We found that CRs started 147.3 ± 7.6 ms after CS onset and that EOs started 99.4 ± 5.2 ms after CS onset (n = 11 conditioned/8 naive neurons; Fig. 4j). Conditioned and naive V_m responses started 126.9 ± 6.3 ms and 123.7 ± 4.7 ms after CS onset, respectively. On a trial-by-trial basis, V_m responses started 22.88 ± 13.72 ms *earlier* than CRs in conditioned neurons, whereas V_m responses started 25.23 ± 6.92 ms *later* than EOs in naive neurons ($F_{(3,34)}$ = 8.964, P < 0.001, V_m vs. CR: P = 0.044, V_m vs. EO: P = 0.039, one-way ANOVA with Šídák's multiple comparisons test; Fig. 4i). These data indicate that the timing underlying eyelid movements becomes temporally precise to precede the conditioned eyelid movement. The response in naive AIP neurons could represent an efference copy of the command or feedback signal related to the EO, but likely does not represent an early signal that generates the EO itself.

Vₘ responses follow spontaneous eyelid movements

Since we found that depolarizing V_m responses may play a causative role in CR generation, we asked

how this compares to spontaneous eyelid openings or closures occurring in between stimuli.

Spontaneous eyelid movements likely do not coincide with movements in other parts of the body,

thereby providing us with a measure for the level of eyelid specificity of V_m responses. ROC analysis

revealed that V_m responses during spontaneous eyelid movements occurred in almost all conditioned

(91.67%) and more than half of naive (58.33%) neurons, thereby confirming that we were largely

targeting eyelid-specific AIP neurons (Supplementary Fig. 8a). 75% of all neurons showed responses

only during openings or closures but not during both. Furthermore, openings (52.38%) and closures

(47.62%) were comparably represented amongst the responders. There was no difference in V_m

responses between conditioned and naive mice (V_m amplitude: $F_{(3,11)}$ = 0.308, P = 0.819, one-way

ANOVA, two-tailed; eyelid amplitude $F_{(3,3.564)}$ = 1.098, P = 0.456, Brown-Forsythe ANOVA, two-tailed,

n = 6 conditioned/7 naive neurons), so we pooled the data from both groups of mice. V_m depolarizing

responses to spontaneous eyelid openings and closures, which were comparable in eyelid amplitude

(opening 15.55 ± 1.28 %UR, closure: 19.7 ± 5.95 %UR, $t_{(13)}$ = 0.728, P = 0.479, unpaired *t*-test, two-

tailed, n = 8 opening/7 closure neurons), also had comparable V_m amplitudes (opening: 7.12 ± 1.83

mV, closure: 5.6 ± 1.18 mV, $t_{(13)}$ = 0.674, P = 0.512, unpaired *t*-test, two-tailed; Supplementary Fig.

8b,c). Interestingly, across all pooled occurrences the amplitudes of V_m depolarizations and eyelid

movements did not significantly correlate, neither for openings (*r* = 0.066, P = 0.418, Spearman, two-

tailed, n = 155 openings/8 neurons) nor closures (*r* = 0.028, P = 0.791, Spearman, two-tailed, n = 91

closures/7 neurons; Supplementary Fig. 8e).

Overall, spontaneous eyelid movements started 26.14 ± 8.83 ms earlier than V_m

depolarizations ($t_{(9)}$ = 2.995, P = 0.008, one-sample *t*-test, one-tailed, n = 10 neurons) and this was

consistent for both openings and closures ($t_{(8)}$ = 0.757, P = 0.471, unpaired *t*-test, two-tailed, n = 6

opening/4 closure neurons; Supplementary Fig. 8d). For neurons that showed V_m hyperpolarizations,

we observed that eyelid movements started 34.91 ± 11.14 ms earlier than V_m hyperpolarizations ($t_{(3)}$

= 3.135, P = 0.026, one-sample *t*-test, one-tailed), a lag similar to V_m depolarizations ($t_{(12)}$ = 0.563, P =

0.584, unpaired *t*-test, two-tailed). Interestingly, these lags were almost identical to timings observed

during EOs, where EOs started 25.23 ± 6.92 ms earlier than V_m responses. In contrast, as discussed

earlier, CRs started 22.88 ± 13.72 ms after V_m responses ($F_{(2,28)}$ = 8.779, P = 0.001, spontaneous vs.

naive EO: P < 0.001, one-way ANOVA with Šídák's multiple comparisons test; Supplementary Fig. 8f).

Spike activity during conditioned response trials ramps up during ongoing CRs

To investigate how V_m responses are translated into spike activity during EBC we performed juxtasomal loose-patch recordings in conditioned mice, while recording CR and NR trials. Out of 112 recorded neurons (average spike rate 72.97 ± 3.08 Hz, range 3.6-165.3 Hz), 50% of the neurons responded with significant changes in spike rate to the CS. Of these responders 76.8% showed spike facilitation and 23.2% showed spike suppression (Supplementary Fig. 9a). CS-responding neurons were located in both the ipsilateral (52.7% of n=55 neurons) and contralateral (33.3% of n=33 neurons) hemisphere relative to the US. Neurons that responded with spike facilitation to the CS showed on average significantly increased spike rates during CR trials (CR: 48.22 ± 5.78 Hz, NR: 31.12 ± 4.95 Hz, Z = -341 P < 0.001, Wilcoxon, two-tailed, n = 30 neurons), while the timing of peak spike rate was comparable among trials (CR: 202.6 ± 9.98 ms, NR: 176.2 ± 11.43 ms, Z = -137, P = 0.083, Wilcoxon, two-tailed; Supplementary Fig. 9b-d). Similar to V_m recordings, average spike rate traces revealed that not only on CR, but also on NR trials, transient increases in spike rates occur. Yet again, only during the late phase of the CS-US interval these differences become prominent, temporally coinciding with the sustained phase of V_m depolarizations in CS-activated neurons. We correlated the number of spikes during the last 125 ms of the CS-US period on CR trials with eyelid amplitudes across spike facilitation neurons and found a significant correlation between CR amplitude and spike count (r = 0.218, P < 0.001, 648 trials in 30 neurons; Supplementary Fig. 9e). We also tested whether spikes during the first 125 ms correlate with CR amplitude, which it did (r = 0.139, P < 0.001, 648 trials in 30 neurons), although this correlation was weaker than that for spikes during the second half of the CS-US interval (Z = -2.299, P = 0.022, Fisher's z-transformation). Together, these data argue for a close translation from subthreshold to spike activity, where spikes during the ongoing CR, coinciding with the sustained phase of V_m depolarizations, are best correlated with the size of the conditioned eyelid response.

Decoding of eyelid behavior from V_m responses and spike data

To determine whether CS-evoked eyelid responses can be decoded from trial-by-trial V_m population data, we performed linear discriminant analysis (LDA) on spike-removed V_m traces, using a 10-fold cross-validation procedure to train and test a classifier. Conditioned (n = 199 CR trials and n = 265 NR trials in 15 neurons) and naive traces (n = 113 EO trials and n = 214 NR trials in 15 neurons) were used to decode CR vs. NR trials, and EO vs. NR trials, respectively. This analysis retains the temporal dynamics of the decoding accuracy across different phases of CS-US trials. We found that the decoder

performed significantly above chance level for both naive and conditioned traces during the CS-US interval (Fig. 5a). Significant above-chance decoding was observed starting 107.1 and 161.7 ms after the onset of the CS, for naive and conditioned traces, respectively. Furthermore, decoding accuracy from naive traces was significantly higher than conditioned between 123.1 and 201.7 ms after the CS (Fig. 5a, gray area). We also performed LDA on pooled spiking data from the juxtasomally recorded CN neurons in conditioned mice for comparison (n = 1444 CR trials and n = 414 NR trials in 112 neurons). The decoder performed significantly above chance between 155 and 245 ms after the CS (Fig. 5b, gray area). Peak decoding accuracy however was significantly lower than naive and conditioned V_m decoding ($H_{(2)}$ = 37.82, P < 0.001, naive V_m vs. cond V_m: P = 0.003, naive V_m vs. cond spikes: P < 0.001, cond V_m vs. cond spikes: P = 0.05, Kruskal-Wallist test with Dunn's multiple comparisons test; Fig. 5c). Overall, peak decoding accuracy occurred at 107.6 ms for naive V_m, 205.9 ms for conditioned V_m and 220 ms after CS onset for conditioned spike data (Fig. 5d).

We also asked how decoding accuracy changes as a function of group size. Decoding was performed on pooled trial data from groups of 15 to 3 neurons, where for each group we randomly selected a unique subset of neurons for 30 iterations. As expected, for both groups the decoder performed best with the maximal group size and accuracy declined with smaller group sizes (Supplementary Fig. 10a,b). The time the decoder performed significantly above chance declined significantly as a function of group size for both groups ($F_{(1,348)}$ = 4.016, P = 0.046, 2-way repeated measure ANOVA). The same trend was observed for variation in decoding accuracy ($F_{(1,348)}$ = 237.2, P < 0.001, 2-way repeated measure ANOVA), but not for mean decoding accuracy during the period of above-chance significance ($F_{(1,348)}$ = 1.435, P = 0.232, 2-way repeated measure ANOVA; Supplementary Fig. 10c-e). In summary, while V_m and spike data from both groups can sufficiently inform a decoder to distinguish trial types, decoding based on naive traces leads to the highest decoding accuracies covering the longest time span.

Encoding of the unconditioned stimulus in AIP neurons

We next asked what are the V_m dynamics underlying the US, a 10 ms air-puff directed at the left eye. The US is thought to mainly activate neurons in the inferior olive, which leads to complex spikes in PCs through CF activity[45], which also have collaterals that innervate the CN[46,47]. We found that all recorded conditioned and naive CN neurons showed significant V_m responses to the US, with V_m depolarizations as the most common response (cond: 80% US-activated, 13.3% US-suppressed, 6.7% US activated/suppressed; pseudo: 86.7% US-activated, 13.3% US-suppressed; Supplementary Fig. 11a,b). Typically, if neurons showed V_m depolarizations after the CS, they also did after the US, although some

neurons without CS-response did show US responses. Since there was significant variation in US-

evoked V_m depolarizations as well as UR amplitudes between groups, our dataset was insufficient to

detect a difference in V_m area (effect size d = 0.325, Df = 19, power = 0.175; Supplementary Fig. 11c,d).

However, comparing CR vs. NR trials on paired trials within the conditioned group revealed that

preceding CRs led to reduced US-evoked V_m depolarizations (amplitude: t_{10} = 2.368, P = 0.039, paired

t-test, two-tailed, area: t_{10} = 2.057, P = 0.067, paired *t*-test, two-tailed, n = 11 neurons; Supplementary

Fig. 11e,f). A similar effect of a preceding EO on US-evoked V_m depolarizations was not observed for

naive neurons (Supplementary Fig. 11g,h). Thus, a preceding CR influences V_m encoding of the US,

consistent with the protective function of the conditioned eyelid closure.

Discussion

In this study we provide novel insights into the synaptic and neuronal coding mechanisms during cerebellar learning in the AIP, the part of the cerebellar nuclei that is crucial for both the acquisition and expression of conditioned EBC behavior[48–53]. We show that stimulating MF afferents to AIP as a CS can effectively drive learning, with conditioned mice exhibiting well-timed CRs that are adaptive to changing CS-US delays. Chemogenetically silencing cerebellar granule cells selectively impairs learning to a cortical optogenetic CS, but leaves learning to a MF-AIP CS intact. Interestingly, learning with MF-AIP stimulation as a CS is not modulated by locomotor activity, in contrast to learning with cortical MF stimulation[25]. Taken together, these results strongly suggest that there is a capacity for well-timed learning within the AIP, without the need to involve the cerebellar cortex. The idea that the AIP is capable of generating learning without the participation of the cerebellar cortex has been previously put forward[54–56], but this is the first study that employs a targeted genetic approach to separate these pathways with mice achieving similarly high levels of EBC performance compared to those trained with a visual CS or optogenetic MF-cortex stimulation.

We further show that learning leads to structural changes in both excitatory MF and inhibitory inputs to the AIP, while MF input to the granule cell layer remains unchanged. At the same time we find that optogenetic stimulation of the PN neurons, the main source of MF input to the cerebellum, elicits larger eyelid movements after conditioning. By performing *in vivo* whole-cell recordings in awake behaving mice during EBC, we show how these learning-induced changes shape the V_m dynamics of AIP neurons. We find that conditioned neurons sustain a depolarized V_m state until US onset, while naive mice show transient V_m depolarizations. Importantly, after conditioning the V_m response becomes temporally precise in that it starts earlier than eyelid closures, away from the initial delay between V_m response and eyelid openings in naive mice. This indicates that conditioned AIP neurons could play a role in initiating the CR, whereas naive neurons likely process an efference copy of the eyelid opening with movements initiated by other brain areas[43,44]. This does not rule out the possibility that AIP activity modulates the ongoing CR or EO, which is likely since eyelid movements and V_m responses are well-correlated in a substantial subset of AIP neurons.

Juxtasomal spike recordings predominantly show increased spike responses during the second half of the CS-US interval, which temporally coincides with the sustained V_m depolarization as well as the upward ramping phase of CRs. Both V_m and spike data are sufficient to inform a decoder to separate movement from no-movement trials during similar periods of the CS-US interval, despite the fact that a considerable proportion of AIP neurons shows CR-independent V_m responses and some show no detectable CS-responses. This response heterogeneity may reflect a combination of the

cellular diversity of the CN[57] and differences in input patterns and activity. AIP neurons may also
differentially encode aspects of the conditioned movement, or even movements of multiple body
parts during defensive behavior[42]. Nevertheless, an open question remains to what extent CS
responses correlate with CN cell types and what are the precise contributions of inhibitory and
excitatory inputs to the V_m and spike response. Here we chose to prioritize current-clamp recordings
to measure how AIP neurons behave during EBC without experimenter interference. This approach
did not allow us to distinguish inhibitory versus excitatory input, as possible during voltage-clamp
recordings. We envision a future study employing voltage-clamp recordings in cell type-identified AIP
neurons as a next step in this line of study.

There is broad consensus that CRs emerge as the result of well-timed pauses in PC activity in
response to the CS[14,19], where plasticity in the AIP would be under the control of PC inputs[58]. However,
our results indicate that at least part of the memory trace may be generated and stored directly at the
level of the nuclei[59], at least in part through structural modifications of VGLUT1+ MF and inhibitory
inputs. This would be in agreement with the observation that many AIP neurons gradually increase
their spike rates over the course of learning, coinciding with larger conditioned eyelid movements[22].
Besides changes in the number of inputs, other forms of plasticity may take place in parallel. For
instance, changes in synaptic ultrastructure[60] and transmission, intrinsic excitability[61], postsynaptic re-
localization of receptors or changes in activity from inputs[14], may also take place. This could occur
under the control of perineuronal nets harboring chemo-attractive or repulsive molecules[62]. Our
recordings show the sum total of these changes as they occur during learning, thereby providing a
unique insight into how the output of the cerebellum is conveyed to downstream motor nuclei before
and after learning.

One has to be aware of several potential caveats in our study, *e.g.,* CNO is known to have
potential off-target effects. Although we used a concentration that is well within the range typically
used for hM4Di receptors[27] and we implemented proper controls in the form of non-DREADD-
expressing animals injected with CNO, the potential effects of repeated injections of CNO have not
been well documented. Furthermore, the precise location of our whole-cell recordings could not be
determined *post-hoc*, as the relatively short duration of recordings (149 s median) did not allow for
reliable neurobiotin filling. In response to this limitation we performed surface-to-depth
measurements, microstimulation, fluorescent tracer injections as well as validation of spike activity to
ensure that we targeted the AIP, although this does not ascertain that all recorded neurons were from
AIP. We further observed that mice develop whole-body movements, which are correlated with the
eyelid movement after conditioning, but not pseudo-conditioning. Activity of AIP neurons has been
shown to correlate with control of multiple body parts[42] and thus our estimation of eyelid-related

activity in the AIP may be overestimated. However, it should be noted that similar V_m activity was also

present during spontaneous eyelid movements, which occur isolated from other body movements,

arguing in favor of eyelid-specificity. In the same line of reasoning, one could hypothesize that

optogenetic stimulation of PN neurons may lead to increased whole-body movements besides the

eyelid. To determine whether this is the case and to determine to what extent eyelid-encoding AIP

neurons are also involved in other functions requires a follow-up study.

Nevertheless, our findings shed new light on the longstanding debate regarding the relative

contributions of the cerebellar cortex and nuclei for cerebellum-dependent motor learning. Lesion

and reversible inactivation studies have suggested that these two areas may encode different features

of learning, with plasticity at MF-AIP synapses mediating the expression of motor responses, while

cortical cells regulate their gain and timing[21,53,63–72]. One of the most influential models in the field

posits that temporally specific plasticity in the cerebellar cortex leads, over learning, to well-timed

pausing of PCs, and, consequently, a downstream disinhibition of AIP neurons[63]. Indeed, AIP neurons,

which are like PCs highly intrinsically active[22,40], are unusually sensitive to the differential impact of

synchronized activity of PCs during behavior[73]. Our findings demonstrate how a combination of

structural and physiological changes at the level of MF inputs to the CN may work together with

cerebellar cortical plasticity mechanisms to facilitate efficient associative learning.

Acknowledgements

We are grateful to the following people for their contribution to the study: J.W. Potters, H.J. Boele, S.K.E. Koekkoek, Z. Gao, M.M Ten Brinke, J.K. Spanke, A.C.H.G. IJpelaar, M. Tang, B. Winkelman, A. Court, C. Geelen, M. Mešković, G. Özel, M. Bouw, S.O. Stokman, M. Romijn, S. Dijkhuizen and T. Jacobs. We thank Tracy Pritchett and Ana Machado for maintenance of mouse lines. Catarina Carvalhas assisted with some of the data acquisition. We thank Jovin Jacobs for helpful discussions throughout the project. We thank I. Duguid and P. Chadderton for their constructive comments. This study was enabled by funding from the Netherlands Organization for Scientific Research (NWO-ALW 824.02.001; C.I.D.Z., NWO STEM - VBT 2021 19224 and NWO 863.14.005; C.B.C.), the Dutch Organization for Medical Sciences (ZonMW 91120067; C.I.D.Z.), Medical Neuro-Delta (MD 01092019-31082023; C.I.D.Z.), INTENSE LSH-NWO (TTW/00798883; C.I.D.Z.), ERC-adv (GA-294775 C.I.D.Z.) and ERC-POC (nrs. 737619 and 768914; C.I.D.Z.), the NIN Vriendenfonds for Albinism (C.I.D.Z.), the Dutch NWO Gravitation Program, Dutch Brain Interface Initiative (DBI2; C.I.D.Z.), the Boehringer Ingelheim Fonds, a Howard Hughes Medical Institute International Early Career Scientist Grant #55007413 (M.R.C.), European Research Council Starting (#640093) and Consolidator (866237) Grants (M.R.C.) and a fellowship from the Portuguese Fundação para a Ciência e a Tecnologia SFRH/BD/77686/2011 (C.A.).

Author contributions

R.B., C.A., C.B.C., M.R.C., C.I.D.Z. conceived the project and designed the experiments. R.B. performed and analyzed the awake and anaesthetized whole-cell and juxtasomal recordings, pontine nuclei optogenetics, behavioral trainings for histological investigation, whole-body video recordings and decoding analysis. C.A. performed and analyzed the optogenetic mossy fiber ChR2 and chemogenetic experiments. D.C. performed and analyzed the histological procedures. R.B., C.A., D.C., C.B.C. M.R.C., C.I.D.Z. wrote and edited the manuscript. C.B.C., M.R.C., C.I.D.Z. provided supervision. C.I.D.Z., C.B.C. and M.R.C. acquired the funding.

Declaration of interests

The authors declare no competing interests.

Fig. 1. Mice can learn to optogenetic stimulation of mossy fibers in the cerebellar nuclei as a conditioned stimulus. a, Schematic of a head-fixed mouse on a treadmill with an optic fiber implanted just above the AIP. Optogenetic stimulation of mossy fiber (MF) terminals in the AIP as a conditioned stimulus (CS) was paired with an air-puff to the ipsilateral eye (unconditioned stimulus, US). **b,** Representative histological sample indicating the placement of an optical fiber over the AIP of Thy1-ChR2-YFP mice expressing ChR2-YFP in MFs. **c,** Average eyelid traces of sessions from a representative animal trained to optogenetic stimulation of MF terminals in the AIP (MF-AIP CS). Each trace represents the average of CS-only trials from sessions 1, 2, 4, 6 and 8. **d**, Average %CR learning curves to a MF-AIP CS for animals walking on a self-paced treadmill (slow) and those running at a fixed speed of 0.18 m/s on a motorized treadmill (fast). **e**, Onset session of learning (average CR amplitude of session exceeds 10% of UR) plotted against the average walking speed on the treadmill of animals trained to a MF-AIP CS (blue) and to a MF-CTX CS (purple). Lines represent linear fits (MF-CTX: N=14, slope = -16.8, P < 0.001; MF-AIP: N=12, slope = 1.4, P = 0.564). **f**, Cerebellar circuit diagram with lightning bolts representing the sites of optogenetic stimulation: mossy fiber terminals in the lobule simplex (purple) and MF terminals in the AIP (blue). MF, mossy fiber; Gc, granule cell; PC, Purkinje cell; IN, interneuron; PF, parallel fiber; AIP, anterior interpositus nuclei. **g,** Fluorescence image of a parasagittal cerebellar section of a MF-ChR2- gc-DREADD mouse, indicating ChR2-YFP expression in MFs (yellow fluorescence), and hM4Di receptors (DREADDs) selectively in granule cells (red fluorescence). **h-j,** Average %CR learning curves of granule cell-expressing DREADD mice (darker colors) and respective non-DREADD-expressing controls (lighter colors), injected with CNO prior to

each training session. **h,** Animals trained to a MF-AIP CS. **i,** Animals trained to a MF-CTX CS. **j,** Animals

trained to a natural visual CS. Data is shown as mean ± SEM.

Fig. 2. Learning-induced changes in VGLUT1+ mossy fiber terminals and gephyrin+ inhibitory

terminals. a-b, Expression of VGLUT1 and VGLUT2 in the AIP of a representative conditioned (cond; a)

and pseudo-conditioned (pseudo; b) mouse. **c**, The density of VGLUT1+ puncta in the AIP is higher in

conditioned mice (N = 11) compared to pseudo mice (N = 13). **d**, The density of VGLUT2+ puncta in the

AIP is unchanged after conditioning (cond: N = 4, pseudo: N = 6). **e, f,** Density of VGLUT1+ puncta in

the AIP plotted against the CR amplitude (%UR) at the last training session for conditioned animals.

VGLUT1 density correlates positively with the CR amplitude in the ipsilateral (e) but not in the

contralateral (f) side. **g-h,** Representative pictures of YFP+ mossy fibers and varicosities, which mostly

contain VGLUT1 (arrows), in conditioned (g) and pseudo (h) mice. **i,** The number of varicosities/mm

axon length is increased in conditioned mice (N = 3) when compared to pseudo (N = 3) mice. **j-j,**

Representative pictures of mossy fiber terminals (MFT; or rosettes), labelled by YFP and for the vast

majority containing VGLUT1 (arrows), in the EBC area of the cerebellar cortex. **k,** The size of

YFP+/VGLUT1+ MFTs in conditioned mice (N = 5, n = 613) is not different from that of pseudo mice (N

= 4, n = 761). **l-m,** Expression of gephyrin in the AIP of conditioned (l) and pseudo (m) mice. **n,** The vast majority of gephyrin+ clusters in a putative glutamatergic AIP neuron are apposed to CALB+ terminals (arrows). **o,** Conditioned mice show a significantly higher density of gephyrin+ clusters when compared to pseudo mice (pseudo: N = 6; cond: N = 4). Data is shown as mean ± SEM. *P < 0.05, **P < 0.01.

Fig. 3. Membrane potential responses during the conditioned stimulus in cerebellar nuclei neurons.
a, Schematic of *in vivo* whole-cell recordings of head-fixed mice during EBC. Eyelid responses to CS and US stimuli are recorded using a high-speed camera. Upward deflections in the eyelid trace indicate eyelid closures, downward deflections indicate eyelid openings. CR: conditioned response, EO: eyelid opening, NR: no response, CS: conditioned stimulus, US: unconditioned stimulus, AIP: anterior interpositus nuclei, CCD: charge-coupled device camera. **b**, Recordings were targeted at the AIP. Electrode track is visualized by coating the electrode with Evans blue dye. **c**, Performance of conditioned and naive mice during the recordings. Conditioned mice showed a higher %CR and CR amplitude, while naive mice show a higher %EO and EO amplitude. **d-e**, Example traces of a conditioned and a naive CS-activated neuron. **d**, Conditioned AIP neuron showing V_m depolarizations

during paired, CS only and US only trials, but also during spontaneous eyelid movements (arrowhead). Larger depolarizations are seen on CR trials compared to NR trials. **e**, Naive AIP neuron showing V_m depolarizations during paired and CS only trials, and during spontaneous eyelid movements (arrowhead). Larger depolarizations are seen on EO trials compared to NR trials. **f**, Population eyelid and V_m response averages across all neurons show an eyelid closure and sustained V_m depolarization for conditioned mice and an eyelid opening coinciding with a transient depolarization for naive mice. **g**, Distribution of CS-evoked V_m responses for both groups, showing in both cases that the predominant response classification is CS-activation. **h**, Eyelid and V_m response averages for CS-activated AIP neurons. Gray area indicates the last 40 ms before the US used calculation of area in (i). **i**, Graphs showing quantifications of V_m responses shown in (h). The area during the last 40 ms before the US is significantly higher in conditioned than in naive neurons, indicating a sustained versus transient profile of V_m depolarizations, respectively. Graphs and traces show mean ± SEM. *$P < 0.05$, ***$P < 0.001$, #$P = 0.056$.

Fig. 4. Relationship between membrane potential responses in CS-activated neurons and eyelid movements. a, Average eyelid and V_m traces on conditioned response (CR) and no response (NR) trials of all CS-activated neurons in conditioned mice. **b,** Average V_m traces on CR and NR trials for neurons that show different responses on both types of trials (CR-dependent). Most prominent differences occur during the last 50 ms before the US (gray area). **c,** Same as (b), but for neurons that show similar responses on CR trials compared to NR trials (CR-independent). **d,** Graphs comparing the V_m amplitude and the area within the gray area in (b) and (c). V_m amplitude and area are increased on CR trials for CR-dependent, but not for CR-independent neurons. **e,** Average eyelid and V_m traces on eyelid opening (EO) and no response (NR) trials of all CS-activated neurons in naive mice. **f,** Average V_m traces on EO and NR trials for neurons that show different responses on both types of trials (EO-dependent). Gray area indicates 50 ms before start of the US. **g,** Graphs comparing the V_m amplitude and area during the 50 ms before the US, showing that the amplitude, but not the area, is significantly increased on EO trials compared to NR trials. **h,** Eyelid and V_m traces on CR or EO trials for CS-activated neurons recorded in conditioned or naive mice, respectively, aligned to the onset of the CR/EO (left panels) or the onset of the V_m response (right panels). Traces include a combination of CS and US responses since the CS-US delay is 250 ms and CS-only and paired trials are pooled. **i,** Onset of V_m response versus onset of CR or EO for all conditioned and naive neurons, respectively. On average, V_m responses start earlier than CRs, but later than EOs. **j,** Onsets of V_m responses and EO or CR responses relative to the onset of the CS. As shown in (i), V_m responses in naive mice start after EOs, but V_m responses start

698   before CRs in conditioned mice. Boxplots show median, whiskers show minimum and maximum

699   values. Data is shown as mean ± SEM in all panels, except (j). *P < 0.05, **P < 0.01.

Fig. 5. Decoding of eyelid movements based on conditioned stimulus-evoked neuronal activity. a,

Decoding of CR versus NR trials based on V_m data from all conditioned neurons and decoding of EO

versus NR trials based on V_m data from all naive neurons. Decoding was significantly above chance for

both groups, but during different time periods relative to the CS-US interval. Time periods of

significant decoding are indicated with the colored horizontal bars and dotted lines for each group.

Gray area indicates the period during which decoding based on naive neurons performed significantly

better than decoding based on conditioned neurons. Eyelid and V_m traces are visualized as reference.

b, Decoding of CR versus NR trials based on juxtasomal spike data from all conditioned neurons.

Decoding was significantly above chance within the period indicated with the horizontal bar and

dotted line, coinciding with the upward sloping CR and largest differences in the average spike rates

(lower panels). **c**, Peak decoder accuracy was highest based on V_m traces from naive neurons. Within

the conditioned group decoding based on V_m traces outperformed decoding based on spike data. **d**,

Time of the peak decoding accuracy relative to CS onset. Data is shown as mean ± SEM. *P < 0.05, **P

< 0.01, ***P < 0.001.

Supplementary Fig. 1. Timing of conditioned responses during granule cell inhibition and shifting of inter-stimulus intervals. a-c, Average eyelid traces from granule cell-expressing DREADD mice (darker colors) and respective non-DREADD-expressing controls (lighter colors), injected with CNO prior to training sessions. Averages include CS-only trials from the last session of training. Vertical gray line indicates the expected time of US delivery. **a,** Animals trained to MF-AIP CS. **b,** Animals trained to MF-CTX CS. **c,** Animals trained to a visual CS. **d,** Eyelid peak times comparing DREADD mice versus non-DREADD-expressing controls, conditioned with different types of CS. **e,** Schematic of experimental design: mice were trained to shifting ISIs of 500, 200 and back to 500 ms. **f-g,** Average eyelid traces from mice trained to a MF-AIP CS or a MF-CTX CS. Traces include CS-only trials from sessions with a ISI of 500 ms (blue line), 200 ms (cyan line) and back to 500ms (dashed blue line). **h,** Average CR peak time for MF-AIP CS and MF-CTX CS animals trained to shifting ISIs: 500 ms, 200 ms and 500 ms. No differences were found across ISIs between the two groups (respectively: P = 0.62, P = 0.56, P = 0.95, unpaired *t*-test, two-sided). In d and h, dots represent individual animals. Boxes indicate median and 25th-75th percentiles, and whiskers extend to the most extreme data points. In all other graphs data are shown as mean ± SEM.

Supplementary Fig. 2. Learning curves of conditioned and pseudo-conditioned mice. a, Learning
curves of conditioned (cond) and pseudo-conditioned (pseudo) mice for histological quantifications of
biomarkers during acquisition training. Conditioned, but not pseudo-conditioned mice showed
increases in %CR (main-effect of day: $F_{(9,23.878)}$ = 51.525, P < 0.001, main-effect of group: $F_{(1,24.066)}$ =
383.255, P < 0.001, interaction day x group: $F_{(9,23.878)}$ = 48.090, P < 0.001), CR amplitude (main-effect
of day: $F_{(9,23.325)}$ = 15.710, P < 0.001, main-effect of group: $F_{(1,23.408)}$ = 79.379, P < 0.001, interaction day
x group: $F_{(9,23.325)}$ = 18.365, P < 0.001) and amplitude at US onset (main-effect of day: $F_{(9,23.992)}$ = 158.821,
P < 0.001, main-effect of group: $F_{(1,24.006)}$ = 127.739, P < 0.001, interaction day x group: $F_{(9,23.992)}$ =
153.423, P < 0.001, linear mixed models). **b**, Learning curves of conditioned mice for optogenetic
stimulation of the pontine nuclei (PN), showing increasing %CR ($F_{(3.098,24.79)}$ = 16.27, P < 0.001), CR
amplitude ($F_{(2.885,23.08)}$ = 4.385, P = 0.015) and amplitude at US onset ($F_{(4.139,33.11)}$ = 10.78, P < 0.001, one-
way repeated-measures ANOVA). **c**, Learning curves of conditioned mice for awake whole-cell
recordings, showing increasing %CR ($F_{(12,87.223)}$ = 6.889, P < 0.001), CR amplitude ($F_{(12,72.079)}$ = 3.6, P <
0.001) and amplitude at US onset ($F_{(12,84.439)}$ = 6.069, P < 0.001, linear mixed models). After acquisition
training mice were placed in the electrophysiology setup and trained for >3 days to maintain high
levels of performance (day before recording). **d**, Learning curves of conditioned mice for awake
juxtasomal recordings, showing increasing %CR ($F_{(9,49.188)}$ = 7.134, P < 0.001) and amplitude at US onset
($F_{(9,47.846)}$ = 4.599, P < 0.001, linear mixed models; left panels). Average performance of the same mice
during juxtasomal recordings (right panels), CRs occurred in 57.95 ± 2.67% of trials during recordings
and had amplitudes of 48.38 ± 3.1 %UR. Data is shown as mean ± SEM.

Supplementary Fig. 3. Optogenetic stimulation of pontine nuclei neurons leads mainly to spike facilitation and eyelid movements after conditioning. a, Schematic of optogenetic stimulation of pontine nuclei (PN) neurons giving rise to MFs targeting the cerebellum. AAV-Chronos-GFP was injected bilaterally in the PN and optic fiber implants were placed directly above. b, Expression of GFP+ axons in the PN, anterior interposed nucleus (AIP) and granule cell layer (GCL). No GFP expression in the molecular layer (ML) indicating that no climbing fibers were labeled. c, Juxtasomal recordings (n = 24) showing 62% spike facilitation, 13% spike suppression and 25% no response neurons. Example spike traces show representative spike facilitation and suppression neuron. Histograms show average spike rates across spike facilitation and suppression neurons. Short-latency spike changes occur after PN activation. d, Schematic of dual optogenetic stimulation. AAV-Chronos-GFP and AAV-eArch3.0-YFP were co-injected bilaterally in the PN followed by optic fiber implants. Eyelid movements were recorded while optogenetically stimulating PN neurons. e, Average eyelid movements of mice (N = 4) during optogenetic stimulation of PN neurons with 465 nm (blue) light on day (D) 0, D5 and D10 of conditioning. Area under the curve (AUC) of short-latency eyelid movements (0-150 ms after light onset) showing a significant increase in magnitude of eyelid closure on D10 (arrowhead; AUC at D0 vs. D5 vs. D10: 184.2 ± 13.57 vs. 258.9 ± 57.1 vs. 335.7 ± 55.31, $F_{(2,6)}$ = 6.883, P = 0.028, D0 vs. D10: P =

0.023, one-way repeated-measures ANOVA with Tukey's multiple comparisons test). **f**, Average eyelid

movements of mice (N = 9) during optogenetic stimulation of PN neurons with 590 nm (amber) light

on D0, D5 and D10 of conditioning. AUC of eyelid movements (0-150 ms after light onset) does not

show significant changes over the course of conditioning ($F_{(1.594, 8.764)}$ = 0.2812, P = 0.714, linear mixed

models). Data are shown as mean ± SEM.

Supplementary Fig. 4. Cerebellar nuclei neurons show mainly inhibition during optogenetic

activation of granule cells. a, Schematic of whole-cell recordings of anterior interposed nucleus (AIP)

neurons during optogenetic stimulation of granule cells with 465 nm (blue) light in GABRA6-

Cre::ChR2EYFP transgenic mice. **b**, Expression of ChR2EYFP specifically in granule cells and parallel

fibers in the molecular layer. **c**, AIP neurons show predominantly inhibitory post-synaptic potentials

(86.7%) immediately after onset of optogenetic light. **d**, Example representative neuron showing

inhibitory responses to granule cell activation. Raw traces (black) with average trace (red)

superimposed. Electrical noise artifacts at light onset and offset are clipped.

Supplementary Fig. 5. Membrane potential responses in CS-suppressed cerebellar nuclei neurons.

a, Example trace of a representative conditioned CS-suppressed neuron. V_m hyperpolarizations with temporary cessation of spike activity occur during CS-US paired trials and between trials. Larger average hyperpolarizations appear on conditioned response (CR) trials compared to no response (NR) trials during the CS-US interval (right panels). CS: conditioned stimulus, US: unconditioned stimulus. **b**, Eyelid and V_m response averages for CS-suppressed neurons, separating conditioned and naive neurons. Gray area indicates the last 40 ms before the US used calculation of area in (c). **c**, Graphs

showing quantifications of V_m responses shown in (b). No statistical comparison due to the small

sample size. **d**, Eyelid and V_m response averages for CS-no response neurons, separating conditioned

and naive neurons. These neurons do not show CS-evoked V_m responses, irrespective of eyelid

movements. **e**, Average eyelid and V_m traces on CR and NR trials of CS-suppressed neurons in

conditioned mice. **f**, Normalized V_m amplitude (top panel) and normalized area during the last 50 ms

of the CS-US interval (bottom panel) are not correlated with normalized eyelid amplitude across

pooled trial data from conditioned CS-suppressed neurons (n = 72 trials, N = 2 neurons). **g**, Average

eyelid and V_m traces on eyelid opening (EO) and NR trials of CS-suppressed neurons in naive mice. **h**,

Normalized V_m amplitude (top panel) and normalized area during the last 50 ms of the CS-US interval

(bottom panel) are not correlated with normalized eyelid amplitude across pooled trial data from

naive CS-suppressed neurons (n = 78 trials, N = 3 neurons). Linear regression lines for each individual

neuron (pink) and for pooled trials of all neurons (red) in f and h. Graphs and traces show mean ± SEM.

Supplementary Fig. 6. Membrane potential responses in CS-activated CR-dependent and CR-independent cerebellar nuclei neurons. a, Example trace of a representative conditioned CS-activated CR-independent neuron. V_m depolarizations and spike activity occurs in response to conditioned (CS) and unconditioned stimuli (US), in addition to spontaneous eyelid movements (arrowhead). V_m depolarizations are similar on conditioned response (CR) and no response (NR) trials, and are therefore not dependent on eyelid movement (CR-independent; right panel). An example of a CR-dependent neuron is shown in Fig. 3d. **b**, Normalized V_m amplitude (left panel) and normalized area during the last 50 ms before the US are both significantly correlated with the normalized eyelid amplitude across pooled trial data from conditioned CR-dependent CS-activated neurons (n = 34 trials, N = 3 neurons). **c**, Normalized V_m amplitude (left panel) and normalized area during the last 50 ms before the US are not correlated with the normalized eyelid amplitude across pooled trial data from conditioned CR-independent CS-activated neurons (n = 203 trials, N = 6 neurons). **d**, Normalized V_m amplitude (left panel) and normalized area during the last 50 ms before the US are both significantly correlated with the normalized eyelid amplitude across pooled trial data from naive EO-dependent CS-activated neurons (n = 135 trials, N = 8 neurons). In panels b-d, linear regression lines for each individual neuron are indicated in pink and for pooled trials of all neurons in red.

Supplementary Fig. 7. Mice develop whole-body movements during conditioning. a, Video
recordings of infrared-illuminated mice during conditioning or pseudo-conditioning on a treadmill. **b**,
Whole-body movements visualized as movement index for conditioned (N = 9) and pseudo (N = 6)
mice. Both groups show short-latency movements to conditioned stimulus (CS) onset, both on day 1
(D1) and day 10 (D10; black arrowhead). Conditioned, but not pseudo mice, develop movements
during the period of the CS-US interval where conditioned eyelid movements occur on D10 (red
arrowhead). **c**, Whole-body movements during the unconditioned stimulus (US) does not change
during conditioning or pseudo-conditioning. **d**, Areas under the curve (AUC) on CS-only trials,
comparing D1 versus D10 between groups, show that significant body movements develop on D10 in
conditioned mice. **e**, Standard deviation (SD) of baseline body movements, occurring before CS or US
stimuli, shows a significant increase for both groups on D10. **f**, Body movements in response to the US
do not significantly change. Graphs and traces show mean ± SEM. **P < 0.01.

Supplementary Fig. 8. Membrane potential dynamics during spontaneous eyelid movements. a,

Average eyelid and V_m traces of neurons showing V_m depolarizing responses during spontaneous

eyelid openings and eyelid closures. Traces are aligned to the onset of the eyelid movement (top

panels) or aligned to the V_m onset (bottom panels). Only neurons in which the onsets of V_m responses

and eyelid movements could be determined are included. **b-c**, Graphs showing quantifications of the

absolute eyelid amplitude (b) and V_m amplitude (c), which show no difference in amplitude of the

eyelid movement and V_m response. **d**, Average lag of V_m response relative to the onset of eyelid

opening or closure is comparable. **e**, Normalized V_m amplitude and normalized eyelid amplitude show

no correlation for eyelid openings (n = 155 openings, N = 8 neurons) or eyelid closures (n = 91 closures,

N = 7 neurons). Linear regression lines for each individual neuron are indicated in pink and for pooled

trials of all neurons in red. **f**, Onset of eyelid movements relative to V_m onset, comparing spontaneous

to CS-evoked eyelid movements. Spontaneous eyelid movements as well as CS-evoked eyelid openings

(EO) start before onset of the V_m response. CRs start on average after the V_m response. Boxes and

whiskers show median with minimum and maximum values. All other panels show mean ± SEM.

Supplementary Fig. 9. Spike activity during conditioned response and no response trials in

conditioned mice. a, Schematic of *in vivo* juxtasomal recordings from AIP neurons in head-fixed mice

on a treadmill during EBC (top). Distribution of spike responses to the conditioned stimulus (CS;

bottom) shows that the predominant response is spike facilitation. **b**, Example of a representative

spike facilitation neuron showing increased spike activity during a conditioned response (CR) trial, but

no change in spike activity during a no response (NR) trial. **c,** Average of all spike facilitation neurons

showing eyelid movements (top) on CR and NR trials with normalized spike rates (bottom). **d**, Average

peak spike rates on NR and CR trials. Significantly higher spike rates are observed during CR trials, but

the time of peak spike rate after CS onset is comparable. **e**, Pooled correlation analysis of all CR trials

(n = 648 trials, N = 30 neurons) shows a significant correlation between number of spikes during the

CS-US interval and CR amplitude. Data are shown as mean ± SEM; *** indicates P < 0.001.

Supplementary Fig. 10. Decoding of eyelid movements based on V$_m$ data decreases with decreasing group sizes. a, Average decoding accuracy (30 iterations; CR versus NR trials) over the course of a trial based on V$_m$ data, for different group sizes of conditioned neurons (n = 15 to n = 3 neurons). **b**, Average decoding accuracy (30 iterations; EO versus NR trials) over the course of a trial based on V$_m$ data, for different group sizes of naive neurons (n = 15 to n = 3 neurons). **c**, Total time that above-average decoding was achieved decreases with decreasing group sizes and plateaus to ~25 ms for both groups. **d**, Variability in decoding accuracy, shown as standard error of the mean (SEM) across iterations, increases with decreasing group sizes. **e**, Mean decoding accuracy during the time period of significant above-chance decoding does not significantly decrease with decreasing group sizes. Data are shown as mean ± SEM; *$P < 0.05$ and ***$P < 0.001$.

Supplementary Fig. 11. Membrane potential dynamics and spike coding during the unconditioned

stimulus. a, Population eyelid and V_m response averages across all neurons, showing predominantly

V_m depolarizations in conditioned and naive neurons coinciding with the unconditioned eyelid closure

response (UR). **b**, Distribution of US-evoked V_m responses. For both groups the predominant response

classification is US-activation. **c**, Eyelid and V_m response averages of US-activated neurons. **d**, Graphs

showing quantifications of US responses shown in (c). There are no differences in UR amplitude, V_m

amplitude, peak time and area following the US between conditioned and naive neurons. **e**, Eyelid and

V_m response averages on paired trials of US-activated conditioned neurons, separating CR and NR

trials. **f**, Graphs showing quantifications of US responses shown in (e). A preceding CR significantly

lowers the amplitude of the US-evoked V_m depolarization. The same effect is reflected as a trend in

the V_m area following the US. *$P < 0.05$, #$P = 0.067$. **g**, Eyelid and V_m response averages on paired trials

of US-activated naive neurons, separating EO and NR trials. **h**, Graphs showing quantifications of US

responses shown in (g). There are no differences in UR amplitude, V_m amplitude, peak time and area

following the US between EO and NR trials, indicating that a preceding EO does not influence the US-

evoked V_m response. Graphs and traces show mean ± SEM.

Supplementary Movie 1. Example recording of a CS- and US-activated conditioned neuron. Awake

whole-cell recording (real-time speed) during eyeblink conditioning, showing simultaneous camera

recording of the eye, quantification of eyelid opening and CS and US stimulus signals, followed by a

single paired trial (10x slower).

Supplementary Movie 2. Example recording of a CS-suppressed and US-activated & suppressed

naive neuron. Awake whole-cell recording (real-time speed) during eyeblink conditioning, showing

simultaneous camera recording of the eye, quantification of eyelid opening and CS and US stimulus

signals, followed by a single paired trial (10x slower).

Supplementary Movie 3. Whole-body movements develop only in conditioned mice. Camera

recordings of an example conditioned and pseudo-conditioned mouse, comparing responses to the

conditioned stimulus (CS) on training day 1 vs. day 10. Only conditioned mice develop whole-body

movements after CS onset.

Table 1. Electrophysiological properties of neurons recorded in awake mice.

Group	ID	Mouse ID	Side	Depth (μm from pia)	V$_m$ rest (mV)	Spike amp (mV)	Spike HW (ms)	Rs (MΩ)	Ri (MΩ)	Rm (MΩ)	Cm (pF)
Cond	1	1	Ipsi	n/a	-59.0	29.5	0.3	43.0	111.4	68.3	75.8
	2	2	Ipsi	1910	-60.4	27.9	1.8	24.7	89.5	64.9	35.9
	3	3	Ipsi	1538	-69.8	36.8	1.0	43.7	518.1	474.4	3.5
	4	4	Ipsi	1885	-68.7	23.9	0.8	73.6	621.1	547.5	7.7
	5	5	Ipsi	1741	-64.8	35.3	0.5	36.4	211.0	174.6	161.0
	6	6	Contra	1771	-68.7	22.0	1.0	37.1	980.4	943.3	18.8
	7	6	Contra	1830	-71.1	n/a	n/a	52.1	135.7	83.5	1.4
	8	7	Contra	1956	-61.3	21.0	0.9	57.5	305.8	248.3	62.6
	9	7	Contra	1694	-70.2	38.9	2.1	39.6	840.3	800.8	2.7
	10	8	Contra	1757	-58.4	18.2	0.5	61.2	308.6	247.4	87.1
	11	9	Contra	1701	-69.8	22.2	1.1	42.6	425.5	382.9	5.0
	12	10	Ipsi	1900	-73.2	n/a	n/a	57.3	110.9	53.5	3.8
	13	10	Ipsi	2005	-72.1	n/a	n/a	71.2	119.0	47.8	13.1
	14	11	Ipsi	n/a	-74.7	n/a	n/a	61.8	89.4	27.6	147.5
	15	12	Ipsi	1608	-68.8	n/a	n/a	50.0	125.8	75.8	6.6
Mean ± SEM				1792 ± 38.3	-67.4 ± 1.4	27.6 ± 2.3	1.0 ± 0.2	50.1 ± 3.5	332.8 ± 74.4	282.7 ± 75.1	42.2 ± 13.8
Naive	1	1	Ipsi	2244	-68.1	24.2	0.5	63.1	115.6	52.6	73.9
	2	1	Contra	1600	-70.0	15.2	1.6	74.6	568.2	493.6	5.4
	3	2	Ipsi	1978	-66.7	27.5	0.9	29.1	213.2	184.1	21.6
	4	3	Ipsi	1757	-67.4	26.3	1.0	28.3	190.8	162.5	38.2
	5	4	Contra	n/a	-55.3	32.1	0.3	31.5	57.9	26.4	163.4
	6	5	Ipsi	1712	-65.9	27.7	1.0	37.7	216.9	179.2	44.2
	7	6	Ipsi	1526	-69.2	27.7	1.6	35.4	352.1	316.7	4.6
	8	6	ipsi	1637	-68.9	n/a	n/a	61.8	136.4	74.6	23.8
	9	7	Ipsi	1968	-67.3	14.9	2.2	62.1	317.5	255.3	32.6
	10	8	Contra	2081	-68.9	51.8	1.3	30.2	885.0	854.8	17.0

	11	8	Contra	1903	-68.9	48.9	1.0	20.6	1754.4	1733.8	23.3
	12	9	Contra	1795	-67.3	25.3	0.9	51.8	125.3	73.5	51.7
	13	9	Contra	1775	-74.8	n/a	n/a	49.1	78.4	29.2	4.4
	14	10	Ipsi	n/a	-70.7	40.0	1.1	41.0	452.5	411.5	5.5
	15	11	Ipsi	1850	-73.8	n/a	n/a	53.8	97.5	43.7	90.9
Mean ± SEM				**1813 ± 52.9**	**-68.2 ± 1.1**	**30.1 ± 3.3**	**1.1 ± 0.1**	**44.7 ± 4.1**	**370.8 ± 114.4**	**326.1 ± 116.1**	**40.0 ± 11.0**

Methods

Animals and ethics statement

All animal experimental procedures were conducted in accordance with the institutional animal

welfare committees of the Erasmus Medical Center, the Royal Dutch Academy of Arts and Sciences,

the Champalimaud Centre for the Unknown or the Direcçãoo Geral de Veterinária (Ref. No.

0421/000/000/2015). All experiments adhered to the European guidelines for the care and use of

laboratory animals (Council Directive 86/6009/EEC). Male and female C57Bl/6J mice (4-14 weeks of

age, from institutional colony and from Janvier Laboratories) were socially housed with *ad libitum*

access to food and water. Mice for optogenetic stimulation of the mossy fibers (Fig. 1, Supplementary

Fig. 1) were kept on a reverse light cycle (12:12 hour light/dark) so that all experiments were

performed during the dark period while mice were more active. Mice used for the remaining

experiments were kept on a normal light cycle (12:12 hour light/dark) and experiments were

performed at day time.

Transgenic mouse lines

The Thy1-ChR2-YFP mouse line (B6.Cg-Tg(Thy1-COP4/EYFP)18Gfng/J)[74] was obtained from The

Jackson Laboratory (stock number: 007612). The MF-ChR2- gc-DREADD mouse line was obtained in

two steps. First, Gabra6-Cre mice (B6.D2-Tg(Gabra6-cre)B1Lfr/Mmucd)[75] obtained from MMRRC

(stock number: 015966-UCD) were crossed with R26-LSL-Gi-DREADD mice (B6.129-

Gt(ROSA)26Sor$^{tm1(CAG-CHRM4*,-mCitrine)Ute}$/J)[76] obtained from The Jackson Laboratory (stock number:

026219). Second, Gabra6-Cre::DREADD animals were crossed with Thy1-ChR2-YFP mice to obtain

Gabra6-Cre::Thy1-ChR2::DREADD mice. Gabra6-Cre::ChR2(H134R)-EYFP animals were obtained by

crossing Gabra6-Cre mice with Ai32 mice (B6.Cg-*Gt(ROSA)26Sor$^{tm32(CAG-COP4*H134R/EYFP)Hze}$*/J)[77] obtained

from The Jackson Laboratory (stock number: 024109).

Surgical methods

Pedestal/head plate surgery

Mice were anesthetized with isoflurane (4-5% induction, 0.5-2% maintenance; in 0.2 L/min O_2 and 0.2

L/min air) and placed in a stereotaxic frame (David Kopf Instruments) where their body temperature

was kept at 37 °C using a feedback-controlled heating-pad. Eyes were covered with antimicrobial

ointment (Terra-Cortril, Pfizer) to prevent drying. Skin covering the skull was shaved and a mid-sagittal

incision of approximately 1 cm was made to expose the skull bone, after which local anesthesia was

applied on the skin around the incision (10% Xylocaine, AstraZeneca). Primer (Optibond All-In-One,

Kerr) was applied on the bone and treated with UV light. A 6 x 3 x 5.7 mm aluminum block ('pedestal'),

or a head plate for MF-ChR2 experiments, was attached to a stereotaxic arm and attached to the skull

bone with dental acrylic (Flowline, Heraeus Kulzer or Super Bond, C&B). Skin edges were attached to

the dental acrylic using tissue glue (Histoacryl, Aesculap). Post-operative analgesia was given

subcutaneously with Meloxicam (2 mg/kg Metacam, Boehringer Ingelheim Animal Health) and mice

were placed under an infrared light while recovering from anesthesia. Mice recovered for at least two

days following surgery before subsequent experiments.

Viral injections

Mice were prepared for surgery as described above. A longer mid-sagittal incision was made until the

edge of the occipital bone. A stereotaxic arm was used to mark the location of entry to reach the

lateral pontine nuclei (PN) (AP -3.8 to -4 mm, ML 1 mm, DV 5.5 to 5.8 mm rel. to Bregma), comparable

to previously used coordinates[18]. Two small craniotomies were made using a dental drill (Foredom

Drill K1070-2E, Blackstone Industries) and the dura was carefully removed using a fine forceps

(Dumont #5). For optogenetic stimulation of the PN, mice were injected with AAV1/Syn-Chronos-GFP

(UNC Vector Core, #AV6550) or co-injected with AAV2/hsyn-eArch3.0-EYFP (UNC Vector Core,

#AV5229B). For tracing of fluorescently labeled MFs in the AIP and cerebellar cortex, mice were

injected with AAV9-hSyn-hChR2(H134R)-EYFP (Addgene, #26973-AAV9). The Chronos and eArch3.0

viruses were diluted 4x and a total of 1 µL virus solution was injected in each hemisphere. The ChR2-

EYFP virus was not diluted and 220 nL was injected in each hemisphere. In both cases we used

borosilicate glass pipettes connected to a Nanoject II or Nanoject III Programmable Nanoliter Injector

(1-3 µL/min, Drummond Scientific). Injections of 69 nL were made every 1 min and after the last

injection the pipette was left in place for 2-10 minutes to allow diffusion and was then retracted.

Optic fiber implants

For MF-ChR2 experiments, unilateral optic fibers (Ø200 µm, 0.53 NA, Doric lenses) were lowered into

the brain and positioned at the cortical eyelid region (AP -5.7 mm, ML +1.9 mm, DV 1.5 mm rel. to

Bregma) or just over the AIP (AP -6 mm, ML +1.7 mm, DV 2.1 mm rel. to Bregma)[78] . The implants were

fixed into place using dental cement (Super Bond, C&B). Mice were subsequently monitored during 2

days of recovery. For the remaining experiments, bilateral optic fibers (Ø200 µm,0.39 NA, FT200UMT,

Thorlabs) connected to ceramic zirconia ferrules (MM-FER2007C-2300, Precision Fiber Products) were

lowered into the brain through the same injection track until a depth of 5 mm to enable light delivery

to the PN. Fibers were attached to the skull bone using primer (Optibond All-In-One, Kerr) and dental

cement (Flowline, Heraeus Kulzer). Dental cement was painted using silver conductive paint (RS

Components) to reduce the escape of optogenetic light. Mice were allowed to recover at least 2 weeks

before subsequent experiments.

Craniotomy surgery for in vivo electrophysiology

Mice for awake whole-cell recordings were prepared for surgery as described above. Eyes were protected and kept moist using eye drops (Duodrops, Ceva Santé Animale). Hairs in the neck were removed and an incision was made in the skin covering the neck muscles, after which local anesthesia was applied (10% Xylocaine, AstraZeneca). Neck muscles were removed to expose the occipital bone and two square-shaped bilateral craniotomies were made using a dental drill. Position of craniotomy was above Crus II to access the CN in a 43° angle. A thin layer of UV-curing primer (Optibond All-In-One, Kerr) was applied around the craniotomies and a recording bath was made using dental acrylic (Flowline and Charisma, Heraeus Kulzer). The recording bath was filled with 0.9% NaCl solution to keep the dura moist. Skin and muscles surrounding the bath were attached to the dental cement using tissue glue (Histoacryl, Aesculap). The dura in the craniotomy was carefully removed using a needle tip (30G x ½ inch, Microlance, BD) and fine forceps (Dumont #7 curved). Directly after removing the dura, the recording bath was cleaned with 0.9% NaCl solution and filled with a low viscosity silicone elastomer sealant (Kwik-cast, World Precision Instruments) to protect the brain tissue from the air. Mice were given post-operative analgesia subcutaneously (Metacam 2 mg/kg) and were placed under an infrared light during recovery. Mice were closely monitored for at least two hours before starting the recording session. Gabra6-Cre::ChR2-EYFP mice for anesthetized whole-cell recordings were subjected to a similar craniotomy surgery, but received an intra-peritoneal injection of 75 mg/kg ketamine and 12 mg/kg xylazine instead of isoflurane anesthesia. The depth of anesthesia was ensured by monitoring the heartbeat, whisker movements and reflexes to a tail-pinch. Additional ketamine-xylazine mixture (10% of initial dose) was administered if necessary. Gabra6-Cre::ChR2-EYFP were subjected to whole-cell recordings immediately following the craniotomy surgery.

Eyeblink conditioning

Habituation

Mice for MF-ChR2 experiments were trained on an experimental setup previously described[25]. Head-fixed mice habituated for >3 days to walk at the target speed on the motorized treadmill until they walked without signs of distress. Speed of the treadmill was controlled by a DC motor with an encoder (Maxon). For all other experiments mice were handled and habituated for 3-5 days with increasing durations (15-60 min). Habituations were done in sound- and light-isolating boxes (Neurasmus) equipped with a cylindrical, non-motorized treadmill and a horizontal bar for head fixation[71]. Eyelid movements were measured using magnet distance measuring technique (MDMT)[79] or using a high-speed camera[22]. In order to place a magnet under the eye for MDMT mice were shortly anesthetized

with isoflurane (5% in 0.5 L/min O_2 and 0.5 L/min air). Hairs under the left eyelid were removed under

isoflurane anesthesia on day 2. From day 3 onward a 1.5 x 0.5 x 0.7 mm neodymium magnet

(Supermagnete) was placed under the left eye using super glue (Bison International), after which they

were placed on the treadmill and woke up in the dark environment of the box. Eyelid movements

were recorded through a magnetic sensor that was placed above the left eye. During the habituation

session, 20 CS trials and 2 US trials with an inter-trial-interval (ITI) between 10-12 ± 2 sec were

presented to familiarize the mouse with the stimuli. The CS consisted of a 260 ms light flash delivered

through a green/blue LED placed ~7 cm in front of the mouse. The US consisted of a 10 ms corneal air-

puff (35-40 psi) delivered through a P20 pipette tip positioned ~5 mm from the left eye. The US airflow

was controlled by a Milli-Pulse Pressure Injector (MPPI-3, Applied Scientific Instrumentation). CS and

US stimuli were identical in camera boxes, but no magnet or magnet sensor was used. Mice also did

not undergo isoflurane anesthesia in these boxes.

1033

Acquisition

Mice for MF-ChR2 experiments were subjected to 6 to 10 days of acquisition sessions, consisting of

the presentation of 90% CS-US paired trials and 10% CS-only trials. Each session consisted of 110 trials,

separated by a randomized ITI of 10-15s. In each trial, the inter-stimulus interval (ISI) between the CS

and US was 300ms, 500ms for the timing experiments, and both stimuli co-terminated. The US was an

air-puff (30-50 psi, 50 ms) controlled by a Picospritzer (Parker) and delivered via a 27G needle

positioned 0.5 cm away from the cornea of the right eye, positioned so that it induced a full eye blink.

The visual CS was a white light LED positioned 2 cm directly in front of the mouse. For all other

experiments acquisition trainings lasted 5-10 days and consisted of one training session daily. A

session was comprised of 200 CS-US (paired) trials, 20 CS trials and 20 US trials, presented as a 20x

repeated block of 1US-10paired-1CS. Trials followed a standardized format of 0-500 ms baseline, 500-

760 ms CS (paired and CS trials), 750-760 ms US (paired and US trials), 760-2000 ms post-stimuli

period. MDMT data was captured at 1017.26 Hz using custom-written LabVIEW software (National

Instruments)[79]. Trainings were started at least 10 min after placing mice in the box to allow full

recovery of isoflurane anesthesia. Trials were initiated only if the eye was >75% opened.

1049

1050 *CS-US timing experiments*

1051 Mice were trained with a long ISI of 500 ms for 10 consecutive sessions, then switched to a shorter

1052 ISI of 200ms for the duration of 3 sessions, and finally switched back to 3 sessions of 500ms ISI. At

1053 the end of each block of sessions with a particular ISI we performed a test session where 50% of the

1054 trials were CS-only. These trials were included in our timing analysis.

1055

1056 *Granule cell DREADD experiments*

1057 Gabra6-Cre::Thy1-ChR2::DREADD mice received intra-peritoneal injections of CNO (5 mg/kg; Tocris

1058 Bioscience, #4936) 60-90 min before starting eyeblink experiments.

1059

Eyeblink conditioning during electrophysiology

Mice for *in vivo* juxtasomal and whole-cell recordings were first trained in boxes using MDMT, after

which they were transferred to an electrophysiology setup equipped with a camera. Mice were then

trained for at least 3 more days on this setup before performing recording sessions, to ensure they

reached maximum performance. A 250 fps CCD camera (scA640-120gc, Basler) was directed at the left

eye, which was illuminated with an infrared light. Camera frames were transformed into a data vector

based on pixel thresholding within a region of interest overlaying the eye and the eyelid. The resulting

data was sampled at 2441 Hz. Stimuli were triggered by custom-made software utilizing TDT System

3 (Tucker Davis Technologies) and NI-PXI (National Instruments) processors. During the US the TTL

onset was corrected for the delay between TTL onset and air-puff arrival at the cornea (13 ms). To

avoid distortion of the camera signal caused by reflection of the infrared light from the whiskers, a

thin layer of water-proof black mascara was applied on the left whiskers before mice were head-fixed.

Timings and parameters of stimuli were identical to those used during the acquisition training

sessions.

1074

Eyeblink data analysis

For MF-ChR2 experiments videos from each trial were analyzed offline with custom-written MATLAB

software (MathWorks). Distance between eyelids was calculated frame-by-frame by thresholding the

grayscale image of the eye and extracting the count of pixels that constitute the minor axis of the

elliptical shape that delineates the eye. Eyelid traces were normalized for each session, ranging from

1 (full blink) to 0 (eye fully open). Trials were classified as CRs if the eyelid closure reached at least 0.1

(in normalized pixel values) and occurred between 100 ms after CS onset and US onset. The average

running for each animal was calculated by summing the average speed of each session (total distance

run divided by session duration) and dividing by the total number of learning sessions, usually 20.

Running speed for trial was calculated by dividing the distance run in the ITI preceding the current trial

by the elapsed time. For all other experiments, MDMT data was processed using custom LabVIEW

software (Neurasmus). Data was subjected to several quality checks: (i) trials with an unstable baseline

were removed (trials with values exceeding 5x SD during the baseline period) and (ii) sessions with a

large variation in UR amplitude (>0.5 x CV) were excluded. Finally, sessions from which more than 75%

of trials were removed were excluded to ensure accurate behavioral measurements. Similarly to the

MF-ChR2 experiments, CRs were defined as eyelid closures during the last 200 ms of the CS-US interval

that exceeded 10% of UR amplitude. UR amplitude was calculated as the average of full eyelid closures
during US only trials and used to normalize the eyelid movements. Camera data, including data from
electrophysiology experiments, were analyzed using custom-made MATLAB scripts. Acquired data
were preprocessed by filtering with a Gaussian low-pass filter (50 Hz cutoff frequency) and trials with
unstable baselines (values >5 x SD of the baseline period) were excluded. CRs were defined as eyelid
closures during the last 200 ms of the CS-US interval exceeding 10% of the UR amplitude. In some
cases, CRs were defined as eyelid closures exceeding values >5 x SD of baseline values during the last
200 ms of the CS-US interval. CR percentage was calculated by dividing the number of CR trials by the
total amount of eligible trials, multiplied by 100%. CR amplitude was calculated by dividing the
maximum eyelid closure during the CS-US interval by the average UR amplitude, multiplied by 100%.
CR amplitude at US onset represents the amplitude of eyelid closure at US onset (250 ms after CS
onset. CR or EO onsets were manually detected on a trial-by-trial basis in a blind manner and represent
the first data point before a continuously rising (CR) or decreasing (EO) signal.

1104

Spontaneous eyelid movement analysis
Spontaneous eyelid movements were identified during periods of recordings after removing the 2 s
windows surrounding eyeblink trials. Eyelid movements were detected in Gaussian filtered (5 Hz cut-
off) eyelid traces, as values of the first derivative that exceeded 3 x SD of the filtered trace. The
'findpeaks' MATLAB function with a minimum peak distance of 500 ms and minimum peak
prominence of $1x10^{-5}$ (a.u.) returned the positions of significant eyelid movements (openings or
closures). Only isolated movements without openings or closures within a 500 ms window before the
maximum velocity of the movement were included. Spontaneous eyelid onset was determined to be
the first data point exceeding 2 x SD of the second eyelid derivative of the trace. Onset of V_m
depolarizations or hyperpolarizations were manually determined and were defined as the time point
of the last minimum before a continuously increasing signal or decreasing signal, respectively. V_m
responses were tested for significance by ROC analysis on amplitudes surrounding the maximum
eyelid velocity and preceding baselines.

1118

Microelectrode stimulation and Evans Blue labeling
Microelectrode stimulation was performed to determine the recording location targeting the
eyeblink-encoding region of the CN. Eyelid movements were induced using an 80-µm-diameter
platinum iridium monopolar electrode (100 KΩ; Alpha Omega), lowered into the brain with a 43° angle
until a depth of 1500-1600 µm was reached. Electrodes were advanced in steps of 100 µm while
currents were generated (ISO-flex Stimulus Isolater, A.M.P.I.) in the range of 1–15 µA (200 ms pulse

trains; 250 µs biphasic pulses; 500 Hz). The depth at which reliable eyelid contractions could be observed with low current amplitudes (<10 µA) was identified as the eyeblink-encoding region of the CN. To mark this area in the CN, the electrode was retracted, coated with Evans Blue solution (5% in saline, Sigma-Aldrich) and reinserted using identical position and coordinates. Mice were transcardially perfused using 4% paraformaldehyde (PFA, Sigma-Aldrich) in phosphate buffered saline (PBS) after a lethal dose of pentobarbital (Nembutal) and brains were collected for histology.

In vivo electrophysiology

Whole-cell and juxtasomal recordings were made using glass electrodes with a tip diameter of 1-2 µm and a resistance of 4-8 MΩ. Electrodes were heat-pulled on a P-1000 micropipette puller (Sutter Instrument) from filamented borosilicate glass capillaries (1.5 mm OD, 0.86 mm ID, Harvard Apparatus). Electrodes were back-filled with intracellular solution, containing (in mM): 10 KOH, 3.48 $MgCl_2$, 4 NaCl, 129 K-Gluconate, 10 HEPES, 17.5 glucose 4 Na_2ATP, and 0.4 Na_3GTP (295 - 305 mOsm; pH 7.2). The intracellular solution was supplemented with 0.5% Neurobiotin (Vector Labs). Electrophysiological recordings were made using a Multiclamp 700B amplifier (Axon Instruments, Molecular Devices) and digitized at 100 or 50 kHz with a Digidata 1440 digitizer (Axon Instruments). Electrodes were mounted with a 43° angle on a pipette holder connected to a head stage (CV-7B, Axon Instruments). Pipette movements were controlled by a micromanipulator (SM7, Luigs und Neumann, Ratingen, GE). Pipettes were targeted at the AIP region of the CN and entered the surface of the cerebellum 1.5-2.0 mm lateral and -1.5 mm below Bregma, as previously described[40]. Electrodes were advanced while maintaining high pressure to a depth of 1500-1600 µm relative to the brain surface, after which the pressure was lowered to 15-20 mbar.

During whole-cell recordings, electrodes were advanced in steps of 2 µm while monitoring the current during 10 mV steps at 33 Hz in voltage-clamp mode without holding potential. All recorded neurons reached a seal resistance of at least 1 GΩ before establishing whole-cell configuration. Several protocols were run shortly after break-in to characterize the neuron, including a seal test protocol (10 mV steps at -65 mV) and a current step protocol (200 to -200 pA) in current clamp mode after compensation. Following these protocols, continuous recordings were made without current injection in current clamp mode while CS, US and CS-US paired trials were given (naive: 1:1:1 ratio in a randomized order, conditioned: repetitive sequences of 1US-5paired-1CS to maintain conditioned behavior throughout the experiment). Trial ITI was 12 ± 2 s and no threshold for eyelid opening was used to maximize the number of trials. Camera recordings of the eyelid were made simultaneously and synchronized with electrophysiology by recording TTL pulses controlling the CS and the US. Recordings were made until the neuron was lost or the recording quality became insufficient. Only

1159   one recording session was performed following the craniotomy surgery with a duration less than 4

1160   hours. During juxtasomal recordings, electrodes were advanced until action potentials ('spikes'; >1

1161   mV) of a single neuron could be identified. Then, CS, US and CS-US trials were given while spiking

1162   responses were recorded. Juxtasomal recordings were made once per day, for 1-3 sessions each with

1163   a duration of less than 4 hours.

1164

Electrophysiology data analysis

Calculation of electrophysiological parameters

Data were processed in Clampfit (v10.5, Axon Instruments) and MATLAB (R2011b and R2022a,

MathWorks). Series resistance (R_s) and input resistance (R_i) were calculated during the seal test in

voltage-clamp (holding of -70 mV), based on the peak current and steady-state current during a 10

mV voltage step, respectively. Membrane resistance (R_m) represents R_i subtracted with R_s. Membrane

time constant (tau) was estimated by fitting a single exponential function on the decay phase of the

seal test or calculating the time between 100-37% of peak-to-steady state amplitude of the 10 mV

voltage step. Membrane capacitance was calculated based on tau, R_s and R_m. Resting membrane

potential (V_m rest) was estimated based on R_s and the current necessary to maintain a holding

potential of -70 mV in voltage-clamp. Spikes in V_m recordings were detected and analyzed using

Clampfit. Spikes were detected during threshold search on the first trace in the current step protocol

that induced spikes. For some neurons we were unable to analyze spikes, because our current step

protocol did not induce spikes. Since we prioritized V_m measurements during EBC and spike rates were

recorded juxtasomally in other recordings, we did not further drive the neuron to spike threshold.

1180

Exclusion and quality criteria whole-cell recordings

Recordings were discarded if R_s exceeded 100 MΩ. Only traces that had a limited drift in V_m were

analyzed. Some recordings were affected by an incorrect V_m offset directly visible when switching to

current-clamp, caused by an issue with the reference electrode. Since we analyzed V_m responses

relative to baseline, we included these neurons. Most recordings were without current injection, but

a few neurons received limited current (-25 to -200 pA) to maintain V_m at original values directly after

break-in. Average recording depth of all neurons (1812.4 ± 33.4 μm) was consistent with the depth of

the CN[40]. Neurons were recorded in both hemispheres (table 1) and we did not find a difference in CS-

evoked V_m amplitude between sides for either group (cond: P = 0.77, naive: P = 0.571, pooled: 0.7,

unpaired t-test).

1191

Spike analysis of juxtasomal recordings

Spikes were detected using a custom-written MATLAB spike analysis program (B.H.J. Winkelman,
Netherlands Institute for Neuroscience, Amsterdam), after which spikes were sorted based on spike
waveform characteristics. Only data with spikes from single units were included in the analysis. Based
on spike times we calculated: spiking frequency (= number of spikes / duration of time period) and
coefficient of variation of spiking (CV = standard deviation (all inter-spike-intervals (ISIs) / average (all
ISIs)). Spontaneous spike calculations were based on time periods between trials.

1199

Processing of V_m and eyelid data

Electrophysiology and eyelid data were loaded into MATLAB using the 'abf2load' (H. Hentschke,
University of Tuebingen and F. Collman, Princeton) and the 'TDTbin2mat' function, respectively, and
further analyzed using custom MATLAB code. In short, TTL timings were used to synchronize both
types of data. In V_m traces spikes were removed by median filtering (10 ms window). Trial epochs were
normalized to baseline (250-0 ms) before the CS. CS-evoked responses were obtained by pooling
paired and CS-only trials, and CS-responses were analyzed until US onset (250 ms after CS onset). No
separate analysis of CS only trials was performed because of the low number compared to the number
of CS-US paired trials. To establish whether a neuron showed responses to a stimulus we used non-
parametric ROC analysis on trial-by-trial amplitudes during the baseline (250-0 ms before CS onset)
vs. the period of interest (CS: 50-250 ms after CS; US: 0-500 ms after US). The overlap between
amplitude distributions was quantified as area under the ROC curve (AUC). Bootstrap analysis (1000
repeats) with shuffled data from those distributions was used to test for statistical significance
between AUC distributions. Amplitude and peak time of the CS-evoked V_m response was determined
on a trial-by-trial basis 50-250 ms after CS onset. Area was calculated as the sum of values over time
periods indicated in the text. Amplitude, peak time and area of the US-evoked V_m response was
determined on a trial-by-trial basis between 0-500 ms after US onset. Data per neuron were obtained
by averaging trial quantifications. Onset of V_m depolarizations or hyperpolarizations were manually
determined and were defined as the time point of the last minimum before a continuously increasing
signal or decreasing signal, respectively. Onsets of CRs or EOs were manually determined in a similar
way, while the experimenter was blind to the coinciding V_m trace.

1221

Whole-body video recordings

Videos were acquired using iPi Recorder 4 software at 60 fps (800x600 pixels) using an infrared-
integrated camera (ELP-USBFHD05MT-KL36IR, ELP). Videos were converted to AVI format using iPi
Mocap Studio 4 and analyzed in MATLAB. The sum value of the first derivative between two
subsequent frames was calculated as a measure of movement. The trace was filtered using a Gaussian

filter (25 Hz cut-off) and normalized using the median and the max values across the recording. CS

onsets were identified by reflection of the LED on the wall of the box. Amplitude and area were

calculated between 0-250 ms after CS onset (CS quantifications) and 0-500 ms after US onset (US

quantifications).

1231

Optogenetics

During MF-ChR2 experiments, optogenetic channels were stimulated using 473 nm laser light (LRS-

0473-PFF-00800-03, Laserglow Technologies), controlled by custom-written LabVIEW (National

Instruments) software. Laser power was adjusted for each mouse and controlled for each experiment

using a power meter (Thorlabs) at the beginning and end of each session. With these stimulation

parameters, stimulation is likely to be highly localized (predicted spread from cortex to AIP sites of

<1% based on https://web.stanford.edu/group/dlab/optogenetics). For activation of ChR2 in mossy

fibers as a CS, laser power was lowered below the threshold for detectable eyelid movement upon

repeated stimulus presentation. The optogenetic CS consisted of stimulation that lasted for 350 ms

(or 550 ms in the timing experiments) delivered in trains of 2 ms pulses and was paired with a co-

terminating 50 ms air-puff US. Laser power ranged from 0.20 to 1 mW, corresponding to intensities

(power output per unit area) of 6 to 30 mW/mm2 for stimulation within the cerebellar cortex, and of

3 to 20 mW/mm2 for stimulation in the AIP.

In experiments using optogenetic stimulation of PN neurons 465 nm light-induced activation

of Chronos was achieved using a 5W LED (60 lm, LZ1-B200, LED Engin Inc) connected to a custom-

made LED driver, as described previously[80]. This resulted in a light output of 1-2.5 mW at the tip of

the optic fiber implants. LEDs were controlled by TTL pulses from the amplifier and these pulses were

used to synchronize electrophysiology and camera recordings. Silver conductive paint and aluminum

foil coating was used to reduce the escape of optogenetic light from the fibers and equipment. The

AUC for the optogenetically-evoked eyelid responses was calculated as the absolute sum of the

normalized-to-baseline eyelid trace 0-150 ms after LED onset. Optogenetic stimulation of

ChR2(H134R) in Gabra6-Cre::ChR2EYFP mice was achieved using a 465 nm 5W LED (60 lm, LZ1-B200,

LED Engin Inc) mounted on a moveable arm. After establishing a whole-cell recording, the LED was

positioned close to the cerebellum to activate granule cells.

1256

Histological procedures

To examine fiber placement after MF-ChR2 experiments and to verify hM4Di-DREADD expression,

mice were deeply anesthetized using a ketamine-xylazine mixture and perfused transcardially with 4%

paraformaldehyde and their brains were removed. Sagittal or coronal sections (50 μm thick) were cut

using a vibratome. Slices from the DREADD transgenic mice were stained with a rabbit monoclonal

anti-HA antibody (1:400, Cell Signaling, #3724) and DAPI (abcam, #ab104139). Slices were then

mounted on glass slides with Mowiol mounting medium (Merck Millipore Calbiochem). For all other

histological quantifications, mice were deeply anesthetized with pentobarbital (Nembutal) and

transcardially perfused with 100 ml 4% paraformaldehyde (PFA) in 0.12 M phosphate buffer. Brains

were dissected and post-fixed overnight at 4°C, then cryoprotected in 0.12 M phosphate buffer

containing 30% sucrose at 4°C until they sank. Cerebella were cut on a cryostat into 25 μm-thick

coronal or sagittal floating sections and collected in phosphate-buffered saline (PBS). Primary

antibodies were incubated overnight at 4°C in PBS containing 0.25% Triton X-100, and 5% fetal calf

serum. Primary antibodies were: mouse anti-NeuN (1:500, Millipore), rabbit anti-GFP, which also

detects YFP (1:1000, Chemicon), mouse anti-calbindin (1:1500, Swant), guinea-pig anti-VGLUT1

(1:500, Synaptic Systems), rabbit anti-VGLUT1 (1:1000, Synaptic Systems), guinea pig anti-VGLUT2

(1:1000, Synaptic Systems) and mouse anti-gephyrin (1:500, Synaptic Systems). After washing,

sections were then incubated for 1 h at room temperature with one of the following fluorophore-

conjugated secondary antibodies or streptavidin: donkey anti-mouse Cy3 (ThermoFisher Scientific),

goat anti-chicken Alexa Fluor 488 (ThermoFisher Scientific), donkey anti-guinea pig Cy3 (ThermoFisher

Scientific), donkey anti-rabbit Alexa Fluor 647 (ThermoFisher Scientific), donkey anti-rabbit Alexa Fluor

488 (ThermoFisher Scientific), streptavidin-488 (ThermoFisher Scientific) or streptavidin-Cy3 (Jackson

Immunoresearch). For each immunohistochemical reaction, slices from all experimental conditions

were processed together and incubation times were kept constant. After processing sections were

mounted on microscope slides with Tris-glycerol supplemented with 10% Mowiol (Merck Millipore

Calbiochem).

1283

Imaging

Fluorescent images were acquired using a confocal microscope (SP5 and SP8, Leica Microsystems) or

an upright Zeiss Axio Imager M2 microscope using a 10x objective. Confocal images were taken at a

resolution of 1024 x 1024 dpi and a 50 Hz speed. Laser intensity, gain and offset were maintained

constant in each analysis. Quantitative evaluations were made by a blind experimenter using Image J

software. Adobe Photoshop 6.0 (Adobe Systems) was used to adjust image contrast and assemble the

final plates.

1291

Quantification of density of glutamatergic terminals

To estimate the density of VGLUT1 and VGLUT2-positive axon terminals, at least three coronal sections

containing the AIP (Bregma: -6.00, -6.12 mm) were selected for each animal. In each section three 1

μm-thick confocal images/side were captured under a 63x objective. A "maximum intensity" function

was applied to the Z-projection and the ''analyze particle'' function was used to estimate the density

of boutons (number of terminals/mm^2), after selecting the automatic threshold and using the

"watershed" function. Only particles with a size between 0.4 um^2 and 3 um^2 were included in the

analysis. The average/animal in the ipsilateral and contralateral side was calculated (VGLUT1:

conditioned mice, N = 11; pseudo-conditioned mice, N = 13; VGLUT2: conditioned mice, N = 4; pseudo-

conditioned mice, N = 6). To correct for unwanted differences in different batches of mice, values were

normalized to the values of the pseudo-conditioned group in each batch of mice.

1303

Quantification of density of axonal varicosities

The density of axonal varicosities in the AIP was evaluated in the images used for VGLUT1

quantifications (see paragraph above) where YFP+ axons could be detected by measuring in each

image the length of individual YFP+ axonal segments and the number of enlargements on each

segment (conditioned mice: N = 3, n = 322 axonal segments; pseudo-conditioned mice: N = 3, n = 381

axonal segments). The density for each side was calculated by dividing the total number of varicosities

by the total length.

1311

Quantification of gephyrin-positive puncta

Single 0.4 μm thick-confocal images of the AIP (ipsilateral and contralateral side) in coronal sections

(Bregma: -6.00, -6.12 mm) were collected under a 63x objective with 2.5x zoom. At least three sections

were selected/mouse. On such images the density of gephyrin-positive puncta around the soma of

excitatory neurons (identified by their size, *i.e.* > 240 μm[33]) was evaluated by Fiji and expressed as

number/μm neuronal membrane. Gephyrin-positive puncta were manually identified, as they were

clearly visible as discreet fluorescent puncta. At least 15 neurons/mouse were quantified (conditioned

mice: N = 4; n = 84; pseudo-conditioned mice: N = 6, n = 110). For each neuron the number of gephyrin-

positive puncta apposed to calbindin-positive terminals was quantified and the % of those puncta (out

of the total number of puncta) was calculated. To correct for unwanted differences in different

batches of mice, gephyrin densities were normalized to the values of the pseudo-conditioned group

in each batch of mice.

1324

Quantification of size of rosettes

Pictures including YFP+ rosettes in the eyeblink area (i.e. at the bottom of the primary fissure)

consisted of thirteen 0.5 um-thick confocal steps acquired under a 63x objective. Three pictures were

taken to cover the eyeblink area in each side. The "maximum intensity" function in Fiji was applied to

the Z-projection and the area of individual rosettes was quantified (conditioned mice: N = 5, n = 613;

pseudo-conditioned mice: N = 4, n = 761).

1331

Decoding analysis

Linear discriminant analysis (10 fold cross-validation) was performed on pooled trial data with a

varying number of trials for each neuron (cond: n = 199 CR and n = 265 NR trials in 15 neurons; naive:

n = 113 EO and n = 214 NR trials in 15 neurons). For whole-cell data decoding was performed on spike-

removed, normalized and Gaussian filtered (10 ms cut-off) V_m traces. For spike data, spikes were

binned into 5 ms time intervals, Gaussian filtered (10 ms cut-off) and normalized to baseline rate. The

dataset was then divided in a train and test set (9:1 ratio) and decoding was performed using the

'*classify*' function in MATLAB (custom MATLAB code provided by M. Tang, Australian National

University, 2022). Accuracy represented the percentage of accurately classified trials for each

sampling point or time bin. To test whether the decoder performed above chance level, we performed

a cluster-corrected permutation test with 2000 permutations in MATLAB (E. Spaak, Oxford University,

2015). To calculate decoding accuracy for decreasing group sizes, we performed 30 iterations of the

decoding analysis where in each iteration a random subsample of the total dataset was selected using

the '*randsample*' function in MATLAB. Quantifications and traces for each group size were calculated

by averaging across 30 iterations. Time period in which above-average decoding took place was

calculated as the average time the cluster-test was significant.

1348

Statistics

Statistical analysis was carried out using the Statistics toolbox of MATLAB (R2011b and R2022a,

MathWorks, CA, USA), SPSS (v22, IBM, NY, USA), GraphPad Prism 5 and 9 (GraphPad Software Inc.,

CA, USA). Normality of distributions was assessed using Shapiro-Wilk test. Homogeneity of variance

was assessed using Bartlett's test. If the assumption of normality was violated, non-parametric

alternatives were performed. For the correlation between walking speed vs. onset session we used

linear regression analysis and a unpaired t-test. To compare average peak timings for the different ISIs

we used a unpaired t-test. For MF-ChR2 experiments, mice were randomly assigned to specific

experimental groups without bias and no animals were excluded. Behavioral performance was

analyzed using a repeated-measures ANOVA or linear mixed models with the maximum likelihood

method. Group and session/day was modelled as fixed effects, and %CR, CR amplitude and CR

amplitude at US onset were modelled as dependent variables. We assessed the fit of the model by

running the analysis with the unstructured, diagonal and first-order autoregressive repeated

covariance types, after which we choose the covariance type with the lowest Akaike's information

criterion (AIC) value. Correlations between eyelid and V_m or spike data were performed using

Spearman correlation, where data were normalized to the maximum value per neurons. Differences

in correlations were tested using the Fisher's Z-transformation (J. Decoster & A-M. Leistico, University

of Virginia, 2005). Distributions of CS responses between conditioned and naive mice were tested

using the chi-square test. Histological data were analyzed using a paired t-test (when comparing

ipsilateral and contralateral sides), unpaired t-test (when comparing conditioned with pseudo-

conditioned mice), Pearson correlation analysis and Kolmogorov-Smirnov test (when comparing

cumulative frequencies). Data are reported as mean ± standard error of the mean (SEM), unless

indicated otherwise. Statistical power was calculated using Gpower (version 3.1.9.4). Data are shown

in figures as mean ± SEM., unless indicated otherwise. Differences were considered significant at $*P <$

0.05, $**P < 0.01$ and $***P < 0.001$.

1374

1375   **Data availability**

1376   A source data file supporting the findings of this study will be provided upon acceptance of the

1377   manuscript. Data that support the findings of this study are available from the corresponding

1378   authors upon reasonable request.

1379

1380   **Code availability**

1381   All MATLAB code used to generate the results will be uploaded to GitHub and be made freely

1382   available upon acceptance of the manuscript.

References

1. Gormezano, I., Schneiderman, N., Deaux, E. & Fuentes, I. Nictitating Membrane: Classical Conditioning and Extinction in the Albino Rabbit. *Science* **138**, 33–34 (1962).
2. Schneiderman, N., Fuentes, I. & Gormezano, I. Acquisition and Extinction of the Classically Conditioned Eyelid Response in the Albino Rabbit. *Science* **136**, 650–652 (1962).
3. Steinmetz, J. E., Rosen, D. J., Chapman, P. F., Lavond, D. G. & Thompson, R. F. Classical Conditioning of the Rabbit Eyelid Response With a Mossy-Fiber Stimulation CS: I. Pontine Nuclei and Middle Cerebellar Peduncle Stimulation. *Behavioral Neuroscience* **100**, 878–887 (1986).
4. Hesslow, G., Svensson, P. & Ivarsson, M. Learned movements elicited by direct stimulation of cerebellar mossy fiber afferents. *Neuron* **24**, 179–185 (1999).
5. Silva, N. T., Ramírez-Buriticá, J., Pritchett, D. L. & Carey, M. R. Neural instructive signals for associative cerebellar learning. 2022.04.18.488634 Preprint at https://doi.org/10.1101/2022.04.18.488634 (2022).
6. Sillitoe, R. V., Fu, Y. & Watson, C. Chapter 11 - Cerebellum. in *The Mouse Nervous System* (eds. Watson, C., Paxinos, G. & Puelles, L.) 360–397 (Academic Press, 2012). doi:10.1016/B978-0-12-369497-3.10011-1.
7. Luo, Y. & Sugihara, I. Cerebellar afferents originating from the medullary reticular formation that are different from mossy, climbing or monoaminergic fibers in the rat. *Brain Research* **1566**, 31–46 (2014).
8. Wu, H. S., Sugihara, I. & Shinoda, Y. Projection patterns of single mossy fibers originating from the lateral reticular nucleus in the rat cerebellar cortex and nuclei. *The Journal of comparative neurology* **411**, 97–118 (1999).
9. Beitzel, C. S., Houck, B. D., Lewis, S. M. & Person, A. L. Rubrocerebellar Feedback Loop Isolates the Interposed Nucleus as an Independent Processor of Corollary Discharge Information in Mice. *J. Neurosci.* **37**, 10085–10096 (2017).
10. Shinoda, Y., Sugiuchi, Y., Futami, T. & Izawa, R. Axon collaterals of mossy fibers from the pontine nucleus in the cerebellar dentate nucleus. *Journal of Neurophysiology* **67**, 547–560 (1992).
11. Ekerot, C.-F. & Kano, M. Long-term depression of parallel fibre synapses following stimulation of climbing fibres. *Brain Research* **342**, 357–360 (1985).
12. Marr, D. A theory of cerebellar cortex. *J. Physiol.* **202**, 437–470 (1969).
13. Schonewille, M. *et al.* Purkinje Cell-Specific Knockout of the Protein Phosphatase PP2B Impairs Potentiation and Cerebellar Motor Learning. *Neuron* **67**, 618–628 (2010).
14. Ten Brinke, M. M. *et al.* Evolving Models of Pavlovian Conditioning: Cerebellar Cortical Dynamics in Awake Behaving Mice. *Cell Reports* **13**, 1977–1988 (2015).
15. Boele, H.-J. *et al.* Impact of parallel fiber to Purkinje cell long-term depression is unmasked in absence of inhibitory input. *SCIENCE ADVANCES* 9 (2018).
16. Johansson, F., Carlsson, H. A. E., Rasmussen, A., Yeo, C. H. & Hesslow, G. Activation of a Temporal Memory in Purkinje Cells by the mGluR7 Receptor. *Cell Reports* **13**, 1741–1746 (2015).
17. Gao, Z. *et al.* Excitatory Cerebellar Nucleocortical Circuit Provides Internal Amplification during Associative Conditioning. *Neuron* **89**, 645–657 (2016).

18. Boele, H.-J., Koekkoek, S. K. E., De Zeeuw, C. I. & Ruigrok, T. J. H. Axonal Sprouting and
Formation of Terminals in the Adult Cerebellum during Associative Motor Learning.
Journal of Neuroscience **33**, 17897–17907 (2013).
19. Jirenhed, D.-A., Bengtsson, F. & Hesslow, G. Acquisition, Extinction, and Reacquisition of
a Cerebellar Cortical Memory Trace. *Journal of Neuroscience* **27**, 2493–2502 (2007).
20. Berthier, N. E. & Moore, J. W. Cerebellar Purkinje cell activity related to the classically
conditioned nictitating membrane response. *Experimental brain research* **63**, 341–350
(1986).
21. McCormick, D. A. & Thompson, R. F. Neuronal responses of the rabbit cerebellum during
acquisition and performance of a classically conditioned nictitating membrane-eyelid
response. *The Journal of neuroscience* **4**, 2811–2822 (1984).
22. Ten Brinke, M. M. *et al.* Dynamic modulation of activity in cerebellar nuclei neurons
during pavlovian eyeblink conditioning in mice. *eLife* **6**, (2017).
23. Morcuende, S., Delgado-García, J.-M. & Ugolini, G. Neuronal Premotor Networks
Involved in Eyelid Responses: Retrograde Transneuronal Tracing with Rabies Virus from
the Orbicularis Oculi Muscle in the Rat. *J Neurosci* **22**, 8808–8818 (2002).
24. Pacheco-Calderón, R., Carretero-Guillén, A., Delgado-García, J. M. & Gruart, A. Red
Nucleus Neurons Actively Contribute to the Acquisition of Classically Conditioned Eyelid
Responses in Rabbits. *J. Neurosci.* **32**, 12129–12143 (2012).
25. Albergaria, C., Silva, N. T., Pritchett, D. L. & Carey, M. R. Locomotor activity modulates
associative learning in mouse cerebellum. *Nature Neuroscience* **21**, 725–735 (2018).
26. Armbruster, B. N., Li, X., Pausch, M. H., Herlitze, S. & Roth, B. L. Evolving the lock to fit
the key to create a family of G protein-coupled receptors potently activated by an inert
ligand. *Proceedings of the National Academy of Sciences* **104**, 5163–5168 (2007).
27. Roth, B. L. DREADDs for Neuroscientists. *Neuron* **89**, 683–694 (2016).
28. Koekkoek, S. K. E. Cerebellar LTD and Learning-Dependent Timing of Conditioned Eyelid
Responses. *Science* **301**, 1736–1739 (2003).
29. Hioki, H. *et al.* Differential distribution of vesicular glutamate transporters in the rat
cerebellar cortex. *Neuroscience* **117**, 1–6 (2003).
30. Geisler, S., Derst, C., Veh, R. W. & Zahm, D. S. Glutamatergic Afferents of the Ventral
Tegmental Area in the Rat. *J. Neurosci.* **27**, 5730–5743 (2007).
31. Leergaard, T. B. & Bjaarlie, J. G. Topography of the complete corticopontine projection:
From experiments to principal maps. *Frontiers in Neuroscience* **1**, 211–223 (2007).
32. Uusisaari, M. & Knöpfel, T. Functional Classification of Neurons in the Mouse Lateral
Cerebellar Nuclei. *The Cerebellum* **10**, 637–646 (2011).
33. Uusisaari, M., Obata, K. & Knopfel, T. Morphological and Electrophysiological Properties
of GABAergic and Non-GABAergic Cells in the Deep Cerebellar Nuclei. *Journal of
Neurophysiology* **97**, 901–911 (2007).
34. Krueger-Burg, D., Papadopoulos, T. & Brose, N. Organizers of inhibitory synapses come
of age. *Current Opinion in Neurobiology* **45**, 66–77 (2017).
35. Tyagarajan, S. K. & Fritschy, J.-M. Gephyrin: a master regulator of neuronal function?
Nat Rev Neurosci **15**, 141–156 (2014).
36. Celio, M. R. Calbindin D-28k and parvalbumin in the rat nervous system. *Neuroscience*
35, 375–475 (1990).
37. Scotti, A. L. Calbindin D28k in the olivocerebellar projection. A light and electron
microscope study. *J Anat* **187**, 649–659 (1995).

38. Berthier, N. E. & Moore, J. W. Activity of deep cerebellar nuclear cells during classical
conditioning of nictitating membrane extension in rabbits. *Experimental brain research*
83, 44–54 (1990).
39. Uusisaari, M. & De Schutter, E. The mysterious microcircuitry of the cerebellar nuclei.
The Journal of Physiology **589**, 3441–3457 (2011).
40. Canto, C. B., Witter, L. & De Zeeuw, C. I. Whole-Cell Properties of Cerebellar Nuclei
Neurons In Vivo. *PloS one* **11**, e0165887 (2016).
41. Yarden-Rabinowitz, Y. & Yarom, Y. *In vivo* analysis of synaptic activity in cerebellar nuclei
neurons unravels the efficacy of excitatory inputs. *The Journal of Physiology* (2017)
doi:10.1113/JP274115.
42. Heiney, S. A., Wojaczynski, G. J. & Medina, J. F. Action-based organization of a cerebellar
module specialized for predictive control of multiple body parts. *Neuron* **109**, 2981-
2994.e5 (2021).
43. Gong, S., DeCuypere, M., Zhao, Y. & LeDoux, M. S. Cerebral cortical control of orbicularis
oculi motoneurons. *Brain Research* **1047**, 177–193 (2005).
44. Suzuki, Y., Kiyosawa, M., Mochizuki, M., Ishiwata, K. & Ishii, K. The Pre-Supplementary
and Primary Motor Areas Generate Rhythm for Voluntary Eye Opening and Closing
Movements. *Tohoku J. Exp. Med.* **222**, 97–104 (2010).
45. Freeman, J. H. & Steinmetz, A. B. Neural circuitry and plasticity mechanisms underlying
delay eyeblink conditioning. *Learning & Memory* **18**, 666–677 (2011).
46. De Zeeuw, C. I., Van Alphen, A. M., Hawkins, R. K. & Ruigrok, T. J. H. Climbing fiber
collaterals contact neurons in the cerebellar nuclei that provide a GABAergic feedback to
the inferior olive. *Neuroscience* **80**, 981–986 (1997).
47. Nicholson, D. A. & Freeman, J. H., Jr. Selective Developmental Increase in the Climbing
Fiber Input to the Cerebellar Interpositus Nucleus in Rats. *Behavioral Neuroscience* **118**,
1111–1116 (2004).
48. Bracha, V., Webster, M. L., Winters, N. K., Irwin, K. B. & Bloedel, J. R. Effects of muscimol
inactivation of the cerebellar interposed-dentate nuclear complex on the performance
of the nictitating membrane response in the rabbit. *Experimental brain research* **100**,
453–468 (1994).
49. Clark, G. A., McCormick, D. A., Lavond, D. G. & Thompson, R. F. Effects of lesions of
cerebellar nuclei on conditioned behavior and hippocampal neuronal responses. *Brain*
Research **291**, 125–136 (1984).
50. Lavond, D. G. & Steinmetz, J. E. Acquisition of classical conditioning without cerebellar
cortex. *Behavioral Brain Research* **33**, 113–164 (1989).
51. Lavond, D. G., Hembree, T. L. & Thompson, R. F. Effect of kainic acid lesions of the
cerebellar interpositus nucleus on eyelid conditioning in the rabbit. *Brain Research* **326**,
179–182 (1985).
52. Ohyama, T., Nores, W. L., Medina, J. F., Riusech, F. A. & Mauk, M. D. Learning-Induced
Plasticity in Deep Cerebellar Nucleus. *Journal of Neuroscience* **26**, 12656–12663 (2006).
53. Yeo, C. H., Hardiman, M. J. & Glickstein, M. Classical conditioning of the nictating
membrane of the rabbit. I. Lesions of the cerebellar nuclei. *Experimental Brain Research*
60, 87–98 (1985).
54. Ohyama, T., Nores, W. L. & Mauk, M. D. Stimulus generalization of conditioned eyelid
responses produced without cerebellar cortex: implications for plasticity in the
cerebellar nuclei. *Learning & Memory* **10**, 346–354 (2003).

55. Steinmetz, A. B. & Freeman, J. H. Localization of the cerebellar cortical zone mediating acquisition of eyeblink conditioning in rats. *Neurobiology of Learning and Memory* **114**, 148–154 (2014).

56. Medina, J. F., Nores, W. L., Ohyama, T. & Mauk, M. D. Mechanisms of cerebellar learning suggested by eyelid conditioning. *Current opinion in neurobiology* **10**, 717–724 (2000).

57. Uusisaari, M. Y. & Knöpfel, T. Diversity of Neuronal Elements and Circuitry in the Cerebellar Nuclei. *The Cerebellum* **11**, 420–421 (2012).

58. Ivarsson, M. & Hesslow, G. Bilateral control of the orbicularis oculi muscle by one cerebellar hemisphere in the ferret. *Neuroreport* **4**, 1127–1130 (1993).

59. Medina, J. F., Garcia, K. S. & Mauk, M. D. A Mechanism for Savings in the Cerebellum. *The Journal of Neuroscience* **21**, 4081–4089 (2001).

60. Weeks, A. C. W. *et al.* Eye-blink conditioning is associated with changes in synaptic ultrastructure in the rabbit interpositus nuclei. *Learning & Memory* **14**, 385–389 (2007).

61. Wang, D. *et al.* Changes in membrane properties of rat deep cerebellar nuclear projection neurons during acquisition of eyeblink conditioning. *Proceedings of the National Academy of Sciences* **115**, E9419–E9428 (2018).

62. Carulli, D. *et al.* Cerebellar plasticity and associative memories are controlled by perineuronal nets. *Proc Natl Acad Sci USA* **117**, 6855–6865 (2020).

63. Perrett, S. P., Ruiz, B. P. & Mauk, M. D. Cerebellar cortex lesions disrupt learning-dependent timing of conditioned eyelid responses. *The Journal of Neuroscience* **13**, 1708–1718 (1993).

64. Thompson, R. F. The neurobiology of learning and memory. *Science* **233**, 941–947 (1986).

65. Chapman, P. F., Steinmetz, J. E., Sears, L. L. & Thompson, R. F. Effects of lidocaine injection in the interpositus nucleus and red nucleus on conditioned behavioral and neuronal responses. *Brain Research* **537**, 149–156 (1990).

66. Steinmetz, J., Lavond, D., Ivkovich, D., Logan, C. & Thompson, R. Disruption of classical eyelid conditioning after cerebellar lesions: damage to a memory trace system or a simple performance deficit? *J. Neurosci.* **12**, 4403–4426 (1992).

67. Clark, R. E., Zhang, A. A. & Lavond, D. G. Reversible lesions of the cerebellar interpositus nucleus during acquisition and retention of a classically conditioned behavior. *Behavioral Neuroscience* **106**, 879–888 (1992).

68. Krupa, D. J. & Thompson, R. F. Reversible inactivation of the cerebellar interpositus nucleus completely prevents acquisition of the classically conditioned eye-blink response. *Learning & Memory* **3**, 545–556 (1997).

69. Garcia, K. S. & Mauk, M. D. Pharmacological analysis of cerebellar contributions to the timing and expression of conditioned eyelid responses. *Neuropharmacology* **37**, 471–480 (1998).

70. Aksenov, D. P., Serdyunkova, N., Irwin, K. & Bracha, V. GABA Neurotransmission in the Cerebellar Interposed Nuclei: Involvement in Classically Conditioned Eyeblinks and Neuronal Activity. *Journal of Neurophysiology* **91**, 719–727 (2004).

71. Heiney, S. A., Wohl, M. P., Chettih, S. N., Ruffolo, L. I. & Medina, J. F. Cerebellar-Dependent Expression of Motor Learning during Eyeblink Conditioning in Head-Fixed Mice. *Journal of Neuroscience* **34**, 14845–14853 (2014).

72. Raymond, J. L., Lisberger, S. G. & Mauk, M. D. The Cerebellum: A Neuronal Learning Machine? *Science, New Series* **272**, 1126–1131 (1996).

73. Zheng, N. & Raman, I. M. Synaptic Inhibition, Excitation, and Plasticity in Neurons of the
Cerebellar Nuclei. *The Cerebellum* **9**, 56–66 (2010).
74. Arenkiel, B. R. *et al.* In Vivo Light-Induced Activation of Neural Circuitry in Transgenic
Mice Expressing Channelrhodopsin-2. *Neuron* **54**, 205–218 (2007).
75. Fünfschilling, U. & Reichardt, L. F. Cre-mediated recombination in rhombic lip
derivatives. *genesis* **33**, 160–169 (2002).
76. Zhu, H. *et al.* Cre-dependent DREADD (Designer Receptors Exclusively Activated by
Designer Drugs) mice. *Genesis* **54**, 439–446 (2016).
77. Madisen, L. *et al.* A toolbox of Cre-dependent optogenetic transgenic mice for light-
induced activation and silencing. *Nat Neurosci* **15**, 793–802 (2012).
78. Ohmae, S. & Medina, J. F. Climbing fibers encode a temporal-difference prediction error
during cerebellar learning in mice. *Nature Neuroscience* **18**, 1798–1803 (2015).
79. Koekkoek, S. K. E., Den Ouden, W. L., Perry, G., Highstein, S. M. & De Zeeuw, C. I.
Monitoring kinetic and frequency-domain properties of eyelid responses in mice with
magnetic distance measurement technique. *Journal of neurophysiology* **88**, 2124–2133
(2002).
80. Witter, L., Canto, C. B., Hoogland, T. M., de Gruijl, J. R. & De Zeeuw, C. I. Strength and
timing of motor responses mediated by rebound firing in the cerebellar nuclei after
Purkinje cell activation. *Frontiers in Neural Circuits* **7**, (2013).
1585