Mutations in *NAKED-ENDOSPERM* IDD genes reveal functional interactions with *SCARECROW* and a maternal influence on leaf patterning in C$_4$ grasses

Thomas E. Hughes[1], Olga Sedelnikova[1,2], Mimi Thomas[1] & Jane A. Langdale[1*]

[1]Department of Biology, University of Oxford, South Parks Road, Oxford OX13RB
[2]Current address: Syngenta Jealott's Hill International Research Centre, Bracknell RG42 6EY

Running title: SCR and NKD interactions in grass leaves

Summary statement: Mutations in *NKD* IDD genes enhance loss of function *scr* phenotypes in the leaves of C$_4$ grasses maize and *Setaria viridis* but not in the C$_3$ grass rice, and reveal a maternal effect on cell-type patterning in leaves that are initiated during embryogenesis.

Key words: patterning, SCARECROW, NAKED-ENDOSPERM, development, veins

* Corresponding author: jane.langdale@biology.ox.ac.uk

ABSTRACT

Leaves comprise a number of different cell-types that are patterned in the context of either the epidermal or inner cell layers. In grass leaves, two distinct anatomies develop in the inner leaf tissues depending on whether the leaf carries out C_3 or C_4 photosynthesis. In both cases a series of parallel veins develops that extends from the leaf base to the tip but in ancestral C_3 species veins are separated by a greater number of intervening mesophyll cells than in derived C_4 species. We have previously demonstrated that the GRAS transcription factor SCARECROW (SCR) regulates the number of photosynthetic mesophyll cells that form between veins in the leaves of the C_4 species maize, whereas it regulates the formation of stomata in the epidermal leaf layer in the C_3 species rice. Here we show that SCR is required for inner leaf patterning in the C_4 species *Setaria viridis* but in this species the presumed ancestral stomatal patterning role is also retained. Through a comparative mutant analysis between maize, setaria and rice we further demonstrate that loss of NAKED-ENDOSPERM (NKD) INDETERMINATE DOMAIN (IDD) protein function exacerbates loss of function *scr* phenotypes in the inner leaf tissues of maize and setaria but not rice. Specifically, in both setaria and maize, *scr;nkd* mutants exhibit an increased proportion of fused veins with no intervening mesophyll cells. Thus, combined action of SCR and NKD may control how many mesophyll cells are specified between veins in the leaves of C_4 but not C_3 grasses. Finally, we identified a maternal effect in maize whereby maternally derived NKD can affect patterning of cells in leaf primordia that are initiated during embryogenesis. Together our results provide insight into the evolution of cell patterning in grass leaves, demonstrate a novel patterning role for IDD genes in C_4 leaves and suggest that NKD can influence embryonic leaf development non-cell autonomously from the surrounding maternal tissue.

INTRODUCTION

Understanding how cell patterning is genetically regulated is a key challenge in developmental biology. In grass leaves, two distinct cellular anatomies underpin photosynthesis. In grasses such as rice (*Oryza sativa*) that carry out C_3 photosynthesis, widely spaced parallel veins are encircled by non-photosynthetic bundle-sheath (BS) cells which are themselves separated by up to ten photosynthetic mesophyll (M) cells. By contrast, in grasses that perform C_4 photosynthesis, such as maize (*Zea mays*) and green millet (*Setaria viridis*), parallel veins are surrounded by concentric layers of BS and M cells, both of which are photosynthetic. This arrangement of cell-types is referred to as 'Kranz' because the two cell-types form wreaths around the veins and Kranz is German for wreath (Haberlandt, 1896). Notably Kranz anatomy evolved from C_3-type anatomy on multiple independent occasions (Grass Phylogeny Working Group II, 2012; Sage et al., 2011), each occurrence generating leaves with higher vein densities than in the ancestral form because BS cells are separated by fewer M cells. To date, very few regulators of cell-patterning in inner leaf tissues have been identified in C_3 or C_4 grass species.

We previously demonstrated that duplicate genes encoding the GRAS transcription factor SCARECROW (SCR) regulate cell divisions in the innermost ground meristem layer of maize leaf

primordia to determine the number of M cells that form between veins (Hughes et al., 2019). In double *Zmscr1;Zmscr1h* mutants, the majority of BS cells are separated by one rather than two M cells, and in some cases BS cells are fused with no intervening M cells. In addition, many veins develop ectopic sclerenchyma either ad- or abaxially and some veins are surrounded by additional BS cells that are not in contact with the vasculature. Intriguingly, no such patterning perturbations were observed in inner leaf tissues when *SCR* orthologs were mutated in rice (Hughes & Langdale, 2022). Instead, loss of function mutants in rice fail to develop stomata on the leaf surface (Hughes & Langdale, 2022; Wu et al., 2019). The distinction between mutant phenotypes in maize and rice leaves raises the possibility that the deployment of SCR function in inner leaf tissues was associated with the evolution of Kranz anatomy, with the patterning of stomata in the epidermis being the ancestral role in grass leaves.

Although the function of SCR itself could differ between maize and rice, for example by targeting different downstream genes, the distinct patterning roles observed could alternatively result from species-specific differences in SCR interacting proteins. Little is known about how SCR-mediated patterning is regulated in monocots but in Arabidopsis roots, SCR functions with another GRAS transcription factor (SHORTROOT (SHR)) and with several INDETERMINATE DOMAIN (IDD) C2H2 zinc-finger transcription factors (Laurenzio et al., 1996; Long et al., 2015; Nakajima et al., 2001; Ogasawara et al., 2011; Welch et al., 2007). The IDD genes (also referred to as BIRD genes), act both to modulate *SCR* and *SHR* gene expression and to co-regulate the expression of downstream target genes (Welch et al., 2007). Despite the existence of many IDD genes in grass genomes, and the demonstration that *SCR* and *SHR* have patterning functions in maize and rice, links between *SCR* and IDD genes in the patterning of either root or leaf cell-types in monocots have not yet been established.

We previously proposed that the *NAKED-ENDOSPERM* (*NKD*) *IDD* genes (originally referred to as *ZmJAY* genes) may function during Kranz patterning in maize (Fouracre et al., 2014; Sedelnikova et al., 2018). This proposal was based on the fact that *ZmNKD1* and *ZmNKD2* transcripts accumulate during early leaf development at the relevant stages of Kranz patterning (Wang et al., 2013), both transcripts accumulate specifically in mesophylls of mature leaves (Chang et al., 2012; Li et al., 2010) and both have known patterning functions in aleurone development (Gontarek et al., 2016; Yi et al., 2015). Building on this suggestion and on the outstanding questions raised above, here we have tested two hypotheses. First, that SCR function is required for patterning inner leaf tissues in C_4 grasses and second that NKD is a component of the patterning pathway. By characterizing *scr* mutants in the C_4 species *Setaria viridis* and carrying out a comparative analysis of *scr;nkd* mutants in maize, *Setaria viridis* and rice, we found evidence to support both hypotheses. In addition, we identified a maternal effect of NKD on the patterning of leaves formed during embryogenesis in maize.

RESULTS

SCR patterns both epidermal and inner tissues in leaves of *Setaria viridis*

To determine whether the distinct patterning roles of SCR in the rice epidermis versus the maize inner leaf reflect C_3- versus C_4-specific functions, we generated loss-of-function mutants in the C_4 species *Setaria viridis* (hereafter referred to as setaria) using CRISPR (Figure 1A, Figure S1). As in maize and rice, two *SCR* genes are present in setaria (*SvSCR1*- Sevir.7G316501 and *SvSCR2*- Sevir.8G008100), each pair the result of a within species duplication (Hughes et al., 2019). In gene edited T2 lines, a proportion of plants were identified that were extremely stunted, paler than wild-type and did not survive beyond 3-4 weeks after germination (Figure 1B-D). These plants appeared at the expected segregation ratios of 1/4 or 1/16 depending on whether the parental plant was *Svscr1/+;Svscr2* (or *Svscr1;Svscr2/+*) or *Svscr1/+;Svscr2/+* respectively. In all cases, these phenotypically abnormal plants were confirmed to be homozygous for both *Svscr1* and *Svscr2*, with phenotypically wild-type plants always being heterozygous or wild-type for one of the *SCR* genes. Because single *Svscr* mutants displayed no growth perturbations (Figure S2) and single mutants in both rice and maize were not associated with any patterning defects, all further analyses were undertaken with homozygous double mutants. The perturbed growth phenotype exhibited by *Svscr1;Svscr2* plants was far more severe than that seen in maize *Zmscr1;Zmscr1h* mutants (where plants can be grown for 6-8 weeks in the greenhouse without issue) (Hughes et al., 2019), but was similar to the phenotype observed in *Osscr1;Osscr2* mutants of rice (Hughes & Langdale, 2022). We therefore reasoned that, as in rice, *SCR* may play a role in stomatal patterning in setaria. Epidermal impressions of *Svscr1;Svscr2* mutant leaves confirmed this hypothesis, revealing an absence of stomata on both the abaxial and adaxial leaf surfaces (Fig. 2A-F). Therefore, the stomatal patterning role first identified in rice is not a C_3-specific function.

To determine whether *SCR* also plays a role in inner leaf patterning in setaria, transverse sections of wild-type and *Svscr1;Svscr2* mutant leaves were examined (Fig. 2G, H). In ME034V wild-type leaves, around 70-80% of veins were separated by two M cells with the rest separated by three (a feature that is more common in setaria than in maize) (Fig. 2I). In contrast, 20-30% of veins in *Svscr1;Svscr2* mutant leaves were separated by just a single M cell (Fig. 2I). These data demonstrate that, as in maize, *SvSCR* genes regulate cell divisions in the ground meristem to determine how many M cells develop between BS cells. Intriguingly, *SvSCR* genes undertake this inner leaf patterning role in addition to a role in stomatal patterning.

A feedback loop links *SCR* and *NKD* IDD gene expression in maize

The distinct roles of *SCR* genes in maize, rice and setaria – patterning inner, epidermal or both tissue layers of the leaf respectively – could be due to species-specific differences in the activity of interacting IDD genes, of which *NKD* genes are potential candidates. To determine whether the expression of *ZmSCR1/h* genes is interconnected with *ZmNKD1/2* gene function in maize (as is the case for *SCR* and IDD genes in Arabidopsis), we first quantified transcript levels of each gene in existing double

Zmscr1-m2;Zmscr1h-m1 (Hughes et al., 2019) and double *Zmnkd1-Ds;Zmnkd2-Ds* (Yi et al., 2015) mutants (Fig. 3A, Figure S3). *ZmSCR1* and *ZmSCR1h* transcripts accumulated at elevated levels in the *Zmnkd1-Ds;Zmnkd2-Ds* mutant, in both cases being increased around two-fold relative to wild-type (Fig. 3C). In contrast, *ZmNKD1* and *ZmNKD2* transcripts accumulated at lower levels in *Zmscr1-m2;Zmscr1h-m1* mutants than in wild-type, with *ZmNKD1* reduced on average to 60% and *ZmNKD2* to 50% of wild-type levels (Fig. 3A). To determine whether the interactions between NKD and SCR are likely to occur within or between cell-types, *in situ* hybridization was carried out to localize transcripts in developing leaf primordia. Figure 3B shows that both *NKD1* and *SCR1* transcripts accumulate in the ground meristem cells that surround developing vascular centres. Together these data suggest that a negative feedback loop controls *ZmSCR1/1h* and *ZmNKD1/2* transcript levels, most likely in a cell-autonomous manner, with ZmSCR1/ZmSCR1h promoting the expression of *ZmNKD1/ZmNKD2*, and ZmNKD1/ZmNKD2 repressing the expression of *ZmSCR1/ZmSCR1h*.

Loss of function *nkd* mutants do not exhibit perturbed leaf development

Given the patterning role of SCR in leaves of maize, setaria and rice, and the interaction between *SCR* and *NKD* gene expression in maize, we next sought to determine the role of *NKD* in leaf development in each of the three species. To this end, we first identified *NKD* orthologs in setaria and rice by constructing a maximum likelihood phylogeny. Figure 3C shows that *NKD* exists as a single copy gene in both setaria and rice (*SvNKD* - Sevir.7G212900, *OsNKD* - LOC_Os04g47860), suggesting that the two copies in maize (*ZmNKD1* and *ZmNKD2*) resulted from the recent whole genome duplication (Swigonova et al., 2004). For phenotypic characterization, the homozygous double *Zmnkd1-Ds;Zmnkd2-Ds* line and newly generated CRISPR loss-of-function mutants in setaria and rice were all compared to wild-type lines from the same genetic background (Fig. 4A, Figure S1 & S4). Homozygous *Zmnkd1-Ds;Zmnkd2-Ds* seed exhibited the characteristic shrunken kernel phenotype that is caused by defective patterning of the aleurone layer but no altered seed phenotype was observed in *Svnkd* or *Osnkd* mutants (Figure S5). Overall plant growth was normal in maize, setaria and rice *nkd* mutants (Fig. 4B) and no leaf patterning perturbations were observed (Fig. 4C-J, Figure S6). In *Zmnkd1-Ds;Zmnkd2-Ds* mutants, no changes relative to wild-type were observed in leaf vein-density or in the ratio of rank-1 to rank-2 intermediate veins (Fig. 4G & 4H), traits that are both altered in *Zmscr1;Zmscr1h* mutants. Furthermore, there was no change in the number of M cells separating veins (the most penetrant phenotype in equivalent *scr* mutants) in either maize or setaria *nkd* mutants (Fig. 4I & 4J). Taken together, these results indicate that *NKD* is not necessary for normal leaf development in maize, setaria or rice.

Loss of *NKD* gene function enhances *scr* mutant phenotypes in maize and setaria but not in rice

To determine whether a role for *NKD* in leaf patterning could be revealed in the absence of SCR function, quadruple *scr1;scr1h;nkd1;nkd2* mutants of maize and triple *scr1;scr2;nkd* mutants of setaria

and rice were generated (Figure 5, Figure S1 & S4). An initial assessment of overall growth phenotypes revealed that perturbations in three independent triple *Osscr1;Osscr2;Osnkd* mutant lines were similar to those seen in double *Osscr1;Osscr2* mutants (Fig. 5B-D). By contrast, triple *Svscr1;Svscr2;Svnkd* mutants displayed more severe perturbations than double *Svscr1;Svscr2* mutants (Fig. 5E-I). In the progeny of *Svscr1;Svscr2/+;Svnkd/+* plants from two independent lines, the resultant triple mutants were smaller and died faster than double *Svscr1;Svscr2* mutant siblings (Fig. 5H & 5I). This was consistent in lines fixed for *Svnkd*, where only 8/20 and 8/34 triple mutants from two independent *Svscr1;Svscr2;Svnkd* lines survived 22 days after sowing, compared to 14/15 and 13/17 *Svscr1;Svscr2* double mutants. In maize, quadruple *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* mutants appeared similar to double *Zmscr1;Zmscr1h* mutants, with pale, drooping leaves and a shorter stature than either wild-type or double *Zmnkd1-Ds;Zmnkd2-Ds* mutant plants (Fig. 5J-O). However, germination of quadruple mutants was often inconsistent, and seedlings appeared weaker and more susceptible to fungal contamination. Mutant combinations thus demonstrate that NKD and SCR act synergistically in maize and setaria but not in rice.

To determine the impact of SCR and NKD interactions on leaf development, inner leaf patterning was examined in quadruple (maize) and triple (setaria and rice) mutants. Notably, the complete penetrance of the stomatal phenotype in double *scr1;scr2* mutants of setaria and rice precluded an assessment of interactions with *NKD* during stomatal development but no changes in stomatal patterning were observed in single *Svnkd* or *Osnkd* mutants (Fig. S6). In inner leaf tissues, the number of M cells between veins were the same in three independent triple *Osscr1;Osscr2;Osnkd* mutant lines of rice as in corresponding double *Osscr1;Osscr2* mutants, with the modal number of M cells being five in all cases (Fig. 6A-C). By contrast, fused veins (whereby the BS cells of adjacent veins are in contact with no intervening M cells) were observed in triple *Svscr1;Svscr2;Svnkd* mutants of setaria but not in double *Svscr1;Svscr2* mutants (Fig. 6D-F). Such fusions were not found in every triple mutant examined, with examples observed in 10/22 samples across two experiments with two independent lines (Fig 6F & Fig. S7). However, only one occurrence of a fused vein was observed in twenty-five double *Svscr1;Svscr2* mutant samples. Fused veins were also observed in quadruple mutants of maize (Fig. 7A-H). Quadruple mutants did not exhibit higher vein density or an increase in veins associated with sclerenchyma when compared to double *Zmscr1-m2;Zmscr1h-m1* mutants (Fig. 7I & 7J) but the increase in the number of veins fused to adjacent veins was consistent and statistically significant (Fig. 7F, 7H, 7K-M). Furthermore, whereas only two adjacent veins fused in double *Zmscr1-m2;Zmscr1h-m1* mutants, groups of three or more fused veins were often seen in quadruple mutants (Fig. 7L). These fused veins resulted in a large increase in the number of veins separated by no M cells, with only ~20% of veins separated by the normal two M cells in quadruple mutants, compared to just over 40% in double *Zmscr1-m2;Zmscr1h-m1* mutants. In both maize and setaria the proportion of veins separated by only one M cell was similar in double and quadruple/triple mutants (Fig. 6F & 7M). To ensure that these differences were not caused by heterozygosity at unlinked loci following the cross

between the *Zmscr1-m2;Zmscr1h*-m1 and *Zmnkd1-Ds;Zmnkd2-D* double mutants, we also compared *Zmscr1-m2;Zmscr1h-m1* mutants pre- and post- outcross and found no difference in the proportion of fused veins (Fig. S8A). Furthermore, we validated that the phenotype was consistent across two further quadruple mutants from crosses using independent *Zmnkd1-Ds;Zmnkd2-Ds* plants (Fig. S8B). Together these results demonstrate that in the C_4 species maize and setaria, but not in the C_3 species rice, *NKD* genes function with *SCR* to determine whether, and how many, M cells are positioned between veins.

Evidence for a NKD-mediated maternal effect on leaf patterning in maize

To assess whether ZmNKD1 and ZmNKD2 function redundantly in the context of leaf patterning, we quantified the percentage of fused veins found in plants homozygous for *Zmscr1;Zmscr1h* but with a range of *ZmNKD1* and *ZmNKD2* genotypes (Fig. 8A). These genotypes were generated by self-pollinating plants that had at least one wild-type copy of either *ZmNKD1* or *ZmNKD2.* Surprisingly, some genotypes displayed more fused veins than others. For example, triple *Zmscr1;Zmscr1h;Zmnkd1;ZmNKD2* mutants derived from either *Zmscr1/+;Zmscr1h;Zmnkd1/+;Zmnkd2/+* or *Zmscr1/+;Zmscr1h/+;Zmnkd1;ZmNKD2* parents exhibited a higher number of fused veins than the three quadruple mutants derived from *Zmscr1/+;Zmscr1h;Zmnkd1/+;Zmnkd2/+* or *Zmscr1/+;Zmscr1h;Zmnkd1/+;Zmnkd2* parents (Fig. 8A, Table S1). Given that phenotyping was undertaken on leaf 3, which is initiated and partially patterned in the embryo, we hypothesised that the inconsistencies observed may reflect a maternal effect. Because all parental plants had a wild-type copy of *SCR1*, any inconsistencies must be mediated by NKD, with NKD protein in the parental plant either directly or indirectly compensating for the lack of NKD in the developing embryo.

To further evaluate the hypothesis that NKD may have a maternal influence on patterning of embryonic leaves, we examined the progeny of a plant which had the genotype *Zmscr1-m2/+;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds*. The progeny of this homozygous *nkd1/nkd2* parent segregated ¼ for quadruple mutants and in leaf 3 of each of the three quadruple mutant seedlings examined a greater percentage of fused veins was observed than in leaf 3 of the quadruple mutant seedlings that had originated from a parent with at least one wild-type copy of *NKD1* and/or *NKD2* (Fig. 8A). To further investigate this effect, we compared cross sections of quadruple mutant embryos from a *Zmscr1-m2/+;Zmscr1h;nkd1/+;nkd2/+* (*nkd* heterozygous) parent and a *Zmscr1-m2/+;Zmscr1h;nkd1;nkd2* (*nkd* homozygous) parent, with wild-type, *Zmscr1-m2;Zmscr1h-m1* and *Zmnkd1-Ds;Zmnkd2-Ds* embryos (Fig. 8B-H). Transverse sections from the tip of the meristem (Fig. 8B), where P1-P4 leaf primordia (where P4 will go on to form leaf 1, P3 leaf 2 etc.) are visible, were examined. Due to the anticipated challenges of identifying fused veins in P4 leaf primordia, where cell-division and differentiation is ongoing, we first quantified vein density to test whether it could be used as a proxy for an increase in fused veins. However, no consistent change in vein density was observed between genotypes (Fig.

8C). Sections were therefore carefully examined for evidence of vein fusion by three independent people, two of whom had no knowledge of which sections corresponded to which genotype. All three observers identified fused veins in what were revealed to be the quadruple mutants from a homozygous *nkd* parent, but not in the quadruple mutants from the heterozygous *nkd* parent. Taken together, these results support the hypothesis that there is a maternal effect of NKD on embryonic leaf development. However, progeny of all homozygous *Zmnkd1-Ds;Zmnkd2-Ds* plants develop in the context of a shrunken kernel which has a disordered aleurone comprised of multiple cell layers. As such, the fused vein phenotype in leaf 3 of quadruple mutant progeny could result as an indirect consequence of the altered kernel structure as opposed to a direct consequence of loss of NKD function. To evaluate whether NKD has a direct role, we examined leaf 6 of quadruple mutants that were derived from a homozygous *nkd* parent. These leaves, which are patterned post-germination, also had an increased proportion of fused veins relative to double *Zmscr1;Zmscr1h* mutants (Fig. S8C-F), demonstrating a role for NKD in leaf patterning beyond embryogenesis. Although this observation does not preclude the possibility that the maternal effect is a consequence of physical alterations to the aleurone, collectively these data suggest that NKD functions alongside SCR to pattern both embryonic and post-embryonic leaves in maize.

DISCUSSION

SCARECROW has recently emerged as a regulator of monocot leaf development (Hughes et al., 2019; Hughes & Langdale, 2022; Slewinski et al., 2012; Wu et al., 2019). Here we have demonstrated that unlike in maize and rice where SCR has distinct functions in inner leaf (maize) or stomatal (rice) patterning, in the C_4 monocot *Setaria viridis* it undertakes both patterning functions (Fig. 1 & 2). Prior to this study, no regulators that function alongside SCR in monocot leaf patterning had been identified. Here we show that *NKD IDD* genes play such a role, at least in C_4 monocots (Fig. 3). Although loss of function *nkd* mutants in maize, setaria and rice show no obvious perturbations in leaf development (Fig. 4), when mutations are combined with loss of function *scr* mutations, synergistic interactions are observed in both maize and setaria, but not in rice (Fig. 5-7). Most strikingly, *scr;nkd* mutants in maize and setaria exhibit an increase in the proportion of fused veins, whereby the BS cells of adjacent vascular bundles are in direct contact with no intervening M cells (Fig. 6 & 7). Intriguingly, the presence or absence of this phenotype in embryonic leaves suggests that the expression of *NKD* in maternal tissue may influence the development of leaves in the embryo (Fig. 8). Taken together, our results provide insight into the evolution of SCR function in monocot leaves and identify another component of the patterning pathway.

We previously proposed that the role of SCR in patterning inner leaf tissues of maize may represent a C_4-specific function, and that the stomatal role in rice may be a C_3-specific function (Hughes & Langdale, 2022). The analysis presented here argues against the latter because *Svscr1;Svscr2* mutants of the C_4 species *Setaria viridis* do not develop stomata. However, inner leaf patterning

perturbations were also observed in *Svscr1;Svscr2* mutants, and thus it remains plausible that the inner leaf patterning function is C_4-specific. In this scenario, the ancestral role of SCR would be in epidermal patterning, with recruitment into Kranz patterning occurring in those monocots that evolved the C_4 pathway. This situation contrasts with that seen in eudicots where Arabidopsis (C_3) *scr* mutants develop normal stomata but have perturbations in BS cell identity (Cui et al., 2014; Dhondt et al., 2010). On the basis of gene expression patterns and mutant phenotypes in rice, it has been proposed that SCR functions in the epidermis to interpret a positional signal (likely SHR) emanating from veins in the inner leaf (Nunes et al., 2020; Schuler et al., 2018). In this way, stomatal files are correctly positioned in rows adjacent to underlying veins. It is difficult to envisage how this could happen in setaria if SCR is also expressed in the inner leaf as in maize, particularly given that the canonical model for SCR function dictates that SCR prevents SHR movement beyond the cell layer in which SCR accumulates (Cui et al., 2007; Gallagher et al., 2004; Koizumi et al., 2012; Nakajima et al., 2001). How epidermal and inner leaf patterning processes are both mediated through the SCR pathway in setaria is a question for future research.

Another outstanding question from this study is why the severity of the fused vein phenotype is much higher in maize than in setaria *scr;nkd* mutants. In maize, many quadruple mutants exhibited more than 30% of leaf veins that were fused to an adjacent vein, whereas in setaria these incidents were rarer, and no examples were found of more than two veins fused together. One possible explanation is that because *scr;nkd* mutants in setaria have no stomata they are much weaker than those in maize. As a consequence, fewer plants survived and inner leaf phenotypes were necessarily assessed on those survivors, possibly biasing the analysis towards mutants that were less severely affected. An alternative possibility is that the mutant alleles in setaria are hypomorphic rather than null. The successful edits in *SvSCR* and *SvNKD* genes were positioned towards the 3' end of the sequence, affecting the final 20% and 40% of the encoded proteins respectively. Whereas we can be confident that the *Svscr* alleles are null due to the conserved phenotype of *Svscr1;Svscr2* mutants with rice and maize, it is possible that the *Svnkd* alleles retain some wild-type function and that a more severe *Svscr;Svnkd* vein clustering phenotype would be observed if edits towards the 5' end of the gene had been successful. If neither of these technical explanations is correct, the differences observed between maize and setaria must represent biological differences between the two species, which is plausible given that they are separated by millions of years of evolution and clearly deploy the SCR pathway in different developmental contexts.

IDD genes have long been known to modulate the SCR/SHR pathway in Arabidopsis. For example, JACKDAW is known to physically interact with the SCR-SHR complex (Welch et al., 2007), and it has been suggested that this interaction with an IDD C2H2 transcription factor partner enables SCR-SHR to bind DNA and activate expression of downstream targets (Hirano et al., 2017). There are IDD orthologs in monocot genomes, but very little functional insight into their roles. This may be in part due

to the extensive functional redundancy in grass genomes which combined with the challenges of generating loss-of-function lines in monocots makes functional analysis challenging. However, it was recently found that IDD12 and IDD13 act alongside SHR in rice to regulate ground meristem proliferation in leaves (Liu et al., 2022). This finding is intriguing because we find here that *NKD* plays a similar role alongside *SCR* in C$_4$ inner leaf patterning, but that neither gene functions in the inner leaf of rice (Figure 6, Hughes & Langdale, 2022). It may be that different IDD genes fine-tune the action of both SCR and SHR by controlling the activation of distinct downstream targets in different developmental contexts.

The accumulation of *SCR* and *NKD* transcripts in the ground meristem cells between developing veins, and the fused veins observed in both maize and setaria *scr;nkd* mutants imply two possible mechanisms for SCR and NKD function. The first is that the combined action of SCR and NKD promotes cell division in the ground meristem and then specifies those cells as mesophyll. In this model, in the absence of SCR and NKD ground meristem cells do not divide and/or are mis-specified such that veins develop in direct contact with each other. This would be analogous to the role played by SCR in the arabidopsis root, where loss of gene function leads to loss and mis-specification of the endodermal cell-layer (Laurenzio et al., 1996). The second is that SCR and NKD act to inhibit the formation of veins in regions between existing veins. In their absence, veins are formed in regions between already specified veins, resulting in the runs of fused veins observed in maize and setaria *scr;nkd* mutants. This mechanism may be analogous to the lateral inhibition mechanism that patterns trichomes and stomata on the leaf surface (Schellmann et al., 2002; Torii, 2012; Zeng et al., 2020), whereby the fused veins represent a clustering phenotype caused by absence of the inhibitor. Because SCR and NKD do not accumulate in developing veins, in this scenario it is likely that they would interpret an inhibitory signal emanating from adjacent specified veins. In either case, the inner leaf perturbations observed in *scr* but not *nkd* mutants suggest that SCR can compensate for loss of NKD function in the inner leaf patterning pathway whereas NKD cannot compensate for loss of SCR function, a relationship that is similar to that seen in other pathways where SCR and IDD proteins interact (Long et al., 2015; Ogasawara et al., 2011; Welch et al., 2007).

NKD has a well-defined role in the endosperm of maize, where it acts to suppress proliferation of the aleurone layer – the aleurone is a single cell layer in wild-type endosperm whereas in *nkd1;nkd2* mutants it is multi-layered (Yi et al., 2015). Within the endosperm, *NKD1* and *NKD2* are expressed in both the aleurone and the starchy endosperm, and loss of function of either gene is compensated for by upregulation of the other (Yi et al., 2015). Although low levels of *NKD* gene expression have also been reported in the developing embryo (Yi et al., 2015), the data reported here suggest that maternally expressed *NKD* may play a role in the patterning of embryonic leaves. Without further experimentation it is impossible to determine the mechanistic basis of this non cell-autonomous effect. Perturbations to leaf patterning in embryos developing within *nkd1;nkd2* mutant seed could either be an indirect

consequence of disrupted signalling through the mutant aleurone layer or a direct consequence of loss of activity of NKD and/or its downstream targets. Speculation at this stage would be premature but it is interesting to note that most of the maternal effect genes reported to date encode epigenetic regulators or small molecules (peptides or miRNAs) as opposed to transcription factors (reviewed in Ingram, 2020). Taken together, the data presented here suggest that NKD is a key regulator of leaf patterning in C_4 grasses, acting with SCR to promote M cell specification or repress vein specification, and influencing the patterning of embryonic leaves in a non-cell autonomous manner.

MATERIALS & METHODS

Phylogenetics

Primary transcript proteomes from *Zea mays* (B73), *Sorghum bicolor*, *Setaria italica*, *Setaria viridis*, *Oryza sativa*, *Brachypodium distachyon*, *Brachypodium stacei*, *Ananas comosus*, *Arabidopsis thaliana*, *Solanum lycopersicum* and *Physcomitrella patens* were downloaded from Phytozome12 (Goodstein et al., 2012). The ZmNKD1 (GRMZM2G129261) primary protein sequence was used as a query in a BLASTp search (e-value of 1e-3) against these proteomes. The top 100 results were aligned using MAFFT-linsi (Katoh & Standley, 2013) and the alignment was then used to generate a maximum likelihood phylogeny using IQtree (Hoang et al., 2018; Trifinopoulos et al., 2016).

Plant material and growth conditions

Maize inbred line W22 and *Zmnkd1-Ds;Zmnkd2-Ds* (introgressed into W22) seed were obtained from Phil Becraft, Iowa State University. *Zmscr1-m2;Zmscr1h-m1* seed were generated and reported previously (Hughes et al. 2019). W22 was used as the wild-type control in all maize experiments except for *in situ* hybridization experiments where the inbred line B73 was used. *Setaria viridis* accession ME034V was used as the wild-type control for all setaria experiments and *Oryza sativa spp japonica* cv. Kitaake was used as the wild-type control for all rice experiments. *Osscr1;Osscr2* seed were generated and reported previously (Hughes & Langdale, 2022).

Maize and setaria plants for phenotypic analysis were grown in a greenhouse in Oxford, UK with 16 h light/ 8 h dark cycle, daytime temperature 28 °C and night-time temperature 20 °C, with supplemental light provided when natural light levels were below 120 µmol photon m^{-2} s^{-1}(Hughes et al., 2019). Maize seed were germinated and grown as described previously (Hughes et al., 2019). Setaria seed were treated for at least 3 weeks in damp sphagnum moss (Zoo-Med Laboratories Inc) at 4°C prior to germination to break dormancy. Seed were then germinated on damp paper towels in sealed petri dishes, in a growth cabinet with the same conditions as the greenhouse. After seven days, seedlings were transferred to Sinclair compost in 60 well modular trays for growth in the greenhouse. Plants grown for seed propagation were re-potted after 4 weeks into 7.5cm pots with the same compost, for growth to maturity. Rice seed were sterilised prior to germination on ½ MS media and subsequently

transferred to a hydroponic growth system in 50 ml falcon tubes, as described previously (Hughes & Langdale, 2022).

Generation of maize quadruple mutant and genotyping

Pollen from *Zmscr1-m2/+;Zmscr1h-m1* plants (double mutants do not produce pollen or ears) was used to cross to *Zmnkd1-Ds;Zmnkd2-Ds* homozygous ears. 50% of the F1 plants were thus heterozygous for all four mutant alleles, and these plants were self-pollinated. In the subsequent F2, kernels with the shrunken *nkd1/nkd2* phenotype were selected for self-pollination. F3 seed derived from individuals with *Zmscr1/+;Zmscr1h;Zmnkd1;Zmnkd2* or *Zmscr1;Zmscr1h/+;Zmnkd1;Zmnkd2* genotypes were then used for quadruple mutant analysis. Other mutant combinations were derived from parental genotypes as indicated in Table S1.

Genomic DNA for genotyping was extracted either using a modified SDS 96-well plate method or a CTAB method depending on the number of samples to be processed, as described previously (Hughes et al 2019). PCR genotyping assays were used to track the genotype at each locus through the generations. The *Zmscr1-m2* and *Zmscr1h-m1* alleles were genotyped as described previously (Hughes et al. 2019). The expected *Ds* insertion site in the *Zmnkd1-Ds* allele was confirmed by two separate PCR assays, one using two primers (nkd1-F, TATCTTATCCGTCGATGCGTTG and nkd1-R, TCGGTCATGGCATCCTGCCTCCG) that flanked the insertion site and produced an amplicon for the wild-type *ZmNKD1* sequence, and another using one flanking primer (nkd1-F) and a second primer nested within the Ds transposon sequence (W22-Ds-R1, GGAGCTGGCCATATTGCAGTCATC) that produced an amplicon when the *Zmnkd1-Ds* sequence was present. The *Zmnkd2-Ds* allele referred to here was previously described as *nkd2-Ds0766* and believed, like *Zmnkd1-Ds,* to be a loss-of-function allele caused by the insertion of a *Ds* transposon. However, attempts to amplify from the *Ds* transposon sequence to the region flanking the insertion failed. Instead, we demonstrated that loss-of-function of this allele was due to a deletion at the 5' end of the gene, with amplification (primers ZmNKD2-F2, CTTGTTGCCGTTGTTGATTG and ZmNKD2-R2, GTGCCATGTGGCTCCTATTT) yielding a fragment that was 786bp smaller in homozygous *Zmnkd2-Ds* plants than in wild-type (Fig. 4A, Figure S9). This deletion is believed to be caused by a chromosomal rearrangement (Phil Becraft, personal communication). As expected, *Zmnkd1-Ds;Zmnkd2-Ds* seed always exhibited the shrunken kernel phenotype characteristic of *Zmnkd1;Zmnkd2* double mutants. PCR amplifications were undertaken using GoTaq DNA polymerase (Promega) and PCR cycles of 95 °C for 5 min; 35 cycles of 95 °C for 30 s, 57-64 °C for 30 s and 72 °C for 60-90 s; and 72 °C for 5 min. Betaine (1 M; Sigma Aldrich) was added to all reactions to aid amplification of regions with high-GC content.

Generation of rice and setaria CRISPR lines

Rice and setaria *NKD* orthologs were identified from the phylogeny presented in Figure 3. Setaria *SCR* orthologs were identified from a previously published phylogeny (Hughes et al. 2019). The *OsNKD*

sequence was obtained from phytozome V12 and two guide RNAs targeting the 5' end of the *coding* sequence were designed using CRISPOR (Concordet & Haeussler, 2018) (OsNKD-g59: CTTCGGGATTAGCAGGGATG, OsNKD-g72: GTGATATCAGCGGCTTAATC). Guide RNAs targeting *OsSCR1* and *OsSCR2* were designed and used previously (Hughes & Langdale, 2022), and two of these (OsSCR1-g397: TCCACCCAAGCCGTACTAGG, OsSCR2-g507: CGAGGTCGGGGTTACATGGC) were used in this study. Guides were cloned as described previously into four constructs targeting either *OsNKD* (EC17821: OsNKD-g59 and EC17822: OsNKD-g72) or *OsSCR1, OsSCR2* and *OsNKD* (EC17827: OsSCR1-g397, OsSCR2-g507, OsNKD-g59 and EC17828: OsSCR1-g397, OsSCR2-g507, OsNKD-g72). *SvSCR1, SvSCR2* and *SvNKD* sequences were obtained from phytozome V12, however, as the ME034V accession used for transformation is not the same as the sequenced accession, we first amplified the target regions of the ME034V gene sequences. Surprisingly, we found that the ME034V *SvSCR1* and *SvSCR2* sequences more closely matched the published *Setaria italica* sequences, and the ME034V *SvNKD* sequence was intermediate between the published *viridis* and *italica* sequences. Confirmed ME034V sequences were used in subsequent guide RNA design.

Due to a low editing efficiency of guide RNAs in *Setaria viridis* (Daniela Vlad, personal communication)*, multiple guides were designed against each gene (Figure S1). Because *SvSCR1* and *SvSCR2* have high sequence similarity, a common set of six guides were designed to target both genes simultaneously. For both *SvSCR* and *SvNKD*, guides were positioned towards the 5' end of the gene to make complete loss-of-function alleles more likely. However, at least one guide was positioned closer to the 3' end of the gene both to increase the chance of obtaining mutations and to enable the potential excision of most of the gene sequence. To minimise the number of constructs for transformation, guides were cloned into polycistronic expression modules separated by tRNA spacers (Hahn et al., 2020). Two such polycistronic guide arrays were assembled, one with the six *SvSCR* guides and one with the four *SvNKD* guides (Figure S1), both driven by the rice U3 promoter. These arrays were then combined into three constructs using the same Golden Gate cloning system described previously for rice (Hughes & Langdale, 2022): C402052 (SvSCR array), C402053 (SvNKD array) and C402054 (SvSCR and SvNKD arrays). Three identical constructs were cloned that included the GRF-GIF1 fusion previously shown to enhance rice and wheat transformation efficiency (Debernardi et al., 2020). No enhancement in setaria transformation was observed here, however, one line that was generated from one of these constructs (C402057: SvSCR and SvNKD arrays) was used for subsequent phenotypic analysis after the construct (and thus GRF-GIF1 fusion) were segregated away from the mutations.

Constructs were transformed into Kitaake rice seed using agrobacterium strain EHA105 as described previously (Hughes & Langdale, 2022). Constructs were transformed into setaria ME034V using a modified transformation protocol (Finley et al., 2021). In brief, ME034V seed were dehulled and

sterilised with 10% bleach plus 0.1% tween for 3 minutes. Seed were placed on callus induction media (CIM) (4.3 g/L MS salts, 40 g maltose, 35 mg/L $ZnSO_4 \cdot 7H2O$, 0.6 mg/L $CuSO_4 \cdot 5H_2O$, 0.5 mg/L kinetin, 2 mg/L 2,4-D 4 g/L Gelzan, pH 5.8) and after 4-5 weeks embryonic callus was moved to fresh media and gelatinous callus removed. After a further 3 weeks, gelatinous callus was again removed and the remaining callus moved to fresh media for 1-2 weeks prior to transformation. Constructs were cloned into agrobacterium strain AGL1 and grown in liquid media to an OD of ~0.6. Agrobacteria were then collected by centrifugation and resuspended in 50ml liquid CIM media with the addition of 40 µm acetosynringone and 0.02 % synperonic acid. Around 100 pieces of calli were added to this suspension and incubated for 5 minutes with occasional rocking prior to drying and transfer to fresh CIM plates with filter paper placed on top of the media. Calli were co-cultivated for 3 days at 22°C before transfer to CIM selective plates (CIM containing 150 mg/L timentin and 40mg/L hygromycin for 16 days at 24°C in the dark. Calli were then transferred to plant regeneration media (PRM) (4.3 g/L MS salts, 20 g/L sucrose, 7 g/L Phytoblend, 2 mg/L kinetin, 150 mg/L timentin, 15 mg/L hygromycin, pH 5.8) and moved to fresh media every 2 weeks. Emerging shoots were dissected from calli and moved to rooting media (RM) (2.15 g/L MS salts, 30 g/L sucrose, 7 g/L Phytoblend, 150 mg/L timentin, 20 mg/L hygromycin, pH 5.7). Shoots that survived this stage were transferred to compost for genotyping.

Genomic DNA for rice and setaria T0 genotyping was obtained using the same methods described for maize. T0 seedlings were first screened using primers that amplified a fragment of the hygromycin gene (Hughes & Langdale, 2022) to assess transformation success. Gene specific primers were then used to amplify and sequence the region targeted for editing. In rice, all guides induced successful edits, however, in the case of construct EC17822 only a few transformed plants were obtained and thus edited plants from EC17821 were taken forward for *Osnkd* single mutant characterization. In setaria, only one guide for both *SCR* and *NKD* (SvSCR-ex2g49: GAGCAGGACCTGAGCCACTC and SvNKD-ex3g438: CATGAGTCCATGGAACGGCT) was found to consistently yield successful edits. In both cases this guide was positioned closer to the 3' end of the gene. In the case of *SCR*, out-of-frame mutations created by the successful guide resulted in the final ~20% of the protein sequence being either nonsense or prematurely truncated by an early stop codon, whereas for *NKD* the final ~40% of the protein was affected. As was the case when generating rice *scr* mutants (Hughes & Langdale, 2022), no plants were identified in the T0 generation that had out-of-frame mutations in all four *SCR* alleles, with all screened plants having at least one unedited copy of one of the *SCR* genes. Interestingly, some T0 plants exhibited pale sectors in leaves, but only in plants transformed with the *SCR* guide array. All *Svnkd* mutations corresponded to the same C deletion (*Svnkd-m1*), and plants homozygous for this edit were viable. For both rice and setaria, T1 lines were screened to identify mutated plants that had segregated away from the construct. From this analysis, two independent *Osnkd* lines, three *Osscr1;Osscr2;Osnkd* lines (alongside two previously generated *Osscr1;Osscr2* lines), two *Svscr1;Svscr2* lines, two *Svnkd* lines and three *Svscr1;Svscr2;Svnkd* lines were prioritized for phenotypic characterization in the T2 and T3 generation.

Leaf phenotyping

Fully expanded leaves were used to obtain transverse leaf sections of maize, setaria and rice. For maize and setaria a region from the midpoint along the proximal-distal axis was cut and placed in molten 7% agar. Once solidified, blocks were trimmed down and mounted using superglue for sectioning on a vibratome. Sections of 30-60 μm were cleared in either 3:1 ethanol: acetic acid (maize) or 70% ethanol (setaria) for 10 minutes. *Svscr1;Svscr2* and *Svscr1;Svscr2;Svnkd* leaves were sufficiently pale for leaf anatomy to be visible without any clearing. Sections were transferred to slides and imaged using a Leica DMRB microscope with a DFC7000T camera under either brightfield or UV illumination using Leica LASX image analysis software. Rice leaf transverse sections were obtained by first fixing a segment from the midpoint along the proximal-distal axis of fully expanded leaf 5 in 3:1 ethanol: acetic acid for 30 mins, before transferring to 70% ethanol. Samples were wax infiltrated, sectioned at 10 μm using a Leica RM2135 rotary microtome and stained with Safranin O and Fastgreen as described previously (Hughes & Langdale, 2022). Stomatal impressions of setaria leaves were obtained and imaged under phase-contrast microscopy using the same dental resin and nail varnish method described previously (Hughes & Langdale, 2022).

Seed phenotyping

Setaria seed were and dehulled fixed in FAA (10% formaldehyde, 5% acetic acid, 50% ethanol) for 2 months prior to sectioning. Seed were then embedded in paraffin as described previously (Hughes & Langdale, 2022) and sectioned into 20μm sections using a Leica RM2135 rotary microtome. The resultant sections were incubated in Histoclear (2 x 10 min), and then re-hydrated through 100%, 95%, 90%, 80%, 70%, 50%, 30% and 10% (w/v) ethanol, all at room temperature. Sections were then rinsed in distilled H_2O, stained for 5 seconds in 0.05% (w/v) Toluidine Blue (50 mM citrate buffer, pH 4.4), and then rinsed in distilled H_2O again, before finally being mounted using a drop of entellen (Merck Millipore). Slides were imaged under brightfield using the same microscope described above. Rice and setaria whole seed were photographed using a Leica S9i stereo microscope.

***In situ* hybridization**

In situ hybridization was carried out using wax-embedded shoot apices as described by (Schuler et al., 2018), with digoxygenin (DIG)-labelled RNA probes designed to specifically detect either *ZmSCR1* or *ZmNKD1* transcripts. The *ZmSCR1* probe was a 108 bp region towards the end of the first exon, which shared 78% identity with the corresponding region of *ZmSCR1h* (Hughes et al., 2019). The *ZmNKD1* probe was a 116 bp region of the 5'UTR ending 12 nucleotides upstream of the ATG. The probe shared 59% identity with the corresponding region of *ZmNKD2* and with the stringency conditions used was expected to be specific for *ZmNKD1*. Post-hybridization washes were undertaken with 0.005 x SSC buffer made from a 20x SSC stock (3M NaCl, 0.3M Na$_3$citrate), calculated to ensure stringency.

Maize kernel genotyping and embryo imaging

The endosperm of kernels from segregating seed packets was first chipped and used for genotyping using the CTAB DNA extraction method described above, except samples were homogenized prior to the addition of extraction buffer. The resultant genomic DNA was used as template in PCR amplifications to identify homozygous mutants for phenotypic characterization. Kernels of interest were then soaked overnight in water, to enable the removal of the embryo the following day. Embryos were fixed in FAA (10% formaldehyde, 5% acetic acid, 50% ethanol) and vacuum infiltrated before incubation overnight in fresh FAA. Samples were transferred to 50% and then 70% ethanol the next day, and wax infiltrated as described previously (Hughes & Langdale, 2022). 10 μm transverse sections were obtained using a microtome just beneath the tip of the meristem where all primordia are visible, and stained with Safranin O and Fastgreen as described above. Slides were imaged using brightfield illumination and the whole region of interest obtained using the Leica XY builder software.

qPCR

RNA was extracted using the RNeasy kit (Qiagen) from whole maize shoots of both *Zmscr1-m2;Zmscr1-m1* and *Zmnkd1-Ds;Zmnkd2-Ds* mutants and the respective wild-type lines, 7 days after germination. RNA was DNase treated (TURBO DNase, Thermo Fisher) and used as template to generate cDNA (Maxima first strand cDNA synthesis kit, Thermo Fisher). Primers were designed to amplify short fragments of *ZmSCR1, ZmSCR1h, ZmNKD1* and *ZmNKD2* gene sequences as well as two housekeeping genes (*ZmCYP* and *ZmEF1α*) verified previously as suitable for use in maize qRT-PCR (Fig. S3A) (Lin et al., 2014). RT-PCR was used to verify that a single product of the correct size was amplified. Quantitative RT-PCR amplifications were undertaken using SYBR-Green (Thermo Fisher) with cycling conditions 95 °C for 10 min, then 40 cycles of 95 °C for 15 s and 60 °C for 1 min. Melt curves were obtained by heating the resultant product from 60 °C to 95 °C to confirm that a single product was amplified for each primer pair (Fig. S3B). Amplification efficiencies were assessed using a dilution series of cDNA template with >80% taken to be suitable. In all cases three technical replicates with Ct range <0.5 were obtained from each sample and all comparisons were run on the same 96-well plate. Ct values were obtained using the qPCR miner algorithm (Zhao & Fernald, 2005) and fold-change between wild-type and mutants was calculated using the $2^{-\Delta\Delta CT}$ method (Livak & Schmittgen, 2001). A combined wild-type average was used to compare each individual wild-type to indicate the range of the wild-type data. Mutant samples were then compared with the same overall wild-type average. Values <1 indicate a relative reduction compared with wild-type whereas values >1 indicate a relative increase compared to wild-type.

AUTHOR CONTRUBUTIONS

T.E.H and J.A.L conceived and designed the experiments. O.S carried out the *in situ* hybridization experiment and M.T. analyzed the embryonic leaf phenotypes in maize and the aleurone phenotype in

setaria. T.E.H carried out all of the remaining experiments and analysed the data. T.E.H and J.A.L wrote the manuscript.

ACKNOWLEDGEMENTS

The authors thank Phil Becraft, Erik Vollbrecht, Hao Wu, Ruaridh Sawers and Ruben Rellan Alvarez for enabling maize genetics in the field in Iowa and Mexico; John Baker for plant photography; Roxaana Clayton, Julie Bull and Lizzie Jamison for technical support; Matthew Karadzas for initiating the qRT-PCR experiments; Sophie Johnson, Chiara Perico, Daniela Vlad, Sovanna Tan, Julia Lambret-Frotte and Maricris Zaidem for discussion throughout the experimental work and during manuscript preparation.

COMPETING INTERESTS

No competing interests declared

FUNDING

This work was funded by the Bill and Melinda Gates Foundation C_4 Rice grant awarded to the University of Oxford (2015-2019; OPP1129902) and by a BBSRC sLoLa grant (BB/P003117/1)

FIGURE LEGENDS

Figure 1. Mutations in *SvSCR1* and *SvSCR2* affect plant growth. A) Sequences of mutant *Svscr1* and *Svscr2* alleles in two independent lines (C402052-6 & C402054-7). Wild-type (WT) sequence is shown on top, with the sequence of the mutant allele shown beneath. Guide sequences are depicted in blue, and mismatches between the WT and mutant sequence indicated in red. **B-D)** Photos of WT ME034V (B), *Svscr1-m3;Svscr2-m4* (C) and *Svscr1-m2-Svscr2-m1* (D) plants taken 20 days after sowing. Scale bars: 10cm.

Figure 2. *Svscr1;Svscr2* mutant plants have no stomata and fewer mesophyll cells separating veins. A-F) Stomatal impressions of wild-type (WT) ME034V (A,D), *Svscr1-m1;Svscr2-m4* (B,E) and *Svscr1-m2;Svscr2-m1* (C,F) leaf 3 from either the abaxial (A-C) or adaxial (D-F) surface. Stomata are false coloured orange. Scale bars: 100μm. **G-H)** Transverse sections imaged using UV illumination of WT ME034V (G) and *Svscr1-m1;Svscr2-m4* (H) leaf 4, taken from the mid-point along the proximal-distal axis. The number of mesophyll cells between each pair of veins is indicated above the leaf. Scale bars: 100μm. **I)** Histograms summarizing the mean number of mesophyll cells separating veins in WT ME034V and two independent *Svscr1;Svscr2* mutants. The number of biological replicates is indicated above each plot and letters in the top right corner of each plot indicate statistically different groups ($P≤0.05$, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype.

Figure 3. *NKD* and *SCR* transcripts accumulate in the same spatial domain with levels determined by a feedback loop. A) Quantitative RT-PCR of *ZmSCR1* and *ZmSCR2* transcripts in the *Zmnkd1-Ds;Zmnkd2-Ds* mutant, and *ZmNKD1* and *ZmNKD2* in the *Zmscr1-m2;Zmscr1h-m2* mutant. Open circles are individual biological replicates, black crosses indicate the mean for each genotype. Statistical significance as calculated on log-transformed fold change data by Student's *t*-test (two-tailed) indicated above each plot: *$P≤0.05$; **$P≤0.01$; ***$P≤0.001$. **B)** *In-situ* hybridization to *ZmNKD1* and *ZmSCR1* in maize wild-type B73 apices. In each image the P2 primordium is outlined in blue, P3 in red, P4 in purple and P5 in green, as indicated in the adjacent cartoon diagram. Darker purple signal represents successful hybridization to each transcript of interest. Scale bars: 50μm. **C)** Maximum likelihood phylogeny of the *NKD* genes in monocots. The *NKD* clade is highlighted in green, and the adjacent monocot clade in purple. Bootstrap values are displayed at each branch of the phylogeny.

Figure 4. *nkd* loss of function mutations do not perturb growth in maize, rice or setaria. A) Cartoon depictions of loss of function *nkd* alleles in maize (*Zm*), setaria (*Sv*) and rice (*Os*). In each case, 5' and 3' untranslated regions are depicted in purple, introns in orange and coding regions in green. Transposon insertions are indicated by a yellow triangle. For setaria and rice, CRISPR guide sequences are indicated in blue and maroon respectively, with the sequence of the mutant allele

beneath with edits highlighted in red. **B)** Whole plant phenotypes of maize, setaria and rice *nkd* mutants. Photos were taken 31 days (maize), 20 days (setaria) or 14 days (rice) after sowing. Scale bars: 10cm. **C-F)** Transverse sections of maize wild-type (WT) W22 (C), *Zmnkd1-Ds;Zmnkd2-Ds* (D), setaria WT ME034V (E) and *Svnkd-m1* (F), imaged under either brightfield (maize) or UV illumination (setaria). Images were taken at the mid-point along the proximal-distal axis of either leaf 5 (maize) or leaf 4 (setaria). Scale bars: 100μm. **G-H)** Quantification of vein density (G) and the ratio of rank1:rank2 intermediate veins (H) of WT W22 (blue) and *Zmnkd1-Ds;Zmnkd-Ds* (green) mutants. Open circles indicate distinct biological replicates, black crosses indicate the mean for each genotype. Statistical significance as calculated by Student's *t*-test (two-tailed) indicated above each plot: n.s. *P*>0.05. **I-J)** Histograms summarising the mean number of mesophyll cells separating veins in WT W22 versus *Zmnkd1-Ds;Zmnkd2-Ds* (I) and WT ME034V versus two independent *Svnkd-m1* lines (J). Error bars are the standard error of the mean, and the number of biological replicates is indicated above each plot. Letters in the top right corner of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype. WT is coloured blue and *nkd* mutants green.

Figure 5. Loss of function *nkd* mutations enhance growth perturbations in *scr* mutants of maize and setaria but not rice. A) Sequences of mutant *scr1;scr2;nkd* alleles in three independent lines of both rice and setaria. Wild-type sequences are shown above with guides highlighted in either maroon (rice) or blue (setaria). Edits are highlighted in red. Note *Svscr1-m2* allele is included in Figure 1. **B-O)** Whole plant phenotypes of rice (B-D), setaria (E-I) and maize (J-O). Images were taken 14, 20 or 21 days after sowing for rice, setaria and maize respectively. Plants in (H) and (I) are from segregating families in each case, such that *Svscr1;Svscr2* and *Svscr1;Svscr2;Svnkd* in each panel are siblings. Scale bars: 10cm.

Figure 6. Loss of function *nkd* mutations induce the formation of fused leaf veins in *scr* mutants of setaria but not rice. A-B) Transverse sections of *Osscr1-m7;Osscr2-m3* (A) and *Osscr1-m7;Osscr2-m9;Osnkd-m3* (B) mutant leaves, taken at the midpoint along the proximal-distal axis of leaf 5. **C)** Histograms summarizing the mean number of mesophyll cells separating veins in two independent *Osscr1;Osscr2* (yellow) and three independent *Osscr1;Osscr2;Osnkd* (purple) mutant lines. Error bars are standard error of the mean. Sample sizes (n=) are biological replicates. **D-E)** Transverse sections of *Svscr1-m1;Svscr2-m4* (D) and *Svscr1-m2;Svscr2-m1;Svnkd-m1* (E) mutant leaves, taken at the midpoint along the proximal-distal axis of leaf 4, imaged under UV illumination. The number of mesophyll cells separating veins is displayed above each region. In (E) the inset shows an example of a fused vein with no separating mesophyll cells. **F)** Histograms summarizing the mean number of mesophyll cells separating veins in wild-type (WT) ME034V (blue), *Svscr1-m1;Svscr2-m4* (yellow) and *Svscr1-m4;Svscr2-m1;Svnkd-m1* (purple) mutants. Error bars are standard error of the mean. Sample sizes (n=) are biological replicates and letters in the top right corner (beneath sample

sizes) of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype. Scale bars: 100 μm.

Figure 7. *Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2* quadruple mutants have a striking increase in the number of fused leaf veins compared to *Zmscr1;Zmscr1h* double mutants. A-H) Transverse cross sections of wild-type (WT) W22 (A & C), *Zmnkd1-Ds;Zmnkd2-Ds* (B & D), *Zmscr1-m2;Zmscr1h-m1* (E & G) and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* (F & H) leaf 3, from the mid-point along the proximal distal axis and imaged under either brightfield (A-B, E-F) or UV (C-D, G-H) illumination. Scale bars: 100 μm. **I-L)** Strip charts summarizing quantification of vein density (I), the ratio of rank 1: rank 2 intermediate veins (J), the % of fused veins (K) and the % of veins formed in runs of ≥3 fused veins (L). Open circles indicate measurements from independent biological replicates, and black crosses indicate the mean for each genotype. The number of biological replicates (n=) is indicated above each plot and letters at the top of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD). **M)** Histograms summarizing the mean number of mesophyll cells separating veins in WT W22 (blue), *Zmnkd1-Ds;Zmnkd2-Ds* (green), *Zmscr1-m2;Zmscr1h-m1* (yellow) and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* (purple) mutants. Error bars are standard error of the mean. Sample sizes (n=) are biological replicates and letters in the top right corner (beneath sample sizes) of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype.

Figure 8. Embryonic leaf phenotypes reveal a NKD-mediated maternal effect. A) Quantification of the % of fused veins in leaf 3 of various genotypes. Open circles are data points from distinct biological replicates, red crosses indicate the mean for each genotype. Note that WT have no fused veins (see Fig. 7K). **B)** Transverse section of a wild-type (WT) W22 embryo taken across the tip of the meristem, such that P1-P4 are visible. Scale bar: 100 μm. **C)** Quantification of vein density in P4 primordia. Open circles indicate measurements from independent biological replicates, and black crosses indicate the mean for each genotype. **D-H)** Transverse sections of P4 primordia from WT W22 (D), *Zmnkd1-Ds;Zmnkd2-Ds* (E), *Zmscr1-m2;Zmscr1h-m1* (F), *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* from a heterozygous *Zmscr1-m2+;Zmscr1h-m1;Zmnkd1-Ds /+;Zmnkd2-Ds/+* parent (G) and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* from a homozygous *Zmscr1-m2/+;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* parent (H). Fused veins are noticeable in (H) (black arrows). Scale bars: 50 μm.

SUPPLEMENTARY INFORMATION

Figure S1. Summary of setaria CRISPR design. A) Cartoon summaries of *SvSCR1/2* and *SvNKD* genes with guide positions indicated by black arrows above each gene sequence. 5' and 3' untranslated regions are depicted in purple, coding sequences in green and introns in orange. **B)** Guide RNA names and sequences. Guides that successfully edited are highlighted green. **C)** Cartoon summary of Level 1 and Level 2 Golden Gate constructs designed for this study. Promoters are indicated by arrows, and expression modules by rectangles. **D)** Genotyping primers used in this study.

Figure S2. *Svscr1* and *Svscr2* single mutants are phenotypically indistinguishable from wild-type. A-E)** Images of wild-type (WT) ME034V (A), *Svscr1-m1* (B), *Svscr1-m2* (C), *Svscr2-m1* (D) and *Svscr2-m2* (E) plants taken 21 days after sowing. Scale bars: 10cm.

Figure S3. Primers used for quantitative RT-PCR. A) Primer sequences used for quantitative RT-PCR of each gene in this study. **B)** Melt-curves for each primer pair.

Figure S4. Summary of rice CRISPR design. A) Cartoon depiction of *OsNKD* with guide positions indicated above the gene with black arrows. 5' and 3' untranslated regions are depicted in purple, coding sequences in green and introns in orange. **B)** Guide RNA sequences used in this study. **C)** Cartoon summary of Level 2 Golden Gate constructs used to generate *Osnkd* (17821) and *Osscr1;Osscr2;Osnkd* (17827 and 17828) mutants. **D)** Primer sequences and restriction digests used to genotype *OsNKD* editing events.

Figure S5. *nkd* mutants do not have perturbed seed development in setaria or rice. A-D)** Images of setaria (A & B) and rice (C & D) seed with (A & C) or without (B & D) the husk for both wild-type (WT) (top rows) and *nkd* mutants (bottom rows). **E-G)** Cross sections of setaria WT ME034V (E), *Svnkd-m1* (line 1) (F) and *Svnkd-m1* (line 2) (G) mature seed. The aleurone layer is indicated by Al in each image.

Figure S6. *Osnkd* and *Svnkd* mutants do not exhibit perturbed leaf development. A-D)** Stomatal impressions of the abaxial surface of wild-type (WT) Kitaake rice (A), *Osnkd-m6* (B), WT setaria ME034V (C) and *Svnkd-m1* (D) leaves. Rice images are taken of leaf 5 and setaria images of leaf 3. Stomata are false coloured orange. Scale bars: 100 μm. **E, F)** Cross sections of WT Kitaake (E) and *Osnkd-m6* (F) leaf 5 taken from the midpoint along the proximal-distal axis. Scale bars: 200 μm.

Figure S7. *Svscr1;Svscr2;Svnkd* mutant leaves exhibit occasional fused veins with no intervening mesophyll cells.** Quantification of the number of M cells in five setaria genotypes: wild-type ME034V, *Svscr1-m1;Svscr2-m3*, *Svscr1-m2;Svscr2-m1*, *Svscr1-m4;Svscr2-m1;Svnkd-m1* and

Svscr1-m1;Svscr2-m2;Svnkd-m1. Quantification undertaken on leaf 4. Bars are the standard error of the mean. Biological replicates (n=) are indicated and letters in the top right corner of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of M cells in each genotype.

Figure S8. Quadruple *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* mutants exhibit increased fused leaf veins in a variety of contexts. A) Quantification of the percentage of fused veins in leaf 3 of *Zmscr1-m2;Zmscr1h-m1* mutants pre- and post-outcross to *Zmnkd1-Ds;Zmnkd2-Ds*. Mutants phenotyped post-outcross were also segregating for *Zmnkd1-Ds* and *Zmnkd2-Ds* (genotypes labelled on figure). **B)** Quantification of the percentage of fused veins in leaf 3 of wild-type (WT) W22, *Zmscr1-m2;Zmscr1h-m1* and two additional quadruple mutants from two outcrosses using independent *Zmnkd1-Ds;Zmnkd2-Ds* plants. **C-E)** Transverse sections of leaf 6 from WT W22, *Zmscr1-m2;Zmscr1h-m1* and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* lines. In (E) fused veins are indicated by arrows. Scale bars: 100 µm. **F)** Quantification of the percentage of fused veins in WT W22, *Zmscr1-m2;Zmscr1h-m1* and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* leaf 6. In (A), (B) and (F) open circles are data points from biological replicates, and black crosses indicate the mean for each genotype. Letters above each genotype indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD).

Figure S9. *Zmnkd2-Ds* genotyping. Genotyping of the *Zmnkd2-Ds* allele revealing a 786bp deletion.

Table S1. Parental and zygotic genotypes of different combinations of *scr* and *nkd* mutant alleles in maize, along with the presence or absence of fused vein phenotypes.

REFERENCES

Chang, Y.-M., Liu, W.-Y., Shih, A. C.-C., Shen, M.-N., Lu, C.-H., Lu, M.-Y. J., Yang, H.-W., Wang, T.-Y., Chen, S. C.-C., Chen, S. M., Li, W.-H., & Ku, M. S. B. (2012). Characterizing regulatory and functional differentiation between maize mesophyll and bundle sheath cells by transcriptomic analysis. *Plant Physiology*, *160*(1), 165–177. https://doi.org/10.1104/pp.112.203810

Concordet, J.-P., & Haeussler, M. (2018). CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Research*, *46*(W1), W242–W245. https://doi.org/10.1093/NAR/GKY354

Cui, H., Kong, D., Liu, X., & Hao, Y. (2014). SCARECROW, SCR-LIKE 23 and SHORT-ROOT control bundle sheath cell fate and function in Arabidopsis thaliana. *The Plant Journal : For Cell and Molecular Biology*, *78*(2), 319–327. https://doi.org/10.1111/tpj.12470

Cui, H., Levesque, M. P., Vernoux, T., Jung, J. W., Paquette, A. J., Gallagher, K. L., Wang, J. Y., Blilou, I., Scheres, B., & Benfey, P. N. (2007). An evolutionarily conserved mechanism delimiting SHR movement defines a single layer of endodermis in plants. *Science (New York, N.Y.)*, *316*(5823), 421–425. https://doi.org/10.1126/science.1139531

Debernardi, J. M., Tricoli, D. M., Ercoli, M. F., Hayta, S., Ronald, P., Palatnik, J. F., & Dubcovsky, J. (2020). A GRF–GIF chimeric protein improves the regeneration efficiency of transgenic plants. *Nature Biotechnology 2020 38:11*, *38*(11), 1274–1279. https://doi.org/10.1038/s41587-020-0703-0

Dhondt, S., Coppens, F., de Winter, F., Swarup, K., Merks, R. M. H., Inzé, D., Bennett, M. J., & Beemster, G. T. S. (2010). SHORT-ROOT and SCARECROW regulate leaf growth in Arabidopsis by stimulating S-phase progression of the cell cycle. *Plant Physiology*, *154*(3), 1183–1195. https://doi.org/10.1104/pp.110.158857

Finley, T., Chappell, H., & Veena, V. (2021). Agrobacterium-Mediated Transformation of Setaria viridis, a Model System for Cereals and Bioenergy Crops. *Current Protocols*, *1*(5). https://doi.org/10.1002/CPZ1.127

Fouracre, J. P., Ando, S., & Langdale, J. A. (2014). Cracking the Kranz enigma with systems biology. *Journal of Experimental Botany*, *65*(13), 3327–3339. https://doi.org/10.1093/jxb/eru015

Gallagher, K. L., Paquette, A. J., Nakajima, K., & Benfey, P. N. (2004). Mechanisms regulating SHORT-ROOT intercellular movement. *Current Biology : CB*, *14*(20), 1847–1851. https://doi.org/10.1016/J.CUB.2004.09.081

Gontarek, B. C., Neelakandan, A. K., Wu, H., & Becraft, P. W. (2016). NKD Transcription Factors Are Central Regulators of Maize Endosperm Development. *The Plant Cell*, *28*(12), 2916–2936. https://doi.org/10.1105/TPC.16.00609

Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., Putnam, N., & Rokhsar, D. S. (2012). Phytozome: a comparative platform for green

plant genomics. *Nucleic Acids Research*, *40*(D1), D1178–D1186. https://doi.org/10.1093/nar/gkr944

Grass Phylogeny Working Group II. (2012). New grass phylogeny resolves deep evolutionary relationships and discovers C4 origins. *New Phytologist*, *193*(2), 304–312. https://doi.org/10.1111/j.1469-8137.2011.03972.x

Haberlandt, G. (1896). *Physiologische Pflanzenanatomie* (E. Wilhelm, Ed.).

Hahn, F., Korolev, A., Sanjurjo Loures, L., & Nekrasov, V. (2020). A modular cloning toolkit for genome editing in plants. *BMC Plant Biology*, *20*(1), 1–10. https://doi.org/10.1186/S12870-020-02388-2/FIGURES/6

Hirano, Y., Nakagawa, M., Suyama, T., Murase, K., Shirakawa, M., Takayama, S., Sun, T., & Hakoshima, T. (2017). Structure of the SHR–SCR heterodimer bound to the BIRD/IDD transcriptional factor JKD. *Nature Plants*, *3*(3), 17010. https://doi.org/10.1038/nplants.2017.10

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*, *35*(2), 518–522. https://doi.org/10.1093/molbev/msx281

Hughes, T. E., & Langdale, J. A. (2022). SCARECROW is deployed in distinct contexts during rice and maize leaf development. *Development (Cambridge, England)*, *149*(7). https://doi.org/10.1242/DEV.200410/274715/AM/SCARECROW-IS-DEPLOYED-IN-DISTINCT-CONTEXTS-DURING

Hughes, T. E., Sedelnikova, O. v., Wu, H., Becraft, P. W., & Langdale, J. A. (2019). Redundant SCARECROW genes pattern distinct cell layers in roots and leaves of maize. *Development (Cambridge, England)*, *146*(14). https://doi.org/10.1242/dev.177543

Ingram, G. C. (2020). Family plot: the impact of the endosperm and other extra-embryonic seed tissues on angiosperm zygotic embryogenesis. *F1000Research*, *9*. https://doi.org/10.12688/F1000RESEARCH.21527.1

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/MOLBEV/MST010

Koizumi, K., Hayashi, T., & Gallagher, K. L. (2012). SCARECROW reinforces SHORT-ROOT signaling and inhibits periclinal cell divisions in the ground tissue by maintaining SHR at high levels in the endodermis. *Plant Signaling & Behavior*, *7*(12), 1573. https://doi.org/10.4161/PSB.22437

Laurenzio, L. di, Wysocka-Diller, J., & Malamy, J. (1996). The SCARECROW Gene Regulates an Asymmetric Cell Division That Is Essential for Generating the Radial Organization of the Arabidopsis Root. *Cell*, *86*(3), 423–433. http://www.sciencedirect.com/science/article/pii/S0092867400801154

Li, P., Ponnala, L., Gandotra, N., Wang, L., Si, Y., Tausta, S. L., Kebrom, T. H., Provart, N., Patel, R., Myers, C. R., Reidel, E. J., Turgeon, R., Liu, P., Sun, Q., Nelson, T., & Brutnell, T. P. (2010). The

developmental dynamics of the maize leaf transcriptome. *Nature Genetics*, *42*(12), 1060–1067. https://doi.org/10.1038/ng.703

Lin, Y., Zhang, C., Lan, H., Gao, S., Liu, H., Liu, J., Cao, M., Pan, G., Rong, T., & Zhang, S. (2014). Validation of Potential Reference Genes for qPCR in Maize across Abiotic Stresses, Hormone Treatments, and Tissue Types. *PLoS ONE*, *9*(5), e95445. https://doi.org/10.1371/journal.pone.0095445

Liu, Q., Teng, S., Deng, C., Wu, S., Li, H., Wu, J., Cui, X., Zhang, Z., Paul Quick, W., Sun, X., Lu, T., Sun sunxuehui, X., & Brutnell, T. P. (2022). A SHR-IDD-PIN regulatory network mediates minor vein differentiation in rice. *BioRxiv*, 2022.09.22.509058. https://doi.org/10.1101/2022.09.22.509058

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. *Methods*, *25*(4), 402–408. https://doi.org/10.1006/meth.2001.1262

Long, Y., Smet, W., Cruz-Ramírez, A., Castelijns, B., de Jonge, W., Mähönen, A. P., Bouchet, B. P., Perez, G. S., Akhmanova, A., Scheres, B., & Blilou, I. (2015). Arabidopsis BIRD Zinc Finger Proteins Jointly Stabilize Tissue Boundaries by Confining the Cell Fate Regulator SHORT-ROOT and Contributing to Fate Specification. *The Plant Cell*, *27*(4), 1185–1199. https://doi.org/10.1105/tpc.114.132407

Nakajima, K., Sena, G., Nawy, T., & Benfey, P. N. (2001). Intercellular movement of the putative transcription factor SHR in root patterning. *Nature*, *413*(6853), 307–311. https://doi.org/10.1038/35095061

Nunes, T. D. G., Zhang, D., & Raissig, M. T. (2020). Form, development and function of grass stomata. *The Plant Journal*, *101*(4), 780–799. https://doi.org/10.1111/TPJ.14552

Ogasawara, H., Kaimi, R., Colasanti, J., & Kozaki, A. (2011). Activity of transcription factor JACKDAW is essential for SHR/SCR-dependent activation of SCARECROW and MAGPIE and is modulated by reciprocal interactions with MAGPIE, SCARECROW and SHORT ROOT. *Plant Molecular Biology*, *77*(4–5), 489–499. https://doi.org/10.1007/s11103-011-9826-5

Sage, R. F., Christin, P.-A., & Edwards, E. J. (2011). The C4 plant lineages of planet Earth. *Journal of Experimental Botany*, *62*(9), 3155–3169. https://doi.org/10.1093/jxb/err048

Schellmann, S., Schnittger, A., Kirik, V., Wada, T., Okada, K., Beerman, A., Thumfahrt, J., Jürgens, G., & Hülskamp, M. (2002). TRIPTYCHON and CAPRICE mediate lateral inhibition during trichome and root hair patterning in Arabidopsis. *The EMBO Journal*, *21*(19), 5036. https://doi.org/10.1093/EMBOJ/CDF524

Schuler, M. L., Sedelnikova, O. v., Walker, B. J., Westhoff, P., & Langdale, J. A. (2018). SHORTROOT-Mediated Increase in Stomatal Density Has No Impact on Photosynthetic Efficiency. *Plant Physiology*, *176*(1), 757–772. https://doi.org/10.1104/PP.17.01005

Sedelnikova, O. v., Hughes, T. E., & Langdale, J. A. (2018). Understanding the Genetic Basis of C_4 Kranz Anatomy with a View to Engineering C_3 Crops. *Annual Review of Genetics*, *52*(1), 249–270. https://doi.org/10.1146/annurev-genet-120417-031217

Slewinski, T. L., Anderson, A. A., Zhang, C., & Turgeon, R. (2012). Scarecrow plays a role in establishing Kranz anatomy in maize leaves. *Plant & Cell Physiology*, *53*(12), 2030–2037. https://doi.org/10.1093/pcp/pcs147

Swigonova, Z., Lai, J., Ma, J., Ramakrishna, W., Llaca, V., Bennetzen, J. L., & Messing, J. (2004). On the tetraploid origin of the maize genome. *Comparative and Functional Genomics*, *5*(3), 281–284. https://doi.org/10.1002/cfg.395

Torii, K. U. (2012). Two-dimensional spatial patterning in developmental systems. *Trends in Cell Biology*, *22*(8), 438–446. https://doi.org/10.1016/J.TCB.2012.06.002

Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., & Minh, B. Q. (2016). W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research*, *44*(W1), W232–W235. https://doi.org/10.1093/nar/gkw256

Wang, P., Kelly, S., Fouracre, J. P., & Langdale, J. A. (2013). Genome-wide transcript analysis of early maize leaf development reveals gene cohorts associated with the differentiation of C4 Kranz anatomy. *The Plant Journal*, *75*(4), 656–670. https://doi.org/10.1111/tpj.12229

Welch, D., Hassan, H., Blilou, I., Immink, R., Heidstra, R., & Scheres, B. (2007). Arabidopsis JACKDAW and MAGPIE zinc finger proteins delimit asymmetric cell division and stabilize tissue boundaries by restricting SHORT-ROOT action. *Genes & Development*, *21*(17), 2196–2204. https://doi.org/10.1101/gad.440307

Wu, Z., Chen, L., Yu, Q., Zhou, W., Gou, X., Li, J., & Hou, S. (2019). Multiple transcriptional factors control stomata development in rice. *New Phytologist*, *223*(1), 220–232. https://doi.org/10.1111/NPH.15766

Yi, G., Neelakandan, A. K., Gontarek, B. C., Vollbrecht, E., & Becraft, P. W. (2015). The naked endosperm genes encode duplicate INDETERMINATE domain transcription factors required for maize endosperm cell patterning and differentiation. *Plant Physiology*, *167*(2), 443–456. https://doi.org/10.1104/pp.114.251413

Zeng, S. M., Lo, E. K. W., Hazelton, B. J., Morales, M. F., & Torii, K. U. (2020). Effective range of non-cell autonomous activator and inhibitor peptides specifying plant stomatal patterning. *Development (Cambridge)*, *147*(17). https://doi.org/10.1242/DEV.192237/266967/AM/EFFECTIVE-RANGE-OF-NON-CELL-AUTONOMOUS-ACTIVATOR

Zhao, S., & Fernald, R. D. (2005). Comprehensive algorithm for quantitative real-time polymerase chain reaction. *Journal of Computational Biology*, *12*(8), 1047–1064. https://doi.org/10.1089/cmb.2005.12.1047

Figure 1. Mutations in *SvSCR1* and *SvSCR2* affect plant growth. A) Sequences of mutant *Svscr1* and *Svscr2* alleles in two independent lines (C402052-6 & C402054-7). Wild-type (WT) sequence is shown on top, with the sequence of the mutant allele shown beneath. Guide sequences are depicted in blue, and mismatches between the WT and mutant sequence indicated in red. **B-D)** Photos of WT ME034V (B), *Svscr1-m3;Svscr2-m4* (C) and *Svscr1-m2-Svscr2-m1* (D) plants taken 20 days after sowing. Scale bars: 10cm.

Figure 2. *Svscr1;Svscr2* mutant plants have no stomata and fewer mesophyll cells separating veins. A-F) Stomatal impressions of wild-type (WT) ME034V (A,D), *Svscr1-m1;Svscr2-m4* (B,E) and *Svscr1-m2;Svscr2-m1* (C,F) leaf 3 from either the abaxial (A-C) or adaxial (D-F) surface. Stomata are false coloured orange. Scale bars: 100µm. **G-H)** Transverse sections imaged using UV illumination of WT ME034V (G) and *Svscr1-m1;Svscr2-m4* (H) leaf 4, taken from the mid-point along the proximal-distal axis. The number of mesophyll cells between each pair of veins is indicated above the leaf. Scale bars: 100µm. **I)** Histograms summarizing the mean number of mesophyll cells separating veins in WT ME034V and two independent *Svscr1;Svscr2* mutants. The number of biological replicates is indicated above each plot and letters in the top right corner of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the average number of mesophyll cells in each genotype.

Figure 3. *NKD* **and** *SCR* **transcripts accumulate in the same spatial domain with levels determined by a feedback loop. A)** Quantitative RT-PCR of *ZmSCR1* and *ZmSCR2* transcripts in the *Zmnkd1-Ds;Zmnkd2-Ds* mutant, and *ZmNKD1* and *ZmNKD2* in the *Zmscr1-m2;Zmscr1h-m2* mutant. Open circles are individual biological replicates, black crosses indicate the mean for each genotype. Statistical significance as calculated on log-transformed fold change data by Student's *t*-test (two-tailed) indicated above each plot: *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$. **B)** *In-situ* hybridization to *ZmNKD1* and *ZmSCR1* in maize wild-type B73 apices. In each image the P2 primordium is outlined in blue, P3 in red, P4 in purple and P5 in green, as indicated in the adjacent cartoon diagram. Darker purple signal represents successful hybridization to each transcript of interest. Scale bars: 50μm **C)** Maximum likelihood phylogeny of the *NKD* genes in monocots. The *NKD* clade is highlighted in green, and the adjacent monocot clade in purple. Bootstrap values are displayed at each branch of the phylogeny.

Figure 4. *nkd* loss of function mutations do not perturb growth in maize, rice or setaria. A) Cartoon depictions of loss of function *nkd* alleles in maize (*Zm*), setaria (*Sv*) and rice (*Os*). In each case, 5' and 3' untranslated regions are depicted in purple, introns in orange and coding regions in green. Transposon insertions are indicated by a yellow triangle. For setaria and rice, CRISPR guide sequences are indicated in blue and maroon respectively, with the sequence of the mutant allele beneath with edits highlighted in red. **B)** Whole plant phenotypes of maize, setaria and rice *nkd* mutants. Photos were taken 31 days (maize), 20 days (setaria) or 14 days (rice) after sowing. Scale bars: 10cm. **C-F)** Transverse sections of maize wild-type (WT) W22 (C), *Zmnkd1-Ds;Zmnkd2-Ds* (D), setaria WT ME034V (E) and *Svnkd-m1* (F), imaged under either brightfield (maize) or UV illumination (setaria). Images were taken at the mid-point along the proximal-distal axis of either leaf 5 (maize) or leaf 4 (setaria). Scale bars: 100μm. **G-H)** Quantification of vein density (G) and the ratio of rank1:rank2 intermediate veins (H) of WT W22 (blue) and *Zmnkd1-Ds;Zmnkd-Ds* (green) mutants. Open circles indicate distinct biological replicates, black crosses indicate the mean for each genotype. Statistical significance as calculated by Student's *t*-test (two-tailed) indicated above each plot: n.s. *P*>0.05. **I-J)** Histograms summarising the mean number of mesophyll cells separating vascular bundles in WT W22 versus *Zmnkd1-Ds;Zmnkd2-Ds* (I) and WT ME034V versus two independent *Svnkd-m1* lines (J). Error bars are the standard error of the mean, and the number of biological replicates is indicated above each plot. Letters in the top right corner of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype. WT is coloured blue and *nkd* mutants green.

Figure 5. Loss of function *nkd* mutations enhance growth perturbations in *scr* mutants of maize and setaria but not rice. A) Sequences of mutant *scr1;scr2;nkd* alleles in three independent lines of both rice and setaria. Wild-type sequences are shown above with guides highlighted in either maroon (rice) or blue (setaria). Edits are highlighted in red. Note *Svscr1-m2* allele is included in Figure 1. **B-O)** Whole plant phenotypes of rice (B-D), setaria (E-I) and maize (J-O). Images were taken 14, 20 or 21 days after sowing for rice, setaria and maize respectively. Plants in (H) and (I) are from segregating families in each case, such that *Svscr1;Svscr2* and *Svscr1;Svscr2;Svnkd* in each panel are siblings. Scale bars: 10cm.

Figure 6. Loss of function *nkd* mutations induce the formation of fused leaf veins in *scr* mutants of setaria but not rice. A-B) Transverse sections of *Osscr1-m7;Osscr2-m3* (A) and *Osscr1-m7;Osscr2-m9;Osnkd-m3* (B) mutant leaves, taken at the midpoint along the proximal-distal axis of leaf 5. **C)** Histograms summarizing the mean number of mesophyll cells separating veins in two independent *Osscr1;Osscr2* (yellow) and three independent *Osscr1;Osscr2;Osnkd* (purple) mutant lines. Error bars are standard error of the mean. Sample sizes (n=) are biological replicates. **D-E)** Transverse sections of *Svscr1-m1;Svscr2-m4* (D) and *Svscr1-m2;Svscr2-m1;Svnkd-m1* (E) mutant leaves, taken at the midpoint along the proximal-distal axis of leaf 4, imaged under UV illumination. The number of mesophyll cells separating veins is displayed above each region. In (E) the inset shows an example of a fused vein with no separating mesophyll cells. **F)** Histograms summarizing the mean number of mesophyll cells separating veins in wild-type ME034V (blue), *Svscr1-m1;Svscr2-m4* (yellow) and *Svscr1-m4;Svscr2-m1;Svnkd-m1* (purple) mutants. Error bars are standard error of the mean. Sample sizes (n=) are biological replicates and letters in the top right corner (beneath sample sizes) of each plot indicate statistically different groups ($P \leq 0.05$, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype. Scale bars: 100 μm.

Figure 7. *Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2* quadruple mutants exhibit a significant increase in the number of fused leaf veins compared to *Zmscr1;Zmscr1h* double mutants. A-H) Transverse cross sections of wild-type (WT) W22 (A & C), *Zmnkd1-Ds;Zmnkd2-Ds* (B & D), *Zmscr1-m2;Zmscr1h-m1* (E & G) and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* (F & H) leaf 3, from the mid-point along the proximal distal axis and imaged under either brightfield (A-B, E-F) or UV (C-D, G-H) illumination. Scale bars: 100 μm. **I-L)** Strip charts summarising quantification of vein density (I), the ratio of rank 1: rank 2 intermediate veins (J), the % of fused veins (K) and the % of veins formed in runs of ≥3 fused veins (L). Open circles indicate measurements from independent biological replicates, and black crosses indicate the mean for each genotype. The number of biological replicates (n=) is indicated above each plot and letters at the top of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD). **M)** Histograms summarizing the mean number of mesophyll cells separating veins in WT W22 (blue), *Zmnkd1-Ds;Zmnkd2-Ds* (green), *Zmscr1-m2;Zmscr1h-m1* (yellow) and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* (purple) mutants. Error bars are standard error of the mean. Sample sizes (n=) are biological replicates and letters in the top right corner (beneath sample sizes) of each plot indicate statistically different groups (*P*≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of mesophyll cells in each genotype.

Figure 8. Embryonic leaf phenotypes reveal a NKD-mediated maternal effect. A) Quantification of the % of fused veins in various genotypes. Open circles are data points from distinct biological replicates, red crosses indicate the mean for each genotype. Note that WT have no fused veins (see Fig. 7K). **B)** Transverse section of a wild-type (WT) W22 embryo taken across the tip of the meristem, such that P1-P4 are visible. Scale bar: 100 μm. **C)** Quantification of vein density in P4 primordia. Open circles indicate measurements from independent biological replicates, and black crosses indicate the mean for each genotype. **D-H)** Transverse sections of P4 primordia from WT W22 (D), *Zmnkd1-Ds;Zmnkd2-Ds* (E), *Zmscr1-m2;Zmscr1h-m1* (F), *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* from a heterozygous *NKD* parent (G) and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* from a homozygous *nkd* parent (H). Fused veins are noticeable in (H) (black arrows). Scale bars: 50 μm.

Figure S1. Summary of setaria CRISPR design. A) Cartoon summaries of *SvSCR1/2* and *SvNKD* genes with guide positions indicated by black arrows above each gene sequence. 5' and 3' untranslated regions are depicted in purple, coding sequences in green and introns in orange. **B)** Guide RNA names and sequences. Guides that successfully edited are highlighted green. **C)** Cartoon summary of Level 1 and Level 2 Golden Gate constructs designed for this study. Promoters are indicated by arrows, and expression modules by rectangles. **D)** Genotyping primers used in this study.

Figure S2. ***Svscr1*** **and** ***Svscr2*** **single mutants are phenotypically indistinguishable from wild-type. A-E)** Images of wild-type ME034V (A), *Svscr1-m1* (B), *Svscr1-m2* (C), *Svscr2-m1* (D) and *Svscr2-m2* (E) plants taken 21 days after sowing. Scale bars: 10cm.

A

Gene	Accession/ref	Forward primer sequence (5'-3')	Reverse primer sequence (5'-3')	Product size
ZmSCR1	GRMZM2G131516	GGTGTCGATCGCGTATCCTT	TAGAACGCGCATGAAGAGCA	98bp
ZmSCR1h	GRMZM2G015080	GGCAGATCAACCTTGTTTAGCC	AGCCTTGACCTCAGCTCATG	103bp
ZmNKD1	GRMZM2G129261	AATGAGGGCGTGAATTGCAG	TGCATGCATCCTGACACAAG	149bp
ZmNKD2	GRMZM5G884137	GCCAACAAGCCTTTCCATGG	AGTTCCCGCCAAAAATCCCT	241bp
ZmCYP	Lin et al. (2014)	CTGAGTGGTGGTCTTAGT	AACACGAATCAAGCAGAG	100bp
ZmEF1α	Lin et al. (2014)	TGGGCCTACTGGTCTTACTACTGA	ACATACCCACGCTTCAGATCCT	135bp

B

Figure S3. Primers used for quantitative RT-PCR. A) Primer sequences used for quantitative RT-PCR of each gene in this study. **B)** Melt-curves for each primer pair.

Figure S4. Summary of rice CRISPR design. A) Cartoon depiction of *OsNKD* with guide positions indicated above the gene with black arrows. 5' and 3' untranslated regions are depicted in purple, coding sequences in green and introns in orange. **B)** Guide RNA sequences used in this study. **C)** Cartoon summary of Level 2 Golden Gate constructs used to generate *Osnkd* (17821) and *Osscr1;Osscr2;Osnkd* (17827 and 17828) mutants. **D)** Primer sequences and restriction digests used to genotype *OsNKD* editing events.

Figure S5. *nkd* **mutants do not have perturbed seed development in setaria or rice. A-D)** Images of setaria (A & B) and rice (C & D) seed with (A & C) or without (B & D) the husk for both wild-type (WT) (top rows) and *nkd* mutants (bottom rows). **E-G)** Cross sections of setaria WT ME034V (E), *Svnkd-m1* (line 1) (F) and *Svnkd-m1* (line 2) (G) mature seed. The aleurone layer is indicated by Al in each image.

Figure S6. *Osnkd* and *Svnkd* mutants do not exhibit perturbed leaf development. A-D) Stomatal impressions of the abaxial surface of wild-type (WT) Kitaake rice (A), *Osnkd-m6* (B), WT setaria ME034V (C) and *Svnkd-m1* (D) leaves. Rice images are taken of leaf 5 and setaria images of leaf 3. Stomata are false coloured orange. Scale bars: 100 μm. **E, F)** Cross sections of WT Kitaake (E) and *Osnkd-m6* (F) leaf 5 taken from the midpoint along the proximal-distal axis. Scale bars: 200 μm.

Figure S7. *Svscr1;Svscr2;Svnkd* mutant leaves exhibit occasional fused veins with no intervening mesophyll cells. Quantification of the number of mesophyll cells in five setaria genotypes: Wild-type ME034V, *Svscr1-m1;Svscr2-m3*, *Svscr1-m2;Svscr2-m1*, *Svscr1-m4;Svscr2-m1;Svnkd-m1* and *Svscr1-m1;Svscr2-m2;Svnkd-m1*. Quantification undertaken on leaf 4. Bars are the standard error of the mean. Biological replicates (n=) are indicated and letters in the top right corner of each plot indicate statistically different groups (P≤0.05, one-way ANOVA and Tukey's HSD) calculated using the mean number of M cells in each genotype.

Figure S8. Quadruple *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* mutants exhibit increased fused leaf veins in a variety of contexts. A) Quantification of the percentage of fused veins in leaf 3 of *Zmscr1-m2;Zmscr1h-m1* mutants pre- and post-outcross to *Zmnkd1-Ds;Zmnkd2-Ds*. Mutants phenotyped post-outcross were also segregating for *Zmnkd1-Ds* and *Zmnkd2-Ds* (genotypes labelled on figure). **B)** Quantification of the percentage of fused veins in leaf 3 of wild-type (WT) W22, *Zmscr1-m2;Zmscr1h-m1* and two additional quadruple mutants from two outcrosses using independent *Zmnkd1-Ds;Zmnkd2-Ds* plants. **C-E)** Transverse sections of leaf 6 from WT W22, *Zmscr1-m2;Zmscr1h-m1* and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* lines. In (E) fused veins are indicated by arrows. Scale bars: 100 μm. **F)** Quantification of the percentage of fused veins in WT W22, *Zmscr1-m2;Zmscr1h-m1* and *Zmscr1-m2;Zmscr1h-m1;Zmnkd1-Ds;Zmnkd2-Ds* leaf 6. In (A), (B) and (F) open circles are data points from biological replicates, and black crosses indicate the mean for each genotype. Letters above each genotype indicate statistically different groups (P≤0.05, one-way ANOVA and Tukey's HSD).

Figure S9. *Zmnkd2-Ds* **genotyping.** Genotyping of the *Zmnkd2-Ds* allele revealing a 786bp deletion.

TABLE S1. Parental and zygotic genotypes of different combinations of *scr* and *nkd* mutant alleles in maize, along with the presence or absence of fused vein phenotypes

Plant	Genotype	Genotype	Experiment	Fused_veins	NKD1 in parent	NKD2 in parent
280-6-1	*Zmscr1;Zmscr1h;ZmNKD1;ZmNKD2*	A	Sep-20	6.35	het	het
280-6-12	*Zmscr1;Zmscr1h;Zmnkd1/+;ZmNKD2*	B	Sep-20	14.29	het	het
280-6-5	*Zmscr1;Zmscr1h;Zmnkd1/+;ZmNKD2*	B	Sep-20	16.33	het	het
283-4-5	*Zmscr1;Zmscr1h;Zmnkd1/+;ZmNKD2*	B	Jul-20	15.69	het	WT
283-4-17	*Zmscr1;Zmscr1h;Zmnkd1/+;ZmNKD2*	B	Jul-20	13.64	het	WT
280-6-9	*Zmscr1;Zmscr1h;Zmnkd1/+;Zmnkd2/+*	C	Sep-20	13.73	het	het
280-6-10	*Zmscr1;Zmscr1h;ZmNKD1;Zmnkd2*	D	Sep-20	27.27	het	het
284-3-1	*Zmscr1;Zmscr1h;ZmNKD1;Zmnkd2*	D	Jul-20	0.00	het	mutant
280-6-7	*Zmscr1;Zmscr1h;Zmnkd1;ZmNKD2*	E	Sep-20	46.67	het	het
278-2-16	*Zmscr1;Zmscr1h;Zmnkd1;ZmNKD2*	E	Sep-20	44.23	mutant	WT
283-4-16	*Zmscr1;Zmscr1h;Zmnkd1;ZmNKD2*	E	Jul-20	30.30	het	WT
282-3-16	*Zmscr1;Zmscr1h;Zmnkd1/+;Zmnkd2*	F	Sep-20	26.09	het	mutant
282-3-27	*Zmscr1;Zmscr1h;Zmnkd1/+;Zmnkd2*	F	Sep-20	15.87	het	mutant
280-6-2	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2/+*	G	Sep-20	17.39	het	het
280-6-11	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2/+*	G	Sep-20	34.15	het	het
280-6-4	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2*	H	Sep-20	12.90	het	het
282-3-3	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2*	H	Sep-20	14.29	het	mutant
282-3-8	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2*	H	Sep-20	24.32	het	mutant
OX20 282-3-9-m1	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2*	I	Mar-22	31.91	mutant	mutant
OX20 282-3-9-m3	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2*	I	Mar-22	25.00	mutant	mutant
OX20 282-3-9-m6	*Zmscr1;Zmscr1h;Zmnkd1;Zmnkd2*	I	Mar-22	28.81	mutant	mutant

Parental genotypes

280-6	*scr1/+;scr1h;nkd1/+;nkd2/+*
283-4	*scr1/+;scr1h;nkd1/+;NKD2*
278-2	*scr1/+;scr1h/+;nkd1;NKD2*
282-3	*scr1/+;scr1h;nkd1/+;nkd2*
282-3-9	*scr1/+;scr1h;nkd1;nkd2*