Relatedness coefficients and their applications

for triplets and quartets of genetic markers

8 Kermit Ritland

9 Biodiversity Research Center

10 University of British Columbia

12 RUNNING HEAD: Relatedness with marker triplets and quartets

14   ABSTRACT

Relatedness coefficients which seek the identity-by-descent of genetic markers are

described. The markers are in groups of two, three or four, and if four, can consist of two

pairs. It is essential to use cumulants (not moments) for four-marker-gene probabilities, as

the covariance of homozygosity, used in 4-marker applications, can only be described with

cumulants. A covariance of homozygosity between pairs of markers arises when

populations follow a mixture distribution. Also, the probability of four markers all

identical-by-descent equals the normalized fourth cumulant. In this paper, a "genetic

marker" generally represents either a gene locus or an allele at a locus.

Applications of three marker coefficients mainly involve conditional regression, and

applications of four marker coefficients can involve identity disequilibrium. Estimation of

relatedness using genetic marker data is discussed. However, three- and four-marker

estimators suffer from statistical and numerical problems, including higher statistical

variance, complexity of estimation formula, and singularity at some intermediate allele

frequencies.

INTRODUCTION

Relatedness is a general term for the level of genetic similarity between individuals, and is measured by the sharing alleles identical-by-descent (MALÉCOT 1948; PAMILO AND CROZIER 1982). Relatedness is quantified with gene identity coefficients, which characterize both the pattern and the frequency of identity-by-descent. The unit of observation is normally a pair of marker loci, and the object of estimation is the kinship coefficient or the coefficient of relationship. In this paper, the unit of observation is extended to triplets and quartets of genes, allowing the opportunity to characterize additional parameters of population structure.

Relatedness may be estimated with genetic markers (MORTON *et al.* 1971) and for pairs of marker genes, many computer programs are available for estimation of relatedness (WANG 2014), in particular for "pairwise relationship", such as the *r* of QUELLER AND GOODNIGHT (1989). However, the equations for pairwise relationship are not extendable to three or four genes, as the covariances and higher moments need to be defined in new ways.

With more than two markers the situation becomes much more complex. Estimators for three and four marker-gene measures of relatedness have recently been proposed. SAMANTA (2009) provided the first estimator for three genes. ACKERMAN *et al.* (2017) examined all four genes and described the estimation of seven of eight coefficients of relatedness. Multiallelic data have information about all eight coefficients, but they used a biallelic model which provides just seven degrees of freedom, constraining their space of estimates. In addition, they did not use cumulants and their moments are normalized differently than will be here.

Cumulants are of use in certain problems in quantitative genetics (BURGER 1991;

TURELLI AND BARTON 1994) and compared to cumulants have more useful theoretical

properties (KENDALL *et al.* 1977), but ordinary moments are sufficient for two and three-

gene fixation indices. RITLAND (1987) found cumulants instead of moments were an

essential component of four-gene fixation indices. Fourth order cumulants are needed to

specify the probability of gene identity for all four genes, and also to describe identity

disequilibrium as the "covariance of covariances". As an example of the necessity of

cumulants, for population gene frequency p, the fourth central moment for four genes,

denoted $(X_1, X_2, X_3$ and $X_4)$, is $E[(X_1 - p)(X_2 - p)(X_3 - p)(X_4 - p)] = \sigma_{12}^2 \sigma_{34}^2 + \sigma_{13}^2 \sigma_{24}^2 +$

$\sigma_{23}^2 \sigma_{13}^2 + \kappa_4$ where σ_{ij}^2 is the covariance of X_i and X_j and κ_4 is a fourth-order cumulant

which does not appear in moment-based treatments.

The probabilities developed in the paper are used to estimate relatedness in a

population with polymorphic genetic markers. Any numbers of alleles at a single locus are

allowed in the equations, but practically, one has two and at most three alleles with SNP

types of data, and for tractability, we use a three-allele model to examine a special situation

with four genes. These models and estimation procedures are readily applicable to the

emerging mountains of genome data.

71   APPLICATIONS OF MARKERS

Definitions. Between pairs of genetic markers, the *coefficient of relationship*

measures the degree of consanguinity (e.g., the probability that markers are identical by

descent, termed ibd). The coefficient relationship equals twice the *kinship coefficient.* The

inbreeding coefficient is the probability that a pair of markers within one individual are ibd.

With more than two markers, the coefficient of relationship is more broadly defined with

groups of markers (two, three or four). With four markers, there are nine *modes of*

Jacquard's gene identity, with ibd genes connected by lines. The *normalized central moment*

gives the probability of ibd of all markers. At the level of four markers, *cumulants* are

necessary to describe *identity disequilibrium* (the excess of identity between marker pairs).

The covariance of cumulants forms the machinery of higher order interactions.

Relatedness and two markers. The two-marker coefficient of relatedness is used for

many inferences with genetic markers, mainly involving pairs of genes sampled between

two individuals ("coefficient of relationship" or pairs of genes sampled within one

individual (the "inbreeding coefficient"). Analysis of data with two marker measures are

ubiquitous (WANG 2014) and the two-marker probabilities are often incorporated into

probabilities of groups of three and four genes.

Regression and three markers. The three-marker relationship coefficient is the

probability that the three marker loci have alleles all identical-by-descent (Figure 1a). This

coefficient, G, is usually combined with two-allele coefficients for biological interpretable

parameters, useful at least for problems involving mating systems or kin selection. In the

theory of mating system estimation, the "effective selfing rate" is the genetically equivalent

rate of selfing caused by all types of biparental inbreeding (RITLAND 1985); the effective

selfing rate of individual A equals $2R$-G, where R is the relatedness between mates and G

the third moment involving the two maternal and single paternal allele (Figure 1b). In the

theory of kin selection, the regression coefficient of relatedness is used and, properly, a

three-gene model is needed (Figure 1c), as shown by MICHOD AND HAMILTON (1980), where

their Equation 18 depends upon whether the reference genotype is homozygous (18a) vs.

heterozygous (18b). Note that both the effective selfing rate and the regression coefficients

of relationship can be asymmetrical when inbreeding coefficients differ between the two

relatives.

Identity disequilibrium and four markers. Between two diploid individuals, there are

15 patterns of gene identity (Liu 2005). A pair of individuals can share two, three or four

markers, and at each level, allelic similarities can describe aspects of relatedness. After

Jacquard (1966); Jacquard (2012), for four marker genes and two individuals, there are

nine condensed identity modes, denoted as Δ_i (Figure 2a). There are eight independent

parameters of relatedness: three pairwise measures (F_A, F_B, R), two three-way measures

(G_A, G_B), and three four-marker measures (F_{AB}, R_{AB}, H). At the highest level, the measures

are much different as the four-marker measures F_{AB}, R_{AB} are covariances (not identities).

The four-marker parameter, H, is the probability that four markers are identical. As well,

the quantities must be defined as cumulants. Cumulants equal moments up to order three,

but fourth-order moments do not equal fourth order cumulants.

The fourth central moment equals $\kappa_4 + F_{AB} + 2R_{AB}$ (note the covariances between

second moments enter this expression) and the normalized cumulant κ_4 (equivalent to *H)*

equals the probability of identity of all four marker genes. While the variance is a measure

of the spread of the distribution, kurtosis is a measure of the "peakedness" of the

distribution of random variables, and infrequent extreme deviations contributing

excessively to this statistic (De la Rosa and Moreno Muñoz 2008).

While applications of three and four gene measures are in their infancy, at least, the

skew and kurtosis as measured by higher moments can help remove bias in DNA forensics

caused by genotyping error (Weir 1994).

In the four-marker model, many possibilities exist about attaching meaning to each

of the Δ_i. One example is the progeny-pair model (Ritland and Leblanc 2004) where A and

B are two progeny of the same mother plant (Figure 2b). At another more abstract level,

two of the genes can be markers and two are quantitative trait loci (Figure 2c). If identity

disequilibrium is present, the regression of phenotypic similarity (QTL) on estimated

relationship (markers) gives an estimate of heritability "in the field" (Ritland 2000).

129   TWO, THREE AND FOUR MARKER PROBABILITIES

At any level of comparison, associations are measured as the frequency of a given

configuration (allele "state") divided by the denominators in Table 1. These denominators

are termed "normalization constants", and are the maximum possible value of the

numerator. Some of these normalized measures of association arise naturally in the

derivations below.

1. Probabilities of two-marker relationship

From Equation (7) of Ritland (1987), which follows Kendall *et al.* (1977 eq. 13.36),

the frequency of gametes with allele i and with allele j is

$$f_{ij} = \kappa_i \kappa_j + \kappa_{ij}$$

138   The two-marker coefficient of relationship can be estimated from the frequencies of

139   each allele in a sample. For any given allele, say A_i, it derived by equating the observed

140   frequency of homozygotes to that expected by the above equation

$$f_{ii} = \mathrm{E}[A_i A_i] = p_i^2 + p_i(1 - p_i)R$$

$$f_{ij} = \mathrm{E}[A_i A_j] = 2p_i p_j(1 - p_i)R$$

The likelihood of the data, given R, $L(R) = \prod_{ij} f_{ij}{}^{X_{ij}}$. Solving for R gives estimators based

upon pairs of alleles A,

$$\hat{R}_{ii} = \frac{f_{ii} - p_i^2}{p_i(1 - p_i)}$$

The estimate of R for allele i, estimates are combined across alleles as

$$\hat{R} = \sum_i w_i \hat{R}_{ii} \qquad\qquad (1)$$

where the weights w_i sum to unity.

These weights are found by finding the w_i that minimize $\mathbf{w}^T \mathbf{V} \mathbf{w}$, where $\mathbf{w}$ is an n

element vector of weights, and $\mathbf{V}$ the nxn variance-covariance matrix of allele-specific

estimates (for details see RITLAND (1996)). The weights require prior specification of true

relatedness. With zero prior R, the weight for allele A_i is $w_i = \frac{1 - p_i}{n - 1}$. An m-allele locus

receives the weight $(n_m\text{-}1)$, giving the estimator for r given by equation 5 in RITLAND (1996).

Probabilities of three-marker relationship

The three-marker relationship coefficient is the probability that three sampled

marker genes are all identical-by-descent. From Equation (7) of RITLAND (1987), which

follows KENDALL *et al.* (1977 eq. 13.36), The joint frequency of markers i, j and k is

$$f_{ijk} = \boldsymbol{\kappa}_i \boldsymbol{\kappa}_j \boldsymbol{\kappa}_k + \boldsymbol{\kappa}_i \boldsymbol{\kappa}_j + \boldsymbol{\kappa}_i \boldsymbol{\kappa}_k + \boldsymbol{\kappa}_j \boldsymbol{\kappa}_k + \boldsymbol{\kappa}_{ijk}$$

158   This written in conventional population genetic terms as

$$f_{ijk} = p_i p_j p_k + p_i p_j v_f + (p_k p_l + p_j p_k) v_r + w_{ijk}$$

159   Where alleles i and j are from one individual and allele k from a second individual. The

160   cumulants are written in bold face to emphasize they have a random component that may

161   covary.

162   From Equation (7) of RITLAND (1987), there are three primary patterns

$$f_{iii} = \mathrm{E}[A_i A_i A_i] = p_i^3(1 - F - 2R + 2G) + p_i^2(F + 2R - 3G) + p_i G$$

$$f_{iij} = \mathrm{E}[A_i A_i A_j] = p_i^2 p_j(1 - F - 2R + 2G) + p_i p_j(F - G)$$

$$f_{ijk} = \mathrm{E}[A_i A_j A_k] = p_i p_j p_k(1 - F - 2R + 2G) \qquad (2)$$

Where the order is irrelevant ($A_i A_i A_j$, $A_j A_i A_j$ and $A_i A_j A_j$ are equivalent). The

genotype frequencies are mixtures of marker gene identity: G is the probability that all

three markers are ibd, R-G is the probability of ibd of one pair of markers, $F + 2R - 3G$ for

two pairs, and and 1-F-2R+2G is the probability of no ibd among the three markers).

Solving for G in Equation (7) gives three probabilities involving G,

$$\hat{G}_{iii} = \frac{f_{iii} - p_i^2(1 - p_i)(F + 2R) - p_i^3}{p_i(1 - p_i)(1 - 2p_i)}$$

$$\hat{G}_{iij} = \frac{f_{iij} - p_i(1 - p_i)p_j(F + 2R) - p_i^2 p_j}{p_i p_j(2p_i - 3)}$$

$$\hat{G}_{ijk} = \frac{f_{ijk} - p_i p_j p_k(F + 2R)}{2 p_i p_j p_k} \qquad (3)$$

G is a normalized third central moment and the normalization constant depends

upon the pattern of subscript.

Each allele can provide an estimate of G, denoted $\hat{G}_i$, and its weighted estimate

across possible alleles i is

$$\hat{G} = \sum_i w_i \hat{G}_i . \qquad (4)$$

This represents a "linear estimator" of G. The weights are derived in the appendix. The

best alternative to linear estimation is maximum likelihood.

Probabilities of four-marker relationship

From Equation (7) of Ritland (1987), which follows Kendall *et al.* (1977 eq. 13.36),

$$f_{ijkl} = \kappa_i\kappa_j\kappa_k\kappa_l + \kappa_i\kappa_j\kappa_{kl} + \kappa_i\kappa_k\kappa_{jl} + \kappa_i\kappa_l\kappa_{jk} + \kappa_j\kappa_k\kappa_{il} + \kappa_j\kappa_l\kappa_{ik} + \kappa_k\kappa_l\kappa_{ij}$$

$$+ \kappa_i\kappa_{jkl} + \kappa_j\kappa_{ikl} + \kappa_k\kappa_{ijl} + \kappa_l\kappa_{ijk} + \kappa_{ij}\kappa_{kl} + \kappa_{ik}\kappa_{jl} + \kappa_{il}\kappa_{jk} + \kappa_{ijkl}$$

181   The cumulants κ_i are similar to moments and covariances and may have a random

182   component that may covary with other cumulants. The subscripts indicate alleles. The

183   recursion equation is

$$f_{ijkl} = p_i p_j p_k p_l + (p_i p_j + p_k p_l)v_f + (p_i p_l + p_j p_k + p_j p_l + p_k p_l)v_r$$

$$+ p_i \omega_{jkl} + p_j \omega_{ikl} + p_k \omega_{ijl} + p_l \omega_{jkl} + v_{ij}^2 v_{kl}^2 + v_{ik}^2 v_{jl}^2 + v_{il}^2 v_{jk}^2$$

$$+ Cov(v_{ij}, v_{kl}) + Cov(v_{ik} v_{jl}) + Cov(v_{il} v_{jk}) + \kappa_{ijkl} \tag{5}$$

Where the *v* terms are second order covariances. When there is a mixture model (which

creates the covariances), each subpopulation m, contributes to the mean cumulant across

pooled m. The term $\kappa_{i,m}\kappa_{j,m}\kappa_{k,m}\kappa_{l,m}$ contributes to all 18 population level moments, the

term $\kappa_{i,m}\kappa_{j,m}\kappa_{kl,m}$ contributes to six population level moments, and so on. However, the

quantitative extent of these contributions are complex and beyond treatment here.

Regardless, that subpopulation cumulants "distill" to the same assortment of cumulants,

albeit with perhaps slightly different values. where the covariance terms are across the

mixture terms m. This is a finite mixture model, needed when a single component

distribution is inadequate (McLachlan *et al.* 2019). These can get complex but Withers *et*

al. (2015) does provide the first known expressions for cumulants used available computer

technology (an equation solver), not available in 1987 to KR. Cumulants are allowed to

vary across the mixture, and that this results in effective covariance between second-order

cumulants. The expectations taken across m result in changes to the above expression due

to associations among the *ijkl* across *m*, that causes the cumulants to be associated in a

certain way, since for example, $E[\kappa_{i,m}\kappa_{j,m}] \neq \kappa_i \kappa_j$.

The associations between *pairs* of markers is termed identity disequilibrium. If one

pair of alleles is heterozygous, it is more likely the second pair is also heterozygous. This is

a four-gene marker measure that has been neglected due to inordinate attention to linkage

disequilibrium. Identity disequilibrium has classically been characterized as the excess of

homozygosity above that expected from the squared gene frequencies (HILL 1975) (OHTA

1980). The identity excess is closely correlated to the expectation of the total squared

linkage disequilibrium (TAKAHATA 1982). Some of the problem is that haploid gametes are

not directly assayed but rather imputed (VITALIS AND COUVET 2001b).

We can add a cumulant to the equation for the probability of identity-by-state. From

equation 3.78 in KENDALL *et al.* (1977), the moments about the mean for two squared

random variables (the alleles present at each locus) equals

$$E[p_i^2 p_j^2] = \kappa_{22} + \kappa_{20}\kappa_{02} + 2\kappa_{11}^2$$

Whose form corresponds to $3\sigma_r^4$ for the fourth central moment with the difference that a

the cumulant κ_{22} is added. VITALIS AND COUVET (2001a) and others have given estimator for

identity disequilibrium which omits this cumulant.

The four-allele case introduces higher-order associations and brings with it new

statistical problems. Among four alleles, two new measures arise. The first is termed H

and is the probability that all four alleles are identical-by-descent. The other two have not

been recognized in the literature, perhaps because they invoke the existence of cumulants,

which differ from the corresponding moments with products of four or more variates.

The first, termed R_{AB}, is the probability that both alleles in the first relatives are

identical-by-descent to both alleles in the second relative. The second, termed F_{AB}, is the

probability that both individuals have both marker genes identical-by-descent. Thus, the

three unique four-allele measures are

223
$$H$$

$$F_{AB} = F_A F_B + Cov(F_A, F_B)$$

224
$$R_{AB} = R^2 + Cov(R, R^c)$$

225   (6)

226   the covariances between second moments, $Cov(F_A, F_B)$ and $Cov(R_{AB}, R'_A B')$, exist only

227   when the distribution of gene frequency follows a mixture distribution where

228   subpopulations vary for F and R.

229   We can rewrite Equation (5) as

$$f_{ijkl} = (2 - \delta_{ij})(2 - \delta_{kl}) \left[p_i p_j p_k p_l \right.$$

$$+ (\delta_{jl} p_i p_k p_j + \delta_{jk} p_i p_l p_j + \delta_{il} p_j p_k p_i + \delta_{ik} p_j p_l p_i - 4 p_i p_j p_k p_l) R$$

231
$$+ (\delta_{ik}\delta_{jl} p_i p_j + \delta_{il}\delta_{jk} p_i p_j - \delta_{jl} p_i p_k p_j - \delta_{jk} p_i p_l p_j - \delta_{il} p_j p_k p_i - \delta_{ik} p_j p_l p_i + 2 p_i p_j p_k p_l) R_{AB}.$$

$$+ p_k p_l (\delta_{ij} p_i - p_i p_j) F_A + p_i p_j (\delta_{kl} p_k - p_k p_l) F_B$$

$$+ (\delta_{ij}\delta_{kl} p_i p_k - \delta_{kl} p_i p_j\, p_k - \delta_{ij} p_i p_k p_l + p_i p_j p_k p_l) F_{AB}$$

$$+ 2(\delta_{ijl} p_k p_{ijl} + \delta_{ijk} p_l p_{ijk} - p_i p_k p_{jl} - p_i p_l p_{jk} - p_j p_k p_{il} - p_j p_l p_{ik} - 2 p_k p_l p_{ij} + 4 p_i p_j p_k p_l) G_A$$

$$+ 2(\delta_{jkl} p_i p_j + \delta_{ikl} p_i p_k - p_i p_k p_{jl} - p_i p_l p_{jk} - p_j p_k p_{il} - p_j p_l p_{ik} - 2 p_i p_j p_{kl} + 4 p_i p_j p_k p_l) G_B$$

$$+ \begin{pmatrix} p_{ijkl} - p_i p_{jkl} - p_j p_{ikl} - p_k p_{ijl} - p_l p_{ijk} + 2 p_i p_j p_{kl} + 2 p_i p_k p_{jl} + 2 p_i p_l p_{jk} \\ + 2 p_j p_k p_{il} + 2 p_j p_l p_{ik} + 2 p_k p_l p_{ij} - p_{ij} p_{kl} - p_{ik} p_{jl} - p_{il} p_{jk} - 6 p_i p_j p_k p_l) \end{pmatrix} H$$

232   (7)

where, for shorthand, $p_{ij} = \delta_{ij} p_i$.

In this expression, there are eight relationship coefficients (R_{AB}, R_{ABAB}, F_A, F_B, F_{AB}, G_A,

G_B, H), which in principle will specify eight different classes of marker genotypes. This

probability of four alleles, f_{ijkl}, is then fitted to the observed frequencies in a sample.

For equations that solve for all eight parameters, the choice is somewhat arbitrary

but a natural set of eight classes, in which identity-by-state mirrors the identity-by-descent,

is: $A_iA_iA_iA_i$ (all identical by state, or "ibs"), $A_iA_kA_iA_k$, $A_iA_iA_kA_k$, (two pairs ibs), $A_iA_iA_iA_k$,

$A_iA_kA_kA_k$, (one triplet ibs) and $A_iA_iA_jA_k$, $A_iA_jA_kA_k$ and $A_iA_jA_jA_k$. (one pair ibs between A and

B). Thus we seek the expected frequencies in the vector $(f_{iiii}, f_{ijij}, f_{iijj}, f_{iiij}, f_{iijk}, f_{ijik})$.

The frequency of $A_iA_iA_iA_i$, is obtained from Eq (7) where all $\delta=1$ and all marker

frequencies are p_i:

$$f_{iiii} = p_i\left[p_i^3 + p_i^2q_i(F_A + F_B + 4R) + p_iq_i^2(F_{AB} + 2R_{AB}) + 2p_iq_i(1 - 2p_i)(G_A + G_B)\right.$$
$$\left. + q_i(1 - 6p_iq_i)H\right]$$

244   Likewise, the frequency that A and B are both heterozygous for A_i and A_j is, irrespective of

245   order or phase,

$$f_{ijij} = 4p_ip_j\left[p_ip_j(1 - F_A - F_B - F_{AB}) + (p_i + p_j - 4p_ip_j)R + (1 - p_i - p_j + 2p_ip_j)R_{AB}\right.$$
$$\left. + (-p_i - p_j + 4p_ip_j)(G_A + G_B) - (1 - 2p_i - 2p_j + 6p_ip_j)H\right]$$

246   and homozygous for alternative alleles A_i and A_j is

$$f_{iijj} = 2p_ip_j\left[p_ip_j(1 - 4R + 2R_{AB}) + q_ip_jF_A + p_iq_jF_B + (1 - p_i - p_j+p_ip_j)F_{AB}\right.$$
$$\left. - 2p_j(1 - 2p_i)G_A - 2p_i(1 - 2p_j)G_B - (1 - 2p_i - 2p_j + 6p_ip_j)H\right]$$

247   for triplets of identity-by-state f_{iiij}

$$f_{iiij} = 2p_ip_j\left[p_i^2 + p_iq_iF_A - p_i^2F_B - p_iq_iF_{AB} + (p_i + p_j - 4p_ip_j)R - (p_i + p_j - 2p_ip_j)R_{AB}\right.$$
$$\left. + (1 - 4p_i + 4p_ip_j)G_A - 2p_i(1 - 2p_j)G_B - (1 - 6p_iq_i)H\right]$$

248   Finally, a single allele pair *ii* can be shared only within individual *A*,

$$f_{iijk} = 2p_i p_j p_k [p_i(1 - 4R + 2R_{AB} - F_B + F_{AB}) + q_i F_A - 2(1 - 2p_i)G_A + 4p_i G_B + 2(1 - 3p_i)H$$

249   or shared only once between *A* and *B*:

$$f_{ijik} = 4p_i p_j p_k [p_i(1 - F_A - F_B + F_{AB}) + (1 - 4p_i)R - (1 - 2p_i)R_{AB} - (1 - 4p_i)(G_A + G_B)$$
$$+ 2(1 - 3p_i)H$$

The expressions for f_{ijjj} and f_{ijkk} are obtained by symmetry, and the expression for f_{ijjk} is

summed over all four pairings of *j* between *A* and *B*: $f_{ijjk}, f_{ijkj}, f_{jijk}$, and f_{jikj}.

The appendix gives the 8x8 matrix of probabilities of observing the marker gene

frequencies *f* given the relatedness coefficients. Of course, this depends upon the particular

array of *f*'s used (there are others than the above). In this case, the determinant of the

matrix is $512 p_i^8 \, p_j^7 p_k^3 (p_i - 1)\left(96 p_i^7 - 160 p_i^6 - 4p_i^5(24p_j^2 - 16p_j - 9) + 3p_i^4(32p_j^2 - \right.$

$80p_j + 47) - p_i^3(132p_j^2 - 302p_j + 147) + p_i^2(89p_j^2 - 154p_j + 59) + p(1 - p_j)(23p_j -$

$\left.11) + (p_j - 1)(2p_j - 1)\right)$. That it is non-zero indicates all 8 parameters are jointly

estimable, but a linear approach which uses residuals to simplify things is needed at this

point.

Joint values of H and identity disequilibrium

$$\begin{bmatrix} f'_{AAAA} \\ f'_{AAaa} \end{bmatrix} = \begin{bmatrix} p(1 - p)(1 - 6p(1 - p)) & p^2(1 - p)^2 \\ -2pq(1 - 2p - 2q + 6pq) & 2p^2 q^2 \end{bmatrix} \begin{bmatrix} H \\ F_{ab} \end{bmatrix}$$

262   whose solution is

$$\widehat{H} = \frac{q^2 f'_{AAAA} - (1 - p)^2 f'_{AAaa}}{pq(1 - p)(1 - 2p)(1 - p - q)}$$

$$\widehat{F_{ab}} = \frac{q(1 - 2p - 2q + 6pq)f'_{AAAA} - (1 - p)(1 - 6p(1 - p))f'_{AAaa}}{pq(1 - p)(1 - 2p)(1 - p - q)}$$

263   Joint estimates of H and joint identity disequilibrium

264   $$\begin{bmatrix} f_{AAAA} \\ f_{AaAa} \\ f_{AAaa} \end{bmatrix} = \begin{bmatrix} p(1 - p)(1 - 6p(1 - p)) & p^2(1 - p)^2 & 2p^2(1 - p)^2 \\ -4pq(1 - 2p - 2q + 6pq) & 4p^2q^2 & 4pq((1 - p)(1 - q) + pq) \\ -2pq(1 - 2p - 2q + 6pq) & 2pq(1 - p - q + pq) & 4p^2q^2 \end{bmatrix} \begin{bmatrix} H \\ F \\ R \end{bmatrix}$$

$$\widehat{H} = \frac{2q(1 - p - q + 3pq)f'_{AAAA} - p(1 - p)^2(f'_{AAaa} + f'_{AaAa})}{2pq(1 - p)(1 - p - q)(1 - 3p)}$$

$$\widehat{F_{ab}} = \frac{2q(1 - 2p - 2q + 6pq)f'_{AAAA} + (1 - p)(2p^2 - 4p + 1)f'_{AAaa} + p(1 - 2p)f'_{AaAa}}{2pq(1 - p)(1 - p - q)(1 - 3p)}$$

$$\widehat{R_{abab}} = \frac{4q(1 - 2p - 2q + 6pq)f'_{AAAA} + (1 - p)p(1 - 2p)f'_{AAaa} - (4p^2 - 5p + 1)f'_{AaAa}}{2pq(1 - p)(1 - p - q)(1 - 3p)}$$

The denominator shows that at least three alleles required in the population, and marker

frequencies of p_i=1/3 are non-informative.

Identity disequilibrium has classically been characterized as the excess of

homozygosity above that expected from the squared gene frequencies, as proposed by HILL

(1975) and OHTA (1980). The identity excess is closely correlated to the expectation of the

total squared linkage disequilibrium (TAKAHATA 1982). Some of the problem is that

haploid gametes are not directly assayed but rather imputed (VITALIS AND COUVET 2001b).

The "classic" procedure for estimating identity disequilibrium involves comparing

observed vs. expected double homozygotes. From VITALIS AND COUVET (2001a), one such

estimator can be written in the form,

$$\widehat{R_{AB}} = \frac{af_{AAAA} - bf_{AA,aa}}{c}$$

where $a = 6(6pq - 2p - 2q + 1), b = (1 - 6pq), \ c = 6pq(1 - p - q)(1 - 3p)$. The

numerator is positive as a is always larger than b, but denominator can be either positive

or negative and negative values occurs when b is larger an a. Such is life with four genes. It

is also amazing the four-gene frequencies f_{AAAA} and $f_{AA,aa}$ are directly used in an effective

two gene estimator, as f_{AAAA} is the observed double homozygotes and $f_{AA,aa}$ represents

homozygotes expected with no zygotic association. Equally amazing is that VITALIS AND

COUVET (2001a) and others have given estimators for identity disequilibrium which omits

this cumulant.

285   DISCUSSION AND CONCLUSION

A main feature of higher-order relatedness is the covariance of homozygosity

between pairs of marker genes, this is effectively a covariance of second moments. Such a

"covariance of covariance" arises when pedigrees occur in a mixture distribution

(MCLACHLAN *et al.* 2019). Such a distribution generates the genomic variation of

homozygosity necessary for the existence of covariance of homozygosity between

individuals at specific loci. The simplest mixture distribution is that of two populations

with gene frequency $p+a$ and $p-a$; in this case covariance of heterozygosity, after mixing in

equal proportions, equals $a^4 + 6a^2 p^2$.

Another feature of higher-order relatedness is that the four-marker coefficient of

gene identity must be described with cumulants and not moments. As an example of the

necessity of cumulants, for population gene frequency p, the fourth central moment for four

markers, denoted (X_1, X_2, X_3 and X_4), is $\sigma_{12}^2 \sigma_{34}^2 + \sigma_{13}^2 \sigma_{24}^2 + \sigma_{23}^2 \sigma_{13}^2 + \kappa_4$ where σ_{ij}^2 is the

covariance of X_i and X_j and κ_4 is a fourth-order cumulant which does not appear in the

moment. Some type of term (not involving the product of variances) is needed for κ_4 and it

could be any rational number. In summary, incorporating cumulants into four marker

measures only requires some value X in the expansion of the fourth central moment

$\sigma_{12}^2\sigma_{34}^2 + \sigma_{13}^2\sigma_{24}^2 + \sigma_{23}^2\sigma_{13}^2 + X$ and this X is numerically estimated in the same way as the

lower order cumulant terms.

Cumulants do have useful properties for models of quantitative traits, the most

important is that the cumulant of the sum of two random variables X+Y is

M(X+Y)=M(X)+M(Y); differential equations for models of selection on quantitative traits

that involve cumulants are simpler than models involving moments (BURGER 1991; TURELLI

AND BARTON 1994). This cumulant will also be key in deriving a marker-based estimator for

Q_{st} (RITLAND IN PREP) and for a portrayal of higher order population structure that separately

accounts for both the correlation of relationship and the squared linkage disequilibrium

(RITLAND IN PREP).

We give probabilities of relationship for a homogenous population of just one

generation. Such populations are most commonly assayed in genomics, however it should

be noted that the levels of nucleotide variation (for SNPs) is not high and loci with more

than two alleles are uncommon; in fact, only about 5% of human SNPs are triallelic (CAO *et*

al. 2015) although microsatellites and other types of repeat markers show greater

variation. Reconstruction of pedigree relationship has traditionally involved cumbersome

graph-tracing algorithms, and simpler recursive methods which require at least two

generations of records (KARIGL 1981; THOMPSON 1988; WHITTEMORE AND HALPERN 1994).

Also, current recursive methods (ZHENG *et al.* 2018) assume a known pedigree (KIRKPATRICK

et al. 2018). This is somewhat like estimating Q_{st} with current methods, where aspects of

the pedigree must be known.

Normalization constants

Relatedness coefficients are obtained by calculating the pairwise covariance of

relatives and dividing it by a normalization constant that converts the covariance into a

correlation. This constant is the maximum possible value that the covariance can take. For

cases where pairwise comparisons involve the frequency of identical genotypes, it is simple

to calculate as a binomial variance. For the two-marker relationship coefficient as

described in Equation (1), the maximum covariance between two genotypes, conditioned

upon observing allele *i*, is $\mathrm{E}[A_iA_i] - \mathrm{E}[A_i]^2 = p_i(1\text{-}p_i)$ when R=1. Likewise, the three marker

coefficient has a normalization constant of $p_i(1\text{-}p_i)(1\text{-}2p_i)$ for $A_iA_iA_i$., and the four-marker

coefficients are normalized by $p_i(1\text{-}p_i)(1\text{-}6p_i(1\text{-}p_i))$ for $A_iA_iA_iA_i$. The normalization

constants for combinations of alleles falls out of the analyses.

ACKERMAN *et al.* (2017) provided a different set of normalization constants for three

and four marker measures than given here. In their Equation 6, they normalized the third

central moment by the geometric mean gene frequency of the three central moments

(rather than by $p(1\text{-}p)(1\text{-}2p)$ as done here). Their justification was a similarity of this

"third moment correlation" to a bivariate correlation formula. Their normalization

constant for the four-marker coefficient (Equation 8) involves a parameter α that mixes the

unknown proportions of the two types of higher order identity (all identical vs. 2 pairs

identical), resulting in an inference that may be subject to biases.

Estimation

Table 2 gives example estimates of R, G and H at a single locus. As expected, the

variances decrease with numbers of alleles. While the variances for R are reasonable, those

for H and the variances for the identity disequilibrium R_{AB} are quite large. Interestingly,

and not remarkably, the variances for G are all over the place and its is not even estimate in

one case (p=0.5).

Likelihood requires numerical solutions which introduces complications, as the

numerical solution is normally iterated until convergence. We used the Expectation-

Maximization method which has slow convergence. When using likelihood, I found the

number of marker loci needed for adequate convergence was about 20 loci for three-gene

coefficients and roughly double at 30-50 loci for four gene coefficients. Interestingly, it was

found that loci with fewer alleles are more likely to give convergent estimates because the

problem with non-convergence arises when relatives do not share the same marker allele.

Calculating the probabilities of higher-order relationship poses an interesting set of

obstacles. Equation solvers such as Derive or Mathematica help with the derivation and

interpretation of complicated formulae. The coefficients can also undefined at certain

intermediate frequencies (p=1/2 or 1/3) and show high statistical variance about those

frequencies (RITLAND 1987).

Possible future approaches

The complications of correctly estimating population structure are discussed and

treated by WEIR AND GOUDET (2017). They developed two-marker moment estimators that

can describe the "relativity" and this requires an explicit reference population. They

develop their estimator in a multilevel approach (within individuals, between individual

within populations, and between populations) which promoted a unified treatment of

relatedness and population structure. Clearly, further progress will depend upon adequate

definitions and applications of models.

"Relatedness mapping" (ALBRECHTSEN *et al.* 2009) uses relatedness to identifying

causative mutations, using the principle that affected individuals share higher relatedness

about the mutation. A somewhat related activity is "IBD mapping", in which segments of

identity-by-descent (IBD) present in high-density genomic data are used to map casual

variants (BROWNING AND BROWNING 2012). However, the data by itself only reveals the

presence of the variant.

Other fields have adopted the use of cumulants, which may show new approaches

that population genetics can undertake. The central 4th cumulant has been used to detect

early stages of termite infestation, as it can separate termite alarm signals from

background noise (DE LA ROSA AND MORENO MUÑOZ 2008). Advances in electrophysiological

and imaging techniques are used to study the synchrony of neuron cell firings in the brain

(STAUDE AND ROTTER 2010), and have highlighted the need for correlation measures that go

beyond simple pairwise analyses, taking advantage of the "interaction property" of higher

order cumulants as measures of correlation (STAUDE *et al.* 2010). In information systems, a

"covariance of covariance" approach for individual pixels has been developed for image

description and classification (SERRA *et al.* 2014).

ACKNOWLEDGMENTS Joe Felsenstein provided many helpful comments on my first and

only sabbatical. This work was supported by NSERC grants to KR

Table 1. Examples of the statistical variances of relationship coefficients when estimates

are based upon a single locus and when the true values are zero ("x" denotes not

estimable). See equations 3 and 6 for definitions of G and H.

──

Array of p	F	G_A	G_B	ϕ_{XY}	H
0.6,0.3,0.1	2.93	73.5	74.3	12.96	7.63
0.5,0.3,0.2	2.32	x	x	11.66	3.66
0.4,0.3,0.2,0.1	1.58	0.19	7.79	13.05	2.18

399   ──

Table 2. Allele states and denominators of cumulants.

____

____

Allele states	Expectation	Denomi
Two alleles		
$i = j$	$p_i^2 + p_i(1 - p_i)F$	$p_i(1 -$
$i \neq j$	$2p_i p_j(1 - F)$	$-2p_i$
Three alleles		
$i = j = k$	$p_i^3 + p_i^2(1 - p_i)(F + 2R) + p_i G$	$p_i(1 - p_i)($
$i = j \neq k$	$p_i^2 p_k(2R - 2G)$	$-p_i p_k(1$
$i \neq j \neq k$	$p_i p_j p_k(1 - F - 2R + 2G)$	$2p_i p_j$
Four alleles		
$i = j = k = l$	$p_i^2 q_i^2(F_{AB} + 2R_{AB}) + q_i(1 - 6p_i q_i)H$	$p_i(1 - p_i)(1 - 6p_i$
$i = j = k \neq l$	$2p_i p_j(F_{AB} + (p_i + p_j - 2p_i p_j)R_{AB} - (1 - 6p_i q_i)H)$	$p_i p_k(1 - 2p_i - 2p_k$
$i = j \neq k = l$	$2p_i p_j(2p_i p_j R_{AB} + (1 - p_i - p_j + p_i p_j)F_{AB} - (1 - 2p_i - 2p_j + 6p_i p_j)H)$	$p_i p_k(1 - 2p_i - 2p_k$
$i = j \neq k \neq l$	$2p_i p_j p_k(\lceil p_i(2R_{AB} + F_{AB}) + 2(1 - 3p_i)H)$	$2p_i p_j p_k(1 -$
$i \neq j \neq k \neq l$	$p_i p_j p_k p_l(1 - 4R - 2F + 4G - 8H)$	$-6p_i p_j p_k l$

FIGURE LEGENDS

Figure 1. Three cases where three-gene modes of gene identity are used. (A) Effective selfing model involves two genes of one parent, (B) progeny pair model, (C) Altruist-recipient. Identical genes are linked by lines. In the effective selfing model, two of the genes are from the maternal parent and the other gene is the paternal contribution.

Figure 2. Two cases where four-gene modes of identity are used. (A) The general case where each of the nine identity modes are inferred (B) A model to fit progeny pairs to mating system parameters (C) The inference of heritability in the field, where M is the marker and Q the quantitative trait.

Figure 1

Figure 2.

Literature cited

Ackerman, M. S., P. Johri, K. Spitze, S. Xu, T. G. Doak *et al.*, 2017 Estimating Seven Coefficients of Pairwise Relatedness Using Population-Genomic Data. Genetics 206**:** 105-118.

Albrechtsen, A., T. Sand Korneliussen, I. Moltke, T. van Overseem Hansen, F. C. Nielsen *et al.*, 2009 Relatedness mapping and tracts of relatedness for genome☐wide data in the presence of linkage disequilibrium. Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society 33**:** 266-274.

Browning, S. R., and B. L. Browning, 2012 Identity by Descent Between Distant Relatives: Detection and Applications. Annual Review of Genetics 46**:** 617-633.

Burger, R., 1991 Moments, cumulants, and polygenic dynamics. Journal of mathematical biology 30**:** 199-213.

Cao, M., J. Shi, J. Wang, J. Hong, B. Cui *et al.*, 2015 Analysis of human triallelic SNPs by next-generation sequencing. Ann Hum Genet 79**:** 275-281.

de la Rosa, J. J. G., and A. Moreno Muñoz, 2008 Higher-order cumulants and spectral kurtosis for early detection of subterranean termites. Mechanical Systems and Signal Processing 22**:** 279-294.

Gardner, A., S. A. West and G. Wild, 2011 The genetical theory of kin selection. Journal of Evolutionary Biology 24**:** 1020-1043.

Hill, W. G., 1975 Linkage disequilibrium among multiple neutral alleles produced by mutation in finite population. Theor Popul Biol 8**:** 117-126.

Jacquard, A., 1966 Logique du calcul des coefficients d'identité entre deux individus. Population (french edition)**:** 751-776.

Jacquard, A., 2012 *The genetic structure of populations*. Springer Science & Business Media.

Karigl, G., 1981 A recursive algorithm for the calculation of identity coefficients. Annals of Human Genetics 45**:** 299-305.

Kendall, M. G., A. Stuart and J. K. Ord, 1977 *The advanced theory of statistics.* , London, Griffin.

Kirkpatrick, B., S. Ge and L. Wang, 2018 Efficient computation of the kinship coefficients. Bioinformatics 35**:** 1002-1008.

Liu, W. W. B., 2005 Genotypic probabilities for pairs of inbred individuals. Philosophical Transactions of the Royal Society B 360**:** 1379-1385.

Malécot, G., 1948 *Mathématiques de l'hérédité*. Masson, Paris.

McLachlan, G. J., S. X. Lee and S. I. Rathnayake, 2019 Finite mixture models. Annual review of statistics and its application 6**:** 355-378.

Michod, R. E., and W. D. Hamilton, 1980 Coefficients of relatedness in sociobiology. Nature 288**:** 694-697.

Morton, N. E., S. Yee, D. E. Harris and R. Lew, 1971 Bioassay of kinship. Theoretical Population Biology 2**:** 507-524.

Ohta, T., 1980 Linkage disequilibrium between amino acid sites in immunoglobulin genes and other multigene families. Genetical research 36**:** 181-197.

Pamilo, P., and R. H. Crozier, 1982 Measuring genetic relatedness in natural populations: Methodology. Theoretical Population Biology 21**:** 171-193.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira *et al.*, 2007 PLINK: a tool set for whole-genome association and population-based linkage analyses. The American journal of human genetics 81**:** 559-575.

Queller, D. C., 1992 Quantitative Genetics, Inclusive Fitness, and Group Selection. The American Naturalist 139**:** 540-558.

Queller, D. C., and K. F. Goodnight, 1989 Estimating relatedness using genetic markers. Evolution 43**:** 258-275.

Ritland, K., 1984 The effective proportion of self-fertilization with consanguineous matings in inbred populations. Genetics 106**:** 139-152.

Ritland, K., 1985 The genetic mating structure of subdivided populations I. Open-mating model. Theoretical Population Biology 27**:** 51-74.

Ritland, K., 1987 Definition and Estimation of Higher-Order Gene Fixation Indices. Genetics 117**:** 783-793.

Ritland, K., 1996 Estimators for pairwise relatedness and individual inbreeding coefficients. Genetics Research 67**:** 175-185.

Ritland, K., 2000 Marker-inferred relatedness as a tool for detecting heritability in nature. Molecular Ecology 9**:** 1195-1204.

Ritland, K., and M. Leblanc, 2004 Mating system of four inbreeding monkeyflower (*Mimulus*) species revealed using 'progeny☐pair' analysis of highly informative microsatellite markers. Plant Species Biology 19**:** 149-157.

Samanta, S., Y. J. Li and B. S. Weir, 2009 Drawing inferences about the coancestry coefficient. Theoretical Population Biology 75**:** 312-319.

Serra, G., C. Grana, M. Manfredi and R. Cucchiara, 2014 Covariance of covariance features for image classification, pp. 411-414 in *Proceedings of International Conference on Multimedia Retrieval*.

Staude, B., S. Grün and S. Rotter, 2010 Higher-order correlations and cumulants, pp. 253-280 in *Analysis of parallel spike trains*. Springer.

Staude, B., and S. Rotter, 2010 Higher-order correlations in non-stationary parallel spike trains: statistical modeling and inference. Frontiers in computational neuroscience 4**:** 16.

Takahata, N., 1982 Linkage disequilibrium, genetic distance and evolutionary distance under a general model of linked genes or a part of the genome. Genetical Research 39**:** 63-77.

Thompson, E. A., 1988 Two-locus and Three-locus Gene Identity by Descent in Pedigrees. Mathematical Medicine and Biology: A Journal of the IMA 5**:** 261-279.

Turelli, M., and N. H. Barton, 1994 Genetic and statistical analyses of strong selection on polygenic traits: what, me normal? Genetics 138**:** 913-941.

Vitalis, R., and D. Couvet, 2001a Estimation of Effective Population Size and Migration Rate From One- and Two-Locus Identity Measures. Genetics 157**:** 911-925.

Vitalis, R., and D. Couvet, 2001b Two-locus identity probabilities and identity disequilibrium in a partially selfing subdivided population. Genetics Research 77**:** 67-81.

Wang, J., 2014 Marker-based estimates of relatedness and inbreeding coefficients: an assessment of current methods. Journal of Evolutionary Biology 27**:** 518-530.

Weir, B. S., 1994 The effects of inbreeding on forensic calculations. Annu Rev Genet 28**:** 597-621.

Weir, B. S., and J. Goudet, 2017 A unified characterization of population structure and relatedness. Genetics 206**:** 2085-2103.

Whittemore, A. S., and J. Halpern, 1994 Probability of Gene Identity by Descent: Computation and Applications. Biometrics 50**:** 109-117.

Withers, C. S., S. Nadarajah and S. H. Shih, 2015 Moments and Cumulants of a Mixture. Methodology and Computing in Applied Probability 17**:** 541-564.

Zheng, C., M. P. Boer and F. A. van Eeuwijk, 2018 Recursive algorithms for modeling genomic ancestral origins in a fixed pedigree. G3: Genes, Genomes, Genetics 8**:** 3231-3245.