Dissecting the functional contributions of different neuronal subtypes in the dorsal striatum to perseverative behaviour in ephrin-A2/A5[-/-] mice

Maitri Tomar [1,2], Jennifer Rodger [1,2*], Jessica Moretti [1,2]

[1] School of Biological Sciences, University of Western Australia, Crawley WA, Australia

[2] Perron Institute for Neurological and Translational Science, Nedlands WA, Australia

*Corresponding author:

E-mail: jennifer.rodger@uwa.edu.au (JR)

Keywords: Habit formation; Dorsal striatum; Nucleus accumbens; rTMS; Ephrin; c-Fos

Abstract

Overreliance on habit is linked with disorders such as drug addiction and obsessive-

compulsive disorder and there is increasing interest in the use of repetitive transcranial

magnetic stimulation (rTMS) to alter neuronal activity in the relevant pathways and reduce

relapse or accelerated shift towards habit formation. Here we studied the brains of ephrin-

A2A5$^{-/-}$ mice, which previously showed perseverative behaviour in progressive-ratio tasks,

associated with low cellular activity in nucleus accumbens. We investigated if rTMS

treatment had altered the hierarchical recruitment of brain regions from ventral to dorsal

striatum associated with abnormal habit formation in these mice.

Brain sections of mice that underwent progressive-ratio tasks with and without low

intensity rTMS (LI-rTMS) were taken from a previous study. We take advantage of the

previous characterisation of perseverative behaviour to investigate the contribution of

different neuronal subtypes and striatal regions. Striatal regions were stained for neuronal

activation with c-Fos and for medium spiny neurons with DARPP32. Qualitative analysis

was carried out for other neuronal subtypes in the striatum - GABAergic, parvalbumin-

expressing and cholinergic interneurons.

Contrary to our hypothesis, we found neuronal activity in ephrin-A2A5$^{-/-}$ mice still

reflected goal-directed behaviour. However, we saw that the dorsolateral striatum contributed

more to total striatal activity in untreated ephrin-A2/A5$^{-/-}$ mice. This supported our hypothesis

that ephrin-A2/A5$^{-/-}$ mice have greater c-Fos activity in habit-associated striatal regions. LI-

rTMS in ephrin-A2A5$^{-/-}$ mice also appeared to delay the shift from goal-directed to habitual

behaviour as suggested by increased activation in dorsomedial striatum and nucleus

accumbens.

1   Introduction

Habits form over a period of time and require a shift from goal-directed behaviour to

habitual behaviour. However, sometimes a faulty shift can contribute to disorders such as

drug addiction (Everitt and Robbins, 2005) and obsessive-compulsive disorder (OCD) (Gillan

et al., 2011). The formation of habit is generally defined as a shift from response-outcome

associations, where actions are primarily mediated by the value associated with the outcome,

to stimulus-response associations where, over a period of extended training and repetition,

there is a pattern of outcome devaluation and the behaviour becomes more automatic,

independent from the outcome of the action (Dickinson et al., 1995). This behavioural shift

from goal-directed to habitual behaviour involves dopaminergic pathways and a hierarchical

recruitment of brain regions from ventral to dorsal striatum (Segovia et al., 2012; Liu et al.,

2013).

A potential tool to manipulate this hierarchical recruitment is repetitive transcranial

magnetic stimulation (rTMS), a non-invasive brain stimulation technique that is able to

modulate neuronal activity (Aydin-Abidin et al., 2008) and various neurotransmitters,

including dopamine (Keck et al., 2002; Moretti et al., 2020). rTMS is approved for treatment

of depression (O'Reardon et al., 2007) and OCD (Carmi et al., 2018). Understanding more

about the mechanisms of rTMS, could be useful in refining rTMS use.

To explore the potential of rTMS to repair abnormal circuitry, a transgenic ephrin-A2/A5[-]

[/-] mouse strain has previously been used to measure structural and functional plasticity

induced in abnormal neural pathways. Ephrins are membrane-bound ligands of Eph tyrosine

kinase receptors and are important in cell migration and axon guidance during development

(Wilkinson, 2001). As a result, ephrin- A2/A5[-/-] mice, which lack the ephrin-A2 and ephrin-

A5 ligands, have disorganized axonal projections in neuronal circuits throughout the brain

due to disrupted Eph/ephrin signalling. These mice have abnormal visual topography and

visuomotor behaviour which are both partially rescued by low-intensity rTMS (LI-rTMS)

(Rodger et al., 2012; Makowiecki et al., 2014; Poh et al., 2018). In addition, the mice have

reduced dopaminergic innervation of the striatum (Sieber et al., 2004; Cooper et al., 2009)

associated with abnormal behavioural patterns in goal-directed behaviour (Poh et al., 2018)

but the effects of LI-rTMS on this phenotype are less well understood.

A previous study (Moretti et al., 2021) explored whether the different goal-directed

behaviour of ephrin-A2/A5$^{-/-}$ mice compared to wild type was due to abnormal motivation by

comparing performance of both strains in a progressive ratio (PR) tasks. PR tasks are a good

measure of motivation as they involve an increasing instrumental response requirement for a

reward. Highly motivated animals will continue to respond consistently, while animals with

low motivation stop or slow their responding (Hodos, 1961; Aberman et al., 1998). The

potential ameliorating effect of chronic excitatory LI-rTMS delivered during the PR task was

also examined. Unexpectedly, the results did not support a motivation phenotype: ephrin

A2/A5$^{-/-}$ mice showed perseverative behaviour in the PR task, with a greater number of

responses compared to wildtype mice despite increasing task difficulty (Moretti et al., 2021).

Compared to wildtype mice, ephrin-A2A5$^{-/-}$ mice also showed reduced c-Fos expression in

the nucleus accumbens (NAc) after two weeks of the PR task (Moretti et al., 2021), which

may reflect an accelerated dorsal shift in neuronal activity characteristic of habitual behaviour

(Segovia et al., 2012). Interestingly, chronic excitatory LI-rTMS delivered to ephrin-A2/A5$^{-/-}$

mice during the PR task did not significantly alter behaviour, but did ameliorate the reduced

c-Fos expression in the ventral striatum of ephrin-A2/A5$^{-/-}$ mice, resulting in expression that

was similar to wildtype levels (Moretti et al., 2021). Therefore, in ephrin-A2/A5$^{-/-}$ mice, LI-

rTMS may mitigate abnormal activity along the mesolimbic pathway associated with

behavioural inflexibility, by delaying an accelerated shift in neuronal activity toward the

dorsal striatum.

To understand if altered dorsal striatum activity in ephrin-A2/A5$^{-/-}$ mice was

responsible for perseverative behaviour in these mice and if rTMS could delay an accelerated

shift from goal-directed behaviour to habit, here we used tissue from the previous study

(Moretti et al., 2021) to identify and compare activation of neuronal populations across the

entire striatum, investigating the dorsomedial striatum (DMS), dorsolateral striatum (DLS)

and NAc (core and shell) using c-Fos immunohistochemistry. We compared c-Fos expression

in different regions of the striatum in wildtype and ephrin-A2A5$^{-/-}$ mice that either received

active LI-rTMS or sham stimulation. In addition, since a low percentage of c-Fos expression

colocalised with medium spiny neurons (MSNs) was found in the NAc (Moretti et al., 2021),

we also identified the other neuronal subtypes in the striatum that expressed c-Fos including

GABAergic and cholinergic interneurons. Our in depth characterisation of cell activation

following LI-rTMS in the striatum in behaviourally characterised animals may provide

insights into the use of TMS for disorders such as addiction and OCD where the shift of habit

formation is faulty.

2   Materials and methods

2.1   Animal tissue

All experiments were approved by The University of Western Australia Animal Ethics

Committee (AEC 100/1639). Sagittal mouse brain sections (40 µm) from 11 adult (8-24-

week-old) wildtype C57BL6/J mice (sham: 6; rTMS: 5) and 10 ephrin-A2/A5$^{-/-}$ mice (sham:

6; rTMS: 4) (Moretti et al., 2021) were used. The ephrin- A2/A5$^{-/-}$ mice was backcrossed onto

C57BL6/J mice for >20 generations, bread from heterozygous ephrin-A2$^{-/-}$A5$^{+/-}$ parents and

genotyped at weaning (Feldheim et al., 2000). In the previous study, following habituation to

the LI-rTMS coil and training in the operant box, mice performed one week of an exponential

PR task (required responses increase under an exponential equation) followed by one week of

PR7 (requirement started at 7 responses and increased by 7 with each reinforcement) (Moretti

et al., 2021). Mice received 14 days of biomimetic high frequency stimulation (BHFS) LI-

rTMS or sham during the first 10 minutes of the daily PR task and were euthanized 90

minutes after the beginning of the final PR task to correspond with the peak of c-Fos

expression (Moretti et al., 2021). A summary of the experimental design from Moretti et al.

(2021) is reproduced in Figure 1. The previous study showed perseverative behaviour in

untreated ephrin- A2/A5$^{-/-}$ mice and low c-Fos expression in the NAc which improved

following LI-rTMS (Moretti et al., 2021). This suggested an accelerated shift from goal-

directed to habit in untreated ephrin- A2/A5$^{-/-}$ mice which could have been delayed by LI-

rTMS (Moretti et al., 2021). However, in the previous study neuronal activation in dorsal

striatum was not investigated. In the current study we use additional tissue from the same

animals as Moretti et al. (2021) to confirm the shift in neuronal activity from regions

involved in goal-directed behaviour to habit and if LI-rTMS delayed it on a cellular level.

Figure 1. Outline of the experimental paradigm performed by Moretti et al. (2021).

Reproduced from (Moretti et al., 2021) with permission from Behavioural Brain Research.

2.2 Immunofluorescence

Every fifth sagittal brain section which contained dorsal striatum was identified using the

mouse brain atlas (Paxinos and Franklin, 2012). These sections were stained for three

different antibodies: c-Fos (rabbit polyclonal c-Fos antibody, 1:5000, Abcam, ab190289), a

marker for neuronal activity (Bullitt, 1990), cAMP-regulated phosphoprotein-32 kD

(DARPP32) (purified mouse anti-DARPP32, 1:2000, BD Transduction Laboratories,

611520), a marker for MSNs (Anderson and Reiner, 1991), and glutamate decarboxylase 67

(GAD67) (anti-Goat, 1:750, R&D systems, AF2086-SP), a marker for GABAergic neurons

(Lazarus et al., 2015). For confirmatory analysis of active parvalbumin (PV)-expressing

GABAergic interneurons and cholinergic interneurons, two sections from one animal

belonging to each experimental group were stained for c-Fos, PV (mouse monoclonal anti-

PV antibody, 1:5000, Millipore Sigma, p3088) and ChAT (anti-choline acetyltransferase

antibody, 1:200, Millipore Sigma, AB144P). Hoescht (1:2000, Invitrogen) was used to stain

all nuclei.

Free-floating sections were washed 3x in PBS (5 min each), permeabilised by washing in

0.1% Triton-X in PBS (PBS-T) for 15 min and incubated for 2 h in a blocking buffer of 2%

bovine serum albumin (Sigma) and 3% normal donkey serum (Sigma) diluted in 0.1% PBS-

T. Then, sections were incubated with primary antibodies in blocking buffer overnight at

4°C. Sections were washed in PBS-T 3x (10 min each) and incubated for 2 h with secondary

antibodies diluted in blocking buffer to 1:600 (donkey anti-rabbit IgG Alexa Fluor 488,

Invitrogen, Thermo Fisher, A21206; donkey anti-mouse IgG Alexa Fluor 555, Invitrogen,

Thermo Fisher, A21202; donkey anti-goat IgG H&L Alexa Fluor 647, ab150131). Sections

were then washed in PBS-T 3x (10 min each) and incubated for 10 min in Hoescht in PBS

(1:1000, Invitrogen). Lastly, the sections were washed in PBS 3x (10 min each) and mounted

on gelatin-subbed slides, coverslipped with mounting medium (Dako, Glostrup, Denmark)

and sealed with nail polish. Slides were stored at 4°C in a light-controlled environment until

imaging.

2.3   Imaging and quantification

Commonly used mouse brain atlases (Lein et al., 2007; Paxinos and Franklin, 2012) do

not show detailed segmentation of the dorsal striatum (Chon et al., 2019). Therefore, in the

present study the two regions of the dorsal striatum (DMS and DLS) were identified by

creating a custom sagittal atlas that combined the location information of dorsal striatum

sections from a mouse brain atlas (Chon et al., 2019) annotated onto a 3D magnetic

resonance imaging (MRI) data of P56 mouse brain atlas using ITK-SNAP (Yushkevich et al.,

2006) software (Fig 2).

Fig 2. Identification of regions of interest on a mouse brain atlas. (A) Location of DMS

and DLS was marked in red and green respectively on a coronal plane in the software, ITK-

SNAP which produced corresponding images in the horizontal (B) and sagittal planes (C).

Scale bar = 5 mm. CP, coronal plane; SP, sagittal plane; HP, horizontal plane.

Sections were imaged through z-stacks (2 µm apart) on a Nikon C2 confocal microscope

(Nikon, Tokyo, Japan). 10x magnification was used to identify the regions of interest. 8-11

fields of view (FOV) for DLS and 6-10 FOV for DMS across 5-7 sections each animal were

used to image at 40x magnification to identify c-Fos, DARPP32 and GAD67 markers. For the

confirmatory analysis, 2 FOV for DLS and DMS and 1 FOV for NAc across 2 sections from

each experimental group was imaged at 40x magnification.

Cell counts were performed using stereological principles using ImageJ software and

were carried out blinded to experimental group after all images were captured. The total

number of c-Fos single labelled cells, c-Fos$^+$/DARPP32$^+$ double labelled cells, c-

Fos$^+$/GAD67$^+$/DARPP32$^-$ double labelled cells, c-Fos$^+$/PV$^+$ double labelled cells and c-

Fos$^+$/ChAT$^+$ double labelled cells were identified and counted for each image (Fig 3). The

combination of DARPP32$^+$ and GAD67$^+$ double labelling was not assessed since DARPP32$^+$

cells are also GAD67$^+$ because they are GABAergic in nature. c-Fos$^+$ cells not colocalized

with DARPP32 were classified as unspecified c-Fos.

Fig 3. Representative images taken at 40x magnification showing different marker

expressions. (A) expression of c-Fos$^+$ cells (green) (B) expression of GAD67$^+$ cells (yellow)

and DARPP32$^+$ cells (red) (C) expression of c-Fos$^+$ (green) /DARPP32$^+$ (red) colocalised

cells (D) expression of c-Fos$^+$ (green) /GAD67$^+$ (yellow) colocalised cells (E) expression of c-

Fos$^+$ (green) /ChAT$^+$ (yellow) colocalised cells (F) expression of c-Fos$^+$ (green) /PV$^+$ (red)

colocalised cells. Scale bar = 50 µm. GAD67, Glutamate decarboxylase 67; cAMP-regulated

phosphoprotein-32 kD, DARPP32.

For confirmatory analysis, c-Fos-labelled cells colocalised with GAD67, PV and

ChAT were counted and analysed to explain unspecified c-Fos counts. During imaging,

sections from one ephrin-A2/A5$^{-/-}$-rTMS animal was excluded due to poor staining (ephrin-

A2/A5$^{-/-}$-rTMS *n = 3*).

2.4   Statistical analysis

For analysis, the total c-Fos cells/mm^3 and percentage of c-Fos cells colocalised with

DARPP32, GAD67, PV and ChAT were calculated for each animal. Statistical analysis was

not done on counts showing c-Fos colocalization with PV and ChAT due to low sample size

but assessed qualitatively.

Assumptions of normality and homogeneity were checked. For DLS counts, the

assumptions of normality were not met for percentage c-Fos-GAD67 and total c-Fos

cells/mm^3 counts (p<0.05). Homogeneity of variance was checked, and it was met for all data

sets (Levene's test, p>0.05). For DMS counts, the assumptions of normality were not met for

percentage c-Fos-DARPP32, percentage c-Fos-GAD67 and total c-Fos cells/mm^3 (p<0.05).

The assumption of homogeneity of variance was not met for percentage c-Fos-DARPP32

data (Levene's test, p<0.05) but corrected with logarithmic transformation.

To compare whether cell counts changed across groups, multiple two-way analysis of

variance (ANOVAs) were used with the independent variables of strain and treatment. Each

ANOVA had the dependent variable of either total c-Fos cells/mm^3, percentage c-Fos-

DARPP32 or percentage unspecified c-Fos cell count data and were run separately for both

DLS and DMS. ANOVA was also performed for c-Fos$^+$/GAD67$^+$/DARPP32$^-$ cell counts for

the DLS region. For the DMS regions, due to small sample size, Ephrin-rTMS group was

excluded from statistical analysis for c-Fos$^+$/GAD67$^+$/DARPP32$^-$ data set and student t-tests

were performed to compare remaining experimental groups.

Additional data for percentage of c-Fos$^+$/GAD67$^+$/DARPP32$^-$ cell counts were also

obtained for NAc core and shell if the regions were present in the stained tissue (NAc core,

Group: wildtype-sham $n=5$, wildtype-rTMS $n=4$, ephrin-A2/A5$^{-/-}$-sham $n=4$, ephrin-A2/A5$^{-/-}$

-rTMS $n=2$; NAc shell, Group: wildtype-sham $n=6$, wildtype-rTMS $n=5$, ephrin-A2/A5$^{-/-}$-

sham $n=4$, ephrin-A2/A5$^{-/-}$-rTMS $n=3$). This was combined with total c-Fos cells/mm^3 and

percentage c-Fos-DARPP32 cell counts for NAc core and shell from the previous study

(Moretti et al., 2021). The percentage of c-Fos-GAD67 data set in NAc core and shell

assumptions of normality and heterogeneity. Therefore, non-parametric Mann-Whitney U

tests were performed to compare the percentage of c-Fos$^+$/GAD67$^+$/DARPP32$^-$ cell counts

between sham and treatment for both wildtype and ephrin-A2A5$^{-/-}$ groups.

3   Results

3.1   Total c-Fos cells/mm^3

In DLS, there was no overall effect of strain ($F_{(1,16)}=1.040$, $p=0.323$, $\eta^2=0.058$) or

treatment ($F_{(1,16)}=0.973$, $p=0.339$, $\eta^2=0.054$) on c-Fos cells/mm^3. The strain*treatment

interaction was also not significant ($F_{(1,16)}=0.0007$, $p=0.979$, $\eta^2=0.00$) (Fig 4). Similarly, in

DMS there was no significant difference in cell counts between strain ($F_{(1,16)}=0.902$,

$p=0.356$, $\eta^2=0.046$) and no overall effect of treatment was observed ($F_{(1,16)}=1.342$, $p=0.264$,

$\eta^2=0.068$). No significant strain*treatment interaction was observed ($F_{(1,16)}=1.378$, $p=0.258$,

$\eta^2=0.070$) (Fig 4).

Fig 4. Graphs showing total c-Fos$^+$ cells/mm^3 counts for each group in (A) DLS and (B)

DMS. No significant difference in total c-Fos$^+$ cells/mm^3 cells was found between groups in

both DLS and DMS (p>0.05). Points represent individual animal counts. Error bars represent

$\pm$ SEM.

3.2   c-Fos colocalization in MSNs

In DLS, there was no overall effect of strain (F(1,16)=0.100, p=0.756, η^2=0.006) or

treatment (F(1,16)=0.001, p=0.976, η^2=0) on percentage of c-Fos-DARPP32 colocalization.

The strain*treatment interaction was also not significant (F(1,16)=0.376, p=0.548, η^2=0.023)

(Fig 5). Similarly, in DMS there was no significant difference in percentage of colocalization

between strains (F(1,16)=0.0461, p=0.833, η^2=0.002) and no overall effect of treatment was

observed (F(1,16)=1.2556, p=0.279, η^2=0.056). However, there was a significant

strain*treatment interaction (F(1,16)=5.1062, p=0.038, η^2=0.228). To understand the

interaction, we followed up with pairwise post-hoc comparisons using Tukey's test. There

was an increase in percentage of c-Fos-DARPP32 colocalization after treatment in ephrin-

A2/A5$^{-/-}$ mice compared to ephrin-sham group (Fig 5), however the effect did not survive

multiple comparison Tukey correction (uncorrected p=0.041, t(16)=-2.225, Tukey corrected

p=0.159) despite the significant omnibus interaction. All other groups were non-significantly

different from each other.

Fig 5. Graphs showing percentage of c-Fos$^+$ cells colocalized with DARPP32$^+$ cells for

each group in (A) DLS and (B) DMS. Following stimulation, a significant increase was

only seen in c-Fos-DARPP32 colocalization in DMS in ephrin-A2/A5$^{-/-}$ mice in the omnibus

test but not in pairwise post-hoc comparisons. Points represent individual animal counts.

Error bars represent ± SEM.

3.3   Percentage of unspecified c-Fos cells

In DLS, there was no overall effect of strain ($F(1,16)=1.9052$, $p=0.186$, $\eta^2=0.105$) or

treatment ($F(1,16)=0.0309$, $p=0.863$, $\eta^2=0.002$) on the percentage of unspecified c-Fos cells.

The strain*treatment interaction was also not significant ($F(1,16)=0.2026$, $p=0.659$,

$\eta^2=0.011$) (Fig 6). Similarly in DMS, there was no overall effect of strain ($F(1,16)=0.482$,

$p=0.497$, $\eta^2=0.026$) or treatment ($F(1,16)=0.270$, $p=0.611$, $\eta^2=0.015$) on the percentage of

unspecified c-Fos cells. The strain*treatment interaction was also not significant

($F(1,16)=1.738$, $p=0.206$, $\eta^2=0.094$) (Fig 6).

Fig 6. Graphs showing percentage of unspecified c-Fos[+] cells for each group in (A) DLS

and (B) DMS. No significant difference in percentage of c-Fos[+] cells that are single-labelled

(c-Fos[+]/GAD67[-]/DARPP32[-]) was found between groups in both DLS and DMS ($p > 0.05$).

Points represent individual animal counts. Error bars represent ± SEM.

3.3.1 Confirmatory analysis for unspecified c-Fos-labelled cells

Across all experimental groups, approximately 60% and 40% of c-Fos labelled cells

in the ventral and dorsal striatum, respectively were unidentified (i.e., not MSNs). Therefore,

the sections were first stained for GABAergic interneurons to identify unspecified c-Fos

labelled cells. Qualitative observation suggests that in DMS there were more active

GABAergic interneurons as compared to DLS (Fig 7) and NAc where the percentage of

colocalization across all groups were less than 1% and 2%, respectively.

Fig 7. Graphs showing percentage of c-Fos$^+$ cells colocalized with GAD67$^+$-DARPP32$^-$

cells for each group in (A) DLS and (B) DMS. No significant difference in percentage of c-

Fos$^+$ cells colocalised with GAD67$^+$-DARPP32$^-$ cells were found between groups in both

DLS and DMS (p>0.05). Points represent individual animal counts. Error bars represent ±

SEM.

For percentage of c-Fos$^+$/GAD67$^+$/DARPP32$^-$ colocalization in DLS, no overall effect

of strain (F(1,14)=0.058, p=0.814, η^2=0.004) or treatment (F(1,14)=0.061, p=0.809,

η^2=0.004) was observed. The strain*treatment interaction was also not significant

(F(1,14)=0.088, p=0.771, η^2=0.006) (Fig 7). In the DMS, there was no significant change in

c-Fos$^+$/GAD67$^+$/DARPP32$^-$ colocalization between Wildtype-Sham and Wildtype-rTMS

mice (t(9)=-0.864, p=0.410), and there was no significant difference between strains,

Wildtype-Sham and Ephrin-Sham (t(8)=-1.32, p=0.222). Qualitatively, c-Fos$^+$ that is

colocalised with GAD67$^+$decreased following LI-rTMS in ephrin-A2A5$^{-/-}$ mice as seen in Fig

7.

In NAc core, there was no significant difference in the percentage of c-

Fos$^+$/GAD67$^+$/DARPP32$^-$ colocalization following treatment in both wildtype (U=8.5,

p=0.771) and ephrin-A2A5$^{-/-}$ mice (U=2, p=0.289). For the same data set, there was no

significant difference between strains, Wildtype-Sham and Ephrin-Sham (U=6, P=0.240).

Similarly, in NAc shell there was no significant difference in the percentage of c-

Fos$^+$/GAD67$^+$/DARPP32$^-$ colocalization following treatment in both wildtype (U=12,

p=0.597) and ephrin-A2A5$^{-/-}$ mice (U=5, p=0.825). There was also no significant strain

difference, Wildtype-Sham and Ephrin-Sham (U=11, p=0.896).

As the percentage of c-Fos colocalised with GAD67-labelled cells were also low in

both dorsal and ventral striatum, additional confirmatory tests were done for PV-expressing

GABAergic interneurons and cholinergic interneurons. Qualitative analysis suggests there

were more active PV-expressing GABAergic interneurons in DMS as compared to DLS and

NAc. In DMS, a slight increase in active PV-expressing GABAergic interneurons was

observed in ephrin-A2A5$^{-/-}$ mice following treatment, whereas a slight decrease was seen in

wildtype mice following treatment. No active cholinergic interneurons were observed in both

DMS and NAc across all groups. In DLS, only about 2% c-Fos$^+$ cells were colocalised with

cholinergic interneurons in the wildtype-sham group, in all other groups no active cholinergic

interneurons were observed.

3.4   Comparison of c-Fos activity across regions and groups

To further explore a potential change in activity distribution across regions in the

overall striatum, we examined whether the proportion of c-Fos activity per regions differed

between groups. We ran a 4*4 contingency χ^2 test with c-Fos counts per region (NAc shell,

NAc core, DMS and DLS) compared across experimental groups. The initial 4*4 contingency

χ^2 test was significant (χ^2 (9, N=666524) =8030, p<0.001, V=0.0634). We then ran follow up

2*4 contingency χ^2 test to understand where the difference was, comparing the proportion of

striatal c-Fos activity per region for each group with the Wildtype-Sham group. Proportions

of striatal c-Fos activity per region differed in comparison with Wildtype-Sham for all

groups, although the test may be overly sensitive due to the large sample size (Wildtype-

rTMS: χ^2 (3, N=436329) =642, p<0.001, V = .0384); Ephrin-Sham: χ^2 (3, N=335837) = 2156,

p<0.001, V = .0801; Ephrin-rTMS: χ^2 (3, N=384288) =2446, p<0.001, V = .0798). Although

there are significant differences, the effect sizes for each comparison are small.

To further understand where the difference in striatal activity per region occurred, we

performed region-by-region comparison for the percentage total striatal c-Fos activity present

for each region across groups (Fig 8). Comparison between each experimental group and

Wildtype-Sham used multiple Mann-Whitney U tests. Statistics are presented in Table 1.

Despite significant comparisons on the group level, no regions appeared to be statistically

different, however some effect sizes when comparing ephrin-A2A5$^{-/-}$ groups with Wildtype-

Sham were moderate (0.3-0.5) suggesting that the proportion of c-Fos in several regions for

ephrin-A2/A5$^{-/-}$ mice tended to be larger than Wildtype-Sham.

Fig 8. Average region-by-region proportion of total striatal activity in the experimental

groups.

Table 1. Region by region Mann-Whitney U comparisons for proportion of striatal c-

Fos

Group comparison (Wildtype-Sham vs.)	Mann-Whitney U	p	Effect Size (Rank biserial correlation)	Mean Difference	95% Confidence Interval	
					Lower	Upper
Wildtype-rTMS						
Region						
DLS	14	0.926	0.0667	0.271	-2	4
DMS	11	0.537	0.267	-5.5	-22	21
NAc Core	14.5	1	0.0333	0.00368	-18	14
NAc Shell	12	0.662	0.200	5.5	-11	17
Ephrin-Sham						
Region						
DLS	11.5	0.333	0.361	-3.88	-21	3
DMS	15	0.699	0.167	-5	-24	20
NAc Core	11	0.31	0.389	9.5	-9	27
NAc Shell	17	0.936	0.056	3	-13	13
Ephrin-rTMS						
Region						
DLS	4.5	0.276	0.500	2	-2	6
DMS	7	0.714	0.222	4.5	-20	42
NAc Core	9	1	0	-0.0727	-23	17
NAc Shell	6	0.548	0.333	-4.5	-21	8

4   Discussion

Evidence of a habitual behavioural responding pattern in ephrin-A2/A5$^{-/-}$ mice led to

the hypothesis that these mice would display a shift in neuronal activity towards striatal

regions involved in habit formation (Moretti et al., 2021). However, contrary to the

hypothesis, neuronal activation was mostly seen in the regions of the striatum that are

involved in goal-directed behaviour. We also hypothesized that the LI-rTMS change in c-Fos

densities seen previously in ephrin-A2/A5$^{-/-}$ mice NAc would extend to the dorsal striatum. In

partial support of the hypothesis, we did not observe a change in total c-Fos expression after

LI-rTMS in ephrin-A2/A5$^{-/-}$ mice, but in the DMS there was an increase in MSN activation

following stimulation. Although the proportion of c-Fos labelling across the striatum was

significantly different in all experimental groups compared to control animals, there was not

one specific region driving this difference.

4.1   Neuronal activation assessed by c-Fos expression was highest in regions involved
in goal-directed behaviour

The overall proportion of c-Fos + cells was significantly different between Wildtype-

Sham group and all other groups. However, comparing the difference in proportions across

regions did not identify specific regions that were significantly different to the Wildtype-

Sham group. Nonetheless, the biggest effect sizes were associated with the difference

between Wildtype-Sham and the Ephrin groups, suggesting less activity in the DLS of

Ephrin-rTMS while the comparisons with Ephrin-Sham animals suggest less activity in the

NAc and greater activity in the DLS compared to Wildtype-Sham animals. Therefore, it

appears that there is an increased proportion of dorsal striatum activity in Ephrin-Sham

animals compared to Wildtype animals, and this difference is reduced following LI-rTMS.

This is in line with our hypothesis that LI-rTMS may delay an accelerated shift from goal-

directed behaviour to habit seen in ephrin-A2/A5$^{-/-}$ mice.

While looking at the change in c-Fos activity in specific striatal regions, it was found

that in both sham and treatment groups of ephrin-A2/A5[-/-] mice, DMS had more c-Fos

expression than DLS. Although DMS is a part of the nigrostriatal pathway, it is a transition

zone and is versatile in its function as it is involved in both stimulus-response and action-

outcome learning (Corbit et al., 2001) Therefore, DMS shows more activity when the

behaviour is goal-directed, and this activity slowly decreases with over-training and shifts to

DLS for formation of habits (Lipton et al., 2019). Similarly, there is a progression of c-Fos

expression during training in a fixed-ratio (FR) task, from higher c-Fos activity in NAc shell

on the first day of the task, towards the NAc core, DMS and finally to DLS as training

progressed (Segovia et al., 2012). In our study, the major proportion ofc-Fos+ cells remained

in the NAc core and DMS, suggesting that behaviour in all mice remained goal-directed (Fig

8). However, the Ephrin-Sham group showed signs of higher relative DLS activity, and less

NAc core activity compared to Wildtype-Sham group which supports our hypothesis that

behaviour may have shifted towards habit formation in ephrin-A2/A5[-/-] mice. A caveat is that,

unlike previous studies (Segovia et al., 2012) , using c-Fos as a proxy for neuronal activation

reflects only the final timepoint, after the completion of PR task. Also, in a PR task, unlike in

a FR schedule, the response requirement changes over time so mice cannot generally predict

how much work is required to obtain a reward based on previous trials. Therefore, although

we can assume a similar hierarchical progression of activity within the striatum, the variable

nature of the PR task could impact the behavioural and cellular response differently than

suggested for the FR framework (Segovia et al., 2012).

However, we conclude that the perseverative and inflexible behaviour in the ephrin-

A2/A5[-/-] mice used in this study (Moretti et al., 2021) is likely not due to habit formation,

because neuronal activity in these mice is mostly localised in regions responsible for goal-

directed behaviour. However, previous studies have shown other behavioural patterns in

ephrin-A knockout mice that suggest compulsive or perseverative tendencies. Self-injurious

compulsive grooming behaviour similar to some behaviours seen in autism spectrum disorder

has been reported in a different strain of ephrin-A knockout mice and attributed to

abnormalities in the corticostriatal pathway (Wurzman et al., 2015) Although we saw

repetitive responding pattern in the ephrin-A2/A5$^{-/-}$ mice in our study, we did not observe

self-injurious compulsive grooming. Further evidence of perseverative behaviour in ephrin-

A2$^{-/-}$ mice comes from maintenance of a responding pattern during a reversal task, suggesting

insensitivity to the reward (Arnall et al., 2010). This perseverative behaviour was broadly

attributed to abnormalities in the orbitofrontal cortex and the circuitry involving the thalamus,

striatum and the prefrontal cortex. Therefore, perturbations in ephrin-A signalling appear to

be consistently linked to compulsive and perseverative behaviour, suggesting that although

dopamine abnormalities play a key role in such behaviours, it may be that disrupted

organization of neural circuits may be an additional mechanism underpinning perseverative

and compulsive behavioural patterns. Neurological disorders often have genetic and

developmental factors contributing to their presentation (Ting and Feng, 2011). Therefore,

future studies with ephrin-A2/A5$^{-/-}$ mice could give insight into how disruption of neural

circuit organization during development have behavioural impacts (Pasquale, 2005; North et

al., 2013). Additionally, applying LI-rTMS and its history of successful network

reorganization in ephrin models (Rodger et al., 2012; Makowiecki et al., 2014; Poh et al.,

2022) to more established models of neuropsychiatric disorders, could be an interesting

avenue to elucidate therapeutic potential and mechanisms of rTMS.

4.2   c-Fos colocalization with MSNs and interneurons in the striatum

MSNs form the major population of striatal neurons (95%) with the remaining 5%

being interneurons (Melzer et al., 2017). These interneurons are quite diverse and have been

classified into two major classes, cholinergic interneurons and GABAergic interneurons

(Kawaguchi et al., 1995). In our study, we saw an increase in active MSNs (identified by c-

Fos-DARPP32 colocalization) in DMS of LI-rTMS-stimulated but not sham stimulated

ephrin-A2/A5$^{-/-}$ mice. This suggests that LI-rTMS specifically enhances DMS MSN activity,

which may contribute to delaying the shift in striatal activity to DLS. If LI-rTMS can

potentially delay an accelerated shift to habitual behaviour, it would be interesting to

investigate whether higher dose or more prolonged rTMS treatment could reverse the

hierarchical recruitment of striatal regions by shifting activity from the DLS towards regions

that are involved in early acquisition of a behaviour, perhaps through activation of DMS

MSNs. Such an effect would be relevant to reversing DLS activity associated with

overreliance on habits as seen in drug seeking behaviours (Willuhn et al., 2012).

Although MSNs account for the majority of striatal neurons, our qualitative analysis

across all experimental groups suggests that roughly 60% and 40% of c-Fos single-labelled

cells in the NAc and dorsal striatum respectively, were not MSNs. Therefore, as an

exploratory analysis we stained for GABAergic and cholinergic interneurons to identify the

nature of the unspecified c-Fos single-labelled cells. In all groups, DMS had more c-Fos^{+}-

GAD67^{+}/DARPP32^{-} colocalization and c-Fos^{+}-PV-expressing GABAergic interneurons

compared to DLS which could mean more interneuron-related inhibitory activity in the DMS.

However, we saw no significant changes in GAD67 activity between groups. In NAc core

and shell, across all experimental groups we found that less than 2% of the c-Fos^{+} cells that

were colocalised with GAD67^{+}/DARPP32^{-} cells. Therefore, although there were no major

group differences with GABAergic interneuron activity, the modulatory role of GABAergic

interneurons is likely most influential in the DMS due to the higher proportion of active

neurons that are GABAergic interneurons. Lastly, in our qualitative analysis we did not find

any evidence to support activation of cholinergic interneurons in either dorsal or ventral

striatum.

As we tested for all neuronal populations in the striatum, the identity of the high

number of unspecified c-Fos$^+$ cells is still not clear, as together the staining accounted for

only around 40% and 50% of c-Fos$^+$ cells in the ventral and dorsal striatum, respectively.

There is a possibility that these unspecified c-Fos$^+$ cells could be glial cells, however, a

confirmatory test in the ventral striatum found that most c-Fos$^+$ cells were colocalised with

neurons (Moretti et al., 2021).

4.3   Relevance

Although the small sample size of the study limits our conclusions, the possibility of

rTMS delaying habit formation is exciting. It shines light on potential mechanisms of rTMS

as a therapy for disorders such as drug addiction and OCD. rTMS is already used to treat

OCD, and there is evidence for anti-craving effects for cocaine (Camprodon et al., 2007) and

nicotine (Amiaz et al., 2009). The anti-craving effects of rTMS have been thought to be

linked to the ability of rTMS to modulate reward circuitry in the brain. For example, rTMS

may counteract addiction-related plasticity by increasing dopamine release and dopamine

receptor density, as well as modulating glutamatergic receptor expression (Moretti et al.,

2020). Ephrin-A2/A5$^{-/-}$ mice have low dopamine levels in the striatum possibly due to

reduced dopaminergic innervation (Cooper et al., 2009; Sheleg et al., 2013). Future studies

could look at whether rTMS can alter these abnormal dopaminergic innervations in ephrin-

A2/A5$^{-/-}$ mice to normalise dopamine levels in the striatum. Linking this to changes in

perseverative behaviour in ephrin-A2/A5$^{-/-}$ mice following rTMS could add to evidence that

can support rTMS as a therapy for disorders with overreliance on habits.

4.4   Conclusion

The present study did not fully support the assumptions made by our previous study

(Moretti et al., 2021) in which it was hypothesized that neuronal activity in ephrin-A2/A5$^{-/-}$

mice with repetitive and perseverative responding behaviour would have shifted to the striatal

region involved in habit formation. Rather, in this study, it was found that activity remained

dominant in regions involved in goal-directed behaviours. However, the relative proportion

of c-Fos activity in the striatum slightly shifted towards DLS activation in untreated ephrin-

A2/A5$^{-/-}$ mice when compared to other experimental groups. This was in line with our

hypothesis that ephrin-A2/A5$^{-/-}$ mice would have greater c-Fos activity in striatal regions

associated with habitual behaviour and that LI-rTMS would reduce this difference. As a

whole, rTMS seems to partly delay the shift from goal-directed behaviour to habit in ephrin-

A2/A5$^{-/-}$ mice, and this could be due to the changes in MSN activity in the DMS. Future

studies could perform electrophysiological recordings from the striatum to map any causal

relationship. Overall, LI-rTMS shows some promise for delaying the shift from goal-directed

behaviour to habit and/or compulsive behaviour, on a cellular level. This delay could be

further investigated in more clinically relevant animal models as a mechanism behind clinical

rTMS effects seen in disorders like drug addiction and OCD.

5   Acknowledgements

We thank Dr. Bhedita Seewoo and Mr. Samuel J Bolland for their guidance on creating the

mouse brain atlas for this study.

6   Author contributions

Maitri Tomar: Methodology, Writing - original draft, Writing – review & editing, Formal

analysis, Investigation, Visualization. **Jennifer Rodger.** Writing – review & editing,

Supervision. **Jessica Moretti.** Conceptualization, Writing – review & editing, Methodology,

Formal analysis, Visualization, Supervision.

7   Declaration of competing interests

The authors declare no conflict of interest.

8   References

Aberman, J. E., Ward, S. J., and Salamone, J. D. (1998). Effects of dopamine antagonists and

accumbens dopamine depletions on time-constrained progressive-ratio performance.

Pharmacol. Biochem. Behav. 61, 341–348. doi: 10.1016/S0091-3057(98)00112-9.

Amiaz, R., Levy, D., Vainiger, D., Grunhaus, L., and Zangen, A. (2009). Repeated high-

frequency transcranial magnetic stimulation over the dorsolateral prefrontal cortex

reduces cigarette craving and consumption. *Addiction* 104, 653–660. doi:

10.1111/j.1360-0443.2008.02448.x.

Anderson, K. D., and Reiner, A. (1991). Immunohistochemical localization of DARPP-32 in

striatal projection neurons and striatal interneurons: implications for the localization of

D1-like dopamine receptors on different types of striatal neurons. *Brain Res.* 568, 235–

243. doi: 10.1016/0006-8993(91)91403-N.

Arnall, S., Cheam, L. Y., Smart, C., Rengel, A., Fitzgerald, M., Thivierge, J. P., et al. (2010).

Abnormal strategies during visual discrimination reversal learning in ephrin-A2-/-mice.

Behav. Brain Res. 209, 109–113. doi: 10.1016/j.bbr.2010.01.023.

Aydin-Abidin, S., Trippe, J., Funke, K., Eysel, U. T., and Benali, A. (2008). High- and low-

frequency repetitive transcranial magnetic stimulation differentially activates c-Fos and

zif268 protein expression in the rat brain. *Exp. Brain Res.* 188, 249–261. doi:

10.1007/s00221-008-1356-2.

Bullitt, E. (1990). Expression ofC-fos-like protein as a marker for neuronal activity following

noxious stimulation in the rat. *J. Comp. Neurol.* 296, 517–530. doi:

10.1002/cne.902960402.

Camprodon, J. A., Martínez-Raga, J., Alonso-Alonso, M., Shih, M. C., and Pascual-Leone,

A. (2007). One session of high frequency repetitive transcranial magnetic stimulation

(rTMS) to the right prefrontal cortex transiently reduces cocaine craving. *Drug Alcohol*

Depend. 86, 91–94. doi: 10.1016/j.drugalcdep.2006.06.002.

Carmi, L., Alyagon, U., Barnea-Ygael, N., Zohar, J., Dar, R., and Zangen, A. (2018). Clinical

and electrophysiological outcomes of deep TMS over the medial prefrontal and anterior

cingulate cortices in OCD patients. *Brain Stimul.* doi: 10.1016/j.brs.2017.09.004.

Chon, U., Vanselow, D. J., Cheng, K. C., and Kim, Y. (2019). Enhanced and unified

anatomical labeling for a common mouse brain atlas. *Nat. Commun.* 10, 5067. doi:

10.1038/s41467-019-13057-w.

Cooper, M. A., Kobayashi, K., and Zhou, R. (2009). Ephrin-A5 regulates the formation of the

ascending midbrain dopaminergic pathways. *Dev. Neurobiol.* 69, 36–46. doi:

10.1002/dneu.20685.

Corbit, L. H., Muir, J. L., and Balleine, B. W. (2001). The role of the nucleus accumbens in

instrumental conditioning: Evidence of a functional dissociation between accumbens

core and shell. *J. Neurosci.* 21, 3251–3260. doi: 10.1523/jneurosci.21-09-03251.2001.

Dickinson, A., Balleine, B., Watt, A., Gonzalez, F., and Boakes, R. A. (1995). Motivational

control after extended instrumental training. *Anim. Learn. Behav.* 23, 197–206. doi:

10.3758/BF03199935.

Everitt, B. J., and Robbins, T. W. (2005). Neural systems of reinforcement for drug addiction:

From actions to habits to compulsion. *Nat. Neurosci.* 8, 1481–1489. doi:

10.1038/nn1579.

Feldheim, D. A., Kim, Y.-I. I., Bergemann, A. D., Frisén, J., Barbacid, M., and Flanagan, J.

G. (2000). Genetic analysis of ephrin-A2 and ephrin-A5 shows their requirement in

multiple aspects of retinocollicular mapping. *Neuron* 25, 563–574. doi: 10.1016/S0896-

6273(00)81060-0.

Gillan, C. M., Papmeyer, M., Morein-Zamir, S., Sahakian, B. J., Fineberg, N. A., Robbins, T.

W., et al. (2011). Disruption in the Balance Between Goal-Directed Behavior and Habit

Learning in Obsessive-Compulsive Disorder. *Am. J. Psychiatry* 168, 718–726. doi:

10.1176/appi.ajp.2011.10071062.

Hodos, W. (1961). Progressive Ratio as a Measure of Reward Strength. *Science (80-.).* 134,

943–944. doi: 10.1126/science.134.3483.943.

Kawaguchi, Y., Wilson, C. J., Augood, S. J., and Emson, P. C. (1995). Striatal interneurones:

chemical, physiological and morphological characterization. *Trends Neurosci.* 18, 527–

535. doi: 10.1016/0166-2236(95)98374-8.

Keck, M. E., Welt, T., Muller, M. B., Erhardt, A., Ohl, F., Toschi, N., et al. (2002).

Repetitive transcranial magnetic stimulation increases the release of dopamine in the

mesolimbic and mesostriatal system. *Neuropharmacology* 43, 101–109. doi:

S0028390802000692 [pii].

Lazarus, M. S., Krishnan, K., and Huang, Z. J. (2015). GAD67 Deficiency in Parvalbumin

Interneurons Produces Deficits in Inhibitory Transmission and Network Disinhibition in

Mouse Prefrontal Cortex. *Cereb. Cortex* 25, 1290–1296. doi: 10.1093/cercor/bht322.

Lein, E. S., Hawrylycz, M. J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., et al. (2007).

Genome-wide atlas of gene expression in the adult mouse brain. *Nature* 445, 168–176.

doi: 10.1038/nature05453.

Lipton, D. M., Gonzales, B. J., and Citri, A. (2019). Dorsal Striatal Circuits for Habits,

Compulsions and Addictions. *Front. Syst. Neurosci.* 13. doi: 10.3389/fnsys.2019.00028.

Liu, H.-S., Chefer, S., Lu, H., Guillem, K., Rea, W., Kurup, P., et al. (2013). Dorsolateral

caudate nucleus differentiates cocaine from natural reward-associated contextual cues.

Proc. Natl. Acad. Sci. 110, 4093–4098. doi: 10.1073/pnas.1207531110.

Makowiecki, K., Harvey, A. R., Sherrard, R. M., and Rodger, J. (2014). Low-Intensity

Repetitive Transcranial Magnetic Stimulation Improves Abnormal Visual Cortical

Circuit Topography and Upregulates BDNF in Mice. *J. Neurosci.* 34, 10780–10792. doi:

10.1523/JNEUROSCI.0723-14.2014.

Melzer, S., Gil, M., Koser, D. E., Michael, M., Huang, K. W., and Monyer, H. (2017).

Distinct Corticostriatal GABAergic Neurons Modulate Striatal Output Neurons and

Motor Activity. *Cell Rep.* 19, 1045–1055. doi: 10.1016/j.celrep.2017.04.024.

Moretti, J., Poh, E. Z., Bolland, S. J., Harvey, A. R., Albrecht, M. A., and Rodger, J. (2021).

Concurrent LI-rTMS induces changes in c-Fos expression but not behavior during a

progressive ratio task with adult ephrin-A2A5-/- mice. *Behav. Brain Res.* 400, 113011.

doi: 10.1016/j.bbr.2020.113011.

Moretti, J., Poh, E. Z., and Rodger, J. (2020). rTMS-Induced Changes in Glutamatergic and

Dopaminergic Systems: Relevance to Cocaine and Methamphetamine Use Disorders.

Front. Neurosci. 14. doi: 10.3389/fnins.2020.00137.

North, H. A., Clifford, M. A., and Donoghue, M. J. (2013). "Til Eph Do Us Part":

Intercellular Signaling via Eph Receptors and Ephrin Ligands Guides Cerebral Cortical

Development from Birth Through Maturation. *Cereb. Cortex* 23, 1765–1773. doi:

10.1093/cercor/bhs183.

O'Reardon, J. P., Solvason, H. B., Janicak, P. G., Sampson, S., Isenberg, K. E., Nahas, Z., et

al. (2007). Efficacy and Safety of Transcranial Magnetic Stimulation in the Acute

Treatment of Major Depression: A Multisite Randomized Controlled Trial. *Biol.*

Psychiatry 62, 1208–1216. doi: 10.1016/j.biopsych.2007.01.018.

Pasquale, E. B. (2005). Eph receptor signalling casts a wide net on cell behaviour. *Nat. Rev.*

Mol. Cell Biol. 6, 462–475. doi: 10.1038/nrm1662.

Paxinos, G., and Franklin, K. (2012). *Paxinos and Franklin's the Mouse Brain in Stereotaxic*

Coordinates. 4th ed. Cambridge, MAq: Academic Press.

Poh, E. Z., Green, C., Agostinelli, L., Penrose-Menz, M., Karl, A. K., Harvey, A. R., et al.

(2022). Manipulating the Level of Sensorimotor Stimulation during LI-rTMS Can

Improve Visual Circuit Reorganisation in Adult Ephrin-A2A5-/- Mice. *Int. J. Mol. Sci.*

23. doi: 10.3390/ijms23052418.

Poh, E. Z., Harvey, A. R., Makowiecki, K., and Rodger, J. (2018). Online LI-rTMS during a

Visual Learning Task⬚: differential Impacts on Visual Circuit and Behavioural

Plasticity in Adult Ephrin-A2A5 Mice. *eNeuro* 5, ENEURO.0163-17.2018. doi:

10.1523/ENEURO.0163-17.2018.

Rodger, J., Mo, C., Wilks, T., Dunlop, S. A., and Sherrard, R. M. (2012). Transcranial pulsed

magnetic field stimulation facilitates reorganization of abnormal neural circuits and

corrects behavioral deficits without disrupting normal connectivity. *FASEB J.* 26, 1593–

1606. doi: 10.1096/fj.11-194878.

Segovia, K. N., Correa, M., Lennington, J. B., Conover, J. C., and Salamone, J. D. (2012).

Changes in nucleus accumbens and neostriatal c-Fos and DARPP-32 immunoreactivity

during different stages of food-reinforced instrumental training. *Eur. J. Neurosci.* 35,

1354–1367. doi: 10.1111/j.1460-9568.2012.08036.x.

Sheleg, M., Yochum, C. L., Wagner, G. C., Zhou, R., and Richardson, J. R. (2013). Ephrin-

A5 deficiency alters sensorimotor and monoaminergic development. *Behav. Brain Res.*

236, 139–147. doi: 10.1016/j.bbr.2012.08.032.

Sieber, B. A., Kuzmin, A., Canals, J. M., Danielsson, A., Paratcha, G., Arenas, E., et al.

(2004). Disruption of EphA/ephrin-A signaling in the nigrostriatal system reduces

dopaminergic innervation and dissociates behavioral responses to amphetamine and

cocaine. *Mol. Cell. Neurosci.* 26, 418–428. doi: 10.1016/j.mcn.2004.03.009.

Ting, J. T., and Feng, G. (2011). Neurobiology of obsessive–compulsive disorder: insights

into neural circuitry dysfunction through mouse genetics. *Curr. Opin. Neurobiol.* 21,

842–848. doi: 10.1016/j.conb.2011.04.010.

Wilkinson, D. G. (2001). Multiple roles of eph receptors and ephrins in neural development.

Nat. Rev. Neurosci. 2, 155–164. doi: 10.1038/35058515.

Willuhn, I., Burgeno, L. M., Everitt, B. J., and Phillips, P. E. M. (2012). Hierarchical

recruitment of phasic dopamine signaling in the striatum during the progression of

cocaine use. *Proc. Natl. Acad. Sci.* 109, 20703–20708. doi: 10.1073/pnas.1213460109.

Wurzman, R., Forcelli, P. A., Griffey, C. J., and Kromer, L. F. (2015). Repetitive grooming

and sensorimotor abnormalities in an ephrin-A knockout model for Autism Spectrum

Disorders. *Behav. Brain Res.* 278, 115–128. doi: 10.1016/j.bbr.2014.09.012.

Yushkevich, P. A., Piven, J., Hazlett, H. C., Smith, R. G., Ho, S., Gee, J. C., et al. (2006).

User-guided 3D active contour segmentation of anatomical structures: Significantly

improved efficiency and reliability. *Neuroimage* 31, 1116–1128. doi:

10.1016/j.neuroimage.2006.01.015.

A

B