Structure of the photosynthetic Calvin-Benson-Bassham sedoheptulose-1,7-bisphosphatase SBPase from the model microalga *Chlamydomonas reinhardtii*

Théo Le Moigne[1,2], Martina Santoni[1,3], Lucile Jomat[1], Stéphane D. Lemaire[1], Mirko Zaffagnini[3], Nicolas Chéron[4], Julien Henri[1,*]

1. Sorbonne Université, CNRS, Laboratoire de Biologie Computationnelle et Quantitative UMR 7238, Institut de Biologie Paris-Seine, 4 Place Jussieu, Paris F-75005, France

2. Faculty of Sciences, Doctoral School of Plant Sciences, Université Paris-Saclay, Saint-Aubin F-91190, France

3. Department of Pharmacy and Biotechnology, University of Bologna, Via Irnerio 42, Bologna I-40126, Italy

4. Ecole Normale Supérieure, CNRS, Sorbonne Université, Laboratoire Pasteur UMR 8640, Département de Chimie de l'Ecole Normale Supérieure rue d'Ulm, F-Paris 75005, France

*corresponding author

e-mail: julien.henri@sorbonne-universite.fr

Phone: +33144276606

ORCID: Théo Le Moigne 0000-0003-2313-3052; Stéphane Lemaire 0000-0001-6442-0547, Mirko Zaffagnini 0000-0001-5115-0859, Nicolas Chéron 0000-0002-4090-5897; Julien Henri 0000-0003-0772-8881

Abstract

The Calvin-Benson-Bassham cycle (CBBC) performs carbon fixation in photosynthetic organisms. Among the eleven enzymes that participate in the pathway, sedoheptulose-1,7-bisphosphatase (SBPase) is expressed in photo-autotrophs and contributes to the regeneration of ribulose-1,5-bisphosphate, the carbon fixation co-substrate used by ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco). While SBPase is structurally similar to fructose-1,6-bisphosphatase (FBPase) involved in both neoglucogenesis and the CBBC, it exclusively functions in the CBBC and is indispensable for building a productive cycle. In this study we report the first structure of an SBPase from a chlorophyte, the model unicellular green microalga *Chlamydomonas reinhardtii*. By combining experimental and computational structural analyses, we describe the topology, conformations and quaternary structure of *Chlamydomonas reinhardtii* SBPase. We identify active site residues and locate sites of redox- and phospho-post-translational modifications that contribute to enzymatic functions. Finally, we observe that *Cr*SBPase adopts distinct oligomeric states that may dynamically contribute to the control of its activity.

Introduction

Photosynthetic carbon fixation is performed by eleven enzymes that collectively operate the Calvin-Benson-Bassham cycle (CBBC) (Calvin 1962, Johnson 2016), wherein Rubisco plays the prominent role in catalyzing fixation of inorganic carbon onto a pentose acceptor: ribulose-1,5-bisphosphate (RuBP). The net gain of the CBBC is one triose in the form of glyceraldehyde-3-phosphate (G3P) for every three Rubisco catalyzed carboxylation reactions (Johnson and Alric 2013). Regeneration of RuBP is ensured by the other ten enzymes of the CBBC, among which phosphoribulokinase (PRK) catalyzes the phosphorylation of ribulose-5-phosphate (R5P) and sedoheptulose-1,7-bisphosphatase (SBPase) catalyzes the hydrolysis of sedoheptulose-1,7-bisphosphate (SBP) into sedoheptulose-7-phosphate (S7P). Remaining CBBC enzymes are paralogs of those involved in cytoplasmic glycolysis, neoglucogenesis or the pentose-phosphate pathway. Hence, Rubisco, PRK, and SBPase are expressed

in photosynthetic organisms and fulfil biochemical functions so far uniquely attributed to the photosynthetic CBBC. The CBBC is perfectly conserved in plants and cyanobacteria since its enzymes are expressed in oxygenic prokaryotes and plant chloroplasts, which consist in subcellular compartments that evolved from ancestral endosymbiosis between cyanobacteria and heterotrophic eukaryotic cells. For instance, the analysis of PRK from *Synechococcus* PCC 6301, *Arabidopsis thaliana*, and *Chlamydomonas reinhardtii* display high structural similarities (Gurrieri, Del Giudice et al. 2019, Wilson, Hayer-Hartl et al. 2019).

Native SBPase was purified from wheat chloroplasts (Woodrow 1982) and corn leaves (Nishizawa and Buchanan 1981), allowing quantitative assays of its catalytic function and the *in vitro* confirmation of magnesium-dependent activation and activity modulation by pH and redox switch mechanisms (Anderson 1974, Woodrow and Walker 1980, Cadet and Meunier 1988). Redox control is exerted through a dithiol/disulfide interchange dependent upon the ferredoxin-thioredoxin cascade arising from the activity of the illuminated photosystems (Schürmann and Buchanan 1975). In tobacco leaves extracts, SBPase reaches its maximal activity after 10 minutes illumination (Zimmer, Swart et al. 2021). This delayed activation kinetics is consistent with the slow reduction pattern of *Arabidopsis* SBPase *in vitro* (Yoshida and Hisabori 2018).

The crystal structure of SBPase from the moss *Physcomitrium patens* (*Pp*SBPase) has been solved at a resolution of 1.3 Å (Protein Data Bank entry : 5iz3) (Gutle, Roret et al. 2016). *Pp*SBPase fold is highly similar to that of moss FBPase, yet the protein was shown to distinctively oligomerize as a homodimer and to form a disulfide bridge in a different motif than that previously described for FBPase (Chiadmi, Navaza et al. 1999). In spite of their homology, the sequences of FBPase and SBPase seem to have evolved different regulatory properties that allow different contributions to the CBBC. Recently, it was proposed that the CBBC can bypass fructose-1,6-bisphosphate (FBP) altogether, by only relying on SBP, glycolysis and the oxidative pentose-phosphate pathway paralogs to regenerate the substrate of Rubisco (Ohta 2022). Engineered SBPase overexpression enhances photosynthesis and growth in plants (Lefebvre, Lawson et al. 2005, Feng, Han et al. 2007, Feng, Wang

et al. 2007, Rosenthal, Locke et al. 2011, Simkin, Lopez-Calcagno et al. 2017) and in the green

microalga *Chlamydomonas reinhardtii* (Hammel, Sommer et al. 2020), indicating that SBPase appears

as a metabolic bottleneck for the CBBC along with Rubisco. Based on these evidences, it is plausible

to consider that SBPase is a good target to improve the photosynthetic capacity by means of

metabolic engineering coupled to synthetic biology (Kubis and Bar-Even 2019), especially in the

experimentally convenient multi-omic model alga Chlamydomonas (Mettler, Muhlhaus et al. 2014,

Schmollinger, Mühlhaus et al. 2014).

In Chlamydomonas, SBPase is encoded by a single nuclear gene (Cre03.g185550), and the protein

(Uniprot entry P46284) is addressed to the chloroplast by a 54-residue amino-terminal peptide

(Emanuelsson, Brunak et al. 2007, Almagro Armenteros, Salvatore et al. 2019). SBPase abundance in

the algal cell was quantified by mass spectrometry as 1.1-1.3 ± 0.3 amol/cell, representing 0.15 % of

total cell proteins (Hammel, Sommer et al. 2020). In Chlamydomonas protein extracts, SBPase was

shown to undergo post-translational modifications by phosphorylation of serines and threonines

(Werth, McConnell et al. 2017), and redox modifications of cysteine thiols (Zaffagnini, Bedhomme et

al. 2012, Morisse, Zaffagnini et al. 2014, Pérez-Pérez, Mauriès et al. 2017). However, the mechanisms

by which post-translational modifications affect the catalytic activity of CrSBPase is still lacking. Here,

we describe the crystal structures of *Cr*SBPase in two redox states, assay its activity *in vitro*, and

investigate the correlation between the redox state and the oligomerization of the protein. Our

results suggest that a redox control over oligomer equilibrium could represent a novel mechanism to

regulate enzymatic activities in the CBBC.

Material and methods

Chemical reagents were purchased from Merck (Darmstadt, Germany), chromatography material

was provided by Cytiva (Vélizy-Villacoublay, France), crystallization solutions and consumable were

purchased from Qiagen (Hilden, Germany), Hampton Research (Aliso Viejo, CA USA), and SPT Labtech

(Melbourn, UK).

Cloning and mutagenesis of *Cr*SBPase expression plasmids

The sequence coding for full-length mature SBPase was amplified by reverse transcriptase

polymerase chain reaction (RT-PCR) from *Chlamydomonas reinhardtii* total ribonucleic acids extract

and inserted at sites 5'-NdeI and BamHI-3' into expression vector pET3c modified for the fusion of an

amino-terminal hexa-histidine tag in the recombinant protein, yielding plasmid pSL-175-His$_6$-

*Cr*SBPase-WT. PCR primers were 5'-TCTCGC<u>CATATG</u>GCCGTTCTGACCCAGGCC-3' (forward, NdeI site

underlined) and 5'-CGGGTG<u>GGATCC</u>TTAGGCAGCCACCTTCTC-3' (reverse, BamHI site underlined). The

encoded protein encompasses 331 amino acids, with a molar mass of 36,176.7 Da and a theoretical

extinction coefficient of 24,722.5 M^{-1}.cm^{-1}. Unless otherwise stated, amino acid residues numbering

is that of GenBank entry DAA79954.1. Point mutants of cysteines 115 and 120 singly substituted with

serines were generated by site-directed PCR mutagenesis with primers listed in supplementary table

1, producing plasmids pSL-181-His6-*Cr*SBPase-C115S, and pSL-182-His6-*Cr*SBPase-C120S.

Recombinant *Cr*SBPase purification

Plasmids pSL-175, -181, and -182 were used to transform *Escherichia coli* BL21(DE3). The

transformants were grown at 37 °C in 2YT supplemented with 150 μg/mL ampicillin. When cultures

reached an OD$_{600}$ of ~0.6, T7-dependent *Cr*SBPase overexpression was induced by the addition of

isopropyl-β-D-thiogalactopyranoside (IPTG) at 0.2 mM for 3 h.

Harvested cell pellets were resuspended in 20 mmol.L^{-1} Tris pH 7.9, 100 mmol.L^{-1} NaCl (buffer A) and

lyzed by 1 sec / 1 sec pulsed sonication for 3 min. The soluble fraction of the lysate was separated by

20 min centrifugation at 20,000 *g* and loaded on 2 mL Ni-NTA resin for His-tag mediated affinity

chromatography. The resin was washed in four steps with 25 mL buffer A, 25 mL buffer A

supplemented with 10, 20, and 30 mmol.L^{-1} imidazole, and *Cr*SBPase was eluted with 10 mL buffer A

supplemented with 100, 200, and 300 mmol.L^{-1} imidazole. Pooled eluates containing *Cr*SBPase were

loaded on HiLoad Superdex 200 26/600 size-exclusion chromatography column and eluted in buffer

A. Protein purity was assessed by electrophoresis on 12% acrylamide gel under denaturing and

reducing conditions (supplementary figure 1). Peak fractions were assembled and concentrated by

ultrafiltration on Amicon units of MWCO 30,000 Da cut-off. Final concentrations of purified proteins

were in the 1-10 $mg.mL^{-1}$ range as measured by Nanodrop ND-2000 spectrophotometer.

Western blot

Polyclonal antibodies were generated by immunization of rabbits with pure recombinant *Cr*SBPase

(Genecust, Boynes France). Analyzed fractions were separated by denaturing polyacrylamide

electrophoresis and transferred to nitrocellulose membrane for detection with primary antibody

subsequently detected by secondary anti-rabbit antibodies coupled to horseradish peroxidase.

Detection was done with commercial ECL peroxidase assay (GE Healthcare, Chicago IL USA) with a

Chemidoc (Bio-Rad, Hercules CA USA).

Crystallization and structure determination

Pure recombinant His6-*Cr*SBPase-WT concentrated at 3-6 mg/mL was tested for crystallization on

screens JCSG I-IV in 200 nL sitting drops. Single crystals were obtained and optimized with mother

liquor 0.1 $mol.L^{-1}$ sodium HEPES pH=7.5, 2% (v/v) polyethylene glycol 400, 2.0 $mol.L^{-1}$ ammonium

sulfate, flash-frozen in mother liquor supplemented with 25% glycerol and tested for X-ray

diffraction. Complete X-ray diffraction datasets were collected at SOLEIL beamline Proxima-2A and

allowed the determination of the crystal structure by model replacement with 5iz3, the structure of

ortholog SBPase from the moss *Physcomitrium patens.* Model building in COOT and refinement in the

PHENIX software suite yielded structure 7b2o (Emsley, Lohkamp et al. 2010) (Adams, Afonine et al.

2011) (Liebschner, Afonine et al. 2019). After we obtained our experimental structure, the European

Bioinformatics Institute (Hinxton, UK) communicated high-accuracy prediction models computed by

Deepmind ALPHAFOLD2 (Jumper, Evans et al. 2021), including a prediction of *Cr*SBPase structure

(https://alphafold.ebi.ac.uk/entry/P46284) that matches 7b2o crystal structure with RMSD=0.453 Å.

Hence, X-ray crystallography and ALPHAFOLD independently contribute to the proposed structural

analysis.

In order to determine the structure of CrSBPase in a reduced state, crystals were grown in the

presence of 10 mmol.L^{-1} of the reducing agent tris-(2-carboxyethyl)phosphine (TCEP) in condition

JCSG IV E8/56 0.2 M lithium sulfate, 0.1 M Tris pH 7.0, 1.0 M sodium/potassium tartrate. Crystals

were cryo-protected in mother liquor supplemented with 25% ethylene glycol, flash-frozen and

tested for X-ray diffraction at SOLEIL beamline Proxima-2A. A complete dataset was collected that

allowed the determination of the structure by molecular replacement with 7b2o as a template.

Among the eight CrSBPase copies of the asymmetric unit, two had their disulfides absent as a

consequence of the TCEP treatment while the others six subunits were essentially identical to

untreated structures in entry 7b2o.

Enzymatic assays

The catalytic activity of 10-200 nmol.L^{-1} pure recombinant CrSBPase was spectrophotometrically

assayed in buffer A by coupling the hydrolysis of 1 M fructose-1,6-bisphosphate (FBP) into fructose-6-

phosphate (F6P) to the reduction of 0.5 M NADP^{+} into NADPH through isomerization of F6P into

glucose-6-phosphate (G6P) by 0.5 U.mL^{-1} phosphoglucose isomerase (PGI) and the oxidation of G6P

into 6-phosphogluconolactone (6PGL) by 0.1-0.5 U.mL^{-1} glucose-6-phosphate dehydrogenase

(G6PDH). Reporter assay is coupled in a molar ratio FBP:F6P:G6P:6PGL of 1:1:1:1 to the oxidation of

one molar equivalent of NADP^{+} into NADPH that we recorded over 0-20 minutes by measuring the

absorbance at 340 nm using a UVIKON spectrophotometer. Pre-treatment with 0-100 mmol.L^{-1} DTT,

0-30 mmol.L^{-1} MgSO$_4$, and 0-15 µmol.L^{-1} recombinant CrTRX-f2 supplemented with 1 mM DTT

(Lemaire, Tedesco et al. 2018) was carried out for 30 min at room temperature before adding the

enzyme in the assay solution.

Titration of accessible cysteine thiols

Quantification of free reactive cysteines on recombinant CrSBPase was performed with 5,5′-dithio-

bis-(2-nitrobenzoic acid) (DTNB, Ellman's reagent) mixed with the purified recombinant protein in a

200:1 molar ratio in a 1 mL reaction volume containing 100 mM Tris-HCl, pH 7.9. Reaction of DTNB

with solvent-exposed cysteine thiols produces 2-nitro-5-thiobenzoate (TNB⁻) that we quantified by

monitoring the absorbance at 412 nm and a molar extinction coefficient of TNB⁻ of 14.15 $mM^{-1}.cm^{-1}$.

We calculated the number of free reactive cysteines per CrSBPase monomer by dividing the molar

concentration of TNB⁻ by that of the protein at equilibrium (t >60 min).

Molecular dynamics

Molecular dynamics (MD) simulations were performed with the Gromacs package (v2021.4). A dimer

of CrSBPase was solvated in a rhombic dodecahedron box with at least 8 Å between the protein

atoms and the box edges, which corresponds to ~64000 atoms in total (~18000 water molecules) and

a triclinic box of 97x97x68 $Å^{3}$. The box was neutralized with 18 sodium atoms to make it neutral. The

starting conformation was the crystallized structure of oxidized CrSBPase, and we protonated C115

and C120 to manually create the reduced state. The protein was described with the Amber14SB force

field and water with the TIP3P model. Non-bonding interactions were described with standard

parameters: van der Waals interactions had a cut-off of 8 Å, and electrostatic interactions were

computed with PME with default parameters with a separation between spaces at 8 Å. Bonds

containing a hydrogen atom were restrained with the LINCS algorithm with default parameters. After

an energy minimization, the system was equilibrated under the NPT conditions with a δt=1 fs

timestep, the velocity-rescale thermostat and the Berendsen barostat. Velocities were generated at

100 K and the system was heated up to 300 K in 400 ps before performing 100 ps at 300 K. We then

ran simulations of 20 μs under the NPT conditions with a δt=2 fs timestep, the velocity-rescale

thermostat and the Parrinello-Rahman barostat.

Small-angle X-ray scattering

Pure recombinant *Cr*SBPase was injected on Agilent HPLC BioSEC-3 300 column in buffer A, in line

with the X-ray beam capillary at SOLEIL beamline SWING. *Cr*SBPase was untreated or pre-treated

with 10 mmol.L^{-1} of reducing agent dithiothreitol (DTTred) or oxidizing agent trans-1,2-dithiane-4,5-

diol (DTTox). Diffusion curves were analyzed with ATSAS software to determine the molecular mass

and oligomeric state of the protein.

Results

Crystal structure of *Cr*SBPase

Pure recombinant *Cr*SBPase crystallized in space group P2$_1$2$_1$2 and provided a complete X-ray

diffraction dataset at a resolution of 3.09 Å. The asymmetric unit is composed of six polypeptide

chains packing as dimers of dimers (see below) and four water molecules. All chains align within a

root mean square deviation (RMSD) <0.25 Å and can be considered identical. Three hundred and

nine out of the 331 residues were built in the electron density, the unmodelled sequences are in the

disordered extensions at the amino- and carboxy-termini of the model. *Cr*SBPase folds into two

distinct domains (figure 1A-B), an amino-terminal domain composed of a mixed β-sheet formed by

strands 3-9 and a carboxy-terminal domain composed of a mixed β-sheet formed by strands 10-13.

α-helices 1-3 pack on the amino-terminal end of the seven-strand sheet, while α-helices 4,5, and 8

pack on the carboxy-terminal end of the four-strand sheet. α-helices 6 and 7 are sandwiched in

between the two sheets. A short antiparallel β-sheet is constituted of 4-residue strands 1 and 2

separated by a β-turn and preceded by loop A$_{113}$SCAGTAC$_{120}$. This motif projects to solvent from α-

helices 2 and 3. Residues 113-130 forming the loop and β-hairpin (LBH) motif are positioned away

from the core of the protein and will be further discussed in the next sections.

CrSBPase catalytic pocket

*Cr*SBPase is structurally similar to its ortholog from the moss *Physcomitrium patens* (Gutle, Roret et al. 2016). SBPase is a homolog of FBPase, another enzyme involved in the CBBC in photosynthetic organisms and also ubiquitously involved in neoglucogenesis. Cytosolic isoforms of FBPase from non-plant sources have been extensively studied (Moorhead, Hodgson et al. 1994, Lee and Hahn 2003). For instance, the crystal structure of human muscle FBPase complexed with its substrate (FBP) was solved (PDB entry: 5l0a, to be published). We aligned the structure *Cr*SBPase onto that of human FBPase:FBP complex, confirming their similarity (RMSD = 0.824 Å) with the notable exception of the LBH motif which is absent in the human enzyme. In a 4-Å distance to FBPase co-crystallized with FBP we aligned 12 residues that are likely available for FBP or SBP binding by *Cr*SBPase (figure 1C): E155, D173, D176, G177, Y287, G289, G290, M291, K317, L318, R319, and E323. These residues are exposed to the solvent in a continuous surface (figure 1D). Water molecule 401 could be modelled at the centre of the putative SBPase catalytic pocket, in the vicinity of E155, D173, D176, R319, and E323, in the catalytic pocket predicted from alignment with human FBPase bound to FBP. W401 specifically is in bonding distance to catalytic residues D178, D181, and E328. W401 is likely to react with the substrate SBP during hydrolysis.

Sites of redox post-translational modifications

Ten cysteines could be located in the model. Previous redox-based proteomic studies established that SBPase cysteines 115, 120, 222, 231, 355, and 362 are redox-modified by thioredoxin-dependent dithiol/disulfide exchange (Lemaire, Guillon et al. 2004, Pérez-Pérez, Mauriès et al. 2017), S-glutathionylation on unidentified cysteines (Zaffagnini, Bedhomme et al. 2012), or S-nitrosylation at C355 and C362 (Morisse, Zaffagnini et al. 2014). C222, C231, C355, and C362 are putative targets of thioredoxin. DTNB-based thiol titration on the native purified SBPase indicates that 4.0 ± 0.6 cysteines are likely reactive owing to their relative exposure to solvent in a free thiol form. Amino acid residue accessible surface area and accessibility (ASA) calculation is consistent with DTNB

titration, with C149, C153, C231, and C355 exposed to the solvent (ASA respective areas: 0.154 Å²,

0.113 Å², 0.308 Å², and 10.083 Å²), and the six others either shielded from solvent (C169, C222, C340,

and C362) or engaged in a disulfide bridge (C115, C120) (Ahmad, Gromiha et al. 2004). In *Cr*SBPase

crystal structure, C222 and C231 are positioned on neighbouring strands of the amino-terminal

domain, exposing their thiol 3.5 Å away from each other (Figure 1E). The formation of an

intramolecular disulfide bridge at this cysteines pair is possible if local main chain rearrangements

allow a closer contact. C362 thiol is positioned 10.5 Å away from the nearest thiol group of C340.

Additionally, C340 and C362 are exposed on an opposite side of the carboxy-terminal β-sheet,

disfavouring the formation of an internal disulfide bridge. C355 points its side chain towards C153

and C169 with a thiol-thiol distance of 4.6 Å and 10.1 Å, respectively. Although not observed in our

model, a C355-C153/C169 dithiol-disulfide exchange is thus plausible, as described for plant FBPase

(Jacquot, Lopez-Jaramillo et al. 1997). Local rearrangements that occur upon FBPase disulfide

reduction causes a larger scale relaxation of the active site required to reach maximal catalytic

potential (Chiadmi, Navaza et al. 1999) as kinetically characterized for *Arabidopsis* (Yoshida and

Hisabori 2018) and *Nicotiana* enzymes (Zimmer, Swart et al. 2021). In SBPase, the FBPase-type

regulatory disulfide is not observed but the C115-C120 bridge in the LBH motif may regulate the

enzyme by alternative conformational rearrangements. Indeed, the disulfide-bonded cysteines

constrain loop $A_{113}SCAGTAC_{120}$ into a compact lasso conformation that could open into a relaxed

conformation hypothetically activating catalysis.

Sites of post-translational phosphorylations

*Cr*SBPase was reported to be phosphorylated from algae extracts at residues T112, S114, T118, S123,

T310, S311, and T313 (Wagner, Gessner et al. 2006, Werth, McConnell et al. 2017, McConnell, Werth

et al. 2018). Consistently with the proximity of these residues in two sectors of the sequence

(supplementary figure 2), the candidate modification sites are located in two contiguous regions in

the three-dimensional model (figure 1F). T112, S114, T118, and S123 are exposed to the solvent of

the amino-terminal domain. The residues belong to or are located nearby the $A_{113}SCAGTAC_{120}$ motif

where phosphorylations may cross-signal with redox modifications on C115 and C120. T310, S311,

and T313 are located on the other edge of the enzyme, at the solvent exposed tip of the carboxy-

terminal domain. Their co-localization at the $T_{310}SPT_{313}$ motif likely facilitates a coordinated

phosphorylation of the three residues by the same kinase. Because both groups of potential

phospho-sites are located at 17-18 Å from the catalytic pocket (H_2O 401 taken as a reference point),

there is no straightforward mechanism by which such modifications would exert a control over

*Cr*SBPase activity. Analysis of phospho-mimicking mutants along with the identification of the actual

kinases/phosphatases couples that control SBPase phosphorylation state would allow   to

comprehend the role of this putative regulatory mechanism.

_Cr_SBPase enzymatic activity

Recombinant *Cr*SBPase was assayed for its capacity to catalyze FBP hydrolysis into F6P (Gutle, Roret

et al. 2016), a proxy for the hydrolysis of SBP into S7P that we could not test by lack of available

substrate and kinetic reporter method. Previous studies demonstrated that plant SBPase requires the

cofactor Mg^{2+} and a chemical reductant (DTT) to be fully activated *in vitro* (Anderson 1974, Woodrow

and Walker 1980, Cadet and Meunier 1988). In agreement with this, we found a total absence of

activity in the untreated CrSBPase (data not shown), whereas pre-incubation of the enzyme with DTT

and Mg^{2+} resulted in a strong activation (figure 2). By testing different concentrations of $MgSO_4$ (3,

10, 15, and 20 mM) and DTT (1, 10, 25, 50, and 100 mM), we established that 10 mM of DTT and 10

mM $MgSO_4$ are the optimal condition to attain the maximum specific activity which corresponds to

12.2 ± 2.2 μmol(NADPH) min^{-1} mg(SBPase)$^{-1}$ (figure 2A and 2B). Unexpectedly, DTT concentrations

above 10 mM had an inhibitory effect, causing a drastic drop in activity that probably resulted from

structural effects leading to denaturation or aggregation of the enzyme (figure 2A). In contrast, we

observed no such inhibitory effect in the presence of magnesium at concentrations above 10 mM

(figure 2B). Under physiological conditions, the light-dependent redox regulation of plant SBPase is

specifically exerted by f-class thioredoxins (Gutle, Roret et al. 2016), which are reduced by the

ferredoxin-thioredoxin cascade (Schürmann and Buchanan 1975). The chloroplast TRX-f2 from

Chlamydomonas was thus tested and validated for its capacity to activate CrSBPase by reduction

(figure 2C). The maximum activity, corresponding to 12.2 ± 2.2 µmol(NADPH) min-1 mg(SBPase)-1,

was obtained by pre-incubating the enzyme with 1 µM CrTRX-f2 supplemented with 1 mM DTT and

10 mM $MgSO_4$, and maintained even at higher CrTRX-f2 concentrations (*i.e.*, 5 and 10 µM) (figure

2C).

Redox controlled dynamics of *Cr*SBPase

*Cr*SBPase crystallized under reducing treatment (10 mmol.L^{-1} TCEP). Among the eight subunits in the

asymmetric unit, C115-C120 disulfide is absent in two chains (subunits C coloured in magenta and E

coloured in salmon, figure 3D and 3F), while the other six subunits still have the C115-C120 disulfide

bridge (figure 3B,C,E,G,H,I). All other cysteines are unaffected by the reducing treatment. These six

subunits present mildly variable conformations of the $A_{113}SCAGTAC_{120}$ lasso conformation similar to

the untreated oxidized structure of *Cr*SBPase (figure 3A). Reduction of the C115-C120 disulfide

increases local disorder (supplementary figure 3), and the main chain electron density became

uninterpretable over a few residues for subunits C (residues 115-118, 124-127), and E (residues 119-

127).

Molecular dynamics (MD) simulations of the reduced structure were performed by starting from a

dimer of the crystallized oxidized form and forcing residues C115 and C120 to be reduced in a thiol

state (-SH). We then ran two independent replicas of 20 µs of MD. In addition, we also ran two

independent replicas of 2 µs of MD in the oxidized form. In one of the replicas of the reduced form

(MD1), the protein structure barely changed as observed in the SG(C115)/SG(C120) distance that

plateaued at ~7 Å, as well as in the RMSD with respect to the starting conformation that stayed

around 2 Å (see supplementary figures 7-8 for plots and 12 for images of the structures). However, in

the other replica of the reduced form (MD2), significant changes of conformation appeared: the

SG(C115)/SG(C120) distance ended up at ~10 Å with peaks at 20 Å and the RMSD ended up at 3.7

and 2.8 Å (supplementary figures 7-8). The difference between MD1 and MD2 is stronger when one

focuses on the RMSD of the mobile motif (residues 112 to 131, supplementary figure 9). We

observed that after 17 µs of MD all chains adopted an equilibrated conformation and then fluctuated

around that conformation (supplementary figure 9); representative structures extracted from MD

were thus obtained from a clustering on the time frame (or block) 17-20 µs.

The two chains A and B from MD2 adopt a different final conformation (see figure 4A for the overlap

with the crystallized oxidized structure and figure 4B for the overlap with the crystallized reduced

structure), and we cannot directly conclude on which one is the most stable. Moreover, the residues

115 to 128 of each chain from MD1/2 do not overlap well with the crystallized reduced structure

which means that neither of the two chains conformations from MD simulations has converged

towards the crystallized state. However, we can observe in the reduced crystallized structure that

residues 134-148 form an α-helix which is also present in chain B from MD2 whereas this structure is

not kept in chain A from MD2 (see green chain on the right of figures 4A-B). In addition, mobile motif

from chain B seems to open towards the solvent in a similar way with what was observed for FBPase

in the reduced state. Thus, we conclude that chain B from MD2 is more representative of the true

reduced conformation than chain A from MD2. If we compare the mobility of chain B from MD2 in

the reduced state with one chain in the oxidized state (also chain B from MD2), we can see that the

mobile motif is overall more flexible in the reduced state, which is especially true for residues around

C120 (residues 118 to 123, supplementary figure 10). We then analyzed the RMSD of residues 112 to

131 from chain B during MD2 (supplementary figure 11): residues 112, 130 and 131 barely move

which is expected since they belong or are directly connected to the two α-helixes flanking the

mobile motif. At ~6 µs of the trajectory almost all other residues (with the exception of C120) display

an abrupt change of conformation, with a different magnitude though: starting from ~2 Å, the RMSD

jumped to 3 Å for some residues and to 15 Å for other residues (supplementary figures 8, 9, and 11).

At ~13.5 µs, another change occurs which involves C120 but not all the other residues. Thus, we

conclude that the opening of the motif happens in a concerted way and not sequentially residue per

residue.

Oligomeric states of *Cr*SBPase

Size-exclusion profile of the purified native protein is polydisperse with at least three distinct species

of *Cr*SBPase (supplementary figure 1A,C). Notably, we observed that the protein elutes as a main

peak of 79 kDa, close to the mass of a homodimer (theoretical mass from monomer sequence:

36,176.7 Da). This peak is broadened by a later-eluting shoulder, with the apparent molecular mass

of the tailing species at 48 kDa suggesting the presence of a monomeric state of *Cr*SBPase. The third

species elutes at an apparent molecular mass of 145 kDa, that fits with a homotetramer. According

to the absorbance of each peak and considering that they represent all states of purified and oxidized

*Cr*SBPase in solution, we estimate a repartition of species of 10:53:37 for the

tetramer:dimer:monomer mixture (total = 100). Re-injection of the dimer peak fraction over size-

exclusion chromatography yielded a similar mixed profile, supporting a dynamic equilibrium over the

time of the experiment (>2 h) (data not shown).

*Cr*SBPase was crystallized in space groups $P2_12_12$ or $P12_11$ with respectively 6 and 8 protomers in the

asymmetric unit. Proteins Interfaces Structures and Assemblies (PISA) analysis reveals that

protomers of both crystals pack as homodimers with extensive subunit interface of 4 200 Å² in a

similar manner (figure 4A) (Krissinel and Henrick 2007). In the asymmetric unit of CrSBPase

crystallized in reducing conditions (7zuv) or within crystallographic neighbouring units of CrSBPase in

oxidized state (7b2o), *Cr*SBPase further packs into a homotetramer, a dimer of dimers (figure 4B).

The apparent molecular mass of SBPase extracted from *Chlamydomonas* cultures cultivated in TAP

medium under 40 µE.m^{-2}.s^{-1} illumination is comprised between 130 and 40 kDa with the homodimer

representing the predominant state as supported by immunoblot analysis (supplementary figure 4).

To further investigate the oligomeric dynamics of CrSBPase, we performed small-angle X-ray

scattering (SAXS) analysis of untreated or reduced enzyme *in vitro* which reveals an increase in the

protein radius of gyration from 39 to 62 Å upon reduction (supplementary figure 5). This observation

implies a change in the oligomeric state of CrSBPase correlating with the redox state. Besides, after a

week at 4 °C, the purified protein partially precipitates into an amorphous aggregate that treatment

with 10 mmol.L^{-1} TCEP turns into liquid-liquid phase separations (supplementary figure 6). Since the

LBH redox module is positioned at the homodimer interface (figure 5C), we postulated that the

reduction of the C115-C120 disulfide bond and the subsequent local conformation changes depicted

by MD affect the oligomer equilibrium of the enzyme. As a matter of fact, we observed that protein

mutants C115S or C120S, which are unable to form the C115-C120 disulfide bond, elute in a different

oligomer ratio than wild-type CrSBPase (supplementary figure 1C). Indeed, we observed a relative

accumulation of monomer and tetramer species at the expense of the dimer state in both mutants.

This suggests that C115-C120 is involved in the stabilization of the dimer. Meanwhile, the activity of

the two Cys mutants still responded to DTT/Mg-dependent activation resembling the response of the

wild-type protein. No activity was detected for native and untreated forms, while specific activities of

C115S and C120S mutants upon treatment with DTT and MgSO$_4$ were respectively 96% and 107%

with respect to wild-type maximum activity (13.3 ± 2.5 and 14,9 ± 0,9 µmol(NADPH) min^{-1}

mg(SBPase)$^{-1}$ for C115S and C120S mutants, respectively). Based on these data, we can thus conclude

that C115-C120 disulfide bond is not the sole thiol-dependent regulatory mechanism involved in the

modulation of CrSBPase activity and that the redox dependency of CrSBPase catalysis likely depends

on other cysteine pairs. Future studies are required to shed light on the identity of cysteine residues

that contribute to CrSBPase catalytic modulation other than C115 and C120.

Conclusions and perspectives

We report the first crystal structure of an SBPase from the microalga *Chlamydomonas reinhardtii*, a

model for the molecular and cell biology of plants. SBPase is a photosynthetic enzyme involved in the

regenerative phase of the CBBC cycle proved to be the target of multiple regulations that we map

onto our structural model. The native folding of CrSBPase proved to be highly similar to that of

SBPase from the moss *Physcomitrium patens* and to FBPase from chloroplast or cytoplasmic origins.

Notwithstanding its propensity to form crystals, recombinant CrSBPase encompasses a range of

oligomeric states, dominated by a homodimeric form susceptible to exchange for monomers,

tetramers and higher-order assemblies. This oligomer equilibrium is dynamic and correlates with the

presence of a disulfide bridge at the vicinity of the monomer-monomer interface, in the SBPase-

specific LBH motif. We postulate that the reversible reduction/oxidation of the C115-C120 pair

modulates the oligomerization of CrSBPase, through the induction of a local disorder that we

observed in the crystal structure of the protein under reducing conditions and that correlates with

computations of molecular dynamics.

How this oligomeric exchange impacts the actual activity of the enzyme is an open question but

should be aligned with previous reports on the allosteric character of the structurally similar FBPase

enzyme which follows dimer/tetramer exchanges (Barciszewski, Wisniewski et al. 2016) with a Hill

cooperativity coefficient close to 2 representing positive cooperativity in the context of a homodimer

(Giudici-Orticoni, Buc et al. 1990). Lastly, systematic proteome mapping in the chloroplast of

Chlamydomonas localized SBPase in a region surrounding the Rubisco pyrenoid nearby five other

CBBC enzymes: phosphoglycerate kinase 1 (PGK), glyceraldehyde-3-phosphate dehydrogenases

(GAP1, GAP3), fructose-bisphosphate aldolase (FBA3), ribulose-5-phosphate 3-epimerase (RPE1), and

phosphoribulokinase (PRK) (Wang, Patena et al. 2022). Based on our structure-function study on

recombinant SBPase, we can hypothesize that upon reductive activation SBPase is co-addressed to

this liquid partition of the stroma where it contributes to CBBC reactions at optimized rates.

Tables.

Table 1. Crystallographic data collections and models building statistics.

	7b2o (oxidized)	7zuv (partially reduced)
Wavelength (Å)	0.9801	0.9801
Resolution range (Å)	46.62 - 3.095 (3.206 - 3.095)	48.54 - 3.11 (3.221 - 3.11)
Space group	P 2_1 2_1 2	P 1 2_1 1
Unit cell (Å, °)	178.224 183.652 75.196 90 90 90	53.774 163.462 172.765 90 91.939 90
Total reflections	615155 (55989)	376091 (32376)
Unique reflections	45740 (4245)	53456 (5213)
Multiplicity	13.4 (12.9)	7.0 (6.2)
Completeness (%)	99.34 (94.20)	99.65 (97.17)
Mean I/sigma(I)	12.02 (1.72)	8.54 (1.30)
Wilson B-factor (Å²)	64.82	68.45
R-merge	0.495 (1.877)	0.2492 (1.502)
R-meas	0.5144 (1.954)	0.2691 (1.64)
R-pim	0.1386 (0.5343)	0.1009 (0.6477)
CC1/2	0.983 (0.623)	0.988 (0.479)
CC*	0.996 (0.876)	0.997 (0.805)
Reflections used in refinement	45607 (4242)	53417 (5193)
Reflections used for R-free	1986 (183)	1983 (191)
R-work	0.1942 (0.3071)	0.1963 (0.3239)
R-free	0.2390 (0.3482)	0.2328 (0.3740)
CC(work)	0.944 (0.788)	0.956 (0.726)
CC(free)	0.912 (0.649)	0.949 (0.596)
Number of non-hydrogen atoms	14194	18714
Macromolecules	14190	18704
Solvent	4	10
Protein residues	1859	2451
RMS(bonds) (Å)	0.005	0.004
RMS(angles) (°)	0.75	0.73
Ramachandran favored (%)	94.26	94.80
Ramachandran allowed (%)	5.52	4.70
Ramachandran outliers (%)	0.22	0.50
Rotamer outliers (%)	0.13	0.55
Clashscore	4.88	6.60
Average B-factor (Å²)	67.00	68.41
macromolecules (Å²)	67.01	68.40
solvent (Å²)	49.59	94.37

435 Statistics for the highest-resolution shell are shown in parentheses.

Figures legends

Figure 1. Crystal structure of CrSBPase.

(**A**) Topology is displayed as cartoon with main chain coloured from blue (amino-terminus) to red (carboxy-terminus). (**B**) Rotated view of A. by x-axis over -90°. (**C**) Putative active site residues inferred from alignment with FBPase bound to FBP (5l0a, human muscle fructose-1,6-bisphosphatase E69Q mutant in active R-state in complex with fructose-1,6-bisphosphate) are represented in sticks. (**D**) Surface representation in the B. orientation with putative active site residues coloured in cyan. Water molecule H_2O 401 oxygen is represented as a red sphere. (**E**) Cysteines site chains are represented in spheres. (**F**) Threonines and serines side chains reported to be the target of phosphorylations are represented in spheres.

Figure 2. Functional characterization of CrSBPase *in vitro*.

Reported enzymatic activity of CrSBPase assayed for (**A**) reduction by dithiothreitol (DTT), (**B**) Magnesium sulfate ($MgSO_4$), and (**C**) recombinant thioredoxin f2 from *Chlamydomonas reinhardtii* (*Cr*TRX-f2).

Figure 3. Crystallographic structures of *Cr*SBPase under reducing treatment.

(**A-I**) Aligned structures of *Cr*SBPase protomers without redox treatment (**A**, 7b2o chain A) or in the presence of 10 mmol.L^{-1} TCEP reducing agent (**B-I**, 7zuv chains A-H).

Figure 4. Molecular dynamics simulation of *Cr*SBPase after reduction.

(**A**) Overlap of the crystallographic structure of oxidized SBPase and representative structures of equilibrated reduced SBPase during molecular dynamics simulation 2 (MD2). For structures extracted from MD, only residues 109 to 148 are displayed since most of the other residues are closely overlapping those of the crystallographic structure. (**B**) Overlap of the crystallographic structure of

reduced SBPase and representative structures of equilibrated reduced SBPase during MD2. For

structures extracted from MD, only residues 109 to 148 are displayed.

Figure 5. Oligomeric structure of CrSBPase.

(**A**) Asymmetric unit dimer of untreated *Cr*SBPase (7b2o). Chains are represented in cartoon and

coloured cyan (chain B) and salmon (chain E). (**B**) Asymmetric unit homotetramer under reducing

treatment (7zuv). Chains A (cyan), C (magenta), F (white), and H (orange) belong to the same

asymmetric unit. (**C**) Close-up view on figure A homodimer interface. Loop 113-ASCAGTAC-120 from

chain B (in cyan) is in 5-Å distance of neighbouring chain E (in salmon). C115 and C120 are bonded by

a disulfide bridge. (**D**) Surface representation of 7b2o homodimer, coloured according to

conservation score from teal (lowest conservation) to purple (highest conservation) among 150

aligned homologs.

Supplementary information

Supplementary table 1. Primers used for point mutagenesis.

5'-forward C115S	CCGCACCGCCTCGAGCGCCGGTACCGCCTGCGTG
3'-reverse C115S	CACGCAGGCGGTACCGGCGCTCGAGGCGGTGCGG
5'-forward C120S	GCGCCGGTACCGCCAGCGTGAACAGCTTCGGCG
3'-reverse C120S	CGCCGAAGCTGTTCACGCTGGCGGTACCGGCGC

Supplementary figure 1. Purification of recombinant *Cr*SBPase.

A. Size-exclusion chromatography profile recorded at absorbance λ= 280 nm on Superdex 200 26/600

GL column. B. Electrophoresis of affinity-chromatography fractions on 12% acrylamide gel in

denaturing and reducing condition. Gel was stained with Coomassie blue. Mw: molecular mass

standards ladder. C. Size-exclusion chromatograms of *Cr*SBPase wild-type (blue), mutant C115S

(orange), mutant C120S (gray).

Supplementary figure 2. SBPase multiple sequences alignment.

*Cr*SBPase mature sequence was used for a BLAST search. Retrieved homolog sequences were aligned

and coloured-coded according to residue conservation. Illustration made with EndScript (Robert and

Gouet 2014).

Supplementary figure 3. Related to figure 3. Local disorder of the $A_{113}SCAGTAC_{120}$ loop.

Main chain was traced according to crystallographic b-factor, with large orange sections representing

high b-factor values and thin blue sections representing low b-factors. A-I. Aligned structures of

*Cr*SBPase protomers without redox treatment (A, 7b2o chain A) or in the presence of 10 mmol.L^{-1}

TCEP reducing agent (B-I, 7zuv chains A-H).

Supplementary figure 4. Size-exclusion fractionation of *Chlamydomonas* cell extracts.

A. *Chlamydomonas* cell culture was harvested, lysed and the soluble fraction of the lysate was loaded

on Superose6 16/600 size-exclusion column. Chromatography fractions were analyzed by western

blot with anti-*Cr*SBPase primary antibodies. First membrane was loaded with fractions eluted from

40 to 80 mL. Second membrane was loaded with fractions eluted from 80 to 120 mL. M lane is

loaded with molecular mass standards ladder. Recombinant *Cr*SBPase was loaded on last lane.

Supplementary figure 5. Size-exclusion chromatography coupled to small angle X-rays scattering

(SEC-SAXS) of *Cr*SBPase.

A. X-ray scattering curve log(I) = f(s) for pure *Cr*SBPase untreated (dark blue) or treated (light blue)

with 10 mmol.L^{-1} dithiothreitol (DTT). B. Radius of gyration of untreated protein was computed by

ATSAS Primus Guinier Wizard as Rg = 39.14 ± 0.06 Å (Franke, Petoukhov et al. 2017). C. Radius of

gyration of DTT-treated protein was computed as Rg = 62.47 ± 0.38 Å.

Supplementary figure 6. Phase separation of reduced *Cr*SBPase precipitates.

*Cr*SBPase at 2 mg/mL formed amorphous precipitates after storage at 4 °C for a week. The

suspension was sedimented by centrifugation at 20,000 *g* for 20 minutes and a microliter drop of the

supernatant or of the pellet were treated with 10 mmol.L^{-1} reducing agent TCEP and visualized by

binocular optical microscope. A. Untreated precipitate. B. TCEP-treated precipitate. C. Untreated

supernatant. D. TCEP-treated supernatant.

Supplementary figures 7-11. Computational detail of molecular dynamic simulation.

Supplementary figure 7.

SG(Cys115)/SG(Cys120) distances along the MD trajectories for each chain (left: oxidized state, right:

reduced state). Light colours: raw data saved every 100 ps, dark colours: moving average on 20ns

blocs.

Supplementary figure 8.

RMSD of main chain along the MD trajectories for each chain (left: oxidized state, right: reduced

state). Light colours: raw data saved every 100 ps, dark colours: moving average on 20 ns blocs.

Supplementary figure 9.

RMSD of main chain of the mobile motif in the reduced state (residues 112 to 131) along the MD

trajectories for each chain (left: RMSD with respect to the starting conformation of MD, right: RMSD

with respect to the conformation at 20 µs). Light colours: raw data saved every 100 ps, dark colours:

moving average on 20 ns blocs.

Supplementary figure 10.

RMSF per residue for each chain (top row: full sequence, bottom row: zoom on residues 110 to 170;

left column: oxidized state, middle column: reduced state, right column: comparison of one chain

from each state).

Supplementary figure 11.

RMSD of main chain of each residue from the mobile motif along the trajectory of chain B from MD2.

Data are moving average on 200 ns blocs.

Supplementary figure 12. Structures of CrSBPase retrieved from molecular dynamics simulations 1

and 2.

Two independent replicas of 20 µs were performed. In one of them (MD1), the overall structure of

both chains of the enzyme barely changed. To illustrate this, we present in supplementary figure 12

the overlap between the crystallographic oxidized structure and representative structures extracted

from MD (based on clustering the last 3 µs since each chain does not display significative changes

after 17 µs). A. Overlap of the crystallographic structure of oxidized CrSBPase and representative

structures of equilibrated reduced SBPase during MD1. For structures extracted from MD, only

residues 111 to 132 are displayed since the other residues are closely overlapping those of the

crystallographic structure. B. Overlap of the crystallographic structure of oxidized SBPase and

representative structures of equilibrated reduced CrSBPase during MD2. For structures extracted

from MD, only residues 109 to 148 are displayed since most of the other residues are closely

overlapping those of the crystallographic structure.

Authors contributions

JH, NC: Conceptualization; JH, NC, MS, TLM: Data curation; JH, NC, MZ: Formal analysis; JH: Funding

acquisition; JH, NC, MZ, SDL, MS, TLM: Investigation; JH, NC, MZ, MS, TLM: Methodology; JH, SDL:

Project administration; JH, NC, SDL: Resources; NC: Software; JH, MZ: Supervision; JH, NC, MZ, SDL,

MS, TLM: Writing - original draft; JH, NC, MZ, SDL, MS, TLM: Writing - review & editing.

Acknowledgements

We acknowledge the Institut de Biologie Physico-Chimique (CNRS, FR 505) for access to the

crystallization facility. We acknowledge SOLEIL for provision of synchrotron radiation facilities at

beamlines SWING, PROXIMA-1, and PROXIMA-2A. We thank Guillaume Q. Robert for his contribution

to protein purification, crystallization under reducing treatment, and model building. Plasmids pSL-

175, -181, and -182 were cloned by Dr. Laure Michelet. This work was funded by the grant

CALVINTERACT from the Agence Nationale de la Recherche (ANR-19-CE11-0009). Martina Santoni

received funding from the University of Bologna (Bologna, Italy) and Sorbonne University (Paris,

France) exchange program ERASMUS+. This work was performed using HPC resources from

GENCI–IDRIS (Grant 2021-077156). The authors express their gratitude to the financial support of the

AAP AIN INSB CNRS.

Conflicts of interest statement

The authors report no conflict of interest relevant to the conduct or report of this research.

Bibliography

Adams, P. D., P. V. Afonine, G. Bunkóczi, V. B. Chen, N. Echols, J. J. Headd, L. W. Hung, S. Jain, G. J.
Kapral, R. W. Grosse Kunstleve, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, R. J. Read, D. C.
Richardson, J. S. Richardson, T. C. Terwilliger and P. H. Zwart (2011). "The Phenix software for
automated determination of macromolecular structures." Methods **55**(1): 94-106.
Ahmad, S., M. Gromiha, H. Fawareh and A. Sarai (2004). "ASAView: database and tool for solvent
accessibility representation in proteins." BMC Bioinformatics **5**: 51.
Almagro Armenteros, J. J., M. Salvatore, O. Emanuelsson, O. Winther, G. von Heijne, A. Elofsson and
H. Nielsen (2019). "Detecting sequence signals in targeting peptides using deep learning." Life Sci
Alliance **2**(5).
Anderson, L. E. (1974). "Activation of pea leaf chloroplast sedoheptulose 1,7-diphosphate
phosphatase by light and dithiothreitol." Biochem Biophys Res Commun **59**(3): 907-913.
Barciszewski, J., J. Wisniewski, R. Kolodziejczyk, M. Jaskolski, D. Rakus and A. Dzugaj (2016). "T-to-R
switch of muscle fructose-1,6-bisphosphatase involves fundamental changes of secondary and
quaternary structure." Acta Crystallogr D Struct Biol **72**(Pt 4): 536-550.
Cadet, F. and J. C. Meunier (1988). "pH and kinetic studies of chloroplast sedoheptulose-1,7-
bisphosphatase from spinach (Spinacia oleracea)." Biochem J **253**(1): 249-254.
Calvin, M. (1962). "The path of carbon in photosynthesis." Science **135**(3507): 879-889.
Chiadmi, M., A. Navaza, M. Miginiac-Maslow, J. P. Jacquot and J. Cherfils (1999). "Redox signalling in
the chloroplast: structure of oxidized pea fructose-1,6-bisphosphate phosphatase." EMBO J **18**(23):
6809-6815.
Emanuelsson, O., S. Brunak, G. von Heijne and H. Nielsen (2007). "Locating proteins in the cell using
TargetP, SignalP and related tools." Nat Protoc **2**(4): 953-971.
Emsley, P., B. Lohkamp, W. G. Scott and K. Cowtan (2010). "Features and development of Coot." Acta
Crystallogr D Biol Crystallogr **66**(Pt 4): 486-501.
Feng, L., Y. Han, G. Liu, B. An, J. Yang, G. Yang, Y. Li and Y. Zhu (2007). "Overexpression of
sedoheptulose-1,7-bisphosphatase enhances photosynthesis and growth under salt stress in
transgenic rice plants." Funct Plant Biol **34**(9): 822-834.
Feng, L., K. Wang, Y. Li, Y. Tan, J. Kong, H. Li, Y. Li and Y. Zhu (2007). "Overexpression of SBPase
enhances photosynthesis against high temperature stress in transgenic rice plants." Plant Cell Rep
26(9): 1635-1646.
Franke, D., M. V. Petoukhov, P. V. Konarev, A. Panjkovich, A. Tuukkanen, H. D. T. Mertens, A. G.
Kikhney, N. R. Hajizadeh, J. M. Franklin, C. M. Jeffries and D. I. Svergun (2017). "ATSAS 2.8: a
comprehensive data analysis suite for small-angle scattering from macromolecular solutions." J Appl
Crystallogr **50**(Pt 4): 1212-1225.
Giudici-Orticoni, M. T., J. Buc, M. Bidaud and J. Ricard (1990). "Thermodynamics of information
transfer between subunits in oligomeric enzymes and kinetic cooperativity. 3. Information transfer
between the subunits of chloroplast fructose bisphosphatase." Eur J Biochem **194**(2): 483-490.
Gurrieri, L., A. Del Giudice, N. Demitri, G. Falini, N. V. Pavel, M. Zaffagnini, M. Polentarutti, P. Crozet,
C. H. Marchand, J. Henri, P. Trost, S. D. Lemaire, F. Sparla and S. Fermani (2019). "Arabidopsis and
Chlamydomonas phosphoribulokinase crystal structures complete the redox structural proteome of
the Calvin-Benson cycle." Proc Natl Acad Sci U S A **116**(16): 8048-8053.
Gutle, D. D., T. Roret, S. J. Muller, J. Couturier, S. D. Lemaire, A. Hecker, T. Dhalleine, B. B. Buchanan,
R. Reski, O. Einsle and J. P. Jacquot (2016). "Chloroplast FBPase and SBPase are thioredoxin-linked
enzymes with similar architecture but different evolutionary histories." Proc Natl Acad Sci U S A
113(24): 6779-6784.
Hammel, A., F. Sommer, D. Zimmer, M. Stitt, T. Mühlhaus and M. Schroda (2020). "Overexpression of
Sedoheptulose-1,7-Bisphosphatase Enhances Photosynthesis in Chlamydomonas reinhardtii and Has
No Effect on the Abundance of Other Calvin-Benson Cycle Enzymes." Front Plant Sci **11**: 868.

Jacquot, J. P., J. Lopez-Jaramillo, M. Miginiac-Maslow, S. Lemaire, J. Cherfils, A. Chueca and J. Lopez-
Gorge (1997). "Cysteine-153 is required for redox regulation of pea chloroplast fructose-1,6-
bisphosphatase." FEBS Lett **401**(2-3): 143-147.

Johnson, M. P. (2016). "Photosynthesis." Essays Biochem **60**(3): 255-273.

Johnson, X. and J. Alric (2013). "Central carbon metabolism and electron transport in
Chlamydomonas reinhardtii: metabolic constraints for carbon partitioning between oil and starch."
Eukaryot Cell **12**(6): 776-793.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates,
A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-
Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M.
Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K.
Kavukcuoglu, P. Kohli and D. Hassabis (2021). "Highly accurate protein structure prediction with
AlphaFold." Nature **596**(7873): 583-589.

Krissinel, E. and K. Henrick (2007). "Inference of macromolecular assemblies from crystalline state." J
Mol Biol **372**(3): 774-797.

Kubis, A. and A. Bar-Even (2019). "Synthetic biology approaches for improving photosynthesis." J Exp
Bot **70**(5): 1425-1433.

Lee, S. W. and T. R. Hahn (2003). "Light-regulated differential expression of pea chloroplast and
cytosolic fructose-1,6-bisphosphatases." Plant Cell Rep **21**(6): 611-618.

Lefebvre, S., T. Lawson, O. V. Zakhleniuk, J. C. Lloyd, C. A. Raines and M. Fryer (2005). "Increased
sedoheptulose-1,7-bisphosphatase activity in transgenic tobacco plants stimulates photosynthesis
and growth from an early stage in development." Plant Physiol **138**(1): 451-460.

Lemaire, S. D., B. Guillon, P. Le Maréchal, E. Keryer, M. Miginiac-Maslow and P. Decottignies (2004).
"New thioredoxin targets in the unicellular photosynthetic eukaryote *Chlamydomonas reinhardtii*."
Proc Natl Acad Sci U S A **101**(19): 7475-7480.

Lemaire, S. D., D. Tedesco, P. Crozet, L. Michelet, S. Fermani, M. Zaffagnini and J. Henri (2018).
"Crystal Structure of Chloroplastic Thioredoxin f2 from Chlamydomonas reinhardtii Reveals Distinct
Surface Properties." Antioxidants (Basel) **7**(12).

Liebschner, D., P. V. Afonine, M. L. Baker, G. Bunkóczi, V. B. Chen, T. I. Croll, B. Hintze, L. W. Hung, S.
Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R. J. Read, J. S. Richardson,
D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H. Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L.
L. Videau, C. J. Williams and P. D. Adams (2019). "Macromolecular structure determination using X-
rays, neutrons and electrons: recent developments in Phenix." Acta Crystallogr D Struct Biol **75**(Pt
10): 861-877.

McConnell, E. W., E. G. Werth and L. M. Hicks (2018). "The phosphorylated redox proteome of
Chlamydomonas reinhardtii: Revealing novel means for regulation of protein structure and function."
Redox Biol **17**: 35-46.

Mettler, T., T. Muhlhaus, D. Hemme, M. A. Schottler, J. Rupprecht, A. Idoine, D. Veyel, S. K. Pal, L.
Yaneva-Roder, F. V. Winck, F. Sommer, D. Vosloh, B. Seiwert, A. Erban, A. Burgos, S. Arvidsson, S.
Schonfelder, A. Arnold, M. Gunther, U. Krause, M. Lohse, J. Kopka, Z. Nikoloski, B. Mueller-Roeber, L.
Willmitzer, R. Bock, M. Schroda and M. Stitt (2014). "Systems Analysis of the Response of
Photosynthesis, Metabolism, and Growth to an Increase in Irradiance in the Photosynthetic Model
Organism Chlamydomonas reinhardtii." Plant Cell **26**(6): 2310-2350.

Moorhead, G. B., R. J. Hodgson and W. C. Plaxton (1994). "Copurification of cytosolic fructose-1,6-
bisphosphatase and cytosolic aldolase from endosperm of germinating castor oil seeds." Arch
Biochem Biophys **312**(2): 326-335.

Morisse, S., M. Zaffagnini, X. H. Gao, S. D. Lemaire and C. H. Marchand (2014). "Insight into protein S-
nitrosylation in Chlamydomonas reinhardtii." Antioxid Redox Signal **21**(9): 1271-1284.

Nishizawa, A. N. and B. B. Buchanan (1981). "Enzyme regulation in C4 photosynthesis. Purification
and properties of thioredoxin-linked fructose bisphosphatase and sedoheptulose bisphosphatase
from corn leaves." J Biol Chem **256**(12): 6119-6126.

Ohta, J. (2022). "A novel variant of the Calvin-Benson cycle bypassing fructose bisphosphate." <u>Sci Rep</u>
12(1): 3984.
Pérez-Pérez, M. E., A. Mauriès, A. Maes, N. J. Tourasse, M. Hamon, S. D. Lemaire and C. H. Marchand
(2017). "The Deep Thioredoxome in Chlamydomonas reinhardtii: New Insights into Redox
Regulation." <u>Mol Plant</u> **10**(8): 1107-1125.
Robert, X. and P. Gouet (2014). "Deciphering key features in protein structures with the new
ENDscript server." <u>Nucleic Acids Research</u> **42**(W1): W320-W324.
Rosenthal, D. M., A. M. Locke, M. Khozaei, C. A. Raines, S. P. Long and D. R. Ort (2011). "Over-
expressing the C(3) photosynthesis cycle enzyme Sedoheptulose-1-7 Bisphosphatase improves
photosynthetic carbon gain and yield under fully open air CO(2) fumigation (FACE)." <u>BMC Plant Biol</u>
11: 123.
Schmollinger, S., T. Mühlhaus, N. R. Boyle, I. K. Blaby, D. Casero, T. Mettler, J. L. Moseley, J. Kropat, F.
Sommer, D. Strenkert, D. Hemme, M. Pellegrini, A. R. Grossman, M. Stitt, M. Schroda and S. S.
Merchant (2014). "Nitrogen-Sparing Mechanisms in Chlamydomonas Affect the Transcriptome, the
Proteome, and Photosynthetic Metabolism." <u>Plant Cell</u> **26**(4): 1410-1435.
Schürmann, P. and B. B. Buchanan (1975). "Role of ferredoxin in the activation of sedoheptulose
diphosphatase in isolated chloroplasts." <u>Biochim Biophys Acta</u> **376**(1): 189-192.
Simkin, A. J., P. E. Lopez-Calcagno, P. A. Davey, L. R. Headland, T. Lawson, S. Timm, H. Bauwe and C.
A. Raines (2017). "Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-
bisphophate aldolase and the photorespiratory glycine decarboxylase-H protein increases CO(2)
assimilation, vegetative biomass and seed yield in Arabidopsis." <u>Plant Biotechnol J</u> **15**(7): 805-816.
Wagner, V., G. Gessner, I. Heiland, M. Kaminski, S. Hawat, K. Scheffler and M. Mittag (2006).
"Analysis of the phosphoproteome of Chlamydomonas reinhardtii provides new insights into various
cellular pathways." <u>Eukaryot Cell</u> **5**(3): 457-468.
Wang, L., W. Patena, K. A. Van Baalen, Y. Xie, E. R. Singer, S. Gavrilenko, M. Warren-Williams, L. Han,
H. R. Harrigan, V. Chen, V. T. N. P. Ton, S. Kyin, H. H. Shwe, M. H. Cahn, A. T. Wilson, J. Hu, D. J.
Schnell, C. D. McWhite and M. Jonikas (2022). "A Chloroplast Protein Atlas Reveals Novel Structures
and Spatial Organization of Biosynthetic Pathways." <u>bioRxiv</u>: 2022.2005.2031.493820.
Werth, E. G., E. W. McConnell, T. S. Gilbert, I. Couso Lianez, C. A. Perez, C. K. Manley, L. M. Graves, J.
G. Umen and L. M. Hicks (2017). "Probing the global kinome and phosphoproteome in
Chlamydomonas reinhardtii via sequential enrichment and quantitative proteomics." <u>Plant J</u> **89**(2):
416-426.
Wilson, R. H., M. Hayer-Hartl and A. Bracher (2019). "Crystal structure of phosphoribulokinase from
Synechococcus sp. strain PCC 6301." <u>Acta Crystallogr F Struct Biol Commun</u> **75**(Pt 4): 278-289.
Woodrow, I. E. (1982). "Sedoheptulose-1,7-bisphosphatase from wheat chloroplasts." <u>Methods</u>
<u>Enzymol</u> **90 Pt E**: 392-396.
Woodrow, I. E. and D. A. Walker (1980). "Light-mediated activation of stromal sedoheptulose
bisphosphatase." <u>Biochem J</u> **191**(3): 845-849.
Yoshida, K. and T. Hisabori (2018). "Determining the Rate-Limiting Step for Light-Responsive Redox
Regulation in Chloroplasts." <u>Antioxidants (Basel)</u> **7**(11).
Zaffagnini, M., M. Bedhomme, C. H. Marchand, S. Morisse, P. Trost and S. D. Lemaire (2012). "Redox
regulation in photosynthetic organisms: focus on glutathionylation." <u>Antioxid Redox Signal</u> **16**(6):
567-586.
Zimmer, D., C. Swart, A. Graf, S. Arrivault, M. Tillich, S. Proost, Z. Nikoloski, M. Stitt, R. Bock, T.
Mühlhaus and A. Boulouis (2021). "Topology of the redox network during induction of
photosynthesis as revealed by time-resolved proteomics in tobacco." <u>Sci Adv</u> **7**(51): eabi8307.
725

Figure 1.

Figure 2.

A

B

C

Figure 3.

Figure 4.

A

Oxidized structure (7B2O)
MD2, Chain A
MD2, Chain B

B

Reduced structure
MD2, Chain A
MD2, Chain B

Figure 5.