The antibacterial T6SS of *Agrobacterium tumefacien* promotes disease occurrence

and influences crown gall microbiota composition

Si-Chong Wang[1,2], Ai-Ping Chen[1], Shu-Jen Chou[1], Chih-Horng Kuo[1*], Erh-Min Lai[1*]

[1] Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan,

[2] Department of Life Science, National Central University, Taoyuan, Taiwan

Running title: T6SS on tumorigenesis and gallobiome

*Corresponding authors

Erh-Min Lai: emlai@gate.sinica.edu.tw

Chih-Horng Kuo: chkuo@gate.sinica.edu.tw

Institute of Plant and Microbial Biology, Academia Sinica

128, Sec. 2, Academia Road, Nankang, Taipei, Taiwan 11529

ORCID

CHK   https://orcid.org/0000-0002-2857-0529

EML   https://orcid.org/0000-0003-3630-8683

SCW   https://orcid.org/0000-0002-6192-8844

Keywords: *Agrobacterium tumefaciens*, crown gall, type VI secretion system,

microbiota, 16S rRNA gene amplicon sequencing

ABSTRACT

The type VI secretion system (T6SS) is deployed by many proteobacteria to secrete
effector proteins into bacterial competitors for competition or eukaryotic cells for
pathogenesis. *Agrobacterium tumefacien*, a soil-borne phytopathogen causing crown
gall disease on various plant species, deploys its T6SS to attack closely- and distantly-
related bacterial species *in vitro* and *in planta*. Current evidence suggests that the *A.*
tumefaciens T6SS is not essential for pathogenesis under direct inoculation but it
remains unknown whether T6SS influences natural disease incidence and microbial
community of crown galls. To address these two key questions, we established a soil
inoculation method on wounded tomato seedlings that mimics natural infections and
performed 16S rRNA gene amplicon sequencing. By comparing the *A. tumefaciens*
C58 wild-type strain with two T6SS mutants, we demonstrate that the T6SS
influences disease occurrence and microbiota composition within crown galls. Based
on multiple inoculation trials across seasons, all three strains could induce tumors but
the mutants had significantly lower disease incidences. The season of inoculation
played a more important role than the T6SS in shaping the gall microbiota (i.e.,
gallobiome). The influence of T6SS was evident in summer, in which two bacterial
families (Sphingomonadaceae and Burkhoderiaceae) were enriched in the gallobiome
induced by the T6SS mutants. Further *in-vitro* competition and colonization assay
demonstrated the T6SS-mediated antagonism to a tomato isolate *Sphingomonas* sp.
R1. In conclusion, this study informs that the *A. tumefaciens* T6SS may promote
tumorigenesis in natural infection process and provide competitive advantages in gall-
associated microbiota.

IMPORTANCE

T6SS is widespread in the agrobacterial genome and used by *Agrobacterium*
tumefaciens for interbacterial competition. *A. tumefaciens* is a soil inhabitant and
opportunistic bacterial pathogen causing crown gall disease in wide range of plants.
Current evidence indicates that T6SS is not required for gall formation when the
agrobacteria are inoculated directly on plant wounding sites. However, in terms of
disease ecology, agrobacteria may need to compete with other bacteria in bulk soil to
gain access to plant wounds and influence microbial community inside crown galls.
To date, the role of T6SS in agrobacterial pathogenesis and gallobiome under a more
natural setting remains unknown. In this study, we successfully developed a soil
inoculation method coupled with a Blocker-mediated Bacterial 16S rRNA gene
amplicon Enrichment Sequencing (BBRES) to address these two important questions.

We provided evidence that T6SS promotes disease occurrence and influences crown

gall microbiota composition by interbacterial competition.

INTRODUCTION

Many proteobacteria, including pathogens and commensals, deploy the type VI

secretion system (T6SS) for antagonism or pathogenesis (1, 2) . It is a protein

translocation apparatus used to inject effectors into target cells, mainly in a contact-

dependent manner. Based on the destination and biological function of known

effectors (3), T6SS mainly functions as an antibacterial weapon used by bacteria to

inhibit or kill the competing bacterial species, thus impacting their survival or shaping

microbiota in their ecological niche (4).

Previous studies on the microbiome associated with animal guts or plants indicated

that the T6SS genes are enriched in these communities, suggesting T6SS may be

important for niche competition (5-8). Metagenomic analysis of human gut

microbiota revealed a role of T6SS for the domination of the gut

symbiont *Bacteroides fragilis* by targeting other members of microbiome *in vitro* and

in vivo (6, 7). A recent study further showed that a murine pathogen *Citrobacter*

rodentium and resident commensal Enterobacteriaceae share the same strategy by

using T6SS for niche competition in the murine gastrointestinal tract (9). T6SS is also

deployed by plant pathogens to gain competitive growth advantage *in planta* as well

as for beneficial bacteria to prevent or reduce disease symptoms caused by competing

pathogens (10-12). Comparative metagenomic analysis of microbiota between T6SS$^+$

and T6SS$^-$ bacterial strains were also carry out. The gut microbiota of pests infected

by *Pseudomonas protegens,* a plant-beneficial bacterium capable of invading insect

pests, showed that T6SS has no significant impact on microbiota diversity at

phylum/class level but affects the abundance of Enterobacteriaceae (13). These

studies collectively suggest that T6SS is a potent antibacterial weapon used by

invading pathogens or resident bacteria to gain competitive advantage in their

ecological niche. However, the knowledge into the degree of T6SS in shaping

microbiota and the molecular mechanisms of interbacterial competition in complex

microbial community are limited.

Agrobacteria are a group of bacteria capable of inducing crown gall or hairy root

disease on plants by transferring a piece of DNA named T-DNA from bacteria into

plants via the type IV secretion system (T4SS) (14). T6SS is highly conserved in the

Agrobacterium tumefaciens species complex and plays a role in interbacterial

competition (15-17). In *A. tumefaciens,* the T6SS is encoded by a gene cluster

consisting of an *imp* operon encoding components for T6SS and an *hcp* operon

encoding the puncturing device and effectors. Using *A. tumefaciens* reference strain

C58, we previously discovered that it uses Tde DNase effectors as the major

antibacterial weapon to gain competitive growth advantage to its siblings lacking the

cognate immunity protein or other bacterial species by co-infiltration into leaves of

Nicotiana benthamiana (10). Interestingly, agrobacteria with incompatible effector-

immunity (EI) pairs exhibit strong antagonism at the between species level, while

only weak or non-detectable effect at the within species level (15). Moreover, higher

T6SS-mediated killing outcome was observed when nutrient is scarce, as opposed to

nutrient rich condition (18). Thus, genetic and environmental factors beyond EI pairs

also contributes to interbacterial competition.

To date, agrobacterial T6SS has been demonstrated as an antibacterial weapon (15-

17). No evidence for its role in promoting virulence when tumor assays were

conducted in sterile condition or directly inoculated on the stems of various plant

species including tomato plants (19). Considering that agrobacteria may need to

compete with other bacteria in bulk soil or rhizosphere to gain access to plant wounds

for inciting crown galls, we reasoned that T6SS may influence microbial community

and pathogenesis under a more natural setting. To address these questions, we

developed a soil inoculation protocol to mimic natural infection on wounded tomato

seedlings across seasons for evaluating the impact of the *A. tumefaciens* T6SS on

disease occurrence and crown gall microbiota (termed gallobiome). Based on 16S

rRNA amplicon sequencing analysis, followed by *in-vitro* competition and

colonization assays, the results informed that the T6SS may help *A. tumefaciens* to

maintain a stable colonization for competitive advantage on plant surface for effective

infection, leading to higher disease incidence.

RESULTS

T6SS mutants of *A. tumefaciens* C58 induced lower crown gall disease

incidences, which was also affected by environmental factors

Agrobacterium tumefaciens C58 wild type (WT) and two T6SS mutants with deletion

of essential T6SS genes, Δ*tssL* and Δ*tssB* respectively, were used to study the effects

of T6SS in tumorigenesis using soil inoculation method. Nine batches of inoculation

experiments across different seasons were conducted in this study (Table 1). In total,

139 crown galls were collected, including 70 induced by WT, 41 induced by Δ*tssL*,

and 28 induced by Δ*tssB*. The results showed that all three strains are capable of

inducing tumors, but the disease incidences of Δ*tssL* and Δ*tssB* were significantly

lower than WT (Table 1, Fig. 1A). There was no significant weight difference among

the crown galls induced by different strains (Fig. 1B). It is notable that the disease

incidences of those inoculated in June and July were significantly lower than those in

October and November, resulting in an inverse correlation between disease incidence

and temperature across seasons for both WT and T6SS mutants throughout the year

(Fig. 1C, 1D). These results indicated that both the strains and month of inoculation

affected the disease incidence under soil inoculation condition.

Round I of 16S rRNA amplicon sequencing: Initial trial

To determine the gallobiome composition, crown galls with similar weights across

seasons were selected for 16S rRNA gene amplicon sequencing. For Round I, six

crown gall DNA samples, three induced by WT and three by Δ*tssL* were amplified

with two commonly used 16S rRNA gene primer sets V3-4 and V5-7 (Table S1). The

Illumina sequencing produced 140,830 and 601,299 reads from V3-4 and V5-7 sets,

respectively. However, the majority (99.1% in V3-4 and 93.6% for V5-7) of these

reads were derived from plant chloroplast and mitochondria. After removing these

host contaminations, only 1,230 and 38,320 bacterial reads remained (Table 2**).**

Round II of 16S rRNA amplicon sequencing: Optimization

Due to the severe host contamination observed in Round I that used the universal 16S

rRNA gene primers, we evaluated the performance of different 16S rRNA gene

primers with blockers (3' modified oligonucleotides with C3 spacer) that could

prevent the amplification of tomato chloroplast and mitochondrial rRNA genes (20-

22).

Based on the alignment of bacterial, tomato chloroplast and mitochondrial 16S rRNA

gene sequences (Fig. S1), we designed the primer sets and cognate blockers against

different variable region in 16S rRNA genes (Table S1 and S2). The new primer sets

and blockers were tested using the three WT-induced crown gall DNA samples from

Round I. After sequencing and data processing, the result indicated that the blockers

have variable effectiveness in reducing host contaminations (Table 3 and Fig. 2A**).**

For V3-4 and V5-7 primer sets, adding blockers increased the bacterial reads to 5.3-

41.5% of total reads, compared to only 0.2 -1.9% of bacterial reads per sample

without blockers. For V1-3 and V6-8 primer sets, adding blockers only increased the

bacterial reads to 1.6-5.6% of total reads per sample. The alpha rarefaction curves

based on the counts of operational taxonomic units (OTUs) indicated that, for those

crown gall samples, the curves would approach saturation when the sequencing depth

was over about 10,000 reads per sample (Fig. 2B). According to the bar plot of

bacterial composition at family levels, we could observe higher number of families

identified from the same crown galls amplified by adding blockers (Fig. 3A).

However, the result of Principal Coordinate Analysis (PCoA) indicated that adding

blockers did not cause significant biases in the inferred microbiota composition (Fig.

3B). However, the use of different primer sets resulted in significant difference in the

inferred microbiota composition (Fig. 3B, 3C), which has been reported previously

(23).

Overall, adding blockers at the PCR step increased bacterial read counts and the

resolution of gallobiome. Because the use of V5-7 primers and cognate blockers

obtained the highest number and percentage of bacterial reads, V5-7 set-up with

blockers were selected for 16 rRNA gene amplicon sequencing of crown galls

induced by WT and T6SS mutants.

Round III of 16S rRNA amplicon sequencing: Impact of T6SS on gallobiome

Among all of 139 crown galls collected, 53 tumors in the range of 0.06 – 0.56 g (24

induced by WT, 16 by Δ*tssL,* and 13 by Δ*tssB*) were used for the analysis. We

obtained an average of 6,9980 ± 26,452 (mean ± SD) reads per sample, ranged from

9,814 to 116,072 reads per sample. Before the analysis, the amplicon sequence

variants (ASVs) were clustered into 99% OTUs and the singletons were removed.

Based on the alpha rarefaction curves and the minimal read counts of the samples we

got (Fig. S2), diversity analysis was conducted with datasets subsampling at 9800

reads per sample. The result of PCoA showed that the gallobiome induced by WT and

two T6SS mutant strains were not significantly different (Fig. 4A). Most of the

variation between samples were contributed by different seasons (July vs.

October/November) (Fig. 4B). After splitting and re-analyzing the dataset based on

the month of inoculation, we observed the difference between the gallobiomes

associated with WT and two T6SS mutants in July (Fig. 4C). In contrast, no

difference was detected in those galls induced in October or November (Fig. 4D and

4E).

The alpha diversity indices including observed OTUs, Shannon index, and Pielou's

evenness index had no significant difference between the gallobiomes associated with

WT and T6SS mutants (Fig. 5A, 5B and 5C). Interestingly, we found that, in galls

induced in July, the WT gallobiome showed significantly higher Faith's phylogenic

diversity than the gallobiome associated with Δ*tssL*; and in galls induced in

November, the Δ*tssL* gallobiome showed higher diversity than the WT gallobiome

(Fig. 5D). The gallobiome associated with Δ*tssB* also exhibited higher diversity than

the WT gallobiome but the difference was not statistically significant.

While there were variations in bacterial compositions among crown galls, the top 10

bacterial family in gallobiomes were quite consistent, including: Rhizobiaceae,

Comamonadaceae, Chitinophagaceae, Methyphliaceae, Caulobacteraceae,

Sphingomonadaceae, Microbacreriaceae, Xanthobacteraceae, Oxalobacteraceae, and

Pseudomonaceae (Fig. 5E). The bacterial family Rhizobiaceae, which *A. tumefaciens*

belongs to, was found to be highly variable and accounted for 5-85% of the entire

community. The most abundant OTU in this Rhizobiaceae dataset was 100% matched

to 16S rRNA genes of *A. tumefaciens* C58, suggesting that the abundance of this OTU

likely indicated to the relative abundance of *A. tumefaciens* WT or mutant inoculum

in crown galls. When the weight of crown galls was plotted against the relative

abundance of this OTU in gallobiomes, we found that the abundance of Rhizobiaceae

in July was dramatically lower than those in October and November (Fig. 5F), but

there was no significant difference observed among different strains. Furthermore, a

positive correlation between weight of crown galls and relative abundance of *A.*

tumefaciens OTU was observed (Fig. 5G).

236   Sphingomonadaceae and Burkholderiacea were more abundant in the

237   gallobiomes induced by the T6SS mutants in July

Next, we analyzed the species-level OTU and family with differential abundance

(DA) between gallobiome associated with WT and T6SS mutants in July. Two OTUs

belonged to Sphingomonadaceae, named as SphinOTU1 and SphinOTU2 here, and

Burkhoderiaceae family were significantly enriched in the gallobiomes induced by

T6SS mutants (Fig. 6A). SphinOTU1 was only present in gallobiomes in July but not

in October/November, and SpinOTU2 was present in both July and October but not in

November (Fig 6B). Neither SpinOTU1 or SpinOTU2 was identified in gallobiomes

induced by the WT in July. Burkhoderiaceae was enriched in gallobiomes induced by

the T6SS mutants in July, but no consistent enrichment in October and November

(**Fig 6C**). Although SphinOTU1 and SphinOTU2 were only present in gallobiomes

induced by T6SS mutants in July; at family level, Sphingomonadaceae was not

enriched in T6SS-dependent manner in any months (Fig. 6C).

A. tumefaciens **exhibits T6SS antagonism to a** *Sphingomonas* **sp. isolate**

The T6SS-dependent differential abundance of two *Sphingomonas* OTUs in

gallobiomes motivated us to investigate whether *A. tumefaciens* exhibits T6SS

antibacterial activity to *Sphingomonas*. A *Sphingomonas* sp. strain R1 isolated from

the tomato rhizosphere in one soil inoculation experiment was used for *in vitro* and *in*

planta interbacterial competition assays. The results showed that the colony forming

unit (CFU) of *Sphingomonas* sp. R1 was lower after co-culture with *A. tumefaciens*

C58 WT than with T6SS mutants (Fig. 7A). Each of *A. tumefaciens* C58 strains was

also mixed at 1:1 ratio with *Sphingomonas* sp. R1 to determine their colonization

efficiency and occupation by counting CFU from wounded tomato stems and crown

galls by soil inoculation. The results showed a reduced colonization efficiency of

Sphingomonas sp. R1 when co-inoculated with WT as compared to that with Δ*tssL*

and Δ*tssB* or R1 only, although not statistically different between WT and T6SS

mutants (Fig. 7B). No difference on the recovered CFU among *A. tumefaciens* WT

and T6SS mutants inoculated alone or with *Sphingomonas* sp. R1. However, we

noticed that the colonization efficiency of those T6SS mutants was more variable as

compared to those of WT from the combined data set (Fig. 7C). Surprisingly,

Sphingomonas sp. R1 was not always present in crown galls at 28 day-post-

inoculation (dpi), in which R1 was only recovered from one out of three independent

experiments (Fig. S3). The abundance of *A. tumefaciens* C58 ranged from 10^4 to 10^7

CFU per gall and no consistent difference could be observed between the WT and the

T6SS mutants. Together, these results suggest that *A. tumefaciens* C58 exhibits weak

but detectable antibacterial activity to inhibit the *in vitro* growth and plant

colonization efficiency of *Sphingomonas* sp. R1. The T6SS may help *A. tumefaciens*

to maintain stable colonization on plant surface for effective infection, leading to

higher disease incidence.

DISCUSSION

In this study, we designed experiments to address whether T6SS affects *Agrobacterium*

tumorigenesis and gallobiome. By establishing a soil inoculation platform to evaluate

the disease incidence of *A. tumefaciens* on wounded stems of tomato seedlings across

seasons, we revealed that T6SS exhibits positive impact on stable colonization and

disease occurrence efficiency but no impacts on tumor weight. Furthermore, by

developing an effective protocol named as <u>B</u>locker-mediated <u>B</u>acterial 16S r<u>R</u>NA gene

amplicon <u>E</u>nrichment <u>S</u>equencing (BBRES) that minimizes host contamination, we

found that seasons or environmental factors, but not T6SS are the major driver in

shaping gallobiome compositions.

The evidences that *A. tumefaciens* T6SS promotes tumorigenesis by soil inoculation

(Table 1, Fig. 1) but not under direct inoculation on tomato stems (19) suggest that the

T6SS is not directly involved in virulence. Instead, *A. tumefaciens* deploys its

antibacterial activity to increase the occurrence of infection in the presence of relatively

complex microbial communities in soil or rhizosphere. Once the tumor is induced,

T6SS is not required for tumor development since no significant tumor weight was

observed between WT and T6SS mutants (Fig. 1B). The presence of T6SS has no

impact on the population size of agrobacteria, consistent with a recent study that T6SS

genes were not identified as fitness genes by Tn-seq screen in crown galls (24). The

T6SS of *A. tumefaciens* C58 also appears not the key factor to shape or maintain the

gallobiome, suggesting that *A. tumefaciens* may use other ways for niche competition

in the tumors. It is generally believed that the agrobacterial strain inciting tumors and

later become residents of its population have privilege to access the specific opines

synthesized by its T-DNA (25). This "opine concept" has not been experimentally

validated but recent study showed that the ability of agrobacteria to trap opines can

acquire a competitive advantage over the siblings which cannot utilize opine (26).

Therefore, it would be interesting to investigate whether opine synthesis and other

fitness genes would impact the crown gall microbiota in which T6SS may

synergistically enhance the niche occupation.

The findings that disease incidence and crown gall weight in reverse correlation

with temperature (Fig. 1C) are expected. Early studies demonstrated that crown gall

tumor sizes were dramatically decreased when the host plants were inoculated at 28-

30°C, with no galls were formed at 31°C or above (27, 28). In addition, VirB/D4

T4SS complex and its associated T-pilus are more stable or produced at higher levels

at 19°C than 28°C whereas T4SS-mediated plasmid conjugation is deficient at 28°C

(29-33). The increased tumor weight when inoculated in fall/winter is also correlated

with a higher relative abundance of agrobacterial OTU (Fig. 5G), suggesting the

active proliferation of agrobacterial population inside crown galls. Furthermore, our

data showed that crown galls produced in summer have distinct microbiota as

compared to those from fall/winter (Fig. 4B), which echoes previous studies on

Agrobacterium vitis that crown gall microbiota of grapevine is significantly different

in summer (34). Interestingly, while there is no global difference of gallobiome

induced by WT or T6SS mutants inoculated in fall/winter, we found that the two

Sphingomonas OTUs and the family Burkholderiaceae were only present or

significantly enriched in the gallobiome induced by T6SS mutants in summer (Fig. 6).

By counting viable agrobacterial inoculum, we showed the *A. tumefaciens* exhibits a

T6SS-mediated antibacterial activity to *Sphingomonas* sp. R1 *in vitro* and in

rhizosphere (Fig. 7A, B) but no significant difference of R1 abundance between the

tumors co-infected with *A. tumefaciens* WT or T6SS mutants (Fig. S3). Accordingly,

we also observed more stable colonization of *A. tumefaciens* WT than T6SS mutants

on wounded tomato stems via soil inoculation (Fig. 7C). To this end, it remains to be

determined whether the absence or reduced abundance of *Sphingomonas* and

Burkholderiaceae of gallobiome induced by WT as compared to those by T6SS

mutants are due to antibacterial activity conferred by WT T6SS. However, current

evidence suggests that *A. tumefaciens* may use T6SS to gain competitive advantage

by its antibacterial activity against other bacterial species residing in the soil or

rhizosphere and therefore increases its success in inciting tumors on infected plants.

Besides Rhizobiaceae that agrobacteria belong to, other bacterial families were also

occupied considerable proportions of crown gall microbiota, such as

Comamonadaceae, members of which are commonly found in soil and water as well

as plant-associated bacteria *Variovorax* shown to be important for root growth of

Arabidopsis (35, 36). Bacteria belong to Xanthobacteraceae and Pseudomonaceae, the

two families highly associated with plants as either pathogens or commensals, also

had high relative abundance in the crown galls. Some *Pseudomonas* species such as *P.*

fluorescens and *P. putida* are capable to utilize opine (37, 38), which may explain why

bacterial species other than the agrobacterial inoculum could also colonize and

propagate in high relative abundance in crown galls.

We found that the predominance of plant DNA in crown gall samples led to the low

resolution of endophytic bacterial microbiota survey via standard 16S rRNA gene

amplicon sequencing due to the co-amplification of chloroplast and mitochondrial

sequences. Although the use of 16S rRNA gene primers containing mismatches to 16S

rRNA gene of plant chloroplast or mitochondria could potentially reduce host

contamination in endophytic microbiota studies (23, 39), the "bacteria-specific"

primers did not work well in previous studies (20) and resulted in > 90% host reads of

our initial attempt (Round I) (Table 2). When adding blockers that could specifically

anneal to chloroplast or mitochondrial DNA but not be extended by DNA polymerase

during PCR, we successfully reduced the host contamination and obtained sufficient

sequencing depth for this study. Our results support the effectiveness of blockers in

increasing bacterial reads of endophyte community (20-22). The newly developed

BBRES protocol for gallobiome may be applicable to microbiota associated with

tomato, or even other plants after modifying the blocker sequences.

To date, collective plant microbiota studies suggest the enrichment of T6SS genes in

microbiota and as a protected trait for niche competition in microbiota (5, 8).

However, the role and impact of T6SS in different plant and microbiome contexts

may be different. Our study suggest that agrobacteria may use T6SS to gain

competitive advantage at initial stage of infection but likely not critical for its

proliferation once they established their niche inside the crown galls. Future work to

investigate the spatial distribution of agrobacterial cells and other endophytes inside

crown gall or wounded site of plants together with visualization of T6SS dynamics is

important to dissect the role of T6SS during agrobacterial life cycle.

MATERIALS AND METHODS

Bacterial strains and growth condition

Bacterial strains and plasmids used in this study are listed in Table S3. *Agrobacterium*

tumefaciens strain C58 wild type and two T6SS mutants (i.e., Δ*tssB* and Δ*tssL*) (40)

were first streaked on 523 medium agar plates (41) and incubate at 25 °C for 48 hr.

Sphingomonas sp. R1 was grown on R2A medium without soluble starch and

pyruvate (42). *Escherichia coli* was grown on lysogeny broth (LB) medium at 37°C.

Freshly grown colony was inoculated into corresponding broth for overnight culture.

The antibiotics and concentration used are: 10 μg/ml gentamycin for *E. coli,* 50 μg/ml

gentamycin for *A. tumefaciens*, and 20 μg/ml kanamycin and 50μg/ml gentamycin for

Sphingomonas sp. R1.

Construction of strains enable antibiotics selection

Primers used for construction are listed in Table S4. The DNA fragment containing

gentamycin resistance (Gm^R) gene and green fluorescence protein *gfp*(S65T) were

amplified from pRL662::GFP(S65T) with the primer pair "BclI-GFP-GmR-F-25" and

"BclI-GFP-GmR-R-25", and then cloned into XbaI digested pJQ-COM. The newly

constructed plasmid was named pJQ-com-GmR-GFP, which enabled the generation of

Gm^R gene and *gfp* (S65T) knocked in at the upstream of *actC* (*atu5330*) of *A.*

tumefaciens genome after double crossover. The knock-in strains were generated in *A.*

tumefaciens C58, Δ*tssB*, and Δ*tssL* background. The knock-in event and excision of

pJQ200ks backbone were confirmed by GFP signal, gentamycin resistance, and PCR

with the primer pair "3'sacB-5'GmR pJQ200 F" and "3'sacB-5'GmR pJQ200 R",

which target the fragment between *sacB* and Gm^R gene on pJQ200ks. The plasmids

pRL66::GFP(S65T) and pBBR::mCherry conferring kanamycin resistance were each

transformed into *Sphingomonas* sp. R1 through electroporation. The procedure of

electroporation, doble crossover via pJQ200ks were performed as described

previously (43).

Plant material and growth condition

Seeds of *Solanum lycopersicum* (tomato) cultivar Known-You 301 from KNOWN-

YOU SEED CO., LTD (Kaohsiung, Taiwan) were germinated and grown in

unsterilized potting mixture (Jiffy® Premium Fine Peat Substrate, perlite, and

vermiculite, mixed in 4:1:1 ratio, v:v:v) in the greenhouse EL329

(N25.043047890856812, E121.61135464167913, Greenhouse building, Academia

Sinica, Taipei, Taiwan).

Soil inoculation

Soil inoculation method was performed essentially following the method used for *A.*

tumefaciens on pea and walnut (44, 45). Overnight (14-16 hr) culture of *A.*

tumefaciens strain in 12 ml of 523 broth were centrifugated at $6,000 \times g$ for 10 min.

The pellets were washed once by using 10 ml of sterile saline (0.9% NaCl in H_2O).

The washed pellets were centrifuged and resuspended in sterile saline with OD_{600}

adjusted to 1. Bacterial suspension was mixed into unsterilized soil (1 ml suspension

per 100 g of soil), which is expected to result in $\sim 10^7$ colony-forming units (CFUs)/g

soil. Tomato seedlings with two true leaves (two- to three-week-old) were wounded at

the site between primary root and cotyledons by a fire-sterilized sewing needle,

planted into the inoculated soil, and grown in the greenhouse. Each strain was

inoculated with 14 to 20 tomato seedlings, and the inoculation with saline were used

as a negative control

Harvest, surface sterilization and storage of crown gall samples

Two months after the inoculation, tomato plants were harvested to examine gall

formation on the wounded site. The disease incidence was calculated by dividing the

number of plants with visible gall formation by the total number of inoculated plants.

The sections with crown gall were cut out and sterilized in 35 ml of sterilization

solution (3% NaOCl, 0.01% tween-20) for 30s, then transferred to 35 ml of 70%

ethanol and washed in 35 ml of sterile H_2O for three times (46). The sterilization was

evaluated by spreading 100 µl liquid from the third wash onto 523 agar plate for 2

days to make sure that there was no growth of bacteria. After sterilization, the crown

galls were dissected from each of the plant segments in sterile petri dish, wrapped in

sterile aluminum foil, frozen by liquid nitrogen, and stored at -80 °C prior to DNA

extraction.

DNA extraction form crown galls

439 Crown gall samples were homogenized by pestle and mortar with liquid nitrogen.

440 0.25 g of homogenized tissue were transferred to the beating tube of DNeasy®

441 PowerSoil® Kit (Qiagen, Germany) following the manufacturer's instructions.

442 Concentrations of extracted DNA were determined by NanoDrop™ 1000 (Thermo

443 Fisher Scientific, United State) and Qubit dsDNA HS (Invitrogen, United State).

16S rRNA gene amplification and sequencing

Three rounds of amplicon sequencing were conducted in this study. The round I was

conducted according to established protocols of soil microbiota studies using 16S

rRNA gene variable regions (23). Each primer with a partial adaptor sequence was

compatible to Illumina Truseq Combinatorial dual (CD) index system. The first PCR

amplifications were carried out in following 25 µl reaction: 5 ng DNA template, 2×

KAPA HiFi HotStart DNA Polymerase ReadyMix (Roche, Switzerland), and 0.2 µM

for each of the forward and reverse primers. The first PCR program and reaction

details for V3-4 and V5-7 are listed in **Table S5**. The PCR products were purified by

Ampure XP beads (Thermo Fisher Scientific Inc, Sweden). The V5-7 PCR products

underwent BluePippin (Sage Science) size-selection to remove tomato amplicon. To

attach dual indices and full-length Illumina sequencing adapters, a second round of

PCR amplifications were employed for 50 µl reactions with purified or size-selection

PCR products, 2× KAPA HiFi HotStart DNA Polymerase ReadyMix (Roche,

Switzerland), and 0.5 µM for each of the forward and reverse primers. All the library

products of second PCR were purified and pooled. Pooled libraries were loaded onto

an Illumina MiSeq V3 flow cell (Illumina, United States) and for 2×300 bp paired-

end sequencing.

The round II of amplicon sequencing was conducted to test the effect of adding PCR

blockers on reducing plant chloroplast and mitochondrial sequences. Three gall

samples (1115-W21, 1115-W22 and 1115-W25) were used as the DNA templates to

provide biological replicates. For each DNA sample, four primer pairs targeting

different variable regions of 16S rRNA gene were used, including: V1-3, V3-4, V5-7,

and V6-8 (Table S1). Each primer contained a partial adaptor sequence of the Illumina

Truseq Combinatorial dual (CD) index system. Cognate 3' C3 spacer modified

oligonucleotides (blockers) which would block the amplification of 16S rRNA genes

of tomato chloroplasts and mitochondria were designed according to a previous study

(20) and applied to one of the two PCRs for each sample-primer combination (Table

S2). Procedures for PCR and Illumina sequencing were based on those used for round

I, but the amount of DNA template was increased to 25 ng to reduce PCR cycle time

and the annealing temperature was adjusted for blockers. More detailed technical

information is provided in Table S4.

Based on results from the first two rounds of amplicon sequencing, an optimized

protocol that utilizes blockers and targets V5-7 regions was used for round III. All 53

crown gall samples were included in this final round.

Raw read processing and microbiota analysis

The Illumina raw reads in FASTQ format were imported into QIIME2 release 2020.8

(47). The primer region of imported sequences was trimmed using DADA2 (48). The

denoised and merged paired-end reads were used to construct output feature tables

containing amplicon sequence variants (ASVs) ID and counts in different samples.

For amplicon sequencing round III, the ASVs were further clustered into operational

taxonomic units (OTUs) based on 99% sequence identity using the "qiime vsearch

cluster-features-de-novo" function (49) and the singletons were removed by using the

"qiime feature-table filter-features" function prior to the downstream analysis. The

taxonomic information was inferred for representative OTU sequences via pre-trained

q2-feature-classifier (50) and classify-sklearn naïve Bayes classifier based on SILVA

version 138 SSU Ref NR 99 full-length sequences dataset (51). The sequences

assigned as chloroplasts, mitochondria, eukaryotes, archaea, or unknown were

removed. After the filtering, a phylogenetic tree of eubacterial sequences was inferred

using the q2-fragment-insertion plug-in (52-55) based on the SILVA 128 SEPP

reference database. Weighted Unifrac (56) analysis of those datasets was performed

based on insertion tree; the distance matrix and principal coordinate analysis (PCoA)

plot were generated using QIIME2. For round III, phylogenic and diversity analysis

was conducted using "qiime diversity core-metrics-phylogenetic" and the sub-

sampling depth was set to 9,800 reads/sample.

Differential abundance analysis of OTU

Analysis of composition of microbiomes (ANCOM) was applied to identify OTUs

that have significantly different relative abundance in the gallobiome associated with

A. tumefaciens C58 WT and T6SS-dificient mutants using the function "qiime

composition ancom" in QIIME2. The feature table collapsed into family level was

imported and transformed by centred log-ratio transformation. The null hypothesis of

ANCOM is that the abundance of two OTUs were not different between two study

groups. After performing all comparison between each OTU in two study groups, the

times that the null hypothesis was rejected is called *w*. OTUs having *w* values that are

at 70 percentile or higher are considered as significant (57, 58).

Interbacterial competition assay *in vitro* and *in planta*

For interbacterial competition *in vitro,* the bacterial suspension was adjusted to OD_{600}

equals to 3.0 in 0.9% NaCl (w/v) after overnight culture. For the *in vitro* competition,

the attacker strains (*A. tumefaciens* C58 WT, Δ*tssL* or Δ*tssB*) were further diluted into

OD_{600} equals to 1, and the prey *Sphingomonas* sp. R1 containing pRL662::GFP was

further diluted into OD_{600} equal to 0.3 or 0.1 for mixing with the attacker in equal

volume for different density ratio. Two spots of 10 µl mixture were spotted on

Agrobacterium kill-triggering medium(18). After co-incubated for 16 h in 25 ℃, the

colonies were scraped and resuspended in 1 ml 0.9% NaCl (w/v), and spread via

exponential mode of easySpiral Dilute® (Ref. 414 000, INTERSCIENCE, France) on

523 media containing 30 µg/ml gentamycin to recover the prey.

For *A. tumefaciens* colonization with or without *Sphingomonas* sp. R1, each of *A.*

tumefaciens C58 strains (WT, Δ*tssL* or Δ*tssB*) with Gm^R-GFP knock-in was mixed

with *Sphingomonas* sp. R1 containing pBBR::mCherry at equal OD_{600} =1.0. The 5 µl

mixture was inoculated on the needle-wounded stem of tomato seedling with first pair

of true leaves. Tomato seedlings were harvested at 10 days post-inoculation (dpi). For

each plant, the 1 cm wounded segment was cut and rinsed by sterile water before

being transferred into a sterilized Eppendorf tube containing 1 ml of 0.9% NaCl. After

vortex at max speed for 3 minutes, the liquid was considered as the rhizosphere

sample. The samples were diluted and plated on media with gentamycin for CFU

calculation.

When counting the CFU of *A. tumefaciens* C58 and *Sphingomonas* sp. R1 in the

tumor, the crown galls were collected after growing in the greenhouse for three

weeks. The 1-cm tomato stem section containing the gall in middle was cut and

homogenized with 1 ml 0.9% NaCl (w/v) by sterilized mortar and pestle. The

homogenized tissue was diluted 10-fold and spread via exponential mode of

easySpiral Dilute® on 523 media containing 50 ppm gentamycin to recover the

attacker and prey. After incubation for two days at 25°C, the colonies were counted

by using the Scan® 500 Automatic colony counter (Ref 436 000, INTERSCIENCE,

France; software version 8.6.1) to calculate CFUs.

Statistics and visualization

GraphPad Prism 8 was used to perform t-test on sequencing read counts, one-way

ANOVA on CFU counts, two-way ANOVA followed by Turkey's multiple

comparison test on crown gall disease incidence and alpha diversity index, Kruskai-

Wallis test on tumor weight, Spearman correlation on gall weight and relative

abundance of *A. tumefaciens* OTU. The alpha-rarefaction curves were plotted using

Phyloseq (59) in R version 3.6.2. ADONIS in "qiime diversity beta-group-

significance" of QIIME2 (60, 61) was used to test if groups of samples have

significantly different microbiota composition. The taxonomic composition plots were

generated using qiime2R (v0.99.35, https://github.com/jbisanz/qiime2R) in R studio

2.

DATA AVAILABILITY

All raw data sets are available in the National Center for Biotechnology Information

(NCBI) under BioProject accession PRJNA894311.

AUTHOR CONTRIBUTION

Conceptualization: SCW, CHK, EML

Funding acquisition: CHK, EML

Investigation: SCW, CHK, EML

Methodology: SCW, APC, SJC, CHK

Project administration: EML

Supervision: CHK, EML

Validation: SCW, CHK

Visualization: SCW, CHK

Writing – original draft: SCW, CHK, EML

Writing – review & editing: SCW, APC, SJC, CHK, EML

ACKNOWLEDGEMENTS

The authors thank the members of Lai and Kuo labs for helpful discussion. We

also acknowledge Lay-Sun Ma for constructing pRL662∷GFP(S65T) and Stanton

Gelvin for providing pBBR∷mCherry. The Illumina sequencing library preparation

was carried out in the Genomic Technology Core (Institute of Plant and Microbial

Biology, Academia Sinica). We thank the Illumina MiSeq sequencing service

provided by the Genomics Core (Institute of Molecular Biology, Academia Sinica).

FUNDING

Research in the Lai lab was supported by Institute of Plant and Microbial Biology,

Academia Sinica, Academia Sinica and Academia Sinica Investigator Award to EML

(grant no. AS-IA-107-L01). Research in the Kuo lab was supported by Academia

Sinica and the Ministry of Science and Technology of Taiwan (MOST 109-2628-B-

001-012; 110-2628-B-001-020; 111-2628-B-001-019). The funders had no role in

study design, data collection and interpretation, or the decision to submit the work for

publication.

REFERENCES

1.   Coulthurst S. 2019. The Type VI secretion system: a versatile bacterial

weapon. Microbiology (Reading) 165:503-515.

2.   Jurenas D, Journet L. 2021. Activity, delivery, and diversity of Type VI secretion

effectors. Mol Microbiol 115:383-394.

3.   Lien YW, Lai EM. 2017. Type VI Secretion Effectors: Methodologies and

Biology. Front Cell Infect Microbiol 7:254.

4.   Gallegos-Monterrosa R, Coulthurst SJ. 2021. The ecological impact of a

bacterial weapon: microbial interactions and the Type VI secretion system.

FEMS Microbiol Rev 45.

5.   Bulgarelli D, Garrido-Oter R, Munch PC, Weiman A, Droge J, Pan Y, McHardy

AC, Schulze-Lefert P. 2015. Structure and function of the bacterial root

microbiota in wild and domesticated barley. Cell Host Microbe 17:392-403.

6.   Wexler AG, Bao Y, Whitney JC, Bobay LM, Xavier JB, Schofield WB, Barry NA,

Russell AB, Tran BQ, Goo YA, Goodlett DR, Ochman H, Mougous JD, Goodman

AL. 2016. Human symbionts inject and neutralize antibacterial toxins to

persist in the gut. Proc Natl Acad Sci U S A 113:3639-44.

7.   Verster AJ, Ross BD, Radey MC, Bao Y, Goodman AL, Mougous JD, Borenstein

E. 2017. The Landscape of Type VI Secretion across Human Gut Microbiomes

Reveals Its Role in Community Composition. Cell Host Microbe 22:411-419

e4.

8.   Vogel CM, Potthoff DB, Schafer M, Barandun N, Vorholt JA. 2021. Protective

role of the Arabidopsis leaf microbiota against a bacterial pathogen. Nat

Microbiol 6:1537-1548.

9.   Serapio-Palacios A, Woodward SE, Vogt SL, Deng W, Creus-Cuadros A, Huus

KE, Cirstea M, Gerrie M, Barcik W, Yu H, Finlay BB. 2022. Type VI secretion

systems of pathogenic and commensal bacteria mediate niche occupancy in

the gut. Cell Rep 39:110731.

10.   Ma LS, Hachani A, Lin JS, Filloux A, Lai EM. 2014. *Agrobacterium tumefaciens*

deploys a superfamily of type VI secretion DNase effectors as weapons for

interbacterial competition in planta. Cell Host Microbe 16:94-104.

11.   Bernal P, Allsopp LP, Filloux A, Llamas MA. 2017. The *Pseudomonas putida*

T6SS is a plant warden against phytopathogens. ISME J 11:972-987.

12.   Bernal P, Llamas MA, Filloux A. 2018. Type VI secretion systems in plant-

associated bacteria. Environ Microbiol 20:1-15.

13.   Vacheron J, Pechy-Tarr M, Brochet S, Heiman CM, Stojiljkovic M, Maurhofer

M, Keel C. 2019. T6SS contributes to gut microbiome invasion and killing of an

herbivorous pest insect by plant-beneficial *Pseudomonas protegens*. ISME J

13:1318-1329.

14.   Hwang HH, Yu M, Lai EM. 2017. *Agrobacterium*-Mediated Plant

Transformation: Biology and Applications, vol 15. American Society of Plant

Biologists.

15.   Wu CF, Santos MNM, Cho ST, Chang HH, Tsai YM, Smith DA, Kuo CH, Chang JH,

Lai EM. 2019. Plant-Pathogenic *Agrobacterium tumefaciens* Strains Have

Diverse Type VI Effector-Immunity Pairs and Vary in In-Planta

Competitiveness. Molecular Plant-Microbe Interactions 32:961-971.

16.   Wu CF, Weisberg AJ, Davis EW, 2nd, Chou L, Khan S, Lai EM, Kuo CH, Chang JH.

2021. Diversification of the Type VI Secretion System in Agrobacteria. mBio

12:e0192721.

17.   Chou L, Lin YC, Haryono M, Santos MNM, Cho ST, Weisberg AJ, Wu CF, Chang

JH, Lai EM, Kuo CH. 2022. Modular evolution of secretion systems and
virulence plasmids in a bacterial species complex. BMC Biol 20:16.

18. Yu M, Wang YC, Huang CJ, Ma LS, Lai EM. 2020. *Agrobacterium tumefaciens*
Deploys a Versatile Antibacterial Strategy to Increase its Competitiveness. J
Bacteriol doi:10.1128/JB.00490-20.

19. Wu HY, Chung PC, Shih HW, Wen SR, Lai EM. 2008. Secretome Analysis
Uncovers an Hcp-Family Protein Secreted via a Type VI Secretion System in
Agrobacterium tumefaciens. J Bacteriol 190:2841-2850.

20. Lefevre E, Gardner CM, Gunsch CK. 2020. A novel PCR-clamping assay
reducing plant host DNA amplification significantly improves prokaryotic
endo-microbiome community characterization. Fems Microbiology Ecology
96.

21. Fitzpatrick CR, Lu-Irving P, Copeland J, Guttman DS, Wang PW, Baltrus DA,
Dlugosch KM, Johnson MTJ. 2018. Chloroplast sequence variation and the
efficacy of peptide nucleic acids for blocking host amplification in plant
microbiome studies. Microbiome 6:144.

22. Arenz BE, Schlatter DC, Bradeen JM, Kinkel LL. 2015. Blocking primers reduce
co-amplification of plant DNA when studying bacterial endophyte
communities. Journal of Microbiological Methods 117:1-3.

23. Thijs S, Op De Beeck M, Beckers B, Truyens S, Stevens V, Van Hamme JD,
Weyens N, Vangronsveld J. 2017. Comparative Evaluation of Four Bacteria-
Specific Primer Pairs for 16S rRNA Gene Surveys. Front Microbiol 8:494.

24. Torres M, Jiquel A, Jeanne E, Naquin D, Dessaux Y, Faure D. 2022.
Agrobacterium tumefaciens fitness genes involved in the colonization of plant
tumors and roots. New Phytol 233:905-918.

25. Guyon P, Chilton MD, Petit A, Tempe J. 1980. Agropine in "null-type" crown
gall tumors: Evidence for generality of the opine concept. Proc Natl Acad Sci U
S A 77:2693-7.

26. Lang J, Vigouroux A, Planamente S, El Sahili A, Blin P, Aumont-Nicaise M,
Dessaux Y, Morera S, Faure D. 2014. *Agrobacterium* uses a unique ligand-
binding mode for trapping opines and acquiring a competitive advantage in
the niche construction on plant host. PLoS Pathog 10:e1004444.

27. Riker AJ. 1926. Studies on the influence of some environmental factors on the
development of crown gall. Journal of Agricultural Research 32:0083-0096.

28. Braun AC. 1947. Thermal Studies on the Factors Responsible for Tumor
Initiation in Crown Gall. American Journal of Botany 34:234-240.

29. Fullner KJ, Nester EW. 1996. Temperature affects the T-DNA transfer
machinery of *Agrobacterium tumefaciens*. J Bacteriol 178:1498-504.

30. Fullner KJ, Lara JC, Nester EW. 1996. Pilus assembly by *Agrobacterium* T-DNA
transfer genes. Science 273:1107-9.

31. Lai EM, Kado CI. 1998. Processed VirB2 is the major subunit of the
promiscuous pilus of *Agrobacterium tumefaciens*. J Bacteriol 180:2711-7.

32. Lai EM, Kado CI. 2000. The T-pilus of *Agrobacterium tumefaciens*. Trends
Microbiol 8:361-9.

33. Baron C, Domke N, Beinhofer M, Hapfelmeier S. 2001. Elevated temperature
differentially affects virulence, VirB protein accumulation, and T-pilus
formation in different *Agrobacterium tumefaciens* and *Agrobacterium vitis*
strains. J Bacteriol 183:6852-61.

34. Faist H, Keller A, Hentschel U, Deeken R. 2016. Grapevine (Vitis vinifera)
Crown Galls Host Distinct Microbiota. Appl Environ Microbiol 82:5542-52.

35. Willems A, De Ley., J. , Gillis, M. Kersters, D. K. . 1991. Comamonadaceae, a
New Family Encompassing the *Acidovorans* rRNA Complex, Including
Variovoraxparadoxus gen. nov., comb. nov. for Alcaligenes paradoxus (Davis
1969). INTERNATIONJOAULRNAOLF SYSTEMATBICACTERIOLO 41.

36. Finkel OM, Salas-Gonzalez I, Castrillo G, Conway JM, Law TF, Teixeira P, Wilson
ED, Fitzpatrick CR, Jones CD, Dangl JL. 2020. A single bacterial genus
maintains root growth in a complex microbiome. Nature 587:103-108.

37. Moore LW, Chilton WS, Canfield ML. 1997. Diversity of opines and opine-
catabolizing bacteria isolated from naturally occurring crown gall tumors.
Applied and Environmental Microbiology 63:201-207.

38. Beauchamp CJ, Dion P, Kloepper JW, Antoun H. 1991. Physiological
Characterization of Opine-Utilizing Rhizobacteria for Traits Related to Plant
Growth-Promoting Activity. Plant and Soil 132:273-279.

39. Beckers B, Op De Beeck M, Thijs S, Truyens S, Weyens N, Boerjan W,
Vangronsveld J. 2016. Performance of 16s rDNA Primer Pairs in the Study of
Rhizosphere and Endosphere Bacterial Microbiomes in Metabarcoding
Studies. Front Microbiol 7:650.

40. Lin JS, Ma LS, Lai EM. 2013. Systematic Dissection of the *Agrobacterium* Type
VI Secretion System Reveals Machinery and Secreted Components for
Subcomplex Formation. PLoS One 8:e67647.

41. Kado CI, and Heskett, M.G. . 1970. Selective media for isolation of
Agrobacterium, Carynebacterium, Erwinia, Pseudomonas, and
Xanthomonas. . Phytopathology 60:969-976.

42. Massa S, Caruso M, Trovatelli F, Tosques M. 1998. Comparison of plate count
agar and R2A medium for enumeration of heterotrophic bacteria in natural
mineral water. World Journal of Microbiology & Biotechnology 14:727-730.

43.   Ma LS, Lin JS, Lai EM. 2009. An IcmF family protein, ImpLM, is an integral
inner membrane protein interacting with ImpKL, and its walker a motif is
required for type VI secretion system-mediated Hcp secretion in
Agrobacterium tumefaciens. J Bacteriol 191:4316-29.
44.   Hawes MC, Smith LY. 1989. Requirement for chemotaxis in pathogenicity of
Agrobacterium tumefaciens on roots of soil-grown pea plants. J Bacteriol
171:5668-71.
45.   Yakabe LE, Parker SR, Kluepfel DA. 2012. Role of Systemic *Agrobacterium*
tumefaciens Populations in Crown Gall Incidence on the Walnut Hybrid
Rootstock 'Paradox'. Plant Disease 96:1415-1421.
46.   McPherson MR, Wang P, Marsh EL, Mitchell RB, Schachtman DP. 2018.
Isolation and Analysis of Microbial Communities in Soil, Rhizosphere, and
Roots in Perennial Grass Experiments. Jove-Journal of Visualized Experiments
doi:ARTN e57932
10.3791/57932.
47.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA,
Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K,
Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodriguez AM,
Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM,
Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons
SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H,
Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang
KB, Keefe CR, Keim P, Kelley ST, Knights D, et al. 2019. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2.
Nat Biotechnol 37:852-857.
48.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nat
Methods 13:581-3.
49.   Rognes T, Flouri T, Nichols B, Quince C, Mahe F. 2016. VSEARCH: a versatile
open source tool for metagenomics. PeerJ 4:e2584.
50.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA,
Gregory Caporaso J. 2018. Optimizing taxonomic classification of marker-gene
amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome
6:90.
51.   Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies
J, Ludwig W, Glockner FO. 2014. The SILVA and "All-species Living Tree Project
(LTP)" taxonomic frameworks. Nucleic Acids Res 42:D643-8.
52.   Janssen S, McDonald D, Gonzalez A, Navas-Molina JA, Jiang L, Xu ZZ, Winker K,

Kado DM, Orwoll E, Manary M, Mirarab S, Knight R. 2018. Phylogenetic
Placement of Exact Amplicon Sequences Improves Associations with Clinical
Information. mSystems 3.

53. Matsen FA, Hoffman NG, Gallagher A, Stamatakis A. 2012. A format for
phylogenetic placements. PLoS One 7:e31009.

54. Eddy SR. 2011. Accelerated Profile HMM Searches. PLoS Comput Biol
7:e1002195.

55. Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-
likelihood and Bayesian phylogenetic placement of sequences onto a fixed
reference tree. BMC Bioinformatics 11:538.

56. Lozupone C, Knight R. 2005. UniFrac: a new phylogenetic method for
comparing microbial communities. Appl Environ Microbiol 71:8228-35.

57. Lin H, Peddada SD. 2020. Analysis of microbial compositions: a review of
normalization and differential abundance analysis. NPJ Biofilms Microbiomes
6:60.

58. Mandal S, Van Treuren W, White RA, Eggesbo M, Knight R, Peddada SD. 2015.
Analysis of composition of microbiomes: a novel method for studying
microbial composition. Microb Ecol Health Dis 26:27663.

59. McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. PLoS One
8:e61217.

60. Anderson MJ. 2001. A new method for non-parametric multivariate analysis
of variance. Austral Ecology 26:32-46.

61. Dixon P. 2003. VEGAN, a package of R functions for community ecology.
Journal of Vegetation Science 14:927-930.

FIGURE LEGENDS

Figure 1. Tumorigenesis assay of *A. tumefaciens* C58 and its T6SS-deficienct

mutants.

For each experiment, 14-20 wounded tomato seedlings were grown in soil containing

one of the strains tested (i.e., C58 WT, Δ*tssL*, and Δ*tssB*) and collected at 60 days post

inoculation (dpi). (A) Disease incidence plotted by strain and color-coded according

to the month of inoculation (light yellow, June; orange, July; light blue, October; dark

blue, November). Lines and error bars indicate mean ± SD. Statistical significance

was tested using two-way ANOVA followed by Tukey's multiple comparisons; $p =$

0.025 and 0.027 when comparing the wild type to Δ*tssL* and Δ*tssB*, respectively. (B)

Weight distribution of the galls collected. No significant difference among the three

strains ($p = 0.15$, Kruskai-Wallis test). (C) Correlation plot of disease incidence and

temperature of daily average. (D) Crown galls generated through soil inoculation,

example from inoculation on 11/15/2018. N indicated no gall formation in some of

inoculated plants.

Figure 2. Optimizing 16S rRNA gene amplification by using PCR blockers.

Three crown gall samples (named as 21W, 22W and 25W) induced by *A. tumefaciens*

C58 wild type were used for analysis. V1-3, V3-4, V5-7 and V6-8 represent different

primer pairs targeting different variable regions on 16S rRNA gene. (A) The OTU

counts of 16 rRNA gene amplicons after filtering non-bacterial OTUs. (ns, not

statistically significant; *, $p < 0.05$, student's t-test). The bar indicated mean of OTUs

± SD. (B) Alpha rarefaction curves of the observed bacterial OTUs based on datasets

from amplicon sequencing run II. Sample size in X-axis indicates different sub-

sampling depth of each dataset.

Figure 3. Gallobiome composition when using different primers amplified with

or without blockers. (A) Different primer sets (V1-3, V3-4, V5-7 and V6-8**)** for

analysis of three crown gall samples (21W, 22W and 25W, see Table S6) induced by

A. tumefaciens C58 wild type were indicated. Top 10 observed families were listed.

"-": amplification without blockers; "+": amplification with blockers. Principle

coordinate analysis (PCoA) plots of the bacterial composition of the datasets were

generated based on the weighted Unifrac matrix, and color coded by (B) blocker

usage (p-value = 0.068, R^2 = 0.026, ADONIS), (C) different primer sets (p-value =

0.001, R^2 = 0.524, ADONIS) and (D) different crown gall samples (p-value = 0.001,

R^2 = 0.238, ADONIS).

Figure 4. Principle coordinate analysis (PCoA) of bacterial composition in 53

crown galls induced by *A. tumefaciens* C58 WT and T6SS mutants.

The plots were drawn based on weighted Unifrac matrix of the bacterial communities.

Crown galls induced by different C58 strains (WT, Δ*tssB*, Δ*tssL*) (A) or inoculated in

different months (B) were labeled with different colors in each panel. The datasets

were further split based on the month of inoculation, and the PCoA plots of

gallobiomes from (C) July, (D) October and (E) November were indicated above.

Statistical differences in clustering were evaluated via ADONIS permutation test, and

the corresponding p-values and R^2 are indicated.

Figure 5. Alpha diversity and composition of crown galls generated in different

months by *A. tumefaciens* C58 WT and T6SS mutants.

(A) Observed OTU, (B) Shannon index, (C) evenness, and (D) Faith's phylogenic

diversity of each sample generated in different months was indicated. Lines and error

bars indicate mean ± SD. Two-way ANOVA followed by Tukey's multiple

comparisons test were performed, and the asterisks indicated the statistically

significant difference between two strains (p-value < 0.05). (E) Bacterial composition

of 53 crown galls. For each strain, the samples are grouped by the month of

inoculation. Top-ten most abundant families were listed and colored differently, and

the remaining ones are combined into "Others". (F) Relative abundance of

Rhizobiaceae in galls inoculated in different months and strains. (G) Scatter plot of

relative abundance of *A. tumefaciens* OTU in gallobiome - weight of the crown galls.

A positive correlation between weight of crown gall and relative abundance of *A.*

tumefaciens C58 was observed (Spearman r = 0.333, p = 0.01).

Figure 6. Analysis of composition of microbiomes (ANCOM) for gallobiomes in

July.

The ANCOM volcano plots were drawn based on the centred-log-ratio-transformed

abundance of (A) OTUs or bacterial families. The taxa that exhibit differential

abundance (DA) between gallobiomes induced by WT and T6SS mutants are

indicated by red arrows. The two DA OTUs belonging to *Sphingomonadaceae* were

named SphinOTU1 and SphinOTU2, respectively. The relative abundance of focal

taxa was shown in panels (B) and (C). Lines and error bars indicate mean ± SD.

Fig. 7 Competition between *A. tumefaciens* C58 and *Sphingomonas* sp. R1 *in vitro*

and on wounded site of tomato stems.

Each of the three *A. tumefaciens* strains (i.e., C58 WT, Δ*tssL*, and Δ*tssB*) was

individually tested for competition against *Sphingomonas* sp. R1 isolated from

rhizosphere of tomato. (A) The *in vitro* competition results based on *A. tumefaciens*:

Sphingomonas sp. R1 at 1:1 and 10:1 ratio, and the survival of *Sphingomonas* sp. R1

was plotted; lines and error bars indicate mean ± SD. Based on ANOVA followed by

Tukey's multiple comparison test, C58 significantly reduced *Sphingomonas* sp. R1

CFUs when mixed in 10:1 ratio ($p = 0.0153$) but not in 1:1 ratio ($p = 0.2753$). (B) The

A. tumefaciens alone or co-infected with *Sphingomonas* sp. R1 at 1:1 ratio in soil

were recovered from the surface of wounded stem segments at 10 dpi, the CFU of *A.*

tumefaciens and *Sphingomonas* sp. R1 were plotted. T-test indicated the CFU

differences of *Sphingomonas* sp. R1 inoculation alone or co-infection with T6SS

mutants as compared to WT with indicated *p* values. ANOVA test indicated no

significant difference among means ($p = 0.0945$ and 0.0551 for *A. tumefaciens* and

Sphingomonas sp. R1 CFUs, respectively). (C) The colonies of *A. tumefaciens* strains

C58 WT, Δ*tssL*, and Δ*tssB* recovered from the surface of wounded stem segments

following the soil inoculation procedure were plotted. Different symbols indicated the

outcome from different batches of colonization assay. F-test indicated the variance of

Δ*tssL* and Δ*tssB* were significantly different when compared to C58 WT ($F =$

0.000467 and $8.04398E-05$, respectively). Brown-Forsythe and Welch ANOVA test

indicated no significant difference among means ($p = 0.4847$). Line indicates median.

Figure S1. DNA sequence alignment of 16S rRNA genes of selected bacterial

strains and tomato KnowYou 301 chloroplast and mitochondria.

The 16S rRNA gene sequences of chloroplasts and mitochondria in *Solanum*

lycopersicum (tomato) cultivar Known-You 301 were obtained from this study. All of

the bacterial 16S rRNA gene sequences were accessed from Reference Sequence

(RefSeq) database in National Center of Biotechnology Information (NCBI). The
species, strain name, and accession number of 16S rRNA gene were listed at on the
left of each sequence. The alignment of sequences mentioned above were aligned in
MEGAX via ClustalW multiple alignment with gap opening penalty 15.00 and gap
extension penalty 6.66 (default). The levels of shaded blue color reflect the degree of
identity. The regions of each primer and blocker were underlined. The sequences
highlighted by red, black, yellow and green are the annealing region of primer sets for
V1-3, V3-4, V5-7 and V6-8 in 16S rRNA gene, respectively. And the gray framed
sequences in tomato chloroplast and mitochondrial 16S rRNA genes are the blockers
for cognate primers.

Figure S2. Alpha rarefaction curves of the observed bacterial OTUs based on
amplicon sequencing of 53 gall samples in sequencing run III.
The rarefaction curves were plotted after clustering DADA2-output ASVs into 99%
OTUs. Sample size in X-axis indicates different sub-sampling depth of each dataset;
Species in Y-axis indicates the number of observed OTUs under certain sub-sampling
depth. Label of each curves indicates the sample ID in metadata.

Figure S3. Competition of *A. tumefaciens* and *Sphingomonas* sp. R1 in tomato
gall on stem at 28 dpi.
Each of the three *A. tumefaciens* strains (i.e., C58 WT, Δ*tssL*, and Δ*tssB*) and R1 were
mixed at 1:1 ratio then inoculated on wounded tomato stem. The galls were harvested
and homogenized for plating at 28 dpi. *A. tumefaciens* C58 strains and *Sphingomonas*
sp. R1 were recover on 523 medium plates containing proper antibiotics. R1 could
only be recovered from galls in batch 3. The *p*-value of ANOVA against CFU
numbers of C58 for each batch is 0.786, 0.0023 and 0.230; the *p*-value of ANOVA
against CFU numbers of R1 was 0.6413. The significance of CFU numbers of C58 in
second batch is due to the low CFU counts of Δ*tssL* single inoculation group, which
could not be replicated in other two batches.

Table 1. Incidences of crown gall disease on wounded tomato stems by soil inoculation

Batch	Date of inoculation	Disease incidence (diseased/total inoculated seedlings)		
		C58 WT[a]	Δ*tssL*[b]	Δ*tssB*[b]
1	2018/11/15	0.86 (12/14)	0.86 (12/14)	N.D.*
2	2018/12/20	0.30 (6/20)	0.45 (9/20)	N.D.
3	2019/06/10	0.10 (2/20)	0.05 (1/20)	0.05 (1/20)
4	2019/07/04	0.35 (7/20)	0.35 (7/20)	0.26 (5/19)
5	2019/07/22	0.30 (6/20)	0.00 (0/20)	0.10 (2/20)
6	2019/07/29	0.27 (4/15)	0.27 (4/15)	0.14 (2/14)
7	2019/10/05	0.35 (7/20)	0.25 (5/20)	0.30 (6/20)
8	2019/10/30	0.50 (8/16)	0.25 (4/16)	0.38 (6/16)
9	2010/11/11	0.94 (15/16)	0.56 (9/16)	0.50 (8/16)

[a], [b]: Two-way ANOVA followed by Tukey's multiple comparisons test on disease incidences of different strains indicated the disease incidence of Δ*tssL* and Δ*tssB* was significantly lower than *A. tumefaciens* C58 WT (*p*-value are 0.0225 and 0.0227 respectively). *: Not Determined.

Table 2. Sum of read counts in six tumor samples before and after filtering non-bacterial reads from amplicon sequencing run I for protocol test

16S rRNA regions	Before filtering	Bacterial reads	Percentage of non-bacterial DNA
V3-4	140,830	1,230	99.1%
V5-7	601,299	38,320	93.6%

Table 3 The read count before and after filtering non-bacterial ASVs from the three crown gall DNA samples (named as W21, W22, and W25) induced by *A. tumefaciens* C58 wild type

Sample-id	Dataset-id	Read before filtering	Reads after filtering chloroplast and mitochondrial DNA	Percentage of non-host reads
1115-21W	T21-v1-3-B	176,544	9,914	5.6%
1115-22W	T22-v1-3-B	183,475	5,608	3.1%
1115-25W	T25-v1-3-B	152,387	2,404	1.6%
1115-21W	T21-v1-3-A	185,107	853	0.5%
1115-22W	T22-v1-3-A	247,240	379	0.2%
1115-25W	T25-v1-3-A	178,980	220	0.1%
1115-21W	T21-v3-4-B	239,243	38,462	16.1%
1115-22W	T22-v3-4-B	246,063	19,218	7.8%
1115-25W	T25-v3-4-B	209,991	11,185	5.3%
1115-21W	T21-v3-4-A	200,185	979	0.5%
1115-22W	T22-v3-4-A	205,966	796	0.4%
1115-25W	T25-v3-4-A	193,350	426	0.2%
1115-21W	T21-v5-7-B	129,422	53,715	41.5%
1115-22W	T22-v5-7-B	140,012	29,078	20.8%
1115-25W	T25-v5-7-B	135,692	18,052	13.3%
1115-21W	T21-v5-7-A	150,015	2,792	1.9%
1115-22W	T22-v5-7-A	139,974	1,507	1.1%
1115-25W	T25-v5-7-A	133,035	786	0.6%
1115-21W	T21-v6-8-B	92,167	3,691	4.0%
1115-22W	T22-v6-8-B	92,053	3,197	3.5%
1115-25W	T25-v6-8-B	82,361	2,077	2.5%
1115-21W	T21-v6-8-A	121,145	1,604	1.3%
1115-22W	T22-v6-8-A	138,186	1,273	0.9%
1115-25W	T25-v6-8-A	121,820	708	0.6%
	NC-v3-4-B	1,012	979	96.7%

Figure 1. Tumorigenesis assay of *A. tumefaciens* C58 and its T6SS-deficienct

mutants.

For each experiment, 14-20 wounded tomato seedlings were grown in soil containing

one of the strains tested (i.e., C58 WT, Δ*tssL*, and Δ*tssB*) and collected at 60 days post

inoculation (dpi). (A) Disease incidence plotted by strain and color-coded according

to the month of inoculation (light yellow, June; orange, July; light blue, October; dark

blue, November). Lines and error bars indicate mean ± SD. Statistical significance

was tested using two-way ANOVA followed by Tukey's multiple comparisons; *p* =

0.025 and 0.027 when comparing the wild type to Δ*tssL* and Δ*tssB*, respectively. (B)

Weight distribution of the galls collected. No significant difference among the three

strains (*p* = 0.15, Kruskai-Wallis test). (C) Correlation plot of disease incidence and

temperature of daily average. (D) Crown galls generated through soil inoculation,

example from inoculation on 11/15/2018. N indicated no gall formation in some of

inoculated plants.

Figure 2. Optimizing 16S rRNA gene amplification by using PCR blockers.

Three crown gall samples (named as 21W, 22W and 25W) induced by *A. tumefaciens*

C58 wild type were used for analysis. V1-3, V3-4, V5-7 and V6-8 represent different

primer pairs targeting different variable regions on 16S rRNA gene. (A) The OTU

counts of 16 rRNA gene amplicons after filtering non-bacterial OTUs. (ns, not

statistically significant; *, $p < 0.05$, student's t-test). The bar indicated mean of OTUs

± SD. (B) Alpha rarefaction curves of the observed bacterial OTUs based on datasets

from amplicon sequencing run II. Sample size in X-axis indicates different sub-

sampling depth of each dataset.

Figure 3. Gallobiome composition when using different primers amplified with

or without blockers. (A) Different primer sets (V1-3, V3-4, V5-7 and V6-8**)** for

analysis of three crown gall samples (21W, 22W and 25W, see Table S6) induced by

A. tumefaciens C58 wild type were indicated. Top 10 observed families were listed.

"-": amplification without blockers; "+": amplification with blockers. Principle

coordinate analysis (PCoA) plots of the bacterial composition of the datasets were

generated based on the weighted Unifrac matrix, and color coded by (B) blocker

usage (p-value = 0.068, R^2 = 0.026, ADONIS), (C) different primer sets (p-value =

0.001, R^2 = 0.524, ADONIS) and (D) different crown gall samples (p-value = 0.001,

R^2 = 0.238, ADONIS).

Figure 4. Principle coordinate analysis (PCoA) of bacterial composition in 53 crown galls induced by *A. tumefaciens* C58 WT and T6SS mutants.

The plots were drawn based on weighted Unifrac matrix of the bacterial communities. Crown galls induced by different C58 strains (WT, Δ*tssB,* Δ*tssL*) (A) or inoculated in different months (B) were labeled with different colors in each panel. The datasets were further split based on the month of inoculation, and the PCoA plots of gallobiomes from (C) July, (D) October and (E) November were indicated above. Statistical differences in clustering were evaluated via ADONIS permutation test, and the corresponding *p*-values and R^2 are indicated.

Figure 5. Alpha diversity and composition of crown galls generated in different months by *A. tumefaciens* C58 WT and T6SS mutants.

(A) Observed OTU, (B) Shannon index, (C) evenness, and (D) Faith's phylogenic diversity of each sample generated in different months was indicated. Lines and error bars indicate mean ± SD. Two-way ANOVA followed by Tukey's multiple comparisons test were performed, and the asterisks indicated the statistically

significant difference between two strains (*p*-value < 0.05). (E) Bacterial composition

of 53 crown galls. For each strain, the samples are grouped by the month of

inoculation. Top-ten most abundant families were listed and colored differently, and

the remaining ones are combined into "Others". (F) Relative abundance of

Rhizobiaceae in galls inoculated in different months and strains. (G) Scatter plot of

relative abundance of *A. tumefaciens* OTU in gallobiome - weight of the crown galls.

A positive correlation between weight of crown gall and relative abundance of *A.*

tumefaciens C58 was observed (Spearman r = 0.333, *p* = 0.01).

Figure 6. Analysis of composition of microbiomes (ANCOM) for gallobiomes in

July.

The ANCOM volcano plots were drawn based on the centred-log-ratio-transformed

abundance of (A) OTUs or bacterial families. The taxa that exhibit differential

abundance (DA) between gallobiomes induced by WT and T6SS mutants are

indicated by red arrows. The two DA OTUs belonging to *Sphingomonadaceae* were

named SphinOTU1 and SphinOTU2, respectively. The relative abundance of focal

taxa was shown in panels (B) and (C). Lines and error bars indicate mean ± SD.

Fig. 7 Competition between *A. tumefaciens* C58 and *Sphingomonas* sp. R1 *in vitro* and on wounded site of tomato stems.

Each of the three *A. tumefaciens* strains (i.e., C58 WT, Δ*tssL*, and Δ*tssB*) was

individually tested for competition against *Sphingomonas* sp. R1 isolated from

rhizosphere of tomato. (A) The *in vitro* competition results based on *A. tumefaciens*:

Sphingomonas sp. R1 at 1:1 and 10:1 ratio, and the survival of *Sphingomonas* sp. R1

was plotted; lines and error bars indicate mean ± SD. Based on ANOVA followed by

Tukey's multiple comparison test, C58 significantly reduced *Sphingomonas* sp. R1

CFUs when mixed in 10:1 ratio ($p = 0.0153$) but not in 1:1 ratio ($p = 0.2753$). (B) The

A. tumefaciens alone or co-infected with *Sphingomonas* sp. R1 at 1:1 ratio in soil

were recovered from the surface of wounded stem segments at 10 dpi, the CFU of *A.*

tumefaciens and *Sphingomonas* sp. R1 were plotted. T-test indicated the CFU

differences of *Sphingomonas* sp. R1 inoculation alone or co-infection with T6SS

mutants as compared to WT with indicated *p* values. ANOVA test indicated no

significant difference among means ($p = 0.0945$ and 0.0551 for *A. tumefaciens* and

Sphingomonas sp. R1 CFUs, respectively). (C) The colonies of *A. tumefaciens* strains

C58 WT, Δ*tssL*, and Δ*tssB* recovered from the surface of wounded stem segments

following the soil inoculation procedure were plotted. Different symbols indicated the

outcome from different batches of colonization assay. F-test indicated the variance of

Δ*tssL* and Δ*tssB* were significantly different when compared to C58 WT ($F =$

0.000467 and $8.04398E-05$, respectively). Brown-Forsythe and Welch ANOVA test

indicated no significant difference among means ($p = 0.4847$). Line indicates median.

Figure S1. DNA sequence alignment of 16S rRNA genes of selected bacterial strains and tomato KnowYou 301 chloroplast and mitochondria.

The 16S rRNA gene sequences of chloroplasts and mitochondria in *Solanum lycopersicum* (tomato) cultivar Known-You 301 were obtained from this study. All of the bacterial 16S rRNA gene sequences were accessed from Reference Sequence (RefSeq) database in National Center of Biotechnology Information (NCBI). The species, strain name, and accession number of 16S rRNA gene were listed at on the

left of each sequence. The alignment of sequences mentioned above were aligned in

MEGAX via ClustalW multiple alignment with gap opening penalty 15.00 and gap

extension penalty 6.66 (default). The levels of shaded blue color reflect the degree of

identity. The regions of each primer and blocker were underlined. The sequences

highlighted by red, black, yellow and green are the annealing region of primer sets for

V1-3, V3-4, V5-7 and V6-8 in 16S rRNA gene, respectively. And the gray framed

sequences in tomato chloroplast and mitochondrial 16S rRNA genes are the blockers

for cognate primers.

Figure S2. Alpha rarefaction curves of the observed bacterial OTUs based on

amplicon sequencing of 53 gall samples in sequencing run III.

121   The rarefaction curves were plotted after clustering DADA2-output ASVs into 99%

122   OTUs. Sample size in X-axis indicates different sub-sampling depth of each dataset;

123   Species in Y-axis indicates the number of observed OTUs under certain sub-sampling

124   depth. Label of each curves indicates the sample ID in metadata.

Figure S3. Competition of *A. tumefaciens* and *Sphingomonas* sp. R1 in tomato

gall on stem at 28 dpi.

Each of the three *A. tumefaciens* strains (i.e., C58 WT, Δ*tssL*, and Δ*tssB*) and R1 were

mixed at 1:1 ratio then inoculated on wounded tomato stem. The galls were harvested

and homogenized for plating at 28 dpi. *A. tumefaciens* C58 strains and *Sphingomonas*

sp. R1 were recover on 523 medium plates containing proper antibiotics. R1 could

only be recovered from galls in batch 3. The *p*-value of ANOVA against CFU

numbers of C58 for each batch is 0.786, 0.0023 and 0.230; the *p*-value of ANOVA

against CFU numbers of R1 was 0.6413. The significance of CFU numbers of C58 in

second batch is due to the low CFU counts of Δ*tssL* single inoculation group, which

could not be replicated in other two batches.

Table S1. 16S rRNA gene primer sets used in this study

Targeted Region	Name	Sequence (5'-3')	Size of bacterial amplicons (bp)	Size of tomato amplicons (bp)*
Round I				
V3-V4	341F	CCTACGGGNGGCWGCAG	444	396 (cp); 395 (mt)
	785R	GACTACHVGGGTATCTAATCC		
V5-V7	799F	AACMGGATTAGATACCCKG	394	376(cp); 742(mt)
	1193R	ACGTCATCCCCACCTTCC		
Round II				
V1-V3	68F	TNANACATGCAAGTCGRRCG	425	440 (cp); 474(mt)
	518R	WTTACCGCGGCTGCTGG		
V3-V4	**341F**	CCTACGGGNGGCWGCAG	429	408 (cp); 391(mt)
	688R	CGCTTTCGHDCCTCAGYGTCA	405	
V5-V7	**819F**	GTCCACVCCSTAAACGWTG	553	551 (cp); 915(mt)
	1276R	RCGATTACTAGCGAHTCC		
V6-V8	**895F**	CRCCTGGGGAGTRCRG	530	527 (cp); 894 (mt)
	1391R	GACGGGCGGTGTGTRCA		

*cp, chloroplast; mt, mitochondrion

Table S2. Corresponding blockers (3' modified oligonucleotides with C3 spacer)

Targeted Region	Name*	Sequence	Tm (℃)
V1	68f_blocker_mt	GTCGAACGTTGTTTTCGGGGAG	62
V1	68f_blocker_cp	GTCGGACGGGAAACACG	61
V4	688ra_blocker_mt	CGTCGGTAGGGACCCAGAGAGCT	69
V4	688ra_blocker_cp	TGTCAGTGTCGGCCCAGCAGAGT	69
V5	819f_blocker_mt	ACGATGAGTGTTCGCCCTTG	61
V5	819f_blocker_cp	AAACGATGGATACTAGGTGCTGT	60
V6	895f_blocker_mt	AGTACGGTCGCAAGACCG	61
V6	895f_blocker_cp	GGAGTACGTTCGCAAGAATG	60

* cp, chloroplast; mt, mitochondrion

Table S3. Bacteria strains and plasmid in this study

Strain/plasmid	Characteristics	EML No.	Reference/ Source
Escherichia coli **DH10B**	Host for DNA cloning	455	Invitrogen
Agrobacterium tumefaciens **C58**	Wild-type, virulent strain containing nonpaline-type Ti-plasimid, pTiC58	530	Eugene Nester [1]
	$\Delta tssL$, deletion mutant of *tssL* gene (Atu4333)	1073	[2]
	$\Delta tssB$, deletion mutant of *tssB* gene (Atu4342)	1109	[2]
	Gm^R-GFP knocked-in chromosome	4094	This study
	$\Delta tssL$, Gm^R-GFP knocked-in chromosome	4097	This study
	$\Delta tssB$, Gm^R-GFP knocked-in chromosome	4100	This study
Sphingomonas sp. **R1**	Wild type, isolated from tomato rhizosphere	4107	This study
pJQ-COM	Gm^R, for generating DNA knock-in adjacent to gene *ActC* in C58	2236	[3]
pJQ-com-Gm^r GFP	Gm^R, for generate Gm^R and GFP knocked-in C58	4091	This study
pRL662::GFP (S65T)	Gm^R, constitutively expressing GFP (S56T)	3375	This study
pBBR1MCS2-mCherry	Km^R, constitutively expressing mCherry	3022	Stanton B. Gelvin

Table S4. Primers for plasmid construction

Name	Sequence	Reference
BclI-GFP-GmR-F-25	AAAAATGATCAGTGAGCGCGCGTAATA CGACTCAC	This study
BclI-GFP-GmR-R-25	AAAAATGATCAGGGTACCGAGCTCGAA TTGACATAAG	This study
3'sacB-5'GmR pJQ200 F	TGCGCCAAGCTTCCTGCTGAACATC	This study
3'sacB-5'GmR pJQ200 R	TTGAGCAGCCGCGTAGTGAGATCTA	This study

Table S5. Condition of 16S rRNA gene amplification

1st PCR Amplification v1-3

Recipe A		Program			Recipe B		Program		
input 25 ng	6.5	95℃ 5 min	1 cycle		input 25 ng	6.5	95℃ 5 min	1 cycle	
H2O		98℃ 30 sec			H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	12.5	55℃ 30 sec	12cycle		2X Kapa HiFi PCR mix	12.5	55℃ 30 sec	15cycle	
68f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec			68f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec		
518r primer 10uM (final 0.2uM)	0.5	72℃ 5 min	1 cycle		518r primer 10uM (final 0.2uM)	0.5	72℃ 5 min	1 cycle	
H2O	5	4℃ ∞	1 cycle		68f_blocking_mito 20uM	2.5	4℃ ∞	1 cycle	
					518r_blocking_chloro_20uM	2.5			
Total	25				Total	25			
Beads purificatin, 1:1					Beads purificatin, 1:1				

2nd Library Amplification

Recipe		Program		
1st purified PCR products	20	98℃ 2 min	1 cycle	
H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	25	60℃ 30 sec	5 cycle	
Truseq-HT-D5xx_F (10uM)	2.5	72℃ 60 sec		
Truseq-HT-D7xx_R (10uM)	2.5	72℃ 5 min	1 cycle	
Total	50	10℃ ∞	1 cycle	
25uL/tube				
Beads purificatin, 1:1				

1st PCR Amplification v3-4

Recipe A		Program			Recipe B		Program		
input 25 ng	6.5	95℃ 5 min	1 cycle		input 25 ng	6.5	95℃ 5 min	1 cycle	
H2O		98℃ 30 sec			H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	12.5	56℃ 30 sec	18cycle		2X Kapa HiFi PCR mix	12.5	56℃ 30 sec	18 cycle	
341f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec			341f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec		
688ar primer 10uM (final 0.2uM)	0.5	72℃ 5 min	1 cycle		688ar primer 10uM (final 0.2uf)	0.5	72℃ 5 min	1 cycle	
H2O	5	4℃ ∞	1 cycle		688ar_blocking_mito 20uM	2.5	4℃ ∞	1 cycle	
					688ar_blocking_chloro_20uM	2.5			
Total	25				Total	25			
Beads purificatin, 1:1					Beads purificatin, 1:1				

2nd Library Amplification

Recipe		Program		
1st purified PCR products	20	98℃ 2 min	1 cycle	
H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	25	60℃ 30 sec	5 cycle (A) / 10 cycle (B)	
Truseq-HT-D5xx_F (10uM)	2.5	72℃ 60 sec		
Truseq-HT-D7xx_R (10uM)	2.5	72℃ 5 min	1 cycle	
Total	50	10℃ ∞	1 cycle	
25uL/tube				
Beads purificatin, 1:1				

1st PCR Amplification v5-7

Recipe A		Program			Recipe B		Program		
input 25 ng	6.5	95℃ 5 min	1 cycle		input 25 ng	6.5	95℃ 5 min	1 cycle	
H2O		98℃ 30 sec			H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	12.5	56℃ 30 sec	20cycle		2X Kapa HiFi PCR mix	12.5	56℃ 30 sec	24cycle	
819f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec			819f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec		
1276r primer 10uM (final 0.2uM)	0.5	72℃ 5 min	1 cycle		1276r primer 10uM (final 0.2uf)	0.5	72℃ 5 min	1 cycle	
H2O	5	4℃ ∞	1 cycle		819f_blocking_mito 20uM	2.5	4℃ ∞	1 cycle	
					819f_blocking_chloro_20uM	2.5			
Total	25				Total	25			
Beads purificatin, 1:1					Beads purificatin, 1:1				

2nd Library Amplification

Recipe		Program		
1st purified PCR products	20	98℃ 2 min	1 cycle	
H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	25	60℃ 30 sec	5 cycle	
Truseq-HT-D5xx_F (10uM)	2.5	72℃ 60 sec		
Truseq-HT-D7xx_R (10uM)	2.5	72℃ 5 min	1 cycle	
Total	50	10℃ forever	1 cycle	
25uL/tube				
Beads purificatin, 1:1				

1st PCR Amplification v6-8

Recipe A		Program			Recipe B		Program		
input 25 ng	6.5	95℃ 5 min	1 cycle		input 25 ng	6.5	95℃ 5 min	1 cycle	
H2O		98℃ 30 sec			H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	12.5	50℃ 30 sec	11cycle		2X Kapa HiFi PCR mix	12.5	50℃ 30 sec	12cycle	
895f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec			895f primer 10uM (final 0.2uM)	0.5	72℃ 60 sec		
1391r primer 10uM (final 0.2uM)	0.5	72℃ 5 min	1 cycle		1391r primer 10uM (final 0.2uf)	0.5	72℃ 5 min	1 cycle	
H2O	5	4℃ ∞	1 cycle		895f_blocking_mito 20uM	2.5	4℃ ∞	1 cycle	
					895f_blocking_chloro_20uM	2.5			
Total	25				Total	25			
Beads purificatin, 1:1					Beads purificatin, 1:1				

2nd Library Amplification

Recipe		Program		
1st purified PCR products	20	98℃ 2 min	1 cycle	
H2O		98℃ 30 sec		
2X Kapa HiFi PCR mix	25	60℃ 30 sec	5 cycle	
Truseq-HT-D5xx_F (10uM)	2.5	72℃ 60 sec		
Truseq-HT-D7xx_R (10uM)	2.5	72℃ 5 min	1 cycle	
Total	50	10℃ forever	1 cycle	
25uL/tube				
Beads purificatin, 1:1				

Table S6. Metadata of harvested crown galls

#SampleID	Strain*	Year of inoculation	Month od inoculation	Day of inoculation	Year of harvest	Month of harvest	Day of harvest	Weight (g)
1115-11L	Δ*tssL*	2018	11	15	2019	1	18	1.80
1115-12L[a]	Δ*tssL*	2018	11	15	2019	1	18	0.81
1115-13L[a]	Δ*tssL*	2018	11	15	2019	1	18	0.62
1115-14L[a]	Δ*tssL*	2018	11	15	2019	1	18	0.96
1115-15L	Δ*tssL*	2018	11	15	2019	1	18	0.40
1115-16L	Δ*tssL*	2018	11	15	2019	1	18	0.40
1115-17L	Δ*tssL*	2018	11	15	2019	1	18	0.07
1115-18L	Δ*tssL*	2018	11	15	2019	1	18	0.16
1115-19W	WT	2018	11	15	2019	1	18	2.00
1115-20W	WT	2018	11	15	2019	1	18	1.27
1115-21W[b]	WT	2018	11	15	2019	1	18	0.56
1115-22W[b]	WT	2018	11	15	2019	1	18	0.56
1115-23W	WT	2018	11	15	2019	1	18	0.42
1115-24W	WT	2018	11	15	2019	1	18	0.37
1115-25W[b]	WT	2018	11	15	2019	1	18	0.82
1115-26W	WT	2018	11	15	2019	1	18	0.22
1115-27W	WT	2018	11	15	2019	1	18	0.10
1115-28W	WT	2018	11	15	2019	1	18	0.08
1115-29W	WT	2018	11	15	2019	1	18	0.05
1115-30W	WT	2018	11	15	2019	1	18	0.02
0704-1W	WT	2019	7	4	2019	9	2	0.12
0704-2W	WT	2019	7	4	2019	9	2	0.12
0704-3W	WT	2019	7	4	2019	9	2	0.09
0704-4W	WT	2019	7	4	2019	9	2	0.06
0704-5W	WT	2019	7	4	2019	9	2	0.11
0704-6W	WT	2019	7	4	2019	9	2	0.14
0704-7W	WT	2019	7	4	2019	9	2	0.27
0704-1L	Δ*tssL*	2019	7	4	2019	9	2	0.10
0704-2L	Δ*tssL*	2019	7	4	2019	9	2	0.34
0704-3L	Δ*tssL*	2019	7	4	2019	9	2	0.12
0704-4L	Δ*tssL*	2019	7	4	2019	9	2	0.07
0704-5L	Δ*tssL*	2019	7	4	2019	9	2	0.06
0704-6L	Δ*tssL*	2019	7	4	2019	9	2	0.10
0704-7L	Δ*tssL*	2019	7	4	2019	9	2	0.13

#SampleID	Strain*	Year of inoculation	Month od inoculation	Day of inoculation	Year of harvest	Month of harvest	Day of harvest	Weight (g)
0704-1B	ΔtssB	2019	7	4	2019	9	2	0.11
0704-2B	ΔtssB	2019	7	4	2019	9	2	0.28
0704-3B	ΔtssB	2019	7	4	2019	9	2	0.07
0704-4B	ΔtssB	2019	7	4	2019	9	2	0.22
0704-5B	ΔtssB	2019	7	4	2019	9	2	0.06
0722-1W	WT	2019	7	22	2019	9	28	0.03
0722-2W	WT	2019	7	22	2019	9	28	0.04
0722-3W	WT	2019	7	22	2019	9	28	0.07
0722-4W	WT	2019	7	22	2019	9	28	0.05
0722-5W	WT	2019	7	22	2019	9	28	0.13
0722-6W	WT	2019	7	22	2019	9	28	0.05
0722-1B	ΔtssB	2019	7	22	2019	9	28	0.01
0722-2B	ΔtssB	2019	7	22	2019	9	28	0.01
0729-1W	WT	2019	7	29	2019	9	29	0.03
0729-2W	WT	2019	7	29	2019	9	29	0.05
0729-3W	WT	2019	7	29	2019	9	29	0.04
0729-4W	WT	2019	7	29	2019	9	29	0.05
0729-5W	WT	2019	7	29	2019	9	29	0.04
0729-6W	WT	2019	7	29	2019	9	29	0.01
0729-7W	WT	2019	7	29	2019	9	29	0.05
0729-8W	WT	2019	7	29	2019	9	29	0.08
0729-9W	WT	2019	7	29	2019	9	29	0.03
0729-1L	ΔtssL	2019	7	29	2019	9	29	0.01
0729-2L	ΔtssL	2019	7	29	2019	9	29	0.05
0729-3L	ΔtssL	2019	7	29	2019	9	29	0.08
0729-4L	ΔtssL	2019	7	29	2019	9	29	0.03
0729-1B	ΔtssB	2019	7	29	2019	9	29	0.09
0729-2B	ΔtssB	2019	7	29	2019	9	29	0.07
1005-1W	WT	2019	10	5	2019	12	13	0.27
1005-2W	WT	2019	10	5	2019	12	13	0.16
1005-3W	WT	2019	10	5	2019	12	13	0.09
1005-4W	WT	2019	10	5	2019	12	13	0.04
1005-5W	WT	2019	10	5	2019	12	13	0.01
1005-6W	WT	2019	10	5	2019	12	13	0.09
1005-7W	WT	2019	10	5	2019	12	13	0.01
#SampleID	Strain*	Year of inoculation	Month od inoculation	Day of inoculation	Year of harvest	Month of harvest	Day of harvest	Weight (g)

#SampleID	Strain*	Year of inoculation	Month od inoculation	Day of inoculation	Year of harvest	Month of harvest	Day of harvest	Weight (g)
1005-1L	ΔtssL	2019	10	5	2019	12	13	0.23
1005-2L	ΔtssL	2019	10	5	2019	12	13	0.32
1005-3L	ΔtssL	2019	10	5	2019	12	13	0.09
1005-4L	ΔtssL	2019	10	5	2019	12	13	0.04
1005-5L	ΔtssL	2019	10	5	2019	12	13	0.01
1005-1B	ΔtssB	2019	10	5	2019	12	13	0.23
1005-2B	ΔtssB	2019	10	5	2019	12	13	0.56
1005-3B	ΔtssB	2019	10	5	2019	12	13	0.16
1005-4B	ΔtssB	2019	10	5	2019	12	13	0.05
1005-5B	ΔtssB	2019	10	5	2019	12	13	0.06
1005-6B	ΔtssB	2019	10	5	2019	12	13	0.03
1005-7B	ΔtssB	2019	10	5	2019	12	13	0.06
1030-1W	WT	2019	10	30	2020	1	9	0.34
1030-2W	WT	2019	10	30	2020	1	9	0.12
1030-3W	WT	2019	10	30	2020	1	9	0.05
1030-4W	WT	2019	10	30	2020	1	9	0.03
1030-5W	WT	2019	10	30	2020	1	9	0.05
1030-6W	WT	2019	10	30	2020	1	9	0.11
1030-7W	WT	2019	10	30	2020	1	9	0.02
1030-8W	WT	2019	10	30	2020	1	9	0.01
1030-1L	ΔtssL	2019	10	30	2020	1	9	0.50
1030-2L	ΔtssL	2019	10	30	2020	1	9	0.07
1030-3L	ΔtssL	2019	10	30	2020	1	9	0.03
1030-4L	ΔtssL	2019	10	30	2020	1	9	0.01
1030-1B	ΔtssB	2019	10	30	2020	1	9	0.17
1030-2B	ΔtssB	2019	10	30	2020	1	9	0.06
1030-3B	ΔtssB	2019	10	30	2020	1	9	0.08
1030-4B	ΔtssB	2019	10	30	2020	1	9	0.00
1030-5B	ΔtssB	2019	10	30	2020	1	9	0.02
1030-6B	ΔtssB	2019	10	30	2020	1	9	0.01
1111-1W	WT	2019	11	11	2020	1	14	0.26
1111-2W	WT	2019	11	11	2020	1	14	0.19
1111-2.1W	WT	2019	11	11	2020	1	14	0.04
1111-3W	WT	2019	11	11	2020	1	14	0.25
1111-3.1W	WT	2019	11	11	2020	1	14	0.04

#SampleID	Strain*	Year of inoculation	Month od inoculation	Day of inoculation	Year of harvest	Month of harvest	Day of harvest	Weight (g)
1111-4W	WT	2019	11	11	2020	1	14	0.13
1111-5W	WT	2019	11	11	2020	1	14	0.01
1111-6W	WT	2019	11	11	2020	1	14	0.04
1111-7W	WT	2019	11	11	2020	1	14	0.05
1111-8W	WT	2019	11	11	2020	1	14	0.11
1111-8.1W	WT	2019	11	11	2020	1	14	0.09
1111-8.2W	WT	2019	11	11	2020	1	14	0.09
1111-9W	WT	2019	11	11	2020	1	14	0.02
1111-10W	WT	2019	11	11	2020	1	14	0.07
1111-11W	WT	2019	11	11	2020	1	14	0.02
1111-12W	WT	2019	11	11	2020	1	14	0.04
1111-13W	WT	2019	11	11	2020	1	14	0.27
1111-13.1W	WT	2019	11	11	2020	1	14	0.18
1111-13.2W	WT	2019	11	11	2020	1	14	0.12
1111-14W	WT	2019	11	11	2020	1	14	0.04
1111-15W	WT	2019	11	11	2020	1	14	0.01
1111-1L	ΔtssL	2019	11	11	2020	1	14	0.21
1111-2L	ΔtssL	2019	11	11	2020	1	14	0.04
1111-2.1L	ΔtssL	2019	11	11	2020	1	14	0.02
1111-3L	ΔtssL	2019	11	11	2020	1	14	0.07
1111-3.1L	ΔtssL	2019	11	11	2020	1	14	0.03
1111-4L	ΔtssL	2019	11	11	2020	1	14	0.03
1111-5L	ΔtssL	2019	11	11	2020	1	14	0.08
1111-6L	ΔtssL	2019	11	11	2020	1	14	0.08
1111-7L	ΔtssL	2019	11	11	2020	1	14	0.14
1111-8L	ΔtssL	2019	11	11	2020	1	14	0.01
1111-9L	ΔtssL	2019	11	11	2020	1	14	0.03
1111-1B	ΔtssB	2019	11	11	2020	1	14	0.18
1111-2B	ΔtssB	2019	11	11	2020	1	14	0.09
1111-3B	ΔtssB	2019	11	11	2020	1	14	0.03
1111-4B	ΔtssB	2019	11	11	2020	1	14	0.03
1111-5B	ΔtssB	2019	11	11	2020	1	14	0.02
1111-6B	ΔtssB	2019	11	11	2020	1	14	0.03

#SampleID	Strain*	Year of inoculation	Month od inoculation	Day of inoculation	Year of harvest	Month of harvest	Day of harvest	Weight (g)
1111-7B	Δ*tssB*	2019	11	11	2020	1	14	0.01
1111-8B	Δ*tssB*	2019	11	11	2020	1	14	0.01

* *Agrobacterium tumefaciens* C58 wilt type.

[a] Used in round I sequencing.

[b] Used in both round I and II sequencing.

Table S7. Crown gall metadata used for analysis of amplicon sequencing round III

#SampleID	Strain	Year_of_inoculation	Month_of_inoculation	Weight	Seasons	Trials	Bactch
#q2:types	categorical	categorical	categorical	numeric	categorical	categorical	categorical
1115-L12	tssL	2018	Nov	0.81	Winter	181115	Batch 1
1115-L13	tssL	2018	Nov	0.62	Winter	181115	Batch 1
1115-L14	tssL	2018	Nov	0.96	Winter	181115	Batch 1
1115-W21	C58 WT	2018	Nov	0.56	Winter	181115	Batch 1
1115-W22	C58 WT	2018	Nov	0.56	Winter	181115	Batch 1
1115-W25	C58 WT	2018	Nov	0.82	Winter	181115	Batch 1
0704-W1	C58 WT	2019	Jul	0.12	Summer	190704	Batch 4
0704-W2	C58 WT	2019	Jul	0.12	Summer	190704	Batch 4
0704-W3	C58 WT	2019	Jul	0.09	Summer	190704	Batch 4
0704-W4	C58 WT	2019	Jul	0.06	Summer	190704	Batch 4
0704-W5	C58 WT	2019	Jul	0.11	Summer	190704	Batch 4
0704-W6	C58 WT	2019	Jul	0.14	Summer	190704	Batch 4
0704-W7	C58 WT	2019	Jul	0.27	Summer	190704	Batch 4
0704-L1	tssL	2019	Jul	0.1	Summer	190704	Batch 4
0704-L2	tssL	2019	Jul	0.34	Summer	190704	Batch 4
0704-L3	tssL	2019	Jul	0.12	Summer	190704	Batch 4
0704-B1	tssB	2019	Jul	0.11	Summer	190704	Batch 4
0704-B2	tssB	2019	Jul	0.28	Summer	190704	Batch 4
0704-B3	tssB	2019	Jul	0.07	Summer	190704	Batch 4
1005-W1	C58 WT	2019	Oct	0.27	Summer	191005	Batch 7
1005-W2	C58 WT	2019	Oct	0.16	Summer	191005	Batch 7
1005-W3	C58 WT	2019	Oct	0.09	Summer	191005	Batch 7
1005-W6	C58 WT	2019	Oct	0.09	Summer	191005	Batch 7
1005-L1	tssL	2019	Oct	0.23	Summer	191005	Batch 7
1005-L2	tssL	2019	Oct	0.32	Summer	191005	Batch 7
1005-L3	tssL	2019	Oct	0.09	Summer	191005	Batch 7
1005-B1	tssB	2019	Oct	0.23	Summer	191005	Batch 7
1005-B2	tssB	2019	Oct	0.56	Summer	191005	Batch 7
1005-B3	tssB	2019	Oct	0.16	Summer	191005	Batch 7
1005-B5	tssB	2019	Oct	0.06	Summer	191005	Batch 7
1005-B7	tssB	2019	Oct	0.06	Summer	191005	Batch 7
1030-W1	C58 WT	2019	Oct	0.34	Winter	191030	Batch 8
1030-W2	C58 WT	2019	Oct	0.12	Winter	191030	Batch 8
1030-W6	C58 WT	2019	Oct	0.11	Winter	191030	Batch 8
1030-L1	tssL	2019	Oct	0.5	Winter	191030	Batch 8

#SampleID	Strain	Year_of_inoculation	Month_of_inoculation	Weight	Seasons	Trials	Bactch
1030-L2	tssL	2019	Oct	0.07	Winter	191030	Batch 8
1030-B1	tssB	2019	Oct	0.17	Winter	191030	Batch 8
1030-B2	tssB	2019	Oct	0.06	Winter	191030	Batch 8
1030-B3	tssB	2019	Oct	0.08	Winter	191030	Batch 8
1111-W1	C58 WT	2019	Nov	0.26	Winter	191111	Batch 9
1111-W2	C58 WT	2019	Nov	0.19	Winter	191111	Batch 9
1111-W3	C58 WT	2019	Nov	0.25	Winter	191111	Batch 9
1111-W4	C58 WT	2019	Nov	0.13	Winter	191111	Batch 9
1111-W8	C58 WT	2019	Nov	0.11	Winter	191111	Batch 9
1111-W10	C58 WT	2019	Nov	0.07	Winter	191111	Batch 9
1111-W13	C58 WT	2019	Nov	0.27	Winter	191111	Batch 9
1111-L1	tssL	2019	Nov	0.21	Winter	191111	Batch 9
1111-L3	tssL	2019	Nov	0.07	Winter	191111	Batch 9
1111-L5	tssL	2019	Nov	0.08	Winter	191111	Batch 9
1111-L6	tssL	2019	Nov	0.08	Winter	191111	Batch 9
1111-L7	tssL	2019	Nov	0.14	Winter	191111	Batch 9
1111-B1	tssB	2019	Nov	0.18	Winter	191111	Batch 9
1111-B2	tssB	2019	Nov	0.09	Winter	191111	Batch 9

References

1. Wood, D.W., et al., The genome of the natural genetic engineer *Agrobacterium tumefaciens* C58. Science, 2001. **294**(5550): p. 2317-23.

2. Lin, J.S., L.S. Ma, and E.M. Lai, Systematic Dissection of the *Agrobacterium* Type VI Secretion System Reveals Machinery and Secreted Components for Subcomplex Formation. PLoS One, 2013. **8**(7): p. e67647.

3. Liu, A.C., et al., A citrate-inducible gene, encoding a putative tricarboxylate transporter, is downregulated by the organic solvent DMSO in *Agrobacterium tumefaciens*. J Appl Microbiol, 2008. **105**(5): p. 1372-1383.