Diverse genetic architectures on the Z chromosome underlie the two rules of speciation in *Papilio* butterfly hybrids

Tianzhu Xiong[1,*], Shreeharsha Tarikere[1,4], Neil Rosser[1], Xueyan Li[2], Masaya Yago[3], and James Mallet[1]

[1]*Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA*
[2]*State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan, China*
[3]*The University Museum, The University of Tokyo, Tokyo, Japan*
[4]*Current Address: Department of Earth Sciences, Carleton University, Ottawa, ON, Canada*
[*]*Corresponding Author:* `txiong@g.harvard.edu`

October 28, 2022

Abstract

Two empirical rules arise from the incompatibility of interspecific hybrids: Haldane's Rule predicts that the chromosomally heterogametic sex (XY or ZW) is more unfit after hybridization; the large-X/Z effect posits that sex chromosomes play a major role in incompatibility. Classical theories on these two rules rely on evidence mainly from taxa with male heterogamety, while female heterogamety received little investigation. Here, we reveal the genetic architectures of the two rules in hybrids between the butterflies *Papilio bianor* and *Papilio dehaanii*, where the female is the heterogametic sex. In these crosses, hybrid females suffer from both body size abnormality and ovary dysgenesis, while males appear normal and fertile. Curiously, abnormal size in females is mapped to a continuum of Z-linked polygenes, each acting quantitatively with small phenotypic effects. This polygenic system, perhaps spanning the entire Z chromosome, also correctly predicts weaker incompatibility effects in males. For ovary dysgenesis, the underlying genetic architecture can be monogenic or polygenic with different maternal backgrounds. Most peculiarly, when comparing ovary dysgenesis in certain maternal backgrounds between *Papilio* and *Heliconius*, we find that F_1 recombination on the Z chromosome often rescues incompatibilities among backcross individuals, while a non-recombined Z chromosome almost always produces strong ovary defects regardless of ancestry. These results suggest that high fitness in these maternal backgrounds requires a balance between the total quantities of introgression on autosomes and the Z chromosome. Our study highlights that, in addition to incompatibility factors with large effects, genomically dispersed polygenes are also abundant in creating butterfly reproductive isolation.

1. Introduction

Speciation is a complex and a stochastic process, yet it obeys empirical rules across sexually reproducing lineages [1]. One such rule (Haldane's Rule) states that among the offspring between different species, the sex with a greater fitness cost is the heterogametic sex [2, 3]. Another rule of speciation is the so-called large-X/Z effect on hybrid incompatibility [1, 4]. Haldane's Rule is entirely phenomenological, but it holds firmly across many phylogenetically diverse organisms [5, 6, 7]. Whether its wide pertinence emerges from a common set of genetic mechanisms is an open question [8]. The large-X/Z effect has also been suggested as a robust signature, but its evidence is less direct unless incompatibility factors can be genetically mapped [9, 10, 11, 12, 13].

The most compelling evidence on the genetic and molecular basis of Haldane's Rule comes from studies in mammals and *Drosophila*, both of which have X/Y sex determination, and in which hybrid males develop more incompatibility [14, 15]. These findings converge upon several genetic mechanisms of Haldane's Rule: First, dominance theory posits that the single X chromosome in males might expose recessive genes that are deleterious in a hybrid genetic background, thus increasing the likelihood of incompatibility [4, 16]. Second, the heterogametic sex might speed up the evolution of sex chromosomes, and Haldane's Rule can be produced from potentially a combination of processes involving haploid selection (faster-X theory) [17], sex-specific selection (faster-male theory) [18], and sex chromosome conflict (meiotic drive theory) [19, 20]. Accelerated sex chromosome evolution also provides a natural explanation for the large-X/Z effect. However, as these mechanisms have direct evidence mostly from male-heterogametic taxa, whether they offer a general explanation for the two rules of speciation or are specific to each system is unknown. For instance, it has been suggested that the intrinsic sensitivity of spermatogenesis to hybrid disruption can explain a higher incidence of male sterility [21], but this is not applicable to female-heterogametic taxa.

In Lepidoptera (butterflies and moths), the female is the heterogametic sex (with Z/W sex chromosomes), and hybrid females are more prone to abnormal phenotypes than males [22]. Knowing the genetic basis of this direction of the two rules will allow us to test if proposed genetic mechanisms based on male-heterogamety can be extended to explain the same phenomenon in female-heterogametic taxa. To date, little is known about the genomic basis of hybrid incompatibilities in Lepidoptera, except for a few studies using sparse genetic markers [23, 24, 25] and one recent whole-genome QTL study of *Heliconius* ovary dysgenesis [26]. Nonetheless, these studies robustly show that disruption is often linked to the Z chromosome, conforming to the large-X/Z effect. Here, we investigate hybrid incompatibility between a pair of Asian swallowtail butterflies, *Papilio bianor* and *Papilio dehaanii* that produce abnormal females upon hybridization. The two focal species are closely related and are almost parapatric in NW China (Fig. 1A). All interspecific crosses in this species complex, including the island species *P. ryukyuensis*, produce fertile males but sterile females [27, 28]. In addition, abnormal body size is also frequently observed in hybrids among related *Papilio* species [29]. To test for the basis of hybrid incompatibility in this system, we investigate the genetic archi-

Figure 1: The study system. **(A)** The distribution and species relationship of the *Papilio bianor* complex. Scale bar=1000km. **(B)** Top: Pupal weight variation among pure species, F_1 hybrids, and backcross individuals. Horizontal bars in each category represent the mean and its standard error, while the vertical line represents the mean +/-standard deviation. Bottom: changes in the mean pupal weight between F_1 and pure species in each sex. Error bars are standard errors of $\langle W_{F1} - W_{pure} \rangle$. We test if changes are significantly different between males and females of the same comparison, and report its p-value by calculating Z-score. **(C)** The cross scheme. **(D)** The crossover distribution in F_1 males. Chromosome 14 is excluded due to intrachromosomal assembly problems. **(E)** The cumulative distribution of recombination breakpoints along the Z chromosome is indistinguishable from a uniform distribution (p-value is from a Kolmogorov-Smirnov test).

tecture underlying two important traits: body size (pupal weight) and female reproduction (ovary dysgenesis).

2. Results and Discussion

2.1 Haldane's Rule between *P. bianor* and *P. dehaanii* involves asymmetrically inherited elements

To investigate phenotypic abnormalities among interspecific hybrids, we produced reciprocal F_1 crosses between the two species, and then constructed three types of backcrosses by mating F_1 males with pure females (Fig. 1C). Cross notations follow the order (female × male). For instance, "B(BD)" is equivalent to "*bianor* ♀× (*bianor* ♀× *dehaanii* ♂) ♂", where "B" and "D" stand for *bianor* and *dehaanii*, respectively.

To study body size, pupal weight (denoted as W) within three days of pupation is treated as a proxy for size. We find that F_1 females are significantly smaller than pure females from both species when the mother is *dehaanii* (cross direction DB), but they are within the range of pure females if the mother is *bianor* (cross direction BD) (Fig. 1B, top). For F_1 males, the change in pupal weight (compared to pure males) is smaller or equal to that of F_1 females (Fig. 1B, bottom). This result suggests that abnormal size in F_1 hybrids is much more prominent in females. We therefore interpret female-biased F_1 abnormal size in a broad sense as an example of Haldane's Rule. Since reciprocal crosses affect size in opposite ways, asymmetrically inherited genetic elements must be involved.

To study female reproduction in hybrids, we dissected ovaries across the entire pedigree and determined all major ovary phenotypes (Fig. 2). To our surprise, while F_1 females with a *bianor* mother have almost empty ovaries and cannot lay eggs (Fig. 2C,J), F_1 females in the reciprocal cross develop and lay superficially normal eggs (Fig. 2B,I). However, these eggs do not hatch even when the female is mated, and the distribution of yolk in a follicle is also slightly different from that of control (compare sheath-removed figures in Fig. 2A and Fig. 2B). As we only determine ovary phenotypes rather than female fertility *per se*, ovaries with regularly spaced and spherical follicles are subsequently all classified as "Normal", regardless of actual fertility. This is a reasonable decision as variation in ovary phenotype is often large and can be easily scored under a microscope (Fig. 2A-L). Overall, ovary phenotypes are strongly asymmetric between the reciprocal crosses, which suggests the involvement of asymmetrically inherited genetic elements in Haldane's Rule of this system. For this reason, we map phenotypes conditioning on a fixed maternal background.

2.2 Male meiotic crossover on the Z chromosome is obligatory, spatially uniform, and with strong interference

Recombination determines the ancestry of a hybrid chromosome and is crucial to our subsequent analysis. To infer haplotypes and crossover patterns across the pedigree, we perform whole-genome low-coverage sequencing in backcross individuals ($\sim$1×), while F_1 and parents are sequenced to higher depths (>5× and >30×, respectively). Prior to crossover analysis, we use linkage information from all families to manually correct assembly errors in the original reference genome of *P. bianor* [30]

Figure 2: Ovary phenotypes. Monochrome confocal images for each channel are in Fig. S1 and S2 (WGA: membranes; Hoechst: DNA; Phalloidin: F-actin). A schematic diagram of butterfly ovarioles is shown at bottom right. **(A,H)** Phenotype Normal in a pure individual. The confocal image shows follicles with their sheath removed. **(B,I)** Phenotype Normal in F_1 females (cross direction DB). Sheath is retained in the confocal image. Follicles are mostly indistinguishable from those of pure females, but the distribution of yolk in late-stage oocytes is overly scattered. It is not classified as a separate phenotype. **(C,J)** Phenotype Empty in F_1 females (cross direction BD). Sheath is retained in the confocal image, and very little tissue remains inside the sheath. **(D,K)** Phenotype Diminished. Sheath is retained in the confocal image, and a substantial amount of tissue remains, but without any discernible structures of normal follicles. **(E-G,L)** Phenotype Jammed (including its subtypes). Sheath is removed in the confocal image, and follicle cells are merged into a tube-like structure with many nurse cells and oocytes. Subtype canonical: the entire ovariole collapses into a single bag of oocytes and nurse cells. Subtype oblong: similar to "canonical", but the ovariole is elongated. Subtype ambiguous: some parts of the ovariole have merged follicles; other parts contain isolated follicles.

except for chromosome 14, where our manual correction cannot resolve all visible errors (Materials and Methods). As such errors affect the inference of recombination breakpoints, we do not report crossover patterns on chromosome 14.

We infer the crossover pattern in F_1 males by counting estimated recombination breakpoints across all backcross offspring (female meiosis in Lepidoptera is achiasmatic). Most F_1 males have at least one crossover per chromosome pair per meiosis, but the degree of crossover interference varies among chromosomes (Fig. 1D). Double crossovers occur as frequently as over 40% (chromosome 8), but are almost absent in many chromosomes. Importantly, the Z chromosome in this system has almost no double crossovers, and its recombination breakpoints are approximately uniformly distributed along the chromosomal axis (Fig. 1E). This indicates that recombination on the Z chromosome can be approximated by a model where the first obligatory crossover occurs uniformly at random along the chromosome, while the second crossover is permanently suppressed. We apply this model to the study of abnormal size below.

2.3 Abnormal size is determined by the entire Z chromosome

Pupal weight in backcross individuals (measured in backcrosses of type B(BD) and D(DB)) has a broad distribution (Fig. 1B). At first sight, such a distribution might be easily associated with many loci across the genome or extreme developmental/environmental stochasticity. Curiously, the Z chromosome is the only chromosome controlling pupal weight variation among backcross females in either direction, and no autosomal locus is significantly associated (Fig. 3A,3B). A single-locus QTL scan (1-QTL) on the Z chromosome can explain over 50% of the phenotypic variance, and a two-locus additive QTL scan (2-QTL) can explain over 70% (Fig. 3D), so variation of abnormal size in hybrid females is determined mostly by the Z chromosome.

Rather than inferring putative QTL, we argue that this genetic architecture is best described by a Z-chromosome continuum where many quantitative effects of small magnitude are uniformly distributed along the chromosomal axis (Fig. 3C). We refer to this as the Z-spanning polygenic system. Since different parts of the chromosome have identical effects under this assumed architecture, pupal weight will depend only on the average Z chromosome ancestry, f_D and f_B, corresponding to the fraction of the Z chromosome inherited from *dehaanii* and *bianor*, respectively. If effects are additive, this polygenic model is simply

$$W_{B(BD)} = w_B + \alpha_D f_D + \epsilon$$
$$W_{D(DB)} = w_D - \alpha_B f_B + \epsilon$$

(1)

where W is the pupal weight of a backcross female, w is the intercept, α is the net effect of Z-linked introgression, and ϵ contains non-genetic fluctuation. We will use f throughout the paper to refer to the fraction of introgression, and its subscript specifies the context. The relationship between pupal weight and average Z chromosome ancestry in our data is shown in Fig. S9.

Figure 3: Analysis of abnormal size. **(A, B)** The LOD score of genome-wide 1-QTL scans on female pupal weight. **(C)** The Z-spanning polygenic model posits that every infinitesimal introgression on the Z chromosome has a fixed infinitesimal effect on pupal weight. **(D)** The highest explanatory power (R^2) of 1-QTL regressions, 2-QTL regressions, and the regression to the average ancestry on the Z chromosome. **(E)** Dividing the Z chromosome into multiple blocks and performing multiple regression does not strongly improve the explanatory power. **(F, G)** The relative loss of predictive power in a 1-QTL scan on the Z chromosome. **(H, I)** The relative loss of predictive power in a 2-QTL scan on the Z chromosome.

Initial evidence supporting this polygenic architecture is as follows. First, this polygenic model (Equation 1) achieves a similar or even better explanatory power than the best QTL models (Fig. 3D), suggesting that perhaps the entire Z chromosome contributes to the phenotype. Second, to test if different parts of the Z chromosome could have different effects, we divide the Z chromosome into K equal-sized blocks ($K = 1, 2, \cdots, 10$), then perform multiple regression between pupal weight and the vector containing average ancestries on each of the K blocks. One should expect the total explanatory power to rise significantly with K if different blocks have very different effects due to increased degrees of freedom in regression. However, we find that using a single block (the entire chromosome) is almost as powerful as using multiple blocks (Fig. 3E). Effects are thus likely distributed evenly on the Z chromosome. This reasoning might seem at odds with the fact that both 1-QTL and 2-QTL scans peak in the interior of the Z chromosome (Fig. S8). However, we show below that such dome-shaped QTL results are exactly predicted by the proposed Z-spanning polygenic system.

2.4 Strong evidence for the Z-spanning polygenic architecture

The reason that 1-QTL scans produce peaks near the center of the chromosome is that the backcross generation still has long blocks of identical local ancestry. Since the chromosome center is the region that is the least distant from all other positions on the chromosome, and the crossover process on the Z chromosome is approximately spatially uniform, local ancestry at the chromosome center will thus correlate best with average chromosomal ancestry (f_D, f_B). Consequently, markers near the center will also be the most informative about any phenotype determined by the proposed polygenic model (Equation 1), while markers away from the center will lose phenotypic information. This is analogous to the "mid-domain effect" in species richness gradients [31, 32]. To quantify how much phenotypic information is lost by assuming a single marker, we calculate the relative loss of predictive power between a 1-QTL scan on a specific marker versus the polygenic model (Materials and Methods):

$$\frac{R^2_{\text{1QTL}}}{R^2_{\text{Z-ancestry}}} = \frac{3}{8}\left[1 + 2l\left(1 - l\right)\right]^2,$$ (2)

where l $(0 \leq l \leq 1)$ is the relative marker position on the Z chromosome. Since $R^2_{\text{Z-ancestry}}$ is a constant, this equation has no degree of freedom besides l, and it depicts precisely a dome-shaped predictive power of a 1-QTL scan peaking at the chromosome center ($l = 0.5$). It provides a good fit to observed 1-QTL results in both cross directions (Fig. 3F,3G). Similarly, for a 2-QTL scan, the relative loss of predictive power for two markers is

$$\frac{R^2_{\text{2QTL}}}{R^2_{\text{Z-ancestry}}} = \frac{1}{8 - 4|l_1 - l_2|}\left\{6\left|l_1 - l_2\right|\left(l_1 + l_2 - 1\right)^2 \right.$$
$$\left. + 3\left[1 + 2l_1\left(1 - l_1\right)\right]\left[1 + 2l_2\left(1 - l_2\right)\right]\right\},$$ (3)

where l_1 and l_2 are the relative positions of two markers on the Z chromosome. This equation shows that the two most informative markers are located near $l_1 \approx 0.27$ and $l_2 \approx 0.73$—about a quarter into the chromosome from both ends. Again, the equation is a good fit to the results of 2-QTL scans (Fig. 3H,3I). From this, we speculate that, by calculating the same quantity for 3-QTL scans, this polygenic model will still be a good fit, and it is confirmed in Fig. S10.

Despite being polygenic, the proposed architecture is mathematically parsimonious because it is a linear model with just one predictive variable (mean chromosomal ancestry) (Eq. 1), while still carrying almost the complete spatial information of QTL scans. Thus, we conclude that abnormal size is best explained by many Z-linked polygenes, each having small phenotypic effects, evenly smeared across the entire chromosome.

Figure 4: Analysis of ovary dysgenesis. For maternally *bianor* hybrids: **(A)** Phenotype Empty is dominantly controlled by Locus E on the Z chromosome. The LOD plot shows both the score for the canonical Empty phenotype as well as the score when we include a few ambiguous individuals that are classified as Empty. **(B)** If the Z chromosome is purely *bianor*, introgression on Locus R suppresses abnormal phenotypes. For maternally *dehaanii* hybrids: **(C)** If the Z chromosome is purely *dehaanii* in backcross females, ovaries develop the Diminished phenotype. **(D)** If the Z chromosome is purely *bianor* in backcross females, ovaries are mostly Jammed.

2.5 A Z-linked system shapes ovary dysgenesis in maternally *bianor* hybrids

In F_1 females of type BD, ovaries are small and empty (phenotype Empty), and only a trace amount of tissue remains in each ovariole (Fig. 2C,J). The locus (E) responsible for Empty is mapped to a 0.8-1Mb interval on the Z chromosome with a simple genetic mechanism (Fig. 4A): the *dehaanii* allele at locus E is sufficient to induce Empty regardless of other genetic variation across the genome.

Other ovary phenotypes in this maternal background include Normal and various types of Jammed, where multiple oocytes and a myriad of nurse cells congregate into a single tube-shaped follicle (Fig. 2E-G,L). We remove the effect of locus E, together with other possible Z-linked factors by selecting individuals where the Z chromosome has no introgression from *dehaanii*. Conditioning on this subset of individuals, we further identified a locus (R) on chromosome 8 that suppresses pheno-

type Jammed when it is heterozygous in ancestry (Fig. 4B). Conversely, if locus *R* has a homozygous *bianor* genotype, approximately half of the offspring develop Jammed, but the sample size becomes too low to map its associated loci. Overall, the genetic basis of ovary dysgenesis in maternally *bianor* hybrids is controlled mainly by a small number of genomic regions with large effects.

2.6 Recombination on the Z chromosome rescues ovary dysgenesis in maternally *dehaanii* backcrosses

Figure 5: The relationship between phenotypic defect and chromosomal ancestry. Across panels, f_Z and f_A are the proportions of introgressed ancestry on the Z chromosome and autosomes, respectively. **(A)** Average Z chromosome ancestry is very informative in ovarian variation in cross direction D(DB). **(B)** The "ancestry-balance" model posits that the probability of observing normal phenotypes is highest when the amount of introgression on autosomes and the Z chromosome reaches a critical balance (not necessarily equal). **(C)** Between *Heliconius pardalinus butleri* and *H. p. sergestus*, oogenesis in maternally *butleri* backcross females is better when Z-linked introgression is at intermediate levels. **(D)** The probability of observing the Normal phenotype conditioning on the difference between f_Z and f_A in cross direction D(DB). **(E)** In cross direction B(BD), female pupal weight is less different from that of pure *bianor* females when Z-linked introgression is at intermediate levels. **(F)** In cross direction D(DB), deviation of female pupal weight from that of pure *dehaanii* females is smaller with less amount of Z-linked introgression.

In the opposite maternal background (F_1 and backcrosses with *dehaanii* mothers), ovary dysgenesis yields a number of counter-intuitive results. In this maternal background, ovaries fall into three phenotypic categories: Diminished, Normal, and Jammed (including its subtypes). Overall, the Z chromosome is still the most predictive chromosome with significant LOD peaks (Fig. S11). However, two pieces of evidence suggest ovary phenotypes are influenced also by autosomes. First, if the Z chromosome has pure *dehaanii* ancestry, backcross females all have the Diminished pheno-

type (Fig. 4C,5A). Comparing these individuals to pure *dehaanii* females show that autosome-only introgression from *bianor* at the level of a backcross individual is sufficient to induce Diminished. In other words, Diminished is stably caused by the existence of certain levels of introgression on autosomes, rather than the exact location of introgression, which is evident from autosomal LOD scores (Fig. 4C). Second, if the Z chromosome has pure *bianor* ancestry, almost all backcross females develop abnormal ovaries as well, and most of them have the Jammed phenotype (Fig. 4D,5A). Similarly, comparing these individuals against F_1 females of type DB (which have Normal ovaries and a Z chromosome inherited from *bianor*) leads to the result that abnormality is caused by many autosomal factors (the LOD score in Fig. 4C). Both cases suggest a polygenic role of autosomes in producing ovary dysgenesis in a pure-Z background. Nonetheless, Normal ovaries do exist among backcross females but they almost always require a recombined Z chromosome (Fig. 5A). Furthermore, two individuals with Z chromosomes recombined in opposite ways can be both Normal (Fig. S12). This quantitative rescue effect of recombination on the Z chromosome is strikingly similar to the only other genome-wide mapping of ovarian dysgenesis in Lepidoptera (*Heliconius*) [26], which we discuss below.

2.7 Z-autosome ancestry balance as a precondition for high fitness

In *Heliconius*, F_1 females between *H. pardalinus butleri* and *H. pardalinus sergestus* develop empty ovaries [26]. When F_1 males are backcrossed to *H. p. butleri*, female offspring have the most normal ovary phenotypes when the Z chromosome contains about 1/4 introgression from either end (Fig. 5C). Autosomes must contribute to ovary dysgenesis in D(DB) and *Heliconius* females, because ovaries are strongly defective with autosome-only introgression ($f_A \gg 0, f_Z = 0$), while increasing introgression on the Z chromosome will ameliorate autosome-induced defects until reaching a turning point (Fig. 5D). The same phenomenon in abnormal size seems to occur in B(BD) females (Fig. 5E), but not in D(DB) females (Fig. 5F). If Z chromosomes with recombined ancestry yield the best phenotype, autosomal introgression must bring in a large quantity of partially dominant incompatibility factors so that phenotypic defect stably appears when there is no Z-linked introgression. It also implies that Z-linked introgression contains many other factors balancing those on autosomes quantitatively. The outcome of this assumed process is an "ancestry-balance" precondition between Z and autosomes for the appearance of normal phenotypes (Fig. 5B). Nonetheless, Z-autosome ancestry balance does not necessarily hold for all traits in all crosses, because some traits can be masked by Mendelian factors of large effects (Locus *E* in ovary dysgenesis), and some traits are only weakly affected by autosome-only introgression (abnormal size in D(DB), although this particular example can also fit the "ancestry-balance" model by choosing a highly skewed balance at $f_Z - f_A = -1/4$). In short, if this precondition is valid, phenotypes will segregate by average ancestries between Z and autosomes and it indicates the involvement of highly dispersed genomic factors for incompatibility.

2.8 Male homogamety ameliorates size abnormality

The converse of the problem underlying Haldane's Rule is to explain weaker incompatibility effects in the homogametic sex. In our system, unlike F_1 males, backcross males in both directions have a significant variation in pupal weight (Fig. 1B), but the spread is visibly smaller than that of females. It is congruent with the homogametic sex being less affected by hybridization, but what could be its genetic mechanism? Here we show analytically that this phenomenon is explicable by the Z-spanning polygenic architecture in male butterflies.

The genetic model outlined by Eq. 1 has a single net effect α combining both the effect of native alleles (α_0) and the effect of introgressed alleles (α_1). If we split these contributions into two terms, and let $\langle \Delta W \rangle$ be the expected deviation from normal pupal weight, then

$$\langle \Delta W \rangle_{\text{Female}} = \underbrace{\alpha_0(1-f)}_{\text{Native}} + \underbrace{\alpha_1 f}_{\text{Introgressed}} \tag{4}$$

In Fig. 5E and 5F, extreme ancestries on the Z chromosome ($f = 1$ and $f = 0$) will push ΔW to two opposite extremes (B(BD)) or to only one extreme (D(DB)). It implies that native and introgressed alleles do not act in the same direction on phenotypes ($\alpha_0 \alpha_1 \leq 0$).

Now we apply this equation to backcross males with a slight variation. Backcross males have one copy of the pure Z chromosome and another copy that could be partially introgressed. For regions on the Z chromosome homozygous for native ancestry, the effect is the same as for females (α_0), since dosage compensation in Lepidoptera partially and symmetrically suppresses both Z chromosomes in males [33, 34]. For regions heterozygous in ancestry, let h_0 and h_1 be the dominance of native and introgressed alleles, respectively. Again, we assume that dosage compensation exists along regions heterozygous in ancestry (perhaps imperfectly) so that h_0 and h_1 capture not only dominance in the ordinary sense, but also the reduction in effect sizes due to partial expression suppression. Then:

$$\langle \Delta W \rangle_{\text{Male}} = \underbrace{\alpha_0(1-f)}_{\text{Homozygous}} + \underbrace{(h_0\alpha_0 + h_1\alpha_1)f}_{\text{Heterozygous}} \tag{5}$$

The genetic variance in either sex is induced by variation in f, and its ratio between the two sexes is:

$$\begin{aligned}
\frac{V_{g,\text{Male}}}{V_{g,\text{Female}}} &= \left[\frac{h_1\alpha_1 - (1-h_0)\alpha_0}{\alpha_1 - \alpha_0} \right]^2 \\
&= \left[\frac{h_1|\alpha_1| + (1-h_0)|\alpha_0|}{|\alpha_1| + |\alpha_0|} \right]^2 \leq 1
\end{aligned} \tag{6}$$

This inequality is simply guaranteed by our observation that alleles of opposite ancestry have mostly opposite phenotypic effects among backcrosses ($\alpha_0 \alpha_1 \leq 0$). Even though backcross males were not sequenced, Eq. 6 provides a generic qualitative prediction that the genetic variance of

male pupal weight should be smaller than that of females. What we observe is shown in Table 1. It clearly shows a weaker effect of Z-linked polygenes on male size. Interestingly, dominance theory ($h_0 = h_1 = 0$) becomes a particular case of this equation, and if it holds, we can solve for the phenotypic effect of introgressed alleles relative to that of native ones:

$$\left.\frac{\alpha_1}{\alpha_0}\right|_{h_0=h_1=0} = 1 - \sqrt{\frac{V_{g,\text{Female}}}{V_{g,\text{Male}}}} \tag{7}$$

If dominance theory does not hold, and effects are fully additive with complete but perhaps asymmetric dosage compensation, then $h_0 + h_1 = 1$, which leads to a constant ratio of genetic variance, and

$$\left. h_1 \right|_{h_1+h_0=1} = \sqrt{\frac{V_{g,\text{Male}}}{V_{g,\text{Female}}}} \tag{8}$$

Under this assumption of complete dosage compensation on additive effects, the ratio now reflects which ancestry has a biased expression along heterozygous regions. In the absence of such bias ($h_1 = 1/2$), dosage compensation will be symmetric between two haplotypes along regions heterozygous in ancestry, and the ratio of genetic variance will be fixed at 0.25. In our data, assuming the additive model, B(BD) males will show more equalized contributions between Z-linked alleles of different ancestries (h_1 is closer to 0.5), while D(DB) males are more affected by the contribution of introgressed alleles on Z chromosomes. Finally, we can rule out the classical dominance theory in D(DB), because native alleles on the Z chromosome have very weak phenotypic effects ($\alpha_0 \approx 0$, see Fig. 5F), which will force the ratio of V_g between males and females to be zero if $h_0 = h_1 = 0$. Taken together, we show that male homogamety can ameliorate size abnormality under quite general conditions in this particular system. This further validates our proposal of the Z-spanning polygenic architecture.

Table 1: The ratio of genetic variance between male and female pupal weight among backcross individuals

Cross direction	D(DB)	B(BD)
Observed ratio of V_g	0.37	0.23
95% Confidence interval of V_g	(0.18, 0.62)	(0.05, 0.46)
α_1/α_0 (dominance theory)	N.A.	-1.09
h_1 (complete dosage compensation on additive effects)	0.61	0.48

2.9 Relevance to existing theories

We proceed to test if genetic architectures underlying abnormal size and ovary dysgenesis can be explained by existing theories of Haldane's rule and the large-X/Z effect. Dominance theory is in

principle compatible with Lepidoptera as it does not rely on sex-specific mechanisms. For abnormal size, we have shown in the previous section that dominance theory is a particular case of the more general reasoning based on the quantitative genetic basis of pupal weight variation. In our reasoning, the critical assumption is the partial suppression ($h_0, h_1 \in [0, 1)$) of mostly opposite allelic effects ($\alpha_0 \alpha_1 \leq 0$) between different ancestries on Z chromosomes in the homogametic sex.

For ovary dysgenesis, the effect of autosomal introgression is not completely recessive. For instance, introgression causing Diminished ovaries in maternally *dehaanii* females (yellow tracts in Fig. 4C) is fully heterozygous in ancestry and should contain at least additive factors. In maternally *bianor* females, introgression at locus *R* dominantly suppresses dysgenesis when the Z chromosome contains no introgression (Fig. 4B). We are aware that it is difficult to detect introgressed alleles with recessive effects using this backcross scheme, but our result contrasts with findings in *Drosophila* hybrids, where almost all autosomal introgressions act recessively in hybrid incompatibility [15]. However, it is difficult to judge the recessivity of Z-linked female sterility factors because oogenesis is a female-specific trait, and the female only has a single Z chromosome. Consequently, we cannot rule out dominance theory in ovary dysgenesis.

Since the Z chromosome is involved in almost all cases of hybrid disruption, and its effect is not localized to specific regions except for Locus *E*, can it be explained by increased differentiation on the entire Z chromosome relative to the autosomes? We find no evidence for this faster-Z theory. On the Z chromosome, most sequence windows (50kb) are less divergent between the two species than the average divergence on autosomes (Fig. S13). This echoes existing findings showing mixed support for faster-Z in Lepidoptera [35, 36, 37]. Theories specifically tailored for X/Y systems (the faster-male theory, the spermatogenesis theory, see [18]) do not apply to butterflies as females are heterogametic. Finally, although our data cannot rule out the possibility that female meiotic drive disrupts oogenesis, it is unknown how this process might occur in Lepidoptera females because their chromosomes are truly holokinetic (unlike *C. elegans*, which has "mini-centromeres")[38, 39], and most documented sex-ratio drives in Lepidoptera are caused by *Wolbachia* infection [40].

2.10 Why are incompatibility factors so dispersed? A hypothesis based on epigenetic regulation

One of our key findings is that many factors of perhaps individually small effects are widely dispersed across autosomes or on the Z chromosome. Consequently, average chromosomal ancestry is often more informative of phenotypes than any particular locus. This pattern is similar to the polygenic threshold model of hybrid incompatibility in *Drosophila*, where abnormal phenotypes depend more on the total quantity of introgression than where introgression occurs in the genome [41, 42, 43]. As an attempt to explain our results under a coherent framework, we introduced in Fig. 5B a hypothesis positing that normal phenotypes require average autosomal ancestry and average Z chromosome ancestry to achieve a certain balance. Here, we explain why such balance might involve epigenetic regulation, particularly dosage compensation mechanisms specific to Lep-

idoptera.

Although we cannot test for mechanisms of Z-autosomes ancestry balance using our current data, we can think of two extreme scenarios for the identity of underlying factors: First, the balance might be between identical sequences widely distributed across the genome, perhaps repetitive DNA and their regulators. But it seems unlikely that this would have a highly uniform effect on the Z chromosome to additively determine hybrid size. Second, the balance could be entirely epigenetic. Abnormal size in hybrids that is asymmetric between reciprocal crosses exists in other systems, and it is often linked to epigenetic regulation such as imprinting [44, 45]. In our case, it must be related to some process that occurs between autosomes and the Z chromosome with a broad genomic basis. It is tempting to consider dosage compensation as a candidate mechanism because it affects the expression of the entire genome as well as specific genes. The details of dosage compensation might have diverged between the two species (e.g., the absolute level of genomic expression, species-specific epigenetic factors for dosage compensation etc.), while expression is still fully compensated between the two sexes within each lineage [34]. In Lepidoptera, dosage compensation is typically achieved by partial suppression of both Z chromosomes in males, and might also involve some up-regulation of the Z chromosome in females [33]. To illustrate the effect of hybridization, consider a hybrid female with only Z-linked introgression. Suppose this Z-linked region introgressed from a different species is more sensitive to up-regulation than its native homolog, so its presence increases the total expression of the Z chromosome and ruptures regular levels of dosage compensation. If hybrids are also fully compensated for dosage, then they might up-regulate autosomes to match the expression on the Z chromosome, which creates a quantitative effect on expression levels all over autosomes proportional to the percentage of introgression on the Z chromosome. On the other hand, if only autosomes have introgression, the expression of the native Z must change accordingly to achieve dosage compensation in hybrids. If the amount of introgression is balanced between autosomes and Z so that both are affected in the same way, there is perhaps no effect on the default dosage compensation program. This process might provide an explanation for the highly dispersed genetic architecture.

This hypothesis makes a number of untested assumptions, but it provides a phenomenological fit for our observation. In ovary dysgenesis among D(DB) females, since Diminished is a stable ovary phenotype over many configurations of autosomal introgression (Fig. 4C), it is more likely to be directly caused by the misexpression of a fixed set of factors, for instance, a set of oogenesis genes on the pure Z chromosome. The indirect cause is then 15%-30% introgression on autosomes rupturing the global outcome of dosage compensation when the Z chromosome contains no introgression. This would explain the dispersed nature of autosomal factors. If recombination occurs on the Z chromosome, it allows the Z chromosome to achieve a critical level of introgression matching that of autosomes in backcross individuals. It consequently pushes the system towards normal levels of dosage compensation, which explains the rescuing effect of recombination.

The same logic also applies to size abnormality. If size is affected mainly by the expression of a

few essential genes, such as those in the insulin signaling pathways or juvenile hormones [46, 47], the level of misexpression will be proportional to the difference between autosomal introgression and Z-linked introgression. Since the Z chromosome is much more variable in mean ancestry compared to that over all autosomes in the backcross generation, it will explain why the Z chromosome is the most informative of pupal weight, and it will also explain why the entire Z chromosome is involved rather than its local regions. This hypothesis is also congruent with the reduction of abnormal size in F_1 males: Autosomes and two Z chromosomes in F_1 males inherit half from each species so that ancestry balance is more likely to be attained; In F_1 females, the only Z chromosome has pure ancestry, while autosomes have equalized contribution, creating strong ancestry imbalance between Z and autosomes, which might rupture dosage compensation. Finally, our reasoning for the reduction of abnormal size in backcross males also involves dosage compensation features unique to Lepidoptera (partial repression of both Z chromosomes). A future test for this hypothesis should incorporate genome-wide, allele-specific expression data on hybrids.

3. Conclusion

The two rules of speciation have many genetic mechanisms, and most of them are based on male-heterogametic systems. Our study reveals that Haldane's Rule and the large-Z effect in this butterfly system have a highly polygenic basis on *both* autosomes and the Z chromosome, but narrow regions of large effects do exist in some particular cases. The rescue effect of recombination on the Z chromosome in several traits suggests a balancing process between autosomes and the Z chromosome, the nature of which awaits to be revealed.

4. Materials and Methods

4.1 Breeding

Lineages of *P. dehaanii* were purchased directly from a butterfly farm in Qingdao (Shandong Province, China), exclusively sourced from a small local population. Lineages of *P. bianor* were collected in the field from Ningbo (Zhejiang Province, China) for breeding in 2020, 2021, and a few individuals were collected in the field from Kunming (Yunnan Province, China) for breeding in 2019. All crosses were done by hand-pairing. Eggs were collected by putting females in small cages with host plants under fluorescent light. The following host plants were used throughout the project: *Tetradium daniellii*, *Zanthoxylum bungeanum*, *Z. ailanthoides*, *Z. beecheyanum*, *Z. simulans*, *Choisya ternata*, *Phellodendron amurense*. Juveniles were kept in greenhouse conditions (approximately 20°C~35°C), with a combination of natural and greenhouse lights to maintain at least 10 hours of illumination per day. Adults for dissection were immediately put into a 5°C room after eclosion to reduce activity. Otherwise, they were fed with sugar water once a day, and females were subsequently kept in the dark, while males were in an illuminated environment to facilitate hand-pairing.

4.2 Phenotyping ovaries

Ovaries were dissected from females within five days of eclosion in $1\times$ PBS solution. The ovariole sheath was manually removed, and most images were taken using the internal camera of a Leica EZ4 HD microscope (pictures of a few specimens were taken by a cellphone through the eyepiece of a Zeiss Stemi 2000 microscope). Since ovary phenotypes are categorical, we established all major categories by defining the most obvious and the most frequent phenotypes across all dissected ovaries, and it was confirmed later by confocal imaging that they indeed have large qualitative differences. Phenotype Jammed is variable in terms of the fraction and the position of Jammed follicles. To reduce subjective opinions in scoring different kinds of Jammed, we lumped them into a single category in the QTL analysis. A small number of ovaries have ambiguous phenotypes due to one of the following reasons: 1) Different ovarioles develop different phenotypes; 2) Some part of the ovary is lost in dissection; 3) Extremely rare phenotypes that are hard to classify into existing categories. These ambiguous individuals, mostly from cross direction D(DB), were assigned multiple categories. When such uncertainty affects analyses, we randomly select a phenotype from previously assigned categories on ambiguous individuals, perform analyses, and repeat the same procedure many times to control for uncertainty in phenotype.

4.3 Staining and confocal imaging of ovaries

Dissected ovaries were fixed in 4% Paraformaldehyde solution in $1\times$ PBS for 20 minutes at room temperature. The ovaries were washed for 15 minutes each in 0.1% PBTx ($1\times$ PBS, 0.1% Triton-X 100), 1% PBTx, 2% PBTx, 0.01% Saponin (Sigma Aldrich 47036) in PBS and Block solution ($1\times$ PBS, 0.3% Triton-X 100, 0.5% Normal Goat Serum). The ovaries were then stained for 12 hours using the following reagents at 1:500 dilution in blocking solution: Hoechst 33342 (10 mg/ml, Thermo Fisher H3570), Wheat Germ Agglutinin-647 (WGA, Thermo Fisher W32466) and Rhodamine Phalloidin (Thermo Fisher R415). Stained ovaries were washed four times for 15 minutes each in 0.1% PBTx followed by a final PBS wash. After the washes, ovaries were mounted in equal volumes of PBS and Vectashield mountant (Vector labs H1900) on a slide.

The ovary samples were imaged by acquiring Z-section images on a Zeiss LSM 880 laser scanning confocal microscope at Harvard Center for Biological Imaging. The microscope was equipped with an Argon laser and a He/Ne 633 nm laser. Zeiss Plan-Apochromat $10\times$/0.45 M27 or $20\times$/0.8 M27 objective lenses were used for imaging. All images were 1024×1024 pixels in size and were acquired using PMT detectors. Images were acquired at excitation/emission wavelengths of 405/450 nm for Hoechst 33342, 561/610 nm for Rhodamine Phalloidin, and 633/696 nm for WGA.

4.4 DNA extraction and sequencing

Samples from the cross were preserved in either pure ethanol or RNAlater at -20C prior to DNA extraction. For extraction, we used E.Z.N.A Tissue DNA kits (Omega Bio-tek, Inc.). Whole-genome library preparation was performed using Illumina DNA 1/4 reactions kits at Harvard University

Bauer Core, and subsequently sequenced altogether on a single lane of Illumina NovaSeq S4. Autosomal coverage varies among individuals: backcrosses-1×, parents-5×, grandparents-30×∼60×. Raw reads were trimmed with Cutadapt-3.4 [48] to remove adapters (CTGTCTCTTATACACATCT), and subsequently mapped to the reference genome of *P. bianor* using the BWA-0.7.17 MEM algorithm. Duplicated reads were marked using Picard-2.25.7 [49]. For samples from the cross, we used BCFtools-1.9 [50] to pile up reads with very light quality filtering, and to call variants with associated genotype likelihoods. VCF files produced by the variants caller will be used for linkage analysis.

4.5 Linkage analysis

For quality control, we first calculated kinship coefficients among individuals using NgsRelate-2 [51] and corrected the pedigree position of a few individuals (Fig. S3). Lep-Map-3 was used for all subsequent linkage analysis [52]. First, VCF files and the pedigree were combined in module ParentCall2, and we imputed haplotype structure along the reference genome with module Order-Markers2. We also generated de novo marker orders using the same module. Comparing de novo marker orders against the order in the original reference genome revealed some intra-chromosome assembly problems. We corrected these problems below.

4.6 Linkage-based reference genome correction

For a pre-correction reference genome, we plotted the de novo marker order against genomic order and discovered some assembly errors that affect the order and orientation among PacBio scaffolds on chromosomes. Using inferred haplotypes in the grandparent phase, we calculated the correlation of ancestry between each pair of markers, which should be a decreasing function of marker distance due to recombination. This information helps correct large-scale errors and generate an intermediate genome. To correct for smaller errors, we inferred the de novo marker order on the intermediate genome and compared it against the genome itself. This extra step corrected the position/orientation of several smaller scaffolds. We were able to correct all visible errors except for those on chromosome 14, where an orientation problem appears to occur *within* a PacBio scaffold and we were unable to determine its breakpoint. (This may be due to a different population used here compared to the reference population). The de novo marker order generated on this final genome yields mostly a collinear relationship to the genomic order (Fig. S4). Finally, haplotypes in the grandparent phase were re-inferred on the corrected genome for all subsequent analyses.

4.7 Crossover analysis

To infer crossover frequency, we calculated the number of recombination breakpoints in each paternal haplotype among all backcross individuals (For all paternal haplotypes, see Fig. S5, S6, S7). No chromosome has more than two breakpoints except for chromosome 14 (one individual). We excluded chromosome 14 from this crossover analysis due to the aforementioned assembly prob-

lem. Let n_0, n_1, and n_2 be the number of haplotypes having 0, 1, or 2 recombination breakpoints for a given chromosome. The maximum likelihood estimate of crossover frequency is as follows (see "SI-Crossover analysis" for derivation). First, calculate:

$$
\begin{aligned}
c_0 &= (n_0 - n_1 + n_2)/(n_0 + n_1 + n_2) \\
c_1 &= (2n_1 - 4n_2)/(n_0 + n_1 + n_2) \\
c_2 &= 4n_2/(n_0 + n_1 + n_2)
\end{aligned}
\tag{9}
$$

If c_0, c_1, c_2 are all nonnegative, they are the frequencies of having 0, 1, or 2 crossovers. If $c_0 < 0$, the adjusted estimate is

$$
\begin{aligned}
c_0^* &= 0 \\
c_1^* &= (n_0 - n_2)/(n_0 + n_2) \\
c_2^* &= 2n_2/(n_0 + n_2)
\end{aligned}
\tag{10}
$$

4.8 QTL analysis

One-dimensional QTL analysis was performed with R-package qtl2 [53]. Ovary phenotypes were always mapped one at a time using a binary trait logistic mapper (i.e., phenotype of interest = 1, other phenotypes = 0). This approach is suitable for unordered categorical traits such as ovary morphology. For pupal weight, we introduced brood as a covariate to control for seasonal variation in pupal weight due to diet and environmental factors. LOD-scores were always estimated on 1,000 random permutations of phenotypes. To calculate the explanatory power of QTL on pupal weights (R^2_{1QTL}, R^2_{2QTL}), we performed linear regression of brood-corrected pupal weight on marker ancestry (one marker for a 1-QTL scan or two additive markers for a 2-QTL scan), and explanatory power is defined as the R^2 of the regression model.

4.9 The Z-spanning polygenic system

Let $f \in [0, 1]$ be the fraction of Z chromosome coming from a foreign species, and let W be the random variable describing pupal weight. The Z-spanning polygenic system postulates that, on average, each infinitesimal quantity of introgression on Z has a fixed infinitesimal effect on pupal weight. In other words,

$$
\mathrm{d}\langle W \rangle = \alpha\,\mathrm{d}f
\tag{11}
$$

In the form of linear regression, this is often written as

$$
W = w + \alpha f + \epsilon
\tag{12}
$$

where w is the average pupal weight without Z-linked introgression, and ϵ is a noise variable with mean zero. Among a group of individuals, the explanatory power (R^2) of this model is simply

$\alpha^2 \mathrm{Var}(f)/\mathrm{Var}(W)$. Let $p(l)$ be the marker ancestry (native=0) at position l on a chromosome of unit length. The Z-spanning polygenic model implies the following Markov Chain: $\{p(l)\}_{l=0}^1 \to f \to W$. Thus, ancestry p at a particular position l will be much less informative about W than f. The regression model for a one-marker QTL analysis is $W \sim p$, while regression with the polygenic model is $W \sim f$. The loss of predictive power by using $W \sim p$ relative to $W \sim f$ is simply the squared coefficient of (multiple) correlation:

$$\frac{R^2_{W \sim p}}{R^2_{W \sim f}} = \rho^2_{p,f} = \frac{\mathrm{Cov}^2(p,f)}{\mathrm{Var}(p)\mathrm{Var}(f)} \tag{13}$$

where $\rho_{p,f}$ is the correlation coefficient between p and f. This result is written as $R^2_{\mathrm{1QTL}}/R^2_{\mathrm{Z\text{-}ancestry}}$ in the main text. For a two-marker additive model $W \sim p_1 + p_2$, the relative loss is analogously

$$\frac{R^2_{W \sim p_1 + p_2}}{R^2_{W \sim f}} = \begin{bmatrix} \rho_{p_1,f} \\ \rho_{p_2,f} \end{bmatrix}^\top \begin{bmatrix} 1 & \rho_{p_1,p_2} \\ \rho_{p_2,p_1} & 1 \end{bmatrix}^{-1} \begin{bmatrix} \rho_{p_1,f} \\ \rho_{p_2,f} \end{bmatrix} \tag{14}$$

These correlation coefficients will depend on marker positions as well as the crossover process. In our case, since crossover on the Z chromosome is obligatory, spatially uniform, and with strong interference, it is straightforward to model one round of crossover for backcross females and calculate ρ. Then we get

$$\rho_{p(l),f} = \frac{1}{2}\sqrt{\frac{3}{2}}\left[1 + 2l(1-l)\right]$$
$$\rho_{p(l_1),p(l_2)} = 1 - |l_2 - l_1| \tag{15}$$

Applying these values to Equations 13 and 14 leads to the final result of Equation 2 and 3

For the variance analysis of male pupal weight, we assume that non-genetic fluctuation (ϵ) has the same magnitude between males and females, and we estimated $\mathrm{Var}(\epsilon)$ to be used in estimating $V_{g,\mathrm{Male}}$.

4.10 Posterior probabilities for ovary phenotypes based on $f_Z - f_A$

Let f_Z, f_A be the fraction of introgression from *bianor* into maternally *dehaanii* individuals on Z and autosomes, respectively. Let $x = f_Z - f_A$ be their difference. For ovary phenotype k from the set {Diminished, Normal, Jammed}, the posterior probability of observing k conditioning on a given range of x is (by Bayes theorem):

$$\mathbb{P}(k|x_i \leq x \leq x_{i+1}) = \frac{\mathbb{P}(x_i \leq x \leq x_{i+1}|k)\mathbb{P}(k)}{\sum_k \mathbb{P}(x_i \leq x \leq x_{i+1}|k)\mathbb{P}(k)} \tag{16}$$

where $\mathbb{P}(k)$ is calculated from the entire dataset. Due to the uncertainty of ovary phenotypes, we re-sampled phenotypes for ambiguous individuals and repeated the estimation for 1,000 times and reported the average value of $\mathbb{P}(k|x_i \leq x \leq x_{i+1})$ and its standard error.

5. Data Availability

Raw reads are deposited into the NCBI Sequence Read Archive (BioProject: PRJNA892033). Scripts are available from:

https://github.com/tzxiong/2022_Papilio_HybridIncompatibilityMapping

6. Author Contributions

T.X. and J.M. designed the project. T.X. established the cross and performed analysis. S.T. performed ovary staining, confocal imaging, and contributed to ovary phenotype classification. N.R. provided and analyzed *Heliconius* data. M.Y. and X.L. provided parental phenotypic data and key genomic resources. T.X., J.M. and S.T. wrote the original draft. All authors contributed to the improvement of the manuscript.

7. Acknowledgement

T.X. was funded by a studentship from the Department of Organismic and Evolutionary at Harvard University, the Quantitative Biology Initiatives at Harvard, the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard (award number #1764269), and a Grant in Aid of Research from Sigma Xi (The Scientific Research Honor Society). We thank Naomi Pierce and Adam Cotton for providing background information on the study system; Janet Sherwood, Shui Xu, Yuchen Zheng, Jinbo Hu, and Anastasios Kougionis for their assistance and knowledge in breeding/sourcing host plants and butterflies; Cassandra Extavour for discussing oogenesis; John Wakeley, Robin Hopkins, Liang Qiao, Sarah Dendy, Nathaniel Edelman, Shuzhe Guan, Fernando Seixas, and Yuttapong Thawornwattana for their support during the project.

References

[1] Coyne, J. A. Two rules of speciation. *Speciation and its consequences* 180–207 (1989).

[2] Haldane, J. B. Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics* **12**, 101–109 (1922).

[3] Orr, H. A. Haldane's rule. *Annual Review of Ecology and Systematics* 195–218 (1997).

[4] Turelli, M. & Orr, H. A. Dominance, epistasis and the genetics of postzygotic isolation. *Genetics* **154**, 1663–1679 (2000).

[5] Presgraves, D. C. & Orr, H. A. Haldane's rule in taxa lacking a hemizygous X. *Science* **282**, 952–954 (1998).

[6] Schilthuizen, M., Giesbers, M. & Beukeboom, L. Haldane's rule in the 21st century. *Heredity* **107**, 95–102 (2011).

[7] Delph, L. F. & Demuth, J. P. Haldane's rule: genetic bases and their empirical support. *Journal of Heredity* **107**, 383–391 (2016).

[8] Orr, H. A. Haldane's rule has multiple genetic causes. *Nature* **361**, 532–533 (1993).

[9] Presgraves, D. C. Sex chromosomes and speciation in *Drosophila*. *Trends in Genetics* **24**, 336–343 (2008).

[10] Presgraves, D. C. Evaluating genomic signatures of "the large X-effect" during complex speciation. *Molecular Ecology* **27**, 3822–3830 (2018).

[11] Larson, E. L., Keeble, S., Vanderpool, D., Dean, M. D. & Good, J. M. The composite regulatory basis of the large X-effect in mouse speciation. *Molecular Biology and Evolution* **34**, 282–295 (2017).

[12] Kitano, J. *et al.* A role for a neo-sex chromosome in stickleback speciation. *Nature* **461**, 1079–1083 (2009).

[13] Irwin, D. E. Sex chromosomes and speciation in birds and other ZW systems. *Molecular Ecology* **27**, 3831–3851 (2018).

[14] Davies, B. *et al.* Re-engineering the zinc fingers of PRDM9 reverses hybrid sterility in mice. *Nature* **530**, 171–176 (2016).

[15] Masly, J. P. & Presgraves, D. C. High-resolution genome-wide dissection of the two rules of speciation in *Drosophila*. *PLoS Biology* **5**, e243 (2007).

[16] Orr, H. A. & Turelli, M. Dominance and Haldane's rule. *Genetics* **143**, 613 (1996).

[17] Charlesworth, B., Coyne, J. A. & Barton, N. H. The relative rates of evolution of sex chromosomes and autosomes. *The American Naturalist* **130**, 113–146 (1987).

[18] Wu, C.-I. & Davis, A. W. Evolution of postmating reproductive isolation: the composite nature of Haldane's rule and its genetic bases. *The American Naturalist* **142**, 187–212 (1993).

[19] Frank, S. A. Divergence of meiotic drive-suppression systems as an explanation for sex-biased hybrid sterility and inviability. *Evolution* **45**, 262–267 (1991).

[20] Hurst, L. D. & Pomiankowski, A. Causes of sex ratio bias may account for unisexual sterility in hybrids: a new explanation of Haldane's rule and related phenomena. *Genetics* **128**, 841–858 (1991).

[21] Tao, Y. & Hartl, D. L. Genetic dissection of hybrid incompatibilities between *Drosophila simulans* and *D. mauritiana*.: III. Heterogeneous accumulation of hybrid incompatibilities, degree of dominance, and implications for Haldane's Rule. *Evolution* **57**, 2580–2598 (2003).

[22] Presgraves, D. C. Patterns of postzygotic isolation in Lepidoptera. *Evolution* **56**, 1168–1183 (2002).

[23] Jiggins, C. D. *et al.* Sex-linked hybrid sterility in a butterfly. *Evolution* **55**, 1631–1638 (2001).

[24] Naisbit, R. E., Jiggins, C. D., Linares, M., Salazar, C. & Mallet, J. Hybrid sterility, Haldane's rule and speciation in *Heliconius cydno* and *H. melpomene*. *Genetics* **161**, 1517–1526 (2002).

[25] Kost, S., Heckel, D. G., Yoshido, A., Marec, F. & Groot, A. T. A Z-linked sterility locus causes sexual abstinence in hybrid females and facilitates speciation in *Spodoptera frugiperda*. *Evolution* **70**, 1418–1427 (2016).

[26] Rosser, N. *et al.* Complex basis of hybrid female sterility and Haldane's rule in *Heliconius* butterflies: Z-linkage and epistasis. *Molecular Ecology* **31**, 959–977 (2022).

[27] Ae, A. S. A study of the *Papilio bianor* Group mainly based on hybridization (Lepidoptera, Papilionidae). *Tyo Ga* **41**, 13–19 (1990).

[28] Kitahara, H. & Shirai, K. Crossing experiments with *Papilio okinawensis* Fruhstorfer from Okinawa Island and *P. dehaanii* C. & R. Felder from central Honshu, Japan (Lepidoptera, Papilionidae). *Lepidoptera Science* **69**, 85–91 (2018).

[29] Ae, S. A. A study of hybrids between Japanese and Himalayan *Papilio* butterflies. *Special Bulletin of Lepidopterological Society of Japan* **2**, 75–107 (1966).

[30] Lu, S. *et al.* Chromosomal-level reference genome of Chinese peacock butterfly (*Papilio bianor*) based on third-generation DNA sequencing and Hi-C analysis. *GigaScience* **8**, giz128 (2019).

[31] Lees, D. C., Kremen, C. & Andriamampianina, L. A null model for species richness gradients: bounded range overlap of butterflies and other rainforest endemics in Madagascar. *Biological Journal of the Linnean Society* **67**, 529–584 (1999).

[32] Colwell, R. K. & Lees, D. C. The mid-domain effect: geometric constraints on the geography of species richness. *Trends in Ecology & Evolution* **15**, 70–76 (2000).

[33] Rosin, L. F., Chen, D., Chen, Y. & Lei, E. P. Dosage compensation in *Bombyx mori* is achieved by partial repression of both Z chromosomes in males. *Proceedings of the National Academy of Sciences* **119**, e2113374119 (2022).

[34] Huylmans, A. K., Macon, A. & Vicoso, B. Global dosage compensation is ubiquitous in Lepidoptera, but counteracted by the masculinization of the Z chromosome. *Molecular Biology and Evolution* **34**, 2637–2649 (2017).

[35] Rousselle, M., Faivre, N., Ballenghien, M., Galtier, N. & Nabholz, B. Hemizygosity enhances purifying selection: lack of fast-Z evolution in two Satyrine butterflies. *Genome Biology and Evolution* **8**, 3108–3119 (2016).

[36] Pinharanda, A. *et al.* Sexually dimorphic gene expression and transcriptome evolution provide mixed evidence for a fast-Z effect in *Heliconius*. *Journal of Evolutionary Biology* **32**, 194–204 (2019).

[37] Mongue, A. J., Hansen, M. E. & Walters, J. R. Support for faster and more adaptive Z chromosome evolution in two divergent Lepidopteran lineages. *Evolution* **76**, 332–345 (2022).

[38] Drinnenberg, I. A., deYoung, D., Henikoff, S. & Malik, H. S. Recurrent loss of CenH3 is associated with independent transitions to holocentricity in insects. *eLife* **3**, e03676 (2014).

[39] Senaratne, A. P. *et al.* Formation of the CenH3-deficient holocentromere in Lepidoptera avoids active chromatin. *Current Biology* **31**, 173–181 (2021).

[40] Ahmed, M. Z., Araujo-Jnr, E. V., Welch, J. J. & Kawahara, A. Y. *Wolbachia* in butterflies and moths: geographic structure in infection frequency. *Frontiers in Zoology* **12**, 1–9 (2015).

[41] Naveira, H. F. Location of X-linked polygenic effects causing sterility in male hybrids of *Drosophila simulans* and *D. mauritiana*. *Heredity* **68**, 211–217 (1992).

[42] Liénard, M. A., Araripe, L. O. & Hartl, D. L. Neighboring genes for DNA-binding proteins rescue male sterility in *Drosophila* hybrids. *Proceedings of the National Academy of Sciences* **113**, E4200–E4207 (2016).

[43] Presgraves, D. C. & Meiklejohn, C. D. Hybrid sterility, genetic conflict and complex speciation: lessons from the *Drosophila simulans* clade species. *Frontiers in Genetics* 1002 (2021).

[44] Vrana, P. B. *et al.* Genetic and epigenetic incompatibilities underlie hybrid dysgenesis in *Peromyscus*. *Nature genetics* **25**, 120–124 (2000).

[45] Brekke, T. D. *et al.* X chromosome-dependent disruption of placental regulatory networks in hybrid dwarf hamsters. *Genetics* **218**, iyab043 (2021).

[46] Jindra, M., Palli, S. R. & Riddiford, L. M. The juvenile hormone signaling pathway in insect development. *Annual Review of Entomology* **58**, 181–204 (2013).

[47] Okamoto, N. & Yamanaka, N. Nutrition-dependent control of insect development by insulin-like peptides. *Current Opinion in Insect Science* **11**, 21–30 (2015).

[48] Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

[49] Picard toolkit. https://broadinstitute.github.io/picard/ (2019).

[50] Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008 (2021).

[51] Korneliussen, T. S. & Moltke, I. NgsRelate: a software tool for estimating pairwise relatedness from next-generation sequencing data. *Bioinformatics* **31**, 4009–4011 (2015).

[52] Rastas, P. Lep-MAP3: robust linkage mapping even for low-coverage whole genome sequencing data. *Bioinformatics* **33**, 3726–3732 (2017).

[53] Broman, K. W. *et al.* R/qtl2: software for mapping quantitative trait loci with high-dimensional data and multiparent populations. *Genetics* **211**, 495–502 (2019).