Full Title:

The mesodermal source of fibronectin is required for heart morphogenesis and cardiac

outflow tract elongation by regulating cell shape, polarity, and mechanotransduction in

the second heart field.

Short Title:

Role of mesodermal fibronectin in cardiac development

Authors:

Cecilia Arriagada and Sophie Astrof*

Affiliation:

Department of Cell Biology and Molecular Medicine, Cardiovascular Research Institute,

Rutgers Biomedical and Health Sciences, 185 South Orange Ave, Newark, NJ, 07103,

USA.

*Author for correspondence: Sophie Astrof, Ph.D.

22   Phone: 617-429-8295

23   E-mail: sophie.astrof@rutgers.edu

Abstract

Failure in the elongation of the cardiac outflow tract results in congenital heart disease

due to ventricular septum defects and misalignment of the great vessels. The cardiac

outflow tract lengthens by the accretion of progenitors derived from the second heart field

(SHF). SHF cells in the splanchnic mesoderm are exquisitely regionalized and organized

into an epithelial-like layer forming the dorsal pericardial wall (DPW). Tissue tension within

the DPW, cell polarity and proliferation are requisite processes regulating the addition of

SHF-derived cells to the heart and elongation of the cardiac outflow tract. However, genes

regulating these processes are not completely understood. Using conditional

mutagenesis in the mouse, we show that fibronectin (*Fn1)* synthesized by the SHF is a

central regulator of the epithelial architecture in the DPW. Fn1 expression is enriched in

the anterior DPW and mediates outflow tract elongation by regulating cell polarity, shape,

cohesion, and mechanotransduction. Our studies establish that Fn1 synthesized

specifically by the mesoderm coordinates multiple cellular behaviors in the anterior DPW

requisite for the elongation of the cardiac outflow tract and embryonic viability.

Introduction

Congenital heart disease (CHD) is one of the most common birth defects. An estimated 40,000 newborns in the U.S. are affected by CHD each year (Tsao et al., 2022). Therefore, it is important to understand the cellular and molecular mechanisms regulating cardiac morphogenesis and the alterations that cause CHD. Several sources of cardiac progenitor cells have been identified (Zhang et al., 2021). Progenitors in the first heart field mainly contribute to the left ventricle, while progenitors in the second heart field (SHF) contribute to the right ventricle, the outflow tract (OFT), and the atria. Correct formation and elongation of the OFT are necessary for the proper morphogenesis of the arterial pole of the heart, e.g., for intraventricular septation and correct positioning of the great vessels relative to the cardiac chambers (Cortes et al., 2018). Defects in these processes are among the most common causes of CHD (Tsao et al., 2022). OFT forms as a single tube connecting the heart with the systemic circulation. In the mouse, the OFT elongates between embryonic day (E) 8.5 and E10.75 of development by the addition of the SHF cells to the nascent heart tube (Meilhac and Buckingham, 2018). The aberrant contribution of SHF cells to the OFT results in defective elongation of the cardiac OFT leading to aberrant septation and alignment of the arterial pole of the heart.

SHF cells in the dorsal pericardial wall (DPW) (Fig S1A), directly behind the forming heart, exhibit epithelial properties such as apicobasal polarity (Cortes et al., 2018; Francou et al., 2014). Aberrations in SHF cell polarity and cohesion in the DPW result in shortened OFT and CHD (Cortes et al., 2018; Li and Wang, 2018). For instance, *Tbx1*, a key transcription factor mutated in 22q11 deletion syndrome, mediates OFT elongation by

regulating SHF cell polarity, organization of the actin cytoskeleton, and epithelial tension

in the DPW (Francou et al., 2017; Francou et al., 2014). Similarly, mediators of non-

canonical Wnt signaling such as *Vngl2, Wnt5a,* and *Dvl* mediate OFT elongation by

regulating apicobasal polarity of SHF cells in the DPW, actin cytoskeleton, and cell

intercalation behaviors important for the movement of SHF cells in the DPW to join the

OFT (Li et al., 2019; Li et al., 2016; Li and Wang, 2018; Ramsbottom et al., 2014; Sinha

et al., 2015; Sinha et al., 2012).

In addition to being polarized, SHF cells in the DPW are regionalized along the anterior-

posterior and left-right axes (Bajolle et al., 2008; Kelly, 2023). For example, *Osr1, Hoxb1*,

and *Wnt5a* are expressed in the posterior DPW, while *Tbx1* is expressed in the anterior

(Stefanovic et al., 2020). This regionalization is important for OFT elongation and

morphogenesis (Li and Wang, 2018; Stefanovic et al., 2020). We noticed that the

expression of the extracellular matrix (ECM) glycoprotein *Fn1* was enriched in the anterior

DPW at the protein and *mRNA* levels. In our previous studies, we found that cardiac OFTs

were short in Fn1-null mutants (Mittal et al., 2013). Conditional mutagenesis indicated

that the expression of Fn1 and its major receptor integrin $\alpha5\beta1$ in the *Isl1* lineage was

required for OFT elongation (Chen et al., 2015). These experiments underscore the

important role of Fn1 in heart development. However, mechanisms by which Fn1

regulates OFT elongation have not yet been investigated.

Prior to heart formation, *Fn1* mRNA is broadly expressed in the paraxial and lateral plate

mesoderm of early (1-3 somite) murine embryos (Pulina et al., 2011). During heart

development (6-23 somites), *Fn1* mRNA continues to be expressed in the splanchnic

mesoderm, while its expression in differentiated cardiomyocytes is nearly lost (Mittal et

al., 2010). Mesodermal populations expressing *Fn1 mRNA* contain cardiac progenitors.

Since heart development is affected in global Fn1-null embryos and embryos with

conditional deletion of Fn1 in the *Isl1* lineage (Chen et al., 2015; Mittal et al., 2013), we

hypothesized that Fn1 synthesized by cardiac progenitors regulates heart development.

To investigate this hypothesis, we used the *Mesp1$^{Cre/+}$* knock-in strain to conditionally

ablate Fn1 expression in the anterior mesoderm. In this manuscript, we show that

mesodermal Fn1 coordinates multiple cellular behaviors regulating the accretion of SHF

cells from the DPW during OFT elongation. The deletion of the mesodermal Fn1 results

in aberrant cellular architecture in the anterior DPW cells contiguous with the elongating

OFT. Our studies indicate that Fn1 secreted by the mesoderm mediates OFT elongation

by regulating cardiac progenitor cell shape, apicobasal polarity, and

mechanotransduction.

Results

Mesodermal synthesis of Fn1 is required for cardiogenesis and embryonic viability.

The expression of the transcription factor Mesp1 marks the earliest known cardiac

progenitors; descendants of Mesp1-expressing cells contribute to all mesodermally-

derived cells of the heart (Saga et al., 1996; Saga et al., 1999). Therefore, we used the

Mesp1$^{Cre/+}$ strain to conditionally ablate *Fn1* expression from heart precursors. Although

Mendelian proportions of all genotypes were recovered at E9.5, we found some Fn1$^{flox/-}$;

Mesp1$^{Cre/+}$ mutant embryos at E9.5 with mild cardiac defects (S1B-D1 Fig). However, only

50% of the expected number of Fn1$^{flox/-}$; *Mesp1*$^{Cre/+}$ embryos were recovered at E10.75

(S1E-G Fig). All the recovered mutants at E10.75 had prominent cardiac edema,

defective heart, and were degenerating (S1E-F1 Fig). No Fn1$^{flox/-}$; *Mesp1*$^{Cre/+}$ mutants

were recovered after E13.5 days of gestation (S1G Fig). These data indicated that the

expression of *Fn1* in the Mesp1 lineage is required for heart development and embryonic

viability.

Mesodermal Fn1 regulates the elongation of the cardiac OFT.

The shape of the heart in E10.75 mutants suggested that mesodermal Fn1 played a role

in the development of the cardiac OFT (S1E-F1 Fig). To test this hypothesis, we first

examined embryos at E8.5 (8-12 somites), when the OFT begins to form. At this stage of

development, we did not find alterations in any cardiac structures, and the nascent OFTs

in mutants and controls were of similar lengths (S2 Fig). The heart remained similar in

shape in control and mutant embryos at 18-19 somites, and the OFT length was

comparable among control and mutant genotypes at this stage (Fig. 1A-D). However, at

20-25 somites, OFT length in mutant embryos was significantly shorter than in controls,

and both the proximal and the distal portions of the OFT were affected at this stage (Fig.

1E-H). In control embryos, the right and the left ventricles align in the plane perpendicular

to the embryonic anterior-posterior axis (Fig. 1I). In mutants with shortened OFT, this

alignment is distorted and the right ventricle is shifted anteriorly due to the decreased

length of the OFT (Li et al., 2016). The alignment of the right ventricle is similarly distorted

in Fn1$^{flox/-}$; *Mesp1*$^{Cre/+}$ mutants (Fig. 1J-K). Together, these data indicate that mesodermal

Fn1 mediates cardiogenesis by regulating OFT elongation.

Mesodermal Fn1 regulates the epithelial architecture of the dorsal pericardial wall.

Cardiac OFT elongates through the addition of cardiac precursors from the SHF in the

first and second pharyngeal arches and the splanchnic mesoderm of the DPW (Cortes et

al., 2018; Li and Wang, 2018; Meilhac and Buckingham, 2018). The length of the OFT

remained comparable among controls and mutants by the ~19th somite stage of

development, suggesting that the addition of SHF cells from the pharyngeal arches was

not affected by the deletion of the mesodermal source of Fn1 (Meilhac and Buckingham,

2018). However, cells in the DPW of E8.5 mutants were disorganized (Fig. 2). Instead of

a single layer of epithelial-like cells, as seen in control Fn1$^{flox/+}$; *Mesp1*$^{Cre/+}$ embryos, cells

in the anterior DPW of Fn1$^{flox/-}$; *Mesp1*$^{Cre/+}$ mutants were multilayered (compare Fig. 2A-

A1 with Fig. 2E-E1). In controls, DPW cells appeared ellipsoid, regularly oriented along

the dorsal-ventral axis, and neatly stacked along the entire length of the DPW (Fig. 2A-

D). In the mutants, SHF cells were more spherical in the anterior DPW (Fig. 2E-H). To

quantify differences in cell shapes in 2D and 3D, we measured cell circularity and ellipticity

in controls and mutants (Fig. 2I-L). Circularity is the ratio of cell width to cell length.

Elongated cells have circularity indexes that deviate from 1, while cuboid or spherical cells

have circularity indexes close to 1. Differences in circularity indexes among SHF cells in

the anterior DPW were statistically significant between controls and mutants (Fig. 2I).

However, in the posterior DPW, the circularity of control and mutant cells were

comparable (Fig. 2J). To determine if 3D cell shape was also affected in the mutants, we

segmented individual SHF cells in the anterior DPW and measured ellipticity. Ellipticity

was higher in control embryos than in mutants, indicating that in the absence of

mesodermal Fn1, SHF cells in the anterior DPW were more spherical than in controls (Fig

2K). The volume of individual SHF cells was not affected in the mutants at this stage (Fig.

2L). These differences in 2D and 3D cell shapes were observed at E8.5 and preceded

the shortening of the cardiac OFT. The aberrant epithelial architecture of the anterior

DPW persisted at E9.5, at the time when defective OFT elongation became detectable in

the mutants (S3A-K Fig). In addition to altered cell shape, cell volume was also decreased

in E9.5 mutants (S3L Fig). Together, these data suggest that alterations in SHF cell shape

in the anterior DPW give rise to defective OFT elongation in the mutants.

Mesodermal Fn1 regulates SHF cell orientation and apicobasal polarity in the

anterior DPW.

To determine mechanisms by which mesodermal Fn1 regulates the cellular architecture

of the DPW, we tested the hypothesis that Fn1 was important for apicobasal cell polarity.

At E8.5, the Golgi apparatus was positioned toward the apical surface of SHF cells along

the entire length of the DPW (Fig. 3A-A4). In contrast, in the mutants, the orientation of

the Golgi apparatus was disorganized in the anterior DPW, and in some SHF cells, the

Golgi apparatus faced toward the basal side (Fig. 3B-B1, B3, 3C, and 3D). However, in

the posterior DPW, the Golgi apparatus was positioned toward the apical cell side in

controls and mutants alike (compare Fig. 3A2 with 3B2, quantified in Fig.3A4, B4, and

Fig. 3E).

Similar to disorganized Golgi orientation, markers of the apical cell surface were re-

distributed in the mutants and were also present at the basal surface of SHF cells in the

anterior DPW (Fig. 4). These proteins included podocalyxin (compare Fig. 4A1 with Fig.

4B1 quantified in Fig. C and Fig. 4A2 with Fig. 4B2 quantified in Fig. 4D) and atypical

PKC ζ (aPKC ζ) (compare Fig. 4E-F with Fig. 4H-I quantified in Fig. 4K). Basal cell

properties of SHF cells were not as distorted as apical in the mutants: the expression of

laminin γ1 was disorganized only in 1 of 3 E8.5 mutants (Fig. 4G, J) and cellular

protrusions on the basal side of SHF cells were present in equal numbers in controls and

mutants at E9.5 (S4 Fig). These studies suggest the expansion of the apical surface of

SHF cells in the anterior DPW, while the basal cell surface was properly specified.

Changes in apical cell polarity were observed at E8.5 before defects in the OFT

elongation were detectable; however, by E9.5, apicobasal cell polarity recovered in the

mutants (S5 Fig). Altogether, these data show that changes in SHF cell shape and polarity

within the anterior DWP preceded changes in the OFT elongation and were likely causes

of defective OFT elongation in the mutants (Cortes et al., 2018; Li and Wang, 2018).

Mesodermal Fn1 regulates cell cohesion in the SHF and collective cell migration.

Adherens junctions form between the apical and basal cell surfaces (Nelson, 2003).

Therefore, we examined the localization of N-Cadherin in the DPW and found that N-

cadherin levels were downregulated in the DPW of the mutants (Fig. 5A-D1). To quantify

these changes, we used mouse embryo fibroblasts (MEFs) as a model system and

generated four lines of MEFs (S6 Fig): 1) Fn1[flox/+] MEFs, in which wild-type Fn1 protein

can be produced from both *Fn1* alleles, 2) Fn1[flox/-] MEFs, in which wild-type Fn1 protein

is generated from the floxed *Fn1* allele, and "-" *Fn1* allele is a null allele, 3) Fn1[/+] MEFs,

in which the floxed region of *Fn1* allele was resulting in one Fn1-null allele (Δ) (Sakai et

al., 2001) and one wild-type *Fn1* allele in these MEFs, and 4) Fn1$^{\Delta/-}$ MEFs, which are

Fn1-protein-null (S6 Fig). Immunofluorescence and quantitative Western blot analyses

showed that total and cell-surface levels of N-Cadherin were downregulated in Fn1$^{\Delta/-}$

MEFs (Fig. 5E-O1). We observed similar levels of N-cadherin between Fn1$^{flox/+}$ and Fn1$^{\Delta/+}$

MEFs, indicating that the decrease in N-cadherin in Fn1$^{\Delta/-}$ MEFs was not due to and non-

specific effect of the Cre infection (Fig. 5E-G and Fig5H-O1). Cell cohesion is important

for the collective cell migration (Friedl and Mayor, 2017). Since N-Cadherin was

downregulated in Fn1$^{\Delta/-}$ MEFs (Fig. 5E-K), we used this model to determine whether the

synthesis of Fn1 by MEFs was important for their collective cell migration. Monolayer

scratch assay showed that, indeed, scratch wounds made in Fn1$^{\Delta/-}$ MEF monolayers

closed slower than in Fn$^{flox/+}$, Fn$^{flox/-}$, or Fn1$^{\Delta/+}$ MEFs (S7 Fig). Downregulation of N-

Cadherin in the cardiac progenitors within the anterior DPW causes their precautious

differentiation (Soh et al., 2014). Consistent with these findings, we observed an

expanded expression domain of the sarcomeric myosin heavy chain (assayed by the MF-

20 antibody) in the anterior DPW of our mutants (S8 Fig). Together, these studies suggest

that mesodermal Fn1 regulates cell cohesion and collective cell migration of SHF cells,

facilitating the maintenance of the progenitor cell state in the DPW and the cellular

movement from the DPW to the OFT during its elongation.

Mesodermal Fn1 regulates mechanotransduction in the dorsal pericardial wall.

Nuclear localization of YAP is regulated by the mechanical tension in the DPW and is

important for the proliferation of DPW cells and OFT elongation (Francou et al., 2017).

YAP is a central transducer of the mechanical signals (Dupont, 2016). Defective shape

and orientation of SHF cells in the anterior DPW of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants suggested

that mutant SHF cells in the anterior DPW are deficient in generating, sensing, or

responding to mechanical cues. The remodeling of the actin cytoskeleton and nuclear

localization of YAP are known to be facilitated by cell adhesion to Fn1 *in vitro* (Kim and

Gumbiner, 2015; Singh et al., 2010). To test whether mesodermal Fn1 was required for

cytoskeletal remodeling and nuclear localization of YAP in the DPW, we first stained

control and Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutant embryos to detect filamentous (F)-actin. We found

that F-actin organization at E9.5 was disrupted specifically in the anterior DPW of the

mutants (Fig. 6): The orientation of F-actin filaments was disorganized (Fig. 6A-J) and F-

actin filaments were shorter in the anterior DPW (Fig. 6K). The length and orientation of

F-actin in the posterior and middle DPW were comparable between genotypes (S9 Fig).

Changes in cell shape and actomyosin cytoskeleton serve are faithful readouts of altered

mechanical tension in epithelial tissues (Luciano et al., 2022). Cells in the anterior DPW

are apically constricted and have a smaller apical surface area than cells in the posterior

DPW (Francou et al., 2017). Consistent with these findings, we observed the enrichment

of F-actin and an activated form of Myosin Light Chain 2 phosphorylated on Ser19

(pMLC), capable of driving the actomyosin contractility (Somlyo and Somlyo, 1994) at the

apical cell surface of the DPW (Fig. 7A-B2, E-F2). In contrast, apical enrichment of F-

actin and pMLC was lost in Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants (Fig. 7, compare Fig. 7B1 with

7D1 and Fig. 7F1 with 7H1). These changes were concordant with decreased nuclear

YAP localization in the anterior DPW of mutants (Fig. 8, compare Fig. 8B1 with 8C1,

quantified in Fig. 8D). Nuclear localization of YAP in the posterior DPW was not affected

(compare Fig. 8B2 with 8C2, quantified in Fig. 8E).

Cell adhesion to Fn1 regulates the nuclear localization of YAP (Kim and Gumbiner, 2015).

YAP, in turn, is a central integrator of mitogenic, biomechanical, and cell polarity cues

important for cell proliferation (Borreguero-Munoz et al., 2019; Fan et al., 2013). Inhibition

of YAP causes decreased proliferation of DPW cells in developing embryos (Francou et

al., 2017). Consistent with these studies, cell proliferation (but not cell survival) was

decreased in the DPW and the heart of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants, while the proliferation

of adjacent non-mesodermal tissues was not affected (Fig. 9, S10-S12 Figs). Altogether,

altered (more rounded) cell shape, the re-distributed of actomyosin cytoskeleton from the

apical surface, decreased cell volume, and decreased levels of nuclear YAP are all

indicators of an altered mechanical microenvironment (Perez Gonzalez et al., 2018) in

the anterior DPW of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants.

Mesodermal Fn1 is the essential source of Fn1 in the anterior DPW.

We next asked whether defects observed in the DPW of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants were

due to decreased levels of Fn1 or were caused by the absence of Fn1 protein made

specifically in the mesoderm. *Fn1 mRNA* and protein are dynamically and non-uniformly

expressed in developing embryos (Chen et al., 2015; Mittal et al., 2010), and previous

studies demonstrated cell-autonomous roles of Fn1 during embryonic development,

angiogenesis, and cell migration (Cseh et al., 2010; Turner et al., 2017; Wang and Astrof,

2016; Zimmerman et al., 2017). Cells in the DPW and the pharyngeal endoderm are

closely apposed in mid-gestation embryos, and *Fn1* mRNA is synthesized by both tissues

(Fig. 10A-B2). At E8.5, Fn1 protein is present in the DPW and the heart of controls, but

it is downregulated in Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants (S13A-H Fig).

Although there were no differences in Fn1 protein levels between the anterior and

posterior DPW in control embryos at E8.5 (the ratio of Fn1 protein in aDPW to that in

pDPW is close to 1, S13E Fig), at E9.5, *Fn1* mRNA was enriched nearly 2-fold in the

anterior DPW relative to the posterior (Fig. 10A-C). This enrichment is particularly well-

visualized in control Fn1$^{flox/+}$; *Mesp1$^{Cre/+}$* embryos (Fig. 10D-F2). In these embryos, one

allele of Fn1 is deleted in the mesoderm, resulting in a ~50% decrease in *Fn1 mRNA* in

the DPW relative to the wild-type (Fig. 10G). In Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants, levels of *Fn1*

mRNA in the entire DPW were low (Fig. 10H-I2 and 10G, J). However, *Fn1 mRNA*

remained synthesized by the endoderm (Fig. 10I1-I2, yellow arrowheads). The loss of

Fn1 mRNA in the mesoderm but not in the endoderm is consistent with the known

specificity of Cre expression in the *Mesp1$^{Cre/+}$* strain (Saga et al., 1996; Saga et al., 1999)

Similar to *Fn1* mRNA , Fn1 protein was also enriched in the anterior DPW relative to the

posterior (Fig. 10K-L). Despite the efficient downregulation of *Fn1 mRNA* in the

mesoderm of the DPW, *Fn1* mRNA synthesis from the endoderm resulted in a similar

accumulation of Fn1 protein in the anterior DPW in controls and mutants at E9.5 (Fig.

10M-P). Moreover, the Fn1 protein was localized between the germ layers and enriched

in the anterior DPW in controls and mutant alike (Fig. 10L). But despite the presence of

similar levels of Fn1 protein, Fn1 derived from the endoderm did not rescue defects in the

architecture of the DPW in the mutants.

To rule out the possibility that differences in Fn1 protein levels contributed to defects in

the DPW in Fn1$^{flox/-}$; $Mesp1^{Cre/+}$mutants, we deleted Fn1 using $Sox17^{2A-iCre/+}$ strain, in

which Cre recombinase is expressed in the endoderm and endothelium (Engert et al.,

2009). In Fn1$^{flox/-}$; $Sox17^{2A-iCre/+}$mutants, *Fn1 mRNA* was downregulated in the endoderm,

but it was still synthesized by the SHF cells of the DPW (compare S14A-A1 Fig with S14D-

D1 Fig). And although the levels of Fn1 protein underlying the DPW were substantially

reduced in Fn1$^{flox/-}$; $Sox17^{2A-iCre/+}$mutants (compare S14B-B1 Fig with S14E-E1 Fig), the

epithelial organization and the expression of N-cadherin were not as affected in Fn1$^{flox/-}$;

$Sox17^{2A-iCre/+}$mutants as they were in Fn1$^{flox/-}$; $Mesp1^{Cre/+}$mutants (S14C-C1, S14F-F1

Fig). Together, these data indicate that 1) Fn1 synthesized by the mesoderm rather than

total levels of Fn1 are required for the proper cellular architecture of the DPW, 2) the

presence of Fn1 protein synthesized by the endoderm is not sufficient to rescue the

absence of mesodermal Fn1, and 3) that synthesis of Fn1 by the DPW is specifically

required for the proper cellular architecture and mechanotransduction in the anterior DPW

to mediate OFT elongation, heart development, and embryonic viability (Fig. 11).

Discussion

Here we show that Fn1 synthesized by the mesoderm is required for proliferation, cell

shape, cell polarity, orientation, and the orderly stacking of SHF cells in the anterior DPW.

In Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$*mutants, defects in these properties occur prior to the observable

deficiencies in the OFT elongation, indicating that defects in DPW architecture cause

aberrant OFT elongation in the absence of Fn1 synthesized by the mesoderm.

Cell adhesion to Fn1 facilitates the remodeling of the actomyosin cytoskeleton and

synergizes with growth factor signaling to promote the sustained activation of Erk, the

expression of cyclin D1, and cell cycle progression (Bohmer et al., 1996; Roovers and

Assoian, 2003; Walker et al., 2005). While cellular architecture was mainly affected in the

anterior DPW in Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$*embryos, cell proliferation was affected along the

entire length of the DPW as well as in the heart, where the mesoderm is the sole source

of Fn1 protein. Therefore, it is possible that in the DPW and the heart, Fn1 regulates

proliferation via multiple mechanisms; in the anterior DPW, the defect in cell proliferation

may be secondary to changes in actomyosin cytoskeleton, cell shape, and the nuclear

localization of YAP, while in the posterior DPW and the heart, Fn1 may regulate

proliferation directly.

At the early stages of OFT formation, at E8.5, *Fn1* mRNA and protein are synthesized by

the pharyngeal endoderm and the DPW. At this stage, the Fn1 protein is uniformly

localized at the basal side along the entire length of the DPW, between the mesoderm

and the endoderm. Fn1 protein in this region is decreased in Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$*mutants.

It is possible, then, that the onset of defects in cell proliferation and DPW architecture

results from the diminished levels of Fn1 at E8.5 and that the anterior DPW cells are

especially sensitive to Fn1 levels for maintaining the proper cellular architecture.

However, the depletion of Fn1 protein in Fn1$^{flox/-}$; *Sox17$^{2A-iCre/+}$* mutants, where Fn1 was

ablated in the endoderm, was even more severe than in Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants.

Despite the dramatic decrease in the levels of Fn1 protein underlying the DPW in Fn1$^{flox/-}$

; *Sox17$^{2A-iCre/+}$*, the cellular architecture of the DPW and the expression of N-Cadherin at

cell-cell junctions in Fn1$^{flox/-}$; *Sox17$^{2A-iCre/+}$* mutants were comparable to controls. These

results suggest that the decrease in Fn1 levels at E8.5 cannot solely explain cellular

defects in the anterior DPW of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutants.

The OFT and DPW are exquisitely regionalized, with distinct genetic programs regulating

OFT elongation and the development of the arterial pole of the heart (Bajolle et al., 2006;

Bajolle et al., 2008; Kelly, 2023; Li and Wang, 2018; Rammah et al., 2022; Ramsbottom

et al., 2014; Stefanovic et al., 2020; Theveniau-Ruissy et al., 2008; Xia et al., 2019). We

found that Fn1 is an important factor in this regionalization and provided additional

evidence that cellular architecture along the anterior-posterior axis is differentially

regulated.

In the caudal portion of the DPW, actin cytoskeletal dynamics and cell intercalation are

regulated by non-canonical Wnt signaling mediated by *Wnt5a* and *Dvl* (Li et al., 2019; Li

et al., 2016; Sinha et al., 2015; Sinha et al., 2012). Cell intercalations in the posterior

DPW lead to the overall displacement of the DPW cells toward the anterior pole, "pushing"

them into the OFT, facilitating the OFT elongation (Li and Wang, 2018). In the OFT

immediately adjacent to the anterior DPW, *Vngl2* and *Hand2* regulate cell-cell and cell-

ECM interactions mediating intercellular intercalations and the lengthening of the OFT

(Ramsbottom et al., 2014; Xia et al., 2019). In the anterior DPW, cellular architecture is

regulated by *Tbx1,* where its expression is enriched (Francou et al., 2014; Stefanovic et

al., 2020). Deletion of *Tbx1* leads to defects in cell-extracellular matrix interactions, cell

shape, organization, polarity, and actin cytoskeletal dynamics (Alfano et al., 2019;

Francou et al., 2014), resulting in the altered tension in the DPW and defective OFT

elongation (Francou et al., 2017). Similar to *Tbx1*, *Fn1* mRNA synthesis and protein

localization are enriched in the anterior DPW at E9.5, where Fn1 regulates cell shape,

orientation, and biomechanical signaling. It is interesting that *Tbx1* regulates apical

constriction in the anterior DPW (Francou et al., 2017). Here, we show that *Fn1* regulates

the apical enrichment of actomyosin fibrils. Apical constriction is a known regulator of the

epithelial morphogenesis (Sawyer et al., 2010). Removal of the OFT leads to the

relaxation of cellular tension in the DPW (Francou et al., 2017); thus, we speculate that

the apical constriction at the anterior DPW is important for the generation of tissue tension

and, possibly, for the transitioning of DPW cells through the bent region connecting the

anterior DPW and the OFT.

The ablation of mesodermal Fn1 led to cell rounding, decreased cell volume,

downregulation of N-cadherin, and disrupted apical enrichment of actomyosin cables,

which altogether indicate diminished mechanical tension in the anterior DPW (Perez

Gonzalez et al., 2018). Nuclear accumulation of YAP was also disrupted in the mutants.

Cell adhesion to Fn1 regulates the activity of the Hippo pathway and promotes nuclear

localization of YAP (Kim and Gumbiner, 2015). However, the nuclear localization of YAP

is also affected by cell shape, and nuclear YAP is diminished upon cell rounding (Dupont,

2016). In Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$*embryos, defects in cell shape and orientation are already

noted at E8.5 before the nuclear accumulation of YAP is detectable in these cells

(Francou et al., 2017). Therefore, our current experiments cannot discern whether

signaling by Fn1 signaling in the DPW regulates YAP nuclear translocation directly or

indirectly through the regulation of cell shape.

In *Xenopus Laevis* embryos, disruption of cadherin-mediated adhesions interferes with

Fn1 fibrillogenesis, resulting in tissue thickening and the shortening of the anterior-

posterior axis due to defects in cell shape, intercellular intercalations, and convergent

extension (Dzamba et al., 2009; Hirsh et al., 2018; Weber et al., 2011). Conversely,

disruption of Fn1 fibrillogenesis in fish and frog embryos leads to defective planar cell

polarity, cell shape, cadherin-mediated adhesion, and cell intercalation (Garavito-Aguilar

et al., 2010; Marsden and DeSimone, 2001; Marsden and DeSimone, 2003; Rozario et

al., 2009; Trinh and Stainier, 2004; Trinh et al., 2005). Our data are consistent with these

findings, indicating an evolutionary-conserved role of Fn1 and the crosstalk between cell-

cell and cell-ECM adhesions during tissue morphogenesis and cardiac development.

Although Fn1 synthesis is abolished in the mesoderm of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* embryos, the

levels of Fn1 protein at the basal side of the DPW recovered by E9.5, likely due to the

continued synthesis of *Fn1* mRNA by the adjacent endoderm. However, despite similar

levels of Fn1 protein underlying the DPW and the ~2-fold enrichment of Fn1 protein in the

anterior DPW of Fn1$^{flox/-}$; *Mesp1$^{Cre/+}$* mutant embryos, Fn1 synthesized by the endoderm

(and by any other tissues) did not rescue defects in N-cadherin localization and cellular

architecture of the anterior DPW in these mutants, causing their demise by E10.75.

These results suggest that Fn1 synthesized by the mesoderm regulates epithelial

properties of the anterior DPW in a cell-autonomous manner. Altogether, our studies

support the notion that during morphogenesis, Fn1 regulates cell behaviors locally, in a

cell-autonomous manner, e.g., (Cseh et al., 2010; Turner et al., 2017; Wang and Astrof,

2016), and that Fn1 expressed by the anterior mesoderm is a central player coordinating

multiple cellular behaviors regulating the development of the DPW and the elongation of

the cardiac OFT.

Materials and Methods

Antibodies:

See Supplemental Table 1 for antibody information.

Ethics Statement

All experiments involving vertebrate animals were approved by the Rutgers University Institutional Animal Care and Use Committee and were performed following federal guidelines for the humane care of animals.

Mouse strains:

C57BL/6J mice (cat # 0664) and Rosa$^{mTmG/mTmG}$ mice, Gt(ROSA)26Sor$^{tm4(ACTB-tdTomato,-EGFP)Luo}$ mice (Cat# 037456) (Muzumdar et al., 2007), were purchased from Jackson Laboratories. Fn1$^{flox/flox}$ mice were a gift from Reinhardt Fassler (Sakai et al., 2001). Mesp1$^{Cre/+}$ knock-in mice (Saga et al., 1996; Saga et al., 1999) were obtained from Dr. Yumiko Saga. Fn1$^{+/-}$ strain (George et al., 1997; George et al., 1993) was a gift from Richard Hynes. Sox17$^{2A-iCre}$ mice were a gift from Heicko Lickert (Engert et al., 2009). For experiments, Fn1$^{+/-}$ mice were crossed with Mesp1$^{Cre/+}$ knock-in mice to generate Fn1$^{+/-}$; Mesp1$^{Cre/+}$ males which were propagated by mating with C57BL/6J females. To obtain embryos, Fn1$^{+/-}$; Mesp1$^{Cre/+}$ males were crossed with Fn1$^{flox/flox}$; Rosa$^{mTmG/mTmG}$ females, and pregnant females were dissected at days specified in the figures. Mice and embryos were genotyped according to published protocols (Engert et al., 2009; George et al., 1993; Saga et al., 1999). Embryos were staged by counting somites.

Cell culture and generation of cell lines:

Mouse embryo fibroblasts (MEFs) were isolated from E13.5 embryos resulting from the

mating of Fn1$^{+/-}$ mice with Fn1$^{flox/flox}$; ROSA$^{mTmG/mTmG}$ mice.

MEFs were cultured on plates pre-coated with 0.1% gelatin (Sigma, #G1890) dissolved

in H_2O and maintained in complete medium, which consisted of Dulbecco's Modified

Eagle's medium (DMEM) with 4.5 g/L glucose, L-glutamine & sodium pyruvate (Corning,

#10-013-CM) supplemented with 100 U/ml penicillin, 100 μg/ml streptomycin (Gibco,

#15140-122), 1% v/v GlutaMax (L-glutamin, Gibco, #35050061) and 10% v/v fetal bovine

serum (FBS, Gemini Biosciences, #100-106) in a humidified incubator with 5% CO_2 at

37°C.

To generate MEFs lacking Fn1 (Fn1-null), or control, Fn1-heterozygous (Fn1-het) MEFs,

Fn1$^{flox/+}$; ROSA$^{mTmG/+}$ and Fn1$^{flox/-}$; ROSA$^{mTmG/+}$ MEFs were infected with

adenoviruses encoding Cre recombinase, Ad-Cre-IRED-GFP (Vector Biolabs, #1710).

Cre-induced recombination not only leads to the ablation of Fn1 by recombining the floxed

Fn1 sequence but also induces the expression of membrane-tethered GFP in these cells.

For this experiment, $3x10^4$ MEFs were plated on gelatin-coated wells of 6-well plates. 24

hours later, cells were rinsed twice with Phosphate Buffered Saline 1X Sterile Solution

(PBS, RPI Research Products, # P10400-500) and infected with Cre adenovirus diluted

to the multiplicity of infection of 500 in DMEM without FBS. 20 hours post-infection, the

medium was replaced by the complete medium. Although each of these cell types carries

one ROSA^{mTmG} allele, only cells infected with Ad-Cre are constitutively and permanently

marked by the expression of GFP due to the recombination of the reporter locus. GFP

expression from the adenovirus is transient and dissipates after three days post-infection.

After expansion, Ad-Cre-treated cells were sorted to generate Fn1$^{\Delta/-}$ (Fn1-null, GFP+),

Fn1$^{flox/-}$ (Fn1-heterozygous, GFP-), Fn1$^{\Delta/+}$ (Fn1-heterozygous, GFP+), and Fn1$^{flox/+}$ (Fn1-

wild type, GFP-) MEF lines. Western blot analysis was used to assay Fn1 expression.

MEF protein lysates:

Cells were plated in 6-well dishes at a density of 2×10^5 cells per well. After 48 hours, cells

were washed twice with cold PBS supplemented with 0.1 mM CaCl$_2$ and 1 mM

MgCl$_2$ (PBS-Ca^{2+}/Mg^{2+}) and lysed with 400 μl of RIPA lysis buffer (50 mM Tris-Cl, 150

mM NaCl, 2 mM EDTA, 1% v/v NP-40, 0.5% w/v sodium deoxycholate, 0.1% w/v SDS,

1X protease inhibitor cocktail (Cell Signaling Technology, #5871), pH 8.0).

The samples were centrifuged at 16,000 × g for 15 min at 4°C.

Protein concentration was determined using BCA protein assay (Pierce™ BCA Protein

Assay Kit, #23225). Then, lysis buffer was added to equalize protein concentration across

samples. 4xNuPAGE™ LDS Sample buffer (Thermo Scientific, #NP0008) with 10% β-

mercaptoethanol (Sigma, #63689) was added to each sample to the final concentration

of 1X NuPAGE and 2.5% β-mercaptoethanol. Samples were heated to 95°C for 5 minutes

prior to loading on gels.

Immunoblotting and densitometry quantification:

For immunoblotting, samples were resolved using 4-12% Tris-Glycine gel (Invitrogen,

#XP04120BOX) at 130 V in 1x Tris-Glycine SDS Running Buffer (Invitrogen, # LC2675)

for 1 hour 30 minutes. Proteins were then transferred to nitrocellulose membranes with a

0.2 µm pore size (Bio-Rad, # 1620122) and 1x Tris-Glycine, 15% methanol transfers

buffer (10x Tris-Glycine buffer: 30g Tris Base. 144 g Glycine in 1L of ddH20, pH: 8.5).

The transfer was carried out at 90 V for 2 hours in ice.

After the transfer, the nitrocellulose membranes were blocked with Intercept Blocking

Buffer (Li-Cor, # 927-60001) for 1 hour. Primary antibodies were diluted in Intercept

Antibody Diluent (Li-Cor, # 927-65001) and incubated overnight at 4°C. After 24h, the

nitrocellulose membranes were washed three times with PBS containing 0.1% Tween for

10 min. We used IRDye 680RD Donkey anti-Rabbit IgG (#926-68073), and IRDye

800CW Donkey anti-Mouse IgG Secondary Antibody (#926-32212) as secondary

antibodies in a 1:15000 dilution in Intercept Antibody Diluent (Li-Cor, # 927-65001). The

nitrocellulose membranes were incubated with the secondary antibodies for 1 hour at

room temperature. Finally, membranes were washed three times with PBS 0.1% Tween

for 10 minutes and imaged using Li-Cor Odyssey 9120 Gel Imaging System (#ODY-

2425). The density of unsaturated pixels was determined using ImageJ software (version

2.1.0/1.53c4). For each condition, protein bands were quantified from at least three

independent experiments.

Cell surface biotinylation:

3.5×10^5 cells were plated on gelatin-coated 10 cm plates. After 48 hours of growth, cells

were washed twice with ice-cold PBS supplemented with 0.1 mM $CaCl_2$ and 1 mM $MgCl_2$

(PBS-Ca^{2+}/Mg^{2+}), followed by incubation with 1 mM EZ-Link Sulfo-NHS-LC-Biotin

(Thermo Scientific, #21335) in PBS-Ca^{2+}/Mg^{2+} for 30 min at 4°C. The biotinylation

reaction was quenched by incubating cells with Tris-buffered solution (50 mM Tris-HCl,

pH 7.4) for 10 min at 4°C. After two subsequent washes with PBS-Ca^{2+}/Mg^{2+}, cells were

lysed with 600 µl of RIPA lysis buffer (50 mM Tris-Cl, 150 mM NaCl, 2 mM EDTA, 1% v/v

NP-40, 0.5% w/v sodium deoxycholate, 0.1% w/v SDS, 1X protease inhibitor cocktail (Cell

Signaling Technology, #5871), pH 8.0). To remove insoluble material, cell extracts were

clarified by centrifugation at 16,000 x g for 15 min at 4°C. Soluble biotinylated protein

concentration was determined using BCA protein assay (Pierce™ BCA Protein Assay Kit,

#23225). 300 µg of protein per sample was incubated with 40 µl of Neutravidin-Agarose

beads (Thermo Scientific, # 29202) pre-washed with RIPA lysis buffer. After 1 hour of

incubation at 4°C, the Neutravidin-Agarose beads were washed 3 times with RIPA lysis

buffer containing 1X protease inhibitor cocktail. 120 µl of 1X NuPAGE™ LDS Sample

buffer (Thermo Scientific, #NP0008) with 2.5% β-mercaptoethanol (Sigma, #63689) and

1X protease inhibitor cocktail was added to beads and resolved by SDS-PAGE (Arriagada

et al., 2020).

Immunofluorescence microscopy:

$2x10^4$ cells were plated on gelatin-coated #1.5 round glass coverslips (Fisher Scientific,

t#12-545-81) in 24-well dishes. After for 48 hours, cells were washed with PBS that was

pre-warmed to 37°C for 5 minutes and fixed with 4% PFA (Fisher Scientific # 50-980-487,

diluted in PBS) for 20 minutes at room temperature. Then, cells were washed three times

with PBS for 5 minutes each and permeabilized with PBS containing 0.1% Triton-X 100

(PBST) for 15 minutes. Cells were blocked with 5% Donkey serum (Sigma, #d9663) in

PBST (blocking solution) for 30 minutes. After blocking, cells were incubated with the

primary antibodies diluted in blocking solution overnight at 4°C. Following 3 washes in

PBST for 10 minutes each, cells were incubated with secondary antibodies diluted in

blocking solution for 60 minutes at room temperature. Finally, cells were washed three

times with PBST for 10 minutes, mounted using a 50% v/v Glycerol, 50% v/v Methanol

solution.

Wound healing migration assay:

1×10^4 cells were plated on gelatin-coated 8-well ibidi dishes (Ibidi, #80827) for 72 hours.

When cells were confluent, the monolayer was scratched with a sterile 10 µl pipette tip.

Cell debris was washed out with PBS, and incubated with imaging medium 100 µl

FluoroBrite DMEM (Thermo Fisher Scientific, #A1896701) supplemented with 100 U/ml

penicillin, 100 µg/ml streptomycin (Gibco, #15140-122), 1% v/v GlutaMax (L-glutamin,

Gibco, #35050061) and 2% v/v fetal bovine serum (FBS, Gemini Biosciences, #100-106).

Images were acquired right after performing the scratch and then after 14 hours. % of

wound closure was measured. % of Wound closure = *$100*(A_{t=0} - A_{t=14})/ A_{t=0})$*

Where $A_{t=0}$ is the cell-free area at t=0 and $A_{t=14}$ is the cell-free area at t=14 hrs.

Whole Mount Immunofluorescence staining:

Timed matings were set up to obtain embryos at specific embryonic stages. Vaginal plugs

were checked daily, and noon of the day the plug was found was an embryonic day (E)

0.5 of gestation. Embryo dissections were aided using a Zeiss Stemi 2000C

stereomicroscope. Embryos were dissected into ice-cold PBS; yolk sacs were kept for

genotyping. After dissection, embryos were fixed with 4% PFA overnight at 4°C. Then,

embryos were washed two times with ice-cold PBS and permeabilized with PBS

containing 0.1% Triton- X 100 (PBST) overnight at 4°C. Embryos were blocked with 10%

Donkey serum in PBST (blocking solution) overnight at 4°C. After blocking, embryos were

incubated with the primary antibodies diluted in blocking solution for 3 days at 4°C.

Following four 1 hour-washes with PBST, embryos were incubated with secondary

antibodies diluted in blocking solution for 3 days at 4°C. Finally, after four washes with

PBST for 1 hour each, embryos were embedded in 1% agarose (Bio-Rad Laboratories,

#1613101), and cleared using methanol and Benzyl Alcohol (Sigma, # B-1042)/ Benzyl

Benzoate (Sigma, #B-6630) as described (Ramirez and Astrof, 2020). For imaging,

embryos were a placed between two #1.5 coverslips (VWR, #16004-312) separated by a

rubber spacer (Grace Bio Labs, # 664113).

Whole Mount Immunofluorescence in situ hybridization: After dissection, embryos were

fixed with 4% PFA overnight at 4°C and washed twice with PBS. *In situ hybridization*

assay carried out using the RNAscope® Multiplex Fluorescent v2 reagents (Advanced

Cell Diagnostics, # 323110) as previously described (Nomaru et al., 2021). We used a

C1 probe for Fn1 (#408181).

Embryo Cryosection and immunofluorescence staining:

After dissection, embryos were fixed with 4% PFA overnight at 4°C and washed twice

with PBS. Embryos were then incubated with 10% sucrose in PBS for 4 hours at room

temperature and then with 20% sucrose in PBS overnight at 4°C with agitation. Embryos

were then incubated with 30% sucrose in PBS for 4 hours at room temperature, and then

with a 1:1 mixture of Tissue-Tek O.C.T. Compound (O.C.T, Sakura, #4583) and 30%

sucrose solution for 1 hour at room temperature with agitation. Finally, embryos were

placed into 100% O.C.T for 10-20 minutes without agitation. Embryos were then

positioned in fresh 100% OCT solution in plastic molds and frozen by dipping the mold

into 2-methyl butane chilled on dry ice. 8 μm-thick sections were obtained using a Leica

CM1950 Cryostat. For long term storage, slides were kept at -80°C. For

immunofluorescence, we used an ImmEdge Hydrophobic Pen (VWR, 101098-065) to

draw a hydrophobic barrier around tissue sections. Tissue sections were permeabilized

by washing with PBS containing 0.05% Tween-20 (Sigma, cat # P7949) 3 times for 5

minutes. Then, slides were placed in a humidified chamber, and sections were incubated

with a blocking buffer containing 5% donkey serum in PBS-0.05% Tween for 30 minutes

at room temperature. The blocking buffer was then removed and replaced with primary

antibodies diluted in blocking buffer and incubated with sections overnight at 4°C. After

washing the slides 3 times for 5 minutes with PBS-0.05% Tween; sections were incubated

with secondary antibodies for 1 hour at room temperature. Finally, slices were washed 3

times for 5 minutes each with PBS-0.05%Tween and mounted using a 50% v/v Glycerol,

50% v/v Methanol solution. Sections stained with SiR-Actin (Cytoskeleton, #CY-SC001,

2μM), samples were mounted with Vectashield Antifade Mounting Medium (Vector

laboratories, #H-1000).

TUNEL assay:

TUNEL assay was performed using the In Situ Cell Death Detection Kit from Roche (cat

#11684795910) following manufacturer's instructions.

Imaging, quantifications, and statistical analyses:

Embryos were imaged using confocal microscopy using Nikon A1R microscope with 20x

CFI Apo LWD Lambda S water immersion objective (#MRD77200) or 25x CFI Plan Apo

Lambda S silicone oil objectives (#MRD73250). Cultured cells were imaged using

confocal microscopy and the Plan Fluor 567 40x Oil immersion objective (numerical

aperture 1.3, # MRH01401). 3D reconstructions, surfacing, and quantifications were

performed using IMARIS software (Bitplane, USA). Fluorescence intensity and angles

were measured using Fiji software and plotted using Orient software, version 3.1.1. Cell

ellipticity was measured using Imaris as described (Andrews et al., 2021). Statistical

analyses were performed using Prism 9, version 9.4.1 (GraphPad Software, LLC).

Embryo protein extract:

Embryos were lysed with 100 μL of SDS lysis buffer (0.1 M Tris-HCl, 0.2% SDS, 5mM

EDTA pH 8 containing 1X protease inhibitor cocktail and homogenized by drawing the

lysates 10 times though a 27G syringe. 33.3 μl of 4X NuPAGE™ LDS Sample buffer

containing 10% β-mercaptoethanol were added to each sample, samples were incubated

at 95°C for 5 minutes.

Acknowledgments: We thank Sam Russo for help with mouse colony maintenance and

genotyping. We thank members of the Astrof lab for their careful reading of the manuscript

and for helpful comments and suggestions throughout this work.

Figures Legends

Figure 1. The expression of Fn1 in Mesp1 lineage is required for the OFT

elongation.

Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at

E9.5 and staged according to somite number. **A-D.** Embryos between 18-19 somites. n=5

controls and 4 mutants. **E-H.** Embryos between 20-25 somites, n= 5 control and 7 mutant

embryos. Imaris was used to surface distal OFT (yellow), proximal OFT (turquoise), and

the left ventricle and atria (pink). **C-D, G-H.** The length of the distal and proximal OFT in

each embryo was normalized to the height of the left ventricle to control for embryo size.

I-J. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were

dissected at E9.5 (20-25s) and stained with DAPI and the MF-20 antibody. **K.**

Quantification of angles in controls and mutants as shown in **I-J**. n= 9 controls and 13

mutants. Means (horizontal bars) and standard deviations (error bars) are displayed; p

was calculated using a 2-tailed, unpaired Student's t-test. OFT-outflow tract, MF-20

antibody detects sarcomeric myosin heavy chain. All scale bars : 200 mm

Figure 2. The expression of Fn1 in the mesoderm is important for the regulation of cell shape in the anterior DPW at E8.5. **A.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **E.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E8.5 (9-11s) and stained to detect GFP (green) and nuclei (DAPI, blue). Regions marked by dashed rectangles in **A, B** are magnified in **A1-E1** and **A2-E2**. **B-C, F-G.** Outlines of aDPW and pDPW cells. **D** and **H**. 3D reconstruction of aDPW cells. **I-L.** Cell shape. Small dots mark data points from each cell; large dots mark the average per embryo. Data points from one embryo are marked with the same color. **I-J**. Cell circularity. **I.** 320 cells from 5 control and 367 cells from 5 mutant embryos were analyzed. **J.** 311 cells from 5 control and 326 cells from 5 mutant embryos were analyzed. **K-L.** Ellipticity and volume of aDPW cells; 51 cells from 5 control and 50 cells from mutant embryos were analyzed. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. OFT-outflow tract, aDPW-anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, A-atrium, V-ventricle.

Figure 3. Expression of Fn1 in the mesoderm regulates Golgi apparatus orientation in the anterior DPW. A-A2. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **B-B2.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E8.5 (9-11s) and stained to detect GFP (green), GM130 (white), and nuclei (DAPI, blue). Boxes in **A, B** are magnified in the panels **A1-A2,** and **B1-B2.** Arrows point to Golgi. **A3, A4, B3, B4**. Semicircular histograms represent the distribution of the Golgi orientation angles in DPW cells that were measured as indicated in **C**. N=total number of cells analyzed in 3 controls and 3 mutants. **D, E**. Quantification of Golgi polarization. Golgi was considered polarized if the angle between

the Golgi and the plane of the endoderm fell between 85^0 and 95^0. OFT-outflow tract,

aDPW-anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, A-atrium,

V-ventricle

Figure 4. Expression of Fn1 in the mesoderm regulates cell polarity in the anterior

DPW. A. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **B.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were

dissected at E8.5 (9-11s), and stained to detect GFP (green), podocalyxin (white), and

nuclei (DAPI, blue). Boxes in **A** and **B** were expanded in **A1** and **B1** to show the anterior

DPW and in **A2** and **B2** to show the posterior DPW. Arrows in **A1** point to cells with apical

localization of podocalyxin. Arrows in **B1** point to defective basolateral localization of

podocalyxin in mutants. **A2, B2.** Podocalyxin is localized apically in the posterior DPW,

arrows. **C-D** Quantification of cells with defective podocalyxin localization in the anterior

and posterior DPW. **C.** 148 cells in the aDPW from 3 controls and 224 cells in aDPW from

3 mutants were analyzed. **D.** 149 cells in the pDPW from 3 controls and 182 cells in the

pDPW from 3 mutants were analyzed. Each small dot represents data from one slice, and

each large dot is an average of all slices per embryo. Data from the same embryo is

marked by the same color. **E-G.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **H-J.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$

(Mutant) embryos were dissected at E8.5 (7-8s), and stained for GFP (green), aPKC,

laminin g1, and nuclei (DAPI, blue). Arrows in **F** point to apical localization of aPKC in

control DPW cells. **I.** Arrows point to defective basolateral localization of aPKC in mutants.

Arrowheads in **G** point to basal localization of laminin g1 in controls and in **J** to

disorganized localization of laminin g1 in one of three mutants. **K.** Quantification of cells

in which aPKC was re-distributed away from the apical surface. DPW cells from 5 different

slices were evaluated in 3 control and 3 mutant embryos. Each small dot represents data

from one slice, and each large dot is an average of all slices per embryo. Data from the

same embryo is marked by the same color. Means (horizontal bars) and standard

deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired

Student's t-test.   OFT-outflow tract, aDPW-anterior dorsal pericardial wall, pDPW-

posterior dorsal pericardial wall, A-atrium, V-ventricle.

Figure 5. Mesodermal Fn1 regulates the expression of N-Cadherin at cell-cell

junctions in the DPW. A-B. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **C-D.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$

(Mutant) embryos were dissected at E9.5 (21-26s) and stained to detect GFP (green), N-

cadherin (white) and nuclei (DAPI, blue). Junctional N-Cadherin is enriched in the anterior

DPW (boxed region in **B**). Boxes in **B**-**D** are expanded in **B1** and **D1.** Arrows point to

DPW. **E.** Western blot analysis of total cell lysates and cell surface proteins. Positions of

molecular weight markers are indicated on the left. **F-G**. Densitometrical quantification of

immunoblot signals. N = 4 for each genotype. Each experiment is represented by a dot

of a different color. Means (horizontal bars) and standard deviations (error bars) are

displayed; 2-tailed, unpaired Student's t-tests were used to determine p values. **H-K.**

Immunostaining to detect GFP (green), N-cadherin (white), and nuclei (DAPI, blue) in

Fn1$^{flox/-}$, Fn1$^{\Delta/-}$ (Fn1-null), Fn1$^{flox/+}$ and Fn1$^{\Delta/+}$ MEFs. Boxes in **I-K** and in **M-O** are

expanded in **I1-K1 and M1-O1.** Arrows point to N-cadherin at cell borders in control MEFs

(**I1, M1, and O1**) and to the absence of N-cadherin at cell borders in Fn1$^{\Delta/-}$ MEFs (**K1**).

DPW- dorsal pericardial wall, OFT-outflow tract, A-atrium.

Figure 6. Mesodermal Fn1 regulates the length and organization of F-actin in the anterior DPW. A-B1. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **C-D1.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (21-26s), and transverse cryosections (a plain of sections shown by the dashed line in **E**) from the anterior DPW were stained to detect GFP (green), F-actin (white), and nuclei (DAPI, blue). Yellow arrows point to F-actin in the DPW in **B1** and **D1**. Yellow arrowheads point to apically enriched F-actin in **B1** and re-distributed F-actin in **D1**. **E.** Schematic representation of different regions in the DPW. **F-H.** Traces of F-actin in control and mutants. **G-I.** Angles made by F-actin filaments relative to cells' apical surface were measured. **J.** Quantification of actin polarization. An F-actin filament was considered polarized if it was between 85 and 95 degrees relative to the apical surface of the DPW. **K** Quantification of the F-actin length in the anterior DPW; 235 cells from 3 controls and 417 cells from 5 mutants were examined. Small dots mark the F-actin length of individual cells, and large dots mark the average per slice. Dots from the same embryo are displayed in the same color. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. DPW-dorsal pericardial wall, A-atrium, NT-Neural tube, AVC-atrioventricular canal, OFT-outflow tract, V-ventricle.

Figure 7 Mesodermal Fn1 regulates the expression of pMLC in the anterior DPW. A-B2. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and C-D. Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (21-26s), and sagittal cryosections were stained to detect GFP (green), F-actin (white) and nuclei (DAPI, blue). Boxes in **A-B** were expanded in **B1** and **B2** to show the anterior and posterior DPW, respectively. Boxes in **C-D** were expanded, in **D1**

and **D2**, to show the anterior and posterior DPW, respectively. Arrows point to the apical

enrichment of F-actin in the anterior and posterior of controls and the lack of such

enrichment in the anterior DPW of the mutants (**D1**). 6 controls and 3 mutants were

analyzed. **E-H2**. $Fn1^{flox/+}$; $Mesp1^{Cre/+}$ (Control) and **C-D.** $Fn1^{flox/-}$; $Mesp1^{Cre/+}$ (Mutant)

embryos were dissected at E9.5 (21-26s) and sagittal cryosections were stained to detect

GFP (green), pMLC (white) and nuclei (DAPI, blue). Boxes in **E-F** are expanded in **F1**

and **F2** to show the anterior and posterior DPW, respectively. Boxes in **G-H** are expanded

in **H1** and **H2** to show anterior and posterior DPW, respectively. Arrows in **F1** indicate

cells with apical localization of pMLC. Arrows in **H1** show a defective basolateral

localization of pMLC in mutants. **F2, H2**. pMLC is localized apically in the posterior DPW.

8 controls and 3 mutants were analyzed.

Figure 8. Mesodermal Fn1 regulates nuclear localization of YAP. A. Ventral view of

E9.5 embryo. YAP localization was assayed in the DPW between the left and the right

limits (marked by the vertical lines).

B-B2. $Fn1^{flox/+}$; $Mesp1^{Cre/+}$ (Control) and **C-C2.** $Fn1^{flox/-}$; $Mesp1^{Cre/+}$ (Mutant) embryos were

dissected at E9.5 (23-26s) and stained to detect nuclei (DAPI, magenta) and YAP (cyan).

Regions outlined by dashed rectangles are expanded in **B1, C1** (anterior DPW) and **B2,**

C2 (posterior DPW). **D-E**. Quantification of the intensity ratio between nuclear and

cytosolic YAP in the aDPW and pDPW. 7-8 optical slices from 4 controls and 7-8 optical

slices from 4 mutant embryos were analyzed. Each small dot represents one slice, and

each large dot is an average of all slices in one embryo. Data from the same embryo is

marked by the same color. **F.** Summary: YAP is enriched in the nuclei in the presence of

both the mesodermal (red lines) and endodermal (blue lines) Fn1. In the absence of

mesodermal Fn1, nuclear translocation of YAP are decreased. aDPW- anterior dorsal

pericardial wall, pDPW-posterior dorsal pericardial wall, OFT-outflow tract.

Figure 9. Mesodermal Fn1 is required for DPW cell proliferation. A-C. Fn1$^{flox/+}$;

Mesp1$^{Cre/+}$ (Control) and **D-F** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5

(20-25s) and stained for PHH3 (white), and nuclei (DAPI, blue) E9.5 (20-25s). **A,B, D, E.**

DPW (magenta) and PHH3^{+} cells in the DPW (white) were segmented using Imaris. **C,F.**

Proportion of PHH3+ cells in the aDPW (magenta) and pDPW (turquoise) were quantified

in **G-H**. Each dot marks one embryos. n=4 control and 6 mutant embryos. Means

(horizontal bars) and standard deviations (error bars) are displayed, 2-tailed, unpaired

Student's t test was used to determine p values. aDPW-anterior dorsal pericardial wall,

pDPW- posterior dorsal pericardial wall, PHH3- Phosphorylated Histone H3.

Figure 10. Expression patterns of *Fn1 mRNA* and protein in controls and mutants.

A-J. Expression of *Fn1 mRNA*. Embryos were dissected at E9.5 (23-26s) and labeled

with DAPI (magenta) and anti-sense Fn1 probe (turquoise). **A-B2** Fn1$^{flox/+}$; Mesp1$^{+/+}$

(designated as WT), **C.** *Fn1 mRNA* intensity ratio between aDPW and pDPW, **D-F2.**

Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) **G.** *Fn1 mRNA* intensity in the anterior (**H**) and posterior (**J**)

DPW was normalized to that in the endoderm. 6 different slices from 3 WT, 4 control and

9 mutant embryos were analyzed. **H-I2.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos. Regions

outlined by the dotted rectangles in **B, E,** and **I** are expanded in panels to the right and

color-coded according to fluorescence intensity (scale is on the right). White arrows point

to DPW. Yellow arrowheads point to the endoderm. **K.** Ventral view of the DPW, Fn1

expression is color-coded according to fluorescence intensity. **L.** Ratio of Fn1 protein

intensity shows similar enrichment of Fn1 in the anterior DPW of controls and mutants.

M, M1. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **N, N1.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos

were dissected at E9.5 (22-26s) and stained to detect Fn1 protein (white) and nuclei

(DAPI, blue). Fn1 protein expression in the DPW (**M, N**) is color-coded according to

fluorescence intensity in **M1, N1**. **O-P.** Fn1 protein levels underlying the anterior DPW are

unchanged in the mutants. Fn1 intensity in the aDPW and pDWP regions outlined by the

red dash lines in **M1-N1** was normalized by DAPI intensity; 7 optical slices from 8 control

and 10 mutant embryos were evaluated. Each small dot represents one slice, and each

large dot is an average of all slices per embryo. Data from the same embryo is marked

by the same color. Means (horizontal bars) and standard deviations (error bars) are

displayed, 2-tailed, unpaired Student's t-test was used to determine p values. aDPW-

anterior dorsal pericardial wall, pDPW- posterior dorsal pericardial wall, OFT-outflow tract,

A-atrium, LV-left ventricle.

Figure 11. The central role of the mesodermal Fn1 in regulating the elongation of

the cardiac outflow tract. Fn1 synthesized by the mesoderm mediates the elongation

of the cardiac outflow tract by regulating cell shape, polarity, proliferation, and

mechanotransduction in the anterior dorsal pericardial wall (aDPW).

Figure S1. Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ mutant embryos do not survive past E10.75 of

gestation.

A. Schematic representation of a sagittal section. **B-B1** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and

C-D1 Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos dissected at E9.5. **E-E1.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$

(Control) and **F-F1** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos dissected at E10.75. **G.**

Genotype distributions of embryos isolated at different stages of gestation. dOFT-distal

outflow tract, pOFT- proximal outflow tract, aDPW-anterior dorsal pericardial wall, pDPW-

posterior dorsal pericardial wall, A-atrium.

Figure S2. The initial formation of the cardiac outflow tract is not affected by the

deletion of Fn1 in the mesoderm at E8.5. A-B. Illustration of our method for measuring

OFT dimensions. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) embryo at E9.5. Imaris was used to

surface the Distal OFT (yellow), Proximal OFT (turquoise) and the left ventricle and atria

(pink). White lines show how the length of OFT segments and LV were measured. **C-C1**

Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **D-D1** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were

dissected at E8.5. Imaris was used to surface the OFT (yellow) and the left ventricle and

atria (pink). **E.** The length of the OFT was normalized to the height of the left ventricle to

control for embryo size. 11 control and 11 mutant embryos were analyzed. Each dot

corresponds to one embryo. Means (horizontal bars) and standard deviations (error bars)

are displayed p was calculated using 2-tailed, unpaired Student's t-test. OFT- outflow

tract.

Figure S3. Mesodermal Fn1 regulates DPW cell shape at E9.5. **A.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$

(Control) and **E.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (21-25s)

and stained to detect GFP (green) and nuclei (DAPI, blue). Regions marked by dashed

rectangles in **A, B** are magnified in **A1-E1** and **A2-E2**. **B-C, F-G.** Outlines of aDPW and

pDPW cells. **D** and **H.** 3D reconstruction of aDPW cells. **I-L.** Cell shape analyses. Small

dots mark a data point from each cell; large dots mark the average per embryo. Data

points from one embryo are marked with the same color. **I-J.** Cell circularity. **I.** 396 cells

from 5 control and 436 cells from 5 mutant embryos were analyzed. **J.** 390 cells from 5

control and 419 cells from 5 mutant embryos were analyzed. **K-L.** Ellipticity and volume

of aDPW cells. n=10 cells from 5 control and 10 cells from 5 mutant embryos were

analyzed. Means (horizontal bars) and standard deviations (error bars) are displayed; p

was calculated using a 2-tailed, unpaired Student's t-test. OFT-outflow tract, aDPW-

anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, A-atrium.

Figure S4. Mesodermal depletion of Fn1 does not alter the number of basal

protrusions in the aDPW. **A.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **B.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$

(Mutant) embryos were dissected at E9.5 (21-26s), and transverse cryosections from the

anterior DPW were stained to detect GFP (black). Red rectangles in **A, B** are expanded

in **A1** and **B1**. Arrows point to protrusion extending from the basal membranes of aDPW

cells. **C.** Quantification of the number of protrusions normalized for the length of the DPW.

n= 3 control and 4 mutant embryos. 2 – 7 sections per embryo were analyzed. Small dots

mark the number of protrusions in each section. Large dots mark the mean in each

embryo. Data from the same embryo is marked with the same color. P was calculated

using a 2-tailed, unpaired Student's t-test. aDPW- anterior dorsal pericardial wall. P was

calculated using a 2-tailed, unpaired Student's t-test.

Figure S5. mesodermal Fn1 do not regulates cell polarity in the DPW at E9.5 . A. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **B.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (19-24s), and stained to detect GFP (green), podocalyxin (white), and nuclei (DAPI, blue). Boxes in **B** and **D** were expanded in **B1** and **D1**. Arrows show apical localization of podocalyxin in control and mutant embryos. **E** Quantification of cells with defective podocalyxin localization in the anterior and posterior DPW. P was calculated using a 2-tailed, unpaired Student's t-test; p$\geqslant$0.05 indicates a non-significant difference between the two groups. DPW- dorsal pericardial wall, A- atrium, NT- Neural tube, OFT- outflow tract.

Figure S6. Generation of MEF lines . A-B. Schematic representation of resultant alleles upon Cre recombination in Fn1$^{flox/-}$;ROSA$^{mTmG/+}$ and Fn1$^{flox/+}$; ROSA$^{mTmG/+}$ MEFs. **A.** To delete Fn1, we infected Fn1$^{flox/-}$;ROSA$^{mTmG/+}$ MEFs with adenoviruses encoding Cre recombinase, generating Fn1$^{\Delta/-}$ (Fn1-null) and Fn1$^{flox/-}$ (control cells). **B.** Fn1$^{flox/+}$; ROSA$^{mTmG/+}$ MEFs were treated with adenoviruses encoding Cre recombinase to generate control Fn1$^{\Delta/+}$ MEFs. **C.** After infection, cells were expanded and sorted. Membrane-tagged GFP-expressing cells are in green, and membrane-tagged td-Tomato-expressing cells are in red. **D.** Western blot using anti-Fn1 antibody shows that we successfully ablated Fn1 in Fn1$^{\Delta/-}$ (Fn1-null) MEFs. β-actin was used as a loading control. **E.** Densitometry quantification of Fn1 levels relative to β-actin.

Figure S7. Monolayer scratch assay. A-D. Cells imaged immediately after the scratch wound (time=0 hrs). **A1-D1.** Cells were imaged 14 hours after the scratch (time=14 hrs).

White lines mark the area remaining cell-free at t=14 hours. **E**. To quantify % wound

closure, cell-free area at t=14 ($A_{t=14}$) was subtracted from cell-free area at t=0 ($A_{t=0}$), and

the % wound closure was calculated as $100^*(A_{t=0} - A_{t=14})/ A_{t=0}$); n=6 independent

experiments. Means (horizontal bars) and standard deviations (error bars) are displayed;

p was calculated using a 2-tailed, unpaired Student's t-test.

Figure S8. Deletion of Fn1 in Mesp1 lineage causes ectopic DPW differentiation. **A-**

C. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **D-F** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were

dissected at E9.5 (21-26s), and stained to detect GFP (green), sarcomeric myosin heavy

chain (MF-20, white), and nuclei (DAPI, blue). Arrow in **E** points to cells in the aDPW

ectopically expressing sarcomeric myosin heavy chain. **C, F.** Ventral views of the DPW.

3D reconstruction of the DPW (pink) using Imaris. MF-20 staining in the DPW is marked

by the white surface. **G**. Schematic representation of the method used to quantify MF-20

staining. **H.** The length of the MF20 staining relative to the DPW length in each analyzed

section is plotted.Eight optical sections from 5 control and eight optical sections 8 mutant

embryos were analyzed. Small dots mark data from each slice; Large dots mark means.

Data from the same embryo is displayed with the same color. A 2-tailed, unpaired.

Student's t-test was used to determine p values. DPW- dorsal pericardial wall, OFT-

outflow tract, A- atrium. MF-20- sarcomeric myosin heavy chain.

Figure S9. The organization of F-actin cytoskeleton in the posterior DPW does not

depend on the mesodermal Fn1. A. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and Fn1$^{flox/-}$;

Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (21-26s), and transverse

cryosections (dashed line in **A**) from the middle of the DPW were stained to detect GFP

(green), -actin (white), and nuclei (DAPI, blue). Yellow arrows point to F-actin

cytoskeleton. **F**-**G.** Quantification of F-actin angles in the middle DPW was measured

relative to the apical cell side. 270 cells from 4 controls and 123 cells from 3 mutants were

analyzed. **H.** Percent of actin filaments oriented between 85^0 and 95^0 to the base of the

DPW**.** 260 cells from 4 controls and 121 cells from 3 mutants were analyzed. **I.**

Quantification of F-actin length in middle DPW cells. **J – M**. $Fn1^{flox/+}$; $Mesp1^{Cre/+}$ (Control)

and $Fn1^{flox/-}$; $Mesp1^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (21-26s), and

transverse cryosections (dashed line in **A**) from the posterior of the DPW were stained as

in **B-E**. **N-O.** Quantification of F-actin angles in the posterior DPW was measured relative

to the apical cell side. 200 cells from 3 controls and 210 cells from 2 mutants were

analyzed as in **F-G**. **P.** Actin organization in cells of the posterior DPW was measured as

in **H**. **Q**. Quantification of F-actin length in the posterior DPW cells. Small dots mark data

from each cell. Large dots are means from each optical slice. Data from the same embryo

is displayed with the same color. Means (horizontal bars) and standard deviations (error

bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. $p < 0.05$

is considered significant. DPW- dorsal pericardial wall, A- atrium, NT- Neural tube, OFT-

outflow tract, V- ventricle.

Figure S10. Mesodermal Fn1 regulates DPW cell proliferation but not cell survival.

A-C. $Fn1^{flox/+}$; $Mesp1^{Cre/+}$ (Control) and $Fn1^{flox/-}$; $Mesp1^{Cre/+}$ (Mutant) embryos were

dissected and stained for PHH3 (white), and nuclei (DAPI, blue). **A-B.** E8.5 (11-12s)

embryos. DPW (magenta) and $PHH3^+$ cells in the DPW (white) were segmented using

Imaris. **C**. Quantification of PHH3 $^+$ cells in the DPW of E8.5 embryos, n=3 control and 4

mutant embryos were analyzed. **D, E, F, G.** Endoderm (orange) and PHH3$^+$ cells in the

endoderm (white) were segmented using Imaris. **D-E.** E8.5 embryos (11-12s) and **F-G.**

E9.5 embryos (20-25). **H**. Quantification of PHH3$^+$ cells in the endoderm of E8.5 embryos.

n=3 for each genotype. **I**. Quantification of PHH3$^+$ cells in the endoderm of E9.5 embryos

n=4 for each genotype. **J-J1.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **K-K1.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$

(Mutant) embryos were dissected at E9.5 (21-26s), sectioned in the transverse

orientation, stained to detect GFP (green), nuclei (DAPI, blue), and apoptosis using the

TUNEL assay (magenta). Boxes in **J** and **K** are expanded in **J1** and **K1**. Arrows point to

apoptotic cells (TUNEL$^+$) in the DPW. **M.** Quantification of TUNEL$^+$ cells in the DPW. 2-

7 slices per embryo were analyzed. Small dots indicate the number of TUNEL$^+$ cells in

one slice. Large dots mark the mean in each embryo. Data from the same embryo is

marked with the same color. Means (horizontal bars) and standard deviations (error bars)

are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. DPW- dorsal

pericardial wall, PHH3- Phosphohistone H3, TUNEL- Terminal deoxynucleotidyl

transferase (TdT) dUTP Nick-End Labeling.

Figure S11. Mesodermal Fn1 regulates OFT myocardial cell proliferation at E9.5. A,

D. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **B, E.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were

dissected and stained to detect PHH3 (white) and nuclei (DAPI, blue) at E8.5 and E9.5.

A-B. E8.5 embryos (11-12s). Myocardial cells in the OFT were segmented in yellow and

the PHH3$^+$ cells in the OFT were segmented in white using Imaris. **D-E.** E9.5 embryos

(20-25s). Myocardial cells in the distal OFT were segmented in yellow, and proximal OFT

– in turquoise, and the PHH3+ cells in each segment were marked in white using Imaris.

C, F, G. The number of PHH3$^+$ cells was normalized for the total number of nuclei in the OFT. **C**. Quantification of PHH3$^+$ cells in the OFT of E8.5 embryos. n= 3 control and 4 mutant embryos. **F-G**. Quantification of PHH3$^+$ cells in the distal and proximal OFT of E9.5 embryos. n=4 control and 6 mutant embryos. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. OFT- outflow tract, PHH3 – phosphorylated Histone H3.

Figure S12. Deletion of mesodermal Fn1 does not alter OFT cell survival or NCC cell migration. A-C. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control, n=3) and **D-F** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant, n=3) embryos were dissected at E9.5 (21-26s), and transverse cryosections were used to perform TUNEL assay (magenta), detect Mesp1 lineage by the expression of GFP (green), an NCC marker TFAP2α (white), and nuclei (DAPI, blue). Boxes in **B** and **E** are expanded in **B1, B2** and **E1, E2,** where white arrows indicate apoptotic cells (magenta) in the OFT. Boxes in **C** and **F** are expanded in **C1** and **F1** where arrows point to NCCs (TFAP2α + cells) in the PA. **G.** Quantification of the number of TUNEL$^+$ cells in the OFT per section. 2 - 7 sections per embryo were analyzed. Small dots in the plots mark the number of TUNEL$^+$ cells in the OFT in each section. Large dots mark the mean in each embryo. **H.** Quantification of the number TFAP2α+ cells in the second pharyngeal arch. 2 – 7 sections per embryo were analyzed. Small dots mark the number of TFAP2α+ cells in each section. Large dots mark the mean in each embryo. P was calculated using a 2-tailed, unpaired Student's t-test. OFT- outflow tract, NCC cells- neural cardiac crest cells, TUNEL- Terminal deoxynucleotidyl transferase (TdT) dUTP Nick-End Labeling, TFAP2α -Transcription Factor AP-2 Alpha.

Figure S13. Efficient downregulation of Fn1 protein in the mesoderm of Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos. A-B2. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **C-D2.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E8.5 (8-12s) and stained to detect Fn1 protein (white) and nuclei (DAPI, blue). **E.** Ratio of Fn1 protein intensity shows similar levels Fn1 in the anterior and posterior DPW embryos. **F-H.** Fn1 protein levels decrease in the DPW and left ventricle in mutants. Fn1 intensity in the aDPW, pDWP and left ventricle regions outlined by the red dash lines in **B1-2** and **D1-D2** was normalized by DAPI intensity; 5 optical slices from 5 control and 5 mutant embryos were evaluate. **I-M.** E9.5 **I-J.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **K-L.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (22-26s) and stained to detect Fn1 protein (white) and nuclei (DAPI, blue). Regions outlined by rectangles are expanded in **J1** and **L1**. **M.** Fn1 protein expression in the left ventricle. 7 optical slices from 8 control and 10 mutant embryos were evaluate. Each small dot represents one slice, and each large dot is an average of all slices per embryo. Data from the same embryo is marked by the same color. Means (horizontal bars) and standard deviations (error bars) are displayed, 2-tailed, unpaired Student's t-test was used to determine p values. aDPW-anterior dorsal pericardial wall, pDPW- posterior dorsal pericardial wall, OFT-outflow tract, A-atrium, LV-left ventricle.

Figure S14. Fn1 synthesized by the endoderm is not required for the epithelial organization or N-Cadherin expression in the DPW. A-C. Fn1$^{flox/+}$ Sox17$^{Cre/+}$ (Control) and **D-F.** Fn1$^{flox/-}$ Sox17$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (20-26s). In **A** and **D,** embryos were labeled to detect *Fn1 mRNA* (turquoise). **A1** and **D1** are magnifications

of the DWP. White and yellow arrows indicate the mesoderm and endoderm, respectively.

In **B** and **E** embryos were stained to detect Fn1 protein (green), and nuclei (DAPI blue).

B1 and **E1** are expanded from **B** and **E**. Arrows point to Fn1 protein staining in the DPW.

In **C** and **F** embryos were stained to detect N-cadherin (white). **C1** and **F1** are expanded

from **C** and **F.** Arrows point to N-cadherin expression at SHF cell-cell junctions in the

DPW. OFT- outflow tract, DPW- dorsal pericardial wall, A- atrium.

Literature Cited

Alfano, D., A. Altomonte, C. Cortes, M. Bilio, R.G. Kelly, and A. Baldini. 2019. Tbx1
regulates extracellular matrix-cell interactions in the second heart field. *Hum Mol Genet*.
28:2295-2308.

Andrews, T.G.R., W. Ponisch, E.K. Paluch, B.J. Steventon, and E. Benito-Gutierrez.
2021. Single-cell morphometrics reveals ancestral principles of notochord development.
Development. 148.

Arriagada, C., V.A. Cavieres, C. Luchsinger, A.E. Gonzalez, V.C. Munoz, J. Cancino,
P.V. Burgos, and G.A. Mardones. 2020. GOLPH3 Regulates EGFR in T98G
Glioblastoma Cells by Modulating Its Glycosylation and Ubiquitylation. *Int J Mol Sci*. 21.

Bajolle, F., S. Zaffran, R.G. Kelly, J. Hadchouel, D. Bonnet, N.A. Brown, and M.E.
Buckingham. 2006. Rotation of the myocardial wall of the outflow tract is implicated in
the normal positioning of the great arteries. *Circ Res*. 98:421-428.

Bajolle, F., S. Zaffran, S.M. Meilhac, M. Dandonneau, T. Chang, R.G. Kelly, and M.E.
Buckingham. 2008. Myocardium at the base of the aorta and pulmonary trunk is
prefigured in the outflow tract of the heart and in subdomains of the second heart field.
Dev Biol. 313:25-34.

Bohmer, R.M., E. Scharf, and R.K. Assoian. 1996. Cytoskeletal integrity is required
throughout the mitogen stimulation phase of the cell cycle and mediates the anchorage-
dependent expression of cyclin D1. *Mol Biol Cell*. 7:101-111.

Borreguero-Munoz, N., G.C. Fletcher, M. Aguilar-Aragon, A. Elbediwy, Z.I. Vincent-
Mistiaen, and B.J. Thompson. 2019. The Hippo pathway integrates PI3K-Akt signals
with mechanical and polarity cues to control tissue growth. *PLoS Biol*. 17:e3000509.

Chen, D., X. Wang, D. Liang, J. Gordon, A. Mittal, N. Manley, K. Degenhardt, and S.
Astrof. 2015. Fibronectin signals through integrin alpha5beta1 to regulate
cardiovascular development in a cell type-specific manner. *Dev Biol*. 407:195-210.

Cortes, C., A. Francou, C. De Bono, and R.G. Kelly. 2018. Epithelial Properties of the
Second Heart Field. *Circ Res*. 122:142-154.

Cseh, B., S. Fernandez-Sauze, D. Grall, S. Schaub, E. Doma, and E. Van Obberghen-
Schilling. 2010. Autocrine fibronectin directs matrix assembly and crosstalk between

cell-matrix and cell-cell adhesion in vascular endothelial cells. *J Cell Sci*. 123:3989-
3999.

Dupont, S. 2016. Role of YAP/TAZ in cell-matrix adhesion-mediated signalling and
mechanotransduction. *Exp Cell Res*. 343:42-53.

Dzamba, B.J., K.R. Jakab, M. Marsden, M.A. Schwartz, and D.W. DeSimone. 2009.
Cadherin adhesion, tissue tension, and noncanonical Wnt signaling regulate fibronectin
matrix organization. *Dev Cell*. 16:421-432.

Engert, S., W.P. Liao, I. Burtscher, and H. Lickert. 2009. Sox17-2A-iCre: a knock-in
mouse line expressing Cre recombinase in endoderm and vascular endothelial cells.
Genesis. 47:603-610.

Fan, R., N.G. Kim, and B.M. Gumbiner. 2013. Regulation of Hippo pathway by
mitogenic growth factors via phosphoinositide 3-kinase and phosphoinositide-
dependent kinase-1. *Proc Natl Acad Sci U S A*. 110:2569-2574.

Francou, A., C. De Bono, and R.G. Kelly. 2017. Epithelial tension in the second heart
field promotes mouse heart tube elongation. *Nat Commun*. 8:14770.

Francou, A., E. Saint-Michel, K. Mesbah, and R.G. Kelly. 2014. TBX1 regulates
epithelial polarity and dynamic basal filopodia in the second heart field. *Development*.
141:4320-4331.

Friedl, P., and R. Mayor. 2017. Tuning Collective Cell Migration by Cell-Cell Junction
Regulation. *Cold Spring Harb Perspect Biol*. 9.

Garavito-Aguilar, Z.V., H.E. Riley, and D. Yelon. 2010. Hand2 ensures an appropriate
environment for cardiac fusion by limiting Fibronectin function. *Development*. 137:3215-
3220.

George, E.L., H.S. Baldwin, and R.O. Hynes. 1997. Fibronectins are essential for heart
and blood vessel morphogenesis but are dispensable for initial specification of
precursor cells. *Blood*. 90:3073-3081.

George, E.L., E.N. Georges-Labouesse, R.S. Patel-King, H. Rayburn, and R.O. Hynes.
1993. Defects in mesoderm, neural tube and vascular development in mouse embryos
lacking fibronectin. *Development*. 119:1079-1091.

Hirsh, G.D., B.J. Dzamba, P.R. Sonavane, D.R. Shook, C.M. Allen, and D.W.
DeSimone. 2018. Multiple functions for the catenin family member plakoglobin in
cadherin-dependent adhesion, fibronectin matrix assembly, and *Xenopus* gastrulation
movements. *bioRxiv*:318774.

Kelly, R.G. 2023. The heart field transcriptional landscape at single-cell resolution. *Dev*
Cell. 58:257-266.

Kim, N.G., and B.M. Gumbiner. 2015. Adhesion to fibronectin regulates Hippo signaling
via the FAK-Src-PI3K pathway. *J Cell Biol*. 210:503-515.

Li, D., A. Angermeier, and J. Wang. 2019. Planar cell polarity signaling regulates
polarized second heart field morphogenesis to promote both arterial and venous pole
septation. *Development*. 146.

Li, D., T. Sinha, R. Ajima, H.S. Seo, T.P. Yamaguchi, and J. Wang. 2016. Spatial
regulation of cell cohesion by Wnt5a during second heart field progenitor deployment.
Dev Biol. 412:18-31.

Li, D., and J. Wang. 2018. Planar Cell Polarity Signaling in Mammalian Cardiac
Morphogenesis. *Pediatr Cardiol*. 39:1052-1062.

Luciano, M., M. Versaevel, E. Vercruysse, A. Procès, Y. Kalukula, A. Remson, A.
Deridoux, and S. Gabrielea. 2022. Appreciating the role of cell shape changes in the
mechanobiology of epithelial tissues. *Biophysics Reviews*. 3:011305.

Marsden, M., and D.W. DeSimone. 2001. Regulation of cell polarity, radial intercalation
and epiboly in Xenopus: novel roles for integrin and fibronectin. *Development*.
128:3635-3647.

Marsden, M., and D.W. DeSimone. 2003. Integrin-ECM interactions regulate cadherin-
dependent cell adhesion and are required for convergent extension in Xenopus. *Curr*
Biol. 13:1182-1191.

Meilhac, S.M., and M.E. Buckingham. 2018. The deployment of cell lineages that form
the mammalian heart. *Nat Rev Cardiol*. 15:705-724.

Mittal, A., M. Pulina, S.Y. Hou, and S. Astrof. 2010. Fibronectin and integrin alpha 5 play
essential roles in the development of the cardiac neural crest. *Mech Dev*. 127:472-484.

Mittal, A., M. Pulina, S.Y. Hou, and S. Astrof. 2013. Fibronectin and integrin alpha 5 play
requisite roles in cardiac morphogenesis. *Dev Biol*. 381:73-82.

Muzumdar, M.D., B. Tasic, K. Miyamichi, L. Li, and L. Luo. 2007. A global double-
fluorescent Cre reporter mouse. *Genesis*. 45:593-605.

Nelson, W.J. 2003. Adaptation of core mechanisms to generate cell polarity. *Nature*.
422:766-774.

Nomaru, H., Y. Liu, C. De Bono, D. Righelli, A. Cirino, W. Wang, H. Song, S.E. Racedo,
A.G. Dantas, L. Zhang, C.L. Cai, C. Angelini, L. Christiaen, R.G. Kelly, A. Baldini, D.
Zheng, and B.E. Morrow. 2021. Single cell multi-omic analysis identifies a Tbx1-
dependent multilineage primed population in murine cardiopharyngeal mesoderm. *Nat
Commun*. 12:6645.

Perez Gonzalez, N., J. Tao, N.D. Rochman, D. Vig, E. Chiu, D. Wirtz, and S.X. Sun.
2018. Cell tension and mechanical regulation of cell volume. *Mol Biol Cell*. 29:0.

Pulina, M.V., S.Y. Hou, A. Mittal, D. Julich, C.A. Whittaker, S.A. Holley, R.O. Hynes, and
S. Astrof. 2011. Essential roles of fibronectin in the development of the left-right
embryonic body plan. *Dev Biol*. 354:208-220.

Ramirez, A., and S. Astrof. 2020. Visualization and Analysis of Pharyngeal Arch Arteries
using Whole-mount Immunohistochemistry and 3D Reconstruction. *J Vis Exp*.

Rammah, M., M. Theveniau-Ruissy, R. Sturny, F. Rochais, and R.G. Kelly. 2022.
PPARgamma and NOTCH Regulate Regional Identity in the Murine Cardiac Outflow
Tract. *Circ Res*. 131:842-858.

Ramsbottom, S.A., V. Sharma, H.J. Rhee, L. Eley, H.M. Phillips, H.F. Rigby, C. Dean,
B. Chaudhry, and D.J. Henderson. 2014. Vangl2-regulated polarisation of second heart
field-derived cells is required for outflow tract lengthening during cardiac development.
PLoS Genet. 10:e1004871.

Roovers, K., and R.K. Assoian. 2003. Effects of rho kinase and actin stress fibers on
sustained extracellular signal-regulated kinase activity and activation of G(1) phase
cyclin-dependent kinases. *Mol Cell Biol*. 23:4283-4294.

Rozario, T., B. Dzamba, G.F. Weber, L.A. Davidson, and D.W. DeSimone. 2009. The
physical state of fibronectin matrix differentially regulates morphogenetic movements in
vivo. *Dev Biol*. 327:386-398.

Saga, Y., N. Hata, S. Kobayashi, T. Magnuson, M.F. Seldin, and M.M. Taketo. 1996.
MesP1: a novel basic helix-loop-helix protein expressed in the nascent mesodermal
cells during mouse gastrulation. *Development*. 122:2769-2778.

Saga, Y., S. Miyagawa-Tomita, A. Takagi, S. Kitajima, J. Miyazaki, and T. Inoue. 1999.
MesP1 is expressed in the heart precursor cells and required for the formation of a
single heart tube. *Development*. 126:3437-3447.

Sakai, T., K.J. Johnson, M. Murozono, K. Sakai, M.A. Magnuson, T. Wieloch, T.
Cronberg, A. Isshiki, H.P. Erickson, and R. Fassler. 2001. Plasma fibronectin supports
neuronal survival and reduces brain injury following transient focal cerebral ischemia but
is not essential for skin-wound healing and hemostasis. *Nat Med*. 7:324-330.

Sawyer, J.M., J.R. Harrell, G. Shemer, J. Sullivan-Brown, M. Roh-Johnson, and B.
Goldstein. 2010. Apical constriction: a cell shape change that can drive morphogenesis.
Dev Biol. 341:5-19.

Singh, P., C. Carraher, and J.E. Schwarzbauer. 2010. Assembly of fibronectin
extracellular matrix. *Annu Rev Cell Dev Biol*. 26:397-419.

Sinha, T., D. Li, M. Theveniau-Ruissy, M.R. Hutson, R.G. Kelly, and J. Wang. 2015.
Loss of Wnt5a disrupts second heart field cell deployment and may contribute to OFT
malformations in DiGeorge syndrome. *Hum Mol Genet*. 24:1704-1716.

Sinha, T., B. Wang, S. Evans, A. Wynshaw-Boris, and J. Wang. 2012. Disheveled
mediated planar cell polarity signaling is required in the second heart field lineage for
outflow tract morphogenesis. *Dev Biol*. 370:135-144.

Soh, B.S., K. Buac, H. Xu, E. Li, S.Y. Ng, H. Wu, J. Chmielowiec, X. Jiang, L. Bu, R.A.
Li, C. Cowan, and K.R. Chien. 2014. N-cadherin prevents the premature differentiation
of anterior heart field progenitors in the pharyngeal mesodermal microenvironment. *Cell
Res*. 24:1420-1432.

Somlyo, A.P., and A.V. Somlyo. 1994. Signal transduction and regulation in smooth
muscle. *Nature*. 372:231-236.

Stefanovic, S., B. Laforest, J.P. Desvignes, F. Lescroart, L. Argiro, C. Maurel-Zaffran, D.
Salgado, E. Plaindoux, C. De Bono, K. Pazur, M. Theveniau-Ruissy, C. Beroud, M.
Puceat, A. Gavalas, R.G. Kelly, and S. Zaffran. 2020. Hox-dependent coordination of
mouse cardiac progenitor cell patterning and differentiation. *Elife*. 9.

Theveniau-Ruissy, M., M. Dandonneau, K. Mesbah, O. Ghez, M.G. Mattei, L. Miquerol,
and R.G. Kelly. 2008. The del22q11.2 candidate gene Tbx1 controls regional outflow
tract identity and coronary artery patterning. *Circ Res*. 103:142-148.

Trinh, L.A., and D.Y. Stainier. 2004. Fibronectin regulates epithelial organization during
myocardial migration in zebrafish. *Dev Cell*. 6:371-382.

Trinh, L.A., D. Yelon, and D.Y. Stainier. 2005. Hand2 regulates epithelial formation
during myocardial diferentiation. *Curr Biol*. 15:441-446.

Tsao, C.W., A.W. Aday, Z.I. Almarzooq, A. Alonso, A.Z. Beaton, M.S. Bittencourt, A.K.
Boehme, A.E. Buxton, A.P. Carson, Y. Commodore-Mensah, M.S.V. Elkind, K.R.
Evenson, C. Eze-Nliam, J.F. Ferguson, G. Generoso, J.E. Ho, R. Kalani, S.S. Khan,
B.M. Kissela, K.L. Knutson, D.A. Levine, T.T. Lewis, J. Liu, M.S. Loop, J. Ma, M.E.
Mussolino, S.D. Navaneethan, A.M. Perak, R. Poudel, M. Rezk-Hanna, G.A. Roth, E.B.
Schroeder, S.H. Shah, E.L. Thacker, L.B. VanWagner, S.S. Virani, J.H. Voecks, N.Y.
Wang, K. Yaffe, S.S. Martin, E. American Heart Association Council on, C. Prevention
Statistics, and S. Stroke Statistics. 2022. Heart Disease and Stroke Statistics-2022
Update: A Report From the American Heart Association.
Circulation:CIR0000000000001052.

Turner, C.J., K. Badu-Nkansah, and R.O. Hynes. 2017. Endothelium-derived fibronectin
regulates neonatal vascular morphogenesis in an autocrine fashion. *Angiogenesis*.
20:519-531.

Walker, J.L., A.K. Fournier, and R.K. Assoian. 2005. Regulation of growth factor
signaling and cell cycle progression by cell adhesion and adhesion-dependent changes
in cellular tension. *Cytokine Growth Factor Rev*. 16:395-405.

Wang, X., and S. Astrof. 2016. Neural crest cell-autonomous roles of fibronectin in
cardiovascular development. *Development*. 143:88-100.

Weber, G.F., M.A. Bjerke, and D.W. DeSimone. 2011. Integrins and cadherins join
forces to form adhesive networks. *J Cell Sci*. 124:1183-1193.

Xia, M., W. Luo, H. Jin, and Z. Yang. 2019. HAND2-mediated epithelial maintenance
and integrity in cardiac outflow tract morphogenesis. *Development*. 146.

Zhang, Q., D. Carlin, F. Zhu, P. Cattaneo, T. Ideker, S.M. Evans, J. Bloomekatz, and
N.C. Chi. 2021. Unveiling Complexity and Multipotentiality of Early Heart Fields. *Circ*
Res. 129:474-487.

Zimmerman, S.P., S.B. Asokan, B. Kuhlman, and J.E. Bear. 2017. Cells lay their own
tracks - optogenetic Cdc42 activation stimulates fibronectin deposition supporting
directed migration. *J Cell Sci*. 130:2971-2983.
1175

Figure 1

Figure 1. The expression of Fn1 in Mesp1 lineage is required for the OFT elongation.

Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 and staged according to somite number. **A-D.** Embryos between 18-19 somites. n=5 controls and 4 mutants. **E-H.** Embryos between 20-25 somites, n= 5 control and 7 mutant embryos. Imaris was used to surface distal OFT (yellow), proximal OFT (turquoise), and the left ventricle and atria (pink). **C-D, G-H.** The length of the distal and proximal OFT in each embryo was normalized to the height of the left ventricle to control for embryo size.

I-J. Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (20-25s) and stained with DAPI and the MF-20 antibody. **K.** Quantification of angles in controls and mutants as shown in **I-J**. n= 9 controls and 13 mutants. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. OFT-outflow tract, MF-20- antibody detects sarcomeric myosin heavy chain. All scale bars : 200 μm

Figure 2

Figure 2. The expression of Fn1 in the mesoderm is important for the regulation of cell shape in the anterior DPW at E8.5. **A.** Fn1$^{flox/+}$; Mesp1$^{Cre/+}$ (Control) and **E.** Fn1$^{flox/-}$; Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E8.5 (9-11s) and stained to detect GFP (green) and nuclei (DAPI, blue). Regions marked by dashed rectangles in **A, B** are magnified in **A1-E1** and **A2-E2**. **B-C, F-G.** Outlines of aDPW and pDPW cells. **D** and **H.** 3D reconstruction of aDPW cells. **I-L.** Cell shape analyses. Small dots mark data points from each cell; large dots mark the average per embryo. Data points from one embryo are marked with the same color. **I-J**. Cell circularity. **I.** 320 cells from 5 control and 367 cells from 5 mutant embryos were analyzed. **J.** 311 cells from 5 control and 326 cells from 5 mutant embryos were analyzed. **K-L.** Ellipticity and volume of aDPW cells; 51 cells from 5 control and 50 cells from mutant embryos were analyzed. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. OFT-outflow tract, aDPW-anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, A-atrium, V-ventricle

Figure 3

Figure 3. Expression of Fn1 in the mesoderm regulates Golgi apparatus orientation in the anterior DPW. A-A2. Fn1[flox/+];Mesp1[Cre/+] (Control) and **B-B2.** Fn1[flox/-]; Mesp1[Cre/+] (Mutant) embryos were dissected at E8.5 (9-11s) and stained to detect GFP (green), GM130 (white), and nuclei (DAPI, blue). Boxes in **A, B** are magnified in the panels to the right. Arrows point to Golgi. **A3, A4, B3, B4.** Semicircular histograms represent the distribution of the Golgi orientation angles in SHF cells that were measured as is indicated in **C** . N=total number of cells analyzed in 3 controls and 3 mutants. **D, E.** Quantification of Golgi polarization. Golgi was considered polarized if the angle between the Golgi and the plane of the endoderm fell between 85^0 and 95^0. OFT-outflow tract, aDPW-anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, A-atrium, V-ventricle.

Figure 4

Figure 4. Expression of Fn1 in the mesoderm regulates cell polarity in the anterior DPW. A. Fn1^{flox/+}; Mesp1^{Cre/+} (Control) and **B.** Fn1^{flox/-}; Mesp1^{Cre/+} (Mutant) embryos were dissected at E8.5 (9-11s), and stained to detect GFP (green), podocalyxin (white), and nuclei (DAPI, blue). Boxes in **A** and **B** were expanded in **A1** and **B1** to show the anterior DPW and in **A2** and **B2** to show the posterior DPW. Arrows in **A1** point to cells with apical localization of podocalyxin. Arrows in **B1** point to defective basolateral localization of podocalyxin in mutants. **A2, B2.** Podocalyxin is localized apically in the posterior DPW, arrows. **C-D** Quantification of cells with defective podocalyxin localization in the anterior and posterior DPW. **C.** 148 cells in the aDPW from 3 controls and 224 cells in aDPW from 3 mutants were analyzed. **D.** 149 cells in the pDPW from 3 controls and 182 cells in the pDPW from 3 mutants were analyzed. Each small dot represents data from one slice, and each large dot is an average of all slices per embryo. Data from the same embryo is marked by the same color. **E-G.** Fn1^{flox/+}; Mesp1^{Cre/+} (Control) and **H-J.** Fn1^{flox/-}; Mesp1^{Cre/+} (Mutant) embryos were dissected at E8.5 (7-8s), and stained for GFP (green), aPKC, laminin γ1, and nuclei (DAPI, blue). Arrows in **F** point to apical localization of aPKC in control DPW cells. **I.** Arrows point to defective basolateral localization of aPKC in mutants. Arrowheads in **G** point to basal localization of laminin γ1 in controls and in **J** to disorganized localization of laminin γ1 in one of three mutants. **K.** Quantification of cells in which aPKC was re-distributed away from the apical surface. DPW cells from 5 different slices were evaluated in 3 control and 3 mutant embryos. Each small dot represents data from one slice, and each large dot is an average of all slices per embryo. Data from the same embryo is marked by the same color. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. OFT-outflow tract, aDPW-anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, A-atrium, V-ventricle.

Figure 5

Figure 5. Mesodermal Fn1 regulates the expression of N-Cadherin at cell-cell junctions in the DPW. A-B. Fn1flox/+;Mesp1Cre/+ (Control) and **C-D.** Fn1flox/-; Mesp1Cre/+ (Mutant) embryos were dissected at E9.5 (21-26s) and stained to detect GFP (green), N-cadherin (white) and nuclei (DAPI, blue). Junctional N-Cadherin is enriched in the anterior DPW (boxed region in **B**). Boxes in **B-D** are expanded in **B1 and D1.** Arrows point to DPW. **E.** Western blot analysis of total cell lysates and cell surface proteins. Positions of molecular weight markers are indicated on the left. **F-G**. Densitometrical quantification of immunoblot signals. N = 4 for each genotype. Each experiment is represented by a dot of a different color. Means (horizontal bars) and standard deviations (error bars) are displayed, and a 2-tailed, unpaired Student's t-test was used to determine p values. **H-K.** Immunostaining to detect GFP (green), N-cadherin (white) and nuclei (DAPI, blue) in Fn1flox/-, Fn1Δ/- (Fn1-null), Fn1flox/+, and Fn1Δ/+ MEFs. Boxes in **I-K** and in **M-O** are expanded in **I1-K1 and M1-O1.** Arrows point to N-cadherin at cell junctions in control MEFs (**I1, M1, and O1**) and to the absent N-cadherin in Fn1-null MEFs (**K1**). DPW- dorsal pericardial wall, OFT-outflow tract, A-atrium.

Figure 6. Mesodermal Fn1 regulates the length and organization of F-actin in the anterior DPW. A-B1. Fn1[flox/+]; Mesp1[Cre/+] (Control) and **C-D1.** Fn1[flox/-]; Mesp1[Cre/+] (Mutant) embryos were dissected at E9.5 (21-26s), and transverse cryosections (a plain of sections shown by the dashed line in **E**) from the anterior DPW were stained to detect GFP (green), F-actin (white), and nuclei (DAPI, blue). Yellow arrows point to F-actin in the aDPW in **B1** and **D1**. Yellow arrowheads point to apically enriched F-actin in B1 and re-distributed F-actin in **D1**. **E.** Schematic representation of the different regions in the DPW. **F-H.** Traces of F-actin in control and mutants. **G-I.** Angles made by F-actin filaments relative to cells' apical surface were measured. **J.** Quantification of actin polarization. An actin filament was considered polarized if was between 85 and 95 degrees relative to the apical surface of the DPW. **K** Quantification of the F-actin length in the anterior DPW; 235 cells from 3 controls and 417 cells from 5 mutants were examined. Small dots mark the F-actin length of individual cells, and large dots mark the average per slice. Dots from the same embryo are displayed in the same color. Means (horizontal bars) and standard deviations (error bars) are displayed; p was calculated using a 2-tailed, unpaired Student's t-test. DPW-dorsal pericardial wall, A-atrium, NT-Neural tube, AVC-atrioventricular canal, OFT-outflow tract, V-ventricle.

Figure 7

Figure 7 Mesodermal Fn1 regulates the expression of pMLC in the anterior DPW. A-B2. Fn1^{flox/+}; Mesp1^{Cre/+} (Control) and **C-D.** Fn1^{flox/-}; Mesp1^{Cre/+} (Mutant) embryos were dissected at E9.5 (21-26s), and sagittal cryosections were stained to detect GFP (green), F-actin (white) and nuclei (DAPI, blue). Boxes in **A-B** were expanded in **B1** and **B2** to show the anterior and posterior DPW, respectively. Boxes in **C-D** were expanded, in **D1** and **D2**, to show the anterior and posterior DPW, respectively. Arrows point to the apical enrichment of F-actin in the anterior and posterior of controls and to the lack of such enrichment in the anterior DPW of the mutants (**D1**). 6 controls and 3 mutants were analyzed. **E-H2.** Fn1^{flox/+}; Mesp1^{Cre/+} (Control) and **C-D.** Fn1^{flox/-}; Mesp1^{Cre/+} (Mutant) embryos were dissected at E9.5 (21-26s) and sagittal cryosections were stained to detect GFP (green), pMLC (white) and nuclei (DAPI, blue). Boxes in **E-F** are expanded in **F1** and **F2** to show the anterior and posterior DPW, respectively. Boxes in **G-H** are expanded in **H1** and **H2** to show anterior and posterior DPW, respectively. Arrows in **F1** indicate cells with apical localization of pMLC. Arrows in **H1** show a defective basolateral localization of pMLC in mutants. **F2, H2.** pMLC is localized apically in the posterior DPW. 8 controls and 3 mutants were analyzed.

Figure 8

Figure 8. Mesodermal Fn1 regulates nuclear localization of YAP. A. Ventral view of E9.5 embryo. YAP localization was assayed in the DPW between the left and the right limits (marked by the vertical lines).

B-B2. Fn1[flox/+]; Mesp1[Cre/+] (Control) and **C-C2.** Fn1[flox/-]; Mesp1[Cre/+] (Mutant) embryos were dissected at E9.5 (23-26s) and stained to detect nuclei (DAPI, magenta) and YAP (cyan). Regions outlined by dashed rectangles are expanded in **B1, C1** (anterior DPW) and **B2, C2** (posterior DPW). **D-E**. Quantification of the intensity ratio between nuclear and cytosolic YAP in the aDPW and pDPW. 7-8 optical slices from 4 controls and 7-8 optical slices from 4 mutant embryos were analyzed. Each small dot represents one slice, and each large dot is an average of all slices in one embryo. Data from the same embryo is marked by the same color. **F.** Summary: YAP is enriched in the nuclei in the presence of both the mesodermal (red lines) and endodermal (blue lines) Fn1. In the absence of mesodermal Fn1, nuclear translocation of YAP are decreased. aDPW- anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, OFT-outflow tract.

Figure 9

Figure 9. Mesodermal Fn1 is required for DPW cell proliferation. A-C. Fn1$^{flox/+}$;Mesp1$^{Cre/+}$ (Control) and **D-F** Fn1$^{flox/-}$;Mesp1$^{Cre/+}$ (Mutant) embryos were dissected at E9.5 (20-25s) and stained for PHH3 (white), and nuclei (DAPI, blue) E9.5 (20-25s). **A,B, D, E**. DPW (magenta) and PHH3$^+$ cells in the DPW (white) were segmented using Imaris. **C,F**. Proportion of PHH3+ cells in the aDPW (magenta) and pDPW (turquoise) were quantified in **G-H**. Each dot marks one embryos. n=4 control and 6 mutant embryos. Means (horizontal bars) and standard deviations (error bars) are displayed, 2-tailed, unpaired Student's t test was used to determine p values. aDPW-anterior dorsal pericardial wall, pDPW- posterior dorsal pericardial wall, PHH3- Phosphorylated Histone H3.

Figure 10

Figure 10. Expression patterns of *Fn1 mRNA* and protein in controls and mutants. **A-J.** Expression of *Fn1 mRNA*. Embryos were dissected at E9.5 (23-26s) and labeled with DAPI (magenta) and anti-sense Fn1 probe (turquoise). **A-B2** Fn1^{flox/+}; Mesp1^{+/+} (designated as WT), **C.** *Fn1 mRNA* intensity ratio between aDPW and pDPW, **D-F2.** Fn1^{flox/+}; Mesp1^{Cre/+} (Control) **G.** *Fn1 mRNA* intensity in the anterior (**H**) and posterior (**J**) DPW was normalized to that in the endoderm. 6 different slices from 3 WT, 4 control and 9 mutant embryos were analyzed. **H-I2.** Fn1^{flox/-};Mesp1^{Cre/+} (Mutant) embryos. Regions outlined by the dotted rectangles in **B, E,** and **I** are expanded in panels to the right and color-coded according to fluorescence intensity (scale is on the right). White arrows point to DPW. Yellow arrowheads point to the endoderm. **K.** Ventral view of the DPW, Fn1 expression is color-coded according to fluorescence intensity. **L.** Ratio of Fn1 protein intensity shows similar enrichment of Fn1 in the anterior DPW of controls and mutants. **M, M1.** Fn1^{flox/+}; Mesp1^{Cre/+} (Control) and **N, N1.** Fn1^{flox/-};Mesp1^{Cre/+} (Mutant) embryos were dissected at E9.5 (22-26s) and stained to detect Fn1 protein (white) and nuclei (DAPI, blue). Fn1 protein expression in the DPW (**M, N**) is color-coded according to fluorescence intensity in **M1, N1**. **O-P.** Fn1 protein levels underlying the anterior DPW are unchanged in the mutants. Fn1 intensity in the aDPW and pDWP regions outlined by the red dash lines in **M1-N1** was normalized by DAPI intensity; 7 optical slices from 8 control and 10 mutant embryos were evaluated. Each small dot represents one slice, and each large dot is an average of all slices per embryo. Data from the same embryo is marked by the same color. Means (horizontal bars) and standard deviations (error bars) are displayed, 2-tailed, unpaired Student's t-test was used to determine p values. aDPW-anterior dorsal pericardial wall, pDPW-posterior dorsal pericardial wall, OFT-outflow tract, A-atrium, LV-left ventricle.

Figure 11

Figure 11. The central role of the mesodermal Fn1 in regulating the elongation of the cardiac outflow tract. Fn1 synthesized by the mesoderm mediates the elongation of the cardiac outflow tract by regulating cell shape, polarity, proliferation, and mechanotransduction in the anterior dorsal pericardial wall (aDPW).