Distinct *Wolbachia* localization patterns in oocytes of diverse host species reveal multiple strategies of maternal transmission

Yonah A. Radousky[1,*], Michael T.J. Hague[2,*,†], Sommer Fowler[1], Eliza Paneru[1], Adan Codina[1], Cecilia Rugamas[1], Grant Hartzog[1], Brandon S. Cooper[2], William Sullivan[1]

[1]Department of Molecular, Cell, and Developmental Biology, University of California Santa Cruz, Santa Cruz, California, United States of America

[2]Division of Biological Sciences, University of Montana, Missoula, Montana, United States of America

[*]Co-first authors

[†]Corresponding Author:
Michael T.J. Hague
Division of Biological Sciences, University of Montana
32 Campus Dr. HS 104, Missoula, MT 59812
michael.hague@mso.umt.edu

ABSTRACT

A broad array of endosymbionts radiate through host populations via vertical transmission, yet much remains unknown concerning the cellular basis, diversity and routes underlying this transmission strategy. Here we address these issues, by examining the cellular distributions of *Wolbachia* strains that diverged up to 50 million years ago in the oocytes of 18 divergent *Drosophila* species. This analysis revealed three *Wolbachia* distribution patterns: 1) a tight clustering at the posterior pole plasm (the site of germline formation); 2) a concentration at the posterior pole plasm, but with a significant bacteria population distributed throughout the oocyte; 3) and a distribution throughout the oocyte, with none or very few located at the posterior pole plasm. Examination of this latter class reveals *Wolbachia* accesses the posterior pole plasm during the interval between late oogenesis and the blastoderm formation. We also find that one *Wolbachia* strain in this class concentrates in the posterior somatic follicle cells that encompass the pole plasm of the developing oocyte, suggesting these are the source of *Wolbachia* that ultimately occupy the germline. In contrast, strains in which *Wolbachia* concentrate at the posterior pole plasm generally exhibit no or few *Wolbachia* in the follicle cells associated with the pole plasm. Phylogenomic analysis indicates that closely related *Wolbachia* strains tend to exhibit similar patterns of posterior localization, suggesting that specific localization strategies are a function of *Wolbachia*-associated factors. Previous studies revealed that endosymbionts rely on *one* of two distinct routes of vertical transmission: continuous maintenance in the germline (germline-to-germline) or a more circuitous route via the soma (germline-to-soma-to-germline). Here we demonstrate that *Wolbachia* strains infecting *Drosophila* species maintain the diverse arrays of cellular mechanisms necessary for *both* of these distinct transmission routes. This characteristic may account for its ability to infect and spread globally through a vast range of host insect species.

INTRODUCTION

A large class of endosymbionts rely on efficient vertical transmission to ensure their maintenance and spread through host populations. Many insect endosymbionts achieve this by a strict association with the host germline through successive generations. That is, during oogenesis, endosymbionts target the extreme posterior of the oocyte, the site of germline formation. In contrast to this strict germline-to-germline route of maternal transmission, other endosymbionts occupy the germline via invasion from neighboring somatic cells (Russell et al., 2019). Occupation of the germline via a somatic route requires that the endosymbiont has the capacity to undergo cell-to-cell transmission.

Among all known endosymbionts, maternally transmitted *Wolbachia pipientis* is the most common, infecting 30 to 60% of all insect species (Hilgenboecker et al., 2008; Zug and Hammerstein, 2012). Hosts acquire *Wolbachia* from other species via horizontal (non-sexual) transfer or from sister species via introgressive transfer (O'Neill et al., 1992; Raychoudhury et al., 2009; Gerth and Bleidorn, 2016; Conner et al., 2017; Turelli et al., 2018; Cooper et al., 2019). Following establishment, maternal transmission rates in conjunction with *Wolbachia* effects on host fitness and reproduction determine *Wolbachia* spread and prevalence in host populations (Hoffmann et al., 1990; Turelli and Hoffmann, 1995; Cooper et al., 2017; Cogni et al., 2021).

While sometimes imperfect, the success and prevalence of *Wolbachia* ultimately depends on vertical transmission through the female (Hoffmann et al., 1990; Turelli and Hoffmann, 1995; Carrington et al., 2011). The ability of *Wolbachia* to establish and persist in diverse insect species indicates that it has evolved to usurp highly conserved host processes and components. For example, studies of *w*Mel *Wolbachia* in model organism *D. melanogaster* suggest that *Wolbachia* engage conserved host motor proteins Dynein and Kinesin in order to target the extreme posterior of the oocyte, the site of germline formation (Ferree et al., 2005; Serbus and Sullivan, 2007; Russell et al., 2018).

To study the vertical transmission of *Wolbachia* in *Drosophila* we take advantage of the wealth of knowledge regarding the molecular and cell biology of *Drosophila* oogenesis (for a review, see Bastock and St Johnston, 2008). *Drosophila* oocytes contain two clusters of ovarioles, each containing two to three germline stem cells and various stages of developing oocytes. The stem cells undergo asymmetric mitotic division in which one daughter cell undergoes self-renewal and the other differentiates into a cytoblast. The cytoblast undergoes four rounds of mitotic divisions resulting in 16 interconnected cells, one of which becomes the oocyte nucleus while the others differentiate into polyploid nurse cells. The developing oocyte is encapsulated by a monolayer of somatically derived follicle cells resulting in the formation of discrete egg chambers as the maturing oocyte progresses through the ovariole. During the late stages of oogenesis, the nurse cell cytoplasm is expelled into the oocyte, resulting in a rapid doubling of the oocyte volume and concomitant shrinking of nurse cell volume (Theurkauf and Hazelrigg, 1998; Bastock and St Johnston, 2008). This is followed by posterior pole plasm formation and establishment of the germline (Bilinski et al., 2017).

Wolbachia distribution and localization in insect oocytes and early embryos has been well characterized in *w*Mel-infected *D. melaongaster* (Kose and Karr, 1995; Veneti et al., 2004; Ferree et al., 2005; Serbus et al., 2008; Ramalho et al., 2018; Russell et al., 2018; Guo et al., 2019). Early

in oogenesis, *w*Mel *Wolbachia* are evenly distributed throughout all the cells in the chamber. At this stage, microtubules emanate from the oocyte into the nurse cells (Theurkauf, 1994; Grieder et al., 2000). By associating with the minus-end directed microtubule motor protein Dynein, *w*Mel *Wolbachia* are transported from the nurse cells and concentrate at the anterior end of the oocyte (Ferree et al., 2005). As the oocyte matures, microtubules undergo a dramatic reorganization such that they originate from the oocyte cortex with their plus-ends oriented toward the posterior pole (Theurkauf et al., 1992; Steinhauer and Kalderon, 2006). This reorganization is accompanied by a release and dispersal of *w*Mel *Wolbachia* from the anterior pole. Then, *w*Mel *Wolbachia* associate with the plus-end directed motor protein Kinesin for transport to the posterior pole and the site of germline formation (Serbus and Sullivan, 2007). There, *Wolbachia* (*w*Mel) associate with pole plasm determinants maintaining their position at the site of germline formation (Serbus and Sullivan, 2007) in spite of extensive cytoplasmic streaming (Serbus et al., 2005).

Thus, *w*Mel *Wolbachia* location and migration patterns provide insights into the host factors with which *Wolbachia* engages throughout oogenesis. The *D. simulans*-associated *w*Ri strain exhibits a similar anterior distribution in early oogenesis, and then becomes distributed across the entire mature oocyte. However, unlike *w*Mel, it does not concentrate at the posterior pole of the oocyte (Veneti et al., 2004; Serbus and Sullivan, 2007). This suggests *w*Ri engages host Dynein and Kinesin, but not pole plasm determinants. *w*Mel *Wolbachia* strains introgressed into a *D. simulans* host background exhibit a concentration in the pole plasm, indicating the *Wolbachia* genome rather than the host genome drives oocyte localization properties (Veneti et al., 2004; Serbus and Sullivan, 2007).

With the exception of *w*Mel and *w*Ri, little is known of the extent to which distribution patterns in the oocyte and transmission routes vary among diverse *Wolbachia* strains and host species. Here, we fill this important knowledge gap, by examining 18 *Drosophila* host species infected with *Wolbachia* strains that diverged up to 50 million years ago. These studies also provide a direct read-out of *Wolbachia* and host motor protein/pole plasm interactions that are highly conserved and those that vary among species. This work complements and extends previous analyses revealing *Wolbachia* strain-dependent localization patterns in the *Drosophila* oocyte and embryo (Veneti *et al.*, 2004; Toomey et al., 2013).

We discovered that *Wolbachia* distribution patterns in the *Drosophila* oocytes fall into three categories: (1) the vast majority of *Wolbachia* clusters at the posterior pole; (2) a concentration at the posterior pole, but with a large portion of *Wolbachia* distributed throughout the entire oocyte; and (3) *Wolbachia* distributed throughout oocyte, with none or very few located at the posterior pole. This latter class is particularly interesting as it suggests *Wolbachia* must enter the oocyte via a somatic route for transmission to the next generation. We also provide evidence that the neighboring somatic follicle cells are the likely source of *Wolbachia* entering the germline. Together, these findings indicate that *Wolbachia* are capable of achieving efficient vertical transmission either via strict maintenance in the germline from one generation to the next or via invasion of the germline through neighboring somatic cells. Thus, *Wolbachia* must maintain the diverse arrays of cellular mechanisms necessary for *both* of these distinct transmission routes. We identify candidate surface proteins that may be responsible for these diverse localization patterns in host oocytes.

RESULTS

***Wolbachia* navigates the developing *Drosophila* oocyte in distinct stages**

As described above, *w*Mel *Wolbachia* migration and distribution throughout oogenesis in *D. melanogaster* has been well documented (Kose and Karr, 1995; Veneti et al., 2004; Serbus and Sullivan, 2007; Serbus et al., 2008; Ramalho et al., 2018; Russell et al., 2018; Guo et al., 2019). As a reference for our analysis of *Wolbachia* oocyte distributions in a range of *Drosophila* species, the key distribution patterns of *w*Mel throughout *D. melanogaster* oocytes are depicted in **Figure 1**.

Stage 1 includes the germarium, a structure consisting of the germline stem cells and their cystoblast daughter cells that will differentiate into a mature oocytes. Progression to stage 2, involves the cystoblast undergoing four rounds of mitotic division to produce a syncytium of 16 cells connected by cytoplasmic bridges and surrounded by a layer of somatically derived follicle cells (Bastock and St Johnston, 2008). Propidium Iodine (PI)-stained *Wolbachia* (red) are clearly observed as puncta in the cytoplasm of the infected, but not the uninfected 16 cell cyst (**Figure 1 Stages 1-2, Arrows**). Actin is labeled in green.

During stage 3, the cyst has become polarized with one of the cells differentiating into an oocyte and the others becoming polyploid nurse cells. *Wolbachia* are clearly seen in the infected oocyte cytoplasm anteriorly concentrated at two cytoplasmic bridges that connect to the nurse cells (**Figure 1, Stage 3, Arrows**). Previous studies indicate that *Wolbachia* rely on the extensive complex of microtubules that radiate through the oocyte/nurse cell complex undergoing Dynein mediated transport to the anterior region of the oocyte (Ferree et al., 2005).

During stages 4 and 5, the nurse cell/oocyte complex becomes oblong and is fully encapsulated by follicle cells. *Wolbachia* are clearly observed in both the oocyte and nurse cell cytoplasm (**Figure 1, Stages 4 and 5, Arrows**). The *Wolbachia* in the oocyte are tightly associated with the anterior cortex.

During stage 6, the number of follicle cells increases and they are more densely packed in a monolayer encompassing the oocyte. Here we discovered a previously unrecognized aspect of *Wolbachia* dynamics and localization: the number of anteriorly localized *Wolbachia* is greatly diminished at this stage (**Figure 1, Stage 6, Arrow and Supplemental Figure 1**). Possibly *Wolbachia* in the oocyte are degraded or migrate through the ring canals back into the nurse cells (from whence they came). If the latter, it may be *Wolbachia* are prematurely engaging with host Kinesin (see below).

At stage 7, *Wolbachia* are again observed in the oocyte. However, rather than being tightly associated with the anterior cortex, *Wolbachia* are evenly distributed throughout the oocyte (**Figure 1, Stage 7, Arrow**). Up through stage 6**,** the microtubules are organized with their minus-end located at the posterior pole of the oocyte with their plus-ends extending through the ring canals into the nurse cell complex (Mahowald and Strassheim, 1970). However, during stage 7, the microtubules reorient such that their plus-ends extend toward the posterior pole (Cha et al., 2001; Cha et al., 2002; Steinhauer and Kalderon, 2006). Accordingly, the plus-end motor protein Kinesin is required to establish the even distribution of *Wolbachia* throughout the maturing oocyte

(Serbus and Sullivan, 2007). Stage 8 heralds the beginning of vitellogenesis, the process of yolk formation and deposition of nutrients into the oocyte (Cummings et al., 1971). *Wolbachia* are near, but not yet located at, the posterior pole (**Figure 1, Stage 8, Arrow**). At stage 9, anterior follicle cell migration occurs (Rørth, 2002). At this stage, *Wolbachia* displays an archlike posterior cortical localization. (**Figure 1, Stage 9, Arrow**). By stage 10, the oocyte accounts for approximately half the volume of the egg chamber. Nurse cell dumping will shortly follow, leading into stage 11-12 with the oocyte rapidly growing in size (Mahajan-Miklos and Cooley, 1994). At stage 10, there is a distinct concentration of *Wolbachia* along the posterior cortex (**Figure 1, Stage 10, Arrow**).

As described above, much is known concerning *w*Mel *Wolbachia* dynamics and localization patterns during oocyte development in *D. melanogaster*, as well as the host factors with which *Wolbachia* interacts (Serbus et al., 2008; Russell et al., 2019). These studies provide an outline of the molecular and cellular mechanisms driving *Wolbachia* vertical transmission. They also raise the question of whether the mechanisms and strategies observed in *D. melanogaster* are conserved across diverse *Wolbachia* strains and *Drosophila* species. To explore this issue, the *w*Mel *Wolbachia* distribution in *D. melanogaster* serves as the basis for our comparison of 19 other *Wolbachia* strains. Bayesian phylograms depicting phylogenetic relationships among the *Wolbachia* strains and the *Drosophila* species are shown in **Figure 4 and Supplemental Figure 2**. Our analysis comprised 18 A-group *Wolbachia* strains, including six *w*Mel-like strains (Cooper et al., 2019; Hague et al., 2020a) and five *w*Ri-like strains (Turelli et al., 2018), as well as the B-group *Wolbachia* strain *w*Mau, which diverged from A-group *Wolbachia* up to 50 million years ago (Meany et al., 2019). These *Wolbachia* strains infect 18 divergent *Drosophila* host species spanning three species groups (Suvorov et al., 2022).

An initial concentration of *Wolbachia* at the oocyte anterior is a conserved feature

As described above, *w*Mel *Wolbachia* enter the oocyte through a complex of ring canals connecting the nurse cells to the oocyte and concentrate at the anterior cortex. Initially observed in *D. melanogaster*, the number of *Wolbachia* increases dramatically while at the anterior (Ferree et al., 2005). This concentration is at least in part due to import of *Wolbachia* from the nurse cells through the ring canals into the oocyte. The increase may result from transport from the nurse cells, or from *Wolbachia* replication at this anterior site, or both. To determine if anterior localization is a conserved aspect of *Wolbachia*'s navigation through the developing oocyte, we first characterized this trait in eight *Wolbachia* strains in their respective hosts: *w*Pan in *D. pandora*, *w*Nik in *D. nikananu*, *w*Cha in *D. chauvacae*, *w*Ha in *D. simulans*, *w*Tsa in D. *tsacasi*, *w*Boc in *D. bocki*, *w*Ana in *D. ananassae*, and *w*Mel in *D. melanogaster*. All eight strains exhibited a distinct anterior localization during mid-oogenesis (**Figure 2**). It is interesting that this localization occurs during stage 5 of oogenesis, which, in *D. melanogaster,* is prior to the localization of all known anterior axis determinants. Thus, *Wolbachia* likely relies on as yet undiscovered anteriorly concentrated factor(s). The functional significance of this anterior localization remains unknown. Given the dramatic increase in *Wolbachia* abundance, the cytoplasmic environment of this anterior region may be rich in membrane and other components necessary for *Wolbachia* replication.

Wolbachia exhibits three distinct distributions with respect to posterior localization in the mature _Drosophila_ oocyte

In _D. melanogaster_, oocyte microtubules undergo a dramatic rearrangement during stages 7-8 of mid-oogenesis (Theurkauf et al., 1992). Microtubule minus-ends, originally concentrated at the posterior pole with their plus-ends extending anteriorly through the ring canals, reorganize such that the microtubule minus-ends orient toward anteriorly. Concomitant with this microtubule rearrangement is the release of _w_Mel _Wolbachia_ from the oocyte anterior resulting in the dispersal of the bacteria throughout the entire length of the oocyte with a fraction concentrating at the posterior pole (Ferree et al., 2005). The posterior dispersal of _Wolbachia_ requires the plus-end motor protein Kinesin (Serbus and Sullivan, 2007). Maintenance of those _Wolbachia_ that reach the posterior pole rely on a stable association with key pole plasm components (Serbus et al., 2011).

We next examined the oocytes of 18 _Drosophila_ species infected with 19 diverse _Wolbachia_ strains ($N = 72$ total oocytes) to test the hypothesis that oocyte posterior distributions vary. Unlike the conserved anterior localization, _Wolbachia_ exhibit three distinct patterns of posterior localization in stage 10 oocytes. As shown in **Figure 3** and **Supplemental Figure 3A,** eight _Wolbachia_ strains exhibit a localization pattern similar to that observed for _w_Mel in _D. melanogaster_ with the bulk of _Wolbachia_ evenly distributed throughout the oocyte, and a significant fraction concentrated at the posterior pole. We refer to this pattern as Posteriorly Localized.

Our analysis revealed four _Wolbachia_ strains manifested a second pattern in which the vast majority of _Wolbachia_ localized to a distinct cluster at the posterior pole and only few _Wolbachia_ are distributed throughout the remainder of the oocyte (**Figure 3, Supplemental Figure 3B**). In contrast to the pattern described above, rather than being distributed in a wide arc along the posterior pole, they are tightly centered at the extreme posterior region of the oocyte (**Figure 3B**). We refer to this pattern as Posteriorly Clumped.

Finally, seven _Wolbachia_ strains exhibit a third pattern in which _Wolbachia_ are distributed throughout the entire oocyte but there is no bacteria concentration at the posterior pole (**Figure 3 and Supplemental Figure 3C).** The lack of the _Wolbachia_ at the posterior pole was unexpected as this was thought to be necessary for incorporation into the germline of the next generation. We refer to this pattern as Dispersed. **Table 1** shows a quantification of the _Wolbachia_ distribution patterns in the examined species**.**

Quantification of _Wolbachia_ oocyte distribution patterns

This work establishes three distinct classes (Posteriorly Localized, Posteriorly Clumped, and Dispersed) that qualitatively describe different _Wolbachia_ localization patterns in host oocytes. We used one-way ANOVAs to test whether these three classes differ statistically in quantitative estimates of _Wolbachia_ abundance in the whole oocyte, at the posterior region, and at the posterior cortex (**Supplemental Figure 4**). We defined the posterior region as the posterior 12.5% of the oocyte, and the posterior cortex as the narrow cortical region of vasa expression (Russell et al., 2018), and we then measured total _Wolbachia_ abundance in each region as the Corrected Total Cell Fluorescence (CTCF). We found that _Wolbachia_ abundance differs significantly among the

three classes (one-way ANOVA, $F_{(2,69)}$ = 6.2, P = 0.003), with Tukey multiple comparisons indicating that the Clumped class has significantly more *Wolbachia* in whole oocytes than the Localized group (P = 0.002). The three classes also differed significantly in the amount of *Wolbachia* in the posterior region ($F_{(2,69)}$ = 23.09, P < 0.001), such that the Dispersed class has significantly fewer *Wolbachia* at the posterior that the Clumped (P < 0.001) and Localized (P < 0.001) groups. Finally, the three classes also differ in the amount of *Wolbachia* at the posterior cortex ($F_{(2,64)}$ = 38.12, P < 0.001). Here, multiple comparisons indicate that all three class significantly differ in levels of *Wolbachia* at the posterior cortex: the Clumped class has the most *Wolbachia*, followed by Localized, and then Dispersed (**Supplemental Figure 4**). Mean estimates of *Wolbachia* abundance (CTCF) for each *Wolbachia* strain and host species are shown in **Table 1**.

Wolbachia abundance in the oocyte posterior has strong phylogenetic signal

We hypothesized that the diversity of *Wolbachia* localization patterns may be due to *Wolbachia*-associated factors, implying that closely related *Wolbachia* strains would exhibit similar localization patterns in host oocytes. We used the phylogram describing the evolutionary relationships among *Wolbachia* strains to test whether patterns of *Wolbachia* localization exhibit a phylogenetic signal using Pagel's λ (Pagel, 1999; Hague et al., 2020b; Hague et al., 2021) (**Figure 4**). A λ value of 1 would be consistent with trait evolution that entirely agrees with the *Wolbachia* phylogeny (i.e., strong phylogenetic signal supporting a role of *Wolbachia* factors), whereas a value of 0 would indicate that trait evolution occurs independently of phylogenetic relationships (Pagel, 1999; Freckleton et al., 2002). Most notably, we found that *Wolbachia* abundance (CTCF) in the posterior region exhibits a strong, significant phylogenetic signal (λ = 0.982 [0, 0.998], P = 0.021). *Wolbachia* abundance at the posterior cortex also has a high, but non-significant λ value (λ = 0.959 [0, 0.995], P = 0.683). Here, the large confidence intervals and our simulations suggest that a larger *Wolbachia* phylogeny (N = 25, 50 *Wolbachia* strains) is required to detect a significant signal of λ > 0 (**Supplemental Figure 5**). These results highlight differences in patterns of *Wolbachia* localization between the *w*Mel- and *w*Ri-like *Wolbachia* clades (**Figure 3**). Closely related *w*Mel-like *Wolbachia* tend to occur at a higher abundance in the posterior region of oocytes, whereas the *w*Ri-like strains are generally more dispersed. We found no evidence that *Wolbachia* abundance in the whole oocyte exhibits a phylogenetic signal (λ < 0.001 [0, 0.753], P = 1).

The correlation between *Wolbachia* localization in oocytes and phylogenetic divergence implies that factors in the *Wolbachia* genome determine *Wolbachia* concentration at the oocyte posterior. Because these distributions likely involve direct interactions between *Wolbachia* and the host cytoplasm, we focused on six major *Wolbachia* surface proteins: the three *wsp* paralogs (WD_1063/*wsp*, WD_0009/*wspB*, WD_0489/*wspC*), as well as WD_0501, WD_1041, WD_1085. We tested whether sequence divergence at any of these candidate loci predicts *Wolbachia* localization patterns. Specifically, we tested whether gene trees of the *Wolbachia* surface proteins (**Supplemental Figure 6**) exhibit a phylogenetic signal, using the methods described above. One surface protein (WD_1085) stood out with especially strong evidence of a phylogenetic signal, with significant departures from λ = 0 for *Wolbachia* abundance at the posterior region (λ = 0.974 [0, 0.998], P = 0.001) and the posterior cortex (λ = 0.942 [0, 0.994], P = 0.028). In addition, *Wolbachia* abundance at the posterior region also exhibited significant phylogenetic signal on the

gene tree of WD_0501 (λ = 0.992 [0.521, 1], P = 0.022). All tests for phylogenetic signal using *Wolbachia* gene trees are summarized in **Supplemental Table 1**.

To gain insight into WD_1085 and WD_0501 cellular function, we ectopically expressed codon-optimized forms of the genes encoding these proteins in yeast using a Galactose-inducible promoter. Under normal growth conditions, ectopic expression of both genes inhibited growth, with WD_1085 exhibiting a more pronounced inhibition (**Supplemental Figure 7**, **Top Panel**, **Bottom Row**). Because *Wolbachia* exhibits a close association with microtubules, we also assayed whether ectopic expression of these genes enhanced the growth defects in situations in which the microtubule spindle assembly checkpoint was compromised by a mad1 mutation (Luo et al., 2018), or when the microtubules were compromised directly through the addition of benomyl. As indicated in the bottom right panel, ectopic expression of WD_1085 significantly inhibited growth when both the microtubules and spindle checkpoint were compromised (**Supplemental Figure 7**). These results suggest that WD_1085 perturbs microtubule function.

We recently found that the surface protein WD_0009 (*wspB*) is pseudogenized due to a premature stop codon in a tropical *w*Mel variant sampled from Australia, which seems to influence *w*Mel abundance in stage 10 oocytes when hosts are reared in the cold (Hague et al., 2022). Here, we did not find that the *wspB* gene tree exhibits phylogenetic signal related to *Wolbachia* oocyte abundance (**Supplemental Table 1**); however, we note that *wspB* appears to be psuedogenized in a number of other *Wolbachia* strains in addition to tropical *w*Mel. We found derived deletions of varying length in the *wspB* sequences of *w*Ha and *w*Bai beginning at nucleotide position 256, both of which produce frame shifts that generate multiple downstream stop codons (**Supplemental Figure 8**). In addition, *w*Boc and *w*Leo share a large insertion starting at bp position 351. Because the insertions contain a contig break, we were unable to resolve the full length of the insertion. Lastly, *w*Bic contains a 104 bp insertion at bp position 647 that creates multiple downstream stop codons. These results suggest that the surface protein *wspB* has become psuedogenized at least four times in different *Wolbachia* lineages. Notably, the *Wolbachia* strains with a putatively pseudogenized version of *wspB* do not differ from the other strains in mean *Wolbachia* abundance in whole oocytes (Wilcoxon rank sum test, W = 36, P = 0.964), the posterior region (W = 37, P = 0.893), or the posterior cortex (W = 41, P = 0.622). This suggests that psuedogenization of *wspB* does not influence the diverse patterns of *Wolbachia* localization observed here. To gain insight into *wspB* cellular function, we ectopically expressed this gene (and the other *wsp* paralogs; WD_1063/*wsp* and WD_0489/*wspC)* in yeast using the Galactose expression system described above. Using this system, we found that ectopic expression of codon optimized *wspB* and *wsp* significantly inhibited yeast growth **(Supplemental Figure 9)**. Ectopic expression of *wspC* kills the cells. Based on similar ectopic expression studies, the suppression is likely due to *Wolbachia* proteins interacting with and disrupting the function of yeast components and cellular processes that are conserved and performing similar functions in insect cells.

Lastly, we ran similar phylogenetic analyses using the host phylogeny (as opposed to *Wolbachia*) to test whether host-associated factors might also contribute to the diverse *Wolbachia* distribution patterns in fly oocytes (**Supplemental Figure 10**). We generally found that *Wolbachia* localization patterns do not exhibit a phylogenetic signal on the host tree. *Wolbachia* abundance in whole oocytes (λ = 0.365 [0, 0.856], P = 0.656), at the posterior region (λ < 0.001 [0, 0.658], P = 1), and at the posterior cortex (λ < 0.001 [0, 0.637], P = 1) did not exhibit phylogenetic signal using the

host phylogram, implying that host-associated factors are not as important in determining diverse *Wolbachia* localization patterns in oocytes.

Vertical transmission in some species likely relies on *Wolbachia* entering the germline from neighboring somatic tissues

Because *Wolbachia* strains with a Dispersed localization do not localize to the pole, one would expect maternal transmission to be compromised. However, previous studies demonstrate that some of these *Wolbachia*, such as the B-group strain *w*Mau in *D. mauritiana*, have high maternal transmission rates under standard laboratory conditions (i.e., constant 25°C), equivalent to that of *w*Mel in *D. melanogaster* (Meany et al., 2019; Hague et al., 2022). These findings suggest that, in addition to strict germline-to-germline transmission, *Wolbachia* may utilize alternative routes of maternal transmission. Because numerous vertically inherited endosymbionts are transmitted via cell-to-cell transmission from the soma to the germline (Frydman, 2006; Russell et al., 2019), we explored whether *Wolbachia* that infect *Drosophila* exploits this strategy.

To determine the developmental stage during which somatic *Wolbachia* invade the germline in Dispersed species, we focused on B-group *w*Mau in *D. mauritiana.* Initially we examined the female germline (oocytes) of 3rd instar larva for the presence of *Wolbachia*. For comparison, we also examined the oocytes of *w*Mel-infected *D. melanogaster* (a strain with a Posteriorly Localized concentration of *Wolbachia*) 3rd-instar larvae using anti-FtsZ to mark the *Wolbachia* and anti-vasa to mark the germline. As shown in **Figure 5**, *Wolbachia* is abundantly present in ring-like patterns in many of the vasa-positive cells in the center of the developing oocyte. Imaging the oocytes of *w*Mau-infected *D. mauritiana* revealed a similar pattern with *Wolbachia* concentrated in the germline stem cells at the center of the oocyte (**Figure 5**). Images of *w*Ri in *D. simulans* as well as *w*Bic in *D. bicornuta*—strains with a Dispersed localization and Posteriorly Clumped localization in mature oocytes, respectively—displayed the same ring-like structure in the oocyte germline of 3rd instar larva, although they appeared to have a much higher titer than *w*Mau or *w*Mel (**Figure 5**). This indicates that in these species, in which vertical transmission occurs with a few or no *Wolbachia* present at the site of germline formation in mature adult oocytes, *Wolbachia* are able to occupy the germline at some point between the final stages of oocyte maturation and development of the fertilized egg into a 3rd instar larva. Presumably the source of the *Wolbachia* is via invasion from neighboring somatic cells.

To determine if *Wolbachia* occupied the germline prior to the 3rd instar larval stage, we examined the newly formed germline of late blastoderm and cellularized embryos. At this stage, newly formed germline cells, known as the pole cells, form a distinct cluster of cells at the extreme posterior to the embryo. In all the four strains described above, using PI or anti-*Wolbachia* staining, *Wolbachia* are readily found in these posterior germline cells (**Figure 6).** Imaging *w*Mau in *D. mauritiana* early blastoderm embryos revealed the presence of *Wolbachia* in the posterior pole cells. This observation narrows the time window in which *Wolbachia* invades and occupies the germline from between the final stages of oocyte maturation to early blastoderm formation. Images taken of *D. mauritiana* in one-hour old embryos also showed *w*Mau positioned in the posterior pole, narrowing the developmental window of *Wolbachia* germline occupation between the final stages of oocyte maturation to one hour after fertilization **(Figure 6).**

These findings motivated us to examine *w*Mau in *D. mauritiana* oocytes just prior to egg deposition. Previous studies found that in many infected *Drosophila* species, *Wolbachia* is concentrated in the most posterior positioned follicle cells (Kamath et al., 2018). These somatically-derived cells, known as polar cells, directly contact the oocyte posterior pole plasm. In accord with Kamath et al. (2018) studies, we find that *Wolbachia* are sporadically distributed in subsets of more anterior positioned follicle cells encompassing the oocyte. *w*Mau presence in the polar follicle cells is a consistent feature in every *D. mauritiana* oocyte examined (**Figure 7).** These observations suggest that *Wolbachia* in the polar follicle cells is the likely somatic source of germline *w*Mau in *D. mauritiana.*

To determine the extent to which *Wolbachia* polar follicle cell localization is conserved, we analyzed *Wolbachia* localization in the polar and more anterior follicle cells for seven *Wolbachia* strains with the following patterns: Dispersed (N = 3), Posteriorly Localized (N = 3), and Posteriorly Clumped (N = 1) (**Figure 7**). We scored for the presence of follicle cell *Wolbachia* in each oocyte (N = 75) using the following criteria: the presence of *Wolbachia* in either both or one of the posterior polar follicle cells (**Figure 7A)** and the presence of *Wolbachia* in the more anteriorly localized follicle cells (**Figure 7B**). This analysis revealed that in a high percentage of oocytes from Dispersed strains (a lack of the *Wolbachia* at the oocyte posterior), *Wolbachia* are frequently present in the posterior polar follicle cells. In contrast, with the exception of wMel, oocytes examined from strains in which *Wolbachia* are posteriorly concentrated (the Posteriorly Localized and Clumped strains), few possess *Wolbachia* in the posterior follicle cells. We found that the seven *Wolbachia* strains exhibit significant variation in whether *Wolbachia* cells are found in the polar follicle cells ($\chi^2_{(12)}$ = 65.15, P < 0.001) (**Figure 7A**). In particular, the divergent B-group strain *w*Mau stood out with a large proportion of oocytes containing *Wolbachia* in the two polar follicle cells. The A-group *w*Mel strain also had a large portion of oocytes with *Wolbachia* in one or two of the polar cells. We also found that the *Wolbachia* strains differ in whether *Wolbachia* cells are found in the other more anterior follicle cells ($\chi^2_{(6)}$ = 43.29, P < 0.001) (**Figure 7B**). Again, the *w*Mau and *w*Mel strains had a large portion of oocytes with *Wolbachia* in the other more anterior follicle cells encompassing the oocyte. Of note, for *w*Mau in *D. mauritiana*, every oocyte had at least one of their polar cells filled with *Wolbachia*, observationally at a higher titer (**Figure 7**). This suggests that in the Dispersed strains, vertical transmission may be achieved via invasion of *Wolbachia* from the polar follicle cells, particularly for the divergent B-Group *w*Mau strain.

DISCUSSION

Vertical transmission through the female lineage is a common strategy for many endosymbionts. These endosymbionts achieve high efficiencies of maternal transmission either through direct germline-to-germline transmission or invasion of the maternal germline through neighboring somatic cells (Russell et al., 2019). Because the former involves a continuous presence in the germline while the latter requires navigating a soma-to-germline passage, it is expected that each requires distinct interactions with and manipulations of host cellular processes. Thus, it would be expected that a given endosymbiont would have evolved to utilize one, but not both of these strategies. Here we explore this issue by examining cellular aspects of *Wolbachia* vertical transmission. In contrast to expectations, our comprehensive analysis demonstrates that *Wolbachia* rely on both strict germline-to-germline and soma-to-germline vertical transmission strategies. Which strategy employed varies among *Wolbachia* strains.

The initial anterior *Wolbachia* localization in the developing oocyte appears to be a conserved aspect of *Wolbachia* transmission, as all eight strains examined exhibit this localization. It is striking that this localization occurs prior to any known anterior determinant, indicating *Wolbachia* may associate with an as yet undiscovered host factor concentrated at the anterior. That this localization is conserved suggests that it has a functionally significant, but currently unknown role in *Wolbachia* transmission. While anteriorly localized, the amount of *Wolbachia* dramatically increases. Whether *Wolbachia* replication, transport from the nurse cells, or both are responsible for this increase is unknown. The anterior location may be a rich source of membrane for *Wolbachia* replication. The anterior localization also results in a high concentration of *Wolbachia* that closely associated with, and perhaps influences, the oocyte nucleus.

To our surprise, we discovered that immediately after the release of *w*Mel *Wolbachia* from its anterior position in the oocyte (Stage 6), the amount of *Wolbachia* in the oocyte dramatically decreases (see **Figure 1, Stage 6** and **Supplemental Figure 1**). This suggests that the *Wolbachia* are transported back into the nurse cell complex. The functional significance of this clearing of *Wolbachia* from the oocyte is unknown. The retreat of *Wolbachia* occurs during the time when the anteriorly positioned oocyte and neighboring follicle cells signal one another in order to establish the dorsal ventral axis (Merkle et al., 2020). Perhaps *Wolbachia* exits the oocyte in order to not disrupt this process. Given the orientation of microtubules at this stage, it is likely *Wolbachia* rapidly exits the oocyte through an association with the plus-end directed motor protein Kinesin.

Following microtubule reorientation and the return of *Wolbachia* to the oocyte, *Wolbachia* spreads posteriorly throughout the entire oocyte relying on the host motor protein Kinesin (Serbus and Sullivan, 2007). All of the *Wolbachia* strains examined are distributed toward the posterior regions suggesting the interaction with Kinesin is also conserved. How *Wolbachia* engages these motor proteins is unknown. Surprisingly, none of the known host Kinesin linker proteins are utilized by *Wolbachia,* suggesting *Wolbachia* may interact directly with Kinesin (Russell et al., 2018).

It is after this stage in which *Wolbachia* is distributed throughout the oocyte, that we observe variability in the 19 *Wolbachia* strains with each falling into one of three distinct classes: two distinct classes in which *Wolbachia* concentrate at the posterior (Posteriorly Localized and Posteriorly Clumped) and one class in which *Wolbachia* fail to exhibit a posterior concentration (Dispersed). The former two classes are distinguished by whether the vast majority (Posteriorly Clumped) or only a small fraction of the *Wolbachia* localize to the posterior pole (Posteriorly Localized).

The Posteriorly Localized class, in which only a small fraction localize to the posterior pole, can be explained by previous work demonstrating that *w*Mel is a weak competitor for Kinesin; both *Wolbachia* and germplasm components rely on Kinesin for transport to the posterior pole (Russell et al., 2018). It is thought that *Wolbachia* has evolved to be a weak competitor because interference with the transport of essential germplasm components and thus germplasm formation may be disadvantageous for *Wolbachia* proliferation. The other posterior class (Posteriorly Clumped) is strikingly similar to that observed for *w*Mel in *D. melanogaster* in which the plus-end motor protein Kinesin is over-expressed and excess amounts of *Wolbachia* are transported to the posterior pole (Russell et al., 2018). It may be that the Posteriorly Clumped strains occur in host species with naturally much higher levels of host Kinesin or that these specific *Wolbachia* strains are much better competitors for host Kinesin. The latter is supported by our phylogenetic analyses, but

functional work will be required to formally test this hypothesis. In particular, it would be interesting to investigate whether there are instances in which *Wolbachia* disrupts the A-P axis (anterior-posterior) because of its accumulation at the posterior pole.

The posterior pole concentration among *Wolbachia* strains may also differ due to variability in the ability of *Wolbachia* to stably associate with the pole plasm. Previous studies demonstrated that *Wolbachia* posterior pole concentration requires intact pole plasm (Serbus and Sullivan, 2007). For example, in *oskar* mutations, a key pole plasm determinant, *w*Mel *Wolbachia* fails to accumulate at the posterior region. As described, the oocyte experiences tremendous cytoplasmic streaming requiring posterior components to be anchored directly or indirectly to the cortex (Quinlan, 2016). This variability could be due to differences in the ability of *Wolbachia* to compete with other pole plasm components such as mitochondria for binding sites. Alternatively, the host species may vary in the extent to which their pole plasm accommodates *Wolbachia*. These alternatives can be explored through trans-infection studies.

The overlapping, but distinct, *Wolbachia* oocyte localization patterns suggest corresponding overlapping and distinct interactions with host factors among the three *Wolbachia* classes of posterior localization. This variation in the host proteins with which *Wolbachia* engages, is in contrast to other endosymbionts such as *Listeria*. *Listeria* relies on a surface protein, which binds and polymerizes host actin, propelling it both within and between host cells (Kühn and Enninga, 2020). Because the interaction between ActA and actin is essential for cell-to-cell transmission, natural variants of this interaction and transmission strategy have not been discovered. Thus, the variation in *Wolbachia* oocyte localization suggests variation in the host protein interactions facilitates multiple strategies of *Wolbachia* vertical transmission.

Two classes in which *Wolbachia* concentrates in the posterior pole (Posteriorly Localized and Posteriorly Clumped) provide a ready explanation for the cellular mechanisms by which it is vertically transmitted through generations. In every generation, *Wolbachia* targets the site of germline formation in the developing oocyte. This also explains why strains exhibiting this pattern, such as *w*Mel, exhibit efficient transmission strategies under typical laboratory conditions (Hague et al., 2022). The third, Distributed, class that exhibits no-to-low *Wolbachia* in the posterior germplasm of the mature oocyte is puzzling. This would be expected to result in embryos lacking *Wolbachia* in the germline. However, within hours after fertilization, *Wolbachia* is clearly present in the germlines of both sexes (**Figure 6**). This implies that in these strains *Wolbachia* are capable of invading the germline from neighboring somatic cells. In filarial nematodes, *Wolbachia* germline invasion from the soma has been directly observed (Landmann et al., 2012). While germline invasion via cell-to-cell transmission has not been directly observed in insects, a number of lines of evidence indicate that it likely occurs. In *Drosophila* cell culture, *Wolbachia* efficiently undergoes cell-to-cell transmission (White et al., 2017; White et al., 2017). *Wolbachia* injected into the abdomen of adult *Drosophila* females migrate to and occupy the germline and follicle stem cells (Frydman et al., 2006). Collections directly from nature reveal some strains with developing oocytes lacking *Wolbachia* (Casper-Lindley et al., 2011). In these strains, all of the mature oocytes were infected, suggesting an alternate route of infection via neighboring somatic cells.

As previously hypothesized, given the posterior follicle cells are directly adjacent to the pole plasm, we suspect *Wolbachia* present in the latter are the source of germline *Wolbachia* (Kamath

et al., 2018). As found in female filarial nematodes, this requires that *Wolbachia* undergo cell-to-cell transmission (Landmann et al., 2012). Our results are in accord with and complement previous studies demonstrating *Wolbachia* concentrated in the posterior polar follicle cells in a number of *Wolbachia*-infected species (Kamath et al., 2018). For example, in *w*Mau-infected *D. mauritiana*, a Dispersed class, there is a striking concentration of *Wolbachia* in the posterior follicle pole cells directly adjacent to the pole plasm. Thus, *Wolbachia* in these somatic cells are well positioned to invade the pole plasm becoming incorporated into the germline of the next generation (**Figures 7 and 8**). In contrast, in the Posteriorly Localized and Clumped strains, we found no to few *Wolbachia* in the posterior polar follicle cells. Among the *Wolbachia* strains examined, *w*Mel stands out as an exception. Not only is it concentrated at the posterior pole of the oocyte, but there is also a large *Wolbachia* concentration in the posterior polar follicle cells. Thus, *w*Mel may maintain both robust germline-to-germline and soma-to-germline modes of transmission. Accordingly, it has an extremely high efficiency of vertical transmission under laboratory conditions (but see Hague et al. 2022). Thus, *Wolbachia* may be a rarity among endosymbionts, exhibiting two distinct modes of vertical transmission (**Figure 8**).

Surprisingly, literature surveys reveal that germline invasion of endosymbionts from the soma every generation is the most common form of vertical inheritance (Russell et al., 2019). Vertical transmission via a continuous presence in the germline is much less common. One explanation is that hosts maintains mechanisms preventing endosymbiont occupation during the formative stages of germline development. In instances in which the endosymbiont is maintained in the germline, this may evolve into an obligate relationship in which development of the germline depends on the presence of the endosymbiont (Sullivan, 2017). Examples of this include the leafhopper (*E. plebeus*), in which the endosymbiont is required for normal embryonic development (Sander, 1968). In addition, removal of *Wolbachia* from some insects results in increased apoptosis and abnormal oocyte development (Dedeine et al., 2001; Pannebakker et al., 2007).

Examining a diversity of *Wolbachia* strains and host species enabled us to test whether *Wolbachia* oocyte abundance and localization are correlated with the evolutionary relationships of the *Wolbachia* strains or host species. This analysis revealed a strong correlation between *Wolbachia* posterior abundance and *Wolbachia* phylogenetic relationships, suggesting that factors intrinsic to *Wolbachia* contribute to posterior localization patterns. Generally, we find that closely related *w*Mel-like *Wolbachia* occur at higher abundance at the oocyte posterior, whereas *w*Ri-like *Wolbachia* exhibit a more dispersed distribution (**Figure 4**). This conclusion is in accord with previous *Wolbachia* transplantation studies demonstrating that posterior localization is determined by *Wolbachia* rather than host factors (Veneti et al., 2004; Serbus and Sullivan, 2007). It is likely that the composition of *Wolbachia* surface proteins play a major role in the extent of posterior localization.

In an effort to identify specific *Wolbachia* factors promoting posterior localization, we performed the same phylogenetic analysis using gene trees of the six predicted *Wolbachia* surface proteins. This analysis revealed that two candidate surface proteins, particularly WD_1085, exhibited a strong phylogenetic signal associated with *Wolbachia* localization patterns. While we highlight these loci as potential candidates, we note that our results should be interpreted with caution since the surface protein WD_1085 could be in linkage with causal loci. Nonetheless, *Wolbachia* surface proteins are generally considered to be strong candidates for interactions with host cells (Werren et al., 2008; Baldridge et al., 2016; Hague et al., 2022).

We did not find that *Wolbachia* localization patterns exhibit a significant phylogenetic signal on the host phylogeny, suggesting that host-associated factors are less important in determining *Wolbachia* localization patterns. Interestingly, this result contrasts with the transplantation studies described above that indicate hosts may also play a significant role in determining *Wolbachia* abundance (Veneti et al., 2004). In addition, genome-wide screens reveal host factors play a major role in determining *Wolbachia* intracellular abundance (White et al., 2017; Grobler et al., 2018). Finally, nutrients and other environmental factors influence *Wolbachia* abundance in the oocyte and maternal transmission rates for closely related *w*Mel-like *Wolbachia* (Serbus et al., 2015; Hague et al., 2020a; Hague et al., 2022).

MATERIALS AND METHODS

Drosophila **Stocks**

All stocks were grown on standard brown food (Sullivan et al., 2000) at 25°C with a 12h light/dark cycle. Uninfected *Drosophila* stocks were generated by tetracycline-curing of the infected stock (Serbus et al., 2015). Listed in Table 1 are the 19 different *Wolbachia* strains we examined, which infect 18 different *Drosophila* host species. This included six *w*Mel-like strains (Cooper et al., 2019; Hague et al., 2020a), five *w*Ri-like strains (Turelli et al., 2018), seven other A-group *Wolbachia*, and the B-group strain *w*Mau that diverged from A-group *Wolbachia* up to 46 million years ago (Meany et al., 2019). Each genotype was generated as an iso-female line by sampling a single gravid female from the field and placing her individually into a vial.

Ovary, embryo and larva fixation and staining

Newly eclosed flies were transferred to fresh food for 3-5 days for aging. Approximately 10 females were dissected for each slide preparation. Ovaries were fixed using a modification of previously published procedures (Brendza et al., 2000; Russell et al., 2018). Ovaries were removed and separated in phosphate buffer solution and then fixed in a 200 μl devitellinizing solution (2% paraformaldehyde and 0.5% v/v NP40 in 1x PBS) mixed with 600 μl heptane for 20 min at room temperature, on a shaker. After removal of the organic layer by brief centrifugation for sample isolation, the ovaries were washed 3 times with PBS-T (0.1% Triton X-100 in 1x PBS) along with three five-minute washes. Samples were treated with RNAse A (10 mg/ml) and left overnight at room temperature. Samples were then washed again a few times with PBS-T and then stained with a dilute solution of phalloidin for actin staining on a rotator for 1 hour. Samples were washed again with 3 quick PBS-T rounds and then solution changes with PBS-T over two hours. 60uL of propidium iodide (PI) in mounting media was added to the sample after removal of the wash solution and left again overnight. Ovaries were then mounted and carefully separated out again for ease of imaging and removal of excess mature eggs. Slides were coated in nail polish and stored at -20°C until imaged.

Embryos were collected for 1-3 hours on plates and fixed in equal volume 32% paraformaldehyde and heptane and fixed as previously described (Sullivan et al., 2000). After extraction with methanol, embryos were blocked for one hour in 5% PBST-BSA (2.5g bovine serum albumin fraction V in 50ml PBS-T) at room temperature. Embryos were then incubated with anti-WD_0009 or anti-Ftfz at 1:500 overnight at 4°C. After three washes with 1% PBST-BSA over one hour, embryos were incubated in Alexa 488 Anti-Rabbit at 1:500 for one hour at room temperature.

Samples were then washed three more times over one hour with 1% PBST-BSA. Embryos were mounted with media containing DAPI and stored at -20°C until imaged.

Larval fat bodies were dissected and fixed from 3rd instar larvae using a modified version of a published protocol (Maimon and Gilboa, 2011). Fat bodies were dissected in 1X PBS then fixed in 5% paraformaldehyde in PBS for 20 minutes with gentle agitation. Fat bodies were washed for five minutes, 10 minutes, then 45 minutes with cold 1% NP40 in PBS. After washes, fat bodies were blocked in 5% PBST-BSA (2.5g bovine serum albumin fraction V in 50ml PBS-T) for one hour at room temperature. Fat bodies were then incubated with anti-FtsZ at 1:500 overnight on a rocker at 4°C, washed three times over one hour with 1% PBST-BSA, then incubated with Alexa 488 Anti-Rabbit at 1:500 at room temperature for one hour. After three additional washes over one hour with 1% PBST-BSA, fat bodies were mounted onto a slide with media containing DAPI. Larval ovaries were isolated from fat bodies using 0.1mm needles before sealing the slide with nail polish. Samples were then stored at -20°C.

Confocal microscopy

Imaging was performed on an inverted Leica DMI6000 SP5 scanning confocal microscope. Optical sections of a Nyquist value of 0.38 were used. A variety of zooms were used to optimize image viewing, with most being set at 1.5X. Propidium iodide was excited with the 514 and 543 nm lasers, and emission from 550 to 680 nm was collected. GFP was imaged with the 488 nm laser, and emission from 488 to 540 nm was collected. Alexa 633 was imaged with the 633 laser, and emission from 606 to 700 nm was collected. All imaging was performed at room temperature. Images were acquired with Leica Application Suite Advanced Fluorescence software.

Wolbachia **quantification and analysis**

Following Russell et al. (2018), we used the polygon selection tool to select three different regions of the oocyte (see Fig. S1 in Russell et al. 2018): the whole oocyte, the posterior region, and the posterior cortex. The "area" and "integrated density" were measured for each region. Additionally, the average of a quadruplicate measure of "mean gray value" beside the stained oocytes was measured as background fluorescence. The Corrected Total Cell Fluorescence (CTCF) was calculated as integrated density – (area x background mean gray value) for each region.

All statistical analyses were performed in R. We first tested whether our three qualitative classes (Posteriorly Clumped, Posteriorly Localized, Dispersed) differed significantly in quantitative estimates of *Wolbachia* fluorescence (CTCF). We used a one-way ANOVA to test if the three groups differed based on their mean CTCF values from the whole oocyte, the posterior region, and the posterior cortex. Distribution and leverage analyses indicated that a transformation of $x' = \log(x +1)$ was needed to meet assumptions of normality. We tested for significance using an F test with type III sum of squares using the "Anova" function in the *car* package (Fox and Weisberg, 2019). We conducted Tukey multiple comparisons among the three groups using the "TukeyHSD" function.

Genomic data

In order to perform the phylogenomic analysis and generate phylograms, we obtained *Wolbachia* sequences from publicly available genome assemblies, which included *w*Mel (Wu et al., 2004),

*w*Ri (Klasson et al., 2009), *w*Ha (Ellegaard et al., 2013), *w*Aura, *w*Tria, *w*Pan (Turelli et al., 2018), *w*Ana (Salzberg et al., 2005), and *w*Rec (Metcalf et al., 2014). All other *Wolbachia* sequences included in this study (*w*Bai, *w*Bic, *w*Boc, *w*Burl, *w*Cha, *w*Curt [*w*Tsa], *w*Nik, *w*Seg, *w*Tris, *w*Leo) were obtained using Illumina sequencing and previously described methods (Hague et al., 2020b). We obtained publicly available host sequences for *D. melanogaster* (Hoskins et al., 2015), *D. simulans* (Hu et al., 2013), *D. ananassae* (Clark et al., 2007), *D. pandora* (Turelli et al., 2018), *D. mauritiana* (Meany et al., 2019), *D. auraria, D. triaurari, D. baimaii, D. bicornuta* (aff. *bicornuta*), *D. bocki*, *D. burlai*, *D. chauvacae* (cf. *chauvacae*), *D. curta* (aff. *tsacasi*), *D. nikananu*, *D. seguyi*, *D. tristis*, and *D. leontia* (Conner, 2021). A *D. recens* genome assembly was kindly provided by Kelly Dyer and Rob Unckless. We note that three of the *montium* host species listed above have had recent updates to their taxonomic nomenclature: *D. bicornuta* has been renamed to aff. *bicornuta*, *chauvacae* has been renamed to cf. *chauvacae*, and *curta* has been renamed to aff. *tsacasi*. See Conner et al. (2021) for a full discussion of *montium* species names and relationships.

Wolbachia phylogenetic analysis

Raw Illumina reads from the newly sequenced *Wolbachia* strains were trimmed using Sickle version 1.33 (Joshi and Fass, 2011) and assembled using ABySS version 2.0.2 (Jackman et al., 2017). *K* values of 71, 81, and 91 were used, and scaffolds with the best nucleotide BLAST matches to known *Wolbachia* sequences with *E*-values less than 10^{-10} were extracted as the draft *Wolbachia* assemblies. For each strain, we chose the assembly with the highest N50 and the fewest scaffolds (**Supplemental Table 2**). The newly assembled genomes and the previously published assemblies were annotated using Prokka version 1.11, which identifies homologs to known bacterial genes (Seemann, 2014). To avoid pseudogenes and paralogs, we only used genes present in a single copy with no alignment gaps in all of the genome sequences. Genes were identified as single copy if they uniquely matched a bacterial reference gene identified by Prokka. By requiring all homologs to have identical length in all of the *Wolbachia* genomes, we removed all loci with indels. A total of 66 genes totaling 43,275 bp met these criteria. We then estimated a Bayesian phylogram using RevBayes 1.0.8 under the GTR + Γ + I model partitioned by codon position (Höhna et al., 2016). Four independent runs were performed, which all converged on the same topology. All nodes were supported with Bayesian posterior probabilities of 1.

Similar methods were used to generate gene trees for candidate *Wolbachia* loci putatively involved in host interactions during oogenesis. This included the three *wsp* paralogs (WD_1063 [*wsp*/RS04815], WD_0009 [*wspB*/RS00060], WD_0489 [*wspC*/RS06475]) and three other *Wolbachia* surface proteins (WD_1085 [bamA/RS04910], WD_0501 [p44/MSP2/RS0225], WD_1041 [peptidase M2/RS04710]). Here, the WD_XXXX locus tags refer to the *w*Mel genome assembly (Wu et al., 2004).The protein-coding sequences for each locus were extracted from the assemblies using BLAST and the *w*Mel reference sequences. Notably, we found evidence of insertions in the WD_0009 (*wspB*) sequences of *w*Boc, *w*Leo, and *w*Bic (see Results). We removed these large insertions from the alignment to generate the *wspB* gene tree. In addition, BLAST identified WD_0489 (*wspC*) in *w*Mel-like *Wolbachia* and closely related *w*Bai, *w*Leo, and *w*Boc, but yielded no hits in the other *Wolbachia* strains (*w*Nik, *w*Tsa, *w*Pan, *w*Ana, *w*Tria, *w*Aura, *w*Ri, *w*Mau), indicating that *wspC* is either too diverged from the *w*Mel reference for BLAST recognition or alternatively *wspC* is only present in *w*Mel-related *Wolbachia* strains. Sequences

for each gene were aligned with MAFFT 7 (Katoh and Standley, 2013). We then used RevBayes and the GTR + Γ + I model partitioned by codon position to generate gene trees for each locus.

Host phylogenetic analysis

Host phylogenies were generated using the same nuclear genes implemented in Turelli et al. (2018): *aconitase, aldolase, bicoid, ebony, enolase, esc, g6pdh, glyp, glys, ninaE, pepck, pgi, pgm, pic, ptc, tpi, transaldolase, white, wingless,* and *yellow*. We used BLAST with the *D. melanogaster* coding sequences to extract orthologs from the genomes of each host species. Sequences were then aligned with MAFFT 7. Finally, we used RevBayes and the GTR + Γ + I model partitioned by codon position and gene to accommodate potential variation in the substitution process among genes, as described in Turelli et al. (2018). Our initial phylogenetic analysis produced a polytomy in the *melanogaster* species group regarding the relationships among the *montium*, *melanogaster*, and *ananassae* subgroups (**Supplemental Figure 10**). These relationships have been resolved in previous genomic studies involving additional taxa (Turelli et al., 2018; Suvorov et al., 2022); therefore, we generated a constrained host tree in RevBayes that enforced previously defined relationships among the three subgroups: ((*montium*, *melanogaster*), *ananassae*).

Tests for phylogenetic signal

The resulting phylograms were used to test whether cellular *Wolbachia* abundance in oocytes exhibits phylogenetic signal on either the *Wolbachia* or host phylogenies. We used our quantitative estimates of *Wolbachia* abundance to test for phylogenetic signal using Pagel's lambda (λ) (Pagel, 1999). Here, we used total *Wolbachia* fluorescence in whole oocytes, the posterior region, and the posterior cortex (log-transformed CTCF) as continuous characters to calculate maximum likelihood values of Pagel's λ. A Pagel's λ of 0 indicates that character evolution occurs independently of phylogenetic relationships, whereas λ = 1 is consistent with a Brownian motion model of character evolution. We used the "fitContinuous" function in GEIGER (Harmon et al., 2008) and a likelihood ratio test to compare our fitted value to a model assuming no phylogenetic signal (λ = 0). We also used a Monte Carlo-based method to generate 95% confidence intervals surrounding our estimates using 1,000 bootstrap replicates in the *pmc* package (Boettiger et al., 2012). When applicable, we conducted additional analyses to evaluate whether the number of taxa in our phylogeny (*N* = 19 *Wolbachia* strains, *N* = 18 host species) limited our ability to detect significant departures from λ = 0 (Hague et al., 2020b; Hague et al., 2021). Small phylogenies are likely to generate near-zero values simply by chance, not necessarily because the phylogeny is unimportant for trait evolution (Boettiger et al., 2012). To evaluate whether larger phylogenies increase the accuracy of estimation, we simulated trees with an increasing number of *Wolbachia* strains/host species (*N* = 25, 50) and our empirical estimates using the "sim.bdtree" and "sim.char" functions in the *geiger* R package (Harmon et al., 2008). We then re-estimated confidence intervals using the larger simulated trees.

Our tests for phylogenetic signal using the constrained host phylogram are presented in the main text; however, we note that our analysis using the unconstrained host tree (with a polytomy at the base of the *melanogaster* species group) produced similar results. Using the unconstrained tree, we found that *Wolbachia* abundance in whole oocytes (λ = 0.782 [0, 0.987], *P* = 0.469), the posterior region (λ = 0.951 [0, 1], *P* = 0.377), and the posterior cortex (λ < 0.001 [0, 0.563], *P* = 1) did not exhibit significant phylogenetic signal.

Ectopic expression of *Wolbachia* surface proteins in yeast

SC-Leucine yeast media was prepared as described previously (Rose et al., 1990). Where present, benomyl was added to media from a 10mg/ml stock in DMSO to a final concentration of 2.5 ug/ml. The yeast expression assays are a modified version of the protocol described by Sheehan et al. (2016). *Wolbachia* genes were codon optimized for expression in *S. cerevisiae* and synthesized by Twist Bioscience (South San Francisco). The *wsp*, *wspB* and *wspC* genes were gifts of the Free Genes project (https://stanford.freegenes.org/). These genes were inserted into pGH448 using the New England Biolabs Golden Gate kit according to the manufacturer's instruction. This placed the *Wolbachia* genes under control of the *GAL1* promoter and *CYC1* transcriptional terminator. This plasmid, which will be described in more detail elsewhere, carries a CEN/ARS element and also contains the kanamycin resistance and *LEU2* genes for selection in bacteria and yeast respectively.

The indicated plasmids were transformed into yeast strain GHY1934 [genotype] and GHY1551 [genotype]. Both of these strains are isogenic to S288C and are *GAL2+*. Transformed cells were grown to saturation in SC-Leucine liquid media containing 2% glucose. These cells were counted, diluted to $1x10^{-7}$ cells per ml, five-fold serial diluted and spotted to SC-Leucine plates with either 2% glucose or 2% galactose as the carbon source and incubated at the times and temperatures indicated in the figure legends.

ACKNOWLEDGEMENTS

We thank Dr. Benjamin Abrams (UCSC Life Sciences Microscopy Center, RRID: SCR_021135) for his technical support and assistance with microscopy experiments. We thank Dr. Shelbi Russell for their helpful advice on imaging protocols and analysis. We thank Dr. Irene Newton for kindly supplying the anti-FtsZ antibody. We thank Will Conner for guidance with bioinformatic analyses and Tim Wheeler for laboratory assistance. We thank Drew Endy and the Free Genes project for the gift of *Wolbachia* genes and advice on construction of pGH448. Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM139595 to W. Sullivan and R35GM124701 to B.S. Cooper. This work was also supported by an National Science Foundation (NSF) grant (1456535) to W. Sullivan and an NSF CAREER Award (2145195) to B.S. Cooper. We thank the Big Creek Reserve for use of their facilities. The microscopic analysis was enabled through the generous support of the IBSC microscopy facility at UC Santa Cruz.

FIGURES

Figure 1. Confocal micrographs of *D. melanogaster* (*w*Mel) oocytes tracked through early oogenesis. Samples are stained with propidium iodide (PI) for DNA (red) and phalloidin for actin (green). Representative stages of development (1-10) are shown for *w*Mel infected (left panels) and uninfected (right panels). Grayscale zoomed-in images around the nucleus of the oocyte are shown in the third column, with arrows highlighting the *Wolbachia* in the oocyte. Orange arrow highlights *Wolbachia* in the nurse cells. Scale bars all set to 25 uM.

Figure 2. Conserved *Wolbachia* anterior localization during early oogenesis (stages 4-5). *Wolbachia* and host DNA stained with PI (red) and actin with phalloidin (green) shown in the left most column. Greyscale of PI channel shown in middle. 4X zoom-in around the oocyte shown on the right. All of the eight *Wolbachia* strains examined concentrate at the anterior cortex of the oocyte at this stage. Red arrows highlight *Wolbachia*'s anterior localization positioning for (A) *w*Pan in *D. pandora*, (B) *w*Nik in *D. nikananu*, (C) *w*Cha in *D. chauvacae*, (D) *w*Ha in *D. simulans*, (E) *w*Tsa in D. *tsacasi*, (F) *w*Boc in *D. bocki*, (G) *w*Ana in *D. ananassae*, and (H) *w*Mel in *D. melanogaster*. Oocytes are approximately 50-75 uM in size.

Figure 3. Three distinct patterns of posterior *Wolbachia* localization during late stage oogenesis (A-C). Confocal micrographs of *Drosophila* oocytes DNA-stained with PI (red) and actin-stained with phalloidin (green) show representative examples of (A) Posteriorly Localized *w*Tris in *D. tristis*, (B) Posteriorly Clumped *w*Bic in *D. bicornuta*, and (C) Dispersed localization of *w*Tria in *D. triauraria*. Second row depicts single channel images of PI staining. Third row depicts an enlarged PI-stained image of the posterior region of each oocyte. Scale bars set at 25uM.

Figure 4. (A) Estimated Bayesian phylogram for the 19 A- and B-group *Wolbachia* strains included in tests for phylogenetic signal. (B) Mean estimates of *Wolbachia* abundance (log-transformed CTCF) in the whole oocyte, posterior region, and the posterior cortex. *Wolbachia* abundance in the posterior region has significant phylogenetic signal ($\lambda = 0.982$ [0, 0.998], $P = 0.021$), suggesting *Wolbachia*-associated factors determine *Wolbachia* abundance in the oocyte posterior. (C) Estimated Bayesian phylogram of *Drosophila* host species using 20 conserved single-copy genes. *Wolbachia* and host divergence times in millions of years (MYA) are reproduced from Meany et al. (2019) and Suvorov et al. (2022), respectively. The *Wolbachia* and host tree are generally discordant (see **Supplemental Figure 2**), as expected with frequent horizontal *Wolbachia* acquisition (Turelli et al. 2018).

Figure 5. *Wolbachia*'s presence around the 3rd instar larval germline, in multiple *Drosophila* hosts. *w*Mel *Wolbachia* stained with anti-FtsZ can be seen for *D. melanogaster*, in panel (A) shown in pink. Cells are marked with DAPI (blue). (A') Grayscale channel of anti-FtsZ staining. (B) Uninfected ovary for *D. melanogaster*, (B') grayscale channel for uninfected antibody binding. Germline location marked with anti-vasa shown in green (C), grayscale channel (C'). *w*Mau *Wolbachia* localization in the ovary for *D. mauritiana* shown in panel (D), grayscale channel (D'). *w*Ri and *D. simulans* shown in panel (E), grayscale (E'). Panel (F), *w*Bic and *D. bicornuta*, (F') grayscale channel. Scale bar shown at 25 uM.

Figure 6. (A) 1 hour *D. mauritiana* (*w*Mau) embryo stained with PI. (A') Greyscale of PI channel. (B) staining of 2-3 hour developed *D. mauritiana* (*w*Mau) embryo with PI. (B') PI of greyscale channel. (C) PI and anti-vasa (green) staining of *w*Mel *Wolbachia* in 2-3 hour *D. melanogaster* embryo. (C') Greyscale of PI channel. (D) Anti-FtsZ/DAPI staining for *w*Ri *Wolbachia* in *D. simulans*. (D') Greyscale of anti-FtsZ channel. Scale bars set at 25uM.

Figure 7. (A) *Wolbachia* quantification in the polar follicle cells during oogenesis stages 9-10. *Wolbachia* were quantified in midsections of each oocyte that displays unique posterior localization patterns: Dispersed, Posteriorly Localized, and Posteriorly Clumped. For each strain, the count is shown for the number of oocytes observed with *Wolbachia* present in both polar follicle cells (orange), only one polar cell (blue), and neither polar cell (grey). (B) Counts of oocytes observed with *Wolbachia* present anywhere in the follicle cells (pink) or with *Wolbachia* absent (grey). (C) Representative anti-FtsZ staining of *w*Mau in *D. mauritiana* (Dispersed class) present in both polar follicle cells (corresponding to orange labels above). (D) Representative staining of *w*Mel in *D. melanogaster* (Posteriorly Localized class) present in only one polar cell (blue labels above). (E) Representative staining of *w*Bic in *D. bicornuta* (Posteriorly Clumped class) absent in both polar cells (grey labels above). (F) Representative staining of *w*Mel in *D. melanogaster* (Posteriorly Localized class) broadly present in anterior follicle cells (pink labeling above). (G-H) Greyscale anti-FtsZ channel of respective composite images above. (K-N) 5X zoom of polar cells, with circles highlighting representative staining in the area. Scale bar set at 25uM. (O) Schematic of categories used for description of follicle cell *Wolbachia*. Dispersed localization pattern shown, both polar cells infected, with *Wolbachia* broadly present in follicle cells. (P) Posteriorly Localized pattern shown, one polar cell infected, with *Wolbachia* broadly present in follicle cells. (Q) Posteriorly Clumped pattern shown, with no *Wolbachia* present in the polar cells or other follicle cells.

Figure 8. Multiple routes of *Wolbachia* vertical transmission. (A) In *Wolbachia* strains that concentrate at the posterior pole plasm, vertical transmission likely occurs through a continuous maintenance in the germline. (B) In strains where no or few *Wolbachia* localize to the posterior pole plasm, vertical transmission likely occurs through *Wolbachia* invading the germline from neighboring somatic cells**.** **(**C) In theory, vertical transmission could occur through a symbiont invading the oocyte from neighboring somatic cells never associating with the germline.

TABLES

Table 1. Qualitative and quantitative quantification of cellular *Wolbachia* abundance in stage 10 oocytes. The number of confocal images (*N*) used to generate means is shown for each *Wolbachia*-infected host species.

Wolbachia strain	Host species	Localization class	*N*	Whole oocyte mean CTCF	Posterior mean CTCF	Posterior cortex mean CTCF
*w*Rec	*D. recens*	Clumped	3	482174	300753	58453
*w*Tris	*D. tristis*	Localized	4	223048	129013	105641
*w*Seg	*D. seguyi*	Dispersed	5	58552	11402	557
*w*Cha	*D. chauvacae*	Localized	5	108588	42636	9786
*w*Burl	*D. burlai*	Localized	4	72806	53754	36944
*w*Mel	*D. melanogaster*	Localized	2	191451	136174	92341
*w*Bic	*D. bicornuta*	Clumped	3	90114	83621	54900
*w*Ri	*D. simulans*	Dispersed	5	333505	90664	20494
*w*Pan	*D. pandora*	Localized	3	45423	6486	1890
*w*Ana	*D. ananassae*	Localized	4	41578	16843	12927
*w*Tria	*D. triauraria*	Dispersed	2	487897	1809	345
*w*Aura	*D. auraria*	Dispersed	5	266478	3699	645
*w*Ha	*D. simulans*	Dispersed	5	82941	3488	807
*w*Bai	*D. baimaii*	Dispersed	2	11359	0	0
*w*Boc	*D. bocki*	Clumped	6	137664	98339	82408
*w*Leo	*D. leontia*	Clumped	4	997045	685260	713918
*w*Tsa	*D. tsacasi*	Localized	4	61314	36478	20620
*w*Nik	*D. nikananu*	Localized	3	44764	27099	13014
*w*Mau	*D. mauritiana*	Dispersed	3	69362	799	322

SUPPLEMENTAL INFORMATION

Supplemental Figure 1. (A) Representative image of the reduction of *w*Mel *Wolbachia* titer between stages 4/5 and 6 in *D. melanogaster* oogenesis. *Wolbachia* stained with anti-FtsZ shown in pink. Scale bars set at 25 uM. (B) Greyscale DAPI channel. (C) Greyscale of anti-FtsZ channel. (A'-C') 5X zoom around the nucleus of the oocyte during stage 4 shown for each channel. (A''-C'') 4X zoom around the nucleus during stage 6 for each respective channel seen above. At least 10 oocytes across 3 slides were imaged with Z-stacks for a visual description of this trend.

Supplemental Figure 2. Bayesian phylograms depicting the evolutionary relationships among *Wolbachia* strains diverged up to 46 million years (left) and *Drosophila* host species diverged up to 50 million years (right). *w*Mel- and *w*Ri-like clades of closely related *Wolbachia* are labeled on the left. Grey lines pair *Wolbachia* strains with their *Drosophila* host species, which highlights patterns of topological discordance due to introgressive and horizontal transfer of *Wolbachia* among host species. The *Wolbachia* phylogram was estimated using 66 full-length and single-copy genes (43,275 bp) of equal length. All nodes have posterior probabilities >0.95. The host phylogram was estimated using 20 conserved single-copy genes (see Materials and Methods regarding host tree topology). *Wolbachia* and host divergence times in millions of years (MYA) are reproduced from Meany et al. (2019) and Suvorov et al. (2022), respectively.

Supplemental Figure 3. Representative images of *Wolbachia* posterior localization patterns. Confocal micrographs of *Drosophila* oocytes DNA-stained with PI (red) and actin-stained with phalloidin (green) show representative examples of infected *Drosophila* species with different *Wolbachia* localization classes. Second columns depict single greyscale channel images of PI staining. Third columns depict an enlarged PI-stained image of the posterior region of each oocyte. Panels A, B and C are grouped into *Wolbachia* strains that exhibit a Posteriorly Localized, Posteriorly Clumped, or Dispersed localization pattern respectively. Scale bar set at 25 uM.

A

B

C

Supplemental Figure 4. Cellular *Wolbachia* abundance in stage 10 oocytes measured as *Wolbachia* fluorescence due to propidium iodide (CTCF). Points are color coded by *Wolbachia* localization class (Posteriorly Clumped, Posteriorly Localized, Dispersed), with unique shapes indicating each *Wolbachia* strain and host species. Asterisks indicate significant differences among groups based on one-way ANOVAs and Tukey comparisons using *P*-values <0.05 adjusted for multiple comparisons. Above, a schematic is shown for fluorescent quantification of different regions of the oocyte using the PI greyscale channel: the whole oocyte, the posterior region, and the posterior cortex.

Supplemental Figure 5. Distribution of maximum likelihood estimates of λ from 1,000 bootstrap replicates based on the *Wolbachia* phylogram and *Wolbachia* abundance at the oocyte posterior cortex (log-transformed CTCF). The bootstrap analysis of our *Wolbachia* phylogram is shown to the left ($N = 19$ *Wolbachia* strains). To the right are simulated phylogenies with an increasing number of *Wolbachia* strains included ($N = 25, 50$). For simulated trees, character evolution was simulated with our λ estimate of 0.959 using the "sim.bdtree" and "sim.char" functions in the *Geiger* R package (Harmon et al. 2008). For each graph, fitted λ values for the original phylogeny are shown above with a vertical dashed line. Note that fitted λ values for the simulated phylogenies differ slightly from $\lambda = 0.959$, because "sim.char" uses a Brownian-motion model to simulate character evolution along the phylogeny. Below each graph, the mean estimate of λ from the 1,000 replicates ($\bar{\lambda}$) is shown with associated 95% confidence intervals. Small phylogenies (e.g., $N = 19$) are likely to generate many near-zero λ values by chance, not necessarily because the phylogeny is unimportant for trait evolution (Boettiger et al. 2012). As the number of strains in our analysis increases ($N = 25, 50$), bootstrapped estimates of λ cluster around the true λ value fitted to the original phylogeny.

Supplemental Figure 6. Gene trees of *Wolbachia* surface proteins. All nodes have Bayesian posterior probabilities of >0.95 unless otherwise noted (see only *wspC*).

Supplemental Figure 7. To gain insight into WD_0501 (RS0225/p44/Msp2) and WD_1085 (RS04910/bamA) cellular function, we ectopically expressed these genes in yeast using a Galactose inducible promoter. Under normal growth conditions, ectopic expression of both genes inhibited growth, with WD_1085 (RS04910/bamA) exhibiting a more pronounced inhibition. Growth inhibition of WD_1085 was dramatically increased when the integrity of microtubules was compromised with the yeast placed in the mad1 spindle assembly checkpoint background or the microtubules were compromised directly through the addition of benomyl. These results suggest that WD_1085 may directly or indirectly interact with host microtubules.

Supplemental Figure 8. Full amino acid alignment of *wspB* (WD_0009) homologs in *w*Mel and the *Wolbachia* strains with a putatively pseudogenized version of the surface protein. Stop codons are indicated by asterisks and insertions of varying length are indicated by "XXX".

```
wMel reference    MISKKTLAVTAFALLLSQQSFASETEGFYFGSGYYGQYLNNTSVLKTSTTGIKNLSINDR    60
wHa               MISKKTLAVTAFALLLSQQSFASETEGFYFGSRYYGQYLNNTSVLKTSTTGIKNLSINDR    60
wBai              MISKKTLAVTALALLLSQQSFASETEGFYFGSGYYGQFFNGMGELKTSTTDIKNLSINDR    60
wBoc              MISKKTLAVTALALLLSQQSFASETEGFYFGSGYYGQFFNGMGELKTSTTDIKNLSINDR    60
wLeo              MISKKTLAVTALALLLSQQSFASETEGFYFGSGYYGQFFNGMGELKTSTTDIKNLSINDR    60
wBic              MISKKTLAVTAFALLLSQQSFASETEGFYFGSGYYGQYLNNTSVLKTSTTGIKNLSINDR    60

wMel reference    GAQNTEGQSLSEYKGDYNPPFAANVAFGYTGELGNNSYRAELEGMYSSVKVD---NIGLT   117
wHa               GAQNTEGQSLSEYKGDYNPPLLQMWHLVTQENWVTTAIGLNWKGCILL*KWI---ILV*Q   117
wBai              DAQNTKGQSLSEYKGDYNPPL------VIQGNWVTTAIGLNWKGCILL*KWI---ILV*Q   111
wBoc              GAQNTKGQSLSKYKGDYNPPFAANVALGYTGELGNNSCRAELEGMYSSVKVDXXXILV*Q   120
wLeo              GAQNTKGQSLSKYKGDYNPPFAANVALGYTGELGNNSCRAELEGMYSSVKVDXXXILV*Q   120
wBic              GAQNTEGQSLSEYKGDYNPPFAANVAFGYTGELGNNSYRAELEGMYSSVKVD---NIGLT   117

wMel reference    SSQITVSYLKETGEDPDKETYLYSAAVSHDQIENISVMANVYHHWKSDRFSFSPYVGIGI   177
wHa               VAK*LFHT*RRLVRILIKKLISIVLQLVMTKLRTYL*WQMFIIIGKVTVSLFLLTLVLGS   177
wBai              VAK*LFHI*RTLVRMLIKKLISIVLQLIMTKLRTYL*WQMFIIIGKVTVSLFLLTLVLES   171
wBoc              VVK*LFHT*RTLVRMLVKKLVSIVLQLIMTKLKTCL*WQMFIIIGRVTVSLFLLMLGLES   180
wLeo              VVK*LFHT*RTLVRMLVKKLVSIVLQLIMTKLKTCL*WQMFIIIGRVTVSLFLLMLGLES   180
wBic              SSQITVSYLKETGEDPNKETYLYSAAVSHDQIENISVMANVYHNWKSDRFSFSPYVGIGI   177

wMel reference    GATRMTMFEKPSIRPAGQLKAGFDYRINEDVN---MHIGYRGFGAIGSDIKLTAKRLGQV   234
wHa               VQQE*RCLKNRQ*DPQVN*KLALTIA*TKM*I---CISDIEVLVLLVAILSLQQKG*DKW   234
wBai              VQQE*RCLKNRQ*DPQVK*KLALTIT*TKM*I---CISDIEVLVLLVAILSLQQKS*DKW   228
wBoc              VQQE*QCLKNRQ*DPQVN*KLALIIT*TKM*I---CISDIEVLVLLVVLLVMVLNLQRMR   237
wLeo              VQQE*QCLKNRQ*DPQVN*KLALIIT*TKM*I---CISDIEVLVLLVVLLVMVLNLQRMR   237
wBic              GATRMTMFEKPSIRPAGQLKAGFDYRINEDV*XXXCISDIEVLVLLVAILSLQQKG*DKW   237

wMel reference    VDDPNNDKKKK------LNPSSGSKVTEEINIGNQLFHTHGIEAGLTFHFASKA        282
wHa               *TTLIMIKKRS------LILAQVAK*LRK*I*VINYFIHTV*RLVLLSILPAKL        282
wBai              *TTLIMIKKRS------LILAQVAKYSISHNSPNRS*EGRINSTTKGFE*RNV-        275
wBoc              LEKL*MILITGIKRSLMIRVNQQMQQKKI*I*VIHRFTHTV*RLALLSTLPAKL        291
wLeo              LEKL*MILITGIKRSLMIRVNQQMQQKKI*I*VIHRFTHTV*RLALLSTLPAKL        291
wBic              *TTLIMIKKRS------LILAQVAKYSTSRNSPNRIRYVLMAMVEQ*HIMICSD        285
```

Supplemental Figure 9. Predicted *Wolbachia* surface proteins WD_1063 (*wsp*/RS04815), WD_0009 (*wspB*/RS00060), and WD_0489 (*wspC*/RS06475) were assayed for their effect on eukaryotic cellular processes through ectopic expression in yeast. Note that galactose induction of gene expression of *wspC* kills the cells, while expression of *wsp* and *wspB* inhibits growth.

Supplemental Figure 10. (A) Estimated Bayesian phylogram of host species based on 20 nuclear loci. All nodes have posterior probabilities >0.95. (B) Bayesian phylogram estimated using similar methods, but with a constrained topology to account for previously resolved relationships among subgroups within the *melanogaster* species group ((*montium*, *melanogaster*) *ananassae*) based on Turelli et al., (2018) and Suvorov et al., (2022). (C) Mean estimates of *Wolbachia* abundance (log-transformed CTCF) in the whole oocyte, posterior region, and the posterior cortex. For *D. simulans*, the mean values include both *w*Ri- and *w*Ha-infected individuals.

Supplemental Table 1. Estimates of phylogenetic signal (Pagel's λ) with 95% confidence intervals (CI) for *Wolbachia* surface proteins.

Locus	Whole Oocyte				Posterior				Posterior Cortex			
	λ	Lower CI	Upper CI	*P*	λ	Lower CI	Upper CI	*P*	λ	Lower CI	Upper CI	*P*
WD_1063 (*wsp*)	0.000	0.000	0.578	1.000	0.945	0.000	0.993	0.071	0.000	0.000	0.633	1.000
WD_0009 (*wspB*)	0.000	0.000	0.401	1.000	0.000	0.000	0.374	1.000	0.279	0.000	0.720	0.655
WD_0489 (*wspC*)	0.210	0.000	0.947	0.659	0.000	0.000	0.845	1.000	0.000	0.000	0.811	1.000
WD_0501	0.000	0.000	0.597	1.000	0.992	0.521	1.000	0.022	0.000	0.000	0.536	1.000
WD_1041	0.000	0.000	0.692	1.000	0.000	0.000	0.555	1.000	0.000	0.000	0.484	1.000
WD_1085	0.000	0.000	0.553	1.000	0.974	0.000	0.998	0.001	0.942	0.000	0.994	0.028

Supplemental Table 2. Scaffold count, N50, and total assembly size of each new *Wolbachia* assembly.

Host species	*Wolbachia*	Scaffold Count	N50	Total Assembly Size
D. baimaii	*w*Bia	209	9004	1135209
D. bicornuta	*w*Bic	140	16267	1192611
D. bocki	*w*Boc	227	7115	1090928
D. burlai	*w*Burl	128	20905	1248003
D. chauvacae	*w*Cha	406	4802	1300201
D. tsacasi	*w*Tsa	59	29597	1132109
D. nikananu	*w*Nik	83	22663	1257108
D. seguyi	*w*Seg	201	8880	1174927
D. tristis	*w*Tris	114	20171	1264866
D. leontia	*w*Leo	186	9149	1096851

Supplemental Table 3. Summary of results for *Wolbachia* surface proteins.

Gene	Significant Phylogenetic Signal	Ectopic Expression
WD_1063 (*wsp*/RS04815)	None	Growth inhibition
WD_0009 (*wspB*/RS00060)	None	Growth inhibition
WD_0489 (*wspC*/RS06475)	None	Lethality
WD_1085 (RS04910)	Oocyte posterior, posterior cortex	Growth inhibition and sensitive to microtubule checkpoints/inhibitors
WD_0501 (RS0225)	Oocyte posterior	Slight growth inhibition
WD_1041 (RS04710)	None	Not tested

REFERENCES

BALDRIDGE, G.D., LI, Y.G., WITTHUHN, B.A., HIGGINS, L., MARKOWSKI, T.W., BALDRIDGE, A.S., and FALLON, A.M., 2016, Mosaic composition of ribA and wspB genes flanking the virB8-D4 operon in the *Wolbachia* supergroup B-strain, *w*Str: Arch Microbiol, v. 198, p. 53-69, doi: 10.1007/s00203-015-1154-8.

BASTOCK, R., and ST JOHNSTON, D., 2008, *Drosophila* oogenesis: Curr Biol, v. 18, p. R1082-7, doi: 10.1016/j.cub.2008.09.011.

BILINSKI, S.M., JAGLARZ, M.K., and TWORZYDLO, W., 2017, The Pole (Germ) Plasm in Insect Oocytes: Results Probl Cell Differ, v. 63, p. 103-126, doi: 10.1007/978-3-319-60855-6_5.

BOETTIGER, C., COOP, G., and RALPH, P., 2012, Is your phylogeny informative? Measuring the power of comparative methods: Evolution, v. 66, p. 2240-51, doi: 10.1111/j.1558-5646.2011.01574.x.

BRENDZA, R.P., SERBUS, L.R., DUFFY, J.B., and SAXTON, W.M., 2000, A function for kinesin I in the posterior transport of oskar mRNA and Staufen protein: Science, v. 289, p. 2120-2, doi: 10.1126/science.289.5487.2120.

CARRINGTON, L.B., LIPKOWITZ, J.R., HOFFMANN, A.A., and TURELLI, M., 2011, A re-examination of *Wolbachia*-induced cytoplasmic incompatibility in California *Drosophila simulans*: PLoS One, v. 6, p. e22565, doi: 10.1371/journal.pone.0022565.

CASPER-LINDLEY, C., KIMURA, S., SAXTON, D.S., ESSAW, Y., SIMPSON, I., TAN, V., and SULLIVAN, W., 2011, Rapid fluorescence-based screening for *Wolbachia* endosymbionts in *Drosophila* germ line and somatic tissues: Appl Environ Microbiol, v. 77, p. 4788-94, doi: 10.1128/AEM.00215-11.

CHA, B.J., KOPPETSCH, B.S., and THEURKAUF, W.E., 2001, In vivo analysis of *Drosophila* bicoid mRNA localization reveals a novel microtubule-dependent axis specification pathway: Cell, v. 106, p. 35-46, doi: 10.1016/s0092-8674(01)00419-6.

CHA, B.J., SERBUS, L.R., KOPPETSCH, B.S., and THEURKAUF, W.E., 2002, Kinesin I-dependent cortical exclusion restricts pole plasm to the oocyte posterior: Nat Cell Biol, v. 4, p. 592-8, doi: 10.1038/ncb832.

CLARK, A.G., EISEN, M.B., SMITH, D.R., BERGMAN, C.M., OLIVER, B., MARKOW, T.A., KAUFMAN, T.C., KELLIS, M., GELBART, W., IYER, V.N., POLLARD, D.A., SACKTON, T.B., LARRACUENTE, A.M., SINGH, N.D., ABAD, J.P., ABT, D.N., ADRYAN, B., AGUADE, M., AKASHI, H., ANDERSON, W.W., AQUADRO, C.F., ARDELL, D.H., ARGUELLO, R., ARTIERI, C.G., BARBASH, D.A., BARKER, D., BARSANTI, P., BATTERHAM, P., BATZOGLOU, S., BEGUN, D., BHUTKAR, A., BLANCO, E., BOSAK, S.A., BRADLEY, R.K., BRAND, A.D., BRENT, M.R., BROOKS, A.N., BROWN, R.H., BUTLIN, R.K., CAGGESE, C., CALVI, B.R., BERNARDO DE CARVALHO, A., CASPI, A., CASTREZANA, S., CELNIKER, S.E., CHANG, J.L., CHAPPLE, C., CHATTERJI, S., CHINWALLA, A., CIVETTA, A., CLIFTON, S.W., COMERON, J.M., COSTELLO, J.C., COYNE, J.A., DAUB, J., DAVID, R.G., DELCHER, A.L., DELEHAUNTY, K., DO, C.B., EBLING, H., EDWARDS, K., EICKBUSH, T., EVANS, J.D., FILIPSKI, A., FINDEISS, S., FREYHULT, E., FULTON, L., FULTON, R., GARCIA, A.C., GARDINER, A., GARFIELD, D.A., GARVIN, B.E., GIBSON, G., GILBERT, D., GNERRE, S.,

GODFREY, J., GOOD, R., GOTEA, V., GRAVELY, B., GREENBERG, A.J., GRIFFITHS-JONES, S., GROSS, S., GUIGO, R., GUSTAFSON, E.A., HAERTY, W., HAHN, M.W., HALLIGAN, D.L., HALPERN, A.L., HALTER, G.M., HAN, M.V., HEGER, A., HILLIER, L., HINRICHS, A.S., HOLMES, I., HOSKINS, R.A., HUBISZ, M.J., HULTMARK, D., HUNTLEY, M.A., JAFFE, D.B., JAGADEESHAN, S., JECK, W.R., JOHNSON, J., JONES, C.D., JORDAN, W.C., KARPEN, G.H., KATAOKA, E., KEIGHTLEY, P.D., KHERADPOUR, P., KIRKNESS, E.F., KOERICH, L.B., KRISTIANSEN, K., KUDRNA, D., KULATHINAL, R.J., KUMAR, S., KWOK, R., LANDER, E., LANGLEY, C.H., LAPOINT, R., LAZZARO, B.P., LEE, S.J., LEVESQUE, L., LI, R., LIN, C.F., LIN, M.F., LINDBLAD-TOH, K., LLOPART, A., LONG, M., LOW, L., LOZOVSKY, E., LU, J., LUO, M., MACHADO, C.A., MAKALOWSKI, W., MARZO, M., MATSUDA, M., MATZKIN, L., MCALLISTER, B., MCBRIDE, C.S., MCKERNAN, B., MCKERNAN, K., MENDEZ-LAGO, M., MINX, P., MOLLENHAUER, M.U., MONTOOTH, K., MOUNT, S.M., MU, X., MYERS, E., NEGRE, B., NEWFELD, S., NIELSEN, R., NOOR, M.A., O'GRADY, P., PACHTER, L., PAPACEIT, M., PARISI, M.J., PARISI, M., PARTS, L., PEDERSEN, J.S., PESOLE, G., PHILLIPPY, A.M., PONTING, C.P., POP, M., PORCELLI, D., POWELL, J.R., PROHASKA, S., PRUITT, K., PUIG, M., QUESNEVILLE, H., RAM, K.R., RAND, D., RASMUSSEN, M.D., REED, L.K., REENAN, R., REILY, A., REMINGTON, K.A., RIEGER, T.T., RITCHIE, M.G., ROBIN, C., ROGERS, Y.H., ROHDE, C., ROZAS, J., RUBENFIELD, M.J., RUIZ, A., RUSSO, S., SALZBERG, S.L., SANCHEZ-GRACIA, A., SARANGA, D.J., SATO, H., SCHAEFFER, S.W., SCHATZ, M.C., SCHLENKE, T., SCHWARTZ, R., SEGARRA, C., SINGH, R.S., SIROT, L., SIROTA, M., SISNEROS, N.B., SMITH, C.D., SMITH, T.F., SPIETH, J., STAGE, D.E., STARK, A., STEPHAN, W., STRAUSBERG, R.L., STREMPEL, S., STURGILL, D., SUTTON, G., SUTTON, G.G., TAO, W., TEICHMANN, S., TOBARI, Y.N., TOMIMURA, Y., TSOLAS, J.M., VALENTE, V.L., VENTER, E., VENTER, J.C., VICARIO, S., VIEIRA, F.G., VILELLA, A.J., VILLASANTE, A., WALENZ, B., WANG, J., WASSERMAN, M., WATTS, T., WILSON, D., WILSON, R.K., WING, R.A., WOLFNER, M.F., WONG, A., WONG, G.K., WU, C.I., WU, G., YAMAMOTO, D., YANG, H.P., YANG, S.P., YORKE, J.A., YOSHIDA, K., ZDOBNOV, E., ZHANG, P., ZHANG, Y., ZIMIN, A.V., BALDWIN, J., ABDOUELLEIL, A., ABDULKADIR, J., ABEBE, A., ABERA, B., ABREU, J., ACER, S.C., AFTUCK, L., ALEXANDER, A., AN, P., ANDERSON, E., ANDERSON, S., ARACHI, H., AZER, M., BACHANTSANG, P., BARRY, A., BAYUL, T., BERLIN, A., BESSETTE, D., BLOOM, T., BLYE, J., BOGUSLAVSKIY, L., BONNET, C., BOUKHGALTER, B., BOURZGUI, I., BROWN, A., CAHILL, P., CHANNER, S., CHESHATSANG, Y., CHUDA, L., CITROEN, M., COLLYMORE, A., COOKE, P., COSTELLO, M., D'ACO, K., DAZA, R., DE HAAN, G., DEGRAY, S., DEMASO, C., DHARGAY, N., DOOLEY, K., DOOLEY, E., DORICENT, M., DORJE, P., DORJEE, K., DUPES, A., ELONG, R., FALK, J., FARINA, A., FARO, S., FERGUSON, D., FISHER, S., FOLEY, C.D., FRANKE, A., FRIEDRICH, D., GADBOIS, L., GEARIN, G., GEARIN, C.R., GIANNOUKOS, G., GOODE, T., GRAHAM, J., GRANDBOIS, E., GREWAL, S., GYALTSEN, K., HAFEZ, N., HAGOS, B., HALL, J., HENSON, C., HOLLINGER, A., HONAN, T., HUARD, M.D., HUGHES, L., HURHULA, B., HUSBY, M.E., KAMAT, A., KANGA, B., KASHIN, S., KHAZANOVICH, D., KISNER, P., LANCE, K., LARA, M., LEE, W., LENNON, N., LETENDRE, F., LEVINE, R., LIPOVSKY, A., LIU, X., LIU, J., LIU, S., LOKYITSANG, T., LOKYITSANG, Y., LUBONJA, R., LUI, A., MACDONALD, P., MAGNISALIS, V., MARU, K., MATTHEWS, C., MCCUSKER, W., MCDONOUGH, S., MEHTA, T., MELDRIM, J., MENEUS, L., MIHAI, O., MIHALEV, A., MIHOVA, T., MITTELMAN, R., MLENGA, V., MONTMAYEUR, A., MULRAIN, L., NAVIDI, A., NAYLOR, J., NEGASH, T., NGUYEN, T., NGUYEN, N., NICOL, R., NORBU, C., NORBU, N., NOVOD, N., O'NEILL, B.,

OSMAN, S., MARKIEWICZ, E., OYONO, O.L., PATTI, C., PHUNKHANG, P., PIERRE, F., PRIEST, M., RAGHURAMAN, S., REGE, F., REYES, R., RISE, C., ROGOV, P., ROSS, K., RYAN, E., SETTIPALLI, S., SHEA, T., SHERPA, N., SHI, L., SHIH, D., SPARROW, T., SPAULDING, J., STALKER, J., STANGE-THOMANN, N., STAVROPOULOS, S., STONE, C., STRADER, C., TESFAYE, S., THOMSON, T., THOULUTSANG, Y., THOULUTSANG, D., TOPHAM, K., TOPPING, I., TSAMLA, T., VASSILIEV, H., VO, A., WANGCHUK, T., WANGDI, T., WEIAND, M., WILKINSON, J., WILSON, A., YADAV, S., YOUNG, G., YU, Q., ZEMBEK, L., ZHONG, D., ZIMMER, A., ZWIRKO, Z., ALVAREZ, P., BROCKMAN, W., BUTLER, J., CHIN, C., GRABHERR, M., KLEBER, M., MAUCELI, E., MACCALLUM, I., and CONSORTIUM, D.G., 2007, Evolution of genes and genomes on the *Drosophila* phylogeny: Nature, v. 450, p. 203-18, doi: 10.1038/nature06341.

COGNI, R., DING, S.D., PIMENTEL, A.C., DAY, J.P., and JIGGINS, F.M., 2021, *Wolbachia* reduces virus infection in a natural population of *Drosophila*: Commun Biol, v. 4, p. 1327, doi: 10.1038/s42003-021-02838-z.

CONNER, W., 2021, A phylogeny for the *Drosophila* montium species group: A model clade for comparative analyses: Molecular Phylogenetics and Evolution, v. 158, p. 107061.

CONNER, W.R., BLAXTER, M.L., ANFORA, G., OMETTO, L., ROTA-STABELLI, O., and TURELLI, M., 2017, Genome comparisons indicate recent transfer of *w*Ri-like *Wolbachia* between sister species *Drosophila suzukii* and *D. subpulchrella*: Ecol Evol, v. 7, p. 9391-9404, doi: 10.1002/ece3.3449.

COOPER, B.S., GINSBERG, P.S., TURELLI, M., and MATUTE, D.R., 2017, *Wolbachia* in the *Drosophila yakuba* Complex: Pervasive Frequency Variation and Weak Cytoplasmic Incompatibility, but No Apparent Effect on Reproductive Isolation: Genetics, v. 205, p. 333-351, doi: 10.1534/genetics.116.196238.

COOPER, B.S., VANDERPOOL, D., CONNER, W.R., MATUTE, D.R., and TURELLI, M., 2019, *Wolbachia* acquisition by *Drosophila yakuba*-clade hosts and transfer of incompatibility loci between distantly related *Wolbachia*: Genetics, v. 212, p. 1399-1419, doi: 10.1534/genetics.119.302349.

CUMMINGS, M.R., BROWN, N.M., and KING, R.C., 1971, The cytology of the vitellogenic stages of oogenesis in *Drosophila melanogaster*. 3. Formation of the vitelline membrane: Z Zellforsch Mikrosk Anat, v. 118, p. 482-92, doi: 10.1007/BF00324615.

DEDEINE, F., VAVRE, F., FLEURY, F., LOPPIN, B., HOCHBERG, M.E., and BOULETREAU, M., 2001, Removing symbiotic *Wolbachia* bacteria specifically inhibits oogenesis in a parasitic wasp: Proc Natl Acad Sci U S A, v. 98, p. 6247-52, doi: 10.1073/pnas.101304298.

ELLEGAARD, K.M., KLASSON, L., NÄSLUND, K., BOURTZIS, K., and ANDERSSON, S.G., 2013, Comparative genomics of *Wolbachia* and the bacterial species concept: PLoS Genet, v. 9, p. e1003381, doi: 10.1371/journal.pgen.1003381.

FERREE, P.M., FRYDMAN, H.M., LI, J.M., CAO, J., WIESCHAUS, E., and SULLIVAN, W., 2005, *Wolbachia* utilizes host microtubules and Dynein for anterior localization in the *Drosophila* oocyte: PLoS Pathog, v. 1, p. e14, doi: 10.1371/journal.ppat.0010014.

FOX, J., and WEISBERG, S., 2019, An R companion to applied regression: SAGE.

FRECKLETON, R.P., HARVEY, P.H., and PAGEL, M., 2002, Phylogenetic analysis and comparative data: a test and review of evidence: Am Nat, v. 160, p. 712-26, doi: 10.1086/343873.

FRYDMAN, H.M., LI, J.M., ROBSON, D.N., and WIESCHAUS, E., 2006, Somatic stem cell niche tropism in *Wolbachia*: Nature, v. 441, p. 509-12, doi: 10.1038/nature04756.

GERTH, M., and BLEIDORN, C., 2016, Comparative genomics provides a timeframe for *Wolbachia* evolution and exposes a recent biotin synthesis operon transfer: Nat Microbiol, v. 2, p. 16241, doi: 10.1038/nmicrobiol.2016.241.

GRIEDER, N.C., DE CUEVAS, M., and SPRADLING, A.C., 2000, The fusome organizes the microtubule network during oocyte differentiation in *Drosophila*: Development, v. 127, p. 4253-64.

GROBLER, Y., YUN, C.Y., KAHLER, D.J., BERGMAN, C.M., LEE, H., OLIVER, B., and LEHMANN, R., 2018, Whole genome screen reveals a novel relationship between *Wolbachia* levels and *Drosophila* host translation: PLoS Pathog, v. 14, p. e1007445, doi: 10.1371/journal.ppat.1007445.

GUO, Y., GONG, J.T., MO, P.W., HUANG, H.J., and HONG, X.Y., 2019, *Wolbachia* localization during *Laodelphax striatellus* embryogenesis: J Insect Physiol, v. 116, p. 125-133, doi: 10.1016/j.jinsphys.2019.05.006.

HAGUE, M.T.J., MAVENGERE, H., MATUTE, D.R., and COOPER, B.S., 2020a, Environmental and genetic contributions to imperfect *w*Mel-like *Wolbachia* transmission and frequency variation: Genetics, v. 215, p. 1117-1132, doi: 10.1534/genetics.120.303330.

HAGUE, M.T.J., CALDWELL, C.N., and COOPER, B.S., 2020b, Pervasive effects of *Wolbachia* on host temperature preference: mBio, v. 11, doi: 10.1128/mBio.01768-20.

HAGUE, M.T.J., SHROPSHIRE, J.D., CALDWELL, C.N., STATZ, J.P., STANEK, K.A., CONNER, W.R., and COOPER, B.S., 2022, Temperature effects on cellular host-microbe interactions explain continent-wide endosymbiont prevalence: Curr Biol, v. 32, p. 878-888.e8, doi: 10.1016/j.cub.2021.11.065.

HARMON, L.J., WEIR, J.T., BROCK, C.D., GLOR, R.E., and CHALLENGER, W., 2008, GEIGER: investigating evolutionary radiations: Bioinformatics, v. 24, p. 129-31, doi: 10.1093/bioinformatics/btm538.

HILGENBOECKER, K., HAMMERSTEIN, P., SCHLATTMANN, P., TELSCHOW, A., and WERREN, J.H., 2008, How many species are infected with *Wolbachia*?--A statistical analysis of current data: FEMS Microbiol Lett, v. 281, p. 215-20, doi: 10.1111/j.1574-6968.2008.01110.x.

HOFFMANN, A.A., TURELLI, M., and HARSHMAN, L.G., 1990, Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*: Genetics, v. 126, p. 933-48, doi: 10.1093/genetics/126.4.933.

HOSKINS, R.A., CARLSON, J.W., WAN, K.H., PARK, S., MENDEZ, I., GALLE, S.E., BOOTH, B.W., PFEIFFER, B.D., GEORGE, R.A., SVIRSKAS, R., KRZYWINSKI, M., SCHEIN, J., ACCARDO, M.C., DAMIA, E., MESSINA, G., MÉNDEZ-LAGO, M., DE PABLOS, B., DEMAKOVA, O.V., ANDREYEVA, E.N., BOLDYREVA, L.V., MARRA, M., CARVALHO, A.B., DIMITRI, P., VILLASANTE, A., ZHIMULEV, I.F., RUBIN, G.M., KARPEN, G.H., and CELNIKER, S.E.,

2015, The Release 6 reference sequence of the *Drosophila melanogaster* genome: Genome Res, v. 25, p. 445-58, doi: 10.1101/gr.185579.114.

HU, T.T., EISEN, M.B., THORNTON, K.R., and ANDOLFATTO, P., 2013, A second-generation assembly of the *Drosophila simulans* genome provides new insights into patterns of lineage-specific divergence: Genome Res, v. 23, p. 89-98, doi: 10.1101/gr.141689.112.

HÖHNA, S., LANDIS, M.J., HEATH, T.A., BOUSSAU, B., LARTILLOT, N., MOORE, B.R., HUELSENBECK, J.P., and RONQUIST, F., 2016, RevBayes: Bayesian Phylogenetic Inference Using Graphical Models and an Interactive Model-Specification Language: Syst Biol, v. 65, p. 726-36, doi: 10.1093/sysbio/syw021.

JACKMAN, S.D., VANDERVALK, B.P., MOHAMADI, H., CHU, J., YEO, S., HAMMOND, S.A., JAHESH, G., KHAN, H., COOMBE, L., WARREN, R.L., and BIROL, I., 2017, ABySS 2.0: resource-efficient assembly of large genomes using a Bloom filter: Genome Res, v. 27, p. 768-777, doi: 10.1101/gr.214346.116.

JOSHI, N., and FASS, J., 2011, Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files.

KAMATH, A.D., DEEHAN, M.A., and FRYDMAN, H.M., 2018, Polar cell fate stimulates *Wolbachia* intracellular growth: Development, v. 145, doi: 10.1242/dev.158097.

KATOH, K., and STANDLEY, D.M., 2013, MAFFT multiple sequence alignment software version 7: improvements in performance and usability: Mol Biol Evol, v. 30, p. 772-80, doi: 10.1093/molbev/mst010.

KLASSON, L., WESTBERG, J., SAPOUNTZIS, P., NÄSLUND, K., LUTNAES, Y., DARBY, A.C., VENETI, Z., CHEN, L., BRAIG, H.R., GARRETT, R., BOURTZIS, K., and ANDERSSON, S.G., 2009, The mosaic genome structure of the *Wolbachia w*Ri strain infecting *Drosophila simulans*: Proc Natl Acad Sci U S A, v. 106, p. 5725-30, doi: 10.1073/pnas.0810753106.

KOSE, H., and KARR, T.L., 1995, Organization of *Wolbachia* pipientis in the *Drosophila* fertilized egg and embryo revealed by an anti-*Wolbachia* monoclonal antibody: Mech Dev, v. 51, p. 275-88, doi: 10.1016/0925-4773(95)00372-x.

KÜHN, S., and ENNINGA, J., 2020, The actin comet guides the way: How *Listeria* actin subversion has impacted cell biology, infection biology and structural biology: Cell Microbiol, v. 22, p. e13190, doi: 10.1111/cmi.13190.

LANDMANN, F., BAIN, O., MARTIN, C., UNI, S., TAYLOR, M.J., and SULLIVAN, W., 2012, Both asymmetric mitotic segregation and cell-to-cell invasion are required for stable germline transmission of *Wolbachia* in filarial nematodes: Biol Open, v. 1, p. 536-47, doi: 10.1242/bio.2012737.

LUO, Y., AHMAD, E., and LIU, S.T., 2018, MAD1: Kinetochore Receptors and Catalytic Mechanisms: Front Cell Dev Biol, v. 6, p. 51, doi: 10.3389/fcell.2018.00051.

MAHAJAN-MIKLOS, S., and COOLEY, L., 1994, Intercellular cytoplasm transport during *Drosophila* oogenesis: Dev Biol, v. 165, p. 336-51, doi: 10.1006/dbio.1994.1257.

MAHOWALD, A.P., and STRASSHEIM, J.M., 1970, Intercellular migration of centrioles in the germarium of *Drosophila melanogaster*. An electron microscopic study: J Cell Biol, v. 45, p. 306-20, doi: 10.1083/jcb.45.2.306.

MAIMON, I., and GILBOA, L., 2011, Dissection and staining of *Drosophila* larval ovaries: J Vis Exp, doi: 10.3791/2537.

MEANY, M.K., CONNER, W.R., RICHTER, S.V., BAILEY, J.A., TURELLI, M., and COOPER, B.S., 2019, Loss of cytoplasmic incompatibility and minimal fecundity effects explain relatively low *Wolbachia* frequencies in *Drosophila mauritiana*: Evolution, v. 73, p. 1278-1295, doi: 10.1111/evo.13745.

MERKLE, J.A., WITTES, J., and SCHÜPBACH, T., 2020, Signaling between somatic follicle cells and the germline patterns the egg and embryo of *Drosophila*: Curr Top Dev Biol, v. 140, p. 55-86, doi: 10.1016/bs.ctdb.2019.10.004.

METCALF, J.A., JO, M., BORDENSTEIN, S.R., and JAENIKE, J., 2014, Recent genome reduction of *Wolbachia* in *Drosophila recens* targets phage WO and narrows candidates for reproductive parasitism: PeerJ, v. 2, p. e529, doi: 10.7717/peerj.529.

O'NEILL, S.L., GIORDANO, R., COLBERT, A.M., KARR, T.L., and ROBERTSON, H.M., 1992, 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects: Proc Natl Acad Sci U S A, v. 89, p. 2699-702, doi: 10.1073/pnas.89.7.2699.

PAGEL, M., 1999, Inferring the historical patterns of biological evolution: Nature, v. 401, p. 877-84, doi: 10.1038/44766.

PANNEBAKKER, B.A., LOPPIN, B., ELEMANS, C.P., HUMBLOT, L., and VAVRE, F., 2007, Parasitic inhibition of cell death facilitates symbiosis: Proc Natl Acad Sci U S A, v. 104, p. 213-5, doi: 10.1073/pnas.0607845104.

QUINLAN, M.E., 2016, Cytoplasmic Streaming in the *Drosophila* Oocyte: Annu Rev Cell Dev Biol, v. 32, p. 173-195, doi: 10.1146/annurev-cellbio-111315-125416.

RAMALHO, M.O., VIEIRA, A.S., PEREIRA, M.C., MOREAU, C.S., and BUENO, O.C., 2018, Transovarian Transmission of *Blochmannia* and *Wolbachia* Endosymbionts in the Neotropical Weaver Ant *Camponotus textor* (Hymenoptera, Formicidae): Curr Microbiol, v. 75, p. 866-873, doi: 10.1007/s00284-018-1459-3.

RAYCHOUDHURY, R., BALDO, L., OLIVEIRA, D.C., and WERREN, J.H., 2009, Modes of acquisition of *Wolbachia*: horizontal transfer, hybrid introgression, and codivergence in the *Nasonia* species complex: Evolution, v. 63, p. 165-83, doi: 10.1111/j.1558-5646.2008.00533.x.

RUSSELL, S.L., CHAPPELL, L., and SULLIVAN, W., 2019, A symbiont's guide to the germline: Curr Top Dev Biol, v. 135, p. 315-351, doi: 10.1016/bs.ctdb.2019.04.007.

RUSSELL, S.L., LEMSEFFER, N., and SULLIVAN, W.T., 2018, *Wolbachia* and host germline components compete for kinesin-mediated transport to the posterior pole of the *Drosophila* oocyte: PLoS Pathog, v. 14, p. e1007216, doi: 10.1371/journal.ppat.1007216.

RØRTH, P., 2002, Initiating and guiding migration: lessons from border cells: Trends Cell Biol, v. 12, p. 325-31, doi: 10.1016/s0962-8924(02)02311-5.

SALZBERG, S.L., DUNNING HOTOPP, J.C., DELCHER, A.L., POP, M., SMITH, D.R., EISEN, M.B., and NELSON, W.C., 2005, Serendipitous discovery of *Wolbachia* genomes in multiple *Drosophila* species: Genome Biol, v. 6, p. R23, doi: 10.1186/gb-2005-6-3-r23.

SANDER, 1968, Developmental physiological studies on embryonal mycetoma *Euscelis plebejus* I. Removal of and abnormal combination of unicellular components of symbiotic sysems: Dev Biol, v. 17, p. 16-38.

SEEMANN, T., 2014, Prokka: rapid prokaryotic genome annotation: Bioinformatics, v. 30, p. 2068-9, doi: 10.1093/bioinformatics/btu153.

SERBUS, L.R., CASPER-LINDLEY, C., LANDMANN, F., and SULLIVAN, W., 2008, The genetics and cell biology of *Wolbachia*-host interactions: Annu Rev Genet, v. 42, p. 683-707, doi: 10.1146/annurev.genet.41.110306.130354.

SERBUS, L.R., CHA, B.J., THEURKAUF, W.E., and SAXTON, W.M., 2005, Dynein and the actin cytoskeleton control kinesin-driven cytoplasmic streaming in *Drosophila* oocytes: Development, v. 132, p. 3743-52, doi: 10.1242/dev.01956.

SERBUS, L.R., FERRECCIO, A., ZHUKOVA, M., MCMORRIS, C.L., KISELEVA, E., and SULLIVAN, W., 2011, A feedback loop between *Wolbachia* and the *Drosophila* gurken mRNP complex influences *Wolbachia* titer: J Cell Sci, v. 124, p. 4299-308, doi: 10.1242/jcs.092510.

SERBUS, L.R., and SULLIVAN, W., 2007, A cellular basis for *Wolbachia* recruitment to the host germline: PLoS Pathog, v. 3, p. e190, doi: 10.1371/journal.ppat.0030190.

SERBUS, L.R., WHITE, P.M., SILVA, J.P., RABE, A., TEIXEIRA, L., ALBERTSON, R., and SULLIVAN, W., 2015, The impact of host diet on *Wolbachia* titer in *Drosophila*: PLoS Pathog, v. 11, p. e1004777, doi: 10.1371/journal.ppat.1004777.

SHEEHAN, K.B., MARTIN, M., LESSER, C.F., ISBERG, R.R., and NEWTON, I.L., 2016, Identification and Characterization of a Candidate *Wolbachia pipientis* Type IV Effector That Interacts with the Actin Cytoskeleton: MBio, v. 7, doi: 10.1128/mBio.00622-16.

STEINHAUER, J., and KALDERON, D., 2006, Microtubule polarity and axis formation in the *Drosophila* oocyte: Dev Dyn, v. 235, p. 1455-68, doi: 10.1002/dvdy.20770.

SULLIVAN, W., 2017, *Wolbachia*, bottled water, and the dark side of symbiosis: Mol Biol Cell, v. 28, p. 2343-2346, doi: 10.1091/mbc.E17-02-0132.

SULLIVAN, W., ASHBURNER, M., and HAWLEY, R.S., 2000, *Drosophila* Protocols: Cold Spring Harbor Press.

SUVOROV, A., KIM, B.Y., WANG, J., ARMSTRONG, E.E., PEEDE, D., D'AGOSTINO, E.R.R., PRICE, D.K., WADDELL, P., LANG, M., COURTIER-ORGOGOZO, V., DAVID, J.R., PETROV, D., MATUTE, D.R., SCHRIDER, D.R., and COMEAULT, A.A., 2022, Widespread introgression across a phylogeny of 155 *Drosophila* genomes: Curr Biol, v. 32, p. 111-123.e5, doi: 10.1016/j.cub.2021.10.052.

THEURKAUF, W.E., 1994, Microtubules and cytoplasm organization during *Drosophila* oogenesis: Dev Biol, v. 165, p. 352-60, doi: 10.1006/dbio.1994.1258.

THEURKAUF, W.E., and HAZELRIGG, T.I., 1998, In vivo analyses of cytoplasmic transport and cytoskeletal organization during *Drosophila* oogenesis: characterization of a multi-step anterior localization pathway: Development, v. 125, p. 3655-66.

THEURKAUF, W.E., SMILEY, S., WONG, M.L., and ALBERTS, B.M., 1992, Reorganization of the cytoskeleton during *Drosophila* oogenesis: implications for axis specification and intercellular transport: Development, v. 115, p. 923-36.

TOOMEY, M.E., PANARAM, K., FAST, E.M., BEATTY, C., and FRYDMAN, H.M., 2013, Evolutionarily conserved *Wolbachia*-encoded factors control pattern of stem-cell niche tropism in *Drosophila* ovaries and favor infection: Proc Natl Acad Sci U S A, v. 110, p. 10788-93, doi: 10.1073/pnas.1301524110.

TURELLI, M., COOPER, B.S., RICHARDSON, K.M., GINSBERG, P.S., PECKENPAUGH, B., ANTELOPE, C.X., KIM, K.J., MAY, M.R., ABRIEUX, A., WILSON, D.A., BRONSKI, M.J., MOORE, B.R., GAO, J.J., EISEN, M.B., CHIU, J.C., CONNER, W.R., and HOFFMANN, A.A., 2018, Rapid Global Spread of *w*Ri-like *Wolbachia* across Multiple *Drosophila*: Curr Biol, v. 28, p. 963-971.e8, doi: 10.1016/j.cub.2018.02.015.

TURELLI, M., and HOFFMANN, A.A., 1995, Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations: Genetics, v. 140, p. 1319-38.

VENETI, Z., CLARK, M.E., KARR, T.L., SAVAKIS, C., and BOURTZIS, K., 2004, Heads or tails: host-parasite interactions in the *Drosophila-Wolbachia* system: Appl Environ Microbiol, v. 70, p. 5366-72, doi: 10.1128/AEM.70.9.5366-5372.2004.

WERREN, J.H., BALDO, L., and CLARK, M.E., 2008, *Wolbachia*: master manipulators of invertebrate biology: Nat Rev Microbiol, v. 6, p. 741-51, doi: 10.1038/nrmicro1969.

WHITE, P.M., PIETRI, J.E., DEBEC, A., RUSSELL, S., PATEL, B., and SULLIVAN, W., 2017, Mechanisms of Horizontal Cell-to-Cell Transfer of *Wolbachia* spp. in *Drosophila melanogaster*: Appl Environ Microbiol, v. 83, doi: 10.1128/AEM.03425-16.

WHITE, P.M., SERBUS, L.R., DEBEC, A., CODINA, A., BRAY, W., GUICHET, A., LOKEY, R.S., and SULLIVAN, W., 2017, Reliance of *Wolbachia* on High Rates of Host Proteolysis Revealed by a Genome-Wide RNAi Screen of *Drosophila* Cells: Genetics, v. 205, p. 1473-1488, doi: 10.1534/genetics.116.198903.

WU, M., SUN, L.V., VAMATHEVAN, J., RIEGLER, M., DEBOY, R., BROWNLIE, J.C., MCGRAW, E.A., MARTIN, W., ESSER, C., AHMADINEJAD, N., WIEGAND, C., MADUPU, R., BEANAN, M.J., BRINKAC, L.M., DAUGHERTY, S.C., DURKIN, A.S., KOLONAY, J.F., NELSON, W.C., MOHAMOUD, Y., LEE, P., BERRY, K., YOUNG, M.B., UTTERBACK, T., WEIDMAN, J., NIERMAN, W.C., PAULSEN, I.T., NELSON, K.E., TETTELIN, H., O'NEILL, S.L., and EISEN, J.A., 2004, Phylogenomics of the reproductive parasite *Wolbachia* pipientis *w*Mel: a streamlined genome overrun by mobile genetic elements: PLoS Biol, v. 2, p. E69, doi: 10.1371/journal.pbio.0020069.

ZUG, R., and HAMMERSTEIN, P., 2012, Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected: PLoS One, v. 7, p. e38544, doi: 10.1371/journal.pone.0038544.