Distinct *Wolbachia* localization patterns in oocytes of diverse host species reveal multiple strategies of maternal transmission

Yonah A. Radousky[1,*], Michael T.J. Hague[2,*,†], Sommer Fowler[1], Eliza Paneru[1], Adan Codina[1], Cecilia Rugamas[1], Grant Hartzog[1], Brandon S. Cooper[2], William Sullivan[1]

[1]Department of Molecular, Cell, and Developmental Biology, University of California Santa Cruz, Santa Cruz, California, United States of America

[2]Division of Biological Sciences, University of Montana, Missoula, Montana, United States of America

[*]Co-first authors

[†]Corresponding Author:
Michael T.J. Hague
Division of Biological Sciences, University of Montana
32 Campus Dr. HS 104, Missoula, MT 59812
michael.hague@mso.umt.edu

ABSTRACT

A broad array of endosymbionts radiate through host populations via vertical transmission, yet
much remains unknown concerning the cellular basis, diversity and routes underlying this
transmission strategy. Here we address these issues, by examining the cellular distributions of
Wolbachia strains that diverged up to 50 million years ago in the oocytes of 18 divergent
Drosophila species. This analysis revealed three *Wolbachia* distribution patterns: 1) a tight
clustering at the posterior pole plasm (the site of germline formation); 2) a concentration at the
posterior pole plasm, but with a significant bacteria population distributed throughout the oocyte;
3) and a distribution throughout the oocyte, with none or very few located at the posterior pole
plasm. Examination of this latter class indicates *Wolbachia* accesses the posterior pole plasm
during the interval between late oogenesis and the blastoderm formation. We also find that one
Wolbachia strain in this class concentrates in the posterior somatic follicle cells that encompass
the pole plasm of the developing oocyte. In contrast, strains in which *Wolbachia* concentrate at the
posterior pole plasm generally exhibit no or few *Wolbachia* in the follicle cells associated with the
pole plasm. Taken together, these studies suggest that for some *Drosophila* species, *Wolbachia*
invade the germline from neighboring somatic follicle cells. Phylogenomic analysis indicates that
closely related *Wolbachia* strains tend to exhibit similar patterns of posterior localization,
suggesting that specific localization strategies are a function of *Wolbachia*-associated factors.
Previous studies revealed that endosymbionts rely on *one* of two distinct routes of vertical
transmission: continuous maintenance in the germline (germline-to-germline) or a more circuitous
route via the soma (germline-to-soma-to-germline). Here we provide compelling evidence that
Wolbachia strains infecting *Drosophila* species maintain the diverse arrays of cellular mechanisms
necessary for *both* of these distinct transmission routes. This characteristic may account for its
ability to infect and spread globally through a vast range of host insect species.

INTRODUCTION

A large class of endosymbionts rely on efficient vertical transmission to ensure their maintenance and spread through host populations. Many insect endosymbionts achieve this by a strict association with the host germline through successive generations. That is, during oogenesis, endosymbionts target the extreme posterior of the oocyte, the site of germline formation. In contrast to this strict germline-to-germline route of maternal transmission, other endosymbionts occupy the germline via invasion from neighboring somatic cells (Russell et al., 2019). Occupation of the germline via a somatic route requires that the endosymbiont has the capacity to undergo cell-to-cell transmission.

Among all known endosymbionts, maternally transmitted *Wolbachia pipientis* is the most common, infecting 30 to 60% of all insect species (Hilgenboecker et al., 2008; Zug and Hammerstein, 2012; Weinert et al., 2015). Hosts acquire *Wolbachia* from other species via horizontal (non-sexual) transfer or from sister species via introgressive transfer (O'Neill et al., 1992; Raychoudhury et al., 2009; Gerth and Bleidorn, 2016; Conner et al., 2017; Turelli et al., 2018; Cooper et al., 2019). Following establishment, maternal transmission rates in conjunction with *Wolbachia* effects on host fitness and reproduction determine *Wolbachia* spread and prevalence in host populations (Hoffmann et al., 1990; Turelli and Hoffmann, 1995; Cooper et al., 2017; Cogni et al., 2021).

While sometimes imperfect, the success and prevalence of *Wolbachia* ultimately depends on vertical transmission through the female (Hoffmann et al., 1990; Turelli and Hoffmann, 1995; Carrington et al., 2011; Hague et al., 2022). The ability of *Wolbachia* to establish and persist in diverse insect species indicates that it has evolved to usurp highly conserved host processes and components. For example, studies of *w*Mel *Wolbachia* in model organism *D. melanogaster* suggest that *Wolbachia* engage conserved host motor proteins Dynein and Kinesin in order to target the extreme posterior of the oocyte, the site of germline formation (Ferree et al., 2005; Serbus and Sullivan, 2007; Russell et al., 2018).

To study the vertical transmission of *Wolbachia* in *Drosophila* we take advantage of the wealth of knowledge regarding the molecular and cell biology of *Drosophila* oogenesis (for a review, see Bastock and St Johnston, 2008). *Drosophila* oocytes contain two clusters of ovarioles, each containing two to three germline stem cells and various stages of developing oocytes. The stem cells undergo asymmetric mitotic division in which one daughter cell undergoes self-renewal and the other differentiates into a cytoblast. The cytoblast undergoes four rounds of mitotic divisions resulting in 16 interconnected cells, one of which becomes the oocyte nucleus while the others differentiate into polyploid nurse cells. The developing oocyte is encapsulated by a monolayer of somatically derived follicle cells resulting in the formation of discrete egg chambers as the maturing oocyte progresses through the ovariole. During the late stages of oogenesis, the nurse cell cytoplasm is expelled into the oocyte, resulting in a rapid doubling of the oocyte volume and concomitant shrinking of nurse cell volume (Theurkauf and Hazelrigg, 1998; Bastock and St Johnston, 2008). This is followed by posterior pole plasm formation and establishment of the germline (Bilinski et al., 2017).

Wolbachia distribution and localization in insect oocytes and early embryos has been well characterized in *w*Mel-infected *D. melaongaster* (Kose and Karr, 1995; Veneti et al., 2004; Ferree

et al., 2005; Serbus et al., 2008; Ramalho et al., 2018; Russell et al., 2018; Guo et al., 2019). Early
in oogenesis, *w*Mel *Wolbachia* are evenly distributed throughout all the cells in the chamber. At
this stage, microtubules emanate from the oocyte into the nurse cells (Theurkauf, 1994; Grieder et
al., 2000). By associating with the minus-end directed microtubule motor protein Dynein, *w*Mel
Wolbachia are transported from the nurse cells and concentrate at the anterior end of the oocyte
(Ferree et al., 2005). As the oocyte matures, microtubules undergo a dramatic reorganization such
that they originate from the oocyte cortex with their plus-ends oriented toward the posterior pole
(Theurkauf et al., 1992; Steinhauer and Kalderon, 2006). This reorganization is accompanied by a
release and dispersal of *w*Mel *Wolbachia* from the anterior pole. Then, *w*Mel *Wolbachia* associate
with the plus-end directed motor protein Kinesin for transport to the posterior pole and the site of
germline formation (Serbus and Sullivan, 2007). There, *Wolbachia* (*w*Mel) associate with pole
plasm determinants maintaining their position at the site of germline formation (Serbus and
Sullivan, 2007) in spite of extensive cytoplasmic streaming (Serbus et al., 2005).

Thus, *w*Mel *Wolbachia* location and migration patterns provide insights into the host factors with
which *Wolbachia* engages throughout oogenesis. The *D. simulans*-associated *w*Ri strain exhibits
a similar anterior distribution in early oogenesis, and then becomes distributed across the entire
mature oocyte. However, unlike *w*Mel, it does not concentrate at the posterior pole of the oocyte
(Veneti et al., 2004; Serbus and Sullivan, 2007). This suggests *w*Ri engages host Dynein and
Kinesin, but not pole plasm determinants. *w*Mel *Wolbachia* strains introgressed into a *D. simulans*
host background exhibit a concentration in the pole plasm, indicating the *Wolbachia* genome rather
than the host genome drives oocyte localization properties (Poinsot et al., 1998; Veneti et al., 2004;
Serbus and Sullivan, 2007).

With the exception of *w*Mel and *w*Ri, little is known of the extent to which distribution patterns in
the oocyte and transmission routes vary among diverse *Wolbachia* strains and host species. Here,
we fill this important knowledge gap, by examining 18 *Drosophila* host species infected with
Wolbachia strains that diverged up to 50 million years ago. These studies also provide a direct
read-out of *Wolbachia* and host motor protein/pole plasm interactions that are highly conserved
and those that vary among species. This work complements and extends previous analyses
revealing *Wolbachia* strain-dependent localization patterns in the *Drosophila* oocyte and embryo
(Veneti *et al.*, 2004; Toomey et al., 2013).

We discovered that *Wolbachia* distribution patterns in the *Drosophila* oocytes fall into three
categories: (1) the vast majority of *Wolbachia* clusters at the posterior pole; (2) a concentration at
the posterior pole, but with a large portion of *Wolbachia* distributed throughout the entire oocyte;
and (3) *Wolbachia* distributed throughout oocyte, with none or very few located at the posterior
pole. This latter class is particularly interesting as it suggests *Wolbachia* likely enters the oocyte
via a somatic route for transmission to the next generation. We also provide evidence that
neighboring somatic follicle cells are the likely source of *Wolbachia* entering the germline.
Together, these findings indicate that *Wolbachia* are capable of achieving efficient vertical
transmission either via strict maintenance in the germline from one generation to the next or via
invasion of the germline through neighboring somatic cells. Thus, *Wolbachia* must maintain the
diverse arrays of cellular mechanisms necessary for *both* of these distinct transmission routes. We
identify candidate surface proteins that may be responsible for these diverse localization patterns
in host oocytes.

RESULTS

Wolbachia navigates the developing *Drosophila* oocyte in distinct stages

As described above, *w*Mel *Wolbachia* migration and distribution throughout oogenesis in *D. melanogaster* has been well documented (Kose and Karr, 1995; Veneti et al., 2004; Serbus and Sullivan, 2007; Serbus et al., 2008; Ramalho et al., 2018; Russell et al., 2018; Guo et al., 2019). As a reference for our analysis of *Wolbachia (w*Mel*)* oocyte distributions in a range of *Drosophila* species, the key distribution patterns of *w*Mel throughout *D. melanogaster* oocytes are depicted in **Figure 1**.

Stage 1 includes the germarium, a structure consisting of the germline stem cells and their cystoblast daughter cells that will differentiate into a mature oocytes. Progression to stage 2, involves the cystoblast undergoing four rounds of mitotic division to produce a syncytium of 16 cells connected by cytoplasmic bridges and surrounded by a layer of somatically derived follicle cells (Bastock and St Johnston, 2008). Propidium Iodine (PI)-stained *Wolbachia* (red) are clearly observed as puncta in the cytoplasm of the infected, but not the uninfected 16 cell cyst (**Figure 1 Stages 1-2, Arrows**). Actin is labeled in green.

During stage 3, the cyst has become polarized with one of the cells differentiating into an oocyte and the others becoming polyploid nurse cells. At this stage, *Wolbachia* rely on Dynein-mediated microtubule transport (Ferree et al., 2005) and are clearly seen in the infected oocyte cytoplasm anteriorly concentrated at two cytoplasmic bridges that connect to the nurse cells (**Figure 1, Stage 3, Arrows**).

During stages 4 and 5, the nurse cell/oocyte complex becomes oblong and is fully encapsulated by follicle cells. *Wolbachia* are clearly observed in both the oocyte and nurse cell cytoplasm (**Figure 1, Stages 4 and 5, Arrows**). The *Wolbachia* in the oocyte are tightly associated with the anterior cortex.

During stage 6, the number of follicle cells increases and they are more densely packed in a monolayer encompassing the oocyte. Here we discovered a previously unrecognized aspect of *Wolbachia* dynamics and localization: the number of anteriorly localized *Wolbachia* is greatly diminished at this stage (**Figure 1, Stage 6, Arrow**). This pattern was consistent across all 10 of the stage 6 oocytes we observed in our analysis (see **Supplemental Figure 1**).

At stage 7, *Wolbachia* are again observed in the oocyte. However, rather than being tightly associated with the anterior cortex, *Wolbachia* are evenly distributed throughout the oocyte (**Figure 1, Stage 7, Arrow**). During stage 7, the microtubules reorient such that their plus-ends extend toward the posterior pole (Cha et al., 2001; Cha et al., 2002; Steinhauer and Kalderon, 2006). Accordingly, the plus-end motor protein Kinesin is required to establish the even distribution of *Wolbachia* throughout the maturing oocyte (Serbus and Sullivan, 2007). Stage 8 heralds the beginning of vitellogenesis, the process of yolk formation and deposition of nutrients into the oocyte (Cummings et al., 1971). *Wolbachia* are near, but not yet located at, the posterior pole (**Figure 1, Stage 8, Arrow**). At stage 9, anterior follicle cell migration occurs (Rørth, 2002). At this stage, *Wolbachia* displays an archlike posterior cortical localization**. (Figure 1, Stage 9, Arrow**). By stage 10, there is a distinct concentration of *Wolbachia* along the posterior cortex, the

site of germline formation (**Figure 1, Stage 10, Arrow**). Nurse cell dumping will shortly follow,
leading into stage 11-12 with the oocyte rapidly growing in size (Mahajan-Miklos and Cooley,
1994).

The *w*Mel strain in *D. melanogaster* provides an outline of the molecular and cellular mechanisms
driving *Wolbachia* vertical transmission. It also raises the question of whether the mechanisms and
strategies observed in *D. melanogaster* are conserved across diverse *Wolbachia* strains and
Drosophila species. To explore this issue, the *w*Mel distribution in *D. melanogaster* serves as the
basis for our comparison of 18 other *Wolbachia* strains. Bayesian phylograms depicting
phylogenetic relationships among the *Wolbachia* strains and the *Drosophila* species are shown in
Figure 4 and **Supplemental Figure 2**. Our analysis comprised 18 A-group *Wolbachia* strains,
including six *w*Mel-like strains (Cooper et al., 2019; Hague et al., 2020a) and five *w*Ri-like strains
(Turelli et al., 2018), as well as the B-group *Wolbachia* strain *w*Mau, which diverged from A-
group *Wolbachia* up to 50 million years ago (Meany et al., 2019). These *Wolbachia* strains infect
18 divergent *Drosophila* host species spanning three species groups (Suvorov et al., 2022).

An initial concentration of *Wolbachia* at the oocyte anterior is a conserved feature

As described above, *w*Mel *Wolbachia* enter the oocyte through a complex of ring canals
connecting the nurse cells to the oocyte and concentrate at the anterior cortex at stage 3-5. Initially
observed in *D. melanogaster*, the number of *Wolbachia* increases dramatically while at the anterior
(Ferree et al., 2005). To determine if anterior localization is a conserved aspect of *Wolbachia*'s
navigation through the developing oocyte, we first characterized this trait in eight *Wolbachia*
strains in their respective hosts: *w*Pan in *D. pandora*, *w*Nik in *D. nikananu*, *w*Cha in *D. chauvacae*,
*w*Ha in *D. simulans*, *w*Tsa in D. *tsacasi*, *w*Boc in *D. bocki*, *w*Ana in *D. ananassae*, and *w*Mel in
D. melanogaster. All eight strains exhibited a distinct anterior localization during mid-oogenesis
(Figure 2). It is interesting that this localization occurs during stage 5 of oogenesis, which, in *D.
melanogaster,* is prior to the localization of all known anterior axis determinants. Thus, *Wolbachia*
likely relies on as yet undiscovered anteriorly concentrated factor(s).

Wolbachia exhibits three distinct distributions with respect to posterior localization in the
mature *Drosophila* oocyte

Concomitant with microtubule rearrangement during stages 7-8, *w*Mel *Wolbachia* are released
from the oocyte anterior resulting in the dispersal of the bacteria throughout the entire length of
the oocyte with a fraction concentrating at the posterior pole (Ferree et al., 2005). The posterior
dispersal of *Wolbachia* requires the plus-end motor protein Kinesin (Serbus and Sullivan, 2007).
Maintenance of those *Wolbachia* that reach the posterior pole rely on a stable association with key
pole plasm components (Serbus et al., 2011).

We next examined the oocytes of 18 *Drosophila* species infected with 19 diverse *Wolbachia*
strains ($N = 72$ total oocytes) to test the hypothesis that oocyte posterior distributions vary. Unlike
the conserved anterior localization, *Wolbachia* exhibit three distinct patterns of posterior
localization in stage 10 oocytes. As shown in **Figure 3** and **Supplemental Figure 3A,** eight
Wolbachia strains exhibit a localization pattern similar to that observed for *w*Mel in *D.*
melanogaster with the bulk of *Wolbachia* evenly distributed throughout the oocyte, and a large
fraction concentrated at the posterior pole. We refer to this pattern as Posteriorly Localized.

Our analysis revealed four *Wolbachia* strains manifested a second pattern in which the vast majority of *Wolbachia* localized to a distinct cluster at the posterior pole and only few *Wolbachia* are distributed throughout the remainder of the oocyte (**Figure 3, Supplemental Figure 3B**). In contrast to the pattern described above, rather than being distributed in a wide arc along the posterior pole, they are tightly centered at the extreme posterior region of the oocyte (**Figure 3B**). We refer to this pattern as <u>Posteriorly Clumped</u>.

Finally, seven *Wolbachia* strains exhibit a third pattern in which *Wolbachia* are distributed throughout the entire oocyte but there are few to no bacteria concentration at the posterior pole (**Figure 3 and Supplemental Figure 3C**). The lack of the *Wolbachia* at the posterior pole was unexpected as this was thought to be necessary for incorporation into the germline of the next generation. We refer to this pattern as <u>Dispersed</u>.

Quantification of *Wolbachia* oocyte distribution patterns

We used one-way ANOVAs to test whether the three classes (Posteriorly Localized, Posteriorly Clumped, and Dispersed) differ statistically in quantitative estimates of *Wolbachia* abundance in the whole oocyte, at the posterior region, and at the posterior cortex (**Supplemental Figure 4**). We defined the posterior region as the posterior 12.5% of the oocyte, and the posterior cortex as the narrow cortical region of Vasa expression (Russell et al., 2018), and we then measured total *Wolbachia* abundance in each region as the Corrected Total Cell Fluorescence (CTCF). We found that *Wolbachia* abundance differs significantly among the three classes (one-way ANOVA, $F_{(2,69)}$ = 6.2, P = 0.003), with Tukey multiple comparisons indicating that the Clumped class has significantly more *Wolbachia* in whole oocytes than the Localized group (P = 0.002). The three classes also differed significantly in the amount of *Wolbachia* in the posterior region ($F_{(2,69)}$ = 23.09, P < 0.001), such that the Dispersed class has significantly fewer *Wolbachia* at the posterior that the Clumped (P < 0.001) and Localized (P < 0.001) groups. Finally, the three classes also differ in the amount of *Wolbachia* at the posterior cortex ($F_{(2,64)}$ = 38.12, P < 0.001). Here, multiple comparisons indicate that all three classes significantly differ in levels of *Wolbachia* at the posterior cortex: the Clumped class has the most *Wolbachia*, followed by Localized, and then Dispersed (**Supplemental Figure 4**). Mean estimates of *Wolbachia* abundance (CTCF) for each *Wolbachia* strain and host species are shown in **Table 1**.

***Wolbachia* abundance in the oocyte posterior has strong phylogenetic signal**

We hypothesized that the diversity of *Wolbachia* localization patterns may be due to *Wolbachia*-associated factors, implying that closely related *Wolbachia* strains would exhibit similar localization patterns in host oocytes. We used the phylogram describing the evolutionary relationships among *Wolbachia* strains to test whether patterns of *Wolbachia* localization exhibit a phylogenetic signal using Pagel's λ (Pagel, 1999; Hague et al., 2020b; Hague et al., 2021) (**Figure 4**). A λ value of 1 would be consistent with trait evolution that entirely agrees with the *Wolbachia* phylogeny (i.e., strong phylogenetic signal supporting a role of *Wolbachia* factors), whereas a value of 0 would indicate that trait evolution occurs independently of phylogenetic relationships (Pagel, 1999; Freckleton et al., 2002). Most notably, we found that *Wolbachia* abundance (CTCF) in the posterior region exhibits a strong, significant phylogenetic signal (λ = 0.982 [0, 0.998], P = 0.021). *Wolbachia* abundance at the posterior cortex also has a high, but non-significant λ value (λ = 0.959 [0, 0.995], P = 0.683). Here, the large confidence intervals and our

simulations suggest that a larger *Wolbachia* phylogeny ($N = 25$, 50 *Wolbachia* strains) is required
to detect a significant signal of $\lambda > 0$ (**Supplemental Figure 5**). These results highlight differences
in patterns of *Wolbachia* localization between the *w*Mel- and *w*Ri-like *Wolbachia* clades (**Figure
3**). Closely related *w*Mel-like *Wolbachia* tend to occur at a higher abundance in the posterior region
of oocytes, whereas the *w*Ri-like strains are generally more dispersed. We found no evidence that
Wolbachia abundance in the whole oocyte exhibits a phylogenetic signal ($\lambda < 0.001$ [0, 0.753], *P*
= 1).

The correlation between *Wolbachia* localization in oocytes and phylogenetic divergence implies
that factors in the *Wolbachia* genome determine *Wolbachia* abundance at the oocyte posterior.
Because these distributions likely involve direct interactions between *Wolbachia* and the host
cytoplasm, we focused on six major *Wolbachia* surface proteins: the three *wsp* paralogs
(WD_1063/*wsp*, WD_0009/*wspB*, WD_0489/*wspC*), as well as WD_0501, WD_1041, WD_1085.
We tested whether sequence divergence at any of these candidate loci predicts *Wolbachia*
localization patterns. Specifically, we tested whether gene trees of the *Wolbachia* surface proteins
(**Supplemental Figure 6**) exhibit a phylogenetic signal, using the methods described above. One
surface protein (WD_1085) stood out with especially strong evidence of a phylogenetic signal,
with significant departures from $\lambda = 0$ for *Wolbachia* abundance at the posterior region ($\lambda = 0.974$
[0, 0.998], *P* = 0.001) and the posterior cortex ($\lambda = 0.942$ [0, 0.994], *P* = 0.028). In addition,
Wolbachia abundance at the posterior region also exhibited significant phylogenetic signal on the
gene tree of WD_0501 ($\lambda = 0.992$ [0.521, 1], *P* = 0.022). All tests for phylogenetic signal using
Wolbachia gene trees are summarized in **Supplemental Table 1**.

Motivated by previous successful analysis of *Wolbachia* protein function through ectopic
expression in yeast, we expressed codon-optimized forms of WD_1085 and WD_0501 in yeast
using a Galactose-inducible promoter (Sheehan et al., 2016; Rice et al., 2017). Under normal
growth conditions, ectopic expression of both genes inhibited growth, with WD_1085 exhibiting
a more pronounced inhibition (**Supplemental Figure 7**). Because *Wolbachia* exhibits a close
association with microtubules, we also assayed whether ectopic expression of these genes
enhanced the growth defects in situations in which the microtubule spindle assembly checkpoint
was compromised by a mad1 mutation (Luo et al., 2018), or when the microtubules were
compromised directly through the addition of benomyl. As indicated in **Supplemental Figure 7**,
ectopic expression of WD_1085 significantly inhibited growth when both the microtubules and
spindle checkpoint were compromised. These results suggest that ectopic WD_1085 expression
perturbs microtubule function.

We recently found that the surface protein WD_0009 (*wspB*) is pseudogenized due to a premature
stop codon in a tropical *w*Mel variant sampled from Australia, which seems to influence *w*Mel
abundance in stage 10 oocytes when hosts are reared in the cold (Hague et al., 2022). Here, we did
not find that the *wspB* gene tree exhibits phylogenetic signal related to *Wolbachia* oocyte
abundance (**Supplemental Table 1**); however, we note that *wspB* appears to be psuedogenized in
a number of other *Wolbachia* strains in addition to tropical *w*Mel. We found derived deletions of
varying length in the *wspB* sequences of *w*Ha and *w*Bai beginning at nucleotide position 256, both
of which produce frame shifts that generate multiple downstream stop codons (**Supplemental
Figure 8**). In addition, *w*Boc and *w*Leo share a large insertion starting at bp position 351. Because
the insertions contain a contig break, we were unable to resolve the full length of the insertion.
Lastly, *w*Bic contains a 104 bp insertion at bp position 647 that creates multiple downstream stop

codons. These results suggest that the surface protein *wspB* has become psuedogenized at least four times in different *Wolbachia* lineages. Notably, the *Wolbachia* strains with a putatively pseudogenized version of *wspB* do not differ from the other strains in mean *Wolbachia* abundance in whole oocytes (Wilcoxon rank sum test, $W = 36$, $P = 0.964$), the posterior region ($W = 37$, $P = 0.893$), or the posterior cortex ($W = 41$, $P = 0.622$). This suggests that psuedogenization of *wspB* does not influence the diverse patterns of *Wolbachia* localization observed here. To gain insight into *wspB* cellular function, we ectopically expressed this gene (and the other *wsp* paralogs; WD_1063/*wsp* and WD_0489/*wspC)* in yeast using the Galactose expression system described above. Using this system, we found that ectopic expression of codon optimized *wspB* and *wsp* significantly inhibited yeast growth **(Supplemental Figure 9)**. Ectopic expression of *wspC* kills the cells. Based on similar ectopic expression studies (Sheehan et al., 2016), the suppression is likely due to *Wolbachia* proteins interacting with and disrupting the function of yeast components and cellular processes that are conserved and performing similar functions in insect cells.

Lastly, we ran similar phylogenetic analyses using the host phylogeny (as opposed to *Wolbachia*) to test whether host-associated factors might also contribute to the diverse *Wolbachia* distribution patterns in fly oocytes (**Supplemental Figure 10**). We generally found that *Wolbachia* localization patterns do not exhibit a phylogenetic signal on the host tree. *Wolbachia* abundance in whole oocytes ($\lambda = 0.365$ [0, 0.856], $P = 0.656$), at the posterior region ($\lambda < 0.001$ [0, 0.658], $P = 1$), and at the posterior cortex ($\lambda < 0.001$ [0, 0.637], $P = 1$) did not exhibit phylogenetic signal using the host phylogram, implying that host-associated factors are not as important in determining diverse *Wolbachia* localization patterns in oocytes.

Vertical transmission in some host species likely relies on *Wolbachia* entering the germline from neighboring somatic tissues

Because *Wolbachia* strains with a Dispersed localization do not localize to the pole, one would expect maternal transmission to be compromised. However, previous studies demonstrate that some of these *Wolbachia*, such as the B-group strain *w*Mau in *D. mauritiana*, have high maternal transmission rates under standard laboratory conditions (i.e., constant 25°C), equivalent to that of *w*Mel in *D. melanogaster* (Meany et al., 2019; Hague et al., 2022). These findings suggest that, in addition to strict germline-to-germline transmission, *Wolbachia* may utilize alternative routes of maternal transmission. Because numerous vertically inherited endosymbionts are transmitted via cell-to-cell transmission from the soma to the germline (Frydman, 2006; Russell et al., 2019), we explored whether *Wolbachia* that infect *Drosophila* exploits this strategy.

To determine the developmental stage during which somatic *Wolbachia* might invade the germline in Dispersed species, we focused on B-group *w*Mau in *D. mauritiana.* Initially we examined the female germline (oocytes) of 3rd instar larva for the presence of *Wolbachia*. For comparison, we also examined the oocytes of *w*Mel-infected *D. melanogaster* (a strain with a Posteriorly Localized concentration of *Wolbachia*) 3rd-instar larvae using anti-FtsZ to mark the *Wolbachia* and anti-Vasa to mark the germline. As shown in **Figure 5**, *Wolbachia* are abundantly present in ring-like patterns in many of the Vasa-positive cells in the center of the developing oocyte. Imaging the oocytes of *w*Mau-infected *D. mauritiana* revealed a similar pattern with *Wolbachia* concentrated in the developing female germline of 3rd instar larva (**Figure 5**). Images of *w*Ri in *D. simulans* as well as *w*Bic in *D. bicornuta*—strains with a Dispersed localization and Posteriorly Clumped localization in mature oocytes, respectively—displayed the same ring-like structure in

the developing oocyte of 3^{rd} instar larva, although they appeared to have a much higher titer than
*w*Mau or *w*Mel (**Figure 5**). This indicates that in these species, in which vertical transmission
occurs with a few or no *Wolbachia* present at the posterior site of germline formation in mature
adult oocytes, *Wolbachia* are able to occupy the germline at some point between the final stages
of oocyte maturation and development of the fertilized egg into a 3^{rd} instar larva. Presumably the
source of the *Wolbachia* is via invasion from neighboring somatic cells.

To determine if *Wolbachia* occupied the germline prior to the 3^{rd} instar larval stage, we examined
the newly formed germline of late blastoderm and cellularized embryos. At this stage, newly
formed germline cells, known as the pole cells, form a distinct cluster of cells at the extreme
posterior to the embryo. In all the four strains described above, using PI or anti-*Wolbachia* staining,
Wolbachia are readily found in these posterior germline cells (**Figure 6).** Imaging *w*Mau in *D.*
mauritiana early blastoderm embryos revealed the presence of *Wolbachia* in the posterior pole
cells. This observation narrows the time window in which *Wolbachia* invades and occupies the
germline from between the final stages of oocyte maturation to early blastoderm formation. Images
taken of *D. mauritiana* in one-hour old embryos also showed *w*Mau positioned in the posterior
pole, narrowing the developmental window of *Wolbachia* germline occupation between the final
stages of oocyte maturation to one hour after fertilization **(Figure 6).** Interestingly, Veneti et al.
(2004) reported that *w*Mau exhibits an anterior localization pattern during *D. mauritiana*
embryogenesis; however, we found no such pattern for any of the embryos examined.

Our findings motivated us to examine *w*Mau in *D. mauritiana* oocytes just prior to egg deposition.
Previous studies found that in many infected *Drosophila* species *Wolbachia* is concentrated in the
most posterior positioned follicle cells (Kamath et al., 2018). These somatically-derived cells,
known as polar cells, directly contact the oocyte posterior pole plasm. In accord with Kamath et
al. (2018), we find that *Wolbachia* are sporadically distributed in subsets of more anterior
positioned follicle cells encompassing the oocyte. *w*Mau presence in the polar follicle cells is a
consistent feature in every *D. mauritiana* oocyte examined (**Figure 7).** Given their position, these
*w*Mau-infected cells are likely the polar follicle cells. These observations raise the possibility that
Wolbachia in the polar follicle cells may be a somatic source of germline *w*Mau in *D. mauritiana.*
In spite of much effort, however, we have not been able to find cytological evidence for this
hypothesis.

To determine the extent to which *Wolbachia* polar follicle cell localization is conserved, we
analyzed *Wolbachia* localization in the polar and more anterior follicle cells for seven *Wolbachia*
strains with the following patterns: Dispersed ($N = 3$), Posteriorly Localized ($N = 3$), and
Posteriorly Clumped ($N = 1$) (**Figure 7**). We scored for the presence of follicle cell *Wolbachia* in
each oocyte ($N = 75$) using the following criteria: the presence of *Wolbachia* in either both or one
of the posterior polar follicle cells (**Figure 7A)** and the presence of *Wolbachia* in the more
anteriorly localized follicle cells (**Figure 7B**). This analysis revealed that in a high percentage of
oocytes from Dispersed strains (a lack of the *Wolbachia* at the oocyte posterior), *Wolbachia* are
frequently present in the posterior polar follicle cells. In contrast, with the exception of wMel,
oocytes examined from strains in which *Wolbachia* are posteriorly concentrated (the Posteriorly
Localized and Clumped strains), few possess *Wolbachia* in the posterior follicle cells. The seven
Wolbachia strains exhibit significant variation in whether *Wolbachia* cells are found in the polar
follicle cells ($\chi^2_{(12)} = 65.15$, $P < 0.001$) (**Figure 7A**). In particular, the divergent B-group strain
*w*Mau stood out with a large proportion of oocytes containing *Wolbachia* in the two polar follicle

cells. The A-group *w*Mel strain also had a large portion of oocytes with *Wolbachia* in one or two
of the polar cells. We also found that the *Wolbachia* strains differ in whether *Wolbachia* cells are
found in the other more anterior follicle cells ($\chi^2_{(6)}$ = 43.29, *P* < 0.001) (**Figure 7B**). Again, the
*w*Mau and *w*Mel strains had a large portion of oocytes with *Wolbachia* in the other more anterior
follicle cells encompassing the oocyte. Of note, for *w*Mau in *D. mauritiana*, every oocyte had at
least one of their polar cells filled with *Wolbachia*, observationally at a higher titer (**Figure 7**).
This suggests that in the Dispersed strains, vertical transmission may be achieved via invasion of
Wolbachia from the polar follicle cells, particularly for the divergent B-Group *w*Mau strain.

DISCUSSION

Vertical transmission through the female lineage is a common strategy for many endosymbionts.
These endosymbionts achieve high efficiencies of maternal transmission either through direct
germline-to-germline transmission or invasion of the maternal germline through neighboring
somatic cells (Russell et al., 2019). Because the former involves a continuous presence in the
germline while the latter requires navigating a soma-to-germline passage, it is expected that each
requires distinct interactions with and manipulations of host cellular processes. Thus, it would be
expected that a given endosymbiont would have evolved to utilize one, but not both of these
strategies. Here we explore this issue by examining cellular aspects of *Wolbachia* vertical
transmission. In contrast to expectations, our comprehensive analysis demonstrates that *Wolbachia*
likely rely on both strict germline-to-germline and soma-to-germline vertical transmission
strategies. Which strategy employed varies among *Wolbachia* strains.

The initial anterior *Wolbachia* localization in the developing oocyte appears to be a conserved
aspect of *Wolbachia* transmission, as all eight strains examined exhibit this localization. It is
striking that this localization occurs prior to any known anterior determinant, indicating *Wolbachia*
may associate with an as yet undiscovered host factor concentrated at the anterior. That this
localization is conserved suggests that it has a functionally significant, but currently unknown role
in *Wolbachia* transmission. While anteriorly localized, the amount of *Wolbachia* dramatically
increases. Whether *Wolbachia* replication, transport from the nurse cells, or both are responsible
for this increase is unknown. The anterior location may be a rich source of membrane for
Wolbachia replication. The anterior localization also results in a high concentration of *Wolbachia*
that closely associated with, and perhaps influences, the status of the oocyte nucleus.

To our surprise, we discovered that immediately after the release of *w*Mel *Wolbachia* from its
anterior position in the oocyte (stage 6), the amount of *Wolbachia* in the oocyte dramatically
decreases (see **Figure 1, Stage 6** and **Supplemental Figure 1**). This suggests that the *Wolbachia*
are transported back into the nurse cell complex. The functional significance of this clearing of
Wolbachia from the oocyte is unknown. The retreat of *Wolbachia* occurs during the time when the
anteriorly positioned oocyte nucleus and neighboring follicle cells signal one another in order to
establish the dorsal ventral axis (Merkle et al., 2020). Perhaps *Wolbachia* exits the oocyte in order
to not disrupt this process. Given the orientation of microtubules at this stage, it is likely *Wolbachia*
rapidly exits the oocyte through an association with the plus-end directed motor protein Kinesin.

Following microtubule reorientation and the return of *Wolbachia* to the oocyte, *Wolbachia* spreads
posteriorly throughout the entire oocyte relying on the host motor protein Kinesin (Serbus and
Sullivan, 2007). All of the *Wolbachia* strains examined are distributed toward the posterior regions

suggesting the interaction with Kinesin is also conserved. How *Wolbachia* engages these motor proteins is unknown. Surprisingly, none of the known host Kinesin linker proteins are utilized by *Wolbachia*, suggesting *Wolbachia* may interact directly with Kinesin (Russell et al., 2018).

It is after this stage in which *Wolbachia* is distributed throughout the oocyte, that we observe variability among the *Wolbachia* strains, with each falling into one of three distinct classes: two distinct classes in which *Wolbachia* concentrate at the posterior (Posteriorly Localized and Posteriorly Clumped) and one class in which *Wolbachia* fail to exhibit a posterior concentration (Dispersed). The former two classes are distinguished by whether the vast majority (Posteriorly Clumped) or only a small fraction of the *Wolbachia* localize to the posterior pole (Posteriorly Localized). Here, our phylogenomic analyses revealed a strong correlation between *Wolbachia* posterior abundance and *Wolbachia* phylogenetic relationships. Generally, we find that closely related *w*Mel-like *Wolbachia* occur at higher abundance at the oocyte posterior, whereas *w*Ri-like *Wolbachia* exhibit a more dispersed distribution (**Figure 4**). These results suggest that factors intrinsic to *Wolbachia* help determine posterior localization patterns, which is consistent with previous *Wolbachia* transplantation studies demonstrating that posterior localization is determined by *Wolbachia* rather than host factors (Poinsot et al., 1998; Veneti et al., 2004; Serbus and Sullivan, 2007). Below, we discuss how specific factors might contribute to variation in *Wolbachia* localization patterns.

The Posteriorly Localized class, in which only a small fraction localize to the posterior pole, can be explained by previous work demonstrating that *w*Mel is a weak competitor for Kinesin; both *Wolbachia* and germplasm components rely on Kinesin for transport to the posterior pole (Russell et al., 2018). It is thought that *Wolbachia* has evolved to be a weak competitor because interference with the transport of essential germplasm components and thus germplasm formation may be disadvantageous for *Wolbachia* proliferation. The other posterior class (Posteriorly Clumped) is strikingly similar to that observed for *w*Mel in *D. melanogaster* in which the plus-end motor protein Kinesin is over-expressed and excess amounts of *Wolbachia* are transported to the posterior pole (Russell et al., 2018). It may be that the Posteriorly Clumped strains are much better competitors for host Kinesin. Alternatively, these strains could occur in host species with naturally much higher levels of host Kinesin, although our phylogenetic analysis did not support a role of host factors. Functional work will be required to formally test these hypotheses. In particular, it would be interesting to investigate whether there are instances in which *Wolbachia* disrupts the A-P axis (anterior-posterior) because of its accumulation at the posterior pole.

The posterior pole concentration among *Wolbachia* strains may also differ due to variability in the ability of *Wolbachia* to stably associate with the pole plasm. Previous studies demonstrated that *Wolbachia* posterior pole concentration requires intact pole plasm (Serbus and Sullivan, 2007). For example, in *oskar* mutations, a key pole plasm determinant, *w*Mel *Wolbachia* fails to accumulate at the posterior region (Hadfield and Axton, 1999; Serbus and Sullivan, 2007). As described, the oocyte experiences tremendous cytoplasmic streaming requiring posterior components to be anchored directly or indirectly to the cortex (Quinlan, 2016). This variability could be due to differences in the ability of *Wolbachia* to compete with other pole plasm components such as mitochondria for binding sites. Alternatively, the host species may vary in the extent to which their pole plasm accommodates *Wolbachia*. These alternatives can be explored through trans-infection studies.

It is likely that the composition of *Wolbachia* surface proteins play a major role in the extent of
posterior localization. Our phylogenetic analyses revealed that the gene trees of two candidate
surface proteins, particularly WD_1085, are strongly associated with *Wolbachia* localization
patterns. The WD_1085 protein has sequence and structural homology to the bacterial outer
membrane protein BamA, which consists of a large periplasmic domain attached to a 16-stranded
β-barrel domain (**Supplemental Figure 11**). BamA is the central subunit of the β-barrel
assembly machinery (BAM), which is essential for outer membrane protein biogenesis (Noinaj et
al., 2013; Doyle and Bernstein 2019). Given this finding, it is intriguing that we find
overexpression of BamA is sensitive to disruptions in microtubules, as these cytoskeletal
elements are essential for outer membrane biogenesis in mitochondria (Mado et al., 2019). It
may be that microtubules are also required for *Wolbachia* membrane biogenesis. While we
highlight these loci as potential candidates, we note that our results should be interpreted with
caution since the WD_1085 gene could be in linkage with other causal loci. Nonetheless,
Wolbachia surface proteins are generally considered to be strong candidates for interactions with
host cells (Werren et al., 2008; Baldridge et al., 2016; Hague et al., 2022).

Interestingly, our phylogenomic analyses did not find a correlation between host phylogenetic
relationships and *Wolbachia* localization patterns, suggesting that host-associated factors are less
important in determining *Wolbachia* posterior localization. This result contrasts with
transplantation studies that indicate hosts may also play a significant role in determining
Wolbachia abundance (Veneti et al., 2004). In addition, genome-wide screens reveal host factors
play a major role in determining *Wolbachia* intracellular abundance (White et al., 2017; Grobler
et al., 2018). Thus, variation in the host proteins that *Wolbachia* engages could still plausibly
influence *Wolbachia* localization patterns. This would contrast with other endosymbionts such as
Listeria, which relies on a surface protein that binds and polymerizes host actin to propel the
bacteria within and between host cells (Kühn and Enninga, 2020). Because the interaction between
ActA and actin is essential for cell-to-cell transmission, natural variants of this interaction and
transmission strategy have not been discovered. In addition to *Wolbachia* and host factors,
nutrients and other environmental factors have also been shown to influence *Wolbachia* abundance
and localization in host oocytes for closely related *w*Mel-like *Wolbachia* strains (Serbus et al.,
2015; Hague et al., 2020a; Hague et al., 2022).

Two classes in which *Wolbachia* concentrates in the posterior pole (Posteriorly Localized and
Posteriorly Clumped) provide a ready explanation for the cellular mechanisms by which it is
vertically transmitted through generations. In every generation, *Wolbachia* targets the site of
germline formation in the developing oocyte. This also explains why strains exhibiting this pattern,
such as *w*Mel, exhibit efficient transmission strategies under typical laboratory conditions (Hague
et al., 2022). The third, Distributed class, that exhibits no-to-low *Wolbachia* in the posterior
germplasm of the mature oocyte is puzzling. This would be expected to result in embryos lacking
Wolbachia in the germline. However, within hours after fertilization, *Wolbachia* is clearly present
in the germlines of both sexes (**Figure 6**). This implies that either these *Wolbachia* strains are
capable of invading the germline from neighboring somatic cells or a few undetected *Wolbachia*
are delivered late in oogenesis to the posterior pole. Support for the first hypothesis comes from
the fact that in filarial nematodes *Wolbachia* germline invasion from the soma has been directly
observed (Landmann et al., 2012). While germline invasion via cell-to-cell transmission has not
been directly observed in insects, a number of lines of evidence indicate that it likely occurs. In
Drosophila cell culture, *Wolbachia* efficiently undergoes cell-to-cell transmission (White et al.,

2017; White et al., 2017). *Wolbachia* injected into the abdomen of adult *Drosophila* females
migrate to and occupy the germline and follicle stem cells (Frydman et al., 2006). Collections
directly from nature reveal some strains with developing oocytes lacking *Wolbachia* (Casper-
Lindley et al., 2011). In these strains, all of the mature oocytes were infected, suggesting an
alternate route of infection via neighboring somatic cells.

As previously hypothesized, given the posterior follicle cells are directly adjacent to the pole
plasm, we propose *Wolbachia* present in these follicle cells may be the source of germline
Wolbachia (Kamath et al., 2018). Our results are in accord with and complement previous studies
demonstrating *Wolbachia* concentrated in the posterior polar follicle cells in a number of
Wolbachia-infected species (Kamath et al., 2018). For example, in *w*Mau-infected *D. mauritiana*,
a Dispersed class, there is a striking concentration of *Wolbachia* in the posterior follicle pole cells
directly adjacent to the pole plasm. Thus, *Wolbachia* in these somatic cells are well positioned to
invade the pole plasm becoming incorporated into the germline of the next generation (**Figures 7**
and 8). In contrast, in the Posteriorly Localized and Clumped strains, we found no to few
Wolbachia in the posterior polar follicle cells. We note that invasion of *Wolbachia* from posterior
follicle cells into the germline would require *Wolbachia* to cross the seemingly impenetrable newly
formed vitelline membrane, which could suggest alternative sources of germline *Wolbachia* (we
thank the reviewer for this insight). Nonetheless, taken together the data strongly argue for a
somatic-to-germline route of vertical transmission in some *Drosophila* species.

Among the *Wolbachia* strains examined, *w*Mel stands out as an exception. Not only is it
concentrated at the posterior pole of the oocyte, but there is also a large *Wolbachia* concentration
in the posterior polar follicle cells. Thus, *w*Mel may maintain both robust germline-to-germline
and soma-to-germline modes of transmission. Accordingly, it has a high efficiency of vertical
transmission under laboratory conditions (but see Hague et al. 2022). Notably, we found that *w*Mel
and *w*Mau are also both quite prevalent in the more anteriorly located follicle cells of stage 10
oocytes (**Figure 7**), raising the possibility that vertical transmission could even occur without
Wolbachia ever directly associating with the host germline (i.e., soma-to-soma transmission;
Figure 8).

Surprisingly, literature surveys reveal that germline invasion of endosymbionts from the soma
every generation is the most common form of vertical inheritance (Russell et al., 2019). Vertical
transmission via a continuous presence in the germline is much less common. One explanation is
that hosts maintains mechanisms preventing endosymbiont occupation during the formative stages
of germline development. In instances in which the endosymbiont is maintained in the germline,
this may evolve into an obligate relationship in which development of the germline depends on
the presence of the endosymbiont (Sullivan, 2017). Examples of this include the leafhopper (*E.*
plebeus), in which the endosymbiont is required for normal embryonic development (Sander,
1968). In addition, removal of *Wolbachia* from some insects results in increased apoptosis and
abnormal oocyte development (Dedeine et al., 2001; Pannebakker et al., 2007). Here we find that
Wolbachia exhibit evidence of both the soma-to-germline and germline-to-germline transmission
strategies (**Figure 8**), suggesting that *Wolbachia* may stand out as a rarity among endosymbionts.

MATERIALS AND METHODS

Drosophila **Stocks**

All stocks were grown on standard brown food (Sullivan et al., 2000) at 25°C with a 12h light/dark
cycle. Uninfected *Drosophila* stocks were generated by tetracycline-curing of the infected stock
(Serbus et al., 2015). Listed in Table 1 are the 19 different *Wolbachia* strains we examined, which
infect 18 different *Drosophila* host species. This included six *w*Mel-like strains (Cooper et al.,
2019; Hague et al., 2020a), five *w*Ri-like strains (Turelli et al., 2018), seven other A-group
Wolbachia, and the B-group strain *w*Mau that diverged from A-group *Wolbachia* up to 46 million
years ago (Meany et al., 2019). Each genotype was generated as an iso-female line by sampling a
single gravid female from the field and placing her individually into a vial.

Ovary, embryo and larva fixation and staining

Newly eclosed flies were transferred to fresh food for 3-5 days for aging. Approximately 10
females were dissected for each slide preparation. Ovaries were fixed using a modification of
previously published procedures (Brendza et al., 2000; Russell et al., 2018). Ovaries were removed
and separated in phosphate buffer solution and then fixed in a 200 μl devitellinizing solution (2%
paraformaldehyde and 0.5% v/v NP40 in 1x PBS) mixed with 600 μl heptane for 20 min at room
temperature, on a shaker. After removal of the organic layer by brief centrifugation for sample
isolation, the ovaries were washed 3 times with PBS-T (0.1% Triton X-100 in 1x PBS) along with
three five-minute washes. Samples were treated with RNAse A (10 mg/ml) and left overnight at
room temperature. Samples were then washed again a few times with PBS-T and then stained with
a dilute solution of phalloidin for actin staining on a rotator for 1 hour. Samples were washed again
with 3 quick PBS-T rounds and then solution changes with PBS-T over two hours. 60uL of
propidium iodide (PI) in mounting media was added to the sample after removal of the wash
solution and left again overnight. Ovaries were then mounted and carefully separated out again for
ease of imaging and removal of excess mature eggs. Slides were coated in nail polish and stored
at -20°C until imaged.

Embryos were collected for 1-3 hours on plates and fixed in equal volume 32% paraformaldehyde
and heptane and fixed as previously described (Sullivan et al., 2000). After extraction with
methanol, embryos were blocked for one hour in 5% PBST-BSA (2.5g bovine serum albumin
fraction V in 50ml PBS-T) at room temperature. Embryos were then incubated with anti-WD_0009
or anti-Ftfz at 1:500 overnight at 4°C. After three washes with 1% PBST-BSA over one hour,
embryos were incubated in Alexa 488 Anti-Rabbit at 1:500 for one hour at room temperature.
Samples were then washed three more times over one hour with 1% PBST-BSA. Embryos were
mounted with media containing DAPI and stored at -20°C until imaged.

Larval fat bodies were dissected and fixed from 3rd instar larvae using a modified version of a
published protocol (Maimon and Gilboa, 2011). Fat bodies were dissected in 1X PBS then fixed
in 5% paraformaldehyde in PBS for 20 minutes with gentle agitation. Fat bodies were washed for
five minutes, 10 minutes, then 45 minutes with cold 1% NP40 in PBS. After washes, fat bodies
were blocked in 5% PBST-BSA (2.5g bovine serum albumin fraction V in 50ml PBS-T) for one
hour at room temperature. Fat bodies were then incubated with anti-FtsZ at 1:500 overnight on a
rocker at 4°C, washed three times over one hour with 1% PBST-BSA, then incubated with Alexa
488 Anti-Rabbit at 1:500 at room temperature for one hour. After three additional washes over one
hour with 1% PBST-BSA, fat bodies were mounted onto a slide with media containing DAPI.
Larval ovaries were isolated from fat bodies using 0.1mm needles before sealing the slide with
nail polish. Samples were then stored at -20°C.

Confocal microscopy

Imaging was performed on an inverted Leica DMI6000 SP5 scanning confocal microscope. Optical sections of a Nyquist value of 0.38 were used. A variety of zooms were used to optimize image viewing, with most being set at 1.5X. Propidium iodide was excited with the 514 and 543 nm lasers, and emission from 550 to 680 nm was collected. GFP was imaged with the 488 nm laser, and emission from 488 to 540 nm was collected. Alexa 633 was imaged with the 633 laser, and emission from 606 to 700 nm was collected. All imaging was performed at room temperature. Images were acquired with Leica Application Suite Advanced Fluorescence software.

***Wolbachia* quantification and analysis**

Following Russell et al. (2018), we used the polygon selection tool to select three different regions of the oocyte (see Fig. S1 in Russell et al. 2018): the whole oocyte, the posterior region, and the posterior cortex. The "area" and "integrated density" were measured for each region. Additionally, the average of a quadruplicate measure of "mean gray value" beside the stained oocytes was measured as background fluorescence. The Corrected Total Cell Fluorescence (CTCF) was calculated as integrated density – (area x background mean gray value) for each region.

All statistical analyses were performed in R. We first tested whether our three qualitative classes (Posteriorly Clumped, Posteriorly Localized, Dispersed) differed significantly in quantitative estimates of *Wolbachia* fluorescence (CTCF). We used a one-way ANOVA to test if the three groups differed based on their mean CTCF values from the whole oocyte, the posterior region, and the posterior cortex. Distribution and leverage analyses indicated that a transformation of $x' = \log(x + 1)$ was needed to meet assumptions of normality. We tested for significance using an F test with type III sum of squares using the "Anova" function in the *car* package (Fox and Weisberg, 2019). We conducted Tukey multiple comparisons among the three groups using the "TukeyHSD" function.

Genomic data

In order to perform the phylogenomic analysis and generate phylograms, we obtained *Wolbachia* sequences from publicly available genome assemblies, which included *w*Mel (Wu et al., 2004), *w*Ri (Klasson et al., 2009), *w*Ha (Ellegaard et al., 2013), *w*Aura, *w*Tria, *w*Pan (Turelli et al., 2018), *w*Ana (Salzberg et al., 2005), and *w*Rec (Metcalf et al., 2014). All other *Wolbachia* sequences included in this study (*w*Bai, *w*Bic, *w*Boc, *w*Burl, *w*Cha, *w*Curt [*w*Tsa], *w*Nik, *w*Seg, *w*Tris, *w*Leo) were obtained using Illumina sequencing and previously described methods (Hague et al., 2020b). Briefly, tissue samples for genomic data were extracted using a DNeasy Blood & Tissue kit (Qiagen). DNA was cleaned using Agencourt AMPure XP beads (Beckman Coulter, Inc.) following manufacturers' instructions, and eluted in 50 µl 1 × TE buffer for shearing. DNA was sheared using an E220 Focused Ultrasonicator (Covaris Inc.) to a target size of 400 bp. We prepared libraries using NEBNext® Ultra™ II DNA Library Prep with Sample Purification Beads (New England BioLabs). We indexed samples using NEBNext® Multiplex Oligos for Illumina® (Index Primers Set 3 & Index Primers Set 4), and 10 µL of each sample were shipped to Novogene (Sacramento, CA, USA) for sequencing using Illumina HiSeq 4000, generating paired-end 150 bp reads.

We obtained publicly available host sequences for *D. melanogaster* (Hoskins et al., 2015), *D.*
simulans (Hu et al., 2013), *D. ananassae* (Clark et al., 2007), *D. pandora* (Turelli et al., 2018), *D.*
mauritiana (Meany et al., 2019), *D. auraria, D. triaurari, D. baimaii, D. bicornuta* (aff.
bicornuta), *D. bocki*, *D. burlai*, *D. chauvacae* (cf. *chauvacae*), *D. curta* (aff. *tsacasi*), *D. nikananu*,
D. seguyi, *D. tristis*, and *D. leontia* (Conner, 2021). A *D. recens* genome assembly was kindly
provided by Kelly Dyer and Rob Unckless. We note that three of the *montium* host species listed
above have had recent updates to their taxonomic nomenclature: *D. bicornuta* has been renamed
to aff. *bicornuta*, *chauvacae* has been renamed to cf. *chauvacae*, and *curta* has been renamed to
aff. *tsacasi*. See Conner et al. (2021) for a full discussion of *montium* species names and
relationships.

***Wolbachia* phylogenetic analysis**

Raw Illumina reads from the newly sequenced *Wolbachia* strains were trimmed using Sickle
version 1.33 (Joshi and Fass, 2011) and assembled using AbySS version 2.0.2 (Jackman et al.,
2017). *K* values of 71, 81, and 91 were used, and scaffolds with the best nucleotide BLAST
matches to known *Wolbachia* sequences with *E*-values less than 10^{-10} were extracted as the draft
Wolbachia assemblies. For each strain, we chose the assembly with the highest N50 and the fewest
scaffolds (**Supplemental Table 2**). The newly assembled genomes and the previously published
assemblies were annotated using Prokka version 1.11, which identifies homologs to known
bacterial genes (Seemann, 2014). To avoid pseudogenes and paralogs, we only used genes present
in a single copy with no alignment gaps in all of the genome sequences. Genes were identified as
single copy if they uniquely matched a bacterial reference gene identified by Prokka. By requiring
all homologs to have identical length in all of the *Wolbachia* genomes, we removed all loci with
indels. A total of 66 genes totaling 43,275 bp met these criteria. We then estimated a Bayesian
phylogram using RevBayes 1.0.8 under the GTR + Γ + I model partitioned by codon position
(Höhna et al., 2016). Four independent runs were performed, which all converged on the same
topology. All nodes were supported with Bayesian posterior probabilities of 1.

Similar methods were used to generate gene trees for candidate *Wolbachia* loci putatively involved
in host interactions during oogenesis. This included the three *wsp* paralogs (WD_1063
[*wsp*/RS04815], WD_0009 [*wspB*/RS00060], WD_0489 [*wspC*/RS06475]) and three other
Wolbachia surface proteins (WD_1085 [bamA/RS04910], WD_0501 [p44/MSP2/RS0225],
WD_1041 [peptidase M2/RS04710]). Here, the WD_XXXX locus tags refer to the *w*Mel genome
assembly (Wu et al., 2004).The protein-coding sequences for each locus were extracted from the
assemblies using BLAST and the *w*Mel reference sequences. Notably, we found evidence of
insertions in the WD_0009 (*wspB*) sequences of *w*Boc, *w*Leo, and *w*Bic (see Results). We removed
these large insertions from the alignment to generate the *wspB* gene tree. In addition, BLAST
identified WD_0489 (*wspC*) in *w*Mel-like *Wolbachia* and closely related *w*Bai, *w*Leo, and *w*Boc,
but yielded no hits in the other *Wolbachia* strains (*w*Nik, *w*Tsa, *w*Pan, *w*Ana, *w*Tria, *w*Aura, *w*Ri,
*w*Mau), indicating that *wspC* is either too diverged from the *w*Mel reference for BLAST
recognition or alternatively *wspC* is only present in *w*Mel-related *Wolbachia* strains. Sequences
for each gene were aligned with MAFFT 7 (Katoh and Standley, 2013). We then used RevBayes
and the GTR + Γ + I model partitioned by codon position to generate gene trees for each locus.

Host phylogenetic analysis

Host phylogenies were generated using the same nuclear genes implemented in Turelli et al.
(2018): *aconitase, aldolase, bicoid, ebony, enolase, esc, g6pdh, glyp, glys, ninaE, pepck, pgi, pgm,*
pic, ptc, tpi, transaldolase, white, wingless, and *yellow*. We used BLAST with the *D. melanogaster*
coding sequences to extract orthologs from the genomes of each host species. Sequences were then
aligned with MAFFT 7. Finally, we used RevBayes and the GTR + Γ + I model partitioned by
codon position and gene to accommodate potential variation in the substitution process among
genes, as described in Turelli et al. (2018). Our initial phylogenetic analysis produced a polytomy
in the *melanogaster* species group regarding the relationships among the *montium, melanogaster,*
and *ananassae* subgroups (**Supplemental Figure 10**). These relationships have been resolved in
previous genomic studies involving additional taxa (Turelli et al., 2018; Suvorov et al., 2022);
therefore, we generated a constrained host tree in RevBayes that enforced previously defined
relationships among the three subgroups: ((*montium, melanogaster*), *ananassae*).

Tests for phylogenetic signal

The resulting phylograms were used to test whether cellular *Wolbachia* abundance in oocytes
exhibits phylogenetic signal on either the *Wolbachia* or host phylogenies. We used our quantitative
estimates of *Wolbachia* abundance to test for phylogenetic signal using Pagel's lambda (λ) (Pagel,
1999). Here, we used total *Wolbachia* fluorescence in whole oocytes, the posterior region, and the
posterior cortex (log-transformed CTCF) as continuous characters to calculate maximum
likelihood values of Pagel's λ. A Pagel's λ of 0 indicates that character evolution occurs
independently of phylogenetic relationships, whereas $\lambda = 1$ is consistent with a Brownian motion
model of character evolution. We used the "fitContinuous" function in GEIGER (Harmon et al.,
2008) and a likelihood ratio test to compare our fitted value to a model assuming no phylogenetic
signal ($\lambda = 0$). We also used a Monte Carlo-based method to generate 95% confidence intervals
surrounding our estimates using 1,000 bootstrap replicates in the *pmc* package (Boettiger et al.,
2012). When applicable, we conducted additional analyses to evaluate whether the number of taxa
in our phylogeny ($N = 19$ *Wolbachia* strains, $N = 18$ host species) limited our ability to detect
significant departures from $\lambda = 0$ (Hague et al., 2020b; Hague et al., 2021). Small phylogenies are
likely to generate near-zero values simply by chance, not necessarily because the phylogeny is
unimportant for trait evolution (Boettiger et al., 2012). To evaluate whether larger phylogenies
increase the accuracy of estimation, we simulated trees with an increasing number of *Wolbachia*
strains/host species ($N = 25, 50$) and our empirical estimates using the "sim.bdtree" and "sim.char"
functions in the *geiger* R package (Harmon et al., 2008). We then re-estimated confidence intervals
using the larger simulated trees.

Our tests for phylogenetic signal using the constrained host phylogram are presented in the main
text; however, we note that our analysis using the unconstrained host tree (with a polytomy at the
base of the *melanogaster* species group) produced similar results. Using the unconstrained tree,
we found that *Wolbachia* abundance in whole oocytes ($\lambda = 0.782$ [0, 0.987], $P = 0.469$), the
posterior region ($\lambda = 0.951$ [0, 1], $P = 0.377$), and the posterior cortex ($\lambda < 0.001$ [0, 0.563], $P = 1$)
1) did not exhibit significant phylogenetic signal.

Structural analysis of *Wolbachia* candidate genes

701 We used HHPred to identify protein domains of WD_1085 and WD_0501 using the COPe70_2.07,
702 Pfam-A_v33.1, COG_KOG_v1.0, and SMART_v6 databases (Zimmermann et al., 2018; Gabler

et al., 2020). Here we used default setting, including an E-value cutoff for MSA generation of 1e-3, a minimum coverage of MSA hits of 20%, and minimum probability in hitlist of 20%, and a maximum of 250 target hits. As expected, both WD_1085 and WD_0501 yielded multiple hits of varying length for domains associated with outer membrane proteins that form beta barrel structures (results summarized in **Supplemental Table 3**). We used SignalP-6.0 to identify the signal peptide cleavage sites for WD_1085 between aa residues 25 and 26 (Teufel et al., 2022) and removed the signal peptide for homology modeling. SignalP was unable confidently identify the signal peptide sequence for WD_0501, so we used the full sequence for homology modeling. We then used I-TASSER to model the three-dimensional structure of each protein through sequence homology (Yang and Zhang 2015; Zhang et al., 2017). For WD_1085, we used PDB:4K3B with an amino acid sequence identity of 23% in the threaded aligned region as a template (Noinaj et al., 2103). For WD_0501, we used PDB:2MLH with an amino acid sequence identity of 17% in the threaded aligned region as a template (Fox et al., 2014). PyMOL was used for image generation (Schrodinger 2015). Results are displayed in **Supplemental Figure S11**.

Ectopic expression of *Wolbachia* surface proteins in yeast

SC-Leucine yeast media was prepared as described previously (Rose et al., 1990). Where present, benomyl was added to media from a 10mg/ml stock in DMSO to a final concentration of 2.5 ug/ml. The yeast expression assays are a modified version of the protocol described by Sheehan et al. (2016). *Wolbachia* genes were codon optimized for expression in *S. cerevisiae* and synthesized by Twist Bioscience (South San Francisco). The *wsp*, *wspB* and *wspC* genes were gifts of the Free Genes project (https://stanford.freegenes.org/). These genes were inserted into pGH448 using the New England Biolabs Golden Gate kit according to the manufacturer's instruction. This placed the *Wolbachia* genes under control of the *GAL1* promoter and *CYC1* transcriptional terminator. This plasmid, which will be described in more detail elsewhere, carries a CEN/ARS element and also contains the kanamycin resistance and *LEU2* genes for selection in bacteria and yeast respectively.

The indicated plasmids were transformed into yeast strain GHY1934 [genotype] and GHY1551 [genotype]. Both of these strains are isogenic to S288C and are *GAL2+*. Transformed cells were grown to saturation in SC-Leucine liquid media containing 2% glucose. These cells were counted, diluted to 1×10^{-7} cells per ml, five-fold serial diluted and spotted to SC-Leucine plates with either 2% glucose or 2% galactose as the carbon source and incubated at the times and temperatures indicated in the figure legends.

DATA AVAILABILITY STATEMENT

Newly generated genomic data are available on GenBank (BioProject PRJNA907328). All other data are publicly available on Dryad (doi:10.5061/dryad.v6wwpzh0q).

ACKNOWLEDGEMENTS

We thank Dr. Benjamin Abrams (UCSC Life Sciences Microscopy Center, RRID: SCR_021135) for his technical support and assistance with microscopy experiments. We thank Dr. Shelbi Russell for their helpful advice on imaging protocols and analysis. We thank Dr. Irene Newton for kindly supplying the anti-FtsZ antibody. We thank Will Conner for guidance with bioinformatic analyses

and Tim Wheeler for laboratory assistance. We thank Drew Endy and the Free Genes project for

the gift of *Wolbachia* genes and advice on construction of pGH448. Research reported in this

publication was supported by the National Institute of General Medical Sciences of the National

Institutes of Health under Award Number R35GM139595 to W. Sullivan and R35GM124701 to

B.S. Cooper. This work was also supported by the National Science Foundation (NSF) grant

(1456535) to W. Sullivan and an NSF CAREER Award (2145195) to B.S. Cooper. We thank the

Big Creek Reserve for use of their facilities. The microscopic analysis was enabled through the

generous support of the IBSC microscopy facility at UC Santa Cruz.

FIGURES

Figure 1. Confocal micrographs of *D. melanogaster* (*w*Mel) oocytes tracked through early oogenesis. Samples are stained with propidium iodide (PI) for DNA (red) and phalloidin for actin (green). Representative stages of development (1-10) are shown for *w*Mel infected (left panels) and uninfected (right panels). Grayscale zoomed-in images around the nucleus of the oocyte are shown in the third column, with red arrows highlighting *Wolbachia* in the oocyte. Corresponding arrows in the sixth column show the equivalent area in uninfected oocytes. The magenta double arrow in the stage 5 infected oocyte (second column) highlights *Wolbachia* in the nurse cells. Scale bars all set to 25 uM.

Figure 2. Conserved *Wolbachia* anterior localization during early oogenesis (stages 4-5). *Wolbachia* and host DNA stained with PI (red) and actin with phalloidin (green) shown in the left most column. Greyscale of PI channel shown in middle. 4X zoom-in around the oocyte shown on the right. All of the eight *Wolbachia* strains examined concentrate at the anterior cortex of the oocyte at this stage. Red arrows highlight *Wolbachia*'s anterior localization positioning for (A) *w*Pan in *D. pandora*, (B) *w*Nik in *D. nikananu*, (C) *w*Cha in *D. chauvacae*, (D) *w*Ha in *D. simulans*, (E) *w*Tsa in *D. tsacasi*, (F) *w*Boc in *D. bocki*, (G) *w*Ana in *D. ananassae*, and (H) *w*Mel in *D. melanogaster*. Oocytes are approximately 50-75 uM in size.

Figure 3. Three distinct patterns of posterior *Wolbachia* localization during late stage oogenesis (A-C). Confocal micrographs of *Drosophila* oocytes DNA-stained with PI (red) and actin-stained with phalloidin (green) show representative examples of (A) Posteriorly Localized *w*Tris in *D. tristis*, (B) Posteriorly Clumped *w*Bic in *D. bicornuta*, and (C) Dispersed localization of *w*Tria in *D. triauraria*. Second row depicts single channel images of PI staining. Third row depicts an enlarged PI-stained image of the posterior region of each oocyte. Scale bars set at 25uM.

Figure 4. *Wolbachia* abundance in the posterior region of host oocytes exhibits phylogenetic signal on the *Wolbachia* phylogeny. (A) Estimated Bayesian phylogram for the 19 A- and B-group *Wolbachia* strains included in tests for phylogenetic signal. (B) Mean estimates of *Wolbachia* abundance (log-transformed CTCF) in the whole oocyte, posterior region, and the posterior cortex. *Wolbachia* abundance in the posterior region has significant phylogenetic signal ($\lambda = 0.982$ [0, 0.998], $P = 0.021$), suggesting *Wolbachia*-associated factors determine *Wolbachia* abundance in the oocyte posterior. (C) Estimated Bayesian phylogram of *Drosophila* host species using 20 conserved single-copy genes. *Wolbachia* and host divergence times in millions of years (MYA) are reproduced from Meany et al. (2019) and Suvorov et al. (2022), respectively. The *Wolbachia* and host tree are generally discordant (see **Supplemental Figure 2**), as expected with frequent horizontal *Wolbachia* acquisition (Turelli et al. 2018).

Figure 5. *Wolbachia*'s presence around the 3rd instar larval germline, in multiple *Drosophila* hosts. *w*Mel *Wolbachia* stained with anti-FtsZ can be seen for *D. melanogaster*, in panel (A) shown in pink. Cells are marked with DAPI (blue). (A') Grayscale channel of anti-FtsZ staining. (B) Uninfected ovary for *D. melanogaster*, (B') grayscale channel for uninfected antibody binding. Germline location marked with anti-Vasa shown in green (C), grayscale channel (C'). *w*Mau *Wolbachia* localization in the ovary for *D. mauritiana* shown in panel (D), grayscale channel (D'). *w*Ri and *D. simulans* shown in panel (E), grayscale (E'). Panel (F), *w*Bic and *D. bicornuta*, (F') grayscale channel. Scale bar shown at 25 uM.

Figure 6. *Wolbachia*'s presence in the newly formed germline cells of late blastoderm and cellularized embryos**.** (A) 1 hour *D. mauritiana* (*w*Mau) embryo stained with PI. (A') Greyscale of PI channel. (B) staining of 2-3 hour developed *D. mauritiana* (*w*Mau) embryo with PI. (B') PI of greyscale channel. (C) PI and anti-Vasa (green) staining of *w*Mel *Wolbachia* in 2-3 hour *D. melanogaster* embryo. (C') Greyscale of PI channel. (D) Anti-FtsZ/DAPI staining for *w*Ri *Wolbachia* in *D. simulans*. (D') Greyscale of anti-FtsZ channel. Scale bars set at 25uM.

Figure 7. The A-group *w*Mel and B-group *w*Mau *Wolbachia* strains are frequently found in the polar follicle cells, as well as other follicle cells. (A-C) Schematic of categories used for description of follicle cell *Wolbachia*. (A) Dispersed localization pattern shown, both polar cells infected, with *Wolbachia* broadly present in follicle cells. (B) Posteriorly Localized pattern shown, one polar cell infected, with *Wolbachia* broadly present in follicle cells. (C) Posteriorly Clumped pattern shown, with no *Wolbachia* present in the polar cells or other follicle cells. (D) *Wolbachia* quantification in the polar follicle cells during oogenesis stages 9-10. *Wolbachia* were quantified in midsections of each oocyte that displays unique posterior localization patterns: Dispersed, Posteriorly Localized, and Posteriorly Clumped. For each strain, the count is shown for the number of oocytes observed with *Wolbachia* present in both polar follicle cells (orange), only one polar cell (blue), and neither polar cell (grey). (E) Counts of oocytes observed with *Wolbachia* present anywhere in the follicle cells (pink) or with *Wolbachia* absent (grey). (F) Representative anti-FtsZ staining of *w*Mau in *D. mauritiana* (Dispersed class) present in both polar follicle cells (corresponding to orange labels above). (G) Representative staining of *w*Mel in *D. melanogaster* (Posteriorly Localized class) present in only one polar cell (blue labels above). (H) Representative staining of *w*Bic in *D. bicornuta* (Posteriorly Clumped class) absent in both polar cells (grey labels above). (I) Representative staining of *w*Mel in *D. melanogaster* (Posteriorly Localized class) broadly present in anterior follicle cells (pink labeling above). (J-M) Greyscale anti-FtsZ channel of respective composite images above. (N-Q) 5X zoom of polar cells, with circles highlighting representative staining in the area. Scale bar set at 25uM.

Figure 8. Multiple routes of *Wolbachia* vertical transmission. (A) In *Wolbachia* strains that concentrate at the posterior pole plasm, vertical transmission likely occurs through a continuous maintenance in the germline. (B) In strains where no or few *Wolbachia* localize to the posterior pole plasm, vertical transmission likely occurs through *Wolbachia* invading the germline from neighboring somatic cells**. (**C) In theory, vertical transmission could occur through a symbiont invading the oocyte from neighboring somatic cells never associating with the germline.

TABLES

Table 1. Qualitative and quantitative quantification of cellular *Wolbachia* abundance in stage 10 oocytes. The number of confocal images (*N*) used to generate means is shown for each *Wolbachia*-infected host species. *Wolbachia* abundance was measured in each region of the oocyte as corrected total cell fluorescence (CTCF).

Wolbachia strain	Host species	Localization class	*N*	Whole oocyte mean CTCF	Posterior mean CTCF	Posterior cortex mean CTCF
*w*Rec	*D. recens*	Clumped	3	482174	300753	58453
*w*Tris	*D. tristis*	Localized	4	223048	129013	105641
*w*Seg	*D. seguyi*	Dispersed	5	58552	11402	557
*w*Cha	*D. chauvacae*	Localized	5	108588	42636	9786
*w*Burl	*D. burlai*	Localized	4	72806	53754	36944
*w*Mel	*D. melanogaster*	Localized	2	191451	136174	92341
*w*Bic	*D. bicornuta*	Clumped	3	90114	83621	54900
*w*Ri	*D. simulans*	Dispersed	5	333505	90664	20494
*w*Pan	*D. pandora*	Localized	3	45423	6486	1890
*w*Ana	*D. ananassae*	Localized	4	41578	16843	12927
*w*Tria	*D. triauraria*	Dispersed	2	487897	1809	345
*w*Aura	*D. auraria*	Dispersed	5	266478	3699	645
*w*Ha	*D. simulans*	Dispersed	5	82941	3488	807
*w*Bai	*D. baimaii*	Dispersed	2	11359	0	0
*w*Boc	*D. bocki*	Clumped	6	137664	98339	82408
*w*Leo	*D. leontia*	Clumped	4	997045	685260	713918
*w*Tsa	*D. tsacasi*	Localized	4	61314	36478	20620
*w*Nik	*D. nikananu*	Localized	3	44764	27099	13014
*w*Mau	*D. mauritiana*	Dispersed	3	69362	799	322

SUPPLEMENTAL INFORMATION

Supplemental Figure 1. (A) Representative image of the reduction of *w*Mel *Wolbachia* titer between stages 4/5 and 6 in *D. melanogaster* oogenesis. *Wolbachia* stained with anti-FtsZ shown in pink. Scale bars set at 25 uM. (B) Greyscale DAPI channel. (C) Greyscale of anti-FtsZ channel. (A'-C') 5X zoom around the nucleus of the oocyte during stage 4 shown for each channel. (A''-C'') 4X zoom around the nucleus during stage 6 for each respective channel seen above. At least 10 oocytes across 3 slides were imaged with Z-stacks for a visual description of this trend.

Supplemental Figure 2. Bayesian phylograms depicting the evolutionary relationships among *Wolbachia* strains diverged up to 46 million years (left) and *Drosophila* host species diverged up to 50 million years (right). *w*Mel- and *w*Ri-like clades of closely related *Wolbachia* are labeled on the left. Grey lines pair *Wolbachia* strains with their *Drosophila* host species, which highlights patterns of topological discordance due to introgressive and horizontal transfer of *Wolbachia* among host species. The *Wolbachia* phylogram was estimated using 66 full-length and single-copy genes (43,275 bp) of equal length. All nodes have posterior probabilities >0.95. The host phylogram was estimated using 20 conserved single-copy genes (see Materials and Methods regarding host tree topology). *Wolbachia* and host divergence times in millions of years (MYA) are reproduced from Meany et al. (2019) and Suvorov et al. (2022), respectively.

Supplemental Figure 3. Representative images of *Wolbachia* posterior localization patterns. Confocal micrographs of *Drosophila* oocytes DNA-stained with PI (red) and actin-stained with phalloidin (green) show representative examples of infected *Drosophila* species with different *Wolbachia* localization classes. Second columns depict single greyscale channel images of PI staining. Third columns depict an enlarged PI-stained image of the posterior region of each oocyte. Panels A, B and C are grouped into *Wolbachia* strains that exhibit a Posteriorly Localized, Posteriorly Clumped, or Dispersed localization pattern respectively. Scale bar set at 25 uM.

A

B

C

Supplemental Figure 4. Cellular *Wolbachia* abundance in stage 10 oocytes measured as *Wolbachia* fluorescence due to propidium iodide (CTCF). Points are color coded by *Wolbachia* localization class (Posteriorly Clumped, Posteriorly Localized, Dispersed), with unique shapes indicating each *Wolbachia* strain and host species. Asterisks indicate significant differences among groups based on one-way ANOVAs and Tukey comparisons using *P*-values <0.05 adjusted for multiple comparisons. Above, a schematic is shown for fluorescent quantification of different regions of the oocyte using the PI greyscale channel: the whole oocyte, the posterior region, and the posterior cortex.

Supplemental Figure 5. Distribution of maximum likelihood estimates of λ from 1,000 bootstrap replicates based on the *Wolbachia* phylogram and *Wolbachia* abundance at the oocyte posterior cortex (log-transformed CTCF). The bootstrap analysis of our *Wolbachia* phylogram is shown to the left ($N = 19$ *Wolbachia* strains). To the right are simulated phylogenies with an increasing number of *Wolbachia* strains included ($N = 25, 50$). For simulated trees, character evolution was simulated with our λ estimate of 0.959 using the "sim.bdtree" and "sim.char" functions in the *Geiger* R package (Harmon et al. 2008). For each graph, fitted λ values for the original phylogeny are shown above with a vertical dashed line. Note that fitted λ values for the simulated phylogenies differ slightly from $\lambda = 0.959$, because "sim.char" uses a Brownian-motion model to simulate character evolution along the phylogeny. Below each graph, the mean estimate of λ from the 1,000 replicates ($\bar{\lambda}$) is shown with associated 95% confidence intervals. Small phylogenies (e.g., $N = 19$) are likely to generate many near-zero λ values by chance, not necessarily because the phylogeny is unimportant for trait evolution (Boettiger et al. 2012). As the number of strains in our analysis increases ($N = 25, 50$), bootstrapped estimates of λ cluster around the true λ value fitted to the original phylogeny.

Supplemental Figure 6. Gene trees of *Wolbachia* surface proteins. All nodes have Bayesian posterior probabilities of >0.95 unless otherwise noted (see only *wspC*).

Supplemental Figure 7. To gain insight into WD_0501 (RS0225/p44/Msp2) and WD_1085 (RS04910/bamA) cellular function, we ectopically expressed these genes in yeast using a Galactose inducible promoter. Under normal growth conditions, ectopic expression of both genes inhibited growth, with WD_1085 (RS04910/bamA) exhibiting a more pronounced inhibition. Growth inhibition of WD_1085 was dramatically increased when the integrity of microtubules was compromised with the yeast placed in the mad1 spindle assembly checkpoint background or the microtubules were compromised directly through the addition of benomyl. These results suggest that WD_1085 may directly or indirectly interact with host microtubules.

Supplemental Figure 8. Full amino acid alignment of *wspB* (WD_0009) homologs in *w*Mel and the *Wolbachia* strains with a putatively pseudogenized version of the surface protein. Stop codons are indicated by asterisks and insertions of varying length are indicated by "XXX".

```
wMel reference    MISKKTLAVTAFALLLSQQSFASETEGFYFGSGYYGQYLNNTSVLKTSTTGIKNLSINDR    60
wHa               MISKKTLAVTAFALLLSQQSFASETEGFYFGSRYYGQYLNNTSVLKTSTTGIKNLSINDR    60
wBai              MISKKTLAVTALALLLSQQSFASETEGFYFGSGYYGQFFNGMGELKTSTTDIKNLSINDR    60
wBoc              MISKKTLAVTALALLLSQQSFASETEGFYFGSGYYGQFFNGMGELKTSTTDIKNLSINDR    60
wLeo              MISKKTLAVTALALLLSQQSFASETEGFYFGSGYYGQFFNGMGELKTSTTDIKNLSINDR    60
wBic              MISKKTLAVTAFALLLSQQSFASETEGFYFGSGYYGQYLNNTSVLKTSTTGIKNLSINDR    60

wMel reference    GAQNTEGQSLSEYKGDYNPPFAANVAFGYTGELGNNSYRAELEGMYSSVKVD---NIGLT   117
wHa               GAQNTEGQSLSEYKGDYNPPLLQMWHLVTQENWVTTAIGLNWKGCILL*KWI---ILV*Q   117
wBai              DAQNTKGQSLSEYKGDYNPPL------VIQGNWVTTAIGLNWKGCILL*KWI---ILV*Q   111
wBoc              GAQNTKGQSLSKYKGDYNPPFAANVALGYTGELGNNSCRAELEGMYSSVKVDXXXILV*Q   120
wLeo              GAQNTKGQSLSKYKGDYNPPFAANVALGYTGELGNNSCRAELEGMYSSVKVDXXXILV*Q   120
wBic              GAQNTEGQSLSEYKGDYNPPFAANVAFGYTGELGNNSYRAELEGMYSSVKVD---NIGLT   117

wMel reference    SSQITVSYLKETGEDPDKETYLYSAAVSHDQIENISVMANVYHHWKSDRFSFSPYVGIGI   177
wHa               VAK*LFHT*RRLVRILIKKLISIVLQLVMTKLRTYL*WQMFIIIGKVTVSLFLLTLVLGS   177
wBai              VAK*LFHI*RTLVRMLIKKLISIVLQLIMTKLRTYL*WQMFIIIGKVTVSLFLLTLVLES   171
wBoc              VVK*LFHT*RTLVRMLVKKLVSIVLQLIMTKLKTCL*WQMFIIIGRVTVSLFLLMLGLES   180
wLeo              VVK*LFHT*RTLVRMLVKKLVSIVLQLIMTKLKTCL*WQMFIIIGRVTVSLFLLMLGLES   180
wBic              SSQITVSYLKETGEDPNKETYLYSAAVSHDQIENISVMANVYHNWKSDRFSFSPYVGIGI   177

wMel reference    GATRMTMFEKPSIRPAGQLKAGFDYRINEDVN---MHIGYRGFGAIGSDIKLTAKRLGQV   234
wHa               VQQE*RCLKNRQ*DPQVN*KLALTIA*TKM*I---CISDIEVLVLLVAILSLQQKG*DKW   234
wBai              VQQE*RCLKNRQ*DPQVK*KLALTIT*TKM*I---CISDIEVLVLLVAILSLQQKS*DKW   228
wBoc              VQQE*QCLKNRQ*DPQVN*KLALIIT*TKM*I---CISDIEVLVLLVVLLVMVLNLQRMR   237
wLeo              VQQE*QCLKNRQ*DPQVN*KLALIIT*TKM*I---CISDIEVLVLLVVLLVMVLNLQRMR   237
wBic              GATRMTMFEKPSIRPAGQLKAGFDYRINEDV*XXXCISDIEVLVLLVAILSLQQKG*DKW   237

wMel reference    VDDPNNDKKKK------LNPSSGSKVTEEINIGNQLFHTHGIEAGLTFHFASKA        282
wHa               *TTLIMIKKRS------LILAQVAK*LRK*I*VINYFIHTV*RLVLLSILPAKL        282
wBai              *TTLIMIKKRS------LILAQVAKYSISHNSPNRS*EGRINSTTKGFE*RNV-        275
wBoc              LEKL*MILITGIKRSLMIRVNQQMQQKKI*I*VIHRFTHTV*RLALLSTLPAKL        291
wLeo              LEKL*MILITGIKRSLMIRVNQQMQQKKI*I*VIHRFTHTV*RLALLSTLPAKL        291
wBic              *TTLIMIKKRS------LILAQVAKYSTSRNSPNRIRYVLMAMVEQ*HIMICSD        285
```

Supplemental Figure 9. Predicted *Wolbachia* surface proteins WD_1063 (*wsp*/RS04815), WD_0009 (*wspB*/RS00060), and WD_0489 (*wspC*/RS06475) were assayed for their effect on eukaryotic cellular processes through ectopic expression in yeast. Note that galactose induction of gene expression of *wspC* kills the cells, while expression of *wsp* and *wspB* inhibits growth.

Supplemental Figure 10. (A) Estimated Bayesian phylogram of host species based on 20 nuclear loci. All nodes have posterior probabilities >0.95. (B) Bayesian phylogram estimated using similar methods, but with a constrained topology to account for previously resolved relationships among subgroups within the *melanogaster* species group ((*montium*, *melanogaster*) *ananassae*) based on Turelli et al., (2018) and Suvorov et al., (2022). (C) Mean estimates of *Wolbachia* abundance (log-transformed CTCF) in the whole oocyte, posterior region, and the posterior cortex. For *D. simulans*, the mean values include both *w*Ri- and *w*Ha-infected individuals.

Supplemental Figure 11. Illustrations of the homology models for the *Wolbachia* surface proteins WD_1085 and WD_0501.

Supplemental Table 1. Estimates of phylogenetic signal (Pagel's λ) with 95% confidence intervals (CI) for *Wolbachia* surface proteins.

Locus	Whole Oocyte				Posterior				Posterior Cortex			
	λ	Lower CI	Upper CI	*P*	λ	Lower CI	Upper CI	*P*	λ	Lower CI	Upper CI	*P*
WD_1063 (*wsp*)	0.000	0.000	0.578	1.000	0.945	0.000	0.993	0.071	0.000	0.000	0.633	1.000
WD_0009 (*wspB*)	0.000	0.000	0.401	1.000	0.000	0.000	0.374	1.000	0.279	0.000	0.720	0.655
WD_0489 (*wspC*)	0.210	0.000	0.947	0.659	0.000	0.000	0.845	1.000	0.000	0.000	0.811	1.000
WD_0501	0.000	0.000	0.597	1.000	0.992	0.521	1.000	0.022	0.000	0.000	0.536	1.000
WD_1041	0.000	0.000	0.692	1.000	0.000	0.000	0.555	1.000	0.000	0.000	0.484	1.000
WD_1085	0.000	0.000	0.553	1.000	0.974	0.000	0.998	0.001	0.942	0.000	0.994	0.028

Supplemental Table 2. Scaffold count, N50, and total assembly size of each new *Wolbachia* assembly.

Host species	*Wolbachia*	Scaffold Count	N50	Total Assembly Size
D. baimaii	*w*Bai	209	9004	1135209
D. bicornuta	*w*Bic	140	16267	1192611
D. bocki	*w*Boc	227	7115	1090928
D. burlai	*w*Burl	128	20905	1248003
D. chauvacae	*w*Cha	406	4802	1300201
D. tsacasi	*w*Tsa	59	29597	1132109
D. nikananu	*w*Nik	83	22663	1257108
D. seguyi	*w*Seg	201	8880	1174927
D. tristis	*w*Tris	114	20171	1264866
D. leontia	*w*Leo	186	9149	1096851

Supplemental Table 3. Summary of domain annotation results from HHPred for WD_1085 and WD_0501.

Query name	Number	Hit	Name	Prob	E-value	Aligned cols	Target length	Query residues	Domain length
WD_1085	1	COG4775	BamA; Outer membrane protein assembly factor BamA [Cell wall/membrane/envelope biogenesis]	100	3.10E-63	733	766	25-778	753
WD_1085	2	COG0729	TamA; Outer membrane translocation and assembly module TamA [Cell wall/membrane/envelope biogenesis]	100	2.00E-42	505	594	227-778	551
WD_1085	3	COG2831	FhaC; Hemolysin activation/secretion protein [Intracellular trafficking, secretion, and vesicular transport]	100	2.90E-39	490	554	237-778	541
WD_1085	4	KOG2602	Predicted cell surface protein homologous to bacterial outer membrane proteins [General function prediction only]	100	1.00E-31	433	457	315-778	463
WD_1085	5	PF19143.3	Omp85_2 ; OMP85 superfamily	99.94	2.60E-23	308	360	419-778	359
WD_1085	6	COG4775	BamA; Outer membrane protein assembly factor BamA [Cell wall/membrane/envelope biogenesis]	99.93	6.20E-21	569	766	25-637	612
WD_1085	7	COG4775	BamA; Outer membrane protein assembly factor BamA [Cell wall/membrane/envelope biogenesis]	99.92	2.80E-22	310	766	93-433	340
WD_1085	8	PF01103.26	Omp85 ; Omp85 superfamily domain	99.92	5.40E-22	306	424	454-778	324
WD_1085	9	PF03865.16	ShlB ; Haemolysin secretion/activation protein ShlB/FhaC/HecB	99.87	2.10E-19	285	310	416-742	326
WD_1085	10	COG4775	BamA; Outer membrane protein assembly factor BamA [Cell wall/membrane/envelope biogenesis]	99.61	2.50E-13	232	766	24-275	251
WD_0501	1	PF05736.14	OprF ; OprF membrane domain	99.78	3.00E-16	134	186	159-303	144
WD_0501	2	COG3637	LomR; Opacity protein and related surface antigens [Cell wall/membrane/envelope biogenesis]	99.75	1.20E-15	135	199	166-303	137
WD_0501	3	COG3047	OmpW; Outer membrane protein W [Cell wall/membrane/envelope biogenesis]	99.74	1.80E-15	144	213	159-303	144
WD_0501	4	PF06316.14	Ail_Lom ; Enterobacterial Ail/Lom protein	99.7	1.20E-14	179	199	3-303	300
WD_0501	5	PF19573.2	DUF6089 ; Domain of unknown function (DUF6089)	99.67	3.60E-14	186	226	4-303	299
WD_0501	6	PF13505.9	OMP_b-brl ; Outer membrane protein beta-barrel domain	99.65	8.40E-14	175	209	7-303	296
WD_0501	7	SCOP_d1p4ta_	f.4.1.1 (A:) Outer membrane protein NspA {Neisseria meningitidis [TaxId: 487]}	CLASS: Membrane and cell surface protei	99.64	2.10E-13	154	155	95-303	208
WD_0501	8	PF01389.20	OmpA_membrane ; OmpA-like transmembrane domain	99.62	2.60E-13	173	202	95-303	208
WD_0501	9	SCOP_d1qjpa_	f.4.1.1 (A:) Outer membrane protein A (OMPA) transmembrane domain {Escherichia coli [TaxId: 562]}	CLASS: Membrane and	99.61	4.60E-13	167	171	94-303	209
WD_0501	10	PF03922.17	OmpW ; OmpW family	99.6	3.40E-13	164	201	96-303	207

Supplemental Table 4. Summary of results for *Wolbachia* surface proteins.

Gene	Description	Significant Phylogenetic Signal	Ectopic Expression
WD_1063 (wsp/RS04815)	surface antigen Wsp	None	Growth inhibition
WD_0009 (*wspB*/RS00060)	surface antigen Wsp paralog	None	Growth inhibition
WD_0489 (*wspC*/RS06475)	surface antigen, Wsp paralog	None	Lethality
WD_1085 (RS04910)	outer membrane protein assembly factor BamA	Oocyte posterior, posterior cortex	Growth inhibition and sensitive to microtubule checkpoints/inhibitors
WD_0501 (RS0225)	P44/Msp2 family outer membrane protein	Oocyte posterior	Slight growth inhibition
WD_1041 (RS04710)	peptidase M2	None	Not tested

REFERENCES

BALDRIDGE, G.D., LI, Y.G., WITTHUHN, B.A., HIGGINS, L., MARKOWSKI, T.W., BALDRIDGE, A.S., and FALLON, A.M., 2016, Mosaic composition of ribA and wspB genes flanking the virB8-D4 operon in the *Wolbachia* supergroup B-strain, *w*Str: Arch Microbiol, v. 198, p. 53-69, doi: 10.1007/s00203-015-1154-8.

BASTOCK, R., and ST JOHNSTON, D., 2008, *Drosophila* oogenesis: Curr Biol, v. 18, p. R1082-7, doi: 10.1016/j.cub.2008.09.011.

BILINSKI, S.M., JAGLARZ, M.K., and TWORZYDLO, W., 2017, The Pole (Germ) Plasm in Insect Oocytes: Results Probl Cell Differ, v. 63, p. 103-126, doi: 10.1007/978-3-319-60855-6_5.

BOETTIGER, C., COOP, G., and RALPH, P., 2012, Is your phylogeny informative? Measuring the power of comparative methods: Evolution, v. 66, p. 2240-51, doi: 10.1111/j.1558-5646.2011.01574.x.

BRENDZA, R.P., SERBUS, L.R., DUFFY, J.B., and SAXTON, W.M., 2000, A function for kinesin I in the posterior transport of oskar mRNA and Staufen protein: Science, v. 289, p. 2120-2, doi: 10.1126/science.289.5487.2120.

CARRINGTON, L.B., LIPKOWITZ, J.R., HOFFMANN, A.A., and TURELLI, M., 2011, A re-examination of *Wolbachia*-induced cytoplasmic incompatibility in California *Drosophila simulans*: PLoS One, v. 6, p. e22565, doi: 10.1371/journal.pone.0022565.

CASPER-LINDLEY, C., KIMURA, S., SAXTON, D.S., ESSAW, Y., SIMPSON, I., TAN, V., and SULLIVAN, W., 2011, Rapid fluorescence-based screening for *Wolbachia* endosymbionts in *Drosophila* germ line and somatic tissues: Appl Environ Microbiol, v. 77, p. 4788-94, doi: 10.1128/AEM.00215-11.

CHA, B.J., KOPPETSCH, B.S., and THEURKAUF, W.E., 2001, In vivo analysis of *Drosophila* bicoid mRNA localization reveals a novel microtubule-dependent axis specification pathway: Cell, v. 106, p. 35-46, doi: 10.1016/s0092-8674(01)00419-6.

CHA, B.J., SERBUS, L.R., KOPPETSCH, B.S., and THEURKAUF, W.E., 2002, Kinesin I-dependent cortical exclusion restricts pole plasm to the oocyte posterior: Nat Cell Biol, v. 4, p. 592-8, doi: 10.1038/ncb832.

CLARK, A.G., EISEN, M.B., SMITH, D.R., BERGMAN, C.M., OLIVER, B., MARKOW, T.A., KAUFMAN, T.C., KELLIS, M., GELBART, W., IYER, V.N., POLLARD, D.A., SACKTON, T.B., LARRACUENTE, A.M., SINGH, N.D., ABAD, J.P., ABT, D.N., ADRYAN, B., AGUADE, M., AKASHI, H., ANDERSON, W.W., AQUADRO, C.F., ARDELL, D.H., ARGUELLO, R., ARTIERI, C.G., BARBASH, D.A., BARKER, D., BARSANTI, P., BATTERHAM, P., BATZOGLOU, S., BEGUN, D., BHUTKAR, A., BLANCO, E., BOSAK, S.A., BRADLEY, R.K., BRAND, A.D., BRENT, M.R., BROOKS, A.N., BROWN, R.H., BUTLIN, R.K., CAGGESE, C., CALVI, B.R., BERNARDO DE CARVALHO, A., CASPI, A., CASTREZANA, S., CELNIKER, S.E., CHANG, J.L., CHAPPLE, C., CHATTERJI, S., CHINWALLA, A., CIVETTA, A., CLIFTON, S.W., COMERON, J.M., COSTELLO, J.C., COYNE, J.A., DAUB, J., DAVID, R.G., DELCHER, A.L., DELEHAUNTY, K., DO, C.B., EBLING, H., EDWARDS, K., EICKBUSH, T., EVANS, J.D., FILIPSKI, A., FINDEISS, S., FREYHULT, E., FULTON, L., FULTON, R., GARCIA, A.C., GARDINER, A., GARFIELD, D.A., GARVIN, B.E., GIBSON, G., GILBERT, D., GNERRE, S.,

GODFREY, J., GOOD, R., GOTEA, V., GRAVELY, B., GREENBERG, A.J., GRIFFITHS-JONES, S., GROSS, S., GUIGO, R., GUSTAFSON, E.A., HAERTY, W., HAHN, M.W., HALLIGAN, D.L., HALPERN, A.L., HALTER, G.M., HAN, M.V., HEGER, A., HILLIER, L., HINRICHS, A.S., HOLMES, I., HOSKINS, R.A., HUBISZ, M.J., HULTMARK, D., HUNTLEY, M.A., JAFFE, D.B., JAGADEESHAN, S., JECK, W.R., JOHNSON, J., JONES, C.D., JORDAN, W.C., KARPEN, G.H., KATAOKA, E., KEIGHTLEY, P.D., KHERADPOUR, P., KIRKNESS, E.F., KOERICH, L.B., KRISTIANSEN, K., KUDRNA, D., KULATHINAL, R.J., KUMAR, S., KWOK, R., LANDER, E., LANGLEY, C.H., LAPOINT, R., LAZZARO, B.P., LEE, S.J., LEVESQUE, L., LI, R., LIN, C.F., LIN, M.F., LINDBLAD-TOH, K., LLOPART, A., LONG, M., LOW, L., LOZOVSKY, E., LU, J., LUO, M., MACHADO, C.A., MAKALOWSKI, W., MARZO, M., MATSUDA, M., MATZKIN, L., MCALLISTER, B., MCBRIDE, C.S., MCKERNAN, B., MCKERNAN, K., MENDEZ-LAGO, M., MINX, P., MOLLENHAUER, M.U., MONTOOTH, K., MOUNT, S.M., MU, X., MYERS, E., NEGRE, B., NEWFELD, S., NIELSEN, R., NOOR, M.A., O'GRADY, P., PACHTER, L., PAPACEIT, M., PARISI, M.J., PARISI, M., PARTS, L., PEDERSEN, J.S., PESOLE, G., PHILLIPPY, A.M., PONTING, C.P., POP, M., PORCELLI, D., POWELL, J.R., PROHASKA, S., PRUITT, K., PUIG, M., QUESNEVILLE, H., RAM, K.R., RAND, D., RASMUSSEN, M.D., REED, L.K., REENAN, R., REILY, A., REMINGTON, K.A., RIEGER, T.T., RITCHIE, M.G., ROBIN, C., ROGERS, Y.H., ROHDE, C., ROZAS, J., RUBENFIELD, M.J., RUIZ, A., RUSSO, S., SALZBERG, S.L., SANCHEZ-GRACIA, A., SARANGA, D.J., SATO, H., SCHAEFFER, S.W., SCHATZ, M.C., SCHLENKE, T., SCHWARTZ, R., SEGARRA, C., SINGH, R.S., SIROT, L., SIROTA, M., SISNEROS, N.B., SMITH, C.D., SMITH, T.F., SPIETH, J., STAGE, D.E., STARK, A., STEPHAN, W., STRAUSBERG, R.L., STREMPEL, S., STURGILL, D., SUTTON, G., SUTTON, G.G., TAO, W., TEICHMANN, S., TOBARI, Y.N., TOMIMURA, Y., TSOLAS, J.M., VALENTE, V.L., VENTER, E., VENTER, J.C., VICARIO, S., VIEIRA, F.G., VILELLA, A.J., VILLASANTE, A., WALENZ, B., WANG, J., WASSERMAN, M., WATTS, T., WILSON, D., WILSON, R.K., WING, R.A., WOLFNER, M.F., WONG, A., WONG, G.K., WU, C.I., WU, G., YAMAMOTO, D., YANG, H.P., YANG, S.P., YORKE, J.A., YOSHIDA, K., ZDOBNOV, E., ZHANG, P., ZHANG, Y., ZIMIN, A.V., BALDWIN, J., ABDOUELLEIL, A., ABDULKADIR, J., ABEBE, A., ABERA, B., ABREU, J., ACER, S.C., AFTUCK, L., ALEXANDER, A., AN, P., ANDERSON, E., ANDERSON, S., ARACHI, H., AZER, M., BACHANTSANG, P., BARRY, A., BAYUL, T., BERLIN, A., BESSETTE, D., BLOOM, T., BLYE, J., BOGUSLAVSKIY, L., BONNET, C., BOUKHGALTER, B., BOURZGUI, I., BROWN, A., CAHILL, P., CHANNER, S., CHESHATSANG, Y., CHUDA, L., CITROEN, M., COLLYMORE, A., COOKE, P., COSTELLO, M., D'ACO, K., DAZA, R., DE HAAN, G., DEGRAY, S., DEMASO, C., DHARGAY, N., DOOLEY, K., DOOLEY, E., DORICENT, M., DORJE, P., DORJEE, K., DUPES, A., ELONG, R., FALK, J., FARINA, A., FARO, S., FERGUSON, D., FISHER, S., FOLEY, C.D., FRANKE, A., FRIEDRICH, D., GADBOIS, L., GEARIN, G., GEARIN, C.R., GIANNOUKOS, G., GOODE, T., GRAHAM, J., GRANDBOIS, E., GREWAL, S., GYALTSEN, K., HAFEZ, N., HAGOS, B., HALL, J., HENSON, C., HOLLINGER, A., HONAN, T., HUARD, M.D., HUGHES, L., HURHULA, B., HUSBY, M.E., KAMAT, A., KANGA, B., KASHIN, S., KHAZANOVICH, D., KISNER, P., LANCE, K., LARA, M., LEE, W., LENNON, N., LETENDRE, F., LEVINE, R., LIPOVSKY, A., LIU, X., LIU, J., LIU, S., LOKYITSANG, T., LOKYITSANG, Y., LUBONJA, R., LUI, A., MACDONALD, P., MAGNISALIS, V., MARU, K., MATTHEWS, C., MCCUSKER, W., MCDONOUGH, S., MEHTA, T., MELDRIM, J., MENEUS, L., MIHAI, O., MIHALEV, A., MIHOVA, T., MITTELMAN, R., MLENGA, V., MONTMAYEUR, A., MULRAIN, L., NAVIDI, A., NAYLOR, J., NEGASH, T., NGUYEN, T., NGUYEN, N., NICOL, R., NORBU, C., NORBU, N., NOVOD, N., O'NEILL, B.,

OSMAN, S., MARKIEWICZ, E., OYONO, O.L., PATTI, C., PHUNKHANG, P., PIERRE, F., PRIEST, M., RAGHURAMAN, S., REGE, F., REYES, R., RISE, C., ROGOV, P., ROSS, K., RYAN, E., SETTIPALLI, S., SHEA, T., SHERPA, N., SHI, L., SHIH, D., SPARROW, T., SPAULDING, J., STALKER, J., STANGE-THOMANN, N., STAVROPOULOS, S., STONE, C., STRADER, C., TESFAYE, S., THOMSON, T., THOULUTSANG, Y., THOULUTSANG, D., TOPHAM, K., TOPPING, I., TSAMLA, T., VASSILIEV, H., VO, A., WANGCHUK, T., WANGDI, T., WEIAND, M., WILKINSON, J., WILSON, A., YADAV, S., YOUNG, G., YU, Q., ZEMBEK, L., ZHONG, D., ZIMMER, A., ZWIRKO, Z., ALVAREZ, P., BROCKMAN, W., BUTLER, J., CHIN, C., GRABHERR, M., KLEBER, M., MAUCELI, E., MACCALLUM, I., and CONSORTIUM, D.G., 2007, Evolution of genes and genomes on the *Drosophila* phylogeny: Nature, v. 450, p. 203-18, doi: 10.1038/nature06341.

COGNI, R., DING, S.D., PIMENTEL, A.C., DAY, J.P., and JIGGINS, F.M., 2021, *Wolbachia* reduces virus infection in a natural population of *Drosophila*: Commun Biol, v. 4, p. 1327, doi: 10.1038/s42003-021-02838-z.

CONNER, W., 2021, A phylogeny for the *Drosophila* montium species group: A model clade for comparative analyses: Molecular Phylogenetics and Evolution, v. 158, p. 107061.

CONNER, W.R., BLAXTER, M.L., ANFORA, G., OMETTO, L., ROTA-STABELLI, O., and TURELLI, M., 2017, Genome comparisons indicate recent transfer of *w*Ri-like *Wolbachia* between sister species *Drosophila suzukii* and *D. subpulchrella*: Ecol Evol, v. 7, p. 9391-9404, doi: 10.1002/ece3.3449.

COOPER, B.S., GINSBERG, P.S., TURELLI, M., and MATUTE, D.R., 2017, *Wolbachia* in the *Drosophila yakuba* Complex: Pervasive Frequency Variation and Weak Cytoplasmic Incompatibility, but No Apparent Effect on Reproductive Isolation: Genetics, v. 205, p. 333-351, doi: 10.1534/genetics.116.196238.

COOPER, B.S., VANDERPOOL, D., CONNER, W.R., MATUTE, D.R., and TURELLI, M., 2019, *Wolbachia* acquisition by *Drosophila yakuba*-clade hosts and transfer of incompatibility loci between distantly related *Wolbachia*: Genetics, v. 212, p. 1399-1419, doi: 10.1534/genetics.119.302349.

CUMMINGS, M.R., BROWN, N.M., and KING, R.C., 1971, The cytology of the vitellogenic stages of oogenesis in *Drosophila melanogaster*. 3. Formation of the vitelline membrane: Z Zellforsch Mikrosk Anat, v. 118, p. 482-92, doi: 10.1007/BF00324615.

DEDEINE, F., VAVRE, F., FLEURY, F., LOPPIN, B., HOCHBERG, M.E., and BOULETREAU, M., 2001, Removing symbiotic *Wolbachia* bacteria specifically inhibits oogenesis in a parasitic wasp: Proc Natl Acad Sci U S A, v. 98, p. 6247-52, doi: 10.1073/pnas.101304298.

DIRKS, U., GRUBER-VODICKA, H.R., LEISCH, N., BULGHERESI, S., EGGER, B., LADURNER, P. and OTT, J.A., 2012, Bacterial symbiosis maintenance in the asexually reproducing and regenerating flatworm Paracatenula galateia: PLoS One, v. 7, p.e34709, doi: 10.1371/journal.pone.0034709.

DOYLE, M.T. and BERNSTEIN, H.D., 2019, Bacterial outer membrane proteins assemble via asymmetric interactions with the BamA β-barrel: Nat Commun, v. 10, p.1-13, doi: 10.1038/s41467-019-11230-9.

ELLEGAARD, K.M., KLASSON, L., NÄSLUND, K., BOURTZIS, K., and ANDERSSON, S.G., 2013, Comparative genomics of *Wolbachia* and the bacterial species concept: PLoS Genet, v. 9, p. e1003381, doi: 10.1371/journal.pgen.1003381.

FERREE, P.M., FRYDMAN, H.M., LI, J.M., CAO, J., WIESCHAUS, E., and SULLIVAN, W., 2005, *Wolbachia* utilizes host microtubules and Dynein for anterior localization in the *Drosophila* oocyte: PLoS Pathog, v. 1, p. e14, doi: 10.1371/journal.ppat.0010014.

FOX, D.A., LARSSON, P., LO, R.H., KRONCKE, B.M., KASSON, P.M. and COLUMBUS, L., 2014, Structure of the neisserial outer membrane protein Opa60: Loop flexibility essential to receptor recognition and bacterial engulfment: J Am Chem Soc, v. 136, p. 9938-9946, doi: 10.1021/ja503093y.

FOX, J., and WEISBERG, S., 2019, An R companion to applied regression: SAGE.

FRECKLETON, R.P., HARVEY, P.H., and PAGEL, M., 2002, Phylogenetic analysis and comparative data: a test and review of evidence: Am Nat, v. 160, p. 712-26, doi: 10.1086/343873.

FRYDMAN, H.M., LI, J.M., ROBSON, D.N., and WIESCHAUS, E., 2006, Somatic stem cell niche tropism in *Wolbachia*: Nature, v. 441, p. 509-12, doi: 10.1038/nature04756.

GABLER, F., NAM, S.Z., TILL, S., MIRDITA, M., STEINEGGER, M., SÖDING, J., LUPAS, A.N. and ALVA, V., 2020, Protein sequence analysis using the MPI bioinformatics toolkit: Curr Protoc Bioinformatics, v. 72, p. e108, doi: 10.1002/cpbi.108.

GERTH, M., and BLEIDORN, C., 2016, Comparative genomics provides a timeframe for *Wolbachia* evolution and exposes a recent biotin synthesis operon transfer: Nat Microbiol, v. 2, p. 16241, doi: 10.1038/nmicrobiol.2016.241.

GRIEDER, N.C., DE CUEVAS, M., and SPRADLING, A.C., 2000, The fusome organizes the microtubule network during oocyte differentiation in *Drosophila*: Development, v. 127, p. 4253-64.

GROBLER, Y., YUN, C.Y., KAHLER, D.J., BERGMAN, C.M., LEE, H., OLIVER, B., and LEHMANN, R., 2018, Whole genome screen reveals a novel relationship between *Wolbachia* levels and *Drosophila* host translation: PLoS Pathog, v. 14, p. e1007445, doi: 10.1371/journal.ppat.1007445.

GUO, Y., GONG, J.T., MO, P.W., HUANG, H.J., and HONG, X.Y., 2019, *Wolbachia* localization during *Laodelphax striatellus* embryogenesis: J Insect Physiol, v. 116, p. 125-133, doi: 10.1016/j.jinsphys.2019.05.006.

HADFIELD, S.J. and AXTON, J.M., 1999, Germ cells colonized by endosymbiotic bacteria: Nature, v. 402, p. 482-482, doi: 10.1038/45002.

HAGUE, M.T.J., MAVENGERE, H., MATUTE, D.R., and COOPER, B.S., 2020a, Environmental and genetic contributions to imperfect *w*Mel-like *Wolbachia* transmission and frequency variation: Genetics, v. 215, p. 1117-1132, doi: 10.1534/genetics.120.303330.

HAGUE, M.T.J., CALDWELL, C.N., and COOPER, B.S., 2020b, Pervasive effects of *Wolbachia* on host temperature preference: mBio, v. 11, doi: 10.1128/mBio.01768-20.

HAGUE, M.T.J., SHROPSHIRE, J.D., CALDWELL, C.N., STATZ, J.P., STANEK, K.A., CONNER, W.R., and COOPER, B.S., 2022, Temperature effects on cellular host-microbe interactions

explain continent-wide endosymbiont prevalence: Curr Biol, v. 32, p. 878-888.e8, doi: 10.1016/j.cub.2021.11.065.

HARMON, L.J., WEIR, J.T., BROCK, C.D., GLOR, R.E., and CHALLENGER, W., 2008, GEIGER: investigating evolutionary radiations: Bioinformatics, v. 24, p. 129-31, doi: 10.1093/bioinformatics/btm538.

HILGENBOECKER, K., HAMMERSTEIN, P., SCHLATTMANN, P., TELSCHOW, A., and WERREN, J.H., 2008, How many species are infected with *Wolbachia*?--A statistical analysis of current data: FEMS Microbiol Lett, v. 281, p. 215-20, doi: 10.1111/j.1574-6968.2008.01110.x.

HOFFMANN, A.A., TURELLI, M., and HARSHMAN, L.G., 1990, Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*: Genetics, v. 126, p. 933-48, doi: 10.1093/genetics/126.4.933.

HOSKINS, R.A., CARLSON, J.W., WAN, K.H., PARK, S., MENDEZ, I., GALLE, S.E., BOOTH, B.W., PFEIFFER, B.D., GEORGE, R.A., SVIRSKAS, R., KRZYWINSKI, M., SCHEIN, J., ACCARDO, M.C., DAMIA, E., MESSINA, G., MÉNDEZ-LAGO, M., DE PABLOS, B., DEMAKOVA, O.V., ANDREYEVA, E.N., BOLDYREVA, L.V., MARRA, M., CARVALHO, A.B., DIMITRI, P., VILLASANTE, A., ZHIMULEV, I.F., RUBIN, G.M., KARPEN, G.H., and CELNIKER, S.E., 2015, The Release 6 reference sequence of the *Drosophila melanogaster* genome: Genome Res, v. 25, p. 445-58, doi: 10.1101/gr.185579.114.

HU, T.T., EISEN, M.B., THORNTON, K.R., and ANDOLFATTO, P., 2013, A second-generation assembly of the *Drosophila simulans* genome provides new insights into patterns of lineage-specific divergence: Genome Res, v. 23, p. 89-98, doi: 10.1101/gr.141689.112.

HÖHNA, S., LANDIS, M.J., HEATH, T.A., BOUSSAU, B., LARTILLOT, N., MOORE, B.R., HUELSENBECK, J.P., and RONQUIST, F., 2016, RevBayes: Bayesian Phylogenetic Inference Using Graphical Models and an Interactive Model-Specification Language: Syst Biol, v. 65, p. 726-36, doi: 10.1093/sysbio/syw021.

JACKMAN, S.D., VANDERVALK, B.P., MOHAMADI, H., CHU, J., YEO, S., HAMMOND, S.A., JAHESH, G., KHAN, H., COOMBE, L., WARREN, R.L., and BIROL, I., 2017, ABySS 2.0: resource-efficient assembly of large genomes using a Bloom filter: Genome Res, v. 27, p. 768-777, doi: 10.1101/gr.214346.116.

JOSHI, N., and FASS, J., 2011, Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files.

KAMATH, A.D., DEEHAN, M.A., and FRYDMAN, H.M., 2018, Polar cell fate stimulates *Wolbachia* intracellular growth: Development, v. 145, doi: 10.1242/dev.158097.

KATOH, K., and STANDLEY, D.M., 2013, MAFFT multiple sequence alignment software version 7: improvements in performance and usability: Mol Biol Evol, v. 30, p. 772-80, doi: 10.1093/molbev/mst010.

KLASSON, L., WESTBERG, J., SAPOUNTZIS, P., NÄSLUND, K., LUTNAES, Y., DARBY, A.C., VENETI, Z., CHEN, L., BRAIG, H.R., GARRETT, R., BOURTZIS, K., and ANDERSSON, S.G., 2009, The mosaic genome structure of the *Wolbachia w*Ri strain infecting *Drosophila simulans*: Proc Natl Acad Sci U S A, v. 106, p. 5725-30, doi: 10.1073/pnas.0810753106.

KOSE, H., and KARR, T.L., 1995, Organization of *Wolbachia* pipientis in the *Drosophila* fertilized egg and embryo revealed by an anti-*Wolbachia* monoclonal antibody: Mech Dev, v. 51, p. 275-88, doi: 10.1016/0925-4773(95)00372-x.

KÜHN, S., and ENNINGA, J., 2020, The actin comet guides the way: How *Listeria* actin subversion has impacted cell biology, infection biology and structural biology: Cell Microbiol, v. 22, p. e13190, doi: 10.1111/cmi.13190.

LANDMANN, F., BAIN, O., MARTIN, C., UNI, S., TAYLOR, M.J., and SULLIVAN, W., 2012, Both asymmetric mitotic segregation and cell-to-cell invasion are required for stable germline transmission of *Wolbachia* in filarial nematodes: Biol Open, v. 1, p. 536-47, doi: 10.1242/bio.2012737.

LUO, Y., AHMAD, E., and LIU, S.T., 2018, MAD1: Kinetochore Receptors and Catalytic Mechanisms: Front Cell Dev Biol, v. 6, p. 51, doi: 10.3389/fcell.2018.00051.

MADO, K., CHEKULAYEV, V., SHEVCHUK, I., PUURAND, M., TEPP, K. and KAAMBRE, T., 2019, On the role of tubulin, plectin, desmin, and vimentin in the regulation of mitochondrial energy fluxes in muscle cells: Am J Physiol Cell Physiol, v. 316, p. C657-C667, doi: 10.1152/ajpcell.00303.2018.

MAHAJAN-MIKLOS, S., and COOLEY, L., 1994, Intercellular cytoplasm transport during *Drosophila* oogenesis: Dev Biol, v. 165, p. 336-51, doi: 10.1006/dbio.1994.1257.

MAHOWALD, A.P., and STRASSHEIM, J.M., 1970, Intercellular migration of centrioles in the germarium of *Drosophila melanogaster*. An electron microscopic study: J Cell Biol, v. 45, p. 306-20, doi: 10.1083/jcb.45.2.306.

MAIMON, I., and GILBOA, L., 2011, Dissection and staining of *Drosophila* larval ovaries: J Vis Exp, doi: 10.3791/2537.

MEANY, M.K., CONNER, W.R., RICHTER, S.V., BAILEY, J.A., TURELLI, M., and COOPER, B.S., 2019, Loss of cytoplasmic incompatibility and minimal fecundity effects explain relatively low *Wolbachia* frequencies in *Drosophila mauritiana*: Evolution, v. 73, p. 1278-1295, doi: 10.1111/evo.13745.

MERKLE, J.A., WITTES, J., and SCHÜPBACH, T., 2020, Signaling between somatic follicle cells and the germline patterns the egg and embryo of *Drosophila*: Curr Top Dev Biol, v. 140, p. 55-86, doi: 10.1016/bs.ctdb.2019.10.004.

METCALF, J.A., JO, M., BORDENSTEIN, S.R., and JAENIKE, J., 2014, Recent genome reduction of *Wolbachia* in *Drosophila recens* targets phage WO and narrows candidates for reproductive parasitism: PeerJ, v. 2, p. e529, doi: 10.7717/peerj.529.

NOINAJ, N., KUSZAK, A.J., GUMBART, J.C., LUKACIK, P., CHANG, H., EASLEY, N.C., LITHGOW, T. and BUCHANAN, S.K., 2013, Structural insight into the biogenesis of β-barrel membrane proteins: Nature, v. 501, p. 385-390, doi: 10.1038/nature12521.

O'NEILL, S.L., GIORDANO, R., COLBERT, A.M., KARR, T.L., and ROBERTSON, H.M., 1992, 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects: Proc Natl Acad Sci U S A, v. 89, p. 2699-702, doi: 10.1073/pnas.89.7.2699.

PAGEL, M., 1999, Inferring the historical patterns of biological evolution: Nature, v. 401, p. 877-84, doi: 10.1038/44766.

PANNEBAKKER, B.A., LOPPIN, B., ELEMANS, C.P., HUMBLOT, L., and VAVRE, F., 2007, Parasitic inhibition of cell death facilitates symbiosis: Proc Natl Acad Sci U S A, v. 104, p. 213-5, doi: 10.1073/pnas.0607845104.

POINSOT, D., BOURTZIS, K., MARKAKIS, G., SAVAKIS, C. and MERÇOT, H., 1998, *Wolbachia* transfer from *Drosophila melanogaster* into *D. simulans*: host effect and cytoplasmic incompatibility relationships: Genetics, v. 150, p.227-237, doi: 10.1093/genetics/150.1.227.

QUINLAN, M.E., 2016, Cytoplasmic Streaming in the *Drosophila* Oocyte: Annu Rev Cell Dev Biol, v. 32, p. 173-195, doi: 10.1146/annurev-cellbio-111315-125416.

RAMALHO, M.O., VIEIRA, A.S., PEREIRA, M.C., MOREAU, C.S., and BUENO, O.C., 2018, Transovarian Transmission of *Blochmannia* and *Wolbachia* Endosymbionts in the Neotropical Weaver Ant *Camponotus textor* (Hymenoptera, Formicidae): Curr Microbiol, v. 75, p. 866-873, doi: 10.1007/s00284-018-1459-3.

RAYCHOUDHURY, R., BALDO, L., OLIVEIRA, D.C., and WERREN, J.H., 2009, Modes of acquisition of *Wolbachia*: horizontal transfer, hybrid introgression, and codivergence in the *Nasonia* species complex: Evolution, v. 63, p. 165-83, doi: 10.1111/j.1558-5646.2008.00533.x.

RICE, D.W., SHEEHAN, K.B. and NEWTON, I.L., 2017, Large-scale identification of *Wolbachia pipientis* effectors: Genome Biol Evol, v. 9, p. 1925-1937, doi: 10.1093/gbe/evx139.

RUSSELL, S.L., CHAPPELL, L., and SULLIVAN, W., 2019, A symbiont's guide to the germline: Curr Top Dev Biol, v. 135, p. 315-351, doi: 10.1016/bs.ctdb.2019.04.007.

RUSSELL, S.L., LEMSEFFER, N., and SULLIVAN, W.T., 2018, *Wolbachia* and host germline components compete for kinesin-mediated transport to the posterior pole of the *Drosophila* oocyte: PLoS Pathog, v. 14, p. e1007216, doi: 10.1371/journal.ppat.1007216.

RØRTH, P., 2002, Initiating and guiding migration: lessons from border cells: Trends Cell Biol, v. 12, p. 325-31, doi: 10.1016/s0962-8924(02)02311-5.

SALZBERG, S.L., DUNNING HOTOPP, J.C., DELCHER, A.L., POP, M., SMITH, D.R., EISEN, M.B., and NELSON, W.C., 2005, Serendipitous discovery of *Wolbachia* genomes in multiple *Drosophila* species: Genome Biol, v. 6, p. R23, doi: 10.1186/gb-2005-6-3-r23.

SANDER, 1968, Developmental physiological studies on embryonal mycetoma *Euscelis plebejus* I. Removal of and abnormal combination of unicellular components of symbiotic sysems: Dev Biol, v. 17, p. 16-38.

SEEMANN, T., 2014, Prokka: rapid prokaryotic genome annotation: Bioinformatics, v. 30, p. 2068-9, doi: 10.1093/bioinformatics/btu153.

SERBUS, L.R., CASPER-LINDLEY, C., LANDMANN, F., and SULLIVAN, W., 2008, The genetics and cell biology of *Wolbachia*-host interactions: Annu Rev Genet, v. 42, p. 683-707, doi: 10.1146/annurev.genet.41.110306.130354.

SERBUS, L.R., CHA, B.J., THEURKAUF, W.E., and SAXTON, W.M., 2005, Dynein and the actin cytoskeleton control kinesin-driven cytoplasmic streaming in *Drosophila* oocytes: Development, v. 132, p. 3743-52, doi: 10.1242/dev.01956.

SERBUS, L.R., FERRECCIO, A., ZHUKOVA, M., MCMORRIS, C.L., KISELEVA, E., and SULLIVAN, W., 2011, A feedback loop between *Wolbachia* and the *Drosophila* gurken mRNP complex influences *Wolbachia* titer: J Cell Sci, v. 124, p. 4299-308, doi: 10.1242/jcs.092510.

SERBUS, L.R., and SULLIVAN, W., 2007, A cellular basis for *Wolbachia* recruitment to the host germline: PLoS Pathog, v. 3, p. e190, doi: 10.1371/journal.ppat.0030190.

SERBUS, L.R., WHITE, P.M., SILVA, J.P., RABE, A., TEIXEIRA, L., ALBERTSON, R., and SULLIVAN, W., 2015, The impact of host diet on *Wolbachia* titer in *Drosophila*: PLoS Pathog, v. 11, p. e1004777, doi: 10.1371/journal.ppat.1004777.

SHEEHAN, K.B., MARTIN, M., LESSER, C.F., ISBERG, R.R., and NEWTON, I.L., 2016, Identification and Characterization of a Candidate *Wolbachia pipientis* Type IV Effector That Interacts with the Actin Cytoskeleton: MBio, v. 7, doi: 10.1128/mBio.00622-16.

SCHRODINGER LLC, 2015, The PyMOL molecular graphics system, version 1.8.

STEINHAUER, J., and KALDERON, D., 2006, Microtubule polarity and axis formation in the *Drosophila* oocyte: Dev Dyn, v. 235, p. 1455-68, doi: 10.1002/dvdy.20770.

SULLIVAN, W., 2017, *Wolbachia*, bottled water, and the dark side of symbiosis: Mol Biol Cell, v. 28, p. 2343-2346, doi: 10.1091/mbc.E17-02-0132.

SULLIVAN, W., ASHBURNER, M., and HAWLEY, R.S., 2000, *Drosophila* Protocols: Cold Spring Harbor Press.

SUVOROV, A., KIM, B.Y., WANG, J., ARMSTRONG, E.E., PEEDE, D., D'AGOSTINO, E.R.R., PRICE, D.K., WADDELL, P., LANG, M., COURTIER-ORGOGOZO, V., DAVID, J.R., PETROV, D., MATUTE, D.R., SCHRIDER, D.R., and COMEAULT, A.A., 2022, Widespread introgression across a phylogeny of 155 *Drosophila* genomes: Curr Biol, v. 32, p. 111-123.e5, doi: 10.1016/j.cub.2021.10.052.

TEUFEL, F., ALMAGRO ARMENTEROS, J.J., JOHANSEN, A.R., GÍSLASON, M.H., PIHL, S.I., TSIRIGOS, K.D., WINTHER, O., BRUNAK, S., VON HEIJNE, G. and NIELSEN, H., 2022, SignalP 6.0 predicts all five types of signal peptides using protein language models: Nat Biotechnol, v. 40, p. 1023-1025, doi: 10.1038/s41587-021-01156-3.

THEURKAUF, W.E., 1994, Microtubules and cytoplasm organization during *Drosophila* oogenesis: Dev Biol, v. 165, p. 352-60, doi: 10.1006/dbio.1994.1258.

THEURKAUF, W.E., and HAZELRIGG, T.I., 1998, In vivo analyses of cytoplasmic transport and cytoskeletal organization during *Drosophila* oogenesis: characterization of a multi-step anterior localization pathway: Development, v. 125, p. 3655-66.

THEURKAUF, W.E., SMILEY, S., WONG, M.L., and ALBERTS, B.M., 1992, Reorganization of the cytoskeleton during *Drosophila* oogenesis: implications for axis specification and intercellular transport: Development, v. 115, p. 923-36.

TOOMEY, M.E., PANARAM, K., FAST, E.M., BEATTY, C., and FRYDMAN, H.M., 2013, Evolutionarily conserved *Wolbachia*-encoded factors control pattern of stem-cell niche tropism in *Drosophila* ovaries and favor infection: Proc Natl Acad Sci U S A, v. 110, p. 10788-93, doi: 10.1073/pnas.1301524110.

TURELLI, M., COOPER, B.S., RICHARDSON, K.M., GINSBERG, P.S., PECKENPAUGH, B., ANTELOPE, C.X., KIM, K.J., MAY, M.R., ABRIEUX, A., WILSON, D.A., BRONSKI, M.J., MOORE, B.R., GAO, J.J., EISEN, M.B., CHIU, J.C., CONNER, W.R., and HOFFMANN, A.A., 2018, Rapid Global Spread of *w*Ri-like *Wolbachia* across Multiple *Drosophila*: Curr Biol, v. 28, p. 963-971.e8, doi: 10.1016/j.cub.2018.02.015.

TURELLI, M., and HOFFMANN, A.A., 1995, Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations: Genetics, v. 140, p. 1319-38.

VENETI, Z., CLARK, M.E., KARR, T.L., SAVAKIS, C., and BOURTZIS, K., 2004, Heads or tails: host-parasite interactions in the *Drosophila-Wolbachia* system: Appl Environ Microbiol, v. 70, p. 5366-72, doi: 10.1128/AEM.70.9.5366-5372.2004.

WEINERT, L.A., ARAUJO-JNR, E.V., AHMED, M.Z. and WELCH, J.J., 2015, The incidence of bacterial endosymbionts in terrestrial arthropods: Proc Royal Soc B, v. 282, p. 20150249, doi: 10.1098/rspb.2015.0249.

WERREN, J.H., BALDO, L., and CLARK, M.E., 2008, *Wolbachia*: master manipulators of invertebrate biology: Nat Rev Microbiol, v. 6, p. 741-51, doi: 10.1038/nrmicro1969.

WHITE, P.M., PIETRI, J.E., DEBEC, A., RUSSELL, S., PATEL, B., and SULLIVAN, W., 2017, Mechanisms of Horizontal Cell-to-Cell Transfer of *Wolbachia* spp. in *Drosophila melanogaster*: Appl Environ Microbiol, v. 83, doi: 10.1128/AEM.03425-16.

WHITE, P.M., SERBUS, L.R., DEBEC, A., CODINA, A., BRAY, W., GUICHET, A., LOKEY, R.S., and SULLIVAN, W., 2017, Reliance of *Wolbachia* on High Rates of Host Proteolysis Revealed by a Genome-Wide RNAi Screen of *Drosophila* Cells: Genetics, v. 205, p. 1473-1488, doi: 10.1534/genetics.116.198903.

WU, M., SUN, L.V., VAMATHEVAN, J., RIEGLER, M., DEBOY, R., BROWNLIE, J.C., MCGRAW, E.A., MARTIN, W., ESSER, C., AHMADINEJAD, N., WIEGAND, C., MADUPU, R., BEANAN, M.J., BRINKAC, L.M., DAUGHERTY, S.C., DURKIN, A.S., KOLONAY, J.F., NELSON, W.C., MOHAMOUD, Y., LEE, P., BERRY, K., YOUNG, M.B., UTTERBACK, T., WEIDMAN, J., NIERMAN, W.C., PAULSEN, I.T., NELSON, K.E., TETTELIN, H., O'NEILL, S.L., and EISEN, J.A., 2004, Phylogenomics of the reproductive parasite *Wolbachia* pipientis *w*Mel: a streamlined genome overrun by mobile genetic elements: PLoS Biol, v. 2, p. E69, doi: 10.1371/journal.pbio.0020069.

YANG, J. and ZHANG, Y., 2015, I-TASSER server: new development for protein structure and function predictions: Nucleic Acids Res, v. 43, p. W174-W181, doi: 10.1093/nar/gkv342.

ZHANG, C., FREDDOLINO, P.L. and ZHANG, Y., 2017, COFACTOR: improved protein function prediction by combining structure, sequence and protein–protein interaction information: Nucleic Acids Res, v. 45, p. W291-W299, doi: 10.1093/nar/gkx366.

ZIMMERMANN, L., STEPHENS, A., NAM, S.Z., RAU, D., KÜBLER, J., LOZAJIC, M., GABLER, F., SÖDING, J., LUPAS, A.N. and ALVA, V., 2018, A completely reimplemented MPI

bioinformatics toolkit with a new HHpred server at its core: J Mol Biol, v. 430, p. 2237-2243, doi: 10.1016/j.jmb.2017.12.007.

ZUG, R., and HAMMERSTEIN, P., 2012, Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected: PLoS One, v. 7, p. e38544, doi: 10.1371/journal.pone.0038544.