Title

How density dependence, genetic erosion, and the extinction vortex impact evolutionary rescue

Running head: Density dependence & evolutionary rescue

Authors

Scott W. Nordstrom[1,2], Ruth A. Hufbauer[3], Laure Olazcuaga[4], Lily F. Durkee[3] and Brett A.

Melbourne[1, *]

[1]Department of Ecology & Evolutionary Biology, University of Colorado, Boulder, Boulder, CO,

80309

[2]BioFrontiers Institute, University of Colorado, Boulder, Boulder, CO, 80303

[3]Department of Agricultural Biology and Graduate Degree Program in Ecology, Colorado State

University, Fort Collins, CO 80523

[4]Department of Agricultural Biology, Colorado State University, Fort Collins, CO 80523

[*] For correspondence: brett.melbourne@colorado.edu

Abstract

Following severe environmental change that reduces mean population fitness below

replacement, populations must adapt to avoid extinction, a process called evolutionary rescue.

Models of evolutionary rescue demonstrate that initial size, genetic variation, and degree of

maladaptation influence population fates. However, many models feature populations that

grow without negative density dependence or with constant genetic diversity despite

precipitous population decline, assumptions likely to be violated in conservation settings. We

examined the influences of density-dependent growth and erosion of genetic diversity

simultaneously on populations adapting to novel environmental change using stochastic,

individual-based simulations. Density dependence decreased the probability of rescue and

increased the probability of extinction, especially in large and initially well-adapted populations

that previously have been predicted to be at low risk. Increased extinction occurred shortly

following environmental change, as populations under density dependence experienced more

rapid decline and reached smaller sizes. Populations that experienced evolutionary rescue lost

genetic diversity through drift and adaptation, particularly under density dependence.

Populations that declined to extinction entered an extinction vortex, where small size increased

drift, loss of genetic diversity, and the fixation of maladaptive alleles, hindered adaptation, and

kept populations at small densities where they were vulnerable to extinction *via* demographic

stochasticity.

Introduction

Adaptation to novel environments is often necessary for populations to persist in this

current era of large-scale anthropogenic change and habitat alteration. Novel or sudden

environmental change can abruptly render a population poorly adapted to its habitat. When

the environmental change is so severe that mean population fitness falls below the

replacement rate, the population will go extinct if it does not adapt. This process of populations

adapting to severe change in environmental conditions sufficiently to avoid extinction is called

evolutionary rescue (Gomulkiewicz and Holt 1995, Carlson et al. 2014).

Gomulkiewicz and Holt (1995) formalized the idea of evolutionary rescue, highlighting the

"U"-shaped demographic trajectory where populations initially decline due to maladaptation to

novel environmental conditions, followed by adaptation that allows populations to avoid

extinction and return to their original size. They found that evolutionary rescue was most likely

to occur in large populations with high genetic diversity and a relatively low degree of

maladaptation to their environment (i.e., population mean phenotype close to that favored by

the novel environment) at the time of the environmental change. Subsequent work in

controlled experiments has confirmed the importance of these processes to evolutionary

rescue (e.g., Bell and Gonzalez 2009, Agashe et al. 2011, Hufbauer et al. 2015).

Models examining rescue often rely on assumptions of density-independent growth and

constant additive genetic variance (Kopp and Matuszewski 2014), potentially making the

findings less applicable to many conservation targets. For example, Gomulkiewicz and Holt's

(1995) model assumes that population growth is density independent, and that populations can

grow unconstrained after rescue. This assumption will often be violated in conservation settings

where habitat loss, degradation, or other environmental change limits resource availability and

increases intraspecific competition. Additionally, Gomulkiewicz and Holt's (1995) model (and

others, e.g., Chevin and Lande 2010) rely on quantitative genetics results from Lande (1976),

which assume that population size is infinite and therefore that additive genetic variance

remains constant over time. However, threatened populations experiencing rapid decline are

likely to experience erosion of genetic diversity (Frankham et al. 1999, Bijlsma and Loeschke

2011, Pauls et al. 2013) that may decrease their ability to adapt to novel conditions (Houle

1992), possibly hindering rescue.

Incorporating demographic and genetic processes such as negative density dependence

and genetic erosion will improve the accuracy of predictions from models of evolutionary

rescue for populations and species of conservation concern. Negative density dependence such

as that caused by intraspecific competition should constrain evolutionary rescue by accelerating

population declines and slowing population recovery from reduced size (Chevin and Lande

2010). Some models of rescue and environmental tracking have included constrained

population size by including a ceiling-type carrying capacity (e.g., Lande 1993, Orr and Unckless

2014, Lyberger et al. 2021), but growth below carrying capacity in these models was density

independent, failing to capture how intraspecific competition or other negative density-

dependent processes depress population growth in declining and recovery phases of rescue.

Orr and Unckless (2008) analyzed a density-dependent model of adaptation with a single locus

trait, finding that density dependence impedes rescue by reducing the survival of novel

beneficial mutants. However, they did not investigate effects of density dependence on

adaptation from standing genetic variation. Relaxing the assumption of density independence,

Chevin and Lande (2010) found that density dependence impedes rescue, but their model held

genetic diversity constant, possibly ignoring effects of drift for populations reaching low

density. A population's additive genetic variance for a trait under selection in a population

influences its rate of adaptation to novel conditions (Lande 1976, Houle 1992); loss of this

genetic diversity as populations decline in response to environmental change should slow rates

of adaptation, increasing the time until rescue or making eventual rescue less likely.

Populations that are small at the onset of environmental change are especially likely to

experience genetic erosion through drift, inbreeding, and the fixation of maladaptive alleles

(Hedrick and García-Dorado 2016). Boulding and Hay (2001) and Barfield and Holt (2016)

modeled populations allowing genetic diversity to change over time, finding that drift could

occur potentially slowing adaptation, but they did not focus on interactions of these factors

with competition that constrains population growth rates.

Modeling these demographic and genetic processes simultaneously is important, as

they can interact and reinforce each other, creating a positive feedback loop of demographic

and genetic decline that populations in the process of evolutionary rescue may be drawn into

(Fig. 1). In the conservation biology literature this positive feedback loop is called an "extinction

vortex" (Gilpin and Soulé 1986). In an extinction vortex (Fig. 1) small populations experience

drift and inbreeding, reducing fitness directly *via* increased genetic load and indirectly by

erosion of genetic variation and reducing the rate of future adaptation (Frankham et al. 1999),

keeping populations small or causing them to decline further to a smaller size. This in turn leads

to further genetic erosion or increased genetic load, reduced fitness, and further declines in

population size until inevitably the population is highly vulnerable to extinction *via* stochasticity

in births and deaths, demographic heterogeneity, and sampling variation in sex ratios

(Melbourne and Hastings 2008). This feedback loop potentially opposes or resists the

evolutionary rescue process, where adaptation increases fitness, increasing population growth

rates to the point where population size stabilizes and eventually returns to original size.

Negative density dependence has the potential to disrupt evolutionary rescue and draw

populations back towards the vortex by accelerating reductions in, or slowing increases in,

population growth rate, fitness, and genetic diversity (Fig. 1). The original conceptualization of

an extinction vortex does not necessarily assume a degraded or altered habitat where

adaptation is necessary for persistence (Gilpin and Soulé 1986). Perhaps for this reason, to our

knowledge the extinction vortex has not been explicitly studied in the context of evolutionary

rescue and adaptation to novel environments. Yet, these interactive, combined effects of small

population size could be of particular importance in evolutionary rescue, as by definition

populations undergoing rescue initially decline to smaller sizes, which could initiate entry into

an extinction vortex. The extinction vortex also necessitates understanding how density

dependence and genetic erosion interact with other important determinants of the success of

rescue, e.g., the initial conditions with respect to population size, genetic diversity, and degree

of maladaptation (Gomulkiewicz and Holt 1995, Bell and Gonzalez 2009, Agashe et al. 2011).

For example, large populations adapting to sudden environmental change might never

approach the extinction vortex under density-independent population growth but might more

often enter the vortex under density dependence. Ultimately, under realistic conditions in

natural settings, evolutionary rescue might often involve a population escaping from an

extinction vortex.

Here, we relax the assumptions of density independence and constant genetic diversity

to examine their simultaneous effects on evolutionary rescue and the extinction vortex,

establishing a connection between these two major concepts in conservation biology. We do

this by combining stochastic Ricker population growth and a genetic framework with a finite-

locus quantitative trait in the same individual-based model. We simulate both rapid (15

generations) and longer (50 generations) timescales. We address two main objectives. First, we

assess how negative density dependence interacts with initial conditions with respect to

population size, additive genetic variance, and degree of genetic maladaptation in the novel

environment to influence trajectories of population size, probabilities of extinction and rescue,

additive genetic variance and fitness over time. Second, we test for the presence of an

extinction vortex and evaluate how negative density dependence influences this vortex.

Methods

Model description

We explored evolutionary rescue and extinction using stochastic simulations. This

approach allowed us to incorporate several stochastic demographic and genetic processes that

are important for rescue or extinction (Melbourne and Hastings 2008, Carlson et al. 2014).

While extinction is straightforward to identify, the classical criterion of evolutionary rescue

(population growth rate above replacement) is more difficult to operationalize for individual

populations because rescued populations may decline in size or growth rate due to

demographic or genetic stochasticity, inbreeding, and genetic load and revert to an unrescued

state. We operationalize rescue using two separate criteria: (1) expected number of offspring

per individual ($\bar{R}_t$, as defined below) greater than 1 for three consecutive generations

(hereafter "fitness-based rescue") and (2) populations exceeding their initial size for three

consecutive generations (hereafter "size-based rescue").

We derive a discrete-time individual-based model with diploid genetic inheritance that

has general features of the semelparous life history strategy common to many sexually

reproducing organisms. Our genetic submodel is a finite quantitative locus model with random

and independent segregation of alleles, similar to that of Boulding and Hay (2001). Our

demographic submodel is based on an expanded stochastic Ricker model (Ricker 1954,

Melbourne and Hastings 2008). We use this model because it is a common form of density

dependence while also accurate for the population dynamics of the model organism *Tribolium*

castaneum in a discrete-time laboratory culture (Melbourne and Hastings 2008) thus

complementing current and future experiments of rescue using this system (e.g., Agashe et al.

2011, Hufbauer et al. 2015, Stewart et al. 2017), effectively grounding our simulations in

concrete details of biology.

Each individual has an intrinsic fitness determined by genetic and environmental

components. The genetic component is determined by each of *m* biallelic diploid loci. Each

allele-copy contributes a positive $(1/(2\sqrt{m}))$ or negative $(-1/(2\sqrt{m}))$ amount to the genotype

of individual *i*, g_i, which is the sum of allele-copies across all loci. Parental alleles are segregated

independently to offspring during reproduction. The additive genetic variance in the population

is thus

$$\sigma_a^2 \;=\; 2\sum_{j=1}^{m}\frac{p_j(1-p_j)}{m} \;=\; 2\left(p'\,(1-p') \;-\; \sigma_p^2\right)$$

(1)

where p_j is the frequency of the positive allele at locus *j* across the population and p' and σ_p^2 are

respectively the mean and variance of allele frequencies across all loci (see Lynch and Walsh

1998 and Supporting Information A). An individual's phenotype, z_i, is the genotype, g_i, plus a

deviation due to environmental variation drawn from a normal distribution with mean zero and

variance σ_e^2. Intrinsic fitness, W_i, the expected fitness of an individual in the absence of density

effects, is determined by Gaussian selection (Lande 1976, Gomulkiewicz and Holt 1995) such

that $W_i = W_{max} \, exp(-(z_i - \theta)^2 \, / \, 2w^2)$, where θ is the optimal phenotype, w^2 is the width of

the fitness landscape and is inversely proportional to selection strength , and W_{max} is the

maximum intrinsic fitness (i.e., intrinsic fitness of an individual with the optimal phenotype).

The expected number of offspring of an individual, R_i, is determined according to a Ricker

function that reduces fitness due to density dependence such that $R_i = W_i e^{-\alpha N_t}$, where α is

the strength of density dependence and N_t is the population size at time t (Ricker 1954). The

realized fitness of an individual is a stochastic process as described next.

Each individual is born with a randomly determined sex assigned with equal probability.

Each generation, each female mates with a male chosen uniformly at random from the

population. In practice, mating with multiple males during a single generation occurs in female

T. castaneum (Pai and Yan 2020); however, our model makes the simplifying assumption that

each female mates with exactly one male while males can mate with any number of females,

including zero. This assumption corresponds to well-documented sperm competition and the

fertilization precedence of the last male mate *in T. castaneum* (e.g., Arnaud et al. 2001). Each

female produces a Poisson-distributed number of offspring with mean $2R_i$. Generations are not

overlapping, i.e., only one breeding generation is present at a time. As such, if all females have

zero offspring, extinction is guaranteed in the next time step. We use non-overlapping

generations both for comparison with other models of adaptation and evolutionary rescue

(e.g., Gomulkiewicz and Holt 1995, Boulding and Hay 2001, Chevin and Lande 2010) and

because it provides a tractable setup for experimental validation using the *Tribolium* model

system (e.g., Hufbauer et al. 2015, Stewart et al. 2017).

Experimental Design, Simulations, and Analysis

We simulated populations with m = 25 loci and an environmental change that moved

the phenotypic optimum from its prior state of 0 to θ = 2.8. Because θ > 0, the environment in

our simulations favors a shift toward positive alleles. Populations were founded with positive

and negative alleles assigned to each individual's loci at random with equal probability such

that across simulation trials the expected population mean genotype was zero, although due to

sampling there was variation in initial population mean genotype across trials. Thus, to reach

the new phenotypic optimum θ = 2.8, a typical population needed to change from 25 positive

allele copies (out of 50) per individual to 39 allele copies per individual (see Supporting

Information B). All simulations were run with maximum intrinsic fitness W_{max} = 2, non-genetic

phenotypic variance component σ_e^2 = 0.5, and w^2 = 3.5 as the width of the fitness landscape

Parameter choices are justified in full in Supporting Information C; briefly, parameters were

chosen to resemble observations from the *T. castaneum* model system (e.g., Wade et al. 1996,

Hufbauer et al. 2015) while also being realistic for a variety of natural populations (e.g., Hone et

al. 2010, Bonnet et al. 2022). We chose our fitness landscape width (w^2) based on analysis of

the model of Gomulkiewicz and Holt (1995) demonstrating that rescue is only likely to occur

when the width of the fitness landscape is 1-10 times the amount of phenotypic variance

(Supporting Information C). With these parameter values, mean intrinsic fitness in the typical

population immediately following the environmental change was approximately 0.65 such that

populations were sufficiently challenged to the point that both rescue and extinction were

possible on rapid timescales and resembled observed population dynamics in previous studies

with *T. castaneum* (e.g., Hufbauer et al. 2015, Stewart et al. 2017). All parameter values are

summarized in Supporting Information, Table D1.

To evaluate the effects of negative density dependence on evolutionary rescue and its

potential interactions with initial population size and amount of genetic diversity, we

conducted a 2 x 2 x 2 factorial simulation experiment (yielding eight treatments, Table 1).

Populations were simulated with the following parameter values: Negative density dependence

absent (α = 0; hereafter "density-independent populations") or present (α = 0.0035, hereafter

"density-dependent populations", value from Melbourne and Hastings 2008); initial size N_0 =

100 or N_0 = 20, hereafter referred to as "large" and "small" populations in reference to their

initial state rather than their eventual size; initial genetic diversity high ($\sigma_a^2 \approx$ 0.5, heritability

approx. 0.5) or low ($\sigma_a^2 \approx$ 0.25, heritability approx. 0.33, Lynch and Walsh 1998), hereafter

"high diversity" and "low diversity" populations, again in reference to their initial state rather

than their eventual level of diversity. These initial heritabilities are similar to those found in *T.*

castaneum by Wade et al. (1996), who found heritability of fitness of approximately 0.44 in

outbred lines and approximately 0.25 in lines subject to two generations of inbreeding. To

simulate populations with initial low diversity, at the beginning of each simulation, six loci were

set to fixation for the positive (adaptive) allele and six were set to fixation for the negative

(maladaptive) allele, leaving only 13 loci (approximately half of the genome) available for

selection to act upon while leaving the expected mean population genotype unchanged

compared to the high-diversity treatments. Stochasticity in the founding of populations

produced sampling variation in the initial mean population genotype; we took advantage of this

sampling variation to also analyze a third important determinant of a population's fate, initial

degree of maladaptation (Gomulkiewicz and Holt 1995, Barfield and Holt 2016), and its

interactions with density dependence as described in paragraphs below. A subtle point is that

expected initial additive genetic variance, $(\sigma_a^2)_0$, across replicate populations is slightly less

than the target values because of the sampling variation in allele frequencies associated with

founding the populations (Eq. 1). Likewise, due to sampling variation in initial allele frequencies,

there is more variation in initial mean population genotypes and phenotypes among small

populations than among large populations.

To assess adaptation and extinction on short time scales, we ran 4000 simulations

lasting fifteen generations for each treatment. All simulations were performed in R, version

4.1.2 (R Core Team, 2020). In each simulation, for each generation we tracked population size

(N_t), mean genotype ($\bar{g}_t$), mean phenotype ($\bar{z}_t$), mean intrinsic fitness ($\overline{W}_t$), where all means

are across individuals within a population, as well as additive genetic variance ($(\sigma_a^2)_t$), and the

proportion of loci at fixation within the population for both alleles. Throughout our analysis, we

will use an overbar to represent a mean across individuals within one population in one

timestep. We define adaptation as an increase in $\overline{W}_t$. Because fitness is optimized at a positive

genotypic value (i.e., $\theta > 0$) and fitness is a monotonic function of maladaptive genetic load,

adaptation can be equivalently defined as an increase in $\bar{g}_t$ or a reduction in genetic load, $(\theta -$

$\bar{g}_t)$.

As a first evaluation of how density dependence, population size, and genetic diversity

influence rescue and extinction over 15 generations, we visualized mean population size over

time for each treatment group. This visualization shows whether the U-shaped population

trajectory characteristic of rescue (Gomulkiewicz and Holt 1995) occurs, and if so, how it differs

by treatment. All mean population sizes were estimated including extinct populations as size

zero, thus incorporating the possibility of extinction into expected population size in each time

step. To quantify the average dynamics of the model we also estimated and visualized the

following response variables, averaged over the ensemble of all populations within a

treatment: mean population genotype ($\langle \bar{g}_t \rangle$), mean intrinsic fitness ($\langle \bar{W}_t \rangle$), mean additive

genetic variance ($\langle \sigma_a^2 \rangle_t$), and number of loci at fixation for positive and negative alleles across

all populations in each generation. To differentiate ensemble means from population means for

a single trial, we use angle brackets ($\langle \rangle$) to denote means of population-level means across an

ensemble of simulations. Because these variables are undefined for extinct populations (which

have no genotype, fitness, or additive genetic variance), estimating unconditional expectations

for each generation was impossible, and thus we estimated these means conditioned on

survival or extinction at the end of the simulation. Differences in ensemble means for these

variables between density-independent and density-dependent populations may be due

directly to density dependence, but they may also be due indirectly to non-random extinctions.

For example, a population with low mean intrinsic fitness may persist under density

independence but go extinct under density dependence, lowering the ensemble mean

population fitness for extant populations in a density-independent treatment but not affecting

the ensemble mean of extant populations in a density-dependent treatment, confounding

comparisons between the treatments. As such, we only use plots of variables conditioned on

survival/extinction for qualitative assessment, noting general patterns and interpreting them

with caution.

To assess the probabilities of extinction and rescue on longer timescales, we ran 1000

simulations lasting 50 generations for each treatment. Due to the computational intensity of

simulating large populations, we stopped a trial upon population size exceeding 10000. Because

of this censoring, we did not analyze trajectories of demographic or genotypic state variables in

these longer simulations, instead focusing only on the probabilities of extinction and rescue.

We evaluated extinction in each generation in two ways: (1) instantaneous probability of

extinction (the probability of extinction in generation t conditioned on surviving to t) and (2)

cumulative probability of extinction over time (the probability of extinction in or prior to t). We

also used two separate criteria of rescue (defined above): (1) fitness-based rescue and (2) size-

based rescue. We also estimated time to rescue across treatments, defining time to rescue as

the first of the three generations where the criterion was met (e.g., a population exceeding

original size in generations 10-12 would be considered rescued in generation 10).

We estimated treatment effect sizes on the probabilities of extinction and rescue in the

50-generation trials using Bayesian generalized linear models (GLMs) in the R package

rstanarm, 2.21.1 (Goodrich et al. 2020). We modeled both probabilities of extinction and rescue

for each simulated population as a Bernoulli response and each of our three treatments

(density dependence, initial size, genetic diversity) as categorical variables. Additionally, we

estimated effects of initial degree of maladaptation, an important determinant of extinction

risk (Gomulkiewicz and Holt 1995, Barfield and Holt 2016) by including the mean population

genetic load in the initial generation ($\theta - \bar{g}_0$, henceforth, "maladaptation") as a continuous

explanatory variable. This was possible for our stochastic model because sampling variation in

the founding of populations produced random among-trial variation in mean initial genotype.

We estimated the effects of negative density dependence and its interactions with other

variables by fitting a model with a four-way interaction and examining effect sizes as described

in Supporting Information E. The uncertainty around these estimates was sufficiently low and

could be arbitrarily improved by running more simulations. An effect size for a categorical

variable (for example, initial population size) gives the effect of changing that variable's

parameter on the log odds of extinction (or rescue) while holding all other parameters

constant, while an effect size for maladaptation gives the change in the log odds of extinction

(rescue) associated with switching five alleles from the positive allele to the negative allele in all

individuals.

To test for the presence of an extinction vortex we first reversed and re-centered time

for each extinct population to give τ, the time until extinction, such that populations had a

common origin ($\tau = 0$) with consistent meaning across populations (Fagan and Holmes 2006).

We define τ such that $\tau = 0$ is the first time step where the population is extinct (i.e., population

size of 0 or 1), $\tau = 1$ is the last time step in which the population size is above 0 or 1, etc. The

extinction vortex is characterized by accelerating declines in population size, genetic variation,

and/or rate of adaptation as τ approaches zero (Gilpin and Soulé 1986, Fagan and Holmes

2006). Specifically, we calculated the rate of population growth ($r_\tau = ln(N_\tau/N_{\tau+1})$), the

proportional loss of genetic diversity ($v_\tau = 1 - ((\sigma_a^2)_\tau/(\sigma_a^2)_{\tau+1})$), and the rate of adaptation

$(1 - k_\tau)$. k_τ is defined as in Gomulkiewicz and Holt (1995) such that $k_\tau = (\theta - \bar{g}_\tau)/(\theta - $

$\bar{g}_{\tau+1}$); as ($\bar{g}_\tau - \bar{g}_{\tau+1}$) approaches zero (i.e., as the rate of genotypic change approaches zero),

k_τ approaches 1 and $1 - k_\tau$ approaches 0. If adaptation proceeds at a constant rate, then k_τ

will be constant over time. We use these normalized rates of change instead of non-normalized

state variables (e.g., r_τ instead of N_τ) to facilitate comparisons across treatments and to remove

potential biases due to time until extinction. We estimated the ensemble means $\langle r_\tau \rangle$, $\langle v_\tau \rangle$, and

$1 - \langle k_\tau \rangle$ for all extinct populations in each treatment group using the 15-generation

simulations, where all state variables were tracked for each population. Following Fagan and

Holmes (2006), we considered decreasing $\langle r_\tau \rangle$ (accelerating decline in population size),

increasing $\langle v_\tau \rangle$ (accelerating loss of genetic diversity), or decreasing ($1 - \langle k_\tau \rangle$) (slowing rate of

adaptation) in the lead up to extinction (i.e., as τ approaches zero) to be evidence of an

extinction vortex. Likewise, we assessed how the shape of the extinction vortex varies between

density-dependent and density-independent populations qualitatively by noting differences in

the mean trajectories of the respective experimental treatments. We did not estimate rates for

τ = 0 because r = -∞ for a population at size zero and because additive genetic variance and

mean genotype are undefined for extinct populations.

Results

Population size

For all combinations of genetic diversity and initial population size, in each generation

density-dependent populations were smaller on average than corresponding density-

independent populations (compare black and purple lines, Fig. 2). As populations declined, the

minimum average size (bottom of the "U") reached by density-dependent populations was

smaller than that of density-independent populations across treatments, with larger effects in

large populations (two-thirds reductions in minimum size) than small populations (one-third

reduction). Density-dependent populations were also smaller on average at the end of 15

generations; populations with low diversity had slightly larger reductions in final size (92-95%

reduction) than populations with high diversity (85 - 90% reduction). Regardless of starting

population size, density-dependent populations declined for longer and took longer to recover

their original size (dotted line, Fig. 2), if at all.

Extinction and rescue

These differences in population size and growth rates over 15 generations translated

into more frequent extinctions for populations with density-dependent growth over the course

of the 50-generation simulations (Fig. 3). For both density-dependent and density-independent

populations, most extinctions occurred in the first 20 generations (Fig. 3*A*,*B*). Density

dependence increased extinction probabilities primarily in large populations (Table 2): among

large populations, density dependence increased the cumulative odds of extinction 5.0-fold for

high-diversity populations and 3.6-fold for low-diversity populations (Fig. 3*Ci*,*ii*). Among small

populations, density dependence increased the cumulative odds of extinction 1.5-fold and 2.0-

fold for high- and low-diversity populations, respectively (exponentiated effect sizes, Table 2,

Fig. 3*Ciii*,*iv*). Greater initial maladaptation (i.e., a mean initial genotype further from θ) was

associated with increased extinction risk across all treatments (Table 2, Fig. 3*D*). The effect of

maladaptation on extinction was stronger for large populations than small populations (Table

2); in large populations with high diversity in particular, only the most strongly maladapted

populations had considerable extinction risk (Fig. 3*Di*). Rather than strongly affecting the

relationship between maladaptation and extinction risk (i.e., the slope of the regression line,

Fig. 3*D*), density dependence increased extinction risk for all genotypes and only marginally

affected the relationship between genotype and maladaptation (i.e., raised the intercept while

slightly lowering the slope, Table 2, Fig. 3*D*). The combined result was that in large populations

with density-independent growth, only populations that were highly maladapted were at risk of

going extinct, but under density-dependent growth all populations had substantial extinction

risk (Fig. 3*D*).

The results for probability of rescue were qualitatively identical to those for probability

of extinction but with directions switched (Table 3 and Table F1, Figure F1). Likewise, the results

were qualitatively similar regardless of the type of rescue (i.e., fitness-based rescue or size-

based rescue). For both types, the probability of rescue declined with density dependence and

density dependence had stronger effects in large populations than small ones. Density

dependence increased mean rescue time only slightly (less than 1.5 generations) for all

population treatments except for large populations, for which density dependence increased

the time to return to initial size by 9-10 generations (Fig. F2). Notably, many small populations

were rescued instantaneously, i.e., they achieved three generations either exceeding initial size

or $\bar{R} > 1$ (or both) either instantly or after one time step (Table 3, Fig. 3*A*). These

"instantaneous rescues" happened primarily in small populations, comprising up to 16% of

rescue events observed in small population treatments. Both the overall frequency and the

proportion of total rescues comprising these instantaneous rescues decreased with low genetic

diversity and with density dependence. Pooling across all large treatment populations, only four

populations achieved instantaneous rescue of either type, comprising less than 1% of rescue

episodes for all large treatments. Extinctions occurred after both fitness- and size-based rescue

(Table 3). Extinction after rescue was more common for small populations than large ones, low-

diversity populations than high-diversity ones, as well as more common under density

dependence than density independence for both size- and fitness-based rescue.

Genetic variance, fixation, and fitness

Over the first 15 generations, mean additive genetic variance, $\langle(\sigma_a^2)_t\rangle$, declined for all

treatments (Fig. 4*A*). Surviving density-dependent populations lost more genetic variation than

density-independent populations (Fig. 4*A*, solid black *vs.* solid purple lines); this effect was

greater in large populations. Across treatment and throughout time, populations that went

extinct had lower mean genetic variance prior to extinction than populations that survived (Fig.

4*A*, solid *vs.* dashed lines). This loss of genetic variation was associated with fixation for both

positive (adaptive) and negative (maladaptive) alleles (Fig. 4*B,C*). Populations that went extinct

typically had on average more loci at fixation for either allele than surviving populations (Fig.

4*B,C*, solid *vs.* dashed lines), although in large populations this difference did not appear until

the fifth generation, when populations approached minimum size. Of populations that survived,

density-dependent populations had higher rates of fixation for the positive allele than density-

independent populations (Fig. 4*B*, solid black *vs.* solid purple lines). Conversely, surviving

density-dependent and density-independent populations both typically had few loci at fixation

for the negative allele (Fig. 4C). Differences in the trajectories over time of genotype and

intrinsic fitness between density-dependent and density-independent populations conditioned

on survival or extinction were slight (Figs. G2, G3).

The extinction vortex

Extinct populations in all treatments within the 15-generation simulations had negative

mean population growth rate ($\langle r_\tau \rangle$, Fig. 5A) and a non-zero mean loss of genetic diversity ($\langle v_\tau \rangle$,

Fig. 5B) for each time step. Declines in population growth rates and rate of loss of genetic

diversity accelerated, as expected for an extinction vortex (Gilpin and Soulé 1986, Fagan and

Holmes 2006) in the three generations prior to extinction ($\tau \leq 3$), with populations losing an

average of 17% of their standing additive genetic variance in the final generation preceding

extinction. In contrast, the rate of adaptation ($1 - \langle k_\tau \rangle$) decreased steadily over time in all

treatments rather than accelerating immediately prior to extinction (Fig. 5C). In the three

generations prior to extinction trends in density-dependent and density-independent

populations were similar. Four or more generations prior to extinction ($\tau > 3$), large density-

dependent populations had faster population decline (i.e., more negative $\langle r_\tau \rangle$) than large

density-independent populations, for which the rate of population decline was constant over

time (Fig. 5A). In contrast, density-dependent and density-independent populations had similar

rates of loss of genetic diversity (all differences in $\langle v_\tau \rangle$ between density-dependent and density-

independent treatments within 0.019, Fig. 5B). In the generation prior to extinction,

populations in all treatments had a mean adaptation rate of approximately zero (i.e., $\langle k_{\tau=1} \rangle \approx$

1 and $\langle \bar{g}_{\tau=1} \rangle \approx \langle \bar{g}_{\tau=2} \rangle$). Figures of non-normalized state variables conditioned on generation of

extinction, revealing qualitatively similar results, are available in Supporting Information H.

Discussion

We found that evolutionary rescue is constrained by the combined effects of negative

density dependence and loss of genetic variation. For all treatments, negative density

dependence decreased population growth rates, causing populations to decline more quickly,

reach lower densities, and stay at low densities for longer. This accelerated the loss of additive

genetic variance and slowed rates of adaptation. Ultimately, these demographic and genetic

processes often reinforced each other to produce an extinction vortex, precluding evolutionary

rescue. Here we discuss the effects of negative density dependence, loss of genetic variation,

and the extinction vortex on evolutionary rescue.

How negative density dependence constrains rescue

Although negative density dependence increased extinction risk for all treatments (Fig. 3,

Table 2) and correspondingly reduced incidence of evolutionary rescue (Table 3), it had

disproportionate effects on large populations with relatively low maladaptation to the novel

environment. In the density-independent model of Gomulkiewicz and Holt (1995) and in

laboratory experiments in density-independent settings (Bell and Gonzalez 2009, Hufbauer et

al. 2015) large and better-adapted populations had minimal extinction risk. Our model agrees

with these previous findings: extinction was rare in large, density-independent populations that

were not strongly maladapted (Fig. 3, Table 1). Likewise, our results agree with Chevin and

Lande (2010) that density dependence hastens initial population declines and slows subsequent

population recovery during rescue (Figs. 2, 3*A*). We extend and refine these previous findings

by highlighting that density dependence increased extinction risk and reduced evolutionary

rescue more for large populations than small ones, narrowing the considerable advantage in

survival odds that large initial size confers in density-independent settings (Fig. 3*C*, Table 2).

Furthermore, under density dependence even large populations somewhat adapted to the

novel environment might go extinct (Fig. 3*D*). Thus, populations considered at low risk by

classic, density-independent models face a non-trivial threat of extinction from environmental

change under more realistic density-dependent population growth.

The effects of negative density dependence on population size, population growth rates,

and extinction primarily occurred during the initial generations after environmental change.

Interestingly, negative density dependence had only minimal effects on extinction and rescue

once populations were sufficiently large. Populations with mean intrinsic fitness greater than

one that were large enough to avoid extinction *via* demographic stochasticity approached a

carrying capacity (Fig. 3*A*). In this event, when populations again declined in size through

stochasticity, they regained positive growth rates as the effects of density dependence

weakened, allowing them to return to carrying capacity. On the timescales we modeled, no

extinctions occurred during the period after rescued populations reached carrying capacity

(generations 20 – 50 in our simulations). However, over longer timescales, populations that

fluctuate around a carrying capacity might be susceptible to drift (Frankham et al. 1999) or have

a low supply of beneficial mutations (Orr and Unckless 2008, Osmond and de Mazancourt

2013), reducing future evolutionary potential (Forester et al. 2022). Furthermore, if there is

future environmental change that necessitates further adaptation, populations at a constrained

size (e.g., a carrying capacity) will begin adaptation from a smaller size than populations with

unconstrained growth.

While our model shows that density dependence hinders adaptation through increased

genetic erosion, this result is sensitive to how competition interacts with selection. In our

model, density dependence acts on all individuals equally; that is, the reduction in fitness an

individual faces under competition does not depend on its phenotype. However, if the outcome

of competition depends on fitness, for example by more-fit individuals excluding less-fit

individuals from resources, then competition could aid, rather than hinder, rescue by

accelerating the rate of adaptation (Schluter et al. 1985, Osmond and de Mazancourt 2013).

This explains an apparent contradiction of our results with those of Vinton and Vasseur (2020),

who found that density dependence assisted the ability of populations to track a moving

environment; their model also featured density dependence that led to stronger demographic

penalties for more maladapted individuals. Furthermore, adaptation may even strengthen

competition by increasing per capita resource use and lowering carrying capacities (Bisschop et

al. 2022). Intraspecific competition can also increase the severity of inbreeding depression (Yun

and Agrawal 2014); in a case where purging removes few deleterious alleles, this severe

inbreeding depression can reduce population growth rates (Bozzuto et al. 2019), further

endangering populations facing genetic erosion as they adapt to novel conditions.

How genetic erosion constrains rescue

Adaptation to environmental change was accompanied by a loss of genetic diversity and

potential for adaptation not captured in most models (but see Boulding and Hay 2001, Barfield

and Holt 2016). Quantitative genetic models often assume populations are large for analytical

tractability (Lande 1976); under this assumption fixation and loss of alleles are rare as there is

only marginal sampling variation and thus minimal drift. However, population decline is by

definition part of the rescue process, and our model demonstrates that additive genetic

variance is lost even in populations that survive environmental change (Fig 4*A*). This loss of

genetic diversity was associated with slowing rates of adaptation, contributing to the prolonged

time until rescue. This loss was partly driven by fixation of both adaptive and maladaptive

alleles, typically occurring when populations were at their smallest size (Fig. 4*B,C*). This matches

theoretical results of Boulding and Hay (2001) as well as empirical results of Rêgo et al. (2019)

in a seed beetle adapting to a novel host plant. Likewise, laboratory studies with *T. castaneum*

adapting to novel conditions found that accumulating genetic load in populations likely slows

the rate of adaptation (Stewart et al. 2017) and that more genetic diversity likely assists

population recovery through higher rates of recombination (Agashe et al. 2011). Models of

evolutionary rescue should thus include genetic erosion, as environmental stress (Nowak et al.

2009) and inbreeding due to bottlenecking (Frankham et al. 1999) are known to erode standing

genetic variation (reviewed in Biljsma and Loeschke 2011). Barfield and Holt (2016) were

among the first to model heritability as a function of population size. They similarly found a

long-term decline in heritability (analogous to additive genetic variance in our model) in extinct

populations, which they attributed to favored alleles approaching fixation. However, we did not

find evidence of the initial transient increase in heritability they reported; in our simulations

genetic variance declined most in early generations as populations were shrinking, driven by

fixation of not just adaptive but also of maladaptive alleles (Fig. 4). It is possible this difference

is due to our simulations being initialized with both alleles present at roughly equal frequencies,

which maximizes additive genetic variance (Equation 1). We ran additional simulations with the

negative (maladaptive) alleles at higher initial frequencies and found that genetic variance

remained approximately constant for 2-3 generations before declining, a result more similar to,

yet still distinct from, Barfield and Holt's (2016) results (Supporting Information I).

Our simulations were on short timescales compared to other studies (e.g., 50,000 time

steps in Osmond and de Mazancourt 2013) but reductions in population size and genetic

diversity could be important in the long term even after populations have been rescued. An

effective population size on the order of hundreds is needed to reduce longer-term extinction

risk due to accumulation of deleterious alleles (Lynch et al. 1995, Whitlock 2000, Frankham et

al. 2014). Specifically, populations need to spend long durations at large size to offset episodes

of drift and genetic erosion experienced even over short periods (Whitlock 2000), e.g., during

the "trough" or low point in population size following environmental change. In our

simulations, not only were density-dependent populations held at carrying capacity, potentially

precluding this re-establishment of variation, but they also spent more time at smaller densities

before being rescued (Figs. 2, 3*A*), leading to a greater erosion of variation (Fig. 4). Indeed, in

our simulations we saw as many as 40% of populations in a treatment group that were

"rescued" under our criteria go extinct within 50 generations, with stronger effects in

populations with initially low genetic diversity (Table 3). Furthermore, population size and

remaining standing genetic variation are important for persisting through future environmental

changes. The erosion of genetic variation, particularly of populations that experience severe

decline, is akin to a genetic bottleneck that would hinder the population's ability to adapt to

future environmental changes (Frankham et al. 1999, Whitlock 2000), particularly for changes

that are unidirectional or predictable (Lande and Shannon 1996). In our simulations, we

observed fixation of the positive allele in surviving populations (Fig. 4*B*), suggesting that

adaptive fixations did not guarantee extinction. However, given a future environmental change

favoring the other allele, its loss following the first environmental change event would hinder

subsequent adaptation (Lyberger et al. 2021).

How an extinction vortex impedes evolutionary rescue

Our model shows that challenging environments can draw populations into an extinction

vortex (Gilpin and Soulé 1986). This vortex was characterized by multiple generations of

accelerating decreases in population growth rate (Fig. 5*A*) and loss of genetic diversity (Fig. 5*B*),

the latter associated with a slowing rate of adaptation (Fig. 5*C*). Deterministic models often

feature a critical density below which extinction due to demographic stochasticity is very likely

(e.g., Gomulkiewicz and Holt 1995, Chevin and Lande 2010); our stochastic model allowed us to

observe this process more mechanistically as we directly modeled the small-population

processes. Rather than change the rate of population decline in the generations immediately

preceding extinction, density dependence widens the opening to the vortex, as populations are

more likely to quickly decline to small size several generations before extinction (Fig. 5*A*). Gilpin

and Soulé's (1986) concept of the extinction vortex did not assume a population is initially

severely maladapted to its environment, only that small population size increases susceptibility

to drift and inbreeding. In contrast, evolutionary rescue is defined by populations facing severe

maladaptation that pushes mean fitness below the rate of replacement. Thus, rather than an

extinction vortex arising by a population slowly becoming less fit due to drift, a population

experiencing environmental change immediately becomes less fit and also faces an extinction

vortex *via* drift and stochasticity that further reduces its ability to sufficiently adapt to the new

environment (Fig. 5*B,C*). For managers of small, at-risk populations that are susceptible to

environmental change and extinction, alternative strategies such as assisted migration may be

needed to pull populations out of the self-reinforcing demographic and genetic processes of the

vortex (Hufbauer et al. 2015, Bell et al. 2019).

Rapid rescue in small populations due to founder sampling variation

One surprising result was that upon environmental change, some small populations already

had a large proportion of phenotypes adapted to the environmental change due to sampling

variation, allowing them to be rescued very quickly (Fig. 3*D*, Table 3). In contrast, large

populations were rarely rescued immediately (Table 3) because the average initial phenotype

was less variable across populations (Fig. 3*D*). This appears to run counter to theoretical

arguments that larger populations are more likely to have novel variation compared to small

populations due to increased likelihood of individuals from the tail of the phenotypic

distribution being present in the population (Bell 2013). However, distinguishing between

variation in the population mean and variation among individuals within a population shows

that both phenomena are possible. In small populations, mean population phenotypes may be

initially closer to the new optimum due to sampling variation and drift (Fig. 3*D*) and this relative

abundance of adapted phenotypes can rapidly move the population to this new optimum.

Large populations are more likely to have individuals close to the new environmental optimum,

but it may take several generations for their alleles to dominate the mean population

phenotype due to the large number of maladapted individuals also initially present. Although

small populations were more likely to experience these very rapid rescue episodes, in the long

run larger populations were more likely to be rescued (Table 3). In part, this is because

sampling variation in small populations was equally likely to lead to an over-representation of

maladaptive alleles leading to a large disadvantage following environmental change (Fig. 3*D*).

Furthermore, considering that on the whole rescue is less likely to occur in small populations

compared to larger ones, the fact that some small populations were rescued due to a lucky

initial state underscores the degree to which small populations with unfavorable initial

phenotypes are at a disadvantage when the environment changes.

Implications for conservation biology

We introduce a more biologically realistic model for a more complete and applicable

understanding of population rescue after environmental change, providing better prediction of

potential outcomes and guidance for conservation and management. In our model, we

demonstrate that negative density dependence impedes the process of persisting in a novel

environment, and that large and well-adapted populations previously predicted to be at

minimal risk (Gomulkiewicz and Holt 1995, Bell and Gonzalez 2009, Barfield and Holt 2016) do,

after all, face non-trivial risk of extinction under density dependence. For managers, one

prescription is to increase carrying capacities, which will alleviate reductions in growth rates as

populations recover from environmental change. This can be done by increasing resource and

habitat availability. Although our simulated experimental design only featured density

dependence as a binary treatment, in natural settings it is a continuous variable that managers

can sometimes control. Because expected time to extinction increases exponentially with

carrying capacity (Lande 1993), decreasing the strength or degree of competition could

drastically improve the long-term fates, particularly of large populations previously perceived as

having low extinction risk. A corollary is that, in addition to other predictors of extinction risk

under rescue (Bell and Gonzalez 2009, Kopp and Matuszewski 2014), the carrying capacity or

degree of intraspecific competition should be considered when assessing risks to populations

subject to novel habitat alteration, degradation, or reduction. Furthermore, managers

implementing strategies to maximize evolutionary potential (Forester et al. 2022) should

consider how competition and population size may influence adaptation under environmental

or climate change. These predictions highlight the effects of density dependence and loss of

genetic diversity on populations exposed to environmental change, which is crucial information

in an age of global anthropogenic changes that put populations at unprecedented risk of

extinction.

Acknowledgements

The authors would like to thank Robert Holt for helpful suggestions following an early

presentation on work from this project and Ruth Shaw for feedback on Figure 1. This work was

supported by NSF award 1930222 to BAM and 1930650 to RAH. SWN was also supported by

NSF IGERT award 114807 while working on this project and LFD was also supported by NSF

Graduate Research Fellowship number 006784.

Data and code availability

Data and simulation code to produce data and analysis will be archived on Dryad upon

acceptance of the manuscript. Code and output files are also stored on GitHub at the address

https://github.com/melbourne-lab/evo_rescue_ndd_erosion.

References

Agashe, D., Falk, J.J., and Bolnick, D.I. 2011. Effects of founding genetic variation on adaptation

to a novel resource. Evolution, 65(9): 2481 – 2491.

Arnaud, L., Gage, M.J.G., and Haubruge, E. 2001. The dynamics of second- and third-male

fertilization precedence in *Tribolium castaneum*. Entomologia Experimentalis et Applicata,

99(1): 55 - 64.

Barfield, M., and Holt, R.D. 2016. Evolutionary rescue in novel environments: towards

improving predictability. Evolutionary Ecology Research, 17: 771 – 786.

Bell, G. 2013. Evolutionary rescue and the limits of adaptation. Philosophical Transactions of the

Royal Society of Biology, 368: 20120080.

Bell, G., and Gonzalez, A. 2009. Evolutionary rescue can prevent extinction following

environmental change. Ecology Letters, 12: 942 – 948.

Bell, D.A., Robinson, Z.L., Funk. W.C., Fitzpatrick, S.W., Allendorf, F.W., Tallmon, D.A., and

Whiteley, A.R. 2019. The exciting potential and remaining uncertainties of genetic rescue.

Trends in Ecology and Evolution, 34(12): doi:10.1016/j.tree.2019.06.006.

Bijlsma, R., and Loeschcke, V. 2012. Genetic erosion impedes adaptive responses to stressful

environments. Evolutionary applications, 5: 117 – 129.

Bisschop, K., Alzate, A., Bonte, D., and Etienne, R.S. 2022. The demographic consequences of

adaptation: evidence from experimental evolution. The American Naturalist, 199(6): 729 – 742.

Bonnet, T., et al. 2022. Genetic variance in fitness indicates rapid contemporary adaptive

evolution in wild animals. Science, 376(6596): 1012 - 1016.

Boulding, E.G., and Hay, T. 2001. Genetic and demographic parameters determining population

persistence after a discrete change in the environment. Heredity, 86: 313 – 324.

Bozzuto, C., Biebach, I., Muff, S., Ives, A.R., and Keller, L.F. 2019. Inbreeding reduces long-term

growth of Alpine ibex populations. Nature Ecology and Evolution, 3: 1359 – 1364.

Carlson, S.M., Cunningham, C.J., and Westley, P.A.H. 2014. Evolutionary rescue in a changing

world. Trends in Ecology and Evolution, 29(9) 521 – 530.

Chevin, L-M., and Lande, R. 2010. When do plasticity and genetic evolution prevent extinction

of a density-regulated population? Evolution, 64(4): 1143 – 1150.

Fagan, W.F., and Holmes, E.E. 2006. Quantifying the extinction vortex. Ecology Letters, 9: 51 –

60.

Forester, B.R., Beever, E.A., Darst, C., Szymanski, J., and Funk. W.C. 2022. Linking evolutionary

potential to extinction risk: Applications and future directions. Frontiers in Ecology and

Evolution, 1 - 9. http://doi.org/10.1002/fee.2552

Frankham, R., Lees, K., Montgomery, M.E., England, P.R., Lowe, E.H., and Briscoe, D.A. 1999. Do

population size bottlenecks reduce evolutionary potential? Animal Conservation, 2(4): 255 –

260.

Frankham, R., Bradshaw, C.J.A., and Brook, B.W. 2014. Genetics in conservation management:

revised recommendations for the 50/500 rules, Red List criteria, and population viability

analysis. Biological Conservation, 170: 56 – 623.

Gilpin, M.E., and Soulé, M.E. 1986. Minimum viable populations: processes of extinction. In:

Conservation Biology: The Science of Scarcity and Diversity, ed. Soulé, M.E. Sinauer Associates,

Sunderland, MA.

Gomulkiewicz, R., and Holt, R.D. 1995. When does evolution by natural selection prevent

extinction? Evolution, 49(1): 201 – 207.

Goodrich, B., Gabry, J., Ali, I., and Brilleman, S. 2020. Rstanarm: Bayesian applied regression

modeling via Stan. R package, version 2.21.1. https://mc-stan.org/rstanarm.

Hedrick, P.W., and García-Dorado, A. 2016. Understanding inbreeding depression, purging, and

genetic rescue. Trends in Ecology and Evolution, 31(12): 940 – 952.

Hone, J., Duncan, R.P. and Forsyth, D.M. 2010. Estimations of maximum annual growth rates

(r_m) of mammals and their application in wildlife management. Journal of Applied Ecology,

47(3): 507 - 514.

Houle, D. 1992. Comparing evolvability and variability of quantitative traits. Genetics, 130: 195

– 204.

Hufbauer, R.A., Szűcs, M., Kayson, E., Youngberg, C., Koontz, M.J., Richards, C., Tuff, T., and

Melbourne, B.A. 2015. Three types of rescue can avert extinction in a changing environment.

PNAS, 112(33): 10557 – 10562.

Kopp, M., and Matuszewski, S. 2014. Rapid evolution of quantitative traits: theoretical

perspectives. Evolutionary Applications, 7(1): 169 – 191.

Lande, R. 1976. Natural selection and random genetic drift in phenotypic evolution. Evolution,

30: 314 – 334.

Lande, R. 1993. Risks of population extinction from demographic and environmental

stochasticity and random catastrophes. The American Naturalist, 142(6): 911 – 927.

Lande, R., and Shannon, S. 1996. The role of genetic variation in adaptation and population

persistence in a changing environment. Evolution, 50(1): 434 – 437.

Lyberger, K.P., Osmond, M.M., and Schreiber, S.J. 2021. Is evolution in response to extreme

events good for population persistence? The American Naturalist, 198(1): 44 – 52.

Lynch, M., and Walsh, B. 1998. *Genetics and the analysis of quantitative traits.* Sinauer

Associates, Sunderland, MA.

Lynch, M., Conery, J., and Bürger, R. 1995. Mutational meltdowns in sexual populations.

Evolution, 49(6): 1067 – 1080.

Melbourne, B.A., and Hastings, A. 2008. Extinction risk depends strongly on factors contributing

to stochasticity. Nature, 454: 100 – 103.

Nowak, C., Vogt, C., Pfenninger, M., Schwenk, K., Oehlmann, J., Streit, B., and Oetken, M. 2009.

Rapid genetic erosion in pollutant-exposed experimental chironomid populations.

Environmental Pollution, 157: 881 – 886.

Orr, H.A., and Unckless, R.L. 2008. Population extinction and the genetics of adaptation. The

American Naturalist, 172(2): 160 – 169.

Orr, H.A., and Unckless, R.L. 2014. The population genetics of evolutionary rescue. PLoS

Genetics, 10(8): e1004551.

Osmond, M.M., and de Mazancourt, C. 2013. How competition affects evolutionary rescue. Phil

Trans R Soc B., 368: 20120085.

Pai, A., and Yan, G. 2020. Long-term study of female multiple mating indicates direct benefits in

Tribolium castaneum. Entomologia Experimentalis et Applicata, 168(5): 398 – 406.

Pauls, S.U., Nowak, C., Bálint, M., and Pfenninger, M. 2013. The impact of global climate change

on genetic diversity within populations and species. Molecular Ecology, 22: 925 – 946.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rêgo, A., Messina, F.J., and Gompert, Z. 2019. Dynamics of genomic change during evolutionary

rescue in the seed beetle Callosobruchus maculatus. Molecular Ecology, 28: 2136 – 2154.

Ricker, W.E. 1954. Stock and recruitment. Journal of Fisheries of the Board of Canada, 11: 559 –

623.

Schluter, D., Price, T.D., and Grant, P.R. 1985. Ecological character displacement in Darwin's

finches. Science, 227(4690): 1056 – 1059.

Stewart, G.S., Morris, M.R., Genis, A.B., Szűcs, M. Melbourne, B.A., Tavener, S.J., and Hufbauer,

R.A. 2017. The power of evolutionary rescue is constrained by genetic load. Evolutionary

Applications, 10: 731 – 741.

Vinton, A.C., and Vasseur, D.A. 2020. Evolutionary tracking is determined by differential

selection on demographic rates and density dependence. Ecology and Evolution, 10: 5725 –

5736.

Wade, M.J., Shuster, S.M., and Stevens, L. 1996. Inbreeding: its effect on response to selection

for pupal weight and the heritable variance in fitness in the flour beetle, *Tribolium castaneum*.

Evolution, 50(2): 723 – 733.

Whitlock, M.C. 2000. Fixation of new alleles and the extinction of small populations: drift load,

beneficial alleles, and sexual selection. Evolution, 54(6): 1855 – 1861.

Yun, L. and Agrawal, A.F. 2014. Variation in the strength of inbreeding depression across

environments: effects of stress and density dependence. Evolution, 68(2): 3599 – 3606.

Figure 1: Event sequence diagram of evolutionary rescue (blue, ovals) and two connected extinction vortices (red, squares) that populations may face following harmful environmental change. The effects of negative density dependence, potentially impeding rescue and drawing populations into a vortex, are indicated in purple.

Figure 2: Mean population size in each generation for all treatments (n = 4000 for each treatment). Means in each generation include extinct populations as zeros. Shaded areas are twice the standard error on each side of the mean.

Figure 3: Population size and extinction characteristics of 1000 simulations per treatment group run for up to 50 generations. (A) mean population size of 25 randomly selected trials for each treatment. Dashed lines are extinct populations. Simulations were truncated for populations surpassing size 10000. (B) Instantaneous probability of extinction in each generation, conditioned on survival until that generation. Ribbons represent twice the standard error on both sides of the estimate. (C) Cumulative probability of extinction over time. Areas under the curve are shaded to highlight the difference between density-dependent and density-independent populations. (D) 200 posterior estimates per treatment of extinction probability as a function of initial degree of maladaptation, as predicted by a generalized linear model (Table 2). Marks on x-axis indicate the distribution of initial mean population genotypes for those treatments.

Figure 4: (A) Mean additive genetic variance and the mean number of loci at fixation (out of 25) for (B) positive and (C) negative alleles over time in surviving and extinct populations. Shaded areas represent twice the standard error on each side of the mean (the standard error is often too small to be visible on this plot).

Figure 5: Rates of change as a function of time to extinction (τ), demonstrating the presence of an extinction vortex, in (A) population growth, (B) loss of genetic variation, and (C) rate of adaptation. Small τ represents time steps closer to the time of a population's extinction. Shaded areas represent twice the standard error on both sides of the mean.

Tables

Table 1 Parameter values simulated for the three-way factorial simulation experiment

Factor of interest	Treatment	Value	Name in main text
Density dependence	Absent	$\alpha = 0$	density-independent populations
	Present	$\alpha = 0.0035$	density-dependent populations
Genetic diversity	High	$\sigma_a^2 \approx 0.5$	high-diversity populations
	Low	$\sigma_a^2 \approx 0.25$	low-diversity populations
Initial population size	Large initial size	$N_0 = 100$	large populations
	Small initial size	$N_0 = 20$	small populations

Table 2 Effect sizes of treatment variables changed while others are held constant on log odds of extinction in 50 generations and accompanying 95% credible intervals.

Variable changed	Held constant	Effect size (95% CI)
Density independent to dependent	Large, high diversity	1.60 (1.36, 1.85)
	Large, low diversity	1.27 (1.08, 1.46)
	Small, high diversity	0.41 (0.21, 0.61)
	Small, low diversity	0.71 (0.44, 1.00)
Large to small	High diversity, DI	2.82 (2.58, 3.08)
	High diversity, NDD	1.63 (1.44, 1.83)
	Low diversity, DI	2.13 (1.92, 2.35)
	Low diversity, NDD	1.58 (1.31, 1.85)
High to low diversity	Large, DI	1.67 (1.43, 1.91)
	Large, NDD	1.34 (1.15, 1.53)
	Small, DI	0.99 (0.77, 1.21)
	Small, NDD	1.29 (1.03, 1.56)
Increased maladaptation	Large, high variance, DI	6.65 (4.38, 8.99)
	Large, high variance, NDD	6.04 (4.16, 7.99)
	Large, low variance, DI	7.24 (4.77, 9.79)
	Large, low variance, NDD	6.19 (3.82, 8.55)
	Small, high variance, DI	3.76 (2.90, 4.63)
	Small, high variance, NDD	3.39 (2.40, 4.40)
	Small, low variance, DI	3.99 (2.46, 5.50)
	Small, low variance, NDD	4.48 (2.67, 6.34)

Table 3 Summary of rescue probabilities, proportion of rescue events occurring instantly, times to rescue, and extinction probabilities after rescue 50 generation simulations for both types of rescue.

Size and diversity	Prob. rescue		Prop. (number) instant rescue†		Mean time to rescue		Prob. extinct after rescue	
	DI	NDD	DI	NDD	DI	NDD	DI	NDD
	Fitness-based rescue							
Large, high	0.92	0.71	0.00 (3)	0.00 (0)	5.1	6.7	0.04	0.12
Large, low	0.69	0.41	0.00 (0)	0.00 (0)	8.3	9.2	0.10	0.24
Small, high	0.44	0.36	0.16 (68)	0.06 (23)	4.0	4.7	0.21	0.27
Small, low	0.24	0.16	0.04 (11)	0.01(2)	5.7	7.0	0.31	0.39
	Size-based rescue							
Large, high	0.89	0.62	0.00 (1)	0.00 (0)	11.2	20.3	0.00	0.00
Large, low	0.62	0.26	0.00 (0)	0.00 (0)	17.4	27.7	0.00	0.00
Small, high	0.37	0.29	0.16 (61)	0.05 (15)	6.3	7.9	0.06	0.07
Small, low	0.19	0.12	0.13 (24)	0.08 (9)	7.9	8.6	0.12	0.17

† proportion instant rescue is the proportion of rescued populations in the treatment where rescue occurred in 0 or 1 generations; the number is the total number of observed populations (out of 1000) in the treatment experiencing instantaneous rescue