1 ORIGINAL ARTICLE

3 Disentangling the role of NAc D1 and D2 cells in hedonic eating

4 Mathilde C. C. Guillaumin, Paulius Viskaitis, Ed Bracey, Denis Burdakov and Daria Peleg-Raibstein

5 Institute for Neuroscience, Department of Health Sciences and Technology, Swiss Federal Institute of
6 Technology, ETH Zurich, 8603 Schwerzenbach, Switzerland

7 Correspondence e-mail : daria-peleg@ethz.ch

10 Abstract

Consumption of highly palatable foods is thought to be influenced by the central reward circuits that
control both the hedonic component ('liking') of food and the motivation ('wanting') to eat it. The
nucleus accumbens (NAc), a key brain center implicated in these processes, contains two main
populations: D1 and D2 dopamine receptor-expressing neurons. The precise contribution of these cells
to these reward-associated behaviors is still unresolved and establishing a causal relationship between
activation of each neuronal subtype and its effect on motivated behaviors has proven challenging. To
understand the forces underlying eating behavior that may lead to overconsumption, we assessed
'liking' and 'wanting' in separate sets of experiments. First, in a free palatable food consumption
paradigm, we analyzed licking microstructure to demonstrate that both NAc D1 and D2 cell activity are
more strongly correlated with 'liking' than 'wanting'. Second, we evaluated motivation to retrieve a
reward ('wanting') in an operant task. Both NAc D1 and D2 cell activity encoded motivation during the
reward-approach phase, after the cue and before consumption. Lastly, we show that opto-excitation
of NAc D1 cells, but not D2 cells, intensifies incentive motivation to overconsume the optostimulation-
paired palatable food in a two-bottle choice task. This demonstrates an important role for NAc D1 cells
in preference formation and promoting consumption of palatable foods. Our results suggest a
fundamental difference: NAc D1 cells promote reinforced behaviors under an operant paradigm or a
Pavlovian association task. In contrast, NAc D2 cells are essential for initiating retrieval of palatable
food but, once consumed, reduced activity of these cells allows prolonged eating episodes.

30 Keywords

31 Motivation, dopamine, D1, D2, nucleus accumbens, reward, fiber photometry, optogenetics, food
32 preference

Introduction

Overweight and obesity, and eating disorders generally, have increased in recent decades (Hoek, 2016; Tavolacci et al., 2021). Thus, it is crucial to better understand the factors that contribute to the development of different eating disorders. In the modern obesogenic environment, palatability and availability of foods play a major role in the development of disorders of overeating (Corsica and Hood, 2011; King, 2013). Food consumption is not only motivated by the homeostatic need to monitor, restore and maintain energy balance. Homeostatic signals can also be overridden, causing us to engage in non-homeostatic eating that is driven by the sensory and/or hedonic properties of palatable foods with high fat and/or sugar contents. This 'hedonic' eating, activates brain regions involved in reward, resulting in increased motivation to consume food, and increased energy consumption (Frank et al., 2019; Leigh and Morris, 2018; Volkow et al., 2011) despite a state of satiety (Zheng and Berthoud, 2008).

The 'incentive salience theory' of reward has provided a valuable framework for more than a decade for investigating the role of food hedonics in the context of appetitive behavior in humans and animals (Berridge and Robinson, 1998, 2003). This theory postulates that reward is not a unitary process but comprises of an affective pleasure component referred to as 'liking' and a non-affective motivational component referred to as 'wanting'. 'Liking' is the hedonic component that reflects the immediate experience or anticipation of eating a pleasurable food (Mela, 2006). 'Wanting' is the incentive motivation (reward-seeking) that can lead to increased appetite, food cravings and also to overconsumption of food (Berridge and Robinson, 1998; Berthoud and Morrison, 2008; Finlayson et al., 2007). 'Incentive salience theory' has been studied in the framework of the regulation of food intake, specifically as a model for interpreting studies linking the function of the reward system with food intake or obesity. Looking at these psychological traits in eating behaviors (Berridge, 1996, 2009; Finlayson and Dalton, 2012), overeating in some individuals might reflect the abnormal functioning of either the 'wanting' or 'liking' mechanisms, and there are indications from subjects with eating disorders that 'liking' and 'wanting' of obesogenic foods can also be dissociated (Benton, 2010).

It is widely known that dopaminergic neurotransmission in the nucleus accumbens (NAc) plays a key role in behaviors related to natural rewards and drugs of abuse (Hyman and Malenka, 2001; Kelley and Berridge, 2002; Koob and Le Moal, 2001; Robinson and Berridge, 2003; Soares-Cunha et al., 2016b). In the NAc, two important classes of cells are medium spiny neurons expressing D1 dopamine receptors (D1 cells) and those expressing D2 dopamine receptors (D2 cells) (Surmeier et al., 1996). The precise contribution of D1 and D2 cells in reward-associated behaviors is still unresolved, and the establishment of a causal relationship between activation of each neuronal subtype and its effect on motivated behaviors has proven challenging. D1 cell activation is canonically considered to be related to positive rewarding events leading to persistent reinforcement, while activation of D2 cells is thought to facilitate aversion (Kupchik et al., 2015; Lobo et al., 2010; Trifilieff et al., 2013). However, the current literature is controversial regarding the functionality of these cells and their role in motivation-related behaviors (Natsubori et al., 2017; Soares-Cunha et al., 2016a; Soares-Cunha et al., 2020; Yang et al., 2018).

Therefore, increased understanding of which cells in the dopaminergic system underly brain substrates of 'liking' and 'wanting' may open an avenue to comprehending the impact of food rewards on eating behaviors linked to obesity and other eating disorders. In addition, it may provide insight into brain circuits and mechanisms underlying both hedonic and motivational components of reward, including mechanisms that are common to both natural and artificial rewards.

This paper aims to further understand the differential contribution of NAc D1 and D2 cells to different
components of palatable food consumption, namely 'liking' (hedonic/pleasurable) and 'wanting'
(incentive motivational value/desire/craving). We assess 'liking' and 'wanting' from several different
angles using multiple behavioral paradigms, while employing neuronal recording or optogenetic tools
and analytical techniques, enabling us to better understand the forces underlying eating behavior that
may lead to overconsumption.

Results

NAc D1 cells exhibit transiently elevated activity, and D2 cells exhibit suppressed activity, during
periods of free consumption of a palatable food reward

To probe the neural mechanisms underpinning reward consumption, we used *in vivo* fiber photometry
to record GCaMP6s calcium indicator fluorescence selectively expressed in D1 cells (in D1 receptor-cre
mice) or D2 cells (in adenosine A2a receptor gene (Adora2a)-cre mice) (Gerfen et al., 2013) (**Figure 1a**).

To determine how D1 and D2 cells represented solutions with different palatability, mice were first
given free access to water from a bottle (**Figure S1a**). Once they reliably consumed this, they were
allowed to freely consume a highly palatable food (milkshake, **Figure 1b,c**) over the course of five daily
sessions, in the absence of nutrient deficit.

We found that before licking, both D1 and D2 cells transiently increased their activity, which peaked
just after licking onset. In both D1 and D2 cells, the peak was more pronounced in response to
milkshake than water, possibly reflecting hedonic processes (**Figure 1b-d**). However, after
consumption onset, the activity patterns of the two populations differed; D1 cells returned to baseline,
while D2 cell activity was suppressed below the baseline as the licking bout continued (**Figure 1d**). Both
cell types continued to respond to milkshake over subsequent sessions, but the D1 peak reduced over
sessions, and the D2 suppression was exaggerated (**Figure 1e**). These results demonstrate that D1 and
D2 cells do not act as a single homogeneous population during consumption.

Milkshake consumption within a session (total number of licks) was significantly higher than water,
and this was consistent across sessions (**Figure S1c,d**). This indicates more positive hedonic reactions
to milkshake than water (**Figure S1d**). However, the amount consumed provides an incomplete
readout of ingestive behavior (Davis and Smith, 1992). Therefore, to further probe the driving forces
behind these changes in overall consumption, we investigated different aspects of free consumption
of water vs. a palatable food (milkshake). In particular, if it would be possible to differentiate between
'wanting' and 'liking' parameters (Berridge, 2009). We therefore analyzed licking microstructure
(Davis, 1989), since it provides several measures to evaluate pleasure ('liking') and motivation
('wanting') during liquid-food consumption in rodents (Davis and Smith, 1992; Dwyer et al., 2011;
Galistu and D'Aquila, 2013; Johnson, 2018a; Robinson and McCool, 2015). Specifically, increases in the
duration of single licks, licking bout duration and bout size (lick intensity determined by number of licks
in a bout), and decreases in inter-lick interval (ILI), have been suggested to represent increased hedonic
impact and palatability, while increases in bout number and decreases in inter-bout interval represent
increased incentive salience (Davis, 1973; Davis and Smith, 1992; Dwyer, 2012; Higgs and Cooper,
1998; Robinson and McCool, 2015; Smith, 2001)(**Figure S1b**).

We found that changes in 'liking' parameters paralleled greater 'liking' for milkshake over water; lick
duration, bout duration and bout size increased significantly, while inter-lick interval (ILI) decreased
significantly, between the water and first milkshake sessions (**Figure 1f and S1e**). Therefore, as
expected, animals displayed more positive hedonic reactions ('liking') to the milkshake than to water.

Lick duration also kept increasing over milkshake sessions, suggesting an increased hedonic
component of milkshake consumption in later sessions (**Figure 1f, left**).

Bout number increased, and inter-bout interval (IBI) decreased significantly between water and the
first milkshake session, suggesting that mice were more motivated ('wanted') to consume milkshake
(**Figure 1g**). IBI also kept decreasing across milkshake sessions, suggesting 'wanting' for milkshake
increased over sessions (**Figure 1g, right**).

We then performed multiple regression analysis to investigate whether *in vivo* calcium activity of NAc
D1 and D2 cells during consumption of milkshake across all 5 sessions could predict the level of 'liking'
or 'wanting' of the palatable food consumed. This analysis can determine key photometry signal
parameters, among the multiple parameters that exist, to predict specific aspects of palatable food
consumption. In D1 cells, the minimum and maximum signal values were significant predictors of bout
duration ('liking'), explaining around 30% of the variability in bout duration (as shown by the R^2 change
of models 1 and 2, **Figure S2a, top right**), with negative and positive betas, respectively. The R^2 change
values showed that the higher the initial increase (and to a smaller extent, the lower the subsequent
decrease), the longer the bout (**Figure S2a**). In D2 cells, we found that the area under the curve and
the minimum signal during the 3 sec following lick onset were significant predictors of bout duration,
explaining when combined around 40% of the variability in bout duration (**Figure S2b, top right**). More
specifically, the more the D2 cell signal decreases, the longer the bouts are and the higher the 'liking'
is (negative beta coefficients) (**Figure S2b**). Interestingly, in a similar analysis of inter-bout interval, IBI
('wanting' – **Figure S3**), the significant predictors in the photometry signals only explained a very small
amount of IBI variability and were thus poor predictors overall.

Together, these findings show that *ad libitum* consumption of milkshake induces both 'liking' and
'wanting' to eat. These two main components underlying palatable food consumption are more
positively associated with consuming milkshake than water. However, we detected a dissociation
between 'liking' and 'wanting', with the signals of both D1 and D2 cells being more strongly correlated
with 'liking' than 'wanting'. Specifically, the initial increase in NAc D1 cell activity at licking onset and
the sustained decrease in NAc D2 cell activity during eating are good predictors of 'liking' parameters.
149
Increasing NAc D2 cell activity increases consumption by increasing 'wanting' despite a decrease in
'liking'

We next employed optogenetics to investigate how the roles of NAc D1 and D2 cells differ during *ad*
libitum palatable food consumption. Based on previous findings (Kravitz et al., 2012; Lobo et al., 2010),
we initially hypothesized that activation of NAc D1 cells would lead to increased consumption, whereas
activation of NAc D2 cells – counteracting the natural suppression in activity described earlier – would
decrease 'liking'. However, it would not necessarily affect overall intake, depending on its effect on the
'wanting' aspect of consumption. To test our prediction, we injected the optogenetic Cre-dependent
excitatory actuator ChrimsonR or a control GFP virus, into the NAc of D1- or D2(A2a)-cre mice, and
implanted optical fibers bilaterally in the NAc (**Figure 2a**). Consistent with our photometry data and
our initial prediction, optogenetic activation of D2 cells contingent to eating decreased 'liking'
(increased ILI, **Figure 2c**). Unexpectedly, it increased 'wanting' (increased bout number, **Figure 2d**)
leading to increased overall consumption (increased number of licks, **Figure 2b**). This suggests that the
'wanting' aspect triggered by optical activation overrode 'liking' in driving overall consummatory
behavior. Conversely, contrary to our hypothesis, D1 cell activation did not impact overall consumption
via effects on either 'liking' or 'wanting' parameters (**Figure 2b-d**). This would mean that sustained
activity of D1 cells beyond the naturally occurring transient increase described in **Figure 1d-e** does not
promote further 'liking', nor does it induce 'wanting'.

Next, we sought to test whether silencing NAc D1 and D2 cells using the inhibitory opsin eNpHR3.0 could also influence consumption of a palatable food. We hypothesized that D1 cell silencing would induce a decrease in consumption, while D2 cell silencing would prolong consumption. However, silencing D1 cells did not lead to a significant decrease in overall consumption (**Figure 2b**), but induced a significant increase in 'liking' (decreased ILI, **Figure 2c**) with no significant effect on 'wanting' (**Figure 2d**). The influence of inhibitory post-ingestive mechanisms such as satiation, which increase as ingestion proceeds, could have been a confounding factor to D1 cell inhibition, because they may already have reduced lick number and bout number to their floor (D'Aquila, 2010; Sclafani and Ackroff, 2004).

Inhibition of D2 cells did not lead to any significant difference in lick or bout number compared to controls (**Figure 2c,d**), suggesting that a further decrease of the natural suppression in activity observed during consumption (**Figure 1d,e**) may not be possible and therefore may not yield any stronger consummatory effect.

Taken together, these results show that during *ad libitum* consumption, only D2 cell activation leads to an overall increase in eating, and this is mediated by an increase in 'wanting', despite a concomitant decrease in 'liking'. However, these findings are limited because an *ad libitum* paradigm only examines baseline motivation, and is not designed to evaluate motivation to work for a reward. Therefore, we next probed the role of D1 and D2 cells in a different motivational task designed to interrogate 'wanting'.

NAc D1 and D2 cell activity increases with anticipation during a motivational task to retrieve palatable food, which is followed by a D2 cell activity decrease upon consumption onset

To probe 'wanting' further, we paired reward with a cue, which allowed us to investigate the attribution of 'incentive salience' to the cue. We exposed mice to a Pavlovian 'motivation to retrieve a reward' task in which they received up to 15 milkshake rewards announced by a sound cue, which they had to retrieve from a food magazine. This did not require the mice to develop learning strategies. This allowed us to monitor three parameters: the number of entries to the food magazine, the time to retrieve and consume the milkshake after cue onset (latency), and the length of food-magazine entries. (**Figure 3a,b**).

Photometry recordings showed sustained increased activity in D2 cells during the approach phase (after milkshake delivery cue and before consumption; **Figure 3c, right**). D1 cells, however, only transiently increased their activity following milkshake delivery cue (**Figure 3c, left**). An important distinction in the behavior of D1 cell activity must be made: in contrast to the *ad libitum* paradigm where D1 cell signal increased prior to licking onset, here, in a motivational task, the increase in signal was specifically aligned to the reward delivery cue. During food magazine visits, i.e., during milkshake consumption, the neural dynamics observed were similar to the first *ad libitum* study, with a sustained decrease in the activity of D2 cells and only a small initial increase in D1 cells (**Figure 3d**).

As in the first experiment, we were interested to see if the natural D1 and D2 cell activity recorded with photometry holds any predictive information about components of consummatory behavior of a palatable food. Correlation analysis of 'wanting' behavioral parameters and D2 cell activity around food magazine visit after a milkshake delivery cue revealed that latency can be well predicted by the average D2 cell activity before the food magazine visit, and by the standard deviation (Std) of the pre-food magazine visit – **Figure S4a**). The higher the signal before the food magazine visit, the smaller the latency. This correlation analysis also showed that the number of food magazine visits can be well predicted by the median signal before the food magazine visit (**Figure S4b**). The higher the signal pre-food magazine visit, the more visits occurred. The average length of a food magazine visit was also

found to be well predicted by the minimum of D2 cell activity *during* consumption, and by the standard
deviation of the signal before the visit (**Figure S4c**); the lower the minimum, the longer the food
magazine visit.

Therefore, latency and number of visits correlated with D2 cell signals *before* visits occurred, while the
length of a visit was determined by the signals *during* the visit. These data also suggest that suppression
of D2 cell activity *during* feeding favors consumption, which was investigated further in the next
experiment.

Inhibition of D2 or D1 cells reduces motivation to retrieve a reward, while activation of only D1 cells
enhances motivation

Because both D1 and D2 cells were modulated during the motivation task (**Figure 3c,d**), we aimed to
causally test the effect of cell-type-specific activation and inhibition on motivated behavior ('wanting').
We optogenetically manipulated NAc D1 and D2 cells at either reward (milkshake) delivery cue or at
food magazine visit (first visit following a delivery cue) (**Figure 3a,b**). D2 cell silencing at either
milkshake delivery cue, or at food magazine visit significantly decreased number of food magazine
entries (**Figure 3e,f**), and inhibiting D2 cells after milkshake delivery cue significantly decreased the
average length of the subsequent food magazine visit (**Figure 3i**). These results are in line with our
photometry findings (**Figure 3c, right**) suggesting a role for D2 cell activity in the anticipation of a
reward ('wanting'). Optogenetic activation of D1 cells at either milkshake delivery cue or at food
magazine visit significantly increased the number of food magazine entries (**Figure 3e,f**). Moreover,
silencing D1 cells during the reward delivery cue increased the average latency (**Figure 3g**). This means
that inhibiting D1 cells during the approach phase delayed the approach. Thus, the behavioral results
obtained with D1 cell activation were expected based on the photometry findings (**Figure 3c,d, left**):
increase in D1 cell activity may reflect a higher incentive motivation to approach the palatable food,
causing the mice to be more focused on obtaining the reward and being more reactive to the cue.

Together, our data indicate that both D1 and D2 cell activity during the approach phase of a palatable
food (preceding consumption) encode incentive motivation or 'wanting'.

Modulation of NAc D1 and not D2 cells influence hedonic preference learning (hedonic shifting)

We showed in our two previous experimental paradigms that consumption of a palatable food
(milkshake) elicits, as expected, 'liking' responses. We investigated whether pairing milkshake,
constituting a conditioned stimulus (CS), with optogenetic stimulation (unconditioned stimulus, US) of
either D1 or D2 cells could shift natural preference for one flavor to another. We hypothesized that
optogenetic stimulation of either population would establish a learned preference for the least
innately preferred milkshake flavor by increasing the hedonic value ('liking') of the CS (milkshake).

To assess innate preference, mice were first presented with two bottles, each containing a different
flavor of milkshake, but with identical nutritional composition ('pre-test'). The least preferred flavor
was then coupled to optogenetic stimulation of either D1 or D2 cells during a daily conditioning session
over several days, while the preferred flavor was presented unpaired with optogenetic stimulation in
a separate daily session (**Figure 4a**). Following conditioning, we performed a 'choice-test' to determine
if the paired flavor was now able to trigger incentive motivation ('wanting') and become equally, or
possibly more, attractive than the previously preferred, unpaired flavor. During conditioning, increased
consumption from the paired milkshake flavor over the unpaired one would indicate a hedonic change
('liking'). During the choice test, with respect to the 'liking'/'wanting' hypothesis, disruption of a
reward-oriented behavior by optogenetic manipulation of the NAc could, in principle, be due to a
disruption of either 'liking' or 'wanting' or, presumably, a combination of both.

During optogenetic activation of D2 cells, consumption of the paired and unpaired flavors across the conditioning phase did not differ (**Figure 4c**). Note that during this conditioning phase, each flavor was presented in separate sessions, hence the absence of difference between the amounts consumed, unlike the difference observed in the 'pre-test' (**Figure S5a, right**). However, optogenetic stimulation of D1 cells led to a gradual increase in the consumption of the paired flavor, so that the mice consumed higher amounts of milkshake in the late paired conditioning sessions compared to GFP controls (**Figure 4b**). During the choice-test, an increase in consumption of the paired flavor compared to the pre-test levels was noted while a decrease was observed in the unpaired flavor, leading to significantly different 'choice-test – pre-test' deltas between the two flavors (**Figure 4d, left**). There was no significant change in preference in control GFP mice (**Figure 4d, right**), suggesting repeated exposure to the flavors did not alter preference.

In addition, we used lick microstructure analysis to assess potential changes in 'liking' and 'wanting' produced by flavor preference conditioning. During the conditioning sessions, stimulation of D1, but not D2 cells increased 'liking' (increased lick duration and decreased inter-lick duration) over time (**Figure 5a,b, left inserts**). This indicates an increased palatability of the flavor paired with D1 cell optostimulation, which is of relevance because conditioned increases in palatability have been shown to be important for driving overeating (Johnson, 2018b). Moreover, inter-bout interval decreased over conditioning sessions (**Figure 5d, left insert**), indicating an increase in 'wanting', while motivation to consume the unpaired flavor decreased over sessions, as indicated by decreasing bout numbers and initially increasing IBI (**Figure 5c,d, middle inserts**). During the choice-test, when both flavors were once again presented simultaneously (with no optogenetic stimulation on either of the flavors at this stage), licking microstructure indicated an increased 'liking' of the paired flavor (higher lick duration, and lower ILI, **Figure 5a,b, right inserts**) – effects not observable in the unpaired flavor. 'Wanting' was also increased for the paired flavor (increased bout number, decreased IBI, **Figure 5c,d, right inserts**) and decreased for the unpaired flavor (**Figure 5d, right insert**).

These results show an overall preference for the milkshake flavor coupled with D1 cell stimulation, thus demonstrating an important role for D1 cells in promoting the consumption of preferred palatable foods.

Discussion

Our study reveals distinct patterns of NAc D1 and D2 cell activity during different behavioral paradigms intended to assess different psychological processes of hedonic behaviors: hedonia ('liking' a food) and motivation ('wanting' the food). We found that NAc D1 and D2 cells do not act as a single homogeneous population in regulating these processes. In a free palatable food consumption paradigm a dual role of NAc D2 cells is linked to both 'liking' and 'wanting', where increased D2 cell activity signals 'wanting' and decreased activity enhances 'liking', with the magnitude of natural inhibition following palatable food consumption being a good predictor for 'liking'. In contrast, D1 cells promoted 'liking' during free consumption. In the operant task, we showed that NAc D1 cell activity after cue and right after eating onset drives the motivation to retrieve palatable food, while reduced activity of D2 cells after food magazine visits drives the motivation to remain in the magazine. Lastly, we show that NAc D1 cell stimulation induced hedonic shifting in a Pavlovian task and overconsumption of palatable food. This effect was specific to NAc D1 cells, as stimulation of NAc D2 cells during opto-paired food conditioning had no effect across conditioning days, or during the choice test (See results summary, **Figure 6**).

The role of NAc D1 and D2 cells during periods of free consumption of palatable food

The NAc is part of a neural circuit governing consumption of palatable food (Kelley et al., 2005) and
neuronal subpopulations in the NAc shell encode rewarding stimuli and palatability (Roitman et al.,
2005; Villavicencio et al., 2018). Pharmacological manipulations in rodents and in hyper-dopaminergic
knockout mice showed that 'wanting' and 'liking' sucrose could be separated with an increase in
'wanting' but a constant decrease in 'liking' (Berridge and Valenstein, 1991; Pecina et al., 2003; Smith
and Berridge, 2005; Wyvell and Berridge, 2000) and that 'wanting' was increased with prolonged
sucrose consumption (Avena et al., 2005). Here, using photometry and optogenetics, we showed a
differential role between NAc D1 and D2 cells during free consumption of a palatable food. Activity of
NAc D2 cells was linked to both 'liking' and 'wanting' but with opposing actions; high D2 cell activity
signaled 'wanting', and low D2 cell activity enhanced 'liking'. Importantly, the increased activity in D2
cells was similar across milkshake sessions, whereas inhibition of this signaling following the onset of
milkshake consumption was greater after repeated milkshake sessions (**Figure 1e**). A multiple
regression analysis indicated that this magnitude of inhibition (area under the curve) after
consumption was a good predictor for 'liking' (bout duration, **Figure S2b**). Previous findings showed
that during consumption of sucrose, a decrease in firing rate in a subset of NAc neurons immediately
before and during a lick bout is required to initiate and maintain consummatory behavior (Taha and
Fields, 2005; Taha and Fields, 2006). The present study supports the idea that sustained D2 cell
depression maintains the actions resulting in palatable food consumption and hedonic effects
(Carlezon and Thomas, 2009; Krause et al., 2010; Roitman et al., 2005; Taha and Fields, 2005; Taha and
Fields, 2006). We showed that D1 cells on the other hand promote 'liking' right after consumption
onset. D1 receptors have been shown to play a role in incentive learning (Beninger, 1983), and so may
be important during the initial phase of instrumental learning, when reward is novel and unpredictable
(Schultz, 1998) and in the primary response to drugs of abuse (Self, 2014). This may explain why D1
cell activity was highest during the first milkshake session (**Figure 1e**). Supporting this interpretation,
in a multiple regression analysis, the maximum D1 cell signal following lick onset was a significant
predictor of bout duration; the larger the initial increase, the longer the licking bout (**Figure S2a**).
330

331 Contribution of NAc D1 and D2 cells to reward-associated cues

D1 and D2 cells displayed distinct temporal activity during food magazine visits in the motivational
task; D1 cell activity increased after onset of the food delivery cue, whereas D2 cell activity was
increased before, but suppressed after, food magazine entry (**Figure 3c,d**). These results are in line
with previous findings from drug-addiction literature showing that NAc D1 cells increase their activity
before entering a drug-paired chamber, while D2 cells decrease their activity after (Calipari et al.,
2016). Thus it may be that in our task, D1 cell activity drives the motivation to enter the food magazine,
while reduced activity of D2 cells after food magazine visit onset would drive the motivation to remain
in the magazine. This might suggest that D1, but not D2 cell signaling drives reward- and
reinforcement-related behaviors. In particular, we show a causal role of temporally specific cue-
elicited D1 cell signaling, which can further demonstrate the role of this cell type in motivation to
retrieve rewards. Activation of D1 cells when animals entered the food magazine or at cue-onset
increased the number of food magazine visits. This may imply that the D1 cell signals guide animals to
seek reward, whereas suppressing D1 cell activity at either food magazine or cue onset doesn't lead to
any changes in the number of visits compared to the control group. However, inhibition of D1 cells at
cue onset delayed approach. D1 cell activity may thus drive higher incentive motivation to reach the
palatable food (reward) by increasing focus on the cue, and amplifying 'wanting' for a particular
incentive target (the high-caloric palatable food). This could contribute to intense urges to indulge in
those foods, leading to overeating. Attenuating or eliminating D1 cell signals, may decrease the
associative learning of environmental stimuli (reward-associated cue) with reward. Our experiments
were conducted once the animals learnt the association between cue and reward and the neuronal

response that was measured was to the reward-predictive cue (Schultz, 1997; Wise and Rompre,
1989). Pharmacological studies investigating the role of NAc D1 and D2 receptors in response to drugs
of abuse have also shown that both D1 and D2 receptors have a role in associative learning (Graham
et al., 2007). Dopamine responses to cues predicting drug availability can lead to drug seeking behavior
and over time these responses to drug-associated cues are dampened (Graham et al., 2007; Self, 2014).
Our findings that D1 cells in the context of associative learning are in line with previous publications
demonstrating that D1-like receptors are essential in reward related learning including instrumental
learning (Ranaldi and Beninger, 1995; Sutton and Beninger, 1999) as well as driving motivation to
action (Fibiger, 1993). More specifically, systemic administration as well as local infusion into the NAc
of dopamine D1-like receptor antagonists have been shown to attenuate food-reinforced lever
pressing and to dampen the rewarding effects of palatable food (Aberman et al., 1998; Beninger et al.,
1987; Cousins et al., 1994; Koch et al., 2000; McDougall et al., 1991). Furthermore, optogenetic
activation of NAc D1 cells enhanced drug-induced conditioned place preference (Koo et al., 2014; Lobo
et al., 2010). Pharmacological activation of NAc D1 receptors produced similar findings, showing that
D1 cell activity drives incentive motivation to consume food and drug of abuse (Natsubori et al., 2017;
Schmidt et al., 2006; Wakabayashi et al., 2004).

On the other hand, D2 cell silencing at the time of milkshake delivery cue and at the time of food
magazine visit significantly decreased the number of food magazine entries. Also, silencing natural D2
cell activity after milkshake delivery cue also significantly decreased the average length of a
subsequent food magazine visit. These observations suggest a role for D2 cell activity in the
anticipation of a reward ('wanting'). These effects on motivation, although inconsistent with some
previous findings which suggested that D2 cells are associated with negative rather than positive
valence events (Hoek, 2016; Kravitz et al., 2012; Lobo et al., 2010), were in line with other more recent
findings which have shown that D2 cell activation increased motivation to obtain a reward in a
progressive ratio task (Soares-Cunha et al., 2018; Soares-Cunha et al., 2016b).

NAc D1 cell stimulation drives hedonic shifting and overconsumption of palatable food

In the hedonic-shifting choice test after the final optogenetic stimulation pairing, D1- but not D2-cell
activity significantly increased preference for the milkshake flavor that was previously least preferred.
This indicates that D1 cell optostimulation induced a hedonic preference via an association between
the flavor and the rewarding effects of stimulation (**Figure 4d**). This was not observed in control mice.
In addition, during the conditioning sessions of D1 cell optostimulation, an increase in time spent
consuming the opto-paired flavor was observed across sessions (**Figure 4b**). This may reflect a hedonic
experience which alters associative hedonic/'liking' processes by selectively enhancing D1 cell activity.
This facilitates the association between flavor and opto-stimulation and promotes hedonic shifting that
in turn leads to increased consumption of that flavor as observed in the choice test. This effect was
specific for NAc D1 cells; stimulation of NAc D2 cells during conditioning had no effect either across
conditioning days, or on the choice test.

In this experiment we were able to induce a learned shift in preference for a particular flavor when D1
cell optogenetic stimulation was coupled with a less preferred flavor. This is based on a Pavlovian
association formed between NAc D1 cell activation (US) and the least preferred milkshake flavor (CS),
which changed the taste preference to that flavor. This suggests that hedonic 'liking' of a flavor
stimulus can be determined not only by flavor itself, but also by the relevant 'brain' state at the time
of formation of Pavlovian associations, for example by a change in perceived palatability. In a similar
manner, in the context of drug addiction, it was shown that NAc D1 cell activity preceding entry to the
drug-paired compartment in a conditioned place preference test is needed for drug-associative
learning to take place (Calipari et al., 2016; Lobo et al., 2010).

That our findings are not always in line with previous literature may reflect the discordance between previous studies themselves: the precise NAc subregion targeted plays a role in the results obtained (de Jong et al., 2019; Yang et al., 2018), the pattern (frequency, length) of the optogenetic modulation pattern also has an impact on the direction of the effects (Soares-Cunha et al., 2020), and the experimental paradigm also plays a crucial role depending on the cues and responses it entails (Nicola et al., 2004).

Clinical implications for the role of NAc D1 and D2 cell activity in 'liking' versus 'wanting' in overconsumption and eating disorders

Our findings may carry potential implications for human eating disorders. 'Hedonic eating' does not necessarily involve pleasure, but can also be driven by a motivation to eat which can be dissociated from the drive caused by a nutrient deficit (Lowe and Butryn, 2007; Lutter and Nestler, 2009). It is already well known that in humans, high-calorie foods, especially sweet and fatty ones, promote overconsumption. The incentive theory of 'liking'/'wanting' was significant in the study of food intake regulation, modelling associations between the function of the reward system and feeding or obesity. Studies tried to show that different neural mechanisms are responsible for 'wanting' a certain food than 'liking' it. For more than a decade, human studies of obesity have used 'liking' and 'wanting' to try to better understand 'wanting' food can differ in some individuals, and override 'liking' to cause excessive cue-trigged 'wanting' (craving) that will drive overeating and lead to obesity (Berthoud et al., 2011; Devoto et al., 2018; Gearhardt et al., 2014; Schulte et al., 2017; Stice and Yokum, 2016). Incentive sensitization theory, originally postulated for drug addiction (Robinson and Berridge, 1993), was then applied to eating disorders to try to explain how some individuals may experience 'wanting' to eat palatable foods (Berridge, 1996, 2009). Neuroimaging studies suggest that in humans diagnosed with obesity or binge-eating disorders, overeating is triggered by visual cues associated with palatable food (Frankort et al., 2011; Rothemund et al., 2007; Stoeckel et al., 2008). For example, increased brain activity in response to palatable foods was positively correlated with self-reported craving rates or 'wanting' to eat compared to healthy controls (Polk et al., 2017; Simon et al., 2016; Wolz et al., 2017). This evidence is very similar to findings in individuals suffering from drug addiction (Cameron et al., 2017; Carter et al., 2016; Devoto et al., 2018; Fletcher and Kenny, 2018; Volkow et al., 2017; Westwater et al., 2016).

To our knowledge, our study is the first to approach aspects of NAc dependent reward consumption using such a multi-faceted approach. We incorporated observation, inhibition and excitation of NAc D1 and D2 cells, multiple behavioral paradigms investigating different aspects of 'liking' and 'wanting', including analysis of lick microstructure, and multivariate analysis. In summary, we show that NAc D1 cell activity plays a role in reinforced behaviors induced by palatable foods in an operant paradigm or a Pavlovian association test. However, in a palatable food free-access paradigm D1 cells encode only the hedonic property of the palatable food. NAc D2 cells on the other hand, have a role in the drive to retrieve food, showing elevated activity prior to eating onset in a motivational task. Once eating starts, suppressed D2 cell activity allows for prolonged eating episodes. This would support the idea that evolutionary processes led to these two cell populations that can independently and mutually influence hedonic eating. D1 cells are prone to lead to a maladaptive cycle of increased 'liking' and 'wanting', thus driving overeating, while D2 cells cannot reinforce both processes and are faced with a tradeoff between increasing 'wanting' by being more active or allowing 'liking' by remaining silent. These findings, which temporally distinguish between the different psychological processes of food reward and reinforcement, support the incentive salience hypothesis (Berridge, 1996; Berridge and Robinson, 1998). They may assist the design of future studies and improve diagnostic criteria of addiction and eating disorders.

Materials and Methods

Animals and housing conditions

All animal procedures were carried out in accordance with the Animal Welfare Ordinance (TSchV 455.1) of the Swiss Federal Food Safety and Veterinary Office and were approved by the Zurich Cantonal Veterinary Office. Mice were kept on a reversed 12-h/12-h light/dark cycle and provided with standard chow pellets *ad libitum*. In order to avoid food deprivation for behavioral testing, mice were provided with 2% citric acid in water *ad libitum* in their home cage (Urai et al., 2021). All experiments were performed during the dark phase. Both adult male and female mice were used for the experiments (see Table 1) and mice were at least 7 weeks old before surgeries were performed. As no differences were found in the results between sexes, males and females were pooled.

Viruses and surgeries

For specific targeting of D1 cells and D2 cells, D1-cre (B6.FVB(Cg)-Tg(Drd1-cre)EY262Gsat/Mmucd) and Adora2a-cre (Adora2aTg(Adora2a-cre)KG139Gsat, originally obtained from the Jackson Laboratory) mouse lines were used, respectively. For photometry recordings, 6 mice from each line were injected bilaterally with a pAAV9-CAG-Flex.GCaMP6s.WPRE.SV40 virus (Addgene, titre ≥ 1×10^{13} vg/mL) in the medial NAc shell (D1-cre and D2(A2a)-cre mice; bregma +1.4mm, midline ± 1.35mm, depth 3.9 mm, at an angle of 10°) and were then implanted bilaterally with optic fibers (200 µm diameter, NA 0.37, Doric Lenses), using the same coordinates (except that the depth of the fiber tip was reduced to 3.8mm).

For optogenetic modulation, mice (see Table 1) were injected with pAAV9-EF1a-DIO-ChrimsonR-mRuby2-KV2.1-WPRE-8V40 (Addgene, titer: 1×10^{13} vg/mL) or ssAAV-9/2-hEF1a-dlox-eNpHR3.0_iRFP(rev)-dlox-WPRE-hGHp(A) (ETHZ/UZH Viral Vector Facility, titre: 7.2 x 10^{12} vg/mL) or pAAV-flex-GFP (Addgene, titer: 2.5 x 10^{13} vg/mL) for activation, inhibition and control respectively. Viruses and fiber implants were positioned at the same coordinates as the photometry cohort. Virus expression was assessed using standard histology methods.

For stereotaxic surgeries (Kopf Instruments), a small craniotomy of 0.5 mm in diameter was performed above the injection sites using a microdrill. Injections were performed using a Nanoject III injector and glass capillaries (Süd-Laborbedarf Gaunting); 120nL of virus were injected on each side at a rate of 1nL/sec before optic fibers were slowly inserted and fixed to the skull using Super-Bond (Sun Medical Co. Ltd). Mice were allowed 6 weeks to recover before starting experiments.

Table 1 : Mice implanted in each group and line.

	Adora2a-cre	D1-cre
Photometry – GCaMP6s	6 (4m + 2f)	6 (3m + 3f)
Optogenetics – ChrimsonR	5 (3m + 2f)	6 (3m + 3f)
Optogenetics – eNpHR3.0	6 (3m + 3f)	6 (3m + 3f)
Optogenetics – GFP	6 (3m + 3f)	6 (3m + 3f)

Photometry recordings

Photometry recordings were performed using a camera-based bundle-imaging fiber photometry system (Doric Lenses) using interleaved stimulation illumination produced by two LEDs (2 excitation wavelengths: 405 nm and 465 nm / 2 cycles, Resolution 1200 x 1200, Effective Sampling Rate 20 Hz). Fluorescence produced by 405-nm excitation provided a real-time control for motion artifacts. Recordings were performed using the Doric Neuroscience Studio software.

Optogenetics

Optogenetic activation of D1-ChrimsonR and D2-ChrimsonR cells was performed using a red laser (635nm, Laserglow Technologies). Laser output frequency was 4-ms pulses at 20Hz, driven by an Arduino board. Optogenetic inhibition of D1-eNpHR3.0 and D2-eNpHR3.0 cells was achieved with a yellow laser (589 nm, Laserglow Technologies). The laser delivered a constant light when inhibition was triggered and ramped down at the end of an inhibition bout to avoid neuronal activity rebound. D1-GFP and D2-GFP control mice were split evenly between opto-inhibitory and opto-excitatory stimulation patterns.

The lasers were connected to the bilateral fiber implants using a 1x2 fiber-optic rotary joint (Doric Lenses) and yielded an output power of ~8mW at the end of each fiber tip. Laser pulse onset parameters are described in each relevant experiment section.

'Motivation to retrieve a reward' task

Pre-surgery screening: Prior to surgeries, both photometry and optogenetic cohorts mice were screened in this simple reward delivery task. Mice were habituated to the experimental chambers (Bussey-Saksida Mouse Touch Screen Chambers installed in sound-proof chambers, Lafayette Instruments) for 20 min per day for two days. The task schedule was designed and controlled by ABET II software (Lafayette Instrument). On three consecutive days, mice could receive up to 15 milkshake deliveries (Energy milk, strawberry flavor, Emmi Schweiz AG; 20 µL/delivery; to which mice had been previously habituated in the home cage) during a maximum of 30 min. The session ended when whichever of the 15 deliveries or 30 min came first. The inter-trial interval varied pseudo-randomly, 30 ± 10 sec, counted from the moment a mouse exited the food magazine. Only those able to perform the task in 10 min or less were implanted. The house light (white light) was on during the sessions. The tray light was turned on at each milkshake delivery and turned off when the mouse visited the food magazine. Milkshake delivery was associated with a sound cue.

Photometry cohort: The same paradigm used for pre-surgery screening was run again on 3 consecutive days while photometry signals were being recorded (see *Photometry Recordings* section for details).

Optogenetics cohort: Here, the task structure was similar to the pre-surgery screening task, except the mouse performed the task over two days only. The first day was identical to the pre-screening task and served as a baseline. On the second day, the task was identical but 10s of optogenetic stimulation either started from milkshake delivery cue (D1-eNpHR3.0, D2-eNpHR3.0 and GFP controls) or started at food magazine visit onset (D1-ChrismonR, D2-ChrismonR, and GFP controls). After a few-weeks wash-out period this 2-day experiment was repeated, only this time mice which had received optogenetic stimulation during milkshake delivery now received it during the food magazine visit and *vice versa*. In these tasks the activation and inhibition patterns were the same as those described in the *Optogenetics* section, repeated continuously over the 10 sec each stimulation lasted for. Instead of lasers, light was delivered using a PlexBright optogenetic stimulation system (Plexon Inc.) consisting of an Optogenetic Controller (Plexon Inc.) and orange LEDs for activation (compact LED, 620 nm, Plexon Inc.) or lime LEDs for inhibition (compact LED, 550 nm, Plexon Inc.). The output at the end of each patchcord (200/230 µm, 0.5NA, Plexon Inc.) was ~5mW. The patterns were designed in and controlled by Radiant software (Plexon Inc.) which was itself triggered by ABET II software during the task.

Ad libitum **consumption experiment**

Mice were first habituated to spring-based tip bottles (EBECO) in the home cage using water. Habituation to the arenas (20x20cm^2 arenas, Habitest modular mouse test cage, Coulbourn Instruments) and patchcords (200 µm, low-autofluorescence, Doric Lenses) was performed on 2

consecutive days before the experiment started. The test cages were placed in closed, sound-attenuating chambers (Coulbourn instruments) and illuminated using infra-red light only. Licks were detected using a capacitive sensor (Capacitive Touch Breakout, Sparkfun) and recorded at 500Hz throughout the sessions using a National Instruments board (USB-X Series DAQ, NI-6343) and the LabView software (National Instruments). Weights consumed were recorded at the end of each session. Mice were not food deprived for these experiments.

Photometry cohort: On experimental days, mice were placed in the arenas for 20-min long sessions (one session/day for each mouse) with free access to one 'spring' bottle containing water (days 1-3), strawberry milkshake (days 4-8). Concomitant photometry signals were recorded as described in the *Photometry Recordings* section above.

Optogenetics cohort: Over a 20-min session with access to milkshake, laser stimulation started (according to the patterns described in the *Optogenetics* section) every time a mouse started licking and continued for the duration of licking.

Hedonic shifting experiment

For this experiment only ChrimsonR mice (and their respective GFP controls) were used. Mice were habituated in the home-cage to two novel flavors of milkshake which were matched for their calorie and macro-nutrient contents: a chocolate-flavored and a vanilla-flavored milkshake ('Milk Choco Mountain' Protein, 'Vanilla Drive', respectively, Chiefs AG Schweiz).

Pre-test phase: this phase served to test the natural preference of mice for one flavor over the other. Testing was performed over two consecutive days with 20-min sessions in the same Coulbourn arenas as described in the *Ad libitum consumption Experiment* section. Two 'spring' bottles (EBECO), each with a different flavor of milkshake were freely available. The positions of the bottles in the cage were counterbalanced across mice. Licking and amounts consumed were recorded as described in the *Ad libitum consumption experiment* section. The time spent licking at each bottle on the second day of the pre-test phase was used to determine which was each mouse's preferred flavour.

Conditioning phase: Two 10-min conditioning sessions were run, one in the morning, and one in the afternoon for 7 consecutive days. In each of these sessions only one flavor was available at a time, and the order of the sessions was swapped each day (see **Figure 4a**). Licking at the mouse's initially-less-preferred flavor triggered optogenetic activation for as long as the mouse kept licking, using the patterns and equipment described in the *Optogenetics* and *Ad libitum consumption experiment* sections respectively.

Choice-test phase: 24 hours after conditioning, mice were assessed for their preference again. During the choice-test, both flavors were present simultaneously for 20-min with no optogenetic stimulation on either flavor.

Throughout the conditioning and choice-test phases, but not for the initial preference pre-test phase, mice were mildly food-deprived.

Data Analysis

Licking microstructure analysis

Based on previous literature (Johnson, 2018a; Robinson and McCool, 2015), licking behavior was analyzed by looking at the following parameters:

- Single lick durations, defined as an uninterrupted high signal.

- Inter-lick intervals (ILI), defined as an uninterrupted low in the signal of up to 250ms. An upper
boundary was set in accordance to Johnson (2018a), to distinguish them from inter-bout intervals.

- Number of licking bouts, a bout was defined as a licking episode of at least 500ms, allowing for
interruptions within a bout of up to 250ms.

- Licking bout duration, length of a given bout in seconds.

- Licking bout size, number of individual licks within a given bout.

- Inter-bout intervals (IBI), defined as interruptions in licking of more than 250ms allowing for licking
occurrences of up to 250ms, i.e. a short lick isn't considered a bout and thus does not interrupt a given
IBI.

For the regression analysis (see below), licking bouts had a slightly more inclusive definition, starting
with at least 100 ms of licking and allowing for pauses of up to 1 sec.

Multiple regression analysis

Correlations between photometry signals and 'liking'/'wanting' parameters were performed using the
SPSS (IBM Corp. Version 28) linear regression tool. A subset of 8 parameters depicting the photometry
signal around a licking bout onset was selected: maximum, minimum and area under the curve (AUC)
in the 5 secs following lick onset; mean and median signal over the 3 sec before lick onset; mean and
median signal over the 3 sec following lick onset and standard deviation of the non-z-scored signal in
the 5 sec preceding lick onset. Because aiming to fit too many variables in a regression model can lead
to spurious results, we used all parameters simultaneously in a first step only, to assess correlations
between predictors and see which predictors had the highest and most significant correlations with
the outcome variable. If predictors were highly correlated between themselves (Pearson's r >8.5,
VIF>10, tolerance<0.1, checking expression of each predictor in eigen vector base) then only the
predictor with the most significant correlation with the outcome variable was kept. Then, in a second
step, another linear regression was performed keeping only the non-collinear predictors. A final block-
wise (hierarchical) regression was done keeping only the predictors that showed significant
correlations with the outcome variable and/or model coefficients significantly different from 0. This
final block-wise approach – which started with the predictor yielding the highest significant
standardized model parameter (β) and continued in decreasing order of βs – allowed us to evaluate
from which predictor adding one more predictor to the model did not yield better predictive power
(looking at the p value associated to the F-ratio change). Predictors whose addition to the model gave
a significant F-ratio change (i.e., improved the model) are reported in **Figures S2-4**. Case-wise
diagnostics allowed us to evaluate the adequacy of our dataset, namely whether 95% of data points
had standardized residuals within ±2, and around 1% of the sample at most had standardized residuals
over 2.5. Assumptions of the model (absence of heteroscedasticity, linearity) were evaluated by
plotting standardized residuals against standardized predicted values. Normal distribution was
evaluated via the observed vs. predicted cumulative probability plots. The Durbin-Watson index was
used to evaluate whether the model's residuals were independent (index should be as close to 2, and
at least >1 and <3). As our aim here was to evaluate which aspects of the photometry signal (i.e.,
neuronal activity) best predicted 'liking' and/or 'wanting', but not to apply the model to future
datasets, we did not use bootstrapping. Nonetheless, to assess the validity of our model for a general
population, cross-validation was performed using the 'adjusted R^2 value' – a value as close as possible
to R^2 indicated that the model generalizes well to the population beyond the sample data.

'Motivation to retrieve a reward' task analysis

When analyzing photometry signals around MS delivery, trials where the mouse entered the food
magazine within 5 sec after MS delivery cue were not included so that signals did not include MS
consumption, but only MS anticipation (**Figure 3c**). For analyses of signals around food magazine visits,
trials for which the mouse exited the food magazine just after entering it were discarded (**Figure 3d**).
When calculating the average food magazine visit length, visits of less than 2 seconds were excluded
to avoid skewing the mean to short values of very brief – and mostly non-consummatory – visits (**Figure**
3i,j).

Statistical analysis

Data was analyzed using Matlab R2019b (MathWorks). All statistical tests and descriptive statistics
were performed using the SPSS software (IBM Corp. Version 28). Sex was taken as a factor in the
analysis but did not reveal significant differences in all comparisons. Therefore, since sex was not the
primary independent variable of interest in this study, and given the lack of statistical differences, we
pooled data from both sexes for all further analyses. Tests and their results are presented in the figure
legends. Unless otherwise stated, data are presented as mean ± SEM, and a p-value ≤ 0.05 was
considered to indicate significance. In the motivation task (**Figure 3e-j**), outlier trials (defined as points
whose absolute z-score is >3.29) were excluded from the statistical analysis.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

We would like to thank Valeria Franco Betzler for her assistance in histological evaluations. This study
was supported by the Swiss National Science Foundation (grant no. 310030_189110 to DPR and DB)
and the Novartis Foundation for medical-biological Research (grant no. 21A061 to DPR).

Main Figures and Figure Legends

Figure 1. NAc D1 cells exhibit transiently elevated activity, and D2 cells exhibit suppressed activity, during periods of free consumption of palatable food.

(a) Schematic of experimental set-up for photometry recordings and lick detection during an *ad libitum* consumption paradigm. Left: Illustration of fiber placement. Dashed yellow box indicates location of zoom below.

(b) Top: Example photometry recording from a 20-min session with access to milkshake (MS) from a D1-GCaMP6s mouse. Licks indicated by black raster plot above. Bottom: Zoom of photometry signal showing only 2 min indicated by black dotted square above; licks shown above as a black raster plot.

(c) As in (b) but for a D2-GCaMP6s mouse.

(d) Photometry signals around water (H_2O, blue) or MS (red) lick onset, averaged over all licking bouts in D1-GCaMP6s mice (left, N=6) and D2-GCaMP6s mice (right, N=5). MS signals are an average of 5 sessions. Mean

across bouts and animals is shown, shaded area represents the SEM. D1-GCaMP6s mice, one-sample t-tests: max post lick onset water (t(5)=3.972, p=0.011), delta max-min (t(5)=8.457, p<0.001). D2-GCaMP6s mice, one-sample t-tests: max at lick onset water (t(4)=3.437, p=0.026), max at lick onset MS (t(4)=7.848, p=0.001), min post lick onset MS (t(4)=-3.526, p=0.024). Paired-samples t-test: min post lick onset MS vs. water (t(4)=5.407, p=0.006).

(e) As in (d) but comparing the first and last MS sessions. D1-GCaMP6s mice, one-sample t-tests: max post lick onset MS 1 (t(5)=2.687, p=0.043), MS 5 (t(5)=1.856, N.S.). Paired-samples t-test: max MS 1 vs. MS 5 (t(5)=6.257, p=0.002). D2-GCaMP6s mice, one-sample t-tests: max post lick onset MS 1 (t(4)=7.628, p=0.002), MS 5 (t(3)=2.858, N.S.), delta max-min MS 5 (t(5)=4.274, p=0.008). Paired-samples t-test: min post lick onset MS 1 vs. MS 5 (t(3)=3.088, p=0.05).

(f) Lick duration (left) and inter-lick interval (ILI, right) during a session with access to water (blue dot) and 5 sessions with access to MS (red and pink dots). D1- and D2-GCaMP6s cohorts pooled. Mean ± SEM values are shown in color, with individual values from each mouse overlayed in black. One-way RM-ANOVA: lick duration water vs. MS 1 (F(1,17)=4.799, p=0.043), MS 1 to MS 5 (F(4,68)=3.607, p=0.010), MS 1 vs. MS 5 (F(1,17)=9.961, p=0.006); ILI: water vs. MS 1 (F(1,17)=38.238, p<0.001), MS 1 to MS 5 (F(4,68)=0.924, N.S.), N=18.

(g) As in (f) for lick bout number (left) and inter-bout interval (IBI, right). One-way RM-ANOVA: bout number water vs. MS 1 (F(1,17)=45.661, p<0.001), MS 1 to MS 5 (F(4,68)=2.268, N.S.); IBI water vs. MS 1 (F(1,17)=44.170, p<0.001), MS 1 to MS 5 (F(4,68)=5.318, p=0.001), MS 1 vs. MS 5 (F(1,17)=35.626, p<0.001), N=18.

MS: Milkshake. p-values reported on the figures as follows: *p≤0.05, **p<0.01, ***p<0.001

Figure 2. NAc D2 cell natural activity increases consumption by increasing 'wanting' despite a decrease in 'liking'.

(a) Schematic of experimental set-up for closed-loop optogenetic control based on recorded licks in an *ad libitum* consumption paradigm. Left: fiber placement and viruses used.

(b) Total number of licks recorded in a 20-min *ad libitum* session with access to milkshake (MS) in D1-cre and D2(A2a)-cre mice, expressing either ChrimsonR (ChR, red), eNpHR3.0 (eNpHR, orange) or the GFP (blue) reporter. D1: One-way ANOVA (opsin: F(2,15)=2.564, N.S.). D2: One-way ANOVA (opsin: F(2,14)=3.837, p=0.047), independent-samples t-test: ChR vs GFP (t(9)=2.231, p=0.05).

(c) 'Liking' parameters consisting of lick duration (left) and inter-lick interval (ILI, right) across one session with access to MS in D1-cre and D2(A2a)-cre mice. Mean ± SEM values are shown in color, with individual values from each mouse overlayed in black. D1: ILI, one-way ANOVA (opsin: F(2,15)=7.516, p=0.005), post-hoc independent t-test with Bonferroni correction eNpHR vs. GFP (p=0.004). D2: ILI, one-way ANOVA (opsin: F(2,14)=3.049, p=0.080), independent-samples t-test ChR vs GFP (t(9)=2.963, p=0.016). Lick duration: one-way ANOVA (opsin: D2: F(2,14)=0.950, N.S., D1: F(2,15)=1.813, N.S.).

(d) As in (c) for 'wanting' parameters, consisting of number of licking bouts (left) and inter-bout interval (IBI, right). D1: one-way ANOVAs (opsin), Bout number (F(2,15)=2.235, N.S.), IBI: (F(2,15)=3.180, N.S.). D2: Bout number, one-way ANOVA (opsin: F(2,14)=6.642, p=0.009), post-hoc independent-samples t-test ChR vs GFP (p=0.014). IBI, one-way ANOVA (opsin: F(2,14)=2.076, N.S.).

p-values reported on the figures as follows: *p≤0.05, **p<0.01.

Figure 3. NAc D1 and D2 cell activities increase with anticipation during a motivational task & inhibition of D2 or D1 cells reduces motivation to retrieve a reward.

(a) Schematic of the chamber coupled with photometry recordings or with optogenetic control triggered by the recorded behavior (see Methods).

(b) Schematic of a milkshake delivery trial structure.

(c) Photometry signals around milkshake (MS) delivery time (associated with a sound cue) in D1-GCaMP6s (left) and D2-GCaMP6s (right) mice. Values are an average across mice and over all trials over 2 sessions run on two different days. The line represents mean across trials and animals; shaded areas represent the SEM.

(d) As in (c) but showing photometry signals around food magazine (FM) visits.

(e) Number of food magazine entries in D1-cre and D2(A2a)-cre mice expressing ChrimsonR (ChR), eNpHR3.0 (eNpHR) or GFP when optogenetic stimulation occurs during 10sec following MS delivery cue. Each trial is counted as an individual data point. D1, one-way ANOVA (opsin: $F_{(2,265)}=8.630$, $p<0.001$), post-hoc with Bonferroni correction: ChR vs. GFP ($p=0.003$). D2, one-way ANOVA (opsin: $F_{(2,250)}=9.828$, $p<0.001$), post-hoc with Bonferroni correction eNpHR vs. GFP ($p=0.006$).

(f) As in (e) but when optogenetic stimulation occurs during 10sec following the first food magazine entry of each trial. D1, one-way ANOVA (opsin: $F_{(2,264)}=9.597$, $p<0.001$), post-hoc with Bonferroni correction ChR vs. GFP ($p=0.001$). D2, one-way ANOVA (opsin: $F_{(2,252)}=3.403$, $p<0.035$), post-hoc with Bonferroni correction eNpHR vs. GFP ($p=0.032$).

(g) As in (e) but for the latency to enter the FM following MS delivery cue (optogenetic stimulation at MS delivery cue). D1, one-way ANOVA (opsin: $F_{(2,264)}=9.820$, $p<0.001$), post-hoc with Bonferroni correction eNpHR vs. GFP ($p=0.003$). D2, one-way ANOVA (opsin ($F_{(2,246)}=5.158$, $p=0.006$), post-hoc tests N.S.

(h) As in (g) but when optogenetic stimulation occurs during 10sec following the first food magazine visit of each trial. D1: one-way ANOVA (opsin: N.S.). D2: one-way ANOVA (opsin: N.S.).

(i) As in (e) but for the average length of a food magazine visit (optogenetic stimulation at MS delivery cue). D1, N.S. D2, one-way ANOVA (opsin: $F_{(2,209)}=5.408$, $p=0.005$), post-hoc with Bonferroni correction eNpHR vs. GFP ($p=0.004$).

(j) As in (i) but when optogenetic stimulation occurs during 10sec following the first FM visit of each trial. D1: one-way ANOVA (opsin: N.S.). D2: one-way ANOVA (opsin: N.S.).

FM: Food magazine, MS: Milkshake. p-values reported on the figures as follows: *$p \leq 0.05$, **$p<0.01$, ***$p<0.001$.

Figure 4. Modulation of NAc D1 and not D2 cells influence hedonic preference learning (hedonic shifting).

(a) Schematic of the experimental design for lick detection and closed-loop optogenetic activation on the paired-flavor (laser ON during conditioning (Cond.) phase only, yellow-shaded cartoon).

(b) Time spent licking on the paired-flavor spout during conditioning sessions and last choice-test session in D1-cre mice expressing ChrimsonR (ChR) and GFP. One-sided independent-samples t-test: ChR vs. GFP (Cond.7: $t(7)=2.160$, $p=0.034$; Choice-test: $t(7)=1.804$, $p=0.057$).

(c) As in (b) but in D2(A2a)-cre mice. Mixed-design ANOVA: N.S.

(d) Change in preference expressed as the difference (Δ) choice-test – pre-test in time spent licking on a given spout (paired & unpaired) in D1-ChR (left) and D1-GFP (right) mice. ChR, paired-samples t-test paired vs. unpaired flavors: $t(5)=4.754$, $p=0.003$ – one-sided. GFP: $t(2)=1.074$, $p=0.198$ – one-sided.

ChR: ChrimsonR, Cond.: Conditioning. p-values reported on the figures as follows: *$p \leq 0.05$, **$p<0.01$.

‘Liking’ parameters

‘Wanting’ parameters

Figure 5. Activation of NAc D1 cells induces a preference shift *via* an increase in ‘liking’ and ‘wanting’ for the optoactivation-paired flavor.

(a) Single-lick duration across conditioning (Cond.) sessions (left & middle inserts) and pre-test & choice-test sessions (right insert) in the paired and unpaired flavors in D1-ChR mice. Values are plotted (in color) as mean ± SEM, with individual values from each mouse overlayed in black. Conditioning, paired flavor: RM-ANOVA (day: $F_{(6,30)}=3.386$, $p=0.006$ – one-sided), one-sided paired-sample t-test Cond.1 vs. Cond.3: $t_{(5)}=-2.675$, $p=0.022$; unpaired flavor (day: $F_{(6,30)}=0.825$, N.S.). Choice-test vs. pre-test, paired flavor: one-sided paired-sample t-test, $t_{(5)}=-3.188$, $p=0.012$; unpaired flavor, $t_{(5)}=0.310$, N.S.

(b) As in (a) but for inter-lick interval (ILI) durations. Conditioning, paired flavor: RM-ANOVA (day: $F_{(6,30)}=3.870$, p=0.003 – one-sided), Cond.1 vs. Cond.4 (one-sided paired-sample t-test, $t(5)=3.212$, p=0.012); unpaired flavor (day: $F_{(6,30)}=1.334$, N.S.). Choice-test vs. pre-test, paired flavor: one-sided paired-sample t-test, $t(5)=4.422$, p=0.003; unpaired flavor: $t(5)=-0.834$, N.S.

(c) As in (a) but for the number of licking bouts. Conditioning, paired flavor: RM-ANOVA (day: $F_{(6,30)}=0.872$, N.S.); unpaired flavor (day: $F_{(6,30)}=2.588$, p=0.019 – one-sided), Cond.1 vs. Cond.7 (one-sided paired-samples t-test, $t(5)=2.330$, p=0.034). Choice-test vs. pre-test, paired flavor: one-sided paired-samples t-test, $t(5)=-1.986$, p=0.05; unpaired flavor, $t(5)=1.800$, p=0.066.

(d) As in (a) but for the inter-bout interval (IBI) durations. Conditioning, paired flavor: RM-ANOVA (day: $F_{(6,30)}=3.176$, p=0.008 – one-sided), Cond.1 vs. Cond.7 (one-sided paired-samples t-test, $t(5)=2.074$, p=0.046); unpaired flavor (day: $F_{(6,30)}=2.280$, p=0.031 – one-sided), Cond.1 vs. Cond.7 (one-sided paired-samples t-test, $t(5)=0.276$, p=0.397), Cond.1 vs. Cond.2 ($t(5)=-7.664$, p<0.001). Choice-test vs. pre-test, paired flavor: one-sided paired-samples t-test, $t(5)=1.961$, p=0.05; unpaired flavor: $t(5)=-2.471$, p=0.028.

ChR: ChrimsonR, Cond.: Conditioning, IBI: inter-bout interval, ILI: inter-lick interval. p-values reported on the figures as follows: *p≤0.05, **p<0.01.

Figure 6. NAc D1 cells positively induce 'wanting' and 'liking' while D2 cells positively encode 'wanting' but negatively induce 'liking'.

Summary of the findings from the different experimental paradigms to test the distinctive patterns of NAc D1 and D2 cell activity relative to distinct psychological processes of hedonic behaviors: 'liking' (hedonia) and 'wanting' (motivation). (a) Summary of D1 and D2 cell activity in the NAc around food approach and following food consumption onset, and the effect of repeated exposure to the reward. NAc D1 cells show transient activation following a reward-predicting cue (dotted line) and following consumption onset (dotted line). The increase following feeding onset fades with daily repeated exposure to the food reward (orange arrow). NAc D2 cells show a sustained increase in activity following a food-predicting cue and their activity drops following feeding onset; this drop becomes lower during the consumption bout and with repeated daily exposure to the food reward (orange arrow). (b) Hypothetical model of the mutual relationship between behaviors (approaching and consuming rewards) and NAc D1 and D2 cell activity. NAc D1 cell activation induces both 'liking' and 'wanting' processes (yellow arrows), and reversely, D1 cells are naturally activated during behaviors associated with 'wanting' and/or 'liking' (grey arrows). NAc D2 cell activation, on the other hand, leads to increased 'wanting' and decreased 'liking' (yellow arrows). Reversely, behaviors associated with 'wanting' (e.g., following a food-predicting cue and during food approach) lead to an increase in D2 cell activity while behaviors associated with 'liking' (e.g., during food consumption) lead to a sustained decrease in D2 cell activity in the NAc. (c) If considering 'liking' and 'wanting' as a 2D space underlying hedonic eating, our data show that D1 and D2 cells are orthogonal in that space and can themselves form a base of this 2D space. This would support the idea that evolutionary processes led to these two cell populations that can independently and mutually influence hedonic eating. D1 cells are prone to lead to a maladaptive cycle of increased 'liking' and 'wanting', thus driving overeating, while

784 D2 cells cannot reinforce both processes and are faced with a tradeoff between increasing 'wanting' by being
785 more active or allowing 'liking' by remaining silent.

Supplementary figures

Figure S1. An increase in palatability is reflected in hedonic behavior ('liking') parameters.

(a) Top: Example photometry recording from a 20-min session with access to water from a D1-GCaMP6s mouse with licks indicated as a black raster plot above. Below: Zoomed-in photometry signal from the same session and animal showing only 2 min; licks shown above as a black raster plot. Photometry signals are (465nm signal – fitted 405 signal)/fitted 405nm signal. Bottom: As in top graph, but for a D2-GCaMP6s mouse.

(b) Description of lick microstructure parameters associated with 'liking' and 'wanting'.

(c) Lick number during daily 20-min sessions with access to water (H_2O, first session) and milkshake (MS - 5 sessions). Values are plotted (in color) as mean $\pm$ SEM, with individual values from each mouse overlayed in black. D1-GCaMP6s and D2-GCaMP6s mice are pooled, N=18. One-way repeated measures ANOVA: water vs. MS 1 ($F_{(1,17)}$=147.327, p<0.001), MS 1 to MS.5 ($F_{(4,68)}$=19.895, p<0.001), MS 1 vs. MS 5 ($F_{(1,17)}$=11.006, p=0.004).

(d) Lick number over 5-min blocks across daily 20-min sessions with access to water (first session) and MS (5 sessions). Values are plotted (in color) as mean $\pm$ SEM, N=18.

(e) Additional parameters calculated from lick recordings traditionally correlated with 'liking': bout duration (left) and bout size (i.e., number of licks per bout, right) across one session with access to water and 5 sessions with access to MS. D1-GCaMP6s and D2-GCaMP6s mice are pooled. Values are plotted (in color) as mean $\pm$ SEM, with individual values from each mouse overlayed in black, N=18. One-way repeated measures ANOVA, Bout duration: water vs. MS 1 ($F_{(1,17)}$=25.529, p<0.001), MS 1 to MS 5 ($F_{(4,68)}$=4.245, p=0.004), MS 1 vs. MS 5 ($F_{(1,17)}$=0.675, p=0.423); Bout size: water vs. MS 1 ($F_{(1,17)}$=26.825, p<0.001), MS 1 to MS 5 ($F_{(4,68)}$=5.282, p=0.001), MS 1 vs. MS 5 ($F_{(1,17)}$=0.003, p=0.956)

MS: Milkshake, IBI: inter-bout interval, ILI: Inter-lick interval. p-values reported on the figures as follows: **p$\leq$0.01, ***p<0.001.

‘Liking’ : Bout duration

a

D1

Correlation predictor vs. bout duration

R² changes across models

Standardized model coefficients

R=0.58-p=8.9819e-75-D1

	B	Std. B	β
(Constant)	3.383 (b_0)	0.594	
Min	-0.96 (-b_1)	0.05	-0.826***
Max	0.45 (b_2)	0.03	0.666***
Std pre	0.003 (b_3)	0.001	0.196***

R^2=0.340, Adjusted R^2=0.337. ***p<0.001

Bout duration = b_0 - b_1*Min + b_2*Max + b_3*Std pre

b

D2

Correlation predictor vs. bout duration

R² changes across models

Standardized model coefficients

R=0.7-p=2.6294e-69-D2

	B	Std. B	β
(Constant)	7.573 (b_0)	0.784	
AUC	-0.003 (-b_1)	0	-0.498***
Min	-0.785 (-b_2)	0.109	-0.384***
Median post	0.76 (b_3)	0.128	0.287***
Median pre	0.578 (b_4)	0.146	0.174***
Std pre	-0.006 (-b_5)	0.003	-0.081*

R^2=0.494, Adjusted R^2=0.489. *p<0.05, ***p<0.001

Bout duration = b_0 - b_1*AUC - b_2*Min + b_3*Median post + b_4*Median pre - b_5*Std pre

Figure S2. Photometry signal features of NAc D1 and D2 cells around palatable food consumption are good predictors of ‘liking’ of the food consumed.

(a) Multiple regression analysis results using variables extracted from the photometry signals (Max: Maximum value in the 5 sec following lick onset, Min: Minimum value in the 5 sec following lick onset, AUC: area under the curve over the 5 sec following lick onset, Mean pre: mean signal over the 3 sec before lick onset, Mean post: mean signal over the 3 sec following lick onset, Median pre: median signal over the 3 sec before lick onset, Median post: median signal over the 3 sec following lick onset, Std: standard deviation of the non-z-scored signal in the 5 sec preceding lick onset – see Methods) of D1-GCaMP6s mice (N=6) around licking onset to predict licking bout duration. Top left: Pearson's r values of single correlations between predictors and bout duration. Top right: R^2 change evaluating whether a given regression model (Mod.) significantly improves the prediction of bout duration compared to the previous model; the p-value associated with the R^2 change is reported above each bar. The R^2 change indicates whether the amount of variance in the outcome (bout duration) can be explained by the added predictor. For model 1 (Minimum (Min) as the only predictor) the R^2 change shows that using the Min to predict bout duration significantly improves predictions compared to using the grand mean of bout duration. For model 2 (Minimum and Maximum (Max) included as predictors) the R^2 change shows that including the Max of the photometry signal around licking onset in addition to the Min significantly improves the model's performance in predicting bout duration, with similar reasoning for subsequent models. Middle left: Standardized coefficients of the final model; standardized coefficients allow to compare the relative contribution of each predictor, regardless of the units used to measure them (the higher the absolute value of a standardized coefficient, the higher the contribution to the model). The p-values associated with these contributions (i.e., whether they are

834 significant, regardless of their magnitude) is reported above each bar. Middle right: Plot of the actual vs.
835 predicted bout duration with the R and p values reported on top. Bottom: model coefficients (B), standard
836 deviation for the coefficients (Std. B) and standardized model coefficient (β) of each predictor in the final model,
837 of which the equation is reported below the table.

838 (b) As in (a) but for D2-GCaMP6s mice (N=5).
839
840 AUC: area under the curve, Max: maximum, Min: minimum, pre: before licking onset, post: following licking
841 onset, Std: standard deviation. p-values reported on the figures as follows: *p≤0.05, ***p<0.001.

'Wanting': Inter-bout interval (IBI)

842
843
844 **Figure S3. Photometry signal features of NAc D1 and D2 cells around palatable food consumption are not good**
845 **predictors of 'wanting' of the food consumed.**
846
847 (a) As in Figure S2a but for prediction of inter-bout intervals (IBI) in D1-GCaMP6s mice (N=6).
848
849 (b) As in (a) but for D2-GCaMP6s mice (N=5). Note that here although both the AUC and Min predictors were –
850 individually – significantly correlated with IBI (top left), only the model considering the AUC yielded a significant
851 improvement in predicting IBI duration compared to using the grand mean of IBI duration (i.e., no better
852 predicting power of model 2 compared to model 1) (top right and middle right).
853
854 AUC: area under the curve, IBI: inter-bout interval, Min: minimum, Std: standard deviation. p-values reported on
855 the figures as follows: *$p \leq 0.05$, **$p \leq 0.01$, ***$p < 0.001$.

D2 – Photometry signals around FM visit

Figure S4. The decrease in D2 cell activity during consumption favors longer licking bouts.

(a) Multiple regression analysis results using variables extracted from the photometry signals of D1-GCaMP6s mice (N=6) around food magazine (FM) visit onsets (Max: Maximum value in the 5 sec following FM visit onset, Min: Minimum value in the 5 sec following FM visit onset, AUC: area under the curve over the 5 sec following FM

visit onset, Mean pre: mean signal over the 3 sec before FM visit onset, Mean post: mean signal over the 3 sec following FM visit onset, Median pre: median signal over the 3 sec before FM visit onset, Median post: median signal over the 3 sec following FM visit onset, Std: standard deviation of the non-z-scored signal over the 5 sec preceding FM visit onset – see Methods) to predict the latency to enter the FM following a milkshake delivery (see Methods). Top left: Pearson's r values of single correlations between predictors and FM visit latency. Top right: R^2 change evaluating whether a given regression model (Mod.) significantly improves the prediction of FM visit latency compared to the previous model; the p-value associated with the R^2 change is reported above each bar. The R^2 change indicates whether the amount of variance in the outcome (latency to enter the FM) can be explained by the added predictor. For model 1 (Standard deviation of the signal in the 5 sec before FM visit (Std pre) as the only predictor) the R^2 change shows that using the Std pre to predict latency significantly improves predictions compared to using the grand mean of latencies. For model 2 (Std pre and Mean pre included as predictors) the R^2 change shows that including the mean photometry signal over the 3 sec before FM visit in addition to the Std pre significantly improves the model's performance in predicting latency to enter the FM. Middle left: Standardized coefficients of the final model; standardized coefficients allow to compare the relative contribution of each predictor, regardless of the units used to measure them (the higher the absolute value of a standardized coefficient, the higher the contribution to the model). The p-values associated with these contributions (i.e., whether they are significant, regardless of their magnitude) is reported above each bar. Middle right: Plot of the actual vs. predicted FM visit latency with the R and p values reported on top. Bottom: model coefficients (B), standard deviation of the coefficients (Std. B) and standardized model coefficient (β) of each predictor in the final model, of which the equation is reported below the table.

(b) As in (a) but for prediction of number of food magazine visits. Note that here although both the Median pre and Std pre predictors were – individually – significantly correlated with the number of FM visits (top left), only the model considering the Median pre yielded a significant improvement in predicting number of FM visits compared to using the grand mean of the number of visits (i.e., no better predicting power of model 2 compared to model 1) (top right and middle right).

(c) As in (a) but for the average food magazine visit length.

AUC: area under the curve, FM: food magazine, Min: minimum, pre: before FM visit onset, Std: standard deviation. p-values reported on the figures as follows: *p≤0.05, **p≤0.01, ***p<0.001.

Lick Microstructure

Figure S5. Activation of D2 cells does not lead to a shift in flavor preference for the optoactivation-paired flavor.

(a) Difference in the time spent licking the paired and unpaired flavors during the pre-test session of the hedonic shifting experiment in D1-cre (left) and D2(A2a)-cre mice (right) expressing ChrimsonR (ChR) or GFP. One-way ANOVA (opsin), D1: $F_{(1,7)}<0.001$, $p=0.994$, D2: $F_{(1,6)}=0.062$, $p=0.812$.

(b) As in (a) but for the time spent licking each flavor on the choice-test session. One-way ANOVA (opsin), D1: $F_{(1,7)}=1.526$, $p=0.257$, D2: $F_{(1,6)}=0.048$, $p=0.833$.

(c) Single-lick duration calculated from lick recordings indicating 'liking' across the conditioning (Cond.) sessions (left & middle inserts) and pre-test & choice-test sessions (right insert) in the paired and unpaired flavors in D1-GFP mice. Values are plotted (in color) as mean $\pm$ SEM, with individual values from each mouse overlayed in black. RM-ANOVA – one-sided: Conditioning, paired flavor (day: $F_{(6,12)}=1.705$, N.S.), unpaired flavor (day: $F_{(6,12)}=0.405$, N.S.); Choice-test vs. pre-test, paired flavor (test day: $F_{(1,2)}=3.865$, N.S.), unpaired flavor (test day: $F_{(1,2)}=0.224$, N.S.).

(d) As in (c) but for the number of licking bouts. RM-ANOVA – one-sided: Conditioning, paired flavor (day: $F_{(6,12)}=0.815$, N.S.), unpaired flavor (day: $F_{(6,12)}=0.974$, N.S.); Choice-test vs. pre-test, paired flavor (test day: $F_{(1,2)}=12.563$, $p=0.036$), unpaired flavor (test day: $F_{(1,2)}=0.113$, N.S.).

(e) As in (c) but for inter-lick interval (ILI) durations. RM-ANOVA – one-sided: Conditioning, paired flavor (day: $F_{(6,12)}=1.528$, N.S.), unpaired flavor (day: $F_{(6,12)}=1.128$, N.S.); Choice-test vs. pre-test, paired flavor (test day: $F_{(1,2)}=0.044$, N.S.), unpaired flavor (test day: $F_{(1,2)}=3.277$, N.S.).

(f) As in (c) but for the inter-bout interval (IBI) durations. RM-ANOVA – one-sided: Conditioning, paired flavor (day: $F_{(6,12)}=0.689$, N.S.), unpaired flavor (day: $F_{(6,12)}=13.824$, $p=0.033$); Choice-test vs. pre-test, paired flavor (test day: $F_{(1,2)}=1.307$, N.S.), unpaired flavor (test day: $F_{(1,2)}=0.183$, N.S.).

(g) Single-lick duration during the pre-test & choice-test sessions in the paired and unpaired flavors in D2-ChR (left) and D2-GFP (right) mice. Values are plotted (in color) as mean $\pm$ SEM, with individual values from each mouse overlayed in black. RM-ANOVA: Choice-test vs. pre-test, D2-ChR: paired flavor (test day: $F_{(1,4)}=1.307$, N.S.), unpaired flavor (test day: $F_{(1,4)}=0.599$, N.S.); D2-GFP: paired flavor (test day: $F_{(1,2)}=0.517$, N.S.), unpaired flavor (test day: $F_{(1,2)}=0.011$, N.S.).

(h) As in (g) but for the number of licking bouts. RM-ANOVA: Choice-test vs. pre-test, D2-ChR: paired flavor (test day: $F_{(1,4)}=2.538$, N.S.), unpaired flavor (test day: $F_{(1,4)}=0.423$, N.S.); D2-GFP: paired flavor (test day: $F_{(1,2)}=7.367$, N.S.), unpaired flavor (test day: $F_{(1,2)}=0.906$, N.S.).

(i) As in (g) but for inter-lick interval (ILI) durations. RM-ANOVA: Choice-test vs. pre-test, D2-ChR: paired flavor (test day: $F_{(1,4)}=6.971$, N.S.), unpaired flavor (test day: $F_{(1,4)}=1.887$, N.S.); D2-GFP: paired flavor (test day: $F_{(1,2)}=0.141$, N.S.), unpaired flavor (test day: $F_{(1,2)}=0.153$, N.S.).

(j) As in (h) but for the inter-bout interval (IBI) durations. RM-ANOVA: Choice-test vs. pre-test, D2-ChR: paired flavor (test day: $F_{(1,4)}=0.394$, N.S.), unpaired flavor (test day: $F_{(1,4)}=2.593$, N.S.); D2-GFP: paired flavor (test day: $F_{(1,2)}=2.962$, N.S.), unpaired flavor (test day: $F_{(1,2)}=1.342$, N.S.).

ChR: ChrimsonR, Cond.: conditioning, IBI: inter-bout interval, ILI: inter-lick interval. p-values reported on the figures as follows: *p≤0.05.

References

Aberman, J.E., Ward, S.J., and Salamone, J.D. (1998). Effects of dopamine antagonists and accumbens dopamine depletions on time-constrained progressive-ratio performance. Pharmacol Biochem Behav *61*, 341-348.

Avena, N., Bocarsly, M., Lee, C., Grossberg, A., Hackell, R., Schwenker, J., and Hoebel, B. (2005). Behavioral indices of anxiety during sucrose abstinence in rats. Obes Res *13*, A111-A111.

Beninger, R.J. (1983). The role of dopamine in locomotor activity and learning. Brain Res *287*, 173-196.

Beninger, R.J., Cheng, M., Hahn, B.L., Hoffman, D.C., Mazurski, E.J., Morency, M.A., Ramm, P., and Stewart, R.J. (1987). Effects of extinction, pimozide, SCH 23390, and metoclopramide on food-rewarded operant responding of rats. Psychopharmacology (Berl) *92*, 343-349.

Benton, D. (2010). The plausibility of sugar addiction and its role in obesity and eating disorders. Clin Nutr *29*, 288-303.

Berridge, K.C. (1996). Food reward: brain substrates of wanting and liking. Neurosci Biobehav Rev *20*, 1-25.

Berridge, K.C. (2009). 'Liking' and 'wanting' food rewards: brain substrates and roles in eating disorders. Physiol Behav *97*, 537-550.

Berridge, K.C., and Robinson, T.E. (1998). What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience? Brain Res Brain Res Rev *28*, 309-369.

Berridge, K.C., and Robinson, T.E. (2003). Parsing reward. Trends Neurosci *26*, 507-513.

Berridge, K.C., and Valenstein, E.S. (1991). What psychological process mediates feeding evoked by electrical stimulation of the lateral hypothalamus? Behav Neurosci *105*, 3-14.

Berthoud, H.R., Lenard, N.R., and Shin, A.C. (2011). Food reward, hyperphagia, and obesity. Am J Physiol Regul Integr Comp Physiol *300*, R1266-1277.

Berthoud, H.R., and Morrison, C. (2008). The brain, appetite, and obesity. Annu Rev Psychol *59*, 55-92.

Calipari, E.S., Bagot, R.C., Purushothaman, I., Davidson, T.J., Yorgason, J.T., Pena, C.J., Walker, D.M., Pirpinias, S.T., Guise, K.G., Ramakrishnan, C.*, et al.* (2016). In vivo imaging identifies temporal signature of D1 and D2 medium spiny neurons in cocaine reward. Proc Natl Acad Sci U S A *113*, 2726-2731.

Cameron, J.D., Chaput, J.P., Sjödin, A.M., and Goldfield, G.S. (2017). Brain on Fire: Incentive Salience, Hedonic Hot Spots, Dopamine, Obesity, and Other Hunger Games. Annu Rev Nutr *37*, 183-205.

Carlezon, W.A., Jr., and Thomas, M.J. (2009). Biological substrates of reward and aversion: a nucleus accumbens activity hypothesis. Neuropharmacology *56 Suppl 1*, 122-132.

Carter, A., Hendrikse, J., Lee, N., Yücel, M., Verdejo-Garcia, A., Andrews, Z.B., and Hall, W. (2016). The Neurobiology of "Food Addiction" and Its Implications for Obesity Treatment and Policy. Annual Review of Nutrition *36*, 105-128.

Corsica, J.A., and Hood, M.M. (2011). Eating disorders in an obesogenic environment. J Am Diet Assoc *111*, 996-1000.

Cousins, M.S., Wei, W., and Salamone, J.D. (1994). Pharmacological characterization of performance
on a concurrent lever pressing/feeding choice procedure: effects of dopamine antagonist,
cholinomimetic, sedative and stimulant drugs. Psychopharmacology (Berl) *116*, 529-537.

D'Aquila, P.S. (2010). Dopamine on D2-like receptors "reboosts" dopamine D1-like receptor-mediated
behavioural activation in rats licking for sucrose. Neuropharmacology *58*, 1085-1096.

Davis, J.D. (1973). The effectiveness of some sugars in stimulating licking behavior in the rat. Physiol
Behav *11*, 39-45.

Davis, J.D. (1989). The microstructure of ingestive behavior. Ann N Y Acad Sci *575*, 106-119; discussion
120-101.

Davis, J.D., and Smith, G.P. (1992). Analysis of the microstructure of the rhythmic tongue movements
of rats ingesting maltose and sucrose solutions. Behav Neurosci *106*, 217-228.

de Jong, J.W., Afjei, S.A., Pollak Dorocic, I., Peck, J.R., Liu, C., Kim, C.K., Tian, L., Deisseroth, K., and
Lammel, S. (2019). A Neural Circuit Mechanism for Encoding Aversive Stimuli in the Mesolimbic
Dopamine System. Neuron *101*, 133-151 e137.

Devoto, F., Zapparoli, L., Bonandrini, R., Berlingeri, M., Ferrulli, A., Luzi, L., Banfi, G., and Paulesu, E.
(2018). Hungry brains: A meta-analytical review of brain activation imaging studies on food perception
and appetite in obese individuals. Neurosci Biobehav Rev *94*, 271-285.

Dwyer, D.M. (2012). Licking and liking: The assessment of hedonic responses in rodents. Q J Exp Psychol
65, 371-394.

Dwyer, D.M., Lydall, E.S., and Hayward, A.J. (2011). Simultaneous contrast: evidence from licking
microstructure and cross-solution comparisons. J Exp Psychol Anim Behav Process *37*, 200-210.

Fibiger, H. (1993). Mesolimbic dopamine: An analysis of its role in motivated behavior. Seminars in
Neuroscience *5*, 321–327.

Finlayson, G., and Dalton, M. (2012). Current progress in the assessment of 'liking' vs. 'wanting' food
in human appetite. Comment on '"You Say it's Liking, I Say it's Wanting ... ". On the difficulty of
disentangling food reward in man'. Appetite *58*, 373-378.

Finlayson, G., King, N., and Blundell, J.E. (2007). Liking vs. wanting food: importance for human appetite
control and weight regulation. Neurosci Biobehav Rev *31*, 987-1002.

Fletcher, P.C., and Kenny, P.J. (2018). Food addiction: a valid concept? Neuropsychopharmacology *43*,
2506-2513.

Frank, G.K.W., Shott, M.E., and DeGuzman, M.C. (2019). The Neurobiology of Eating Disorders. Child
Adolesc Psychiatr Clin N Am *28*, 629-640.

Frankort, A., Roefs, A., Siep, N., Roebroeck, A., Havermans, R., and Jansen, A. (2011). Reward activity
in satiated overweight women is decreased during unbiased viewing but increased when imagining
taste: An event-related fMRI study. International journal of obesity (2005) *36*, 627-637.

Galistu, A., and D'Aquila, P.S. (2013). Dopamine on D2-like receptors "reboosts" dopamine D1-like
receptor-mediated behavioural activation in rats licking for a isotonic NaCl solution.
Psychopharmacology (Berl) *229*, 357-366.

Gearhardt, A.N., Boswell, R.G., and White, M.A. (2014). The association of "food addiction" with disordered eating and body mass index. Eat Behav *15*, 427-433.

Gerfen, C.R., Paletzki, R., and Heintz, N. (2013). GENSAT BAC cre-recombinase driver lines to study the functional organization of cerebral cortical and basal ganglia circuits. Neuron *80*, 1368-1383.

Graham, D.L., Hoppenot, R., Hendryx, A., and Self, D.W. (2007). Differential ability of D1 and D2 dopamine receptor agonists to induce and modulate expression and reinstatement of cocaine place preference in rats. Psychopharmacology (Berl) *191*, 719-730.

Higgs, S., and Cooper, S.J. (1998). Effects of benzodiazepine receptor ligands on the ingestion of sucrose, intralipid, and maltodextrin: An investigation using a microstructural analysis of licking behavior in a brief contact test. Behavioral Neuroscience *112*, 447-457.

Hoek, H.W. (2016). Review of the worldwide epidemiology of eating disorders. Curr Opin Psychiatry *29*, 336-339.

Hyman, S.E., and Malenka, R.C. (2001). Addiction and the brain: the neurobiology of compulsion and its persistence. Nat Rev Neurosci *2*, 695-703.

Johnson, A.W. (2018a). Characterizing ingestive behavior through licking microstructure: Underlying neurobiology and its use in the study of obesity in animal models. Int J Dev Neurosci *64*, 38-47.

Johnson, A.W. (2018b). Examining the influence of CS duration and US density on cue-potentiated feeding through analyses of licking microstructure. Learn Motiv *61*, 85-96.

Kelley, A.E., Baldo, B.A., Pratt, W.E., and Will, M.J. (2005). Corticostriatal-hypothalamic circuitry and food motivation: integration of energy, action and reward. Physiol Behav *86*, 773-795.

Kelley, A.E., and Berridge, K.C. (2002). The neuroscience of natural rewards: Relevance to addictive drugs. J Neurosci *22*, 3306-3311.

King, B.M. (2013). The modern obesity epidemic, ancestral hunter-gatherers, and the sensory/reward control of food intake. Am Psychol *68*, 88-96.

Koch, M., Schmid, A., and Schnitzler, H.-U. (2000). Role of nucleus accumbens dopamine D1 and D2 receptors in instrumental and Pavlovian paradigms of conditioned reward. Psychopharmacology *152*, 67-73.

Koo, J.W., Lobo, M.K., Chaudhury, D., Labonté, B., Friedman, A., Heller, E., Peña, C.J., Han, M.-H., and Nestler, E.J. (2014). Loss of BDNF Signaling in D1R-Expressing NAc Neurons Enhances Morphine Reward by Reducing GABA Inhibition. Neuropsychopharmacology *39*, 2646-2653.

Koob, G.F., and Le Moal, M. (2001). Drug addiction, dysregulation of reward, and allostasis. Neuropsychopharmacology *24*, 97-129.

Krause, M., German, P.W., Taha, S.A., and Fields, H.L. (2010). A Pause in Nucleus Accumbens Neuron Firing Is Required to Initiate and Maintain Feeding. The Journal of Neuroscience *30*, 4746.

Kravitz, A.V., Tye, L.D., and Kreitzer, A.C. (2012). Distinct roles for direct and indirect pathway striatal neurons in reinforcement. Nat Neurosci *15*, 816-818.

Kupchik, Y.M., Brown, R.M., Heinsbroek, J.A., Lobo, M.K., Schwartz, D.J., and Kalivas, P.W. (2015).
Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections.
Nat Neurosci *18*, 1230-1232.

Leigh, S.J., and Morris, M.J. (2018). The role of reward circuitry and food addiction in the obesity
epidemic: An update. Biol Psychol *131*, 31-42.

Lobo, M.K., Covington, H.E., 3rd, Chaudhury, D., Friedman, A.K., Sun, H., Damez-Werno, D., Dietz, D.M.,
Zaman, S., Koo, J.W., Kennedy, P.J.*, et al.* (2010). Cell type-specific loss of BDNF signaling mimics
optogenetic control of cocaine reward. Science *330*, 385-390.

Lowe, M.R., and Butryn, M.L. (2007). Hedonic hunger: a new dimension of appetite? Physiol Behav *91*,
432-439.

Lutter, M., and Nestler, E.J. (2009). Homeostatic and hedonic signals interact in the regulation of food
intake. J Nutr *139*, 629-632.

McDougall, S.A., Nonneman, A.J., and Crawford, C.A. (1991). Effects of SCH 23390 and sulpiride on the
reinforced responding of the young rat. Behav Neurosci *105*, 744-754.

Mela, D.J. (2006). Eating for pleasure or just wanting to eat? Reconsidering sensory hedonic responses
as a driver of obesity. Appetite *47*, 10-17.

Natsubori, A., Tsutsui-Kimura, I., Nishida, H., Bouchekioua, Y., Sekiya, H., Uchigashima, M., Watanabe,
M., de Kerchove, d., Exaerde, A., Mimura, M.*, et al.* (2017). Ventrolateral Striatal Medium Spiny
Neurons Positively Regulate Food-Incentive, Goal-Directed Behavior Independently of D1 and D2
Selectivity. The Journal of Neuroscience *37*, 2723.

Nicola, S.M., Yun, I.A., Wakabayashi, K.T., and Fields, H.L. (2004). Cue-evoked firing of nucleus
accumbens neurons encodes motivational significance during a discriminative stimulus task. J
Neurophysiol *91*, 1840-1865.

Pecina, S., Cagniard, B., Berridge, K.C., Aldridge, J.W., and Zhuang, X. (2003). Hyperdopaminergic
mutant mice have higher "wanting" but not "liking" for sweet rewards. J Neurosci *23*, 9395-9402.

Polk, S.E., Schulte, E.M., Furman, C.R., and Gearhardt, A.N. (2017). Wanting and liking: Separable
components in problematic eating behavior? Appetite *115*, 45-53.

Ranaldi, R., and Beninger, R.J. (1995). Bromocriptine enhancement of responding for conditioned
reward depends on intact D1 receptor function. Psychopharmacology *118*, 437-443.

Robinson, S.L., and McCool, B.A. (2015). Microstructural analysis of rat ethanol and water drinking
patterns using a modified operant self-administration model. Physiol Behav *149*, 119-130.

Robinson, T.E., and Berridge, K.C. (1993). The Neural Basis of Drug Craving - an Incentive-Sensitization
Theory of Addiction. Brain Res Rev *18*, 247-291.

Robinson, T.E., and Berridge, K.C. (2003). Addiction. Annual Review of Psychology *54*, 25-53.

Roitman, M.F., Wheeler, R.A., and Carelli, R.M. (2005). Nucleus accumbens neurons are innately tuned
for rewarding and aversive taste stimuli, encode their predictors, and are linked to motor output.
Neuron *45*, 587-597.

Rothemund, Y., Preuschhof, C., Bohner, G., Bauknecht, H.C., Klingebiel, R., Flor, H., and Klapp, B.F. (2007). Differential activation of the dorsal striatum by high-calorie visual food stimuli in obese individuals. Neuroimage *37*, 410-421.

Schmidt, H.D., Anderson, S.M., and Pierce, R.C. (2006). Stimulation of D1-like or D2 dopamine receptors in the shell, but not the core, of the nucleus accumbens reinstates cocaine-seeking behaviour in the rat. European Journal of Neuroscience *23*, 219-228.

Schulte, E.M., Smeal, J.K., and Gearhardt, A.N. (2017). Foods are differentially associated with subjective effect report questions of abuse liability. PLoS One *12*, e0184220.

Schultz, W. (1997). Dopamine neurons and their role in reward mechanisms. Curr Opin Neurobiol *7*, 191-197.

Schultz, W. (1998). Predictive reward signal of dopamine neurons. J Neurophysiol *80*, 1-27.

Sclafani, A., and Ackroff, K. (2004). The relationship between food reward and satiation revisited. Physiology & Behavior *82*, 89-95.

Self, D.W. (2014). Diminished role for dopamine D1 receptors in cocaine addiction? Biol Psychiatry *76*, 2-3.

Simon, J.J., Skunde, M., Walther, S., Bendszus, M., Herzog, W., and Friederich, H.C. (2016). Neural signature of food reward processing in bulimic-type eating disorders. Soc Cogn Affect Neurosci *11*, 1393-1401.

Smith, G.P. (2001). John Davis and the meanings of licking. Appetite *36*, 84-92.

Smith, K.S., and Berridge, K.C. (2005). The ventral pallidum and hedonic reward: neurochemical maps of sucrose "liking" and food intake. J Neurosci *25*, 8637-8649.

Soares-Cunha, C., Coimbra, B., David-Pereira, A., Borges, S., Pinto, L., Costa, P., Sousa, N., and Rodrigues, A.J. (2016a). Activation of D2 dopamine receptor-expressing neurons in the nucleus accumbens increases motivation. Nat Commun *7*, 11829.

Soares-Cunha, C., Coimbra, B., Domingues, A.V., Vasconcelos, N., Sousa, N., and Rodrigues, A.J. (2018). Nucleus Accumbens Microcircuit Underlying D2-MSN-Driven Increase in Motivation. eNeuro *5*.

Soares-Cunha, C., Coimbra, B., Sousa, N., and Rodrigues, A.J. (2016b). Reappraising striatal D1- and D2-neurons in reward and aversion. Neurosci Biobehav Rev *68*, 370-386.

Soares-Cunha, C., de Vasconcelos, N.A.P., Coimbra, B., Domingues, A.V., Silva, J.M., Loureiro-Campos, E., Gaspar, R., Sotiropoulos, I., Sousa, N., and Rodrigues, A.J. (2020). Nucleus accumbens medium spiny neurons subtypes signal both reward and aversion. Mol Psychiatry *25*, 3241-3255.

Stice, E., and Yokum, S. (2016). Neural vulnerability factors that increase risk for future weight gain. Psychol Bull *142*, 447-471.

Stoeckel, L.E., Weller, R.E., Cook, E.W., 3rd, Twieg, D.B., Knowlton, R.C., and Cox, J.E. (2008). Widespread reward-system activation in obese women in response to pictures of high-calorie foods. Neuroimage *41*, 636-647.

Surmeier, D.J., Song, W.J., and Yan, Z. (1996). Coordinated expression of dopamine receptors in neostriatal medium spiny neurons. J Neurosci *16*, 6579-6591.

Sutton, M.A., and Beninger, R.J. (1999). Psychopharmacology of conditioned reward: evidence for a
rewarding signal at D1-like dopamine receptors. Psychopharmacology (Berl) *144*, 95-110.

Taha, S.A., and Fields, H.L. (2005). Encoding of palatability and appetitive behaviors by distinct
neuronal populations in the nucleus accumbens. J Neurosci *25*, 1193-1202.

Taha, S.A., and Fields, H.L. (2006). Inhibitions of Nucleus Accumbens Neurons Encode a Gating Signal
for Reward-Directed Behavior. The Journal of Neuroscience *26*, 217.

Tavolacci, M.P., Ladner, J., and Dechelotte, P. (2021). Sharp Increase in Eating Disorders among
University Students since the COVID-19 Pandemic. Nutrients *13*.

Trifilieff, P., Feng, B., Urizar, E., Winiger, V., Ward, R.D., Taylor, K.M., Martinez, D., Moore, H., Balsam,
P.D., Simpson, E.H.*, et al.* (2013). Increasing dopamine D2 receptor expression in the adult nucleus
accumbens enhances motivation. Mol Psychiatr *18*, 1025-1033.

Urai, A.E., Aguillon-Rodriguez, V., Laranjeira, I.C., Cazettes, F., Mainen, Z.F., and Churchland, A.K.
(2021). Citric Acid Water as an Alternative to Water Restriction for High-Yield Mouse Behavior. Eneuro
8.

Villavicencio, M., Moreno, M.G., Simon, S.A., and Gutierrez, R. (2018). Encoding of Sucrose's
Palatability in the Nucleus Accumbens Shell and Its Modulation by Exteroceptive Auditory Cues. Front
Neurosci-Switz *12*.

Volkow, N.D., Wang, G.J., and Baler, R.D. (2011). Reward, dopamine and the control of food intake:
implications for obesity. Trends Cogn Sci *15*, 37-46.

Volkow, N.D., Wise, R.A., and Baler, R. (2017). The dopamine motive system: implications for drug and
food addiction. Nat Rev Neurosci *18*, 741-752.

Wakabayashi, K.T., Fields, H.L., and Nicola, S.M. (2004). Dissociation of the role of nucleus accumbens
dopamine in responding to reward-predictive cues and waiting for reward. Behav Brain Res *154*, 19-
30.

Westwater, M.L., Fletcher, P.C., and Ziauddeen, H. (2016). Sugar addiction: the state of the science.
Eur J Nutr *55*, 55-69.

Wise, R.A., and Rompre, P.P. (1989). Brain dopamine and reward. Annu Rev Psychol *40*, 191-225.

Wolz, I., Sauvaget, A., Granero, R., Mestre-Bach, G., Baño, M., Martín-Romera, V., Veciana de las Heras,
M., Jiménez-Murcia, S., Jansen, A., Roefs, A.*, et al.* (2017). Subjective craving and event-related brain
response to olfactory and visual chocolate cues in binge-eating and healthy individuals. Scientific
Reports *7*, 41736.

Wyvell, C.L., and Berridge, K.C. (2000). Intra-accumbens amphetamine increases the conditioned
incentive salience of sucrose reward: enhancement of reward "wanting" without enhanced "liking" or
response reinforcement. J Neurosci *20*, 8122-8130.

Yang, H., de Jong, J.W., Tak, Y., Peck, J., Bateup, H.S., and Lammel, S. (2018). Nucleus Accumbens
Subnuclei Regulate Motivated Behavior via Direct Inhibition and Disinhibition of VTA Dopamine
Subpopulations. Neuron *97*, 434-449 e434.

Zheng, H., and Berthoud, H.R. (2008). Neural systems controlling the drive to eat: mind versus
metabolism. Physiology (Bethesda) *23*, 75-83.

1169