Title:

Routing States Transition During Oscillatory Bursts and Attentional Selection

Authors: Kianoush Banaie Boroujeni[1,2*], Thilo Womelsdorf[1,3]

Affiliations:[1]Department of Psychology, Vanderbilt University, Nashville, TN 37240, [2]Princeton Neuroscience Institute, Princeton University, Princeton, NJ 08544, [3]Department of Biomedical Engineering, Vanderbilt University, Nashville, TN 37240.

*Corresponding author: Kianoush Banaie Boroujeni. Email: kb9590@princeton.edu

Summary: Fast spatio-temporal dynamics of brain activity subserves the routing of information across distant regions and is integral to flexible cognition, decision-making, and selective attention. This study demonstrates that routing dynamics emerge as 20 ms brief lead and lag relationships of spiking activities across distant brain areas. The direction and magnitude of the lead and lag relationships systematically switched during frequency-specific oscillatory bursts and when attention shifts to visual cues.

Abstract

Neural information routing relies on spatiotemporal activity dynamics across interconnected brain areas. However, it remains unclear how routing states emerge at fast spiking timescales and interact with the slower activity dynamics of larger networks during cognitive processes. Here, we show that localized neural spiking events generate long-range directional routing states with spiking activity in distant brain areas that dynamically switch or amplify during oscillatory bursts, selective attention, and decision-making. Computational modeling and neural recordings

from lateral prefrontal cortex (LPFC), anterior cingulate cortex (ACC), and striatum of nonhuman primates revealed that cross-areal, directional routing states arise within ~20 ms around spikes of single neurons. On average, LPFC spikes led activity in the ACC and striatum by few milliseconds. The routing state was amplified during LPFC beta bursts between the LPFC and striatum and switched direction during ACC theta/alpha bursts between ACC and LPFC. Selective attention amplified the lead of these theta/alpha-specific lead-ensembles in the ACC, while decision-making amplified the lead of ACC and LPFC spiking output over the striatum. Notably, the fast lead/lag relationships of cross-areal neuronal ensembles that were modulated by attention states or decision-making predicted firing rate dynamics of their neurons during those functional states at slower timescales. Overall, our findings demonstrate directional routing of spiking activity across nonhuman primate frontal and striatal areas, as well as the functional and network states that modulate the direction and magnitude of these interactions.

Introduction

Routing states during flexible behaviors depend on precise spatio-temporal dynamics of neuronal activity across interconnected brain areas (Singer, 2013). These dynamics proceed at varying temporal scales ranging from rapid synaptic communication (Markram et al., 1997; Hasenstaub et al., 2005), fast recurrent inhibitory and excitatory neuronal interactions (Hahn et al., 2014 p.20; Palmigiano et al., 2017), and slower firing rate modulation in neural populations (Murray et al., 2014; Luczak et al., 2015). As for spatial scales, they can range from a single neuron to multiple nearby neurons, a local population of neurons, or a distributed network of neurons across distant brain areas (Einevoll et al., 2013; Lewis et al., 2015; Wang, 2022). However, the

relationship between these temporal and spatial scales during dynamic transitions of brain-wide

routing states has remained unclear.

To understand the spatio-temporal dynamics of brain-wide routing states during flexible

behaviors, we need to evaluate the directionality of sequential spiking activity in neuronal

ensembles distributed in distant brain areas. Achieving this goal remains challenging which is

evident by a lack of direct insights about long-range, directional spiking interactions at fast

timescales in behaving animals. First, fast activity coordination of neural activity in extracellular

recordings has been measured directly by cross-correlating spike trains or indirectly quantifying

spike synchronization to LFP in the gamma frequency band (Womelsdorf et al., 2007; Smith and

Kohn, 2008), but these methods are often not reliable and robust in estimating the direction of

spiking interactions. Second, attempts to measure directionality in spiking networks, thus far,

have largely focused on latent variables in higher-order correlational structures of spiking

dynamics over pre-defined neural populations without providing mechanistic insights about fast

directional interactions (Semedo et al., 2019; van Kempen et al., 2021). Third, studying long-

range directional communication between cortical and subcortical areas subserving higher

cognitive functions in the primate and nonhuman primate brain has been limited to spike-LFP

and LFP-LFP measures which provide only indirect insights about cross-areal spike-to-spike

directional interactions within neuronal ensembles (Gregoriou et al., 2009; Salazar et al., 2012;

Siegel et al., 2012; Antzoulatos and Miller, 2014; Womelsdorf et al., 2014; Fries, 2015;

Fiebelkorn et al., 2018).

Here, we overcome these limitations and unravel the spatio-temporal dynamics of neural

activities within and across brain areas of the medial and lateral fronto-striatal network.

Connectivity of prefrontal and striatal areas is hierarchically organized and rapidly engages and

disengages during sequential and flexible behaviors (Jin et al., 2009; Haber and Knutson, 2010).

Key nodes of this network are the anterior striatum (head of the caudate nucleus), lateral

prefrontal cortex (LPFC, area 49/9), and anterior cingulate cortex (ACC, area 24) (Ferry et al.,

2000; Haber and Knutson, 2010). Activity changes in each of these areas correlates with learning

success as evident at the levels of single neuron firing (Antzoulatos and Miller, 2011; Oemisch et

al., 2019; Boroujeni et al., 2020; Banaie Boroujeni et al., 2021) as well as oscillatory bursting of

the local field potential (LFP) (Feingold et al., 2015; Lundqvist et al., 2016). In addition, activity

of single neurons in the LPFC, ACC, and striatum synchronizes to narrow-band frequency

fluctuations of the LFP during flexible behavior and efficient attentional control (Antzoulatos

and Miller, 2014; Womelsdorf et al., 2014; Voloh et al., 2020). These findings have established

that single neuron spiking and network-level LFP activity in the fronto-striatal network correlates

with flexible behaviors, but it has remained unclear how activity modulation at these different

levels relate to each other during the dynamic transitioning of routing states, leaving open (1)

whether neurons show directional interactions in this network, (2) whether directional

interactions may relate to network fluctuations and LFP activity states, and (3) whether

directional routing states vary with behavioral states.

Here, we addressed these questions by analyzing spiking interactions in the fronto-striatal

network in nonhuman primates. We found that the emergence of routing states can be traced

back to the sequential temporal lead and lag of single neuron spiking events in relation to

multiunit activity (MUA) from distant areas. Overall, spike events in the LPFC preceded (led)

MUA increases in the ACC and striatum, while spiking of neurons in the striatum lagged MUA

in the LPFC and ACC. The lead/lag relationships defined directional routing states of the LPFC,

ACC and striatum network that amplified or switched direction when spikes of single neurons

occurred during oscillatory bursts, during attentional shifts, and during decision making.

Results

We first simulated two reciprocally connected spiking networks to validate a method for

quantifying the temporal lead or lag of single neuron spikes relative to multiunit activity at high

(<5 ms) temporal precision. In a second step we characterized the lead- and lag- relationships in

neuronal ensembles recorded extracellularly from the ACC, LPFC, and striatum of nonhuman

primates performing an attention demanding learning task. If not noted otherwise, statistical

results on time series are based on randomization tests with extreme-point multiple comparison

corrections, and all group-level statistics were corrected for multiple comparisons using false

discovery rate (FDR) correction at an alpha level of 0.05.

Directional interactions between spiking networks

To evaluate how neural activity coordinates between distant networks we simulated two

interconnected networks (Izhikevich, 2004) with cell-types and connectivity corresponding

closely to LPFC (network 1) and ACC (network 2) (**Figure 1A**, see **STAR Methods** and **Table**

S2, 3) (Dombrowski et al., 2001; Medalla et al., 2017). Each network was composed of 1000

spiking neurons and contained 125 recording sites for measuring electrical field dynamics of

postsynaptic and transmembrane potentials (see **Figure S1, 2a-c** for an example and network

characteristics) (Einevoll et al., 2013). This enabled us to evaluate spike-triggered multiunit

activity (stMUA) and the role of LFP oscillatory events on stMUA strength and direction. Within

each network, MUAs were near-symmetrically distributed relative to spike events with a zero

time-lag (**Figure 1B**). This contrasted to inter-network interactions. Spikes in the simulated

LPFC (sLPFC) systematically led MUA in the ACC (mACC, hereafter, the notation sX-mY

indicates a pair with spikes *s* from a neuron in area X and multiunit activity *m* in area Y), while

sACC lagged mLPFC (**Figure 1B**, permutation test, P<0.05). We quantified the spike lead and

lag as the *Lead/Lag index,* which is the difference of stMUA in the 20 ms following versus

preceding the spike event. A positive Lead/Lag index value indicates spikes lead MUA, while a

negative value shows that spike events lagged MUA. The results from simulations verified that

the Lead/Lag index values closely matched the asymmetry of anatomical connectivity strengths

between the two networks (**Figure S2D-F**). The Lead/Lag index showed significant non-zero

and positive values (spike-lead) for sLPFC-mACC pairs and negative values (spike-lag) for

sACC-mLPFC pairs (**Figure 1C**, Wilcoxon test, p<0.001).

In each of the simulated networks stMUAs and the LFPs were rhythmically modulated. Spectral

power of the LFP and the stMUA showed peaks at ~20 Hz in the LPFC and at ~14 Hz in the

ACC network (**Figure 1D,E**). The lower frequency in the ACC is due to a higher proportion of

inhibitory interneurons with slower synaptic decay compared to the LPFC (Dombrowski et al.,

2001; Medalla et al., 2017). LFP oscillations emerged as transient bursts (**Figure 1F,G**). Bursts

of increased oscillatory power have been associated with coherent interareal activity and with

information routing across brain areas (Palmigiano et al., 2017; van Ede et al., 2018; Zich et al.,

2020). Consistent with these suggestions we found that the lead and lag pattern of stMUAs

between networks transitioned during oscillatory bursts. Spike events occurring during 14 Hz

burst events in the ACC switched from lagging to leading MUA recorded in LPFC (**Figure 1H**),

while spikes in LPFC started lagging MUA recorded from ACC (**Figure 1I**). The switch from

lagging to leading of spikes in ACC relative to MUA in LPFC during ~14 Hz bursts was

significant (**Figure 1J,** Wilcoxon test, p<0.001), while it was not significant at ~20 Hz (**Figure**

S2G).

Spikes in ACC and LPFC lead striatum, LPFC spikes lead over ACC

We quantified stMUAs in simultaneous recordings from neurons in the ACC, LPFC, and

striatum of nonhuman primates (492, 571 and 534 recording sites, respectively) as detailed

before (Oemisch et al., 2019) (**Figure 2A**). We extracted the spike times of isolated neurons and

the MUA around each spike time at another electrode recorded in the same area (intra-areal) or a

distant area (inter-areal) (**STAR Methods**; **Figure S3** for spike-MUA pair examples). Intra-areal

stMUAs showed near-symmetric activity distributions (**Figure 2B**). Compared to the LPFC and

striatum, the stMUA in the ACC showed a wider peak and a steeper MUA drop after the spike

events (**Figure 2B**, randomization test, P<0.05). Similar to the model predictions (**Figure 1B**),

intra-areal stMUAs were rhythmically modulated with the strongest spectral power around 8-14

Hz (theta/alpha) in ACC and 15-25 Hz (beta) in LPFC (**Figure 2C**, randomization test, P<0.05).

Inter-areal stMUAs had temporal relationships consistent with anatomical data in primates (Ferry

et al., 2000; Medalla and Barbas, 2009). On average, sLPFC led mACC and mSTR, sACC

tended to lead mSTR, while sSTR lagged both, mACC and mLPFC (**Figure 2D**; randomization

test, P<0.05). The red line segments in **Figure 2D** mark the timepoints at which stMUA for the

area-pairs was significantly greater than the stMUA from randomized area combinations

(randomization test, P<0.05), which included -26 to -9 ms time lag for sACC-mSTR, and -19 to -

8 ms for sLPFC-mSTR (randomization test, P<0.05). The same pattern of results was seen when

using as test statistic the time lag with maximal stMUA modulation, which showed that the lead

of sLPFC over mSTR and mACC was evident in MUA peaking around +5 and +4.5 ms

following the spike event, respectively (**Figure 2E**, Wilcoxon test, P=0.005 and P=0.02 for

mSTR and mACC, respectively). In the reverse direction sSTR lagged mACC and mLPFC, with

the average stMUA peaks at -12 and -7 ms before the spike, respectively (Wilcoxon test;

P<0.001). We confirmed these inter-areal spiking interactions by calculating the Lead/Lag index

which also showed a significant lead of sLPFC over mSTR and mACC, as well as a lag of sSTR

relative to mACC and mLPFC (**Figure 2F**,**G**; Wilcoxon test; P<0.001, FDR corrected for

dependent samples). The lead/lag effects were maximal when using ~20 ms wide time windows

for calculating the Lead/Lag index (**Figure S4**) and were evident in both monkeys (**Figure S5**).

The firing rate in all stMUA analyses was controlled by using the same number of spikes for

each spike-MUA pair (see **STAR Methods**).

Spikes in ACC switch during theta/alpha bursts from lagging to leading LPFC

The lead and lag of stMUAs in the model simulations were systematically modulated during

oscillatory bursts (**Figure 1H,J**), which raises the question whether the magnitude or direction of

empirically measured lead/lag relationships also varied during oscillatory events. We detected

reliable bursts in the theta/alpha and beta band in the LFP using a new adaptive width-magnitude

thresholding method (**Figure 3A**, see **STAR Methods**, **Figure S6**). Bursts in the theta/alpha and

beta bands were most prevalent in the ACC and LPFC, respectively (**Figure S6G,J**). We

calculated inter-areal stMUAs separately for spikes occurring during versus outside of LFP

bursts. Intriguingly, during theta/alpha bursts, spikes recorded in ACC were followed by stronger

sACC-mLPFC stMUA amplitudes than spikes outside of the burst and switched from a net lag

(**Figure 2F**) to a net lead over MUA in mLPFC (**Figure 3B**, randomization test, P<0.05). Spikes

in ACC also significantly increased the amplitude of its lead over MUA in STR during

theta/alpha bursts (**Figure 3B**, randomization test, P<0.05). Consistent with this finding, spikes

during theta/alpha bursts in the striatum (sSTR) showed an amplified lag relative to MUA from

LPFC and ACC (**Figure 3B**, randomization test, P<0.05). In the LPFC spike events during

versus outside of theta/alpha bursts were preceded and followed by stronger MUA in the ACC

and striatum (**Figure 3B**, randomization test, P<0.05, red line segments indicate the timepoints at

which in burst stMUA for the area-pairs was significantly greater than the in burst stMUA from

randomized area combinations; randomized permutation test, P<0.05). Calculating the Lead/Lag

indices across area-pairs confirmed that during theta/alpha bursts the lead was amplified for

sACC-mLPFC pairs, and the lag was amplified for sLPFC-mACC pairs (**Figure 3C**, Wilcoxon

test, P=0.007 and P=0.036, respectively).

We next asked whether the burst-dependent switch and amplification of directional interactions

were based on stMUA modulation in leading or lagging spike-MUA pairs. Ensembles of leading

and lagging spike-MUA pairs were extracted by projecting the stMUA profiles of individual

interareal spike-MUA pairs to a principal components (PCs) space (**Figure S7A,B**, see **STAR**

Methods). Across PCs, only one PC (PC2) showed an asymmetric stMUA temporal profile

(**Figure S7C,D**). The asymmetric PC profile suggested that PC2 captured the directional

component of inter-areal interactions, which we corroborated by finding that stMUAs with

positive and negative PC2 loading had leading and lagging stMUA profiles, respectively (**Figure**

S7E). This finding enabled using the sign of the PC2 loading to group spike-MUA pairs as

members of a *lead-ensemble* when they had a negative loading on the PC2, and conversely to

group spike-MUA pairs as members of a *lag-ensemble* when they showed a positive PC2 loading

(See **Table S1** for details on the number of spike-leading and spike-lagging spike-MUA pairs for

each brain area). We found that the switch from lagging to leading of sACC over mLPFC was

largely due to spike-MUA pairs of the lag-ensemble switching to the lead-ensemble during

theta/alpha bursts (**Figure 3D**, Kruskal Wallis test, P=0.002, Wilcoxon test for non-zero

modulation P<0.001). The stMUA directionality of lead-ensembles was unchanged during

theta/alpha bursts across area-pairs (**Figure S7F,G**). **Figures 3E** and **F** illustrates an example

sACC-mLPFC pair with stMUA that switched from lagging outside of bursts (grey) to leading

during theta/alpha bursts (red). In summary, during theta/alpha bursts, spikes in ACC

transitioned in a routing state with LPFC and STR from lagging to leading relative to LPFC and

to a stronger lead over STR (**Figure 3G**). The lag-to-lead switch was largely based on spike-

MUA pairs of the lag-ensemble joining the lead-ensemble during theta/alpha bursts (**Figure 3H**).

Lead of LPFC over striatum is amplified during beta bursts

Similar to theta/alpha, we detected (15-25 Hz) beta bursts in the LFP of each area and quantified

inter-areal stMUAs for spikes occurring during beta bursts (**Figure 4A**, **Figure S6D**,**H-J**).

Comparing stMUAs during beta bursts to non-burst periods showed that during beta bursts, the

lead of sLPFC over mSTR was amplified and, reciprocally, sSTR more strongly lagged mLPFC

and mACC (**Figure 4B**, Wilcoxon test, P<0.001). Consistent with this finding we found a

significant difference of the Lead/Lag index for burst versus non-burst periods for stMUA of

sLPFC-mSTR pairs (**Figure 4C**, p<0.001, Wilcoxon test). Similar to the theta/alpha burst results,

the amplified lead of LPFC was largely due to spike-MUA pairs that lagged outside of bursts but

switched to lead during beta bursts (**Figure 4D**, **Figure S7H,I**). This switch from a lag- to a lead-

state during beta bursts was evident at the level of individual spike-MUA pairs as shown for an

example in **Figure 4E,F** (for more examples see **Figure S8**). In summary, the neuronal spike

output of LPFC amplified its net lead over MUA in the striatum during beta bursts (**Figure 4G**).

This effect that was driven by neurons of the lag-ensemble joining the lead-ensemble during beta

bursts (**Figure 4H**). The same pattern of results was evident in both monkeys (**Figure S9**).

Theta/alpha-tuned lead-ensemble in ACC responds to attention cue

We next asked whether the directional spiking interactions of neuronal ensembles changed

during selective attention states. Neural activity was recorded while monkeys used the color

onset of peripheral stimuli as attention cue to covertly shift attention left or right from a central

fixation point, and subsequently used the motion direction of the attended peripheral stimulus to

plan a saccadic choice up or down from the fixation point (**Figure 5A**) (Oemisch et al., 2019).

The saccadic choice was triggered a few hundred milliseconds later by a transient dimming of

the attended stimulus. We analyzed stMUA's for spike events occurring in a 0.5 s window

following the attention cue onset and found that spikes of ACC neurons showed significantly

amplified lead over MUA in the striatum during this attention period (**Figure 5B, Figure S10A**).

This attention modulation of the lead was evident at 12-30 ms time lag for sACC-mSTR pairs of

the lead-ensemble, at 4-27 ms time lag for the sACC-mSTR pairs of the theta/alpha-tuned lead-

ensemble, and at 15-25 ms time lag when considering all sACC-mSTR spike-MUA pairs

(randomization tests, all $P<0.05$; **Figure 5B**; **Figure S10A**). Notably, however, for the ACC

connection to LPFC (sACC-mLPFC pairs), the amplified lead during the attention period was

limited to the stMUA pairs of the theta/alpha-lead-ensembles with a significant stMUA

difference at 7-22 ms time lag (randomization test, P<0.05; **Figure 5B**). We quantified the

attention effect as the difference of the Lead/Lag index values during the attention cue period

versus irrespective of that period, which confirmed that on average neuronal spiking in the ACC

amplified its lead over the striatum following the onset of the attention cue (Wilcoxon test,

P=0.05; **Figure 5C**). Attention also amplified the lead for the subset of sACC-mSTR pairs that

showed an ACC lead over the striatum (Wilcoxon test, P=0.03; **Figure 5D**) and for sACC-mSTR

pairs that led the striatum during theta/alpha-bursts (Wilcoxon test, P=0.03; **Figure 5E**).

Analyzing the Lead/Lag index also confirmed that attention exclusively amplified the lead of the

theta/alpha-tuned lead-ensemble of sACC over mLPFC (Wilcoxon test, P=0.03; **Figure 5E**;

Figure S10).

We next asked whether neurons of stMUA pairs of the lead and lag-ensembles show differential

firing rate dynamics after the attention cue onset. We next computed the peri-stimulus time

histogram (PSTH) around the time of the attention cue for neurons labeled either as lead or lag

based on their stMUA profiles during theta/alpha bursts. In the ACC, neurons identified by a

theta/alpha-specific lead showed larger and distinct firing rate dynamics than ACC neurons of

stMUA pairs that lagged during theta/alpha bursts (**Figure 5F.**) The firing rates of neurons with a

lead versus lag relationship during theta/alpha bursts were significantly different for sACC-

mSTR pairs at 240-750 ms and for sACC-mLPFC pairs at 360-610 ms following attention-cue

onset (permutation tests, P<0.05). The group of neurons in ACC that showed a theta/alpha-band

specific lead relation can be considered to constitute an *attention modulated theta/alpha-lead*

subnetwork. In contrast to the attentional effect of this theta/alpha-lead subnetwork, neurons

identified by their lead or lag during beta bursts did not show different attentional effects on their

firing rate dynamics (**Figure S10C,D**).

ACC spiking lead LPFC and striatum during a choice

So far, we found that ACC neurons of the theta/alpha-lead subnetwork amplified their lead over

LPFC and striatum in response to an attention cue. A different change of routing states occurred

during the preparation of the choice immediately prior to the saccadic eye movement. In

response to the transient dimming of the peripherally attended stimulus, the stMUAs with

neurons from both, LPFC and ACC, showed an amplified lead over MUA in the striatum

(**Figure 6A**). This amplified lead was evident when considering all spike-MUA pairs (the overall

population level), when considering the subgroup of spike-MUA pairs that led the striatum (the

level of a leading directional subnetwork), and - with a later onset – when considering pairs of

ACC or LPFC neurons that lagged MUA in the striatum (time periods with significantly

amplified lead are shown in **Figure 6A**; for sACC-mSTR pairs: at 9-19 ms and 46-60 ms time

lag in the lead-ensembles and at 54-70 ms time lag in the lag-ensembles; for sLPFC-mSTR pairs:

at 12-75+ ms time lag in the lead-ensembles, and at 25-51 ms time lag in the lag-ensembles,

randomization test, $P<0.05$). In addition, during the choice period, the lead of stMUAs with

spikes from ACC and MUA in LPFC was amplified in the lead-ensembles (at 22-41ms time lag,

randomization test, $P<0.05$; **Figure6A**). We confirmed the finding of an amplified lead during

the choice by computing the Lead/Lag index values for all sACC-mSTR and sLPFC-mSTR

stMUA pairs (Wilcoxon test, $P=0.03$ and $P=0.01$, respectively; **Figure 6B,C**), and for sACC-

mSTR and sLPFC-mSTR stMUAs pairs of the lead-ensembles (Wilcoxon test, $P<0.01$, for both

area pairs, **Figure 6C**). In addition, the Lead/Lag index showed a significantly amplified lead of

sSTR-mLPFC stMUAs of the lead-ensemble (Wilcoxon test, P=0.016, **Figure 6C**).

Similar to the analysis of attention, we asked whether neurons belonging to lead- and lag-

ensembles responded differently during a choice. We found that the ACC neurons that led over

the MUA in the striatum and LPFC showed on average stronger and distinct firing rate dynamics

than neurons with spikes lagging MUA (**Figure6D**). The firing response difference was

significant at -270-750+ ms and 210-430 ms after the onset of the choice-triggering dimming of

the peripheral stimulus for neurons identified by their lead in the sACC-mSTR and sACC-

mLPFC ensembles, respectively (permutation tests, both P<0.05). In LPFC, neurons that lagged

MUA in the striatum showed distinct and stronger firing responses during the choice than

neurons leading MUA in the striatum (**Figure 6E**, 350-680 ms after the dimming choice event,

permutation test, P<0.05). Neurons of lead and lag-ensembles of sLPFC-mACC pairs that their

lead or lag were not modulated during functional states did not show different firing rate

modulation during the choice (**Figure6E**).

314
315

Discussion

Here, we quantified the spatio-temporal dynamics of interareal spiking interactions and found

that directional routing states emerged within ~20 ms time windows. Despite sparse anatomical

connectivity of medial and lateral prefrontal cortex to the striatum (Ferry et al., 2000; Morris et

al., 2016), our methodology identified lead and lag activity patterns between these brain areas

contiguous with their known anatomical connectivity (**Figure 2**)(Ferry et al., 2000; Haber and

Knutson, 2010). On average, spikes in the LPFC led MUA in the ACC and striatum, while spikes

in the striatum lagged MUA in the ACC and LPFC (**Figure 2**). The fast ~20 ms timescale

characterization of these directional spiking interactions in the fronto-striatal network is new for

both, rodents and non-human primates. In rodents cross-areal, fronto-striatal cell assemblies with

fast temporal coordination have been recently described, but without elucidating the

directionality of spiking interactions (Oberto et al., 2021). In primates, previous studies

succeeded to quantify neuronal interactions but with time resolutions of hundreds of milliseconds

and across populations of neurons (Antzoulatos and Miller, 2011; Oemisch et al., 2015). Thus

far, directional networks have been reported mostly in sensory brain areas using spiking activity

patterns of groups of neurons (Semedo et al., 2019; van Kempen et al., 2021), or in higher-order

brain areas by inferring directional neuronal communication using LFP measures (Gregoriou et

al., 2009; Salazar et al., 2012; Antzoulatos and Miller, 2014; Vezoli et al., 2021). The results of

our study not only quantified lead and lag coordination of spiking activity at a fast <20 ms

timescale across distant higher order brain areas, but also provide direct insight about the

contributions of single neurons in forming these directional routing states.

We found that fast timescale spiking interactions were routed through frequency specific

channels that were evident in the LFP dynamics (**Figure 1F-J**). The LPFC led striatal activity

stronger when its spike occurred during 15-25 Hz beta bursts (**Figure 4A-C**). In the ACC, spikes

that occurred during 8-14 Hz theta/alpha bursts switched their directional relation to lead MUA

in the LPFC (**Figure 3A-C**). Both, the amplification and switching effects were primarily driven

by spike-lagging neuronal ensembles switching direction during LFP bursts (**Figure 3D,4D**).

This result indicates that the transitioning between routing states is likely determined by

modulating the timing of outgoing neural activities in spike-lagging neurons within their local

population into a more synchronized state with spike-leading neurons. The network simulation

showed that spectral peaks of LFP activity matched spectral peaks of the stMUA profiles,

suggesting that oscillatory bursts more likely occur at frequencies that correspond to the

resonance frequency of outgoing neuronal spiking activities (**Figure 1D,E**). These resonance

frequencies of outgoing spiking activities reflect synaptic time decays in each senders' brain area

(Hahn et al., 2014). During oscillatory bursts, spikes occur more synchronized at the resonance

frequency of their local population which as a consequence enhances the efficacy to activate

postsynaptic neurons in a recipient area (Kumar et al., 2008; Hahn et al., 2014), and facilitates

successful information routing between brain areas through frequency specific channels, as

previously shown in models (Palmigiano et al., 2017; Hahn et al., 2019) and at the level of LFP-

LFP interactions(Canolty et al., 2010).

Our study suggests that the ACC plays a significant role in mediating the routing of information

during functionally important events. We found that the directionality of spiking interactions

changed transiently during selective attention and decision-making. This functional specificity of

the routing states might reflect a hierarchical organization of fronto-striatal network

communication. The ACC predominated the network by amplifying its lead over the LPFC and

striatum during both decision-making and selective attention (**Figure 5E,6C**). In contrast, the

lead of LPFC spiking output was only amplified during decision making and over the striatum

(**Figure 6A-C**). These results illustrate a functional hierarchy at the level of single neuron

communication states. In this hierarchy, the ACC ranks higher than the LPFC, and both rank

higher than the striatum consistent with previous studies of hierarchical network organization

(Haber and Knutson, 2010; Wang, 2022). According to this view, fronto-striatal network

interactions do not proceed with a default lead or lag relation, but different nodes of the network

can dynamically gain modulate the temporal lead of the network depending on the functional

states. The transient amplification in the lead of ACC output reflects its higher rank in the

network hierarchy and is consistent with a role of the ACC in top-down control of the PFC when

attention is allocated, prediction errors are evaluated, or uncertainty is estimated to determine the

degree of top-down control (Shenhav et al., 2013; Womelsdorf and Everling, 2015; Banaie

Boroujeni et al., 2021). The functionally transient, rather than persistent 'control state' may be

related to the stronger inhibitory activity of the ACC (both intra- and interareal inhibitory

control)(Medalla and Barbas, 2009; Medalla et al., 2017), which imposes high biological costs

for this area to exert its function. Such transient control states – as opposed to stable, or

'persistent' control states - might facilitate the flexible regulation of routing states depending on

rapidly changing task demands (Shenhav et al., 2013; Aben et al., 2020; Boroujeni et al., 2022).

Lastly, we showed that the functional modulation of the lead and lag relation of a neuron's spike

within ~20 ms time windows predicted the firing rate response at a slower ~150 ms timescale.

We found that neurons with functionally gain modulated spike-lead over distant MUA (i.e. the

theta/alpha-leading ensembles during selective attention, and lead-ensembles during a choice)

showed stronger and distinct firing rate dynamics of functional responses when compared with

neurons from lag-ensembles (**Figure 5F,6D**). These results imply that variable timescales of

information processing previously associated with different brain areas (Murray et al., 2014;

Luczak et al., 2015; Wang, 2022) can be more precisely understood in a relative framework that

quantifies the temporal relation of spiking activity in one brain area relative to another brain area.

In such a relative framework, rather than analyzing aggregated functional responses of groups of

neurons in individual brain regions, the precise, cross-areal temporal sequencing of neuronal

activity is the central unit of functional relevance.

Together, our results demonstrate, to our knowledge for the first time the spatio-temporal

dynamics of spiking activity interactions across multiple spatial and temporal scales and between

distant brain areas during complex cognitive functions. We have shown that fast directional

spiking interactions across distant brain areas instantiate routing states that dynamically switch or

amplify during coherent local population activity and when the fronto-striatal network

contributes to selective attention and decision-making. Such fast spatiotemporally characterized

dynamics predict neuronal activity at slower timescales during relevant functional states.

Author Contributions

K.B.B. and T.W. conceived the research. K.B.B. developed the methodology. K.B.B.

conceptualized and performed modeling, analysis, and visualization. T.W. Supervision. K.B.B.

and T.W. wrote the paper.

Data and materials availability: Code and analysis pipelines will be publicly available upon
publication.

Acknowledgments: This work was supported by the National Institute of Mental Health
(R01MH123687), by the National Institute of Biomedical Imaging and Bioengineering
(R01EB028161), and by the Canadian Institutes of Health Research CIHR (Grant
MOP_102482). The funders had no role in study design, data collection and analysis, the
decision to publish, or the preparation of this manuscript.

Competing Interests Declaration: The authors declare no competing interests.
417

418 **Figure Legends**

Figure 1. ACC and LPFC spiking network simulations shows systematic inter-network lead and lag temporal orders switching during 8-14 Hz LFP bursts. (A) Two networks with excitatory (blue) and inhibitory (red) neurons. Black dots are recording sites in each network measuring the transmembrane and postsynaptic potentials used to extract spiking, multiunit activity (MUA), and local field potentials (LFPs). Raster plots of spiking activity for interneurons (red, INs), and bursting and regular excitatory neurons (blue, PNs). **(B)** Average spike-triggered (st)MUA of the simulated networks 1 (LPFC) and 2 (ACC). Intra-network spikes and MUA show near-symmetric temporal relationships with zero-time lag peaks and oscillatory side-lobes. Inter-network stMUAs have asymmetric shape indicating temporal lead (red shading) and lag (blue shading) relationships. **(C)** Median Lead/Lag index values across all spike-MUA pairs. LPFC spikes (sLPFC) led MUA in the ACC (mACC) network. White dots and error bars show median ±SE Lead/Lag index value. **(D,E)** Power spectral density (PSD) of LFP (D) and of the stMUA (E) recorded in each network. **(F)** LFP power spectrogram in the ACC with a theta/alpha burst at ~2-2.5 sec. Black contours mark LFP burst. **(G)** Example beta burst from LPFC. **(H)** Time-resolved Lead/Lag index values for sACC-mLPFC with a theta/alpha burst occurring at ~2-2.5 sec. At the time of the burst sACC switched to lead mLPFC. **(I)** Same format as h for mLPFC-sACC. **(J)** The stMUA differences between spikes occurred during versus outside of LFP theta/alpha bursts for sACC-mLPFC (top) and sLPFC-mACC (bottom). Shades indicate significant stMUA in 20 ms preceding (blue) or following (red) spike events.

436
Figure 2. Temporal orders of spiking activity across fronto-striatal areas. (**A**) Neuronal activity was
simultaneously recorded from ACC (area 24, red), LPFC (areas 46 and 9, blue), and striatum (caudate and ventral
striatum, green). (**B**) Intra-areal stMUA. red/blue signify stMUA values significantly higher/lower than all intra-
areal pairs population. (**C**) Power spectral density over stMUA has stronger power peak at ~16-22 Hz in the LPFC,
and at ~8-14 Hz in the ACC. (**D**) Inter-areal stMUA, Red lines indicate periods when the stMUA of the area pair is
significantly higher than the random assignment of areas to the population of spike-MUA pairs (randomization test,
P<0.05). Blue/red shades indicate significant non-zero MUA within 20 preceding/following the spike time,
respectively. Post-spike MUA was significantly enhanced for spike-MUA pairs: sLPFC-mSTR, sACC-mSTR,
sLPFC-mACC at p<0.001; and for sACC-mLPFC at p=0.03, Wilcoxon test, FDR corrected for dependent samples.
€ Median time lags of the stMUA peaks. Negative/positive values indicate MUA peaked more preceding/following
the spike time (at 0 ms). sLPFC led over mACC and mSTR, mSTR lag behind sACC and sLPFC. Vertical lines
show significant pair-wise comparisons (Kruskal-Wallis test, P<0.001, Dunn's test, P<0.01). Asterisks denote area
pairs with time-lags significantly different from zero. (**F**) Lead-Lag index values for all intra-areal (diagonal) and
inter-areal (off-diagonal) spike-MUA pairs. sSTR lagged behind mLPFC and mACC, while sLPFC led over mSTR
and mACC (Wilcoxon test; p<0.001 for sSTR-mLPFC and sSTR-mACC, p=0.03 for sLPFC-mSTR, FDR corrected
for dependent samples). White dots and SE's show median ± SE. White asterisks denote values significantly
different than zero. P-values are corrected for multiple comparisons (see STAR Methods). (**G**) Schematic summary
of inter-areal interactions. Red/blue arrows indicate spikes lead/lag MUA, respectively. Line thicknesses reflects
lead or lag strengths.
456

459 **Figure 3. ACC spikes switch to lead over LPFC multiunit activity during Theta/alpha (8-14 Hz) bursts. (A)** A
460 wideband LFP example segment with 8-14 Hz bursts in red (top) and bandpass filtered signal (bottom) with
461 envelope in green. **(B)** The stMUA difference for spikes inside versus outside of theta/alpha bursts for area-pair
462 combinations. Red lines indicate periods where in burst stMUA is significantly higher the area-pair than the
463 randomized population. Red and blue shades indicate significant non-zero stMUA difference of theta/alpha burst
464 versus non-burst within 20 ms following/preceding spikes, respectively. **(C)** Similar Lead-Lag index adjacency
465 matrix as in **(Figure 2f)** for stMUA differences of spikes during versus outside of theta/alpha bursts. White dots and
466 SE's show median ± SE. White asterisks denote values significantly different than zero. **(D)** Lead/Lag index values
467 flipped signs during theta/alpha bursts in the sACC-mLPFC lag-ensembles (Kruskal Wallis test, P=0.01, Wilcoxon
468 test, P=0.03 for nonzero ACC-LPFC. Black horizontal lines show significant differences between area-pairs). **(E)**
469 LFP segments for a spike-MUA pair example around spikes (±75 ms), for spikes inside (red) versus outside (grey)
470 of the theta/alpha bursts. **(F)** The spike-MUA pair example in **(E)** switched from a lag to a lead state. Red, grey, and
471 green colors denote stMUAs inside bursts, outside bursts, and their difference, respectively. **(G)** Schematic summary
472 of theta/alpha burst modulated inter-areal interactions. The filled red/blue portion of the lines indicates where spikes
473 lead and lag relative to theta/alpha bursts. **(H)** Schematic summary of lag to lead switch in the sACC-mLPFC during
474 theta/alpha burst.

475
476

477
478

479
Figure 4. LPFC spikes amplified their lead over striatum during beta (15-25 Hz) bursts. (A) A wideband LFP
example segment with beta bursts in blue (top) and bandpass filtered signal (bottom) with envelope in green. **(B)**
Average differences of stMUA's inside versus outside of beta bursts for area-pair combinations. Red lines indicate
periods where in burst stMUA is significantly higher the area-pair than the randomized population. Red and blue
shades indicate significant non-zero stMUA difference of beta burst versus non-burst within 20 ms
following/preceding spikes, respectively. **(C)** Similar Lead-Lag index adjacency matrix as in **(Figure 3C)** for
stMUA differences of spikes during versus outside of beta bursts. White dots and SE's show median ± SE. White
asterisks denote values significantly different than zero. **(D)** Lead/Lag index values was flipped for lag-ensembles in
the sLPFC-mSTR during beta bursts. Black horizontal lines show significant differences between area-pairs. **(E)**
LFP segments for a spike-MUA pair example around spikes (±75 ms), for spikes inside (blue) versus outside (grey)
of the beta bursts. **(F)** Beta burst modulated stMUA for the spike-MUA pair example in **(E)** for spikes inside (blue)
and outside (grey) of the bursts, and their difference (green). **(G)** Schematic summary of beta burst modulated inter-
areal interactions. The filled red/blue portion of the lines indicates where spikes lead and lag relative to beta bursts.
(H) Schematic summary of lead amplification in the sLPFC-mSTR during beta burst.
494
495
496
497

Figure 5. Theta/alpha-subnetworks of neurons in the ACC route attentional states. (A) Monkeys performed a
task requiring covertly attending a peripheral stimulus and making a saccadic response to a transient luminance
change of the attended stimulus to receive reward (see **STAR Methods**). The stimulus color-reward association
reversed after ~30 trials**. (B)** Attention-modulated stMUA for all neurons (black), theta/alpha-lead (red) and
theta/alpha-lag (blue) neuronal ensembles. The sACC showed stronger lead over mSTR in response to an attention
cue on a population level and over mLPFC only in theta/alpha-lead-ensembles. Black/red horizontal lines show
significant nonzero attention-modulated stMUA for all neurons and theta/alpha-lead neuronal ensembles,
respectively. (**C-E**) Attention-modulated lead/lag index for all neurons (**C**), lead-ensembles (**D**), and theta/alpha-
lead-ensembles (**E**). Bars in red are significantly different than zero (Kruskial Wallis test, P<0.01 for all three, and
Wilcoxon test, P<0.05, FDR corrected, see **STAR Methods**). Black horizontal lines denote pairs with significantly
different lead/lag index**. f,** Neuronal spike rate dynamics in response to attention cue for (**F**) the ACC neurons
identified by their temporal orders during theta/alpha bursts relative to the mSTR (left) and the mLPFC (right). **(G)**
Same as in (**F**) but for LPFC neurons identified by their temporal orders during theta/alpha bursts relative to the
mSTR (left) and the mACC (right). The theta/alpha-lead and theta/alpha-lag subnetworks are shown in red and blue
respectively. gray horizontal lines denote time periods were the two subnetworks show different rate response
dynamics (permutation test, P<0.05, see **STAR Methods**). Overall, the ACC neurons identified by their theta/alpha-
lead over striatum showed earlier onset response compared with the LPFC. Error bars denote standard error of the
mean.

519
Figure 6. The ACC predominates the fronto-striatal network response to a choice. **(A)** Choice-
modulated stMUA for all neurons (black), lead (red) and lag (blue) neuronal ensembles. The sACC and sLPFC show
stronger lead over mSTR during a choice on a population level and both the lead- and lag-neuronal ensembles. The
sACC showed stronger lead over mLPFC on a population level and the lead-neuronal ensembles (randomization
test, P<0.05, see **STAR Methods**). Horizontal lines denote significant nonzero stMUA for all neurons (black), lead
(red) and lag (blue) neuronal ensembles, indicating earlier lead-onset for lead neuronal ensembles. **(B,C)** Choice-
modulated lead/lag index for all neurons **(B)**, and lead-ensembles **(C)**. Bars in red are significantly different than
zero. Black horizontal lines denote pairs of significantly different lead/lag index (Kruskial Wallis test, P<0.01 for all
two, and Wilcoxon test, P<0.05, FDR corrected, see **STAR Methods**). **(D),** Neuronal spike rate dynamics in
response to Go-cue for the ACC neurons identified by their temporal orders relative to the mSTR (left) and the
mLPFC (right). **(E)** Same as in **(D)** but for LPFC neurons identified by their temporal orders relative to the mSTR
(left) and the mACC (right). The lead/lag subnetworks are shown in red/blue, respectively. Gray horizontal lines
denote time periods were the two subnetworks showed different rate response dynamics (permutation test, P<0.05,
see **STAR Methods**). Overall, the ACC neurons identified by their lead over striatum showed earlier onset response
compared with the LPFC. The LPFC showed stronger response dynamics in the neurons that lagged the striatum.
Error bars denote standard error of the mean.

References

Aben B, Calderon CB, Bussche EV den, Verguts T (2020) Cognitive Effort Modulates Connectivity between Dorsal Anterior Cingulate Cortex and Task-Relevant Cortical Areas. J Neurosci 40:3838–3848.

Amatrudo JM, Weaver CM, Crimins JL, Hof PR, Rosene DL, Luebke JI (2012) Influence of Highly Distinctive Structural Properties on the Excitability of Pyramidal Neurons in Monkey Visual and Prefrontal Cortices. J Neurosci 32:13644–13660.

Antzoulatos EG, Miller EK (2011) Differences between Neural Activity in Prefrontal Cortex and Striatum during Learning of Novel Abstract Categories. Neuron 71:243–249.

Antzoulatos EG, Miller EK (2014) Increases in Functional Connectivity between Prefrontal Cortex and Striatum during Category Learning. Neuron 83:216–225.

Banaie Boroujeni K, Tiesinga P, Womelsdorf T (2021) Interneuron specific gamma synchronization indexes cue uncertainty and prediction errors in lateral prefrontal and anterior cingulate cortex Haegens S, ed. eLife 10:e69111.

Barral J, D Reyes A (2016) Synaptic scaling rule preserves excitatory–inhibitory balance and salient neuronal network dynamics. Nat Neurosci 19:1690–1696.

Bartos M, Elgueta C (2012) Functional characteristics of parvalbumin- and cholecystokinin-expressing basket cells. The Journal of Physiology 590:669–681.

Benjamini Y, Yekutieli D (2005) False Discovery Rate–Adjusted Multiple Confidence Intervals for Selected Parameters. Journal of the American Statistical Association 100:71–81.

Billeh YN, Cai B, Gratiy SL, Dai K, Iyer R, Gouwens NW, Abbasi-Asl R, Jia X, Siegle JH, Olsen SR, Koch C, Mihalas S, Arkhipov A (2020) Systematic Integration of Structural and Functional Data into Multi-scale Models of Mouse Primary Visual Cortex. Neuron 106:388-403.e18.

Boashash B (1992) Estimating and Interpreting the Instantaneous Frequency of a Signal – Part 1: Fundamentals. In: Proceedings of the Ieee, pp 520–538.

Boroujeni KB, Oemisch M, Hassani SA, Womelsdorf T (2020) Fast spiking interneuron activity in primate striatum tracks learning of attention cues. PNAS 117:18049–18058.

Boroujeni KB, Sigona MK, Treuting RL, Manuel TJ, Caskey CF, Womelsdorf T (2022) Anterior cingulate cortex causally supports flexible learning under motivationally challenging and cognitively demanding conditions. PLOS Biology 20:e3001785.

Canolty RT, Ganguly K, Kennerley SW, Cadieu CF, Koepsell K, Wallis JD, Carmena JM (2010)
Oscillatory phase coupling coordinates anatomically dispersed functional cell assemblies.
PNAS 107:17356–17361.

Chang Y-M, Luebke JI (2007) Electrophysiological Diversity of Layer 5 Pyramidal Cells in the
Prefrontal Cortex of the Rhesus Monkey: In Vitro Slice Studies. Journal of
Neurophysiology 98:2622–2632.

Chicco D, Jurman G (2020) The advantages of the Matthews correlation coefficient (MCC) over
F1 score and accuracy in binary classification evaluation. BMC Genomics 21:6.

Choi Y-S, Koenig MA, Jia X, Thakor NV (2010) Quantifying Time-Varying Multiunit Neural
Activity Using Entropy-Based Measures. IEEE Trans Biomed Eng
57:10.1109/TBME.2010.2049266.

Condé F, Lund JS, Jacobowitz DM, Baimbridge KG, Lewis DA (1994) Local circuit neurons
immunoreactive for calretinin, calbindin D-28k or parvalbumin in monkey prefronatal
cortex: Distribution and morphology: CALCIUM-BINDING PROTEINS IN MONKEY
PREFRONTAL CORTEX. J Comp Neurol 341:95–116.

Dombrowski SM, Hilgetag CC, Barbas H (2001) Quantitative Architecture Distinguishes
Prefrontal Cortical Systems in the Rhesus Monkey. Cerebral Cortex 11:975–988.

Dunn OJ (1961) Multiple Comparisons among Means. Journal of the American Statistical
Association 56:52–64.

Einevoll GT, Kayser C, Logothetis NK, Panzeri S (2013) Modelling and analysis of local field
potentials for studying the function of cortical circuits. Nat Rev Neurosci 14:770–785.

Feingold J, Gibson DJ, DePasquale B, Graybiel AM (2015) Bursts of beta oscillation
differentiate postperformance activity in the striatum and motor cortex of monkeys
performing movement tasks. PNAS 112:13687–13692.

Ferry AT, Öngür D, An X, Price JL (2000) Prefrontal cortical projections to the striatum in
macaque monkeys: Evidence for an organization related to prefrontal networks. Journal
of Comparative Neurology 425:447–470.

Fiebelkorn IC, Pinsk MA, Kastner S (2018) A Dynamic Interplay within the Frontoparietal
Network Underlies Rhythmic Spatial Attention. Neuron 99:842-853.e8.

Freund TF, Katona I (2007) Perisomatic Inhibition. Neuron 56:33–42.

Fries P (2015) Rhythms for Cognition: Communication through Coherence. Neuron 88:220–235.

Gabbott PLA, Bacon SJ (1996) Local circuit neurons in the medial prefrontal cortex (areas
24a,b,c, 25 and 32) in the monkey: I. Cell morphology and morphometrics. Journal of
Comparative Neurology 364:567–608.

Gilman JP, Medalla M, Luebke JI (2017) Area-Specific Features of Pyramidal Neurons—a Comparative Study in Mouse and Rhesus Monkey. Cerebral Cortex 27:2078–2094.

Gorodkin J (2004) Comparing two K-category assignments by a K-category correlation coefficient. Computational Biology and Chemistry 28:367–374.

Gregoriou GG, Gotts SJ, Zhou H, Desimone R (2009) High-Frequency, Long-Range Coupling Between Prefrontal and Visual Cortex During Attention. Science 324:1207–1210.

Haber SN, Knutson B (2010) The Reward Circuit: Linking Primate Anatomy and Human Imaging. Neuropsychopharmacology 35:4–26.

Hahn G, Bujan AF, Frégnac Y, Aertsen A, Kumar A (2014) Communication through Resonance in Spiking Neuronal Networks. PLOS Computational Biology 10:e1003811.

Hahn G, Ponce-Alvarez A, Deco G, Aertsen A, Kumar A (2019) Portraits of communication in neuronal networks. Nat Rev Neurosci 20:117–127.

Hasenstaub A, Shu Y, Haider B, Kraushaar U, Duque A, McCormick DA (2005) Inhibitory Postsynaptic Potentials Carry Synchronized Frequency Information in Active Cortical Networks. Neuron 47:423–435.

Huang NE, Wu Z (2008) A review on Hilbert-Huang transform: Method and its applications to geophysical studies. Reviews of Geophysics 46 Available at: https://onlinelibrary.wiley.com/doi/abs/10.1029/2007RG000228 [Accessed November 16, 2021].

Izhikevich EM (2003) Simple model of spiking neurons. IEEE Transactions on Neural Networks 14:1569–1572.

Izhikevich EM (2004) Which model to use for cortical spiking neurons? IEEE Transactions on Neural Networks 15:1063–1070.

Jin DZ, Fujii N, Graybiel AM (2009) Neural representation of time in cortico-basal ganglia circuits. Proceedings of the National Academy of Sciences 106:19156–19161.

Kaiser M, Hilgetag CC, van Ooyen A (2009) A Simple Rule for Axon Outgrowth and Synaptic Competition Generates Realistic Connection Lengths and Filling Fractions. Cereb Cortex 19:3001–3010.

Karnani MM, Jackson J, Ayzenshtat I, Tucciarone J, Manoocheri K, Snider WG, Yuste R (2016) Cooperative Subnetworks of Molecularly Similar Interneurons in Mouse Neocortex. Neuron 90:86–100.

Kruskal WH, Wallis WA (1952) Use of Ranks in One-Criterion Variance Analysis. Journal of the American Statistical Association 47:583–621.

Kumar A, Rotter S, Aertsen A (2008) Conditions for Propagating Synchronous Spiking and
Asynchronous Firing Rates in a Cortical Network Model. J Neurosci 28:5268–5280.

Legatt AD, Arezzo J, Vaughan HG (1980) Averaged multiple unit activity as an estimate of
phasic changes in local neuronal activity: effects of volume-conducted potentials. Journal
of Neuroscience Methods 2:203–217.

Lewis C, Bosman C, Fries P (2015) Recording of brain activity across spatial scales. Current
Opinion in Neurobiology 32:68–77.

Logothetis NK, Kayser C, Oeltermann A (2007) In Vivo Measurement of Cortical Impedance
Spectrum in Monkeys: Implications for Signal Propagation. Neuron 55:809–823.

Luczak A, McNaughton BL, Harris KD (2015) Packet-based communication in the cortex. Nat
Rev Neurosci 16:745–755.

Lundqvist M, Rose J, Herman P, Brincat SL, Buschman TJ, Miller EK (2016) Gamma and Beta
Bursts Underlie Working Memory. Neuron 90:152–164.

Markram H, Lübke J, Frotscher M, Sakmann B (1997) Regulation of Synaptic Efficacy by
Coincidence of Postsynaptic APs and EPSPs. Science 275:213–215.

Medalla M, Barbas H (2009) Synapses with Inhibitory Neurons Differentiate Anterior Cingulate
from Dorsolateral Prefrontal Pathways Associated with Cognitive Control. Neuron
61:609–620.

Medalla M, Barbas H (2010) Anterior Cingulate Synapses in Prefrontal Areas 10 and 46 Suggest
Differential Influence in Cognitive Control. J Neurosci 30:16068–16081.

Medalla M, Gilman JP, Wang J-Y, Luebke JI (2017) Strength and Diversity of Inhibitory
Signaling Differentiates Primate Anterior Cingulate from Lateral Prefrontal Cortex. J
Neurosci 37:4717–4734.

Medalla M, Luebke JI (2015) Diversity of Glutamatergic Synaptic Strength in Lateral Prefrontal
versus Primary Visual Cortices in the Rhesus Monkey. J Neurosci 35:112–127.

Melchitzky DS, Lewis DA (2008) Dendritic-targeting GABA neurons in monkey prefrontal
cortex: Comparison of somatostatin- and calretinin-immunoreactive axon terminals.
Synapse 62:456–465.

Milstein J, Mormann F, Fried I, Koch C (2009) Neuronal Shot Noise and Brownian 1/f2
Behavior in the Local Field Potential. PLOS ONE 4:e4338.

Morris LS, Kundu P, Dowell N, Mechelmans DJ, Favre P, Irvine MA, Robbins TW, Daw N,
Bullmore ET, Harrison NA, Voon V (2016) Fronto-striatal organization: Defining
functional and microstructural substrates of behavioural flexibility. Cortex 74:118–133.

Murray JD, Bernacchia A, Freedman DJ, Romo R, Wallis JD, Cai X, Padoa-Schioppa C,
Pasternak T, Seo H, Lee D, Wang X-J (2014) A hierarchy of intrinsic timescales across
primate cortex. Nat Neurosci 17:1661–1663.

Oberto VJ, Boucly CJ, Gao H, Todorova R, Zugaro MB, Wiener SI (2021) Distributed cell
assemblies spanning prefrontal cortex and striatum. Current Biology Available at:
https://www.sciencedirect.com/science/article/pii/S0960982221013610 [Accessed
January 8, 2022].

Oemisch M, Westendorff S, Azimi M, Hassani SA, Ardid S, Tiesinga P, Womelsdorf T (2019)
Feature-specific prediction errors and surprise across macaque fronto-striatal circuits.
Nature Communications 10:1–15.

Oemisch M, Westendorff S, Everling S, Womelsdorf T (2015) Interareal Spike-Train
Correlations of Anterior Cingulate and Dorsal Prefrontal Cortex during Attention Shifts. J
Neurosci 35:13076–13089.

Palmigiano A, Geisel T, Wolf F, Battaglia D (2017) Flexible information routing by transient
synchrony. Nat Neurosci 20:1014–1022.

Perin R, Berger TK, Markram H (2011) A synaptic organizing principle for cortical neuronal
groups. PNAS 108:5419–5424.

Pewsey A (2002) Large-Sample Inference for the General Half-Normal Distribution.
Communications in Statistics - Theory and Methods 31:1045–1054.

Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M (2013) Inhibition of inhibition in visual
cortex: the logic of connections between molecularly distinct interneurons. Nat Neurosci
16:1068–1076.

Potjans TC, Diesmann M (2014) The Cell-Type Specific Cortical Microcircuit: Relating
Structure and Activity in a Full-Scale Spiking Network Model. Cerebral Cortex 24:785–
806.

Salazar RF, Dotson NM, Bressler SL, Gray CM (2012) Content-Specific Fronto-Parietal
Synchronization During Visual Working Memory. Science 338:1097–1100.

Semedo JD, Zandvakili A, Machens CK, Yu BM, Kohn A (2019) Cortical Areas Interact through
a Communication Subspace. Neuron 102:249-259.e4.

Shenhav A, Botvinick MM, Cohen JD (2013) The Expected Value of Control: An Integrative
Theory of Anterior Cingulate Cortex Function. Neuron 79:217–240.

Siegel M, Donner TH, Engel AK (2012) Spectral fingerprints of large-scale neuronal
interactions. Nat Rev Neurosci 13:121–134.

Singer W (2013) Cortical dynamics revisited. Trends in Cognitive Sciences 17:616–626.

Smith MA, Kohn A (2008) Spatial and Temporal Scales of Neuronal Correlation in Primary
Visual Cortex. J Neurosci 28:12591–12603.

Supèr H, Roelfsema PR (2005) Chronic multiunit recordings in behaving animals: advantages
and limitations. In: Progress in Brain Research, pp 263–282 Development, Dynamics and
Pathiology of Neuronal Networks: from Molecules to Functional Circuits. Elsevier.
Available at: https://www.sciencedirect.com/science/article/pii/S0079612304470204
[Accessed November 17, 2021].

van Ede F, Quinn AJ, Woolrich MW, Nobre AC (2018) Neural Oscillations: Sustained Rhythms
or Transient Burst-Events? Trends in Neurosciences 41:415–417.

van Kempen J, Gieselmann MA, Boyd M, Steinmetz NA, Moore T, Engel TA, Thiele A (2021)
Top-down coordination of local cortical state during selective attention. Neuron 109:894-
904.e8.

Vezoli J, Vinck M, Bosman CA, Bastos AM, Lewis CM, Kennedy H, Fries P (2021) Brain
rhythms define distinct interaction networks with differential dependence on anatomy.
Neuron Available at:
https://www.sciencedirect.com/science/article/pii/S089662732100725X [Accessed
October 23, 2021].

Voloh B, Oemisch M, Womelsdorf T (2020) Phase of firing coding of learning variables across
the fronto-striatal network during feature-based learning. Nat Commun 11:4669.

Wang X-J (2022) Theory of the Multiregional Neocortex: Large-Scale Neural Dynamics and
Distributed Cognition. Annu Rev Neurosci 45:533–560.

Wilcoxon F (1945) Individual Comparisons by Ranking Methods. Biometrics Bulletin 1:80–83.

Womelsdorf T, Ardid S, Everling S, Valiante TA (2014) Burst Firing Synchronizes Prefrontal
and Anterior Cingulate Cortex during Attentional Control. Current Biology 24:2613–
2621.

Womelsdorf T, Everling S (2015) Long-Range Attention Networks: Circuit Motifs Underlying
Endogenously Controlled Stimulus Selection. Trends in Neurosciences 38:682–700.

Womelsdorf T, Schoffelen J-M, Oostenveld R, Singer W, Desimone R, Engel AK, Fries P
(2007) Modulation of Neuronal Interactions Through Neuronal Synchronization. Science
316:1609–1612.

Zaitsev AV, Gonzalez-Burgos G, Povysheva NV, Kröner S, Lewis DA, Krimer LS (2005)
Localization of Calcium-binding Proteins in Physiologically and Morphologically
Characterized Interneurons of Monkey Dorsolateral Prefrontal Cortex. Cerebral Cortex
15:1178–1186.

Zaitsev AV, Povysheva NV, Gonzalez-Burgos G, Rotaru D, Fish KN, Krimer LS, Lewis DA
742   (2009) Interneuron Diversity in Layers 2–3 of Monkey Prefrontal Cortex. Cerebral
743   Cortex 19:1597–1615.

744   Zich C, Quinn AJ, Mardell LC, Ward NS, Bestmann S (2020) Dissecting Transient Burst Events.
745   Trends in Cognitive Sciences 24:784–788.

746
747

748
749