1

A Mitochondrial Inside-Out Iron-Calcium Signal Identifies Drug Targets for Parkinson's Disease

Vinita Bharat[1], Aarooran S. Durairaj[1], Roeland Vanhauwaert[1], Li Li[1], Colin M. Muir[2,3], Sujyoti Chandra[1], Chulhwan S. Kwak[1], Yann Le Guen[4,5], Pawan Nandakishore[6], Chung-Han Hsieh[1], Stefano E. Rensi[7], Russ B. Altman[7], Michael D. Greicius[4], Liang Feng[2], and Xinnan Wang[1]*

Affiliations:

1. Department of Neurosurgery, Stanford University School of Medicine, Stanford, CA94305, USA;

2. Department of Molecular and Cellular Physiology, Stanford University School of Medicine, Stanford, CA94305, USA;

3. Graduate Program of Molecular and Cellular Physiology, Stanford University School of Medicine, Stanford, CA94305, USA;

4. Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA94305, USA;

5. Institut du Cerveau - Paris Brain Institute - ICM, 75013, Paris, France;

6. Vroom Inc., Houston, TX77042, USA;

7. Department of Bioengineering, Stanford University, Stanford, CA94305, USA.

* Correspondence and lead contact: xinnanw@stanford.edu

Summary

Genetic backgrounds and risk factors among individuals with Parkinson's disease (PD) are highly heterogenous, limiting our ability to effectively detect and treat PD. Discovering a common pathway connecting several PD risk factors may be crucial for finding a cure. Here we present one such pathway. Elevation of iron levels causes inward mitochondrial Ca^{2+} overflow, through an interaction of Fe^{2+} with mitochondrial calcium uniporter (MCU). Enhanced Ca^{2+} influx across the mitochondrial surface causes spatially confined Ca^{2+} elevation at the outer mitochondrial membrane (OMM), which is subsequently sensed by Miro1, a Ca^{2+}-binding protein. A Miro1 blood test distinguishes PD patients from controls and responds to drug treatment. Miro1-based drug screens in PD cells discover FDA-approved T-type Ca^{2+}-channel blockers. Human genetic analysis reveals enrichment of rare variants in T-type Ca^{2+}-channel subtypes associated with PD status. Our results identify a molecular mechanism in PD pathophysiology, and drug targets and candidates coupled with a convenient stratification method.

Main Text

Introduction

Parkinson's disease (PD) is a leading cause of disability, afflicting the aging population. The dopamine (DA)-producing neurons in the substantia nigra are the first to die in PD patients. A bottleneck that hinders our ability to effectively detect and treat PD may be the presence of highly heterogenous genetic backgrounds and risk factors among different patients. More than 90% of the PD cases are considered sporadic with no known causal mutations. Genome-wide association studies (GWAS) have identified over 90 risk loci [1]. Functional studies on known causal genes of familial patients and from cellular and animal PD models have pointed to multiple "cellular risk elements", such as mitochondrial damage, lysosomal dysfunction, immune system activation, neuronal calcium mishandling, and iron

accumulation [2-11]. These distinct genetic and cellular risk factors may confer individual heterogeneity in

disease onset, but also suggest that there are networks and pathways linking these "hubs" in disease

pathogenesis. Identifying their connections could be crucial for finding a cure for PD.

Mitochondria are the center of cellular metabolism and communication. Ions such as calcium and iron,

are not only essential for diverse mitochondrial functions but can be stored inside the mitochondria to

maintain cellular ionic homeostasis. Ion channels and transporters in the plasma and mitochondrial

membranes coordinate for ion uptake, transport, and storage. Calcium ions enter the cell via voltage- or

ligand-gated calcium channels across the cell surface. Inside the cell, they can be taken up by

mitochondria through the outer mitochondrial membrane (OMM) channel, VDAC, and the inner

mitochondrial membrane (IMM) channel, mitochondrial calcium uniporter (MCU) [12], and extruded into

the cytosol through the IMM transporter, NCLX [13]. MCU is a multimeric holocomplex consisting of

additional regulatory subunits, such as essential MCU regulator (EMRE), mitochondrial calcium uptake

1 (MICU1), MICU2, and MCUb [14-16]. Channels complementary to these major mitochondrial calcium

channels also exist [17]. For iron transport, iron bound carrier proteins such as transferrin deliver iron into

the cell through endocytosis and intracellular iron can be taken up by mitochondria via several IMM

transporters [18]. Although individually, dysregulation of iron or calcium homeostasis has been widely

reported in PD models, their mechanistic link and contribution to disease susceptibility remain elusive.

Identifying the cellular causes to neuron death will not only provide more effective disease management

but also shed light on molecular signatures shared by a subset of people affected by the disease. A

convenient, cost-effective method to spot the vulnerable population, even before the symptom onset,

will be extremely valuable for early intervention and preventive medicine. It will improve the efficacy of

clinical trials for testing experimental drugs, by facilitating patient stratification and serving as a

pharmacodynamic marker.

Understanding disease-causing cellular paths will also help us zoom in on rare genetic variants that

contribute to disease etiology but otherwise are difficult to discover through GWAS. Integrating studies

of risk variants with disease models and human tissues could establish a causal link of a biological

pathway to a disease with complex traits such as PD, with the promise of identifying more effective

druggable targets and biomarkers. In this work, we harness the power of combining human genetics,

cellular and in vivo models, and patient's tissues, and identify a mitochondrial inside-out iron-calcium-

Miro axis in PD. Iron accumulation causes calcium overflow into the mitochondria via promoting MCU

channel activity. The resultant subdomain dynamic changes of Ca^{2+} at the OMM can be sensed by

Miro1. This molecular mechanism in PD identifies drug targets and candidates and a companion

stratification method.

Results

Iron promotes the MCU complex assembly

The MCU's Ca^{2+}-import ability can be modulated by the elevation of cytosolic Ca^{2+} and

intramitochondrial oxidants [15,19]. We investigated whether the elevation of iron levels, a hallmark of PD

[10], could also regulate MCU activity. It has been shown that following the stimulation of G-protein-

coupled receptor (GPCR) agonist, the clearance of cytosolic Ca^{2+} is mediated by MCU [19-21]. To

investigate the relation of MCU and iron, we treated HEK293T cells with Fe^{2+}, stimulated these cells the

next day with the GPCR agonist, thrombin, and measured cytosolic and mitochondrial Ca^{2+} levels with

live Calcium Green and Rhod-2 staining, respectively. We found that thrombin triggered intracellular

Ca^{2+} mobilization and elevation, which was comparable between Fe^{2+}-treated and mock control cells

(Figure 1A-B). However, mitochondria in Fe^{2+}-treated cells sustained a significantly larger Ca^{2+}

elevation after thrombin stimulation, as compared to mock control (Figure 1C-D). These results indicate

that iron promotes the mitochondrial Ca^{2+}-import ability.

Because MCU oligomerization affects its localization and potentially its Ca^{2+}-uptake ability [15,19],

we next explored whether iron altered the assembly of the MCU complex. Native-PAGE is a sensitive

method to determine the overall form and amount of a multimeric native protein. The human MCU

oligomer bands from HEK cells migrated between 400-700 KDa in Native-PAGE (Figure 1E; the

negative control in the left lane, *MCU-/-,* showed no signal) [12,22,23]. Importantly, we found that Fe^{2+}

treatment for 22 hours resulted in an increase in the total intensity of the MCU oligomer bands and

appeared to shift some bands upward (Figure 1E), indicating the formation of higher-order oligomers.

Miro1, an OMM protein, also oligomerized and migrated as a single band around 480 KDa in Native-

PAGE, which was not significantly affected by Fe^{2+} treatment (Figure 1E). The EC_{50} of Fe^{2+} added to

the media to cause the MCU response was 9.334 μM (Figure 1F). We confirmed that HEK cells were

morphologically healthy under all concentrations and no significant cell death was observed (Figure

S1A). To corroborate the Native-PAGE result, we performed an alternative method to differentiate

protein complexes based on their molecular weight (MW)–size exclusion chromatography (SEC).

Protein complexes with higher MW are eluted faster than those with lower MW. Detecting MCU from

cell lysates using SEC has been successfully shown [16,19,22]. We found that Fe^{2+} treatment shifted the

MCU elution peaks to the earlier fractions of higher-order oligomers compared to mock control (Figure

1G, S1B; anti-MCU was validated in Figure S1C). As a positive control for this method, we treated cells

with H_2O_2 for 10 minutes and found the shift of the MCU elution peaks to the earlier fractions (Figure

S1D), consistent with previous studies also using SEC [19]. In contrast, the elution pattern of Miro1 was

largely unaltered by any of these treatments (Figure S1E). These data indicate that iron shifts the MCU

complexes to higher-order oligomers and enlarges the total number of these complexes.

To dissect the role of reactive oxygen species (ROS) [19] in contributing to the Fe^{2+}-triggered

MCU function as observed above, we treated HEK cells with an antioxidant, Vitamin C (VitC) [24], which

has no iron-chelating ability [25], or an iron chelator, deferiprone (DFP) [26], together with Fe^{2+}, and

measured MCU oligomerization and mitochondrial Ca^{2+} uptake. DFP, but not VitC, lowered the MCU

oligomer band intensity and mitochondrial Ca^{2+} import in the presence of Fe^{2+} (Figure S1F-G),

indicating that ROS might have a nominal effect on Fe^{2+}-mediated MCU regulation, at least in this

experimental setting.

We attempted to reconstitute the phenomenon of the Fe^{2+}-triggered increase of MCU

oligomerization in vitro. However, a fast speed of Fe^{2+} oxidation caused precipitation of purified MCU

protein. To circumvent this problem, we used an Fe^{2+} mimic, Co^{2+} ion [27]. The data of Co^{2+} might inform

us about properties of Fe^{2+}. We confirmed that Co^{2+} behaved similarly as Fe^{2+} in our functional assays in

HEK cells: Co^{2+} treatment increased MCU oligomerization detected by Native-PAGE (Figure 1H) and

enhanced the mitochondrial Ca^{2+}-uptake ability following thrombin application (Figure 1I), just like Fe^{2+}

(Figure 1C-E). We then conducted fluorescence-detection SEC using purified human MCU protein [15],

and found that Co^{2+} caused the formation of higher-order oligomers of MCU (Figure 1J). As a positive

control for this approach, we reproduced the elution peak of the dimer of tetramer of the MCU

holocomplex caused by Ca^{2+} addition in vitro (Figure 1J) [15]. These results demonstrate that the Fe^{2+}

mimic, Co^{2+}, increased MCU oligomerization in vitro. Collectively, Fe^{2+} and its mimic Co^{2+} promote the

formation of MCU higher-order oligomers and enhance the mitochondrial Ca^{2+}-uptake activity.

Fe^{2+} binds to the MCU complex to promote its oligomerization and channel activity

Our in vitro result in Figure 1J suggested that Co^{2+} bound to MCU to cause the formation of higher-

order oligomers. We further confirmed that Co^{2+} altered MCU oligomerization and thermostability in a

dose-dependent manner (Figure 1K).

We hypothesized that Fe^{2+} also bound to MCU to increase its oligomerization. To explore this

possibility, we immunoprecipitated (IP) endogenous MCU from HEK cells and detected the iron

concentrations in the IP samples. We found significantly more Fe^{2+} ions pulled down with MCU when

HEK cells were treated with Fe^{2+} (Figure 2A). We next examined whether MCU directly bound to Fe^{2+}

by an in vitro binding assay. To minimize protein precipitation caused by Fe^{2+} oxidation, we performed

an alternative to the thermal shift-SEC experiment with Co^{2+} shown in Figure 1K, which could take

hours to complete. We applied Fe^{2+} to the purified MCU complex (MCU, MICU1, MICU2, ERME) [15]

for about 5 minutes and after wash immediately measured iron levels in the complex by inductively

coupled plasma mass spectrometry (ICP-MS). We found that the MCU complex bound to Fe^{2+} in vitro

with a Kd of 449.3 µM (Figure 2B).

We next searched for amino acid residues in the matrix domain of MCU (PDB: 5KUE) predicted

to bind to Fe^{2+} using an in-silico program (http://bioinfo.cmu.edu.tw/MIB/) [28,29], and found 3 amino

acids: 74D, 148D, and 159H (Figure 2Ci). The latter 2 residues were also predicted to bind to Co^{2+}. We

mutated these 3 sites to Alanine (named "MCU-3A"). If MCU bound to Fe^{2+} via some of these sites,

MCU-3A should reduce their binding. Indeed, we detected significantly fewer Fe^{2+} in the

immunoprecipitate of Flag-tagged MCU-3A, as compared to MCU-WT, expressed from HEK cells

without endogenous MCU (*MCU-/-*) treated with Fe^{2+} (Figure 2Cii). We confirmed this result by ICP-

MS (Figure 2Ciii). Thus, MCU may bind to Fe^{2+} through some of these sites.

To determine whether blocking MCU's binding to Fe^{2+} was sufficient to eliminate the Fe^{2+}-

triggered oligomerization of MCU, we expressed MCU-WT or MCU-3A in *MCU-/-* HEK cells to

prevent the interference of endogenous MCU, treated these cells with Fe^{2+}, and ran IPed MCU in

Native-PAGE. As expected, MCU-3A abolished MCU's response to Fe^{2+} treatment: the MCU oligomer

band intensity was no longer increased (Figure 2D). We then live imaged mitochondrial Ca^{2+} dynamics,

as described in Figure 1, in these cells. We consistently observed a larger mitochondrial Ca^{2+} elevation

following thrombin stimulation in MCU-WT-transfected HEK cells treated with Fe^{2+} as compared to no

Fe^{2+} treatment, and MCU-3A blunted the peak increase (Figure 2E-F). Of note, MCU-3A-expressing

cells maintained normal functions of MCU (Figure 2D-H), suggesting no negative impact on cells by

these mutations. In addition, ROS similarly promoted the oligomerization of MCU-WT and MCU-3A

(Figure S1H), consistent with that the cysteine residues for oxidation [19] remain intact in both proteins.

Thus, ROS might have a negligible role in the difference between MCU-WT and 3A triggered by Fe^{2+}

seen in Figure 2D-H. Altogether, our results show that Fe^{2+} promotes MCU oligomerization and its

channel activity through binding to MCU.

Iron reduced NCLX and MCUb protein levels in a manner dependent on the MCU channel

activity

We next examined additional membrane proteins that may assist the MCU function using HEK cells. By

detecting total protein levels using Western blotting, we found a reduction in only MCUb and NCLX but

not the other 12 proteins examined including MCU monomers, when cells were treated with Fe^{2+} for 20-

22 hours (Figure S2), demonstrating selective protein responses without widespread toxicity at this time

point (Figure S1A). MCUb is an inhibitor of MCU [16], and NCLX is an IMM exchanger believed for

mitochondrial Ca^2 extrusion [13,30]. The reduction of both proteins could contribute to the phenotype of

mitochondrial Ca^{2+} overload upon Fe^{2+} treatment observed earlier (Figure 1C-D).

We explored the possibility that the reduction of NCLX and MCUb was secondary to the MCU

activity and resultant Ca^{2+} overload. We took advantage of the *MCU-3A*-expressing *MCU-/-* cells which

eliminated MCU's Fe^{2+}-binding and Fe^{2+}-elicited channel activity (Figure 2). By measuring the Ca^{2+}

extrusion rate we found that Fe^{2+} treatment slowed the efflux rate in *MCU-WT*-expressing *MCU-/-* cells,

which was prevented by MCU-3A (Figure 2G). This result suggests that the Fe^{2+}-triggered efflux delay

may depend on Ca^{2+} overload caused by the increased MCU activity. We then immunoblotted NCLX

and MCUb protein from these cells. We found again the reduction of NCLX and MCUb in *MCU-WT-*

expressing *MCU-/-* cells upon Fe^{2+} treatment and their protein levels were restored in *MCU-3A*

expressing cells (Figure 2H). Hence, these findings indicate that mitochondrial Ca^{2+} overload upon Fe^{2+}

elevation is the primary reason for the reduction of NCLX and MCUb levels.

PD postmortem brains and neurons mirror Fe^{2+}-treated HEK cells

Our data so far have shown that artificially increasing iron levels causes mitochondrial Ca^{2+}

accumulation by promoting the MCU activity. We examined this mechanism in pathophysiological PD

models without the need of adding exogenous iron to the media. We first investigated postmortem brains

of people with sporadic PD without a family history (Table S1). We homogenized the frontal cortex,

where iron elevation was observed in PD patients before measured by ICP-MS [31], and ran the brain

lysate in Native- or SDS-PAGE. Consistent with previous studies, MCU, NCLX, and MCUb were

detected in the human brain [9,32]. We found that the PD group (7 patients) clustered and separated from

the healthy control group (5), with higher intensity of the MCU oligomer bands in Native-PAGE and

lower intensity of both the NCLX and MCUb bands in SDS-PAGE (Figure 3A, S3A), similar to the

observations in HEK cells treated with Fe^{2+} (Figure 1, S2). This unique clustering was not observed in

the Alzheimer's disease (AD) group (5 patients) (Figure S3A). The total iron level in each brain fraction,

measured by ICP-MS, was positively correlated with the corresponding band intensity of MCU

oligomers in Native-PAGE (Figure 3B).

We next employed induced pluripotent stem cells (iPSCs) from one familial patient (PD-SNCA-

A53T) with the A53T mutation in *SNCA* (encodes α-synuclein) and its isogenic wild-type control

(isogenic-WT) [33-36]. We differentiated iPSCs to neurons expressing tyrosine hydroxylase (TH), the rate-

limiting enzyme for DA synthesis as previously described (Method) [33-36]. These patient-derived neurons

display increased expression of endogenous α-synuclein [34]. We stimulated these neurons with thrombin

and measured cytosolic and mitochondrial Ca^{2+} levels as in Figure 1. We found very similar Ca^{2+}

dynamic pattern in PD neurons as in HEK cells treated with Fe^{2+}: PD neurons suffered from more

mitochondrial Ca^{2+} accumulation after thrombin stimulation, as compared to isogenic control (Figure

3C). We next ran the neuronal lysates in Native- and SDS-PAGE. We again observed an upward shift of

the MCU oligomer bands and an increase in the total MCU band intensity in Native-PAGE indicating

the formation of higher-order oligomers, and a reduction of the intensity of NCLX and MCUb bands in

SDS-PAGE in PD neurons compared to control (Figure 3D). We also discovered a significant increase

in the mitochondrial Fe^{2+} levels detected by RPA and Mito-FerroGreen staining in PD neurons (Figure

3E). The phenotypes of mitochondrial Fe^{2+} and Ca^{2+} were observed in older neurons as well (day 40,

Figure S3B). Collectively, our results demonstrate a positive correlation of iron levels with MCU

oligomerization and its channel activity in PD patient-derived neurons and postmortem brains.

Iron functions upstream of calcium to mediate phenotypes of PD neurons

Our findings suggested the possibility that the elevation of iron levels in PD caused mitochondrial Ca^{2+}

overload by the binding of MCU with Fe^{2+} (Figures 1-3). To test this hypothesis, we reduced iron levels

in PD neurons described in Figure 3 with DFP and found that DFP significantly lowered mitochondrial

Ca^{2+} accumulation following thrombin stimulation (Figure 4A), demonstrating a causal relation between

iron elevation and Ca^{2+} overload in the mitochondria. These PD neurons are more vulnerable to

mitochondrial stressors. The treatment of the complex III inhibitor, Antimycin A, at 10 µM for 6 hours

caused acute neuronal cell death leading to the increase of terminal deoxynucleotidyl transferase dUTP

nick end labeling (TUNEL) signals in neurons derived from this PD patient, but not in isogenic control

neurons (Figure S3C) [34-36]. We found that pretreatment with DFP prevented cell death (Figure S3C). We

validated these findings in neurons differentiated from additional iPSC lines, including those from two

sporadic patients (PD-S1, PD-S2; no family history) and two healthy controls (HC-1, HC-2) [33-35], and an

isogenic pair containing a mutant line (Jax-SNCA-A53T) where *SNCA-A53T* was made in iPSCs of a

healthy donor and an isogenic control (Jax-Rev) by reverting *SNCA-A53T* back to wild type (Method).

The inclusion of multiple familial and sporadic lines should minimize line-to-line variation. We

consistently observed key features shown earlier in the PD-SNCA-A53T and isogenic-WT pair:

mitochondrial Fe^{2+} and stimulated-Ca^{2+} levels were elevated (Figure S3D-E), and DFP prevented

mitochondrial Ca^{2+} overload (Figure S3D) and stressor-induced cell death (Figure S3C) in these PD-

related lines. In order to cross-validate the neuroprotection of DFP in vivo without the need of adding a

stressor, we fed DFP to a fly model of PD, which expressed the pathogenic human α-synuclein protein

with the A53T mutation (α-syn-A53T) in DA neurons driven by *TH-GAL4* [34-36]. These flies exhibit PD-

relevant phenotypes such as age-dependent locomotor decline and DA neuron loss [34-36]. In the

protocerebral posterior lateral 1 (PPL1) cluster of the fly brain, the DA neuron number typically ranges

between 11-14, but this number is reduced in PD flies. We found that feeding PD flies with DFP

restored the DA neuron number in aged fly brains and improved the climbing ability (Figure 4B-C).

Collectively, our results show that iron functions upstream of calcium and mediates neurodegeneration

in multiple PD models.

We next utilized the Fe^{2+}-resistant MCU mutant, MCU-3A (Figure 2), to block extra Fe^{2+}-

binding in PD neurons. While expressing MCU-WT in PD neurons caused mitochondrial deformation

including shortening, fragmenting, and swelling, probably due to substantial iron-mediated

mitochondrial Ca^{2+} overload [37], MCU-3A completely prevented this phenotype (Figure 4D). As

mentioned earlier, ROS should similarly oxidize both MCU-WT and 3A proteins [19] (Figure S1H) and

impact those PD neurons. These results suggest that eliminating MCU's binding to Fe^{2+} renders

protection for mitochondria.

Together, our data from both HEK cells and PD models demonstrate a pathological mechanism

whereby iron accumulation inside the mitochondria causes inward Ca^{2+} overflow via a direct binding of

Fe^{2+} with MCU and the resultant augmentation of the MCU activity (Figure 4E).

Local OMM calcium dysregulation is sensed by Miro1 in Parkinson's models

We reasoned that the enlarged Ca^{2+} ion influx through MCU in PD models could alter local Ca^{2+}

dynamics and balance at the OMM (Figure 4E). This subdomain-scale Ca^{2+} change might be sensed by a

nearby OMM protein. Miro1 is a Ca^{2+}-binding protein on the OMM facing the cytosol, and in close

proximity to mitochondrial and ER Ca^{2+} channels [3,38-40]. Thus, Miro1 may directly encounter Ca^{2+} flux

going through mitochondria and sense even spatially confined Ca^{2+} concentration changes. To explore

this possibility, we swapped the cytosolic portion of Miro1 with the fluorescent Ca^{2+} sensor, GCaMP6f,

while leaving Miro1's C-terminal transmembrane and tail domain intact to allow proper OMM

localization [36], named GCaMP-Miro1-Anchor. A cytosolic expressing miRFP670 gene was constructed

in the same vector for background control (full insertion: miRFP670nano_P2A_GCaMP6f-Miro1-

Anchor). This new Ca^{2+} sensor should detect Ca^{2+} ions at the subdomains of the OMM where Miro1 is

located. We confirmed that GCaMP-Miro1-Anchor was colocalized with Miro1 in human neurons

(Figure 4F). The intensity of GCaMP-Miro1-Anchor was increased showing elevated local Ca^{2+}

concentrations in live PD neurons compared to isogenic control, while the intensity of a Ca^{2+} sensor

located to the cytosolic face of the ER membrane (miRFP670nano_P2A_ER-GCaMP6f) was unchanged

(Figure 4G). This increase in the GCaMP-Miro1-Anchor intensity was not caused by increased

mitochondrial volume because the fluorescent intensity of MitoTracker Green, a membrane-potential-

independent dye, was comparable between control and PD neurons (Figure 3E; Figure Legend).

Importantly, inhibiting the MCU activity by RU360 [41] reduced OMM Ca^{2+} elevation detected by

GCaMP-Miro1-Anchor in PD neurons (Figure S3F). This data suggests that enhanced mitochondrial

Ca^{2+} influx could raise spatially restricted Ca^{2+} concentrations at the OMM and that Miro1 may serve as

an ideal sensor for this change (Figure 4E).

We next sought a quantitative measurement of Miro1's Ca^{2+} sensitivity at the OMM, which

could help us develop specific molecular markers to distinguish PD patients with OMM Ca^{2+} elevation.

We had previously found a Miro1 phenotype in PD models and patients' cells. We explored whether this

Miro1 phenotype was a result of Miro1's Ca^{2+} sensing. In a healthy cell, upon mitochondrial

depolarization Miro1 is quickly removed from the OMM and degraded in proteasomes, primed by PD-

related proteins including PINK1, Parkin, and LRRK2, allowing the subsequent mitophagy [33-35,42,43]. We

had shown that Miro1 degradation upon mitochondrial depolarization was delayed in fibroblasts or

neurons derived not only from familial PD patients with mutations in *Parkin* or *LRRK2* but also from

sporadic patients without known mutations, consequently slowing mitophagy and increasing neuronal

sensitivity to stressors [33-35]. The mechanism underlying this unifying Miro1 phenotype in both familial

and sporadic PD patients remained undefined. To explore whether this phenotype resulted from Miro1's

Ca^{2+} binding, we made GFP-tagged human Miro1 protein in both the wild-type (Miro1-WT) form and in

a mutant form where two point mutations were introduced in the two EF-hands of Miro1 (Miro1-KK) to

block Ca^{2+} binding [44]. We expressed GFP-tagged Miro1 (WT or KK) and Mito-dsRed that labeled

mitochondria in iPSC-derived neurons from the PD patient and isogenic control, described earlier. We

had previously discovered that following acute Antimycin A treatment that depolarized the membrane

potential and triggered mitophagy, Miro1 and mitochondria were sequentially degraded in wild-type

neurons [33-35]. We observed the same mitochondrial events in isogenic control axons transfected with

GFP-Miro1-WT by live imaging (Figure 5A-C). In contrast, in PD neuron axons transfected with GFP-

Miro1-WT, the degradation rates of both Miro1 and damaged mitochondria upon Antimycin A treatment

were slowed (Figure 5A-C), consistent with our previous studies [34,35]. Notably, GFP-Miro1-KK

significantly rescued these phenotypes in PD axons: it expedited the degradation rates to the control

level (Figure 5A-C). We validated the result of Miro1 degradation using an enzyme-linked

immunosorbent assay (ELISA) of GFP: at 25-minute post Antimycin A treatment, Miro1-WT was not

degraded in PD neurons and this phenotype was rescued by Miro1-KK (Figure 5D). These data suggest

that delayed Miro1 and damaged mitochondrial degradation rely on Miro1's Ca^{2+} binding in PD neurons.

To confirm the Miro-Ca^{2+} relation in vivo, we generated transgenic flies carrying T7-tagged fly

Miro (DMiro)-WT or DMiro-KK. DMiro is an ortholog of human Miro1 with high sequence similarity.

Both DMiro-WT and DMiro-KK were expressed at comparable levels when the transgenes were driven

by the ubiquitous driver *Actin-GAL4* (Figure S3G). We next crossed these transgenic flies to a fly PD

model described earlier that expressed human α-syn-A53T in DA neurons driven by *TH-GAL4*. We

found that DMiro-KK significantly rescued the phenotypes of age-dependent DA neuron loss and

locomotor decline, compared to DMiro-WT (Figure 5E-F). This result shows that blocking DMiro's

Ca^{2+}-binding is neuroprotective in PD flies. Altogether, we have provided evidence that Miro1 or DMiro

senses Ca^{2+} to mediate several phenotypes in human neuron and fly models of PD.

A high-content Miro1 screening assay identifies a network of Ca^{2+}-related drug hits for PD

To further support that Miro1 was downstream of Ca^{2+} dysregulation in PD, we performed high

throughput (HTP) drug screens in fibroblasts from a sporadic PD patient using the Miro1 phenotype

(delayed degradation upon depolarization) as a readout. If Miro1 sensed local Ca^{2+} elevation to become

more stable on damaged mitochondria (Figure 4G, 5A-C), we expected to see Ca^{2+}-related drug hits to

show up to promote Miro1 degradation. To this end, we established a sensitive immunocytochemistry

(ICC)-based assay that was suitable for HTP screening (Figure 6A, S4, S5, S6A, Table S2, Method). We

performed the primary screens at the Stanford High-Throughput Bioscience Center (HTBC) using 3

drug libraries containing many compounds that have well-defined roles and targets and show efficacy to

treat certain human diseases (Figure S4). Carbonyl cyanide p-trifluoro-methoxyphenyl hydrazone

(FCCP) was chosen as an uncoupler because it had been used successfully in ICC on Miro1 [45]. We

screened the NIH Clinical collection library (377 unique compounds) at a defined concentration of 10

µM with 4 biological repeats for each drug, the Biomol FDA library (175 unique compounds) with two

five-fold doses (1.25, 2.5, 5, 10, 20 µM), and the Sigma LOPAC library (1269 unique compounds) at a

defined concentration of 20 µM. Drug hits from primary screens were then retested in our own

laboratory using fresh compounds at the highest screening concentration with at least 4 biological

replicates and our confocal microscope (Figure S5). In total, we discovered 15 compounds in two

independent experimental settings that reduced Miro1 protein levels following mitochondrial

depolarization in PD fibroblasts (Figure S5C, Table S2 with all drug properties).

Next, we performed a pathway analysis using a knowledge graph-based tool to reveal the

potential cellular pathways connecting Miro1 to each hit compound. Strikingly, we discovered

intracellular Ca^{2+} as a primary shared factor in the hit drug-Miro1 network (Figure 6B, Table S3). Two

drugs, Benidipine and Tranilast, could directly inhibit plasma membrane Ca^{2+} channels. Benidipine is a

blocker of voltage-gated Ca^{2+} channels (L-, N-, T-type), and Tranilast has been proposed to inhibit

ligand-gated Ca^{2+} channels (TRPV2) [46]. Therefore, our results confirm the importance of the Ca^{2+}

pathway for mediating the Miro1 phenotype in PD.

Benidipine rescues Parkinson's-relevant phenotypes in multiple models of PD

We further validated the top hit from our screens, Benidipine, a pan-Ca^{2+}-channel blocker. Using the

same ICC method as in Figure S5, we found that Benidipine reduced Miro1 in a dose-dependent manner

upon depolarization in PD fibroblasts (Figure S6B). To exclude the possibility of any artifacts caused by

our ICC method, we verified our results by Western blotting and observed the same rescue effect:

Benidipine promoted Miro1 degradation after mitochondrial depolarization in PD fibroblasts (Figure

S6C; Figure Legend). We confirmed that Benidipine did not affect *Miro1* messenger RNA (mRNA)

expression detected by reverse transcription quantitative real-time PCR (RT-qPCR) under basal and

depolarized conditions in PD cells (Figure S6D). Neither did Benidipine alter the basal ATP levels

(Figure S6E), nor the mitochondrial membrane potential measured by TMRM staining (Figure S6F).

We next tested Benidipine in the human neuron model of PD described in Figure 3. We

confirmed that Benidipine reduced local Ca^{2+} elevation at the OMM detected by GCaMP-Miro1-Anchor

(Figure S3F) and promoted Miro1 degradation following Antimycin A treatment in PD neurons (Figure

S6G; similar to DFP). This result corroborates that delayed Miro1 removal upon depolarization reflects

OMM Ca^{2+} elevation. Notably, treating these PD neurons with Benidipine at 10 μM for 30 hours

mitigated stressor-induced neuron death: the loss of TH immunostaining and increase of TUNEL

staining after Antimycin A treatment [34,36] were significantly rescued (Figure 7A-B).

In vivo, we fed Benidipine to the fly model of PD without adding any mitochondrial stressors.

Importantly, feeding these flies with 2.5 μM Benidipine from adulthood prevented DA neuron loss in

aged flies (Figure 7C) and improved their locomotor ability (Figure 7D). Taken together, our results

show that either lowering local OMM Ca^{2+} levels by Benidipine or blocking Miro1's Ca^{2+} binding

eliminates the Miro1 defect of slower degradation upon depolarization and additional phenotypes in PD

models (Figure 5-7, S6).

A Miro1 blood test reflects PD status and responds to drug treatment

A challenge in current PD research is the lack of reliable and convenient molecular markers for patient

stratification. The subset of PD patients with Ca^{2+} elevation at the OMM should be sensed by Miro1, and

once Miro1 bound to Ca^{2+} it could be quantitatively measured by a proxy–the slower Miro1 response to

a mitochondrial uncoupler (Figure 5-7, S6). We next explored the potential of this Miro1 proxy as a PD

marker in a peripheral tissue. Although fibroblasts can be readily obtained by a skin biopsy, a blood test

remains the most convenient method. We set out to develop a Miro1 blood test using peripheral blood

mononuclear cells (PBMCs) which can be acquired from a blood draw during routine clinic visit. We

cultured PBMCs from a healthy donor from the Stanford Blood Center (SBC, Table S4) and depolarized

the mitochondrial membrane potential using two different methods: Antimycin A plus Oligomycin [47], or

carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP). We found that both depolarizing approaches

caused the degradation of Miro1 and additional mitochondrial markers in a time-dependent manner,

detected by Western blotting (Figure S7A-B), consistent with other cell types [33,35,48]. These results show

that Miro1 can respond to mitochondrial depolarization in healthy PBMCs, allowing us to further utilize

these cells to develop a Miro1 assay for PD.

To enable high-content screening, we applied an ELISA of Miro1 (Figure S7C-D) to PBMCs

from the same donor with a 6-hour CCCP treatment. We saw a similar Miro1 response to CCCP using

ELISA (SBC, Table S4) as using Western blotting. The reproducibility of the result using ELISA led us

to choose this method and 6-hour CCCP treatment to screen a total of 80 healthy controls and 107 PD

patients (Table S4). Miro1 Ratio (Miro1 protein value with CCCP divided by that with DMSO from the

same person) was significantly higher in PD patients compared to healthy controls (Figure 7E, Table S4),

suggesting that PBMCs from more PD patients may have Ca^{2+} elevation at the OMM and consequently

fail to rapidly remove Miro1 from damaged mitochondria. Hence, Miro1 Ratio by our method may be

used as an indicator of OMM Ca^{2+} changes.

To determine whether our method could be used to classify an individual into a PD or healthy

group, we employed machine learning approaches using our dataset. We trained a logistic regression

model to assess the impact of Miro1 Ratio on PD diagnosis, solely on its own or combined with

additional demographic and clinical parameters (Method). Unified Parkinson's Disease Rating Scale

(UPDRS) is a tool to measure motor and non-motor symptoms of PD which may reflect disease severity

and progression. Using UPDRS, our model yielded an accuracy (an individual was correctly classified

as with PD or healthy) of 81.2% (p<0.000001; area under the Receiver Operator Curve (ROC)–

AUC=0.822), and using Miro1 Ratio, the accuracy was 67.6% (p=0.03; AUC=0.677). Notably, if both

Miro1 Ratio and UPDRS were considered, our model generated an improved accuracy of 87.8% (p=0.02;

AUC=0.878; p= 5.736e-09 compared to UPDRS alone, paired T test on bootstrapped samples; Method),

without the interference of age or sex (Method, Figure S7E-F). Therefore, our results suggest that the

molecular (Miro1 Ratio–OMM Ca^{2+} elevation) and symptomatic (UPDRS) evaluations may reveal

independent information, and that combining both tests could more accurately categorize individuals

with PD and measure their responses to experimental therapies.

To probe the potential utilization of this Miro1 assay in future clinical trials for stratifying

patients with OMM Ca^{2+} dysregulation and monitoring drug efficacy, we treated PBMCs from 4 PD

patients (Table S4) with either of the two compounds known to reduce Miro1, Benidipine (Figure 6-7,

S4-6) and Miro1 Reducer 3 (MR3) [35,36]. Miro1 protein levels upon CCCP treatment were lowered by

each compound in all 4 patients (Figure 7F), showing that the Miro1 marker in PBMCs can respond to

drug treatment. Collectively, our results suggest that Miro1 protein in blood cells may be used to aid in

diagnosis and drug development.

Rare variants in T-type Ca^{2+} channels are associated with PD status

After dissecting the functional impairment of this iron-calcium-Miro axis in PD, we explored its genetic

contribution to PD. Earlier, we showed that chelating iron, blocking MCU's binding to Fe^{2+}, blocking

Miro's binding to Ca^{2+}, or preventing Ca^{2+} entry into the cell all alleviated Parkinson's related

phenotypes (Figures 4-7). We evaluated the genes encoding the protein targets of these approaches,

which are spatially distinct and localized to three subcellular locations: (1) IMM Ca^{2+} channels and

transporters (targeted by Fe^{2+}), (2) the Ca^{2+}-binding protein Miro on the OMM, and (3) plasma

membrane Ca^{2+} channels (targeted by Benidipine and Tranilast) (Figure 7G, Table S5). By analyzing

common variants within or near any of the investigated genes in GWAS reported in [49], we did not

observe significant association with PD clinical status. We next employed the whole-genome

sequencing (WGS) data from the Accelerating Medicines Partnership–Parkinson's Disease (AMP–PD)

(1,168 control; 2,241 PD), and assessed rare non-synonymous and damaging variants using burden

based and SKATO methods. We discovered an enrichment of rare variants in selective T-type Ca^{2+}-

channel subtypes (*Cav3.2*, *3.3*) (cell surface), *Miro2* (OMM), or *NCLX* (IMM) associated with PD status

with a nominal P value (Table S5). Notably, a SKATO Test on all variants of T-type or L-type Ca^{2+}-

channel subtypes showed a significant association with PD status of T-type channels, which survived

multiple comparison correction, but not of L-type channels (Table S5). Together, our analysis unravels

potential genetic predisposition of T-type Ca^{2+} channels to PD.

Inhibiting T-type Ca^{2+} channels promotes Miro1 degradation

Our discovery of the selective accumulation of rare variants in T-type Ca^{2+} channels associated with PD

diagnosis (Table S5) prompted us to functionally validate this finding. We employed a mini drug screen

using the same screening ICC assay described earlier (Figure S4-5) by which we discovered the non-

selective pan-Ca^{2+}-channel blocker, Benidipine. This time, we examined two L-type and three T-type

Ca^{2+}-channel blockers. Intriguingly, we again found a striking selection of T-type versus L-type

channels, in the connection with Miro1 in PD fibroblasts (Figure S7G): only T-type blockers promoted

Miro1 degradation following depolarization, just like Benidipine. Importantly, two of these blockers,

MK-8998 [50, 51] (FDA approved) and ML-218 [52], were already shown in vivo safety and efficacy for

treating PD-associated symptoms in humans or preclinical rodent models. The third blocker, ABT-639

(FDA approved), was trialed in humans for treating pain [53]. By contrast, L-type blockers, including

Isradipine that recently failed a phase III trial for PD [54], did not affect the Miro1 phenotype (Figure S7G).

To genetically inhibit T-type Ca^{2+} channels, we knocked down by siRNA the T-type subtype *Cav3.2*

which showed strongest association with PD diagnosis in the human genetic analysis (Table S5).

Consistently, genetically reducing *Cav3.2* promoted Miro1 degradation upon depolarization in PD

neurons (Figure S7H). These studies implicate an immediate benefit of coupling T-type channel

inhibitor drugs and the Miro1 marker for clinical trials.

Discussion

In this work, we have established a mitochondrial inside-out pathway of iron-calcium-Miro

dysregulation in our PD models (Figure 7G). Elevation of iron concentrations may cause mitochondrial

Ca^{2+} overload by promoting the MCU activity and reducing NCLX levels (Figures 1-3, S2). It is

possible that the enhanced Ca^{2+} influx through the mitochondrial membranes disrupts Ca^{2+} homeostasis

at the OMM (Figure 4G, S3F), which is sensed by the nearby OMM resident, Miro1 (Figure 4E-G, 5-7).

We have shown multiple ways to target this axis. Chelating iron, reducing Ca^{2+} entry into the cell, or

blocking Miro's binding to Ca^{2+} is each neuroprotective (Figure 4, 5, 7). Hence, this ionic axis may be

important for PD pathogenesis and can be leveraged for better detecting and treating the disease.

Although the precise course of PD pathogenesis remains unclear, emerging evidence has demonstrated

its complex and multifactorial nature. Interactions among multiple organs and cell types, systemic

immune activation, and environmental triggers act uniquely in different individuals. Although GWAS

and segregation studies have unveiled many PD risk or causal loci, rare genetic variants may be

particularly important for conferring individual heterogeneity in disease onset and etiology. However,

rare variants associated with PD risk are difficult to discover given the limitations in sample collection

and the unmet need for exome-sequencing as opposed to sparse genotyping on microarrays. In an

individual prone to PD, there might be already mitochondrial malfunction causing mitochondrial

proteins to release labile iron [55], or impairments in systemic iron uptake and circulation causing iron

accumulation at the cellular level. Any predisposed genetic perturbations in this iron-calcium-Miro axis,

which could be due to different rare variants in each person, would further exacerbate its dysfunction.

With or without another trigger such as a virus infection or environmental insult, the tipping point for

disease onset might be reached. Thus, combining complementary genetic and functional studies may

help us better understand the destructive paths leading to the disease and identify network hubs for

therapeutic targeting.

The Miro1 marker could help address the current unmet needs in PD clinical care. There is a high rate of

misdiagnosis of PD at clinic because there is no definitive molecular marker to confirm the disease [56].

The lack of disease-modifying therapies is partially due to no reliable pharmacodynamic biomarkers in

clinical trials [35,57]. Our Miro1 blood test has a potential to serve both as a patient stratification tool and a

pharmacodynamic marker for drug discovery. The exciting link between T-type channel blockers and

Miro1 in PD suggests this Miro1 blood test may be immediately used in clinical trials for drugs already

at the advanced stage, such as MK-8998 [50,51] (trial ID: NCT05642442). We have also provided evidence

that patients' PBMCs responded to drugs that reduce Miro1. This result opens a new door to examining

personalized drug efficacy and dosing by culturing patients' own cells before administrating the drug to

the patient, thus providing multiple layers of stratification to improve success rates of clinical trials [58].

Further studies with independent cohorts are needed to confirm the promise of this test for PD.

Alpha-synuclein aggregation is a pathological hallmark of PD. Emerging evidence has shown that alpha-

synuclein has profound functions at the mitochondria or to impact mitochondria, including

mitochondrial oxidative phosphorylation, dynamics, mitophagy, Ca^{2+}, and iron homeostasis [8,34,59-64].

Iron accumulation, calcium mishandling, and mitophagy impairment have been individually observed in

PD neurons [3-11,33-35]. However, the relation between these cellular phenotypes in PD has remained

obscure. Now we have provided a mechanistic axis connecting them. Fe^{2+} elicits mitochondrial Ca^{2+}

overload through acting on IMM Ca^{2+} channels and transporters. The relatively low affinity of MCU to

Fe^{2+} in vitro (Figure 2B) is consistent with its low affinity to other divalent metals such as Mg^{2+} [30]. This

data suggests that MCU may not be predominantly occupied by these divalent ions under physiological

conditions [30]. By contrast, under pathological conditions with elevated Fe^{2+}, MCU binds to more Fe^{2+}

and cellular detriment ensues. This finding agrees with prior evidence showing that inhibiting MCU

prevents Fe^{2+}-induced cell death [37,65] and Parkinsonian neurodegeneration [2,9]. Further investigations are

needed to dissect how Fe^{2+} regulates MCUb and NCLX levels. Our results (Figure 2G-H) have

suggested that Fe^{2+}-triggered Ca^{2+} efflux delay and the reduction of NCLX and MCUb depend on

mitochondrial Ca^{2+} overload. Thus, it is possible that NCLX and MCUb are targeted by Ca^{2+}-activated

mitochondrial proteases. A recent study has shown that NCLX protein levels are lowered in another

mitochondrial iron-dependent disease, Friedreich's Ataxia (FA), through a possible mechanism

dependent on calpain cleavage [66]. Intriguingly, iron chelators have displayed promising therapeutic

benefits in preclinical models of both PD [67] and FA [68], and is currently in multiple clinical trials for

treating symptoms of PD patients [67], indicating these two diseases may share mechanisms. It's important

to note there might be disadvantages of metal chelators in humans, as a recent phase-II trial of DFP for

PD failed [69]. Our data also suggests Miro1 may link Ca^{2+} and mitophagy phenotypes in PD. In the same

PD neuron models used in this paper, we have previously characterized a delayed mitophagy phenotype

downstream of Miro1 retention on damaged mitochondria [33-35]. Our paper now shows this Miro1

phenotype is dependent on OMM Ca^{2+} elevation because of iron accumulation. In addition to mitophagy

[33,42,70], Miro may have additional functions in regulating mitochondrial quality control, including the

biogenesis of mitochondrial derived vesicles (MDVs) [71] and transcellular mitochondrial transfer [72-76].

When Miro senses spatially restricted Ca^{2+} elevation at the OMM in PD neurons, it may affect multiple

Miro-mediated biological processes. We have previously observed an upregulation of Miro1 protein in

postmortem brains of sporadic PD patients [34]. More studies are needed to unravel the precise roles of

Miro in PD pathogenesis and how Ca^{2+} regulates these roles.

Dopaminergic neurons in the substantia nigra have elaborate axonal branches, intense neuronal activities,

and exposure to metals and neurotoxins, and consequently their mitochondria exhibit enhanced

oxidative stress [6,10,77,78]. We have found that in iPSC-derived neurons with *SNCA* mutations

mitochondrial stressors cause more cell death for TH-positive than TH-negative neurons [34]. These

results suggest that mitochondria in dopaminergic neurons from PD patients may be more susceptible to

mitochondrial iron and calcium overload due to the existing stress in their microenvironment. In contrast,

mitochondria in neurons at the other part of the brain or in non-neuronal cells may experience less stress.

We and others have found that some PD fibroblasts show minor phenotypes at baseline (Figure S6E-F)

[33-35,79].

Limitation of the study:

It's important to note that flies do not contain an alpha-synuclein homolog, and our results may not

account for the physiological roles of human alpha-synuclein. The sample size for the postmortem brain

study is relatively small and a larger cohort of PD and PD-related disorders is needed to confirm the

prevalence and specificity of the phenotype. Re-evaluation of genetic variants in this pathway is needed

in additional larger cohorts to verify their link to PD. Although MCU oligomerization could be enhanced

with a micromolar range of Fe^{2+} added in the media (Figure 1F), the pathophysiological level of free

Fe^{2+} is difficult to determine in vivo.

Acknowledgements

We thank M.F. Tsai for *MCU* KO cells, L. Pallanck for flies, A. Y. Ting for constructs, A. E. Medlock,

O. Ross, and S. Han for discussion, P. Katiyar, Z. T. Cook, D. M. Conradson, and J. Panji for technical

support, Wang lab members for discussion, Stanford SIGMA Facility for ICP-MS (support from

Stanford Doerr School of Sustainability and Stanford Nano Shared Facilities/Stanford Nanofabrication

Facility under National Science Foundation award ECCS-2026822), Stanford HTBC (RRID:

SCR_017794) and D. E. Solow-Cordero for assistance with drug screens, and SBC, Stanford

Alzheimer's Disease Research Center (ADRC), Department of Pathology at University of California

Los Angeles (UCLA), R. N. Alcalay, Columbia University, Banner Sun Health Research Institute Brain

and Body Donation Program, and Michael J. Fox Foundation (MJFF) for providing human tissues and

cells. This work was supported by NINDS (RO1NS089583, RO1NS128040; X.W.), NIGMS

(RO1GM143258; X.W.), NIA (RO1AG060747; M.D.G.), Stanford SPARK and Spectrum SPADA

(X.W.), Sanofi iAward (X.W.), Warren Alpert Foundation (X.W.), MJFF (021146; X.W.), Harrington

Discovery Institute (X.W.), Belgian American Education Foundation (R.V.), Glenn Foundation

Postdoctoral Fellowship (L.L.), and European Union's Horizon 2020 research and innovation program

under Marie Skłodowska-Curie (890650; Y.LG.). Brain and Body Donation Program was supported by

the NINDS (U24 NS072026), NIA (P30 AG19610), Arizona Department of Health Services (211002),

Arizona Biomedical Research Commission (4001, 0011, 05-901 and 1001), and MJFF. Graphic Abstract

was created with BioRender.com.

564 Author contributions

565 V.B., A.S.D., R.V., C.-H.H., S.C., L.L., and C.S.K. performed human cell experiments. R.V. did drug

566 screens. L.L. conducted fly work. C.M.M. purified MCU and did FSEC with assistance from L.F. C.S.K.

567 did molecular cloning. Y.L.-G. and M.D.G. analyzed human genetic data. P.N. analyzed PBMC data.

568 X.W. conceived and supervised the project. V.B., R.V., and X.W. wrote the paper with assistance from

569 all authors.

571 Declaration of interests

The authors declare the following competing interests: X.W. is a co-founder and shareholder of AcureX Therapeutics, and a shareholder of Mitokinin Inc. V.B., L.L., C.-H.H., and R.V. are shareholders of AcureX Therapeutics. P.N. was employed by Vroom Inc. Patents based on this study were filed by Stanford University with X.W., R.V., V.B., L.L., C.-H.H. as inventors. The remaining authors declare no competing interests.

Inclusion and diversity

We support inclusive, diverse, and equitable conduct of research.

STAR METHODS

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to the lead contact Xinnan Wang (xinnanw@stanford.edu).

Materials availability

All reagents generated in this study are available from the lead contact.

Experimental model and study participant details

Human cells and tissues

No human subjects were used in this study. The iPSC work was approved by Stanford Stem Cell Oversight Committee. iPSCs or fibroblasts were purchased under a material transfer agreement (MTA)

from NINDS Human Cell and Data Repository or Jackson Laboratory. PBMCs were obtained under an

MTA from Columbia University and MJFF, and from SBC and Stanford ADRC. Postmortem brain

tissues were obtained from Stanford ADRC, Department of Pathology at UCLA, and Banner Sun Health

Research Institute. Columbia University, Stanford University, UCLA, and Banner Sun Health Research

Institute Institutional Review Board approved study protocols, which ensured consent from donors and

explained the conditions for donating materials for research. Demographic and clinical details of each

subject are in Supplementary Tables. HEK293T cells were obtained from ATCC.

Fly Model

The following fly stocks were used: *Actin-GAL4* (Bloomington Drosophila Stock Center–BDSC), *TH-*

GAL4 (BDSC), *UAS-SNCA^{A53T}* [80], and *UAS-T7-DMiro-WT* [81]. *UAS-T7-DMiro-KK* were generated by

injecting pUAST-T7-DMiro-KK into flies by Bestgene Inc (Chino Hills, CA). All fly stocks and

experiments were kept at 25°C with a 12:12 hrs light:dark cycle and constant humidity (65%) on

standard sugar-yeast-agar (SYA) medium (15 g/l agar, 50 g/l sugar, 100 g/l autolyzed yeast, 6 g/l

nipagin, and 3 ml/l propionic acid) [82]. Flies were raised at standard density in 200 ml bottles unless

otherwise stated. All fly lines were backcrossed 6 generations into a *w^{1118}* background to ensure a

homogeneous genetic background. The ages of flies for each experiment were stated in figures.

Experiments were carried out on mated females [83].

Method details

Constructs

pcDNA3.1-MCU-Flag was purchased from Genescript (MCU_OHu20123D_pcDNA3.1+/C-(K)-DYK).

pcDNA3.1-MCU-3A-Flag was generated by Synbio Technologies by making 3 point mutations (74D,

148D, 159H to A) of pcDNA3.1-MCU-Flag. pRK5-GFP-hMiro1-WT or KK was custom-made by

Synbio Technologies. pDsRed2-Mito was from Clontech. pUAST-T7-DMiro-KK was generated by

introducing E234K, E354K to pUAST-T7-DMiro [81]. Myc-hMiro1-WT and was described in [44].

pLX304-miRFP670nano_P2A_GCaMP6f-Miro1-Anchor   or   pLX304-miRFP670nano_P2A_ER-

GCaMP6f (cytosol) was made by cutting pLX304-miRFP670nano_P2A_EGFP (a gift from Alice Y.

Ting) with BsiW1 and Nhe1, amplifying each gene fragment by PCR, and then ligating these with the

Gibson assembly (HiFi master mix, NEB) method. pcDNA3.1-ER-GCaMP6f (cytosol) was a gift from

Chris Richards (Addgene #182548). miRFP670nano was a gift from Vladislav Verkhusha (Addgene

#127443). All vectors were confirmed by sequencing.

Cell Culture

Human dermal primary fibroblasts isolated from a PD patient (ND39528) and a healthy control

(ND36091) were cultured in DMEM (11995-065, Thermo Fisher) supplemented with 10% fetal bovine

serum (900-108, heat-inactivated, Gemini Bio-Products), 1×Anti Anti (15240096, Thermo Fisher), and

1×GlutaMax (35050061, Thermo Fisher) maintained in a 37°C, 5% CO_2 incubator with humidified

atmosphere. HEK293T cells were cultured in DMEM (11995-065, Thermo Fisher) supplemented with

10% fetal bovine serum (900-108, heat-inactivated, Gemini Bio-Products), 1% Pen-Strep (15140-122,

Gibco), and maintained in a 37°C, 5% CO_2 incubator with humidified atmosphere. Different

concentrations of Fe^{2+} (Ammonium Iron(II) Sulfate Hexahydrate, 203505, Sigma-Aldrich) were added

to the media for 20-24 hours. The MTT assay (ab211091, Abcam) was used according to the manual.

Cells were treated with 10 µM Benidipine (B6813, Sigma-Aldrich), 40 µM CCCP (C2759, Sigma-

Aldrich), 100 µM VitC (A5960, Sigma-Aldrich), or 100 µM DFP (HY-B0568, MedChemExpress).

iPSC Culture, Differentiation, and Transfection

All iPSC lines in this study are summarized below and have been fully characterized by our previous studies [33,34], NINDS Human Cell and Data Repository, Applied StemCell, or Jackson Laboratory.

iPSC line	Source	Etiology	PD mutation	Age	Sex
PD-SNCA-A53T	NINDS (ND50050)	Genetic	*SNCA (A53T)*	51	Female
Isogenic-WT	NINDS (ND50085)	Isogenic control for *PD-SNCA-A53T*	*Wild-type*	51	Female
PD-S1	NINDS (ND39896)	Sporadic	*Unknown*	77	Male
PD-S2	Applied StemCell (ASE-9028)	Sporadic	*Unknown*	82	Female
HC-1	Hsieh et al., 2016	Healthy Control	*Wild-type*	62	Male
HC-2	Stanford Stem Cell Core, Shaltouki et al., 2018	Healthy Control	*Wild-type*	42	Female
Jax-SNCA-A53T	Jackson Laboratory	Mutation in a healthy background	*SNCA (A53T)*	N/A	Male
Jax-Rev	Jackson Laboratory	Isogenic control for *Jax-SNCA-A53T*	*Wild-type*	N/A	Male

Clinical and additional data are in https://stemcells.nindsgenetics.org/
https://www.jax.org/jax-mice-and-services/ipsc/cells-collection

iPSCs were derived to midbrain DA neurons as previously described with minor modifications [33,84-87]. Briefly, neurons were generated using an adaptation of the dual-smad inhibition method. Neural induction was started when iPSCs reached 50-85% confluency, with the use of dual smad inhibitors, dorsormorphin (P5499, Sigma-Aldrich) and TGFβ inhibitor SB431542 (1614, Tocris), and the addition of GSK3β inhibitor CHIR99021 (04-0004, Stemgent) and smoothened agonist SAG (566661, CalBioChem). To get single cells, 12 days after neural induction, half of the medium was replaced with N2 medium with 20 ng ml^{-1} BDNF (450-02, Peprotech), 200 μM ascorbic acid (A5960, Sigma-Aldrich), 500 nM SAG, and 100 ng ml^{-1} FGF8a (4745-F8-050, R&D Systems). On day 16-17, neurons were split and transferred onto Matrigel (354277, Corning)-coated or poly-ornithine and laminin-coated glass

coverslips in a 24-well plate. On day 19-20, medium was switched to N2 medium supplemented with 20

ng ml^{-1} BDNF, 200 μM ascorbic acid, 20 ng ml^{-1} GDNF (450-10, Peprotech), 1 ng ml^{-1} TGFβ3 (AF-

100-36E, Peprotech), and 500 μM dibutyryl-cAMP (D0627, Sigma-Aldrich) for maturation of DA

neurons. Neurons were used at day 21-26 after neuronal induction in most of the experiments, when

about 80-90% of total cells expressed the neuronal marker TUJ-1, and 14-17% of total cells expressed

TH and markers consistent with ventral midbrain neuronal subtypes [34,88].

For transfection, on day 19-20 after neural induction, culture medium was replaced with Opti-MEM

(Gibco) prior to transfection. For 24-well plates, 0.5-1 μg DNA or 1 μl Lipofectamine 2000 (11668-030,

Invitrogen) was diluted in Opti-MEM at room temperature (22°C) to a final volume of 50 μl in two

separate tubes, and then contents of the two tubes were gently mixed, incubated for 20 minutes at room

temperature, and subsequently added onto neurons (75,000-120,000 per well). For 6-well plates, 1.5 μg

DNA or 6 μl Lipofectamine 2000 (11668-030, Invitrogen) was diluted in Opti-MEM to a final volume

of 250 μl in each tube, and mixed, incubated, and applied to neurons (700,000-1,500,000 per well).

After 6 hours, Opti-MEM containing DNA-Lipofectamine complexes was replaced with regular N2

medium with supplements. For siRNA, NT siRNA (D-001910-10, Horizon Discovery), Cav3.2 siRNA

(EQ-006128-00-0002, Horizon Discovery), and delivery media were used according to the manual.

After transfection for 2-3 days, neurons were used. Neurons from 6-well plates were lysed in 400 μl

lysis buffer (300 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 0.2 mM PMSF, Protease Inhibitor

Cocktail, 5 mM EDTA) and used in the Miro1 ELISA kit (EKL54911, Biomatik) following the manual,

or in 300 μl Extraction Buffer supplemented with 0.2 mM PMSF and Protease Inhibitor Cocktail and

used in the GFP ELISA kit (ab229403, Abcam) following the manual.

Neurons were treated with 50-100 µM DFP (HY-B0568, MedChemExpress), 10 µM Benidipine (B6813,

Sigma-Aldrich), 500 nM RU360 (557440, Sigma-Aldrich), 10 µM Antimycin A (A8674, Sigma-

Aldrich), or 100 µM Antimycin A during live imaging.

Primary Drug Screens

Overall scheme: Fibroblasts were plated and cultured for 24 hours. Compound libraries were then

applied for 10 hours. Next, 20 µM FCCP was added for another 14 hours. FCCP is a mitochondrial

uncoupler that depolarizes the mitochondrial membrane potential [45]. Application of mitochondrial

uncouplers had been successfully used in high-throughput screens searching for genes and chemicals in

the mitophagy pathway [67,89,90]. After fixation, cells were stained with anti-Miro1 and 4',6-diamidino-2-

phenylindole, dihydrochloride (Dapi), and imaged under the confocal microscope with the identical

setting (Figure 6A). Anti-Miro1 has been validated in ICC for specifically recognizing endogenous

Miro1 using human fibroblast-derived neurons with *Miro1* RNAi [33]. The Miro1 intensity was

normalized to Dapi from those images. Using this analysis, we observed significant Miro1 reduction

following FCCP treatment in fibroblasts from a healthy control subject but not in a sporadic PD

fibroblast line (Figure S6A), consistent with our previous studies using the same cell lines by Western

blotting or ELISA [35]. We then employed this method to screen 3 chemical libraries (NIH, FDA, and

Sigma) using the PD cell line. The NIH Clinical collection library contains 377 unique compounds. We

screened those drugs at a defined concentration of 10 µM with 4 biological repeats (independent wells)

for each drug. We compared calculated Miro1 levels in PD cells treated with both compound and FCCP

to those treated with FCCP but without any compound. We found that 11 out of 377 unique compounds

reduced Miro1 protein levels following FCCP treatment on or below 3SD (standard deviation) of control

Miro1 values (those from the same PD cell line without compound treatment but with FCCP), in all

biological repeats (Figure S4A, Table S2A) [91]. We next screened the Biomol FDA library with two five-

fold doses (1.25, 2.5, 5, 10, 20 µM). From 175 unique compounds, we identified 3 hits that reduced

Miro1 protein levels following FCCP treatment in a dose-dependent manner (Figure S4B, Table S2A).

Lastly, we screened the Sigma LOPAC library which contains 1269 unique compounds at a defined

concentration of 20 µM. We found that 21 compounds reduced Miro1 protein levels on or below 3SD of

those control Miro1 values from the same PD cell line without compound treatment but with FCCP

(Figure S4C, Table S2A). We ranked all hits in the order of the degree of Miro1 reduction (Table S2A).

It is important to note that from the same screens we also identified compounds that enhanced Miro1

protein levels in PD cells treated with FCCP (Table S2B). Although our purpose was to search for Miro1

reducers in PD models, the same assay could be used to look for Miro1 enhancers in other disease

models.

Method details: Fibroblasts were plated onto clear-bottomed/black-walled 384-well plates (EK-30091,

E&K Scientific) at 2,000 cells/well with a Matrix Wellmate dispenser (Thermo Fisher), and plates were

incubated at 37°C, 5% CO_2 for 24 hours. Next, chemical library compounds at defined concentrations

were added using fully automated liquid handling system (Caliper Life Sciences Staccato system) with a

Twister II robot and a Sciclone ALH3000 (Caliper Life Sciences) integrated with a V&P Scientific pin

tool, and plates were incubated at 37°C, 5% CO_2 for 10 hours. Then, 20 µM of FCCP (C2920, Sigma-

Aldrich) was added to wells and plates were incubated at 37°C, 5% CO_2 for another 14 hours. Cells

were fixed with ice-cold 90% methanol (482332, Thermo Fisher) for 20 minutes at -20°C, then

incubated with blocking buffer (10% normal goat serum–50062Z, Thermo Fisher; 0.5% BSA–BP1600,

Thermo Fisher; 0.2% Triton X-100–T8787, Thermo Fisher) at room temperature for 15 minutes, and

incubated with anti-Miro1 (HPA010687, Sigma-Aldrich) at 1:100 in blocking buffer overnight at 4°C.

Samples were washed with 1×PBS (10010-049, Thermo Fisher) using a Plate Washer multivalve

(ELx405UV, Bio-Tek), incubated with goat anti-Rabbit IgG (H+L) Cross-Adsorbed, Alexa Fluor 488

(A11008, Thermo Fisher) at 1:500 in blocking buffer at 25°C for 2 hours, then washed again with

1×PBS, and finally 1.0 μg/ml Dapi (D1306, Thermo Fisher) in 1×PBS was added and plates were sealed

using PlateLoc (01867.001, Velocity11). All liquids were dispensed using the Multidrop 384 (5840200,

Titertek) unless otherwise stated. Fluorescent signals in plates were automatically imaged with

ImageXpress Micro (Molecular Devices, IXMicro) and data were analyzed with MetaXpress Analysis

(Molecular Devices). One image was taken from one well. Our negative controls without primary anti-

Miro1 generated no Miro1 signals. Only cells positive for both Miro1 and Dapi staining were chosen for

analysis (>99%; total 400-600 cells per image). The Miro1 fluorescent intensity was normalized to that

of Dapi from the same cell and averaged across all cells from the image from one well. Then, the

median of the mean Miro1 intensities of all wells from the same plate was calculated and subtracted

from the mean Miro1 intensity of each well.

Retesting Hits from the Primary Screens

Overall scheme:

To validate the results of the high-content assays, we retested 34 out of the 35 positive Miro1 reducers

identified at the Stanford HTBC in our own laboratory using fresh compounds and our confocal

microscope. We didn't test 1 hit (CV-3988) because it was unavailable for purchase at the time of the

experiments. We applied each of the 34 compounds at the highest screening concentration to the same

sporadic PD line used at the HTBC with at least 4 biological replicates. We also evaluated Miro1 protein

levels without FCCP but with compound treatment, which unveiled the drug effect on Miro1 under the

basal condition in PD fibroblasts. We imaged all samples with the identical imaging setting, and our

negative controls without primary anti-Miro1 yielded almost no signals (background; Table S2C). Four

compounds, A-77636 hydrochloride, Ebselen, GBR-12909 dihydrochloride and Fenbufen, appeared

toxic (Table S2C). We found that 15 out of the initial 34 hit compounds consistently reduced Miro1

protein following FCCP treatment below 2SD of control Miro1 values (those from the same PD cell line

without treatment) (Figure S5, Table S2C).

Method details: Fibroblasts were plated on coverslips (22-293232, Fisher scientific) into 24-well plates

(10861-558, VWR) at 20,000 cells/well and plates were incubated at 37°C, 5% CO_2 for 24 hours. Next,

fresh chemical compounds dissolved in DMSO were added at defined concentrations and plates were

incubated at 37°C, 5% CO_2 for 10 hours. Then, FCCP at 20 µM was added to wells and plates were

incubated at 37°C, 5% CO_2 for another 14 hours. Cells were fixed with ice-cold 90% methanol for 20

minutes at -20°C, incubated with blocking buffer (10% normal goat serum, 0.5% BSA, 0.2% Triton X-

100) at 25°C for 15 minutes, and immunostained with anti-Miro1 (HPA010687, Sigma-Aldrich) at

1:100 in blocking buffer overnight at 4°C. Samples were washed three times with 1×PBS, incubated

with goat anti-Rabbit IgG (H+L) Cross-Adsorbed, Alexa Fluor 488 (A11008, Thermo Fisher) at 1:500

in blocking buffer at 25°C for 2 hours, washed again three times with 1×PBS, and finally samples were

mounted with ProLong™ Glass Antifade Mountant with NucBlue™ (Dapi) Stain (P36983, hard-setting,

Thermo Fisher) on glass slides and allowed to cure overnight. Samples were imaged at 25°C with a

20×/N.A.0.60 oil Plan-Apochromat objective on a Leica SPE laser scanning confocal microscope (JH

Technologies), with identical imaging parameters among different genotypes. Three images were taken

per coverslip. Total 4 biological repeats (coverslips) for each drug. Images were processed with ImageJ

(Ver. 1.48, NIH), and the average Miro1 intensity in the cytoplasmic area was measured using the

Intensity   Ratio   Nuclei   Cytoplasm   Tool   (http://dev.mri.cnrs.fr/projects/imagej-

macros/wiki/Intensity_Ratio_Nuclei_Cytoplasm_Tool). All compound information for retesting is in

Table S2C or in Figure S7.

Pathway Analysis

Putative mechanistic pathways were generated using graph search over a network of over 2 million

relationships between chemicals, diseases, and genes extracted from the abstracts of journal publications

deposited in PubMed [92]. Each chemical hit and Miro1 were used as the source and target inputs for a

weighted shortest path algorithm where paths were only allowed to traverse chemical-gene and gene-

gene relationships. Graph search over the biomedical graph was performed using Neo4J. A subnetwork

associated with each shortest path was generated from its supporting documents and analyzed using a

knowledge graph browser (docs2graph) to curate the set of supporting documents.

Immunocytochemistry and Confocal Microscopy

Adult fly brains were dissected in PBST (0.3% Triton X-100 in PBS) and incubated with fixative

solution (4% formaldehyde in PBST) for 15 minutes at room temperature. Fixed samples were incubated

in blocking solution (10% BSA in PBST) for an hour, and then immunostained with rabbit anti-TH (AB-

152, EMD Millipore) at 1:200 in blocking solution for 36 hours at 4°C on a rotator. After wash, donkey

anti-rabbit 568 (ab175470, Abcam) at 1:1000 was applied in blocking solution for 1 hour at room

temperature in dark. The DA neuron number was counted throughout the Z stack images of each brain.

Representative images were summed stack images.

Human neurons were fixed in 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA) for

15 minutes at room temperature, permeabilized with 0.25% Triton X-100 in PBS for 20 minutes, then

washed twice in PBS (5 minutes each) and rinsed in deionized water. Alexa Fluor 594 picolyl azide

based TUNEL assay was performed according to the manufacturer's instruction (C10618, Invitrogen).

Coverslips were incubated with TdT reaction buffer at 37°C for 10 minutes, followed by TdT reaction

mixture at 37°C for 60 minutes in a humidified chamber. Coverslips were then rinsed with deionized

water, washed with 3% BSA and 0.1% Triton X-100 in PBS for 5 minutes, followed by incubation with

Click-iT plus TUNEL reaction cocktail at 37°C for 30 minutes, and then washed with 3% BSA and 0.1%

Triton X-100 in PBS for 5 minutes. For nuclear counterstain, 4', 6-Diamidino-2-phenylindole (Dapi;

D9542, Sigma-Aldrich) at 0.5 µg/ml was applied at room temperature for 10 minutes in the dark. The

coverslips were washed with PBS twice, and then mounted on slides with Fluoromount-G mounting

medium (SouthernBiotech). For some experiments, Dapi was stained with ProLong™ Glass Antifade

Mountant with NucBlue™ Stain (P36983, hard-setting, Thermo Fisher). Or iPSC-derived neurons were

fixed in 4% paraformaldehyde for 15 minutes, washed twice in PBS (5 minutes each), and then blocked

in PBS with 5% normal goat serum and 0.3% Triton X-100 for 60 minutes. Neurons were then

immunostained with rabbit anti-TH (AB-152, EMD Millipore) at 1:500, rabbit anti-Flag (F7425, Sigma-

Aldrich) at 1:500, or mouse anti-Myc at 1:1,000 (sc-40, Santa Cruz) in antibody buffer (PBS with 1%

BSA and 0.3% Triton X-100) at 4°C overnight, followed by Alexa Fluor 488 fluorochrome conjugated

goat anti-rabbit IgG (A11008, Invitrogen) or Alexa Fluor 568 fluorochrome conjugated goat anti-mouse

IgG (A11004, Invitrogen) at 1:500 in antibody buffer at room temperature for 1 hour. Coverslips were

washed with PBS twice, and then mounted on slides with Fluoromount-G mounting medium. Samples

were imaged at room temperature with a 20×/N.A.0.60 or 63×/N.A.1.30 oil Plan-Apochromat objective

on a Leica SPE laser scanning confocal microscope, with identical imaging parameters among different

genotypes in a blinded fashion. Images were processed with ImageJ (Ver. 1.48, NIH) using only linear

adjustments of contrast and color.

Live Cell Imaging

RPA at 5 µM (ME043.1, Squarix biotechnology), Mito-FerroGreen at 5 µM (M489, Dojindo
Laboratories), MitoTracker Green at 75-100 nM (M7514, Thermo Fisher), or TMRM at 25 nM (T668,
Molecular Probes) was applied to cells on coverslips in culture media or Hanks Balanced Salt Solution
(HBSS) at 37°C for 30 minutes, then coverslips were washed with HBSS and each was placed in a 35-
mm petri-dish containing Hibernate E low-fluorescence medium (HELF, BrainBits/Transnetyx), and
cells were imaged immediately on a heated stage at 37°C with a 63/N.A.0.9 water-immersion objective.
For calcium imaging, Rhod-2 (R1244, Thermo Fisher) and Calcium Green (C3011MP, Thermo Fisher)
at 5 µM were applied to cells on coverslips and incubated at 37°C for 30 minutes and imaged with the
same set-up as above. Thrombin (10602400001, Sigma-Aldrich) was applied at 100 mUnits/ml to the
dish during imaging. 500 µM Fe^{2+} was applied the day before. Time-lapse movies were obtained
continually with a 2 second interval for 8-10 minutes. For Mito-dsRed and GFP-Miro1 imaging, live
neurons were imaged as above and time-lapse movies were obtained continually with a 1 second interval
before and after 100 µM Antimycin (A8674, Sigma-Aldrich) addition.

Fly Behavioral Analysis

DFP (HY-B0568, MedChemExpress) was dissolved in water at 50 mM and added to SYA fly food at a
final concentration of 163 µM. Benidipine (B6813, Sigma-Aldrich) was dissolved in DMSO at 10 mM
and added to SYA fly food at a final concentration of 2.5 µM. The same volume of solvent was added to
control food. Drug treatment was started on newly-eclosed mated females (day 1), except for DFP on
locomotion which was started from embryogenesis. Food was refreshed every 2 days.

The average Performance Index (PI) (negative geotaxis) was evaluated as previously described [83,93,94].

Briefly, adult flies were gently tapped to the base of a modified 25 ml climbing tube and their climbing

progress was recorded after 45 seconds. Two to four populations of flies were assessed, and for each

population, flies were examined 3 times per experiment. The recorded values were used to calculate the

average PI.

Western Blotting and IP

In general, lysates for SDS-PAGE were mixed with 4×Laemmli sample buffer (161-0747, Bio-Rad)

containing 5-10% β-mercaptoethanol, and then boiled for 5 minutes before being loaded into either a

precast 10% polyacrylamide gel (4561034, Bio-Rad) or manually made gel for electrophoresis with

Tris-glycine-SDS running buffer (24.8 mM Tris, 192 mM glycine, 0.1% SDS). For Native-PAGE,

cells were lysed in lysis buffer (300 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 0.5% Digitonin, 5

mM EDTA, 0.2 mM PMSF, Protease Inhibitor Cocktail–P8340, Sigma-Aldrich), and centrifuged at

16,200 g 4°C for 1 hour. Lysates were mixed with 2×Native Tris-Glycine sample buffer (LC2673,

Thermo Fisher) and loaded directly into a precast 10% polyacrylamide gel (4561034, Bio-Rad) with

Tris-glycine Native running buffer (24.8 mM Tris, 192 mM glycine). After electrophoresis,

nitrocellulose membranes (1620115, Bio-Rad) were used in semi-dry transfer with Bjerrum Schafer–

Nielsen buffer (48 mM Tris, 39 mM glycine, 20% methanol (v/v), pH 9.2). Transferred membranes

were first blocked in TBST (TBS with 0.01-0.05% Tween-20) with 5% milk for 1 hour at room

temperature, and then immunoblotted with the following primary antibodies in TBST with 5% milk at

4°C overnight: mouse anti-Miro1 (WH0055288M1, Sigma-Aldrich) at 1:1,000, rabbit anti-GAPDH

(5174S, Cell Signaling Technology) at 1:1,000, rabbit anti-MCU (HPA016480, Sigma-Aldrich) at 1:500,

rabbit anti-Flag (F7425, Sigma-Aldrich) at 1:750-1,000, rabbit anti-NCLX (ab136975, Abcam) at

1:1,000, rabbit anti-MCU (26312-1-AP, Proteintech) at 1:1,000, rabbit anti-Miro2 (14016S, Cell

Signaling Technology) at 1:1,000, mouse anti-ATP5β (ab14730, Abcam) at 1:1,000, rabbit anti-MCUb

(AP12355b, Abgent) at 1:1,000, rabbit anti-β-Actin (4967S, Cell Signaling Technology) at 1:1,000,

rabbit anti-Mitofusin2 (9482S, Cell Signaling Technology) at 1:1,000, mouse anti-MIC60 (ab110329,

Abcam) at 1:500, mouse anti-MIC19 (TA803454, Origene) at 1:1,000, rabbit anti-MICU1 (HPA037479,

Sigma-Aldrich) at 1:1,000, rabbit anti-MICU2 (ab101465, Abcam) at 1:1,000, rabbit anti-FECH (14466-

1-AP, Proteintech) at 1:500, rabbit anti-EMRE (HPA032117, Sigma-Aldrich) at 1:500, rabbit anti-OPA1

(80471S, Cell Signaling Technology) at 1:1,000, mouse anti-T7 (69622, Novagen) at 1:1,000, and rabbit

anti-VDAC (4661S, Cell Signaling Technology) at 1:1,000. Secondary HRP-conjugated goat anti-rabbit

or mouse IgG antibodies (Jackson ImmunoResearch Laboratory) were used at 1:3,000–10,000 for 1 hour

at room temperature. West Dura ECL Reagent (34075, GE Healthcare) or SuperSignal^{TM} West Pico

PLUS Chemiluminescent Substrate (Thermo fisher scientific) was added. Membranes were scanned

using a Bio-Rad ChemiDoc XRS system.

HEK cells (0.3×10^6/well in a 6-well plate) were plated, grown overnight, and next day when 30-40%

confluent, transfected with 1 μg of DNA, using the calcium phosphate transfection protocol. After 24

hours of transfection, media was changed. After 48 hours of transfection, cells were treated without or

with 500 μM Fe^{2+} (Ammonium Iron(II) Sulfate Hexahydrate, 203505, Sigma-Aldrich) for 24 hours

followed by lysing with 300 μl of lysis buffer (300 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 0.2

mM PMSF, Protease Inhibitor Cocktail) supplemented with 5 mM EDTA or 500 μM Fe^{2+}. For each

condition, 3 wells of a 6-well plate were pooled, followed by centrifugation at 16,200 g for 10 minutes

at 4°C. Part of the supernatant (110 μl out of 900 μl total) was reserved as 'Input'. The remaining 790 μl

was incubated with 10 μl of anti-Flag (F7425, Sigma-Aldrich) for 2 hours on a rotator at 4°C, and then

combined with 60 μl of washed Protein A–Sepharose beads (17-0780-01, GE Healthcare) for 1 hour on

a rotator at 4°C. Beads were then washed five times with lysis buffer. Residual buffer was removed from

the last wash and the beads were mixed with 100 μl of 2×Tris-Glycine Native Sample Buffer (LC2673,

Thermo Fisher) or Laemmli buffer (then boiled), and loaded into a precast gel (4561034, Bio-Rad). For

each gel running, approximately 5% (45 μl out of total 900 μl lysate) of total lysates (Input) and 45% of

total immunoprecipitated proteins were loaded.

Postmortem Brain Lysing for Western Blotting

A piece of postmortem brain (1 mm^3) was cut and put in a 1.5 ml tube, and lysed in 500 μl of lysis

buffer (300 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 0.5% Digitonin, Protease Inhibitor

Cocktail, 0.2 mM PMSF). Tissue was sonicated and then centrifuged at 16,200 g 4°C for 1 hour.

Supernatant was collected.

RT-qPCR

Total RNA was extracted from at least one million cells per sample using TRIzol (Gibco) according to

the manufacturer's instructions. Concentrations of total RNA were measured using a NanoDrop. 1 μg of

total RNA was then subjected to DNA digestion using DNase I (Ambion), immediately followed by

reverse transcription using the iScript Reverse Transcription Supermix (1708841, BIO-RAD). qPCR

was performed using the StepOnePlus^{TM} instrument (Thermo Fisher) and SYBR Green Supermix (172-

5270, Bio-Rad) following the manufacturer's instructions. Human *GAPDH* was amplified as an internal

standard. Expression levels were analyzed by the StepOne Software (Version 2.2.2). The relative

expression level was divided by that of *GAPDH* from the same experiment. Each experiment was

repeated 3-4 times in duplicate.

ATP Levels

ATP concentrations were determined using the Roche ATP Bioluminescence Assay Kit HS II (no. 11699709001, Sigma). Briefly, cell lysate was homogenized in 150 μl of ice-cold lysis buffer as above using a Kontes pellet pestle. Lysate was then boiled for 5 minutes and centrifuged at 20,000 g 4°C for 1 minute. Cleared lysate was diluted 1:200 in dilution buffer and loaded with 10 μl of luciferase. Luminescence was immediately measured using a FlexStation 3 (Molecular Devices). Total protein amounts were measured using the Bicinchoninic Acid (BCA) Protein Assay Kit (23227, Thermo Scientific). The ATP level in each sample was normalized to the total protein amount.

SEC

For control and H_2O_2 treatment (10 mM for 10 minutes), HEK cells were lysed in lysis buffer (1% Triton X-100, 50 mM Tris pH7.5, 300 mM NaCl, 5 mM EDTA, 1:1000 Protease inhibitor cocktail, 0.2 mM PMSF). For Fe^{2+} treatment, HEK cells were treated with 5 mM Fe^{2+} (Ammonium Iron(II) Sulfate Hexahydrate, 203505, Sigma-Aldrich) for 24 hours, and lysed in the same buffer above where EDTA was replaced with 5 mM Fe^{2+}. Cell lysates were injected onto a Superdex 200 Increase 10/300 GL column (GE Healthcare Life Sciences). To slow Fe^{2+}-oxidation and precipitation in equilibration buffer, we preequilibrated the column with low pH buffer and reducing reagent. For the control sample, the column was preequilibrated with 20 mM MES pH 6.0, 300 mM NaCl, 5 mM TCEP, and 0.1% Triton X-100. For the Fe^{2+}-treated sample, the column was preequilibrated with 20 mM MES pH 6.0, 300 mM NaCl, 5 mM TCEP, 2 mM Fe^{2+}, and 0.1% Triton X-100. For the H_2O_2-treated sample, the column was preequilibrated with 20 mM Tris pH 7.4, 300 mM NaCl, and 0.1% Triton X-100.

Fluorescence SEC: For each sample, 5 µg purified human MCU was incubated either on ice or at 35°C

in a thermal cycler for 15 minutes. After the incubation, all samples were centrifuged in a tabletop

centrifuge at 21,000 g at 4°C for 10 minutes to remove precipitated protein. The supernatant was then

injected onto a Superdex 200 Increase 10/300 GL column preequilibrated with 20 mM Tris pH 7.4, 150

mM NaCl, 0.01% lauryl maltose neopentyl glycol (LMNG), and 0.001% cholesteryl hemisuccinate

(CHS).

Iron Detection Assay

HEK cells from one well of a 6-well plate were lysed in 300-400 µl lysis buffer (1% Triton X-100, 50

mM Tris pH7.5, 300 mM NaCl, 5 mM EDTA, 1:1000 Protease inhibitor cocktail, 0.2 mM PMSF), and 3

wells were pooled for each IP, followed by centrifugation at 16,200 g 4°C for 10 minutes. For Fe^{2+}

treatment, HEK cells were treated with 500 µM Fe^{2+} (Ammonium Iron(II) Sulfate Hexahydrate, 203505,

Sigma-Aldrich) for 20-21 hours, and lysed in the same buffer where EDTA was replaced with 500 µM

Fe^{2+}. Supernatant was then incubated with 7 µl rabbit anti-MCU (26312-1-AP, ProteinTech), 7 µl rabbit

IgG (2729S, cell signaling), or 5 µl rabbit anti-Flag (F7425, Sigma-Aldrich) for 2-3 hours at 4°C, and

incubated with Protein A–Sepharose beads (17-0780-01, GE Healthcare) for another hour at 4°C. To

detect iron levels, the colorimetric iron detection assay kit (ab83366, Abcam) was used according to the

manufacturer's instruction. Briefly, 180 µl Iron Assay Buffer was added to the beads with IP, and 100 µl

of mixed buffer-beads was added to each well of total 2 wells. Next, 5 µl Iron Reducer was added to the

well for iron (II+III), and 5 µl of Iron Assay Buffer was added to the well for iron (II), incubated at 37°C

for 30 minutes. Then, 100 µl Iron Probe was added to each well, incubated at 37°C for 60 minutes,

protected from light. Samples were measured immediately by an InfiniteM200 Pro TECAN plate reader

at 593 nm.

ICP-MS

Iron concentration determinations were performed on an Agilent 8900 triple quadrupole inductively

coupled plasma mass spectrometer (ICP-MS) attached to an Agilent SPS-4 autosampler. The sample

introduction system also included a standard Scott double pass cooled spray chamber operated at 2'C, a

2.5mm i.d. Agilent glass torch, and a 200 μl/min Micromist nebulizer. The ICP-MS was fitted with the

high sensitivity lens (s-lens), nickel cones and operated in ammonia (NH_3) gas mode. Ammonia gas in

the reaction cell of the triple quadrupole diminished the presence of argide including ArO which could

form an isobaric interference with iron at mass 56. The large argide peak could also cause high

backgrounds and more complex interferences at mass 57, another mass frequently monitored for iron

concentrations. In ammonia gas mode, mass 56 and 57 backgrounds were dramatically reduced,

resulting in an increase in the signal to noise ratio. This improved the iron detection limit in measured

solutions on our instrument from about 1.8 ppb to 0.04 ppb. All reported concentrations were

determined at mass 57 in ammonia gas mode.

Before analysis, all solutions were further diluted by 50% by teeing in an internal standard solution of 1

μl/l indium (High Purity Standards) prepared with 2% v/v nitric acid to correct for instrumental drift.

Procedural blanks were run in parallel with samples and averaged 48 pg contributing to less than 4% of

the analytical signal and were regarded as negligible. Reproducibility was estimated from repeated

analyses (n=3) of each sample in a single session.

PMBC Analysis

PBMCs were thawed in a 37°C water bath and plated into a 6-well plate (353046, Falcon) with RPMI

media (11875-093, Gibco), supplemented with 10% fetal bovine serum (900-108, heat-inactivated,

Gemini Bio-Products), 1% Pen-Strep (15140-122, Gibco), 1×GlutaMax (35050061, Thermo Fisher), and

maintained in a 37°C, 5% CO_2 incubator with humidified atmosphere. After a 1-hour recovery in the

incubator, cells were then trypsinized with 0.25% Trypsin-EDTA (25200-056, Gibco) and collected in a

cell suspension to be counted via TC20 Cell Counter (Bio-Rad).

Western blotting: After counting, the cell suspension (total 8 million cells) was equally split into 8

groups and replated with 3 ml media per well, 2 groups remaining untreated (to serve as controls), 3

groups treated with a combination of 20 μM Antimycin A (A8674, Sigma-Aldrich) and 2 μM

Oligomycin (75351, Sigma-Aldrich) (AO), and 3 groups treated with 40 μM CCCP (C2759, Sigma-

Aldrich). The untreated groups, serving as the 0-hour treatment in a time course, were then immediately

pelleted and lysed with 150 μl lysis buffer (1% Triton X-100, 50 mM Tris, 300 mM NaCl, 5 mM EDTA,

0.2 mM PMSF, 1:1,000 Protease Inhibitor Cocktail) and stored at -80°C. At each of the following time

points after treatment – 1 hour, 6 hours, and 14 hours – 1 group from each of the treatments (AO or

CCCP) was similarly pelleted, lysed with lysis buffer, and stored at -80°C. Lysates collected were run in

SDS-PAGE (30 μl loaded) and blotted with the following primary antibodies: mouse anti-Miro1

(WH0055288M1, Sigma-Aldrich) at 1:1,000, rabbit anti-GAPDH (5174S, Cell Signaling Technology) at

1:1,000, mouse anti-ATP5β (ab14730, Abcam) at 1:1,000, rabbit anti-VDAC (4661S, Cell Signaling

Technology) at 1:1,000, and rabbit anti-β-Actin (4967S, Cell Signaling Technology) at 1:1,000.

ELISA: After counting, the cell suspension was equally split and replated with 3 ml media per well

containing either 40 μM CCCP or a commensurate volume of DMSO for 6 hours. PBMC vials were

provided in duplicate by the MJFF, so the cell suspensions did not need to be split equally – they kept

separate throughout the counting and replating process. For PBMCs that received drug treatment, the

cell suspension was equally split into 3 wells. One well was pretreated with 10 µM Benidipine or MR3

(1915758622, MCULE) for 18 hours, and then this well and a second well were treated with 40 µM

CCCP for 6 hours. The same volume of DMSO was applied to control wells. The minimum cell number

needed for ELISA was 400,000 per well. Next, cells were scraped gently with a cell lifter and collected

into suspension. The cells were then pelleted and lysed with lysis buffer (1% Triton X-100, 50 mM Tris,

300 mM NaCl, 5 mM EDTA, 0.2 mM PMSF, 1:1,000 Protease Inhibitor Cocktail) and stored at -80°C.

The Miro1 protein level in each sample collected was determined using the RHOT1 ELISA Kit 96T

(EKL54911, Biomatik) according to the manufacturer's manual, and normalized to the total protein

level in the same sample using the BCA Protein Assay Kit (23227, Thermo Scientific). For the ELISA

kit, the specificity and stability were validated by Biomatik. The dynamic detection range (0.625-40

ng/ml), sensitivity (lower limit of detection–LLOD; 0.112 ng/ml), and precision (inter- and intra-assay)

were determined by both Biomatik and us [48] (Figure S7), and the results were comparable. OD values

for ELISA and BCA were measured on an InfiniteM200 Pro TECAN plate reader at 450 and 562 nm

respectively, in duplicate. All experiments were performed in a double-blinded fashion, with PD status

unknown to the experimenter.

1016

1017   Machine Learning: Anova analysis revealed a significant association of Miro1 Ratio with PD status, but

1018   not with other demographic and clinical parameters. We then trained a machine learning model to

1019   classify if a subject was with PD or not. The general form of the logistic regression model was

1020   represented by the following equation:

$$p(C_1|x) = \frac{1}{e^{-(\beta_0 + \sum_{i=1}^{N} \beta_i x_i)}}$$

1021   The goal of the model training was to learn the model parameters   such that we could predict the

1022   posterior probability   .   represented the positive class "PD+" status and   represented the input

1023   of the model. We set a threshold value above which all predictions belonged to the positive class, i.e.,

Left panel showed the trained model for the probability of "PD+" with "Miro1 Ratio" as the only feature. The decision boundary for the probability was set to 0.57. This value maximized the accuracy and AUC score for this model. Above this threshold all predictions were considered as "PD+" and the ones below were considered as "PD-". Right panel showed the trained model for "UPDRS". Please note that in both figures the probability of "PD+" was not zero when Input was zero.

1024   "PD+", and below which they belonged to the "PD-" class.

1025

1026   We trained three logistic regression models with three different inputs. The first model used "Miro1

1027   Ratio" as the only input, the second one had "UPDRS" as the only input, and the third model combined

1028   "Miro1 Ratio" and "UPDRS" as inputs. All three models were trained using the glm function in R.

1029

1030   We broke the total subjects into a 60% training set and a 40% testing set. Data was cleaned by

1031   converting assigning column values to appropriate data types and omitting rows of data where no

1032   "Miro1 Ratio" or "UPDRS" values were present. The target variable, PD status, was labeled and

1033   encoded where 0 represented a subject without PD and 1 represented a PD positive subject.

1034

1035   The table below showed the   and the corresponding p-values.

Model	features	Estimate	Std Error	z.value	Pr
Miro1 Ratio only	(Intercept)	-0.76541	0.500231	-1.5301	0.125991
	Miro1_Ratio	1.006977	0.460706	2.185727	0.028836
UPDRS only	(Intercept)	-1.32384	0.340155	-3.89189	0.0000995
	UPDRS	0.13644	0.027073	5.039669	4.66E-07
Miro1 Ratio + UPDRS	(Intercept)	-3.45836	1.030444	-3.35618	0.00079
	UPDRS	0.193639	0.062998	3.073723	0.002114
	Miro1_Ratio	2.002893	0.863175	2.320378	0.02032
	UPDRS:Miro1_Ratio	-0.04647	0.048114	-0.96573	0.334178

1036

The estimate column represented β_i for each model. Since the "Miro1 Ratio" only and the "UPDRS"

only models had a single feature, there were only two β_i values present. β_0 represented the intercept and

β_1 represented "Miro1 Ratio" value or the "UPDRS" value depending on the model. We observed that

the β_i estimate for the Miro1 Ratio model was significant ($p<0.05$) suggesting that the model predictions

were statistically significant. Similarly, for UPDRS the β_1 value was also significant. Both the UPDRS

and Miro1 Ratio estimates were significant in the combined model, however the interaction term

between them was non-significant.

1044

1045 Once the models were trained, they were tested by evaluating the accuracy and AUC scores. The table

1046 below presented the accuracy and AUC scores for all models:

Models	Accuracy	AUC
Miro1 Ratio only	0.676	0.677
UPDRS only	0.812	0.822

Models	Accuracy	AUC
Miro1 Ratio + UPDRS	0.878	0.878

1047

1048   We observed that the Miro1 Ratio only model performed worse with an accuracy of 67.6% than the

1049   UPDRS only model (81.2%). However, together it yielded a higher score of 87.8%.

1050

1051   We then added "Age" and "Sex" as additional inputs to the combined mode which did not affect the

1052   accuracy.

1053

1054   To determine the statistical significance between "Miro1 Ratio + UPDRS" and "UPDRS only", we

1055   generated 100 samples (datasets) with replacement from each test set,

Distribution of AUC Scores

predicted outcomes for each sample, and calculated the AUC scores

for each sample. In line with standard bootstrapping sampling

practice, we used the test set size as the sample size for generating

each sample. Following the same process for both models it yielded

two sample distributions, one of each model. Figure shows the box

plot for both these distributions.

1062

Welch Two Sample T-test Result:

data: auc_scores_sampling_udprs_only and auc_scores_sampling_miro_udprs

t = -6.0947,

df = 195.48,

p-value = 5.736e-09 alternative hypothesis: true difference in means is not equal to 0 95 percent

confidence interval: -0.04893487, -0.02500783

sample estimates: mean of x: 0.8529143 mean of y : 0.8898856

1070

Here is a breakdown of the results:

t = -6.0947: This is the t-statistic value. The negative sign indicates that the mean of

auc_scores_sampling_udprs_only is less than that of auc_scores_sampling_miro_udprs. df = 195.48:

This is the degree of freedom for the test. The Welch T-test adjusts the degrees of freedom based on the

variances of the two samples, making it more robust when the variances are unequal. p-value = 5.736e-

09: This is the probability of observing a t-statistic as extreme as the one calculated (or more extreme) if

the true difference in means was zero (i.e., under the null hypothesis). A very small p-value like this one

indicates strong evidence against the null hypothesis. 95 percent confidence interval: -0.04893487 -

0.02500783: This interval provides a range of values for the true difference in means between the two

samples. Since the entire interval is negative, it suggests that the mean of

auc_scores_sampling_udprs_only is consistently lower than the mean of

auc_scores_sampling_miro_udprs. sample estimates: These are the sample means for the two groups.

The mean of auc_scores_sampling_udprs_only (mean of x) is 0.8529143, and the mean of

auc_scores_sampling_miro_udprs (mean of y) is 0.8898856.

1085

1086 Code is at https://data.mendeley.com/datasets/wtjcvm3cm3/1

1087

Human Genetics

Based on our experimental data showing the importance of the calcium-Miro axis in PD models, the

following calcium related genes were investigated for genetic association: *CACNA1S, CACNA1C,*

CACNA1D, CACNA1B, CACNA1G, CACNA1H, CACNA1I, RHOT1, RHOT2, MCU, CCDC109B,

MICU1, MICU2, SMDT1, SLC8B1, TRPV1, TRPV2, TRPV3, TRPV4 (Table S5). First, to identify

whether common variants regulating the expression (eQTL) or splicing (sQTL) of these genes are

associated with PD risk, we queried the genome-wide association study (GWAS) summary statistics

from [49]. Genome-wide significant association (if any) was then intersected with eQTL/sQTL

information from GTEx [95]. Due to the absence of association, colocalization was not formally tested.

Second, to test the association of rare variants in these genes we performed Burden and SKATO Test, as

implemented in the R package *SKAT* v2.2.4, considering variants with a CADD (Combined Annotation

Dependent Depletion) score [96] above 20 to only include variants most likely to have an impact on

protein function. Additionally, we grouped variants in *CAC* genes by channel type, particularly grouping

T-type (*CACNA1G, CACNA1H, CACNA1I*) and L-type (*CACNA1S, CACNA1C, CACNA1D*). For rare

variant analysis, we obtained whole-genome sequencing (WGS) data from the Accelerating Medicines

Partnership–Parkinson's Disease (AMP–PD), composed of the following cohorts: New Discovery of

Biomarkers (BioFIND), the Harvard Biomarker Study (HBS), the Parkinson's Progression Markers

Initiative (PPMI) [97], and the Parkinson's Disease Biomarkers Program (PDBP). European ancestry

individuals and principal components to account for genetic ancestry were computed as described in [98],

leading to a total of 1,168 controls (45.4% males) and 2,241 cases (62.8% males). Significance was

assessed separately for Burden and SKATO Test, though in genes where deleterious missense variants

were all associated with the phenotype in the same direction (either increasing or decreasing risk) these

two tests led to similar results.

1111

References

1.   Diaz-Ortiz, M.E., Seo, Y., Posavi, M., Carceles Cordon, M., Clark, E., Jain, N., Charan, R.,
Gallagher, M.D., Unger, T.L., Amari, N., et al. (2022). GPNMB confers risk for Parkinson's
disease through interaction with alpha-synuclein. Science *377*, eabk0637.
10.1126/science.abk0637.

2.   Apicco, D.J., Shlevkov, E., Nezich, C.L., Tran, D.T., Guilmette, E., Nicholatos, J.W., Bantle,
C.M., Chen, Y., Glajch, K.E., Abraham, N.A., et al. (2021). The Parkinson's disease-associated
gene ITPKB protects against alpha-synuclein aggregation by regulating ER-to-mitochondria
calcium release. Proceedings of the National Academy of Sciences of the United States of
America *118*. 10.1073/pnas.2006476118.
3.   Lee, K.S., Huh, S., Lee, S., Wu, Z., Kim, A.K., Kang, H.Y., and Lu, B. (2018). Altered ER-
mitochondria contact impacts mitochondria calcium homeostasis and contributes to
neurodegeneration in vivo in disease models. Proceedings of the National Academy of Sciences
of the United States of America *115*, E8844-E8853. 10.1073/pnas.1721136115.
4.   Tabata, Y., Imaizumi, Y., Sugawara, M., Andoh-Noda, T., Banno, S., Chai, M., Sone, T.,
Yamazaki, K., Ito, M., Tsukahara, K., et al. (2018). T-type Calcium Channels Determine the
Vulnerability of Dopaminergic Neurons to Mitochondrial Stress in Familial Parkinson Disease.
Stem cell reports *11*, 1171-1184. 10.1016/j.stemcr.2018.09.006.
5.   Buttner, S., Faes, L., Reichelt, W.N., Broeskamp, F., Habernig, L., Benke, S., Kourtis, N., Ruli,
D., Carmona-Gutierrez, D., Eisenberg, T., et al. (2013). The Ca2+/Mn2+ ion-pump PMR1 links
elevation of cytosolic Ca(2+) levels to alpha-synuclein toxicity in Parkinson's disease models.
Cell death and differentiation *20*, 465-477. 10.1038/cdd.2012.142.
6.   Surmeier, D.J., Obeso, J.A., and Halliday, G.M. (2017). Selective neuronal vulnerability in
Parkinson disease. Nature reviews. Neuroscience *18*, 101-113. 10.1038/nrn.2016.178.
7.   Kim, J.W., Yin, X., Jhaldiyal, A., Khan, M.R., Martin, I., Xie, Z., Perez-Rosello, T., Kumar, M.,
Abalde-Atristain, L., Xu, J., et al. (2020). Defects in mRNA Translation in LRRK2-Mutant
hiPSC-Derived Dopaminergic Neurons Lead to Dysregulated Calcium Homeostasis. Cell stem
cell *27*, 633-645 e637. 10.1016/j.stem.2020.08.002.
8.   Angelova, P.R., Choi, M.L., Berezhnov, A.V., Horrocks, M.H., Hughes, C.D., De, S., Rodrigues,
M., Yapom, R., Little, D., Dolt, K.S., et al. (2020). Alpha synuclein aggregation drives
ferroptosis: an interplay of iron, calcium and lipid peroxidation. Cell death and differentiation *27*,
2781-2796. 10.1038/s41418-020-0542-z.
9.   Verma, M., Callio, J., Otero, P.A., Sekler, I., Wills, Z.P., and Chu, C.T. (2017). Mitochondrial
Calcium Dysregulation Contributes to Dendrite Degeneration Mediated by PD/LBD-Associated
LRRK2 Mutants. The Journal of neuroscience : the official journal of the Society for
Neuroscience *37*, 11151-11165. 10.1523/JNEUROSCI.3791-16.2017.
10.   Belaidi, A.A., and Bush, A.I. (2016). Iron neurochemistry in Alzheimer's disease and Parkinson's
disease: targets for therapeutics. Journal of neurochemistry *139 Suppl 1*, 179-197.
10.1111/jnc.13425.
11.   Vuuren, M.J.V., Nell, T.A., Carr, J.A., Kell, D.B., and Pretorius, E. (2020). Iron Dysregulation
and Inflammagens Related to Oral and Gut Health Are Central to the Development of
Parkinson's Disease. Biomolecules *11*. 10.3390/biom11010030.
12.   Baughman, J.M., Perocchi, F., Girgis, H.S., Plovanich, M., Belcher-Timme, C.A., Sancak, Y.,
Bao, X.R., Strittmatter, L., Goldberger, O., Bogorad, R.L., et al. (2011). Integrative genomics
identifies MCU as an essential component of the mitochondrial calcium uniporter. Nature *476*,
341-345. 10.1038/nature10234.
13.   Palty, R., Silverman, W.F., Hershfinkel, M., Caporale, T., Sensi, S.L., Parnis, J., Nolte, C.,
Fishman, D., Shoshan-Barmatz, V., Herrmann, S., et al. (2010). NCLX is an essential component
of mitochondrial Na+/Ca2+ exchange. Proceedings of the National Academy of Sciences of the
United States of America *107*, 436-441. 10.1073/pnas.0908099107.

14.   Fan, C., Fan, M., Orlando, B.J., Fastman, N.M., Zhang, J., Xu, Y., Chambers, M.G., Xu, X.,
1163   Perry, K., Liao, M., and Feng, L. (2018). X-ray and cryo-EM structures of the mitochondrial
1164   calcium uniporter. Nature *559*, 575-579. 10.1038/s41586-018-0330-9.
1165   15.   Fan, M., Zhang, J., Tsai, C.W., Orlando, B.J., Rodriguez, M., Xu, Y., Liao, M., Tsai, M.F., and
1166   Feng, L. (2020). Structure and mechanism of the mitochondrial Ca(2+) uniporter holocomplex.
1167   Nature *582*, 129-133. 10.1038/s41586-020-2309-6.
1168   16.   Lambert, J.P., Luongo, T.S., Tomar, D., Jadiya, P., Gao, E., Zhang, X., Lucchese, A.M.,
1169   Kolmetzky, D.W., Shah, N.S., and Elrod, J.W. (2019). MCUB Regulates the Molecular
1170   Composition of the Mitochondrial Calcium Uniporter Channel to Limit Mitochondrial Calcium
1171   Overload During Stress. Circulation *140*, 1720-1733. 10.1161/CIRCULATIONAHA.118.037968.
1172   17.   Patron, M., Tarasenko, D., Nolte, H., Kroczek, L., Ghosh, M., Ohba, Y., Lasarzewski, Y.,
1173   Ahmadi, Z.A., Cabrera-Orefice, A., Eyiama, A., et al. (2022). Regulation of mitochondrial
1174   proteostasis by the proton gradient. The EMBO journal *41*, e110476.
1175   10.15252/embj.2021110476.
1176   18.   Paul, B.T., Manz, D.H., Torti, F.M., and Torti, S.V. (2017). Mitochondria and Iron: current
1177   questions. Expert Rev Hematol *10*, 65-79. 10.1080/17474086.2016.1268047.
1178   19.   Dong, Z., Shanmughapriya, S., Tomar, D., Siddiqui, N., Lynch, S., Nemani, N., Breves, S.L.,
1179   Zhang, X., Tripathi, A., Palaniappan, P., et al. (2017). Mitochondrial Ca(2+) Uniporter Is a
1180   Mitochondrial Luminal Redox Sensor that Augments MCU Channel Activity. Molecular cell *65*,
1181   1014-1028 e1017. 10.1016/j.molcel.2017.01.032.
1182   20.   Berridge, M.J., Bootman, M.D., and Roderick, H.L. (2003). Calcium signalling: dynamics,
1183   homeostasis and remodelling. Nature reviews. Molecular cell biology *4*, 517-529.
1184   10.1038/nrm1155.
1185   21.   Clapham, D.E. (2007). Calcium signaling. Cell *131*, 1047-1058. 10.1016/j.cell.2007.11.028.
1186   22.   Tomar, D., Dong, Z., Shanmughapriya, S., Koch, D.A., Thomas, T., Hoffman, N.E., Timbalia,
1187   S.A., Goldman, S.J., Breves, S.L., Corbally, D.P., et al. (2016). MCUR1 Is a Scaffold Factor for
1188   the MCU Complex Function and Promotes Mitochondrial Bioenergetics. Cell Rep *15*, 1673-
1189   1685. 10.1016/j.celrep.2016.04.050.
1190   23.   Ghosh, S., Basu Ball, W., Madaris, T.R., Srikantan, S., Madesh, M., Mootha, V.K., and Gohil,
1191   V.M. (2020). An essential role for cardiolipin in the stability and function of the mitochondrial
1192   calcium uniporter. Proceedings of the National Academy of Sciences of the United States of
1193   America *117*, 16383-16390. 10.1073/pnas.2000640117.
1194   24.   Michailides, C., and Velissaris, D. (2021). Common anti-oxidant vitamin C as an anti-infective
1195   agent with remedial role on SARS-CoV-2 infection. An update. Monaldi Arch Chest Dis *91*.
1196   10.4081/monaldi.2021.1808.
1197   25.   Chen, A.Y., Lu, J.M., Yao, Q., and Chen, C. (2016). Entacapone is an Antioxidant More Potent
1198   than Vitamin C and Vitamin E for Scavenging of Hypochlorous Acid and Peroxynitrite, and the
1199   Inhibition of Oxidative Stress-Induced Cell Death. Med Sci Monit *22*, 687-696.
1200   10.12659/msm.896462.
1201   26.   Munson, M.J., Mathai, B.J., Ng, M.Y.W., Trachsel-Moncho, L., de la Ballina, L.R., Schultz,
1202   S.W., Aman, Y., Lystad, A.H., Singh, S., Singh, S., et al. (2021). GAK and PRKCD are positive
1203   regulators of PRKN-independent mitophagy. Nature communications *12*, 6101. 10.1038/s41467-
1204   021-26331-7.
1205   27.   Billesbolle, C.B., Azumaya, C.M., Kretsch, R.C., Powers, A.S., Gonen, S., Schneider, S.,
1206   Arvedson, T., Dror, R.O., Cheng, Y., and Manglik, A. (2020). Structure of hepcidin-bound

ferroportin reveals iron homeostatic mechanisms. Nature *586*, 807-811. 10.1038/s41586-020-
2668-z.
28.   Lin, Y.F., Cheng, C.W., Shih, C.S., Hwang, J.K., Yu, C.S., and Lu, C.H. (2016). MIB: Metal
Ion-Binding Site Prediction and Docking Server. J Chem Inf Model *56*, 2287-2291.
10.1021/acs.jcim.6b00407.
29.   Lu, C.H., Lin, Y.F., Lin, J.J., and Yu, C.S. (2012). Prediction of metal ion-binding sites in
proteins using the fragment transformation method. PloS one *7*, e39252.
10.1371/journal.pone.0039252.
30.   Lee, S.K., Shanmughapriya, S., Mok, M.C.Y., Dong, Z., Tomar, D., Carvalho, E., Rajan, S.,
Junop, M.S., Madesh, M., and Stathopulos, P.B. (2016). Structural Insights into Mitochondrial
Calcium Uniporter Regulation by Divalent Cations. Cell Chem Biol *23*, 1157-1169.
10.1016/j.chembiol.2016.07.012.
31.   Fernandez, B., Olmedo, P., Gil, F., Fdez, E., Naaldijk, Y., Rivero-Rios, P., Bracher, F., Grimm,
C., Churchill, G.C., and Hilfiker, S. (2022). Iron-induced cytotoxicity mediated by
endolysosomal TRPML1 channels is reverted by TFEB. Cell Death Dis *13*, 1047.
10.1038/s41419-022-05504-2.
32.   Jadiya, P., Kolmetzky, D.W., Tomar, D., Di Meco, A., Lombardi, A.A., Lambert, J.P., Luongo,
T.S., Ludtmann, M.H., Pratico, D., and Elrod, J.W. (2019). Impaired mitochondrial calcium
efflux contributes to disease progression in models of Alzheimer's disease. Nature
communications *10*, 3885. 10.1038/s41467-019-11813-6.
33.   Hsieh, C.H., Shaltouki, A., Gonzalez, A.E., Bettencourt da Cruz, A., Burbulla, L.F., St Lawrence,
E., Schule, B., Krainc, D., Palmer, T.D., and Wang, X. (2016). Functional Impairment in Miro
Degradation and Mitophagy Is a Shared Feature in Familial and Sporadic Parkinson's Disease.
Cell stem cell *19*, 709-724. 10.1016/j.stem.2016.08.002.
34.   Shaltouki, A., Hsieh, C.H., Kim, M.J., and Wang, X. (2018). Alpha-synuclein delays mitophagy
and targeting Miro rescues neuron loss in Parkinson's models. Acta Neuropathol *136*, 607-620.
10.1007/s00401-018-1873-4.
35.   Hsieh, C.H., Li, L., Vanhauwaert, R., Nguyen, K.T., Davis, M.D., Bu, G., Wszolek, Z.K., and
Wang, X. (2019). Miro1 Marks Parkinson's Disease Subset and Miro1 Reducer Rescues Neuron
Loss in Parkinson's Models. Cell Metab *30*, 1131-1140 e1137. 10.1016/j.cmet.2019.08.023.
36.   Li, L., Conradson, D.M., Bharat, V., Kim, M.J., Hsieh, C.H., Minhas, P.S., Papakyrikos, A.M.,
Durairaj, A.S., Ludlam, A., Andreasson, K.I., et al. (2021). A mitochondrial membrane-bridging
machinery mediates signal transduction of intramitochondrial oxidation. Nat Metab.
10.1038/s42255-021-00443-2.
37.   Lee, D.G., Park, J., Lee, H.S., Lee, S.R., and Lee, D.S. (2016). Iron overload-induced calcium
signals modulate mitochondrial fragmentation in HT-22 hippocampal neuron cells. Toxicology
365, 17-24. 10.1016/j.tox.2016.07.022.
38.   Niescier, R.F., Hong, K., Park, D., and Min, K.T. (2018). MCU Interacts with Miro1 to
Modulate Mitochondrial Functions in Neurons. The Journal of neuroscience : the official journal
of the Society for Neuroscience *38*, 4666-4677. 10.1523/JNEUROSCI.0504-18.2018.
39.   Gottschalk, B., Klec, C., Leitinger, G., Bernhart, E., Rost, R., Bischof, H., Madreiter-Sokolowski,
C.T., Radulovic, S., Eroglu, E., Sattler, W., et al. (2019). MICU1 controls cristae junction and
spatially anchors mitochondrial Ca(2+) uniporter complex. Nature communications *10*, 3732.
10.1038/s41467-019-11692-x.
40.   Katona, M., Bartok, A., Nichtova, Z., Csordas, G., Berezhnaya, E., Weaver, D., Ghosh, A.,
Varnai, P., Yule, D.I., and Hajnoczky, G. (2022). Capture at the ER-mitochondrial contacts

licenses IP(3) receptors to stimulate local Ca(2+) transfer and oxidative metabolism. Nature
communications *13*, 6779. 10.1038/s41467-022-34365-8.

41.   Garcia-Rivas Gde, J., Carvajal, K., Correa, F., and Zazueta, C. (2006). Ru360, a specific
mitochondrial calcium uptake inhibitor, improves cardiac post-ischaemic functional recovery in
rats in vivo. Br J Pharmacol *149*, 829-837. 10.1038/sj.bjp.0706932.

42.   Wang, X., Winter, D., Ashrafi, G., Schlehe, J., Wong, Y.L., Selkoe, D., Rice, S., Steen, J.,
LaVoie, M.J., and Schwarz, T.L. (2011). PINK1 and Parkin target Miro for phosphorylation and
degradation to arrest mitochondrial motility. Cell *147*, 893-906. 10.1016/j.cell.2011.10.018.

43.   Lopez-Domenech, G., Howden, J.H., Covill-Cooke, C., Morfill, C., Patel, J.V., Burli, R.,
Crowther, D., Birsa, N., Brandon, N.J., and Kittler, J.T. (2021). Loss of neuronal Miro1 disrupts
mitophagy and induces hyperactivation of the integrated stress response. The EMBO journal *40*,
e100715. 10.15252/embj.2018100715.

44.   Wang, X., and Schwarz, T.L. (2009). The mechanism of Ca2+ -dependent regulation of kinesin-
mediated mitochondrial motility. Cell *136*, 163-174. 10.1016/j.cell.2008.11.046.

45.   Birsa, N., Norkett, R., Wauer, T., Mevissen, T.E., Wu, H.C., Foltynie, T., Bhatia, K., Hirst, W.D.,
Komander, D., Plun-Favreau, H., and Kittler, J.T. (2014). Lysine 27 ubiquitination of the
mitochondrial transport protein Miro is dependent on serine 65 of the Parkin ubiquitin ligase.
The Journal of biological chemistry *289*, 14569-14582. 10.1074/jbc.M114.563031.

46.   Darakhshan, S., and Pour, A.B. (2015). Tranilast: a review of its therapeutic applications.
Pharmacol Res *91*, 15-28. 10.1016/j.phrs.2014.10.009.

47.   Ordureau, A., Paulo, J.A., Zhang, J., An, H., Swatek, K.N., Cannon, J.R., Wan, Q., Komander,
D., and Harper, J.W. (2020). Global Landscape and Dynamics of Parkin and USP30-Dependent
Ubiquitylomes in iNeurons during Mitophagic Signaling. Molecular cell *77*, 1124-1142 e1110.
10.1016/j.molcel.2019.11.013.

48.   Nguyen, D., Bharat, V., Conradson, D.M., Nandakishore, P., and Wang, X. (2021). Miro1
Impairment in a Parkinson's At-Risk Cohort. Front Mol Neurosci *14*, 734273.
10.3389/fnmol.2021.734273.

49.   Nalls, M.A., Blauwendraat, C., Vallerga, C.L., Heilbron, K., Bandres-Ciga, S., Chang, D., Tan,
M., Kia, D.A., Noyce, A.J., Xue, A., et al. (2019). Identification of novel risk loci, causal
insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association
studies. The Lancet. Neurology *18*, 1091-1102. 10.1016/S1474-4422(19)30320-5.

50.   Papapetropoulos, S., Lee, M.S., Boyer, S., and Newbold, E.J. (2019). A Phase 2, Randomized,
Double-Blind, Placebo-Controlled Trial of CX-8998, a Selective Modulator of the T-Type
Calcium Channel in Inadequately Treated Moderate to Severe Essential Tremor: T-CALM Study
Design and Methodology for Efficacy Endpoint and Digital Biomarker Selection. Front Neurol
10, 597. 10.3389/fneur.2019.00597.

51.   Margaret S. Lee, S.P., Michelle S. Higgin, Muralikrishna Duvvuri, Bruce Rehlaender, Evan
NEWBOLD (2020) Treating essential tremor using (r)-2-(4-isopropylphenyl)-n-(1-(5-(2,2,2-
trifluoroethoxy)pyridin-2-yl)ethyl)acetamide. patent WO2020072773A1, 2020-04-09.

52.   Xiang, Z., Thompson, A.D., Brogan, J.T., Schulte, M.L., Melancon, B.J., Mi, D., Lewis, L.M.,
Zou, B., Yang, L., Morrison, R., et al. (2011). The Discovery and Characterization of ML218: A
Novel, Centrally Active T-Type Calcium Channel Inhibitor with Robust Effects in STN Neurons
and in a Rodent Model of Parkinson's Disease. ACS Chem Neurosci *2*, 730-742.
10.1021/cn200090z.

53.   Patel, R., Montagut-Bordas, C., and Dickenson, A.H. (2018). Calcium channel modulation as a
target in chronic pain control. Br J Pharmacol *175*, 2173-2184. 10.1111/bph.13789.

54.   Parkinson Study Group, S.-P.D.I.I.I.I. (2020). Isradipine Versus Placebo in Early Parkinson
1300   Disease: A Randomized Trial. Ann Intern Med *172*, 591-598. 10.7326/M19-2534.
1301   55.   Kelly, C.J., Couch, R.K., Ha, V.T., Bodart, C.M., Wu, J., Huff, S., Herrel, N.T., Kim, H.D.,
1302   Zimmermann, A.O., Shattuck, J., et al. (2023). Iron status influences mitochondrial disease
1303   progression in Complex I-deficient mice. eLife *12*. 10.7554/eLife.75825.
1304   56.   Hess, C.W., and Okun, M.S. (2016). Diagnosing Parkinson Disease. Continuum (Minneap Minn)
1305   *22*, 1047-1063. 10.1212/CON.0000000000000345.
1306   57.   Bartels, T. (2016). Conformation-Specific Detection of alpha-Synuclein: The Search for a
1307   Biomarker in Parkinson Disease. JAMA Neurol. 10.1001/jamaneurol.2016.4813.
1308   58.   Bharat, V., and Wang, X. (2020). Precision Neurology for Parkinson's Disease: Coupling Miro1-
1309   Based Diagnosis With Drug Discovery. Movement disorders : official journal of the Movement
1310   Disorder Society *35*, 1502-1508. 10.1002/mds.28194.
1311   59.   Choi, M.L., Chappard, A., Singh, B.P., Maclachlan, C., Rodrigues, M., Fedotova, E.I.,
1312   Berezhnov, A.V., De, S., Peddie, C.J., Athauda, D., et al. (2022). Pathological structural
1313   conversion of alpha-synuclein at the mitochondria induces neuronal toxicity. Nature
1314   neuroscience *25*, 1134-1148. 10.1038/s41593-022-01140-3.
1315   60.   Ramezani, M., Wagenknecht-Wiesner, A., Wang, T., Holowka, D.A., Eliezer, D., and Baird, B.A.
1316   (2023). Alpha Synuclein Modulates Mitochondrial Ca (2+) Uptake from ER During Cell
1317   Stimulation and Under Stress Conditions. bioRxiv. 10.1101/2023.04.23.537965.
1318   61.   Erustes, A.G., Guarache, G.C., Guedes, E.D.C., Leao, A., Pereira, G., and Smaili, S.S. (2022).
1319   alpha-Synuclein Interactions in Mitochondria-ER Contacts: A Possible Role in Parkinson's
1320   Disease. Contact (Thousand Oaks) *5*, 25152564221119347. 10.1177/25152564221119347.
1321   62.   Melo, T.Q., Copray, S., and Ferrari, M.F.R. (2018). Alpha-Synuclein Toxicity on Protein Quality
1322   Control, Mitochondria and Endoplasmic Reticulum. Neurochem Res *43*, 2212-2223.
1323   10.1007/s11064-018-2673-x.
1324   63.   Paillusson, S., Gomez-Suaga, P., Stoica, R., Little, D., Gissen, P., Devine, M.J., Noble, W.,
1325   Hanger, D.P., and Miller, C.C.J. (2017). alpha-Synuclein binds to the ER-mitochondria tethering
1326   protein VAPB to disrupt Ca(2+) homeostasis and mitochondrial ATP production. Acta
1327   Neuropathol *134*, 129-149. 10.1007/s00401-017-1704-z.
1328   64.   Devi, L., Raghavendran, V., Prabhu, B.M., Avadhani, N.G., and Anandatheerthavarada, H.K.
1329   (2008). Mitochondrial import and accumulation of alpha-synuclein impair complex I in human
1330   dopaminergic neuronal cultures and Parkinson disease brain. The Journal of biological chemistry
1331   *283*, 9089-9100. 10.1074/jbc.M710012200.
1332   65.   Kamer, K.J., Sancak, Y., Fomina, Y., Meisel, J.D., Chaudhuri, D., Grabarek, Z., and Mootha,
1333   V.K. (2018). MICU1 imparts the mitochondrial uniporter with the ability to discriminate
1334   between Ca(2+) and Mn(2+). Proceedings of the National Academy of Sciences of the United
1335   States of America *115*, E7960-E7969. 10.1073/pnas.1807811115.
1336   66.   Britti, E., Delaspre, F., Tamarit, J., and Ros, J. (2021). Calpain-Inhibitors Protect Frataxin-
1337   Deficient Dorsal Root Ganglia Neurons from Loss of Mitochondrial Na(+)/Ca(2+) Exchanger,
1338   NCLX, and Apoptosis. Neurochem Res *46*, 108-119. 10.1007/s11064-020-03020-3.
1339   67.   Clark, E.H., Vazquez de la Torre, A., Hoshikawa, T., and Briston, T. (2020). Targeting
1340   mitophagy in Parkinson's disease. The Journal of biological chemistry.
1341   10.1074/jbc.REV120.014294.
1342   68.   Llorens, J.V., Soriano, S., Calap-Quintana, P., Gonzalez-Cabo, P., and Molto, M.D. (2019). The
1343   Role of Iron in Friedreich's Ataxia: Insights From Studies in Human Tissues and Cellular and
1344   Animal Models. Front Neurosci *13*, 75. 10.3389/fnins.2019.00075.

69.   Devos, D., Labreuche, J., Rascol, O., Corvol, J.C., Duhamel, A., Guyon Delannoy, P., Poewe,
W., Compta, Y., Pavese, N., Ruzicka, E., et al. (2022). Trial of Deferiprone in Parkinson's
Disease. N Engl J Med *387*, 2045-2055. 10.1056/NEJMoa2209254.
70.   Chan, N.C., Salazar, A.M., Pham, A.H., Sweredoski, M.J., Kolawa, N.J., Graham, R.L., Hess, S.,
and Chan, D.C. (2011). Broad activation of the ubiquitin-proteasome system by Parkin is critical
for mitophagy. Human molecular genetics *20*, 1726-1737. 10.1093/hmg/ddr048.
71.   Konig, T., Nolte, H., Aaltonen, M.J., Tatsuta, T., Krols, M., Stroh, T., Langer, T., and McBride,
H.M. (2021). MIROs and DRP1 drive mitochondrial-derived vesicle biogenesis and promote
quality control. Nature cell biology. 10.1038/s41556-021-00798-4.
72.   Melentijevic, I., Toth, M.L., Arnold, M.L., Guasp, R.J., Harinath, G., Nguyen, K.C., Taub, D.,
Parker, J.A., Neri, C., Gabel, C.V., et al. (2017). C. elegans neurons jettison protein aggregates
and mitochondria under neurotoxic stress. Nature *542*, 367-371. 10.1038/nature21362.
73.   Davis, C.H., Kim, K.Y., Bushong, E.A., Mills, E.A., Boassa, D., Shih, T., Kinebuchi, M., Phan,
S., Zhou, Y., Bihlmeyer, N.A., et al. (2014). Transcellular degradation of axonal mitochondria.
Proceedings of the National Academy of Sciences of the United States of America *111*, 9633-
9638. 10.1073/pnas.1404651111.
74.   Saha, T., Dash, C., Jayabalan, R., Khiste, S., Kulkarni, A., Kurmi, K., Mondal, J., Majumder,
P.K., Bardia, A., Jang, H.L., and Sengupta, S. (2022). Intercellular nanotubes mediate
mitochondrial trafficking between cancer and immune cells. Nat Nanotechnol *17*, 98-106.
10.1038/s41565-021-01000-4.
75.   Rosina, M., Ceci, V., Turchi, R., Chuan, L., Borcherding, N., Sciarretta, F., Sanchez-Diaz, M.,
Tortolici, F., Karlinsey, K., Chiurchiu, V., et al. (2022). Ejection of damaged mitochondria and
their removal by macrophages ensure efficient thermogenesis in brown adipose tissue. Cell
Metab. 10.1016/j.cmet.2022.02.016.
76.   Ahmad, T., Mukherjee, S., Pattnaik, B., Kumar, M., Singh, S., Kumar, M., Rehman, R., Tiwari,
B.K., Jha, K.A., Barhanpurkar, A.P., et al. (2014). Miro1 regulates intercellular mitochondrial
transport & enhances mesenchymal stem cell rescue efficacy. The EMBO journal *33*, 994-1010.
10.1002/embj.201386030.
77.   Pacelli, C., Giguere, N., Bourque, M.J., Levesque, M., Slack, R.S., and Trudeau, L.E. (2015).
Elevated Mitochondrial Bioenergetics and Axonal Arborization Size Are Key Contributors to the
Vulnerability of Dopamine Neurons. Current biology : CB *25*, 2349-2360.
10.1016/j.cub.2015.07.050.
78.   Surmeier, D.J., Guzman, J.N., Sanchez-Padilla, J., and Goldberg, J.A. (2010). What causes the
death of dopaminergic neurons in Parkinson's disease? Prog Brain Res *183*, 59-77.
10.1016/S0079-6123(10)83004-3.
79.   Ambrosi, G., Ghezzi, C., Sepe, S., Milanese, C., Payan-Gomez, C., Bombardieri, C.R.,
Armentero, M.T., Zangaglia, R., Pacchetti, C., Mastroberardino, P.G., and Blandini, F. (2014).
Bioenergetic and proteolytic defects in fibroblasts from patients with sporadic Parkinson's
disease. Biochimica et biophysica acta *1842*, 1385-1394. 10.1016/j.bbadis.2014.05.008.
80.   Trinh, K., Moore, K., Wes, P.D., Muchowski, P.J., Dey, J., Andrews, L., and Pallanck, L.J.
(2008). Induction of the phase II detoxification pathway suppresses neuron loss in Drosophila
models of Parkinson's disease. The Journal of neuroscience : the official journal of the Society
for Neuroscience *28*, 465-472. 10.1523/JNEUROSCI.4778-07.2008.
81.   Tsai, P.I., Course, M.M., Lovas, J.R., Hsieh, C.H., Babic, M., Zinsmaier, K.E., and Wang, X.
(2014). PINK1-mediated phosphorylation of Miro inhibits synaptic growth and protects
dopaminergic neurons in Drosophila. Scientific reports *4*, 6962. 10.1038/srep06962.

82.   Castillo-Quan, J.I., Li, L., Kinghorn, K.J., Ivanov, D.K., Tain, L.S., Slack, C., Kerr, F., Nespital,
1392   T., Thornton, J., Hardy, J., et al. (2016). Lithium Promotes Longevity through GSK3/NRF2-
1393   Dependent Hormesis. Cell Rep *15*, 638-650. 10.1016/j.celrep.2016.03.041.
1394   83.   Sofola, O., Kerr, F., Rogers, I., Killick, R., Augustin, H., Gandy, C., Allen, M.J., Hardy, J.,
1395   Lovestone, S., and Partridge, L. (2010). Inhibition of GSK-3 ameliorates Abeta pathology in an
1396   adult-onset Drosophila model of Alzheimer's disease. PLoS genetics *6*, e1001087.
1397   10.1371/journal.pgen.1001087.
1398   84.   Byers, B., Cord, B., Nguyen, H.N., Schule, B., Fenno, L., Lee, P.C., Deisseroth, K., Langston,
1399   J.W., Pera, R.R., and Palmer, T.D. (2011). SNCA triplication Parkinson's patient's iPSC-derived
1400   DA neurons accumulate alpha-synuclein and are susceptible to oxidative stress. PloS one *6*,
1401   e26159. 10.1371/journal.pone.0026159.
1402   85.   Kriks, S., Shim, J.W., Piao, J., Ganat, Y.M., Wakeman, D.R., Xie, Z., Carrillo-Reid, L., Auyeung,
1403   G., Antonacci, C., Buch, A., et al. (2011). Dopamine neurons derived from human ES cells
1404   efficiently engraft in animal models of Parkinson's disease. Nature *480*, 547-551.
1405   10.1038/nature10648.
1406   86.   Wichterle, H., Lieberam, I., Porter, J.A., and Jessell, T.M. (2002). Directed differentiation of
1407   embryonic stem cells into motor neurons. Cell *110*, 385-397.
1408   87.   Nguyen, H.N., Byers, B., Cord, B., Shcheglovitov, A., Byrne, J., Gujar, P., Kee, K., Schule, B.,
1409   Dolmetsch, R.E., Langston, W., et al. (2011). LRRK2 mutant iPSC-derived DA neurons
1410   demonstrate increased susceptibility to oxidative stress. Cell stem cell *8*, 267-280.
1411   10.1016/j.stem.2011.01.013.
1412   88.   Sanders, L.H., Laganiere, J., Cooper, O., Mak, S.K., Vu, B.J., Huang, Y.A., Paschon, D.E.,
1413   Vangipuram, M., Sundararajan, R., Urnov, F.D., et al. (2014). LRRK2 mutations cause
1414   mitochondrial DNA damage in iPSC-derived neural cells from Parkinson's disease patients:
1415   reversal by gene correction. Neurobiology of disease *62*, 381-386. 10.1016/j.nbd.2013.10.013.
1416   89.   Hiroyuki Katayama, H.H., Koji Nagasawa, Hiroshi Kurokawa, Mayu Sugiyama, Ryoko
1417   Ando,Masaaki Funata, Nobuyo Yoshida, Misaki Homma, Takanori Nishimura, Megumu
1418   Takahashi, Yoko Ishida, Hiroyuki Hioki, Yoshiyuki Tsujihata, and Atsushi Miyawaki (2020).
1419   Visualizing and Modulating Mitophagy for Therapeutic Studies of Neurodegeneration. Cell *181*,
1420   12. https://doi.org/10.1016/j.cell.2020.04.025.
1421   90.   Hoshino, A., Wang, W.J., Wada, S., McDermott-Roe, C., Evans, C.S., Gosis, B., Morley, M.P.,
1422   Rathi, K.S., Li, J., Li, K., et al. (2019). The ADP/ATP translocase drives mitophagy independent
1423   of nucleotide exchange. Nature *575*, 375-379. 10.1038/s41586-019-1667-4.
1424   91.   Ehrenkaufer, G.M., Suresh, S., Solow-Cordero, D., and Singh, U. (2018). High-Throughput
1425   Screening of Entamoeba Identifies Compounds Which Target Both Life Cycle Stages and Which
1426   Are Effective Against Metronidazole Resistant Parasites. Front Cell Infect Microbiol *8*, 276.
1427   10.3389/fcimb.2018.00276.
1428   92.   Percha, B., and Altman, R.B. (2018). A global network of biomedical relationships derived from
1429   text. Bioinformatics *34*, 2614-2624. 10.1093/bioinformatics/bty114.
1430   93.   Rival, T., Soustelle, L., Strambi, C., Besson, M.T., Iche, M., and Birman, S. (2004). Decreasing
1431   glutamate buffering capacity triggers oxidative stress and neuropil degeneration in the
1432   Drosophila brain. Current biology : CB *14*, 599-605. 10.1016/j.cub.2004.03.039.
1433   94.   Kinghorn, K.J., Castillo-Quan, J.I., Bartolome, F., Angelova, P.R., Li, L., Pope, S., Cocheme,
1434   H.M., Khan, S., Asghari, S., Bhatia, K.P., et al. (2015). Loss of PLA2G6 leads to elevated
1435   mitochondrial lipid peroxidation and mitochondrial dysfunction. Brain : a journal of neurology
1436   *138*, 1801-1816. 10.1093/brain/awv132.

95. Consortium, G.T., Laboratory, D.A., Coordinating Center -Analysis Working, G., Statistical
Methods groups-Analysis Working, G., Enhancing, G.g., Fund, N.I.H.C., Nih/Nci, Nih/Nhgri,
Nih/Nimh, Nih/Nida, et al. (2017). Genetic effects on gene expression across human tissues.
Nature *550*, 204-213. 10.1038/nature24277.
96. Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J., and Kircher, M. (2019). CADD: predicting
the deleteriousness of variants throughout the human genome. Nucleic Acids Res *47*, D886-
D894. 10.1093/nar/gky1016.
97. Nalls, M.A., Keller, M.F., Hernandez, D.G., Chen, L., Stone, D.J., Singleton, A.B., and
Parkinson's Progression Marker Initiative, i. (2016). Baseline genetic associations in the
Parkinson's Progression Markers Initiative (PPMI). Movement disorders : official journal of the
Movement Disorder Society *31*, 79-85. 10.1002/mds.26374.
98. Le Guen, Y., Napolioni, V., Belloy, M.E., Yu, E., Krohn, L., Ruskey, J.A., Gan-Or, Z., Kennedy,
G., Eger, S.J., and Greicius, M.D. (2021). Common X-Chromosome Variants Are Associated
with Parkinson Disease Risk. Ann Neurol *90*, 22-34. 10.1002/ana.26051.

1451

1452

1453

FIGURE LEGENDS

Figure 1. Iron Promotes MCU Oligomerization. (A-D) HEK cells were treated with 500 µM Fe^{2+} for

22 hours, stimulated with thrombin, and mitochondrial (Rhod-2) and cytosolic Ca^{2+} levels (Calcium

Green) were measured. (A, C) Representative traces of Ca^{2+} ions. (B, D) Quantification of the peak

fluorescent intensity normalized to the baseline within each cell. n=20 cells from 4 independent

coverslips. (E) HEK cells were treated with 5 mM Fe^{2+} for 22 hours, run in Native-PAGE, and blotted.

Right: Qualification of the band intensity normalized to the total protein amount measured by a

Bicinchoninic Acid (BCA) assay. Given that MCU oligomers appear between 400-700KDa this region

of the gel is quantified. Same for the rest of the figures. n=5 independent experiments. Please note that

EDTA was present in non-treated controls for all biochemistry experiments. (F) Similar to (E), but with

a range of Fe^{2+} concentrations added to the media. n=4 independent experiments. Data point shows

Mean±S.E.M. (G) Elution profiles of MCU from SEC samples (detected by Western blotting shown in

Figure S1B). (H) HEK cells were treated with 500 µM Co^{2+} for 22 hours and run in Native-PAGE.

Qualification of the band intensity normalized to the total protein amount measured by BCA. n=4

independent experiments. (I) HEK cells treated with 500 μM Co^{2+} for 22 hours were stimulated with

thrombin and mitochondrial Ca^{2+} levels (Rhod-2) were measured. The peak fluorescent intensity

normalized to the baseline is quantified. n=20 cells from 4 independent coverslips. (B, D, E, H, I) Two-

tailed Welch's T Test. (J-K) Fluorescence-detection SEC profiles of purified human MCU. Note that

MCU holocomplex was used for Ca^{2+}. The annotation of the peaks is based on [14,15]. See also Figure S1.

1473

Figure 2. MCU Binds to Fe^{2+}. (A) HEK cells were treated with or without 500 μM Fe^{2+} for 21 hours,

then IPed with anti-MCU, and Fe^{2+} concentrations in the IP samples were detected (normalized to the

total protein amount). Two-tailed paired T Test. n=4 independent experiments. (B) Different

concentrations of Fe^{2+} was added to the purified MCU complex in vitro and after wash the total iron

level in the complex was detected by ICP-MS. Mean±S.E.M. n=3. (Ci) Structural visualization of the

matrix domain of MCU (green) bound to Fe^{2+} (orange) via these sites (pink) generated by PyMol. (Cii)

MCU-/- HEK cells transfected as indicated were treated with 500 μM Fe^{2+} for 20 hours, then IPed with

anti-Flag, and Fe^{2+} concentrations in the IP samples were detected. Two-tailed paired T Test. n=4

independent experiments. (Ciii) ICP-MS measurement of the total iron level in IPed MCU-Flag from

cells as in (Cii), normalized to the total protein amount. n=3. (D) Representative blots of IP with anti-

Flag using cell lysates as indicated, run in Native- or SDS-PAGE. Bottom: Quantification of the band

intensity of MCU-Flag in Native-PAGE normalized to that in SDS-PAGE from the same experiment.

n=4 independent experiments. (E) *MCU-/-* HEK cells transfected as indicated and treated with or

without 500 μM Fe^{2+} for 22 hours were stimulated with thrombin and mitochondrial Ca^{2+} levels (Rhod-2)

were measured. Representative traces of Ca^{2+} ions. (F-G) Based on traces like in (E), the peak

fluorescent intensity normalized to baseline (F) or efflux rate (G) is quantified. n=17 cells from 4

independent coverslips. (H) Cells as in (D-E) were lysed and blotted. Quantification of NCLX or MCUb

band intensity normalized to that of tubulin from the same experiment. n=4 independent experiments.

(D-H) Two-tailed Welch's T Test. Relative to "MCU-WT" or "MCU-3A" without treatment.

1493

Figure 3. PD Models Mirror Fe^{2+}-Treated HEK Cells. (A) Postmortem brains were run in Native- or

SDS-PAGE and blotted. Blots are in Figure S3A. The MCU oligomer bands in Native-PAGE and the

NCLX and MCUb bands in SDS-PAGE (average of 3 replicates) are used in the plot. HC: healthy

control. (B) The total iron level (y axis) in each brain fraction was measured by ICP-MS (normalized to

wet weight, average of 3 repeats), and plotted with the band intensity of MCU oligomers in Native-

PAGE (x axis) from the same brain fraction. Linear fit. (C) iPSC-derived neurons from a PD patient

with *SNCA-A53T* and the isogenic control were stimulated with thrombin and mitochondrial (Rhod-2)

and cytosolic Ca^{2+} levels (Calcium Green) were measured in the somas due to the ease of signal

identification at baseline. Left: Representative traces of Ca^{2+} ions in neurons. Right: Quantification of

the peak fluorescent intensity normalized to the baseline within each cell body. n=15 cell bodies from 3

independent coverslips. (D) iPSC-derived neurons were lysed and blotted as indicated. The same

amount of total protein measured by BCA was loaded. Quantification of the band intensity normalized

to the total protein amount. n=4 independent experiments. (E) Left: Confocal single section images of

live neurons as in (C-D) stained with RPA (quenched by Fe^{2+}) and MitoTracker Green (Mito-Green), or

Mito-FerroGreen (stains Fe^{2+}). Scale bars: 10 μm. Middle: Quantification of the RPA fluorescent

intensity in each soma from inverted images normalized to MitoTracker Green, a membrane-potential-

independent dye whose intensity was similar between control and PD neurons (P=0.2524). n=26 (WT)

and 28 (PD) cell bodies from 3 coverslips. Right: Quantification of the Mito-FerroGreen fluorescent

intensity in each soma subtracting background. n=55 cell bodies from 3 coverslips. Both quantifications

reflect the mitochondrial Fe^{2+} levels. Two-tailed Welch's T Test for all panels. (C-E) Day 21-26 after

neural differentiation. See also Figure S3.

1515

Figure 4. Iron Functions Upstream of Calcium in PD neurons. (A) Similar to Figure 3, iPSC-derived

neurons, with or without treatment of 100 μM DFP for 24 hours, were stimulated with thrombin, and

mitochondrial Ca^{2+} (Rhod-2) was measured. Quantification of the peak fluorescent intensity normalized

to the baseline within each cell body. n=15 cell bodies from 3 independent coverslips. Control data

without DFP treatment are the same as in Figure 3C. (B) The DA neuron number was counted in the

PPL1 cluster of flies with indicated genotypes and conditions. Drug treatment was started from

adulthood (day 1). In the images, each circular structure is one DA neuron cell body stained with anti-

TH. The number is averaged between both sides of the brain. Scale bar: 20 μm. n=6, 9, 8, 7 (from left to

right) flies. (C) The Performance Index was measured in flies. Drug treatment was started from

embryogenesis. n=35, 33, 40, 34 flies (from left to right), 3 independent experiments. (D) Representative

confocal images of neurites (where mitochondrial morphology is easier to discern) transfected with

MCU-Flag and Mito-dsRed. Scale bar: 10 μm. Quantification of the percentage of total neurons with

deformed mitochondria (defined as a neuron without any mitochondrion in neurites >2.5 μm in length).

n=4 coverslips. (A, D) One-Way Anova Post Hoc Tukey Test. (B-C) Two-Way Anova Post Hoc Tukey

Test. (E) Schematic representation of iron triggered mitochondrial calcium influx in PD. IMM: inner

mitochondrial membrane. IMS: mitochondrial intermembrane space. (F) A *PD-SNCA-A53T* iPSC-

derived neuron was transfected as indicated and immunostained with anti-Myc. A neurite is shown for

better visualizing colocalization. Scale bar: 10 μm. (G) Confocal live images of neurons as in (D)

transfected with miRFP670_P2A_GCaMP6f-Miro1-Anchor or miRFP670_P2A_ER-GCaMP6f. Scale

bar: 10 μm. Quantification of the intensity of GCaMP6f normalized to miRFP670 within the same cell

body. n=15 cell bodies from 3 coverslips. Two-tailed Welch's T Test. See also Figure S3.

1537

Figure 5. Miro Senses Calcium to Mediate Several PD Relevant Phenotypes. (A) Representative still

images from live Mito-dsRed and GFP-Miro1 recordings in axons of indicated genotypes, following 100

μM Antimycin A treatment. Scale bar: 10 μm. Mitochondria are more sparsely distributed in axons than

in somas, allowing better visualization of degradation. This dose of Antimycin A was selected to allow

rapid Miro1 and mitochondrial degradation, needed for live imaging to minimize laser scanning time

[33,35]. (B) Degradation rate profiles of the GFP-Miro1 intensity following 100 μM Antimycin A treatment

normalized to the same axonal region at 0 min. (C) Degradation rate profiles of the Mito-dsRed intensity

following 100 μM Antimycin A treatment normalized to the same axonal region at 0 min. (B-C) n=5

axons (one axon per coverslip). Comparison of values at 25 min or 58 min with those at 0 min within the

same condition. In control axons, GFP-Miro1 was degraded faster (significant at 25 min) than Mito-

dsRed (significant at 58 min). One-Way Anova Post Hoc Dunnett's Test. $p < 0.0001$ if comparing "PD,

Miro1-WT" with all the other 3 conditions with Two-Way Anova Post Hoc Tukey Test. (D) iPSC-

derived neurons were transfected as indicated and GFP-Miro1 was detected by an ELISA normalized to

the total protein amount. n=4. Two-tailed Welch's T Test compared between without and with treatment

within each condition. (E) The DA neuron number was counted in the PPL1 cluster of flies with

indicated genotypes as described in Figure 4. Scale bar: 20 μm. n=7, 4, 6, 5 (from left to right). (F) The

Performance Index was measured in flies with indicated genotypes and age. n (from left to right)=49, 47,

39, 47 (day 14); 48, 45, 37, 44 (day 20); 3 independent experiments. (E-F) Two-Way Anova Post Hoc

Tukey Test. See also Figure S3.

1557

Figure 6. HTP Screens Identify Ca^{2+}-Related Drug Hits for PD. (A) Schematic representation of a

custom-designed drug screen for Miro1 in PD fibroblasts. (B) Pathway analysis identified calcium as a

shared factor in the primary hit-Miro1 network. Each individual pathway is generated using a primary

hit and Miro1 as search query and the resulting subnetwork is visualized and curated using docs2graph–

a knowledge-graph browser. The visualization shows a subgraph generated by docs2graph from the

collection of curated supporting documents for each pathway. See also Figure S4-5.

1564

Figure 7. Benidipine Rescues PD Relevant Phenotypes. (A-B) iPSC-derived neurons from a PD

patient with *SNCA-A53T* and the isogenic control, treated as indicated, were immunostained with anti-

TH (A) or TUNEL and Dapi (B), and imaged under a confocal microscope. Scale bars: 100 µm. Below:

Quantification of the density of TH-positive neurons (A) or the percentage of TUNEL-positive neurons

(B). n=20 images from 3 independent coverslips. P values are compared with the far-left bar, except

indicated otherwise. (C) 40-day-old fly brains were immunostained with anti-TH and the DA neuron

number was counted in the PPL1 cluster as described in Figure 4. Scale bar: 20 µm. n=4, 6, 7, 4 (from

left to right). (D) The Performance Index was measured in 30-day-old flies, fed as indicated. n=59, 57,

54, 57 flies (from left to right), 3 independent experiments. (C-D) Drug treatment was started from

adulthood (day 1). (A-B) One-Way Anova Post Hoc Tukey Test. (C-D) Two-Way Anova Post Hoc

Tukey Test. (E) Miro1 protein levels were measured using ELISA in PBMCs treated with DMSO or

40 µM CCCP for 6 hours. Miro1 Ratio is calculated by dividing the Miro1 value treated with CCCP by

that with DMSO from the same subject. Dot plot with Mean±S.E.M. n=80 healthy controls and 107 PD.

Two-tailed Welch's T Test. (F) PBMCs from 4 PD patients were treated with 40 µM CCCP for 6 hours,

or pretreated with 10 µM Benidipine or MR3 for 18 hours and then with 40 µM CCCP for another 6

hours, and Miro1 protein was detected using ELISA. Patient IDs are the same as in Table S4. Two-tailed

paired T Test. See Method for more details. (G) Schematic representation of the iron-calcium-Miro axis

discovered in this study. See also Figure S6-7.

1583
1584

1585
1586
1587
1588