The neuropeptidergic connectome of *C. elegans*

Lidia Ripoll-Sánchez[1,2], Jan Watteyne[3], HaoSheng Sun[4,5], Robert Fernandez[4], Seth R Taylor[6],

Alexis Weinreb[7], Mark Hammarlund[7], David M Miller III[6], Oliver Hobert[4], Isabel Beets[3,†],

Petra E Vértes[2,†], William R Schafer[1,3,†,#*]

1 - Neurobiology Division, MRC Laboratory of Molecular Biology, Cambridge, UK

2 - Department of Psychiatry, Cambridge University, Cambridge, UK

3 - Division of Animal Physiology and Neurobiology, KU Leuven, Leuven, Belgium

4 - Department of Biological Sciences, Howard Hughes Medical Institute, Columbia

University, New York, NY, USA

5 - Department of Cell, Developmental, and Integrative Biology, University of Alabama at

Birmingham, AL, USA

6 - Department of Cell and Developmental Biology, Vanderbilt University School of

Medicine, Nashville, TN, USA

7 - Department of Genetics and Department of Neurobiology, Yale University School of

Medicine, New Haven, CT, USA

† - Senior author

- Lead contact

Correspondence: wschafer@mrc-lmb.cam.ac.uk

Keywords

Neuropeptides, connectomics, networks, neuromodulation, *C. elegans*

Summary

Efforts are currently ongoing to map synaptic wiring diagrams or connectomes in order to understand the neural basis of brain function. However, chemical synapses represent only one type of functionally important neuronal connection; in particular, extrasynaptic, "wireless" signaling by neuropeptides is widespread and plays essential roles in all nervous systems. By integrating single-cell anatomical and gene expression datasets with a biochemical analysis of receptor-ligand interactions, we have generated a draft connectome of neuropeptide signaling in the *C. elegans* nervous system. This connectome is characterized by a high connection density, extended signaling cascades, autocrine foci, and a decentralized topology, with a large, highly interconnected core containing three constituent communities sharing similar patterns of input connectivity. Intriguingly, several of the most important nodes in this connectome are little-studied neurons that are specialized for peptidergic neuromodulation. We anticipate that the *C. elegans* neuropeptidergic connectome will serve as a prototype to understand basic organizational principles of neuroendocrine signaling networks.

Introduction

Understanding how behavior arises from neuronal interactions in the brain is one of the great challenges of modern neuroscience. In recent years, many projects have started to map the synaptic wiring diagrams (or synaptic connectomes) of simple and complex nervous systems to define their architecture at the cellular level[1–4]. Specifically, connectomics has aimed to identify all the individual units of the nervous system (i.e., its neurons) as well as all the connections between these units. In practice this involves reconstructing volumes of brain tissue using electron microscopy (EM), and then analyzing these images to trace neuronal processes and identify pre- and post-synaptic hallmarks of chemical synapses between identified neurons. Currently there are several endeavors of this type that are compiling complete connectomes for organisms important for neuroscience research, including *Drosophila*, *Platynereis*, *Ciona,* zebrafish, and mouse[2,3,5–7].

Although most connectomics research focuses on synaptic connectivity, chemical synapses are only one of several paths through which neurons communicate in the brain. For example, many important interactions between neurons involve volume transmission, by which secreted molecules are released extrasynaptically and activate receptors on neurons that are not wired through synapses or gap junctions to the releasing neuron. Unlike synaptic and gap junction transmission, where signaling is restricted to neurons on either side of the synapse, volume transmission can mediate signaling across distances of microns[8–10]. Such extrasynaptic signaling has been described for classical neurotransmitters as well as monoamines, and is particularly prevalent for neuropeptides, which are released from dense core vesicles outside the synaptic active zone and act on a longer temporal and spatial scale[10,11]. Although these extrasynaptic interactions are largely independent from the synaptic wiring, they play key roles in neural circuits and are thus critical for understanding the neural basis of behavior[12–18].

Neuropeptides carry out particularly complex and important functions in the brains of all organisms[19–21]. Neuropeptides are thought to be among the most ancient neuronal signaling molecules, and their origin may in fact predate the actual evolution of neurons[22,23]. Biologically-active peptides

are usually formed from larger genetically-encoded precursors, which are enzymatically processed

through proteolysis and/or chemical modification[24]. Although some neuropeptides have been shown

to activate ion channels or transmembrane kinases, they typically bind to G protein-coupled receptors

(GPCRs), which act through second messenger pathways to modulate diverse aspects of neuronal

physiology[25]. Neuropeptide-activated GPCRs fall into two main classes, the rhodopsin-like (Class A)

and secretin-like (Class B) families, and within these groups many receptors (e.g.

oxytocin/vasopressin, neuropeptide Y/F, neuromedin U, somatostatin) show significant conservation

across widely divergent animal phyla[26,27]. Neuropeptide systems also play broadly conserved roles in

the control of behavioral states, including those involved in feeding, sleep and arousal, reproductive

behavior, and learning[14,28–33]. In humans, the role of neuropeptides as neuromodulators has made this

signaling system a highly sought-after target for new neuropsychiatric treatments. Currently 50 drugs

acting via GPCRs have been approved by the FDA[34], including several neuropeptide-GPCRs such as:

orexin antagonists for treatment of insomnia[35], a substance P antagonist for treatment of

chemotherapy-induced nausea[36]. Besides new treatments are being developed like a GIP agonist as

treatment for Alzheimer and Parkinson's[37]. These promising therapies and the large number of

remaining neuropeptide-GPCRs in humans[18,38] indicate the therapeutic potential of this molecular

system.

The diversity and extent of neuropeptide signaling implies that the pathways for peptidergic

communication can also be considered as a network. The genomes of all animals encode many, in

some cases hundreds of neuropeptides, along with a similarly large number of GPCRs[18,39–41].

Moreover, in contrast to monoamines, which are typically expressed in only a small subset of neurons

in the brain, neuropeptides are expressed extremely broadly; indeed, recent single-cell transcriptomic

studies indicate that most if not all neurons in the mouse cerebral cortex express one or more

neuropeptides as well as multiple neuropeptide-binding GPCRs[39,42]. These data imply that peptidergic

signaling underpins a dense and pervasive interaction network involving most if not all neurons of an

animal's nervous system[43,44]. Since neuropeptide signaling is thought to be mostly extrasynaptic, the

topology and structure of this wireless neuropeptide connectome is expected to be fundamentally

distinct from that of the wired synaptic connectome[45]. However, in most organisms there is

insufficient gene expression data mapped with single-cell resolution to identified neurons to map the

structure of these extrasynaptic networks.

The nematode *C. elegans* represents an attractive animal to investigate the structure and

organization of neuropeptide signaling networks due to its small and well-characterized nervous

system. *C. elegans* was the first and is currently still the only adult organism with a completely

mapped synaptic neuronal connectome, with each of its 302 neurons and approximately 2300 synaptic

connections identified through electron microscopy (EM) reconstructions[46–48]. Despite its small size,

the *C. elegans* nervous system has been found to share several structural features with that of larger

animals. For example, the *C. elegans* connectome, like those of larger nervous systems, exhibits a

small-world topology, with relatively high clustering paired with relatively short average path

lengths[49,50]. Likewise, the *C. elegans* nervous system is highly modular, with functionally segregated

local clusters of high within-group connectivity[51–54]. Finally, the worm connectome contains a small

number of highly connected hubs, which are interconnected in a core or rich club and facilitate

communication between modules[55]; similar rich club topology has been observed in bigger brains,

including the human cortex[56,57]. Shared structural features are also apparent at the microcircuit level;

for example, feed-forward motifs are significantly overrepresented in the nematode connectome, just

as in the mammalian cortex[58–60]. Thus, insights gained from analysis of neuropeptide signaling

networks in *C. elegans* may also reveal organizational principles that are conserved in larger brains.

Although the *C. elegans* nervous system is anatomically small, at the biochemical level its

neuropeptide signaling pathways show similar complexity and remarkable conservation with humans

and other larger-brained animals. Its genome contains 159 predicted neuropeptide precursor genes

(NPP) (including 40 insulin-like peptides) that produce over 300 different neuropeptides[24,61].

Likewise, approximately 150 *C. elegans* genes encode GPCRs known or predicted to be activated by

neuropeptides[25,62]. These numbers are of a similar order of magnitude to the numbers of neuropeptides

and peptide activated GPCRs in the human genome[18,63]. Each *C. elegans* neuron not only expresses at

least one neuropeptide and at least one neuropeptide receptor, as do vertebrate neurons,[18,39,40,42,43] but

each *C. elegans* neuron class expresses a unique combination of neuropeptide-encoding genes[40], a

notion that illustrates the tremendous potential complexity of neuropeptide signaling. Many *C.*

elegans neuropeptides and cognate receptors are phylogenetically conserved across animal phyla and

have clear human homologues, with some families (such as the RFamide peptides) having undergone

expansion in the nematode lineage[64]. Thus, neuropeptide signaling in nematodes shows surprising

conservation as well as similar diversity to neuropeptide signaling in the human brain, despite vast

differences in neuron number and anatomical complexity.

Here we present a draft connectome of neuropeptidergic signaling in *C. elegans,* built by

integrating gene expression, ligand-receptor interaction, and anatomical datasets. We identified

predicted neuron-to-neuron signaling interactions mediated by 91 neuropeptide-receptor pairs, and by

aggregating these individual neuropeptide-signaling networks generated a comprehensive network of

peptidergic signaling in the *C. elegans* nervous system. This resulting neuropeptidergic connectome

differs significantly in its structure from the previously-characterized wired connectome[47]; for example,

it is denser, contains distinct hubs, high-weight edges, and autocrine signaling foci, and has a

decentralized topology allowing direct communication among a large fraction of neurons. We expect

this nematode neuropeptide connectome will serve as a prototype for understanding the organization of

peptidergic signaling and how it interacts with the wired neural circuitry in other organisms, including

those with much larger brains.

Results

Construction of a neuropeptidergic connectome

To generate a draft neuropeptide connectome, we integrated information from three datasets to infer

potential pathways for neuropeptide signaling between individual *C. elegans* neurons (Figure 1A). To

identify biologically relevant molecular interactions between individual neuropeptides and receptors,

we used data from a large-scale reverse pharmacology screen which tested 55,386 potential

neuropeptide-receptor interactions *in vitro* and identified 461 peptide-GPCR couples encoded by 55

neuropeptide and 57 GPCR genes[40]. To determine which neurons express each neuropeptide and

receptor, we used publicly available single-neuron transcriptome data from the CeNGEN project, which

described single-cell RNA-sequencing transcription profiles of all predicted neuropeptide and peptide-

activated GPCR genes in *C. elegans*[40]. We validated these scRNA datasets with several CRISPR/Cas9-

engineered reporter alleles of ligands and receptors. Finally, we used previously published EM

anatomical reconstructions to assess physical constraints on potential diffusion of neuropeptides to their

target receptors[48,65,66]. In constructing the neuropeptide network, we consider that two neurons (nodes)

are connected by a directed edge if the first neuron expresses a particular neuropeptide ligand and the

second expresses a paired receptor, subject to spatial constraints on signaling (Figure 1B).

For each dataset, it was necessary to threshold for biologically relevant interactions.

For biochemical interactions, the raw dataset contained 461 neuropeptide-receptor pairs that showed

dose-dependent activation with a half-maximal effective concentration (EC_{50}) ranging between 0.1 pM

to 22 μM[64]. At the gene level, these pairs were encoded by 55 neuropeptide and 56 GPCR genes, with

148 unique gene-gene interactions. In assessing neuropeptide-receptor pairings we initially opted for a

conservative EC_{50} threshold of 500 nM, as many neuropeptide-receptor couples with EC_{50} values in this

range have been validated *in vivo*[14,25,28,32,62]. By this criterion, we defined 91 individual neuropeptide-

receptor gene couples, with a large number (51) of the predicted neuropeptide GPCRs having at least

one identified ligand. The ligand-receptor interactions were complex, with several peptides activating

more than one receptor (promiscuous receptors) and several receptors being activated by multiple

peptides encoded by distinct precursor genes (versatile peptides)[64].

We next sought to determine an appropriate threshold for the scRNA-based gene expression

dataset. The transcriptomic data from the CeNGEN project has been differentially thresholded based

on a ground-truth dataset of reliable gene expression from across the whole *C. elegans* nervous

system using fosmid or receptor tagged reporters[40]. Each threshold was given a false discovery rate

and a true positive rate depending on the number of correctly identified cells for a given gene by the

scRNAseq analysis[40]. To determine the most appropriate threshold for our analysis, we obtained

single-copy genomic knock-in reporters for 17 representative neuropeptide precursor genes and 9

representative genes for neuropeptide receptors and characterized their expression patterns

comprehensively using the NeuroPAL marker strain (Figure 2A, B; Supplementary Figure 1). For

both we found that threshold 4, although on some occasions overly stringent, was a good

approximation of the expression pattern (Figure 2; Supplemental Table 1-3). Thus, although this

stringent threshold has lower discovery power and thus may undercount the number of neurons

expressing each gene, its stringency minimizes the likelihood that our resulting network will contain

edges that do not represent authentic paths for potential neuropeptide signaling.

Evaluating spatial constraints on neuropeptide signaling

These biochemical and gene expression datasets allowed us to infer which neurons express

neuropeptide and receptor genes that could mediate a neuromodulatory interaction; however, the

physics of diffusion and the neuroanatomy of the animal might potentially constrain some of these

interactions *in vivo*. We therefore considered several possible models of the neuropeptidergic network

based on different anatomical and spatial constraints (Figure 3B, C). We used EM reconstruction data

for each individual neuron in the *C. elegans* nervous system to identify its anatomical location (Figure

3A, Supplementary Table 4) and the neuropil bundles into which their axons and dendrites

project[48,65,66]. In the first and most permissive model, long-range signaling is permitted and

neuropeptidergic connections can take place between any neurons in the system. In the second ("mid-

range") model, neuropeptidergic connections can occur only between neuronal processes in the same

anatomical area, such as the head (including pharynx), tail and midbody (Figure 3A). In the third

("short-range") model, we only considered potential peptidergic connections between neurons in the

same neuronal process bundle, that is, between neurons whose processes overlap in the nerve ring,

ventral or dorsal cord, or an auxiliary nerve. Finally, it has recently been shown that the nerve ring,

into which more than half the neurons (168) project, can be divided into four layers or strata based on

patterns of physical contact[67,68]. We therefore also considered a fourth model in which neuropeptide

signaling inside the nerve ring is constrained within individual strata and thus between neurons that

make physical contact.

To evaluate each model, we investigated whether the expression patterns of receptors and

ligands were consistent with neuropeptide signaling being restricted as described above (Figure 3B,

C). For example, if neuropeptide signaling occurred only between neurons in physical contact (as in

model 4) we would expect neurons expressing a given receptor to project to the same nerve ring strata

as neurons expressing its ligand. To simplify this analysis, we focused on the neuropeptide-receptor

couples in which the receptor has only one ligand and the neuropeptide only binds to that receptor,

assessing whether, under the constraints of each model, there would be receptor-expressing neurons

that were unable to receive a signal from any neurons expressing its ligand. Conversely, we also

analyzed couples where the ligand activated only one receptor, asking whether under a given model

there were ligand-expressing neurons that could not communicate with cells expressing its receptor.

This analysis appeared to argue against the most restrictive model, in which ligands could not

diffuse between nerve ring strata. Specifically, we observed many examples in which a neuropeptide

receptor was expressed only in strata that did not express its specific ligand (Figure 3B, C, orange

points). For instance, while the *capa-1* neuropeptide precursor gene is expressed only in the ASG

sensory neuron, which projects its axon into the fourth nerve ring stratum, the gene for its receptor

NMUR-1 is expressed in all four strata[32,69](Figure 3B, C). Thus, NMUR-1 receptors in strata 1-3 must

be activated by CAPA-1 peptides that diffuse from their release site in stratum 4. Overall, 37 of the 41

receptors analyzed were expressed in at least one neuron making no contact with a neuron expressing

its ligand. Likewise, 32 of the neuropeptide precursor genes were expressed in at least one neuron

making no contact with neurons expressing its receptor (Figure 3C, yellow points). Since the nerve

ring strata are not separated by glial or other barriers, and experimental evidence[14,32] indicates

neuropeptides can indeed travel between strata, it is reasonable to infer that neuropeptides are not

likely to be constrained by anatomical layers in the nerve ring or other neuronal bundles.

We likewise observed cases in which the expression of a neuropeptide or receptor gene

implies mid-range signaling between distinct nerve bundles in the same body region (Figure 3B, C).

For example, the *frpr-7* receptor gene is expressed in multiple pharyngeal neurons, while its ligand

FLP-1 is released exclusively from the AVK neuron whose processes lie in the nerve ring and ventral

cord[13]. Thus, FRPR-7 receptors in the pharynx must be activated by peptides released by AVK in the

nerve ring, consistent with published evidence that neuropeptides can signal between head neurons

and pharyngeal neurons[13,70]. Indeed, for a majority (14/22) of the receptors expressed in pharyngeal

neurons, their ligand was expressed only outside the pharynx (Figure 3B, red points). Likewise,

receptors such as NPR-3 and TRHR-1 are expressed in the CAN neurons, yet their ligands FLP-15

and NLP-54 respectively are not expressed in any neurons projecting into the canal-associated nerve[71]

(Figure 3B, C, CAN neurons labelled). Thus, we hypothesize that at least some neuropeptide

signaling occurs between different nerve bundles, in particular between the pharynx and the nerve

ring and between CAN and other nerves in the body. We therefore focused our subsequent analysis on

model 2, consisting only of stringently selected short-range interactions, and model 3, which includes

mid-range signaling between nerve bundles in the same body region.

Neuropeptide networks exhibit diverse topologies

Based on these criteria, we first constructed network graphs between neuropeptide-expressing

and receptor-expressing neurons for each of the 91 individual neuropeptide/receptor couples in our

dataset (Supplementary Figure 2-3). These networks were filtered by removing edges between

neurons that did not project an axon or dendrite into the same process bundle (for the short-range

network) or into the same body region (for the mid-range network). In their short-range version, 78 of

these ligand-receptor couples formed a single connected network, whereas 13 formed networks with

two or even three disconnected components (Figure 4B, Supplementary Figure 2). In their mid-range

versions, all 91 couples formed a single connected network (Supplementary Figure 3).

We observed a diverse range of topologies in the networks for individual peptide-receptor

couples. One topological measure in which the different networks varied was their assortativity, or the

extent to which nodes are preferentially connected to nodes of similar degree. Using the conservative

short-range model, we observed that most of the networks, including many of those highly conserved

in evolution, were local networks, in which both ligand and receptor were expressed in relatively

restricted sets of neurons (Figure 4A, Supplementary Figure 2). The neurons in these networks

showed relatively low average degree and often encompassed only a subregion of the total nervous

system. In addition, we found 8 highly disassortative *integrative* networks, with many low-degree

peptide-releasing neurons signaling to relatively few high-degree receptor-expressing neurons (Figure

4A). We also observed 23 disassortative *broadcasting* networks that are characterized by a small

number of high-degree peptide-releasing neurons that signal to many low-degree receiving neurons

with broadly expressed receptors (Figure 4A, Supplementary Figure 2). Interestingly, both integrative

and broadcasting networks involved promiscuous receptors with multiple ligands, specifically FRPR-

8, EGL-6, DMSR-7, and DMSR-1. DMSR-7 and DMSR-1 also figured prominently in a fourth

category of more assortative, pervasive networks, in which both the ligand and receptor show broad

expression and most neurons thus show relatively high degree (Figure 4A, B). Relaxing the spatial

restrictions on our model had relatively modest impact on network topology, with four networks that

are local in the short-range model becoming either integrative (one network) or broadcasting (3

networks) in the mid-range model (Supplementary Figure 3). Thus, the topologies of neuropeptide

networks appear relatively robust to our assumptions about the spatial scope of neuropeptidergic

signaling.

The neuropeptide connectome is a decentralized, dense network

By aggregating the networks from the individual neuropeptide-receptor couples, we next

compiled complete neuropeptide connectome networks based on short-range or mid-range signaling

(Figure 5). Even for the more conservative short-range network, in which signaling was restricted to

neurons with processes overlapping in the same bundle, the density of connections was remarkably

high, with more than a third of all possible connections or edges (0.3437) present in the network.

Allowing mid-range connections (0.4429) or long-range connections (0.5929) increased this density

further (Supplementary Figure 4A, B). By comparison, the *C. elegans* synaptic (0.0251) and

monoamine (0.0236) networks were far less dense, fourteen-fold lower than the short-range

neuropeptide network. We also computed edge weights for the connections in the neuropeptide

networks based to the number of neuropeptide-receptor pairs capable of signaling between two nodes.

We observed that a large number (35% in the short-range network) of neuron pairs were connected by

only a single neuropeptide-receptor interaction, but some (9%) were connected by 6 or more different

peptide-receptor couples (Figure 5). In the most extreme case, we observed that two neurons, the

AVD premotor interneurons and the PQR oxygen-sensing neurons, were linked by 18 different

neuropeptide-receptor pairs, suggesting extraordinarily complex patterns of signaling could exist

between these cells (Figure 5, highlighted). Interestingly many other high-weight, biochemically

complex connections occurred between neurons of the oxygen sensing circuit and were mediated by a

common set of promiscuous receptors (DMSR-1, DMSR-7 and FRPR-8) and versatile neuropeptides

(FLP-4, FLP-9, FLP-10 and FLP-13). These neurons are thus connected by multiple neuropeptidergic

channels of communication, potentially allowing complex regulation by context and experience.

A striking feature of the midrange network in particular was its integration of neurons

disconnected from the synaptic connectome. For example, the 14 neuron classes of the pharyngeal

nervous system form a heavily synaptically interconnected, functionally autonomous network akin to

vertebrate enteric nervous systems, which is topologically isolated from the rest of the nervous

system[65,72,73]. We find that unlike the wired connectome, there are no neuropeptidergic networks that

define circuitry exclusively in the pharynx; to the contrary, pharyngeal neurons are fully integrated

into the somatic nervous system via strong reciprocal interconnectivity (Figure 5). All pharyngeal

neuron classes receive 90% or more of their incoming connections from outside the pharynx. Some

classes of pharyngeal neurons also broadcast extensively to the somatic nervous system, with several

(I1, I3, I4, I5, M5 and NSM) having more than 100 outgoing connections (more than 90% of their

total) to non-pharyngeal neurons. Likewise, it is notable that the CAN neurons, which completely lack

chemical synapses, show strong and reciprocal neuropeptidergic connectivity with the rest of the

nervous system, indicating that this unusual neuron class is well embedded in the neural network

(Figure 5, CAN highlighted).

We next investigated topological features of the aggregate neuropeptide network. In

particular, we analyzed the degree nodes of the network, defined as the number of incoming (in-

degree) and outgoing (out-degree) connections made by each neuronal node. Degree is often an

indicator of functional significance in networks, since high-degree nodes or hubs often play key

functional roles in the brain and other complex systems. As expected from the high density of

neuropeptide signaling, the average degree of both the short-range (Figure 6A) and mid-range

networks (Figure 6B) were significantly higher than for the previously characterized synaptic (Figure

6C) and monoamine (Figure 6D) networks. Moreover, when we analyzed the distribution of degree

across the neuropeptide networks, we observed relatively flat curves with many high-degree nodes;

more than half the neurons in the short-range network and nearly two thirds in the mid-range network

had a degree higher than 200, indicating that they have in and out connections with more than a third

of all other neurons (Figure 6A, B). In contrast, the synaptic network was much more centralized,

with only 10 neurons having degree greater than 50 (Figure 6C); similarly, the monoamine network

had only 18 high (>50) degree neurons (Figure 6D). The best-connected neurons in the neuropeptide

network have exceptionally high in-degree as well as out-degree, indicating that their status as hubs

depends as much on incoming connections as outgoing connections, and as much on their expression

of broadly-signaling neuropeptides as on integrating GPCRs (Figure 6A, B); this contrasts with the

monoamine network, where the hubs are exclusively monoamine-releasing neurons with a high out-

degree (Figure 6D).

The neuropeptide connectome contains a highly connected core with unique peptidergic hubs

A significant feature of many networks is the so-called rich club property. In such networks,

the most highly connected hubs form more connections between themselves than expected based on

their high degree alone; these hubs therefore comprise a rich club or core of the network that

facilitates communication between more peripheral network nodes. The *C. elegans* synaptic

connectome, for example, contains a rich club consisting of 11 premotor interneurons that appear to

play important roles in driving global brain states[55,74]. Remarkably, we found that the neuropeptide

connectome also shows the rich club property, but its rich club consists of 156 neurons (166 in mid-

range network), more than half of the nervous system (Figure 6F). Within the rich club the density of

connections is 0.6834 (p<0.00001), more than double the density of the overall network (0.3427).

Interestingly, compared to the wired synaptic connectome, the neuropeptide connectome as a whole

has significantly higher clustering and reciprocity (i.e. neurons are more likely to connect in both

directions) but lower modularity and disassortativity. This finding further supports the notion of a

decentralized neuropeptide connectome, with a broad-based densely-connected core allowing direct

communication between a large fraction of neuron pairs.

To further investigate the relationship between neuropeptide and synaptic signaling, we

looked at the correlation between synaptic and neuropeptidergic degree (Figure 6G). We observed

that neuropeptide degree and synaptic degree were positively correlated ($p < 0.0001$, $r = 0.53$). We

further observed that neurons with a high synaptic degree were among the most important

neuropeptidergic hubs; for example, all 11 members of the synaptic rich club (DVA, PVCL/R,

AVAL/R, AVBL/R, AVDL/R, AVEL/R) were among the 25 highest-degree nodes in the short-range

neuropeptide network (Figure 6G). However, there were also several neurons of very high degree in

the neuropeptide connectome but unexceptional synaptic degree (Figure 6G); these neurons can be

described as specialized for neuropeptide signaling. Six neurons (AVKL/R, PVQL/R, PVT and PVR)

had higher short-range neuropeptide degree than any of the synaptic rich club neurons (Figure 6A); if

mid-range signaling is considered these same six neurons remain the highest-degree hubs (Figure 6B).

Interestingly, the AVKL/R and PVT neurons are notable for expressing no classical neurotransmitters

or monoamines[75], while the PVQL/R neurons appear anatomically specialized for neuropeptide

signaling due to a preponderance of dense core vesicles[14,76]. AVKL/R, and PVT have been previously

linked to the control of global behavioral states related to sleep and arousal[14,73], but the functions of

the other neuropeptide hub neurons PVQ and PVR are not well-characterized.

The neuropeptide connectome core has a defined mesoscale structure

To further probe the structure of the neuropeptide connectome we investigated whether the

network contained modules or other forms of mesoscale substructure. We first applied standard

methods for modular decomposition, but the high density of the network precluded the identification

of any discrete modules, unless we aggressively filtered out lower-weight edges where neurons were

connected only by a small number of neuropeptide-receptor interactions. However, we hypothesized

that other types of meso-scale structure may be present[52,77]. We wondered for example whether we

could identify subgroups of neurons with similar patterns of incoming and outgoing neuropeptide

connections. We therefore applied dimensionality reduction methods (Principal Component Analysis

(PCA) and t-SNE) to the connectivity matrix of the aggregate neuropeptide networks to identify

groups of neurons that share a connectivity pattern. This analysis highlighted 3 clearly defined groups

of neurons that receive similar incoming connections, along with a more diffuse cloud of neurons with more variable connectivity (Figure 7A, B). Interestingly the three clusters comprise most neurons of the neuropeptidergic network (155 of 302), 76% of which are also members of the neuropeptidergic rich club (112; Figure 7A, B). Interestingly, each group receives a characteristic range of incoming connections, and this alone is enough to significantly sort them into these groups (p<0.0001) (Figure 7D): group 1 neurons have indegree of 134-156 (mean rank 162.2), group 2 neurons have an indegree of 252-282 (mean rank 276.5), group 3 neurons have an indegree of 169-212 (mean rank 222.9), and neurons in the unclustered cloud have an indegree below 169 (mean rank 77) (Figure 7D).

These groups also diverge in the neuron types that form them and to which they are connected (Figure 7C, E). Group 1 is mostly motorneurons (motor core), including those involved in locomotion (VA, DA, VB, and DB)[78] and the grouping of these neurons is driven by inputs from the posterior touch mechanosensory neurons PVM and PLML/R[79] and the interneuron PVWL/R (Figure 7C). Group 2 neurons are a mix of interneurons and motorneurons, including all top neuropeptidergic hubs (hubs core), that receive connections from every other neuron type, with the neuropeptidergic hubs (PQR, PVT and PVR) particularly important drivers for the grouping (Figure 7C). Finally, group 3 neurons are a mix of pharyngeal neurons, interneurons, motorneurons and sensory neurons (sensory core) that receive connections from every other neuron type but motorneurons, with RIR and pharyngeal neurons I5 and I4 driving the cluster (Figure 7C). Interestingly, although the characteristic neural inputs of the groups differed substantially, the most important (highest indegree) neuropeptide-receptor interactions for each group, involved versatile neuropeptides FLP-9, FLP-11 and FLP-16 and promiscuous receptors DMSR-7 and DMSR-1 (Figure 7C), leading to interconnections between groups (Figure 7E). For example, Group 1 forms connections with itself, Group 2 and with the unclustered cloud but not with Group 3; Group 3 forms connections with Group 2 and unclustered cloud neurons but not Group 1, and Group 2 forms connections with all other groups. Thus, Group 2 serves as a link with Groups 1 and 3, which have few direct connections with each other.

Co-expression of neuropeptides with their target receptors generates autocrine pathways

Signaling cascades, in which a neuropeptide receptor is specifically co-expressed with a non-

cognate peptide whose release it controls, are a classic hallmark of neuroendocrine pathways. To

investigate whether the *C. elegans* neuropeptide connectome contains such cascades, we evaluated co-

expression between neuropeptide and receptor genes. We performed a Fisher's test to look for

neuropeptide genes and GPCR genes that are co-expressed in the *C. elegans* nervous system more

often than expected from their individual expression frequencies. We identified 121 peptide-receptor

pairs meeting this criterion (Supplementary Figure 5), only 5% of which corresponded to cognate

neuropeptide-receptor pairs (autocrine connections). Using these peptide-receptor pairs as nodes and

the neuropeptide-receptor interaction data to form edges between them, we built a network of

overrepresented signaling cascades within the larger neuropeptide connectome (Supplementary Figure

5B). This network is fully connected and provides a simplified view of how neuropeptide signaling

pathways interact within the nervous system.

Co-expression of a neuropeptide receptor with its own ligand will generate a self-loop or

autocrine connection, in which release of the peptide can signal back onto the sending cell. If we

consider all cases where a peptide and its receptor are co-expressed in a given neuron, we find that 58%

of the *C. elegans* nervous system harbors putative autocrine peptide connections (Figure 8A and 8B).

Autocrine signals appear most prevalent in neurons categorized as either interneuron or motorneuron

(Figure 8B), although two of the three types with the largest number of autocrine connections (URX

and PQR) are sensory neurons. Promiscuous receptors participate more in autocrine peptide signaling

(Figure 8C, Suppl. Figure S6), the most prominent of which are DMSR-1 and DMSR-7 (Figure 8C).

We observed a strong positive correlation between the number of different autocrine peptide

connections that a neuron harbors to its degree within the neuropeptide network (Figure 8D, left panel),

with some but not all neuropeptide hubs showing a high diversity of autocrine signaling. In contrast,

weak to no correlations were observed between autocrine connection diversity and the synaptic, gap

junction, or monoamine connectomes.

Autocrine connections were especially prevalent in specific circuits in the *C. elegans* nervous

system. For example, both URX and PQR, which have among the highest diversity of autocrine

signaling, are key sensory neurons that mediate *C. elegans'* responses to aversive O_2 levels[80–82]; other

O_2 sensory neurons (AQR and BAG) also co-express ligand-receptor pairs, though to a lesser degree

(Figure 8E, bottom left). As O_2 sensing neurons tonically signal ambient O_2 concentrations[83], autocrine

signaling may play a stabilizing role in maintaining the homeostasis of neuronal activity. Autocrine

neuropeptide signals are also prevalent in the neuromuscular circuit of the ventral nerve cord (VNC –

Figure 8F, bottom right) which mediates locomotion[84]. These autocrine loops are primarily mediated

by eight RFamide-related neuropeptides that activate two promiscuous receptors, DMSR-1 and DMSR-

7, although other receptors such as NPR-5 are also involved (Supplementary Figure 6F). Interestingly,

autocrine signaling is restricted to the excitatory A- and B-class VNC motorneurons which drive

backward and forward locomotion, respectively[84]; in contrast, both D-type inhibitory motor neurons

and the excitatory AS motorneurons, which play modulatory roles in both forward and backward

crawling[85], do not co-express any neuropeptide-GPCR pairs. It should be noted that some autocrine

peptidergic pathways are shared between neighboring excitatory motorneurons in the VNC, raising the

possibility that they might coordinate activity and/or neurosecretion across excitatory motorneurons in

a paracrine manner[86–89]. Heterogeneity in autocrine neuropeptide pathways within the A- and B-class

motorneurons (Supplementary Figure 6F) hints at heterogeneity in the contribution of individual

motorneurons in this process.

Discussion

Neuropeptide signaling forms a complex wireless network

Neuropeptide signaling has long been recognized as critical to brain function, yet despite

recent advances in connectomics, the structures of neuropeptidergic signaling networks are largely

uncharacterized. We have generated a draft neuropeptide connectome by integrating information from

three datasets: a biochemical deorphanization screen matching neuropeptidergic GPCRs with their

ligands, a scRNAseq dataset characterizing the expression patterns of all peptide and receptor genes at

the single neuron level, and an anatomical dataset defining the morphologies and neuron-neuron

contacts of all *C. elegans* neurons. The resulting connectome is remarkably dense; even a short-range

network with the most conservative assumptions about the spatial diffusion of neuropeptides was

more than ten-fold denser than the previously mapped synaptic connectome of *C. elegans*. As we have

also used conservative assumptions regarding gene expression thresholds and ligand-receptor

affinities, and since approximately half the putative neuropeptide-activated GPCRs remain to be

deorphanized, the actual neuropeptide connectome is almost certainly even more dense than the draft

network described here. However, based on sensitivity analysis it seems clear that the overall

structure and topological features of the neuropeptide connectivity are likely to hold even when more

connections are added to the network (Supplementary Figure 4A-B).

A particularly salient feature of the neuropeptide connectome is its decentralized topology,

which contrasts sharply with the more centralized structure of wired neural connectomes. Synaptic

connectomes from worms to humans are characterized by a core of high-degree hubs, which are

interconnected to form a rich club. This rich club, which in *C. elegans* consists of 11 premotor

interneurons, occupies a central position in the connectome, connecting local modules and

coordinating their activity. Rich clubs have been previously linked to important functional properties;

for example, in *C. elegans* the synaptic rich club has been shown to be involved in global brain states

related to locomotion[55,74], and in *Drosophila melanogaster* the synaptic rich club constitutes the

sensorimotor integrative center of the organism[90]. The gap junction and monoamine connectomes of

C. elegans likewise contain relatively small rich clubs of less than 20 neurons of relatively high

degree. In contrast, the neuropeptide connectome contains a rich club of (in the short-range network)

156 neurons, more than half the neurons in the entire nervous system. The neurons in this

neuropeptidergic rich club are extremely well-connected to each other as well as to the rest of the

nervous system; all have a degree greater than 203 (in the mid-range network degree of 251), meaning

they both send and receive direct connections from at least 40% of all neurons. This remarkable

decentralization may be a feature of neuropeptide signaling networks in other organisms[43,44,91].

Implications of neuropeptide network structure for neuronal computation

The decentralized structure of the neuropeptidergic connectome implies that the strategies it

uses for computation and information flow may differ significantly from those employed by the more

centrally-organized synaptic network. Intriguingly, both nematode and mammalian neuronal types

appear to express nearly unique combinations of neuropeptides and receptors, serving as a molecular

bar code for neural identity[40,43]. Thus, the source of a signal may be encoded by the precise

combination of peptides released by the sending neuron. The dynamics of neuropeptide release are

also likely to be critical to the information conveyed by peptidergic signals. For example, acute

release of FLP-20 peptides by mechanosensory neurons has been shown to trigger short-term sensory

locomotor arousal in response to touch stimulation, while chronic release of FLP-20 peptides from the

same neurons mediates long-term cross-modal plasticity in olfactory circuits when the sense of touch

is lost[14]. Future studies of such mechanisms are likely to provide general insights into how

neuropeptide networks encode information in the brain.

Despite its highly dense connectivity, the core of the neuropeptide connectome exhibited a

clear substructure. When input patterns were analyzed using dimensionality reduction methods, the

neurons in the network core could be divided into three clear groups, one containing mainly

pharyngeal and sensory neurons, a second containing the highest degree interneurons, and a third

containing many motorneurons. These three groups are themselves connected in a defined pattern,

with the second group linking the first and third groups which show few direct links with each other.

This organization contrasts in interesting ways with the organization of many synaptic networks, in

which peripheral neurons form modules with high internal connectivity that connect to each other

through the hubs of the rich club core. In the neuropeptide connectome the core itself exhibits a clear

meso-scale organization, forming 3 distinct groups of neurons defined not by unusually high intra-

group connectivity, but by intra-group similarity of incoming and outgoing connections patterns. This

is in line with recent work showing the existence of different types of meso-scale organization[77] and is

also reminiscent of stochastic block modelling approaches previously applied to the *C. elegans*

synaptic connectome[52]. It will be interesting to see if neuromodulatory and synaptic networks from

other nervous systems show a similar diversity on meso-scale structure. We expect that looking

beyond classical modules is likely to become increasingly important as we begin to look at cellular-

scale connectomes of larger organisms, with classes of sensory neurons that are not interconnected but

have similar connectivity profiles and perform similar functions.

Identification of neuropeptide signaling hubs

Although the rich club of the neuropeptide connectome is extensive and encompasses a large

portion of the *C. elegans* nervous system, some neurons within this group are of particularly high

degree and therefore may play key roles in neuromodulatory signaling. Perhaps not surprisingly, the

premotor neurons of the synaptic rich club are also highly connected in the neuropeptide connectome;

all 11 of these neurons show a neuropeptide degree above 400 in the short-range network. Since these

neurons are known to play important roles in driving global brain states[14,74,76] it is logical that they

would also be important targets of neuromodulatory control. In addition, the neuropeptide

connectome contains 6 neurons whose neuropeptide degree is higher than any of the synaptic rich

club (degree > 490). Three of these neurons (PVT and the pair of AVK neurons) are specialized

peptidergic neurons that express no classical neurotransmitter or monoamine and have been linked to

arousal and sleep-like behaviors[14,92,93]. Specialized neuropeptidergic neurons have also been

described in other organisms, and in mice they have been linked to global behaviors such as fear[94].

The other peptidergic hubs (PVR and the pair of PVQ neurons) are tail neurons that extend long

processes to the nerve ring, and in the case of the PVQs (as well as PVT) these processes are rich in

dense-core vesicles[48,65,66]. Since AVK also has a long process (extended from its cell body in the head

through the nerve ring and the ventral cord to the tail) these neurons may be morphologically

specialized for local release of peptides throughout the nervous system. While AVK is known to play

roles in arousal and motor control, the functions of the remaining neuropeptide hubs (or of other

neurons of high neuropeptidergic degree such as PVN and PVP) are largely uncharacterized. Given

their importance in the neuropeptide signaling network, it will be interesting in the future to explore

the roles of these neurons in the control of behavioral states.

It is interesting to note that even in the long-range network, where no spatial restrictions on

neuropeptide diffusion are imposed, the six short and mid-range hub neurons retain their central

importance in the network. This confirms that the high degree of the hubs in the short- and mid-range

networks is not merely an artifact of the spatial conditions we imposed but rather is a consequence of

expressing key combinations of neuropeptides and receptors. Notably, the long-range network

contains additional high-degree nodes that are not hubs in the short- and mid-range networks, in

particular the oxygen-sensing neurons URX, AQR, PQR and BDU and the tail motorneurons DA09

and VA12. One might speculate that these neurons could in fact engage in long-range neuroendocrine

signaling, and that the long-process morphology of the short- and mid-range hubs might allow them to

carry out neuropeptide modulation on a finer temporal or spatial scale. In the future, these questions

may be addressable using *in vivo* probes for neuropeptide receptor signaling[95,96].

Neuropeptide signaling links specific components of the nervous system

In addition to these high degree nodes, the neuropeptide connectome also contains edges of

unusually high weight, representing neuron pairs linked by multiple neuropeptide-signaling pathways.

While many neuron pairs in the network are connected by a single neuropeptide pathway, 17 pairs of

neurons are linked by 15 or more different neuropeptide-receptor couples. Most of these extremely

high-weight edges involve connections between the oxygen-sensing neurons PQR or URXL/R and the

motor circuit (either the DB motorneurons or the AVD premotor interneurons). Why might oxygen-

sensing neurons participate in so many complex neuromodulatory interactions? These neurons

strongly influence locomotor states and their tonic responses to ambient oxygen are influenced by

experience and other sensory cues. Complex neuropeptide signaling may allow feedback between the

oxygen sensors and motorneurons to fine-tune the activity of this circuit across time and space.

The oxygen-sensing neurons and the motor circuit were also found to be important sites for

autocrine signaling, in which a neuropeptide receptor and its ligand are co-expressed in the same

neuron. Autocrine peptide signaling is ubiquitous in other brains as well[97,98], suggesting it supports

important aspects of neural function. For example, autocrine neuropeptide pathways are known to

mediate cell-autonomous feedback[99–104], and are therefore often suggested to maintain neuronal

homeostasis; this may be particularly important to regulate the activity of tonically signaling neurons,

such as the oxygen-sensing neurons. In the motor circuit, ostensibly autocrine signaling may play a

further role by coordinating the activities of neighboring neurons with similar gene expression

patterns. Although proprioceptive and electrical coupling between adjacent motorneurons generates

waves of muscle contraction over adjacent body regions[88], autocrine pathways also have the capacity

to coordinate physiology across neighboring cells[105,106]. Indeed, mutants for several of the

neuropeptide ligands that act in motorneuron autocrine pathways have been shown to have

locomotion defects[107,108], suggesting an important role for these connections in patterning locomotor

behavior.

We also observe a critical role for neuropeptide signaling in linking disconnected components

of the nervous system to the broader network. In particular, the pharyngeal neurons, which form a

wired network analogous (if not homologous) to the vertebrate's enteric nervous system, and the CAN

neurons, whose processes lie in the excretory canal-associated nerve, are virtually completely

unconnected to the somatic synaptic and gap junction connectomes yet are extensively integrated into

the neuropeptide connectome. Both CAN and the pharyngeal neurons express multiple GPCRs whose

ligands are expressed exclusively outside the pharynx or canal-associated nerve, indicating that these

neurons are regulated by peptides released from physically-unconnected processes. Despite their

disconnection from the wired connectomes, the pharynx and CAN neurons carry out essential

physiological functions; indeed, CAN and the pharyngeal neuron M4 are the only neurons whose

ablation is lethal to the animal[109,110]. This provides a means for communication between the wider

nervous system and these isolated but biologically critical neurons.

Prospects for future mapping of wireless brain connectomes

We have described here the draft neuropeptide connectome of *C. elegans*, an animal with 302

neurons. In the future, we plan to refine this connectome, for example by expanding its scope through

deorphanization of neuropeptide receptors whose ligands are currently unknown. Differential

posttranscriptional processing may also lead to the synthesis of different peptides and receptors in

neurons that express the same gene, and alternative patterns of gene expression during development or

in response to environmental cues may likewise alter the structure and function of neuropeptide

signaling networks. Moreover, non-neuronal cells function as both senders and receivers of

neuropeptide signals; with the use of reporter lines for peptides and receptors it should be possible in

the future to incorporate these cells into the neuropeptide connectome and observe plasticity in the

network. We likewise plan to integrate functional information into the connectome map; for example,

in vitro experiments could identify the G-protein pathways downstream of individual receptors, and *in*

vivo sensors could provide empirical data on the spatial scope of neuropeptide signaling pathways.

Together these data will facilitate functional modeling of neuropeptidergic circuits revealed in the

connectome maps.

In principle, the approaches described here should also be applicable to mapping the

neuropeptide signaling networks of animals with much larger brains. Single-cell RNAseq data from

different vertebrate animals indicates that, as in *C. elegans*, most neurons express at least one

neuropeptide precursor and at least one neuropeptide receptor, facilitating dense and potentially

decentralized networks[18,39,43,44]. Although it is currently not possible to precisely link gene expression

clusters with individual neurons in brain circuits, the use of targeted reporters should eventually make

it possible to relate neuropeptide and receptor expression to increasingly detailed synaptic

connectome maps in flies and mice. Basic mechanisms of neuropeptide signaling are shared in all

animals, from nematodes to mammals: neuropeptides are released from dense core vesicles and

diffuse across space to neurons unconnected to the releasing cell by wired synapses. And although the

C. elegans nervous system is anatomically small, at the molecular level its neuropeptide systems are

highly complex and show significant homology to other animals. Thus, the neuropeptide connectome

of *C. elegans* may serve as a prototype to unravel general principles of neuromodulatory network

structure that also apply to much larger brains.

[34,35363718,38]

Methods

Reporter Transgenic Strains

CRISPR based constructs: Thirty-two transcriptional C-terminal GFP reporters of neuropeptides and

neuropeptide receptors were created where GFP was recombineered in the last coding exon of the

gene of interest. Of the 17 total neuropeptide precursor genes that we made reporters for, 8 had only a

T2A::3xNLS::GFP tag, 1 had a T2A::GFP::H2B tag, 6 had both T2A::3xNLS::GFP and

SL2::GFP::H2B tags, and 2 had only an SL2::GFP::H2B tag (Supplementary Table 1). For the GPCR

receptors, 8 gene expression reporters were made all with SL2::GFP::H2B tags (Supplementary Table

2). Most of these constructs were made by SUNY Biotech.

Reporter Analysis

Young adult animals were mounted on 5% agarose pads and immobilized with 100 mM sodium azide

and imaged on a Zeiss LSM880 using a 40X objective lens. GFP expression reporters were identified

at single neuron resolution as described[111]. GFP reporter expression of these constructs, as reported in

Table S1-2 were noted using three categories: moderate to high expression, low and variable

expression, and no detected expression. Additionally, we compared our GFP reporter expression data

to single-cell RNA-seq expression (scRNAseq) data from the CeNGEN project using their standard

thresholds (4 being the most stringent, 1 being the least stringent, and blank be unfiltered)[40].

In our analysis, for each gene and each CeNGEN threshold, we tallied 1) the number of neurons that

showed GFP expression but not scRNA expression, 2) the number of neurons showing both GFP

expression and scRNA expression, and 3) the number of neurons that showed scRNA expression but

no GFP expression (Table S3).

Based on the results of this analysis threshold 4, although in some occasions too conservative, had the

best correlation between GFP reporter and CeNGEN scRNAseq data expression per neuron for the

tested NPP and GPCR genes.

Synaptic and gap junction networks

The synaptic and gap junction networks used in this work were based on the full

hermaphrodite *C. elegans* connectome, containing all 302 neurons. This network was composed from

the somatic connectome[48], updated and released by the Chklovskii lab[47]; and the pharyngeal network

of Albertson and Thomson[65], made available by the Cybernetic *Caenorhabditis elegans* Program

(CCeP). The functional classifications referred to in the text (i.e., *sensory*

neuron, *interneuron*, *motorneuron*) are based on the classification scheme used in WormAtlas[112].

When there is double or triple classification in nerve ring neurons that can be sensory neurons,

interneurons, or motorneurons, the Zhen lab classification[66] was used to select one neuron type. URB

left and right neurons are the only ones in which the WormAtlas and the Zhen lab classification

diverge, leading us to classify them as sensory neuron following the later most recent classification.

DB neurons are identified as motorneurons although WormAtlas indicates that these could also be

interneurons[112]. The gap junction network was modelled as an undirected network with bidirectional

electrical synapses; note however that some gap junctions might be rectifying and thus exhibit

directionality. In the synaptic network reciprocal connections between nodes are considered as two

separate unidirectional connections.

Monoamine network construction

The monoamine network used in this work was made following the same procedure as in (Bentley et

al., 2016). The monoamine expression for the 302 neurons comes from the neurotransmitter atlas of

C. elegans[75] and receptor expression for the 302 neurons comes from the single-cell expression data

from the CeNGEN project (https://www.cengen.org)[40]. We used the expression data at CeNGEN

threshold 4. The interactions between ligand and receptors were previously described[45]. The

adjacency matrix was built using a binary version of the expression data for the 302 neurons. For a

given point $A^M(i,j)$ and for a given monoamine receptor pair M the connection between two neurons is

defined by $A^M(i,j) = Mon^M(i,j) \times Receptor^M(i,j)$. Each monoamine receptor interaction forms an

individual binary network. To get the overall monoamines network we add each individual

monoamine receptor network resulting in a weighted network where the weight indicates the number

of monoamine receptor pairs that connect two nodes. Reciprocal connections between nodes are

considered as two separate unidirectional connections.

Neuropeptide network construction

The neuropeptide network used in this work was made using a similar approach to that used for the

monoamines. The interactions between ligands and receptors were identified using a large-scale *in*

vitro reverse pharmacology pipeline in which over 87% of the predicted peptide GPCRs were

challenged with FMRFamide related peptides (FLP) and non-insulin non-FLP like peptides (NLP)[64].

Neuropeptide precursor and GPCR gene expression for the 302 neurons was extracted from the

single-cell transcriptome data of the CeNGEN project (https://www.cengen.org)[40]. We used the

expression data at CeNGEN threshold 4. The adjacency matrix was built using a binary version of the

expression data for the 302 neurons. For a given point $A^N(i,j)$ and for a given neuropeptide receptor

pair N the connection between two neurons is defined by $A^N(i,j) = NPP^N(i,j) \times GPCR^N(i,j)$. Each

neuropeptide receptor interaction forms an individual binary network. To get the overall

neuropeptides network we add each individual neuropeptide receptor network resulting in a weighted

network where the weight indicates the number of neuropeptide receptor pairs that connect two nodes.

Reciprocal connections between nodes are considered as two separate unidirectional connections.

Neuropeptide network spatial constraining

Neuropeptidergic networks were locally thresholded to filter out connections between neurons that

were anatomically far from each other. The anatomical EM data was obtained from The Mind of the

worm (https://www.wormatlas.org/MoW_built0.92/MoW.html) and other literature[48,65,66]. This data

was used to create a table of locations for each neuronal process, identifying 27 different neuronal

process bundles in the *C. elegans* nervous system as previously defined[48]. This classification was then

used to filter out neuropeptidergic connections based on putative signaling ranges. The stringent

short-range thresholding allows connections only in between neuronal processes that are in the same

process bundle and the pharynx is a separated system were all connections are allowed in between

pharyngeal neurons only. The mid-range stringency thresholding allows connections between neurons

with neuronal processes in the same anatomical area: head (including pharynx and the ventral cord

neurons that are in the ventral ganglion), midbody and tail. And in the unthresholded system all

neuropeptidergic connections are allowed.

Topological network measures

Edge counts, adjacency matrices and reducibility clusters were all computed using binary directed

versions of the networks. The same networks, excluding self-connections (i.e. setting all diagonal

elements to 0), were used to compute all other measures.

Network measures are compared to 100 null model networks generated using the degree-preserving

edge swap procedure from the Brain Connectivity Toolbox for MATLAB[113]. This is performed by

selecting a pair of edges ($A{\rightarrow}B$) ($C{\rightarrow}D$) and swapping them to give ($A{\rightarrow}D$)($C{\rightarrow}B$). If the resulting

edges already exist in the network, another pair of edges is selected instead. Each edge was swapped

10 times to ensure full randomization.

Degree

Degree is the number of edges connected to a given node. Indegree is the number of incoming

connections connected to a given node and outdegree is the number of outgoing connections.

Density

Density d is the fraction of present connections K to possible connections between the given nodes N:

$$d = \frac{K}{N^2 - N}$$

Clustering coefficient

Transitivity defines the ratio of triangles to triplets in the network (where a triple is a single node with

edges running to an unordered pair of others, and a triangle is a fully-connected triple). For a directed

network, this is equivalent to: $T = \frac{\sum_{i \in N} t_i}{\sum_{i \in N}[(k_i^{out} + k_i^{in})(k_i^{out} + k_i^{in} - 1) - 2\sum_{j \in N} A_{ij}A_{ji}]}$ where A is the

adjacency matrix, N is the number of nodes, k^{out} and k^{in} are the out-degree and in-degree, and t_i is the

number of triangles around a node: $t_i = \frac{1}{2}\sum_{j,h \in N}(A_{ij} + A_{ji})(A_{ih} + A_{hi})(A_{jh} + A_{hj})$

Reciprocity

Reciprocity is the fraction of reciprocal edges in the network: $r = \frac{|E^{\leftrightarrow}|}{M}$ where M is the number of edges,

and $|E^{\leftrightarrow}|$ is the number of reciprocal edges: $|E^{\leftrightarrow}| = \sum_{i \neq j} A_{ij}A_{ji}$

Rich-club coefficient

The rich-club coefficient measures the tendency for high-degree nodes in a network to form highly

interconnected communities[55]. These communities can be identified by creating subnetworks for each

degree level k and removing nodes with a degree $\leq k$. Then the rich-club coefficient $\Phi(k)$ for each

subnetwork is defined as the ratio of connections in the subnetwork M_k to the number of maximum

possible connections. For a directed network with no self-connections, where N_k is the number of

remaining nodes, this is given by: $\Phi(k) = \dfrac{M_k}{N_k(N_k - 1)}$

Thus, a fully connected subnetwork at a given degree k has a rich-club coefficient $\Phi(k) = 1$. We

normalize the rich-club coefficient by calculating: $\Phi_{norm}(k) = \dfrac{\Phi(k)}{\langle \Phi_{random}(k) \rangle}$ were $<\Phi_{random}(k)>$ is the

average value of the rich-club coefficient across random networks.

A rich-club exists when $\Phi_{norm}(k) \geq 1$, but in order to get a clear threshold range we use a probabilistic

approach. The threshold range of the rich-club is defined by $\Phi_{norm}(k) \geq 1 + 1\sigma$, where σ is the Standard

Deviation of $\Phi_{random}(k)$ for the 100 random networks.

Dimensionality reduction analysis

t-SNE is an algorithm for dimensionality reduction that facilitates visualizing high dimensionality

data. The analysis described here was performed using the MATLAB *t-sne* function on the adjacency

network of connections. The neuropeptide dimension was reduced, and clustering was performed

based on the pattern of connections due to receptor expression. Different distant measures were tested

to confirm the clustering: Euclidean distance, Chebychev distance, cosine distance and Mahalanobis

distance.

Co-expression analysis

The signaling networks used in this work represent connections between co-occurring genes. Nodes

are defined as pairs of neuropeptide precursor and GPCR genes that co-occur more than expected by

chance as measured by a Fisher's exact test[114] with FDR (false positive rate) correction[115]. The 2x2

contingency table for the Fisher's exact test contains the number of neurons for which both genes co-

occur and the number of neurons in which each NPP and GPCR gene is expressed without co-

occurring with the second gene. Thus, the Fisher's test is defined as:

752
$$p = \frac{(R_1!\,R_2!)(C_1!\,C_2!)}{N!\,\prod_{i,j} n_{i,j}!}$$

753
Where R_1 and R_2 are the row sums, C_1 and C_2 are the column sums, N is the total number of

observations in the contingency table, and n_{ij} is the value in the ith row and jth column of the table

Interactions between nodes are defined by the receptor-ligand interactions that the co-occurring genes

have with genes that co-occur in another node. The interactions between ligand and receptors were

identified using a large-scale *in vitro* reverse pharmacology pipeline[64].

Software

Network measures were computed in MATLAB (v9.8.0.1323502 (R2020a), The MathWorks Inc.

Natick, MA) using the Brain Connectivity Toolbox[113] (v2019-03-03) and the MATLAB/Octave

Networks Toolbox[116]. Clustering and visualization of multilayer plots was performed using

MuxViz[117]. And additional network visualizations were created using Cytoscape[118].

Lead contact

Requests for resources should be directed to the Lead Contact William Schafer (wschafer@mrc-

lmb.cam.ac.uk)

Data and software availability

The scRNAseq data are available at www.cengen.org. The biochemical deorphanization data are

available at https://worm.peptide-gpcr.org/project/neuropeptides/flps/ [64]. Analysis code is available at

https://github.com/LidiaRipollSanchez/Neuropeptide-Connectome.git

Nematode Strains

The strains used in this study are listed in Supplementary Tables 1 and 2.

Acknowledgements

This work was funded by grants from NIH (R01NS110391, to OH, WRS and IB), the MRC (MC-A023-5PB91, to WRS), the Wellcome Trust (WT103784MA, to WRS), MQ: Transforming Mental Health (MQF17_24, to PV) the ERC (950328, to IB), HHMI (to OH), and the Simons Foundation (855199, to RF). This research was also supported in part by the NIHR Cambridge Biomedical Research Centre (BRC-1215-20014) and NIHR Applied Research Centre. The views expressed are those of the author(s) and not necessarily those of any funding bodies. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising from this submission.

We'd like to thank the LMB Visual Aids and Scientific Computing teams for their support, especially Johanna Westmorland for figure 3 design. We thank Tom Shimizu, Erik Larsson Lekholm, Mark Moyle, Caroline Nettekoven, Sarah Morgan, Albert Cardona, Ed Bullmore, Amy Courtney and Mei Zhen for productive discussions. We thank WormAtlas, OpenWorm, Nemanode and WormBase for their resources.

Author contributions

L.R.S., W.R.S., P.E.V., J.W. and I.B. designed research. L.R.S. and J.W. performed analysis. H.S. and R.F. made strains and interpreted images. H.S., R.F., S.R.T., A.W., M.H., D.M.M., O.H. contributed with data and discussion. L.R.S., W.R.S., P.E.V., J.W. and I.B. contributed to writing the paper. All authors reviewed and approved the manuscript.

Declaration of Interests

The authors declare no competing interests

Figure titles and Legends

Figure 1. *In silico* reconstruction of the neuropeptidergic connectome of the *C. elegans* nervous system by combining *in vitro* neuropeptide-GPCR interaction, gene expression and anatomical datasets.

(A) Datasets used to build the network *in silico*. Left: biochemical neuropeptide-GPCR interaction dataset defining 91 neuropeptide - G protein-coupled receptor (GPCR) pairs below

500 nM EC$_{50}$ affinity, coupling 51 GPCRs to 48 neuropeptides[64]. Neuropeptides from the

same precursor gene (NPP) are considered the same. Middle: anatomical and morphological

description of the *C. elegans* hermaphrodite nervous system[48]. Right: single-cell RNAseq

expression of the 302 *C. elegans* neurons, adapted from[40].

(B) Representation of the *C. elegans* nervous system connectome network in which all types of

neurons (pharyngeal, interneuron, CAN neurons, motorneurons and sensory neurons) are

connected through neuropeptidergic connections.

Figure 2. Single-copy genomic knock-in reporters corroborate the expression profiles of neuropeptide

and GPCR genes obtained from scRNAseq. GFP-positive neurons were identified using the

NeuroPAL multicolor transgene [111]Figure pictures all segments (head, tail, vulva and midbody)

showing neuronal expression, within which individual neurons are circled and labeled. Scale bars

represent 10 μm. Tables of expression data and comparisons between cells identified in reporters and

scRNAseq detection thresholds are in Supplementary Tables S1-3.

(A) Fluorescent GFP reporters for the expression of two representative neuropeptide genes, *flp-20*

and *nlp-45,* showing high correalation between reporter expression and CeNGEN scRNAseq

expression. Data for additional neuropeptide gene reporters are shown in Supplementary

Figure S1A.

(B) Fluorescent GFP reporters for the two representative neuropeptide-activated GPCRs,

expression of *tkr-1* and *dmsr-6*, showing high correalation between reporter expression and

CeNGEN scRNAseq expression.[111]. Data for additional GPCR gene reporters are shown in

Supplementary Figure S1B.

Figure 3. Analysis of neuropeptide and receptor expression allows assessment of the spatial scale of

neuropeptide signaling.

(A) Full anatomical description of the *C. elegans* hermaphrodite nervous system[112]. Neuronal

bundles represented in red, and the pharynx in green. Short-range connections are defined as

occurring within the same neuronal bundle (axons and dendrites considered equally). Mid-

range connections occur between neurons whose processes are present in one of following

distinct anatomical areas: head (ranging from nose to deirid commissures including the

ventral and retrovesicular ganglion), midbody (between but not including deirid commissures

and the preanal ganglion), or tail (from and including the preanal ganglion to the tail's tip).

Classifications of individual neurons are in Supplementary Table S4.

(B) Expression matrix for 23 GPCRs activated *in vitro* by a single neuropeptide ligand. Neurons

are sorted by neuron type on the x axis and the expression of each GPCR on the y axis. Colors

indicate the range of diffusion distance required for communication with at least one ligand-

expressing neuron: blue indicates that the receptor expressing neuron makes contact with a

neuron expressing its ligand, either within the same stratum of the nerve ring (dark blue) or in

a thinner neuronal bundle (light blue); orange indicates short-range connections between

strata in the nerve ring (i.e. between neurons in same bundle but not in physical contact); red

indicates connections between neurons in the same anatomical area but not in the same

bundle. Nerve ring stratum as defined[67,6867,6867,68]

(C) Expression matrix for 23 neuropeptide precursor genes with a single target receptor in vitro.

Neurons are sorted by neuron type on the x axis and the expression of each neuropeptide gene

on the y axis. Colors indicate the range of diffusion distance required for communication with

a receptor-expressing neuron, as described above.

Figure 4. Assortativity analysis of 91 individual NPP-GPCR networks shows different topologies

depending on the NPP-GPCR interaction.

(A) Representation of network topologies depending on assortativity, reflecting the preference of

neuropeptides to signal to GPCRs that are expressed in the same number of neurons.

Networks are classified either as local, pervasive, broadcaster or integrative. Bottom left:

scatter plot showing the number of neurons expressing a particular GPCR versus the number

of neurons expressing the corresponding NPP gene for each of the 91 individual networks.

Local networks show restricted NPP and GPCR expression in 50 neurons or less. Pervasive

networks have broad NPP and GPCR expression in more than 50 neurons. Broadcaster

networks show broad GPCR (>50 neurons) but restricted NPP expression (≤50 neurons),

while integrative networks display broad NPP (>50 neurons) and restricted GPCR expression

(≤50 neurons). Filled circles indicate receptor presence in that neuron and empty circles

indicate neuropeptide release from that neuron. Left top: example of local network for CAPA-

1/NMUR-1 that has been linked to learning[32]. Left middle top: example of pervasive network

FLP-18/NPR-5 of which the receptor is implicated within locomotory behavior[119]. Right

middle top: example of broadcaster network FLP-1/FRPR-7 that modulates the interaction

between food sensation and locomotion[13]. Right top: example of integrative network NLP-

47/GNRR-1, the receptor of which is linked to egg-laying behavior [120] and is only activated by

NLP-47 *in vitro* [64,120]. Diagrams of all 91 networks are in Supplementary Figures S2 and S3.

(B) The topology of the network is defined by the individual NPP-GPCR pairing. Networks of the

same receptor can show different topologies depending on its neuropeptide ligand. In each

example, the same receptor can form both pervasive and broadcasting networks (DMSR-2,

DMSR-7) or both local and integrative networks (FRPR-8 and EGL-6) depending on the

activating ligand.

Figure 5. The aggregate neuropeptide connectome in *C. elegans* connects all 302 neurons in a dense

network. Adjacency matrix representation of the aggregate 91 NPP-GPCR pair network considering

short-range (color) and mid-range (gray) diffusion models. Axes represent the number of NPP and

GPCR genes per neuron. Virtually all neurons express at least 1 NPP and 1 GPCR gene, the only

exception of which is the IL1 neuron class that does not show GPCR expression. Neurons can

communicate with each other using up to 18 different NPP-GPCR pathways, a feature conserved in

both short- and mid-range diffusion models. Dense and diverse connections are mainly seen from

oxygen sensing neurons to inter- and motorneurons. 5% of all connections are putative autocrine

connections in which a single neuron co-expresses both NPP and GPCR genes of a cognate pair.

Supplementary Figure 4 shows adjacency matrices for long-range (A) and mid-range (B) models.

Figure 6. Topological analysis of the *C. elegans* nervous system highlights neuropeptide signaling

hubs and a large rich club.

(A) Degree distribution of the short-range model aggregate neuropeptide network. Degree (sum of incoming and outgoing connections) is shown in orange, the indegree (sum of incoming connections) is shown in green and the outdegree (sum of outgoing connections) is shown in purple. . The 10 hubs with the highest degree are indicated, including hubs of the synaptic connectome (AVAL/R, PVCL/R and DVA) as well as 6 neurons with specialised roles in the neuropeptide signaling network (AVKL/R, PVT, PVQL/R, PVR).

(B) Degree distribution of the mid-range model aggregate neuropeptide network. Similar to the short-range model, degree distribution is equally influenced by both in- and outdegree: the 10 hub neurons of highest degree are indicated.

(C) Degree distribution of the synaptic network of *C. elegans*. Compared to the neuropeptide connectome, the synaptic connectome contains a much smaller number of high-degree hub nodes.

(D) Degree distribution of the monoamine network of *C. elegans*. The overall degree distribution shows two distinct phases, correlating with the network having a star-like topology centered around a small core of high degree hubs. This topological feature is further reflected in the low outdegree of most neurons.

(E) Conceptual network representation highlighting: node (neuron), edge (connection between neurons), degree (k, sum of incoming and outgoing connections), hub (highly connected neurons), and rich club (group of hubs that connect more to one another than to the rest of the network).

(F) Rich club of the *C. elegans* neuropeptidergic connectome. The rich club coefficient $\Phi(k)$ for the *C. elegans* neuropeptidergic network is shown in black and the randomized rich club curve $\Phi_{random}(k)$ generated by averaging the rich club coefficients of 100 randomized versions of the *C. elegans* neuropeptidergic network that preserve degree distribution) is depicted in gray. The red curve is the normalized coefficient. $\Phi(k) \geq \Phi_{random}(k) + 10\sigma$ over the range $203 \leq k \leq 298$, indicating the rich club. The large number of connections in the range $k > 298$ make the random networks more unreliable but since $\Phi(k)$ is very close and mostly higher

than $\Phi_{random}(k) + 10\sigma$ we can infer that all neurons where $203 \le k$ are part of the rich club.

This means the short-range aggregate neuropeptide network has a rich club of 156 neurons

(166 in the mid-range, Suppl. Figure 6)

(G) Correlation between synaptic and neuropeptidergic degrees. A positive correlation was

observed over all neurons ($r = 0.54$, $p = 3.1\ e^{-14}$). Dark red dots indicate neurons in the *C.*

elegans neuropeptidergic rich club. Synaptic hubs, which also exhibit a high neuropeptide

degree ($k>400$) (ranking among the top 30 neuropeptide hubs) are highlighted. Also

highlighet are 6 neuropeptidergic hubs, with low synaptic degree ($k < 30$, below the

correlation line), as well as low monoamine ($22 > k \ge 10$) and gap junction ($14 > k \ge 3$)

degrees (Suppl. Figure 6), appear to be specialized for neuropeptide signaling.

Figure 7. Mesoscale structure of the neuropeptide connectome

(A) t-SNE dimensionality reduction of the adjacency matrix of the mid-range aggregate network

(Euclidean distance, perplexity 30). This plot we identifies 3 clear clusters (1, 2, 3) which

encompass 112 of out the 166 mid-range neuropeptide rich club neurons), in addition to a

periphery of loosely clustered neurons. Datapoint markers represent their neuronal

classification.

(B) PCA dimensionality reduction of the adjacency matrix of the mid-range aggregate network

(Euclidean distance, perplexity 30). Neurons in this plot are colored based on that of the

groups defined in (A). A clustering pattern similar to that following t-SNE appears,

highlighting the robustness of these defined groups across dimensionality reduction

techniques. Datapoint markers represent their neuronal classification.

(C) Mid-range aggregate network sorted in both dimensions (sending and receiving neurons)

based on neuronal clusters defined in (A) and (B). Neuronal clusters divisions are highlighted

with discontinuous lines. Group 3 does not receive nor send connections to group 1. All three

groups, specially 2 and 3, send a large number of connections to peripheral neurons while not

receiving significant input from the periphery. Group 2 is the only group that sends and

receives connections to and from all 4 neuronal clusters. The neurons with higher weight of

connections (number of NPP-GPCR pairs used to make the connection) are mainly found in

groups 2 and 3. The neurons defining the patterns that drive the clustering of each group are

shown at the bottom: group 1 is mainly driven by the connections of DA, VC and VA

terminal motorneurons; group 2 is mainly driven by the connections of neuropeptidergic hubs

PVQL/R and PVR, synaptic hubs AVAL/R and PVNL/R neurons; group 3 is mainly driven

by the connections of pharyngeal neurons I5 and I4 and RIR, RICL/R interneurons.

(D) Violin plots showing indegree values for the 3 clusters and the periphery. Indegree cleanly

defines the 3 groups and the periphery (median group 1: 139, median group 2: 267, median

group 3: 198, median periphery: 54), and therefore is the determinant factor in the definition

of these groups. Significances determined by Kruskal-Wallis test followed by Tukey-Kramer

test for multiple comparisons with rank sums. The mean ranks for the 4 groups were shown to

be significantly different (mean rank group 1: 162, mean rank group 2: 276, mean rank group

3: 223, mean rank periphery: 77). n.s. not significant; ***$p \leq 0.001$.

(E) Diagram showing the organization of connections between the 3 defined clusters. Group 1

contains the main motorneurons core. These connect between themselves, with the periphery

and with group 2. Group 2 contains the main interneurons core including the top 10 hubs.

These neurons connect to almost every other neuron in the nervous system. Group 3 contains

the main sensory neurons and pharyngeal neurons core. These neurons connect to group 2

neurons and the periphery but do not connect to group 1 neurons. This structure indicates an

intrinsic organization of the *C. elegans* neuropeptidergic connectome rich club and how it

coordinates the function of the overall neuropeptidergic connectome.

Figure 8. Co-expression between GPCRs and their corresponding neuropeptide ligand(s) facilitates

potential autocrine and paracrine signaling. Additional data on putative autocrine signaling is in

Supplementary Figure S6.

(A) Neuronal expression matrix for both NPP and GPCR genes of the 91 NPP-GPCR pairs. Grey

dots represent neuronal expression of only the NPP (upper panel) or GPCR (lower panel)

gene, which black dots indicating co-expression of both.

(B) Percentage of each neuron type showing peptide autocrine connections (upper panel). The

number of different NPP-GPCR pairs being co-expressed in each neuron type is figured in the

bottom panel. Neurons showing the highest diversity of different NPP-GPCR pairs being co-

expressed are highlighted, including the sensory neurons PQR and URX and the interneuron

PVR.

(C) Scatter plot showing the number of neurons with co-expression for each of the 91 NPP-

GPCRs. Pairs with promiscuous GPCRs, specially DMSR-1 and DMSR-7 are co-expressed in

many different neurons, highlighting their central role in autocrine neuropeptidergic

signaling.

(D) Correlation between number of autocrine connections and neuropeptide (left) or synaptic

(right) degree for each of the 302 neurons. Point shapes indicate degree (round), indegree

(incoming arrow), outdegree (outgoing arrow).

(E) Representation of the autocrine connections in each neuron of the *C. elegans* worm (anterior

part of the worm in the left side, posterior in the right, dorsal upper part, and ventral lower

part). The size of each cell body indicates the number of autocrine NPP-GPCR pairs

expressed in that neuron. Circuits in which autocrine connections are prominent are the

locomotion and oxygen sensing circuit. The colors indicate the neurons in the oxygen sensing

and locomotion circuits that have the largest number of autocrine connections. Left bottom:

oxygen sensing circuits with autocrine connections, URX and PQR are the neurons with the

largest number of connections (arrow size indicates number of NPP-GPCR pairs). Right

bottom: description of the locomotion system with autocrine connections, VA/VB and

DA/DB are excitatory motorneurons that actuate worm crawling and which show many

autocrine connections while inhibitory DD/VD and minor excitatory AS motorneurons do not

have autocrine connections.

997 **Supplementary Figure S1.** Expression patterns of neuropeptide and GPCR reporters. Cells were

998 identified as described in Figure 2. A) Fluorescent GFP reporters for the expression of 17

999 representative neuropeptide precursor genes. B) Fluorescent GFP reporters for the expression of 8

1000 representative GPCR precursor genes. Images of the *trhr-1* expression strain show some bleed-

1001 through signal that does not represent neuronal expression.

1002

1003 **Supplementary Figure S2.** Graphical representations of the 91 NPP-GPCR networks with short-

1004 range connections. Colors given by assortativity like Figure 4; yellow indicates local, green pervasive,

1005 blue integrative and red promiscuous networks.

1006

1007 **Supplementary Figure S3.** Graphical representations of the 91 NPP-GPCR networks with mid-range

1008 connections. Networks are sorted in the same order as in Supplementary Figure 2 for comparison.

1009 Colors given by assortativity as defined in Figure 4, yellow indicates local, green pervasive, blue

1010 integrative and red promiscuous networks. Networks that change assortativity between short and mid-

1011 range spatial models of neuropeptide transmission are highlighted.

1012

1013 **Supplementary Figure 4.** Adjacency matrix representation of the long and mid-range neuropeptide

1014 networks. A) Model 1, aggregate matrix of the long-range neuropeptide connections, the histogram

1015 lining the matrix indicate the number of GPCR or NPP that the neuron expresses. B) Model 2,

1016 aggregate matrix of the mid-range neuropeptide connections.

1017

Supplementary Figure 5. Co-expression analysis of NPP and GPCR genes. A) Hierarchical

clustering matrix showing NPP – GPCR pairs with significant co-expression. The y axis indicates

GPCR and the x axis indicates NPP expression. The significance of their co-expression is indicated by

the legend. B) Adjacency matrix representation of the co-expression network. The nodes represent

NPP – GPCR pairs with significant neuronal coexpression, and edges are formed when the NPP of

one node is the ligand of the GPCR of another node. C) Hierarchical clustering matrix of the NPP vs

NPP co-expression. D) Hierarchical clustering matrix of the GPCR vs. GPCR co-expression. E)

Visual representation of one of the possible signaling cascades based on the co-expression network.

DMSR-3 co-expresses with FLP-34 and NLP-54 releasing neurons, which means that secretion could

be activated by FLP-14 binding to DMSR-3.

1028

Supplementary Figure 6. Autocrine connections sensitivity analysis. A) comparison of co-

expression of cognate NPP-GPCR pairs in the 4 different combinations of CeNGEN and

deorphanization datasets with different stringencies (CeNGEN threshold 2 or 4; EC_{50} threshold 1 µM

or 500 nM), each depicted in a different color. B) Number of NPP-GPCR co-expressed per fraction of

neurons in these 4 different combinations of datasets. C) Number of neurons with co-expression for

the FLP or NLP neuropeptides show that FLP are more prevalent. D) Comparison of number of

autocrine connections between specific and promiscuous NPP-GPCR pairs, showing that promiscuous

pairs tend to have more autocrine connections. E) Correlation of autocrine connections with gap

junction and monoamine networks. F) Number of autocrine connections in each motorneuron for each

of the main promiscuous NPP-GPCR pairs present in those neurons.

1039

1040

References

1.   Eichler, K., Li, F., Litwin-Kumar, A., Park, Y., Andrade, I., Schneider-Mizell, C.M.,

Saumweber, T., Huser, A., Eschbach, C., Gerber, B., et al. (2017). The complete connectome

of a learning and memory centre in an insect brain. Nature *548*, 175–182.

10.1038/nature23455.

2.   Scheffer, L.K., Xu, C.S., Januszewski, M., Lu, Z., Takemura, S., Hayworth, K.J., Huang, G.B.,

Shinomiya, K., Maitlin-Shepard, J., Berg, S., et al. (2020). A connectome and analysis of the

adult Drosophila central brain. Elife *9*, e57443. 10.7554/eLife.57443.

3.   Verasztó, C., Jasek, S., Gühmann, M., Shahidi, R., Ueda, N., Beard, J.D., Mendes, S., Heinz,

K., Bezares-Calderón, L.A., Williams, E., et al. (2020). Whole-animal connectome and cell-

type complement of the three-segmented *Platynereis dumerilii* larv.

4.   Hulse, B.K., Haberkern, H., Franconville, R., Turner-Evans, D., Takemura, S., Wolff, T.,

1053   Noorman, M., Dreher, M., Dan, C., Parekh, R., et al. (2021). A connectome of the Drosophila

1054   central complex reveals network motifs suitable for flexible navigation and context-dependent

1055   action selection. Elife *10*, e66039. 10.7554/eLife.66039.

1056   5.   Ryan, K., Lu, Z., and Meinertzhagen, I.A. (2016). The CNS connectome of a tadpole larva of

1057   Ciona intestinalis (L.) highlights sidedness in the brain of a chordate sibling. Elife *5*, e16962.

1058   10.7554/eLife.16962.

1059   6.   Kunst, M., Laurell, E., Mokayes, N., Kramer, A., Kubo, F., Fernandes, A.M., Förster, D., Dal

1060   Maschio, M., and Baier, H. (2019). A Cellular-Resolution Atlas of the Larval Zebrafish Brain.

1061   Neuron *103*, 21-38.e5. 10.1016/j.neuron.2019.04.034.

1062   7.   Oh, S.W., Harris, J.A., Ng, L., Winslow, B., Cain, N., Mihalas, S., Wang, Q., Lau, C., Kuan,

1063   L., Henry, A.M., et al. (2014). A mesoscale connectome of the mouse brain. Nature *508*, 207–

1064   214. 10.1038/nature13186.

1065   8.   Nässel, D.R. (2009). Neuropeptide signaling near and far: how localized and timed is the

1066   action of neuropeptides in brain circuits? Invertebrate Neuroscience *9*, 57–75.

1067   10.1007/s10158-009-0090-1.

1068   9.   van den Pol, A.N. (2012). Neuropeptide Transmission in Brain Circuits. Neuron *76*, 98–115.

1069   10.1016/j.neuron.2012.09.014.

1070   10.   Atkinson, L.E., Liu, Y., McKay, F., Vandewyer, E., Viau, C., Irvine, A., Rosa, B.A., Li, Z.,

1071   Liang, Q., Marks, N.J., et al. (2021). *Ascaris suum* Informs Extrasynaptic Volume

1072   Transmission in Nematodes. ACS Chem Neurosci *12*, 3176–3188.

1073   10.1021/acschemneuro.1c00281.

1074   11.   Nurrish, S. (2014). Dense Core Vesicle Release: Controlling the Where as Well as the When.

1075   Genetics *196*, 601–604. 10.1534/genetics.113.159905.

1076   12.   Marder, E. (2012). Neuromodulation of Neuronal Circuits: Back to the Future. Neuron *76*, 1–

1077   11. 10.1016/j.neuron.2012.09.010.

1078   13.   Oranth, A., Schultheis, C., Tolstenkov, O., Erbguth, K., Nagpal, J., Hain, D., Brauner, M.,

1079   Wabnig, S., Costa, W.S., McWhirter, R.D., et al. (2018). Food Sensation Modulates

Locomotion by Dopamine and Neuropeptide Signaling in a Distributed Neuronal Network.

Neuron *100*, 1414-1428.e10. 10.1016/j.neuron.2018.10.024.

14.   Chew, Y.L., Tanizawa, Y., Cho, Y., Zhao, B., Yu, A.J., Ardiel, E.L., Rabinowitch, I., Bai, J.,

Rankin, C.H., Lu, H., et al. (2018). An Afferent Neuropeptide System Transmits

Mechanosensory Signals Triggering Sensitization and Arousal in C. elegans. Neuron *99*, 1233-

1246.e6. 10.1016/j.neuron.2018.08.003.

15.   Lin, S., Senapati, B., and Tsao, C.-H. (2019). Neural basis of hunger-driven behaviour in

Drosophila. Open Biol *9*, 180259. 10.1098/rsob.180259.

16.   Ramachandran, S., Banerjee, N., Bhattacharya, R., Lemons, M.L., Florman, J., Lambert, C.M.,

Touroutine, D., Alexander, K., Schoofs, L., Alkema, M.J., et al. (2021). A conserved

neuropeptide system links head and body motor circuits to enable adaptive behavior. Elife *10*,

e71747. 10.7554/eLife.71747.

17.   Melzer, S., Newmark, E.R., Mizuno, G.O., Hyun, M., Philson, A.C., Quiroli, E., Righetti, B.,

Gregory, M.R., Huang, K.W., Levasseur, J., et al. (2021). Bombesin-like peptide recruits

disinhibitory cortical circuits and enhances fear memories. Cell *184*, 5622-5634.e25.

10.1016/j.cell.2021.09.013.

18.   Zhong, W., Barde, S., Mitsios, N., Adori, C., Oksvold, P., Feilitzen, K. von, O'Leary, L.,

Csiba, L., Hortobágyi, T., Szocsics, P., et al. (2022). The neuropeptide landscape of human

prefrontal cortex. Proceedings of the National Academy of Sciences *119*, e2123146119.

10.1073/pnas.2123146119.

19.   Bargmann, C.I. (2012). Beyond the connectome: How neuromodulators shape neural circuits.

BioEssays *34*, 458–465. 10.1002/bies.201100185.

20.   Schoofs, L., de Loof, A., and van Hiel, M.B. (2017). Neuropeptides as Regulators of Behavior

in Insects. Annu Rev Entomol *62*, 35–52. 10.1146/annurev-ento-031616-035500.

21.   Bhat, U.S., Shahi, N., Surendran, S., and Babu, K. (2021). Neuropeptides and Behaviors: How

Small Peptides Regulate Nervous System Function and Behavioral Outputs. Front Mol

Neurosci *14*.

22.   Jékely, G. (2021). The chemical brain hypothesis for the origin of nervous systems.

1108   Philosophical Transactions of the Royal Society B: Biological Sciences *376*, 20190761.

1109   10.1098/rstb.2019.0761.

1110   23.   Yañez-Guerra, L.A., Thiel, D., and Jékely, G. (2022). Premetazoan Origin of Neuropeptide

1111   Signaling. Mol Biol Evol *39*, msac051. 10.1093/molbev/msac051.

1112   24.   van Bael, S., Watteyne, J., Boonen, K., de Haes, W., Menschaert, G., Ringstad, N., Horvitz,

1113   H.R., Schoofs, L., Husson, S.J., and Temmerman, L. (2018). Mass spectrometric evidence for

1114   neuropeptide-amidating enzymes in. Journal of Biological Chemistry *293*, 6052–6063.

1115   10.1074/jbc.RA117.000731.

1116   25.   Frooninckx, L., van Rompay, L., Temmerman, L., van Sinay, E., Beets, I., Janssen, T.,

1117   Husson, S., and Schoofs, L. (2012). Neuropeptide GPCRs in C. elegans. Front Endocrinol

1118   (Lausanne) *3*.

1119   26.   Jékely, G. (2013). Global view of the evolution and diversity of metazoan neuropeptide

1120   signaling. Proceedings of the National Academy of Sciences *110*, 8702–8707.

1121   10.1073/pnas.1221833110.

1122   27.   Rizzo, M.J., and Johnson, E.C. (2020). Homodimerization of Drosophila Class A neuropeptide

1123   GPCRs: Evidence for conservation of GPCR dimerization throughout metazoan evolution.

1124   Biochem Biophys Res Commun *523*, 322–327. 10.1016/j.bbrc.2019.12.019.

1125   28.   Rogers, C., Reale, V., Kim, K., Chatwin, H., Li, C., Evans, P., and de Bono, M. (2003).

1126   Inhibition of Caenorhabditis elegans social feeding by FMRFamide-related peptide activation

1127   of NPR-1. Nat Neurosci *6*, 1178–1185. 10.1038/nn1140.

1128   29.   Beets, I., Janssen, T., Meelkop, E., Temmerman, L., Suetens, N., Rademakers, S., Jansen, G.,

1129   and Schoofs, L. (2012). Vasopressin/Oxytocin-Related Signaling Regulates Gustatory

1130   Associative Learning in *C. elegans*. Science (1979) *338*, 543–545. 10.1126/science.1226860.

1131   30.   Iannacone, M.J., Beets, I., Lopes, L.E., Churgin, M.A., Fang-Yen, C., Nelson, M.D., Schoofs,

1132   L., and Raizen, D.M. (2017). The RFamide receptor DMSR-1 regulates stress-induced sleep in

1133   C. elegans. Elife *6*, e19837. 10.7554/eLife.19837.

1134   31.   Nässel, D.R., and Zandawala, M. (2019). Recent advances in neuropeptide signaling in

1135   Drosophila, from genes to physiology and behavior. Prog Neurobiol *179*, 101607.

1136   10.1016/j.pneurobio.2019.02.003.

1137   32.   Watteyne, J., Peymen, K., van der Auwera, P., Borghgraef, C., Vandewyer, E., van Damme,

1138   S., Rutten, I., Lammertyn, J., Jelier, R., Schoofs, L., et al. (2020). Neuromedin U signaling

1139   regulates retrieval of learned salt avoidance in a C. elegans gustatory circuit. Nat Commun *11*,

1140   2076. 10.1038/s41467-020-15964-9.

1141   33.   Topilko, T., Diaz, S.L., Pacheco, C.M., Verny, F., Rousseau, C. v, Kirst, C., Deleuze, C.,

1142   Gaspar, P., and Renier, N. (2022). Edinger-Westphal peptidergic neurons enable maternal

1143   preparatory nesting. Neuron *110*, 1385-1399.e8. 10.1016/j.neuron.2022.01.012.

1144   34.   Davenport, A.P., Scully, C.C.G., de Graaf, C., Brown, A.J.H., and Maguire, J.J. (2020).

1145   Advances in therapeutic peptides targeting G protein-coupled receptors. Nat Rev Drug Discov

1146   *19*, 389–413. 10.1038/s41573-020-0062-z.

1147   35.   Coleman, P.J., Gotter, A.L., Herring, W.J., Winrow, C.J., and Renger, J.J. (2017). The

1148   Discovery of Suvorexant, the First Orexin Receptor Drug for Insomnia. Annu Rev Pharmacol

1149   Toxicol *57*, 509–533. 10.1146/annurev-pharmtox-010716-104837.

1150   36.   Hargreaves, R., Ferreira, J.C.A., Hughes, D., Brands, J., Hale, J., Mattson, B., and Mills, S.

1151   (2011). Development of aprepitant, the first neurokinin-1 receptor antagonist for the

1152   prevention of chemotherapy-induced nausea and vomiting. Ann N Y Acad Sci *1222*, 40–48.

1153   10.1111/j.1749-6632.2011.05961.x.

1154   37.   Zhang, Z.Q., and Hölscher, C. (2020). GIP has neuroprotective effects in Alzheimer and

1155   Parkinson's disease models. Peptides (N.Y.) *125*, 170184. 10.1016/j.peptides.2019.170184.

1156   38.   Hoyer, D., and Bartfai, T. (2012). Neuropeptides and Neuropeptide Receptors: Drug Targets,

1157   and Peptide and Non-Peptide Ligands: a Tribute to Prof. Dieter Seebach. Chem Biodivers *9*,

1158   2367–2387. 10.1002/cbdv.201200288.

1159   39.   Yao, Z., van Velthoven, C.T.J., Nguyen, T.N., Goldy, J., Sedeno-Cortes, A.E., Baftizadeh, F.,

1160   Bertagnolli, D., Casper, T., Chiang, M., Crichton, K., et al. (2021). A taxonomy of

transcriptomic cell types across the isocortex and hippocampal formation. Cell *184*, 3222-

3241.e26. 10.1016/j.cell.2021.04.021.

40.   Taylor, S.R., Santpere, G., Weinreb, A., Barrett, A., Reilly, M.B., Xu, C., Varol, E.,

Oikonomou, P., Glenwinkel, L., McWhirter, R., et al. (2021). Molecular topography of an

entire nervous system. Cell *184*, 4329-4347.e23. 10.1016/j.cell.2021.06.023.

41.   Li, H., Janssens, J., de Waegeneer, M., Kolluru, S.S., Davie, K., Gardeux, V., Saelens, W.,

David, F.P.A., Brbić, M., Spanier, K., et al. (2022). Fly Cell Atlas: A single-nucleus

transcriptomic atlas of the adult fruit fly. Science (1979) *375*, eabk2432.

10.1126/science.abk2432.

42.   Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J.,

Garren, E., Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct transcriptomic

cell types across neocortical areas. Nature *563*, 72–78. 10.1038/s41586-018-0654-5.

43.   Smith, S.J., Sümbül, U., Graybuck, L.T., Collman, F., Seshamani, S., Gala, R., Gliko, O.,

Elabbady, L., Miller, J.A., Bakken, T.E., et al. (2019). Single-cell transcriptomic evidence for

dense intracortical neuropeptide networks. Elife *8*, e47889. 10.7554/eLife.47889.

44.   Smith, S.J. (2021). Transcriptomic evidence for dense peptidergic networks within forebrains

of four widely divergent tetrapods. Curr Opin Neurobiol *71*, 100–109.

10.1016/j.conb.2021.09.011.

45.   Bentley, B., Branicky, R., Barnes, C.L., Chew, Y.L., Yemini, E., Bullmore, E.T., Vértes, P.E.,

and Schafer, W.R. (2016). The Multilayer Connectome of Caenorhabditis elegans. PLoS

Comput Biol *12*, e1005283. 10.1371/journal.pcbi.1005283.

46.   Cook, S.J., Jarrell, T.A., Brittin, C.A., Wang, Y., Bloniarz, A.E., Yakovlev, M.A., Nguyen,

K.C.Q., Tang, L.T.-H., Bayer, E.A., Duerr, J.S., et al. (2019). Whole-animal connectomes of

both Caenorhabditis elegans sexes. Nature *571*, 63–71. 10.1038/s41586-019-1352-7.

47.   Varshney, L.R., Chen, B.L., Paniagua, E., Hall, D.H., and Chklovskii, D.B. (2011). Structural

Properties of the Caenorhabditis elegans Neuronal Network. PLoS Comput Biol *7*, e1001066.

10.1371/journal.pcbi.1001066.

48.   White, J.G., Southgate, E., Thomson, J.N., and Brenner, S. (1986). The structure of the

1189   nervous system of the nematode *Caenorhabditis elegans*. Philosophical Transactions of the

1190   Royal Society of London. B, Biological Sciences *314*, 1–340. 10.1098/rstb.1986.0056.

1191   49.   Watts, D.J., and Strogatz, S.H. (1998). Collective dynamics of 'small-world' networks. *393*, 3.

1192   50.   Bassett, D.S., and Bullmore, E.T. (2017). Small-World Brain Networks Revisited. The

1193   Neuroscientist *23*, 499–516. 10.1177/1073858416667720.

1194   51.   Reigl, M., Alon, U., and Chklovskii, D.B. (2004). Search for computational modules in the C.

1195   elegans brain. BMC Biol *2*, 25. 10.1186/1741-7007-2-25.

1196   52.   Pavlovic, D.M., Vértes, P.E., Bullmore, E.T., Schafer, W.R., and Nichols, T.E. (2014).

1197   Stochastic Blockmodeling of the Modules and Core of the Caenorhabditis elegans

1198   Connectome. PLoS One *9*, e97584. 10.1371/journal.pone.0097584.

1199   53.   Pathak, A., Chatterjee, N., and Sinha, S. (2020). Developmental trajectory of Caenorhabditis

1200   elegans nervous system governs its structural organization. PLoS Comput Biol *16*, e1007602.

1201   10.1371/journal.pcbi.1007602.

1202   54.   Helm, A., Blevins, A.S., and Bassett, D.S. (2021). The growing topology of the C. elegans

1203   connectome (bioRxiv).

1204   55.   Towlson, E.K., Vertes, P.E., Ahnert, S.E., Schafer, W.R., and Bullmore, E.T. (2013). The Rich

1205   Club of the C. elegans Neuronal Connectome. Journal of Neuroscience *33*, 6380–6387.

1206   10.1523/JNEUROSCI.3784-12.2013.

1207   56.   van den Heuvel, M.P., and Sporns, O. (2011). Rich-Club Organization of the Human

1208   Connectome. Journal of Neuroscience *31*, 15775–15786. 10.1523/JNEUROSCI.3539-11.2011.

1209   57.   Schroeter, M.S., Charlesworth, P., Kitzbichler, M.G., Paulsen, O., and Bullmore, E.T. (2015).

1210   Emergence of Rich-Club Topology and Coordinated Dynamics in Development of

1211   Hippocampal Functional Networks In Vitro. Journal of Neuroscience *35*, 5459–5470.

1212   10.1523/JNEUROSCI.4259-14.2015.

1213   58.   Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., and Alon, U. (2002).

1214   Network Motifs: Simple Building Blocks of Complex Networks. Science (1979) *298*, 824–

1215   827. 10.1126/science.298.5594.824.

1216   59.   Alon, U. (2007). Network motifs: theory and experimental approaches. Nat Rev Genet *8*, 450–

1217   461. 10.1038/nrg2102.

1218   60.   Matelsky, J.K., Norman-Tenazas, R., Davenport, F., Reilly, E.P., and Gray-Roncal, W. (2021).

1219   Circuit motifs and graph properties of connectome development in C. elegans (bioRxiv).

1220   61.   McKay, F.M., McCoy, C.J., Crooks, B., Marks, N.J., Maule, A.G., Atkinson, L.E., and

1221   Mousley, A. (2022). In silico analyses of neuropeptide-like protein (NLP) profiles in parasitic

1222   nematodes. Int J Parasitol *52*, 77–85. 10.1016/j.ijpara.2021.07.002.

1223   62.   Hobert, O. (2013). The neuronal genome of Caenorhabditis elegans. WormBook, 1–106.

1224   10.1895/wormbook.1.161.1.

1225   63.   Foster, S.R., Hauser, A.S., Vedel, L., Strachan, R.T., Huang, X.-P., Gavin, A.C., Shah, S.D.,

1226   Nayak, A.P., Haugaard-Kedström, L.M., Penn, R.B., et al. (2019). Discovery of Human

1227   Signaling Systems: Pairing Peptides to G Protein-Coupled Receptors. Cell *179*, 895-908.e21.

1228   10.1016/j.cell.2019.10.010.

1229   64.   Beets, I., Zels, S., Vandewyer, E., Demeulemeester, J., Caers, J., Baytemur, E., Schafer, W.R.,

1230   Vértes, P.E., Mirabeau, O., and Schoofs, L. (2022). System-wide discovery of neuropeptide-

1231   GPCR pairs in C. elegans. Preprint in bioRxiv. https://doi.org/10.1101/2022.10.30.514428.

1232   65.   Albertson, D.G., and Thomson, J.N. The pharynx of Caenorhabditis elegans. Philos Trans R

1233   Soc Lond B Biol Sci *275*, 299–325. https://doi.org/10.1098/rstb.1976.0085.

1234   66.   Witvliet, D., Mulcahy, B., Mitchell, J.K., Meirovitch, Y., Berger, D.R., Wu, Y., Liu, Y., Koh,

1235   W.X., Parvathala, R., Holmyard, D., et al. (2021). Connectomes across development reveal

1236   principles of brain maturation. Nature *596*, 257–261. 10.1038/s41586-021-03778-8.

1237   67.   Moyle, M.W., Barnes, K.M., Kuchroo, M., Gonopolskiy, A., Duncan, L.H., Sengupta, T.,

1238   Shao, L., Guo, M., Santella, A., Christensen, R., et al. (2021). Structural and developmental

1239   principles of neuropil assembly in C. elegans. Nature *591*, 99–104. 10.1038/s41586-020-

1240   03169-5.

1241   68.   Brittin, C.A., Cook, S.J., Hall, D.H., Emmons, S.W., and Cohen, N. (2021). A multi-scale

1242   brain map derived from whole-brain volumetric reconstructions. Nature *591*, 105–110.

1243   10.1038/s41586-021-03284-x.

69. Maier, W., Adilov, B., Regenass, M., and Alcedo, J. (2010). A Neuromedin U Receptor Acts

with the Sensory System to Modulate Food Type-Dependent Effects on C. elegans Lifespan.

PLoS Biol *8*, e1000376. 10.1371/journal.pbio.1000376.

70. Pocock, R., and Hobert, O. (2010). Hypoxia activates a latent circuit for processing gustatory

information in C. elegans. Nat Neurosci *13*, 610–614. 10.1038/nn.2537.

71. van Sinay, E., Mirabeau, O., Depuydt, G., van Hiel, M.B., Peymen, K., Watteyne, J., Zels, S.,

Schoofs, L., and Beets, I. (2017). Evolutionarily conserved TRH neuropeptide pathway

regulates growth in *Caenorhabditis elegans*. Proceedings of the National Academy of Sciences

114. 10.1073/pnas.1617392114.

72. Cook, S.J., Crouse, C.M., Yemini, E., Hall, D.H., Emmons, S.W., and Hobert, O. (2020). The

connectome of the Caenorhabditis elegans pharynx. J Comp Neurol *528*, 2767–2784.

10.1002/cne.24932.

73. Vidal, B., Gulez, B., Cao, W.X., Leyva-Díaz, E., Reilly, M.B., Tekieli, T., and Hobert, O.

(2022). The enteric nervous system of the C. elegans pharynx is specified by the Sine oculis-

like homeobox gene ceh-34. Elife *11*, e76003. 10.7554/eLife.76003.

74. Kaplan, H.S., Nichols, A.L.A., and Zimmer, M. (2018). Sensorimotor integration in

Caenorhabditis elegans: A reappraisal towards dynamic and distributed computations.

Philosophical Transactions of the Royal Society B: Biological Sciences *373*.

10.1098/rstb.2017.0371.

75. Pereira, L., Kratsios, P., Serrano-Saiz, E., Sheftel, H., Mayo, A.E., Hall, D.H., White, J.G.,

LeBoeuf, B., Garcia, L.R., Alon, U., et al. (2015). A cellular and regulatory map of the

cholinergic nervous system of C. elegans. Elife *4*, e12432. 10.7554/eLife.12432.

76. Niu, L., Li, Y., Zong, P., Liu, P., Shui, Y., Chen, B., and Wang, Z.-W. (2020). Melatonin

promotes sleep by activating the BK channel in C. elegans. Proc Natl Acad Sci U S A *117*,

25128–25137. 10.1073/pnas.2010928117.

77. Betzel, R.F., Medaglia, J.D., and Bassett, D.S. (2018). Diversity of meso-scale architecture in

human and non-human connectomes. Nat Commun *9*, 346. 10.1038/s41467-017-02681-z.

78.   Wen, Q., Po, M.D., Hulme, E., Chen, S., Liu, X., Kwok, S.W., Gershow, M., Leifer, A.M.,

1272   Butler, V., Fang-Yen, C., et al. (2012). Proprioceptive Coupling within Motor Neurons Drives

1273   C. elegans Forward Locomotion. Neuron *76*, 750–761. 10.1016/j.neuron.2012.08.039.

1274   79.   Schafer, W.R. (2015). Mechanosensory molecules and circuits in C. elegans. Pflugers Arch

1275   *467*, 39–48. 10.1007/s00424-014-1574-3.

1276   80.   Fenk, L.A., and de Bono, M. (2017). Memory of recent oxygen experience switches

1277   pheromone valence in *Caenorhabditis elegans*. Proceedings of the National Academy of

1278   Sciences *114*, 4195–4200. 10.1073/pnas.1618934114.

1279   81.   Guillermin, M.L., Carrillo, M.A., and Hallem, E.A. (2017). A Single Set of Interneurons

1280   Drives Opposite Behaviors in C. elegans. Current Biology *27*, 2630-2639.e6.

1281   10.1016/J.CUB.2017.07.023.

1282   82.   Laurent, P., Soltesz, Z., Nelson, G.M., Chen, C., Arellano-Carbajal, F., Levy, E., and de Bono,

1283   M. (2015). Decoding a neural circuit controlling global animal state in C. Elegans. Elife *2015*.

1284   83.   Busch, K.E., Laurent, P., Soltesz, Z., Murphy, R.J., Faivre, O., Hedwig, B., Thomas, M.,

1285   Smith, H.L., and de Bono, M. (2012). Tonic signaling from O2 sensors sets neural circuit

1286   activity and behavioral state. Nat Neurosci *15*, 581–591. 10.1038/nn.3061.

1287   84.   Haspel, G., Deng, L., Harreguy, M.B., and Tanvir, Z. (2020). Elegantly. The Neural Control of

1288   Movement, 3–29. 10.1016/B978-0-12-816477-8.00001-6.

1289   85.   Tolstenkov, O., Van der Auwera, P., Costa, W.S., Bazhanova, O., Gemeinhardt, T.M., Bergs,

1290   A.C.F., and Gottschalk, A. (2018). Functionally asymmetric motor neurons contribute to

1291   coordinating locomotion of Caenorhabditis elegans. Elife *7*. 10.7554/ELIFE.34997.

1292   86.   Butler, V.J., Branicky, R., Yemini, E., Liewald, J.F., Gottschalk, A., Kerr, R.A., Chklovskii,

1293   D.B., and Schafer, W.R. (2015). A consistent muscle activation strategy underlies crawling

1294   and swimming in Caenorhabditis elegans. J R Soc Interface *12*. 10.1098/RSIF.2014.0963.

1295   87.   Faumont, S., Rondeau, G., Thiele, T.R., Lawton, K.J., McCormick, K.E., Sottile, M.,

1296   Griesbeck, O., Heckscher, E.S., Roberts, W.M., Doe, C.Q., et al. (2011). An Image-Free Opto-

1297   Mechanical System for Creating Virtual Environments and Imaging Neuronal Activity in

1298   Freely Moving Caenorhabditis elegans. PLoS One *6*, e24666. 10.1371/journal.pone.0024666.

88.   Wen, Q., Po, M., Hulme, E., Chen, S., Liu, X., Kwok, S., Gershow, M., Leifer, A., Butler, V.,

Fang-Yen, C., et al. (2012). Proprioceptive coupling within motor neurons drives C. elegans

forward locomotion. Neuron *76*, 750–761. 10.1016/J.NEURON.2012.08.039.

89.   Gao, S., Guan, S.A., Fouad, A.D., Meng, J., Kawano, T., Huang, Y.-C., Li, Y., Alcaire, S.,

Hung, W., Lu, Y., et al. (2018). Excitatory motor neurons are local oscillators for backward

locomotion. Elife *7*. 10.7554/eLife.29915.

90.   Worrell, J.C., Rumschlag, J., Betzel, R.F., Sporns, O., and Mišić, B. (2017). Optimized

connectome architecture for sensory-motor integration. Netw Neurosci *1*, 415–430.

10.1162/NETN_a_00022.

91.   Williams, E.A., Verasztó, C., Jasek, S., Conzelmann, M., Shahidi, R., Bauknecht, P.,

Mirabeau, O., and Jékely, G. (2017). Synaptic and peptidergic connectome of a neurosecretory

center in the annelid brain. Elife *6*, e26349. 10.7554/eLife.26349.

92.   Turek, M., Besseling, J., Spies, J.-P., König, S., and Bringmann, H. (2016). Sleep-active

neuron specification and sleep induction require FLP-11 neuropeptides to systemically induce

sleep. Elife *5*, e12499. 10.7554/eLife.12499.

93.   Xu, Y., Zhang, L., Liu, Y., Topalidou, I., Hassinan, C., Ailion, M., Zhao, Z., Wang, T., Chen,

Z., and Bai, J. (2021). Dopamine receptor DOP-1 engages a sleep pathway to modulate

swimming in C. elegans. iScience *24*, 102247. 10.1016/j.isci.2021.102247.

94.   Priest, M.F., Freda, S.N., Badong, D., Dumrongprechachan, V., and Kozorovitskiy, Y. (2021).

Peptidergic modulation of fear responses by the Edinger-Westphal nucleus (bioRxiv).

95.   Kim, M.W., Wang, W., Sanchez, M.I., Coukos, R., von Zastrow, M., and Ting, A.Y. (2017).

Time-gated detection of protein-protein interactions with transcriptional readout. Elife *6*,

e30233. 10.7554/eLife.30233.

96.   Kim, C.K., Cho, K.F., Kim, M.W., and Ting, A.Y. (2019). Luciferase-LOV BRET enables

versatile and specific transcriptional readout of cellular protein-protein interactions. Elife *8*,

e43826. 10.7554/eLife.43826.

97.   Smith, S.J., Smbül, U., Graybuck, L.T., Collman, F., Seshamani, S., Gala, R., Gliko, O.,

1326   Elabbady, L., Miller, J.A., Bakken, T.E., et al. (2019). Single-cell transcriptomic evidence for

1327   dense intracortical neuropeptide networks. Elife 8, 1–35. 10.7554/eLife.47889.

1328   98.   Chen, R., Wu, X., Jiang, L., and Zhang, Y. (2017). Single-Cell RNA-Seq Reveals

1329   Hypothalamic Cell Diversity. Cell Rep 18, 3227–3241. 10.1016/J.CELREP.2017.03.004.

1330   99.   Terauchi, A., Johnson-Venkatesh, B., Bullock, B., Lehtinen, M., and Umemori, H. (2016).

1331   Retrograde fibroblast growth factor 22 (FGF22) signaling regulates insulin-like growth factor

1332   2 (IGF2) expression for activity-dependent synapse stabilization in the mammalian brain. Elife

1333   5. 10.7554/ELIFE.12151.

1334   100.   Ghosh, D.D., Sanders, T., Hong, S., McCurdy, L.Y., Chase, D.L., Cohen, N., Koelle, M.R.,

1335   and Nitabach, M.N. (2016). Neural Architecture of Hunger-Dependent Multisensory Decision

1336   Making in C. elegans. Neuron 92, 1049–1062. 10.1016/j.neuron.2016.10.030.

1337   101.   Root, C.M., Ko, K.I., Jafari, A., and Wang, J.W. (2011). Presynaptic facilitation by

1338   neuropeptide signaling mediates odor-driven food search. Cell 145, 133–144.

1339   10.1016/j.cell.2011.02.008.

1340   102.   Horowitz, L., Brandt, J., and Ringstad, N. (2019). Repression of an activity-dependent

1341   autocrine insulin signal is required for sensory neuron development in C. elegans.

1342   Development 146. 10.1242/DEV.182873.

1343   103.   Mardinly, A., Spiegel, I., Patrizi, A., Centofante, E., Bazinet, J., Tzeng, C., Mandel-Brehm, C.,

1344   Harmin, D., Adesnik, H., Fagiolini, M., et al. (2016). Sensory experience regulates cortical

1345   inhibition by inducing IGF1 in VIP neurons. Nature 531, 371–375. 10.1038/NATURE17187.

1346   104.   Choi, C., Cao, G., Tanenhaus, A., McCarthy, E., Jung, M., Schleyer, M., Shang, Y., Rosbash,

1347   M., Yin, J., and Nitabach, M. (2012). Autoreceptor control of peptide/neurotransmitter

1348   corelease from PDF neurons determines allocation of circadian activity in drosophila. Cell Rep

1349   2, 332–344. 10.1016/J.CELREP.2012.06.021.

1350   105.   Yoshioka, J., Prince, R., Huang, H., Perkins, S., Cruz, F., MacGillivray, C., Lauffenburger, D.,

1351   and Lee, R. (2005). Cardiomyocyte hypertrophy and degradation of connexin43 through

spatially restricted autocrine/paracrine heparin-binding EGF. Proc Natl Acad Sci U S A *102*,

10622–10627. 10.1073/PNAS.0501198102.

106.	Tse, L., and Wong, Y. (2019). GPCRs in Autocrine and Paracrine Regulations. Front

Endocrinol (Lausanne) *10*. 10.3389/FENDO.2019.00428.

107.	Yemini, E., Jucikas, T., Grundy, L.J., Brown, A.E.X., and Schafer, W.R. (2013). A database of

Caenorhabditis elegans behavioral phenotypes. Nat Methods *10*, 877–879.

10.1038/nmeth.2560.

108.	Brown, A.E.X., Yemini, E.I., Grundy, L.J., Jucikas, T., and Schafer, W.R. (2013). A

dictionary of behavioral motifs reveals clusters of genes affecting Caenorhabditis elegans

locomotion. Proceedings of the National Academy of Sciences *110*, 791–796.

10.1073/pnas.1211447110.

109.	Avery, L., and Horvitz, H.R. (1989). Pharyngeal pumping continues after laser killing of the

pharyngeal nervous system of C. elegans. Neuron *3*, 473–485. 10.1016/0896-6273(89)90206-

7.

110.	Forrester, W.C., and Garriga, G. (1997). Genes necessary for C. elegans cell and growth cone

migrations. Development *124*, 1831–1843. 10.1242/dev.124.9.1831.

111.	Yemini, E., Lin, A., Nejatbakhsh, A., Varol, E., Sun, R., Mena, G.E., Samuel, A.D.T.,

Paninski, L., Venkatachalam, V., and Hobert, O. (2021). NeuroPAL: A Multicolor Atlas for

Whole-Brain Neuronal Identification in C. elegans. Cell *184*, 272-288.e11.

10.1016/j.cell.2020.12.012.

112.	Hall, D.H., and Altun, Z.F. (2011). Nervous system, general description. WormAtlas: The

Handbook of C.elegans Anatomy. doi:10.3908/wormatlas.1.18.

113.	Rubinov, M., and Sporns, O. (2010). Complex network measures of brain connectivity: Uses

and interpretations. Neuroimage *52*, 1059–1069. 10.1016/j.neuroimage.2009.10.003.

114.	Fisher, R.A. (1934). Statistical methods for research workers, 5th ed. (Edinburgh).

115.	Storey, J.D. (2002). A direct approach to false discovery rates. J R Stat Soc Series B Stat

Methodol *64*, 479–498. 10.1111/1467-9868.00346.

116.   Bounova, G., and de Weck, O. (2012). Overview of metrics and their correlation patterns for

1380   multiple-metric topology analysis on heterogeneous graph ensembles. Phys Rev E *85*, 016117.

1381   10.1103/PhysRevE.85.016117.

1382   117.   de Domenico, M., Porter, M.A., and Arenas, A. (2015). MuxViz: a tool for multilayer analysis

1383   and visualization of networks. J Complex Netw *3*, 159–176. 10.1093/comnet/cnu038.

1384   118.   Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,

1385   Schwikowski, B., and Ideker, T. (2003). Cytoscape: A Software Environment for Integrated

1386   Models of Biomolecular Interaction Networks. Genome Res *13*, 2498–2504.

1387   10.1101/gr.1239303.

1388   119.   Bhardwaj, A., Thapliyal, S., Dahiya, Y., and Babu, K. (2018). FLP-18 Functions through the

1389   G-Protein-Coupled Receptors NPR-1 and NPR-4 to Modulate Reversal Length in

1390   Caenorhabditis elegans. The Journal of Neuroscience *38*, 4641–4654.

1391   10.1523/JNEUROSCI.1955-17.2018.

1392   120.   Lindemans, M., Liu, F., Janssen, T., Husson, S.J., Mertens, I., Gäde, G., and Schoofs, L.

1393   (2009). Adipokinetic hormone signaling through the gonadotropin-releasing hormone receptor

1394   modulates egg-laying in Caenorhabditis elegans. Proceedings of the National Academy of

1395   Sciences *106*, 1642–1647. 10.1073/pnas.0809881106.

1396

A

HEAD
Nerve ring
Amphid commissures
Anterior ventral nerve cord
Deirid commissures

MIDBODY
Dorsal sublateral left and right
Dorsal Nerve Cord
Ventral sublateral left and right
Ventral Nerve cord
Canal associated left and right

TAIL
Preanal ganglion
Canal associated
Tail tip bundle

B

CAN neurons

GPCR expression

Pharynx | Sensory neurons | Interneurons | Motorneurons

C

CAN neurons

Cognate NPP expression

Pharynx | Sensory neurons | Interneurons | Motorneurons

Type of connection required to explain expression

Mid range | Between strata | Within bundle (outside NR) | Within stratum

A

Local

Pervasive

Broadcasting

Integrative

○ CAPA-1 ● NMUR-1

○ FLP-18 ● NPR-5

○ FLP-1 ● FRPR-7

○ NLP-13 ● B0334.6

B

● DMSR-1

● DMSR-7

FLP-11   FLP-10

FLP-11   FLP-1

● FRPR-8

● EGL-6

FLP-13   FLP-11

FLP-10   FLP-5

GPCR gene expression

AVDL/R

PQR

Pharynx

Sensory
neurons

CAN
neurons

Interneurons

Motorneurons

Sending neurons

Neuropeptide precursor gene expression

Pharynx

Sensory
neurons

Interneurons

Motorneurons

Receiving neurons

short range 1 2 4 6 8 10 12 14 16 18

mid range 1 2 4 6 8 10 12 14 16 18

Edge weight (number of linking NPP-GPCR pairs)

A Model 3 - short-range

Number of connections / Individual neurons

degree, indegree, outdegree

hubs

Top 10 hubs
AVKL/R
PVT
PVQL/R
PVR
DVA
PVCL/R
AVAL/R

B Model 2 - mid-range

Number of connections / Individual neurons

hubs

Top 10 hubs
AVKL/R
PVT
PVQL/R
PVR
DVA
BDUL/R
RID

C Synaptic network

Number of connections / Individual neurons

hubs

Top 10 hubs
AVAR/L
AVBL/R
PVCL/R
AVDR
DVA
AVEL/R

D Monoamine network

Number of connections / Individual neurons

hubs

Top 10 hubs
CEPDL/R
CEPVL/R
ADEL/R
PDEL/R
RIML/R

E node, edge, hub, Rich club, degree (k) sum of connections

F Φ *C.elegans* / Φ Normalised / Neuropeptide degree (k)

G Synaptic degree (k) vs Neuropeptide degree (k); synaptic hubs, neuropeptide hubs

Driving neurons	Group 1	Group 2	Group 3
	DA09	PVQL/R	RIR
	VA12	PVNL/R	I5
	VC06	AVAL/R	I4
	DA08	PVR	RICL/R

A

Expression of NPP/GPCR gene ● Co-expression of NPP-GPCR gene pairs

NPP expression

GPCR expression

Pharynx | Sensory neurons | Interneurons | Motorneurons

B

Autocrine connections by Neuron Type

Neurons (%)

All neurons | Pharyngeal neurons | Sensory neurons | Inter-neurons | Motor-neurons

Neurons (%)

All neurons
Pharyngeal neurons
Sensory neurons
Interneurons
Motorneurons

PQR
PVR
URX

NPP-GPCR gene pairs co-expressed

C

neurons with co-expression

FLP-type
NLP-type
○ Promiscuous GPCR

DMSR-1

DMSR-7

NPP - GPCR gene pairs

D

Correlation with neuropeptide network

Degree

● Total degree
▶ In-degree
◀ Out-degree

AVK
PVQ
PVT
DVA

PVR
PQR
URX

autocrine connections

Correlation with synaptic network

Degree

● Total degree
▶ In-degree
◀ Out-degree

AVAR
AVAL

AVB
PVC
DVA
AVD

PVR
URX
PQR

autocrine connections

E

URX
PVR
PQR

D
A — P
V

different autocrine peptide connections
● 0
○ 1
○ 7
○ 13

Oxygen-sensory neurons

BAG
URX
PQR
AQR
+ others

autocrine peptide

responses to oxygen

dorsal muscle

DD
DA
AS
DB
VA
VB
VD

ventral muscle

forward
backward

→ excitatory synapse
⊣ inhibitory synapse
⟲ autocrine peptide

A

B

A Model 1 - long-range
GPCR gene

B Model 2 - mid-range
GPCR gene

Number of different types of connection

2 4 6 8 10 12 14 16 18 20

A

CeNGEN threshold 2 - EC$_{50}$ 1 μM · CeNGEN threshold 2 - EC$_{50}$ 500 nM · CeNGEN threshold 4 - EC$_{50}$ 1 μM · CeNGEN threshold 4 - EC$_{50}$ 500 nM

Co-expression of NPP-GPCR gene pairs

Pharynx | Sensory neurons | Interneurons | Motorneurons

B

Fraction of neurons

CeNGEN threshold 2 - EC$_{50}$ 1 μM | CeNGEN threshold 2 - EC$_{50}$ 500 nM | CeNGEN threshold 4 - EC$_{50}$ 1 μM | CeNGEN threshold 4 - EC$_{50}$ 500 nM

Number of NPP-GPCR gene pairs co-expressed

— All neurons
—·— Pharyngeal neurons
— Sensory neurons
– – Interneurons
······· Motorneurons

C

\# neurons with co-expression

FLP-type   NLP-type

**

D

\# neurons with co-expression

Specific   Promiscuous

GPCR showing autocrine connections (%)

Specific   Promiscuous

E

Correlation with gap junction network

— Total degree

Degree

\# autocrine connections

AVAL
AVAR
AVBR
AVBL
PVR
URX PQR

Correlation with monoaminergic network

— Total degree
▶ In-degree
◀ Out-degree

Degree

\# autocrine connections

CEP
ADE PDE
URX PVR
PQR

F

DA

VA

DB

VB

DMSR-1
DMSR-7
NPR-5