LSR Targets YAP to Modulate Intestinal Paneth Cell Differentiation

Yanan An[#1], Chao Wang[#1], Baozhen Fan[#2], Ying Li[1], Feng Kong[3,4,5], Chengjun Zhou[6], Zhang Cao[7], Jieying Liu[1], Mingxia Wang[1], Hui Sun[1], Shengtian Zhao[*2,3,4], Yongfeng Gong[*1]

Author affiliation:

Department of Physiology, Binzhou Medical University, Yantai, Shandong, CHINA

Department of Urology, Binzhou Medical University Hospital, Binzhou, Shandong, CHINA

Shandong Provincial Engineering Laboratory of Urologic Tissue Reconstruction, Jinan, Shandong, CHINA

Department of Urology, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong, CHINA

Department of Central Laboratory, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong, CHINA

Department of Pathology, The Second Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong, CHINA

Department of Pathology, Binzhou Medical University Hospital, Binzhou, Shandong, CHINA

[#] These three authors contributed equally to this work

* Corresponding authors:

Yongfeng Gong, MD, PhD, Professor

Department of Physiology

Binzhou Medical University

346 Guanhai Avenue

Yantai, Shandong, CHINA 264003

Phone: +86-535-6913163, +86-18705439006

E-mail: ygong@bzmc.edu.cn

Shengtian Zhao, MD, PhD, Professor

Department of Urology

Binzhou Medical University Hospital

661 Huanghe 2nd Road

Binzhou, Shandong, CHINA 256603

Phone: +86-543-3256802, +86-15153169567

E-mail: zhaoshengtian@sdu.edu.cn

Abstract

Lipolysis-stimulated lipoprotein receptor (LSR) is a multi-functional protein that is best known for its roles in assembly of epithelial tricellular tight junctions and hepatic clearance of lipoproteins. Here, we investigate the function of LSR in intestine biology. By using multiple conditional deletion mouse models and *ex vivo* cultured organoids, we find that LSR elimination in intestinal stem cells results in disappearance of Paneth cell without affecting the differentiation of other cell lineages. Mechanistic studies reveal that LSR deficiency increase abundance and nuclear localization of YAP by modulating its phosphorylation and proteasomal degradation. Intestinal LSR-deficient mice are susceptible to development of necrotizing enterocolitis. In addition, LSR can sense and interpret fatty acid signals derived from dietary lipids and transduce into inactivation of YAP. Thus, this study identifies LSR as an upstream negative regulator of YAP activity and part of the mechanism mediating connection between fat diet and YAP signaling in intestine.

Introduction

Lipolysis-stimulated lipoprotein receptor (LSR), a multi-functional type I transmembrane protein containing an immunoglobulin-like domain (Konno et al., 2020; Masuda et al., 2011), has been linked to a variety of biological processes, molecular functions and cellular compartments. In the liver, LSR recognizes apolipoprotein B/E-containing lipoproteins in the presence of free fatty acids, and is thought to be involved in the hepatic clearance of triglyceride-rich lipoproteins (Narvekar et al., 2009; Yen et al., 1994; Yen et al., 1999). LSR also plays an essential role in organization of tricellular tight junctions that are involved in epithelial barrier function (Masuda et al., 2011; Sugawara et al., 2021). In the brain, LSR is specifically expressed at tricellular tight junctions between central nervous system endothelial cells and plays critical roles for proper blood–brain barrier formation and function (Sohet et al., 2015). LSR is also a target molecule for cell binding and internalization of Clostridium difficile transferase (Hemmasi et al., 2015; Papatheodorou et al., 2011). Although its role as either a tumor promoter or suppressor (or

both) is not established, expression and localization of LSR have been found to be altered in several cancers (Dong et al., 2022; Shimada et al., 2017; Takahashi et al., 2021). The dominant subcellular localization of LSR is on the membrane, however, its localization in the nucleus of human epithelial cells has been reported (Reaves et al., 2017). Taken together, these findings suggest that the subcellular localization, function, and signaling pathways regulated by LSR are tissue- and cell type-specific. However, the specific cell types that express LSR have been difficult to identify, and the functions of LSR in postnatal development and tissue homeostasis have been hampered by the perinatal lethality of *Lsr* null mice (Mesli et al., 2004; Sohet et al., 2015).

Previous studies have revealed that LSR is highly expressed in the intestinal epithelium and localized at the basolateral membrane in addition to tricellular tight junctions of mouse intestine (Sugawara et al., 2021). *In vitro* studies in cultured EpH4 mouse mammary epithelial cells and Madin-Darby Canine Kidney (MDCK) cells indicates that LSR plays essential roles in the plasma membrane seal at tricellular tight junctions and in epithelial barrier function (Masuda et al., 2011; Sugawara et al., 2021). However, roles of LSR in intestinal homeostasis *in vivo* remain to be fully elucidated. Earlier work in Drosophila has established a pivotal link between tricellular tight junction proteins, stem cell behaviour, and intestinal homeostasis (Resnik-Docampo et al., 2017), to what extent this connection is conserved in mammalian systems remains uncertain. Meanwhile, embryonic lethality in LSR-deficient mice highlights the importance of LSR for development (Mesli et al., 2004), but the significance and relevance of this protein in regulating intestinal development and differentiation in mammals are largely unclear. It is also unclear whether LSR can transduce dietary signals and organismal nutritional status into regulation of intestinal homeostasis, given that it is located near the apical part of the epithelium and in contact with the intestinal lumen.

Here, we used a combination of *in vivo* conditional deletion mouse models and *ex vivo* cultured organoids to investigate the role of LSR in the regulation of intestinal homeostasis. We show that absence of LSR in intestinal stem cells (ISCs) affects the development of Paneth cells in small intestine by regulating protein abundance and localization of YAP. An *ex vivo* organoid-forming assay showed that intestinal crypts from *Lsr* knockout mice or induced pluripotent stem cells (iPSCs) with stable *LSR* knockdown formed symmetrical spherical organoids without Paneth cell lineage. Mechanistically, LSR controls YAP

phosphorylation and proteasomal degradation by modulating the binding of YAP to 14-3-3 zeta and

phosphatase protein phosphatase 2A catalytic subunit (PP2Ac). Moreover, the disease relevance of these

findings was indicated by the striking susceptibility of epithelial-specific LSR-deficient mouse to the

development of experimental necrotizing enterocolitis (NEC), a severe and often fatal disease that affects

premature infants, and in which intestinal epithelial LSR was shown to be markedly decreased in mice

and humans. Further, although the relationship between high fat diet (HFD) exposure and YAP activity

has been established (Poon et al., 2013; Wang et al., 2020b; Yuan et al., 2017), the molecular basis

associated with how fat diet affect YAP activity has not been thoroughly examined. This study discovered

that fat diet component, such as palmitic acid (PA), potently inhibit YAP activity through LSR, which

represents a novel regulatory pathway for YAP signaling with important implications for human

responses to dietary constituents.

Results

LSR ablation results in increased numbers of proliferating cells and loss of Paneth cell lineage in

the small intestine

LSR is widely expressed in small intestine and colon which suggested a role for LSR in regulating

homeostasis in the gastrointestinal tract. To bypass the embryonic lethality of constitutive deletion and

investigate the potential role of *Lsr* in intestinal homeostasis, we generated *Lsr$^{loxP/loxP}$* mice, in which the

mutant Lsr allele contains exons 1~2 flanked by loxP sites, on a C57BL/6J background (Fig. S1A). We

used *Lsr$^{loxP/loxP}$;CAG-CreER* mice to achieve global *Lsr* deletion (Lsr$^{-/-}$) upon tamoxifen treatment

initiated at 7–8 weeks of age. We also generated intestinal-epithelium-specific *Lsr*-deficient mice (Lsr^{vill}

^{KO}) by intercrossing *villin-Cre* and *Lsr$^{loxP/loxP}$* mice (Fig. S1B). qRT-PCR and western blot analysis of

LSR expression showed that the transcription and expression of LSR in the intestines of Lsr$^{vill\ KO}$ and

Lsr$^{-/-}$ mice were successfully blocked (Fig. 1, A-D). LSR has been previously reported to be localized at

both the basolateral membrane and tricellular tight junctions (Sugawara et al., 2021), we found that LSR

was expressed throughout the cellular membrane of epithelium in confocal sections, and did not show

remarkable concentration at tricellular tight junctions in the Lsr^{ctrl} mouse intestine (Fig. 1E). This was

further ascertained by the disappearance of LSR signal in intestinal epithelium of Lsr$^{vill\ KO}$ and Lsr$^{-/-}$ mice

(Fig. 1, E and F). In addition, transmission electron microscopy (TEM) showed that the structure of tight

junctions between the intestine epithelial cells (IECs) in Lsr$^{\text{vill KO}}$ mice was not significantly different

from that in Lsr$^{\text{ctrl}}$ mice (Fig. S1C). Further immunostaining of ZO-1 and CLDN1, as an indicator of

bicellular tight junctions' integrity, showed no difference between Lsr$^{\text{vill KO}}$ and Lsr$^{\text{ctrl}}$ small intestine (Fig.

S1, D and E). The tricellular tight junctions' integrity, assessed by tricellulin localization which was

detected as dots at tricellular tight junctions, remained intact in Lsr$^{\text{vill KO}}$ small intestine (Fig. S1F).

Together, these results indicate that the gross barrier function of tight junctions is not affected by the

deletion of LSR in IECs.

LSR deficiency had no effect on the overall structures of small intestine or colon (Fig. 1, G and H).

Surprisingly, the small intestines of Lsr$^{\text{vill KO}}$ mice were significantly thicker than those of Lsr$^{\text{ctrl}}$ mice,

and this phenomenon was consistent at various times during the growth of mice (Fig. 1, G, I and J).

Moreover, we observed significantly longer villi in H&E-stained sections of small intestine of Lsr$^{\text{vill KO}}$

and Lsr$^{-/-}$ mice (Fig. 1, K and L). We speculated that there could be a considerable increase in proliferative

cell numbers in the *Lsr* mutant mice. Immunofluorescence was used to analyze the proliferation-related

proteins Ki67 and proliferating cell nuclear antigen (PCNA) in the small intestine, and the expression of

Ki67 and PCNA in the duodenum, jejunum and ileum was found to be significantly increased in Lsr$^{\text{vill}}$

$^{\text{KO}}$ mice (Fig. 2, A and B). Western blot analysis of Ki67 and PCNA confirmed this result in Lsr$^{\text{vill KO}}$ and

Lsr$^{-/-}$ mice (Fig. 2, C and D). To address if ISCs are responsible for the proliferation in small intestine,

we performed qRT-PCR and immunofluorescence analysis for olfactomedin 4 (OLFM4), a marker

expressed by the Lgr5$^+$ ISCs. Absence of LSR did not change OLFM4 expression of the small intestine

(Fig. 2, E and F and Fig. S1G), indicating that the highly proliferating cells in Lsr$^{\text{vill KO}}$ and Lsr$^{-/-}$ mice

were not ISCs. Since an increased percentage of Ki67 positive cells was predominantly found in the

transit-amplifying zone (Fig. 2A), we speculated that the proliferating cells may be undifferentiated

progenitor cells. In summary, these data demonstrate that in the absence of LSR, ISCs do not contribute

to the population of proliferating cells, presumably representing the transit-amplifying population leading

to increased villus length.

In agreement with histological observations, which indicated the absence of granule-containing cells in

H&E-stained sections of small intestine of Lsr$^{\text{vill KO}}$ and Lsr$^{-/-}$ mice (Fig. 1, K and L), RNA sequencing

(RNA-seq) analysis of small intestinal lysate from in Lsr^vill KO and Lsr^ctrl showed that *Lsr*-knockout

downregulated many genes comprising the Paneth cell signature (*Lyz1, Itln1, Mptx2, Defa5,* and *Cd24*)

(Fig. 2, G and H) and DEFA family members (Fig. S1H). Data were validated by immunofluorescence

(Fig. 2, I and K) and qRT-PCR (Fig. 2J and Fig. S1I) performed on the small intestine from Lsr^vill KO,

Lsr^-/- and control littermates. These results demonstrated that LSR elimination in intestinal epithelium

leads to depletion of Paneth cells. Despite the absence of Paneth cells, expression levels of markers for

other types of cells (EpCAM for IECs, MUC2 and SPDEF for goblet cells, CHGA and NEUROG3 for

enteroendocrine cells, CDX1 and CDX2 for columnar absorptive cells, and LGR5 for stem cells) and

Periodic acid–Schiff (PAS) staining of the goblet cells were not significantly changed (Fig. 2, L and Fig.

S1, J-M). These data collectively suggest that LSR is absolutely required to maintain the Paneth cell

lineage.

LSR is required for Lgr5+ ISCs to Paneth cell differentiation

Paneth cells, like all other intestinal epithelial cell types, can be derived from Lgr5+ ISCs, we conjectured

that LSR might affect the function of Lgr5+ ISCs to differentiate into Paneth cells. Interestingly, we found

that LSR was highly expressed in Lgr5+ ISCs at the bottom of intestinal crypts (Fig. 3A). These results

suggested that LSR may affect the function of Lgr5+ ISCs, thereby making them lose the ability to

differentiate into Paneth cells. We used a *Lgr5-CreERT2* strain to selectively delete *Lsr* in Lgr5+ ISCs in

a tamoxifen-inducible manner (hereafter referred to as Lsr^ISC KO) and to specifically explore the cellular

and molecular consequences of *Lsr* deletion in Lgr5+ ISCs (Fig. 3, B and C). Although the lifespan of

Paneth cells is estimated to be 30 days (Bjerknes and Cheng, 1981), qRT-PCR and immunofluorescence

staining revealed reduced expression of Paneth cell markers (Lyz1, Itln1, Mptx2, Defa5, and Cd24a) in

the small intestine of Lsr^ISC KO mice 15 days after the last dosage of tamoxifen injection, again supporting

the idea that LSR is required for Lgr5+ ISCs to Paneth cell differentiation (Fig. 3, D and E). In line with

the findings in Lsr^vill KO mice, Lsr^ISC KO intestine possessed other intestinal cell types, such as IECs, ISCs,

enteroendocrine cells and goblet cells (Fig. S2, A and B). In Lsr^ISC KO mice, intestinal permeability assay

showed that the concentration of FITC-Dextran 4000 in plasma seemed to increase compared to that of

the Lsr^ctrl mice, but it did not attain a significant difference (Fig. S2C), indicating the phenotype caused

by LSR deficiency is not attributable to the dysfunction of permeability barrier. In contrast to Lsr^vill KO

mice, we found no difference in the proliferative cells in small intestine of Lsr[ISC KO] mice, indicating a

potential direct role of LSR on transit-amplifying population. These results further demonstrated that in

the absence of LSR in Lgr5[+] ISCs, Paneth cells were not formed, but the differentiation of other IEC

types was unaffected.

We also took advantage of crypt organoid culture to confirm the findings. Immunofluorescence analysis

showed that LSR was expressed in organoid culture of Lsr[ctrl] crypts and localized to the cell membrane,

while no LSR expression was observed in Lsr[vill KO] crypt organoids (Fig. 3F). As expected, most crypts

in Lsr[ctrl] mice differentiated and budded into organoids with villi over time (Fig. 3G). However, when

intestinal crypts from Lsr[vill KO] mice were cultured, *Lsr* deletion resulted in the formation of symmetrical

spherical organoids (Fig. 3, F and G). In addition, on day 6, the shape of the spherical organoids did not

change significantly; and the number of organoids gradually decreased (Fig. 3, G-I).

Immunofluorescence staining showed that the distribution and quantity of stem cells (OFLM4) and

epithelial cells (EpCAM) in Lsr[vill KO] organoids were similar to those in Lsr[ctrl] organoids (Fig. 3, J and

K). Interestingly, similar to the above *in vivo* results, the Paneth cell markers (LYZ1 and MPTX2)

exhibited significantly reduced and even complete loss of expression in Lsr[vill KO] organoids (Fig. 3, L and

M). Therefore, our results indicated that the Paneth cell number was positively correlated with the LSR

expression level in both the crypt organoids and the intestine. Although qRT-PCR showed that the levels

of the Paneth cell markers (*Mptx2, Itln1* and *Defa5*) were significantly reduced in Lsr[vill KO] organoids

(Fig. S2D), the numbers of other types of cells did not differ significantly between the Lsr[vill KO] and Lsr[ctrl]

organoids (Fig. 3, N-P and Fig. S2E). Therefore, the organoid culture of *Lsr* deficient crypts truthfully

recapitulated the phenotype of Lsr[vill KO] mice intestine. Next, we knocked down *LSR* in human iPSCs

(Fig. 3Q) and then cultured organoids. Unsurprisingly, we obtained results similar to those described

above (Fig. 3, R-V), indicating a conserved function of LSR in the intestine from mouse to human.

Collectively, these findings support a critical role for LSR in regulating ISCs differentiation to Paneth

cells.

Loss of LSR results in upregulation and activation of YAP

In intestine, excessive YAP can inhibit the differentiation of ISCs into Paneth cells (Gregorieff et al.,

2015), and promote the proliferation of undifferentiated progenitor cells (Camargo et al., 2007). Recent

work by Serra *et al.* revealed that homogeneous activation or suppression of YAP abolished Paneth cell

differentiation and organoid budding, leading to the development of symmetrical spheres that were either

develop into enterocysts or remain as undifferentiated (Serra et al., 2019). The loss of Paneth cells,

proliferation of undifferentiated progenitor cells, and formation of symmetrical spherical organoids

resulting from elimination of *Lsr* in the intestine led us to test whether abnormal YAP activation

contributes to these phenotypes. Confocal imaging showed a distinct colocalization of YAP with LSR in

mouse intestine and Lgr5[+] ISCs (Fig. 4A), indicating that LSR might have a regulatory effect on YAP.

Additionally, in Lsr[-/-], Lsr[vill KO] and Lsr[ISC KO] mouse models, we observed enhanced YAP protein

abundance accompanied by increased mRNA expression of the well-established YAP target genes, such

as *Edn1, Ctgf,* and *Cyr61 (Fan et al., 2022)*, in the small intestine (Fig. 4, B-G), but the mRNA expression

level of *Yap* did not change significantly (Fig. 4, E-G). To test directly whether elevated YAP contributes

to the phenotype caused by LSR deletion, we lowered YAP signaling in Lsr[vill KO] mice with verteporfin

(VP), a pharmacologic inhibitor of YAP-TEAD association, twice per day from postnatal day 7 to day

21. And then immunofluorescence staining of MPTX2 was performed to evaluate the content of Paneth

cells in small intestine. The expression of MPTX2 was found to be significantly increased in small

intestine of VP-treated Lsr[vill KO] mice (Fig. 4H). In addition, qRT-PCR showed that the mRNA levels of

Paneth cell markers (*Lyz1, Itln1, Mptx2* and *Defa5*) were significantly increased in VP-treated Lsr[vill KO]

mice (Fig. S2F). These results indicated that VP treatment restored Paneth cell number. H&E staining

showed that the small intestinal villi hyperproliferation were also partially suppressed in VP-treated Lsr[vill]

[KO] mice (Fig. 4I). The expression of several YAP target genes *Edn1, Ctgf* and *Cyr61* was significantly

suppressed by VP treatment (Fig. 4J), indicating decreased YAP signaling in intestine of VP-treated

Lsr[vill KO] mice. Similar results were observed in Lsr[vill KO] mice treated with lentiviruses carrying the *Yap*

shRNA (Fig. 4, K-M and Fig. S2G). These observations strongly support the causal link between the

LSR-dependent YAP regulation and the altered Paneth cell differentiation in *Lsr*-null mice.

Next, we sought to verify the importance of YAP to the phenotype of *Lsr*-deficient intestine. We

examined the expression of YAP in ISCs *in vitro*, and found that *Lsr* deletion led to significant increases

in YAP expression and nuclear translocation in the ISCs isolated from Lsr[ctrl] and Lsr[ISC KO] mouse intestine

(Fig. 4N). Then the enhanced YAP expression and nuclear translocation were also observed in ISCs

isolated from the Lsr^ctrl and Lsr^vill KO mouse intestine (Fig. S2H). Next, we verified the above results in

Lsr^vill KO intestinal organoids by treating with YAP knockdown lentivirus according to a previously

described method (Goga et al., 2021). Briefly, 12 hours after budding was observed in the Lsr^ctrl

organoids, a timepoint YAP inactivation was observed (Gjorevski et al., 2022), Lsr^vill KO organoids were

randomly divided into YAP knockdown virus treatment group and control virus treatment group. 5 hours

after virus transfection, organoids were seeded in Matrigel and cultured for 3 days before harvesting for

RNA isolation and immunofluorescence staining. The qRT-PCR showed that *Yap* shRNA lentiviral

transduction reduced the levels of *Yap* down to 30% in organoids (Fig. S2I), and rescued de novo crypt

formation (Fig. S2J) and normalized Paneth cell differentiation (Fig. S2K). Taken together, these results

confirmed a role for YAP activation in regulation of ISCs function downstream of LSR.

Wnt/β-catenin signaling pathway was discovered to be required for intestinal homeostasis and Paneth

cell differentiation (Totaro et al., 2018). However, β-catenin staining of crypts in the Lsr^vill KO and Lsr^ISC

^KO mice was similar to that of control crypts (Fig. S2, L and M). Activation of YAP has been reported to

directly inhibit Wnt signaling in the intestine (Cheung et al., 2020; Li et al., 2020), but LSR knockout

did not affect expression of most of the β-catenin-target genes including *Ccnd1, Axin2,* and *Cd44* in both

the intestinal tissue and Lgr5^+ ISCs isolated from Lsr^vill KO and Lsr^ctrl mice (Fig. S2, N and O). This data

may be due to alterations in negative feedback mediators in the Wnt pathway. Further, both ATOH1 and

SOX9, two critical transcription factors for the differentiation of intestinal Paneth cell lineage, were not

significantly altered by *Lsr* ablation in ISCs (Fig. S2, P-R).

14-3-3 zeta and PP2Ac are involved in the regulation of YAP by LSR

The above data clearly indicate a role of LSR on YAP expression and activation, we examined in closer

detail whether Hippo pathway activity, which negatively regulates YAP, is perturbed by LSR ablation.

Immunoblot analysis indicated that LSR ablation actually led to up-regulation of phosphorylated large

tumor suppressor (p-Lats), a component of the mammalian Hippo pathway, which inhibits YAP nuclear

translocation and promotes its proteasomal degradation (Fig. S3, A and B). We conjectured that the

elevation of p-Lats1 is induced by YAP activation via a potential negative feedback mechanism.

Moreover, the increase of YAP protein level was not due to increased transcription, since *Yap* mRNA

was slightly decreased in the small intestine (Fig. 4, F and G). Since Hippo pathway cannot account for

YAP up-regulation, we speculated that *Lsr* deletion may affect the metabolism of YAP and cause its

protein level to increase. Therefore, a protein synthesis inhibitor (cycloheximide, CHX) and a proteasome

inhibitor (MG132) were used to determine the effect of LSR on YAP metabolism. The degradation rate

of YAP in the Lsr[vill KO] group was significantly lower than that in the Lsr[ctrl] group after CHX treatment

(Fig. 5A). Moreover, MG132 blocked the turnover of YAP in the presence of CHX, and deletion of *Lsr*

had an effect similar to that of MG132 on blocking YAP turnover (Fig. 5B). Thus, YAP is degraded via

the proteasome pathway, and deletion of LSR might increase YAP accumulation by suppressing its

proteasomal degradation. Phosphorylation of YAP at Ser381 can increase its ubiquitination, ultimately

leading to proteasomal degradation (Zhao et al., 2010). Decreased phosphorylation of p-YAP at Ser381

(Fig. 5C), as well as decreased YAP ubiquitination (Fig. 5D) was observed in the ISCs isolated from

Lsr[vill KO] mice. These results demonstrated that *Lsr* knockout enhanced YAP stability by decreasing its

phosphorylation at Ser381, thereby blocking its ubiquitination and degradation.

We next sought to determine the molecular mechanism through which LSR regulate the phosphorylation

of YAP. We performed a mass spectrometry analysis of LSR immunocomplexes from ISCs. Results

from three experiments identified a list of 289 cellular proteins that specifically interact with LSR. We

focused on two 14-3-3 proteins, 14-3-3 zeta and 14-3-3 gamma, and two protein phosphatases, protein

phosphatase 1 catalytic subunit alpha (PPP1CA) and PP2Ac (Fig. 5E), because they have been reported

to participate in the regulation of YAP in multiple tissues (Hu et al., 2017; Meng et al., 2016; Qi et al.,

2015; Schlegelmilch et al., 2011; Sun et al., 2021; Yan et al., 2021). It is known that the turnover of YAP

is regulated through its phosphorylation and ubiquitination in a 14-3-3 dependent manner. Remarkably,

coimmunoprecipitation (co-IP) and immunoblot analysis of *in vivo* and *in vitro* samples confirmed that

14-3-3 zeta interacted with both YAP and LSR (Fig. 5F and Fig. S3C), but 14-3-3 gamma, PPP1CA, or

PP2Ac did not interact with YAP or LSR either *in vivo* or *in vitro* (Fig. S3, D-H and Fig. 5G). PPP1CA

and PP2Ac have been reported to be phosphatases that directly and effectively dephosphorylate YAP

(Ser381). An antagonistic and competitive interaction has been reported between PP2Ac and 14-3-3 for

the phosphorylation of YAP (Schlegelmilch et al., 2011). Hence, we tested whether deletion of LSR

results in an increased association of YAP with PP2Ac or PPP1CA. YAP interacted with PP2Ac instead

of PPP1CA and exhibited reduced binding to 14-3-3 zeta in the ISCs from Lsr[vill KO] mouse intestine (Fig.

5H and Fig. S3I) and HEK293 cells transfected with YAP, 14-3-3 zeta and PP2Ac expression plasmids (Fig. S3J), while LSR and YAP did not seem to bind to PP2Ac in the ISCs from Lsr^ctrl mouse intestine or HEK293 cells (Fig. 5G and Fig. S3H). Therefore, YAP could interact with 14-3-3 zeta and PP2Ac respectively, and deletion of LSR might increase the interaction between YAP and PP2Ac by decreasing the YAP/14-3-3 zeta association. To determine whether PP2Ac can efficiently dephosphorylate YAP (Ser381), an *in vitro* phosphatase assay is performed in the presence of PP2Ac. The presence of PP2Ac resulted in diminished YAP (Ser381) phosphorylation and YAP accumulation (Fig. 5I). These results suggested that loss of *Lsr* leads to more efficient association of YAP with PP2Ac, which can potentially dephosphorylate YAP at Ser381 and thereby decrease its degradation.

Loss of *Lsr* exacerbates inflammation in a mouse model of NEC

Paneth cells, which are enriched in the ileum, have a central role in preventing intestinal inflammation (Adolph et al., 2013) and defects in Paneth cells are a hallmark of NEC. Infants who have developed NEC have decreased Paneth cell numbers compared to age-matched controls (Coutinho et al., 1998), and ablation of murine Paneth cells results in a NEC-like phenotype (Lueschow and McElroy, 2020; Lueschow et al., 2018). We speculated that Paneth cell loss caused by LSR deletion may render the immature small intestine susceptible to development of NEC. Hence, to evaluate the role of LSR in NEC, we used 2,4,6-trinitrobenzenesulfonic acid (TNBS) to establish NEC models. Enteral administration of TNBS in 14-day-old mouse pups induced NEC, as revealed by villus disruption, clear separation of lamina propria and edema in the submucosa (Fig. S4A and Fig. 6A), increased mRNA levels of proinflammatory factors, including *IL-1α, IL-2, IL-6, IFN-γ* and *TNF-α* (Fig. 6B), and increased infiltration of inflammatory cells (macrophages and neutrophils) (Fig. S4, B and C and Fig. 6, C and D) in small intestine of mice treated with TNBS in Lsr^ctrl mice. As expected, LSR expression was significantly decreased in the small intestines of mice treated with TNBS (Fig. 6, E and F), and the number of Paneth cells was also significantly reduced (Fig. S4, D and Fig. 6, G and H). Compared with Lsr^ctrl mice, Lsr^vill KO mice showed worsening intestinal injury, represented by more severe transmural injury in the small intestine (Fig. S4E and Fig. 6I), increased mRNA levels of *IL-1α, IL-2, IL-6, IFN-γ, and TNF-α* (Fig. 6J), and increased infiltration of macrophages and neutrophils (Fig. S4, F and G and Fig. 6, K and L) after NEC induction. Collectively, these results implied that the lack of LSR led to more

severe NEC. To further explore the effect of LSR on NEC, we used an AAV10 system with intraperitoneal injection, which has been proven to primarily transduce cells in the intestine, then established the NEC model in mice transduced with AAV encoding mouse *Lsr* (AAV-*Lsr*) or control vector (AAV-CTL). The LSR overexpression AAV decreased the severity of TNBS induced NEC as indicated by histopathological analysis (Fig. S4H and Fig. 6M), suppressed cytokines production (Fig. 6N), decreased inflammatory cells infiltration (Fig. S4, I and J and Fig. 6, O and P), and preserved Paneth cell number (Fig. 6, Q and R) in the small intestine of AAV-*Lsr* treated mice. These results suggest that LSR protects against TNBS induced NEC through upregulation of Paneth cell. To further verify the role of LSR in NEC, we tested intestinal samples from NEC patients. Compared with healthy control, patients with NEC exhibited severe intestinal injury (Fig. 6S) and significantly reduced expression of LSR (Fig. 6T), especially in ISCs (Fig. 6U). Similar to previous reports (Underwood, 2012), infants with NEC have significantly decreased numbers of Paneth cells compared to age-matched controls (Fig. 6V). These results confirmed that LSR plays an important role in the development of NEC and is expected to be a potential therapeutic target for NEC.

LSR mediates the inhibitory effect of PA on YAP

LSR has been described as a lipoprotein receptor activated by free fatty acid, with oleate and palmitate were the most efficient (Mann et al., 1995). Next, we aim to identify if fatty acid could be a potential extracellular signal which are able to modulate the activity of LSR in the intestine. HFD or PA, a main component of the HFD, did not alter the expression of LSR in small intestine (Fig. 7, A-D). However, we observed less abundant YAP and reduced expression of a subset of YAP target genes such as *Edn1, Ctgf,* and *Cyr61* at the mRNA levels in small intestine from HFD and PA-treated mice compared to controls (Fig. 7, E-H). Then we isolated the Lgr5$^+$ ISCs from mouse small intestine and incubated that in serum-free medium supplemented with PA for 24 hours. Results demonstrated that PA reduced expression of YAP (Fig. 7K), partially translocated YAP from the nucleus into cytoplasm (Fig. 7I), and suppressed the expression of many of its target gene (Fig. 7J), indicating YAP inhibition. These data suggest that PA can suppress YAP activity. Next, we examined whether the inhibitory effects of PA on YAP are indeed mediated by LSR. In contrast to wild-type intestine, loss of LSR abrogated the HFD or PA-driven downregulation of YAP and its target genes, *Edn1, Ctgf,* and *Cyr61* (Fig. 7, E-H). Then we

isolated the ISCs from the small intestine of Lsr[ctrl] and Lsr[vill KO] mice and treated the cells with PA. LSR

deficiency attenuated the regulatory effect of PA on YAP protein abundance, as well as the reduction of

YAP target gene expression (Fig. 7, J and K). The results collectively indicate LSR may be able to sense

and interpret fatty acid signals derived from dietary lipids and transduce the inactivation of YAP.

Discussion

Although the abundant expression of LSR has been described in intestine (Mesli et al., 2004; Sugawara

et al., 2021), roles of this receptor in intestinal development, physiology and disease remain unclear. Here,

we generated multiple conditional deletion mouse models to identify roles for LSR in the intestinal

epithelium. We show that genetic ablation of *Lsr* in the intestinal epithelium results in increased numbers

of proliferating cells, as well as a reduction in Paneth cell lineage. We demonstrate that the numbers of

Lgr5[+] ISCs quantified by OLFM4 immunostaining displayed no significant difference between Lsr[vill KO]

and control samples, suggesting that the associated increase in proliferating cells in Lsr[vill KO] crypts is

likely due to increased progenitor cell numbers. Strikingly, elimination of LSR from the intestinal

epithelial compartment results in the almost complete disappearance of Paneth cell lineage, but had no

effect on the number and distribution of other types of cells in the small intestine. The disappearance of

Paneth cell was reproduced in the tamoxifen-inducible Lsr[-/-] and Lsr[ISC KO] mice. An *ex vivo* organoid-

forming assay showed that intestinal crypts from Lsr[vill KO] mice or iPSCs with LSR knockdown formed

symmetrical spherical organoids without Paneth cell lineage, implying that LSR plays a key role in the

regulation of ISC function and differentiation toward Paneth cells in the intestinal epithelium.

YAP is a regulator of intestinal regeneration and ISCs fate (Gjorevski et al., 2022). We here show that

the loss of LSR causes an increased abundance and enhanced nuclear translocation of YAP in Lsr[vill KO]

intestinal epithelium and in Lsr[ISC KO] ISCs. As for now, the molecular events and functional consequences

that follow YAP activation remain controversial (Barry et al., 2013; Cai et al., 2010; Cheung et al., 2020;

Gregorieff et al., 2015; Li et al., 2020; Zhou et al., 2011). Previous work has shown that overexpression

of YAP results in the enhanced proliferation of the stem cell compartment and loss of differentiated cell

types in the small intestine (Camargo et al., 2007). Mice deficient for Hippo components Mst1 and Mst2

exhibited an expansion of stem-like undifferentiated cells and the almost complete disappearance of all

secretory lineages, due to an increased abundance and enhanced nuclear translocation of YAP caused by

the loss of Mst1/Mst2 (Zhou et al., 2011). Recent studies by several group have demonstrated that

removal of Lats1/2, which caused the increase of YAP/TAZ expression and nuclear accumulation,

resulted in expansion of the transit-amplifying population (Guillermin et al., 2021; Li et al., 2020). In

agreement with these findings, the small intestine in Lsr[vill KO] mice was significantly thickened

accompanied by increased number of transit-amplifying cells. However, genetic ablation of *Lsr* only

resulted in loss of Paneth cell lineage, stem cells survived and produced the various cell types of intestinal

epithelium. Although Paneth cell development is tightly controlled by both Wnt and Notch pathways and

their downstream transcription factors, recent studies pointed out that spatial heterogeneities in YAP

activity are required for intestinal tissue morphogenesis, and homogeneous inhibitions or overexpression

of YAP in all cells reduces Paneth cell differentiation (Gjorevski et al., 2022; Li, 2019). We confirm this

finding and observe that homogeneous upregulation of YAP caused by LSR elimination from the

intestinal epithelial compartment results in the almost complete disappearance of Paneth cell lineage.

Loss of Paneth cells was confirmed in organoid cultures derived from *Lsr* knockout mouse crypts or

human iPSCs with stable *LSR* knockdown. Moreover, LSR deletion resulted in the formation of only

symmetrical spherical organoids from mouse crypts or human iPSCs. These phenotypes are similar to

YAP-overexpression organoids, where YAP is homogeneously active in all cells, and neither form Paneth

cells nor display symmetry breaking (Serra et al., 2019). Originally, the transcriptional regulator YAP

can stimulate single stem cells to enter a regenerative state and form a symmetric organoid (Serra et al.,

2019); then, the emergence of Paneth cells can break the symmetry of the organoid, followed by of

budding and the formation of villus structures (Chacon-Martinez et al., 2018; Sato et al., 2009). However,

if YAP is overexpressed and localized in the nucleus, the symmetry cannot be broken, and the organoid

remains spherical (Lukonin et al., 2020). Consistent with this concept, uniform induction in YAP activity,

which also abrogated spatial gradients in YAP activity, resulted in *Lsr* knockout organoids remain

symmetrical and contain no Paneth cells. Bearing in mind the pattern of YAP activity described above,

we reasoned that uniform inhibition of YAP by treatment with *Yap* shRNA lentivirus could not restore

the ability of *Lsr* knockout ISCs to differentiate into Paneth cells. Following YAP targeted knockdown,

however, we observed rescued de novo crypt formation and enhanced Paneth cell number in *Lsr* mutant

organoid. We do not currently know why such a difference exists. One possibility is that although

lentiviral transduction is a highly efficient method to introduce targeted gene in organoids, the number of integrations per cell can be variable in organoids, resulting in heterogeneous knockdown efficiency of YAP within the cell population.

Phosphorylated YAP is an inactive protein localized in the cytoplasm. When stimulated, YAP/TAZ are dephosphorylated and translocate into the nucleus to regulate transcriptional activity (Oh and Irvine, 2009). Our study showed that LSR can affect the function of YAP by affecting its phosphorylation. A previous study confirmed that the Ser381 of YAP is phosphorylated and further primes YAP for phosphorylation by CK1d/e, after which it is ubiquitinated by $SCF^{\beta-TRCP}$ (an E3 ubiquitin ligase) and degraded in the cytoplasm (Azzolin et al., 2014; Lapi et al., 2008; Maehama et al., 2021). Here, we confirmed that LSR affects Ser381 phosphorylation of YAP. When *Lsr* is deleted, diminished YAP degradation accounts for the increase in YAP abundance, in part or in whole.

14-3-3 proteins are a family of linker proteins widely distributed in eukaryotic cells that bind to hundreds of phosphorylated proteins, playing a pivotal role in controlling signaling pathways and regulating the subcellular localization of interacting proteins (Stevers et al., 2018; Xi et al., 2013). In intestine, 14-3-3 family proteins are likely an important part of machinery that regulates IEC differentiation, survival and apoptosis during physiological and pathological conditions (Gomez-Suarez et al., 2016). Protein phosphatase 2A (PP2A) is a major serine/threonine phosphatase ubiquitously expressed in eukaryotic cells and involved in a plethora of cellular processes, including apoptosis, autophagy, cell proliferation, and DNA repair (Dzulko et al., 2020). PP2A catalytic subunit (PP2Ac) is present in all forms of PP2A and is an essential component of PP2A complexes activity (Zolnierowicz, 2000). 14-3-3 and PP2Ac complexes compete for binding to target proteins, thereby regulating the subcellular localization of the target protein and cellular activities (Paroni et al., 2008; Xi et al., 2013). Here, we found that binding of LSR, 14-3-3zeta, and YAP resulted in the formation a stable complex, thus inhibiting the binding of PP2Ac to YAP. When LSR was deleted, PP2Ac competitively bound to YAP and dephosphorylated it, thus inhibiting its degradation and translocating it into the nucleus, where it exerted its effects on transcriptional activity.

NEC is a life-threatening intestinal disease in premature infants (Weis et al., 2021), however, the

pathophysiology of NEC remains incompletely understood (Frost et al., 2017). Although NEC's pathogenesis is multifactorial, Paneth cell depletion or dysfunction has been linked mechanistically to development of NEC (Lueschow and McElroy, 2020). Our data clearly show that Paneth cell deficient mice caused by *Lsr* depletion displayed increased susceptibility to TNBS induced NEC. We found that the expression of LSR was significantly reduced in the intestines of NEC patients, and similar results were also obtained in NEC mouse models. Systemic delivery of LSR using an adenoviral delivery system profoundly inhibited the development and progression of NEC. These results reveal that LSR contributes to intestinal injury and disease progression in NEC, which will potentially advance our understanding of NEC, describe new biomarkers, and lead to novel therapeutic strategies for this multifactorial disease. It would be interesting to investigate the contribution of LSR to other gut inflammatory disorders such as Crohn's disease apart from NEC, because misfunctioning Paneth cells accelerate the progress of these disorders (Wehkamp and Stange, 2020).

Fatty acid such as oleate acid and PA have been shown to be able to activate LSR (Bihain and Yen, 1992). In addition to the importance of fatty acid as a metabolic energy source, it also plays important roles in the regulation of stem cell proliferation and differentiation (Kang et al., 2014). PA has been shown to both accelerate and augment gut mucosal recovery and growth after massive small bowel resection (Sukhotnik et al., 2008). In support of a link between PA and stemness, is the observation that primary organoids exposed *ex vivo* to PA possessed more Lgr5$^+$ ISCs and gave rise to more secondary organoids compared to controls (Beyaz et al., 2016). On the other hand, a series of studies have established YAP as nutrient sensors that integrate nutrient signals to growth regulation (Koo and Guan, 2018; Santinon et al., 2016). HFD feeding results in a strong increase in nuclear localization of YAP/TAZ in white adipose tissue which protects mature adipocytes from cell death during obesity (Wang et al., 2020a). PA can inhibit YAP activity via inducing expression of MST1, which leads to suppression of endothelial angiogenesis (Yuan et al., 2017). Here, we identified LSR mediated the HFD and PA-driven YAP activity suppression in small intestine, indicating LSR may be able to connect nutrients to YAP activity by which it contributes to intestinal homeostasis and certain pathological conditions. Moreover, it is possible that the inhibitory effects of LSR activation on YAP activity are involved in the setting of a HFD impacts tumor initiation and progression. Thus, these findings might further expand the connections between metabolism and YAP. Although it has been proposed that the direct interaction of fatty acid with LSR

might cause a conformational shift to activate LSR (Mann et al., 1995), future research should be devoted

exploring whether fatty acid mediate its effects on YAP through direct interactions with the LSR or via

indirect mechanisms.

YAP signaling activity can be regulated by multiple factors, however, extracellular ligands, cell surface

receptors, and signaling pathways regulating YAP have not been thoroughly examined. Our results

demonstrate that LSR can directly impact the differentiation of ISCs into Paneth cells via regulating the

degradation of YAP in the ISCs (Fig.S5). This study establishes an important regulatory role of LSR in

restricting YAP activity, however, many questions remain. Are there additional direct or indirect effects

of LSR on YAP signaling activity? Are these tissue-specific or universal? Whether LSR can be used to

combat excess YAP activity? Intense studies are currently underway in our laboratory to address those

important questions.

MATERIAL AND METHODS

Cell lines, plasmids, and antibodies

Human HEK293 cells were obtained from ATCC (USA) and iPSCs were obtained from National

Collection of Authenticated Cell Cultures (China), and cultured according to the distributor's

recommendations. The expression plasmids containing mouse Lsr (pCMV6-LSR, NM_001164185,

MR225038), human LSR (pCMV6-LSR, NM_NM_205834, RC207960), mouse Yap (pCMV6-YAP,

NM_001171147, MR226049), mouse 14-3-3 zeta (pCMV6-14-3-3 zeta, NM_011740, MR203074),

mouse 14-3-3gamma (pCMV6-14-3-3 gamma, NM_018871, MR203103), mouse Ppp1ca (pCMV6-

Ppp1ca, NM_031868, MR204856), mouse PP2AC (pCMV6- Pp2ac, NM_019411, MR204384) genes,

and the knockdown plasmids containing YAP mouse shRNA plasmid (pGFP-C-shLenti, 22601) were

purchased from Origene Technologies (USA). The antibodies used in this study are summarized in

Supplemental Table 1.

Animals

All animals were housed in micro-isolator cages and received food and water freely. The laboratory

temperature was 24±1°C, and the relative humidity was 40-60%. The protocol in this study was approved

by the Committee on the Ethics of Animal Care and Use of Binzhou Medical University. The study was

conducted following the Guide for the Care and Use of Animals in Research of the People's Republic of

China. All animal studies were performed under isoflurane anesthesia, and every effort was made to

minimize suffering. C57BL/6J mice were obtained from Vital River Laboratory (China).

For the generation of Lsr$^{loxP/loxP}$ mice, targeting vector was constructed by inserting one SDA (self-

deletion anchor)-flanked neomycin cassette and two loxP sites flanking exons 1~2 of Lsr and then

electroporating into embryonic stem cells from C57BL/6J mice (conducted by Cyagen Biosciences Inc,

China). With one subsequent cross with C57BL/6J animals, the Neo transgene was removed and the

obtained Lsr$^{loxP/-}$ mice were then intercrossed in order to generate the Lsr$^{loxP/loxP}$ mice. Lsr$^{loxP/loxP}$ mice

were crossed with villin-Cre mice (Jackson Laboratory, 021504) (Madison et al., 2002), a mouse Cre line

that drives Cre recombination throughout the intestinal epithelium from embryonic day 12.5, to remove

Lsr from intestinal epithelium (Lsr$^{loxP/loxP}$/villin-Cre$^{+/-}$, *Lsr$^{vill\ KO}$*). Homozygous with villin-Cre negative

litter mates served as controls. *Lsr$^{loxP/loxP}$* mice were crossed with *CAG-CreER* mice (Jackson Laboratory,

004682) (Hayashi and McMahon, 2002) to generate systemic *Lsr* knockout mice *Lsr$^{loxP/loxP}$/CAG-Cre$^{+/-}$*

(*Lsr$^{-/-}$*), homozygous with *CAG-Cre* negative litter mates served as controls. *Lsr$^{loxP/loxP}$* mice were crossed

with *Lgr5-CreERT2* mice (GemPharmatech, T003768) to generate ISC-specific LSR knockout mice

(Lsr$^{loxP/loxP}$/Lgr5-CreERT2$^{+/-}$, *Lsr$^{ISC\ KO}$*), homozygous with *Lgr5-CreERT2* negative litter mates served as

controls. To induce the knockout of genes, tamoxifen suspended in corn oil was administered

intraperitoneally at a concentration of 10 mg/ml and dosed at 100mg/kg once every 24 hours for a total

of 5 consecutive days. 15 days post last tamoxifen injection, tissues were harvested for qRT-PCR,

Western blot, H&E or immunofluorescence staining analysis.

Patient sample

Human intestinal tissues were collected after approval by the Binzhou Medical University Review Board.

De-identified formalin-fixed paraffin-embedded tissue sections of NEC were compared with healthy

tissue margins resected for indications other than NEC (intestinal obstruction or spontaneous intestinal

perforation (n = 10). Written informed consent for research was obtained from donors before tissue

acquisition. The patients were from both sexes, between 4 and 49 days of born, and weighed between 0.9 and 1.15 kg.

Intestinal permeability analysis

Intestinal permeability was evaluated with fluorescein isothiocyanate (FITC)-Dextran 4000 (MCE, HY-128868A, China) test. Briefly, mice were fasted for 6 h and then FITC-dextran 4000 (440 mg/kg body wt) was administered by gavage. Blood was collected through tail vein 0h, 0.5h, 1h, 2h, 4h and 8h after gavage, and serum was isolated. The analysis for the fluorescence intensity of FITC-dextran 4000 in plasm was carried out with a multimode plate reader (PerkinElmer EnSpire Plate Reader, Waltham, MA, USA) (excitation, 485 nm; emission, 520 nm). Standard curves for calculating the FITC-dextran concentration were obtained by diluting FITC-dextran in PBS.

TNBS-induced NEC

2, 4, 6-Trinitrobenzene sulfonic acid (TNBS) were used to develop NEC (MohanKumar et al., 2012). Briefly, TNBS was injected into 14-day-old C57BL/6J mice (n = 8) (50 mg/kg total body weight, dissolved in 30% weight/volume ethanol by gavage and enema). Animals were monitored every 3h and were euthanized if they developed signs of disease or at 48h using CO_2 inhalation followed by cervical dislocation. Small intestine was harvested and processed for histological analyses. As previously described (Namachivayam et al., 2020), intestinal injury was confirmed by histopathological grading and was assigned by a blinded reviewer: grade 0 = no injury; grade 1 = mild injury: disruption of villus tips or mild separation of lamina propria in the small intestine; grade 2 = moderate injury: villus disruption, clear separation of lamina propria and/or edema in the submucosa; grade 3 = severe injury: transmural injury in the small intestine. Others were collected for qRT-PCR and immunofluorescence analysis. Control animals (n=8) received vehicle alone (30% ethanol).

High fat diet and palmitic acid treatment

High fat diet (containing 60% kcal from fats (Trophic Diet, TP23520, China)) was provided to Lsr^{ctrl} mice and Lsr$^{vill\ KO}$ mice beginning at the age of 8 weeks and extending for a period of 4 months. Control

mice were sex and age-matched and fed with standard chow ad libitum.

The Lsr[ctrl] mice and Lsr[vill KO] mice received an intragastric gavage of 0.2 ml 45% palmitic acid (MCE,

HY-N0830, China) dissolved in corn oil or corn oil alone. Mice were given 3 equally sized feedings daily

at 8:00 AM, 1:00 PM, and 6:00 PM. Gavage-fed animals were gradually adapted to this feeding regimen

for 5 days. At the end of day 5, mice were sacrificed 3 hours after the final gavage.

In vivo **administration of recombinant AAV**

AAVrh10 has been proved to be the most efficient rAAV serotype for intestine gene delivery (Polyak et

al., 2012). 7-day-old C57BL/6J mice received with 1×10^8 MOI virus genomic particles of Rh10-

pHBAAV-CMV-MCS-3flag-T2A-ZsGreen (AAV-CTL) or Rh10-pHBAAV-CMV-Lsr-3flag-T2A-

ZsGreen (AAV-*Lsr*) (prepared by Hanbio Biotechnology Inc, China) by intraperitoneal injection (n = 8).

1 week after transduction, mice were treated with TNBS to construct NEC models as described above.

To silence YAP *in vivo*, 1-week-old Lsr[vill KO] mice were injected intraperitoneally with a single dose of 2

$\times$ 10^8 MOI particles of lentivirus/pHBLV-U6-MCS-CMV-ZsGreen-PGK-PURO (CTL) or

lentivirus/pHBLV-U6-YAP-siRNA-CMV-ZsGreen-PGK-PURO (*Yap* shRNA) (prepared by Hanbio

Biotechnology Inc, China) (n = 8). Mice were monitored every day for general wellbeing. 2 weeks after

transduction, mice were euthanized using CO_2 inhalation followed by cervical dislocation. Small

intestine tissues were harvested for RNA isolation and immunofluorescence and histological staining.

In vivo **administration of YAP inhibitor verteporfin**

1-week-old Lsr[vill KO] mice were injected intraperitoneally with verteporfin (100 mg/kg body weight)

every 2 days for 2 weeks, (n = 8). Mice were monitored every day for general wellbeing. After 2 weeks,

mice were euthanized using CO_2 inhalation followed by cervical dislocation. Small intestine tissues were

harvested for RNA isolation and immunofluorescence and histological staining.

Histologic analysis

Human and mouse intestine tissues were fixed overnight at 4°C in 4% paraformaldehyde and embedded

in paraffin. Intestine sections (5 μm thickness) were prepared by a routine procedure, then standard histological staining with H&E and periodic acid-Schiff (PAS). Transmission electron microscopy (TEM) was performed on glutaraldehyde-fixed, epoxy-embedded intestine samples and stained with uranyl acetate and lead citrate.

Immunofluorescence staining

The small intestine (duodenal, jejunal and ileal) cross sections were embedded in OCT. Small intestine cryosections (8 μm) were mounted onto glass slides, fixed in cold methanol or acetone for 20 min at -20°C. For immunofluorescence of mouse or human organoid, organoid on collagen I coated confocal dish were fixed with 4% paraformaldehyde. And then washed three times with PBS for 10 min each time, and blocked for 1 h at room temperature in 10% BSA diluted in PBS (containing 0.01% Tween 20). Then incubated overnight at 4°C with appropriate primary antibodies. Washed as before and incubated using FITC and/or rhodamine (TRITC) conjugated secondary antibodies (Invitrogen), followed by DAPI (Sigma) nuclear staining. And then sealed with ProLong™ Gold antifade reagent and analyzed using a confocal laser microscope (LSM 880 Basic Operation, Germany).

For human Paraffin section (5 μm thickness), immunostaining was preceded by antigen retrieval in EDTA buffer (pH 8.0), and blocked for 1 h at room temperature in 10% BSA. The primary antibodies were anti-human: LSR, MPTX2, OFLM4 and incubated overnight at 4°C. Then incubated with FITC and/or rhodamine (TRITC) conjugated secondary antibodies (Invitrogen) for 1.5 h at room temperature, followed by DAPI (Sigma) nuclear staining. The images were acquired using confocal laser microscope (LSM 880 Basic Operation, Germany).

RNA-sequencing (RNA-seq)

Total RNA was extracted using Trizol reagent (Invitrogen, 15596018) from intestine of Lsr[ctrl] and Lsr[vill] [KO] mice at 4 weeks after birth. RNA-seq was performed on an Illumina Novaseq platform by Novogene (China). Paired-end clean reads were aligned to the mouse reference genome (Ensemble_GRCm38.90) with TopHat (version 2.0.12), and the aligned reads were used to quantify mRNA expression by using HTSeq-count (version 0.6.1). Differential expression analysis of two groups was performed using the

581 DESeq2 R package (1.16.1). The resulting P-values were adjusted using the Benjamini and Hochberg's

582 approach for controlling the false discovery rate. Genes with an adjusted *P-value <0.05* found by

583 DESeq2 were assigned as differentially expressed.

Crypt and Lgr5$^+$ ISCs isolation and culture

Small intestines crypt cultures were prepared using previously described methods with modifications

(Gjorevski et al., 2016; Goga et al., 2021). In short, small intestines were first flushed with PBS, then cut

longitudinally. The tissue was chopped into 2-3 mm pieces, and further washed with cold PBS. The tissue

fragments were incubated in 5 mM EDTA-PBS solution on ice. Then EDTA was removed, the

supernatant fractions enriched for crypts were passed through a 70-micron strainer (BD Biosciences) to

remove the remaining cells of the intestinal villi fraction. The filtered fractions were then centrifuged to

collect the crypt fraction. Crypts isolated from the small intestine of Lsr$^{vill\ KO}$ and Lsr^{ctrl} mice was

resuspended into matrigel with DMEM/F12 medium (1:1 ratio) and plated into 24 well plates (Serra et

al., 2019). Crypt fraction organoids were kept in IntestiCult Organoid Growth Medium (STEMCELL

Technologies) with 100 μg/ml Penicillin-Streptomycin for amplification and maintenance.

For ISCs isolation, crypts were isolated from the small intestine of Lsr$^{vill\ KO}$ and Lsr^{ctrl} mice as described

previously in the methods. ISCs were isolated from the small intestine by immunomagnetic beads. In

short, the crypt suspensions were centrifuged and resuspended in undiluted TrypLE Express (Invitrogen)

supplemented with DNase I (2000 U/μl, Roche), and 10 μM Y27632 (MCE) and incubated at 37°C for

8min. Then, cold DMEM/F12 (STEMCELL Technologies) were added to the samples and triturated

twice, followed by centrifugated and resuspend the cell pellet in DMEM/F12. Positive selection of ISCs

were achieved using magnetic beads (1 μm diameter, Invitrogen) coated with a mouse monoclonal

antibody against LGR5 (invitrogen). For PA treatment, Lgr5$^+$ ISCs isolated from mouse small intestine

were incubated in serum-free medium supplemented with 0.3 mM PA for 24 hours. And then cells were

harvested for RT-PCR, Western blot or immunofluorescence staining analysis.

Lentivirus transduction of organoids

For Lentivirus-Green-PURO-YAP (YAP-KD) virus infection experiments, cells recovery solution

(Corning) was used to harvest organoids and mechanically dissociated in Basal Media using a 10 ml

Pipette for 5-7 times up and down. Then, centrifuged the organoids at $250 \times g$ for 5 min, 4°C, and

resuspend use 400 μl full growth medium containing 1×10^7 MOI YAP-KD Lentivirus, then add to 100

μl matrigel together. The mixture was incubated at 37°C for 5 h, mixing every 30 min. Later, the mixture

was centrifuged at $250 \times g$ for 5 min at 4°C and organoids were seeded in matrigel and cultured for 3

days before harvesting for RNA isolation and immunofluorescence staining.

shRNA molecular cloning, screening and retrovirus production

The shRNA hairpin oligonucleotides, complementary to all the variants of human Lsr mRNA sequences

(NM_015925.7, NM_205834.4, NM_205835.4, NM_001260489.2, NM_001260490.2, and

NM_001385215.1) (Table S2), were synthesized by Sangon Biotechnology Inc, annealed, and cloned

into the moloney murine leukemia retrovirus vector pSIREN (Clontech), downstream of the human small

nuclear ribonucleoprotein U6 promoter to create the *LSR* shRNA constructs. Vesicular stomatitis virus

glycoprotein (VSV-G) pseudo typed retroviruses were produced in HEK293 cells and used to infect target

cells at a titer of 1×10^7 CFU/ml, as described previously (Gong et al., 2014).

Maintenance of iPSCs and *LSR* knockdown

Human iPSCs were maintained on matrigel (Corning) in mTesR1 medium (STEMCELL Technologies).

Cells were passaged approximately every 4-5 days. To passage iPSCs, they were washed with serum free

DMEM/F12 medium (STEMCELL Technologies) and incubated in DMEM/F12 with 1 mg/ml dispase

(Invitrogen) until cell edges started to detach from the dish. And then replaced with mTesR1. pipette

gently up and down to make the cells into small clumps and passaged onto fresh matrigel-coated plates.

For virus transfection experiments, above cells add full growth medium containing 1×10^7 MOI above

virus, the cells were incubated at 37°C for 12 h. Then, the medium was changed with fresh complete

medium and cultured for 48 h before harvesting for RNA isolation and organoid culture.

Differentiation of iPSCs into human intestine organoid

Differentiation of human iPSCs into organoid was run in accordance with the manufacturer's instructions

(STEMCELL Technologies). In brief, 1.5×10^5 iPSCs per ml were plated on matrigel coated tissue

culture plate (Thermo Fisher). Medium was changed to STEMdiff™ Endoderm Basal Medium

(STEMCELL Technologies) containing STEMdiff™ Definitive Endoderm Supplement CJ (STEMCELL

Technologies). On day 3, medium was changed to STEMdiff™ Endoderm Basal Medium containing

STEMdiff™. Gastrointestinal Supplement PK + STEMdiff™. Gastrointestinal Supplement UB

(STEMCELL Technologies). On day 4-9, medium was changed to Intestinal OGM (STEMCELL

Technologies) and incubate at 37°C with 5% CO2 and 95% humidity.

Quantitative real time PCR

The small intestine samples were homogenized in Trizol (Ambion, USA) to get total RNA. RNA was

reverse transcribed by using a reverse transcription system kit according to the instructions of the

manufacturer (Takara, Japan) followed by quantitative real time PCR (qRT-PCR) amplification using

SYBR Green PCR Master Mix (ABI, USA) and the Thermo Fisher QuantStudio3 system. The

Supplemental Table 2 contains the primer sequences used in this study. The expression levels of each

mRNA were calculated after normalizing to those of β-actin. Results were expressed as $2^{-\Delta Ct}$ values with

$\Delta CT = Ct_{gene} - Ct_{\beta\text{-actin}}$.

Western blot analysis

Total protein extracts were obtained by lysing isolated ISCs or small intestine tissues in RIPA lysis buffer

(Sangon Biotech) containing PMSF and phosphatase inhibitors (Sangon Biotech) on ice, supernatants

were collected after centrifugation at 4°C for 10 min at $12,000 \times g$. Then, the protein concentration was

assessed using a BCA protein assay (Takara). Proteins (50 μg) were separated in 10% SDS-PAGE and

then transferred onto PVDF membranes (0.22 μm), and then blocked with 5% skim milk dissolved in

TBST and incubated with the primary antibody at 4°C overnight. HRP-conjugated secondary antibodies

(Thermo Fisher) followed by ECL (Thermo Fisher) incubation allowed protein band detection. The

integration of all blots images was performed on Adobe Illustrator 2021. Image J v1.8.0 was used to

quantify western blotting results.

Co-Immunoprecipitation (Co-IP)

Isolated ISCs or HEK293 cells transfected with expressing vectors for LSR, YAP, and 14-3-3 zeta; LSR,

YAP, and 14-3-3 gamma; LSR, YAP, and PPP1CA; LSR, YAP, and PP2Ac; and YAP, 14-3-3 zeta and

PP2Ac were lysed in 50mM Tris (pH 8.0) by 25–30 repeated passages through a 25-gauge needle,

followed by centrifugation at $5,000 \times g$ for 20 min, 4℃. The membranes of lysis were extracted using

CSK buffer (1% Triton X-100; 150 mM NaCl; 50 mM Tris, pH 8.0; and protease inhibitors). The

membrane extract was precleared by incubation with protein A/G-sepharose (invitrogen) prior to Co-IP.

The precleared membrane extract was incubated for 16 h at 4℃ with anti-LSR, anti-YAP, anti-14-3-3

zeta, anti-14-3-3 gamma, anti-PPP1CA, anti-PP2Ac, anti-mouse IgG, or anti-Rabbit IgG antibodies.

Antibody-bound material was pelleted with protein A/G-sepharose, washed three times with CSK buffer,

and detected by immunoblotting.

LC-MS/MS Analysis

Follow the above Co-IP method, isolated ISCs and extracted the membranes of lysis. The precleared

membrane extract was incubated for 16 h at 4℃ with anti-LSR and anti-mouse IgG (as negative control)

antibodies. Antibody-bound material was pelleted with protein A/G-sepharose, washed three times with

CSK buffer, and performing SDS-PAGE, and then cut the gel and detected.

The separated peptides were analyzed by Q Exactive HF-X mass spectrometer (Thermo Fisher) by

Novogene (China). The resulting spectra from each fraction were searched separately against uniprot

database by the search engines: Proteome Discoverer 2.2 (PD 2.2, Thermo). The search parameters are

set as follows: mass tolerance for precursor ion was 10 ppm and mass tolerance for product ion was 0.02

Da. Carbamidomethyl was specified in PD 2.2 as fixed modifications. Oxidation of methionine (M) and

acetylation of the N-terminus were specified in PD 2.2 as variable modifications. A maximum of 2 missed

cleavage sites were allowed. The identified protein contains at least 1 unique peptide with FDR no more

than 1%. Proteins containing similar peptides that could not be distinguished by MS/MS analysis were

identified as a same protein group.

Data availability

The raw RNA-seq data were deposited in NCBI sequence read archive (SRA) database, with the

684 accession number PRJNA769233. Additional details on protocols and databases/datasets used in this

685 study are available from the corresponding authors on reasonable request.

686 **Statistical analysis**

687 The significance of differences between groups was tested by Prism 8 (GraphPad Software Inc.).

688 Statistical analysis was performed using the unpaired t test to determine differences between two groups

689 and ANOVA to compare data among groups. P-values less than 0.05 were interpreted as statistically

690 significant. All data are presented as mean ± SEM and other details such as the number of replicates and

691 the level of significance is mentioned in figure legends and supplementary information.

692 **Acknowledgments**

693 We thank Ping Li for animal husbandry. We thank the electron microscopic core lab of Shandong

694 University for assistance on electron microscopy imaging. The work is funded by National Natural

695 Science Foundation of China (81670620, 81870485, 81970578, and 81971281), Taishan Scholars

696 Program of Shandong Province (ts20190953), Natural Science Foundation of Shandong Province

697 (ZR2020QH066 and ZR2021QH115), Shandong First Medical University Academic Promotion Program

698 (2020LI001), and Binzhou Medical University Independent Research Program.

699 **Author contributions**

700 Y. A., C. W., and B. F. designed and conducted *in vivo* and *in vitro* experiments, performed data analysis,

701 and wrote the manuscript. Y. L., F. K., C. Z. and Z. C. performed histologic analysis. J. L., M. W. and H.

702 S. performed mice genotyping. Y. G. and S. Z. designed and jointly supervised the study, analyzed the

703 data, and wrote the manuscript.

704 **Competing Interests statement**

705 The authors declare no competing financial interests.

706 **References**

Adolph, T.E., M.F. Tomczak, L. Niederreiter, H.J. Ko, J. Bock, E. Martinez-Naves, J.N. Glickman, M.
Tschurtschenthaler, J. Hartwig, S. Hosomi, M.B. Flak, J.L. Cusick, K. Kohno, T. Iwawaki, S.
Billmann-Born, T. Raine, R. Bharti, R. Lucius, M.N. Kweon, S.J. Marciniak, A. Choi, S.J. Hagen,
S. Schreiber, P. Rosenstiel, A. Kaser, and R.S. Blumberg. 2013. Paneth cells as a site of
origin for intestinal inflammation. *Nature* 503:272-276.

Azzolin, L., T. Panciera, S. Soligo, E. Enzo, S. Bicciato, S. Dupont, S. Bresolin, C. Frasson, G. Basso, V.
Guzzardo, A. Fassina, M. Cordenonsi, and S. Piccolo. 2014. YAP/TAZ incorporation in the
beta-catenin destruction complex orchestrates the Wnt response. *Cell* 158:157-170.

Barry, E.R., T. Morikawa, B.L. Butler, K. Shrestha, R. de la Rosa, K.S. Yan, C.S. Fuchs, S.T. Magness, R.
Smits, S. Ogino, C.J. Kuo, and F.D. Camargo. 2013. Restriction of intestinal stem cell
expansion and the regenerative response by YAP. *Nature* 493:106-110.

Beyaz, S., M.D. Mana, J. Roper, D. Kedrin, A. Saadatpour, S.J. Hong, K.E. Bauer-Rowe, M.E. Xifaras,
A. Akkad, E. Arias, L. Pinello, Y. Katz, S. Shinagare, M. Abu-Remaileh, M.M. Mihaylova, D.W.
Lamming, R. Dogum, G. Guo, G.W. Bell, M. Selig, G.P. Nielsen, N. Gupta, C.R. Ferrone, V.
Deshpande, G.C. Yuan, S.H. Orkin, D.M. Sabatini, and O.H. Yilmaz. 2016. High-fat diet
enhances stemness and tumorigenicity of intestinal progenitors. *Nature* 531:53-58.

Bihain, B.E., and F.T. Yen. 1992. Free fatty acids activate a high-affinity saturable pathway for
degradation of low-density lipoproteins in fibroblasts from a subject homozygous for
familial hypercholesterolemia. *Biochemistry* 31:4628-4636.

Bjerknes, M., and H. Cheng. 1981. The stem-cell zone of the small intestinal epithelium. I. Evidence
from Paneth cells in the adult mouse. *The American journal of anatomy* 160:51-63.

Cai, J., N. Zhang, Y. Zheng, R.F. de Wilde, A. Maitra, and D. Pan. 2010. The Hippo signaling pathway
restricts the oncogenic potential of an intestinal regeneration program. *Genes &*
development 24:2383-2388.

Camargo, F.D., S. Gokhale, J.B. Johnnidis, D. Fu, G.W. Bell, R. Jaenisch, and T.R. Brummelkamp. 2007.
YAP1 increases organ size and expands undifferentiated progenitor cells. *Current biology :*
CB 17:2054-2060.

Chacon-Martinez, C.A., J. Koester, and S.A. Wickstrom. 2018. Signaling in the stem cell niche:
regulating cell fate, function and plasticity. *Development* 145:

Cheung, P., J. Xiol, M.T. Dill, W.C. Yuan, R. Panero, J. Roper, F.G. Osorio, D. Maglic, Q. Li, B. Gurung,
R.A. Calogero, O.H. Yilmaz, J. Mao, and F.D. Camargo. 2020. Regenerative
Reprogramming of the Intestinal Stem Cell State via Hippo Signaling Suppresses
Metastatic Colorectal Cancer. *Cell stem cell* 27:590-604 e599.

Coutinho, H.B., H.C. da Mota, V.B. Coutinho, T.I. Robalinho, A.F. Furtado, E. Walker, G. King, Y.R.
Mahida, H.F. Sewell, and D. Wakelin. 1998. Absence of lysozyme (muramidase) in the
intestinal Paneth cells of newborn infants with necrotising enterocolitis. *Journal of clinical*
pathology 51:512-514.

Dong, X., X.B. Zhang, P. Liu, Y. Tian, L. Li, and P. Gong. 2022. Lipolysis-Stimulated Lipoprotein
Receptor Impairs Hepatocellular Carcinoma and Inhibits the Oncogenic Activity of YAP1
via PPPY Motif. *Front Oncol* 12:

Dzulko, M., M. Pons, A. Henke, G. Schneider, and O.H. Kramer. 2020. The PP2A subunit PR130 is a
key regulator of cell development and oncogenic transformation. *Biochimica et*
biophysica acta. Reviews on cancer 1874:188453.

Fan, S., Y. Gao, A. Qu, Y. Jiang, H. Li, G. Xie, X. Yao, X. Yang, S. Zhu, T. Yagai, J. Tian, R. Wang, F.J.

Gonzalez, M. Huang, and H. Bi. 2022. YAP-TEAD mediates PPAR alpha-induced
752   hepatomegaly and liver regeneration in mice. *Hepatology* 75:74-88.

753   Frost, B.L., B.P. Modi, T. Jaksic, and M.S. Caplan. 2017. New Medical and Surgical Insights Into
754   Neonatal Necrotizing Enterocolitis: A Review. *JAMA pediatrics* 171:83-88.

755   Gjorevski, N., M. Nikolaev, T.E. Brown, O. Mitrofanova, N. Brandenberg, F.W. DelRio, F.M. Yavitt, P.
756   Liberali, K.S. Anseth, and M.P. Lutolf. 2022. Tissue geometry drives deterministic organoid
757   patterning. *Science* 375:eaaw9021.

758   Gjorevski, N., N. Sachs, A. Manfrin, S. Giger, M.E. Bragina, P. Ordonez-Moran, H. Clevers, and M.P.
759   Lutolf. 2016. Designer matrices for intestinal stem cell and organoid culture. *Nature*
760   539:560-564.

761   Goga, A., B. Yagabasan, K. Herrmanns, S. Godbersen, P.N. Silva, R. Denzler, M. Zund, M. Furter, G.
762   Schwank, S. Sunagawa, W.D. Hardt, and M. Stoffel. 2021. miR-802 regulates Paneth cell
763   function and enterocyte differentiation in the mouse small intestine. *Nature*
764   *communications* 12:3339.

765   Gomez-Suarez, M., I.Z. Gutierrez-Martinez, J.A. Hernandez-Trejo, M. Hernandez-Ruiz, D. Suarez-
766   Perez, A. Candelario, R. Kamekura, O. Medina-Contreras, M. Schnoor, V. Ortiz-Navarrete,
767   N. Villegas-Sepulveda, C. Parkos, A. Nusrat, and P. Nava. 2016. 14-3-3 Proteins regulate
768   Akt Thr308 phosphorylation in intestinal epithelial cells. *Cell Death Differ* 23:1060-1072.

769   Gong, Y., M. Yu, J. Yang, E. Gonzales, R. Perez, M. Hou, P. Tripathi, K.S. Hering-Smith, L.L. Hamm,
770   and J. Hou. 2014. The Cap1-claudin-4 regulatory pathway is important for renal chloride
771   reabsorption and blood pressure regulation. *Proceedings of the National Academy of*
772   *Sciences of the United States of America* 111:E3766-3774.

773   Gregorieff, A., Y. Liu, M.R. Inanlou, Y. Khomchuk, and J.L. Wrana. 2015. Yap-dependent
774   reprogramming of Lgr5(+) stem cells drives intestinal regeneration and cancer. *Nature*
775   526:715-718.

776   Guillermin, O., N. Angelis, C.M. Sidor, R. Ridgway, A. Baulies, A. Kucharska, P. Antas, M.R. Rose, J.
777   Cordero, O. Sansom, V.S.W. Li, and B.J. Thompson. 2021. Wnt and Src signals converge
778   on YAP-TEAD to drive intestinal regeneration. *The EMBO journal* 40:e105770.

779   Hayashi, S., and A.P. McMahon. 2002. Efficient recombination in diverse tissues by a tamoxifen-
780   inducible form of Cre: a tool for temporally regulated gene activation/inactivation in the
781   mouse. *Developmental biology* 244:305-318.

782   Hemmasi, S., B.A. Czulkies, B. Schorch, A. Veit, K. Aktories, and P. Papatheodorou. 2015. Interaction
783   of the Clostridium difficile Binary Toxin CDT and Its Host Cell Receptor, Lipolysis-
784   stimulated Lipoprotein Receptor (LSR). *The Journal of biological chemistry* 290:14031-
785   14044.

786   Hu, J.K., W. Du, S.J. Shelton, M.C. Oldham, C.M. DiPersio, and O.D. Klein. 2017. An FAK-YAP-mTOR
787   Signaling Axis Regulates Stem Cell-Based Tissue Renewal in Mice. *Cell stem cell* 21:91-
788   106 e106.

789   Kang, J.X., J.B. Wan, and C. He. 2014. Concise review: Regulation of stem cell proliferation and
790   differentiation by essential fatty acids and their metabolites. *Stem Cells* 32:1092-1098.

791   Konno, T., T. Kohno, S. Kikuchi, H. Shimada, S. Satohisa, T. Saito, M. Kondoh, and T. Kojima. 2020.
792   Epithelial barrier dysfunction and cell migration induction via JNK/cofilin/actin by
793   angubindin-1. *Tissue barriers* 8:1695475.

Koo, J.H., and K.L. Guan. 2018. Interplay between YAP/TAZ and Metabolism. *Cell metabolism*

28:196-206.

Lapi, E., S. Di Agostino, S. Donzelli, H. Gal, E. Domany, G. Rechavi, P.P. Pandolfi, D. Givol, S. Strano,
X. Lu, and G. Blandino. 2008. PML, YAP, and p73 are components of a proapoptotic
autoregulatory feedback loop. *Molecular cell* 32:803-814.

Li, Q., Y. Sun, G.K. Jarugumilli, S. Liu, K. Dang, J.L. Cotton, J. Xiol, P.Y. Chan, M. DeRan, L. Ma, R. Li,
L.J. Zhu, J.H. Li, A.B. Leiter, Y.T. Ip, F.D. Camargo, X. Luo, R.L. Johnson, X. Wu, and J. Mao.
2020. Lats1/2 Sustain Intestinal Stem Cells and Wnt Activation through TEAD-Dependent
and Independent Transcription. *Cell stem cell* 26:675-692 e678.

Li, V.S.W. 2019. Yap in regeneration and symmetry breaking. *Nature cell biology* 21:665-667.

Lueschow, S.R., and S.J. McElroy. 2020. The Paneth Cell: The Curator and Defender of the Immature
Small Intestine. *Frontiers in immunology* 11:587.

Lueschow, S.R., J. Stumphy, H. Gong, S.L. Kern, T.G. Elgin, M.A. Underwood, K.M. Kalanetra, D.A.
Mills, M.H. Wong, D.K. Meyerholz, M. Good, and S.J. McElroy. 2018. Loss of murine Paneth
cell function alters the immature intestinal microbiome and mimics changes seen in
neonatal necrotizing enterocolitis. *PloS one* 13:e0204967.

Lukonin, I., D. Serra, L. Challet Meylan, K. Volkmann, J. Baaten, R. Zhao, S. Meeusen, K. Colman, F.
Maurer, M.B. Stadler, J. Jenkins, and P. Liberali. 2020. Phenotypic landscape of intestinal
organoid regeneration. *Nature* 586:275-280.

Madison, B.B., L. Dunbar, X.T. Qiao, K. Braunstein, E. Braunstein, and D.L. Gumucio. 2002. Cis
elements of the villin gene control expression in restricted domains of the vertical (crypt)
and horizontal (duodenum, cecum) axes of the intestine. *The Journal of biological*
chemistry 277:33275-33283.

Maehama, T., M. Nishio, J. Otani, T.W. Mak, and A. Suzuki. 2021. The role of Hippo-YAP signaling
in squamous cell carcinomas. *Cancer science* 112:51-60.

Mann, C.J., J. Khallou, O. Chevreuil, A.A. Troussard, L.M. Guermani, K. Launay, B. Delplanque, F.T.
Yen, and B.E. Bihain. 1995. Mechanism of activation and functional significance of the
lipolysis-stimulated receptor. Evidence for a role as chylomicron remnant receptor.
Biochemistry 34:10421-10431.

Masuda, S., Y. Oda, H. Sasaki, J. Ikenouchi, T. Higashi, M. Akashi, E. Nishi, and M. Furuse. 2011. LSR
defines cell corners for tricellular tight junction formation in epithelial cells. *Journal of cell*
science 124:548-555.

Meng, Z., T. Moroishi, and K.L. Guan. 2016. Mechanisms of Hippo pathway regulation. *Genes &*
development 30:1-17.

Mesli, S., S. Javorschi, A.M. Berard, M. Landry, H. Priddle, D. Kivlichan, A.J. Smith, F.T. Yen, B.E. Bihain,
and M. Darmon. 2004. Distribution of the lipolysis stimulated receptor in adult and
embryonic murine tissues and lethality of LSR-/- embryos at 12.5 to 14.5 days of gestation.
European journal of biochemistry 271:3103-3114.

MohanKumar, K., N. Kaza, R. Jagadeeswaran, S.A. Garzon, A. Bansal, A.R. Kurundkar, K.
Namachivayam, J.I. Remon, C.R. Bandepalli, X. Feng, J.H. Weitkamp, and A. Maheshwari.
2012. Gut mucosal injury in neonates is marked by macrophage infiltration in contrast to
pleomorphic infiltrates in adult: evidence from an animal model. *American journal of*
physiology. Gastrointestinal and liver physiology 303:G93-102.

Namachivayam, K., K. MohanKumar, D.R. Shores, S.K. Jain, J. Fundora, A.D. Everett, L. He, H. Pan,
S.A. Wickline, and A. Maheshwari. 2020. Targeted inhibition of thrombin attenuates

murine neonatal necrotizing enterocolitis. *Proceedings of the National Academy of*
Sciences of the United States of America 117:10958-10969.

Narvekar, P., M. Berriel Diaz, A. Krones-Herzig, U. Hardeland, D. Strzoda, S. Stohr, M. Frohme, and
S. Herzig. 2009. Liver-specific loss of lipolysis-stimulated lipoprotein receptor triggers
systemic hyperlipidemia in mice. *Diabetes* 58:1040-1049.

Oh, H., and K.D. Irvine. 2009. In vivo analysis of Yorkie phosphorylation sites. *Oncogene* 28:1916-
1927.

Papatheodorou, P., J.E. Carette, G.W. Bell, C. Schwan, G. Guttenberg, T.R. Brummelkamp, and K.
Aktories. 2011. Lipolysis-stimulated lipoprotein receptor (LSR) is the host receptor for the
binary toxin Clostridium difficile transferase (CDT). *Proceedings of the National Academy*
of Sciences of the United States of America 108:16422-16427.

Paroni, G., N. Cernotta, C. Dello Russo, P. Gallinari, M. Pallaoro, C. Foti, F. Talamo, L. Orsatti, C.
Steinkuhler, and C. Brancolini. 2008. PP2A regulates HDAC4 nuclear import. *Molecular*
biology of the cell 19:655-667.

Polyak, S., A. Mach, S. Porvasnik, L. Dixon, T. Conlon, K.E. Erger, A. Acosta, A.J. Wright, M. Campbell-
Thompson, I. Zolotukhin, C. Wasserfall, and C. Mah. 2012. Identification of adeno-
associated viral vectors suitable for intestinal gene delivery and modulation of
experimental colitis. *American journal of physiology. Gastrointestinal and liver physiology*
302:G296-308.

Poon, K., S. Mandava, K. Chen, J.R. Barson, S. Buschlen, and S.F. Leibowitz. 2013. Prenatal Exposure
to Dietary Fat Induces Changes in the Transcriptional Factors, TEF and YAP, Which May
Stimulate Differentiation of Peptide Neurons in Rat Hypothalamus. *Plos One* 8:

Qi, Q., D.Y. Li, H.R. Luo, K.L. Guan, and K. Ye. 2015. Netrin-1 exerts oncogenic activities through
enhancing Yes-associated protein stability. *Proceedings of the National Academy of*
Sciences of the United States of America 112:7255-7260.

Reaves, D.K., K.A. Hoadley, K.D. Fagan-Solis, D.D. Jima, M. Bereman, L. Thorpe, J. Hicks, D.
McDonald, M.A. Troester, C.M. Perou, and J.M. Fleming. 2017. Nuclear Localized LSR: A
Novel Regulator of Breast Cancer Behavior and Tumorigenesis. *Mol Cancer Res* 15:165-
178.

Resnik-Docampo, M., C.L. Koehler, R.I. Clark, J.M. Schinaman, V. Sauer, D.M. Wong, S. Lewis, C.
D'Alterio, D.W. Walker, and D.L. Jones. 2017. Tricellular junctions regulate intestinal stem
cell behaviour to maintain homeostasis. *Nature cell biology* 19:52-59.

Santinon, G., A. Pocaterra, and S. Dupont. 2016. Control of YAP/TAZ Activity by Metabolic and
Nutrient-Sensing Pathways. *Trends in cell biology* 26:289-299.

Sato, T., R.G. Vries, H.J. Snippert, M. van de Wetering, N. Barker, D.E. Stange, J.H. van Es, A. Abo, P.
Kujala, P.J. Peters, and H. Clevers. 2009. Single Lgr5 stem cells build crypt-villus structures
in vitro without a mesenchymal niche. *Nature* 459:262-265.

Schlegelmilch, K., M. Mohseni, O. Kirak, J. Pruszak, J.R. Rodriguez, D. Zhou, B.T. Kreger, V.
Vasioukhin, J. Avruch, T.R. Brummelkamp, and F.D. Camargo. 2011. Yap1 acts downstream
of alpha-catenin to control epidermal proliferation. *Cell* 144:782-795.

Serra, D., U. Mayr, A. Boni, I. Lukonin, M. Rempfler, L. Challet Meylan, M.B. Stadler, P. Strnad, P.
Papasaikas, D. Vischi, A. Waldt, G. Roma, and P. Liberali. 2019. Self-organization and
symmetry breaking in intestinal organoid development. *Nature* 569:66-72.

Shimada, H., S. Abe, T. Kohno, S. Satohisa, T. Konno, S. Takahashi, T. Hatakeyama, C. Arimoto, T.

Kakuki, Y. Kaneko, K.I. Takano, T. Saito, and T. Kojima. 2017. Loss of tricellular tight junction
884   protein LSR promotes cell invasion and migration via upregulation of TEAD1/AREG in
885   human endometrial cancer. *Sci Rep* 7:37049.

886 Sohet, F., C. Lin, R.N. Munji, S.Y. Lee, N. Ruderisch, A. Soung, T.D. Arnold, N. Derugin, Z.S. Vexler,
887   F.T. Yen, and R. Daneman. 2015. LSR/angulin-1 is a tricellular tight junction protein
888   involved in blood-brain barrier formation. *The Journal of cell biology* 208:703-711.

889 Stevers, L.M., E. Sijbesma, M. Botta, C. MacKintosh, T. Obsil, I. Landrieu, Y. Cau, A.J. Wilson, A.
890   Karawajczyk, J. Eickhoff, J. Davis, M. Hann, G. O'Mahony, R.G. Doveston, L. Brunsveld, and
891   C. Ottmann. 2018. Modulators of 14-3-3 Protein-Protein Interactions. *Journal of*
892   *medicinal chemistry* 61:3755-3778.

893 Sugawara, T., K. Furuse, T. Otani, T. Wakayama, and M. Furuse. 2021. Angulin-1 seals tricellular
894   contacts independently of tricellulin and claudins. *The Journal of cell biology* 220:

895 Sukhotnik, I., L. Hayari, Y. Bashenko, E. Chemodanov, J. Mogilner, R. Shamir, F. Bar Yosef, R. Shaoul,
896   and A.G. Coran. 2008. Dietary palmitic acid modulates intestinal re-growth after massive
897   small bowel resection in a rat. *Pediatric surgery international* 24:1313-1321.

898 Sun, B., F.J. Zhong, C. Xu, Y.M. Li, Y.R. Zhao, M.M. Cao, and L.Y. Yang. 2021. Programmed cell death
899   10 promotes metastasis and epithelial-mesenchymal transition of hepatocellular
900   carcinoma via PP2Ac-mediated YAP activation. *Cell death & disease* 12:849.

901 Takahashi, Y., S. Serada, T. Ohkawara, M. Fujimoto, K. Hiramatsu, Y. Ueda, T. Kimura, H. Takemori,
902   and T. Naka. 2021. LSR promotes epithelial ovarian cancer cell survival under energy stress
903   through the LKB1-AMPK pathway. *Biochem Bioph Res Co* 537:93-99.

904 Totaro, A., T. Panciera, and S. Piccolo. 2018. YAP/TAZ upstream signals and downstream responses.
905   *Nature cell biology* 20:888-899.

906 Underwood, M.A. 2012. Paneth cells and necrotizing enterocolitis. *Gut microbes* 3:562-565.

907 Wang, L., S. Wang, Y. Shi, R. Li, S. Gunther, Y.T. Ong, M. Potente, Z. Yuan, E. Liu, and S. Offermanns.
908   2020a. YAP and TAZ protect against white adipocyte cell death during obesity. *Nat*
909   *Commun* 11:5455.

910 Wang, L., S.P. Wang, Y. Shi, R. Li, S. Gunther, Y.T. Ong, M. Potente, Z.Y. Yuan, E.Q. Liu, and S.
911   Offermanns. 2020b. YAP and TAZ protect against white adipocyte cell death during
912   obesity. *Nat Commun* 11:

913 Wehkamp, J., and E.F. Stange. 2020. An Update Review on the Paneth Cell as Key to Ileal Crohn's
914   Disease. *Frontiers in immunology* 11:646.

915 Weis, V.G., A.C. Deal, G. Mekkey, C. Clouse, M. Gaffley, E. Whitaker, C.B. Peeler, J.A. Weis, M.Z.
916   Schwartz, and A. Atala. 2021. Human placental-derived stem cell therapy ameliorates
917   experimental necrotizing enterocolitis. *American journal of physiology. Gastrointestinal*
918   *and liver physiology* 320:G658-G674.

919 Xi, P., L. Zhou, M. Wang, J.P. Liu, and Y.S. Cong. 2013. Serine/threonine-protein phosphatase 2A
920   physically interacts with human telomerase reverse transcriptase hTERT and regulates its
921   subcellular distribution. *Journal of cellular biochemistry* 114:409-417.

922 Yan, H., C.C. Yu, S.A. Fine, A.L. Youssof, Y.R. Yang, J. Yan, D.C. Karg, E.C. Cheung, R.A. Friedman, H.
923   Ying, E.I. Chen, J. Luo, Y. Miao, W. Qiu, and G.H. Su. 2021. Loss of the wild-type KRAS
924   allele promotes pancreatic cancer progression through functional activation of YAP1.
925   *Oncogene* 40:6759-6771.

926 Yen, F.T., C.J. Mann, L.M. Guermani, N.F. Hannouche, N. Hubert, C.A. Hornick, V.N. Bordeau, G.

927    Agnani, and B.E. Bihain. 1994. Identification of a lipolysis-stimulated receptor that is
928    distinct from the LDL receptor and the LDL receptor-related protein. *Biochemistry*
929    33:1172-1180.
930 Yen, F.T., M. Masson, N. Clossais-Besnard, P. Andre, J.M. Grosset, L. Bougueleret, J.B. Dumas, O.
931    Guerassimenko, and B.E. Bihain. 1999. Molecular cloning of a lipolysis-stimulated remnant
932    receptor expressed in the liver. *The Journal of biological chemistry* 274:13390-13398.
933 Yuan, L., Y. Mao, W. Luo, W. Wu, H. Xu, X.L. Wang, and Y.H. Shen. 2017. Palmitic acid dysregulates
934    the Hippo-YAP pathway and inhibits angiogenesis by inducing mitochondrial damage
935    and activating the cytosolic DNA sensor cGAS-STING-IRF3 signaling mechanism. *J Biol*
936    *Chem* 292:15002-15015.
937 Zhao, B., L. Li, K. Tumaneng, C.Y. Wang, and K.L. Guan. 2010. A coordinated phosphorylation by
938    Lats and CK1 regulates YAP stability through SCF(beta-TRCP). *Genes & development*
939    24:72-85.
940 Zhou, D., Y. Zhang, H. Wu, E. Barry, Y. Yin, E. Lawrence, D. Dawson, J.E. Willis, S.D. Markowitz, F.D.
941    Camargo, and J. Avruch. 2011. Mst1 and Mst2 protein kinases restrain intestinal stem cell
942    proliferation and colonic tumorigenesis by inhibition of Yes-associated protein (Yap)
943    overabundance. *Proceedings of the National Academy of Sciences of the United States of*
944    *America* 108:E1312-1320.
945 Zolnierowicz, S. 2000. Type 2A protein phosphatase, the complex regulator of numerous signaling
946    pathways. *Biochemical pharmacology* 60:1225-1235.

Fig. 1. Generation and characterization of conditional Lsr knockout mice.

(A and B) Expression of *Lsr* mRNA assessed by qRT–PCR in small intestines from Lsr^ctrl and Lsr^vill KO mice (A) and Lsr^ctrl and Lsr^-/- mice (B) (n=4). (C and D) Western blot and quantification analysis of LSR in small intestines from Lsr^ctrl and Lsr^vill KO mice (C) and Lsr^ctrl and Lsr^-/- mice (D) (n=3). (E and F) LSR immunofluorescence and quantification analysis of small intestines from Lsr^ctrl and Lsr^vill KO mice (E) and Lsr^ctrl and Lsr^-/- mice (F) (n=6). (G) Longitudinal views of intestine from Lsr^ctrl and Lsr^vill KO mice at 4, 8 and 24 weeks of age. (H) Longitudinal views of intestine from Lsr^ctrl and Lsr^-/- mice. (I) Transverse views of intestine from Lsr^ctrl and Lsr^vill KO littermate mice. (J) Analysis of intestinal wall thickness of

957   Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=3). (K) Representative H&E-stained transverse segments and quantification

958   analysis of Lsr^{ctrl} and Lsr$^{vill\ KO}$ mouse intestines (n=6). (L) Representative H&E-stained small intestine

959   and quantification analysis of Lsr^{ctrl} and Lsr$^{-/-}$ mice (n=6). Scale bars: E, 50 μm; F, 20 μm; K and J, 100

960   μm. *P < 0.05, **P < 0.01, ***P < 0.001.

Fig. 2. LSR ablation results in increased numbers of proliferating cells and loss of Paneth cell lineage in the small intestine.

(A and B) Ki67 (A) and PCNA (B) immunofluorescence and quantification analysis in duodenal, jejunal and ileal segments from Lsr^ctrl and Lsr^vill KO mice (n=6). (C and D) Western blot and quantification analysis of Ki67 and PCNA protein expression in duodenal, jejunal and ileal segments from Lsr^ctrl and

Lsr$^{vill\ KO}$ mice (C) and Lsr^{ctrl} and Lsr$^{-/-}$ mice (D) (n=3). (E and F) Expression of *Olfm4* mRNA assessed

by qRT–PCR in small intestines from Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (E) and Lsr^{ctrl} and Lsr$^{-/-}$ mice (F) (n=4).

(G) Differentially expressed genes (Lsr$^{vill\ KO}$ vs. Lsr^{ctrl}) are shown on a volcano plot (n=4; fold change >

2, *P < 0.05*). (H) Heatmap showing unsupervised hierarchical clustering of Paneth cell-related genes. (I)

MPTX2 immunofluorescence and quantification analysis in duodenal, jejunal and ileal segments from

Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=6). (J) qRT-PCR validating several downregulated Paneth cell markers in

small intestines from Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=4). (K) MPTX2 immunofluorescence and quantification

analysis in duodenal, jejunal and ileal segments from Lsr^{ctrl} and Lsr$^{-/-}$ mice (n=6). (L) qRT–PCR analysis

of the mRNA expression of columnar absorptive cell markers (*Cdx1* and *Cdx2*), goblet cell markers

(*Muc2* and *Spdef*), and enteroendocrine cell markers (*Neurog3* and *Chga*) in small intestines from Lsr^{ctrl}

and Lsr$^{vill\ KO}$ mice (n=8). Scale bars: A, B, I and K, 50 μm. *P < 0.05, **P < 0.01, ***P < 0.001.*

Fig. 3. LSR is required for Lgr5⁺ ISCs to Paneth cell differentiation.

(A) LSR and OLFM4 immunofluorescence in small intestines from Lsr^ctrl mice (arrows indicated colocalization of LSR and OLFM4 in ISCs). (B) Expression of *Lsr* mRNA assessed by qRT–PCR in small intestines from Lsr^ctrl and Lsr^ISC KO mice (n=4). (C) Western blot and quantification analysis of LSR in small intestines from Lsr^ctrl and Lsr^ISC KO mice (n=3). (D) Expression of Paneth cell markers (*Lyz1, Itln1, Mptx2, Defa5* and *Cd24a*) mRNA assessed by qRT–PCR in small intestines from Lsr^ctrl and Lsr^ISC

KO mice (n=4). (E) MPTX2 immunofluorescence and quantification analysis in duodenal, jejunal and

ileal segments from Lsr[ctrl] and Lsr[ISC KO] mice (n=6). (F and G) LSR immunofluorescence (F) and bright

field images (G) of organoids established from intestinal crypts of Lsr[ctrl] and Lsr[vill KO] mice. (H and I)

Quantification of the average number of buds per organoid (H) and percentage of organoid formation per

well (n=3 wells per group) (I). (J-P) OLFM4 (J), EpCAM (K), LYZ1 (L), MPTX2 (M), MUC2 (N),

CHGA (O) and CDX2 (P) immunofluorescence of organoids established from intestinal crypts of Lsr[ctrl]

and Lsr[vill KO] mice. (Q) Expression of *LSR* mRNA assessed by qRT–PCR in control and *LSR* knockdown

human iPSCs (n=3). (R-V) Bright field images (R) and LSR (S) OLFM4 (T), EpCAM (U) and MPTX2

(V) immunofluorescence of organoids differentiated from control and *LSR* knockdown human iPSCs.

Scale bars: A, 10μm; E, S, T and V, 50 μm; F, J, K, L, M, N, O, P and U, 20 μm; G and R, 125 μm. *P <

*0.05, **P < 0.01, ***P < 0.001.*

Fig. 4. Loss of LSR results in upregulation and activation of YAP.

(A) LSR and YAP immunofluorescence in small intestines from Lsr^ctrl and Lsr^ISC KO mice (arrows indicated the ISCs). (B-D) Western blot and quantification analysis for YAP of small intestine lysates from Lsr^ctrl and Lsr^-/- mice (B), Lsr^ctrl and Lsr^vill KO mice (C) and Lsr^ctrl and Lsr^ISC KO mice (D) (n=3). (E-

G) Expression of *Yap, Edn1, Ctgf,* and *Cyr61* mRNA assessed by qRT–PCR in small intestines from Lsr^{ctrl} and Lsr$^{-/-}$ mice (E), Lsr^{ctrl} and Lsr$^{vill\,KO}$ mice (F) and Lsr^{ctrl} and Lsr$^{ISC\,KO}$ mice (G) (n=4). (H) MPTX2 immunofluorescence and quantification analysis in small intestines from Lsr^{ctrl} mice, Lsr$^{vill\,KO}$ mice, and Lsr$^{vill\,KO}$ mice treated with VP (n=6). (I) H&E staining and quantification analysis in small intestine from Lsr^{ctrl} mice, Lsr$^{vill\,KO}$ mice, and Lsr$^{vill\,KO}$ mice treated with VP (n=6). (J) qRT-PCR analysis for YAP targets in small intestine from Lsr^{ctrl} mice, Lsr$^{vill\,KO}$ mice, and Lsr$^{vill\,KO}$ mice treated with VP (n=4). (K) MPTX2 immunofluorescence and quantification analysis in small intestines from Lsr^{ctrl} mice, Lsr$^{vill\,KO}$ mice, and Lsr$^{vill\,KO}$ mice transduced with lentivirus expressing *Yap* shRNA (n=6). (L) H&E staining and quantification analysis in small intestine from Lsr^{ctrl} mice, Lsr$^{vill\,KO}$ mice, and Lsr$^{vill\,KO}$ mice treated with lentivirus expressing *Yap* shRNA (n=6). (M) qRT–PCR analysis for YAP targets in small intestine from Lsr^{ctrl} mice, Lsr$^{vill\,KO}$ mice, and Lsr$^{vill\,KO}$ mice treated with lentivirus expressing *Yap* shRNA (n=4). (N) YAP immunofluorescence and quantification analysis in ISCs isolated from Lsr^{ctrl} and Lsr$^{ISC\,KO}$ mice (n=6). Scale bars: A and N, 10 μm; H and K, 50 μm; I and L, 100 μm. *P < 0.05, **P < 0.01, ***P < 0.001.*

1019

Fig. 5. 14-3-3 zeta and PP2Ac are involved in the regulation of YAP by LSR.

(A and B) Western blot analysis for YAP in isolated ISCs treated with 5 mg/ml CHX for 0-4 h (A) or

pretreated with 10 mM MG132 for 2 h and then treated with 5 mg/ml CHX for 0-4 h (B). And line graph

showing the relative YAP protein levels quantified by YAP/β-actin ratio, which was arbitrarily set to 100%

at the 0 hour point. (C) Western blot and quantification analysis for YAP, p-YAP (Ser381), LGR5 in

isolated ISCs from Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=3). (D) YAP ubiquitination levels and quantification

analysis in isolated ISCs from Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice. Ubiquitinated YAP was first immunoprecipitated

with anti-YAP antibody or anti-p-YAP (S381) antibody and then detected with anti-ubiquitin antibody

(n=3). (E) Mass spectrometry analysis of LSR immunocomplexes precipitated from isolated ISCs. (F)

Co-IP showing that endogenous YAP, LSR, and 14-3-3 zeta interacts with each other in isolated ISCs

from Lsr^ctrl mice. (G) Co-IP showing that endogenous YAP interacts with endogenous LSR, but not

PP2Ac in isolated ISCs from Lsr^ctrl mice. (H) Co-IP showing that endogenous YAP interacts with

endogenous PP2Ac, but not 14-3-3 zeta in isolated ISCs from Lsr^vill KO mice. (I) Western blot and

quantification analysis of YAP and Ser381-phosphorylated YAP in HEK293 cells transfected YAP alone

or together with PP2Ac (n=3). For all IP or Co-IP analysis, antibodies used for immunoprecipitation are

shown above the lanes; antibody for blot visualization is shown on the right. *P < 0.05, **P < 0.01,

***P < 0.001.

1037

Fig. 6. Loss of _Lsr_ exacerbates inflammation in a mouse model of NEC.

(A) Severity of intestinal injury in control and TNBS-treated mice (n=8). (B) Expression of _Il1α, Il2, Il6,_

Ifnγ and _Tnfα_ mRNA assessed by qRT-PCR in small intestines from control and TNBS-treated mice (n=8).

(C and D) Quantification analysis of F4/80 (C) and LY6G (D) mean intensity per villi in small intestines

from control and TNBS-treated mice (n=6). (E) Expression of _Lsr_ mRNA assessed by qRT–PCR in small

intestines from control and TNBS-treated mice (n=4). (F) LSR immunofluorescence and quantification analysis in small intestines from control and TNBS-treated mice (n=6). (G) Quantification analysis of MPTX2$^+$ cell per crypt in small intestines from control and TNBS-treated mice (n=6). (H) Expression of *Lyz1, Itln1, Mptx2* and *Defa5* mRNA assessed by qRT-PCR in small intestines from control and TNBS-treated mice (n=8). (I) Severity of intestinal injury in TNBS-treated Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=8). (J) Expression of *Lsr, Mptx2, Il1α, Il2, Il6, Ifnγ* and *Tnfα* mRNA assessed by qRT–PCR in small intestines from TNBS-treated Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=8). (K and L) Quantification analysis of F4/80 (K) and LY6G (L) mean intensity per villi in small intestines from TNBS-treated Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice (n=6). (M) Severity of intestinal injury in TNBS-treated mice administered with AAV-CTL or AAV-Lsr (n=8). (N) Expression of *Il1α, Il2, Il6, Ifnγ* and *Tnfα* mRNA assessed by qRT–PCR in small intestines from TNBS-treated mice administered with AAV-CTL or AAV-Lsr (n=8). (O and P) Quantification analysis of F4/80 (O) and LY6G (P) mean intensity per villi in small intestines from TNBS-treated mice administered with AAV-CTL or AAV-Lsr (n=6). (Q) Expression of *Lyz1, Itln1, Mptx2, Defa5* and *Lsr* mRNA assessed by qRT–PCR in small intestines from TNBS-treated mice administered with AAV-CTL or AAV-Lsr (n=8). (R) MPTX2 immunofluorescence and quantification analysis in small intestines from TNBS-treated mice administered with AAV-CTL or AAV-Lsr (n=6). (S) Representative H&E-stained small intestinal biopsy specimens of subjects with NEC and a control. (T-V) LSR (T), LSR and OLFM4 (U), and MPTX2 (V) immunofluorescence and quantification analysis in small intestinal biopsy specimens of subjects with NEC and a control (arrows indicated Paneth cells) (n=6). Scale bars: F, R, T and V, 50 μm; S, 100 μm; U, 20 μm. *P < 0.05, **P < 0.01, ***P < 0.001.

1065

Fig. 7. LSR mediates the inhibitory effect of PA on YAP.

(A) LSR immunofluorescence and quantification analysis in small intestines from control or HFD-fed mice (n=6). (B) qRT–PCR analysis in small intestines from control or HFD-fed mice (n=5). (C) LSR immunofluorescence and quantification analysis in small intestines from vehicle control or PA-treated mice (n=6). (D) qRT–PCR analysis in small intestines from vehicle control or PA-treated mice (n=5). (E) YAP immunofluorescence and quantification analysis in small intestines from control or HFD-fed Lsr^ctrl and Lsr^vill KO mice (n=6). (F) Expression of *Edn1, Ctgf* and *Cyr61* mRNA assessed by qRT–PCR in small intestines from control or HFD-fed Lsr^ctrl and Lsr^vill KO mice (n=5). (G) YAP immunofluorescence and quantification analysis in small intestines from Lsr^ctrl and Lsr^vill KO mice treated with vehicle control or

PA (n=6). (H) Expression of *Edn1, Ctgf* and *Cyr61* mRNA assessed by qRT–PCR in small intestines

from Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice treated with vehicle control or PA. (n=5). (I) YAP immunofluorescence

and quantification analysis in ISCs isolated from small intestine of Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice treated with

vehicle control or PA (n=6). (J) Expression of *Edn1, Ctgf* and *Cyr61* mRNA assessed by qRT–PCR in

ISCs isolated from small intestine of Lsr^{ctrl} and Lsr$^{vill\ KO}$ mice treated with vehicle control or PA. (n=3).

(K) Western blot and quantification analysis for YAP of ISCs lysates from small intestine of Lsr^{ctrl} and

Lsr$^{vill\ KO}$ mice treated with vehicle control or PA (n=3). Scale bars: A, E, G and I, 10 μm; C, 20 μm. *P*

*< 0.05, **P < 0.01, ***P < 0.001, #P < 0.05, ##P < 0.01, ###P < 0.001.*