Title: Antigen presentation by class I molecules on brain endothelium dynamically regulates T-cell mediated neuropathology in experimental cerebral malaria

[1,5]Fain, CE, [3,5]Zheng, J, [1]Jin, F, [1]Ayasoufi, K, Wu, Y, [5]Lilley, MT, [1,5]Dropik, AR, [1]Wolf, DM, Rodriguez, RC, [4,5]Aibaidula, A, [1,5]Tritz, ZP, [5]Bouchal, SM, [7]Pewe, LL, [7]Urban, SL, [1,5]Chen, [6]Y, Chang, S, [1]Hansen, MJ, [9]Kachergus, JM, [9]Shi, J, [9]Thompson, EA, [7]Harty, JT, [6]Parney, IF, [8]Sun, J, [1,3]Wu, LJ, [1,2,3]Johnson, AJ*

[1] Mayo Clinic Department of Immunology, Rochester, MN
[2] Mayo Clinic Department of Molecular Medicine, Rochester, MN
[3] Mayo Clinic Department of Neurology, Rochester, MN
[4] Mayo Clinic Department of Molecular Pharmacology and Experimental Therapeutics
[5] Mayo Clinic Graduate School of Biomedical Sciences, Rochester, MN
[6] Mayo Clinic Department of Neurosurgery, Rochester, MN
[7] University of Iowa Department of Pathology, Iowa City, IA
[8] University of Virginia Department of Medicine, Charlottesville, VA
[9] Mayo Clinic Department of Cancer Biology, Jacksonville, FL

Authors have no conflicts or competing interests to disclose.

Author contributions: CEF, JZ, FJ, KA, MTL, ARD, DMW, RCR, AA, and ZPT

assisted with experiments. CEF, JZ, FJ, YW, SMB, YC, JMK, and JS analyzed data.

CEF and AJJ designed experiments and interpreted results. CEF wrote the manuscript.

All authors provided feedback for editing, read manuscript, and agree with data

presented.

*Correspondence to Aaron J. Johnson, PhD
Professor and Consultant
Departments of Immunology, Neurology, and Molecular Medicine
Guggenheim 4-11C
Mayo Clinic
200 First St. SW
Rochester, MN 55905
Johnson.aaron2@mayo.edu
Tel: (507) 266-1973

Abstract

CD8 T cells are implicated in the neuropathology of human cerebral malaria. Using human data as well as the *Plasmodium berghei* ANKA (PbA) model of experimental cerebral malaria (ECM), we sought to define contributions of brain endothelial cell MHC class I antigen (Ag) presentation to CD8 T cells in this pathology. We crossed novel H-2K^b and H-2D^b LoxP mice to Cdh5-Cre mice to achieve targeted deletion of discrete class I molecules on endothelium. RNA transcriptomes of brains during PbA infection are differential between H-2K^b and H-2D^b expressing mice, highlighting differential regulation of neuropathogenesis. Distinct patterns of disease onset, CD8 T cell infiltration, targeted cell death, and regional BBB disruption and vascular leakage are observed. Strikingly, ablation of H-2K^b or H-2D^b from brain endothelial cells resulted in diminished interactions of T cells with brain vasculature and lessened infiltration of activated CD8 T cells, resulting in prevention of blood-brain barrier (BBB) disruption, CNS vascular leakage and death. Overall, endothelium-specific conditional knockout (cKO) mice were protected from development of ECM resulting in survival of the mice. These data show the first *in vivo* evidence that interactions of CD8 T cells and cognate peptide:MHC on brain endothelium regulate development of ECM neuropathology. Therefore, targeting MHC class I interactions therapeutically may hold potential for treatment of cases of severe malaria.

Introduction

Plasmodium falciparum has been classified as the deadliest parasite by the World Health Organization (WHO). This strain is also the most prevalent malaria parasite, accounting for majority of human infections[1]. Despite initiatives to limit mortality, it is estimated that greater than 600,000 people succumbed to malaria in 2020[2]. A severe complication *of Plasmodium falciparum* is severe encephalopathy, termed cerebral malaria (CM), which results in high

mortality despite treatment[3, 4, 5, 6, 7]. CM accounts for approximately 80% of all malaria-related

deaths, and patient admissions with CM are steadily increasing annually[2]. The majority of

cerebral malaria patients are diagnosed by the presence of asexual parasite in a blood smear,

deep coma, and absence of other infectious disease[3, 8]. Other hallmarks of disease can only be

assessed post-mortem such as disruption of BBB tight junction proteins, vascular permeability,

sequestration of parasitized red blood cell (prbc) and leukocytes, cerebral microhemorrhage and

endothelial cell activation[9, 10]. MRI has been gaining traction as a diagnostic tool in patients, as

vascular permeability can be assessed via T1 and T2 weighted MRI[10]. Recently,

immunohistochemical studies of autopsied human brain specimens have revealed that CD8 T

cells have been shown to be tightly associated with brain microvasculature and infiltrate brain

parenchyma, implying an important role for this immune cell type in neuropathogenesis[11].

Plasmodium berghei ANKA (PbA) infection is an established murine model that

recapitulates human disease and CNS pathology, referred to as ECM[12]. As in the human disease,

CD8 T cell responses are strongly implicated in the neuropathology. Importantly, CD8 T cells

are required to have functioning cytotoxic capacity to develop the fatal BBB disruption

associated with ECM[13, 14, 15, 16]. An IFN-γ dependent upregulation of adhesion molecules (ICAM-

1 and VCAM-1) and molecules associated with Ag-presentation (MHC Class I molecules D^b and

K^b and MHC Class II) has been observed on brain and meningeal endothelial cells day 6 of PbA

infection[17]. However, the contribution of specific cell types producing IFN-γ in this pathology

have not been interrogated[17]. Interest in these CD8 T cell-endothelial cell interactions, has

prompted a concerted effort to define the requirement of Ag-specific CD8 T cells in this

neuropathology. Due to the complex life-cycle of the parasite, defined by extensive antigenic

variation throughout each stage, this has proven a difficult task. The engagement of CD8 T cells

with class I molecules has been functionally assessed with bone marrow chimeras of MHC class

I-deficient hosts reconstituted with class I-sufficient hematopoietic presentation[17]. However, it is

known that CD8 T cells raised in these hosts have inherently impaired development and

survival[17, 18]. While this important work narrows in on the significance of non-hematopoietic Ag-

presentation in this pathology, the role of Ag-presentation specifically on brain vasculature has

not yet been addressed. Furthermore, the contribution of discrete class I molecules in ECM

pathology remain undefined.

In these studies, we sought to define the contribution of mouse brain microvascular

endothelial cell (MBMEC) Ag-presentation by discrete class I molecules to the pathogenesis and

lethality of ECM. We hypothesized that conditional ablation of Ag-presentation by H-2K^b and

H-2D^b class I molecules on MBMECs would result in controlled modulation of activation and

entry of parasite-specific CD8 T cells, attenuating pathology. To test this, we crossed

endothelium-specific Cdh5-Cre mice with our novel H-2K^b LoxP and H-2D^b LoxP mice to

create cKOs of discrete class I molecules. In these studies, we employed human and mouse

transcriptomic data to evaluate gene expression in brains in response to plasmodium expression.

We also employed *in vivo* 2-photon imaging to interrogate T cell interactions at the inflamed

vasculature of the brain when individual MHC class I molecules are present or conditionally

deleted on endothelium during PbA infection. Infiltration, activation, and localization of CD8 T

cells and their potential targets were assessed via spectral flow cytometry in these novel

transgenic mice. We demonstrate that interactions with specific class I molecules determined

CD8 T cell activation, infiltration of the brain and localization and onset of cell death and

neuropathology. However, knockout of H-2K^b and H-2D^b in endothelial cells prevented CD8 T

cell interactions at brain vasculature, CD8 T cell activation, BBB disruption, cell death and

neuropathology. These mice did not develop ECM, resulting in almost complete survival.

Together these data demonstrate that CD8 T cells engage H-2K^b or H-2D^b class I molecules on

MBMECs in order to coordinate activation, entry, and induction of neuropathology. These

studies indicate that modulation of brain microvascular endothelial cell class I restricted Ag-

presentation may hold therapeutic benefit in cases of cerebral malaria.

Results

Mouse and human brain microvascular endothelial cells increase antigen presentation

molecules capable of eliciting T cell responses after *Plasmodium* infection. C57BL/6 mice

were intraperitoneally (I.P.) injected with PbA prbcs, resulting in peripheral infection and

manifestation of ECM on days 6-8 (Fig. 1a). Brains were harvested and RNA was extracted from

whole brains of infected C57BL/6 mice and uninfected controls (Fig. 1a). Nanostring analysis

was performed on bulk RNA for analysis of differential gene expression between uninfected and

infected mice (Fig. 1b). Upregulated genes were associated with major immunological pathways,

including Ag-presentation, T cell responses, cytotoxicity, cell activation and apoptosis.

Downregulated genes were associated with markers of homeostasis or components of blood-

brain barrier (Fig. 1b). Gene expression data were analyzed using Ingenuity Pathway Analysis

(IPA). Causal network analysis identified MHC Class I family as the most significant master

regulator in ECM pathology (Supplementary Table 1). MHC Class I family activation led to

downstream activation of the majority of upregulated genes of interest identified in Fig. 1b (Fig.

1c). Canonical pathway analysis identified the most differentially enriched signaling pathways

during ECM, confirming Ag-presentation as a major pathway of interest (Fig. 1d). MBMECs

have been implicated to be involved in Ag-presentation during ECM (Fig. 1e). To interrogate

their role, we employed enzymatic digestion and high-dimensional spectral flow cytometry to

measure expression of Ag-presentation-associated molecules. MHC Class I median fluorescence

intensity is increased on MBMECs during ECM and the percent of expression on MBMECs is

significantly higher during ECM (Fig. 1f). MHC Class II is a well-known marker of antigen

presenting cells (APCs). MHC Class II expression on MBMECs greatly increased from minimal

to high during ECM, implicating their role in Ag-presentation (Fig. 1g). We next interrogated

gene expression signatures of human brain microvascular endothelial cells (HBMECs) from

human donor after exposure to *Plasmodium falciparum*. To this end, we utilized human

microarray dataset from Tripathi et al. (accession GSE9861)[19] in NCBI GEO database. In brief,

isolated HBMECs were co-cultured with *Plasmodium falciparum* prbcs or uninfected red blood

cells in culture and then analyzed (Fig. 1h). We first assessed the distribution of expression

values for each human sample to ensure suitability for comparison (Supplementary Fig. 1a).

After human sample validation we evaluated MHC Class I and MHC Class II expression levels.

MHC Class I expression was significantly increased in HBMECs exposed to *Plasmodium*

falciparum infected red blood cells (Fig. 1i). Human samples which were exposed to infection

were subdivided into two groups 1) high adhesion molecule expression and 2) low adhesion

molecule expression[19] (Supplementary Fig. 1b). MHC Class II expression was shown to be

significantly upregulated only in the high adhesion molecule samples compared to controls (Fig.

1j). Using GEO2R analysis platform, MHC Class I was identified as one of the most

significantly differentially expressed genes and was in the top 5 genes with significant

upregulation on HBMECs after exposure to *Plasmodium* infection (Supplementary Table 2).

When comparing mouse and human genes involved with Ag-presentation we saw that gene

activation and inhibition in these pathways were highly analogous (Supplementary Fig. 2). We

also saw comparable upregulation of homologous genes of interest downstream of MHC Class I

in human HBMECs, consistent with our observations in the mouse model (Fig. 1k). When

performing gene set enrichment analysis (GSEA) on HBMEC transcriptomics data we identified

the molecular signature for T cell activation involved in immune response to be highly enriched

in HBMECs exposed to infection. The enrichment of this gene set indicates that HBMECs

become an APC during *Plasmodium* infection, capable of eliciting this downstream T cell

response (Fig. 1l). Together, these results suggest that in both mouse and human, brain

endothelial cells are performing Ag-presentation to activate downstream T cell effector activities

following *Plasmodium* infection.

Endothelial cell specific K^b conditional knockout mice develop normal peripheral

***Plasmodium* infection and immune responses.** We crossed (class I deficient background, K^b-/-,

D^b-/-) K^b LoxP mice with (class I deficient background, K^b-/-, D^b-/-) Cdh5-Cre mice (Fig. 2a).

The resulting animals are defined as K^b Cre- littermate (with ubiquitous K^b expression), and

Cdh5-Cre K^b cKO (with endothelial-specific ablation of K^b), respectively. At baseline, Cdh5-Cre

K^b cKO mice had significantly reduced protein levels of K^b on brain endothelium, as well as

reduced percentage of brain endothelial cells that express K^b (Fig. 2b-d). T cell development was

intact in Cdh5-Cre K^b cKO and K^b Cre- littermates with no changes in thymic T cell subset

frequencies, mTEC and cTEC frequencies, or thymic weights between groups (Fig. 2e-f and

Supplementary Fig. 3a). Furthermore, no sex specific differences pertaining to T cell

development, overt changes in CD8 T cell repertoire, or change in splenic weights were observed

(Supplementary Fig. 3b-d). To determine the effect of K^b class I molecule ablation on MBMECs

during ECM, adult Cdh5-Cre K^b cKO (~3-month-old) mice and K^b Cre- littermates were I.P.

infected with 1×10^6 PbA prbcs and tissues were harvested at approximately 6-8 days post

infection (dpi) (Fig. 2g). During infection K^b remained significantly reduced on brain

endothelium (Fig. 2h-j). In the spleen, we observed no changes to granulocyte or APC subsets

during infection in Kb Cre- littermates or Cdh5-Cre K^b cKO mice (Fig. 2k-l). Importantly, there

were no differences in the ability to establish blood stage infection in these mice (Fig. 2m).

Populations of immune cells were assessed in peripheral tissues at baseline and during infection

and showed no changes in CD45+ immune cell numbers between Cre- littermates and Cdh5-Cre

K^b cKO spleen, peripheral blood or inguinal LN (Supplementary Fig. 3e). In these peripheral

tissues CD4 and CD8 T cell numbers and activation state remained unchanged between groups in

all peripheral organs, with the exception of a modest reduction in numbers of activated CD8s at

baseline in inguinal LNs in the Cdh5-Cre K^b cKO mice (Supplementary Fig. 3g). However,

during PbA infection, there were no differences in CD8 activation in inguinal LNs (Fig. 2o).

These results demonstrate that our Cdh5-Cre K^b cKO mouse model has efficient deletion of K^b

on vasculature, while maintaining a fully functional peripheral immune response to *Plasmodium*

infection. This allows us to proceed with interrogation of potentially brain-specific

immunomodulatory functions of K^b Ag-presentation by brain endothelial cells in ECM.

Endothelial cell specific D^b conditional knockout mice develop normal peripheral

***Plasmodium* infection and immune responses.**

We crossed (class I deficient background, K^b-/-, D^b-/-) D^b LoxP mice with (class I deficient

background, K^b-/-, D^b-/-) Cdh5-Cre mice (Fig. 3a). The resulting animals are defined as either D^b

Cre- littermate (with ubiquitous D^b expression), or Cdh5-Cre D^b cKO (with endothelial-specific

ablation of D^b). Endothelial cell expression of D^b at baseline in Cdh5-Cre D^b cKO mice was

comparable to Cre- littermate controls. Percentage of class I expression on endothelium was

reduced but did not reach statistical significance in uninfected Cdh5-Cre D^b cKO mice (Fig. 3b-

d). T cell development was intact with no changes in thymic T cell subset frequencies, mTEC

and cTEC frequencies, or in thymic weights between groups (Fig. 3e-f and Supplementary Fig.

4a). There were no differences between males and females in development of thymic T cell

subsets, and no differences between groups in CD8 T cell V beta usage in spleen, or in splenic

weights (Supplementary Fig. 4b-d). To determine the effect of D^b class I molecule ablation on

MBMECs during ECM, adult Cdh5-Cre D^b cKO (~3-month-old) mice were I.P. infected with

PbA and tissues were harvested at onset of severe neurological symptoms, approximately 6-8 dpi

(Fig. 3g). Importantly, during infection D^b expression was significantly reduced on brain

endothelium (Fig. 3h-j). Spleens of these animals had no changes to granulocyte or APC subsets

during infection (Fig. 3k-l) and mice established full blood stage infection in the periphery (Fig.

3m). Assessment of immune cell populations in peripheral tissues at baseline and during

infection revealed no changes in CD45+ immune cell number between D^b Cre- littermates and

Cdh5-Cre D^b cKO in spleen, peripheral blood or inguinal LN (Supplementary Fig. 4e-f). CD4

and CD8 T cell numbers and activation state remained unchanged between groups in all

peripheral organs at baseline (Supplementary Fig. 4f). However, during infection, there was

reduction in number of activated CD8 T cells in inguinal LN in the Cdh5-Cre D^b cKO animals

(Fig. 3n-p). These results demonstrate that our Cdh5-Cre D^b cKO mouse model has sufficient

deletion of D^b on vasculature during PbA infection and a fully functional peripheral immune

response. These results enabled us to proceed with interrogation of potentially brain-specific

immunomodulatory functions of D^b Ag-presentation by brain endothelial cells in ECM.

T cell dynamics at the barrier are modulated by the presence of distinct class I molecules on endothelium. To assess whether brain vasculature was inherently changed by loss of endothelial class I in cKO mice, we performed maximum intensity projections (MIPs) using MRI angiography to illuminate and assess the blood vessels. Cdh5-Cre K^b cKO and Cdh5-Cre D^b cKO were compared to their respective Cre- littermate controls. At baseline, we observed no differences in 3D volume of overall brain vasculature or voxel units (Supplementary Fig. 5a-b and 5c-d). We therefore proceeded to assess the effects of discrete MHC class I molecules on T cell dynamics in brain vasculature. To address this, we employed acute cranial window placement and *in vivo* two-photon imaging after IV injections of anti-Thy1.2-Alexa Fluor 488 to label T cells in vasculature and Texas Red-Dextran to label vessels. 6-7 days following PbA infection, mice underwent live imaging and T cell interactions at the brain vasculature were tracked over time (Fig. 4a). Cre- Littermates with K^b expression on brain vasculature presented with higher numbers of T cells per field of view (FOV) with slowed velocities, indicating interactions with K^b on endothelium (Fig. 4b and 4c). In contrast, Cdh5-Cre K^b cKO mice, which lack K^b on endothelium, had lower numbers of T cells per FOV, with higher velocity indicating less interactions with endothelium (Fig. 4b and 4c). Interestingly, there were no significant differences between the frequency that T cells contacted the vessels between genotypes. However, arrest on vessels was significantly increased when K^b was expressed on the brain vasculature (Fig. 4d-4e and Supplementary Videos 1-2). To investigate these T cell interactions further, we examined the activation state and adhesion molecule expression of MBMECs. During late-stage PbA infection, endothelial cells upregulate adhesion molecules responsible for initiating T cell extravasation[20]. P-selectin, which facilitates activated T cell rolling, did not differ in expression between Cdh5-Cre K^b cKO mice and Cre- littermates (Fig. 4f). ICAM-1 and

VCAM-1 expression, involved in arrest and adhesion, also showed no significant differences

during infection (Fig. 4g-h). When endothelial cells are activated, they upregulate molecules

involved in Ag-presentation[17]. Surprisingly, when K^b class I molecule expression was ablated,

MHC Class II molecules were upregulated on brain endothelium during infection (Fig. 4i).

T cell dynamics in Cre- D^b expressing and Cdh5-Cre D^b cKO mice were similar to those

of the previous model. 6-7 dpi with PbA T cell interactions with vasculature were tracked over

time during intravital imaging (Fig. 4j). Cre- Littermates with D^b present on vasculature showed

high numbers of T cells/FOV. In Cdh5-Cre D^b cKO mice, T cells were greatly reduced per FOV,

indicating fewer T cells slowed down to interact with endothelium when D^b molecule is ablated

on vasculature (Fig. 4k). Meanwhile, D^b expressing Cre- littermate T cells had slower velocities,

indicating more interactions with D^b on endothelium. When endothelial D^b is conditionally

deleted in Cdh5-Cre D^b cKO mice, T cells that interact with vessels had significantly increased

velocities, with reduced slowing and rolling on vasculature (Fig. 4l). Again, there were no

significant differences between the percentage of T cells contacting vessel walls between groups.

However, the frequency of T cell arrests was significantly increased when D^b was expressed by

brain vasculature (Fig. 4m and 4n). (Supplementary Videos 2-3). P-selectin was greatly

increased when D^b was present on brain endothelium during infection, indicating a higher

probability of activated T cell rolling (Fig. 4o). Interestingly, cells/FOV in the D^b expressing Cre-

littermates, had increased cells/FOV compared to K^b Cre- littermates (Fig. 4b and 4k). There

were no significant changes in ICAM-1 or VCAM-1 expression on endothelium when D^b was

conditionally deleted (Fig. 4p-q). As was seen with K^b cKO mice, deletion of D^b class I molecule

on endothelial cells resulted in higher expression of MHC Class II molecules on endothelium in

comparison to Cre- littermates (Fig.4i and 4r). Together, this data demonstrates that H-$2K^b$ and

H-2D^b class I molecules are critical for optimal T cell adhesion and arrest to brain endothelium

during ECM.

CD8 T cell activation and infiltration into brain parenchyma is differentially modulated by

discrete class I molecules during infection. We next addressed if the changes observed in T

cell dynamics at the brain vasculature during PbA infection corresponded with T cell infiltration

into brain parenchyma. First, we wanted to confirm that cervical lymph node (cLN) priming of

brain draining antigens was intact in our mouse models. To test this, we employed Theiler's

murine encephalomyelitis virus (TMEV) and TMEV-OVA. These neurotropic viruses are

intracranially administered and provide a highly reproduceable model of CNS antigen drainage

resulting in cLN priming, CD8 T cell infiltration, and viral clearance[21, 22, 23]. TMEV

immunodominant peptide VP2$_{121-130}$ is presented in the H-2D^b MHC class I molecule whereas

TMEV-OVA model antigen OVA$_{257-264}$ is presented by H-2K^b [23]. On day 7 post infection mice

were perfused and cLNs were analyzed via spectral flow cytometry (Supplementary Fig. 6a). In

mice infected with either TMEV or TMEV-OVA, tetramer staining was unchanged between Cre-

and Cre+ mice indicating that cLN priming in these mice is intact (Supplementary Fig. 6b-c). We

now evaluated immune cell infiltration into the brain in the ECM model, in which the

Plasmodium infection arises in the periphery prior to neuroinflammation in the brain (Fig. 5a).

At moribund, 6-8 d.p.i., mice were euthanized and perfused. Mouse brains were processed into

single cell suspensions and analyzed via spectral flow cytometry. Overall immune populations in

the brains during infection can be visualized in the UMAP projection (Fig. 5b). Data

encompassing CNS infiltration of immune cells were interrogated. Absolute counts of total

CD45+ immune cells in the brain remain unchanged when K^b class I molecules are present on

endothelium versus endothelial cell class I molecule conditional knockout mice (Supplementary

Fig. 6d). Numbers of CD4 T cells in the brain also remain unchanged regardless of K^b expression

by endothelial cells (Fig. 5e). However, when K^b expression is lost on endothelium, CD8 T cell

infiltration of the brain is greatly diminished (Fig. 5f). Since adhesion molecule expression on

endothelium was previously shown to be unchanged, we also examined CD8 T cells for

chemokine receptors involved in trafficking to the CNS and adhesion to the endothelium. Levels

of CXCR3, a receptor shown to be necessary for recruitment of T cells into the brain to establish

ECM[24, 25], was not impacted by ablation of K^b on vasculature (Supplementary Fig. 6f-g). CCR7,

a chemokine receptor known to be upregulated on T cells trafficking to the brain during

neuroinflammation, was also evaluated[26]. During ECM the olfactory region highly upregulates

CCL21, and blockade of CCR7 decreases T cell activation, recruitment and survival[27]. In Cre-

littermates with K^b, we saw no difference in CD8 T cell expression of CCR7 from mice lacking

K^b on vasculature (Supplementary Fig. 6h-i). We next evaluated effector status of CD4 and CD8

T cells in the brain. Interestingly, CD8 T cells that infiltrated the brain during infection produce

significantly less IFN-γ when K^b expression was ablated on vasculature (Fig. 5g). Surprisingly,

although brain infiltrating CD4 T cell numbers did not change, these lymphocytes increasingly

assumed a Th1 phenotype with increased expression of IFN-γ when K^b was ablated on

endothelium (Fig. 5h). In a parallel set of experiments, we investigated whether antigen

recognition plays a role in infiltration and localization of IFN-γ producing CD8 T cells. We

infected mice and intravenously (IV) injected Alexa Fluor-488 conjugated anti-Thy1.2 antibody

3 minutes before euthanasia on day 6 of infection to label cells that have recently infiltrated brain

parenchyma from the bloodstream. It is important to note that K^b expressing Cre- mice are not

yet moribund on day 6 of infection. We also evaluated surface expression of CD11a as a

surrogate marker for cognate antigen-experienced CD8 T cells as previously described [28]. At 6

d.p.i. we show that numbers of Ag-experienced CD8 T cells were not different between K^b

expressing Cre- and Cdh5-Cre K^b cKO mice in meninges or cervical LNs of PbA infected mice.

However, numbers of brain infiltrated Ag-specific CD8 T cells were greatly reduced when K^b

was lost on vasculature (Fig.5i). K^b expressing Cre- mice presented with a significant increase of

CD11a+ CD8 T cells only in the IV- group, indicating that the majority had infiltrated the brain

prior to IV labeling. In Cdh5-Cre K^b cKO mice lacking K^b on vessels, numbers of both IV- and

IV+ CD8 T cells were greatly reduced, demonstrating that recognition of peptides presented by

K^b is required for optimal CD8 T cell infiltration (Fig. 5j). It is known that IFN-γ producing CD8

T cells have enhanced Ag-specific contacts, cytolytic killing and kinetics through paracrine and

autocrine IFN-γ signaling[29]. We therefore employed IV labeling to determine the localization of

IFN-γ producing CD8 T cells 6 d.p.i with PbA. We noted a slight trend towards increased

frequency of IFN-γ producing CD8 T cells in the recently infiltrated IV+ population in both Cre-

and Cre+ animals. However, this difference did not reach significance (Fig5k).

We next assessed the immune profile of the brains of D^b expressing mice in ECM.

Tissues were harvested at moribund, days 6-8 of infection. We observed no difference in the

overall numbers of CD45+ immune cells in the brain between D^b expressing Cre- and Cdh5-Cre

D^b cKO mice (Supplementary Fig. 6e). Surprisingly, we observed no change in CD8 T cell

infiltration between the groups, and numbers of CD4 T cells in the brain were also unchanged

(Fig. 5l-o). Expression of trafficking and adhesion molecules on CD8 T cells were assessed and

levels of CXCR3 and CCR7 were not significantly different between D^b Cre- littermates and

Cdh5-Cre D^b cKO (Supplementary Fig. 6j-m). Although overall brain infiltration of CD8 T cells

was unchanged, these lymphocytes presented with reduced IFN-γ production when D^b was

ablated on vasculature (Fig. 5p). In contrast, no change in CD4 T cell IFN-γ production was

observed (Fig. 5q). When assessing the role of Ag-recognition in infiltration, we observed 6

d.p.i. that numbers of CD11a+ Ag-experienced CD8 T cells were not different between Cre- and

Cdh5-Cre D^b cKO mice in meninges or cervical LNs. However, numbers of brain infiltrating

CD11a+ Ag-experienced CD8 T cells were significantly reduced when D^b was lost on

vasculature (Fig. 5r). We observed that numbers of previously infiltrated, IV- Ag-experienced

CD8 T cells were much lower in D^b Cre- mice than in K^b Cre- mice, although the IV+

populations between Cre-s were more comparable on day 6 of infection (Fig. 5j and 5s). This

suggests that Ag-specific CD8 T cell infiltration may begin at a later timepoint and escalate more

quickly with D^b on endothelium in comparison to K^b. Alternatively, we may not see a difference

between IV- and IV+ if vascular leakage is present as this could result in increased antibody

staining in the parenchyma (Fig. 5s). The previously infiltrated, Ag-experienced CD8 T cells in

the D^b Cre- littermates were significantly increased only when compared to the IV+ population

in the D^b cKO mice (Fig. 5s).

It is important to highlight that total numbers of brain infiltrated CD8 T cells are similar

between Cre- littermates (with either K^b or D^b) at endpoint (Fig. 5f and 5o). However, Ag-

experienced CD11a+ CD8 T cell numbers were much higher in K^b expressing than in D^b

expressing Cre- controls (Fig. 5i and 5r). As per localization of IFN-γ producing CD8 T cells, D^b

expressing Cre- animals had significantly increased frequency of CD8 T cells producing IFN-γ

in the IV labeled population on day 6 of infection, suggesting these cells have increased effector

function when recently infiltrated from vasculature (Fig. 5t). Meanwhile, Cdh5-Cre D^b cKO

animals had no change in frequency of IFN-γ producing CD8 T cells between IV+ and IV-

groups (Fig. 5t).

To further interrogate infiltration, we compared rates of entry of Ag-experienced

(CD11a+) and Ag-inexperienced (CD11a-) CD8 T cells on day 6 of infection. We observed that

CD11a+ Ag-experienced CD8 T cells in mice with K^b or D^b expression had faster infiltration

than mice with conditional deletion of class I molecules on endothelial cells, though this did not

reach significance in D^b cKO mice (Supplementary Fig. 6n-o). CD8 T cells that were CD11a-

(Ag-inexperienced) in K^b expressing mice had low rates of infiltration and there was not a

significance difference in infiltration between K^b Cre- and Cdh5-Cre K^b cKO mice

(Supplementary Fig. 6p). These data suggest that K^b class I molecule is important for Ag-specific

CD8 T cell brain infiltration in ECM. Although no differences were seen in CD11a- Ag-

inexperienced CD8 T cell rates of entry in between Cre- and D^b cKO mice, D^b expressing Cre-

mice rate of CD11a- CD8 T cell entry was double that of K^b expressing Cre- animals

(Supplementary Fig. 6p and q). Importantly, the same number of activated IFN-y producing CD8

T cells end up in the brain of K^b and D^b Cre- littermates at endpoint despite numbers of CD11a+

Ag-experienced CD8 T cells in D^b expressing mice being much lower on day 6 (Fig. 5g and p).

This demonstrates that D^b class I molecule on endothelium is required to initiate IFN-y

production by CD8 T cells to the same extent as K^b, and that each class I molecule on

endothelium differentially modulates the timeline of entry of Ag-specific CD8 T cells.

It is known that most neuropathology and vascular leakage during CM and ECM localize

to three distinct regions of the brain; olfactory, hippocampus and brainstem[16]. We therefore

wanted to assess whether infiltrating CD8 T cells enter brain parenchyma in these regions. Using

2-photon and confocal microscopy, we imaged cell dynamics and performed

immunofluorescence staining on brain tissues at late stages of infection. At 6-7 dpi we observed

CD8 T cells extravasating via post-capillary venules and entering regions of brain parenchyma in

K^b expressing Cre- mice (Supplementary Video 5). In K^b expressing Cre- littermates, we

observed parenchymal CD8 T cells in olfactory, brainstem, and hippocampal regions. However,

when K^b is deleted on vasculature, CD8 T cell infiltration into these regions is greatly decreased

(Supplementary Fig. 6r). When imaging D^b expressing Cre- littermates, we saw CD8 T cell

infiltration into brain parenchyma (Supplementary Video 6). We also confirmed that when D^b

expression is ablated on vasculature, CD8 T cells are still present in brain parenchyma in these

three regions (Supplementary Fig. 6s). These data support the findings of our spectral flow

cytometry data and demonstrate that CD8 T cells are infiltrated into regions of the brain affected

by ECM, except specifically when K^b is deleted on brain vasculature.

Blood-brain barrier disruption is dependent on MHC Class I molecule expression by brain

endothelium. Next, we assessed the contribution of discrete class I molecules on modulating

BBB disruption during ECM. At 6-8 dpi with PbA, mice were administered gadolinium I.P. and

evaluated for vascular permeability in the brain using T1-weighted gadolinium enhanced MRI.

In Cre- mice with K^b class I molecule expression intact, we see significant enhancement in

regions of olfactory bulb, brainstem and hippocampal regions periventricular to the lateral

ventricles (Fig 6a). All areas of leakage were assessed utilizing 3D volumetric analysis with

Analyze Software as previously described[16]. Brain vascular integrity was retained, and

gadolinium leakage was not observed when K^b expression was ablated on vasculature (Fig. 6a-

c). As a secondary method of vascular leakage assessment, we IV injected ~66 kDa FITC-

albumin, one hour prior to tissue harvest. FITC-albumin mean fluorescence intensity was

quantified in our three brain regions of interest. Uninfected and infected wildtype C57BL/6 mice

were included as visual reference controls (Fig. 6d). Cre- littermates with K^b expression had

significant FITC-albumin leakage in regions of olfactory bulb and brainstem. In contrast, Cdh5-Cre K^b cKO animals lacking K^b on vasculature had negligible BBB disruption, indistinguishable from uninfected controls (Fig. 6d). FITC-albumin leakage is quantified in Fig. 6e. PbA infected D^b expressing Cre- Littermates presented with even more pronounced gadolinium-enhancement in regions of interest than K^b expressing Cre- mice as measured by small animal MRI (Fig. 6a-c and 6f-h). Cdh5-Cre D^b cKO mice however, presented with an intact blood brain barrier, with no notable pathology when D^b is lost on vasculature during infection (Fig. 6f-h). In a secondary analysis utilizing FITC-albumin leakage, we observed increased FITC-albumin mean fluorescence intensity in D^b expressing Cre- olfactory bulb and brainstem (Fig. 6i-j). In Cdh5-Cre D^b cKO mice, brain vascular integrity was retained, and no FITC-albumin leakage was observed (Fig. 6i-j). Tight junction proteins are known to be downregulated in regions of BBB disruption[16]. In human endothelial cells exposed to *Plasmodium falciparum* prbcs *ex vivo*, we observed negative enrichment scores consistent with downregulation of tight junction proteins (Supplementary Fig. 7a). We next performed Claudin-5 staining in our K^b and D^b models, interrogating the regions of vascular leakage in the brains, where T cells were also observed to infiltrate (Supplementary Fig. 8r-s). We saw that the vessels in regions of leakage and CD8 T cell infiltration in both K^b and D^b Cre- littermates had a more sparse and punctate staining for Claudin-5, indicating tight junction protein alteration (Supplementary Fig. 7b-c). However, K^b and D^b cKO animals lacking class I molecules on vasculature had intact Claudin-5 expression along vessels, despite the CD8 T cell infiltration seen in the Cdh5-Cre D^b cKO mice (Supplementary Fig. 7b-c). Overall, these results indicate that vascular leakage and disruption of BBB tight junctions occurs in mice via class I molecule restricted antigen presentation by brain endothelium during *Plasmodium* infection.

PbA infected parabionts sharing peripheral circulation demonstrate critical importance of endothelial D^b expression to onset of ECM. Previously, we had noted a decreased number of activated CD8 T cells in inguinal LNs of PbA infected Cdh5-Cre D^b cKO mice (Fig. 3p). In order to further rule out peripheral immune perturbations and fully demonstrate that neuropathology was dependent on CD8 T cell interactions with MBMECs, we performed parabiosis (Fig. 7a). Surgery was performed to conjoin D^b expressing Cre- littermate controls with Cdh5-Cre D^b cKO mice in order to exchange peripheral blood and circulating immune cells. Three weeks of healing and acclimation were observed before both mice were I.P. injected with PbA to induce ECM (Fig. 7b). Our analysis revealed that all peripheral immune activation was similar in the spleen and inguinal LNs of parabionts after infection (Fig.7c-d). In the brain, no differences in CD45+ immune cell frequencies were observed and CD8 and CD4 T cell numbers mirrored that of the previous mouse cohorts with no difference between groups (Fig. 7e). Also consistent with prior cohorts, the Cdh5-Cre D^b cKO parabionts had decreased IFN-γ expression by brain infiltrating CD8 T cells (Fig. 7e). We performed vascular permeability assays in our D^b expressing Cre- to Cdh5-Cre D^b cKO parabionts and determined that the vascular leakage phenotype was consistent with our previous data on D^b Cre- mice and had enhanced FITC-albumin leakage. In contrast, despite having circulating immune cells from D^b expressing Cre- mice, Cdh5-Cre D^b cKO mice resembled uninfected mice in that no vascular permeability was detectable, directly supporting our previous data (Fig. 7f). This observation was further supported by our results employing dual gadolinium enhanced T1-weighted MRI on the parabionts. We observed diffuse enhancement throughout the brain of the Cre- parabiont, indicating leakage. However, the conjoined Cdh5-Cre D^b cKO mice presented with no vascular

leakage as demonstrated by lack of gadolinium-enhancement with T1-weighted MRI and

retainment of FITC-albumin within vasculature (Fig. 7g). This data confirms that peripheral

immunity was not impeded and that changes in infiltration and activation status are truly barrier

and brain-specific and are impacted by class I restricted antigen presentation by MBMECs.

Differential Endothelial Class I expression alters the neuroinflammatory profile of the

brain and targeted cell death during ECM. To define downstream genes discretely regulated

by either K^b class I molecule or D^b class I molecule, we examined RNA transcriptomics data for

differentially regulated genes in PbA infected wildtype C57BL/6, K^b Cre-, Cdh5-Cre K^b cKO, D^b

Cre-, Cdh5-Cre D^b cKO, and uninfected controls of each group. We observed overlaps in gene

expression that were shared between wildtype C57BL/6 mice (with K^b and D^b) and K^b Cre- mice

(K^b only) (Fig. 8a). Gene programs regulated by K^b were genes primarily expressed in anti-

inflammatory A2 astrocytes, primed microglia, and cytokine and interferon signaling (Fig 8a and

Supplementary Table 3). Cdh5-Cre K^b cKO mice had very few differentially regulated genes,

similar to uninfected mice (Supplementary Figure 8a-c). We also noted distinctively differential

expression of genes that overlap between wildtype (with K^b and D^b) and D^b Cre- mice (D^b only)

(Fig. 8a). Gene programs regulated by D^b Cre- mice were indicative of a pro-inflammatory A1

astrocytic response, stage 2 damage-associated microglia, and microglial neurodegenerative

phenotype signatures[30, 31, 32, 33] .We also noted upregulation of genes associated with enhanced

Ag-presentation and cytotoxic T cell responses in the brains of these animals, indicating a strong

T cell response in these animals (Fig. 8a and Supplementary Table 3). Cdh5-Cre D^b cKO mice

had very few differentially regulated genes (Supplementary Figure 8d-f). Overall, the differential

gene expression observed between mice expressing distinct class I molecules further supports the

independent roles each class I contributes to this pathology.

We next wanted to determine what cells are being targeted by infiltrating CD8 T cells

during PbA infection. We infected mice with PbA and harvested tissue early on day 6 before all

mice were moribund. Brains were harvested to stain with a panel of antibodies to detect stages of

cell death. In K^b expressing Cre- littermates, we observed no significant increases in cell death of

MBMECs on day 6 of infection and no differences when K^b is deleted on vasculature (Fig. 8b).

When assessing neuronal cell death however, there were increased frequencies of cells with early

apoptotic markers, but complete neuronal cell death was not detectable by flow at this time.

There were no differences between K^b expressing Cre- littermates and Cdh5-Cre K^b cKO mice

(Fig. 8c). The low levels of endothelial and neuronal cell death may be attributed to the fact that

pathology is delayed in K^b mice in comparison to D^b mice. The K^b Cre- mice have average

clinical scores of ~1.5 on day 6 for ECM neurological signs, whereas D^b Cre- mice have average

clinical scores of >3 day on 6 (Fig. 8d and 8h). This delay in pathology may also translate to a

delay of neuronal and/or endothelial cell targeting as well. To examine this, we performed

another experiment in which all mice were allowed to progress to humane endpoint.

Transcriptomics analysis revealed that neurons do activate cell death pathways at this late

timepoint in K^b Cre- littermates (Fig. 8j). No endothelial cell death pathway activation was

noted. Cdh5-Cre K^b cKO mice had significantly improved clinical scores when K^b on

vasculature was deleted (Fig. 8d). In extended survival studies, we observed nearly complete

survival of Cdh5-Cre K^b cKO mice in comparison to Cre- littermate controls (Fig. 8e).

When assessing D^b mouse cohorts at day 6, we observed a striking increase in MBMEC

cell death in Cre- littermates that have D^b in comparison to Cdh5-Cre D^b cKO mice (Fig. 8f). A

significant increase in neuronal cell death was observed in D^b expressing Cre- littermates in

comparison to Cdh5-Cre D^b cKO mice (Fig. 8g). Similar to our observations in K^b, we observed

increased frequency of neurons with early apoptotic markers in the Cdh5-Cre D^b cKO mice, but

not late-stage neuronal death (Fig. 8g). These data were in agreement with transcriptomics data

showing activation of cell death pathways only in D^b Cre- mice (Fig. 8j). In an extended survival

study, mice with deletion of D^b on vasculature presented with striking improvement in clinical

scores and overall survival (Fig. 8h-i). Overall, neuronal cell death related gene expression was

increased in infected wildtype C57BL/6, K^b expressing Cre- mice and D^b expressing Cre- mice

(Fig. 8j). These studies demonstrate that class I molecule expression by endothelium is required

for the onset of neuronal cell death in the ECM model and that MBMECs undergo cell death late

in pathology following D^b MHC Class I-specific-CD8 interactions at vasculature.

Discussion

The role of Ag-presentation by brain endothelium and the effect on the CD8 T cell

effector response as a putative mechanism of CM have been highly debated in the cerebral

malaria field[3, 25, 34]. However, the role of individual class I molecule presentation by MBMECs to

brain infiltrating CD8 T cells has never been directly tested *in vivo*. In this study, we demonstrate

that targeted deletion of either K^b or D^b class I molecule on endothelium results in reduced T cell

interactions at brain vasculature and subsequently attenuated BBB disruption and

neuropathology. Comparisons to human data supports these findings, as well as other recent

studies which highlighted the importance of CD8 T cell interactions at the vasculature in the

development of CM and ECM[11, 17]. Overall, peripheral infection and all immune responses

develop normally in our mice, with the exception of mild decrease in CD8 T cell activation

marker CD44 at baseline in non-brain draining inguinal lymph nodes in Cdh5-Cre K^b cKO mice.

However, this difference resolves during infection. In Cdh5-Cre D^b cKO animals there was also

a slight decrease in the CD8 T cell activation marker CD44 in non-brain draining inguinal lymph

nodes during PbA infection. We addressed the potential effects these differences could pose in

ECM by performing parabiosis, which efficiently equalizes peripheral immune responses.

Endothelial cell-specific deletion of class I molecule was still protective, effectively ruling out

potential peripheral immune perturbations as having a role in protection from ECM.

CD8 T cells have been observed to interact with brain vasculature during

neuroinflammation[17, 26, 35, 36, 37]. However, few of these molecular interactions have been

successfully interrogated. The most well-characterized of these interactions is the CD8 T cell

slowing, rolling, and arrest on brain endothelium[38, 39, 40, 41, 42]. Using intravital microscopy on live

mice 6-7 dpi with PbA, we observe high numbers of T cells in both K^b and D^b expressing Cre-

littermates exhibiting slowing, rolling and arresting. In contrast, we observed a drop in these T

cell interactions with brain vasculature with the loss of either K^b or D^b class I expression by

endothelial cells. The novelty of these findings, however, lies in the fact that the adhesion

molecule expression is completely unchanged between Cre- and Cre+ animals in the K^b model.

This implies that the complete loss of T cell interactions we visualize by 2-photon are dependent

on the T cell-K^b class I interaction and not adhesion molecules. In D^b expressing Cre- mice, we

observe upregulation of P-selectin on brain endothelium. Surprisingly, despite their essential role

in T-cell rolling in vasculature, absence of E or P-selectin in mice have been experimentally

shown to be unnecessary for T-cell entry into the CNS, suggesting that T cell rolling is not

required for T-cell migration across inflamed BBB [39, 40, 41, 42]. However, most studies focusing on

diapedesis interrogated CD4 T cell interactions. ICAM-1 and VCAM-1 on endothelium was not

altered when class I molecules were ablated on vascular endothelial cells. Interestingly, when

either class I molecule was deleted on endothelium, expression of MHC Class II molecules was

found to be elevated. This may be a compensatory mechanism employed by endothelial cells in

order to establish a more robust CD4 T cell response when the CD8 T cell response is

impaired[43]. The ability of brain endothelium to shift the T cell effector response in the brain is in

agreement with several studies which propose that endothelial cell Ag-presentation may have

important immunomodulatory functions[35, 36, 37, 44].

Observing that deletion of both K^b and D^b class I molecule on endothelial cells resulted in

loss of T cell interactions with brain vasculature, we expected this may result in loss of CD8 T

cell infiltration. Interestingly, we observed differential CD8 T cell infiltration into brain

parenchyma. In Cdh5-Cre K^b cKO mice, we saw a reduction of CD8 T cells infiltrating the brain.

However, in Cdh5-Cre D^b cKO mice, we saw no changes in CD8 T cell infiltration, indicating

that overall CD8 T cell infiltration and D^b-restriction may be uncoupled in this model. Although

the deletion of D^b does not seem to affect overall infiltration of CD8 T cells, lack of D^b does alter

effector status. It is known that PbA strain of *Plasmodium* causes brain specific sequestration of

prbcs and activation of brain microvasculature[25]. However, work describing the PbA antigenic

D^b-restricted peptide SQLLNAKYL showed that when mice are infected with strains of

Plasmodium that do not cause adherence of prbcs to brain endothelium, CD8 T cells still entered

the brain but were not activated, resulting in absence of neurological symptoms in these

animals[45]. This supports our findings that when *Plasmodium* initiates endothelial cell Ag-

presentation, D^b is required for activation of CD8 T cells, but not infiltration during infection.

Furthermore, CD8 T cell TCR binding to cognate peptide presented by both H-2K^b and H-2D^b

on brain endothelium resulted in IFN-γ production by infiltrating CD8 T cells, which in turn,

caused differentially regulated cell death and mortality of the animal. This is supported by other

works showing that IFN-γ production induces a cascade to recruit more CD8 T cells to the

brain[43, 45, 46]. Overall, we demonstrated that endothelial K^b class I molecule interactions with

CD8 T cells is important for infiltration and activation, while endothelial D^b class I molecule

interactions activate CD8 T cells but does not influence diapedesis into parenchyma.

Previous work from our lab and others has hinted at the importance of antigen-specificity

of these infiltrating CD8 T cells in ECM neuropathology[16, 25, 45, 47]. However, definitive support

for this has been elusive and incomplete[17, 45 48]. Studies with transgenic parasites bearing model

epitopes are unable to recapitulate the complexity of the diverse CD8 T cell responses against the

wide array of native malaria Ags seen in mice and humans[49]. The epitopes that are known have

been shown to elicit variable CD8 T cell responses[49]. Although epitope discovery continues to

be an active area of research, there are currently only 83 known T cell epitopes for PbA out of

what is projected to be many thousands[35, 45, 50]. Importantly, a role for BBB damage-associated

molecular patterns (DAMPs) or parenchymal brain draining DAMPs in eliciting CD8 T cell

responses, have also not been ruled out[25, 34, 49, 50]. To date, our novel mouse models are the first

models capable of distinguishing between K^b and D^b class I restricted Ags specifically presented

by brain endothelial cells for interrogation of ECM pathogenesis. To assess the role of cognate

TCR-MHC interactions in CD8 T cell infiltration in this pathology, we used the well-established

surrogate marker strategy to track Ag-experienced CD8 T cells[28]. We saw no differences in Ag-

experienced CD8 T cells in the meninges or cervical LNs in either model. However, Ag-

experienced CD11a+ CD8 T cells were reduced specifically in brain parenchyma when lacking

K^b or D^b on vasculature[28, 51]. We also employed IV labeling of these infiltrating Ag-experienced

CD8 T cells 3 minutes prior to harvesting tissue. The caveat of this work is that IV labeling may

increase in parenchyma with increased vascular permeability, although this does not seem to be

the case in the K^b model. We saw that infiltration is much higher in K^b Cre- littermates than in

Cdh5-Cre K^b cKO mice at day 6, and that most CD8 T cells had previously infiltrated over time

prior to IV administration of labeling antibody. In the D^b model, however, we see comparable

amounts of IV labeled and IV negative cells in the brain indicating that a large influx of CD8 T

cells begins day 6 of pathology. These data suggest that K^b Ag-presentation by brain

endothelium tightly regulates activation and entry. However, D^b Ag-presentation by brain

endothelium does not employ as strict of regulation of CD8 T cell infiltration at the barrier. This

may mean that another predominant interaction, by a currently undefined cell type or chemokine

gradient, may be initiating this brain infiltration. This other interaction, however, does not

sufficiently activate the T cell as is seen when brain endothelium presents on D^b. Overall, these

data suggest that CD8 T cell activation and infiltration are influenced differentially in the D^b

model during ECM.

The chemokine receptor expression of activated T cells shows dynamic plasticity.

Several factors, including strength of antigenic signals, tissue-specific imprinting by priming

DCs, or cytokine milieu can influence the trafficking of T cells[52]. Both CXCR3 and CCR7 are

receptors on CD8 T cells shown to be necessary for recruitment into the brain to establish

ECM[24, 25, 26]. During ECM, olfactory bulb and brainstem are physically and functionally

damaged, with high prbc and leukocyte sequestration in these regions, which coincides with

vascular pathology and neuronal cell death[17, 27]. Specifically, the olfactory region has been

shown to highly upregulate CCL21, while brain endothelial cells highly express CXCL9 and

neurons highly express CXCL10[24, 27, 53]. Blocking their receptors CCR7 and CXCR3 results in

decreased CD8 T cell activation and recruitment to the CNS, respectively, as well as prolonging

survival during ECM[27, 54]. To support that T cells were sufficiently primed in the periphery and

imprinted for CNS-specific trafficking we examined chemokine receptor expression on CD8 T

cells trafficking to the brain. In the brain, frequency of CD8 T cells expressing CXCR3 and

CCR7, and MFIs of receptor expression were unchanged between Cre- and cKO animals in both

models; thereby reinforcing brain-specificity. The majority of data on CD8 T cell infiltration in

ECM focuses on the cortex, as this is the brain region most heavily infiltrated by CD8 T cells

during this pathology[55]. However, no prior reports have shown regional infiltration of areas

affected by vascular permeability. In this work we not only show regional CD8 infiltration to

areas with highest brain pathology, but that CD8 T cells are also present in regions without

vascular leakage, a topic previously debated[17].

CNS vascular leakage was prevalent in both K^b and D^b class I molecule expressing mice.

However, the presentation was distinct. In comparison to wildtype C57BL/6 animals, in which

both class I molecules are expressed, we saw less leakage in the midbrain of K^b expressing

animals, although there was still significant damage and leakage to olfactory and brainstem

regions. Alternatively, D^b expressing animals presented with worsened pathology, more closely

resembling that of the wildtype C57BL/6 mice which express both class I molecules.

Importantly, when either K^b or D^b were ablated specifically from vasculature, there was

complete rescue of vascular permeability. This finding is highly novel and strongly suggests that

antigens are being presented by either K^b or D^b class I molecules, specifically by brain

endothelial cells, to elicit distinct neuropathology associated with ECM. We also show that

Claudin-5 is altered in regions of permeability in Cre- animals, a known hallmark of vascular

pathology in this disease[16, 17]. However, animals lacking class I on vasculature preserved normal

Claudin-5 expression in the regions of interest.

We observed that K^b and D^b Ag-specific activation of recently infiltrated CD8 T cells

expressing IFN-γ correlates with differential vascular permeability in ECM. We therefore wanted

to interrogate whether K^b and D^b class I molecules produced differential activation of

neuroinflammatory pathways. We did observe distinct patterns of neuroinflammation, with

differential activation of astrocytes and changes in microglial and T cell activation states. This

may explain why D^b restricted Ag-presentation on endothelium, though resulting in later

infiltration of Ag-experienced CD8 T cells than K^b restriction on day 6, still resulted in stronger

T cell signaling, more cell death and worsened vascular leakage and survival at endpoint. Taken

together, these data indicate that discrete class I molecules on brain endothelium have

immunomodulatory capacity, capable of altering the neuroinflammatory profile of the brain

during ECM.

Knowing that K^b-restriction and D^b-restriction incite differential neuroinflammatory

profiles in the brain during PbA infection, we next sought to determine whether cellular targets

of the CD8 T cells were also distinct. We demonstrated that 6 dpi, mice with K^b class I

molecules displayed no markers of apoptosis or cell death in vascular endothelium. However,

neurons had early apoptotic markers and transcriptomic data supported that neuronal death is

induced at endpoint. When K^b was deleted from vasculature, endothelial cells were unchanged.

However, neurons no longer acquired the gene signature indicative of neuronal death. This data

suggests that CD8 T cell interactions with K^b-restricted Ag-presenting endothelial cells are not

required for targeted killing of MBMECs but are supportive of apoptotic neuronal cell death late

in pathology. The K^b-CD8 T cell interaction has been shown to modulate infiltration and

activation of CD8 T cells and may be involved in CNS reactivation or serve an imprinting role

which may instruct CD8 T cells to exact their effector functions in other ways or on other cell

types. When assessing D^b Cre- littermates, we observed a large percentage of cell death of

MBMECs which was ameliorated when D^b was deleted on vasculature. There was also an

increased percentage of cell death seen in neurons in D^b expressing Cre- animals which was

either resolved or delayed when D^b was lost on vasculature. This data demonstrates a

requirement for D^b-restricted Ag-presentation to CD8 T cells by the vasculature in order to enact

an effector response resulting in killing of both MBMECs and ultimately neuronal cell death.

The extent to which neuronal cell death is the result of direct killing by CD8 T cells, or through

alternative mechanisms has yet to be explored in these models. In previous work which detailed

CD8 T cell responses during ECM, they demonstrated that K^b and D^b are capable of presenting

similar peptides, though with very different binding affinities[49]. It is also known that receptor

affinities control effector responses and T cell polyfunctionality[45, 56, 57, 58, 59]. This is consistent

with our data showing that endothelial cells die when D^b presents, but not when K^b presents,

dependent upon the differential peptides they display as well as the strength of their interactions.

These results prove to be highly innovative as this is the first time that *in vivo* evidence has

directly shown the effects of K^b or D^b restricted Ag-presentation by brain endothelial cells during

ECM. Follow-up investigations should address direct mechanisms of cell death in more detail.

Alterations of cytokine milieu, the impact of vascular leakage and blood-derived factors on

neuroinflammation and CD8 T cell responsiveness still must be addressed.

In conclusion, our studies provide the first *in vivo* interrogation of the role of Ag-

presentation by individual class I molecules by MBMECs in developing ECM pathology. We

have dissected the contribution of both K^b and D^b to the onset of neuropathology and provide

evidence to support that K^b-restricted Ag engagement on endothelium plays a critical role in

CD8 T cell vascular adhesion, arrest, infiltration, and activation. These events in turn contribute

to blood-brain barrier tight junction downregulation, vascular leakage, and neuronal death.

Meanwhile, D^b-restricted Ag engagement on endothelium plays a role in vascular adhesion and

arrest but is inconsequential for CD8 T cell infiltration into brain parenchyma. However, the

interaction is important for CD8 T cell activation and IFN-γ secretion which is associated with

heightened CD8 T cell motility and toxicity, tight junction downregulation, and vascular

permeability[25, 29, 46, 60]. Strikingly, we show that only CD8 TCR recognition of D^b-restricted

antigens by brain vasculature results in killing of MBMECs and subsequent neuronal cell death.

We contend that these insights into the pathogenesis of cerebral malaria will provide rationale for

therapeutic targeting of these interactions. Anti-malarial drugs can serve to lessen parasite

burden, but show no improvement in neurological symptoms once there is brain involvement[17].

We therefore propose that patients with CM may benefit from endothelial-specific therapeutics

which serve to specifically blockade T cell-class I interaction in addition to standard-of-care anti-

malarial drugs.

Methods

Animals. Male and female C57BL/6J WT (stock no. 000664) and Cdh5-Cre mice (stock no.

033055) were purchased from Jackson Laboratory. H-2K^b LoxP and H-2D^b LoxP mice were

generated by our program as previously described, in the C57BL/6 background [61, 62]. H-2K^b

LoxP and H-2D^b LoxP mice were crossed with Cdh5-Cre mice to generate Cdh5-Cre K^b cKO

and Cdh5-Cre D^b cKO mice that express Cre under the *Cdh5* promoter and undergo endothelial-

specific targeted gene deletion of H-2K^b and H-2D^b molecules. Mice were then bred and

maintained under specific pathogen free conditions at the Mayo Clinic Animal Facility. In the

studies, 15–20-week-old mice were used for all experiments with appropriate age and sex-

matched controls. All experiments were conducted in accordance with guidelines from the

National Institutes of Health and under the approval of the Mayo Clinic Institutional Animal

Care and Use Committee.

Plasmodium **infection.** *Plasmodium berghei* ANKA (PbA) stocks were generously provided by

J. Harty at the University of Iowa and S. Hamilton Hart at the University of Minnesota Medical

School. PbA infection was induced in 15–20-week-old mice via intraperitoneal injection of ~10^6

parasitized red blood cells on day 0. Mice were monitored twice daily until onset of clinical signs

were observed (days 6-8 post infection), and then continuously. Blinded scores were obtained

daily, scoring signs of neurological impairment until endpoint. Mice were ranked for clinical

signs of ECM adapted from Hortle et al[63]: 0, no clinical symptoms; 1, reduced movement and

limp/flaccid tail; 2, rapid breathing and transient hunching posture; 3, hunched, ruffled fur and

loss strength in four limb hang test; 4, severe disease (ataxia, seizures, coma); and 5, death.

Moribund animals were scored 4.5 and humanely euthanized. Mice were classified as having

ECM if displaying symptoms between days 6-8 post-infection. Peripheral parasitemia was

assessed using thin blood smears and Geimsa staining as outlined on the Centers for Disease

Control website and as previously described[64].

Acute TMEV and TMEV-OVA Infection

The Daniel's strain of TMEV was prepared as previously described[62]. TMEV-XhoI-OVA8

(TMEV-OVA) was generated and prepared by our group as previously described[23]. At 10-12

weeks of age, mice were anesthetized with 1-2% isoflurane and infected intracranially (i.c.) with

2 x 10^6 PFU (10μL) of the Daniel's strain of TMEV, or 2 x 10^5 PFU of TMEV-OVA. Virus was

delivered to the right hemisphere of the brain via automatic 1 mL Hamilton syringe (Hamilton

Company, Reno, NV). Mice were euthanized for high-dimensional flow cytometry of tissues at

7-days post infection (dpi).

Magnetic resonance imaging (MRI). MRI capture and analysis was conducted as previously

reported using Bruker Avance II 7 Tesla or Bruker Avance 16.4 Tesla vertical bore small animal

MRI system (Bruker Biospin, Billerica, MA)[65]. Gadolinium (Multihance-Bracco) was

administered intraperitoneally (100µL of 100mg/kg) ~15 minutes before scans. 25 mm and 34

mm volumetric coils were utilized. Animals were anesthetized with continuous isoflurane

administered through nose cone. Respiration and heart rate were monitored with an MRI

compatible vitals monitoring system (model 1030; SA Instruments, Stony Brook, NY).

After axial and sagittal tripilot positioning scans, isotropic 3D T1-weighted scans were employed

to detect gadolinium, with an approximate scan time of 30 minutes. Maximum intensity

projection scans were performed to detect gadolinium, with an approximate scan time of 10

minutes. Scoring was performed by two independent observers, blinded to genotype with <10%

intra-observer variation found. Analyze12.0 software (Biomedical Imaging Resource; Mayo

Clinic, Rochester, MN) was employed to analyze T1 data and to generate 3D maps of

gadolinium enhancement. Maximum intensity projection data was analyzed using Fiji Is Just

ImageJ (FIJI) 2.3.0; Java 1.8.0_172.

Immunofluorescence. Brains used for immunofluorescence staining were either fresh frozen or

paraformaldehyde fixed following mouse euthanasia and transcardial perfusion with ice-cold

PBS. Brains harvested after 2-photon imaging and FITC-albumin injection were harvested

without perfusion and paraformaldehyde fixed. All brain tissues were embedded in Tissue-Tek

OCT Compound, frozen on dry ice and sliced on cryostat with the exception of FITC-albumin

injected brains which were embedded in 4% agarose and sliced with vibratome. All brain slices

were mounted on SuperFrost Plus microscope slides and stored at 4 degrees.

Single-cell suspension. Mice were euthanized via isoflurane per IACUC guidelines and

transcardially perfused with ice-cold PBS. Brains and meninges were harvested and processed as

previously described[66, 67], via dounce homogenization and 30% Percoll gradient or enzymatically

digested using mouse tumor dissociation kit and gentleMACS™ Octo Dissociator (Miltenyi

Biotec). Spleens and lymph nodes were either digested with Collagenase IV/DNAse or not

digested before mechanical disruption. Lymph nodes were dissociated using frosted glass slides

and spleens were dissociated through 70 µM filter. Thymus samples were processed using

mechanical disruption through 70 µM filter. Blood was collected (100µl) into 1 mL heparin

(100U/mL). Spleens underwent ACK RBC lysis for 3 minutes and blood samples for 1minute.

All samples were filtered using 70 µM filters before counting, staining and running on flow

cytometer.

Flow cytometry. After resuspension in PBS, cells were either first stained with either Zombie

UV or Zombie NIR fixable viability dyes (BioLegend) (1:1000) for 20 minutes at room

temperature in the dark or Ghost Red 780 Viability dye (Tonbo) was added to surface stain

master mix. Samples underwent Fc-receptor block and staining for 30 minutes at 4 degrees with

surface staining antibodies. Panels requiring intracellular staining were then fixed and

permeabilized using Cyto-Fast™ Fix/Perm Buffer Set (BioLegend), and then stained with

intracellular antibodies for 30 minutes at room temperature in the dark before washing. All flow

panel staining was performed with this method with the exception of Annexin staining, which

required staining in Annexin V Binding Buffer (Invitrogen) for 15 minutes at room temperature

prior to fixation and permeabilization and intracellular staining. The antibodies that are specific

for surface markers included CD103 Brilliant Violet 421 (121422; BioLegend), NK1.1 Pac Blue

(108722; BioLegend), B220 Pac Blue (103227; BioLegend), CD62L Brilliant Violet 480

(746726; BD Biosciences), CD11b VioGreen (130-113-239; Miltenyi Biotec), CD44 Brilliant

Violet 570 (103037; BioLegend), CD80 Brilliant Violet 605 (104729; BioLegend), Ly6C

Brilliant Violet 650 (128049; BioLegend), H-2K^b/H-2D^b Brilliant Violet 750 (746869; BD

Biosciences), CCR7 Brilliant Violet 785 (120127; BioLegend), I-A/I-E BB515 (565254; BD

Biosciences), CD8a Spark Blue 550 (100780; BioLegend), CD45 PerCP (103130; BioLegend),

Ly6G PerCP-eFluor 710 (46-9668-82; Invitrogen), OX40L PE-Dazzle 594 (108816;

BioLegend), CD40 PE-Cy5 (124618; BioLegend), CD31 PE-Cy7 (25-0311-82; Invitrogen), CD3

Spark NIR 685 (100262; BioLegend), CD11c Alexa Fluor 700 (117320; BioLegend), CD4 APC-

Fire 750 (100568; BioLegend), CD25 Brilliant Violet 421 (102034; BioLegend), TCRβ APC

(20-5961-U100; Tonbo biosciences), c-kit Brilliant Violet 786 (564012; BD Biosciences), CD69

Brilliant Violet 650 (740460; BD Biosciences), Ly51 FITC (108305; BioLegend), EpCAM PE-

Cy7 (118216; BioLegend), CD8a Pac Blue (563898; BD Biosciences), CD4 APC (100516;

BioLegend), CD11a PE (141006; BioLegend), CD16/CD32 (553141; BD Pharmingen), Thy1.2

Alexa Fluor 488 (105316; BioLegend), I-A/I-E Pac Blue (107620; BioLegend), CD8a Brilliant

Violet 510 (100751; BioLegend), CD44 Brilliant Violet 650 (740455; BD Biosciences), Ly6G

Brilliant Violet 711 (46-9668-82; Invitrogen), CD4 Brilliant Violet 786 (563727; BD

Biosciences), Ly6C PerCP (128027; BioLegend), CD45 PE-CE594 (562420; BD Horizon),

CD11b PE-Cy5 (55-0112-U100; Tonbo biosciences), TCRβ PE-Cy7 (60-5961-U100; Tonbo

biosciences), B220 Spark NIR 685 (103268; BioLegend), CD62L APC-Fire 750 (104450;

BioLegend), CD11c Brilliant Violet 605 (117333; BioLegend), Annexin V(25-8103-72;

Invitrogen), B220 APC-Fire 810 (103277; BioLegend), CD26 Brilliant Violet 605 (745125; BD

Biosciences), CD80 Brilliant Violet 711 (104743; BioLegend), I-A/I-E Brilliant Violet 786

(743875; BD Biosciences), CD8a Brilliant Violet 421 (563898; BD Biosciences), CD31 APC-

Fire 750 (102528; BioLegend), CD11a PE-Cy7 (101122; BioLegend), CD31 PE-Dazzle

(102526; BioLegend), CXCR3 Brilliant Violet 421 (126521; BioLegend), CD106 Alexa Fluor

647 (105712; BioLegend), CD62P PE (148306; BioLegend), CD54 Brilliant Violet 711 (116143;

BioLegend), CD8 Brilliant Violet 570 (100740; BioLegend), Zombie NIR viability dye (Cat.

#423105; BioLegend), Zombie UV (423107; BioLegend). To assess cytokine production cells

were stained with intracellular markers after fixation and permeabilization: NeuN APC (100920;

Novus), IFNγ Brilliant Violet 785 (563773; BD Biosciences), IFNγ APC (20-7311-U100; Tonbo

biosciences) IL-12 PE (505204; BioLegend), IFNγ PE (50-7311-U100; Tonbo Biosciences). In

noted experiments, intravenous injection of 2ug of Thy1.2 Alexa Fluor 488 were administered 3

min. before collecting tissues to label T cells in vasculature. All samples were acquired on Cytek

Aurora spectral flow cytometer (Cytek, Fremont, CA). All analysis was performed using FlowJo

software version 10.8.1(FlowJo LLC, Ashland, OR).

***In vivo* two-photon microscopy.** Mice were intravenously injected with 2 µg of Thy1.2 Alexa

Fluor 488 just prior to cranial window surgery, approximately 30 min prior to imaging. Cranial

windows were implanted as described [68] [69]. In brief, under isoflurane anesthesia (3% induction,

1.5–2% maintenance), a circular craniotomy (<5 mm diameter) was made over somatosensory

cortex with the center at about -2.5 posterior and +2 lateral to bregma. A circular glass coverslip

(5 mm diameter, Warner) was secured over the craniotomy using dental cement (Tetric

EvoFlow). A four-point headbar (NeuroTar) was secured over the window using dental cement.

Mice were injected intravenously with intravascular dye 70,000 MW Dextran-Texas Red

(Invitrogen, D1830) 5 minutes before imaging. Alexa Fluor 488 (T cells) and Dextran-Texas

Red (vessels) were excited by a tunable Ti:Sapphire Mai Tai DeepSee laser (Spectra Physics,

CA, USA) at 940 nm and the emission was collected at 525/50 and 630/75 nm, respectively (U-

MF2, Olympus, Japan). An Olympus LUMPLFLN 40x water-immersion objective was used to

acquire image stacks. A 30-μm stack with 2 μm z-resolution and 512 × 512 pixels were acquired

every minute for a period of 10-30 min. Images were analyzed by Fiji ImageJ [70]. To analyze T

cell velocity, each 30-μm stack was first aligned along the z axis using the StackReg plugin

(Rigid Body) and Z-project into one image. Then images of every minute were stacked and

aligned using StackReg plugin (translation). The T cell velocity was measured using the Manual

Tracking plugin tracing for 10 min per field. The arrest coefficient is the percentage of each track

that a cell is immobile (instantaneous velocity <2 μm/min) [71].

Parabiosis: Procedure was performed as previously described. In brief, two female age-matched

mice of similar weight were selected. Mice were either littermates or co-housed for two weeks

prior to surgery. Anesthesia was induced and maintained with Ketamine/Xylazine (100/10

mg/kg; maintenance dose 30/3 mg/kg). Opthalmic ointment was placed on eyes. Mice are tested

for reflex responses to ensure anesthetic depth. For analgesia during surgery, we administered

Carprofen and Buprenorphine-SR subcutaneously (5 mg/kg and 1 mg/kg). Animals were shaved

on the flank and prepped for surgery with Betadine swabs and sterile drapes. Using a sharp

scissor, superficial epidermal incision was continuously made along the flank starting at 0.5 cm

rostral of olecranon to 0.5 cm superior to patella. Skin absorbable 3-0 silk suture were used to

join elbow and knee joints (Ethicon Inc, Somerville, NJ). Simple continuous suture was

performed with absorbable 4-0 vicryl suture (Ethicon Inc, Somerville, NJ) for closer of incision.

Post-operatively we administer 1 ml of 0.9% NaCl fluids subcutaneous, analgesics Carprofen

intraperitoneally and Buprenorphine SR subcutaneously. Mice were placed on heating pad and

observed for a minimum of 3 hours after surgery. Prophylactically, mice were treated with

antibiotic Enrofloxacin (50 mg/kg) 10 days post-operatively. Parabiont pairs underwent a

minimum of 3 weeks of healing and acclimation time before experimental use. Procedure was

performed according to the IACUC guidelines at Mayo Clinic.

Nanostring. Brain dissections were subjected to RNA extraction (Qiagen RNAeasy Mini Kit Cat.

No. 74104), followed by nanodrop concentration and purity analysis. RNA samples were applied

to NanoString nCounter (NanoString Technologies, Seattle, WA) digital RNA quantification,

using the predesigned mouse Glial Profiling Panel codeset. The Glial Profiling Panel reagents were

provided by Nanostring. Reporter CodeSet Lot#RC8131X1. Capture Codeset Lot#CP8131X1.

Nanostring analysis was performed by the Thompson Lab at the Mayo Clinic Comprehensive

Cancer Center at Mayo Clinic Florida. Briefly, 100 ng of total RNA was hybridized to capture-

reporter probe codesets and immobilized on NanoString cartridges. Excess RNA and probes were

removed, and digital barcodes were counted. NanoString nCounter analysis software (v4.0) was

used for counting, quality control, and normalization. Transcript counts were normalized

according to NanoString's recommendations. Normalization parameters were applied with

normalization cutoff of 10. All samples were quality checked for quality and concentration. 13

housekeeping genes, 6 positive control genes, and 8 negative control genes were utilized for

normalization.

869   Human brain microvascular endothelial cells

Human brain endothelial cell microarray data available through public repository at Gene

Expression Omnibus (GEO), accession number GSE9861. Outlined in the publication by Tripathi

et al, HBMECs were isolated from human brain specimens and characterized as described

previously[19, 72]. In brief, HBMECs (passage < 6) from single adult donor were used for the biologic

replicate experiments. For microarray experiments, HBMECs were grown in collagen-coated 6-

well plates until cells achieved about 80% to 90% confluence. The medium was then changed to

RPMI 1640 with 5% fetal bovine serum for 24 hours before coincubation with Pf-IRBCs, at which

time HBMECs are confluent. Endothelial cells mycoplasma infection negative (Vector

Laboratories).

Gene Expression Analyses. Human microarray data with accession number GSE9861[19] was

utilized via Gene Expression Omnibus (GEO), a public functional genomics data repository

supporting MIAME-compliant data submissions. Genes from human and mouse datasets were

identified to be differentially expressed from each in comparison to a control group using either

Ingenuity Pathway Analysis software (Qiagen), or Geo2R[73]. Gene set enrichment analysis

(GSEA) was also performed on human data[74].

Statistics. When comparing results from two groups, an unpaired Student's t-test was used.

When comparing results from two groups of paired mice, a paired Student's t-test was used.

When comparing results from three groups, a one-way analysis of variance (ANOVA) using

Tukey's post-hoc multiple comparisons test was used. A two-way ANOVA when comparing

statistically significant difference between the means of three or more independent groups that

have been split on two variables, a two-way ANOVA using Šidák's multiple comparisons test

was used. All statistical analyses were performed using Prism 6.0 (GraphPad Software) or Excel.

Data are presented as the mean ± SD. and All p-values are shown or ns for non-significant p-

value ($p > 0.05$).

Acknowledgments: We would like to thank Dr. Vanda Lennon for her feedback, Dr. Emma

Goddery and Dr. Roman Khadka for discussions and Ryan Meloche for assisting with logistics

of dual MRI. Some graphics were prepared on Biorender.com. Tetramers were provided by the

NIH tetramer facility. **Funding sources:** R01NS103212 (AJJ), RF1NS122174 (AJJ), and

F31NS116924 (CEF), R01AI167847 (JTH), Nanostring-Glial Panel Grant (CEF).

Financial disclosures: We have no financial interests to disclose.

     Figure Legends

Fig 1. Mouse and human brain microvascular endothelial cells become antigen presenting
cells during cerebral malaria a) Flowchart showing steps of peripheral malaria infection,
development of experimental cerebral malaria and RNA extraction for Nanostring analysis. b)
Volcano plot of differentially expressed genes in whole brains of uninfected C57BL/6 mice
versus PbA infected mice on day 6 of infection. Significantly upregulated genes are indicated on
upper right (purple dots). Significantly downregulated genes are indicated on upper left
(turquoise dots). Horizontal dotted line indicates cutoff of adjusted p-values (-Log10 of adjusted
p-value 1.3 corresponding to a threshold of p-value of 0.05). Vertical dotted line indicates cutoff
of log2 Fold change (-1.5 to 1.5). c) Ingenuity Pathway Analysis (IPA) was performed, and
Causal network analysis identified MHC Class I family as the most significant network master
regulator in experimental cerebral malaria pathology which led to the differential expression of
the genes of interest. Depicted in c) a schematic of the homologous master regulator pathway and
corresponding downstream genes. Coloring of the molecules corresponds to calculated z-scores
(algorithm predicting activation or inhibition). Red indicates increased gene activation, orange
indicates predicted activation of pathways, green indicates decreased gene activation, blue
indicates predicted inhibition of pathways, white indicates a z-score of zero with equal evidence
for activation and inhibition, grey indicates that z-score could not be calculated. d) Canonical
pathway analysis identified significantly differential signaling pathways. P-value of overlap was
adjusted using the Benjamini-Hochberg correction method. Bars are arranged in order of
significance score with threshold (red line) set at p-value 0.05 (corresponding to a negative log
value of 1.3). Coloring of bars corresponds to calculated z-scores (algorithm predicting
activation or inhibition). Orange bars indicate activated pathways, blue indicates inhibited
pathways, white indicates a z-score of zero with equal evidence for activation and inhibition,
grey indicates that z-score could not be calculated. e) Schematic depicting C57BL/6 and naming
convention of mouse brain microvascular endothelial cells (MBMECs). High-dimensional flow
cytometry was performed on mouse brains on day 6 of PbA infection and uninfected age and sex
matched controls. f) Representative histogram of MHC Class I expression on MBMECs and
quantified frequency of MHC Class I high endothelial cells. g) Representative histogram of
MHC Class II expression on MBMECs and quantified frequency. h) Schematic depicting human
subjects and naming convention of human brain microvascular endothelial cells (HBMECs). i-j)
Box plots depicting Log$_2$ expression data from human endothelial cells in published microarray
(GEO Accession: GSE9861). i) MHC Class I Log$_2$ expression in control HBMECs vs.
Plasmodium exposed HBMECs. j) MHC Class I Log$_2$ expression in control HBMECs vs.
Plasmodium exposed HBMECs selected as highly activated (with high expression of adhesion
molecules). k) Log$_2$ Fold change of genes of interest which lie downstream of the MHC class I
signaling pathway in HBMECs during *Plasmodium* falciparum infection. GSEA was performed
on HBMEC transcriptomic data. l) Enrichment plot showing enrichment of molecular signature
for T cell activation involved in immune response pathway. The y-axis represents enrichment
score (ES) and on the x-axis are genes (vertical black lines) represented in gene sets. The green
line connects points of ES and genes. ES is the maximum deviation from zero as calculated for
each gene going down the ranked list and represents the degree of over-representation of a gene
set at the top or the bottom of the ranked gene list. The colored band at the bottom represents the
degree of correlation of genes with the phenotype (red for positive and blue for negative
correlation). Significance threshold set at nominal p-value < 0.05. Gene sets for T cell activation

involved in immune response were significant only in HBMECs during infection. Data were
analyzed with unpaired t test (f, g, i, and j). Box plots present the mean $\pm$ SD. All p-values are
shown or ns for non-significant p-value *(p > 0.05)*.
963
964
965
Fig 2. Normal peripheral infection and immunity in mice with K^b class I molecule
conditionally ablated on vasculature a) Schematic depicted Cre-LoxP strategy. Cdh5-Cre mice
were crossed to transgenic H-2K^b LoxP mice. b) Representative MFI plot comparing C57BL/6,
Cdh5-Cre K^b cKO, and Cre-Littermate expression of MHC I on brain endothelium in naïve mice.
c) Quantification of MHC I MFIs on brain endothelial cells using arbitrary units (A.U.) in naïve
mice. d) Percentage of brain endothelial cells expressing high levels of MHC Class I molecules
in naïve mice. e) Representative flow plot of C57BL/6, Cdh5-Cre K^b cKO, and Cre-Littermate T
cell subsets in thymus. f) Percentage of thymic T cell subsets. g) Schematic representing timeline
of infection for development of ECM. h) Representative MFI plot comparing C57BL/6, Cdh5-
Cre K^b cKO, and Cre-Littermate expression of MHC I on brain endothelium in PbA infected
mice. i) Quantification of MHC I MFIs on brain endothelial cells using arbitrary units (A.U.) in
PbA infected mice. j) Percentage of brain endothelial cells expressing high levels of MHC Class
I molecules in PbA infected mice. k) Percentage of granulocytes in spleens of infected mice. l)
Percentage of antigen-presenting cells in spleens of infected mice. m) Percent parasitemia
quantified by imaging Giemsa-stained thin blood smears in PbA infected mice. n) Representative
UMAPs depicting peripheral organs and immune cell populations and respective heats maps with
CD44 activation marker expression. o) The scatter plots show quantified absolute cell counts of
CD8 T cells, CD4 T cells and CD44hi CD8 T cells and CD4 T cells in the respective organs
depicted in n). Data were analyzed with unpaired t test or 2way ANOVA with multiple
comparisons. Scatterplots and bar plots present the mean $\pm$ SD. All p-values are shown or ns for
non-significant p-value *(p > 0.05)*.
987
Fig 3. Normal peripheral infection and immunity in mice with D^b class I molecule
conditionally ablated on vasculature a) Schematic depicted Cre-LoxP strategy. Cdh5-Cre mice
were crossed to transgenic H-2D^b LoxP mice. b) Representative MFI plot comparing C57BL/6,
Cdh5-Cre D^b cKO, and Cre-Littermate expression of MHC I on brain endothelium in naïve mice.
c) Quantification of MHC I MFIs on brain endothelial cells using arbitrary units (A.U.) in naïve
mice. d) Percentage of brain endothelial cells expressing high levels of MHC Class I molecules
in naïve mice. e) Representative flow plot of C57BL/6, Cdh5-Cre D^b cKO, and Cre-Littermate T
cell subsets in thymus. f) Percentage of thymic T cell subsets. g) Schematic representing timeline
of infection for development of ECM. h) Representative MFI plot comparing C57BL/6, Cdh5-
Cre D^b cKO, and Cre-Littermate expression of MHC I on brain endothelium in PbA infected
mice. i) Quantification of MHC I MFIs on brain endothelial cells using arbitrary units (A.U.) in
PbA infected mice. j) Percentage of brain endothelial cells expressing high levels of MHC Class
I molecules in PbA infected mice. k) Percentage of granulocytes in spleens. l) Percentage of
antigen-presenting cells in spleens. m) Percent parasitemia quantified by imaging Giemsa-
stained thin blood smears in PbA infected red blood cells. n) Scatter plots show quantified
absolute cell counts of CD8 T cells, CD4 T cells and CD44hi CD8 T cells and CD4 T cells in
spleen. o) Scatter plots show quantified absolute cell counts of CD8 T cells, CD4 T cells and
CD44hi CD8 T cells and CD4 T cells in peripheral blood. p) Scatter plots show quantified
absolute cell counts of CD8 T cells, CD4 T cells and CD44hi CD8 T cells and CD4 T cells in

inguinal LN. Data were analyzed with unpaired t test or 2way ANOVA with multiple
comparisons. Scatterplots and bar plots present the mean $\pm$ SD. All p-values are shown or ns for
non-significant p-value *(p > 0.05)*.
1010
Fig 4. T cell dynamics at brain vasculature are dependent on MHC class I expression
during infection a) Representative images of T cells in brain vasculature with blood vessels
labeled with I.V. Dextran-Texas ~5 min. prior to imaging and T cells I.V. labeled with anti-
Thy1.2-Alexa Fluor 488 ~30 min. Inset is representative of T cell tracking over time through
acute cranial window on days 6-7 post PbA infection. T cells (green) were visualized interacting
with blood vessels (red) over periods of 10-30 min. Upper row represents K^b Cre- littermates and
bottom row represents Cdh5-cre K^b cKO mice. Scale bar, 10 µM. b) Scatter plot represents
numbers of T cells visualized over 10 min time periods/FOV. c) T cell velocity was measured
using the Manual Tracking plugin in FIJI. Data are representative of µm/min over periods of 10
min per field and per cell. d) Percentage of T cells contacting vessels was calculated by obtaining
the percent of all visualized T cells that contacted vasculature over many fields of view. Each
data point is representative of the overall average percentage/mouse. e) The arrest coefficient
is the percentage of each cell track that a cell is immobile (instantaneous velocity <2 µm/min).
Each dot represents one cell and its percentage of contacts over time. (n = 5-7 mice per group).
See corresponding S1-S5 movies. High-dimensional flow cytometry was performed to assess
levels of protein expression on brain endothelium day 6 of PbA infection. f) Scatter plot
represents MFI values of P-selectin on brain endothelial cells. g) MFI of ICAM-1on brain
endothelial cells h) MFI of VCAM-1 on brain endothelial cells. i) Percentage and MFI
quantification of MHC Class II molecules on brain endothelial cells with representative
histogram. j) Representative images of T cells in brain vasculature labeled as described above.
Inset is representative of T cell tracking over time through acute cranial window on days 6-7 post
PbA infection. T cells (green) were visualized interacting with blood vessels (red) over periods
of 10-30 min. Upper row represents D^b Cre- littermates and bottom row represents Cdh5-cre D^b
cKO mice. Scale bar, 10 µM. k) Scatter plot represents numbers of T cells visualized over 10
min time periods/FOV. l) T cell velocity was measured using the Manual Tracking plugin in
FIJI. Data are representative of µm/min over periods of 10 min per field and per cell. m)
Percentage of T cells contacting vessels was calculated by obtaining the percent of all visualized
T cells that contacted vasculature over many fields of view. Each data point is representative of
the overall average percentage/mouse. n) The arrest coefficient is the percentage of each cell
track that a cell is immobile (instantaneous velocity <2 µm/min). Each dot represents one cell
and its percentage of contacts over time. (n = 5 mice per group). See corresponding S1-S5
movies. High-dimensional flow cytometry was performed to assess levels of protein expression
on brain endothelium day 6 of PbA infection. o) Scatter plot represents MFI values of P-selectin
on brain endothelial cells. p) MFI of ICAM-1on brain endothelial cells q) MFI of VCAM-1 on
brain endothelial cells. r) Percentage and MFI quantification of MHC Class II molecules on brain
endothelial cells with representative histogram. Data were analyzed with unpaired t test.
Scatterplots and bar plots present the mean $\pm$ SD. All p-values are shown or ns for non-
significant p-value *(p > 0.05)*.
1049
1050   **Fig 5. Modulation of individual class I molecules on brain endothelium results in brain-**
1051   **specific CD8 T cell infiltration changes.** a) BioRender schematic of events in the periphery
1052   leading up to cerebral malaria. b) Representative UMAP key, colorized to highlight the different

cell subsets in the brain. c) Representative UMAPs of K^b Cre- Littermate and Cdh5-cre K^b cKO brains, colorized with heat map of CD8 expression. d) Pie charts with corresponding frequencies of cell subset in K^b Cre- Littermate and Cdh5-cre K^b cKO brains (of CD45+) e) Absolute counts of CD4 T cells in the brains of infected K^b Cre- Littermate and Cdh5-cre K^b cKO brains. f) Absolute counts of CD8 T cells in the brains of infected K^b Cre- Littermate and Cdh5-cre K^b cKO brains. g) Absolute counts of IFN-γ+ CD8 T cells in the brains of infected K^b Cre- Littermates and Cdh5-cre K^b cKOs. h) Absolute counts of IFN-γ+ CD4 T (Th1) cells in the brains of infected K^b Cre- Littermates and Cdh5-cre K^b cKOs. i) Absolute counts of CD11a+ (antigen-experienced) CD8 T cells in the meninges, cLN and brain of infected K^b Cre- Littermates and Cdh5-cre K^b cKOs. Mice were injected with Alexa Fluor 488-anti-Thy1.2 antibody ~3 min. before euthanasia. j) Quantified absolute counts of either IV+ or IV- antigen-experienced CD8 T cells in the brains of infected K^b Cre- Littermates and Cdh5-cre K^b cKOs. k) Frequency of IFN-γ+ CD8 T cells either labeled with IV antibody or negative for IV labeling. l) Representative UMAPs of D^b Cre- Littermate and Cdh5-cre D^b cKO brains, colorized with heat map of CD8 expression. m) Pie charts with corresponding frequencies of cell subset in D^b Cre- Littermate and Cdh5-cre D^b cKO brains (of CD45+) n) Absolute counts of CD4 T cells in the brains of infected D^b Cre- Littermate and Cdh5-cre D^b cKO brains. o) Absolute counts of CD8 T cells in the brains of infected D^b Cre- Littermate and Cdh5-cre D^b cKO brains. p) Absolute counts of IFN-γ+ CD8 T cells in the brains of infected D^b Cre- Littermates and Cdh5-cre D^b cKOs. q) Absolute counts of IFN-γ+ CD4 T (Th1) cells in the brains of infected D^b Cre- Littermates and Cdh5-cre D^b cKOs. r) Absolute counts of CD11a+ (antigen-experienced) CD8 T cells in the meninges, cLN and brain of infected D^b Cre- Littermates and Cdh5-cre D^b cKOs. Mice were injected with Alexa Fluor 488-anti-Thy1.2 antibody ~3 min. before euthanasia. s) Quantified absolute counts of either IV+ or IV- antigen-experienced CD8 T cells in the brains of infected D^b Cre- Littermates and Cdh5-cre D^b cKOs. t) Frequency of IFN-γ+ CD8 T cells either labeled with IV antibody or negative for IV labeling. Data were analyzed with unpaired t test or 2way ANOVA with multiple comparisons. Scatterplots and bar plots present the mean $\pm$ SD. All p-values are shown or ns for non-significant p-value *(p > 0.05)*.

Fig 6. Vascular leakage during PbA infection is dependent on class I molecule expression on endothelium. a) Top – Representative T1-weighted gadolinium enhanced MRI (sagittal slice) from K^b Cre- littermate day 6-8 PbA infection showing vascular leakage in regions of enhancement. Bottom - Representative T1-weighted gadolinium enhanced MRI (sagittal slice) from Cdh5-cre K^b cKO mice day 6-8 PbA infection with no vascular leakage or enhancement noted. Top right – 3D volumetric modeling of whole brain with vascular leakage highlighted in red. Top left – 3D volumetric modeling of whole brain with no apparent vascular leakage noted. b) Area measurements of regions of enhancement. Enhancement was quantified for each individual MRI slice and combined into a cumulative measurement per mouse. 2 blinded individuals performed measurements. Each data point represents the average of the scores for each mouse/scan. c) 3D pixelated volumetric measurement were obtained from all slices and averaged from 2 blinded reviewers as mentioned in b. d) FITC-albumin leakage assay results. Mice were IV injected with 66kDa FITC-conjugated albumin 1 hour prior to euthanasia to visualize regions of blood-brain barrier disruption and vascular leakage. e) Quantification of FITC-albumin mean fluorescence intensity per ROI. f) Top – Representative T1-weighted gadolinium enhanced MRI (sagittal slice) from D^b Cre- littermate day 6-8 PbA infection showing vascular leakage in regions of enhancement. Bottom - Representative T1-weighted gadolinium

enhanced MRI (sagittal slice) from Cdh5-cre D^b cKO mice day 6-8 PbA infection with no
vascular leakage or enhancement noted. Top right – 3D volumetric modeling of whole brain with
vascular leakage highlighted in red. Top left – 3D volumetric modeling of whole brain with no
apparent vascular leakage noted. g) Area measurements of regions of enhancement.
Enhancement was quantified for each individual MRI slice and combined into a cumulative
measurement per mouse. 2 blinded individuals performed measurements. Each data point
represents the average of the scores for each mouse/scan. h) 3D pixelated volumetric
measurement were obtained from all slices and averaged from 2 blinded reviewers as mentioned
in g. i) FITC-albumin leakage assay results. Mice were IV injected with 66kDa FITC-conjugated
albumin 1 hour prior to euthanasia to visualize regions of blood-brain barrier disruption and
vascular leakage. j) Quantification of FITC-albumin mean fluorescence intensity per ROI. Data
were analyzed with unpaired t test or 2way ANOVA with multiple comparisons. Violin plots and
bar plots present the mean $\pm$ SD. All p-values are shown or ns for non-significant p-value *(p >*
0.05).
1113
Fig 7. PbA infected parabionts sharing peripheral circulation demonstrate requirement of
endothelial D^b expression for T cell responses and development of ECM. a) Schematic
depicting parabiosis procedure and respective D^b Cre- littermate and Cdh5-cre D^b cKO mice. b)
Timeline of experiment depicting parabiosis surgery, healing and acclimation time and PbA
infection of parabionts. c) Absolute counts of total CD45+ immune cells, CD8 T cells, CD4 T
cells and activated CD8s and CD4s in the spleens of parabionts day 6 of PbA infection. d)
Absolute counts of total CD45+ immune cells, CD8 T cells, CD4 T cells and activated CD8s and
CD4s in the iLNs of parabionts day 6 of PbA infection. e) Absolute counts of total CD45+
immune cells, CD8 T cells, CD4 T cells and IFN-γ producing CD8s and CD4s in the brains of
parabionts day 6 of PbA infection. f) FITC-albumin was IV injected one hour before euthanasia
of PbA infected parabionts. D^b Cre- littermate and Cdh5-cre D^b cKO parabiont brains are
visualized with FITC leakage in green. g) T1-weighted gadolinium enhanced dual MRI of
parabiont pair was performed. Representative image is one axial slice with visualization of
gadolinium enhancement. Data were analyzed with unpaired t test. Bar plots present the mean $\pm$
SD. All p-values are shown or ns for non-significant p-value *(p > 0.05).*
1129
Fig 8. Discrete class I molecules differentially regulate cell death and neuroinflammation in
the brain during ECM. Nanostring analysis was performed on whole brain isolated RNA of
PbA infected C57BL/6, K^b and D^b Cre- littermates, Cdh5-cre K^b cKO, Cdh5-cre D^b cKO and
uninfected controls of each group (n = 40, n = 4 mice/group). a) Venn diagrams of differentially
expressed genes are visualized for comparison of differential class I regulation by wildtype mice,
mice with only K^b and mice with only D^b expression. High-dimensional spectral flow cytometry
was performed using a staining panel specific for stages of cell death. Frequencies of cells
positive for markers of early apoptosis, late apoptosis and necrosis (late cell death) determined
by annexin V staining vs viability stain gating. b) Frequencies of respective stages of cell death
in Cre- littermates and Cdh5-cre K^b cKO expressed by brain endothelial cells on day 6 PbA
infection. c) Frequencies of respective stages of cell death in Cre- littermates and Cdh5-cre K^b
cKO expressed by mature neurons on day 6 PbA infection. d) Blinded clinical scoring was
conducted daily on Cre- littermates and Cdh5-cre K^b cKO mice infected with PbA. e) Survival
studies were performed on infected Cre- littermates and Cdh5-cre K^b cKO mice. Mice with K^b
deletion had largely improved survival. f) Frequencies of respective stages of cell death in Cre-

littermates and Cdh5-cre D^b cKO expressed by brain endothelial cells on day 6 PbA infection. g)
Frequencies of respective stages of cell death in Cre- littermates and Cdh5-cre D^b cKO expressed
by mature neurons on day 6 PbA infection. h) Blinded clinical scoring was conducted daily on
Cre- littermates and Cdh5-cre D^b cKO mice infected with PbA. i) Survival studies were
performed on infected Cre- littermates and Cdh5-cre D^b cKO mice. Mice with D^b deletion had
vastly improved survival. j) IPA analysis was performed, and gene Expression Log Ration data
was plotted in heat map to visualize changes in expression between groups. Data were analyzed
with unpaired t test or 2way ANOVA with multiple comparisons. For survival, Kaplan Meier
curves were plotted, and Log-rank (Mantel-Cox) curve comparison was performed. Scatter plots
and bar plots present the mean $\pm$ SD. All p-values are shown or ns for non-significant p-value *(p*
> 0.05).
1156

Supplementary Table 1. IPA Causal network analysis identifies MHC Class I family as the
number one master regulator of experimental cerebral malaria pathology. Causal network
analysis was performed using the Ingenuity Pathway Analysis application. Table includes
normalized data from causal network analysis which identifies master regulators of the most
highly enriched signaling pathways in the Nanostring dataset comparing uninfected C57BL/6
brain vs. C57BL/6 brains with ECM. Master regulators of causal networks are listed in order of
significance, with MHC Class I family identified as most significant causal network of
experimental cerebral malaria pathology. P-value of overlap was adjusted using the Benjamini-
Hochberg correction method.
1166

1167 **Supplementary Table 2. Differentially expressed genes in human endothelial cells after**
1168 ***Plasmodium* infection exposure.** Using GEO2R analysis platform MHC Class I was identified
1169 as one of the most significantly differentially expressed genes and was in the top 5 genes with
1170 significant upregulation on HBMECs after *Plasmodium* infection. Listed in order of significance.
1171

1172 **Supplementary Table 3. Patterns of neuroinflammation determined by cell specific gene**
1173 **enrichment during ECM determined through Mouse Glial Profiling Panel**. Table includes
1174 the nCounter Mouse Glial Profiling Panel gene lists, probes and corresponding cell type
1175 annotations detailing gene expression patterns and their corresponding neuroinflammatory
1176 response types.
1177

Supplementary Figure 1. Validation of human data comparison. a) Boxplots show the
distribution of the expression values for each human subject. Medians are centered indicating
that human datasets are suitable for comparison. Value distribution analysis performed by
GEO2R Web Application – NCBI GEO. b) MHC Class II Log$_2$ expression in control HBMECs
vs. *Plasmodium* exposed HBMECs. Data were analyzed with unpaired t test (b). Box plots
present the mean $\pm$ SD. All p-values are shown or ns for non-significant p-value *(p > 0.05)*.
1184

Supplementary Figure 2. Human and mouse antigen presentation pathways and
corresponding genes show similar activation with *Plasmodium* infection. Ingenuity Pathway
analyses of mouse and human antigen presentation pathways are depicted in schematics.
Coloring of the molecules corresponds to calculated z-scores (algorithm predicting activation or
inhibition). Red indicates increased gene activation, orange indicates predicted activation of

pathways, green indicates decreased gene activation, blue indicates predicted inhibition of
pathways, white indicates a z-score of zero with equal evidence for activation and inhibition,
grey indicates that z-score could not be calculated.
1193
Supplementary Figure 3. Peripheral immune system is intact K^b Cre- littermates and
Cdh5-cre K^b cKO mice. a) Percentage of mTECs and cTECs quantified from thymi of naïve
mice across groups and thymic weights of naïve mice across groups. b) Comparison of
percentages of thymic T cell subsets in males vs females across groups in naïve mice. c)
Percentages of respective VBeta TCRs out of CD8 T cells of the spleen in naïve mice. d) Splenic
weights of naïve mice across groups. e) Absolute counts of CD45+ immune cells in respective
peripheral organs of infected and naïve mice. f) Representative UMAPs depicting peripheral
organs and immune cell populations and respective heats maps with CD44 activation marker
expression. g) The scatter plots show quantified absolute cell counts of CD8 T cells, CD4 T cells
and CD44hi CD8 T cells and CD4 T cells in the respective organs depicted in UMAPs. Data
were analyzed with unpaired t test, Ordinary 1way ANOVA with multiple comparisons or 2way
ANOVA with multiple comparisons. Scatterplots and bar plots present the mean $\pm$ SD. All p-
values are shown or ns for non-significant p-value *(p > 0.05)*.
1207
Supplementary Figure 4. Peripheral immune system is intact in D^b Cre- littermates and
Cdh5-cre D^b cKO mice. a) Percentage of mTECs and cTECs quantified from thymi of naïve
mice across groups and thymic weights of naïve mice across groups. b) Comparison of
percentages of thymic T cell subsets in males vs females across groups in naïve mice. c)
Percentages of respective VBeta TCRs out of CD8 T cells of the spleen in naïve mice. d) Splenic
weights of naïve mice across groups. e) Absolute counts of CD45+ immune cells in respective
peripheral organs of infected mice. f) Scatter plots show quantified absolute cell counts of
CD45+ immune cells, CD8 T cells, CD4 T cells and CD44hi CD8 T cells and CD4 T cells in
respective organs of naïve mice. Data were analyzed with unpaired t test, Ordinary 1way
ANOVA with multiple comparisons or 2way ANOVA with multiple comparisons. Scatterplots
and bar plots present the mean $\pm$ SD. All p-values are shown or ns for non-significant p-value *(p*
> 0.05).
1220
Supplementary Figure 5. No changes to brain vasculature at baseline in mice with class I
deletion on endothelium. a) Representative images from maximum intensity projections (MIPs)
of naïve Cdh5-Cre K^b cKO and Cre- littermate controls. Mice were IP injected with gadolinium
15 min. before MRIs were performed. b) Quantification of 3D volume of overall brain
vasculature enhancement and quantification of 3D pixelated voxel units. c) Representative
images from maximum intensity projections (MIPs) of naïve Cdh5-Cre D^b cKO and Cre-
littermate controls. Mice were IP injected with gadolinium 15 min. before MRIs were performed.
d) Quantification of 3D volume of overall brain vasculature enhancement and quantification of
3D pixelated voxel units. Data were analyzed with unpaired t test. Scatterplots present the mean
$\pm$ SD. All p-values are shown or ns for non-significant p-value *(p > 0.05)*.
1231
1232 **Supplementary Figure 6. CD8 T cells infiltrate regions of interest except when K^b is**
1233 **ablated from vasculature.** a) Timeline depicting intracranial infection on Day 0 with TMEV or
1234 TMEV-OVA which is allowed to progress until day 7 when brains are harvested for high-
1235 dimensional spectral flow cytometry. b) Quantification of absolute counts of K^b: $OVA_{257\text{-}}$

$_{264}^+$ CD8 T cells in Cdh5-Cre K^b cKO and Cre- littermate cLNs day 7 of TMEV-OVA infection. c) Quantification of absolute counts of D^b: VP2$^+$ CD8 T cells in Cdh5-Cre K^b cKO and Cre- littermate cLNs day 7 of TMEV infection. d) Absolute counts of CD45+ immune cells in the brains of Cdh5-Cre K^b cKO and Cre- littermate brains during ECM. e) Absolute counts of CD45+ immune cells in the brains of Cdh5-Cre D^b cKO and Cre- littermate brains during ECM. f) Quantified percentage of CD8s that are CXCR3+ in brains of Cdh5-Cre K^b cKO and Cre- littermates during ECM. g) Quantification of median fluorescence intensities of CXCR3 on CD8 T cells in the brains of Cdh5-Cre K^b cKO and Cre- littermates during ECM. h) Quantified percentage of CD8s that are CCR7+ in brains during ECM. i) Quantification of median fluorescence intensities of CCR7 on CD8 T cells in the brain during ECM. j) Quantified percentage of CD8s that are CXCR3+ in brains of Cdh5-Cre D^b cKO and Cre- littermates during ECM. k) Quantification of median fluorescence intensities of CXCR3 on CD8 T cells in the brains of Cdh5-Cre D^b cKO and Cre- littermates during ECM. l) Quantified percentage of CD8s that are CCR7+ in brains during ECM. m) Quantification of median fluorescence intensities of CCR7 on CD8 T cells in the brain during ECM. n) Calculation of rate of infiltration of antigen-experienced CD11a+ CD8 T cells into the brain from IV labeled vasculature, measured in cells/min in infected K^b Cre- littermates and Cdh5-Cre K^b cKOs. o) Calculation of rate of infiltration of antigen-experienced CD11a+ CD8 T cells into the brain from IV labeled vasculature, measured in cells/min in infected D^b Cre- littermates and Cdh5-Cre D^b cKOs. p) Calculation of rate of infiltration of CD11a- CD8 T cells that have not encountered cognate peptide:MHC into the brain from IV labeled vasculature, measured in cells/min in infected K^b Cre- littermates and Cdh5-Cre K^b cKOs. q) Calculation of rate of infiltration of CD11a- CD8 T cells that have not encountered cognate peptide:MHC into the brain from IV labeled vasculature, measured in cells/min in infected D^b Cre- littermates and Cdh5-Cre D^b cKOs. Confocal imaging was performed. All images were acquired at 20X and zoomed in to 400X using FIJI. r) Representative immunofluorescence images of CD8 T cell infiltration in ROIs affected by ECM showing infiltration into these regions in the K^b Cre- littermates but not in the Cdh5-Cre K^b cKOs. Scale bar, 50 µM. s) Representative immunofluorescence images of CD8 T cell infiltration in ROIs affected by ECM showing infiltration into these regions in the D^b Cre- littermates and in the Cdh5-Cre D^b cKOs. Scale bar, 50 µM. Data were analyzed with unpaired t test. Bar plots present the mean $\pm$ SD. All p-values are shown or ns for non-significant p-value *(p > 0.05)*.

Supplementary Figure 7. Tight junction protein alterations occur in human and mice with cerebral malaria. GSEA was performed on HBMEC transcriptomic data. a) Enrichment plot showing a negative enrichment (downregulation) of molecular signature for tight junction gene set. The y-axis represents enrichment score (ES) and on the x-axis are genes (vertical black lines) represented in gene sets. The green line connects points of ES and genes. ES is the maximum deviation from zero as calculated for each gene going down the ranked list and represents the degree of over-representation of a gene set at the top or the bottom of the ranked gene list. The colored band at the bottom represents the degree of correlation of genes with the phenotype (red for positive and blue for negative correlation). Significance threshold set at nominal p-value < 0.05. Gene sets for tight junctions were significantly negatively enriched indicating downregulation in HBMECs during infection. Confocal imaging was performed. All images were acquired at 20X and zoomed in to 300X using FIJI. b) Immunofluorescence staining of tight junction protein Claudin-5 in ROI in PbA infected Cdh5-Cre K^b cKOs and K^b Cre-

littermates brains day 6. Scale bar, 50 µM. c) Immunofluorescence staining of tight junction protein Claudin-5 in ROI in PbA infected Cdh5-Cre D^b cKOs and D^b Cre- littermates day 6. Scale bar, 50 µM.

Supplementary Figure 8. Discrete endothelial class I molecules regulate differential downstream gene expression. a) Venn diagrams of differentially expressed genes are visualized for comparison of differential class I regulation in wildtype, K^b Cre- littermates and Cdh5-cre K^b cKO mice. Volcano plots of differentially expressed genes in whole brains of uninfected versus PbA infected groups of mice at endpoint during ECM. Significantly upregulated genes are indicated on upper right (purple dots). Significantly downregulated genes are indicated on upper left (turquoise dots). Horizontal dotted line indicates cutoff of adjusted p-values (-Log10 of adjusted p-value 1.3 corresponding to a threshold of p-value of 0.05). Vertical dotted line indicates cutoff of log2 Fold change (-1.5 to 1.5). b) Volcano plots of differentially expressed genes in infected K^b Cre- littermates vs Uninfected K^b Cre- littermates. c) Volcano plots of differentially expressed genes in infected Cdh5-cre K^b cKO mice vs Uninfected Cdh5-cre K^b cKO mice. d) Venn diagrams of differentially expressed genes are visualized for comparison of differential class I regulation in wildtype, D^b Cre- littermates and Cdh5-cre D^b cKO mice. e) Volcano plots of differentially expressed genes in infected D^b Cre- littermates vs Uninfected D^b Cre- littermates. f) Volcano plots of differentially expressed genes in infected Cdh5-cre D^b cKO mice vs Uninfected Cdh5-cre D^b cKO mice. P-values of overlap were adjusted using the Benjamini-Hochberg correction method.

References

1.   Organization, W.H. World malaria report 2017. Geneva: WHO; 2017. 2018.

2.   Organization, W.H. World malaria report 2021. (2021).

3.   Idro, R., Marsh, K., John, C.C. & Newton, C.R. Cerebral malaria: mechanisms of brain injury and strategies for improved neurocognitive outcome. *Pediatr Res* **68**, 267-274 (2010).

4.   Severe and complicated malaria. World Health Organization, Division of Control of Tropical Diseases. *Trans R Soc Trop Med Hyg* **84 Suppl 2**, 1-65 (1990).

5.   Mahmoudi, S. & Keshavarz, H. Efficacy of phase 3 trial of RTS, S/AS01 malaria vaccine: The need for an alternative development plan. *Human vaccines & immunotherapeutics* **13**, 2098-2101 (2017).

6.   Matuschewski, K. Vaccines against malaria—still a long way to go. *The FEBS journal* **284**, 2560-2568 (2017).

7.   Rowe, A.K. *et al.* The burden of malaria mortality among African children in the year 2000. *Int J Epidemiol* **35**, 691-704 (2006).

1327

1328 8. Misra, U. *et al.* Cerebral malaria and bacterial meningitis. *Annals of Indian Academy of*
1329   *Neurology* **14**, S35 (2011).
1330

1331 9. Tunon-Ortiz, A. & Lamb, T.J. Blood brain barrier disruption in cerebral malaria: Beyond
1332   endothelial cell activation. *PLoS Pathogens* **15**, e1007786 (2019).
1333

1334 10. Seydel, K.B. *et al.* Brain swelling and death in children with cerebral malaria. *New*
1335   *England Journal of Medicine* **372**, 1126-1137 (2015).
1336

1337 11. Riggle, B.A. *et al.* CD8+ T cells target cerebrovasculature in children with cerebral
1338   malaria. *The Journal of clinical investigation* **130**, 1128-1138 (2020).
1339

1340 12. Ghazanfari, N., Mueller, S.N. & Heath, W.R. Cerebral malaria in mouse and man.
1341   *Frontiers in immunology* **9** (2018).
1342

1343 13. Finley, R., Mackey, L. & Lambert, P. Virulent P. berghei malaria: prolonged survival and
1344   decreased cerebral pathology in cell-dependent nude mice. *The Journal of Immunology*
1345   **129**, 2213-2218 (1982).
1346

1347 14. Yañez, D.M., Manning, D.D., Cooley, A.J., Weidanz, W.P. & Van Der Heyde, H.
1348   Participation of lymphocyte subpopulations in the pathogenesis of experimental murine
1349   cerebral malaria. *The Journal of Immunology* **157**, 1620-1624 (1996).
1350

1351 15. Nitcheu, J. *et al.* Perforin-dependent brain-infiltrating cytotoxic CD8+ T lymphocytes
1352   mediate experimental cerebral malaria pathogenesis. *The Journal of Immunology* **170**,
1353   2221-2228 (2003).
1354

1355 16. Huggins, M.A. *et al.* Perforin expression by CD8 T cells is sufficient to cause fatal brain
1356   edema during experimental cerebral malaria. *Infection and immunity* **85**, e00985-00916
1357   (2017).
1358

1359 17. Swanson, P.A. *et al.* CD8+ T cells induce fatal brainstem pathology during cerebral
1360   malaria via luminal antigen-specific engagement of brain vasculature. *PLoS pathogens*
1361   **12**, e1006022 (2016).
1362

1363 18. Takada, K. & Jameson, S.C. Self–class I MHC molecules support survival of naive CD8
1364   T cells, but depress their functional sensitivity through regulation of CD8 expression
1365   levels. *Journal of Experimental Medicine* **206**, 2253-2269 (2009).
1366

1367 19. Tripathi, A.K., Sha, W., Shulaev, V., Stins, M.F. & Sullivan, D.J., Jr. Plasmodium
1368   falciparum-infected erythrocytes induce NF-kappaB regulated inflammatory pathways in
1369   human cerebral endothelium. *Blood* **114**, 4243-4252 (2009).
1370

1371 20. Combes, V. *et al.* Pathogenic role of P-selectin in experimental cerebral malaria:
1372   importance of the endothelial compartment. *The American journal of pathology* **164**, 781-
1373   786 (2004).
1374

1375 21. Goddery, E.N. *et al.* Microglia and perivascular macrophages act as antigen presenting
1376   cells to promote CD8 T cell infiltration of the brain. *Frontiers in immunology*, 3428
1377   (2021).
1378

1379 22. Suidan, G.L., Mcdole, J.R., Chen, Y., Pirko, I. & Johnson, A.J. Induction of blood brain
1380   barrier tight junction protein alterations by CD8 T cells. *PloS one* **3**, e3037 (2008).
1381

1382 23. Pavelko, K.D. *et al.* Theiler's murine encephalomyelitis virus as a vaccine candidate for
1383   immunotherapy. *PloS one* **6**, e20217 (2011).
1384

1385 24. Campanella, G.S. *et al.* Chemokine receptor CXCR3 and its ligands CXCL9 and
1386   CXCL10 are required for the development of murine cerebral malaria. *Proceedings of the*
1387   *National Academy of Sciences* **105**, 4814-4819 (2008).
1388

1389 25. Ghazanfari, N., Mueller, S.N. & Heath, W.R. Cerebral malaria in mouse and man.
1390   *Frontiers in immunology*, 2016 (2018).
1391

1392 26. Cupovic, J. *et al.* Central nervous system stromal cells control local CD8+ T cell
1393   responses during virus-induced neuroinflammation. *Immunity* **44**, 622-633 (2016).
1394

1395 27. Zhao, H. *et al.* Olfactory plays a key role in spatiotemporal pathogenesis of cerebral
1396   malaria. *Cell host & microbe* **15**, 551-563 (2014).
1397

1398 28. Rai, D., Pham, N.-L.L., Harty, J.T. & Badovinac, V.P. Tracking the total CD8 T cell
1399   response to infection reveals substantial discordance in magnitude and kinetics between
1400   inbred and outbred hosts. *The Journal of Immunology* **183**, 7672-7681 (2009).
1401

1402 29. Bhat, P., Leggatt, G., Waterhouse, N. & Frazer, I.H. Interferon-γ derived from cytotoxic
1403   lymphocytes directly enhances their motility and cytotoxicity. *Cell death & disease* **8**,
1404   e2836-e2836 (2017).
1405

1406 30. Krasemann, S. *et al.* The TREM2-APOE pathway drives the transcriptional phenotype of
1407   dysfunctional microglia in neurodegenerative diseases. *Immunity* **47**, 566-581. e569
1408   (2017).
1409

1410 31. Keren-Shaul, H. *et al.* A unique microglia type associated with restricting development of
1411   Alzheimer's disease. *Cell* **169**, 1276-1290. e1217 (2017).
1412

1413 32. Aguzzi, A., Barres, B.A. & Bennett, M.L. Microglia: scapegoat, saboteur, or something
1414   else? *Science* **339**, 156-161 (2013).
1415

1416 33. Deczkowska, A. *et al.* Disease-associated microglia: a universal immune sensor of
1417 neurodegeneration. *Cell* **173**, 1073-1081 (2018).
1418

1419 34. Pais, T.F. & Penha-Gonçalves, C. Brain endothelium: the "innate immunity response
1420 hypothesis" in cerebral malaria pathogenesis. *Frontiers in immunology* **9**, 3100 (2019).
1421

1422 35. Carman, C.V. & Martinelli, R. T lymphocyte–endothelial interactions: emerging
1423 understanding of trafficking and antigen-specific immunity. *Frontiers in immunology* **6**,
1424 603 (2015).
1425

1426 36. Waisman, A. & Johann, L. Antigen-presenting cell diversity for T cell reactivation in
1427 central nervous system autoimmunity. *Journal of Molecular Medicine* **96**, 1279-1292
1428 (2018).
1429

1430 37. Amersfoort, J., Eelen, G. & Carmeliet, P. Immunomodulation by endothelial cells—
1431 partnering up with the immune system? *Nature Reviews Immunology*, 1-13 (2022).
1432

1433 38. Murphy, K. & Weaver, C. *Janeway's immunobiology*. Garland science, 2016.
1434

1435 39. Engelhardt, B. & Ransohoff, R.M. The ins and outs of T-lymphocyte trafficking to the
1436 CNS: anatomical sites and molecular mechanisms. *Trends in immunology* **26**, 485-495
1437 (2005).
1438

1439 40. Laschinger, M. & Engelhardt, B. Interaction of α4-integrin with VCAM-1 is involved in
1440 adhesion of encephalitogenic T cell blasts to brain endothelium but not in their
1441 transendothelial migration in vitro. *Journal of neuroimmunology* **102**, 32-43 (2000).
1442

1443 41. Marchetti, L. & Engelhardt, B. Immune cell trafficking across the blood-brain barrier in
1444 the absence and presence of neuroinflammation. *Vascular Biology* **2**, H1-H18 (2020).
1445

1446 42. Sathiyanadan, K., Coisne, C., Enzmann, G., Deutsch, U. & Engelhardt, B. PSGL-1 and
1447 E/P-selectins are essential for T-cell rolling in inflamed CNS microvessels but
1448 dispensable for initiation of EAE. *European journal of immunology* **44**, 2287-2294
1449 (2014).
1450

1451 43. Villegas-Mendez, A. *et al.* IFN-γ–producing CD4+ T cells promote experimental cerebral
1452 malaria by modulating CD8+ T cell accumulation within the brain. *The Journal of
1453 Immunology* **189**, 968-979 (2012).
1454

1455 44. Krausgruber, T. *et al.* Structural cells are key regulators of organ-specific immune
1456 responses. *Nature* **583**, 296-302 (2020).
1457

1458 45. Howland, S.W. *et al.* Brain microvessel cross-presentation is a hallmark of experimental
1459 cerebral malaria. *EMBO molecular medicine* **5**, 984-999 (2013).
1460

1461   46.   Belnoue, E. *et al.* Control of pathogenic CD8+ T cell migration to the brain by IFN-γ
1462   during experimental cerebral malaria. *Parasite immunology* **30**, 544-553 (2008).
1463

1464   47.   Suidan, G.L. *et al.* CD8 T cell-initiated vascular endothelial growth factor expression
1465   promotes central nervous system vascular permeability under neuroinflammatory
1466   conditions. *The journal of immunology* **184**, 1031-1040 (2010).
1467

1468   48.   Miyakoda, M. *et al.* Malaria-specific and nonspecific activation of CD8+ T cells during
1469   blood stage of Plasmodium berghei infection. *The Journal of Immunology* **181**, 1420-
1470   1428 (2008).
1471

1472   49.   Poh, C.M., Howland, S.W., Grotenbreg, G.M. & Rénia, L. Damage to the blood-brain
1473   barrier during experimental cerebral malaria results from synergistic effects of CD8+ T
1474   cells with different specificities. *Infection and immunity* **82**, 4854-4864 (2014).
1475

1476   50.   Kalkal, M. *et al.* A comprehensive study of epitopes and immune reactivity among
1477   Plasmodium species. *BMC microbiology* **22**, 1-14 (2022).
1478

1479   51.   Pardy, R.D. *et al.* Analysis of the T cell response to Zika virus and identification of a
1480   novel CD8+ T cell epitope in immunocompetent mice. *PLoS pathogens* **13**, e1006184
1481   (2017).
1482

1483   52.   Mora, J.R. & von Andrian, U.H. T-cell homing specificity and plasticity: new concepts
1484   and future challenges. *Trends in immunology* **27**, 235-243 (2006).
1485

1486   53.   Ioannidis, L.J., Nie, C.Q. & Hansen, D.S. The role of chemokines in severe malaria: more
1487   than meets the eye. *Parasitology* **141**, 602-613 (2014).
1488

1489   54.   Miu, J. *et al.* Chemokine gene expression during fatal murine cerebral malaria and
1490   protection due to CXCR3 deficiency. *The Journal of Immunology* **180**, 1217-1230
1491   (2008).
1492

1493   55.   Strangward, P. *et al.* A quantitative brain map of experimental cerebral malaria
1494   pathology. *PLoS pathogens* **13**, e1006267 (2017).
1495

1496   56.   Solouki, S. *et al.* TCR signal strength and antigen affinity regulate CD8+ memory T
1497   cells. *The Journal of Immunology* **205**, 1217-1227 (2020).
1498

1499   57.   Tan, M. *et al.* T cell receptor binding affinity governs the functional profile of cancer-
1500   specific CD8+ T cells. *Clinical & Experimental Immunology* **180**, 255-270 (2015).
1501

1502   58.   Irving, M. *et al.* Interplay between T cell receptor binding kinetics and the level of
1503   cognate peptide presented by major histocompatibility complexes governs CD8+ T cell
1504   responsiveness. *Journal of Biological Chemistry* **287**, 23068-23078 (2012).
1505

1506 59. Martinez, R.J. & Evavold, B.D. Lower affinity T cells are critical components and active
1507   participants of the immune response. *Frontiers in immunology* **6**, 468 (2015).
1508

1509 60. Tau, G.Z., Cowan, S.N., Weisburg, J., Braunstein, N.S. & Rothman, P.B. Regulation of
1510   IFN-γ signaling is essential for the cytotoxic activity of CD8+ T cells. *The Journal of*
1511   *Immunology* **167**, 5574-5582 (2001).
1512

1513 61. Malo, C.S. *et al.* Non-equivalent antigen presenting capabilities of dendritic cells and
1514   macrophages in generating brain-infiltrating CD8+ T cell responses. *Nature*
1515   *communications* **9**, 1-13 (2018).
1516

1517 62. Tritz, Z.P. *et al.* Conditional Silencing of H-2Db Class I Molecule Expression Modulates
1518   the Protective and Pathogenic Kinetics of Virus-Antigen–Specific CD8 T Cell Responses
1519   during Theiler's Virus Infection. *The Journal of Immunology* **205**, 1228-1238 (2020).
1520

1521 63. Hortle, E. *et al.* KCC1 Activation protects Mice from the Development of Experimental
1522   Cerebral Malaria. *Scientific reports* **9**, 1-11 (2019).
1523

1524 64. Organization, W.H. & Control, C.f.D. *Basic malaria microscopy: tutor's guide.* World
1525   Health Organization, 2010.
1526

1527 65. Johnson, H.L. *et al.* CD8 T cell-initiated blood–brain barrier disruption is independent of
1528   neutrophil support. *The Journal of Immunology* **189**, 1937-1945 (2012).
1529

1530 66. Garcia, L.M.C., Kelcher, A.M.H., Malo, C.S. & Johnson, A.J. Superior isolation of
1531   antigen-specific brain infiltrating T cells using manual homogenization technique.
1532   *Journal of immunological methods* **439**, 23-28 (2016).
1533

1534 67. Rustenhoven, J. *et al.* Functional characterization of the dural sinuses as a neuroimmune
1535   interface. *Cell* **184**, 1000-1016. e1027 (2021).
1536

1537 68. Liu, Y.U. *et al.* Neuronal network activity controls microglial process surveillance in
1538   awake mice via norepinephrine signaling. *Nature neuroscience* **22**, 1771-1781 (2019).
1539

1540 69. Eyo, U.B. *et al.* P2Y12R-Dependent Translocation Mechanisms Gate the Changing
1541   Microglial Landscape. *Cell reports* **23**, 959-966 (2018).
1542

1543 70. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature*
1544   *methods* **9**, 676-682 (2012).
1545

1546 71. Shakhar, G. *et al.* Stable T cell-dendritic cell interactions precede the development of
1547   both tolerance and immunity in vivo. *Nature immunology* **6**, 707-714 (2005).
1548

1549 72. Tripathi, A.K., Sullivan, D.J. & Stins, M.F. Plasmodium falciparum-infected erythrocytes
1550   increase intercellular adhesion molecule 1 expression on brain endothelium through NF-
1551   κB. *Infection and immunity* **74**, 3262-3270 (2006).

1552
1553   73.   Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets—update. *Nucleic*
1554   *acids research* **41**, D991-D995 (2012).
1555
1556   74.   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
1557   interpreting genome-wide expression profiles. *Proceedings of the National Academy of*
1558   *Sciences* **102**, 15545-15550 (2005).
1559
1560
1561

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

b Area of Enhancement (mm³) — 0.0202

Kᵇ Cre- Littermates / Cdh5-cre Kᵇ cKOs

c Voxel Units

Kᵇ Cre- Littermates / Cdh5-cre Kᵇ cKOs

a

Cre-

Cre+

Wildtype Uninfected   Wildtype Infected   Cre- PbA Infected   Cre+ PbA Infected

d

e Fitc-Albumin MFI — 0.0002   ns   0.0167

■ Kᵇ Cre- Littermates
▲ Cdh5-cre Kᵇ cKOs

Olfactory Bulb   Hippocampus   Brainstem

f

Cre-

Cre+

g Area of Enhancement (mm³) — 0.0158

Dᵇ Cre- Littermates / Cdh5-cre Dᵇ cKOs

h Voxel Units

Dᵇ Cre- Littermates / Cdh5-cre Dᵇ cKOs

WT PbA Infected   Cre- PbA Infected   Cre+ PbA Infected

i

j Fitc-Albumin MFI — 0.0013   ns   0.0482

■ Dᵇ Cre- Littermates
▲ Cdh5-cre Dᵇ cKOs

Olfactory Bulb   Hippocampus   Brainstem

Figure 7

Figure 8

Supplementary Figure 1

a

b

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

a Infect i.c. with TMEV OR TMEV-OVA

Day 0
Day1 Day 2 Day 3 Day 4 Day 5 Day6 Day 7
Harvest
TMEV Infection

b TMEV-OVA

c TMEV

d

e

f

g

h

i

j

k

l

m

n

o

p

q

r Kᵇ Cre- Littermate | Cdh5-cre Kᵇ cKO
DAPI/CD8/Claudin-5
Olfactory
Brainstem
Hippocampus

s Dᵇ Cre- Littermate | Cdh5-cre Dᵇ cKO
DAPI/CD8/Claudin-5
Olfactory
Brainstem
Hippocampus

Supplementary Figure 7

a

b

Claudin-5

K^b Cre- Littermates   Cdh5-cre K^b cKOs

c

Claudin-5

D^b Cre- Littermates   Cdh5-cre D^b cKOs

Supplementary Figure 8