Robust expression of transgenes in *Drosophila melanogaster*

Peter V. Lidsky[1*], Sergey E. Dmitriev[2], and Raul Andino[1*].

[1]Department of Microbiology and Immunology, University of California San Francisco, San Francisco,
94158, CA

[2]Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University, Moscow
119234, Russia

*corresponding authors: peter.lidsy@ucsf.edu, raul.andino@ucsf.edu

Abstract

Drosophila is a classic experimental system used in fundamental and biopharmaceutical research.
Manipulating gene expression in the larvae and adult flies can facilitate basic and translational studies.
Here we report a method for robust transgene overexpression in *Drosophila melanogaster*. This approach
is compatible with the UAS/Gal4 gene expression system. The improved expression involves a gene
expression cassette that prevents transgenic mRNA transcript degradation by the nonsense-mediated RNA
decay pathway. The protection is mediated by CrPV and DCV IRES sequences that apparently encode
cryptic polyadenylation sites and NMD inhibitors. Stabilization of the transgene mRNA results in a >40-
fold increase in its levels.

Introduction

Controlling expression levels in the *Drosophila* model is critical for scientific and biotechnological
applications[1–3]. For instance, expressing brighter fluorescent proteins is critical for imaging reporting in
real-time phenotypes. Expression levels of the transgenic proteins may provide a larger dynamic range,
which could better document complex phenotypes. Insect cells are widely used for protein production[4–6].
Increasing interest in insect models for pharmacological and toxicology applications. In addition, the food
industry anticipates genetically modified animals can be used to produce nutritional additives or bioactive
proteins[7–9].

Not only the levels of expression are of great interest. Tools allowing production two or more proteins at
once are often required. Currently, the primary approach for protein co-expression in *Drosophila* is virus-
derived T2A self-processing cassettes[10]. This type of expression cassettes contains *cis*-acting hydrolase
element that enable resolution of the fusion proteins by a mechanisms called ribosome skipping[11]. The
T2A strategy generates proteins that have additional aminoacids at the N- and C-termini of the expressed
protein[10]. Alternatively, viral internal ribosomal entry sites (IRESes) in mammalian systems are equally

popular[12–17]. While IRESes are described in insect viruses[18,19], they have never been developed into

protein tools for protein production. Moreover, IRESes often lead to variable levels of the expressed

proteins [20,21]. In the context of this study we found that cryptic polyadenylation signals localized in the

intragenic IRESes of dicistroviruses. This leads to truncated mRNA that do not expressed the second

protein in the bicistronic construct. We found that these cryptic polyadenylation sites, when placed into

standard pUAST-based[22] constructs, result in unprecedented overexpression of the truncated transcript.

Our data suggest interaction with nonsense-mediated mRNA decay (NMD) machinery as a primary

reason for the overexpression and indicates the presence of in cis NMD inhibitors within the intergenic

IRESes of cricket paralysis virus (CrPV)[23] and drosophila C virus (DCV)[24].

Results

CrPV or DCV IRESes result in a robust overexpression of the upstream coding region. To analyze

the activity of the internal ribosomal entry sites (IRESes) *in vivo*, we engineered bicistronic constructs

harboring CrPV and DCV intergenic IRESes. The first open reading frame (ORF) encoded red

fluorescent protein mCherry translated via a cap-dependent mechanism, while the second IRES-directed

coding region produced EGFP. Control construct encoded a mCherry-EGFP fusion protein, ensuring

equimolar amounts of both reporters. The UAS promoter controlled the transcription initiation. The

mRNA polyadenylation was guided by the SV40 polyadenylation signal (**Fig 1A**). All constructs were

inserted into the 3rd chromosomal 86Fb-attP attachment site using ΦC31 integrase[25]. Transgenic flies

were crossed with the daughterless:Gal4 (da:Gal4) driver line to test IRES-mediated expression. IRES-

driven EGFP expression was not detected in most fly tissues (**Fig 1B**, middle row). In contrast, we found

that the levels of mCherry produced were enormously elevated compared to the in-frame fusion control

(**Fig 1B**, upper row). The level of overexpression was so high that the intense red coloration of the flies

was visible in daylight with a bare eye (**Fig 1B**, bottom row).

Levels of IRES-containing mRNAs are elevated; the transcripts are truncated and resistant to

NMD. To examine the mechanism by which the first cistron is overexpressed, but the second cistron is

not expressed, we examined the role of RNA decay pathways. SV40 polyadenylation cassette (~600 bp)

acts as a transcriptional enhancer, and used in most UAS-based constructs. Mutations affecting nonsense-

mediated RNA decay (NMD) have been reported to increase the levels of transcripts containing SV40

polyadenylation sites[26]. SV40-derived long 3' UTRs are recognized by the NMD machinery, thus

destabilizing the transcripts[26,27]. To test if NMD is responsible for IRES-mediated overexpression, we

crossed *Upf2*-mutant flies with da:Gal4 and bicistronic constructs expressing mCherry and EGFP (**Fig**

2A). $Upf2^{25G}$ mutation abrogates normal NMD function and is associated with late larval or pupal

lethality[26]. However, flies surviving to the third instar larvae stage could be analyzed for RNA and protein

expression (**Fig 2B-C**). First, we evaluated the mCherry levels with a fluorescent microscope (**Fig 2B**).

The mCherry-EGFP fusion construct displayed a significant upregulation in expression in the NMD-null

background (**Fig 2B,** compare fusion WT and mut). In contrast, IRES-encoding constructs did not display

a significant difference in expression levels. Next, we measured the mRNA levels with the use of RT-

qPCR. We used two oligo pair-sets that specifically amplify mCherry (red bars) or EGFP (green bars)

coding region (**Fig 2C**). The mCherry RNA readouts were significantly higher in IRES-containing

constructs compared to the fusion construct (**Fig 2C**). For instance, the RNA amounts in the IRES

construct containing CrPV exceeded the fusion control RNA accumulation by more than 40-fold.

Importantly, the expression of the fusion control construct was upregulated in the NMD-null background

(**Fig. 2C**, compare fusion WT with mut). In contrast, CrPV construct did not display any substantial

upregulation, while the DCV showed a mild, ~2 fold, increase in mCherry RNA levels. In fusion control

samples, but the IRES-containing transcripts the EGFP levels are lower than those corresponding to

mCherry (**Fig2C).** These results suggest that the second cistron is less stabler that the fist cistron in the

bicistronic construct. Similar results were obtained with eye-specific siRNA-mediated knockdowns of

NMD components *Upf1* and *Upf2* (**Fig 2D, E**).

To examine whether the IRES sequence results in RNA truncation and decay, we examine the integrity of

the bicistronic construct by Northern blotting analysis using specific probes directed against either

mCherry or EGFP coding sequences (**Fig 2F, G**). Consistently with the RT-qPCR results, we observed

two bands corresponding to the mCherry sequences, one corresponding to the bicistronic full length

transcript and the other smaller corresponding to mCherry first cistron. These shorter RNAs were

apparently not sensitive to NMD. The full-length product stained with mCherry and EGFP probes was

upregulated by *Upf2* inactivation in the control fusion construct but not in the CrPV IRES encoding

construct. The DCV IRES containing construct showed both truncated and the full-length transcripts,

which slightly increased in the *Upf2*-mutant background. Thus, we concluded bicistronic constructs (i)

undergo truncation of the transcripts and (ii) provide a certain level of resistance to NMD, especially with

CrPV IRES.

To determine the potential mechanism of transcript truncation, we tested if shorter transcripts (that

comprise most of the mCherry-encoded RNA fraction) are polyadenylated. We purified the poly(A)+

RNAs and analyzed these with qPCR. While purification resulted in a ~200-fold decrease in the amounts

of 18S ribosomal RNA (blue bars), no significant effects were found for the mCherry or EGFP-encoding

RNAs (**Fig 3**). Thus, we conclude the truncated mCherry-encoding transcripts we detected are

polyadenylated.

Discussion

The experiments described above show that CrPV and DCV IRESes present in bicistronic constructs

induce mRNA stabilization and a robust increase in production of the protein encoded by the 5' proximal

cistron. Based on these data, we propose the following model (**Fig 4**). In the absence of the IRESes, the

SV40 polyadenylation signal plays a dual role in gene expression: first, by enhancing the transcription,

second, by destabilizing the RNA by making it sensitive to NMD, in accordance with the earlier

findings[26] (**Fig 4A**).

In dicistroviral genomic RNAs, intragenic IRESes are localized immediately downstream of the first CDS

encoding non-structural proteins[28]. Termination at the stop codon of this CDS should activate NMD

machinery. Growing evidence suggests that NMD is also involved in the degradation of many RNA(+)

virus genomes[29–33]. The NMD-inhibiting *cis*-acting RNA structures were described previously in

retroviruses where these structures are located immediately after stop codons[34,35]. Unlike mammals where

NMD detection relies on exon junction complex recognition, in insects NMD degrades the mRNAs that

possess long 3'UTRs[36]. Termination in the first ORF of viral RNA of *Dicistroviruses* should activate the

degradation pathway. The intragenic IRES sequences that take positions downstream from the stop

codons might possess an additional activity to inhibit NMD (**Fig 4B**).

When placed into the intergenic spacer IRES sequences, introduce cryptic polyadenylation sites into

DNA, resulting in the production of truncated transcripts that have short 3'UTR and are resistant to NMD

(**Fig 2, 3**). This retains the enhancer properties of the SV40 region but mitigates its RNA-destabilizing

properties (**Fig 4C**).

In addition, the full-length bicistronic transcripts containing IRESes also displayed a decreased sensitivity

to NMD if compared to the control construct. The corresponding EGFP-specific RT-qPCR readouts were

not elevated (**Fig 2C)** or only slightly elevated (**Fig 2E**) in NMD-deficient strains, in contrast to those in

the case of the control. Consistently, Northern blot analysis revealed that the full-length transcript in the

CrPV IRES-encoding construct is not affected by *Upf2* inactivation, while the DCV IRES-encoding one

is only mildly stimulated (**Fig 2C, F-G**). Thus, the full-length IRES-containing constructs also confer

some resistance to NMD.

Thus, we suggest that CrPV and DCV IRESes, when put into genomic settings, might stabilize mRNA

transcripts by (i) shortening the 3' UTR due to the presense of cryptic polyadenylation signals and (ii)

direct inhibition of NMD machinery.

In summary, the transcription-enhancing SV40 polyadenylation signal combined with the mRNA-

stabilizing CrPV IRES constitutes a novel effective system for high-level protein production in

Drosophila. Several approaches for tuning expression levels in flies have been published so far[37]. These

methods were based on multimerization of transcription factor binding sites[37], translational regulators[38],

or enhancers of mRNA nuclear export[39]. The novelty of our approach consists in improvement of the

mRNA stability. The application of our method is not limited to fluorescent labels but could be

potentially extended to other reporters and biotechnological protein production.

Materials and methods

Cloning and injection. To construct *pUASt-mCherry-CrPV_ires-EGFP-attB* and *pUASt-mCherry-*

DCV_ires-EGFP-attB plasmids, encoding the IRES-based stress reporters, *pUASt-attB* vector was opened

with EcoRI and XbaI enzymes. Fragments encoding EGFP and mCherry were amplified by PCR. EGFP

fragment was digested with BglII and XbaI. mCherry fragment was treated with EcoRI and BsrGI. IRES

sequences were synthesized by Genescript and excised from their vector plasmid with BsrGI and BglII.

Constructs were injected into M{3xP3-RFP.attP}ZH-86Fb docking site strain[25] by BestGene Inc.

Fly lines. Upf2^{25G}/FM7 flies[26] were a kind gift of Prof. Mark M. Metzstein (University of Utah). A

ubiquitous Gal4 driver, *GAL4-da.G32* was on the third chromosome[40]; in the text, it is referenced as da-

Gal4. *y, w, ey-Flp, Act5c>CD2>Gal4* chromosome[41] was ascribed in the text as ey>Gal4 and was kindly

provided by Prof. Hugo Stocker (ETH, Zurich). Fly stocks were kept at 18°C, while the crosses - at 25°C.

Microscopy and image analysis. Images were obtained with Nikon SMZ1500 and Leica MZ16 F.

Images were deconvolved with Helicon Focus software (HeliconSoft).

RNA. Total RNA isolation was performed with TRI-reagent (Sigma) according to manufacturer

recommendations. The poly(A)-enrichment was performed with NEXTflex™ Poly(A) kit purchased from

BIOO Scientific. RT reactions were performed with Maxima RT (ThermoFisher Scientific), while qPCR

reactions - with SensiFAST™ SYBR® No-ROX Kit (Bioline). For oligonucleotide sequences see Table

S1.

Northern blot was performed as described[42]. 10 ug of total RNA per sample was subjected to

electrophoresis on 1% (w/v) agarose gels containing 1.1 mM formaldehyde. The RNA was transferred

overnight by capillarity to an Amersham Hybond-N+ membrane (GE Healthcare Life Sciences) and

covalently bound to the membrane using a Stratalinker UV crosslinker. Northern blots were hybridized

with DNA probes generated by a random-primed labeling reaction with use of Amersham Ready-To-Go

DNA Labelling Beads (GE Healthcare Life Sciences) supplemented by [α-^{32}P]dCTP and a corresponding

PCR fragment as a template. Membranes were exposed overnight to a Phosphor Imager screen at room

temperature.

Acknowledgements

S.E.D. is a member of the Interdisciplinary Scientific and Educational School of Moscow University

"Molecular Technologies of the Living Systems and Synthetic Biology".

References

1. Jennings, B. H. Drosophila – a versatile model in biology & medicine. *Materials Today* **14**, 190–195

(2011).

2. Mirzoyan, Z. *et al.* Drosophila melanogaster: A Model Organism to Study Cancer. *Frontiers in*

Genetics **10**, 51 (2019).

3. Tolwinski, N. S. Introduction: Drosophila-A Model System for Developmental Biology. *J Dev Biol* **5**, 9

(2017).

4. Ikonomou, L., Schneider, Y.-J. & Agathos, S. Insect cell culture for industrial production of

recombinant proteins. *Applied microbiology and biotechnology* **62**, 1–20 (2003).

5. Maiorella, B., Inlow, D., Shauger, A. & Harano, D. Large-scale insect cell-culture for recombinant

protein production. *Bio/technology* **6**, 1406–1410 (1988).

6. Palomares, L. A., Estrada-Moncada, S. & Ramírez, O. T. Production of recombinant proteins.

Recombinant gene expression 15–51 (2004).

7. Patel, S., Suleria, H. A. R. & Rauf, A. Edible insects as innovative foods: Nutritional and functional

assessments. *Trends in Food Science & Technology* **86**, 352–359 (2019).

8. van Huis, A. & Oonincx, D. G. The environmental sustainability of insects as food and feed. A review.

Agronomy for Sustainable Development **37**, 1–14 (2017).

9. Wang, Y.-S. & Shelomi, M. Review of black soldier fly (Hermetia illucens) as animal feed and human

food. *Foods* **6**, 91 (2017).

10. González, M. *et al.* Generation of stable Drosophila cell lines using multicistronic vectors. *Scientific*

Reports **1**, 75 (2011).

11. Brown, J. D. & Ryan, M. D. Ribosome "Skipping": "Stop-Carry On" or "StopGo" Translation. in

Recoding: Expansion of Decoding Rules Enriches Gene Expression (eds. Atkins, J. F. & Gesteland, R.

F.) 101–121 (Springer New York, 2010). doi:10.1007/978-0-387-89382-2_5.

12. Elroy-Stein, O., Fuerst, T. R. & Moss, B. Cap-independent translation of mRNA conferred by

encephalomyocarditis virus 5'sequence improves the performance of the vaccinia

virus/bacteriophage T7 hybrid expression system. *Proceedings of the National Academy of Sciences*

86, 6126–6130 (1989).

13. Jaafar, Z. A. & Kieft, J. S. Viral RNA structure-based strategies to manipulate translation. *Nature*

Reviews Microbiology **17**, 110–123 (2019).

14. Jan, E. Divergent IRES elements in invertebrates. *Virus research* **119**, 16–28 (2006).

15. Mailliot, J. & Martin, F. Viral internal ribosomal entry sites: four classes for one goal. *Wiley*

Interdisciplinary Reviews: RNA **9**, e1458 (2018).

16. Sorokin, I. I. *et al.* Non-Canonical Translation Initiation Mechanisms Employed by Eukaryotic Viral

mRNAs. *Biochemistry (Moscow)* **86**, 1060–1094 (2021).

17. Terenin, I. M., Smirnova, V. V., Andreev, D. E., Dmitriev, S. E. & Shatsky, I. N. A researcher's guide to

the galaxy of IRESs. *Cellular and Molecular Life Sciences* **74**, 1431–1455 (2017).

18. Sasaki, J. & Nakashima, N. Translation initiation at the CUU codon is mediated by the internal

ribosome entry site of an insect picorna-like virus in vitro. *Journal of virology* **73**, 1219–1226 (1999).

19. Wilson, J. E., Pestova, T. V., Hellen, C. U. & Sarnow, P. Initiation of protein synthesis from the A site

of the ribosome. *Cell* **102**, 511–520 (2000).

20. Mizuguchi, H., Xu, Z., Ishii-Watabe, A., Uchida, E. & Hayakawa, T. IRES-Dependent Second Gene

Expression Is Significantly Lower Than Cap-Dependent First Gene Expression in a Bicistronic Vector.

Molecular Therapy **1**, 376–382 (2000).

21. Underhill, M. F., Smales, C. M., Naylor, L. H., Birch, J. R. & James, D. C. Transient Gene Expression

Levels from Multigene Expression Vectors. *Biotechnology Progress* **23**, 435–443 (2007).

22. Brand, A. H. & Perrimon, N. Targeted gene expression as a means of altering cell fates and

generating dominant phenotypes. *development* **118**, 401–415 (1993).

23. Reinganum, C., O'Loughlin, G. T. & Hogan, T. W. A nonoccluded virus of the field crickets

Teleogryllus oceanicus and T. commodus (Orthoptera: Gryllidae). *Journal of Invertebrate Pathology*

16, 214–220 (1970).

24. Jousset, F.-X., Bergoin, M. & Revet, B. Characterization of the Drosophila C virus. *Journal of General*

Virology **34**, 269–283 (1977).

25. Bischof, J., Maeda, R. K., Hediger, M., Karch, F. & Basler, K. An optimized transgenesis system for

Drosophila using germ-line-specific φC31 integrases. *Proceedings of the National Academy of*

Sciences **104**, 3312–3317 (2007).

26. Metzstein, M. M. & Krasnow, M. A. Functions of the nonsense-mediated mRNA decay pathway in

Drosophila development. *PLoS genetics* **2**, e180 (2006).

27. Nelson, J. O., Moore, K. A., Chapin, A., Hollien, J. & Metzstein, M. M. Degradation of Gadd45 mRNA
by nonsense-mediated decay is essential for viability. *Elife* **5**, e12876 (2016).

28. Bonning, B. C. & Miller, W. A. Dicistroviruses. *Annual review of entomology* **55**, 129–150 (2010).

29. Withers, J. B. & Beemon, K. L. The structure and function of the rous sarcoma virus RNA stability
element. *Journal of cellular biochemistry* **112**, 3085–3092 (2011).

30. Fontaine, K. A. *et al.* The cellular NMD pathway restricts Zika virus infection and is targeted by the
viral capsid protein. *MBio* **9**, e02126-18 (2018).

31. Ramage, H. R. *et al.* A combined proteomics/genomics approach links hepatitis C virus infection with
nonsense-mediated mRNA decay. *Molecular cell* **57**, 329–340 (2015).

32. Balistreri, G. *et al.* The host nonsense-mediated mRNA decay pathway restricts mammalian RNA
virus replication. *Cell host & microbe* **16**, 403–411 (2014).

33. Contu, L., Balistreri, G., Domanski, M., Uldry, A.-C. & Mühlemann, O. Characterisation of the Semliki
Forest Virus-host cell interactome reveals the viral capsid protein as an inhibitor of nonsense-
mediated mRNA decay. *PLoS pathogens* **17**, e1009603 (2021).

34. Barker, G. F. & Beemon, K. Rous sarcoma virus RNA stability requires an open reading frame in the
gag gene and sequences downstream of the gag-pol junction. *Mol Cell Biol* **14**, 1986–1996 (1994).

35. Balagopal, V. & Beemon, K. L. Rous Sarcoma Virus RNA Stability Element Inhibits Deadenylation of
mRNAs with Long 3'UTRs. *Viruses* **9**, 204 (2017).

36. Behm-Ansmant, I. & Izaurralde, E. NMD in Drosophila: A Snapshot into the Evolution of. *Nonsense-
Mediated mRNA Decay* 153 (2006).

37. Pfeiffer, B. D. *et al.* Refinement of tools for targeted gene expression in Drosophila. *Genetics* **186**,
735–755 (2010).

38. Zufferey, R., Donello, J. E., Trono, D. & Hope, T. J. Woodchuck hepatitis virus posttranscriptional
regulatory element enhances expression of transgenes delivered by retroviral vectors. *Journal of
virology* **73**, 2886–2892 (1999).

39. Huang, M. T. & Gorman, C. M. Intervening sequences increase efficiency of RNA 3'processing and
accumulation of cytoplasmic RNA. *Nucleic Acids Research* **18**, 937–947 (1990).

40. Wodarz, A., Hinz, U., Engelbert, M. & Knust, E. Expression of crumbs confers apical character on
plasma membrane domains of ectodermal epithelia of Drosophila. *Cell* **82**, 67–76 (1995).

41. Pignoni, F. & Zipursky, S. L. Induction of Drosophila eye development by decapentaplegic.
Development **124**, 271–278 (1997).

42. Saleh, M.-C. *et al.* Antiviral immunity in Drosophila requires systemic RNA interference spread.

Nature **458**, 346–350 (2009).

Figure legends.

Fig 1. Overexpression of first cistron products in IRES-containing constructs. A. Scheme of the

constructs used. **B.** Flies expressing *da:Gal4* and a corresponding construct (two copies of each) were

examined with epifluorescent microscopy. All images are taken under the same conditions. Images in the

bottom row are taken in the daylight with no excitation light.

Fig 2. mRNAs made with IRES-containing constructs are NMD-protected. A.-E. Effects of Upf^{25G}

mutation only slightly affect levels of IRES-containing constructs. **A.** The crossing scheme. Hemizygous

Upf^{25G} or control males were selected for the analysis. **B.** NMD-null animals display elevated expression

of the control construct but not of IRES constructs. All images are taken under the same conditions. **C.**

RT-qPCR analysis of the RNA samples extracted from the animals shown in panel B. Each bar represents

an average of 3 biological replicates. Numbers were first normalized with *rpl12* qPCR counts. Read-outs

from wild-type animals expressing the control constructs are set as ones. A disparity in concentrations

between mCherry and EGFP sequences in IRES-containing construct is evident. **D.- E.** siRNA-mediated

knockdown of NMD activity only slightly affects IRES-containing constructs' expression. **D.** A crossing

scheme of eye-specific siRNA-mediated knockdown of *Upf1* and *Upf2*, the two critical components of the

NMD machinery. Wild-type flies were used as control. **G.** RT-qPCR analysis performed essentially as in

panel E with normalization made with Gal4 readouts instead of *rpl12*. Levels of the control construct in

the wild-type background are set one. mRNAs produced with IRES-containing constructs display only a

minor upregulation compared to control. An average of three independent replicates is shown. **F.** The

Northern blot of the samples from panel **C**. Denatured RNA resolved in agarose gel was probed with

mCherry or EGFP [α-^{32}P]dCTP-labeled fragments. IRES-containing samples display truncated RNA

isoform that accumulates at high levels. The full-length CrPV transcript levels were low and did not

respond to NMD activity, while the full-length DCV transcript levels also accumulated to higher levels

even in the presence of functional NMD. A representative image out of three replicates is shown. FL –

full-length transcript, TR – truncated transcript. rRNA was used as a size marker. **G.** Quantification of

band intensities from panel **F**. Each bar represents an average of 3 biological replicates. Samples from

wild-type animals expressing the control constructs are set as ones.

Fig 3. Truncated transcripts are polyadenylated since they are retained in the mixture upon isolation of the

polyA fraction with the use of poly-T magnetic beads. Normalization as in panel **2C**. An average of three

biological replicates is shown.

Fig 4. A conceptual scheme of IRES-mediated overexpression. A. SV40 polyadenylation signal
enhances transcription but also makes the transcript sensitive to NMD. **B.** Intragenic IRES position within
the virus genome provides an evolutionary rationale for the presence of NMD inhibitors within the
IRESes. **C.** Cryptic polyadenylation signal within the IRES results in transcript truncation and
stabilization. Putative NMD inhibitors might reside within IRESes, thus explaining decreased sensitivity
to NMD inhibition.
294
295