Hearing elliptic movements reveals the imprint of action on prototypical geometries

Etienne Thoret[1,3*], Mitsuko Aramaki[1], Lionel Bringoux[2], Sølvi Ystad[1], and Richard Kronland-Martinet[1]

[1] Aix Marseille Univ, CNRS, UMR7061 Perception Representations Image Sound Music (PRISM), Marseille, France

[2] Aix Marseille Univ, CNRS, UMR7287 Institut des Sciences du Mouvement (ISM), Marseille, France

[3] Institute of Language, Communication & the Brain, Marseille, France

[*] thoret@prism.cnrs.fr

Abstract

Within certain categories of geometric shapes, prototypical exemplars that best characterize the category have been evidenced. These geometric prototypes are classically identified through the visual and haptic perception or motor production and are usually characterized by their spatial dimension. However, whether prototypes can be recalled through the auditory channel has not been formally investigated. Here we address this question by using auditory cues issued from timbre-modulated friction sounds evoking human drawing elliptic movements. Since non-spatial auditory cues were previously found useful for discriminating distinct geometric shapes such as circles or ellipses, it is hypothesized that sound dynamics alone can evoke shapes such as an exemplary ellipse. Four experiments were conducted and altogether revealed that a common elliptic prototype emerges from auditory, visual, and motor modalities. This finding supports the hypothesis of a common coding of geometric shapes according to biological rules with a prominent role of sensory-motor contingencies in the emergence of such prototypical geometry.

Introduction

Our perceptual system stores and categorizes objects from our surroundings around canonical items, called prototypes, that are the most representative of their category. Geometric prototypes have been viewed as mainly emerging from visual experiences (Rosch, 1973; Rosch & Mervis, 1975; Feldman, 2000; Kalénine et al. 2011; Theurel et al. 2012), via the haptic sensory channel (Theurel et al., 2012; Kalénine et al. 2011) and via a motor restitution (Feldman, 2000; Kalénine et al., 2013). Alternatively, the possibility to recall prototypes through visual and haptic modalities or motor output may suggest common processing of these specific shapes. Whether such prototypes only rely on spatial cues that are present in the visual, haptic, and kinetic domains is still questioned. Could such geometric prototypes also be recalled through auditory stimuli that are solely based on dynamic motor cues? Answering this question would provide a new path to understanding the influence of the motor system on human learning, memory, and cognition. This would indeed suggest that human movement dynamics and more generally the sensory-motor contingencies influence the structuration of geometric shape categories during development.

Interestingly, studies revealed the proficient role of the auditory modality to perceive movements and shapes through timbre variations of monophonic sounds (Merer et al. 2008; 2013). More strikingly, Thoret et al. (2014) demonstrated that auditors listening to synthesized monophonic friction sounds corresponding to those produced by the pencil of someone drawing on a paper, were able to recognize specific kinematics characterizing biological motion, in particular the 1/3 power law linking the tangential velocity of the hand movement to the curvature of the drawn shape (Lacquaniti et al., 1983). In addition, they were even able to discriminate geometric shapes, such as a circle, an ellipse, and a line, simply by listening to synthetic friction sounds in which timbre variations revealed the velocity profile of the drawing movement. In follow-up studies, we demonstrated that these acoustic

variations may even distort the visuomotor coupling of biological motions (Thoret et al,

2016a, 2016b). Taken together, these studies suggest that simple geometric shapes can be

evoked through the auditory channel and that the auditory modality may play a significant

role in the perception and production of geometric shapes.

Here we investigated the hypothesis that geometrical prototypes can be recalled

through the auditory modality employing timbre variations of friction sounds evoking the

velocity of biological movements. We focused on a particular shape category, ellipses, that

encompass any closed shape of a conic section contained between a line and a circle. From a

geometric point of view, an ellipse is principally described by its eccentricity representing its

flatness. The eccentricity is a number comprised between 0 and 1: the flatter the ellipse, the

higher the eccentricity. The line and the circle are two specific cases which eccentricities

equal 1 and 0 respectively. From a dynamic point of view and concerning the 1/3 power law,

the accelerations of an elliptic movement increase as the distance between the focal points of

the ellipse increases (i.e., when the ellipse tends towards a line). Ellipses can be distinguished

by ear from circles and lines and do not involve discontinuity movements. Hence, this

geometric shape has been chosen for the present study.

Based on the dynamic model of biological motion described below, four experiments

were designed to examine how participants assessed the prototypical ellipse through different

modalities. The first three experiments aimed at highlighting the geometric prototype from

visual and motor restitutions. The fourth experiment was the cornerstone of this series of

experiments and focused on sounds' ability to evoke an elliptic prototype based on the

kinematics underlying the drawn ellipse. Consistency between this auditory prototypical

ellipse and the visual and motor prototypes would support a common encoding of prototypical

shapes.

Before presenting the four experiments, the dynamic model of elliptic motions used in

the four experiments will be described in the following section. The results of these

experiments are presented together.

A model of biological elliptic motor dynamics

Biomechanical mechanisms involved in graphical production as handwriting have been

extensively investigated. It has been shown that dynamic and geometric properties of elliptic

motions can be modeled by explicit equations. The dynamic approach of movement

production supports the idea that planar hand movements can be modeled by two harmonic

oscillators (x(t), y(t)) whose frequencies, amplitudes and phases evolve over time

(Hollerbach, 1981) with the following system:

89
$$\begin{cases} x(t) = A_x(t)\cos\big(\omega_x(t) + \phi_x(t)\big) \\ y(t) = A_y(t)\cos\big(\omega_y(t) + \phi_y(t)\big) \end{cases}$$

90   where A_x and A_y are the amplitudes, and ω_x and ω_y the frequencies, and ϕ_x and ϕ_y the phases

91   of the two oscillators. In the case of periodic elliptic motions, this model can be simplified by

92   equaling the amplitudes and frequencies of the oscillators ($A(t) = A_x = A_y = A$ and $\omega(t) = \omega_x =$

93      $\omega_y = \omega$):

94
$$\begin{cases} x(t) = A\cos(\omega t) \\ y(t) = A\cos(\omega t + RP) \end{cases} \tag{1}$$

where A is the amplitude, $\omega = \dfrac{2\pi}{T}$ the frequency with T the period of the motion, and RP $= \phi_x$

$- \phi_y$ the relative phase between the oscillators $x(t)$ and $y(t)$.

Hence, this system co-defines the motion dynamics and the geometry, but may also be

used to parameterize only the geometry of the entire ellipse whether or not the ellipse is

considered dynamically. Practically, the eccentricity e of the ellipse and the relative phase RP

are linked by the following relations demonstrated in Appendix A:

101
$$\begin{cases} RP = 2\arctan\sqrt{1-e^2} \\ e = \sqrt{1-\tan^2\left(\dfrac{RP}{2}\right)} \end{cases} \quad (2)$$

It should be noted that this model complies with the biological rules, namely the 1/3 power

law (Lacquaniti et al., 1983), as demonstrated in Appendix A.

This model then enables the generation of biological elliptic motions whose trajectory

can be continuously morphed from a line (RP = 0°) to a circle (RP = 90°) by simply acting on

RP (see Figure 1). It will be used in the four following experiments in order to generate visual

and auditory stimuli complying with biological motion. Motor productions of Experiment 2

will also be analyzed regarding this model.

Finally, the tangential velocity profile was generated according to the equations (1)

and can be explicitly written as follows:

111
$$v(t) = A\frac{2\pi}{T}\sqrt{\sin^2\left(\frac{2\pi}{T}t\right) + \sin^2\left(\frac{2\pi}{T}t + RP\right)} \quad (3)$$

112   Although the mapping was arbitrarily fixed during the experiment, it is noticeable that it also

113   varied according to the period T and the amplitude A of the motion.

RP	0°	15°	30°	45°	60°	75°	90°
e	1	.99	.96	.91	.82	.64	0

116   *Figure 1*. Continuum of different ellipse shapes from a line (left) to a circle (right) with the

117   corresponding relative phases RP and eccentricities e.

Experiment 1 – Static Visual Output

The first experiment aimed to evaluate the prototype of the ellipse from visual restitution (i.e.,

visual output). To that aim, an adjustment protocol was established. Participants were asked

to adjust the eccentricity of static ellipses on a screen to evoke the most representative shape

of this geometric category.

Methods

Participants. Twenty right-handed participants (8 women) with an average age of 30.6

years (SD = 12.8) voluntarily took part in the experiment. All the participants were naive to

the purpose of the experiment. As in all the following experiments, participants gave their

informed consent before the study, and the experiment was approved by the Aix-Marseille

University ethical committee. For this experiment and the three subsequent ones, participants

were selected from various groups, including personal acquaintances, students, engineers,

researchers, and professors at the CNRS Campus Joseph Aiguier in Marseille, as well as

individuals from the Sport Science Faculty of Aix-Marseille University in Marseille, France.

Stimuli. The visual stimuli were white static ellipses displayed on a black background

with different sizes and eccentricities according to the dynamic model previously introduced

(equations (1)). Three sizes defined by the amplitudes $A_1 = 3cm$, $A_2 = 5cm$, $A_3 = 10cm$ were

chosen. The ellipses were rotated counterclockwise by 45° to conform to the preferential

drawing inclination of an ellipse for right-handed persons (Danna et al., 2011).

Apparatus. The participants sat in front a computer screen (DELL 1907fp) with a

resolution of 1280 x 1024 pixels and a frame rate of 60 Hz. The ellipses were displayed at the

center of the screen and the interface was programmed with Max/MSP software

(http://cycling74.com/). Participants modified the ellipse eccentricities by using a MIDI

AKAY MPK keyboard. The experiment took place in a lighted room.

144   *Task.* The participants were asked to adjust the eccentricity of the static ellipse

145 displayed on the screen so as to set the most representative geometric shape of this category.

146 The notion of the most representative ellipse was explained in French to the participants by

147 the following sentences (here translated in English): *Ellipses correspond to any closed shape*

148 *between a line and a circle. When you imagine an ellipse, you may have one particular*

149 *elliptic shape in mind. You will adjust the ellipse on the screen to display the ellipse you think*

150 *is the most prototypical.* The adjustment protocol was based on the one proposed by Carlyon

151 et al. (2010). The ellipse eccentricity was adjusted thanks to 6 different keys defined on the

152 keyboard: "<<<" – "<<" – "<" and ">" – ">>" – ">>>". Depending on the selected key, the

153 ellipse eccentricity was modified with different step sizes. The higher the number of arrows

154 the greater the step size, *i.e.* the modification of the eccentricity. To avoid a possible non-

155 sensorial bias due to the orientation of the arrows (right or left), the action of the keys on the

156 eccentricity differed across participants. Hence, for half of the participants, arrows that

157 pointed to the right increased the eccentricity and for the other half they had the opposite

158 effect. In addition, the step sizes were not the same in both directions. Hence, for a given

159 participant, all the arrows in one direction (right or left) increased for instance the eccentricity

160 by steps of {.01; .02; .03} while the others (left or right) decreased the eccentricity by steps of

161 {-.1; -.001; -.005}. Finally, 11 repetitions for each size starting from 11 different initial

162 eccentricities equally distributed between 0 and 1 were executed. The experiment was then

163 composed of 33 trials, i.e. 3 {Sizes} x 11 {repetitions}, presented according to two pseudo-

164 random series counterbalanced across the participants. The participants were prompted to

165 explore the whole range of possibilities with the large arrows and to refine their adjustment

166 with the smaller ones. For each participant, 33 final eccentricities were collected and the

167 median was computed.

Experiment 2 – Motor Output

The goal of this experiment was to evaluate whether a prototypical eccentricity of an

ellipse can be elicited by a motor output. A motor production task was set up during which

the participants were asked to draw the ellipse which best represents this geometric shape

category.

Methods

Participants. Twenty participants (2 left-handed - 5 women) with an average age of

39.6 years (SD = 11.5) voluntary took part in the experiment and were naive to the purpose of

the study. None of them participated in Experiment 1.

Apparatus. The participants sat in front of a Wacom Intuos5 graphic tablet enabling to

record graphic movements with a spatial precision of 5.10^{-3}mm and a sample rate of 129 Hz.

The data were recorded and collected using an interface programmed with the Max/MSP

software. The experiment was conducted in a lighted room.

Task. The participants were asked to repeatedly and continuously draw the most

representative ellipse of this geometric shape category on the graphic tablet during 50

seconds. The notion of the most representative ellipse was explained with the same sentence

as in Experiment 1. Participants saw their hands during the recording, but no trace was visible

on the graphic tablet. The experiment comprised three sessions of 50 seconds. The

participants were asked to draw *small*, *intermediate*, and *large* ellipses in two different orders

counterbalanced across participants: 1) small – 2) intermediate – 3) large, or conversely. No

template of the ellipses was presented, but the participants could train in advance by drawing

the 3 different ellipses on the graphic tablet during a session preceding the experiment.

Data analysis. For each participant, recordings of the sampled coordinates ($x(t)$, $y(t)$)

of the stylus on the graphic tablet were collected. To eliminate the numerical noise, the raw

data were smoothed with a Savitsky-Golay filter (Savitsky and Golay, 1964) with a 43

samples window and a 3rd-order interpolation. This is equivalent to a low-pass filter with a

cutoff frequency of 8 Hz. A high-pass filter (Butterworth) with a cutoff frequency of .2 Hz

was also applied to eliminate the spatial drift of the participants' hands during their

movements.

The geometric characteristics of the drawn ellipses were analyzed according to the

relative phase. Computing the Hilbert transforms of $x(t)$ and $y(t)$ enabled to estimate the

relative phase RP between two oscillators (Panter, 1965; Smith and Mersereau, 1991) with the

following formula: $RP(t) = |\tilde{x}(t) - \tilde{y}(t)|$, where $\tilde{x}$ and $\tilde{y}$ are the unwrapped Hilbert

transforms of x and y. The median of the relative phases RP were then computed for each

recording and then transformed into eccentricity. For each participant, 3 final eccentricities

were collected and the median was computed.

Experiment 3 – Dynamic Visual Output

Here, the Experiment 1 was reproduced with dynamic visual stimuli. The participants

were asked to calibrate the eccentricity of the elliptic trajectory of a moving spotlight to create

the most representative ellipse.

Methods

Participants. Twenty participants (1 left-handed; 3 women) with an average age of 32

years (SD = 11.4) voluntary took part in the experiment. None of these participants took part

in Experiments 1 & 2.

Stimuli. The visual stimuli were white spotlights displayed on a black background

moving on elliptic paths of different sizes and with different periods according to the dynamic

model defined in equation (1). As in Experiment 1, the elliptic paths were rotated

counterclockwise by 45°. Three periods of the spotlight motion (T_1 = 1.2s, T_2 = 1.5s, T_3 =

1.8s) inducing different spotlight speeds were chosen.

Task. The adjustment protocol and the task were the same as in Experiment 1. For

each (size) x (period) pair, 11 repetitions were performed starting from 11 different initial

eccentricity values equally distributed between 0 and 1. The experiment was composed of 99

trials, i.e. 3 {sizes} x 3 {periods} x 11 {repetitions}, presented according to two pseudo-

random series counterbalanced across the participants. For each participant, 99 final

eccentricities were collected and the median was computed.

Apparatus. The apparatus was the same as in Experiment 1.

Experiment 4 – Auditory Output

Here, participants were submitted to auditory stimuli generated with the dynamic

characteristics of the motor performances recorded in experiment 2 and were asked to modify

this dynamic to evoke a friction sound that conveys the most representative elliptic movement

exclusively from the auditory modality. By using monophonic sounds, only dynamic

information was transmitted to the participants, as opposed to the visual cases in which both

spatial and dynamic cues were contained in the stimuli. Hence, this experiment intended to

show whether geometric prototypes could emerge from the auditory stimulations, which

would signify that spatial information is not strictly needed to evoke a prototype.

Methods

Participants. Twenty participants (9 women) with an average age of 32.1 years (SD =

9.56) voluntary took part in the experiment. None of them participated to the experiments 1, 2

& 3.

Stimuli. Stimuli were monophonic synthetic friction sounds simulating the motion of a

pencil on a rough surface, e.g., a piece of paper. A physically based model which simulates

the physical sound source as the result of successive impacts of a plectrum, here a pencil nail,

on the asperities of a surface, was used (Conan et al., 2014). The surface roughness is

modeled by a noise reflecting the different heights of the surface asperities. Movements on

this surface is then simulated processing this noise according to the velocity profile of the

movement: the faster the movement the more impacts there are. From a signal processing

point of view, such an operation is equivalent to a low-pass filter on the noise with a central

frequency directly linked to the movement velocity. The filtered noise is finally convolved

with an impulse response simulating the resonance of an object such as a table. To control the

synthesis model, we used the 60 motions (3 recordings x 20 participants) recorded in the

motor experiment (Experiment 2).

Task. The participants were asked to modify the sound to evoke an elliptic motion for

which the trajectory was the most representative of an ellipse. They were implicitly adjusting

the delay between the recorded oscillators which implicitly modify the ellipse's roundness.

For each of the 60 movements to adjust, 4 periods of the initial recording were used to

generate the stimuli. The participants were informed when they reached limits, i.e. 0 or 1

eccentricity, and the adjustment protocol was inspired by one proposed by Carlyon and

colleagues (Carlyon et al., 2010). The eccentricity of the ellipse could be modified either with

a large or a small modification of the actual eccentricity by using top-down and left-right-

arrows respectively, by steps of .10/.3 and .05/.2 respectively. In order to avoid an

experimental bias, for each opposite key, e.g., left/right, the roundness was modified with a

different step so that participants could not count the number of taps to adjust the friction

sound. The participants were prompted to explore the whole range of possibilities with the

large arrows and to refine their adjustment with the smaller ones.

Apparatus. Sounds were presented through Sennheiser HD650 headphones and the

sample rate of the soundcard was 44100 Hz with 16-bit resolution. The friction sounds were

real-time synthesized with MATLAB software. The experiment was carried out in a quiet

room. The sound was modified by the participants using the keyboard of the computer.

Data analysis. For each participant, 60 eccentricities values were collected and their

median was computed to characterize the prototypical ellipse.

Results

Coherence between prototypes. We first investigated whether the four different

experiments led to four different prototypes. For each experiment, we ensured the normality

of the different distributions by using a Lilliefors test (all non-significant, Bonferroni

corrected α=.05/4=.0125). It must be noted that for the auditory experiment the Lilliefors test

was only marginally significant because of Bonferroni corrections (p=.0176). Therefore, in

the following we used both parametric and non-parametric approaches. Two statistical tests

were applied: (1) a non-parametric Kruskal-Wallis test with experiment as a factor (df=3; χ^2=

1.14; p=.76); (2) parametric Bonferroni corrected pairwise t-tests between the different

experiments. Results were all non-significant (*df*=19, motor vs. visual static: p=.33, BF_{10}=.35;

motor vs. visual dynamic: p=.84, BF_{10}=.23; motor vs. auditory: p=.92, BF_{10}=.23; visual static

vs. visual dynamic: p=.55, BF_{10}=.27; visual static vs. auditory: p=.25, BF_{10}=.42; visual

dynamic vs. auditory: p=.82, BF_{10}=.24) (Figure 2a). These tests thus supported an absence of

significant difference between the motor, the visual - static and dynamic - and the auditory

prototypes.   *Eccentricity of the prototypes.* Secondly, we investigated the eccentricity of the

prototypes for each experiment. The mean eccentricity values were: 1) motor: Mdn=.89,

iqr=.067, M=.89, SD=.046; 2) visual static: Mdn=.91, iqr=.062, M=.90, SD=.044; 3) visual

dynamic: Mdn=.91, iqr=.121, M=.88, SD=.086; 4) auditory: Mdn=.88, iqr=.049, M=.88,

SD=.061 (Figure 2a). We determined a statistically based confidence interval for each

prototype, one in each experiment. We linearly sampled the eccentricity range between 0 and

1 with 1000 equally spaced eccentricity values. We then ran 1000 one sample t-tests between

the data distribution of each experiment and the 1000 eccentricity values. The results of each

of the 1000 tests characterizes whether the distribution is different from the candidate

eccentricity. For each of the 1000 t-tests, the inverse of the bayes factor $BF_{10,inverse}=1/BF_{10}$

was stored. $BF_{10,inverse}$ can be interpreted exactly as BF_{10}, but for *no difference* with the null

hypothesis, the higher the $BF_{10,inverse}$ the less different the distribution with the candidate

eccentricity. We here assume that $BF_{10,inverse}>3$ characterizes a fair threshold for no difference

with the null hypothesis, i.e. the eccentricity value can be considered in the prototype range

(Jeffreys, 1961; Goodman, 1999; Held & Ott, 2016). As results distributions were globally

localized, the $1/BF_{10}$ curve is an inverse U-shape. For each experiment, we therefore used

$BF_{10,inverse}=3$ as a threshold to determine the lower and upper bound of the confidence

interval. We also computed the eccentricity ecc_{max} leading to the highest $BF_{10,inverse}$. This

provides a third estimation of the elliptic prototype eccentricity for each experiment. This

analysis provides the following results: motor: $ecc_{max}=.883$, $CI_{BF10,inverse}=[.869,.897]$ - visual

static: $ecc_{max}=.899$, $CI_{BF10,inverse}=[.892,.909]$ -   visual   dynamic: $ecc_{max}=.888$,

$CI_{BF10,inverse}=[.871,.905]$ - auditory: $ecc_{max}=.880$, $CI_{BF10,inverse}=[.870,.893]$. To investigate

specifically whether they differ from the known preferential ellipse of eccentricity .91, t-tests

between the distributions and the .91 value were conducted. Results revealed no significant

differences (*df*=19, Bonferroni corrections $\alpha=.05/4=.0125$, motor: p=.1, $BF_{10}=.82$ – visual

static: p=.34, $BF_{10}=.354$ – visual dynamic: p=.25, $BF_{10}=.42$ - auditory: p=.05, $BF_{10}=1.37$).

Altogether, these analyses revealed that the experiments provide comparable prototypes, with

an average eccentricity of .887. These prototypes are slightly below the preferential

intermediate ellipse of eccentricity .91 observed in the literature (see General Discussion).

Figure 2. Averaged eccentricities obtained from the four experiments. The boxplot represents

the median (dot), the 25[th] and 75[th] percentiles (the bottom and top edges of the box). The

whiskers (vertical dashed line) extend to the most extreme data points below 1.5 the

interquartile range. The dashed line indicates the value .91 corresponding to the elliptic

movement attractor.

General Discussion

In this study, we aimed at investigating whether a common prototype of elliptic shapes

elicited across modalities, especially from auditory stimuli, could be identified. We compared

the restituted geometry through different outputs (visual, motor and auditory) across a series

of four experiments. A common elliptic prototype characterized by an eccentricity of .887

was uncovered, hence demonstrating the common encoding of this prototype.

This complements knowledge from previous studies on the existence of prototypical

geometric shapes (Rosch, 1973, 1975; Feldman, 2000; Kalénine et al., 2013). This is also in

line with those from Wamain et al. (2011), who highlighted that the ellipse drawn

spontaneously has a relative phase slightly below 45°, i.e. an eccentricity slightly below.91,

and with those from pure motor experiments aiming at revealing the preferentially drawn

ellipse (Dounskaia et al., 2000; Sallagoïty et al., 2004; Danna et al., 2011) corresponding to

stable states, so-called motor attractors (Kelso, 1986) (cf. Figure 2). Our results showed that

the perceptual prototype – either visual or auditory – is coherent with the motor one, with an

eccentricity value slightly below the intermediate ellipse of .91. This eccentricity difference

may arise from the intention of the participants which were different because of a different

task. Here we asked them to draw the most prototypical ellipse, which wasn't present at all in

the spontaneously drawn ellipses asked in Wamain et al. (2011). This slight difference

suggests that we cannot obviously exclude that other cue also might have shaped the

geometry of the prototype than motor constraint such as visual experience. Interestingly,

when superimposing the prototypical ellipse obtained in the present study with the

prototypical rectangle highlighted by Kalénine et al. (2013), common proportions can be

observed. The length to width ratio for prototypical rectangles and the semi-major to semi-

minor axis ratio for prototypical ellipses are both close to 2.3, which corresponds to .9 in term

of eccentricity.

Above all, these results suggest a common coding of prototypical shape geometry and

more generally further support the role of the sensorimotor loop on our perceptual processes

(O'Regan and Noë, 2001; Varela et al., 1992; O'Regan, 2011). Interestingly,

neuropsychological studies demonstrated that motor schemes are re-activated during

perceptual processes through different sensory modalities (Chao and Martin, 2000; Creem-

Regehr and Lee, 2005; Grafton et al., 1997; Grèzes and Decety, 2002; Kohler et al., 2002;

Bangert et al., 2006; Zatorre et al., 2007), and through motor imagery (for a review see

Jeannerod, 1995 or Grosprêtre, Ruffino and Lebon, 2015). Concerning handwriting, it has

been shown that seeing a letter activates the cortical processes involved when producing the

corresponding script (Longcamp et al., 2003; James and Gauthier, 2006; Longcamp et al.,

2006). Similarly, perceiving a geometric shape may share processes involved when we are

drawing it. In Experiment 4, an elliptic prototype that did not differ from the visual and motor

outputs was elicited from monophonic sounds that only carried dynamic cues. This salient

result shows that the elliptic prototype is not only geometric but also can be transmitted

through a dynamic dimension. Hence without rejecting the necessity of visual experience in

the emergence of prototypical shapes (Theurel et al., 2011; Kalénine et al., 2011), our data

suggest that the prototypical shape is encoded through a common process based on the

underlying covariation between biological kinematics and shape geometry which

characterizes the corresponding drawing movement (Lacquaniti et al., 1983) and its

perception through several sensory channels (Viviani et al., 1992; Viviani and Stucchi, 1989;

Viviani et al., 1987; Thoret et al., 2014). This unified percept (Hommel et al., 2001) can then

be recalled even through the auditory modality, which does not provide any spatial cues.

Further experiments using the same auditory based paradigm, in particular with other

geometric shapes, are needed to precisely assess the central role of the sensorimotor loop in

the emergence of prototypical shape geometries.

Limitations & Perspectives

This study has mainly two methodological limitations. Firstly the investigation was made at

group level and subjects were different in each experiment. This allowed us to strikingly

highlight the existence of a common prototype between a set of different participants in each

modality. This was the main hypothesis of this study and the reason why the elliptic prototype

was primarily investigated at group level. It would be of interest to investigate this issue at

subject level to exclude that such a prototype may vary across modalities for a given set of

participants. However, such a subject level investigation would require a different

experimental procedure as one experiment might influence the result of the others. In this case

it would not be possible to completely disentangle a potential priming between them.

Secondly, the experiment was performed in controlled laboratory experimental conditions

with a population of N=80 (4x20). The generalizability of the result could be improved by

running such a series of experiments on a larger number of participants with online testing

platforms. Lastly, and this is probably the most important conceptual limitation of this study,

we have here investigated the prototype of only one shape, the ellipse, which is a very specific

shape without any discontinuities, and which moreover can be drawn continuously. To

generalize our results, it would be relevant to run the experiment with a more complete set of

shapes such as rectangles or triangles, that have been subject to a certain number of studies in

the past. This would nevertheless require refinements in the sound modeling procedure due to

the discontinuities in the drawing movement of such shapes.

Authors contributions

Conceived and designed the experiments: ET MA LB SY RKM. Performed the experiments:

ET. Analyzed the data: ET MA LB SY RKM. Contributed reagents/materials/analysis tools:

ET. Wrote the paper: ET MA LB SY RKM

Declaration of interests

The authors declare no competing interests.

Acknowledgments

This work was funded by the French National Research Agency (ANR) under the MetaSon:

Métaphores Sonores (Sound Metaphors) project (ANR-10-CORD-0003) in the CONTINT

2010 framework and the SoniMove Project (ANR-14-CE24-0018). E.T. was funded through

an ILCB/BLRI grant no. ANR-16-CONV-0002 (ILCB), ANR-11-LABX-0036 (BLRI) and

the Excellence Initiative of Aix-Marseille University (A*MIDEX). The authors would like to

thank Thomas Bordonné and Joris Agator for their help running the experiments. E.T. is

thankful to Jeremy Danna for his thoughtful suggestions and guidance at the very beginning

of this project.

References

Aramaki, M., Besson, M., Kronland-Martinet, R., & Ystad, S. (2011). Controlling the
perceived material in an impact sound synthesizer. *IEEE Journal on Audio, Speech,*
and Language Processing, 19(2), 301-314

Bangert, M., Peschel, T., Schlaug, G., Rotte, M., Drescher, D., Hinrichs, H., Heinze, H. J. &
Altenmüller, E. (2006). Shared network for auditory and motor processing in
professional pianists: evidence from fMRI conjunction. *Neuroimage*, 30(3), 917-926.

Catavitello, G., Ivanenko, Y. P., Lacquaniti, F., & Viviani, P. (2016). Drawing ellipses in
water: evidence for dynamic constraints in the relation between velocity and path
curvature. *Experimental Brain Research*, 234(6), 1649-1657.

Carlyon, R. P., Macherey, O., Frijns, J. H. M., Axon, P. R., Kalkman, R. K., Boyle, P.,
Baguley, D. M., Briggs, J., Deeks, J. M., Briaire, J. J., Barreau,X., & Dauman, R.
(2010). Pitch Comparisons between Electrical Stimulation of a Cochlear Implant and
Acoustic Stimuli Presented to a Normal-hearing Contralateral Ear. *Journal of the*
Association for Research in Otolaryngology, 11(4), 625-640.

Chao, L. L., & Martin, A. (2000). Representation of manipulable man-made objects in the
dorsal stream. *Neuroimage*, 12(4), 478-484.

Conan, S., Thoret, E., Aramaki, M., Derrien, O., Gondre, C., Ystad, S., & Kronland-Martinet,
R. (2014). An Intuitive Synthesizer of Continuous-Interaction Sounds: Rubbing,
Scratching, and Rolling. *Computer Music Journal*, 38(4), 24-37.

Creem-Regehr, S. H., & Lee, J. N. (2005). Neural representations of graspable objects: are
tools special?. *Cognitive Brain Research*, 22(3), 457-469.

Danna, J., Athènes, S., & Zanone, P. G. (2011). Coordination dynamics of elliptic shape
drawing: Effects of orientation and eccentricity. *Human Movement Science*, 30(4),
698-710.

Danna, J., Paz-Villagrán, V., Gondre, C., Aramaki, M., Kronland-Martinet, R., Ystad, S., &

438   Velay, J. L. (2015). "Let Me Hear Your Handwriting!" Evaluating the Movement

439   Fluency from Its Sonification. *PLoS One*, 10(6), e0128388.

440   Dounskaia, N., Van Gemmert, A. W. A., & Stelmach, G. E. (2000). Interjoint coordination

441   during handwriting-like movements. *Experimental Brain Research*, 135(1), 127-140.

442   Feldman, J. (2000). Bias toward regular form in mental shape spaces. *Journal of

443   Experimental Psychology: Human Perception and Performance*, 26(1), 152-165.

444   Gaver, W. W. (1993). What in the world do we hear?: an ecological approach to auditory

445   event perception. *Ecological Psychology*, 5(1), 1-29

446   Giordano, B. L., Rocchesso, D., & McAdams, S. (2010). Integration of acoustical information

447   in the perception of impacted sound sources: the role of information accuracy and

448   exploitability. *Journal of Experimental Psychology: Human Perception and

449   Performance*, 36(2), 462-476.

450   Grafton, S. T., Fadiga, L., Arbib, M. A., & Rizzolatti, G. (1997). Premotor cortex activation

451   during observation and naming of familiar tools. *Neuroimage*, 6(4), 231-236.

452   Grèzes, J., & Decety, J. (2002). Does visual perception of object afford action? Evidence from

453   a neuroimaging study. *Neuropsychologia*, 40(2), 212-222.

454   Glasberg, B. R., & Moore, B. C. (1990). Derivation of auditory filter shapes from notched-

455   noise data. *Hearing Research*, 47(1), 103-138.

456   Goodman, S. N. (1999). Toward evidence-based medical statistics. 2: The Bayes factor.

457   *Annals of internal medicine*, 130(12), 1005-1013.

458   Grosprêtre, S., Ruffino, C., & Lebon, F. (2015). Motor imagery and cortico-spinal

459   excitability: A review. *European journal of sport science*, 1-8. doi:

460   10.1080/17461391.2015.1024756

Gygi, B., Kidd, G. R., & Watson, C. S. (2004). Spectral-temporal factors in the identification

of environmental sounds. *The Journal of the Acoustical Society of America*, 115(3),

1252-1265.

Held, L., & Ott, M. (2016). How the maximal evidence of p-values against point null

hypotheses depends on sample size. *The American Statistician*, 70(4), 335-341.

Hollerbach, J. M. (1981). An oscillation theory of handwriting. *Biological Cybernetics*, 39(2),

139-156.

Hommel, B., Müsseler, J., Aschersleben, G., & Prinz, W. (2001). Codes and their

vicissitudes. *Behavioral and Brain Sciences*, 24(05), 910-926.

James, K. H., Gauthier, I. (2006). Letter processing automatically recruits a sensory? Motor

brain network. *Neuropsychologia*, 44(14), 2937-2949

Jeffreys, H. (1961). The theory of probability. OUP Oxford.

Jeannerod, M. (1995). Mental imagery in the motor context. *Neuropsychologia*, 33 (11),

1419-1432

Kalénine, S., Pinet, L., & Gentaz, E. (2011). The visual and visuo-haptic exploration of

geometrical shapes increases their recognition in preschoolers. *International Journal*

of Behavioral Development, 35(1), 18-26.

Kalénine, S., Cheam, C., Izard, V., & Gentaz, E. (2013). Adults and 5‑year‑old children

draw rectangles and triangles around a prototype but not in the golden ratio. *British*

Journal of Psychology, 104(3), 400-412.

Kelso, J. A. S. (1986). Pattern formation in speech and limb movements involving many

degrees of freedom. *Experimental Brain Research*, 15, 105-128.

Kohler, E., Keysers, C., Umilta, M.A., Fogassi, L., Gallese, V., & Rizzolatti, G. (2002).

Hearing sounds, understanding actions: action representation in mirror neurons.

Science, 297, 5582, 846-848

Lacquaniti, F., Terzuolo, C., & Viviani, P. (1983). The law relating the kinematic and figural

487   aspects of drawing movements. *Acta Psychologica*, 54(1), 115-130

488   Li, X., Logan, R. J., & Pastore, R. E. (1991). Perception of acoustic source characteristics:

489   Walking sounds. *The Journal of the Acoustical Society of America*, 90(6), 3036-

490   3049.

491   Longcamp, M., Anton, J.L., Roth, M., & Velay, J.L. (2003). Visual presentation of single

492   letters activates a premotor area involved in writing. *Neuroimage*, 19(4), 1492-1500

493   Longcamp, M., Tanskanen, T., & Hari, R. (2006). The imprint of action: Motor cortex

494   involvement in visual perception of handwritten letters. *Neuroimage*, 33(2), 681-688

495   McAdams, S. (1993). Recognition of sound sources and events. In McAdams, S., & Bigand,

496   E. (Eds.): *Thinking in Sound: The cognitive psychology of human audition*, 146–198

497   Merer, A., Ystad, S., Kronland-Martinet, R. & Aramaki, M. (2008). Semiotics of sounds

498   evoking motions: Categorization and acoustic features. In Kronland-Martinet, R.,

499   Ystad, S., & Jensen, K., (Eds.): *CMMR 2007. Sense of Sounds*, 139–158. Springer,

500   LNCS

501   Merer, A., Aramaki, M., Ystad, S., & Kronland-Martinet, R. (2013). Perceptual

502   characterization of motion evoked by sounds for synthesis control purposes. *ACM*

503   *Transactions on Applied Perception (TAP)*, 10(1), doi:10.1145/2422105.2422106

504   Moore, B. C., & Glasberg, B. R. (1983). Suggested formulae for calculating auditory‐filter

505   bandwidths and excitation patterns. *The Journal of the Acoustical Society of*

506   *America*, 74(3), 750-753.

507   O'Regan, J. K., & Noë, A. (2001). A sensorimotor account of vision and visual consciousness.

508   *Behavioral and Brain Sciences*, 24(5), 939-973.

509   O'Regan, J. K. (2011). Why red doesn't sound like a bell: Understanding the feel of

510   consciousness. Oxford University Press.

Panter, P. F. (1965). Modulation, Noise, and Spectral Analysis. New York: McCraw-Hill.

Rosch, E. H. (1973). Natural categories. *Cognitive psychology*, 4(3), 328-350.

Rosch, E. H., & Mervis, C. B. (1975). Family resemblances: Studies in the internal structure

of categories. *Cognitive Psychology*, 7(4), 573-605.

Sallagoïty, I., Athènes, S., Zanone, P. G., & Albaret, J. M. (2004). Stability of coordination

patterns in handwriting: Effects of speed and hand. *Motor Control*, 8, 405-421.

Savitzky, A., & Golay, M. J. (1964). Smoothing and differentiation of data by simplified least

squares procedures. *Analytical Chemistry*, 36(8), 1627-1639.

Smith, M. J., & Mersereau, R. M. (1991). Introduction to digital signal processing: a

computer laboratory textbook. John Wiley & Sons, Inc.

Theurel, A., Frileux, S., Hatwell, Y., & Gentaz, E. (2012). The haptic recognition of

geometrical shapes in congenitally blind and blindfolded adolescents: is there a

haptic prototype effect. *PLoS one*, 7(6), e40251.

Thoret, E., Aramaki, M., Kronland-Martinet, R., Velay, J. L., & Ystad, S. (2014). From sound

to shape: auditory perception of drawing movements. *Journal of Experimental*

Psychology: Human Perception and Performance, 40(3), 983-994.

Thoret, E., Aramaki, M., Bringoux, L., Ystad, S., & Kronland-Martinet, R. (2016a). Seeing

Circles and Drawing Ellipses: When Sound Biases Reproduction of Visual Motion.

PLoS one, 11(4), e0154475.

Thoret, E., Aramaki, M., Bringoux, L., Ystad, S., & Kronland-Martinet, R. (2016b). When

eyes drive hand: Influence of non-biological motion on visuo-motor coupling.

Neuroscience Letters, 612, 225-230.

Varela, F. J., Thompson, E., & Rosch, E. (1992). The embodied mind. MIT Press.

Van Den Doel, K. and Kry, P.G. & Pai, D.K. (2001). FoleyAutomatic: physically-based

sound effects for interactive simulation and animation. *In Proceedings of the 28th*

annual conference on Computer graphics and interactive techniques ACM, 537-544

Viviani, P., Campadelli, P., & Mounoud, P. (1987). Visuo-manual pursuit tracking of human

two-dimensional movements. *Journal of Experimental Psychology: Human*

Perception and Performance, 13(1), 62-78.

Viviani, P., & Stucchi, N. (1989). The effect of movement velocity on form perception:

Geometric illusions in dynamic displays. *Attention, Perception, & Psychophysics*,

46(3), 266-274

Viviani, P., Baud-Bovy, G., & Redolfi, M. (1997). Perceiving and tracking kinesthetic

stimuli: Further evidence of motor–perceptual interactions. *Journal of Experimental*

Psychology: Human Perception and Performance, 23(4), 1232-1252.

Warren, W. H., & Verbrugge, R. R. (1984). Auditory perception of breaking and bouncing

events: a case study in ecological acoustics. *Journal of Experimental Psychology:*

Human Perception and Performance, 10(5), 704-712

Wamain, Y., Tallet, J., Zanone, P. G., & Longcamp, M. (2011). "Biological geometry

perception": visual discrimination of eccentricity is related to individual motor

preferences. *PLoS one*, 6(1), e15995.

Zatorre, R. J., Chen, J. L., & Penhune, V. B. (2007). When the brain plays music: auditory-

motor interactions in music perception and production. *Nature Reviews*

Neuroscience, 8(7), 547-558.