Title

In-vivo characterization of glutamine metabolism identifies therapeutic targets in clear cell renal cell carcinoma

Authors

Akash K. Kaushik[1], Amy Tarangelo[1], Lindsey K. Boroughs[1], Mukundan Ragavan[2], Yuanyuan Zhang[1], Cheng-Yang Wu[3], Xiangyi Li[1], Kristen Ahumada[1], Jui-Chung Chiang[1], Vanina T. Tcheuyap[4], Faeze Saatchi[4], Quyen N. Do[3], Cissy Yong[5], Tracy Rosales[1], Christina Stevens[4], Aparna D. Rao [6,7], Brandon Faubert[8], Panayotis Pachnis[1], Lauren G. Zacharias[1], Hieu Vu[1], Feng Cai[1], Thomas P. Mathews[1], Giannicola Genovese[9], Barbara S. Slusher[10], Payal Kapur[4], Xiankai Sun[3,4,11], Matthew Merritt[12], James Brugarolas[4], Ralph J. DeBerardinis[1,4,13,*]

Affiliations

[1] Children's Research Institute, University of Texas Southwestern Medical Center, USA

[2] Department of Structural Biology, St. Jude Children's Research Hospital, Memphis, TN, USA

[3] Department of Radiology, University of Texas Southwestern Medical Center, Dallas, USA

[4] Kidney Cancer Program, Harold C. Simmons Comprehensive Cancer Center, University of Texas Southwestern Medical Center, USA

[5] Department of Surgery, University of Cambridge, UK

[6] Peter MacCallum Cancer Centre, Melbourne, Victoria 3000, Australia

[7] Sir Peter MacCallum Department of Oncology, The University of Melbourne, Victoria 3010, Australia

[8] Department of Medicine, The University of Chicago, Chicago, IL, USA

[9] Department of Genitourinary Medical Oncology, MD Anderson Cancer Center, Houston, TX, USA

[10] Department of Neurology and Johns Hopkins Drug Discovery, Johns Hopkins School of Medicine, Baltimore, Maryland, USA

[11] Advanced Imaging Research Center, University of Texas Southwestern Medical Center, USA

[12] Department of Biochemistry and Molecular Biology, University of Florida, USA

[13] Howard Hughes Medical Institute

*Correspondence to: ralph.deberardinis@utsouthwestern.edu

Abstract

Targeting metabolic vulnerabilities has been proposed as a therapeutic strategy in renal cell carcinoma (RCC). Here, we analyzed the metabolism of patient-derived xenografts (tumorgrafts) from diverse subtypes of RCC. Tumorgrafts from *VHL*-mutant clear cell RCC (ccRCC) retained metabolic features of human ccRCC and engaged in oxidative and reductive glutamine metabolism. Genetic silencing of isocitrate dehydrogenase-1 or -2 impaired reductive labeling of TCA cycle intermediates in vivo and suppressed growth of tumors generated from tumorgraft-derived cells. Glutaminase inhibition reduced the contribution of glutamine to the tricarboxylic acid cycle and resulted in modest suppression of tumorgraft growth. Infusions with [amide-^{15}N]glutamine revealed persistent amidotransferase activity during glutaminase inhibition, and blocking these activities with the amidotransferase inhibitor JHU-083 also reduced tumor growth in both immune-compromised and immune-competent mice. We conclude that ccRCC tumorgrafts catabolize glutamine via multiple pathways, perhaps explaining why it has been challenging to achieve therapeutic responses in patients by inhibiting glutaminase.

Introduction

Renal cell carcinomas (RCC) are characterized by mutations in genes that regulate intermediary metabolism, resulting in metabolic dependencies that have been proposed as therapeutic targets (*1*). Clear cell RCC (ccRCC), the most common subtype of RCC, almost universally loses the function of the von-Hippel-Lindau (*VHL*) tumor suppressor through mutation and other mechanisms (*2, 3*). The VHL protein is a subunit of an E3 ubiquitin ligase complex that targets the oxygen-labile α-subunits of hypoxia-inducible transcription factors (HIFs) 1 and 2 for degradation; therefore, *VHL* mutation results in sustained HIF transcriptional activity (*4*). Together, HIF-1 and HIF-2 regulate the expression of many genes, including genes involved in central carbon metabolism. HIF-1 activates the expression of glucose transporters, glycolytic enzymes, lactate dehydrogenase, and pyruvate dehydrogenase kinase-1 (*PDK1*) (*5-8*). Expression of these genes induces a phenotype of enhanced glycolysis and suppressed pyruvate oxidation in mitochondria (*9, 10*). Suppression of pyruvate oxidation results from PDK1's inhibitory phosphorylation of the PDH complex, which converts pyruvate to acetyl-CoA in the mitochondria (*6*).

Although the metabolic effects of *VHL* loss on metabolism are well-established in cultured ccRCC cell lines, it is unclear which, if any, effects result in clinically-actionable metabolic vulnerabilities. An important bottleneck in cancer metabolism studies is the relative lack of information about pathway activity in vivo. By infusing cancer patients intra-operatively with [13]C-labeled nutrients and examining [13]C enrichment in metabolites extracted from the tumor after surgery, we previously characterized human tumor metabolism in several

types of cancer (*11-13*). Infusion of [U-¹³C]glucose into a small cohort of ccRCC patients revealed robust ¹³C labeling of glycolytic intermediates but impaired labeling of TCA cycle intermediates in the tumor relative to adjacent kidney (*13*). This analysis revealed a metabolic phenotype consistent with the Warburg effect, as predicted by transcriptomic and metabolomic studies in ccRCC (*3, 14-17*).

It is unknown how human ccRCCs compensate for reduced pyruvate oxidation. In culture, *VHL*-mutant RCC cells use an unusual form of glutamine metabolism to supply the TCA cycle. In this pathway, termed reductive carboxylation, the reversible isoforms of isocitrate dehydrogenases (IDH1 and IDH2) catalyze the NADPH-dependent carboxylation of glutamine-derived α-ketoglutarate to produce isocitrate and citrate (*18, 19*). Citrate is then transported to the cytosol and cleaved by citrate lyase to produce acetyl-CoA, which is used to generate fatty acids and oxaloacetate (OAA), which gives rise to other TCA cycle intermediates and aspartate (*18, 19*). In ccRCC cells, reductive glutamine metabolism is a consequence of *VHL* loss because re-expressing functional *VHL* suppresses the pathway (*19, 20*). Subcutaneous xenografts derived from an established ccRCC cell line demonstrated a small amount of reductive citrate formation, raising the possibility that this pathway might contribute to central carbon metabolism in some tumors (*20*).

Identifying tumors that unequivocally require glutamine as a fuel, even from disease-relevant preclinical models, would be significant because several inhibitors of this pathway have been evaluated as therapeutic agents. Glutaminases and amidotransferases initiate glutamine catabolism by converting glutamine to glutamate.

Glutaminase (GLS) is a mitochondrial enzyme that releases glutamine's amide group as ammonia. CB-839 is a potent and well-tolerated GLS inhibitor with efficacy in some patients with heavily-pretreated RCC (*21, 22*), although recent combination studies failed to achieve primary clinical endpoints (*23*). Amidotransferases convert glutamine to glutamate, while transferring glutamine's amide nitrogen to asparagine and intermediates in the synthesis of purines, pyrimidines, hexosamines, and nicotinamide adenine dinucleotide cofactors (*24*). The experimental amidotransferase inhibitor JHU-083, a prodrug of 6-diazo-5-oxo-L-norleucine (DON), has preclinical efficacy in multiple cancer models, suppressing tumor cell growth and enhancing antitumor immunity (*25*).

In order to identify actionable metabolic liabilities in cancer, animal models that faithfully recapitulate the metabolic features of human tumors are needed. Access to disease-relevant models is particularly important in analyzing glutamine metabolism, as many cancer cell lines use glutamine as a respiratory substrate in culture (*26*), but this does not guarantee glutamine utilization in vivo. In fact, tumors that have been analyzed with isotope tracers generally do not use glutamine as a prominent carbon source for the TCA cycle (*27, 28*). Here, we characterized the metabolic features of a large panel of patient-derived RCC tumorgrafts that retain clinical and genomic features of human tumors (*29, 30*), to determine whether they consume glutamine in vivo and respond to inhibitors of glutamine metabolism.

Results

<u>*VHL*-mutant tumorgrafts recapitulate core metabolic features of primary human ccRCC</u>

To identify tractable metabolic features in RCC, we performed metabolomics in 28 independent RCC tumorgraft lines (Table S1) passaged orthotopically in NOD-SCID mice as described previously (*29, 31*). These tumorgrafts encompass the histological and clinical diversity of aggressive forms of human RCC, including 19 high-grade ccRCCs of which 65% had confirmed *VHL* mutations, 2 each of FH-deficient RCC and papillary RCC (pRCC), 1 translocation RCC (tRCC), and 4 unclassified RCC (uRCC). Twenty-two of these tumors were derived from the primary site and the rest were from either distant (5) or regional lymph node (1) metastases. Overall, 23/28 of the models were derived from treatment-naïve tumors. Among the ccRCC tumorgrafts, 5 contained sarcomatoid features associated with poor outcomes in patients (*32*).

For metabolic analysis, we orthotopically passaged each tumorgraft into multiple NOD-SCID mice. When the tumor diameter reached ~10-15 mm, we harvested tissues from the tumor and contralateral kidney from 2-4 mice from each model. All tissues were harvested within 3 minutes of sacrificing the mouse. We performed targeted metabolomics on 1-3 fragments from each tumor and kidney. A principal component analysis (PCA) of 118 metabolites revealed that all RCC subtypes were metabolically distinct from the contralateral kidney, but this unsupervised analysis did not distinguish RCC subtypes from each other (Fig. S1A). Supervised analysis of the 50 metabolites that best distinguished tumors from kidneys revealed heterogeneity of metabolomic signatures among the ccRCC models, although most of them clustered together (Fig. S1B). As reported in ccRCC samples from patients (*16, 17, 33*), metabolites related to glutathione and glycolysis were among the most altered metabolites in tumors relative to

kidneys. A supervised analysis (variable importance in the projection, VIP) identified elevated glutathione and other glutamine-related metabolites and decreased purine- and pyrimidine-related metabolites in ccRCC tumorgrafts (Fig. S1C). Similar observations have been reported in human ccRCC (*14, 16, 17*), suggesting that ccRCC tumorgrafts retain several metabolic features of human disease.

To identify robust metabolic changes shared between tumorgrafts and human cancer, we compared metabolomics data between ccRCC tumorgrafts and a large, published metabolomics dataset from human ccRCC (*17*). Unsupervised clustering of the 76 metabolites common to both datasets separated most kidney samples from tumor samples (Fig. S1D). Among the 76 metabolites detected on both platforms, 51 differed between tumor and kidney in both studies, and most of these (62%) were either accumulated or depleted in both human and mouse tumors (Fig. 1A). These commonly-perturbed metabolites included several related to central carbon and glutamine metabolism, including glutathione, 2-hydroxyglutarate, lactate, and aspartate (Fig. 1A). We performed a similar analysis between the ccRCC tumorgrafts and patient tumors from UT Southwestern's Kidney Cancer Program (KCP) (*33*) and again observed consistent alterations in many metabolites that differentiated kidneys from tumors (Fig. S2A). The data indicate that some metabolic features of human ccRCC are durable enough to withstand multiple passages in mice.

Next, we compared metabolomic alterations among the different RCC subtypes represented in the tumorgraft panel. Glutathione was markedly elevated in all subtypes

relative to kidney (Fig. 1B). Glucose/fructose (our platform did not distinguish these from each other), lactate, and pyruvate from the glycolytic pathway were elevated in tumorgrafts derived from primary ccRCCs, but only inconsistently in tumorgrafts from metastatic ccRCC and other RCC subtypes (Fig. 1B). Levels of several TCA cycle and related metabolites were also altered in the ccRCC tumorgrafts, with an enhanced abundance of α-ketoglutarate and 2-hydroxyglutarate and a trend toward elevated glutamine and decreased succinate (Fig. 1B, Fig. S2B), as reported in human ccRCC (*17, 33*).

Orthotopic *VHL*-mutant tumorgrafts display reduced pyruvate oxidation relative to the non-malignant kidney.

We previously showed using isotope-labeled glucose infusions in patients that human ccRCCs display reduced contribution of pyruvate to the TCA cycle compared to patient-matched kidney tissue (*13*). To test whether the same phenotype is observed in ccRCC tumorgrafts, we infused [U-^{13}C]glucose via tail vein in anesthetized NOD-SCID mice with orthotopic *VHL*-mutant ccRCC. This technique assesses the fates of carbon from plasma glucose into glucose-dependent pathways in the tissues, including glycolysis and the TCA cycle (Fig. 1C). The primary ccRCC models displayed elevated ^{13}C labeling in pyruvate compared to the contralateral kidney but reduced labeling in TCA cycle-related metabolites (Fig. 1D). M+2 labeling in the TCA cycle intermediates relative to pyruvate m+3 can be used as a surrogate for carbon entry through pyruvate dehydrogenase (PDH), and m+3 labeling in the TCA cycle intermediates relative to pyruvate m+3 can be used as a surrogate for carbon entry via carboxylases, e.g. by pyruvate carboxylase (PC)

(Fig. 1C). The kidneys displayed good propagation of labeling from pyruvate into the TCA cycle via both pathways, but these ratios were suppressed in the tumors (Fig. 1D). This is similar to the labeling pattern observed in ccRCC patients infused with [U-^{13}C]glucose during nephrectomy (*13*). Overall, in both metabolomics and isotope tracing assays, these tumorgrafts recapitulate metabolic features of human ccRCC and provide an opportunity to identify nutrients used by these tumors.

Glutamine fuels the TCA cycle in *VHL*-mutant RCC

The elevation of glutamine-related metabolites and suppressed contribution of glucose to the TCA cycle suggested that glutamine is a carbon source for the TCA cycle in ccRCC tumorgrafts. To test this, we infused [U-^{13}C]glutamine into mice bearing five independent *VHL*-mutant ccRCC tumorgrafts with genetic and histological heterogeneity (Fig. S3A). We used an infusion method that labels 30-40% of glutamine and 10-17% of glutamate in the plasma (Fig. S3B). Glutamine enrichment did not differ between the tumors and adjacent benign or contralateral kidney (Fig. S3C). However, 4/5 tumorgrafts had elevated labeling in glutamate relative to the plasma and non-malignant kidney, suggesting enhanced conversion of glutamine to glutamate in the tumors (Fig. 2A, Fig. S3B). Glutaminase (GLS) was expressed in all tumorgrafts at levels comparable to the kidney (Fig. S3D).

Next, we examined the contributions of glutamine-derived carbon into TCA cycle metabolites, assessing both the oxidative and reductive pathways of glutamine metabolism (Fig. 2B). [U-^{13}C]glutamine labels glutamate and α-KG as m+5. Oxidation of

α-KG results in m+4 labeling in other TCA cycle intermediates, while reductive carboxylation generates m+5 citrate and m+3 in other metabolites (*18*). Most of the tumors had elevated ^{13}C labeling of α-KG compared with kidney tissues (Fig. S3E). All tumors displayed elevated ^{13}C labeling in at least some of the other TCA cycle intermediates as well. XP258 and XP296 demonstrated enhanced labeling along both pathways, with reductive labeling particularly evident in citrate and oxidative labeling characterizing the rest of the cycle (Fig. 2C, D). In both of these models, oxidative labeling exceeded reductive labeling in all metabolites besides citrate, indicating flow around the TCA cycle to OAA, but suppressed formation of citrate via citrate synthase, likely because of reduced PDH activity (*34, 35*). XP955 demonstrated more prominent reductive labeling and the other two lines had less consistent labeling around the cycle (Fig. S3F-H). All five tumors had enhanced citrate m+5 (reductive) labeling relative to the kidney, even XP374, which did not display elevated glutamate labeling relative to the kidney (Fig. 2E). The increased contribution of glutamine is notable in that glutamine catabolism in tumors analyzed by ^{13}C tracing in vivo is generally low (*27, 28*).

To confirm the reductive formation of citrate in these tumors, we infused [1-^{13}C]glutamine. In this labeling scheme, ^{13}C is released by α-KG dehydrogenase but retained in citrate when α-KG becomes carboxylated (Fig. S4A). *VHL*-mutant tumorgrafts (XP258 and XP955) contained enhanced citrate m+1 relative to healthy kidney tissues (Fig. S4A). We also used ^{13}C-NMR to determine the position of ^{13}C labels in malate to confirm the contribution of reductive glutamine metabolism in tumors compared to non-malignant kidney (details in the supplementary text and Fig. S4B-E).

Because the microenvironment influences tumor metabolism (*36, 37*), we asked whether the site of implantation impacts glutamine-dependent labeling. XP955 tumors were implanted into the subcutaneous space and renal capsules in the same mice. Then the mice were infused with [U-^{13}C]glutamine and metabolites were extracted from tumors at both sites. Absolute ^{13}C enrichment in glutamate and TCA cycle intermediates were nearly identical between subcutaneous and orthotopic tumors, with tumors at both sites demonstrating increased labeling in TCA cycle intermediates relative to non-malignant kidney (Fig. 2F). Overall, this experiment indicates that glutamine handling in these tumors results from cancer cell-autonomous properties rather than the microenvironment.

Reductive carboxylation in cultured cells has been ascribed to an increased α-KG/citrate ratio or a shift in the cellular redox ratio to a more reduced state (i.e., decreased NAD$^+$/NADH ratio) (*20, 35*). NAD$^+$ and NADH measurements in these tumors revealed lower NAD$^+$/NADH ratios than in non-malignant kidneys (Fig. 2G, Fig. S5A). Although both α-KG and citrate levels tended to be elevated in tumors, the α-KG/citrate ratio was not markedly different from the kidneys (Fig. S5B).

ccRCC tumorgrafts contained elevated levels of GSH and 2-hydroxyglutarate (Fig. 1B, Fig. S2B), both of which are also abundant in human ccRCC (*14, 17, 20, 33, 38*). Infusion with [U-^{13}C]glutamine resulted in labeling in both metabolites (Fig. S5C, D). In GSH, the fractional enrichment was similarly low in the tumor and kidney, but the accumulation of GSH in the tumors translated into a much higher abundance of labeled GSH. Labeled 2-

HG was present in the tumors but not the kidneys, and was nearly all L-2HG (Fig. S5D, E), as previously reported in human ccRCC (*38*).

L-[5-^{11}C]glutamine positron emission tomography and computed tomography (PET/CT) in ccRCC tumorgrafts

To assess glutamine metabolism using a different method, we performed PET/CT with L-[5-^{11}C]glutamine in mice bearing XP490 tumorgrafts in orthotopic and subcutaneous regions. Given the low soft-tissue contrast of CT, we assessed the anatomical location of the tumors and contralateral kidney with MRI (Fig. S6A). The dynamic distribution of L-[5-^{11}C]glutamine should reflect uptake and retention of glutamine by the tumors as long as the ^{11}C label remains in the tissue. In the kidney, the time-activity curve of ^{11}C radioactivity (percent injected dose per gram, %ID/g) reached its maximum (~20% ID/g) within 1.5 minutes of injection and then dropped to about half after 37.5 minutes, indicating a significant washout of L-[5-^{11}C]glutamine. This observation was similarly reported in patients with metastatic colorectal cancer (*39*). In the tumors, although the peak PET signal was lower than that in kidney, it was retained over the entire time course (Fig. S6B). A significantly increased ratio of signal uptake relative to the peak time point demonstrated higher signal retention in the tumors (both orthotopic and subcutaneous) relative to kidney (Fig. 2H). This retention is consistent with glutamine's extensive participation in metabolism in these tumors.

IDH1 and IDH2 regulate reductive carboxylation in ccRCC

Reductive carboxylation is catalyzed by NADPH-dependent cytosolic IDH1 and mitochondrial IDH2 (Fig. S7A) (*40*). To study the mechanism of reductive carboxylation in vivo and the role of IDH1 and IDH2 in tumorgraft growth, we generated a cell line from the *VHL*-mutant tumorgraft XP258. Like XP258 tumorgrafts, this cell line displayed both oxidative and reductive metabolism of glutamine in culture, and xenografts generated from XP258 cells after several passages in culture displayed nearly identical patterns of glutamine metabolism as the original tumorgrafts (Fig. 3A).

We next used CRISPR-Cas9 to create pools of cells lacking IDH1, IDH2, or both (Fig. 3B). XP258 cells do not form single-cell clones; therefore, we used pooled KO cells that likely contained a small minority of cells still expressing IDH1 and IDH2. Pools lacking IDH1, IDH2, or both proliferated in the culture at the same rate as parental cells (Fig. 3C). Reductive labeling of citrate from [U-^{13}C]glutamine was only marginally changed by IDH1 loss, but IDH2 loss had a more substantial effect, and reductive labeling was nearly eliminated in cells lacking both IDH1 and IDH2 (Fig. 3D). In cells lacking IDH2 or both IDH1 and IDH2, a higher fraction of citrate and malate displayed labeling through the oxidative pathway (Fig. 3D). Loss of IDH2 or both IDH1 and IDH2 also resulted in enhanced labeling of TCA cycle intermediates from [U-^{13}C]glucose (Fig. S7B).

We next examined the role of IDH1 and IDH2 in xenograft metabolism and growth. Xenografts from either IDH1 KO or IDH2 KO had a reduced fraction of citrate formed through reductive carboxylation, and this fraction was nearly eliminated by the loss of both enzymes (Fig. 3E). In contrast to the in vitro experiments (Fig. S7B), glucose

contribution to the TCA cycle remained similar across control, IDH1, and IDH2 KO tumors, with a small increase in citrate M+2 labeling in tumors lacking both IDH1 and IDH2 (Fig. S7C). The loss of IDH1 or IDH2 also reduced the growth of XP258 tumors (Fig. 3F). We could not consistently generate enough palpable tumors from cells lacking both IDH1 and IDH2 to perform xenograft growth experiments, despite their normal proliferation rate in culture. Examination of protein expression at the end of these experiments revealed that most tumors re-expressed at least some IDH1 and IDH2 (Fig. S7D). We confirmed this in a second cell line generated from XP258 tumorgrafts, again noting xenograft growth suppression but some recovery of IDH1 and IDH2 expression by the end of the experiment (Fig. S7E-G).

The glutaminase inhibitor CB-839 moderately decreases glutamine metabolism and growth in ccRCC tumorgrafts

We next examined the effects of CB-839 on tumor metabolism and growth. We generated subcutaneous tumors from XP490, XP258, and XP296 tumorgrafts. XP490 and XP258 were generated from treatment-naïve patients, and XP296 was generated from a patient previously treated with everolimus. CB-839 decreased glutamine's contribution to the TCA cycle along both the oxidative and reductive pathways in XP296 and XP490, but not in XP258 tumors after a 5-hour [U-^{13}C]glutamine infusion (Fig. 4A-C). CB-839 also decreased glutamine's contribution to glutathione and L-2HG in XP296 tumors (Fig. S8A, B). To more carefully assess glutamine metabolism in XP258 tumors, we performed short (5, 15 and 30 minute) infusions of [U-^{13}C,^{15}N]glutamine, reasoning that this would allow us to detect early differences in metabolite labeling that were no longer apparent when

labeling approached the steady state. These infusions revealed low ^{13}C enrichment in glutamate, α-KG, and malate in CB-839 compared to vehicle-treated tumors (Fig. 4D), indicating that CB-839 does alter glutamine metabolism in this model. All three tumorgraft models displayed modest growth suppression upon CB-839 treatment, with most tumors displaying stable volume over three weeks (Fig. 4E-G). Tumor regression was not observed, and the body weights of animals were largely unaffected by CB-839 (Fig. S8C).

Amidotransferase inhibition by JHU-083 decreases glutamine metabolism and tumor growth in *VHL* mutant ccRCC

Despite the effect of CB-839 on carbon transfer from glutamine into TCA cycle intermediates, residual glutamate labeling was noted in all tumors (Fig. 4A-C). This was not unexpected, as amidotransferases and other enzymes convert glutamine to glutamate and might compensate for glutaminase blockade (Fig. S8D) (*24, 41*). To examine the contribution of these reactions, we infused glutamine labeled on the amide nitrogen, [amide-^{15}N]glutamine, into mice with XP296, XP258 and XP490 tumorgrafts, and examined isotope enrichment in metabolites in the presence and absence of CB839. We first assessed labeling in orotate, an intermediate in pyrimidine metabolism that acquires nitrogen from the amide position of glutamine through the activity of the trifunctional CAD (carbamoyl-phosphate synthetase-2/aspartate transcarbamylase/dihydroorotase) enzyme. Orotate was highly labeled in all tumors studied and enrichment tended to be higher in the tumors than the blood, consistent with local production (Fig. S8E-G). Orotate was also labeled in tumors treated with CB839 (Fig. S8E-G), indicating persistent amidotransferase activity during glutaminase inhibition.

Other metabolites were also labeled from [amide-^{15}N]glutamine. Citrulline, a urea cycle intermediate that acquires nitrogen from ammonia via the carbamoylphosphate synthetase-1 and ornithine transcarbamylase activities in liver mitochondria, was also highly labeled. However, citrulline labeling was equivalent between tumor and plasma and was suppressed by CB-839 (Fig. S8E-G). These findings suggest that unlike orotate, tumors acquire labeled citrulline primarily from the blood. CB-839 may suppress citrulline labeling because ammonia generated through glutaminase activity in the liver and elsewhere supplies the urea cycle.

We next studied the effect of amidotransferase inhibition on glutamine metabolism and growth in ccRCC xenografts. Mice with XP258 and XP296 tumors were treated with JHU-083. Infusions with [U-^{13}C, ^{15}N]glutamine were performed in other mice earlier on treatment to determine the metabolic effects of JHU-083. Several metabolites from pathways involving amidotransferases, including asparagine, guanosine and cytosine monophosphate, displayed suppressed labeling upon JHU-083 treatment (Fig. 5A). However, ^{13}C labeling of TCA cycle intermediates was either unaffected or increased by JHU-083 (Fig. 5B). Labeling of these metabolites was suppressed by CB-839, suggesting that JHU-083's effect on amidotransferases is more prominent than its effect on glutaminase under these conditions. Despite the persistent labeling of TCA cycle intermediates, JHU-083 suppressed the growth of XP258 and XP296 tumors (Fig. 5C), with only a small decrease in body weight (Fig. S9A). Because JHU-083 has outstanding anti-tumor efficacy in some immunocompetent models (25), we tested it in two syngeneic

RCC cell lines, 17175 and 10950, which primarily showed oxidative glutamine metabolism in vivo (Fig S9B). JHU-083 essentially eliminated the growth of xenografts derived from these cells in C57BL/6 mice, again with only mild weight loss (Fig. 5D, Fig. S9C).

Discussion

Metabolic phenotypes in human tumors are incompletely defined, and we seldom know how well key metabolic features in human tumors are recapitulated in mice. This presents a challenge in using mouse models to identify metabolic liabilities in human tumors. Human ccRCC metabolism has been well-characterized through metabolomics and isotope labeling studies. These studies suggest that human ccRCC tumors share metabolic features with *VHL*-deficient ccRCC cell lines, including enhanced glycolysis and suppressed pyruvate oxidation (the Warburg effect) (*13, 14, 16, 17*). We extended these comparisons here by characterizing metabolism in a large series of orthotopic patient-derived tumorgrafts that retain genomic and histological features of human RCC (*30*). ccRCC-derived tumorgrafts display metabolic features of human ccRCC, including suppressed contributions of glucose to TCA cycle intermediates and accumulation of glutamine-derived metabolites such as glutathione and L-2HG. Using two types of tracer studies, [^{13}C]glutamine infusion and dynamic [^{11}C]glutamine PET, we find that glutamine is a carbon source for intermediary metabolism in ccRCC tumorgrafts relative to nonmalignant kidney. The PET data indicate retention of glutamine carbon (specifically, carbon 5) within the tumors. Whether this primarily reflects glutamine's contribution to the TCA cycle, or to glutathione, proteins and other macromolecules, is unknown. We also note that glutamine metabolism is an intrinsic characteristic of these tumors rather than simply a consequence of the microenvironment, as both ^{11}C and ^{13}C labeling features

were similar between orthotopic and subcutaneous sites. A recent study that used radiotracers to compare glutamine uptake between cancer cells and immune cells in the tumor microenvironment also concluded that glutamine is preferentially taken up and retained by malignant cells (*42*).

Our ^{13}C labeling approach reports fractional enrichment of metabolites downstream of the tracer after several hours of exposure to ^{13}C-glutamine. Therefore, a limitation of the approach is that it reports glutamine's metabolic fates but not the rates of glutamine-dependent pathways, including glutamine oxidation and reductive carboxylation. Nevertheless, glutamine's prominent contributions to several pathways in the tumors compared to the kidneys raised the possibility of vulnerabilities associated with these pathways. We confirmed this by blocking glutaminase and amidotransferases. Combining these loss-of-function experiments with isotope tracing is informative because it provides direct evidence that the target enzyme contributes to the pathway in vivo.

Numerous studies in cultured cells have demonstrated enhanced reductive labeling of TCA cycle intermediates under conditions of *VHL* loss, hypoxia, PDH suppression, or defects in the TCA cycle or electron transport chain (ETC) (*18, 19, 34, 35*). A small amount of reductive carboxylation was also observed in xenografts of *VHL*-deficient UMCR3 cells, and growth of these tumors was suppressed by BPTES, a preclinical GLS inhibitor (*20*). The tumors in the current study generally show combined oxidative and reductive labeling, with the latter resulting in citrate m+5 during infusion with [U-^{13}C]glutamine. This phenomenon may be related to PDH suppression, as a reduced

citrate:α-KG or NAD$^+$/NADH ratio increases the appearance of reductively labeled citrate (*20, 35*). We do not know whether these tumorgrafts contain a net reductive flux from α-KG to citrate, but the presence of citrate m+5 prompted us to examine the role of IDH1 and IDH2. Both IDH1 and IDH2 contribute to citrate m+5 labeling in vivo. Although these enzymes are dispensable for cell growth in culture, the loss of either one reduces PDX growth. At present, we do not know if reductive citrate formation per se is the key IDH1/2 function in vivo, or whether these enzymes are required for survival during redox stress or glucose deprivation in vivo, as reported in other models of IDH1/2 dependence (*43, 44*). However, the data suggest that dual targeting of these enzymes could produce a therapeutic effect in a subset of ccRCC tumors that demonstrate reductive carboxylation in vivo. A compound that inhibits the oncogenic, mutant forms of both IDH1 and IDH2 has been developed (*45*), so perhaps potent inhibition of both wild-type enzymes could be achieved with a single compound. We also note that although IDH1/2 knockout suppresses the growth of *VHL*-mutant XP258 tumorgrafts, it is unclear at present whether *VHL* loss is sufficient to induce IDH1/2 dependence in vivo.

Because glutamine is a prominent carbon and nitrogen source, there has been a sustained interest in suppressing glutamine catabolism in tumors. Several recent studies have shown preclinical efficacy of targeting glutamine metabolism in vivo by inhibitors of glutamine transporters, glutaminase, and amidotransferases (*25, 41, 46-50*). Phase-1 dose-escalation clinical studies with CB-839 demonstrated good tolerability of the drug and stable disease or partial response in some RCC patients (*22*). However, the CANTATA (NCT03428217) Phase 3 trial revealed that combining CB-839 with

cabozantinib in advanced, metastatic RCC patients pre-treated with checkpoint inhibitors or anti-angiogenic therapies did not improve outcomes over cabozantinib alone (*51*). Our data indicate that tumor glutaminase is inhibited by CB-839 and that the response to glutaminase inhibition may be dependent on how other pathways compensate for glutaminase loss. Glutaminase expression was similar between kidney and tumors, implying that factors beyond protein abundance (e.g., substrate availability, post-translational modifications, etc.) account for glutamine's higher contribution to the TCA cycle in the tumors. This also emphasizes that direct analysis of tumor metabolism by isotope tracing can uncover metabolic liabilities that would not be obvious from gene/protein expression studies. In this case, isotope tracing also provided some insight into how tumors resist CB-839. Upon treatment with CB-839, ccRCC tumorgrafts derived from treatment-naïve primary tumors display modest growth suppression and metabolic derangements. All tumors contained ^{13}C-labeled glutamate and downstream metabolites despite CB-839 therapy, indicating persistent glutamine metabolism in the tumors. Transfer of ^{15}N from [amide-^{15}N]glutamine or [U-^{13}C,^{15}N]glutamine to products of amidotransferase reactions, including asparagine and intermediates from nucleotide metabolism, indicates that this family of enzymes enables residual glutamine catabolism in CB-839-treated tumors.

Since the early 1950s, clinical studies in several kinds of cancer observed stabilization or remission in some patients treated with low-dose DON (*52, 53*). However, clinical development was halted due to inconsistent response and gastrointestinal toxicity in dose-escalation studies (*54*). The recent development of JHU-083, a chemically-modified

prodrug of DON with selective activation in the tumor microenvironment and a reduced toxicity index, has suggested an alternative approach to broad inhibition of glutamine catabolism in cancer. In syngeneic tumors, JHU-083 impairs glutamine catabolism in cancer cells and produces a nutrient microenvironment more amenable to the antitumor effects of cytotoxic T cells and other immune cells (25, 55, 56). JHU-083's efficacy in the immunodeficient tumorgraft models described here may underestimate its overall efficacy in tumors with robust glutamine catabolism. This is particularly relevant for ccRCC, a highly immunogenic tumor type with clinical responses to immunotherapy (57, 58). Our isotope-tracing experiments with [U-^{13}C,^{15}N]glutamine showed decreased ^{15}N-enrichment in asparagine and nucleotide intermediates upon JHU-083 treatment. Importantly, although several studies have reported that JHU-083 decreases glutaminase activity in tumors (55, 59), the effect appears to be different in ccRCC tumorgrafts. In our models, ^{13}C labeling of TCA cycle intermediates is glutaminase-dependent and inhibited by CB-839. As expected, this drug does not impair amidotransferase reactions. On the other hand, JHU-083 inhibits amidotransferase reactions but not ^{13}C labeling of TCA cycle intermediates in steady-state isotope infusion experiments, suggesting that its effect on glutaminase in these models is modest compared to CB-839. Yet both CB-839 and JHU-083 impair tumor growth. To maximize the therapeutic benefit and durability in ccRCC, it may be necessary to impair amidotransferases and glutaminase concurrently. Developing a strategy with an acceptable therapeutic index would likely benefit from defining which amidotransferase activities are most important for ccRCC growth.

Materials and Methods

Patient-derived xenografts (Tumorgrafts)

All mouse experiments were approved by the UT Southwestern Institutional Animal Care and Use Committee (IACUC). All tumorgrafts used in this study were pre-established patient-derived xenografts. For tumor passaging and metabolomics analysis, tumorgrafts were passaged in the left kidney of 4-8 weeks old male/female NOD-SCID mice, as described previously (29-31). Once the tumor was palpable and close to ~ 10mm in diameter, mice were euthanized to collect tumors in ice-cold HBSS buffer with 1% Pen/Strep. For tumor passaging, tumor tissues were dissected aseptically into ~ 8 mm^3 of tissues. Tumor pieces were freshly implanted into the left kidney of male or female mice, and the mice were monitored periodically for tumor growth by palpating the implanted flank. For subcutaneous tumor studies, resected orthotopic tumors were dissected into small pieces (~64 mm^3) and surgically implanted subcutaneously in the right hind region of the mice. Experiments with both orthotopic and subcutaneous tumors were conducted by surgically implanting tumors simultaneously in the left kidney and right hind region of the mice. For the characterization of glutamine metabolism, we focused on PDXs that grew to sufficient tumor size (within 60 days or less) for metabolic and therapeutic experiments.

Metabolomics

For tissue metabolomics, mice were sacrificed and tissues were collected within 3 minutes in liquid nitrogen to minimize metabolite degradation. Tumor and kidney tissues from mice were dissected on dry ice, and ~50mg of the tissues was placed in 1ml ice-cold 80% MeOH solution in HPLC-grade water. Tissues were homogenized using a hand-held automatic sonicator and subjected to three freeze-thaw cycles between liquid

nitrogen and a 37 °C water-bath. Supernatants were collected by centrifuging the homogenized solution at 10,000 rpm for 10 minutes to remove tissue debris, then the supernatants were evaporated overnight in a SpeedVac concentrator (Thermo Savant). Dried samples were reconstituted in 100μl of 0.03% formic acid in HPLC grade water, vortexed, centrifuged, and the supernatant was transferred in LC/MS glass vials containing sample inserts. As described previously (*60*), LC/MS was performed using an AB QTRAP 5500 liquid chromatography/triple quadrupole mass spectrometer (Applied Biosystems SCIEX, Foster City, CA). Chromatograms and peak area of each metabolite were reviewed and integrated using MultiQuant software version 2.1 (Applied Biosystems SCIEX, Foster City, CA). Each metabolite's peak area was selected from either positive or negative modes depending on previously-run standards. Peak areas were corrected to blank samples and then normalized to the total ion count (TIC) of that sample to correct for variation introduced by sample handling and instrumentation. TIC-normalized data were log-transformed and median normalized for multivariate analyses using Metaboanalyst 4.0 (*61*) (http://www.metaboanalyst.ca). Additional statistical analyses were performed using one-way ANOVA and Student's t-test in R software.

Isotope infusions

Isotope infusion experiments were conducted when tumor size was between 10-50 mm diameter for orthotopic tumors and 0.5-1cm diameter for subcutaneous tumors. When the tumors reached these sizes, the mice were fasted for 16 hours and infused in the morning with glucose for 3 hours or glutamine for 5 hours. The mice were anesthetized on a heating pad, and a catheter (connected to the infusion solution and pump) was inserted

in the lateral tail vein. For [U-^{13}C]glucose infusions, we prepared a 750 µl of saline solution containing 2.48 grams of isotope-labeled glucose per kg of body weight. An initial bolus of 125 µl/min was delivered under anesthesia for 1 min, followed by 2.5 µl/min of continuous infusion for 3 hours. For glutamine infusions, we prepared 1500 µl of a saline solution containing 50 U/ml of Heparin with 1.725 grams of isotope-labeled glutamine per kg of body weight. An initial bolus of 150 µl/min was delivered under anesthesia for 1 min, followed by 2.5 µl/min of continuous infusion for 5 hours. Mice were kept under anesthesia throughout the experiment, and blood samples were collected at regular intervals via retro-orbital puncture using microcapillary tubes. At the end of the infusion, mice were euthanized, and tumors and other organs were harvested and snap-frozen in liquid nitrogen. Endpoint blood samples were collected from the heart with an insulin syringe.

For the time-course study with [U-^{13}C,^{15}N]glutamine, we first generated xenografts from XP258 cell lines implanted bilaterally in NOD-SCID mice. Once the tumors were of a measurable size, we randomized the animals for a 7 dose of CB-839 (200 mg/kg twice daily) or vehicle treatment. Isotope infusions were performed on the last day of treatment using a protocol modified from a reported method (62). Specifically, the priming dose of glutamine (100+1.6*(weight in g) microliters)/min was delivered under anesthesia through a tail vein catheter. Infusion rates were then adjusted to 0.1 microliters*(weight in g)/min. Mice were euthanized, and tissues were collected for mass spectrometry analysis at 5, 15 and 30 minutes after the infusion.

Gas-chromatography mass-spectrometry (GC-MS)

Blood samples from all infusion experiments were kept on ice until the last sample was collected. Plasma was collected from blood samples through centrifugation at 5000 rpm for 5 minutes at 4 ⁰C. While 10µl of plasma was directly resuspended in ice-cold 80% MeOH in water, tissues samples of about 5-20mg were homogenized in 80% MeOH in water and subjected to three cycles of free-thaw in liquid nitrogen and a 37 °C water-bath. Tissue and plasma extracts were centrifuged at 10,000 g for 10 min at 4 ⁰C to separate macromolecules and cell-debris. The supernatants were vacuum dried over-night. The dried samples were reconstituted in 40 µl of pyridine containing methoxyamine (10mg/ml). The samples were vortexed, transferred to GC/MS glass vials with inserts, and heated for 10 minutes at 70 °C. Next, we added 80 µl of the derivatization agent (N-(*tert*-butyldimethylsilyl)-N-methyltrifluoroacetamide (MTBSTFA)) and heated the samples for 1 hour at 70 °C on a heating block. Samples were then loaded into an autoinjector, and 1µl of each sample was injected for analysis. Samples were run on either Agilent 6890 gas chromatograph coupled to an Agilent 5973N Mass Selective Detector or an Agilent 5975C Inert XL Mass Selective Detector. The resulting chromatogram and mass isotopologue distributions were analyzed in Agilent MSD ChemStation Data-Analysis software. The area under the curve for mass isotopologues was corrected for natural abundance using in-house methods run in MATLAB

Liquid-chromatography and mass spectrometry (LC-MS/MS)

For the quantification of NAD, NADH$^+$, alpha-ketoglutarate and the analysis of metabolites labeled from [amide-^{15}N]glutamine, or [U-^{13}C,^{15}N]glutamine, we used HILIC column-based separation of metabolites on a Vanquish UHPLC coupled to a QExactive HF-X

hybrid quadrupole orbitrap high-resolution mass spectrometer (HRMS) from Thermo Fisher Scientific (Bremen, Germany), as described previously([63]). Briefly, ~10-20mg tissue samples were extracted in 80% acetonitrile, homogenized, subjected to 3 freeze-thaw cycles as described above, and cleared of macromolecules by centrifugation. Protein concentration was determined in each supernatant using the BCA Assay, and equivalent aliquots of protein-normalized samples were injected into the instrument. The chromatogram was analyzed as described previously [52] using TraceFinder Software from Thermo Fisher Scientific. The ^{13}C and ^{15}N labeled data were corrected for natural abundance using IsoCorrectoR ([64]).

For targeted GSH quantification, metabolites were extracted from tissues homogenized in 80% acetonitrile solution containing N-ethylmaleimide (0.3 M), followed with 3 freeze-thaw. The samples were centrifuged, and the supernatant was analyzed with a SCIEX QTRAP 5500 liquid chromatography/triple quadrupole mass spectrometer. We performed separation of metabolites on a SeQuant® zic®-pHILIC polymeric HPLC column (150×2 mm) in a Nexera Ultra-High-Performance Liquid Chromatography system (Shimadzu Corporation). Mobile phase solvents were 10 mM ammonium acetate aqueous (pH 9.8 adjusted with ammonia water (A), and pure acetonitrile (B). The gradient elution was: 0–18 min, linear gradient 90–55% B and 18–20 min, linear gradient 55-30% B, 20-25 min, 30% B, 25-27 min, linear gradient 30-90% B, then the column was reconditioned for 6 min using 90% B. The flow rate was 0.2 ml/min and the column was operated at 40 °C. Multiple reaction monitoring (MRM) was used to check N-ethylmaleimide derivatized GSH, 433/304 (M+0, CE: 19V), 438/304 (M+5, CE: 19V).

Measurement of D-2HG and L-2HG in tissues

Metabolites were extracted from the tissues samples with 80% methanol-water solution, and the resulting supernatant was dried in a SpeedVac. Dried pellet from unlabeled samples was mixed with [U^{13}C]-D/L-2HG (internal standard for unlabeled samples, Cambridge isotope laboratories, 10 nG in 10 µl acetonitrile). The mixture (or pellet from labeled samples) was dissolved in a diacetyl-L-tartaric anhydride (DATAN) solution (90 µl, 50 mG/ml in freshly mixed 80% acetonitrile/20% acetic acid, DATAN, Acros Organics). The solution was sonicated, warmed to 75 °C for 30 min, cooled to room temperature, and then centrifuged again to collect the supernatant. The supernatant was dried in a SpeedVac, and the pellet was reconstituted into 1.5 mM ammonium formate aqueous solution with 10% acetonitrile (100 µl). For [U-^{13}C]glutamine tracing samples, derivatized [U^{13}C]-D/L-2HG was used as a standard to identify the derivatized peaks. LC/MS analysis was performed on an AB Sciex 5500 QTRAP liquid chromatography/mass spectrometer (Applied Biosystems SCIEX) equipped with a triple quadrupole/iontrap mass spectrometer with electrospray ionization interface and controlled by AB Sciex Analyst 1.6.1 Software. Waters Acquity UPLC HSS T3 column (150 × 2.1 mM, 1.8 µM) column was used for separation. Solvents for the mobile phase were 1.5 mM ammonium formate aqueous (pH 3.6 adjusted with formic acid (A), and pure acetonitrile (B). The gradient elution was: 0–12 min, linear gradient 1–8% B and 12–15 min, 99% B, then the column was washed with 99% B for 5 min before reconditioning it for 3 min using 1% B. The flow rate was 0.25 ml/min and the column was operated at 35 °C. Multiple reaction monitoring (MRM) was used to check 2-hydroxyglutarate-diacetyl tartrate derivatives: 363/147 (M+0,

CE: −14V); 368/152 (M+5, CE: −14V). This method was modified and adapted from previously published work (*65*).

<u>^{13}C Nuclear Magnetic Resonance Spectroscopy</u>

Kidney (43 – 107 mg) and tumor (220 – 462 mg) samples were subjected to extraction in ice cold acetonitrile – isopropanol – water (ACN/IPA/water) by adding the solvent to pre-weighed tissue samples. Zirconium beads were added to the sample suspension and the samples were homogenized using a Fastprep-24 (MP Biomedicals, California, USA). Homogenization was carried out in a "20 seconds ON – 5 minutes OFF" cycle to avoid overheating extracts and repeated at least four times. The homogenates were centrifuged at 10,000xg for 30 minutes at 4°C and the supernatant was lyophilized at room temperature in a Speedvac system (Thermo Scientific, Waltham, MA). The dried sample was dissolved in a 1:1 acetonitrile-water mixture, centrifuged at 10,000xg for 30 minutes at 4°C and lyophilized again.

Samples were prepared for NMR analysis by dissolving lyophilized powder in 54 µl of 50 mM sodium phosphate buffer in D_2O containing 2 mM ethylenediaminetetraacetic acid (EDTA). 6 µl of an internal standard containing 0.02%(w/v) NaN_3 and 0.5 mM deuterated sodium 3 – trimethylsilyl – 1 – propanesulphonate (d6-DSS) was added to the sample solution. The solution was vortexed thoroughly, centrifuged at 10,000xg for 5 minutes and 54 µl of the supernatant was loaded into a 1.5 mm NMR tube. All NMR spectra were acquired using a 14.1 T NMR magnet equipped with a home-built superconducting (HTS) probe (*66*). Parameters included an acquisition time of 1.5 s, a spectral width of 240 ppm,

and ^{1}H decoupling field strength of 4800 Hz. The spectra were processed (zero-filled to 131,072 points, 0.8 Hz exponential line broadening, and polynomial or spline baseline correction as necessary) and referenced by setting the singlet resonance of taurine (N-C1) to 48.4 ppm. Peaks of interest were fitted to a mixed Gaussian/Lorentzian shape and area under the peaks were obtained.

L-[5-^{11}C]glutamine PET/CT and MRI

A 3.5-inch-long catheter of 0.38 mm inner diameter fitted with a 27-gauge needle was inserted into the tail of a tumor-bearing mouse for intravenous injection of L-[5-^{11}C]-glutamine (~ 5 MBq per mouse). Dynamic PET data acquisition was performed on a Mediso NanoScan PET/CT System (Mediso, USA) immediately after the injection for up to 40 minutes, followed by a CT scan with 720 projections. The images were reconstructed using the manufacturer's software and quantitatively analyzed by Inveon Research Workplace (Siemens, USA). MR Imaging (MRI) was performed on a 1 Tesla Desktop magnetic resonance (MR) scanner (M2 Compact, Aspect Imaging, Shoham, Israel) using a mouse volume coil. T2-weighted imaging was performed with a fast spin-echo (FSE; TR/TE = 2500/80 ms) sequence with the mouse in a prone position.

Cell lines derived from XP258 tumorgrafts

Orthotopic tumors were dissected and collected in ice-cold 20 ml transport media (MEM, 50 µg/ml gentamycin, 2.5 µg/ml fungizone, and 1X Pen/Strep). The tissues were washed three times in 15 ml of transport media and minced into small pieces using sterile Swan-Motron blades. Minced tissue was resuspended in 15 mL of enzyme mix (transport media,

collagenase, hyaluronidase, and DNase IV) and incubated for 1 to 2 hours in 5% CO_2 in a 37 °C incubator. Enzyme-digested tissue samples were filtered to separate cells from debris and larger tissue pieces. Freshly-obtained cells were cultured in high glucose DMEM supplemented with 10% FBS, 1% Pen/Strep, 1X non-essential amino acids, 0.8 mg/ml hydrocortisone and 10 µg/ml EGF.

Syngeneic RCC cell lines

We used 17175 and 10950 cell lines, which were generated by the lab of Dr. Giannicola Genovese, MD Anderson Cancer Center. These cells were plated in 0.1% gelatin-coated plates (Cat # ES-006-B, Merck Millipore) in high-glucose-containing DMEM media with 20% FBS and 1% Pen/Strip. To generate tumors, we injected 1 million cells of each cell line mixed 50:50 (v/v) with Matrigel into the flanks of C57BL/6 mice.

Knockout of IDH1 and IDH2 using CRISPR-Cas9

IDH1 and IDH2 knockout pools were generated using the CRISPR-Cas9 system, as described previously (*44*). Briefly, IDH1 sgRNA (ACGTGGAATTGGATCTACAT) and IDH2 sgRNA (ATGAGATGACCCGTATTATC) were cloned into lentiCRISPR V2 system following a published protocol(*67*). These vectors were transfected into HEK239T cells, and media containing viral particles was collected at 48 hours and 72 hours and filtered through a 0.45 µM filter. The tumorgraft-derived XP258 cell lines were transduced with media containing viral particles, then selected in puromycin (2µg/ml) to generate pools of modified cells. IDH1/2 double KO XP258 cells were generated by transducing IDH2 KO cells with viral particles carrying the IDH1 sgRNA.

Immunoblotting

Cells and tissues were homogenized in RIPA buffer containing protease and phosphatase inhibitors. Supernatants were collected by centrifugation at 14,000 RPM for 20 minutes at 4 °C. Protein concentrations were determined using the BCA protein assay. Protein lysates were resolved in SDS-PAGE and transferred to methanol-activated nitrocellulose membranes. The membrane was first blocked with 5% milk and then incubated overnight with primary antibodies against GLS (ThermoFisher, Cat# PA5-40135), IDH1 (Cell Signaling, Cat# 8137 or Abcam Cat# ab239606), IDH2 (Abcam, ab55271), β-Actin (Cat #5125S), or GAPDH (Cat #8884S) from Cell Signaling Technologies.

CB-839 and JHU-083 treatment

Calithera Inc. (San Francisco, CA) provided the solution for CB-839 and vehicle for in vivo experiments. 200 mg/kg of CB-839 or vehicle in 200µl of solution was administered orally every 12 hours for 21-22 days, as described previously (*48*). JHU-083 was a gift from Dr. Barbara Slusher. JHU-083 was diluted in 50mM HEPES-buffered saline such that 100ul contained 1.83mg/kg of JHU-083. Mice with ccRCC tumorgrafts were administered either 100ul of JHU-083 or vehicle, whereas the mice with syngeneic cell line derived tumors were administered 50ul of JHU-083 or vehicle. Mice were dosed with JHU-083 or vehicle via intraperitoneal injection according to a regimen of 5 days on and 2 days off. Tumor growth and body weight were monitored twice weekly. Tumor volume was measured using electronic calipers and calculated using the formula Length x (Width^2)/2. Infusion of [U-^{13}C]glutamine and metabolomics were conducted at the end of 21 days of the CB-839

treatment. Infusion with [Amide-^{15}N]glutamine, [U-^{13}C,^{15}N]glutamine, and [U-^{13}C]glucose were conducted after 5-7 doses of the drugs (JHU-083 or CB-839) or vehicle.

<u>Isotope tracing in vitro</u>

One million cells were plated overnight for each experiment. For isotope-tracing experiments, cells were washed with PBS and then fed with medium containing 100% isotope-labeled tracer of interest with 10% dialyzed FBS. [U-^{13}C]glucose and [U-^{13}C]glutamine labeling was performed over 3-4 hours. After the experiment, the cells were washed once in ice-cold saline solution and 1ml of 80% methanol solution was added. The cells were collected by scraping and subjected to three freeze-thaw cycles followed by centrifugation to remove debris. The supernatant was vacuum dried overnight in a Speedvac and reconstituted for GC-MS analysis as described above.

<u>Statistical analysis</u>

Metaboanalyst 4.0(*61*) and R-software (http://www.metaboanalyst.ca) were used to conduct statistical analysis of the metabolomics data. R software and PRISM were used to conduct one-way ANOVA and Student's t-test, respectively. Data visualization used both PRISM and R Software. Descriptions of individual statistical analyses can be found in the figure legends.

References

1. W. M. Linehan *et al.*, The Metabolic Basis of Kidney Cancer. *Cancer Discov* **9**, 1006-1021 (2019).
2. X. Ma, K. Yang, P. Lindblad, L. Egevad, K. Hemminki, VHL gene alterations in renal cell carcinoma patients: novel hotspot or founder mutations and linkage disequilibrium. *Oncogene* **20**, 5393-5400 (2001).
3. N. Cancer Genome Atlas Research, Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature* **499**, 43-49 (2013).
4. D. Tarade, M. Ohh, The HIF and other quandaries in VHL disease. *Oncogene* **37**, 139-147 (2018).
5. G. L. Semenza *et al.*, Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1. *J Biol Chem* **271**, 32529-32537 (1996).
6. J. W. Kim, I. Tchernyshyov, G. L. Semenza, C. V. Dang, HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab* **3**, 177-185 (2006).
7. B. L. Ebert, J. D. Firth, P. J. Ratcliffe, Hypoxia and mitochondrial inhibitors regulate expression of glucose transporter-1 via distinct Cis-acting sequences. *J Biol Chem* **270**, 29083-29089 (1995).
8. G. L. Semenza, P. H. Roth, H. M. Fang, G. L. Wang, Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1. *J Biol Chem* **269**, 23757-23763 (1994).
9. G. L. Semenza, HIF-1: upstream and downstream of cancer metabolism. *Curr Opin Genet Dev* **20**, 51-56 (2010).
10. G. L. Semenza, HIF-1 mediates the Warburg effect in clear cell renal carcinoma. *J Bioenerg Biomembr* **39**, 231-234 (2007).
11. E. A. Maher *et al.*, Metabolism of [U-13 C]glucose in human brain tumors in vivo. *NMR Biomed* **25**, 1234-1244 (2012).
12. C. T. Hensley *et al.*, Metabolic Heterogeneity in Human Lung Tumors. *Cell* **164**, 681-694 (2016).
13. K. D. Courtney *et al.*, Isotope Tracing of Human Clear Cell Renal Cell Carcinomas Demonstrates Suppressed Glucose Oxidation In Vivo. *Cell Metab* **28**, 793-800 e792 (2018).
14. H. I. Wettersten *et al.*, Grade-Dependent Metabolic Reprogramming in Kidney Cancer Revealed by Combined Proteomics and Metabolomics Analysis. *Cancer Res* **75**, 2541-2552 (2015).
15. C. J. Ricketts *et al.*, The Cancer Genome Atlas Comprehensive Molecular Characterization of Renal Cell Carcinoma. *Cell Rep* **23**, 313-326 e315 (2018).
16. B. Li *et al.*, Fructose-1,6-bisphosphatase opposes renal carcinoma progression. *Nature* **513**, 251-255 (2014).
17. A. A. Hakimi *et al.*, An Integrated Metabolic Atlas of Clear Cell Renal Cell Carcinoma. *Cancer Cell* **29**, 104-116 (2016).
18. A. R. Mullen *et al.*, Reductive carboxylation supports growth in tumour cells with defective mitochondria. *Nature* **481**, 385-388 (2011).
19. C. M. Metallo *et al.*, Reductive glutamine metabolism by IDH1 mediates lipogenesis under hypoxia. *Nature* **481**, 380-384 (2011).

20. P. A. Gameiro *et al.*, In vivo HIF-mediated reductive carboxylation is regulated by citrate levels and sensitizes VHL-deficient cells to glutamine deprivation. *Cell Metab* **17**, 372-385 (2013).
21. F. Meric-Bernstam *et al.*, Phase 1 study of CB-839, a small molecule inhibitor of glutaminase (GLS), alone and in combination with everolimus (E) in patients (pts) with renal cell cancer (RCC). *Journal of Clinical Oncology* **34**, 4568-4568 (2016).
22. J. J. Harding *et al.*, A Phase 1 Dose-Escalation and Expansion Study of Telaglenastat in Patients With Advanced or Metastatic Solid Tumors. *Clin Cancer Res*, (2021).
23. N. M. Tannir *et al.*, CANTATA: Primary analysis of a global, randomized, placebo (Pbo)-controlled, double-blind trial of telaglenastat (CB-839) + cabozantinib versus Pbo + cabozantinib in advanced/metastatic renal cell carcinoma (mRCC) patients (pts) who progressed on immune checkpoint inhibitor (ICI) or anti-angiogenic therapies. *Journal of Clinical Oncology* **39**, 4501-4501 (2021).
24. L. Yang, S. Venneti, D. Nagrath, Glutaminolysis: A Hallmark of Cancer Metabolism. *Annu Rev Biomed Eng* **19**, 163-194 (2017).
25. R. D. Leone *et al.*, Glutamine blockade induces divergent metabolic programs to overcome tumor immune evasion. *Science* **366**, 1013-1021 (2019).
26. R. J. DeBerardinis *et al.*, Beyond aerobic glycolysis: transformed cells can engage in glutamine metabolism that exceeds the requirement for protein and nucleotide synthesis. *Proc Natl Acad Sci U S A* **104**, 19345-19350 (2007).
27. I. Marin-Valencia *et al.*, Analysis of tumor metabolism reveals mitochondrial glucose oxidation in genetically diverse human glioblastomas in the mouse brain in vivo. *Cell Metab* **15**, 827-837 (2012).
28. S. M. Davidson *et al.*, Environment Impacts the Metabolic Dependencies of Ras-Driven Non-Small Cell Lung Cancer. *Cell Metab* **23**, 517-528 (2016).
29. S. Sivanand *et al.*, A validated tumorgraft model reveals activity of dovitinib against renal cell carcinoma. *Sci Transl Med* **4**, 137ra175 (2012).
30. R. Elias *et al.*, A renal cell carcinoma tumorgraft platform to advance precision medicine. *Cell Rep* **37**, 110055 (2021).
31. A. Pavia-Jimenez, V. T. Tcheuyap, J. Brugarolas, Establishing a human renal cell carcinoma tumorgraft platform for preclinical drug testing. *Nat Protoc* **9**, 1848-1859 (2014).
32. R. Gupta *et al.*, Survival outcomes of sarcomatoid renal cell cancer (sRCC) compared to clear cell renal cell cancer (ccRCC): An analysis of SEER data. *Journal of Clinical Oncology* **38**, e17101-e17101 (2020).
33. Y. Zhang *et al.*, Addressing metabolic heterogeneity in clear cell renal cell carcinoma with quantitative Dixon MRI. *JCI Insight* **2**, (2017).
34. K. N. Rajagopalan *et al.*, Metabolic plasticity maintains proliferation in pyruvate dehydrogenase deficient cells. *Cancer Metab* **3**, 7 (2015).
35. S. M. Fendt *et al.*, Reductive glutamine metabolism is a function of the alpha-ketoglutarate to citrate ratio in cells. *Nat Commun* **4**, 2236 (2013).
36. M. O. Yuneva *et al.*, The metabolic profile of tumors depends on both the responsible genetic lesion and tissue type. *Cell Metab* **15**, 157-170 (2012).
37. J. R. Mayers *et al.*, Tissue of origin dictates branched-chain amino acid metabolism in mutant Kras-driven cancers. *Science* **353**, 1161-1165 (2016).

38. E. H. Shim *et al.*, L-2-Hydroxyglutarate: an epigenetic modifier and putative oncometabolite in renal cancer. *Cancer Discov* **4**, 1290-1298 (2014).

39. A. S. Cohen *et al.*, First-in-human PET imaging and estimated radiation dosimetry of L-[5-(11)C]-glutamine in patients with metastatic colorectal cancer. *J Nucl Med*, (2021).

40. S. Tommasini-Ghelfi *et al.*, Cancer-associated mutation and beyond: The emerging biology of isocitrate dehydrogenases in human disease. *Sci Adv* **5**, eaaw4543 (2019).

41. A. Mendez-Lucas *et al.*, Identifying strategies to target the metabolic flexibility of tumours. *Nat Metab* **2**, 335-350 (2020).

42. B. I. Reinfeld *et al.*, Cell-programmed nutrient partitioning in the tumour microenvironment. *Nature* **593**, 282-288 (2021).

43. A. Vaziri-Gohar *et al.*, Limited nutrient availability in the tumor microenvironment renders pancreatic tumors sensitive to allosteric IDH1 inhibitors. *Nat Cancer* **3**, 852-865 (2022).

44. L. Jiang *et al.*, Reductive carboxylation supports redox homeostasis during anchorage-independent growth. *Nature* **532**, 255-258 (2016).

45. Z. Konteatis *et al.*, Vorasidenib (AG-881): A First-in-Class, Brain-Penetrant Dual Inhibitor of Mutant IDH1 and 2 for Treatment of Glioma. *ACS Med Chem Lett* **11**, 101-107 (2020).

46. M. L. Schulte *et al.*, Pharmacological blockade of ASCT2-dependent glutamine transport leads to antitumor efficacy in preclinical models. *Nat Med* **24**, 194-202 (2018).

47. E. H. Shroff *et al.*, MYC oncogene overexpression drives renal cell carcinoma in a mouse model through glutamine metabolism. *Proc Natl Acad Sci U S A* **112**, 6539-6544 (2015).

48. M. I. Gross *et al.*, Antitumor activity of the glutaminase inhibitor CB-839 in triple-negative breast cancer. *Mol Cancer Ther* **13**, 890-901 (2014).

49. R. Romero *et al.*, Keap1 loss promotes Kras-driven lung cancer and results in dependence on glutaminolysis. *Nat Med* **23**, 1362-1368 (2017).

50. H. Ding *et al.*, Activation of the NRF2 antioxidant program sensitizes tumors to G6PD inhibition. *Sci Adv* **7**, eabk1023 (2021).

51. N. M. Tannir *et al.*, Phase 1 study of glutaminase (GLS) inhibitor CB-839 combined with either everolimus (E) or cabozantinib (Cabo) in patients (pts) with clear cell (cc) and papillary (pap) metastatic renal cell cancer (mRCC). *Journal of Clinical Oncology* **36**, 603-603 (2018).

52. G. B. Magill *et al.*, Pharmacological and initial therapeutic observations on 6-diazo-5-oxo-1-norleucine (DON) in human neoplastic disease. *Cancer* **10**, 1138-1150 (1957).

53. A CLINICAL study of the comparative effect of nitrogen mustard and DON in patients with bronchogenic carcinoma, Hodgkin's disease, lymphosarcoma, and melanoma. *J Natl Cancer Inst* **22**, 433-439 (1959).

54. K. M. Lemberg, J. J. Vornov, R. Rais, B. S. Slusher, We're Not "DON" Yet: Optimal Dosing and Prodrug Delivery of 6-Diazo-5-oxo-L-norleucine. *Mol Cancer Ther* **17**, 1824-1832 (2018).

55. M. H. Oh *et al.*, Targeting glutamine metabolism enhances tumor-specific immunity by modulating suppressive myeloid cells. *J Clin Invest* **130**, 3865-3884 (2020).

56. A. R. Hanaford *et al.*, Orally bioavailable glutamine antagonist prodrug JHU-083 penetrates mouse brain and suppresses the growth of MYC-driven medulloblastoma. *Transl Oncol* **12**, 1314-1322 (2019).

57. N. H. Chakiryan *et al.*, Real-World Survival Outcomes Associated With First-Line Immunotherapy, Targeted Therapy, and Combination Therapy for Metastatic Clear Cell Renal Cell Carcinoma. *JAMA Netw Open* **4**, e2111329 (2021).

58. B. I. Rini *et al.*, Pembrolizumab (pembro) plus axitinib (axi) versus sunitinib as first-line therapy for advanced clear cell renal cell carcinoma (ccRCC): Results from 42-month follow-up of KEYNOTE-426. *Journal of Clinical Oncology* **39**, 4500-4500 (2021).

59. A. S. Yamashita *et al.*, The glutamine antagonist prodrug JHU-083 slows malignant glioma growth and disrupts mTOR signaling. *Neurooncol Adv* **3**, vdaa149 (2021).

60. A. R. Mullen *et al.*, Oxidation of alpha-ketoglutarate is required for reductive carboxylation in cancer cells with mitochondrial defects. *Cell Rep* **7**, 1679-1690 (2014).

61. J. Chong, D. S. Wishart, J. Xia, Using MetaboAnalyst 4.0 for Comprehensive and Integrative Metabolomics Data Analysis. *Curr Protoc Bioinformatics* **68**, e86 (2019).

62. S. Y. Bartman CR, Lee WD, TeSlaa T, Jankowski CSR, Wang L, Yang L, Roichman A, Bhatt V, Lan T, Hu Z, Xing X, Lu W, Guo JY, Rabinowitz JD, Slow TCA flux implies low ATP production in tumors. *BioRxiv*, (2021).

63. A. Tasdogan *et al.*, Metabolic heterogeneity confers differences in melanoma metastatic potential. *Nature* **577**, 115-120 (2020).

64. P. Heinrich *et al.*, Correcting for natural isotope abundance and tracer impurity in MS-, MS/MS- and high-resolution-multiple-tracer-data from stable isotope labeling experiments with IsoCorrectoR. *Sci Rep* **8**, 17910 (2018).

65. E. A. Struys, E. E. Jansen, N. M. Verhoeven, C. Jakobs, Measurement of urinary D- and L-2-hydroxyglutarate enantiomers by stable-isotope-dilution liquid chromatography-tandem mass spectrometry after derivatization with diacetyl-L-tartaric anhydride. *Clin Chem* **50**, 1391-1395 (2004).

66. V. Ramaswamy *et al.*, Development of a (1)(3)C-optimized 1.5-mm high temperature superconducting NMR probe. *J Magn Reson* **235**, 58-65 (2013).

67. N. E. Sanjana, O. Shalem, F. Zhang, Improved vectors and genome-wide libraries for CRISPR screening. *Nat Methods* **11**, 783-784 (2014).

Acknowledgments

We are grateful to the patients who provided tissue for the tumorgrafts we studied; to members of the DeBerardinis laboratory for constructive comments on the project and the manuscript; and to Calithera for providing CB-839. This article is subject to HHMI's Open Access to Publications Policy. HHMI laboratory heads

have previously granted a nonexclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their research articles. Pursuant to those licenses, the author-accepted manuscript of this article can be made freely available under a CC BY 4.0 license immediately upon publication. We also thank the Preclinical MRI core service at UTSW, which is supported by CPRIT (RP210099). A portion of this work was performed in the McKnight Brain Institute at the National High Magnetic Field Laboratory's Advanced Magnetic Resonance Imaging and Spectroscopy (AMRIS) Facility, which is supported by National Science Foundation Cooperative Agreement No. DMR-1644779 and the State of Florida.

Funding:

H.H.M.I. Investigator Program (RJD),

NCI Outstanding Investigator Award (R35CA22044901 to RJD),

Kidney Cancer SPORE grant (P50CA19651602 to JB),

2019 Young Investigator Award, Kidney Cancer Association (AKK),

NCI/NIH Predoctoral to Postdoctoral Fellow Transition Award (K00, 4K00CA234650 to AT)

NIH/NCI R01 (R01CA229451 to B.S.S.)

NIH R01 (R01 DK105346, R01 132254 to MM)

Author contributions:
Conceptualization: AKK, RJD
Methodology: AKK, LKB, AT, MR, CYW, XL, KA, VTT, FS, QND, CY, TR, CS, AR, BF, PP, HV, FC, MM, BS, TPM, PK, IP, JB
Investigation: AKK, AT, MR, CYW, LGZ, FC, GG
Visualization: AKK, AT, MR, CYW, QND, XS, RJD
Supervision: RJD
Writing—original draft: AKK, RJD
Writing—review & editing: AKK, AT, MR, XS, MM, JB, RJD

Competing interests: R.J.D. is a scientific advisor for Agios Pharmaceuticals, Nirogy Therapeutics, Droia Ventures and Vida Ventures, and a founder and advisor for Atavistik Bioscience. AKK is a current employee of Pfizer. J.B. is an

employee/paid consultant for Arrowhead, Calithera, Esai, Exelixis, and Johnson & Johnson and reports receiving commercial research grants from Arrowhead. J.B. and X.S. have a patent application on [^{18}F]PT2385. Author B.S.S. is an inventor on patent applications filed by Johns Hopkins Technology Ventures covering novel glutamine antagonists, including JHU083, and their utility which have been licensed to Dracen Pharmaceuticals. B.S.S. is also a co-founder of and a consultant for Dracen Pharmaceuticals. This arrangement has been reviewed and approved by Johns Hopkins University in accordance with their respective conflict of interest policies. All other authors declare they have no competing interests.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Requests for other relevant data should be submitted to the corresponding author, and it can be provided by RJD pending scientific review

Figure Legends

Figure 1. **ccRCC tumorgrafts recapitulate metabolic signatures of human ccRCC**

(A). Correlation plot of metabolites altered in both ccRCC tumorgrafts and human tumors. In both studies, differential metabolites between tumor and kidney were calculated using Student's t-test and FDR corrected (Q value<0.05). T-statistic (T-stat) was used as a surrogate for z-score to generate the correlation plot using ggplot in R. Positive T-stat values indicate metabolite elevation in tumors and negative T-stat values indicate metabolite depletion in tumors. Red circles are metabolites altered in the same direction in tumorgrafts and human ccRCC, and gray triangles are metabolites altered in opposite directions.

(B). Boxplots of relative abundance of metabolites in RCC tumorgrafts of different histological types, including ccRCC, *FH*-deficient RCC, papillary RCC (pRCC),

translocation RCC (tRCC), and unclassified RCC (uRCC). One-way ANOVA coupled with pairwise t-test in R software was used to calculate statistical significance and P values were FDR corrected.

(C). Illustration of carbon flow from [U-^{13}C]glucose. Pyruvate fuels the TCA cycle via pyruvate dehydrogenase (PDH) (dark blue circles) and pyruvate carboxylase (PC) (light blue circles).

(D). Plots showing percent enrichment of metabolites relative to [U-^{13}C]glucose (left); and the ratio of M+2 (middle) or M+3 (right) TCA cycle metabolites relative to pyruvate M+3 as surrogates of PDH and PC activity, respectively. Eight mice bearing three distinct orthotopic tumorgrafts (XP258, n=3; XP374, n=3; XP490, n=2) were infused with [U-^{13}C]glucose for 3 hours. M is the mass of the unlabeled metabolite. P-values were calculated using Student's t-test.

P values: ***<0.001, **<0.01, *<0.05

Figure 2. **Glutamine is a carbon source for the TCA cycle in ccRCC tumorgrafts.**

(A). Relative glutamate M+5 in tumors and kidneys from mice infused with [U-^{13}C]glutamine. One-way ANOVA was used to assess statistical significance.

(B). Oxidative and reductive carbon flow from glutamine (light and dark blue circles, respectively).

(C-D). TCA cycle labeling in XP258 (C) and XP296 (D) tumorgrafts, adjacent kidney (labeled as normal kidney), and contralateral kidney. The experiment was conducted in at least 3 mice per tumorgraft. One-way ANOVA coupled with a pairwise t-test was used to assess significance of ^{13}C enrichment between tissues.

(E)**.** Enrichment of citrate M+5 across all models. The experiment was conducted in a minimum of 3 mice per model. One-way ANOVA coupled with a pairwise t-test was used to assess statistical significance of ^{13}C enrichment in citrate M+5 between tissues.

(F). TCA cycle intermediate labeling in orthotopic tumors, subcutaneous tumors and kidneys. Data are from three mice implanted with both orthotopic and subcutaneous XP955 tumorgrafts. One-way ANOVA was used to assess the statistical significance of ^{13}C enrichment.

(G). NAD$^+$/NADH ratios in tumorgrafts and kidneys. The number (n) of kidneys (K) and tumors (T) are indicated. Student's t-test was used to assess the significance.

(H). ^{11}C signal relative to 1.5 min (time of maximal signal in the kidney) in kidney and tumors. One-way ANOVA was used to determine statistical significance.

P values: ****<0.0001,***<0.001, **<0.01, *<0.05

Figure 3. **IDH1 and IDH2 regulate reductive carboxylation and ccRCC growth.**

(A). Percent enrichment of ^{13}C relative to [U-^{13}C]glutamine in TCA cycle from orthotopic XP258-derived orthotopic tumorgrafts, cell lines, and subcutaneous xenografts. Tumor-bearing mice were infused with [U-^{13}C]glutamine for 5 hours and XP258 cells were labeled in culture with [U-^{13}C]glutamine for 4 hours.

(B). Western blot showing the abundance of IDH1 and IDH2 in single and double knockout (IDH1 + IDH2 KO) cells generated using lentiviral CRISPR-Cas9. EGFP is a control cell line with a gRNA targeting EGFP.

(C). Viable cell content in EGFP, IDH1 KO, IDH2 KO, and IDH1+IDH2 KO XP258 cells assessed using Promega CellTiter-Glo®.

(D). Fractional enrichment of ^{13}C-labeled isotopologues of citrate and malate in cells after 4 hours of culture with [U-^{13}C]glutamine. One-way ANOVA was used to assess the significance of ^{13}C enrichment in citrate and malate (n=3).

(E). Percentage enrichment of citrate M+4 and citrate M+5 relative to [U-^{13}C]glutamine in EGFP, IDH1 KO, IDH2 KO, and IDH1+IDH2 KO XP258 tumors. Mice were infused with [U-^{13}C]glutamine for 5 hours. Data are from three or more xenografts generated subcutaneously in NOD-SCID mice. One-way ANOVA with a pairwise t-test was used to determine the statistical significance.

(F). Growth of EGFP, IDH1 KO, and IDH2 KO subcutaneous XP258 xenografts in NOD-SCID mice (n=10, each cell line). P values were calculated using a mixed-effect two-way ANOVA that assessed the significance of differences in tumor growth over time between EGFP vs. IDH1 KO and EGFP vs. IDH2 KO tumors.

P values: ***<0.001, **<0.01, *<0.05

Figure 4. **CB-839 variably decreases glutamine metabolism and ccRCC growth.**

(A). Percent enrichment of ^{13}C-labeled TCA cycle intermediates relative to [U-^{13}C]glutamine in the oxidative (top) and reductive (bottom) pathway in XP296 tumorgraft-bearing mice treated with 200 mg/kg of CB-839 or vehicle for 21 days. Student's t-test was used to assess the statistical significance of ^{13}C enrichment between CB-839 and vehicle-treated tumors (n≥4).

(B) Similar to A, but for XP490 tumorgrafts.

(C) Similar to A, but for XP258 tumorgrafts.

(D) Percent enrichment of glutamate M+5 or M+6 (as either [U-^{13}C]glutamate or [U-^{13}C, ^{15}N]glutamate), M+5 α-KG and M+4 or M+3 malate in XP258 xenografts from CB-839 and vehicle-treated animals infused with [U-^{13}C, ^{15}N]glutamine for 5 min, 15 min and 30 min. Student's t-test was used to calculate the p values.

(E). Spider plot of growth of subcutaneous XP296 tumorgrafts in NOD-SCID mice treated twice daily with 200 mg/kg of CB-839 or vehicle. Treatment was started when tumors reached 100-200 mm^3. Data are plotted from two independent experiments. The Student's t-test was used to calculate the statistical significance of tumor growth at each time point (n=10, each treatment arm).

(F). Similar to E, but for XP490 tumorgrafts.

(G). Similar to E, but for XP258 tumorgrafts

P values: ***<0.001, **<0.01, *<0.05

Figure 5. **JHU-083 decreases glutamine-dependent nitrogen metabolism and ccRCC growth.**

(A)**.** Percent enrichment of ^{15}N-labeled asparagine (Asn), guanosine, and cytidine monophosphate (CMP) relative to [U-^{13}C,^{15}N]glutamine in XP258 tumorgrafts treated with 1.83 mg/kg of JHU-083 or vehicle for 5 days. Student's t-test was used to assess the statistical significance of ^{15}N enrichment between JHU-083 and vehicle-treated tumors (n≥4).

(B). Percent enrichment of ^{13}C-labeled TCA cycle intermediates relative to [U-^{13}C, ^{15}N]glutamine in XP258 tumorgrafts treated with JHU-083 and vehicle in panel A.

Student's t-test was used to assess the statistical significance of ^{13}C enrichment between JHU-083 and vehicle-treated tumors (n≥4).

(C). Left: Spider plot of growth of subcutaneous XP258 tumorgrafts in NOD-SCID mice treated with 1.83 mg/kg of JHU-083 or vehicle for 25 days (5 days on and two days off treatment). Treatment was started when tumors reached ~50 mm^3. P values were calculated using a mixed-effect two-way ANOVA that assessed the significance of differences in tumor growth over time between Vehicle vs. JHU-083 in both xenografts. Right: Same as left panel, but for XP296 tumorgrafts treated with 1.83 mg/kg of JHU-083 and vehicle for 21 days.

(D) Growth curves of tumors in C57BL/6 mice generated from 17175 (right) and 10950 (left) cell lines treated with JHU-083 and vehicle. Mice were treated with 0.915 mg/kg of JHU-083 or vehicle for 21 days or more. P values were calculated using a mixed-effect two-way ANOVA that assessed the significance of differences in tumor growth over time between Vehicle vs. JHU-083 in both xenografts.

P values: ***<0.001, **<0.01, *<0.05

Figure 1

Figure 2

Figure 3

Figure 5