Excitatory amino acid transporter 1 supports adult hippocampal neural stem cell self-renewal.

Joshua D. Rieskamp[1], Ileanexis Rosado-Burgos[1], Jacob E. Christofi[1], Eliza Ansar[1], Dalia Einstein[1], Ashley E. Walters[1], Valentina Valentini[1,4], John P. Bruno[1,2], Elizabeth D. Kirby[1,2,3*]

[1]Department of Psychology, [2]Department of Neuroscience, [3]Chronic Brain Injury Program, The Ohio State University, USA, [4]Department of Biomedical Sciences, University of Cagliari, Italy

*Corresponding author email: kirby.224@osu.edu

Summary:

Within the adult mammalian dentate gyrus (DG) of the hippocampus, glutamate stimulates neural stem cell (NSC) self-renewing proliferation, providing a link between adult neurogenesis and local circuit activity. Here, we show that glutamate-induced self-renewal of adult DG NSCs requires glutamate transport via excitatory amino acid transporter 1 (EAAT1) to stimulate lipogenesis. Loss of EAAT1 prevented glutamate-induced self-renewing proliferation of NSCs in vitro and in vivo, with little role evident for canonical glutamate receptors. Transcriptomics and further pathway manipulation revealed that glutamate simulation of NSCs relied on EAAT1 transport-stimulated lipogenesis, likely secondary to intracellular Ca^{2+} release and glutamate metabolism. Our findings demonstrate a critical, direct role for EAAT1 in stimulating NSCs to support neurogenesis in adulthood, thereby providing insights into a non-canonical mechanism by which NSCs sense and respond their niche.

Keywords: neural stem cell, adult neurogenesis, glutamate, excitatory amino acid transporter, hippocampus

Introduction

The dentate gyrus of the hippocampus (DG) is one of a few, isolated brain regions that can support resident neural stem cells (NSCs) that generate new neurons in adult mammals. Adult hippocampal NSCs are distinct from developmental counterparts[1–3], and their persistence through adulthood is conserved across most land-born mammals, likely including primates[4–7] (but see:[8,9]). The excitatory neurotransmitter glutamate is a potent stimulator of NSC proliferation in the adult DG, a link which suggests a mechanism for tethering adult neurogenesis to circuit activity, as well as a potential mechanism for regeneration in excitotoxic conditions[10,11].

Recent work shows that glutamate promotes asymmetric, self-renewing NSC divisions[12–14]. However, the molecular mediator by which glutamate stimulates NSCs is unclear. Early studies found that NDMA receptor antagonists actually increase NSC proliferation, rather than decrease it[15–18]. More recently, it was shown that triple transgenic knockout of AMPA, NMDA and metabotropic glutamate receptors in NSCs had no effect on their proliferation[14]. Together, previous work suggests that the NSC response to glutamate is unlikely to rely on simple receptor stimulation.

A major component of glutamate regulation in the adult mammalian brain is the family of excitatory amino acid transporters (EAATs). EAATs are transmembrane transporters that shuttle glutamate from the extracellular to intracellular environment via ion-coupled transport. Robust EAAT expression, in particular EAAT1, is a widely-used marker of adult NSC phenotype[19,20] and

in vivo transport-associated currents suggest these transporters are present and active in the adult DG NSC membrane[21–23]. The functional role of EAATs in NSC regulation has received little attention to date. To better understand how adult NSCs interact with the glutamatergic environment of the adult mammalian DG, we set out to uncover the functional role of EAATs in these cells.

Results

High expression of EAAT1, but not other EAATs, is widespread in NSCs and declines rapidly with differentiation

The EAAT family of glutamate transporters has 5 primary members, EAAT1, 2, 3, 4 and 5 (gene names Slc1a3, Slc1a2, Slc1a1, Slc1a6, Slc1a7). Adult DG NSCs have been reported to express both EAAT1 and EAAT2[21]. Using single cell RNAseq (scRNAseq) data from Shin et al.[24], we confirmed that adult DG NSCs express EAAT1 (Slc1a3) and EAAT2 (Slc1a2) but not the remaining 3 EAATs (Fig S1A,B). EAAT2 expression was ~5.35x lower than EAAT1. To verify the extent of EAAT1 and 2 expression throughout NSCs in vivo, we used immunolabeling in adult EAAT1dsRed and EAAT2GFP transcriptional reporter mice[21] (Fig 1A,B, S1C). ~53% of

Figure 1: NSC express EAAT1 and to a lesser extent EAAT2

A) Cells in the neurogenic cascade along with immunophenotypical markers used to identify radial glial like NSCs (RGL NSC), intermediate progenitor cells (IPCs), neuroblasts (NB), immature neurons (IN) and mature neurons.

B) Immunolabeling for Nestin, Ki67 and DCX in DG from EAAT1DsRed+EAAT2GFP+ adult mice.

C) Proportion of cells expressing EAAT1DsRed+EAAT2GFP+ in adult DG. *p<0.05,**p<0.01 Tukey's posthoc test. Mean ± SEM of N = 4-5 mice.

D) EAAT1 immunolabeling in NestinGFP+ NSC, with Sox2+ nucleus. Merged z-stack on left, 3D reconstruction of z-stack on right.

All scales = 10 μm.

Nestin+ radial glia-like NSCs were positive for both EAAT1dsRed and EAAT2GFP, with another ~27% positive for EAAT1dsRed alone (Fig 1B,C, S1D). Less than 2% of Nestin+ NSCs were positive for EAAT2GFP alone, leaving ~18% of Nestin+ NSCs negative for either transporter. Ki67+ proliferative cells in the subgranular zone, which are primarily composed of intermediate progenitor cells (IPCs), and doublecortin (DCX+) neuroblasts and immature neurons each showed progressively smaller proportions of cells expressing EAATs. Using 3D image reconstruction of EAAT1 immunolabeling in adult NestinGFP mice[25], as well as nestin immunolabeling in EAAT1dsRed mice, we also showed EAAT1 protein putatively co-localizing throughout the NSC cell soma and in the bushy terminals in the molecular layer (Fig 1D, S1E). Together, these data suggest that EAAT1, and to a lesser extent EAAT2, are expressed throughout adult DG NSCs, with rapid downregulation upon differentiation.

Glutamate stimulates adult NSCs via EAAT-mediated transport.

To begin investigating the role of EAATs in NSCs, we used cultured NSCs derived from adult male and female DG[26]. These cultures are composed primarily of quiescent and cycling NSCs when maintained in standard conditions[27]. Using our published bulk RNAseq and liquid chromatography tandem mass spectrometry data from these cells[27], we first confirmed that cultured NSC expression of glutamate transporter and receptor genes was similar to that of in vivo NSCs, with EAAT1 being most abundant, followed by EAAT2, then lower but detectable expression levels of AMPA receptor subunits and metabotropic glutamate receptor transcripts (Fig S2A-C). We also confirmed that glutamate stimulated NSC proliferation, as revealed by incorporation of the thymidine analog ethynyl deoxyurdine (EdU) (Fig 2A,D).

To determine the role of glutamate receptors versus transporters in NSC proliferation, we treated NSCs with several pharmacologic inhibitors. Treatment with a cocktail of inhibitors to block AMPA, NMDA and mGluR5 (NBQX, D-AP5, and ACDPP, referred to together as αGlut R) had no effect on basal NSC proliferation nor their response to glutamate (Fig 2B,E). The panEAAT inhibitor TFB-TBOA (αpanEAAT), in contrast, blocked glutamate-induced proliferation and did so whether glutamate receptors were inhibited or not. Aspartate, an alternate transport substrate for both EAAT1 and 2, also stimulated NSC proliferation in an EAAT-dependent, receptor-independent manner (Fig 2C, S2D). To further explore the role of glutamate receptors in NSC proliferation, we treated NSCs with glutamate receptor agonists: AMPA, KA, NMDA and CHPG (mGluR5 agonist). None of the agonists increased proliferation (Fig 2F, S2E). NMDA slightly decreased proliferation. CHPG (mGluR5) agonist caused massive cell death at its higher dose and had no effect at its lower dose. These findings suggest, unexpectedly, that EAAT transporter activity alone mediates glutamate-induced NSC proliferation with no, or possibly a countering, contribution of glutamate receptors.

Glutamate stimulation of adult NSCs relies on EAAT1 not EAAT2.

To determine which EAATs might be active in NSCs, we measured glutamate clearance from NSC cultures using ultra-high performance liquid chromatography with electrochemical detection (uHPLC-ECD) of conditioned media. Within 20 minutes, NSCs cleared 82% of a 5 μM glutamate pulse from the media (Fig S2F). Treatment with the selective EAAT1 inhibitor (UCPH101, αEAAT1) or a combination of αEAAT1 and the selective EAAT2 inhibitor (WAY213613, αEAAT2) completely prevented glutamate clearance compared to vehicle treated controls (Fig 2G). αEAAT2 alone had no effect on clearance compared to vehicle treated controls. To confirm activity of the αEAAT2 drug, we replicated the experiment with a higher

Figure 2: Glutamate stimulates NSC proliferation via EAAT1.
A-C) EdU+ cell counts after treatment with: glutamate, inhibitors of glutamate receptors (αGlutR, NBQX, D-AP5, and ACDPP), inhibitor of EAATs (αpanEAAT, TFB-TBOA), aspartate, inhibitor of NMDA receptor (αNMDAR). Mean ± SEM of N = 2-5 replicates/experiment, 3-4 independent experiments.
D-E) Representative images of EdU+ labeling in cultured NSCs after glutamate treatment +/- receptor and transporter inhibitors, with Hoechst to label nuclei. Scale = 20 μm.
F) EdU+ cell counts after treatment with glutamatergic agonists. Mean ± SEM of N = 6-12 replicates/experiment, 4 independent experiments.
G) Glutamate in NSC conditioned media as measured by uHPLC after a 5 μM pulse with pretreatment with inhibitors of EAAT1 (UCPH102), EAAT2 (WAY213613) or their combination (EAAT1/2). Mean ± SEM of N = 3 independent experiments.
H) EdU+ cell counts after treatment with glutamate +/- inhibitors of EAAT1 (UCPH102), EAAT2 (WAY213613), or panEAAT (TFB-TBOA). Mean ± SEM of N = 5 replicates/experiment, 3 independent experiments.
All EdU+ counts expressed relative to 0 μM glutamate/vehicle within experiment. *,**,***,****p<0.05, <0.01, 0.001, 0.0001 posthoc comparisons.

dose that also inhibits EAAT1. Higher dose αEAAT2 partially prevented glutamate clearance, suggesting that the compound is active but only shows effects at doses that inhibit EAAT1 (Fig S2G). Similar results were found with an alternative EAAT2 inhibitor (DHK) (Fig S2H). Taken together, these results demonstrate functional glutamate clearance by NSCs and implicate EAAT1 as the dominant transporter.

We next treated cultured NSCs with αEAAT1, αEAAT2 or αpanEAAT and measured proliferation. Both EAAT1 and panEAAT inhibition each prevented glutamate-induced NSC

proliferation while EAAT2 inhibition alone had no effect (Fig 2H, S2I). Throughout these experiments, cells in all groups appeared healthy and showed no observable signs of apoptosis (fragmented nuclei, loss of plate adhesion). We also confirmed that neither αpanEAAT nor αEAAT1 led to signs of cell death using a Sapphire700 IR dye assay (Fig S2J). Together, these data suggest that glutamate stimulates NSC proliferation directly via EAAT1 activity.

EAAT1 knockdown causes cell-autonomous loss of NSC self-renewal in vivo

To examine the role of EAAT1 in DG NSCs in vivo, we used CRISPR interference (CRISPRi), a transcriptional interference strategy based on the CRISPR/cas9 system. In CRISPRi, a catalytically dead Cas9 protein (dCas9) fused to a Kruppel associated box (KRAB) domain represses transcription of a gene targeted by a highly specific single guide (sg)RNA. We designed a U6→sgRNA against EAAT1 (EAAT1 KD) (and a non-target sgRNA (NT)) then inserted it into a lentiviral CRISPRi backbone[28] with dCas9-KRAB-T2A-GFP (Fig 3A). Cultured NSCs infected with the EAAT1 KD CRISPRi vector showed a 53.1% loss of EAAT1 immunoreactivity and significant suppression of proliferation compared to NT infected cells (Fig S3A-C). We stereotaxically infused the EAAT1 KD and NT CRISPRi lentiviral vectors in the DG of adult mice. 1 week later, no loss of EAAT1 protein was detected (Fig S3E,F). 3 weeks later, a 40.6% knockdown of EAAT1 immunoreactivity was detected in EAAT1 KD compared to NT treated controls (Fig S3G,H). We therefore next assessed the effect of EAAT1 loss on the neurogenic cascade at this 3 week timepoint.

In this paradigm, EAAT1 knockdown could alter NSCs via receptor-independent, cell-autonomous loss of glutamate transport, as our in vitro data suggests, or through increased glutamate stimulation of receptors due to extracellular glutamate buildup, or through a combination of the two. To investigate the cell autonomous effect, we compared dCas9-KRAB-GFP+ NSCs (hereafter GFP+ NSCs) to neighboring GFP- NSCs. While changes in ambient glutamate would affect both GFP+ and GFP- NSCs, only the GFP+ NSCs would be lacking cell autonomous glutamate transport. At 3 weeks after viral infusion, EAAT1 KD mice had 53.9% fewer GFP+GFAP+SOX2+ radial glia-like NSCs compared to NT treated controls (Fig 3B,C). In contrast, the neighboring GFP- NSCs in these mice showed no effect of EAAT1 KD. The number of proliferating (EdU+) GFP+GFAP+ radial glia-like NSCs showed the same pattern, with a 63.5% decrease in density of proliferating GFP+ NSCs with EAAT1 KD but no effect on neighboring GFP- proliferating NSCs compared to NT control mice (Fig 3D,E). No differences were found between NT and EAAT1 KD mice in GFP+ or GFP- GFAP-SOX2+ IPC density nor in density of DCX+ neuroblasts/immature neurons that were BrdU-labeled 2 weeks before perfusion (Fig 3F,G, S3I). These findings suggest that loss of EAAT1 causes a cell-autonomous loss of radial glia-like NSC self-renewal.

To determine whether the loss of NSCs observed at 3 weeks after knockdown would result in loss of adult neurogenesis at longer timepoints, we perfused mice 2 months after NT or EAAT1 KD CRISPRi lentiviral infusion (Fig 3H). EAAT1 KD mice showed a 63.0% loss of GFP+ radial glia-like NSCs with no effect on GFP- NSCs compared to NT control mice (Fig 3I,K). Similar to NSCs, GFP+ IPC density was 54.3% lower in EAAT1 KD mice compared to NT control mice, while density of GFP- IPCs did not differ between EAAT1 KD and NT mice (Fig 3J). EAAT1 KD mice also showed a 73.6% reduction in number of GFP+NeuN+ newborn neurons labeled with BrdU 1 month before perfusion and no effect on GFP-NeuN+BrdU+ cells (Fig 3L,M, S3J). These data suggest that loss of EAAT1 results in loss of adult neurogenesis that is restricted to the progeny of NSCs that lost EAAT1 and does not extend to neighboring NSCs and their progeny.

Figure 3: EAAT1 knockdown causes cell-autonomous loss of NSC self-renewal

A) CRISPRi lentiviral vector design and timeline. Inset image show representative GFP+ labeling in DG at 3 weeks after infusion. Scale = 100 μm.

B,K) Representative images of GFAP, SOX2, GFP immunolabeling for quantification of NSCs and IPCs 3 weeks (B) and 2 mo (K) after infusion. GFP+ and GFP- NSCs are outlined in C. IPCs are also shown by arrows (GFP-) or arrowheads (GFP+) in K. Scale = 20 μm. (legend continued on next page)

Figure 3 (cont): EAAT1 knockdown causes cell-autonomous loss of NSC self-renewal
C-D) Density of GFP+ and GFP- GFAP+SOX2+ total (C) and EdU+ (D) radial glia-like NSCs in NT and EAAT1 KD mice 3 weeks after infusion.
E) Representative image of an EdU+GFP+GFAP+SOX2+ radial glia-like NSC 3 weeks after infusion. Scale = 10 μm.
F,G) Density of GFP+ and GFP- GFAP-SOX2+ IPCs (F) and DCX+BrdU+ newborn neuroblasts/immature neurons (G) in NT and EAAT1 KD mice 3 weeks after infusion.
H) Timeline for panels I-M
I,J) Density of GFP+ and GFP- GFAP+SOX2+ radial glia-like NSCs (I) and GFAP-SOX2+ IPCs (J) in NT and EAAT1 KD mice 2 months after infusion.
K) Density of GFP+ and GFP- BrdU+ cells (NeuN+ and NeuN-) in NT and EAAT1 KD mice 2 months after infusion.
L) Representative images of BrdU+ cells.
All densities relative to NT control within GFP type. Arrowhead = BrdU+NeuN+GFO+, Arrow = NeuN+BrdU+, Dashed arrow = BrdU+. Scale = 10 μm.
C,D,F,G: Mean ± SEM of N = 8 individual mice (points shown). *,**p<0.05, 0.01 Sidak's multiple comparisons within GFP label type.
I,J,L: Mean ± SEM of N = 6-8 individual mice (points shown). *,**p<0.05, 0.01 Sidak's multiple comparisons within GFP label type

EAAT-dependent stimulation of fatty acid synthetase is essential for glutamate-induced NSC proliferation.

To gain insight into the molecular mediators of EAAT-dependent stimulation of NSC self-renewal, we performed bulk RNAseq on RNA from cultured NSCs treated with glutamate and/or panEAAT inhibitor (Fig S4A). In the absence of αpanEAAT, glutamate treatment caused significant upregulation of 106 transcripts and significant downregulation of 49 transcripts (Table S1). In contrast, in the presence of αpanEAAT, glutamate treatment yielded no upregulated transcripts and only 1 downregulated transcript. These data suggest robust NSC transcriptional response to glutamate that is prevented by EAAT inhibition.

To understand the functional significance of genes regulated by glutamate transport, we focused on GO biological process analysis of genes upregulated by glutamate that were also downregulated by αpanEAAT (Table S2). We found 22 overlapping GO terms which centered mostly on metabolic processes related to lipid, amino acid, and protein synthesis (Fig 4A). Particularly notable among the cellular processes implicated in EAAT-dependent regulation of NSC gene expression was lipid biosynthetic process. Lipogenesis is essential for NSC self-renewal, with a particularly critical role for fatty acid synthetase (FASN)[29–31]. In our RNAseq data, glutamate induced upregulation of Fasn, an effect that EAAT inhibition blocked (Table S1,S3). Independent replication using qPCR confirmed that glutamate stimulated Fasn expression in an EAAT-dependent manner (Fig 4B). FASN inhibition with the inhibitor orlistat also prevented glutamate-induced NSC proliferation (Fig 4C, S4B). These findings suggest that glutamate transport by EAAT promotes NSC proliferation via stimulation of FASN.

Glutamate stimulation of NSCs requires both transport-associated Ca^{2+} signaling and glutamate metabolism.

We next sought to determine the upstream signals that could mediate NSC proliferation in response to glutamate transport. In Bergmann and Müller glia, glutamate transport can act as an intracellular signaling event by triggering a Ca^{2+}-dependent signaling cascade culminating in phosphorylation and activation of mTOR[32,33], which itself promotes de novo lipogenesis and proliferation in adult stem cells, including NSCs[34,35]. Similar to these specialized glial cells, we found that glutamate induced mTOR phosphorylation in adult DG NSCs and that effect was blocked by EAAT inhibition (Fig 4D, S4C). In Bergmann and Müller glia, the Ca^{2+} signal that

drives mTOR phosphorylation derives from the co-transport of $3Na^+$ with glutamate via EAATs, which causes reversal of the Na^+-Ca^{2+} transporter, NCX, to export Na^+ and import Ca^{2+}. In adult NSCs, though, treatment with either of 2 NCX inhibitors, SEA0400 or YM 244769, had no effect on basal or glutamate-stimulated NSC proliferation (Fig S4D-F). These data suggest that Ca^{2+} influx via NCX reversal is not a likely mediator of EAAT-dependent stimulation of proliferation.

We next examined an alternate source of Ca^{2+} release: the Ca^{2+} stores of the endoplasmic reticulum (ER). Glutamate transport via EAATs is accompanied by net depolarization due to co-transport with $3Na^+$ and $1H^+$ (plus export of $1K^+$). Cellular depolarization is a potent stimulator of

Figure 4: Glutamate stimulates NSC self-renewal via transport-stimulated lipogenesis

A) 22 GO biological processes that were upregulated by glutamate and downregulated panEAAT inhibition (TFBTBOA) in RNAseq. -logPadj shown. N = 3 replicates/group.

B) Fasn mRNA expression as measured by real time qPCR after treatment with glutamate +/- panEAAT inhibition (TFBTBOA). **,****$p<0.01$, 0.0001 Sidak's multiple comparisons. Mean ± SEM of N = 3 replicates/experiment, 3 independent experiments.

C) EdU+ cell counts after treatment with glutamate +/- inhibitor of FASN (orlistat). Mean ± SEM of N = 4 replicates/experiment, 4 independent experiments. *$p<0.05$ Sidak's multiple comparisons.

D) Phosphor-mTOR/total mTOR in NSCs treated with glutamate +/- panEAAT inhibition (TFBTBOA). *,**$p<0.05$, 0.01 Tukey's multiple comparisons. Mean ± SEM of N = 2-3 replicates/experiment, 3 independent experiments.

E) EdU+ cell counts after treatment with glutamate +/- inhibitors of ryanodine receptor (αRYR, dantrolene) and glutamate dehydrogenase (αGDH, EGCG). *,**,***$p<0.05$, 0.01, 0.001 Sidak's multiple comparisons. Mean ± SEM of N = 4 replicates/experiment, 4 independent experiments.

Ca^{2+} release from the endoplasmic reticulum (ER) via the ER-bound ryanodine receptor (RyR)[36,37]. Blocking RyR alone with the inhibitor dantrolene (αRYR) did not significantly affect NSC proliferation (Fig 4E, S4G).

We next considered a role for glutamate as a substrate for metabolism. Conversion of glutamate to α-ketoglutarate directly fuels lipogenesis by contributing to TCA cycle generation of acetyl-CoA, which is a primary substrate for FASN-mediated de novo fatty acid synthesis. Inhibition of glutamate dehydrogenase (αGDH), the key enzyme for conversion of glutamate to α-ketoglutarate, had no effect on NSCs alone, but when combined with RyR inhibition, glutamate-induced proliferation was completely blocked (Fig 4E, S4G). These data suggest that release of ER Ca^{2+} and glutamate substrate metabolism jointly mediate glutamate transport-induced stimulation of NSC self-renewal (Fig S4H).

Discussion

Our findings suggest a functional, cell-autonomous role for EAAT1 in supporting adult hippocampal neurogenesis. Our findings stand in contrast to a canonical glutamate receptor-dependent mechanism for glutamate response reported in most other cell types and suggest that EAAT1 expressed by NSCs directly supports self-renewing proliferation via transport-dependent stimulation of lipogenesis.

Expression of glutamate transporter proteins, particularly EAAT1, is widespread in the radial glia NSCs of the embryonic forebrain[38]. One study in early postnatal DG NSCs concluded that EAATs indirectly suppress proliferation by limiting the activation of mGluRs[39]. This apparent reversal of the roles of EAAT1 and mGluR5 in early postnatal versus what we observed in adult DG NSCs is paralleled by downregulated expression of mGluR5 and upregulated expression of EAAT1 in DG NSCs during the transition to adulthood[40]. This transition could contribute to the shift in cellular behavior of DG NSCs from rapidly proliferating and depleting early in life to more maintenance-focused later in life[41]. Comparative studies of EAAT function in NSCs across different stages of development could shed further light on this hypothesis.

Our data support depolarization-sensitive RyR-mediated Ca^{2+} release as a process downstream of glutamate signaling that can mediate glutamate-induced NSC proliferation but only in combination with glutamate conversion to the TCA cycle substrate α-ketoglutarate. The putative need for glutamate as a metabolic substrate to stimulate NSC self-renewal provides a parsimonious explanation for why NSC depolarization secondary to glutamate receptor activation alone would not be as effective as glutamate itself in stimulating NSC proliferation[14].

Cell metabolic states are becoming increasingly recognized as active players in NSC self-renewal and fate decisions, both as generators of key cellular substrates for growth and as independent initiators of gene expression[42–44] . Several previous studies demonstrate a particularly critical role for fatty acid synthesis in DG NSC self-renewal[29–31]. In agreement with this previous data, we found that FASN activity was required for glutamate-induced NSC proliferation. Cell metabolism therefore represents a potential convergence point for the various glutamate-stimulated, EAAT-dependent processes that we observed in adult DG NSCs.

NSCs may derive glutamatergic input from multiple sources in vivo. NSC apical processes wrap around putatively glutamatergic synaptic terminals in the molecular layer of the DG[45]. Inputs here from hilar mossy cells[14] and hypothalamus[13] both stimulate NSC self-renewal, as does release from local astrocytes[12]. With multiple sources of glutamate, it is tempting to speculate

that changes in hippocampal glutamate signaling that occur during normal development, neurodegenerative pathology, and plasticity-inducing stimuli[46–49] could be transduced by EAAT1 into metabolic states that regulate the population maintenance of NSCs. Future studies could examine whether changes in EAAT expression or function modulate the response of NSCs to these physiological states.

In summary, we show that glutamate transport via EAAT1 cell-autonomously supports DG NSC self-renewal in adult mice. These findings help resolve some discrepancy in the field about the role of glutamate receptors versus glutamate itself in NSC maintenance. They also raise additional questions about the interplay between glutamate signaling and NSC behavior in contexts such as biological aging, neurological disorders, and plasticity-inducing stimuli like exercise and enriched environment[10,50,51].

Limitations of the study

The presented studies only examine NSCs in the adult DG of mice. The NSCs in the adult subventricular zone neurogenic niche also express EAAT1[52], and the role of EAAT1 in this niche remains an open question, to the best of our knowledge. Whether these results extend to other species, including primates, is also not yet clear either. However, a recent scRNAseq profiling of adult macaque hippocampal NSCs show high levels of EAAT1 and EAAT2 expression[53], suggesting the possibility that NSCs in adult primate hippocampus could rely on EAAT-mediated glutamate transport. We also do not resolve which glutamatergic inputs impact NSCs directly via EAAT1, nor how EAATs come in to play in pathological conditions characterized by excess glutamate transmission. Our work also only begins the uncovering of the intracellular signaling between glutamate transport and stimulation of FASN and NSC self-renewal. These steps will require more detailed investigation in the future.

Methods

Animals
All experimental protocols were performed in accordance with institutional guidelines approved by the Ohio State University Institutional Animal Care and Use Committee (and with recommendations of the National Institutes of Health Guide for the Care and Use of Laboratory Animals. For lentiviral infusion, seven week old C57BL/6J mice (JAX # 000664) were obtained from The Jackson Laboratory were allowed to acclimate for 1 week prior to surgery. GLAST/EAAT1-DsRed and GLT-1/EAAT2-GFP mice (Regan et al., 2007) were obtained as a kind gift from Dr. Jeffrey Rothstein and crossed to obtain offspring used in experiments. From the time of weaning, mice were group housed (5 per cage) in ventilated cages with ad libitum access to food and water and maintained on a 12 hour light cycle. Reporter mice aged 8 – 12 weeks were used for tissue staining. Both male and female mice were used in all experiments.

Perfusion and tissue harvest
Mice were injected with a ketamine/xylazine mixture (87.5 mg/kg, 12.5 mg/kg, i.p.) and transcardially perfused with ice-cold PBS. Brains were collected and fixed for 24h in 4% paraformaldehyde in 0.1 M phosphate buffer prior to equilibration for at least 2 days in 30% sucrose in PBS, both at 4°C. They were then sliced on a freezing microtome (Leica) in a 1 in 12 series of 40 μm slices and stored in cryoprotectant at −20°C.

Immunofluorescence and thymidine analog detection in brain tissue

Sections were rinsed 3x with PBS and then incubated in 1% normal donkey serum (Jackson ImmunoResearch), 0.3% Triton X-100 (blocking buffer) for 1 hour before incubating in primary antibodies (Table 9) diluted in blocking buffer overnight at 4°C with rocking. On the second day, sections were rinsed 3x with PBS and incubated in fluorophore-conjugated secondary antibodies diluted 1:500 in blocking buffer for 2 hours before incubation for 10 min in Hoechst 33342 (Fisher, 1:2000 in PBS), except for sections that were instead incubated with AlexaFluor 350 (Invitrogen) secondary antibodies, and a final 3x PBS rinse. They were then mounted on charged glass slides and coverslipped with ProLong Gold fluorescence mounting media.

For EAAT1 immunolabeling, the above procedure was followed except that tissue was pretreated with ice cold methanol for 10 min for antigen retrieval and incubated in primary antibody at 37˚C overnight with shaking to increase antibody penetration.

For nestin immunolabeling, antigen retrieval was performed using citrate buffer, pH = 6.0, 10 min at 95 ˚C prior to blocking and primary antibody incubation.

For detection of thymidine analogs, the staining process above was used with the following modifications. Tissue sections were incubated for 30 minutes in 2 N HCl at 37°C prior to incubation in anti-BrdU primary antibodies. For detection of EdU, adherent cells or tissue sections were permeabilized for 20 minutes in 0.5% Triton X-100 and incubated in the dark for 30 minutes with copper-dependent click reaction mixture (Click Chemistry Tools #1350) to label EdU with Azide-biotin (Click Chemistry Tools #1265) for subsequent Streptavidin-555 (Invitrogen) conjugation.

Antibody dilutions are listed below:

Host	Target	Vendor	Cat #	Dilution
Rabbit	mCherry/DsRed	Abcam	ab167453	1:500
Mouse	mCherry/DsRed	Novus Biologicals	NBP1-96752	1:500
Goat	GFP	Abcam	ab6673	1:1000
Mouse	Nestin	Millipore-Sigma	MAB353	1:100
Rat	KI67	Invitrogen	14-5698-82	1:500
Rabbit	DCX	Cell Signaling	4604	1:500
Rabbit	EAAT1	Abcam	ab416	1:250
Rabbit	GFAP	Dako	ZO334	1:1000
Rat	SOX2	Affymetrix	14-9811	1:1000
Rat	BrdU	BioRad	OT0030	1:500
Mouse	NeuN	Millipore	MAB377	1:500
Rabbit	EAAT1	Abcam	Ab416	1:500

Image acquisition and analysis

All imaging was performed on a Zeiss Axio Observer Z.1 with apotome digital imaging system and Axiocam 506 monochrome camera (Zeiss). Images of cultured NSCs captured using Zen Blue software (Zeiss) were converted to Tag Image File Format (Tiff) for automated cell counting using ImageJ. Tissue sections were imaged using a 20x objective as z-stacks with 20 x 1 µm steps. Images were analyzed using Zen Blue software (Zeiss). RGLs and IPCs, EdU+ cells, DCX+BrdU+ cells and NeuN+BrdU+ cells were counted and/or phenotyped by colocalization with cell type identity markers within the SGZ (or SGZ plus GCL for BrdU+ cells).

Imaging and cell quantification were performed while blind to animal identity. 3D reconstructions were performed using 0.5µm z-stacks taken with a 63x oil objective and then rendered with Imaris software (Oxford Instruments). EAAT1 immunolabeling for quantification of knockdown in brain tissue was done using ImageJ to select GFAP+GFP+ area and measure EAAT1 intensity within that area.

Cell culture

NSCs were derived from the DG of 6 week old C57BL/6J mice (JAX # 000664) mice based on a published protocol[26]. Briefly, mice were euthanized by CO_2 inhalation and brains were rapidly dissected. Bilateral DG were isolated from each brain in ice-cold Neurobasal A media (Gibco) and minced with a scalpel blade prior to 20 min incubation in papain (Roche)/dispase (StemCell Technologies)/DNase at 37 °C. The mixture was then triturated with a P1000 pipette and centrifuged for 2 min at 800 x g. The pellet was resuspended in warm Neurobasal A and mixed with a 22% Percoll suspension prior to centrifugation for 15 min at 450 x g. The resulting pellet was washed several times in warm Neurobasal A prior to resuspension in complete growth media containing 1% B27 supplement (Gibco), 0.5% GlutaMax (Gibco), and 20 ng/mL each of epidermal growth factor (EGF) and basic fibroblast growth factor (FGF2) (Peprotech). NSCs were maintained in adherent monolayer culture on poly-D-lysine (Sigma) and laminin (Invitrogen) coated plates. All experiments used cells between passages 4 and 20 and were replicated in two separate cell lines, one derived from 4 pooled male DG and the other from 4 pooled female DG. Both lines were mycoplasma tested and confirmed to produce neurons and glia upon growth factor withdrawal as previously described (Denninger et al., 2020).

Pharmacological reagents

Reagents and working dilutions are in Table 1 below:

Compound	Vendor	Cat #	Working [ ] (µM)	Diluent
L-glutamic acid (glutamate)	Millipore-Sigma	G8415	5, 10, 100	equimolar NaOH
L-aspartic acid (aspartate)	Tocris	214	10, 100	equimolar NaOH
TFB-TBOA	Tocris	2532	1	DMSO
UCPH-102	Abcam	ab146404	10	DMSO
Dihydrokainic acid (DHK)	Tocris	0111	100	PBS
WAY 213613	Tocris	2652	1, 10	DMSO
NBQX disodium salt	Tocris	1044	10	PBS
D-AP5	Tocris	0106	100	PBS
ACDPP hydrochloride	Tocris	2254	1	DMSO
(S)-AMPA	Tocris	0254	10, 100	PBS
Kainic acid	Millipore-Sigma	420318	10, 100	PBS
NMDA	Tocris	0114	10, 100	PBS
CHPG sodium salt	Tocris	3695	10, 50	PBS
SEA 0400	Tocris	6164	1	DMSO
YM 244769	Tocris	4544	2	DMSO
Dantrolene, sodium salt	Tocris	0507	10	DMSO
EGCG	Tocris	4524	50	DMSO

Proliferation assays

NSCs were plated 5000 cells per well in a 96 well plate and treated with pharmacological inhibitors or vehicle for 15 minutes prior to the addition of glutamate/aspartate or vehicle. Two days later, 20 µM 5-Ethynyl-2′-deoxyuridine (EdU; Click Chemistry Tools 1149) was added to the cell culture media for 30 minutes to label NSCs in S-phase. The cells were then fixed in 4% PFA for 20 minutes prior to undergoing a click reaction to detect EdU. Cells were permeabilized for 20 minutes in 0.5% Triton X-100 and incubated in the dark for 30 minutes with copper-dependent click reaction mixture (Click Chemistry Tools #1350) to label EdU directly with fluorophores (Azide-488 or –647). Cells were stored at 4 °C in the dark until imaging (see above). After imaging, a standard threshold was applied to the channel containing EdU staining for all images, and the analyze particles function was applied to count the cells with EdU signal above threshold. EdU counts are expressed as relative change to control within experiment to facilitate combining multiple experimental replicates. In generally, a minimum of 3 experimental replicates of each EdU proliferation assay was performed, with multiple well replicates per groups in each individual experiment.

Glutamate clearance assay

Glutamate clearance experiments were performed on NSCs grown in 96 well plates to approximately 95% confluency. The growth media was removed and cells were rinsed twice with oxygenated artificial cerebrospinal fluid (aCSF) prepared according to the following recipe (in mM): 116 NaCl, 3.0 KCl, 1.25 NaH2PO4, 23 NaHCO3, 10 glucose, 2.0 MgSO4, and 2.0 $CaCl_2$[54]. Cells were then pre-incubated for 15 minutes with aCSF containing EAAT inhibitors or vehicle before replacing the solution with 50 µL of aCSF containing 5 µM glutamate and EAAT inhibitors or vehicle. Immediately after (T0) and 5, 10, and 20 minutes later, samples (10 µL) were injected and glutamate quantified by uHPLC-ECD.

uHPLC-ECD

Glutamate was quantified using uHPLC-ECD (ALEXYS Neurotransmitter Analyzer;Antec Scientific, USA). Samples were pre-column derivatized with o-phthaldialdehyde (OPA) and sulphite. Chromatographic separation was performed via an Acquity UPLC HSS T3 1.0 x 50 mm, 1.8 µm column (Waters).). The mobile phase A, for separation consisted of 50 mM phosphoric acid, 50 mM citric acid, 0.1 mM EDTA, and 1.5% v/v acetonitrile with a final pH of 3.1. Mobile Phase B, for post-separation flush, consisted of the same mobile phase A but with 50% v/v acetonitrile. The system was operated at a flow rate of 200 µl/min at a pressure of 380 bar. Electrochemical detection used a working potential of 850 mV (vs Ag/AgCl reference) and a range setting of 5 nA/V. Sample peaks were fitted to a glutamate standard curve.

Plasmid design and lentiviral packaging

The CRISPRi lentivirus construct was modified from pLV hU6-sgRNA hUbC-dCas9-KRAB-T2a-GFP (Addgene #71237, a gift from Charles Gersbach[28]), which expresses all necessary CRISPRi machinery (both the dCas9-KRAB and sgRNA) from the same plasmid. The UbC promoter was replaced with an EF1α promoter to create pLV hU6-sgRNA EF1α-dCas9-KRAB-T2A-GFP. Additionally, two Esp3I recognition sites were placed after the U6 promoter for restriction cloning of sgRNA insert sequences. The insert sequences were synthesized as single strand oligonucleotides (Integrated DNA Technologies) in the form of 5' GGACG(N)20 3' and 5' AAAC(N')20C 3' where (N)20 refers to the sequence of the sgRNA and (N')20 is the reverse complement. These oligonucleotides were annealed together and ligated with Esp3I-digested

pLV hU6-sgRNA EF1α-dCas9-KRAB-T2a-GFP to obtain the final constructs for lentivirus packaging. The sgRNA targeting the region -50 to +300 bp relative to the transcriptional start sequence of Slc1a3 were designed using the CRISPRi function of the online Broad Institute GPP portal[55,56]. The top ranked Slc1a3-targeting sgRNA returned by the GPP tool (5'CGGAGTAACAGCTTAGCGAG) and a non-targeting (NT, 5'GCGAGGTATTCGGCTCCGCG) sgRNA[57] were used to create an EAAT1 KD plasmid and an NT control plasmid. These plasmids were packaged as VSV-G pseudotyped, second-generation lentivirus vectors (titers 1.51×10^9 IFU/mL, EAAT1 KD virus; 1.53×10^9 IFU/mL, NT control virus) by Vigene Biosciences, Inc. These plasmids will be made available via addgene upon publication.

CRISPRi KD in vitro
Cultured adult DG NSCs were plated 5000 cells per well in 96 well plates and treated with EAAT1 KD or NT control virus (~MOI 75 - 150). Cells were maintained in culture for 8 – 12 days after virus treatment and were passaged 1 – 2 times during that timeframe. Cells were treated with 100 μM glutamate or vehicle 2 days before a 30 minute pulse of 20 μM EdU and fixation. Immunofluorescence labeling, EdU detection, and imaging were performed as described above. ImageJ was used to manually draw ROIs around GFP+ and GFP- NSC cell bodies in the brightfield channel and measure the average EAAT1 immunofluorescence signal intensity within the ROI. The number of GFP+EdU+ cells was quantified by manually identifying all GFP+ cells and evaluating colocalization with EdU.

Stereotaxic surgery
Mice were anesthetized by isoflurane inhalation (Akorn, 4% induction, 2% maintenance) in oxygen and mounted in the stereotaxic apparatus (Stoelting). The scalp was sterilized with alcohol (Fisher) and betadine (Fisher) before exposing the skull via a single scalpel blade incision. Bilateral bur holes were drilled into the skull, and Hamilton syringes were positioned at A/P -2.0 mm and M/L +2.0 mm before being slowly lowered to a depth of DV –1.9 mm (all coordinates relative to bregma). Mice were administered 0.9 μL EAAT1 KD virus in one hemisphere and 0.9 μL NT control virus in the contralateral hemisphere at a rate of 0.1 μL/min via an automated injector system (Stoelting). The brain hemisphere receiving EAAT1 KD virus was counterbalanced within each timepoint cohort and across sexes. Mice were administered carprofen prior to surgery and daily for 3 days post surgery as well as buprenorphine immediately post surgery. 36 mice total received lentiviral infusion. Cohort 1: 16 (8 male, 8 female) were given 3 injections of EdU (150 mg/kg, i.p.) 4 hours apart 1 week after lentiviral infusion then perfused. GFP expression, indicating proper infusion placement, was confirmed by a blinded observer in 5 NT and 10 EAAT1 KD DGs. Cohort 2: 10 (5 male and 5 female) received a single daily injection of BrdU (150 mg/kg, i.p.; Sigma B5002) on days 6 – 8 after infusion and 3 injections of EdU on day 21 after lentivirus injection. GFP expression was confirmed by a blinded observer in 8 NT and 8 EAAT1 KD DGs. Cohort 3: 10 mice (5 male and 5 female) received a single daily injection of BrdU (150 mg/kg, i.p.; Sigma B5002) 1 month after infusion, 1 mo before 3 EdU injections and perfusion. GFP expression was confirmed by a blinded observer in 8 NT and 6 EAAT1 KD DGs. All cohorts were perfused two hours after the final EdU injection.

Experimental treatment and RNA extraction for RNAseq
NSCs of passage 4 were plated 75,000 cells per well in a 12 well plate and treated with 1 μM TFB-TBOA or vehicle for 15 minutes prior to the addition of 100 μM glutamate or vehicle (n = 3 wells per treatment). After 48 hours, cells were harvested with Accutase (StemCell

technologies) and a subset (approximately 75,000 cells) was pelleted prior to RNA extraction using the NucleoSpin RNA Plus XS kit (Takara) following manufacturer protocol. Assessment of RNA quality with Qubit RNA HS assay kit (Invitrogen) revealed RIN values of at least 8 for all samples.

RNAseq library generation, sequencing, alignment, and transcript quantification
Library generation was performed as previously described[27] using the Clontech SMART-Seq HT (Takara) kit. Purified library products were then submitted to HiSeq 4000 paired-end sequencing (Illumina) with a depth of 15 – 20 million 2 × 150 bp clusters. AdapterRemovalv2.2.0 was used to trim individual FASTQ files for adapter sequences and filter for a minimum quality score of Q20. HISAT2 v2.0.6 was used to exclude reads aligning to a composite reference of rRNA, mtDNA, and PhiX bacteriophage sequences obtained from the NCBI RefSeq. Primary alignment was performed to the mouse reference genome GRCm38p4 using HISAT2. Gene expression values were quantified for all genes described by the GENCODE GeneTransfer Format (GTF) release M14 (mouse) using the featureCounts tool of the Subread package v1.5.1 58 in stranded mode. Raw and processed counts will be made available via GEO upon publication.

Differentially expressed gene (DEG) analysis
DEG analysis was performed using pair-wise comparisons between the following treatment groups: vehicle/vehicle vs glutamate/vehicle, vehicle/vehicle vs vehicle/TFB-TBOA, glutamate/vehicle vs glutamate/TFB-TBOA, and vehicle/TFB-TBOA vs glutamate/TFB-TBOA. For all comparisons, the DESeq R package[58] was used to compute $\log_2$fold changes in gene expression between treatment groups, and genes with a Benjamini-Hochberg adjusted p-value < 0.05 were considered as DEGs.

Gene ontology enrichment (GO) analysis and clustering
GO analysis of DEGs was performed using the ClueGo plugin v.2.5.7[59] for Cytoscape v3.9.0. Upregulated and downregulated DEGs were analyzed separately. DEGs were compared to the GO term annotations within the reference ontology GO_BiologicalProcess-EBI-UniProt-GOA-ACAP-ARAP_08.05.2020_00h00, and a two-sided hypergeometric test was performed with a Bonferroni adjusted p value <0.05 considered as significant enrichment. Similar GO terms were clustered using iterative kappa score grouping.

Western Blot
NSCs were plated 200,000 cells per well in 6 well plates and treated with 1 µM TFB-TBOA or vehicle for 15 minutes prior to the addition of glutamate or vehicle. Two days later, cells were detached by scraping in ice-cold PBS and pelleted by centrifugation at 800 x g for 2 minutes. Pellets were resuspended, vortexed vigorously, and sonicated in ice-cold lysis buffer consisting of RIPA buffer (Fisher P189900), 1% Halt protease and phosphatase inhibitor (Fisher P178440), and 1% EDTA. Protein lysate concentrations were quantified using the Pierce Rapid Gold BCA protein assay kit (Fisher A53225).

Samples were prepared for SDS-PAGE by diluting 10 µg of protein lysate in Laemmli buffer (Biorad 1610747) with 5% β-mercaptoethanol. Samples were heated at 95°C for 10 min and loaded into a 4-15% gradient Tris-glycine gel (Biorad 4561084) immersed in Tris-glycine-SDS running buffer (Biorad 1610772). Samples were run for 1 hour at 120 V and then transferred overnight at 4°C with 30 V to a nitrocellulose membrane (Fisher NC9680617) in Tris-glycine-20% methanol transfer buffer. The membrane was rinsed with TBS, incubated for 1 hour in 5% BSA/TBS/0.1% Tween20 (blocking buffer), and incubated overnight at 4°C in mouse anti-mTOR

(Cell Signaling 4517) and rabbit anti-phospho(p)-mTOR(Ser2448) (Cell Signaling 2971) both diluted 1:1000 in blocking buffer. Membranes were then rinsed 3x in TBS and incubated for 1 hour in anti-mouse IR680 (VWR 102971-020) and anti-rabbit IR800 (VWR 103011-494) both diluted 1:5000 in blocking buffer. Membranes were then rinsed 3x in TBS and imaged in the 680 and 800 channels using an Odyssey CLx (LI-COR).

Immunoreactive bands for mTOR and p-mTOR were analyzed using ImageStudio (LI-COR). A hand-drawn box of consistent size was traced over each band, and signal intensity was measured with background subtraction using the median of the region immediately surrounding each band. The ratio of p-mTOR to total mTOR signal was calculated for each band.

Statistics

Statistical analysis was performed using GraphPad Prism version 9 or higher, except for RNAseq data, which is described above. Generally, if experiments used a 2x2 factor design, 2-way ANOVA was used with error-corrected posthoc tests. If experiments used a 3x2 factor design, 3-way ANOVA was used with error-corrected posthoc tests. If experiments only had 2 groups, non-parametric comparisons were generally used (Mann Whitney). Details of statistical tests are in Table S4.

Acknowledgements

This research was supported by R21 NS123797 from NINDS/NIH to EDK.

Research reported in this publication was supported by The Ohio State University Comprehensive Cancer Center and the National Institutes of Health under grant number P30 CA016058. We thank the Genomics Shared Resource at The Ohio State University Comprehensive Cancer Center, Columbus, OH for RNAseq services. We also thank the Cancer Program IT resource for sponsoring site licenses for GraphPad Prism and Biorender.

Author contributions

Conceptualization, JDR and EDK; Methodology, JDR and EDK; Investigation, JDR, IRB, JEC, EA, DE, AW, and VV; Writing – Original Draft, JDR and EDK; Writing – Reviewing and Editing, JDR and EDK; Supervision, JPB and EDK; Funding Acquisition, EDK.

Declaration of interests

V. Valentini is currently a consultant for Antec Scientific, USA, a role acquired in Oct 2019 after the completion of the relevant uHPLC-ECD studies but before the writing of the manuscript. The remaining authors declare no competing interests.

References

1. Borrett, M.J., Innes, B.T., Tahmasian, N., Bader, G.D., Kaplan, D.R., and Miller, F.D. (2022). A Shared Transcriptional Identity for Forebrain and Dentate Gyrus Neural Stem Cells from Embryogenesis to Adulthood. eNeuro *9*. 10.1523/ENEURO.0271-21.2021.

2. Götz, M., Nakafuku, M., and Petrik, D. (2016). Neurogenesis in the Developing and Adult Brain—Similarities and Key Differences. Cold Spring Harb Perspect Biol *8*, a018853. 10.1101/cshperspect.a018853.

3. Hochgerner, H., Zeisel, A., Lönnerberg, P., and Linnarsson, S. (2018). Conserved properties of dentate gyrus neurogenesis across postnatal development revealed by single-cell RNA sequencing. Nature Neuroscience *21*, 290–299. 10.1038/s41593-017-0056-2.

4. Hao, Z.-Z., Wei, J.-R., Xiao, D., Liu, R., Xu, N., Tang, L., Huang, M., Shen, Y., Xing, C., Huang, W., et al. (2022). Single-cell transcriptomics of adult macaque hippocampus reveals neural precursor cell populations. Nat Neurosci. 10.1038/s41593-022-01073-x.

5. Moreno-Jiménez, E.P., Flor-García, M., Terreros-Roncal, J., Rábano, A., Cafini, F., Pallas-Bazarra, N., Ávila, J., and Llorens-Martín, M. (2019). Adult hippocampal neurogenesis is abundant in neurologically healthy subjects and drops sharply in patients with Alzheimer's disease. Nature Medicine *25*, 554–560. 10.1038/s41591-019-0375-9.

6. Terreros-Roncal, J., Moreno-Jiménez, E.P., Flor-García, M., Rodríguez-Moreno, C.B., Trinchero, M.F., Cafini, F., Rábano, A., and Llorens-Martín, M. (2021). Impact of neurodegenerative diseases on human adult hippocampal neurogenesis. Science, eabl5163. 10.1126/science.abl5163.

7. Tobin, M.K., Musaraca, K., Disouky, A., Shetti, A., Bheri, A., Honer, W.G., Kim, N., Dawe, R.J., Bennett, D.A., Arfanakis, K., et al. (2019). Human Hippocampal Neurogenesis Persists in Aged Adults and Alzheimer's Disease Patients. Cell Stem Cell *24*, 974-982.e3. 10.1016/j.stem.2019.05.003.

8. Franjic, D., Skarica, M., Ma, S., Arellano, J.I., Tebbenkamp, A.T.N., Choi, J., Xu, C., Li, Q., Morozov, Y.M., Andrijevic, D., et al. (2021). Transcriptomic taxonomy and neurogenic trajectories of adult human, macaque, and pig hippocampal and entorhinal cells. Neuron, S0896627321008667. 10.1016/j.neuron.2021.10.036.

9. Sorrells, S.F., Paredes, M.F., Cebrian-Silla, A., Sandoval, K., Qi, D., Kelley, K.W., James, D., Mayer, S., Chang, J., Auguste, K.I., et al. (2018). Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. Nature *555*, 377–381. 10.1038/nature25975.

10. Berg, D.A., Belnoue, L., Song, H., and Simon, A. (2013). Neurotransmitter-mediated control of neurogenesis in the adult vertebrate brain. Development *140*, 2548–2561. 10.1242/dev.088005.

11. Toda, T., and Gage, F.H. (2018). Review: adult neurogenesis contributes to hippocampal plasticity. Cell Tissue Res *373*, 693–709. 10.1007/s00441-017-2735-4.

12. Asrican, B., Wooten, J., Li, Y.-D., Quintanilla, L., Zhang, F., Wander, C., Bao, H., Yeh, C.-Y., Luo, Y.-J., Olsen, R., et al. (2020). Neuropeptides Modulate Local Astrocytes to Regulate Adult Hippocampal Neural Stem Cells. Neuron, S0896627320306048. 10.1016/j.neuron.2020.07.039.

13. Li, Y.-D., Luo, Y.-J., Chen, Z.-K., Quintanilla, L., Cherasse, Y., Zhang, L., Lazarus, M., Huang, Z.-L., and Song, J. (2022). Hypothalamic modulation of adult hippocampal neurogenesis in mice confers activity-dependent regulation of memory and anxiety-like behavior. Nat Neurosci *25*, 630–645. 10.1038/s41593-022-01065-x.

14. Yeh, C.-Y., Asrican, B., Moss, J., Quintanilla, L.J., He, T., Mao, X., Cassé, F., Gebara, E., Bao, H., Lu, W., et al. (2018). Mossy Cells Control Adult Neural Stem Cell Quiescence and Maintenance through a Dynamic Balance between Direct and Indirect Pathways. Neuron 99, 493-510.e4. 10.1016/j.neuron.2018.07.010.

15. Cameron, H., McEwen, B., and Gould, E. (1995). Regulation of adult neurogenesis by excitatory input and NMDA receptor activation in the dentate gyrus. J. Neurosci. 15, 4687–4692. 10.1523/JNEUROSCI.15-06-04687.1995.

16. Gould, E., McEwen, B.S., Tanapat, P., Galea, L.A.M., and Fuchs, E. (1997). Neurogenesis in the Dentate Gyrus of the Adult Tree Shrew Is Regulated by Psychosocial Stress and NMDA Receptor Activation. J. Neurosci. 17, 2492–2498. 10.1523/JNEUROSCI.17-07-02492.1997.

17. Kitayama, T., Yoneyama, M., and Yoneda, Y. (2003). Possible regulation by N-methyl-d-aspartate receptors of proliferative progenitor cells expressed in adult mouse hippocampal dentate gyrus. Journal of Neurochemistry 84, 767–780. 10.1046/j.1471-4159.2003.01567.x.

18. Nacher, J., Rosell, D.R., Alonso-Llosa, G., and McEwen, B.S. (2001). NMDA receptor antagonist treatment induces a long-lasting increase in the number of proliferating cells, PSA-NCAM-immunoreactive granule neurons and radial glia in the adult rat dentate gyrus. European Journal of Neuroscience 13, 512–520. 10.1046/j.0953-816x.2000.01424.x.

19. DeCarolis, N.A., Mechanic, M., Petrik, D., Carlton, A., Ables, J.L., Malhotra, S., Bachoo, R., Götz, M., Lagace, D.C., and Eisch, A.J. (2013). In vivo contribution of nestin- and GLAST-lineage cells to adult hippocampal neurogenesis. Hippocampus 23, 708–719. 10.1002/hipo.22130.

20. Yuzwa, S.A., Borrett, M.J., Innes, B.T., Voronova, A., Ketela, T., Kaplan, D.R., Bader, G.D., and Miller, F.D. (2017). Developmental Emergence of Adult Neural Stem Cells as Revealed by Single-Cell Transcriptional Profiling. Cell Reports 21, 3970–3986. 10.1016/j.celrep.2017.12.017.

21. Regan, M.R., Huang, Y.H., Kim, Y.S., Dykes-Hoberg, M.I., Jin, L., Watkins, A.M., Bergles, D.E., and Rothstein, J.D. (2007). Variations in Promoter Activity Reveal a Differential Expression and Physiology of Glutamate Transporters by Glia in the Developing and Mature CNS. Journal of Neuroscience 27, 6607–6619. 10.1523/JNEUROSCI.0790-07.2007.

22. Renzel, R., Sadek, A.-R., Chang, C.-H., Gray, W.P., Seifert, G., and Steinhäuser, C. (2013). Polarized distribution of AMPA, but not GABAA, receptors in radial glia-like cells of the adult dentate gyrus. Glia 61, 1146–1154. 10.1002/glia.22505.

23. Wang, L.-P., Kempermann, G., and Kettenmann, H. (2005). A subpopulation of precursor cells in the mouse dentate gyrus receives synaptic GABAergic input. Molecular and Cellular Neuroscience 29, 181–189. 10.1016/j.mcn.2005.02.002.

24. Shin, J., Berg, D.A., Zhu, Y., Shin, J.Y., Song, J., Bonaguidi, M.A., Enikolopov, G., Nauen, D.W., Christian, K.M., Ming, G., et al. (2015). Single-Cell RNA-Seq with Waterfall Reveals Molecular Cascades underlying Adult Neurogenesis. Cell Stem Cell 17, 360–372. 10.1016/j.stem.2015.07.013.

25. Mignone, J.L., Kukekov, V., Chiang, A.-S., Steindler, D., and Enikolopov, G. (2004). Neural stem and progenitor cells in nestin-GFP transgenic mice. Journal of Comparative Neurology *469*, 311–324. 10.1002/cne.10964.

26. Babu, H., Claasen, J.-H., Kannan, S., Rünker, A.E., Palmer, T., and Kempermann, G. (2011). A Protocol for Isolation and Enriched Monolayer Cultivation of Neural Precursor Cells from Mouse Dentate Gyrus. Front. Neurosci. *5*. 10.3389/fnins.2011.00089.

27. Denninger, J.K., Chen, X., Turkoglu, A.M., Sarchet, P., Volk, A.R., Rieskamp, J.D., Yan, P., and Kirby, E.D. (2020). Defining the adult hippocampal neural stem cell secretome: In vivo versus in vitro transcriptomic differences and their correlation to secreted protein levels. Brain Research *1735*, 146717. 10.1016/j.brainres.2020.146717.

28. Thakore, P.I., D'Ippolito, A.M., Song, L., Safi, A., Shivakumar, N.K., Kabadi, A.M., Reddy, T.E., Crawford, G.E., and Gersbach, C.A. (2015). Highly specific epigenome editing by CRISPR-Cas9 repressors for silencing of distal regulatory elements. Nature Methods *12*, 1143–1149. 10.1038/nmeth.3630.

29. Knobloch, M., Pilz, G.-A., Ghesquière, B., Kovacs, W.J., Wegleiter, T., Moore, D.L., Hruzova, M., Zamboni, N., Carmeliet, P., and Jessberger, S. (2017). A Fatty Acid Oxidation-Dependent Metabolic Shift Regulates Adult Neural Stem Cell Activity. Cell Reports *20*, 2144–2155. 10.1016/j.celrep.2017.08.029.

30. Knobloch, M., Braun, S.M.G., Zurkirchen, L., von Schoultz, C., Zamboni, N., Araúzo-Bravo, M.J., Kovacs, W.J., Karalay, Ö., Suter, U., Machado, R.A.C., et al. (2013). Metabolic control of adult neural stem cell activity by Fasn-dependent lipogenesis. Nature *493*, 226–230. 10.1038/nature11689.

31. Ramosaj, M., Madsen, S., Maillard, V., Scandella, V., Sudria-Lopez, D., Yuizumi, N., Telley, L., and Knobloch, M. (2021). Lipid droplet availability affects neural stem/progenitor cell metabolism and proliferation. Nat Commun *12*, 7362. 10.1038/s41467-021-27365-7.

32. López-Colomé, A.M., Martínez-Lozada, Z., Guillem, A.M., López, E., and Ortega, A. (2012). Glutamate Transporter-Dependent mTOR Phosphorylation in Müller Glia Cells. ASN Neuro *4*, AN20120022. 10.1042/AN20120022.

33. Martínez-Lozada, Z., Hernández-Kelly, L.C., Aguilera, J., López-Bayghen, E., and Ortega, A. (2011). Signaling through EAAT-1/GLAST in cultured Bergmann glia cells. Neurochemistry International *59*, 871–879. 10.1016/j.neuint.2011.07.015.

34. Cavallucci, V., Fidaleo, M., and Pani, G. (2016). Neural Stem Cells and Nutrients: Poised Between Quiescence and Exhaustion. Trends in Endocrinology & Metabolism *27*, 756–769. 10.1016/j.tem.2016.06.007.

35. Zhu, J., and Thompson, C.B. (2019). Metabolic regulation of cell growth and proliferation. Nat Rev Mol Cell Biol *20*, 436–450. 10.1038/s41580-019-0123-5.

36. Lobos, P., Córdova, A., Vega-Vásquez, I., Ramírez, O.A., Adasme, T., Toledo, J., Cerda, M., Härtel, S., Paula-Lima, A., and Hidalgo, C. (2021). RyR-mediated Ca2+ release elicited by neuronal activity induces nuclear Ca2+ signals, CREB phosphorylation, and Npas4/RyR2

expression. Proceedings of the National Academy of Sciences *118*, e2102265118. 10.1073/pnas.2102265118.

37. Ouardouz, M., Nikolaeva, M.A., Coderre, E., Zamponi, G.W., McRory, J.E., Trapp, B.D., Yin, X., Wang, W., Woulfe, J., and Stys, P.K. (2003). Depolarization-Induced Ca2+ Release in Ischemic Spinal Cord White Matter Involves L-type Ca2+ Channel Activation of Ryanodine Receptors. Neuron *40*, 53–63. 10.1016/j.neuron.2003.08.016.

38. Anthony, T.E., Klein, C., Fishell, G., and Heintz, N. (2004). Radial Glia Serve as Neuronal Progenitors in All Regions of the Central Nervous System. Neuron *41*, 881–890. 10.1016/S0896-6273(04)00140-0.

39. Gilley, J.A., and Kernie, S.G. (2011). Gltl and Glast negatively regulate calcium-dependent proliferation of hippocampal neural progenitor cells and are persistently upregulated after injury. Eur J Neurosci *34*, 1712–1723. 10.1111/j.1460-9568.2011.07888.x.

40. Berg, D.A., Su, Y., Jimenez-Cyrus, D., Patel, A., Huang, N., Morizet, D., Lee, S., Shah, R., Ringeling, F.R., Jain, R., et al. (2019). A Common Embryonic Origin of Stem Cells Drives Developmental and Adult Neurogenesis. Cell *177*, 654-668.e15. 10.1016/j.cell.2019.02.010.

41. Harris, L., Rigo, P., Stiehl, T., Gaber, Z.B., Austin, S.H.L., Masdeu, M. del M., Edwards, A., Urbán, N., Marciniak-Czochra, A., and Guillemot, F. (2021). Coordinated changes in cellular behavior ensure the lifelong maintenance of the hippocampal stem cell population. Cell Stem Cell *0*. 10.1016/j.stem.2021.01.003.

42. Madsen, S., Ramosaj, M., and Knobloch, M. (2021). Lipid metabolism in focus: how the build-up and breakdown of lipids affects stem cells. Development *148*, dev191924. 10.1242/dev.191924.

43. Wani, G.A., Sprenger, H.-G., Ndoci, K., Chandragiri, S., Acton, R.J., Schatton, D., Kochan, S.M.V., Sakthivelu, V., Jevtic, M., Seeger, J.M., et al. (2022). Metabolic control of adult neural stem cell self-renewal by the mitochondrial protease YME1L. Cell Reports *38*. 10.1016/j.celrep.2022.110370.

44. Kandel, P., Semerci, F., Mishra, R., Choi, W., Bajic, A., Baluya, D., Ma, L., Chen, K., Cao, A.C., Phongmekhin, T., et al. (2022). Oleic acid is an endogenous ligand of TLX/NR2E1 that triggers hippocampal neurogenesis. Proceedings of the National Academy of Sciences *119*, e2023784119. 10.1073/pnas.2023784119.

45. Moss, J., Gebara, E., Bushong, E.A., Sánchez-Pascual, I., O'Laoi, R., M'Ghari, I.E., Kocher-Braissant, J., Ellisman, M.H., and Toni, N. (2016). Fine processes of Nestin-GFP–positive radial glia-like stem cells in the adult dentate gyrus ensheathe local synapses and vasculature. PNAS *113*, E2536–E2545. 10.1073/pnas.1514652113.

46. Hascup, K.N., Findley, C.A., Sime, L.N., and Hascup, E.R. (2020). Hippocampal alterations in glutamatergic signaling during amyloid progression in AβPP/PS1 mice. Sci Rep *10*, 14503. 10.1038/s41598-020-71587-6.

47. Segovia, G., Yagüe, A.G., García-Verdugo, J.M., and Mora, F. (2006). Environmental enrichment promotes neurogenesis and changes the extracellular concentrations of

glutamate and GABA in the hippocampus of aged rats. Brain Research Bulletin *70*, 8–14. 10.1016/j.brainresbull.2005.11.005.

48. Świątkiewicz, M., Fiedorowicz, M., Orzeł, J., Wełniak-Kamińska, M., Bogorodzki, P., Langfort, J., and Grieb, P. (2017). Increases in Brain 1H-MR Glutamine and Glutamate Signals Following Acute Exhaustive Endurance Exercise in the Rat. Frontiers in Physiology *8*.

49. Vivar, C., and van Praag, H. (2017). Running Changes the Brain: the Long and the Short of It. Physiology *32*, 410–424. 10.1152/physiol.00017.2017.

50. Wander, C.M., and Song, J. (2021). The neurogenic niche in Alzheimer's disease. Neuroscience Letters *762*, 136109. 10.1016/j.neulet.2021.136109.

51. Rodríguez-Bodero, A., and Encinas-Pérez, J.M. (2022). Does the plasticity of neural stem cells and neurogenesis make them biosensors of disease and damage? Frontiers in Neuroscience *16*.

52. Fuentealba, L.C., Obernier, K., and Alvarez-Buylla, A. (2012). Adult Neural Stem Cells Bridge Their Niche. Cell Stem Cell *10*, 698–708. 10.1016/j.stem.2012.05.012.

53. Wang, W., Wang, M., Yang, M., Zeng, B., Qiu, W., Ma, Q., Jing, X., Zhang, Q., Wang, B., Yin, C., et al. (2022). Transcriptome dynamics of hippocampal neurogenesis in macaques across the lifespan and aged humans. Cell Res *32*, 729–743. 10.1038/s41422-022-00678-y.

Supplementary Figures

Sup Fig 1: Supplement to Fig 1

A) scRNAseq FPKM values for EAAT transcripts from data in Shin et al., 2015. Only cells with psuedotime <0.5 included to reflect NSCs (as opposed to early IPCs). ***$p<0.001$, #$p<0.0001$ vs EAAT1 and 2, Dunn's multiple comparisons

B) FPKM of EAAT1-5 by pseudotime from Shin et al., 2015. Pearson R^2 values.

C) EAAT1DsRed+EAAT2GFP+ fluorescence in the adult DG. Scale = 100 μm.

D) Orthogonal projections of example: EAAT1DsRed+EAAT2GFP+ (double+) Nestin+ radial glia like (RGL) NSC, EAAT1DsRed+EAAT2GFP+ (double+) Ki67+ IPC, EAAT1DsRed-EAAT2GFP- (double-) DCX+ immature neuron. Scale = 10 μm.

E) Nestin immunolabeling in DG of EAAT1DsRed+ co-localized with EAAT1 immunoreactivity (ir). Left is merged z-stack. Right is 3D reconstruction showing an example Nestin+EAAT1DsRed+ radial glia like NSC with EAAT1ir in the cell body and apical terminals.

Sup Fig 2: Supplement to Fig 2
A) FPKM of glutamate receptor genes from Shin et al., 2015 with EAAT1 and EAAT2 included for scale.

B) Transcript length-normalized RNAseq counts of glutamate receptor genes and EAAT1/2 from cultured adult DG NSCs in Denninger et al., 2020.

C) emPAI length-normalized semi-quantitiative spectral counts for glutamate receptor and transporter proteins from cultured adult DG NSCs in Denninger et al., 2020.

D) Representative image of EdU+ labeling in cultured NSCs after L-aspartate treatment +/- NMDA receptor inhibitor (AP5), with Hoechst to label nuclei. Scale = 20 μm.

E) Representative image of EdU+ labeling in cultured NSCs after glutamate receptor agonist tretament, with Hoechst to label nuclei. Scale = 20 μm.

F) Glutamate in conditioned media as measured by uHPLC after a 5 μM pulse in wells with NSCs or empty wells. ***,****$p<0.001$, 0.0001 vs "No Cells" Sidak's multiple comparisons. Mean ± SEM of N = 6 independent experiments.

G) Glutamate in NSC conditioned media as measured by uHPLC after a 5 μM pulse after pre-treatment with an EAAT2-selective (1 μM) and an EAAT1and2 selective (10 μM) dose of WAY213613. *$p<0.05$ vs 1 μM and vs 10 μM. Mean ± SEM of N = 6 independent experiments.

H) Glutamate in NSC conditioned media as measured by uHPLC after a 5 μM pulse after pre-treatment with a panEAAT inhibitor (TFB-TBOA), an EAAT1 inhibitor (UCPH101), or and EAAT2 inhibitor (DHK). *$p<0.05$ Tukey's comparisons within timepoint. Mean ± SEM of N = 3 independent experiments.

I) Representative image of EdU+ labeling in cultured NSCs after treatment with EAAT inhibitors, with Hoechst to label nuclei. Scale = 20 μm.

J) Sapphire IR signal in cultured NSCs after treatment with glutamate +/- panEAAT inhibitor (TFB-TBOA) or EAAT1 inhibitor (UCPH101). Protein transport inhibitor at toxic dose is positive control for presence of apoptosis in Sapphire IR700 assay. ****$p<0.0001$ Dunnett's multiple comparisons each group vs positive control. Mean ± SEM of N = 3-6 replicates/experiment, 2 independent experiments. Top shows representative image of Sapphire IR signal in treatment groups.

Sup Fig 3: Supplement to Fig 3
A) EAAT1 immunoreactive intensity in cultured NSCs infected with NT and EAAT1 KD CRISPRi lentiviral vectors.
    ****p<0.0001 Sidak's multiple comparisons within GFP type. Mean ± SEM of N = 4 independent infections.

B) Representative images of EAAT1 labeling coupled with GFP and brightfield to visualize all NSCs. Scale = 10 μm.

C) EdU+GFP+ cell counts in cultured NSCs infected with NT and EAAT1 KD CRISPRi lentiviral vectors. Mean ± SEM of N = 3 replicates/infection, 3 independent infections.

D) Representative images of EdU labeling coupled with GFP in NSCs infected with NT and EAAT1 KD CRISPRi lentiviral vectors. Scale = 10 μm.

E) Timeline and EAAT1 immunolabeling intensity within GFP+GFAP+ area of mice infused with NT or EAAT1 KD CRISPRi lentiviral vectors in the DG then perfused 1 week later. Mean ± SEM of N = 5-8 individual mice (points shown).

F) Representative image of EAAT1 labeling coupled with GFP and GFAP at 1 week after CRISPRi infusion. Co-labeling becomes white in merged image. Scale = 20 μm.

G) Timeline and EAAT1 immunolabeling intensity within GFP+GFAP+ area of mice infused with NT or EAAT1 KD CRISPRi lentiviral vectors in the DG then perfused 3 weeks later. *$p < 0.05$ Mann Whitney test. Mean ± SEM of N = 8 individual mice (points shown).

H) Representative image of EAAT1 labeling coupled with GFP and GFAP at 3 weeks after CRISPRi infusion. Co-labeling becomes white in merged image. Scale = 20 μm.

I) Representative image of BrdU, GFP, DCX co-labeling 3 weeks after CRISPRi vector infusion. Scale = 10 μm.

J) Representative orthogonal view of a BrdU+NeuN+GFP+ cell in a NT infused mouse and a BrdU+NeuN+GFP- cell in an EAAT1 KD infused mouse. Scale = 10 μm.

Sup Fig 4: Supplement to Figure 4

A) Principle component analysis of RNAseq samples from treated in vitro NSCs.
B) Representative images of EdU labeling in NSCs treated with glutamate +/- FASN inhibitor orlistat. Hoechst as nuclear marker. Scale = 10 μm.
C) Representative blot of phosphor-mTOR and total mTOR in NSCs glutamate +/- panEAAT inhibitor (TFB-TBOA).
D) Representative images of EdU labeling in NSCs treated with glutamate +/- NCX inhibitor SEA0400. Hoechst as nuclear marker. Scale = 10 μm.
E) EdU+ cell counts after treatment with glutamate +/- NCX inhibitor SEA0400. Mean ± SEM of N = 4 replicates/experiment, 4 independent experiments.
F) EdU+ cell counts after treatment with glutamate +/- NCX inhibitor YM244769. *,**,***p<0.05, 0.01, 0.001 Sidak's multiple comparisons. Mean ± SEM of N = 2 replicates/experiment, 4 independent experiments.
G) Representative images of EdU labeling in NSCs treated with glutamate +/- NCX inhibitor YM244769. Hoechst as nuclear marker. Scale = 10 μm.
H) Model of proposed glutamate transport intracellular signaling.