A secreted protease-like protein in *Zymoseptoria tritici* is responsible for avirulence on *Stb9* resistance gene in wheat

Reda AMEZROU[1], Colette AUDÉON[1], Jérôme COMPAIN[2], Sandrine GÉLISSE[1], Aurélie DUCASSE[1], Cyrille SAINTENAC[3], Nicolas LAPALU[1,2], Simon ORFORD[4], Daniel CROLL[5], Joëlle AMSELEM[2], Sabine FILLINGER[1], Thierry C MARCEL[1*]

[1] Université Paris-Saclay, INRAE, UR BIOGER, 91120 Palaiseau, France.

[2] Université Paris-Saclay, INRAE, UR URGI, 78026 Versailles, France.

[3] UCA, INRAE, GDEC, 63000 Clermont-Ferrand, France.

[4] Crop Genetics, John Innes Centre, Norwich NR4 7UH, UK.

[5] University of Neuchâtel, Laboratory of Evolutionary Genetics, 2000 Neuchâtel, Switzerland.

[*] Corresponding author: thierry.marcel@inrae.fr

Running title: The avirulence factor *AvrStb9* from *Zymoseptoria tritici*

Key words: Avirulence, Gene-for-gene interaction, Protease, GWAS, Wheat, *Zymoseptoria tritici*

Word count (main text): 6,396

Summary

- *Zymoseptoria tritici* is the fungal pathogen responsible for Septoria tritici blotch on wheat. Disease outcome in this pathosystem is partly determined by isolate-specific resistance, where wheat resistance genes recognize specific fungal factors triggering an immune response. Despite the large number of known wheat resistance genes, fungal molecular determinants involved in such cultivar-specific resistance remain largely unknown.

- We identified the avirulence factor *AvrStb9* using association mapping and functional validation approaches. Pathotyping *AvrStb9* transgenic strains on *Stb9* cultivars, near isogenic lines and wheat mapping populations, showed that *AvrStb9* interacts with *Stb9* resistance gene, triggering a strong immune response.

- *AvrStb9* encodes an unusually large avirulence gene with a predicted secretion signal and a protease domain. It belongs to a S41 protease family conserved across different filamentous fungi in the Ascomycota class and may constitute a core effector. *AvrStb9* is also conserved among a global *Z. tritici* population and carries multiple amino acid substitutions caused by strong positive diversifying selection.

35 • These results demonstrate the contribution of an 'atypical' conserved effector protein to fungal
36 avirulence and the role of sequence diversification in the escape of host recognition, adding to our
37 understanding of host-pathogen interactions and the evolutionary processes underlying pathogen
38 adaptation.
39

Introduction

Filamentous fungal pathogens are responsible for the most damaging crop diseases (Savary *et al.*, 2019).
Their ability to cause disease is facilitated by fungal proteins called effectors, which are delivered into
the plant apoplast or cytoplasm during the infection process (Lo Presti *et al.*, 2015; Rocafort *et al.*, 2020).
Effectors play critical role in the infection by manipulating plant immune system and other cellular
functions. To counteract fungal infections, plants have evolved efficient defense systems relying on the
recognition of different fungal factors (Jones & Dangl, 2006). The first type of defense involves plant
receptors recognizing generic fungal molecules, called pathogen-associated molecular patterns
(PAMPs). This PAMP-triggered immunity (PTI) provides basal defense against non-adapted pathogens.
The second type of defense is activated with the recognition of fungal effectors by plant receptors,
defined as resistance (*R*) genes. Effector-triggered immunity (ETI) induces a specific and strong immune
response thought to limit the infection (Jones & Dangl, 2006). This interaction typically follows the
gene-for-gene model, where host *R* gene products recognize pathogen effectors, which are then called
avirulence (*Avr*) genes (Flor, 1956). However, ETI exerts a strong evolutionary pressure on pathogen
populations leading to the selection of virulent isolates overcoming host immune response (Lo Presti et
al., 2015). Most of these virulent isolates displayed mutations in the avirulence gene leading to variant
proteins not recognized by the corresponding R protein (Sánchez-Vallet et *al.*, 2018). These observations
emphasize the importance of pathogen *Avr* genes surveillance and the adequate spatiotemporal
deployment of *R* genes in the field.
A common assumption is that most *Avr* genes encode small secreted proteins (SSP), with no predicted
functional domain and no sequence homology in other species (Stergiopoulos & de Wit, 2009;
Franseschetti et *al.*, 2017). It is often overlooked that *Avr* genes may also code for proteins not strictly
falling into this category. For instance, some *Avr* genes from *Cladosporium fulvum* (*Avr2* and *Avr4*),
Magnaporthe oryzae (*Avr-Pita*) or *Melampsora lini* (*AvrM14*) encode proteins that share sequence
similarities with known enzymes or enzyme inhibitors (reviewed in Franseschetti et *al.*, 2017). However,
only few such 'atypical' avirulence factors have been described in literature so far and their
identification will broaden our knowledge on the complex molecular cross-talk between plants and
pathogens.
Zymoseptoria tritici is an ascomycete fungal pathogen that causes the major wheat foliar disease
Septoria tritici blotch (STB). *Z. tritici* is apoplastic with a biphasic infection process (Petit-Houdenot et
al., 2021). After penetration into wheat leaves through stomata, the fungus colonizes the apoplast for 8-
11 days while the host remains symptomless (Battache et *al.*, 2022). This long asymptomatic phase is

followed by a rapid onset of necrotic symptoms, associated with the development of fungal asexual

fruiting bodies called pycnidia, which contain the pycnidiospores that further spread the disease

(Sánchez-Vallet *et al.*, 2015). STB is currently the economically most important wheat disease in Europe

and is mainly controlled using fungicides and genetic resistances (Fones & Gurr, 2015; Petit-Houdenot

et *al.*, 2021). However, this disease remains a problem due to the ability of fungal populations to

overcome control measures (*i.e.*, fungicide resistance, host resistance breakdown) and adapt to climate

changes (Fones & Gurr, 2015). Despite its economic importance, the molecular basis of *Z. tritici*

avirulence remains largely unknown. Only two avirulence genes have been characterized so far,

AvrStb6, recognized by the resistance protein *Stb6* leading to a strong immune response; and *Avr3D1*

which is thought to trigger quantitative resistance in cultivars carrying *Stb7* resistance gene (Zhong *et*

al., 2017; Kema *et al.*, 2018; Meile *et al.*, 2018). Both genes encode SSPs located in highly plastic

genomic regions rich in transposable elements thought to favor their rapid diversification (Zhong *et al.*,

2017; Meile *et al.*, 2018). In addition to *Stb6* and *Stb7*, 21 major isolate-specific STB resistance genes

have been mapped in wheat, but their corresponding avirulence genes remain unknown (Brown *et al.*,

2015; Yang *et al.*, 2018; Langlands-Perry *et al.*, 2022).

Here, we aimed to broaden our understanding of avirulence in fungal pathogens, the mechanisms

underlying gene-for-gene interactions in the *Z. tritici*–wheat pathosystem and the evolutionary processes

related to the escape of host recognition. We first showed that a large protein encoding a predicted

secreted protease, *AvrStb9*, is responsible for avirulence in a subset of fungal strains. We demonstrated

that recognition *AvrStb9* by hosts carrying *Stb9* resistance gene triggers a strong immune response

following the gene-for-gene model. We next showed that *AvrStb9* belongs to a family of proteins with

a predicted protease S41 domain conserved across diverse ascomycetes lineages and that mutations in

AvrStb9, including some in the predicted protease domain, led to variants escaping *Stb9*-mediated

recognition.

Materials and Methods

Fungal material

A total of 103 *Z. tritici* strains were used for GWAS analysis. The strains derive from a French *Z. tritici*

collection collected in 2009 on cultivars 'Apache' (n = 44), 'Premio' (n = 42), 'Soissons' (n = 9),

'Caphorn' (n = 3), 'Alixan' (n = 1), 'Bermude' (n = 1), 'Dinosaur' (n = 1) and 'Garcia' (n = 1); one

isolate was collected in the UK on cultivar 'Humbert'. The two isolates used for genetic mapping of the

resistance and fungal transformation are IPO323 and IPO09593. IPO323 is a Dutch isolate collected on

cultivar 'Arminda' in 1981 and used as the reference strain for *Z. tritici* (Goodwin *et al.*, 2011).

IPO09593 is part of the 103 strains and has been isolated from 'Apache' in 2009. IPO323 is virulent on

cultivars carrying *Stb9* resistance gene while IPO09593 is avirulent on *Stb9*-cultivars. The three

additional isolates used to construct deletion and allele swap transformants, namely IPO09359

('Apache', 2009), IPO09139 ('Apache', 2009) and SYN32 (unknown cultivar, 2006), are avirulent on *Stb9*-cultivars.

Plant material

We performed quantitative trait loci (QTL) mapping in a recombinant doubled haploid (DH) population comprised of 65 lines, developed from the F_1 progeny of the cultivars 'Beaver' and 'Soissons' (Singh *et al.*, 2009). A panel of 220 modern French bread wheat (BW) varieties was used for GWAS analysis; details on the BW panel used here can be found in Touzy *et al.* (2019). We also characterized *Z. tritici* wild-type and mutant strains on near isogenic lines (NILs) carrying or not the *Stb9* resistance gene in the genetic background of cultivar 'Courtot', developed at INRAE GDEC (Clermont-Ferrand, France). The 'Courtot' NILs were developed following three backcrosses starting with F_1 'Courtot' x 'Chinese spring' (carrying a susceptible allele of *Stb9* gene). To keep the *Stb9* susceptible allele in the progenies, each plant was genotyped with single sequence repeat markers wmc317 and barc129 and phenotyped with the *Stb9* avirulent isolate IPO89011. A BC_3F_1 plant heterozygous at the *Stb9* locus was self-fertilized. BC_3F_2 plants homozygous either for the susceptible *Stb9* allele ('Courtot' – Stb9) or for the resistant allele ('Courtot' + Stb9) were selfed and constitute the 'Courtot' NILs.

Pathogenicity assays

Pathogenicity assays were performed following the procedure described by Langlands-Perry *et al.* (2022). Briefly, 16 days old plants were marked to delimit a 75 mm segment on the first true leaf. The segment was inoculated with a paintbrush dipped into a 10^6 spores.mL^{-1} inoculum solution with one drop of Tween®20 added per 15 mL. After inoculation, the plants were watered and covered with transparent polyethylene bags for 72 hours to initiate the infection. The plants were kept in a climate chamber with a 16 h photoperiod (300 µmol.m^{-2}.s^{-1}), 80-90% relative humidity and 18°C night/22°C day. Phenotyping of the fungal population was performed by visually assessing strain virulence on the cultivar 'Soissons' at 21 days post inoculation (dpi). Visual assessments were made of percent of leaf area covered by necrosis (PLACN) and by pycnidia (PLACP) within the inoculated segment of each leaf. Each of the 103 strains was tested in three replicates with three leaves per replicate. Phenotyping of *AvrStb9* mutant strains was performed following the same procedure.

For the host mapping populations, the Beaver/Soissons DH lines were inoculated with the strain IPO09593, and the 220 wheat varieties were inoculated with the strains IPO09593 and IPO323. Here, each interaction was tested in two replicates with three leaves per replicate and visual assessments of PLACN and PLACP were recorded at 14, 20 and 26 dpi. We calculated the area under disease progress curve for necrosis (AUDPCn) and sporulation (AUDPCs) with the formula:

$$\text{AUDPC} = \frac{\sum[(t_{i+1} - t_i) \times (y_i + y_{i+1})]}{2}$$

With, $t_{i+1}-t_i$: number of days between two observations; y_i: PLACN or PLACP at day i; y_{i+1}: PLACN or

PLACP at day i+1. AUDPC were calculated relative to their maximum theoretical value, RAUDPCn

and RAUDPCs respectively. For all assays, disease severity data were averaged between replicates and

used as the input phenotype in GWAS and QTL analyses.

Fungal whole-genome sequencing and variant calling

Fungal spores were obtained after 7 days growth in liquid cultures of yeast extract-peptone-dextrose

Broth (Difco™ YPD Broth), and then freezed-dried during 24 h. High molecular weight DNA was

extracted using a phenol-chloroform based procedure. Illumina paired-end libraries were prepared for

each sample and DNA fragments sequenced with 2x100 bp reads on a HiSeq-2000 sequencing system

at Genewiz, Inc. (formerly Beckman Coulter Genomics). Depth of sequence coverage was variable

depending on the isolate, ranging from 16 to 303 genome equivalents with an average of 61 genome

equivalents. For each strain, reads were trimmed using trimmomatic v0.32 (Bolger *et al.*, 2014) and then

mapped on the reference genome IPO323 (Goodwin *et al.*, 2011) using the mem algorithm from BWA

v0.7.7 (Li & Durbin, 2009) with default options. Samtools v0.1.19 and Picard tools v1.106 (Li *et al.*,

2009) were used to filter reads (optic duplicates, secondary alignments and reads with a mapping quality

< 30) and keep only pairs in which both reads met quality checks. SNP calling was done with Freebayes

v0.9 with the option --report-monomorphic (Garrison & Marth, 2012). All positions detected as low-

complexity regions or transposable elements (TE) were excluded using RepeatMasker 4.0 with default

options and REPET (Flutre *et al.*, 2011), respectively. Quality filters were applied on the gVCF using

in-house scripts. Depth of coverage (DP) calculated at each position follows a normal distribution

around the depth main peak; a DP filter was set at twice the standard deviation from the mean. All the

gVCF were stored into a matrix reduced to keep only the positions where at least one strain has a detected

SNP in a non-filtered position. The SNP matrix was parsed in .tped format, converted to .ped/.map and

filtered for < 0.5 missing data and > 0.1 minor allele frequency (MAF) using PLINK v.1.9 (Chang *et*

al., 2015).

GWAS analyses

For Genome-Wide Association Analysis (GWAS), we used the linear mixed model implemented in the

R package 'GAPIT' (Lipka *et al.*, 2012). To account for population structure and reduce spurious

associations, a kinship matrix using the VanRaden method and the first three principal components

computed from SNP data were included a covariate in the model. We used a Bonferroni threshold of p

< 2.67e-7 (α=0.05) to declare significant SNPs associated with phenotypic variations. Linkage

Disequilibrium (LD) measures (r^2) for the significant SNPs were obtained with PLINK v.1.9 using the

--r2 command and LD blocks were visualized using the R package 'LDheatmap'. SNPs with an $r^2 \geq 0.5$

were considered in strong LD.

Genetic mapping of *Stb9*

A genetic linkage map was available for the Beaver/Soissons DH population and contains 312 polymorphic loci: 121 SSR, 49 AFLP, 135 DArT markers and 7 major genes (Singh *et al.*, 2009). The map covers 1,526 cM with an average density of one marker per 12.6 cM. QTL mapping was conducted using RAUDPCs and RAUDPCn phenotypes obtained from the isolate IPO09593 (*AvrStb9*) with R/qtl software v.1.48-1 (Broman *et al.*, 2003). Linkage analysis was performed for each trait with an initial Simple Interval Mapping (SIM), followed by a Composite Interval Mapping (CIM). For SIM, 1000 genome-wide permutations were used to calculate the significant logarithm of odds (LOD) threshold. Only QTLs that showed *p*-values less than 0.05 were considered significant. Association mapping in the BW panel was done following the exact same procedure as in the fungal population (see GWAS analyses). The 220 BW varieties have been previously genotyped on an Affymetrix Axiom 410K single nucleotide polymorphism (SNP) array (Danguy des Déserts *et al.*, 2021). Procedure of allele calling, filtering and quality check of genotype data are described in Balfourier *et al.*, 2022. For GWAS analysis, we excluded markers with less than 5% minor allele frequency and more than 10% missing data and kept a set of 186,961 SNPs that passed quality control criteria and could be anchored on the IWGSC annotation v2.1 (Zhu *et al.*, 2021). We used filtered SNP data and phenotypes (RAUDPCs and RAUDPCn) from the isolates IPO323 (*avrStb9*) and IPO09593 (*AvrStb9*) to perform marker-trait associations.

Gene annotation and expression analysis

Predicted genes in the candidate region were retrieved from the *Z. tritici* reference genome annotation (Grandaubert *et al.*, 2015). We used RNA-seq data (Rudd *et al.*, 2015) mapping onto the reference genome IPO323 to manually curate *Zt_1_693* (*AvrStb9)* gene model. Canonical and non-canonical splicing sites were confirmed by *de-novo* transcriptome assembly obtained with DRAP (Cabau *et al.*, 2017). The final gene model differed from those from previous annotations due to errors in predicting correctly introns and includes two missing exons. Candidate genes were screened for the presence of a secretion signal using SignalP 3.0 server (Dyrløv Bendtsen *et al.*, 2004). Blastp was used to search for *AvrStb9* orthologous and paralogous proteins against the reference strain IPO323 and closely-related *Zymoseptoria* species. Protein domains were searched using Interproscan and structure homology prediction was performed in the HHpred server; <https://toolkit.tuebingen.mpg.de/tools/hhpred>. For gene expression analysis, RNA-seq data samples from *in vitro* and *in planta* (1, 4, 9, 14 and 21 dpi) conditions were obtained from Rudd *et al.* (2015), and gene expression quantified as fragments per kilobase of exon per million fragments mapped (FPKM) using Cuffdiff v.2.2.1 (Trapnell *et al.*, 2012). *AvrStb9* DNA sequences are deposited in NCBI Genbank (accession numbers OP376566 and OP376567).

Construction of disruption plasmids

For *AvrStb9* disruption, the hygromycin resistance marker (*hph*) was cloned between the 1,000 bp up-

and downstream flanking regions of the ORF, respectively. The following DNA fragments were

generated and cloned into pCAMBIA0380 plasmids (GenBank: AF234290.1). The up- and downstream

flanks were obtained by PCR on gDNA of IPO323 with primer pairs 1&2 and 5&6 (Table S2)

respectively using Q5 Hot Start High fidelity DNA polymerase (NEB, Evry, France). The *hph* was

amplified from pCAMBIA0380 with the primers 3 and 4 (Table S2). pCAMBIA0380 was digested by

*Eco*RI and *Xho*I and used for the cloning of three PCR fragments with Gibson Assembly cloning kit

(NEB, Evry, France) according to the supplier's indications. The ligation product was introduced into

NEB 5α competent *Escherichia coli* cells (NEB, Evry, France), transformants selected on kanamycin

(50 µg.mL^{-1}) containing LB-broth. Colonies were screened by PCR with the primer-pairs used for

cloning. Four positive colonies were chosen for plasmid extraction (Macherey & Nagel, Düren,

Germany) and restriction analysis. The construct was then introduced by electroporation into the

Agrobacterium tumefaciens Agl1 strain. Transformants were selected and isolated on YEB (5 g. L^{-1} beef

extract, 1 g. L^{-1} yeast extract, 5 g. L^{-1} peptone, 5 g. L^{-1} sucrose, 0.5 g. L^{-1} MgCl2, 1.5% agar) with

kanamycin (50 µg. mL^{-1}).

Gene replacement constructs

To swap *AvrStb9* alleles between virulent (IPO323) and avirulent (IPO09593) strains, the following

replacement cassettes were constructed. The respective *AvrStb9* alleles were amplified with primers 11

and 12 (Table S2) from IPO323 or IPO09593 gDNA. These fragments cover the ORF with 1 kb

upstream of the START codon and 360 bp downstream of the STOP codon. The *hph* (amplified with

primers 9 and 10 from pCAMBIA0380 plasmid DNA) and the 1 kb upstream flank of *AvrStb9* gene

(amplified with primers 7 and 8) were ligated to *AvrStb9* alleles into the *EcoRI-XhoI* digested

pCAMBIA0380, with Gibson Assembly cloning kit (NEB, Evry, France). Plasmids were introduced

into NEB 5α competent *E. coli* cells (NEB, Evry, France) and transformants selected on kanamycine

(50 µg.mL^{-1}) containing LB-broth. Colonies were screened by PCR with the primer-pairs used for

cloning. Four positive colonies were chosen for plasmid extraction (Macherey & Nagel, Düren,

Germany) and sequence analysis. These constructs named respectively pR323 and pR9593 were then

introduced by electroporation into the *Agrobacterium tumefaciens* Agl1 strain. Transformants were

selected and isolated on YEB with kanamycin (50 µg.mL^{-1}).

Generation of *Z. tritici* transformants

The transformation procedure was adapted from Zwiers & de Waard (2001) and was performed as

previously described (Omrane *et al.*, 2015). For *AvrStb9* deletion, the following *Z. tritici* strains were

transformed with the disruption plasmid: IPO323, IPO09593, IPO09359, IPO09139 and SYN32.

Transformants were selected and isolated twice on hygromycin (100 µg. mL^{-1}) containing medium. The

replacement of the *AvrStb9* ORF by the *hph* gene was checked by PCR on gDNA with the primer pair

13 and 14. The gene deletion is expected to give a 1.6 kb fragment instead of the native 2.4 kb amplicon.

The total number of isolated and analyzed transformants were: IPO323 (n=144), IPO09593 (n=23),

IPO09359 (n=26), IPO09139 (n=46), SYN32 (n=38). One single transformant produced the expected

PCR profile, namely SYN32_ΔT12.

For *AvrStb9* allele swaps, IPO323 was transformed with plasmids pR323 and pR9593. Transformants

selected and isolated twice on hygromycin containing medium (100 µg. mL^{-1}). The replacement of

AvrStb9 alleles in the transformants was checked by PCR using primers 15 and 16, leading to a single

1.6 kb amplicon in replacement mutants and to an additional 238 bp amplicon in ectopic transformants.

Transformants with the expected single band size were PCR checked with primers 17 and 18 as these

primers allow to distinguish the deletion of the amino acid residues 66-71 in the IPO09593 background

from the IPO323 one, and subsequently submitted to sequence analysis with primers indicated in Table

S2.

Phylogenetic analysis. We used OrthoMCL algorithm to search for *AvrStb9* orthologous and

paralogous protein sequences using a cutoff E-value less than 1e-20 and filtering for low complexity

regions (Li *et al.*, 2003); <https://orthomcl.org/orthomcl/app/>. Of the 148 putative *AvrStb9*

orthologs/paralogs identified through OrthoMCL, we selected 122 by removing redundant information

using the following criteria: (*i*) when multiple identical sequences were present, one was arbitrarily

chosen and retained; and (*ii*) when multiple sequences from different strains of the same species were

present, an arbitrarily chosen strain (the reference strain, if known) was retained. A multiple sequence

alignment was then generated with Clustal Omega v1.2.4 (Sievers *et al.*, 2011) and alignment cleaning

was performed using trimAL with the --gappyout option (Capella-Gutiérrez *et al.*, 2009). We

constructed a maximum likelihood tree of selected *AvrStb9* orthologs/paralogs using RaxML with the

GAMMA JTT model and 100 bootstrap replicates (Stamatakis, 2014). Phylogenetic trees were

visualized using iTOL (Letunic & Bork, 2016); <https://itol.embl.de/>.

Population genetics and selection analyses

In addition to the fungal population used for genetic analyses, we extended our sequence diversity and

selection analyses to include a global *Z. tritici* population originating from four distinct geographic

locations, including the middle East (n=30, Israel), Australia (n=27, Wagga Wagga), Europe (n=33,

Switzerland) and North America (n=56, Oregon, USA). Procedures of *de novo* genome assemblies and

sequence extraction were previously described (Hartmann *et al.*, 2017). We used DnaSP v6 (Rozas *et*

al., 2017) to calculate summary statistics of the population genetic parameters associated with *AvrStb9*.

Sliding window analyses of nucleotide diversity (π) and Tajima's D were conducted in a window length

of 200 bp and a step size of 20 bp. To test the selective pressure acting on *AvrStb9*, we estimated the

rates of non-synonymous (dN) and synonymous (dS) substitutions using CodeML from the PAML

package (Nielsen & Yang, 1998). A dN/dS (ω) ratio of 1 indicated neutrality, whereas ω < 1 or > 1

suggested purifying or diversifying selection, respectively. However, the presence of intragenic
recombination could bias the ω rate testing. To account for these issues, we performed the analysis with
a non-recombining phylogenetic tree generated from the RDP4 software using RaxML with 100
bootstraps (Martin *et al.*, 2015).

Results

Mapping of a candidate gene conferring avirulence on the cultivar 'Soissons'.

'Soissons' is a French bread wheat variety, developed by Florimond Desprez and released in 1987,
which represented 10 to 40% of the French wheat acreage between 1991 and 2002. 'Soissons' is
moderately susceptible to STB (Suffert *et al.*, 2013) but it has been postulated to carry the resistance
gene *Stb9* based on screening with differentially virulent isolates (data not shown). Infection assays of
the fungal population on cultivar 'Soissons' showed that ~12% of the French isolates are avirulent (Fig.
S1). We conducted GWAS to identify the *Avr* gene interacting with the resistance in 'Soissons'.

A total of 103 *Z. tritici* strains were re-sequenced, generating 718,810 SNPs passing quality filters. Using
full SNP data, we investigated the population structure and kinship and found very little genetic
substructure and relatedness in the fungal population (Fig. S2a,b). For instance, the first three PCs
accounted for only 3.11% of the total genetic variation. Using a linear mixed model to associate genetic
variations with pycnidia (PLACP) and necrosis (PLACN) variations on 'Soissons', we identified a
cluster of 26 SNPs on chromosome 1 from 2,092,563 bp to 2,096,143 bp, significantly associated with
PLACN at the Bonferroni threshold (Fig. 1a). The same SNP cluster was also associated with variation
in PLACP but failed to reach the genome-wide significance (Fig. S3). We also detected a significant
association with PLACN variations (p=3.801e-8; position 2,451,010 bp) on a segment of chromosome
7. However, the region falls in a broad section of the chromosome without any *in planta* expressed genes
(Rudd *et al.*, 2015). The significant SNPs on chromosome 1 mapped to a region containing two predicted
genes, *Zt_1_692* and *Zt_1_693*. The lead SNP (p=7.31e-10; position 2,095,552 bp) mapped to the fourth
exon of *Zt_1_693*, but was in strong linkage disequilibrium with other variants spanned across the two
predicted genes (Fig. 1b). *Zt_1_692* is not expressed *in planta* so we were unable to use RNA-seq data
to evaluate the quality of the gene annotation and blastp searches yielded only weak hits. Without any
corroborating evidence for a biological role, we concluded that *Zt_1_692* is likely a pseudogene or an
annotation error. In contrast, *Zt_1_693* is highly expressed *in planta* (Fig. S4), with a peak expression
at 9 days post infection coinciding with the switch from asymptomatic to necrotrophic phase (Fig. 1d).
Further, allele analyses of the lead SNP clearly discriminated between avirulent and virulent phenotypes
(Fig. 1c). Taken together, we considered *Zt_1_693* as the best candidate gene to explain avirulence on
the wheat cultivar 'Soissons'.

Zt_1_693 is involved in a gene-for-gene relationship with *Stb9* resistance gene.

To determine whether *Zt_1_693* is an avirulence factor recognized by an *R* gene in the cultivar 'Soissons', we sought to generate *Zt_1_693* deletion mutants (KO) in the virulent strain IPO323 and four different avirulent strains by targeted gene replacement. We also switched alleles between avirulent (IPO09395) and virulent (IPO323) strains using targeted gene replacement. However, we could not obtain any deletion mutant neither in IPO323 (144 transformants tested), nor in the avirulent strains IPO09593, IPO09359 and IPO09139 (n=95). Only a single deletion mutant was obtained in strain SYN32 out of 32 transformants, producing small colonies and weak symptoms on the susceptible cultivar 'Taichung-29' (Table S3). This suggests that *Zt_1_693* could be essential for *Z. tritici* growth, and pathogenicity. Nevertheless, using allele swapping we were able to generate mutants expressing the virulent and avirulent versions of *Zt_1_693* in IPO323 genetic background, and performed infection assays on cultivars 'Soissons', 'Courtot' which carries the *Stb9* gene and a 'Courtot' near isogenic line lacking *Stb9*.

IPO323 gene replacement transformants carrying the avirulent allele (n=6) from IPO09593 (referred to as *Avrstb9*) were avirulent on 'Soissons' and 'Courtot', but not on 'Courtot' lacking *Stb9* (Fig. 2a). Inversely, gene replacement transformants with the virulent allele (referred to as *avrstb9*) from IPO323 (n=2) resulted in virulence on all tested cultivars, with PLACP values at 26 dpi ranging from 32 to 100% (Fig. 2a). Interestingly, Sanger sequencing of IPO323$_{Avrstb9}$ transformants revealed that four strains were chimeric with the avirulent allele not being introduced in full length, retaining some polymorphisms of the virulent haplotype between residues 276 and 628. All four chimeric strains were virulent on *Stb9*-containing cultivars, highlighting the importance of these polymorphisms in virulence (Table 1; Table S3; Fig. S6). The fact that all IPO323 mutants carrying the complete *AvrStb9* allele were unable to infect *Stb9*-containing cultivars, while mutants expressing the *avrStb9* allele produced symptoms on all three cultivars, suggest that *Zt_1_693* encodes the avirulence factor *AvrStb9*, which triggers complete resistance in cultivars carrying the cognate resistance gene.

To further validate the *AvrStb9-Stb9* gene-for-gene relationship, we performed two independent genetic analyses using a bi-parental mapping population and a panel of wheat varieties. The isolate-specific resistance to IPO09593 (*AvrStb9*) was mapped by a QTL analysis on Beaver/Soissons recombinant double haploid progenies. A single major resistance QTL was detected on the long arm of the wheat chromosome 2B, likely corresponding to the *Stb9* locus (Chartrain *et al.*, 2009). This QTL had a log-likelihood (LOD) of 10.77 and 15.33 for the RAUDPC of necrosis and sporulation, respectively (Fig. 2b). As expected, the resistance allele originated from 'Soissons' suggesting that the cultivar indeed carries the *Stb9* resistance. We confirmed this interaction by performing a GWAS analysis on a wheat panel phenotyped using the virulent and avirulent strains. Association mapping results revealed a strong signal associated with RAUDPCs for the avirulent strain IPO09593 (eight SNPs above the Bonferroni-corrected cutoff) in the telomeric region of chromosome 2B, but not with the virulent strain IPO323 (Fig. 2c). This confirms that IPO09593 triggers a strong resistance from varieties carrying a resistance

QTL in the vicinity of the *Stb9* gene. Additional details on *Stb9*-associated SNPs are provided in the supplemental Table S1. Finally, based on the lead SNP (AX-89505650; position 808,036,267 bp on chromosome 2B), phenotypic and allele analyses showed that ca. 8.2% of the French varieties likely carry *Stb9*.

Altogether, our combined functional and genetic analyses demonstrate that *Zt_1_693* is the avirulence gene *AvrStb9* and is involved in a gene-for-gene relationship with *Stb9* resistance in wheat.

AvrStb9 encodes a secreted protein with a conserved protease domain shared among diverse ascomycetes lineages.

Unlike other known fungal effectors, *AvrStb9* encodes a large protein and shares sequence similarities with different fungal species. The re-annotated *AvrStb9* gene model has five introns and encodes a 727 amino acid long protein with eight cysteine residues (Fig. 4a; 4b). This protein has a predicted signal peptide sequence of 18 amino acids at its N-terminus and two serine-protease domains (C-terminal processing peptidase S41 family; positions 70-726; Fig. 4b) according to InterProScan (Panther entry PTHR-37049, e-value: 2.7 e-64). HHpred searches in protein structure databases identified the secreted protease CPAF as the strongest hit (e-value of 1.5e-27), which is involved in pathogenesis of the human pathogenic bacterium *Chlamydia trachomatis* (Huang et *al.,* 2008).

To investigate whether *AvrStb9* originated before or after *Z. tritci* speciation, we used blastp searches against *Z. tritci* reference genome IPO323 and closely-related species from the *Zymoseptoria* family. We identified *AvrStb9* and its paralogs from IPO323 and all its closest relatives *Z. pseudotritici, Z. ardabiliae* and *Z. brevis* (Fig. S7a). Interestingly, most of these proteins had a predicted signal peptide, and they all displayed a C-terminal processing peptidase S41 domain. Using IPO323 *in planta* gene expression, we found that three out of the five paralogs (*Zt_4_489, Zt_10_226, Zt_11_236*) had an expression profile comparable to *AvrStb9* (Fig. S7b). According to these expression patterns, these genes could potentially play a role in infection.

Using OrthoMCL, we identified 122 putative *AvrStb9* orthologs/paralogs with an average identity of 33.6% and an average e-value of 8e-73 (Table S4). *AvrStb9* orthologs/paralogs are found in a broad range of ascomycetes lineages belonging to Eurotiomycetes, Leotiomycetes, Sordariomycetes, Orbiliomycetes and Dothideomycetes (Fig. 3; Table S4). Most of fungal species (48 out of 65 species) identified from the orthogroup are pathogenic, indicating that the well conserved *AvrStb9* orthologs/paralogs may be involved in host interaction processes. Interestingly, ~87% of these proteases possess a predicted signal peptide.

Diversification of *AvrStb9* leads to evasion of host recognition

Non-synonymous substitutions in avirulence gene sequences often mediate the evasion of host recognition. Analysis of sequence polymorphisms between the avirulent and virulent strains revealed a

6 amino acid deletion and 12 non-synonymous substitutions (Table 1; Fig. 4b). We crossed this data with the fungal population phenotypes and identified eight mutations potentially involved in host evasion for which more than ~94% of strains carrying the mutation were virulent on *Stb9* cultivars (Table 1). Five of these mutations fall into the virulence-domain cited above, namely A431, F473, S525, D565, L628. All of these mutations except D565 are located in one of the two putative protease domains (Fig. 4b), with F473, S525 and L628 being relatively conserved residues among proteins belonging to the CPAF structural family (in 34-57% of the analyzed proteins respectively; data not shown).

In order to investigate *AvrStb9* diversity at a larger scale, we extended our analysis to include four global *Z. tritici* populations. We found 44 different *AvrStb9* alleles in these populations, which translate to 38 different protein variants, mostly population-specific (Fig. 4c). Next, we analyzed *AvrStb9* nucleotide diversity including ~500 bp of up- and downstream sequences from the start and stop codons, respectively. We observed both a drop-in nucleotide diversity and Tajima's D in the *Avr* coding sequence compared to the flanking regions (Fig. 4a). A negative Tajima's D is indicative of an excess of low frequency polymorphisms, indicating population size expansion and/or selective sweeps.

The codon-based model analysis of *AvrStb9* coding sequence was used to investigate the selection pressure exerted on the avirulence gene. Likelihood ratio tests (LRT) for different models indicated that positive selection was highly probable, as selection models (M2 and M8) showed a significantly better data fit compared to the neutral models M1 and M7 (Table S5). Selection analysis revealed 23 residues (3.16% of the protein sequence) under positive diversifying selection (BEB probability > 0.95) in the M2 model. Interestingly, S525F and L628Y, had the strongest positive selection probability (BEB probability of = 1) among the five mutations potentially responsible of the virulence according to the previous analyses (Table 1). We postulate that these adaptive mutations are under strong selective pressure most probably to counteract recognition by the *Stb9* resistance gene.

Discussion

Gene-for-gene interactions in plant pathosystems are described when a host resistance gene recognizes a pathogen's avirulence product that triggers a strong host immune response. Although 23 isolate-specific *Stb* resistance genes have been mapped in wheat, only two have been cloned so far and their corresponding fungal avirulence genes remain largely unknown. Here, we report the identification and characterization of a novel avirulence gene, *AvrStb9*, which triggers complete resistance in wheat cultivars carrying *Stb9* resistance gene. Additionally, we performed a high-resolution mapping of *Stb9* locus and delivered tightly-linked SNP markers for its use in breeding.

AvrStb9 is highly expressed during the asymptomatic phase of *Z. tritici* infection and drastically downregulated upon the switch to the late necrotrophic phase. In a previous study, *AvrStb9* was among the top 100 highly expressed genes during the transition from biotroph to necrotrophy stage (Palma-Guerrero *et al.*, 2017), suggesting a function in host colonization during the asymptomatic phase or the

transition to the necrotrophic phase. Recognition of the *Avr* allele leads to full resistance in cultivars carrying *Stb9* following the gene-for-gene model. A previous study mapped the *Stb9* resistance locus to the long arm of chromosome 2B but the genomic interval remained very large (Chartrain *et al.*, 2009). We used *AvrStb9* strain to trigger wheat resistance and perform a precise mapping of the resistance locus, which revealed that *Stb9* resides in the telomeric region of wheat chromosome 2B in an interval of 79.99-81.10Mb ($r^2 > 0.50$). Only two STB resistance genes have been cloned to date, *Stb6* and *Stb16q*, both encoding receptor-like kinase proteins (Saintenac et *al.*, 2018; 2021), and our findings will help in the cloning of *Stb9* resistance gene.

AvrStb9 is an atypical fungal avirulence gene, since it encodes a relatively large secreted protein (727 aa). Up to now, avirulence genes from filamentous fungal pathogens encode small (< 300 aa) secreted proteins rich in cysteine. Only three exceptions to this rule are the *Ace1* gene from *Magnaporthe grisea* (Collemare *et al.*, 2008) encoding a large polyketide synthase (PKS) fused to a nonribosomal peptide synthetase (NRPS) likely involved in secondary metabolism, and two *AvrMla* genes from *Blumeria graminis hordei* which encode large proteins (Bourras *et al.*, 2016). Furthermore, in contrast to what has been shown for most avirulence genes, *AvrStb9* has a predicted functional domain corresponding to the family of C-terminal processing S41 proteases and show sequence similarities with proteins from diverse fungal species. In plant pathogenic bacteria, some type III secreted effectors injected in host plant cells are similar to catalytic enzymes involved in a variety of reactions such as E3 ligation or proteolysis (Hann *et al.*, 2010). For instance, *Pseudomonas syringae* secretes the cysteine protease *HopN1* to suppress the host innate response and promote virulence (López-Solanilla *et al.*, 2004).

The importance of proteases in fungal pathogenicity has previously been reported (Jashni *et al.*, 2015b; Chandrasekaran *et al.*, 2016). *Fusarium oxysporum* f. sp. *lycopersicum* secretes a serine protease and a metalloprotease that cleave host chitinases to prevent the degradation of fungal cell walls and facilitate host colonization (Jashni *et al.*, 2015a). The maize anthracnose fungus *Colletotrichum graminicola* secretes a highly conserved metalloprotease effector that enhances virulence (Sanz-Martín *et al.*, 2016). Similarly, the rice blast fungus *Magnaporthe oryaze* secretes during infection *Avr-Pita*, an effector protein with a predicted zinc metalloproteases domain that belongs to a family of proteases that include other avirulence genes recognized by the resistance gene *Pi-ta* (Orbach *et al.*, 2000; Khang *et al.*, 2008). Our protein analyses (OrthoMCL, InterProScan, HHpred) indicated that AvrStb9 has sequence and structural features typical of a protease. Protein structure-based homology searches identified the secreted protease CPAF from the human pathogenic bacterium *C. trachomatis* as closest homolog. CPAF degrades host transcription factors and therefore directly helps *Chlamydia* evading the host defenses. We postulate that AvrStb9 could play a similar role in proteolytic degradation of host protein substrates. However, the enzymatic activity of AvrStb9 is yet to be investigated, as our analysis relied primarily on sequence and structure comparisons. Further, the fact that we were hardly able to produce

AvrStb9-disrupted mutants indicates that the gene may play an important role in *Z. tritici* survival, and

that the proteolytic activity likely extends to include substrates essential for fungal growth.

The broad conservation of AvrStb9 in *Z. tritici* world population, its sister species and some other fungal

species suggested that this protein is a core effector shared among these lineages. Irieda et *al.* (2019)

reported the conservation of *NIS1* effector that was initially detected from two *Colletotrichum* spp. in a

wide range of Ascomycota and Basidiomycota lineages. *NIS1* targets plant immune kinases to suppress

PAMP-triggered immunity (Irieda *et al.*, 2019). We hypothesize that S41 proteases may have been

anciently acquired by fungi, and that the expansion of particular serine protease families is associated

with the evolution of fungal lifestyle. Indeed, S41 serine proteases are suspected to have a role in

programmed cell death and pathogenic life style adaptation in fungi (Muszewska *et al.*, 2017). We also

identified multiple paralogs from multiple fungal species, likely resulting from gene duplication events,

highlighting either the redundancy of these genes or their evolution to interact with different substrates.

The functional redundancy of virulence genes has been a technical challenge to study their biological

functions, as disruption mutants often lead to the conclusion that these genes are not essential for

pathogenicity (Plummer *et al.*, 2004; Have *et al.*, 2010). These redundancies may also increase the

organism's genetic robustness against null mutations or act in compensation of a gene loss (Gu *et al.*,

2003). In *Z. tritici* for instance, the different *AvrStb9* paralogs had a secretion signal and show an

expression pattern consistent with a role as effectors. Taken altogether, we hypothesize that *AvrStb9* and

related S41 proteases constitute a set of core effectors gained early in the evolution of fungal-plant

interactions.

Recognition of effectors by host resistance exerts an evolutionary pressure that favors sequence

diversification, effector deletion or acquisition to counteract the immune response. We analyzed

AvrStb9 sequence diversity and found a broad conservation of its protein residues, unlike the other

known *Z. tritici* avirulence genes, *AvrStb6* and *Avr3D1* (Brunner & McDonald, 2018; Meile *et al.*, 2018).

Nevertheless, we identified few substitutions undergoing strong positive diversifying selection.

Additional data from chimeric AvrStb9 transformants, from *Z. tritici* populations and from sequence

conservation among the CPAF family indicate S525 and L628 as probably the most important residues

involved in virulence, although combinations of mutations may be necessary. Based on the structural

model for CPAF function, one may speculate that a functional protease is required to escape host

recognition.

This is corroborated by the fact that, we did not observe in any of the global strains a loss of *AvrStb9* or

other deleterious mutations. Furthermore, *AvrStb9* deletion proved to be nearly impossible (only one

poorly growing transformant had the gene deletion out of the 271 tested). We therefore suggest that

AvrStb9 plays an essential role in the biology of *Z. tritici*. It is tempting to speculate that the functional

AvrStb9 protein actively degrades (through proteolysis) both fungal protein(s) and host defense

protein(s). Additional functional studies, including mutating the S41 protease active sites in virulent and avirulent alleles will shed light on the role of the protease activity of AvrStb9 in its diverse biological activities (fungal growth, wheat infection, recognition by *Stb9*). But also, the function of the wheat resistance factor Stb9 is intriguing. Does it work as classical receptor protein, recognizing the avirulent AvrStb9 protein? Does it inhibit AvrStb9 protease activity or, on the contrary, may the Stb9 resistance protein be the target of AvrStb9 protease?

In conclusion, we identified a new major avirulence factor of *Z. tritici* that triggers a strong immune response in hosts carrying the corresponding resistance gene. Unlike what has been described for most avirulence genes, *AvrStb9* encodes a large protein with a predicted protease domain and likely belongs to a family of conserved fungal S41 proteases. This is an important step in deciphering molecular mechanisms underlying wheat – *Z. tritici* interactions, and shed light on the role of 'atypical' effectors (*i.e.*, non-SSPs) in fungal avirulence. It opens new research avenues to broaden our knowledge on host-pathogen interactions, in particular the characterization of enzyme activity of effectors, which will greatly help our understanding of the biology of protease effectors, commonly found in fungal pathogens.

Data Availability

Genome sequencing data for the 103 French *Z. tritici* isolates is available at NCBI BioProject PRJNA777581 (accessions SRR16762555 to SRR16762657). *AvrStb9* gene sequences are available at NCBI Genbank (accession numbers OP376566 and OP376567).

Acknowledgements

We thank David Gouache, Henriette Goyeau, Gert H.J. Kema, Marc-Henri Lebrun, Frédéric Suffert and Anne-Sophie Walker for their contribution in establishing the fungal population; Johann Confais for his help in the virulence assays; Robert King and Jason Rudd for providing IPO323 RNA-seq data; Florence Cambon, the VégéPôle and GENTYANE platforms for the development of the Courtot NILs; Marc-Henri Lebrun for critical reading of the manuscript and valuable feedback; and Sophie Bouchet and Fanny Hartmann for assistance with the initial GWAS analyses. This work was supported by the French National Research Agency (program BIOADAPT, grant no. ANR-12-ADAP-0009-04), the French Fund to support Plant Breeding (FSOV 2008 H) and by BASF France SAS (V.I.S.A. project). INRAE BIOGER benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007).

Author contributions

T.C.M and S.F. conceived and designed the study. A.D. and S.G. performed virulence assays, C.A. performed the cloning and genetic transformations. S.O. provided DH seeds and linkage mapping data. C.S. provided the 'Courtot' NILs. J.C., N.L., J.A. and R.A performed bioinformatics analysis. D.C.

provided sequence data from the global *Z. tritici* populations. R.A, T.C.M and S.F. analyzed the data and wrote the manuscript. All authors have reviewed the manuscript.

References

Balfourier F, Bouchet S, Robert S, De Oliveira R, Rimbert H, Kitt J, Choulet F, International Wheat Genome Sequencing Consortium, BreedWheat Consortium, Paux E. 2022. Worldwide phylogeography and history of wheat genetic diversity. *Science Advances* **5**: eaav0536.

Battache M, Lebrun MH, Sakai K, Soudière O, Cambon F, Langin T, Saintenac C. 2022. Blocked at the Stomatal Gate, a Key Step of Wheat Stb16q-Mediated Resistance to *Zymoseptoria tritici*. *Frontiers in Plant Science* **13:** 921074.

Bolger AM, Lohse M, Usadel B. **2014**. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Bourras S, McNally KE, Müller MC, Wicker T, Keller B. **2016**. Avirulence genes in cereal powdery mildews: the gene-for-gene hypothesis 2.0. *Frontiers in Plant Science* **7**: 241.

Broman KW, Wu H, Sen Ś, Churchill GA. **2003**. R/qtl: QTL mapping in experimental crosses. *Bioinformatics* **19**: 889–890.

Brown JKM, Chartrain L, Lasserre-Zuber P, Saintenac C. **2015**. Genetics of resistance to *Zymoseptoria tritici* and applications to wheat breeding. *Fungal Genetics and Biology* **79**: 33–41.

Brunner PC, McDonald BA. **2018**. Evolutionary analyses of the avirulence effector AvrStb6 in global populations of *Zymoseptoria tritici* identify candidate amino acids involved in recognition. *Molecular Plant Pathology* **19**: 1836–1846.

Cabau C, Escudié F, Djari A, Guiguen Y, Bobe J, Klopp C. 2017. Compacting and correcting Trinity and Oases RNA-Seq *de novo* assemblies. *PeerJ* **5**: e2988.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. **2009**. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972–1973.

Chandrasekaran M, Thangavelu B, Chun SC, Sathiyabama M. **2016**. Proteases from phytopathogenic fungi and their importance in phytopathogenicity. *Journal of General Plant Pathology* **82**: 233–239.

Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. **2015**. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience* **4**: 7.

Chartrain L, Sourdille P, Bernard M, Brown JKM. **2009**. Identification and location of *Stb9*, a gene

for resistance to Septoria tritici blotch in wheat cultivars Courtot and Tonic. *Plant Pathology* **58**: 547–

555.

Collemare J, Pianfetti M, Houlle A-E, Morin D, Camborde L, Gagey M-J, Barbisan C, Fudal I,

Lebrun M-H, Böhnert HU. **2008**. *Magnaporthe grisea* avirulence gene *ACE1* belongs to an infection-

specific gene cluster involved in secondary metabolism. *New Phytologist* **179**: 196–208.

Danguy des Déserts A, Bouchet S, Sourdille P, Servin B. **2021**. Evolution of recombination

landscapes in diverging populations of bread wheat. *Genome Biology and Evolution* **13**: evab152.

Dyrløv Bendtsen J, Nielsen H, von Heijne G, Brunak S. **2004**. Improved prediction of signal peptides:

SignalP 3.0. *Journal of Molecular Biology* **340**: 783–795.

Flor HH. **1956**. The complementary genic systems in flax and flax rust**Joint contribution from the

Field Crops Research Branch, Agricultural Research Service, United States Department of Agriculture

and the North Dakota Agricultural Experiment Station. In: Demerec MBT-A in G, ed. Academic Press,

29–54.

Flutre T, Duprat E, Feuillet C, Quesneville H. **2011**. Considering transposable element diversification

in *de novo* annotation approaches. *PLOS ONE* **6**: e16526.

Fones H, Gurr S. **2015**. The impact of Septoria tritici blotch disease on wheat: An EU perspective.

Fungal Genetics and Biology **79**: 3–7.

Franceschetti M, Maqbool A, Jiménez-Dalmaroni MJ, Pennington HG, Kamoun S, Banfield

MJ. 2017. Effectors of filamentous plant pathogens: commonalities amid diversity. *Microbiology and*

Molecular Biology Reviews **81:** e00066–16.

Garrison E, Marth G. **2012**. Haplotype-based variant detection from short-read sequencing. *arXiv*:

1207.3907.

Goodwin SB, Ben M'Barek S, Dhillon B, Wittenberg AHJ, Crane CF, Hane JK, Foster AJ, van

der Lee TAJ, Grimwood J, Aerts A, *et al.* 2011. Finished genome of the fungal wheat pathogen

Mycosphaerella graminicola reveals dispensome structure, chromosome plasticity, and stealth

pathogenesis. *PLOS Genetics* **7**: e1002070.

Grandaubert J, Bhattacharyya A, Stukenbrock EH. **2015**. RNA-seq-based gene annotation and

comparative genomics of four fungal grass pathogens in the genus Zymoseptoria identify novel orphan

genes and species-specific invasions of transposable elements. *G3 Genes|Genomes|Genetics* **5**: 1323–

1333.

Gu Z, Steinmetz LM, Gu X, Scharfe C, Davis RW, Li W-H. **2003**. Role of duplicate genes in genetic
robustness against null mutations. *Nature* **421**: 63–66.

Hann DR, Gimenez-Ibanez S, Rathjen JP. **2010**. Bacterial virulence effectors and their activities.
Current Opinion in Plant Biology **13**: 388–393.

Hartmann FE, McDonald BA, Croll D. **2018**. Genome-wide evidence for divergent selection between
populations of a major agricultural pathogen. *Molecular Ecology* **27**: 2725–2741.

Hartmann FE, Sánchez-Vallet A, McDonald BA, Croll D. **2017**. A fungal wheat pathogen evolved
host specialization by extensive chromosomal rearrangements. *The ISME Journal* **11**: 1189–1204.

Have A ten, Espino JJ, Dekkers E, van Sluyter SC, Brito N, Kay J, González C, van Kan JAL.
2010. The *Botrytis cinerea* aspartic proteinase family. *Fungal Genetics and Biology* **47**: 53–65.

Huang Z, Feng Y, Chen D, Wu X, Huang S, Wang X, Xiao X, Li W, Huang N, Gu L, *et al.* **2008.**
Structural basis for activation and inhibition of the secreted chlamydia protease CPAF. *Cell Host*
Microbe **4:** 529-42.

**Irieda H, Inoue Y, Mori M, Yamada K, Oshikawa Y, Saitoh H, Uemura A, Terauchi R, Kitakura
S, Kosaka A,** *et al.* **2019.** Conserved fungal effector suppresses PAMP-triggered immunity by targeting
plant immune kinases. *Proceedings of the National Academy of Sciences* **116**: 496–505.

Jashni MK, Dols IHM, Iida Y, Boeren S, Beenen HG, Mehrabi R, Collemare J, de Wit PJGM.
2015a. Synergistic action of a metalloprotease and a serine protease from *Fusarium oxysporum* f. sp.
lycopersici cleaves chitin-binding tomato chitinases, reduces their antifungal activity, and enhances
fungal virulence. *Molecular Plant-Microbe Interactions®* **28**: 996–1008.

Jashni MK, Mehrabi R, Collemare J, Mesarich CH, de Wit PJGM. **2015b**. The battle in the
apoplast: Further insights into the roles of proteases and their inhibitors in plant–pathogen interactions.
Frontiers in Plant Science **6**: 584.

Jones JDG, Dangl JL. **2006**. The plant immune system. *Nature* **444**: 323–329.

**Kema GHJ, Mirzadi Gohari A, Aouini L, Gibriel HAY, Ware SB, van den Bosch F, Manning-
Smith R, Alonso-Chavez V, Helps J, Ben M'Barek S,** *et al.* **2018**. Stress and sexual reproduction
affect the dynamics of the wheat pathogen effector AvrStb6 and strobilurin resistance. *Nature Genetics*
50: 375–380.

Khang CH, Park S-Y, Lee Y-H, Valent B, Kang S. **2008**. Genome organization and evolution of the
AVR-Pita avirulence gene family in the *Magnaporthe grisea* species complex. *Molecular Plant-Microbe*
Interactions® **21**: 658–670.

Langlands-Perry C, Cuenin M, Bergez C, Ben Krima S, Gélisse S, Sourdille P, Valade R, Marcel TC. **2022**. Resistance of the wheat cultivar 'Renan' to Septoria leaf Blotch explained by a combination of strain specific and strain non-specific QTL mapped on an ultra-dense genetic map. *Genes* **13**: 100.

Letunic I, Bork P. **2016**. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Research* **44**: W242–W245.

Li H, Durbin R. **2009**. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**: 1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Subgroup 1000 Genome Project Data Processing. **2009**. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Li L, Stoeckert Jr CJ, Roos DS. **2003**. OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome research* **13**: 2178–2189.

Lipka AE, Tian F, Wang Q, Peiffer J, Li M, Bradbury PJ, Gore MA, Buckler ES, Zhang Z. **2012**. GAPIT: genome association and prediction integrated tool. *Bioinformatics* **28**: 2397–2399.

Lo Presti L, Lanver D, Schweizer G, Tanaka S, Liang L, Tollot M, Zuccaro A, Reissmann S, Kahmann R. **2015**. Fungal effectors and plant susceptibility. *Annual Review of Plant Biology* **66**: 513–545.

López-Solanilla E, Bronstein PA, Schneider AR, Collmer A. **2004**. HopPtoN is a *Pseudomonas syringae* Hrp (type III secretion system) cysteine protease effector that suppresses pathogen-induced necrosis associated with both compatible and incompatible plant interactions. *Molecular Microbiology* **54**: 353–365.

Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. **2015**. RDP4: Detection and analysis of recombination patterns in virus genomes. *Virus Evolution* **1**.

Meile L, Croll D, Brunner PC, Plissonneau C, Hartmann FE, McDonald BA, Sánchez-Vallet A. **2018**. A fungal avirulence factor encoded in a highly plastic genomic region triggers partial resistance to Septoria tritici blotch. *New Phytologist* **219**: 1048–1061.

Monod M, Capoccia S, Léchenne B, Zaugg C, Holdom M, Jousson O. **2002**. Secreted proteases from pathogenic fungi. *International Journal of Medical Microbiology* **292**: 405–419.

Muszewska A, Stepniewska-Dziubinska MM, Steczkiewicz K, Pawlowska J, Dziedzic A, Ginalski K. **2017**. Fungal lifestyle reflected in serine protease repertoire. *Scientific Reports* **7**: 9147.

Nielsen R, Yang Z. **1998**. Likelihood models for detecting positively selected amino acid sites and applications to the HIV-1 envelope gene. *Genetics* **148**: 929–936.

Omrane S, Sghyer H, Audéon C, Lanen C, Duplaix C, Walker A-S, Fillinger S. **2015**. Fungicide
efflux and the MgMFS1 transporter contribute to the multidrug resistance phenotype in *Zymoseptoria*
tritici field isolates. *Environmental Microbiology* **17**: 2805–2823.

Orbach MJ, Farrall L, Sweigard JA, Chumley FG, Valent B. **2000**. A telomeric avirulence gene
determines efficacy for the rice blast resistance gene *Pi-ta*. *The Plant cell* **12**: 2019–2032.

Palma-Guerrero J, Ma X, Torriani SFF, Zala M, Francisco CS, Hartmann FE, Croll D, McDonald
BA. **2017**. Comparative transcriptome analyses in *Zymoseptoria tritici* reveal significant differences in
gene expression among strains during plant infection. *Molecular Plant-Microbe Interactions®* **30**: 231–
244.

Petit-Houdenot Y, Lebrun MH, Scalliet G. 2021. Understanding plant-pathogen interactions in
Septoria tritici blotch infection of cereals. In: Achieving durable disease resistance in cereals. London,
UK: *Burleigh Dodds Science Publishing*, 263-302.

Plummer KM, Clark SJ, Ellis LM, Loganathan A, Al-Samarrai TH, Rikkerink EHA, Sullivan
PA, Templeton MD, Farley PC. **2004**. Analysis of a secreted aspartic peptidase disruption mutant of
Glomerella cingulata. *European Journal of Plant Pathology* **110**: 265–274.

Rocafort M, Fudal I, Mesarich CH. 2020. Apoplastic effector proteins of plant-associated fungi and
oomycetes. *Current Opinion in Plant Biology* **56:** 9-19.

Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE,
Sánchez-Gracia A. **2017**. DnaSP 6: DNA Sequence Polymorphism analysis of large data sets.
Molecular Biology and Evolution **34**: 3299–3302.

Rudd JJ, Kanyuka K, Hassani-Pak K, Derbyshire M, Andongabo A, Devonshire J, Lysenko A,
Saqi M, Desai NM, Powers SJ, *et al.* 2015. Transcriptome and metabolite profiling of the infection
cycle of *Zymoseptoria tritici* on wheat reveals a biphasic interaction with plant immunity involving
differential pathogen chromosomal contributions and a variation on the hemibiotrophic lifestyle
definition. *Plant Physiology* **167**: 1158–1185.

Saintenac C, Cambon F, Aouini L, Verstappen E, Ghaffary SMT, Poucet T, Marande W, Berges
H, Xu S, Jaouannet M, *et al.* 2021. A wheat cysteine-rich receptor-like kinase confers broad-spectrum
resistance against Septoria tritici blotch. *Nature Communications* **12**: 433.

Saintenac C, Lee W-S, Cambon F, Rudd JJ, King RC, Marande W, Powers SJ, Bergès H, Phillips
AL, Uauy C, *et al.* 2018. Wheat receptor-kinase-like protein Stb6 controls gene-for-gene resistance to
fungal pathogen *Zymoseptoria tritici*. *Nature Genetics* **50**: 368–374.

Sánchez-Vallet A, McDonald MC, Solomon PS, McDonald BA. **2015**. Is *Zymoseptoria tritici* a
hemibiotroph? *Fungal Genetics and Biology* **79**: 29–32.

Sánchez-Vallet, A, Fouché S, Fudal I, Hartmann FE, Soyer JL, Tellier A, Croll D. 2018. The Genome Biology of Effector Gene Evolution in Filamentous Plant Pathogens. *Annual Review of Phytopathology* **56:** 21-40.

Sanz-Martín JM, Pacheco-Arjona JR, Bello-Rico V, Vargas WA, Monod M, Díaz-Mínguez JM, Thon MR, Sukno SA. **2016**. A highly conserved metalloprotease effector enhances virulence in the maize anthracnose fungus *Colletotrichum graminicola*. *Molecular Plant Pathology* **17**: 1048–1062.

Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. **2019**. The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution* **3**: 430–439.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, et al. **2011**. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* **7**: 539.

Singh D, Simmonds J, Park RF, Bariana HS, Snape JW. **2009**. Inheritance and QTL mapping of leaf rust resistance in the European winter wheat cultivar 'Beaver.' *Euphytica* **169**: 253–261.

Stamatakis A. **2014**. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* **30**: 1312–1313.

Stergiopoulos I, de Wit PJGM. **2009**. Fungal Effector Proteins. *Annual Review of Phytopathology* **47**: 233–263.

Suffert F, Sache I, Lannou C. **2013**. Assessment of quantitative traits of aggressiveness in *Mycosphaerella graminicola* on adult wheat plants. *Plant Pathology* **62**: 1330–1341.

Su M-Y, Som N, Wu C-Y, Su S-C, Kuo Y-T, Ke L-C, Ho M-R, Tzeng S-R, Teng C-H, Mengin-Lecreulx D, et al. **2017**. Structural basis of adaptor-mediated protein degradation by the tail-specific PDZ-protease Prc. *Nature Communications* **8**: 1516.

Touzy G, Rincent R, Bogard M, Lafarge S, Dubreuil P, Mini A, Deswarte J-C, Beauchène K, Le Gouis J, Praud S. 2019. Using environmental clustering to identify specific drought tolerance QTLs in bread wheat (*T. aestivum* L.). *Theoretical and Applied Genetics* **132**: 2859-2880.

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. **2012**. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols* **7**: 562–578.

Yang N, McDonald MC, Solomon PS, Milgate AW. **2018**. Genetic mapping of *Stb19*, a new resistance gene to *Zymoseptoria tritici* in wheat. *Theoretical and Applied Genetics* **131**: 2765–2773.

Zhong Z, Marcel TC, Hartmann FE, Ma X, Plissonneau C, Zala M, Ducasse A, Confais J,
Compain J, Lapalu N, *et al.* **2017**. A small secreted protein in *Zymoseptoria tritici* is responsible for
avirulence on wheat cultivars carrying the *Stb6* resistance gene. *New Phytologist* **214**: 619–631.

Zhu T, Wang L, Rimbert H, Rodriguez JC, Deal KR, de Oliveira R, Choulet F, Keeble-Gagnère
G, Tibbits J, Rogers J, *et al.* **2021**. Optical maps refine the bread wheat *Triticum aestivum* cv. Chinese
Spring genome assembly. *The Plant Journal* **107**: 303–314.

Zwiers L-H, de Waard MA. **2001**. Efficient *Agrobacterium tumefaciens*-mediated gene disruption in
the phytopathogen *Mycosphaerella graminicola*. *Current Genetics* **39**: 388–393.

Tables

Table 1. Polymorphisms in AvrStb9 protein sequences between the wild type and transgenic mutant strains. Difference in amino acid residues between the virulent (avrstb9) and avirulent (Avrstb9) phenotypes are highlighted in orange while residues potentially involved in virulence are highlighted in red.

Strain	Mutant ID	Codon position / Amino acid residue													Phenotype (on *Stb9*)
		58	66-71	73	188	276	349	370	431	473	512	525	565	628	
Wild-type strains															
IPO323 (avrstb9)	-	R	PCNSTI	N	Y	T	A	T	A	F	S	S	D	L	Virulent
IPO09593 (Avrstb9)	-	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323 Avrstb9 transformants															
IPO323_Avrstb9 hph#5	E	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323_Avrstb9 hph#6	F	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323_Avrstb9 hph#9	H	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323_Avrstb9 hph#12	K	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323_Avrstb9 hph#14	L	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323_Avrstb9 hph#18	N	Q	Δ	D	V	N	V	N	D	Y	T	F	N	Y	Avirulent
IPO323 Avrstb9 chimeric transformants															
IPO323_Avrstb9 hph#3	I	Q	Δ	D	V	T	A	T	A	F	S	S	D	L	Virulent
IPO323_Avrstb9 hph#11	J	Q	Δ	D	Y	T	A	T	A	F	S	S	D	L	Virulent
IPO323_Avrstb9 hph#7	G	Q	Δ	D	Y	T	A	T	A	F	S	S	D	L	Virulent
IPO323_Avrstb9 hph#17	M	Q	Δ	D	V	N	A	T	A	F	S	S	D	L	Virulent
IPO323 avrstb9 transformants															
IPO323_avrstb9 hph#3	O	R	PCNSTI	N	Y	T	A	T	A	F	S	S	D	L	Virulent

IPO323$_{avrstb9}$ hph#4	P	R	PCNSTI	N	Y	T	A	T	A	F	S	S	D	L	Virulent
Percentage of virulent isolates in the fungal population		87.5	98.86	86.63	98.86	98.86	54.54	88.63	**98.86**	**98.86**	89.77	**98.86**	**98.86**	**95.54**	
Probability of dN/dS>1		0.97	-	0.99	0.99	0.58	0.54	0.59	0.93	0.60	**0.99**	**1.00**	0.65	**1.00**	

 Figures

 (a)

 (b)

Figure 1. GWAS identifies a candidate gene controlling avirulence on the cultivar 'Soissons'. (a) Manhattan plot representing the association of multiple SNPs with PLACN on cultivar 'Soissons'. The X and Y axes indicate positions on chromosome 1 and −log10 (*p*-value) for associations, while the red line represents the Bonferroni threshold (α=0.05). (b) Top: Gene annotations (black arrows) and transposable elements (grey arrow) on the reference genome IPO323. The candidate gene is highlighted in red. Bottom: Pairwise linkage disequilibrium plot around the lead significant SNP. The association peak is associated with several polymorphisms in the gene *Zt_1_693*. (c) The difference in PLACN of isolates carrying the virulent and avirulent alleles (based on the lead SNP). Significance difference from the mean was computed using a two-sided Mann-Whitney test (****, *p*<0.0001). (d) Gene expression profile of *Zt_1_693* measured as fragments per kilobase of exon per million mapped fragments (FPKM) throughout the time course of *Z. tritici* infection on a susceptible wheat cultivar (Rudd *et al.*, 2015).

(a)

(b)   **(c)**

Figure 2. *Zt_1_693* encodes for the avirulence gene *AvrStb9* and triggers *Stb9* resistance. (a) PLACP at 26 dpi
measured after inoculation with the avirulent isolate IPO09593, the virulent isolate IPO323 and IPO323 mutants
expressing the avirulent allele (IPO323*Avrstb9*) or the virulent allele (IPO323*avrstb9*) on cultivars 'Courtot' (*Stb9*), a
'Courtot' mutant lacking *Stb9* and 'Soissons' (details on the phenotype of each mutant is provided in Fig. S6;
Table S6). Statistical significance was determined using a Tukey's HSD test. (b) Genetic mapping of *Stb9*
resistance gene using linkage mapping. Top panel: Linkage map and marker position in cM. Bottom panel:
Logarithm of odds (LOD) score plot of the QTL analysis (Relative AUDPC for sporulation and necrosis in black
and grey, respectively) possibly containing the *Stb9* locus mapped to the long arm of the chromosome 2B. The
horizontal dashed line indicates a suggestive linkage threshold (LOD = 3). The position of the marker in the QTL
peak is highlighted in red. (c) Manhattan plots representing association mapping results of relative AUDPC
sporulation of the avirulent (top) and virulent (bottom) isolates in the bread wheat panel. Multiple SNPs in the
telomeric region of chromosome 2B associated with the avirulent isolate are located in the *Stb9* genomic region.

Figure 3. Phylogenetic tree of selected *AvrStb9* orthologs/paralogs belonging to five distinct classes of
Ascomycota lineages. Leaf labels are color-coded according to class and represent abbreviated names of the fungal
species (Table S4). AvrStb9 is indicated with a black arrow.

(a)

(b)

Figure 4. *AvrStb9* diversity and phylogeny in the global *Z. tritici* population. (a) The top panel shows the *AvrStb9*

gene structure. Middle and low panels are sliding window analyses for nucleotide diversity (π) and Tajima's D

across the *AvrStb9* locus, respectively, with a window size of 200 bp and a step size of 20 bp. The plots show a

reduced nucleotide diversity and a drop in Tajima's D values, likely corresponding to a selective sweep spanning

the region. (b) Schematic diagram of *AvrStb9* protein sequence with an annotation of amino acid substitutions

between virulent and avirulent alleles. Substitutions potentially involved in virulence are highlighted in red.

SP=Signal peptide. (c) A phylogenetic tree of *AvrStb9* protein sequence generated from a global *Z.*

tritici population and the reference isolate IPO323.