LRRK2 regulates synaptic function through modulation of actin cytoskeletal dynamics

Giulia Tombesi[1], Shiva Kompella[2], Giulia Favetta[1], Chuyu Chen[3], Marta Ornaghi[4,12], Yibo Zhao[5], Ester Morosin[1], Martina Sevegnani[5], Adriano Lama[6], Antonella Marte[7,11], Ilaria Battisti[8], Lucia Iannotta[1], Nicoletta Plotegher[1,9], Laura Civiero[1,9,10], Franco Onofri[7,11], Britta J Eickholt[4], Giovanni Piccoli[6], Giorgio Arrigoni[8], Dayne Beccano-Kelly[2,CA], Claudia Manzoni[5], Loukia Parisiadou[3] and Elisa Greggio[1,9,CA]

Author affiliations:

[1]Department of Biology, University of Padova, Italy

[2]School of Pharmacy and UK Dementia Research Institute, Cardiff University UK

[3]Northwest University, Chicago, USA

[4]Institute for Biochemistry, Charité Universitätsmedizin Berlin, Berlin, Germany

[5]School of Pharmacy, University College London, UK

[6]CIBIO, University of Trento, Italy

[7]Experimental Medicine, University of Genova, Italy

[8]Department of Biomedical Sciences, University of Padova, Italy

[9]Centro Studi per la Neurodegenerazione (CESNE), University of Padova, Italy

[10]IRCCS San Camillo Hospital, Venice, Italy

[11]IRCCS, Ospedale Policlinico San Martino, Largo Rosanna Benzi 10, 16132, Genova, Italy

[12] Freie Universität Berlin, Kaiserswerther Str. 16-18, 14195 Berlin, Germany

Corresponding authors: Dayne Beccano-Kelly (Beccano-KellyD@cardiff.ac.uk) and Elisa Greggio (elisa.greggio@unipd.it)

Running title: LRRK2, BDNF, synapse and actin cytoskeleton

Keywords: Parkinson's disease, LRRK2, actin cytoskeleton, drebrin, BDNF, synapse

Abstract

Parkinson's disease (PD) is a multisystemic disorder that manifests through motor and non-motor symptoms. Motor dysfunction is the most debilitating and it is caused by the degeneration of dopamine-producing neurons in the substantia nigra pars compacta (SNpc). Increasing evidence suggests that synapse dysfunction precedes neuronal loss by years. Still, early synaptic alterations in PD remain poorly understood.

Here we integrate literature meta-analysis, proteomics and phosphoproteomics with biochemical, imaging and electrophysiological measurements in neurons and brains from knockout and knockin Lrrk2 mouse models, as well as human iPSC-derived neurons lacking LRRK2. We demonstrate that phosphorylation of LRRK2 at Ser935 and of RAB proteins is induced by brain-derived neurotrophic factor (BDNF) stimulation in differentiated SH-SY5Y cells and primary mouse neurons. Affinity-purification coupled with mass spectrometry (AP-MS/MS) revealed a significant remodelling of the LRRK2 interactome following BDNF treatment, with enhanced association of LRRK2 to a network of actin cytoskeleton-related proteins. Gene-ontology analyses of both literature-curated LRRK2 interactors and phospho-proteome from striatal tissues with elevated LRRK2 activity (G2019S knockin mice) highlight synapse-actin remodelling as major affected pathways.

We further observed that loss of LRRK2 impairs BDNF signaling and alters postsynaptic density architecture. One month-old *Lrrk2* knockout mice display structural alterations in dendritic protrusions, a phenotype that normalizes with age. In human iPSC-derived neurons, BDNF enhances the frequency of miniature excitatory post-synaptic currents (mEPSC) in wild-type but not in LRRK2 knockout neurons, which appear to bypass this regulation through developmental compensation.

Taken together, our study discloses a critical role of LRRK2 in BDNF-dependent synaptic modulation and identifies the synaptic actin cytoskeleton as a convergent site of LRRK2's pathophysiological activity.

Introduction

Synaptic damage and connectome dysfunction are emerging as early pathological events preceding neurodegeneration and the manifestation of clinical symptoms in multiple neurodegenerative disorders, including Parkinson's disease (PD).[1–5] PD is an age-related motor disorder for which no cure is available. The motor symptoms typically manifest as a consequence of the progressive loss of the dopaminergic neurons of the Substantia Nigra pars compacta (SNpc) projecting to the striatum.[6] The degeneration of the nigrostriatal projections precedes the loss of the dopaminergic cell bodies in the SNpc and synaptic failure may be the triggering cause of axonal deterioration.[5,7] Pre- and postsynaptic alterations were described in PD animal models.[5] For example, PD pathology was reported as associated with an evident striatal spine loss, that appeared to be correlated with the degree of dopamine denervation prevalently due to glutamate excitotoxicity in striatal neurons.[8,9]

Accumulating evidence indicate that more than half of the causative genes and risk factors for PD have a function at the synapse. Mutations in *LRRK2* gene represent the most common cause of autosomal dominant late onset familial PD and variations around the *LRRK2* locus increase lifetime risk for PD.[10] Pathogenic mutations are clustered into the enzymatic core of the LRRK2 protein, composed by a Roc/GTPase and a kinase domain, bridged by a COR scaffold.[11–13] The most frequent mutation (G2019S) located in the kinase domain results in a protein with a gain of kinase activity, associated with increased cellular toxicity.[11] Previous findings from our team and other laboratories suggested that LRRK2 sits at the crossroads between cytoskeletal dynamics and vesicular trafficking. LRRK2 activity seems to be modulated by interactions with a diverse array of cytoskeletal and vesicle-associated proteins, and it regulates membrane-trafficking events *via* phosphorylation of a subset of Rab GTPases, including RAB3, RAB8, RAB10, RAB12, RAB35 and RAB43 (reviewed in [14]).[15,16] LRRK2 subcellular localization is dynamically controlled by phosphorylation/dephosphorylation of a cluster of N-terminal serine residues (e.g. Ser935 and Ser910), which determines 14-3-3 protein binding.[17] Phosphorylation of LRRK2 Ser935 and Ser910 is regulated by multiple kinases, including casein kinase 1 alpha (CK1a), IKKs and protein kinase A (PKA),[18–20] and serves as a scaffold for 14-3-3 protein binding and LRRK2 subcellular localization.[17,21] In immune cells, extracellular signals such as Toll-like receptor 4 (TLR4) agonists stimulate LRRK2 Ser935 and Ser910 phosphorylation to positively modulate immune-related responses.[18,22] In contrast, the

extracellular signals that orchestrate Ser935 phosphorylation enhancing LRRK2 activity in neurons are still unknown.

The ability of the neurons to relocate LRRK2 within or nearby the synapse could be particularly important as LRRK2 was suggested to provide the scaffold for the assembly of the signalling cascade effectors to sustain synaptic functions[15,23,24] In particular, LRRK2 has been shown to regulate synaptic vesicles cycling at the presynaptic site through interaction and phosphorylation of a panel of pre-synaptic proteins,[25] such as NSF,[26] synapsin I,[27] EndophylinA,[28] synaptojanin-1[29] and DNAJC6/Auxilin.[30] Interestingly, recent literature supports a role for LRRK2 in the dendritic spine compartment, where it was suggested to influence AMPA receptor trafficking,[31,32] spine morphology and functionality in developing brains.[33–35]

Here, starting from the observation that brain derived neurotrophic factor (BDNF) stimulates LRRK2 Ser935 and RABs phosphorylations in neurons, we used unbiased protein-protein interaction (PPI) and phosphoproteomic investigations to disclose a novel function of LRRK2 in regulating actin-dependent synaptic dynamics in complementary neuronal models.

Results

BDNF treatment rapidly increases LRRK2 phosphorylation at Ser935

Neurotrophic factors plastically shape neuronal activity by binding to their cognate receptor tyrosine kinases.[36] Brain-derived neurotrophic factor (BDNF) is particularly important to modulate neurotransmission in the cortico-striatal circuitry.[37] This circuitry is relevant for PD and presents with the highest LRRK2 expression in comparison with the rest of the brain.[38,39] To address whether LRRK2 activity is stimulated by BDNF, we prepared primary mouse cortical neurons from C57BL/6J wild-type mice and stimulated them with 100 ng/ml of BDNF at days *in vitro* 14 (DIV14) when synapses are fully functional (Fig. 1A) and LRRK2 expression is detectable.[40] Western blot analysis revealed a rapid and transient increase of Ser935 phosphorylation after BDNF treatment (Fig. 1B and 1C). Of interest, BDNF failed to stimulate Ser935 phosphorylation when neurons were pretreated with the LRRK2 inhibitor

MLi-2, indicating that either LRRK2 kinase activity controls the upstream kinases activated by BDNF or that the binding with MLi-2, a type I inhibitor, locks the kinase in a conformation that favors dephosphorylation/hinders phosphorylation during the stimulus.[41]

To gain mechanistic insights into the effects of BDNF stimulation on LRRK2 function, we differentiated neuroblastoma SH-SY5Y cells with 10 µM retinoic acid (RA) treatment for 6 days[42] and measured BDNF response by monitoring Tropomyosin receptor kinase B (TrkB) levels and phosphorylation of AKT and ERK1/2, two major downstream effector pathways of BDNF/TrkB signaling (Fig. 1D).[43] Expression of TrkB was undetectable in undifferentiated SH-SY5Y cells but was induced upon RA differentiation (Fig. 1E). Accordingly, differentiated SH-SY5Y cells respond to 100 ng/ml BDNF stimulation, as evidenced by increased phosphorylation of ERK1/2 and AKT (Fig. 1F-G). LRRK2 expression, low in undifferentiated cells, also increased upon differentiation (Fig. S1A), indicating that this is a suitable neuronal model to study endogenous LRRK2. Notably, BDNF stimulation enhanced Ser935 phosphorylation with kinetics comparable with that observed in primary neurons (Fig. 1H and 1I). To determine whether BDNF-induced Ser935 phosphorylation reflects LRRK2 kinase activation, we assessed phosphorylation of RAB10 at Thr73, an established LRRK2 substrate. As illustrated in Fig. 1J and 1K, BDNF treatment significantly increased RAB10 phosphorylation, supporting the role of BDNF as a physiological activator of LRRK2.

BDNF promotes LRRK2 interaction with actin cytoskeleton components enriched at the postsynapse.

To clarify the implications of BDNF-mediated LRRK2-activation on neuronal function, we used affinity purification coupled with mass spectrometry (AP-MS/MS). As the yields of endogenous LRRK2 purification were insufficient for AP-MS/MS analysis, we generated polyclonal SH-SY5Y cells stably expressing GFP-LRRK2 wild-type or GFP control. Notably, in this model, Ser935 phosphorylation showed only a modest increase upon BDNF treatment (Fig. S1B), and prolonged stimulation led to a reduction of Ser935 phosphorylation below baseline levels (Fig. 1B–C and Fig. 1H-I). We hypothesized that this apparent limited response could be due to high basal levels of Ser935 phosphorylation, approaching a ceiling effect under resting conditions. This is consistent with previous mass spectrometry analyses reporting that Ser935 is nearly fully phosphorylated (~100%) in the brain under basal conditions.[44] To test

whether high basal phosphorylation masks BDNF-induced increases, we pretreated the cells with MLi-2 for 90 minutes to lower Ser935 phosphorylation. After inhibitor washout, cells were stimulated with BDNF at multiple time points. In these conditions, BDNF robustly increased both Ser935 and RAB phosphorylation, peaking at 5–15 minutes and returning to baseline by 60–180 minutes (Fig. S1C). Notably, while the magnitude of pSer935 phosphorylation at the peak was similar to untreated controls - supporting the idea that Ser935 is near saturation under basal conditions - RAB phosphorylation levels significantly exceeded those of untreated cells. This indicates that, unlike Ser935, RAB phosphorylation remains far from saturated under basal conditions and serves as a more dynamic readout of LRRK2 kinase activity. These data further support BDNF as a physiological activator of LRRK2 and validate GFP-LRRK2 SH-SY5Y cells for interrogating stimulus-induced changes in LRRK2 signaling and protein interactions.

Next, we used GFP-trap purification to isolate GFP or GFP-LRRK2 from RA-differentiated cells unstimulated or stimulated with BDNF for 15 mins prior to MS analysis. The unstimulated LRRK2 interactome (GFP-LRRK2 vs. GFP) is large (207 significant hits) and consistent with previous computational analysis of LRRK2 PPIs (Fig. 2A).[45,46] In particular, we confirmed interactions with known LRRK2 binders including 14-3-3 proteins, cytoskeletal proteins and protein translation factors (Supplementary.xls table 1). Since BDNF is synaptically released and influences pre- and post-synaptic mechanisms, we used SynGO[47] to search for LRRK2 interactors that are enriched at the synapse. About one-third (35%) of LRRK2 PPI under unstimulated conditions are SynGO annotated genes; among them, 33% are presynaptic proteins and 66% are post-synaptic proteins (Fig. 2B). The most significant enriched categories were pre- and post-synaptic ribosome and post-synaptic actin cytoskeleton (Fig. 2B).

We next compared unstimulated vs. BDNF-stimulated LRRK2 interactome. As shown in Fig. 2C, 15 minutes of BDNF stimulation reshapes LRRK2 PPI, with a group of interactors increasing binding with LRRK2 (blue dots) and another group decreasing binding (orange dots). Inclusion criteria were: (i) P value < 0.05 and fold change (FC) > 0.5 or FC < -0.5, or (ii) absolute fold change (|FC|) > 1 regardless of the P value. In this way, we increased the discovery power for gene ontology analysis. SynGO gene ontology revealed that 10 out of 19 hits with increased binding after BDNF [BDNF(+)] and 21 out of 50 hits with decreased binding [BDNF(-)] are synaptic proteins (Fig. 2C inset). Among synaptic interactors, the BDNF(+) binders are enriched in actin-cytoskeleton-related GO:BP categories while the BDNF(-) binders are enriched in protein translation GO:BP categories (Fig. 2D). Furthermore,

80% of synaptic BDNF(+) LRRK2 binders (DBN1, ARPC2, ACTR2, ACTR3, CFL1, CALD1, ACTN1, FLNA) and 62% of synaptic BDNF(-) binders (RPS2, RPS4X, RPS7, RPS9, RPS11, RPS13, RPS15A, RPL13, RPL23, RPL26, RPL27A, TUFM, EEF1A2, PCBP1) form functional and physical interaction networks (Fig. 2E-F).

To validate these findings, we performed western blot analysis on the AP samples analyzed by MS. We selected drebrin, encoded by *DBN1*, because it sits at the center of the BDNF(+) network (Fig. 2E, inset SynGO) and, in DIV14 primary neurons, it colocalizes with PSD95, a postsynaptic marker (Fig. S2A), confirming previous observations.[48] As predicted, BDNF treatment increases LRRK2:drebrin interaction by approximately twofold (Fig. 2G and Fig. S2B). To further characterize this interaction, we performed co-immunoprecipitation in HEK293T cells co-expressing Flag-LRRK2 and either full-length YFP-drebrin, its N-terminal (1-256 aa), or C-terminal (256-649 aa) fragments. These experiments confirmed a direct interaction between LRRK2 and drebrin, with the N-terminal domain appearing to mediate the majority of the binding (Fig. S2C). Notably, this region contains both the ADF-H (actin-depolymerizing factor homology) domain and a coiled-coil motif known to associate with actin.[48,49] Finally, we observed a significant decrease in Ser935 phosphorylation and a trend decrease in Rab phosphorylation in *Dbn1* KO mouse brains (Fig. 2I). This finding further strengthen the notion that drebrin forms a complex with LRRK2 that is important for its activity, e.g. upon BDNF stimulation.

We conclude that BDNF stimulation engages LRRK2 into actin cytoskeleton dynamics via its interaction with actin-regulatory proteins. While SH-SY5Y cells may only contain sparse synapses[50], the enrichment of this BDNF-dependent protein network at synapses indicates that this site might be relevant for LRRK2 function. Our findings also suggest that a pool of LRRK2 binds protein translation factors under unstimulated conditions, which redistributes toward actin-rich domains upon BDNF-TrkB signaling.

Convergence on actin and synaptic processes in literature-curated LRRK2 interactome and LRRK2 G2019S striatal phospho-proteome.

We recently investigated the molecular environment surrounding LRRK2 specifically by constructing the protein-protein interaction (PPI) interactome around LRRK2 based on PPIs derived from peer reviewed literature. After filtering the LRRK2 PPIs for reproducibility (keeping only those interactions that have been reported at least twice in literature or at least replicated with 2 different interaction detection methods) we obtained the interactions linking across the LRRK2 interactors thus effectively building the LRRK2 protein interaction network (LRRK2$_{net}$) which describes the protein milieu around LRRK2 in the cellular context (see Zhao et al.[51] for details). The LRRK2$_{net}$ was topologically analyzed to identify the portions (clusters) of the network that are more densely connected locally in comparison with the overall connectivity of the entire network as these topological clusters are likely to represent functional units.[52] Among the 11 topological clusters identified in Zhao et al.[51], we further elaborated on one cluster in particular, containing 41 genes (Fig. 3A) and found that it exhibited significantly GO-BP enriched terms into semantic categories related to: actin polymerization (16 GO-BP terms), synaptic vesicles (5 GO-BP terms), postsynaptic actin-cytoskeleton (2 GO-BP terms), lamellipodium (6 GO-BP terms), cell-neuron projection/size (2 GO-BP terms), DNA recombination/repair (4 GO-BP terms) and uncategorized (6 GO-BP terms) (Fig. 3B and Supplementary.xls table 2). Out of the 41 proteins forming the cluster, 23 are proteins associated with the actin cytoskeleton including the 4 larger nodes (where the dimension of the nodes is directly proportional to the node degree/connections within the network): DBN1/drebrin, CAPZA2 and LIMA whose association with LRRK2 was identified in Meixner et al.[53] and IQGAP1 identified in Liu et al.[54] (Fig. 3C). Given the overrepresentation of synapse-related GO-BP terms, we further performed GO functional enrichment using SynGO and mapped 16 synaptic proteins within the cluster (Fig. 3C). Remarkably, DBN1/drebrin is a key protein in the cluster, as it engages in multiple interactions (16 edges) and sits at the interface between the nodes involved in GO:BPs functions related to the actin cytoskeleton (blue-colored nodes) and to the synapse (red-colored nodes) (Fig. 3C and Fig. S3).

Given the multiple lines of evidence pointing to the synapse-actin couple as a key area for LRRK2 function, and based on the increased LRRK2 activity upon neurotrophin stimulation, we next asked whether LRRK2 kinase activity is also linked to synaptic function. To this aim, we performed an unbiased phospho-proteomic characterization of the striatum, using knockin mice expressing the hyperactive G2019S mutation to probe the contribution of kinase activity. We isolated total striatal proteins from WT and G2019S striata, enriched the phosphopeptides (Fig. 3D) (also see[55]), and normalized the log$_2$ intensity of each phosphopeptide to the log$_2$

intensity of its corresponding total protein. Comparing WT and G2019S samples, we identified 228 differentially regulated phosphopeptides (P value < 0.05 and $|\text{Log}_2$ fold change (FC)$| > 2$) (Fig. 3E and supplementary.xls table 3). Gene ontology analysis of differentially phosphorylated proteins, using stringent term size (<200 genes), revealed enrichment of post-synaptic spines and presynaptic active zones (Fig. 3F). A SynGO analysis confirmed enrichment of both pre- and postsynaptic categories, with particular high significance for terms related to the postsynaptic cytoskeleton (Fig. 3G). Despite starting from whole striatal tissue - which contains a mix of different cellular populations - the strong synaptic enrichment of phosphorylation changes further supports the critical role of this compartment in LRRK2 brain function. By merging the three datasets (the LRRK2 interactome in SH-SY5Y cells, the literature-based synaptic/actin cluster and the phosphoproteomic dataset from G2019S striatum), we found three common hits: Dbn1/drebrin, Eef1d and Mprip (Fig. 3H). In particular, phosphorylation of drebrin at Ser339 was 3.7 fold higher in G2019S mice. This site is located in the N-terminal region of drebrin, which exhibited higher affinity for LRRK2 (Fig. 3E and Fig. S2C). To infer the potential physiological relevance of this phosphorylation, we examined its detection frequency in phosphoproteomic studies (67 observations, https://www.phosphosite.org/proteinAction.action?id=2675&showAllSites=true) and assessed its predicted pathogenicity using AlphaMissense.[56] Although the AlphaFold structural model has low confidence in this region of drebrin, amino acid substitutions at Ser339 are predicted to be non-pathogenic (Fig. S4), supporting the notion that LRRK2-mediated phosphorylation at this site acts as a regulatory modification for drebrin activity.

BDNF-mediated signalling is impaired in LRRK2 knockout neurons.

Having collected multiple lines of evidence linking LRRK2 to actin cytoskeletal proteins enriched at the synapse, and given that BDNF - a neurtrophin known to promote synaptic plasticity - activates LRRK2 (Figure 1), we next asked whether loss of LRRK2 could affect BDNF signalling. To this end, we generated LRRK2 knockout (KO) SH-SY5Y cells with CRISPR/Cas9 editing (Fig. S5 and supplemental file 1), differentiated them with RA and assessed their response to BDNF stimulation. While application of BDNF increased AKT and ERK1/2 phosphorylation in naïve SH-SY5Y, as expected, LRRK2 KO cells exhibited a weaker response (Fig. 4A-B). These results indicate that LRRK2 acts downstream of TrkB and

upstream of AKT and ERK1/2, in line with its rapid and transient phosphorylation kinetic upon BDNF stimulation (Fig. 1B-C).

Likewise, *Lrrk2* WT and KO primary cortical neurons at DIV14 were stimulated with BDNF at different timepoints. Similar to differentiated SH-SY5Y cells, *Lrrk2* WT neurons responded to BDNF by rapidly increasing Akt and Erk1/2 phosphorylation, whilst KO neurons exhibited a significantly reduced phosphorylation of Akt and Erk1 (Fig. 4C-D).

Considering that (i) BDNF induces dendritic spine formation, maturation and structural plasticity,[57,58] (ii) BDNF(+) LRRK2 interactors are actin-related proteins enriched at the postsynapse (Fig. 2) and (iii) loss of LRRK2 causes reduced phosphorylation of BDNF/TrkB downstream kinases AKT and ERK1/2 (Fig. 4A-D), we evaluated BDNF-induced PSD95 sites in the presence or absence of LRRK2, as a potential process affected by LRRK2 deletion. Primary cortical neurons were transfected with GFP filler at DIV4 and treated with BDNF for 24 h at DIV13 following previous protocols[59,60] (Fig. 4E). As shown in figure 4F-G, BDNF treatment significantly increased the density of PSD95-positive puncta in WT cultures ($P <$ 0.001). In contrast, *Lrrk2* KO neurons did not exhibit increased PSD95 puncta upon BDNF. These findings further support a role for LRRK2 within the BDNF signalling cascade and show that the absence of LRRK2 impairs the remodelling of dendrite protrusions in response to BDNF stimulation.

Morphological Alterations of Dendritic Protrusions in Young Lrrk2 Knockout Mice.

Based on *in vitro* results indicating that BDNF stimulates LRRK2 activity and promotes its recruitment to the actin-cytoskeleton, enhancing the number of PSD95-positive dendritic protrusions, we next investigated whether loss of *Lrrk2* results in dendritic spine defects in the mouse brain. During development, dendritic spines are highly dynamic, with a rate of extension and retraction that allows proper and accurate synapse formation and neuronal circuit assembly.[61] By the adulthood, dendritic spines formation, motility, and pruning decrease reaching a relatively stable number of protrusions.[62] Since spines dynamics vary according to the stage of brain development, we performed a longitudinal analysis of young (1 month-old), adult (4 month-old) and aged (18 month-old) *Lrrk2* WT and KO mice in order to capture any age-dependent defect (Fig. 5A). We focused on the dorsal striatum, a region highly enriched in

SPNs that receives excitatory afferents from the cortex and dopaminergic modulation from the SNpc. This region is relevant in the prodromal stages of PD neurodegeneration[63] and expresses high levels of LRRK2[38,64]. Moreover, several studies showed a key role of LRRK2 in shaping the structure and function of excitatory synapses in the striatum.[33,34,65,66] We evaluated the number and morphology of dendritic spines using Golgi-Cox staining (Fig. 5B). The total number of protrusions varied across ages, but there were no differences between the two genotypes (Fig. 5C). Instead, the width and the length of the protrusions were reduced in 1 month-old Lrrk2 KO striata with respect to WT. Specifically, the average neck height was 15% shorter and the average head width was 27% smaller, indicating that the protrusions are smaller in *Lrrk2* KO brains. No differences were observed at 4 and 18 months of age (Fig. 5C). We then classified the morphology of the protrusions into four categories: filopodia and thin protrusions (immature spines) and mushroom and branched protrusions (mature spines). One month-old *Lrrk2* KO animals exhibit a reduced number of filopodia and an increased quantity of thin protrusions, suggesting that loss of Lrrk2 may favor this transition. Instead, the proportion of mature spines (mushroom and branched) remained unaltered. No differences were observed in older mice (Fig. S6).

Since the overall width/height of *Lrrk2* KO protrusions is reduced (Fig. 5C), we next analyzed the postsynaptic ultrastructure using electron microscopy. To increase the experimental power, we used a separate group of animals. We measured the postsynaptic density (PSD) length as it directly correlates with the amount of postsynaptic receptors and signaling proteins, thus providing an indication of the synaptic strength (Fig. 5D). PSD is significantly shorter in 1 month-old *Lrrk2* KO animals compared to WT ($P < 0.001$), while no differences in PSD length are observed in 18 month-old KO mice (Fig. 5E). Taken all these data together, we conclude that subtle defects in the postsynaptic density are present in the striatum of postnatal *Lrrk2* KO mice, but these abnormalities appear to normalize with age.

To assess whether the loss of *Lrrk2* in young mice leads to reduced dendritic protrusion size and postsynaptic density through alterations in BDNF-TrkB signaling or postsynaptic scaffold expression, we measured transcript levels of *Bdnf*, *TrkB*, *Dbn1, Psd95* and *Shank3*. Neither *Bdnf* nor *TrkB* expression is altered in the striatum, cortex or midbrain of 1-month old *Lrrk2* KO as compared to WT mice (Fig. 5F), supporting a mechanism whereby LRRK2 acts downstream of BDNF signaling. Instead *Dbn1*, *Psd95* and *Shank3* expression levels are reduced in *Lrrk2* KO mice across most brain regions analyzed, with the most pronounced decrease observed in the midbrain (Fig. 5F).

Given (i) the central role of *Dbn1*/drebrin in the LRRK2 synaptic network (Fig. 2-3), (ii) its key function as an actin regulatory protein at the synapse and (iii) its expression highly enriched at dendritic spines (Fig. S2A), we assessed drebrin protein levels at the 3 stages of life in whole brain lysates. In agreement with the qPCR results, drebrin levels are significantly lower in 1 month-old *Lrrk2* KO mice but not in older animals (Fig. 5G), further linking altered spine maturation with improper actin-dynamics in the absence of LRRK2.

Taken together, our results indicate that LRRK2 influences dendritic spine structural dynamics during brain development.

LRRK2 knockout in human iPSC-derived neurons increases basal spontaneous activity.

To further explore the connection between LRRK2 and BDNF activity in a cell model more relevant for human disease, we differentiated human induced pluripotent stem cells (hiPSCs) into cortical neurons following established protocols.[67] Wild-type and isogenic LRRK2 knockout (KO) hiPSCs[68] were differentiated in parallel for 50, 70 and 90 days (3 independent culture rounds per genotype), prior to subjecting them to patch clamp to assess spontaneous activity in the absence or presence of acute exposure to BDNF (24 hrs, 50 ng/ml) (Fig. 6A).

Acute BDNF treatment led to a significant increase in the mean frequency of miniature post-synaptic excitatory currents (mEPSC) in WT cultures (Fig. 6B-C). This finding is corroborated by the significant leftward shift in cumulative probability curves of interevent intervals (IEIs) between BDNF-treated neurons versus untreated neurons (Fig. 6D). In contrast, KO neurons already display elevated mEPSC frequency, comparable to that of BDNF-treated WT cultures. No significant change in mEPSC frequency was observed in KO neurons upon BDNF treatment (Fig. 6C-D). Furthermore, BDNF treatment had no effect on the average peak amplitude in either genotypes, before or after BDNF treatment (Fig. 6E-F).

To determine whether this apparent sealing effect of LRRK2 KO is restricted to DIV70 cultures or measurable also in younger or more mature neurons, we quantified the mEPSCs in DIV50 and DIV90 neurons. As illustrated in Fig. S7 (A-B), no significant differences across genotypes or treatments are observed at these timepoints, indicating that the genotype effect is transient during culture maturation.

Since only the frequency, but not the amplitude, of mEPSCs was altered at DIV70, we infer that BDNF primarily affects the presynapse rather than the postsynapse in this experimental paradigm, likely increasing neurotransmitter release in WT neurons. To determine whether the increased release was due to increase in release probability (P_r) or a higher number of synaptic contacts, we measured the number of presynaptic Bassoon colocalization with postsynaptic Homer (Fig. S7C-D). Confocal imaging of synapses combined with IMARIS volume reconstruction showed no increase in the total number of synapses under BDNF stimulation in neither genotype, suggesting that BDNF increases P_r rather than synapse number.

Discussion

Understanding the early events that lead to synaptic dysfunction in PD is crucial to design effective therapeutic approaches. Previous work from our laboratories and other groups have shown that the PD-associated kinase LRRK2 regulates important synaptic processes.[69,70] At the presynapse, LRRK2 controls synaptic vesicles cycle through association with its WD40 domain[23,24,71,72] and phosphorylation of a panel of presynaptic proteins, including N-ethylmaleimide sensitive fusion (NSF)[26], synapsin I[27], auxilin[30], synaptojanin[29] and EndophylinA.[28] It has also been shown to regulate neurotransmitter release via modulation of presynaptic vesicular interacting proteins. At the postsynapse, LRRK2 plays key roles in shaping the function of excitatory synapses in the striatum.[33,34,65,66,70] However, the mechanisms underlying LRRK2 synaptic function are still poorly elucidated. In the present study we provide multiple lines of evidence that LRRK2 affects synaptic function in response to BDNF neurotrophic stimulation, which promotes relocalization of LRRK2 to the actin-cytoskeleton in a phosphorylation-dependent manner.

There are several novel findings in our study. First, we identified BDNF as an extracellular stimulus that increases LRRK2 activity. The phosphorylation kinetics are rapid and transient, and both *Lrrk2* KO neurons and differentiated SH-SY5Y LRRK2 KO cells display reduced activation of downstream AKT and ERK1/2, consistent with LRRK2 acting downstream of TrkB/BDNF and upstream of ERK1/2 and AKT. As previously documented, we observed high Ser935 phosphorylation under basal conditions, in agreement with mass spectrometry determination showing nearly complete phosphorylation of this residue in whole brain.[44] In contrast, phosphorylation of RAB substrates, as well as BDNF-downstream kinases ERK and

AKT, is minimal at baseline. Thus, the modest increase in Ser935 phosphorylation upon BDNF stimulation may reflect a ceiling effect. Prolonged BDNF exposure leads to a reduction in Ser935 phosphorylation below pre-stimulation levels, likely through negative feedback mechanisms such as phosphatase activity or receptor desensitization.[73] In contrast, phosphorylation of RABs did not drop below pre-stimulation levels. This aligns with the low pRAB stoichiometry in the brain[44] and suggest that these targets remain responsive and largely unsaturated under basal conditions. Taken together, these findings highlight the nuanced regulation of LRRK2 activity and indicate a positive correlation between Ser935 phosphorylation and LRRK2 activity under BDNF signalling, similar to previous observations on microglia cells upon inflammatory stimuli.[74] They also imply that phosphorylated LRRK2 remains incompletely active unless further stimulated (e.g., by BDNF), suggesting a multi-step activation mechanism. Thus, Ser935 phosphorylation status does not always correlate with LRRK2 functional output, potentially reflecting distinct regulatory states or compartment-specific activation dynamics.[75]

Using a literature-based network analysis of LRRK2 PPIs, combined with AP-MS/MS in SH-SY5Y cells and phosphoproteomics of hyperactive Lrrk2 G2019S from brain striatum, we identified a highly interconnected cluster of actin and synaptic proteins linked to LRRK2. Both BDNF-activated LRRK2 and hyperactive LRRK2 G2019S mutant are associated with GO terms related to synaptic function and cytoskeletal organization. The prevalence of synaptic components in the phoshoproteomic study is particularly notable considering that it was performed on total striatal lysates composed of an heterogeneous mixture of cell types. This strengthens the importance of LRRK2-mediated signaling in the synaptic compartment.

The central role of synaptic dysfunction in PD neurodegeneration is increasingly recognized[76] and further supported by the latest and largest meta-analysis of PD genome-wide association studies (GWAS), which identified 134 loci as risk factors for PD.[77] The open reading frames closest to the risk signals showed strong enrichment for neuronal genes involved in dendritic spine and synapse biology (Fig. S8 and [77]). Even though these data require functional validation (GWAS directly identifies risk loci but not genes), they may further support synaptic impairment as an hallmark of PD.[78]

Of note, while LRRK2 G2019S cases are often characterized by the absence of Lewy bodies[79,80], two recent studies disclosed the presence of widespread alpha-synuclein oligomers in LRRK2-PD brains, suggesting that early-stage presynaptic alpha-synuclein aggregates may

contribute to synaptic dysfunction in LRRK2-PD.[81,82] Whether these alpha-synuclein oligomers are enriched in specific dopaminergic subtypes is unknown, and future studies should investigate if there is any correlation with the dopaminergic neuronal subpopulations most vulnerable in PD.[83,84]

Further supporting a contribution of LRRK2 on synaptic function, we found that knockout of mouse *Lrrk2* resulted in blunted BDNF-induced synapse maturation *in vitro*. These findings suggest a mechanism whereby BDNF signaling reshapes the LRRK2 interactome to promote its incorporation into a macromolecular complex required for coordinating actin dynamics during synaptic structural plasticity. Interestingly, in human iPSCs-derived cortical neurons, loss of LRRK2 seems to accelerate synaptic maturation: DIV70 KO neurons exhibit high spontaneous synaptic activity equivalent to BDNF-stimulated WT neurons. This indicates that LRRK2 may act to suppress premature maturation and temporally gate BDNF responsiveness, aligning with the differences in spine structural dynamics observed in KO mice. While previous evidence has shown that LRRK2 modulates neurotransmitter release,[22] our current findings imply that these changes can be acutely altered by BDNF signalling with loss of LRRK2 removing this plastic capacity.

BDNF is key to reorganize the actin cytoskeleton during long term potentiation (LTP) through the action of Rho GTPases Rac1 and Cdc42.[85] LRRK2 was previously reported to interact with Rac1[86,87] and with other actin-related proteins.[53] Here we demonstrated that LRRK2 association with actin cytoskeleton in dendritic spines increases upon BDNF stimulation. We validated the BDNF-dependency of the interaction between LRRK2 and drebrin, an actin binding protein that promotes stable F-actin formation in dendritic spines.[48] A LRRK2-drebrin link is supported by multiple lines of evidence from our data, including increased phosphorylation at Ser339 in G2019S knockin striata and reduced drebrin levels in Lrrk2 knockout brains during development, which matched with morphological alteration of dendritic protrusions. Ser339 in human drebrin is located in the N-terminal region, which we identified as important for the interaction with LRRK2. This site lies just before the proline-rich (PP) domain of the protein. The N-terminal region is positioned between the actin-binding domains and the C-terminal Homer-binding sequences, and it plays a key role in protein-protein interactions and cytoskeletal regulation.[88] Notably, this region was shown to interact with adafin (ADFN), a protein involved in cytoskeletal-related processes crucial for synapse formation and function. Adafin regulates puncta adherentia junctions, presynaptic differentiation, and cadherin complex assembly, essential for hippocampal excitatory synapses,

spine formation, and learning and memory processes.[89] Of note, adafin is listed among LRRK2-interacting proteins (https://www.ebi.ac.uk/intact/home), supporting a possible functional relevance of LRRK2-mediated drebrin phosphorylation in adafin-drebrin complex formation.

It has been shown that drebrin knockout results in delayed synaptogenesis and inhibition of postsynaptic PSD95 accumulation, indicating a key role in spine maturation.[48,90,91] During LTP, Ca^{2+} entry through N-methyl-d-aspartate (NMDA) receptors causes drebrin exodus from the spine compartment to allow actin monomers entrance and polymerization to increase the size of the cytoskeleton.[92] We postulate that LRRK2 may favor drebrin exodus at the shaft of the dendrite during plasticity processes. Actin remodeling during spine enlargement is also coordinated by the ARP2/3 complex, which is activated by binding to WAVE family proteins downstream of active Rac1.[93] Strikingly, we found that BDNF stimulates LRRK2 interaction with three out of seven subunits of the ARP2/3 complex (ACTR2, ACTR3 and ARPC2) and LRRK2 was reported to interact with WAVE proteins.[94] Altogether these findings, in agreement with our earlier study,[35] highlight a crucial role for LRRK2 in actin-based dendritic spine remodeling. Clearly, future studies should be directed at clarifying the precise molecular dynamics and kinetics of these processes.

Another interesting aspect disclosed in our study is a developmental phenotype characterized by defects in dendritic protrusions morphogenesis as consequence of loss of LRRK2. Interestingly, this developmental abnormality is rescued with age. This is not surprising considering that the homologous kinase LRRK1 may compensate for LRRK2 deficiency, as suggested by the overt neurodegeneration observed in double *Lrrk1/Lrrk2* KO mice but absent in the single KO[95–97] and the redundant but not overlapping pattern of expression of the two kinases.[98] Indeed, striatal Lrrk2 expression in the rodent brain increases up to postnatal week 4,[35,64,98,99] while Lrrk1 transcript in this region is relatively stable during development.[64]

Several lines of evidence indicate that genetic, biochemical and cellular alterations of BDNF/TrkB signaling are linked to PD. At striatal synapses, BDNF/TrkB signaling enhances both dopamine and glutamate release and ERK1/2 activation.[100,101] BDNF is required for the establishment of the proper number and for the survival of dopaminergic neurons in the SNpc,[102,103] and presynaptic dopamine release in the striatum is enhanced by BDNF.[37] BDNF levels are lower in peripheral tissues, brain, and blood of sporadic PD patients.[104] Moreover, the V66M polymorphism modifies the risk of sporadic and mutant LRRK2-associated PD.[105,106] Decreased BDNF concentration in serum and brain correlates with an increased degeneration

of dopaminergic neurons in PD[107] and with loss of striatal DA transporter (DAT) in patients with striatal dopaminergic neurodegeneration.[108] Thus, robust literature support deregulation of striatal BDNF signaling in PD. Future studies should be directed at investigating possible deregulated mechanisms of BDNF signaling in G2019S mice and hiPSCs-derived neurons with G2019S mutation.

At the onset of PD, loss of dopaminergic axonal terminals largely exceeds loss of cell bodies,[4] implying that early synapse deterioration may be the trigger of axonal degeneration and, ultimately, neuronal death.[109] In contrast to neuronal loss, which is irreversible, disease-associated synaptic dysfunctions could be rescued through the growth of new terminals and/or dendritic spines.[5] Thus, focusing on early, prodromal dysfunction in PD appears key to design effective therapeutic and preventive strategies, and our study uncovered LRRK2 as a promising disease modifying target of early-stage PD.

Materials and methods

Mouse strains

C57Bl/6J Lrrk2 knock-out (KO) and G2019S-Lrrk2 knock-in mice were provided by Dr. Heather Melrose (Mayo Clinics, Florida, USA). Housing and handling of mice were done in compliance with national guidelines. All animal procedures were approved by the Ethical Committee of the University of Padova and the Italian Ministry of Health (license 1041/2016-PR and 105/2019) and followed the guidelines approved by the Northwestern University Animal Care and Use Committee. Approximately equal numbers of males and females were used for every experiment.

Cell cultures

Generation of LRRK2 KO SH-SY5Y CRISPR/Cas9 edited monoclonal cell line.

The KO of LRRK2 was performed using CRISPR/Cas9-mediated genome editing technology following the protocol by Sharma *et al*.[110] Two sgRNAs were selected among those designed by the laboratory of Zhang F (https://www.genscript.com/gRNA-detail/120892/LRRK2-

CRISPR-guide-RNA.html) and one was designed using the online platform Benchling (https://www.benchling.com). All the gRNAs were synthetized by Sigma-Aldrich. The oligos pairs encoding the 20-nt guide sequence were annealed and ligated into the pSpCas9(BB)-2A-Puro (PX459) V2.0 vector (Addgene, Watertown, MA, US) and then amplified in chemically competent E. coli StbI3 cells (ThermoFisher ScientificTM). Human neuroblastoma-derived SH-SY5Y cells were transfected using Lipofectamine 2000 (Invitrogen) and subjected to puromycin selection. The selected cells were diluted to obtain monoclonal cell lines. Approximately one week after plating, the colonies were inspected for a clonal appearance and progressively expanded. Finally, the deletion of LRRK2 was verified in multiple lines by western blot analysis with LRRK2 specific antibody (Fig. S5).

SH-SY5Y cell line maintenance, differentiation and treatments

SH-SY5Y cells (naïve, LRRK2 KO, expressing GFP or GFP-LRRK2 wild type) were cultured in a mixture (1:1) of Dulbecco's Modified Eagle's Medium (DMEM, Biowest) and Ham's F-12 Nutrient Mixture (F12, Biowest), supplemented with 10% (v/v) Fetal Bovine Serum (FBS, Thermo Fisher Scientific) and 1% Penicillin/Streptomycin solution (PS, GIBCO Life Technologies).

To promote N-type (neuronal-like cells) cell differentiation, cells were plated in DMEM/F12 containing 1% PS, 1% FBS and 10μM of all-trans-retinoic acid (RA, Sigma-Aldrich). At regular intervals of 48 hours, RA was newly provided to the medium. Cells were differentiated for 6 days and then subjected to the treatments.

SH-SY5Y cells were starved for 5 hours in DMEM/F12 supplemented with 1% PS and then stimulated with 100ng/mL BDNF (50240-MNAS, Sino Biological) for different time periods (5, 15, 30 and 60 minutes) in serum-free medium. LRRK2 kinase activity was inhibited pre-treating cells with 0.5μM MLi-2 (ab254528, Abcam) for 90 minutes. For the detection of phospho-S935 and phospho-Rabs, 80 µg of total protein were loaded, whereas 30 µg were used for the detection of phospho-AKT and phospho-ERK1/2, as well as for total proteins.

HEK293T cell line maintenance and transfection

HEK293T cells (Life technologies) were cultured in DMEM (Biowest) supplemented with 10% (v/v) FBS (Thermo Fisher Scientific) and 1% PS (GIBCO Life Technologies). Cells were plated in 100mm dishes and, once they reached 80% confluency, they were transfected with polyethyleneimine (PEI, Polysciences) at 1:2 DNA to PEI ratio (v/w). Flag-LRRK2 was co-expressed with YFP-drebrin full-length, N-terminal domain (1-256 aa), or C-terminal domain (256-649 aa).

Primary mouse cortical neuron preparation, transfection and treatments

Primary cortical neurons from Lrrk2 WT and KO C57BL/6J mice were derived from postnatal mouse (P0) exploiting the Papain Dissociation System (Worthington Biochemical Corporation). Cortices were incubated in Papain solution (Papain and DNase solution in Earle's Balanced Salt Solution) for 40 minutes at 37°C. Subsequently, tissue was triturated and centrifugated for 5 minutes at 200g. The supernatant was discarded, and the pellet was resuspended in Stop solution [DNase solution and Trypsin inhibitor solution (15,5mg/mL) in Earle's Balanced Salt Solution]. For 10 minutes the tissue was allowed to precipitate and then the supernatant was pipetted drop-by-drop on 5mL of 10/10 solution (10 μg/mL Trypsin inhibitor and 10 μg/mL BSA in Earle's Balanced Salt Solution) and centrifuged for 10 minutes at 100g. The pellet was resuspended in Neurobasal A medium (GIBCO Life Technologies) supplemented with 2% B27 supplement (GIBCO Life Technologies), 0.5 mM L-glutamine (GIBCO Life Technologies), 100 units/mL penicillin, and 100 μg/mL streptomycin (GIBCO Life Technologies). Cells were diluted and counted in 0.4% Trypan blue. Neurons were plated at 1000-1500 cells/mm^2 onto 6-well plates or at 200 cells/mm^2 on 12mm glass coverslips in 24-well plates and maintained in culture for 14 days prior to the treatments. After 7 days, 50% of the Neurobasal medium was removed and replaced with fresh one.

At DIV14, Lrrk2 WT and KO mature neurons cultured in the 6-well plates for western blot analysis were treated with MLi-2 (0.5μM) for 90 minutes and with BDNF (100ng/mL) for 5, 30, 60 and 180 minutes in Neurobasal completed medium. At DIV4, Lrrk2 WT and KO primary neurons plated in 24-well plates were transfected with lipofectamine (Lipofectamine 2000, Invitrogen) in a 1:2 ratio with DNA (v/w). The transfection was carried in Opti-MEM (GIBCO Life Technologies) for 45 minutes. Transfected neurons were then maintained in

culture for additional 9-10 days. At DIV14, neurons were treated for 24 hours with 100ng/mL BDNF to study the spine maturation process.

Media composition for cortical neuron differentiation from hiPSC

Neuronal maintenance media (NMM): 1:1 mixture of N2 media (DMEM/F12 GlutaMax supplemented with 1X N2, 50 units/mg/ml Penicillin/Streptomycin solution, 1X MEM Non-Essential Amino Acids Solution (100X), 100μM 2-Mercaptoethanol (ThermoFisher Scientific), 1mM sodium pyruvate and 5 μg/ml insulin (Merck)) and B27 media (Neurobasal supplemented with 1X B27 and 2mM L-Glutamine (ThermoFisher Scientific).

Neuronal induction media (NIM): NMM media supplemented with 100nM LDN193189 (Sigma Aldrich) and 10μM SB431542 (Bio-Techne).

hiPSC cell culture, cortical neuron differentiation and treatments

Isogenic human induced-pluripotent stem cell lines (hiPSC) of WT and LRRK2 KO, made and kindly provided by Mark R Cookson's laboratory,[68] were cultured using E8 media (Thermo Fisher Scientific) in a 6-well plate coated with Cultrex. Cortical neuronal differentiation was carried out according to Shi, *et al.*[67] Briefly, upon reaching 100% confluency, media was changed to neuronal induction media (NIM) and is marked as 0 days *in vitro* (DIV0). Daily NIM media change was carried out till DIV12, upon which neuronal precursor cells (NPCs) are generated. On DIV12, NPCs were expanded via passage at 1:2 ratio, supplemented with neuronal maintenance media (NMM) media and 20 ng/ml FGF2 (Bio-Techne). Neurogenesis was initiated on DIV18 with the removal of FGF2 and the cells supplemented with NMM only changing every 2 days. At DIV25-30, neurons were cryopreserved in CryoStor CS10 solution (Stem Cell technologies).

For electrophysiological recordings and immunocytochemistry/immunofluorescence (ICC/IF), thawed neurons were plated on Cultrex coated 10mm coverslips at a density of 60k/cm^2. All cells were treated with 1 ug/ml Mitomycin C (Sigma-Aldrich) for 1 hour at 37°C, 48-72 hours after plating to remove any proliferating cells. Neurons were maintained in NMM until used for experiments. BDNF treatment was carried out with cells incubated with 50 ng/ml BDNF (Bio-Techne) for 24 hours prior to use in electrophysiological recordings and ICC/IF.

Electrophysiological recordings

Whole-cell patch-clamp recordings were performed on cortical neurons at DIV70-75 (mentioned as DIV70 from here on) in voltage clamp at Vh −70 mV and the membrane test function was used to determine intrinsic membrane properties ∼1 min after obtaining whole-cell configuration, as described previously.[111] Briefly, coverslip containing neurons were transferred to warner bath (RC-26G) and maintained at 35 ± 2 °C through constant perfusion using in-line heater and controller (Warner Instruments) with extra cellular solution (ECS) containing (in mM unless stated): 145 NaCl, 3 KCl, 2 $MgCl_2$, 2 $CaCl_2$, 10 glucose, 10 HEPES, pH 7.4, 310 mOsm. Tetrodotoxin (TTX 0.2μM), and picrotoxin (PTX 100μM), were added before use to block sodium and $GABA_A$ currents. Pipette resistance (Rp) of glass electrodes was 4–6 MΩ when filled with (in mM): 130 Csmethanesulfonate, 5 CsCl, 4 NaCl, 2 MgCl2, 5 EGTA, 10 HEPES, 5 QX-314, 0.5 GTP, 10 Na2-phosphocreatine, and 5 MgATP, 0.1 spermine, pH 7.2, 300 mOsm. Data was acquired by Multiclamp700B amplifier and digidata 1550B and signals were filtered at 2kHz, digitized at 10 kHz, and analyzed in Clampfit 10 (MolecularDevices). Tolerance for series resistance (Rs) was <35 MΩ and uncompensated; ΔRs tolerance cut-off was <20%. mEPSCs were analyzed using Clampfit10 (threshold 5pA); all events were checked by eye. Data are presented as mean ± s.e.m. where n is cells from a minimum of 3 separate cultures (culture N in brackets).

Cells and tissues lysis, SDS-PAGE and Western blotting analysis

Immortalized cells, neurons and mouse brain tissues were lysed for 30 minutes on ice in appropriate volume of cold RIPA lysis buffer (Cell Signaling Technologies) supplemented with protease inhibitor cocktail. Protein concentration was assessed by performing BCA assay (Thermo Fisher Scientific). Proteins were solubilized in sample buffer (200mM Tris-HCl pH 6.8, 8% SDS, 400mM DTT, 40% glycerol, q.s. Bromophenol Blue) and resolved by SDS-PAGE on 8% Tris polyacrylamide homemade gels in Tris-glycine-SDS running buffer for the AP-MS experiments, and on 4-20% polyacrylamide gels (GenScript® Express Plus PAGE) in Tris-MOPS-SDS running buffer (GenScript® Running Buffer Powder) for the other cases. Proteins were transferred on PVDF (polyvinylidene difluoride, Bio-Rad) membranes using a semi-dry transfer system (Trans-Blot® Turbo Transfer System, Bio-Rad). Non-specific

binding sites were blocked incubating membranes with 5% non-fat dry milk diluted in 0.1% Tween-20 Tris-buffered saline (TBS-T) for 1 hour at room temperature under agitation. Membranes were subsequently incubated overnight at 4°C with primary antibodies in 5% non-fat dry milk in TBS-T or in 5% BSA in TBS-T. After three washes in TBS-T at room temperature, membranes were incubated for 1 hour at room temperature with horseradish peroxidase (HRP)-conjugated goat anti-mouse or anti-rabbit IgG. After three more washes in TBS-T, proteins were visualized using chemiluminescence (Immobilon ECL western HRP substrate, Millipore). Densitometric analysis was carried out using the Image J software. The antibodies used for western blotting are as follows: rabbit α-LRRK2 (MJFF2 c41-2, ab133474, Abcam, 1:300); rabbit α-phospho-Ser935 LRRK2 (ab133450, Abcam, 1:300); mouse α-phospho-Ser473-AKT (sc-293125, Santa Cruz Biotechnology, 1:500); rabbit α-AKT (9272S, Cell Signaling Technology, 1:1000); rabbit α-phospho [(Thr202/Tyr204, Thr185/Tyr187)-ERK1/2 (12-302, Millipore, 1:1500); rabbit α-ERK1/2 (4695, Cell Signaling Technology, 1:1000); mouse α-GAPDH (CSB-MA000195, 1:5000); mouse α-βIII tubulin (T8578, Sigma-Aldrich, 1:40000); mouse α-DREBRIN (MA1-20377, Thermo Fisher Scientific, 1:500); α-Flag M2-HRP (A8592, Sigma-Aldrich, 1:5000); α-mouse IgG-HRP (A9044, Sigma-Aldrich, 1:80000); α-rabbit IgG-HRP (A9169, Sigma-Aldrich, 1:16000).

Staining on mammalian cells and brain tissue

Immunocytochemistry and confocal microscopy of primary neurons

Mouse primary cortical neurons derived from Lrrk2 WT C57BL/6 and KO pups (P0) were fixed using 4% paraformaldehyde (PFA, Sigma-Aldrich) in PBS1X pH 7.4 for 20 minutes at room temperature and, after three washes in PBS1X, they were subjected to staining protocol. Cells were firstly permeabilized with 0.3% Triton-X in PBS1X for 5 minutes and then saturated in blocking buffer [1% Bovine serum Albumin (BSA) Fraction V, 0.1% Triton-X, 50mM Glycine, 2% goat serum in PBS1X] for 1 hour at room temperature in agitation. The primary and secondary antibodies incubation steps were carried out in working solution (20% blocking buffer in PBS1X) for 1 hour at room temperature. Both incubations were followed by three washes in working solution. Nuclei staining was performed in Hoechst 33258 (Invitrogen, 1:10000 dilution in PBS1X) for 5 minutes and followed by three rinses in PBS1X. After been cleaning in distilled water, coverslips were mounted on microscope slides with Mowiol® mounting medium. Immunofluorescence z-stack images (z-stack thickness: 0.5 μm x 6 = 3 μm)

were obtained on Zeiss LSM700 laser scanning confocal microscope exploiting a 63X oil immersion objective.

The antibodies used for immunocytochemistry are as follows: mouse α-PSD95 (ab2723, Abcam, 1:200), rabbit α-MAP2 (sc-20172, Santa Cruz, 1:200); mouse α-drebrin (MA1-20377, Thermo Fisher Scientific, 1:400); goat anti-mouse-Alexa Fluor 568 (A11004, Invitrogen), goat anti-mouse-Alexa Fluor (A11004, Invitrogen), goat anti-rabbit-Alexa Fluor 488 (A11034, Invitrogen), rabbit-Alexa Fluor 568 (A11036, Invitrogen), goat anti-rabbit-Alexa Fluor 405.

Immunostaining and Image Analysis of hiPCS-derived cortical neurons

For immunocytochemistry DIV70 cells on coverslips were fixed in 4% paraformaldehyde (PFA) for 10 minutes. Cells were then permeabilized and blocked using 0.4% Tween with 10% normal goat serum (NGS), in PBS1X for 1 hour. Primary antibodies were incubated overnight at 4°C in PBST plus 10% NGS. Cells were washed 3× with PBST before 1 hour incubation at room temperature with secondary antibodies (α-guinea pig Alexa-488, α-rabbit Alexa 568, α-chicken 647 along with DAPI) from Molecular probes and Jackson Laboratories). Primary antibodies were chicken anti-microtubule associated protein 2 (MAP2) (Antibodies A85363, dil. 1:1000), rabbit anti-homer 1 (Synaptic Systems 160003, dil. 1:500), and guinea pig anti-bassoon (Synaptic Systems 141004, dil. 1:500). Coverslips were slide mounted with FluorSave (Sigma-Aldrich) and all images were acquired on Leica confocal microscope as 0.45 μm z-stacks at 60× magnification (flattened with the max projection function for cluster analysis). Single 60× images were used for analyses. Excitation and acquisition parameters were constrained across all experimental paired (culture) acquisitions. All images were analyzed using IMARIS microscopy image analysis software (v10.0.0 Oxford Instruments). Data are presented as mean ± s.e.m.

Golgi-Cox staining

Animals were terminally anesthetized and transcardially perfused with 0.9% saline. Half of each brain was incubated with Golgi-Cox solution (Potassium dichromate, Mercuric chloride, Potassium chromate prepared according to Zaqout *et al.*[112]) in the dark at room temperature for 14 days and then transferred in 30% sucrose solution in PBS1X. Brains were cut with a vibratome in 100 μm thick slices. The sections were then blotted by pressing an absorbent paper

moistened with sucrose solution onto the slides and dried for 7-10 minutes. The samples were then subjected to color development procedure consisting in the following steps: 1. two 5-minutes washes in distilled water; 2. 5-minutes dehydration step in 50% ethanol; 3. 10-minutes incubation step in 20% ammonium hydroxide; 4. 5-minutes wash in distilled H_2O; 5. 8-minutes incubation step in 5% sodium thiosulfate at room temperature in the dark; 6. two additional 1-minute rinses in distilled H_2O; 7. dehydration in ascending grades of ethanol (70%, 95% and 100%). After two final 6-minutes incubations with xylene, the slides were covered with Eukitt® mounting medium. Z-stack images (z-stack thickness between 0.5 μm x 30 = 15 μm and 0.5 μm x 80 = 40 μm) were acquired with Zeiss LSM700 laser scanning confocal microscope using 100X/1,40 Oil DIC M27 immersion objective with phase contrast acquisition mode. All images were analyzed using freely available RECONSTRUCT software and according to Risher et al.[113]. 3-4 neurons per mice (n=3 per genotype) were acquired in the dorsal striatum and n≅4 segments per neurons were analyzed.

Electron Microscopy

Coronal brain slices were fixed with 2.5% glutaraldehyde in 0.1M sodium cacodylate pH 7.4 buffer overnight at 4° C, postfixed in 1% osmium tetroxide for 1h at 4° C, ethanol dehydrated, then infiltrated in a mixture of EMbed 812 (Electron Microscopy Sciences) epoxy resin and absolute ethanol 1:1, and then finally embedded in EMbed 812 (Electron Microscopy Sciences) epoxy resin. Ultrathin sections were obtained with a Leica EM UC7 ultramicrotome and subsequently counterstained with uranyl acetate and lead citrate. Samples were then examined with a Tecnai G2 (Fei-Thermo Fisher) transmission electron microscope operating at 120 kV, digital images were acquired using a Veleta (Olympus Soft Imaging Solutions) digital camera. Images were acquired at the level of dorsal striatum in n=4 mice per genotype (1-month old mice) or n=3 mice per genotype (18-months old mice). n≅20 synapses per mice were analyzed in 1-month old mice; n≅40 synapses per mice were analyzed in 18-months old mice. We gratefully thank the DeBio Imaging facility at the University of Padova for their support with the staining procedure.

Quantitative PCR

Total RNA was isolated from midbrain, striatum, and cortex of 1 month-old animals (n=6 animals per group, Lrrk2 WT and Lrrk2 KO) using Animal Tissue RNA Purification Kit (Norgen), according to manufacturer's instruction. After extraction, RNA concentration was quantified with NanoDrop 2000C spectrophotometer (Thermo Fisher Scientific). After DNAse treatment (Thermo Fisher Scientific, according to manufacturer's protocol), complementary DNA (cDNA) was generated using qRT SuperMix (Bimake). The cDNAs were used for quantitative PCR (qPCR) exploiting iTaq Universal SYBR® Green Supermix and CFX96 RealTime System (BioRad) for 40 cycles. All samples were run in triplicated, and transcripts levels normalized against the geometrical means of RPL27, actin, and GAPDH relative abundance (housekeeping genes). Data shown were produced using Bio-Rad CFX Manager software and analyzed according to ddCt algorithm.

Primers (5'-3') used are:

Bdnf FW: GGCTGACACTTTTGAGCACGTC

Bdnf REV: CTCCAAAGGCACTTGACTGCTG

TrkB FW: TGAGGAGGACACAGGATGTTGA

Trkb REV: TTCCAGTGCAAGCCAGTATCTG

Shank3 FW: ACCTTGAGTCTGTAGATGTGGAAG

Shank3 REV: GCTTGTGTCCAACCTTCACGAC

Psd95 FW: GGTGACGACCCATCCATCTTTATC

Psd95 REV: CGGACATCCACTTCATTGACAAAC

Arpc2 FW: GAGTCACAGTAGTCTTCAGCACG

Arpc2 REV: AGGTTCCCTGTGGCTGAAAAGG

Dbn1 FW: AGAAGTCGGAGTCAGAGGTGGA

Dbn1 REV: ATGCCACTCGTTCCTGCTGTCT

Darpp32 FW: AGATTCAGTTCTCTGTGCCCG

Darpp32 REV: GGTTCTCTGATGTGGAGAGGC

housekeeping genes

Rpl27 FW: AAGCCGTCATCGTGAAGAACA

Rpl27 REV: CTTGATCTTGGATCGCTTGGC

GAPDH FW: AGGTCGGTGTGAACGGAT TTG

GAPDH REV: TGTAGACCATGTAGTTGAGGTCA

ACTB FW: CAACGGCTCCGGCATGTG

ACTB REV: CTCTTGCTCTGGGCCTCG

Protein purification from mammalian cells

For the affinity purification (AP) protocol, SH-SY5Y GFP and SH-SY5Y GFP-LRRK2 cells were plated onto 100mm dishes and differentiated for 6 days. SH-SY5Y GFP-LRRK2 cells were treated with BDNF (100ng/mL) or with an equal volume of vehicle for 15 minutes, while SH-SY5Y OE-GFP cells were left untreated. Cells were harvested in lysis buffer (20mM Tris-HCl pH 7.5, 150mM NaCl, 1mM EDTA, 1% Tween 20, 2.5mM sodium pyrophosphate, 1mM β-glycerophosphate, 1mM sodium orthovanadate) supplemented with protease inhibitor cocktail (Sigma-Aldrich). Lysates were incubated end-over-end with GFP-TrapA beads (ChromoTek) overnight at 4°C. Immunocomplexes were subsequently washed ten times with buffers containing a decreasing amount of NaCl and Tween20 and incubated with sample buffer 2X. The eluted proteins were resolved by SDS-PAGE and processed for western blot analysis or Mass Spectrometry (MS).

HEK293T cells were transfected as described above and drebrin constructs were immunoprecipitated with GFP-trap resin following the same protocol used for the SH-SY5Y cells. The eluded samples were resolved by SDS-PAGE and processed for western blot analysis to visualize the co-precipidated LRRK2 using Flag antibodies.

LC-MS/MS and data analysis

Three biological replicates of GFP-trap purifications of SH-SY5Y cells were processed for the proteomics experiments. Gels were stained with Colloidal Coomassie Brilliant Blue (0.25% Brilliant Blue R-250, 40% ethanol, 10% acetic acid in milli-Q water) for at least 1 hour and

then rinsed in destaining solution (10% isopropanol, 10% acetic acid in milli-Q water). Gel slices were cut into small pieces and subjected to reduction with dithiothreitol (DTT 10 mM in 50 mM NH4HCO3, for 1 h at 56 °C), alkylation with iodoacetamide (55 mM in 50 mM NH4HCO3, for 45 min at RT and in the dark), and in-gel digestion with sequencing grade modified trypsin (Promega, 12.5 ng/µL in 50 mM NH4HCO3). Samples were analyzed with a LTQ-Orbitrap XL mass spectrometer (Thermo Fisher Scientific) coupled to a HPLC UltiMate 3000 (Dionex – Thermo Fisher Scientific) through a nanospray interface. Peptides were separated at a flow rate of 250 nL/min using an 11-cm-long capillary column (PicoFrit, 75-µm ID, 15-µm tip, New Objective) packed in house with C18 material (Aeris Peptide 3.6 µm XB C18; Phenomenex). A linear gradient of acetonitrile/0.1% formic acid from 3 to 40% was used for peptide separation and the instrument operated in a data dependent acquisition mode with a Top4 method (one full MS scan at 60,000 resolution in the Orbitrap, followed by the acquisition in the linear ion trap of the MS/MS spectra of the four most intense ions). Two technical replicates were acquired for each biological replicate. Raw data files were analyzed using MaxQuant and Andromeda software package[114] and searched against the human section of the UniProt database (version September 2020, 75093 entries) concatenated with a database of common contaminants found in proteomic experiments. Trypsin was selected as digesting enzyme with up to two missed cleavages allowed, carbamidomethylation of cysteine residues was set as a fixed modification and methionine oxidation as a variable modification. A search against a randomized database was used to assess the false discovery rate (FDR), and data were filtered to remove contaminants and reverse sequences and keep into account only proteins identified with at least two peptides and a FDR ≤ 0.01, both at peptide and protein level.

Intensity values were retrieved for 350 proteins. A single step of data QC was applied to the intensity values: proteins whose intensity value was 0.00 for both the 2 technical replicates in at least 1 of the 3 pull-down experiments performed with LRRK2-GFP were removed. This is a stringent QC step, but it was implemented to remove proteins with no triplicate data available for the LRRK2 pull-down as this could have occurred because of genuine low levels of pull-down (values close to the limit of detection, missing not at random values) or because of technical issues with the single sample (loss/poor quality of sample, missing at random values). QC reduced the number of proteins in the analysis from 350 to 269 entries.

Intensity data were then processed according to Aguilan et al.[115] Briefly, intensity data were log2 transformed and normalized to the median of all proteins within the same experiment. Missing data (0.00 intensity values) were considered missing not at random thus they were

imputed via probabilistic minimum imputation (random values within less than 2.5 standard deviations from the distribution of intensity values per the single experiment, 0.3 max variability).[115,116]

Finally, fold change (GFP-LRRK2 vs GFP and GFP-LRRK2 BDNF treated vs GFP-LRRK2 untreated) and associated p-value (two-tailed, paired t-test) were calculated and visualized.

Phosphoproteomics analysis

Protein processing for MS

Striata from 8 weeks mice were dissected and rapidly homogenized in four volumes of ice-cold Buffer A (0.32 M sucrose, 5 mM HEPES, pH7.4, 1 mM $MgCl_2$, 0.5 mM $CaCl_2$) supplemented with Halt protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific) using a Teflon homogenizer (12 strokes). Homogenized brain extract was centrifuged at 1400g for 10 minutes. Supernatant (S1) was saved and pellet (P1) was homogenized in buffer A with a Teflon homogenizer (five strokes). After centrifugation at 700 g for 10 minutes, the supernatant (S1') was pooled with S1. Pooled S1 and S1' were centrifuged at 13,800 g for 10 minutes to the crude synaptosomal pellet (P2). The crude synaptosomal pellet (P2) was homogenized in buffer B (0.32M sucrose, 6mM Tris, pH 8.0) supplemented with protease and phosphatase inhibitors cocktail with a Teflon homogenizer (five strokes) and was carefully loaded onto a discontinuous sucrose gradient (0.8 M/1 M/1.2 M sucrose solution in 6 mM Tris, pH 8.0) with a Pasteur pippete, followed by centrifugation in a swinging bucket rotor for 2 hours at 82,500g. The synaptic plasma membrane fraction (SPM) in the interphase between 1 M and 1.2 M sucrose fractions was collected using a syringe and transferred to clean ultracentrifuge tubes. 6 mM Tris buffer was added to each sample to adjust the sucrose concentration from 1.2 M to 0.32 M and the samples were centrifuged in a swinging bucket rotor at 200,000g for 30 minutes. The supernatant was removed and discarded. Added to the pellets the lysis solution containing 12 mM sodium deoxycholate, 12 mM sodium lauroyl sarcosinate, 10 mM TCEP, 40 mM CAA, and phosphatase inhibitor cocktail (Millipore-Sigma) in 50 mM Tris·HCl, pH 8.5, and incubated 10 min at 95°C with vigorous shaking. Sonicated the pellets several times with a sonicator probe and boiled again for 5 minutes. Centrifuged at 16,000g for 10 minutes to remove the debris and collected supernatant. The samples were diluted fivefold with 50 mM triethylammonium bicarbonate and analyzed by BCA to determine protein concentration. The

samples were then normalized to 300µg protein in each and digested with 6µg Lys-C (Wako) for 3 hours at 37°C. 6µg trypsin was added for overnight digestion at 37°C. The supernatants were collected and acidified with trifluoroacetic acid (TFA) to a final concentration of 1% TFA. Ethyl acetate solution was added at 1:1 ratio to the samples. The mixture was vortexed for 2 minutes and then centrifuged at 16,000g for 2 minutes to obtain aqueous and organic phases. The organic phase (top layer) was removed, and the aqueous phase was collected, dried completely in a vacuum centrifuge, and desalted using Top-Tip C18 tips (Glygen) according to manufacturer's instructions. The samples were dried completely in a vacuum centrifuge and subjected to phosphopeptide enrichment using PolyMAC Phosphopeptide Enrichment kit (Tymora Analytical) according to manufacturer's instructions, and the eluted phosphopeptides dried completely in a vacuum centrifuge.

LC-MS/MS Analysis

The full phosphopeptide sample was dissolved in 10.5 µL of 0.05% trifluoroacetic acid with 3% (vol/vol) acetonitrile and 10 µL of each sample was injected into an Ultimate 3000 nano UHPLC system (Thermo Fisher Scientific). Peptides were captured on a 2-cm Acclaim PepMap trap column and separated on a 50-cm column packed with ReproSil Saphir 1.8 µm C18 beads (Dr. Maisch GmbH). The mobile phase buffer consisted of 0.1% formic acid in ultrapure water (buffer A) with an eluting buffer of 0.1% formic acid in 80% (vol/vol) acetonitrile (buffer B) run with a linear 90-min gradient of 6–30% buffer B at flow rate of 300 nL/min. The UHPLC was coupled online with a Q-Exactive HF-X mass spectrometer (Thermo Fisher Scientific). The mass spectrometer was operated in the data-dependent mode, in which a full-scan MS (from m/z 375 to 1,500 with the resolution of 60,000) was followed by MS/MS of the 15 most intense ions (30,000 resolution; normalized collision energy - 28%; automatic gain control target (AGC) - 2E4, maximum injection time - 200 ms; 60sec exclusion].

LC-MS Data Processing

The raw files were searched directly against the mouse database with no redundant entries, using Byonic (Protein Metrics) and Sequest search engines loaded into Proteome Discoverer 2.3 software (Thermo Fisher Scientific). MS1 precursor mass tolerance was set at 10 ppm, and MS2 tolerance was set at 20ppm. Search criteria included a static carbamidomethylation of

cysteines (+57.0214 Da), and variable modifications of phosphorylation of S, T and Y residues (+79.996 Da), oxidation (+15.9949 Da) on methionine residues and acetylation (+42.011 Da) at N terminus of proteins. Search was performed with full trypsin/P digestion and allowed a maximum of two missed cleavages on the peptides analyzed from the sequence database. The false-discovery rates of proteins and peptides were set at 0.01. All protein and peptide identifications were grouped and any redundant entries were removed. Only unique peptides and unique master proteins were reported.

Label-free Quantitation Analysis

All data were quantified using the label-free quantitation node of Precursor Ions Quantifier through the Proteome Discoverer v2.3 (Thermo Fisher Scientific). For the quantification of phosphoproteomic data, the intensities of phosphopeptides were extracted with initial precursor mass tolerance set at 10 ppm, minimum number of isotope peaks as 2, maximum ΔRT of isotope pattern multiplets – 0.2 min, PSM confidence FDR of 0.01, with hypothesis test of ANOVA, maximum RT shift of 5 min, pairwise ratio-based ratio calculation, and 100 as the maximum allowed fold change. For calculations of fold-change between the groups of proteins, total phosphoprotein abundance values were added together, and the ratios of these sums were used to compare proteins within different samples.

Bioinformatics

The LRRK2 interactome was constructed following the procedure reported in *Zhao et al.*[117] Briefly, protein interactions reported for LRRK2 in peer-reviewed literature were considered if the interaction was replicated in at least 2 different papers or identified with 2 different interaction detection methods. The obtained LRRK2 protein interaction network was analysed in Zhao *et al.*[51] to identify topological clusters using the FastGreed R Package (based on degree centrality). Here we report one of the identified clusters that is enriched for synaptic functions. Functional enrichment was performed using g:Profiler, g:GOSt (https://biit.cs.ut.ee/gprofiler/gost)[118] with Gene Ontology Biological Processes (GO-BPs) and SynGO (https://www.syngoportal.org/).

Statistical analysis

Statistical analyses and plotting were performed with GraphPad-Prism9. Unpaired, two-tailed t-test or Shapiro-Wilk test were used for experiments comparing two groups; one-way ANOVA was used for experiments comparing three or more groups; two-way ANOVA was used when multiple groups and two factors (i.e., genotype and treatment) were to be compared. Šídák's multiple comparisons test was used when determining statistical significance for comparisons between groups.

Data availability

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary material.

Protein composition of the LRRK2 cluster enriched for synaptic functions can be obtained from Zhao *et al*.[117]

Funding

This work was supported by the University of Padova [STARS Grants, LRRKing-Role of the Parkinson's disease kinase LRRK2 in shaping neurites and synapses, EG), the Michael J. Fox Foundation for Parkinson's Research-LRRK2 challenge (EG, LP and CM), NIH R01 NS097901 (LP), UKRI future Leader Fellowship funding MR/T041129/1 (DBK).

Acknowledgments

We are very grateful to Dr. Mark R Cookson and Alexandra Beilina (NIA, NIH, USA) for providing WT and LRRK2 KO human iPSC lines used for electrophysiological measurements.

Competing interests

The authors report no competing interests.

References

1. Fiala JC, Spacek J, Harris KM. Dendritic spine pathology: Cause or consequence of neurological disorders? *Brain Res Rev*. 2002;39(1):29-54. doi:10.1016/S0165-0173(02)00158-3

2. Hong S, Beja-Glasser VF, Nfonoyim BM, et al. Complement and microglia mediate early synapse loss in Alzheimer mouse models. *Science (1979)*. 2016;352(6286):712-716. doi:10.1126/science.aad8373

3. Milnerwood AJ, Raymond LA. Early synaptic pathophysiology in neurodegeneration: Insights from Huntington's disease. *Trends Neurosci*. 2010;33(11):513-523. doi:10.1016/j.tins.2010.08.002

4. Soukup S, Vanhauwaert R, Verstreken P. Parkinson's disease: convergence on synaptic homeostasis. *EMBO J*. 2018;37(18). doi:10.15252/embj.201898960

5. Picconi B, Piccoli G, Calabresi P. Synaptic dysfunction in Parkinson's disease. *Adv Exp Med Biol*. 2012;970:553-572. doi:10.1007/978-3-7091-0932-8_24/FIGURES/1

6. Kordower JH, Olanow CW, Dodiya HB, et al. Disease duration and the integrity of the nigrostriatal system in Parkinson's disease. *Brain*. 2013;136(8):2419-2431. doi:10.1093/brain/awt192

7. Bellucci A, Antonini A, Pizzi M, Spano PF. The end is the beginning: Parkinson's disease in the light of brain imaging. *Front Aging Neurosci*. 2017;9(OCT). doi:10.3389/fnagi.2017.00330

8. Chidambaram SB, Rathipriya AG, Bolla SR, et al. Dendritic spines: Revisiting the physiological role. *Prog Neuropsychopharmacol Biol Psychiatry*. 2019;92:161-193. doi:10.1016/j.pnpbp.2019.01.005

9.   González-Rodríguez P, Zampese E, Stout KA, et al. Disruption of mitochondrial complex I induces progressive parkinsonism. *Nature*. 2021;599(7886):650-656. doi:10.1038/s41586-021-04059-0

10.   Cookson MR. LRRK2 Pathways Leading to Neurodegeneration. *Curr Neurol Neurosci Rep*. 2015;15(7). doi:10.1007/s11910-015-0564-y

11.   Greggio E, Cookson MR. Leucine-rich repeat kinase 2 mutations and Parkinson's disease: Three questions. *ASN Neuro*. 2009;1(1):13-24. doi:10.1042/AN20090007

12.   Marchand A, Drouyer M, Sarchione A, Chartier-Harlin MC, Taymans JM. LRRK2 Phosphorylation, More Than an Epiphenomenon. *Front Neurosci*. 2020;14. doi:10.3389/fnins.2020.00527

13.   Wallings R, Manzoni C, Bandopadhyay R. Cellular processes associated with LRRK2 function and dysfunction. *FEBS Journal*. 2015;282(15):2806-2826. doi:10.1111/febs.13305

14.   Alessi DR, Pfeffer SR. Leucine-Rich Repeat Kinases. *Annu Rev Biochem*. 2024;93(1):261-287. doi:10.1146/ANNUREV-BIOCHEM-030122-051144

15.   Civiero L, Cogo S, Biosa A, Greggio E. The role of LRRK2 in cytoskeletal dynamics. *Biochem Soc Trans*. 2018;46(6):1653-1663. doi:10.1042/BST20180469

16.   Iannotta L, Greggio E. LRRK2 signaling in neurodegeneration: two decades of progress. Halliday G, Dzamko N, eds. *Essays Biochem*. 2021;65(7):859-872. doi:10.1042/ebc20210013

17.   Nichols RJ, Dzamko N, Morrice NA, et al. 14-3-3 Binding to LRRK2 is disrupted by multiple Parkinson's disease-associated mutations and regulates cytoplasmic localization. *Biochemical Journal*. 2010;430(3):393-404. doi:10.1042/BJ20100483

18. Dzamko N, Inesta-Vaquera F, Zhang J, et al. The IkappaB kinase family phosphorylates the Parkinson's disease kinase LRRK2 at Ser935 and Ser910 during Toll-Like Receptor signaling. *PLoS One*. 2012;7(6). doi:10.1371/journal.pone.0039132

19. Muda K, Bertinetti D, Gesellchen F, et al. Parkinson-related LRRK2 mutation R1441C/G/H impairs PKA phosphorylation of LRRK2 and disrupts its interaction with 14-3-3. *Proc Natl Acad Sci U S A*. 2014;111(1). doi:10.1073/pnas.1312701111

20. Chia R, Haddock S, Beilina A, et al. Phosphorylation of LRRK2 by casein kinase 1α regulates trans-Golgi clustering via differential interaction with ARHGEF7. *Nat Commun*. 2014;5. doi:10.1038/ncomms6827

21. Dzamko N, Deak M, Hentati F, et al. Inhibition of LRRK2 kinase activity leads to dephosphorylation of Ser 910/Ser935, disruption of 14-3-3 binding and altered cytoplasmic localization. *Biochemical Journal*. 2010;430(3):405-413. doi:10.1042/BJ20100784

22. Nazish I, Arber C, Piers TM, et al. Abrogation of LRRK2 dependent Rab10 phosphorylation with TLR4 activation and alterations in evoked cytokine release in immune cells. *Neurochem Int*. 2021;147. doi:10.1016/j.neuint.2021.105070

23. Maas JW, Yang J, Edwards RH. Endogenous Leucine-Rich Repeat Kinase 2 Slows Synaptic Vesicle Recycling in Striatal Neurons. *Front Synaptic Neurosci*. 2017;9(FEB). doi:10.3389/FNSYN.2017.00005

24. Cirnaru MD, Marte A, Belluzzi E, et al. LRRK2 kinase activity regulates synaptic vesicle trafficking and neurotransmitter release through modulation of LRRK2 macro-molecular complex. *Front Mol Neurosci*. 2014;7(MAY). doi:10.3389/FNMOL.2014.00049

25. Pischedda F, Piccoli G. LRRK2 at the pre-synaptic site: A 16-years perspective. *J Neurochem*. 2021;157(2):297-311. doi:10.1111/jnc.15240

26. Belluzzi E, Gonnelli A, Cirnaru MD, et al. LRRK2 phosphorylates pre-synaptic N-ethylmaleimide sensitive fusion (NSF) protein enhancing its ATPase activity and SNARE complex disassembling rate. *Mol Neurodegener*. 2016;11(1):1-16. doi:10.1186/S13024-015-0066-Z/FIGURES/6

27. Marte A, Russo I, Rebosio C, et al. Leucine-rich repeat kinase 2 phosphorylation on synapsin I regulates glutamate release at pre-synaptic sites. *J Neurochem*. 2019;150(3):264-281. doi:10.1111/jnc.14778

28. Soukup SF, Kuenen S, Vanhauwaert R, et al. A LRRK2-Dependent EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Terminals. *Neuron*. 2016;92(4):829-844. doi:10.1016/J.NEURON.2016.09.037

29. Cao M, Wu Y, Ashrafi G, et al. Parkinson Sac Domain Mutation in Synaptojanin 1 Impairs Clathrin Uncoating at Synapses and Triggers Dystrophic Changes in Dopaminergic Axons. *Neuron*. 2017;93(4):882-896.e5. doi:10.1016/J.NEURON.2017.01.019

30. Nguyen M, Krainc D. LRRK2 phosphorylation of auxilin mediates synaptic defects in dopaminergic neurons from patients with Parkinson's disease. *Proc Natl Acad Sci U S A*. 2018;115(21):5576-5581. doi:10.1073/PNAS.1717590115/SUPPL_FILE/PNAS.1717590115.SD02.XLSX

31. Rassu M, Del Giudice MG, Sanna S, et al. Role of LRRK2 in the regulation of dopamine receptor trafficking. *PLoS One*. 2017;12(6). doi:10.1371/journal.pone.0179082

32. Gupta S, Tielemans A, Guevara CA, Huntley GW, Benson DL. Parkinson's-linked LRRK2-G2019S derails AMPAR trafficking, mobility, and composition in striatum with cell-type and subunit specificity. *Proc Natl Acad Sci U S A*. 2024;121(28). doi:10.1073/PNAS.2317833121

33. Chen C, Soto G, Dumrongprechachan V, et al. Pathway-specific dysregulation of striatal excitatory synapses by LRRK2 mutations. *Elife*. 2020;9:1-26. doi:10.7554/eLife.58997

34. Matikainen-Ankney BA, Kezunovic N, Mesias RE, et al. Altered development of synapse structure and function in striatum caused by Parkinson's disease-linked LRRK2-G2019S mutation. *Journal of Neuroscience*. 2016;36(27):7128-7141. doi:10.1523/JNEUROSCI.3314-15.2016

35. Parisiadou L, Yu J, Sgobio C, et al. LRRK2 regulates synaptogenesis and dopamine receptor activation through modulation of PKA activity. *Nat Neurosci*. 2014;17(3):367-376. doi:10.1038/NN.3636

36. Thoenen H. Neurotrophins and neuronal plasticity. *Science (1979)*. 1995;270(5236):593-598. doi:10.1126/science.270.5236.593

37. Baydyuk M, Xu B. BDNF signaling and survival of striatal neurons. *Front Cell Neurosci*. 2014;8(AUG). doi:10.3389/fncel.2014.00254

38. Iannotta L, Biosa A, Kluss JH, et al. Divergent Effects of G2019S and R1441C LRRK2 Mutations on LRRK2 and Rab10 Phosphorylations in Mouse Tissues. *Cells*. 2020;9(11). doi:10.3390/cells9112344

39. Skelton PD, Tokars V, Parisiadou L. LRRK2 at Striatal Synapses: Cell-Type Specificity and Mechanistic Insights. *Cells*. 2022;11(1). doi:10.3390/cells11010169

40. Piccoli G, Condliffe SB, Bauer M, et al. LRRK2 controls synaptic vesicle storage and mobilization within the recycling pool. *Journal of Neuroscience*. 2011;31(6):2225-2237. doi:10.1523/JNEUROSCI.3730-10.2011

41. Kluss JH, Lewis PA, Greggio E. Leucine-rich repeat kinase 2 (LRRK2): an update on the potential therapeutic target for Parkinson's disease. *Expert Opin Ther Targets*. 2022;26(6):537-546. doi:10.1080/14728222.2022.2082937

42. Encinas M, Iglesias M, Liu Y, et al. Sequential treatment of SH-SY5Y cells with retinoic acid and brain-derived neurotrophic factor gives rise to fully differentiated, neurotrophic factor-dependent, human neuron-like cells. *J Neurochem*. 2000;75(3):991-1003. doi:10.1046/j.1471-4159.2000.0750991.x

43. Numakawa T, Suzuki S, Kumamaru E, Adachi N, Richards M, Kunugi H. BDNF function and intracellular signaling in neurons. *Histol Histopathol*. 2010;25(2):237-258. doi:10.14670/HH-25.237

44. Nirujogi RS, Tonelli F, Taylor M, et al. Development of a multiplexed targeted mass spectrometry assay for LRRK2-phosphorylated Rabs and Ser910/Ser935 biomarker sites. *Biochem J*. 2021;478(2):299-326. doi:10.1042/BCJ20200930

45. Manzoni C, Denny P, Lovering RC, Lewis PA. Computational analysis of the LRRK2 interactome. *PeerJ*. 2015;2015(2). doi:10.7717/peerj.778

46. Tomkins JE, Dihanich S, Beilina A, et al. Comparative Protein Interaction Network Analysis Identifies Shared and Distinct Functions for the Human ROCO Proteins. *Proteomics*. 2018;18(10). doi:10.1002/pmic.201700444

47. Koopmans F, van Nierop P, Andres-Alonso M, et al. SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse. *Neuron*. 2019;103(2):217-234.e4. doi:10.1016/j.neuron.2019.05.002

48. Koganezawa N, Hanamura K, Sekino Y, Shirao T. The role of drebrin in dendritic spines. *Molecular and Cellular Neuroscience*. 2017;84:85-92. doi:10.1016/j.mcn.2017.01.004

49. Shirao T, Hanamura K, Koganezawa N, Ishizuka Y, Yamazaki H, Sekino Y. The role of drebrin in neurons. *J Neurochem*. 2017;141(6):819-834. doi:10.1111/JNC.13988

50. Martin ER, Gandawijaya J, Oguro-Ando A. A novel method for generating glutamatergic SH-SY5Y neuron-like cells utilizing B-27 supplement. *Front Pharmacol*. 2022;13. doi:10.3389/FPHAR.2022.943627

51. Zhao Y, Harvey K, Escott-Price V, Lewis PA, Manzoni C. Transcriptomics analyses of the LRRK2 protein interactome reveal distinct molecular signatures for sporadic and LRRK2 Parkinson's Disease. *bioRxiv*. Published online September 13, 2023:2023.09.12.557373. doi:10.1101/2023.09.12.557373

52. Spirin V, Mirny LA. Protein complexes and functional modules in molecular networks. *Proc Natl Acad Sci U S A*. 2003;100(21):12123-12128. doi:10.1073/PNAS.2032324100

53. Meixner A, Boldt K, Van Troys M, et al. A QUICK screen for Lrrk2 interaction partners--leucine-rich repeat kinase 2 is involved in actin cytoskeleton dynamics. *Mol Cell Proteomics*. 2011;10(1). doi:10.1074/MCP.M110.001172

54. Liu Z, Lee J, Krummey S, Lu W, Cai H, Lenardo MJ. The kinase LRRK2 is a regulator of the transcription factor NFAT that modulates the severity of inflammatory bowel disease. *Nat Immunol*. 2011;12(11):1063-1070. doi:10.1038/NI.2113

55. Chen C, Masotti M, Shepard N, et al. LRRK2 mediates haloperidol-induced changes in indirect pathway striatal projection neurons. *Mol Psychiatry*. Published online April 23, 2025. doi:10.1038/S41380-025-03030-Z

56. Cheng J, Novati G, Pan J, et al. Accurate proteome-wide missense variant effect prediction with AlphaMissense. *Science*. 2023;381(6664). doi:10.1126/SCIENCE.ADG7492

57. Kellner Y, Gödecke N, Dierkes T, Thieme N, Zagrebelsky M, Korte M. The BDNF effects on dendritic spines of mature hippocampal neurons depend on neuronal activity. *Front Synaptic Neurosci*. 2014;6(MAR). doi:10.3389/fnsyn.2014.00005

58. Zagrebelsky M, Korte M. Form follows function: BDNF and its involvement in sculpting the function and structure of synapses. *Neuropharmacology*. 2014;76(PART C):628-638. doi:10.1016/j.neuropharm.2013.05.029

59. Lai KO, Wong ASL, Cheung MC, et al. TrkB phosphorylation by Cdk5 is required for activity-dependent structural plasticity and spatial memory. *Nat Neurosci*. 2012;15(11):1506-1515. doi:10.1038/nn.3237

60. Ji Y, Pang PT, Feng L, Lu B. Cyclic AMP controls BDNF-induced TrkB phosphorylation and dendritic spine formation in mature hippocampal neurons. *Nat Neurosci*. 2005;8(2):164-172. doi:10.1038/nn1381

61. Lendvai B, Stern EA, Chen B, Svoboda K. Experience-dependent plasticity of dendritic spines in the developing rat barrel cortex in vivo. *Nature 2000 404:6780*. 2000;404(6780):876-881. doi:10.1038/35009107

62. Koleske AJ. Molecular mechanisms of dendrite stability. *Nat Rev Neurosci*. 2013;14(8):536-550. doi:10.1038/nrn3486

63. Surmeier DJ, Obeso JA, Halliday GM. Selective neuronal vulnerability in Parkinson disease. *Nat Rev Neurosci*. 2017;18(2):101-113. doi:10.1038/nrn.2016.178

64. Westerlund M, Belin AC, Anvret A, Bickford P, Olson L, Galter D. Developmental regulation of leucine-rich repeat kinase 1 and 2 expression in the brain and other rodent and human organs: Implications for Parkinson's disease. *Neuroscience*. 2008;152(2):429-436. doi:10.1016/j.neuroscience.2007.10.062

65. Tong Y, Pisani A, Martella G, et al. R1441C mutation in LRRK2 impairs dopaminergic neurotransmission in mice. *Proc Natl Acad Sci U S A*. 2009;106(34):14622-14627. doi:10.1073/pnas.0906334106

66. Volta M, Beccano-Kelly DA, Paschall SA, et al. Initial elevations in glutamate and dopamine neurotransmission decline with age, as does exploratory behavior, in LRRK2 G2019S knock-in mice. *Elife*. 2017;6. doi:10.7554/ELIFE.28377

67. Shi Y, Kirwan P, Livesey FJ. Directed differentiation of human pluripotent stem cells to cerebral cortex neurons and neural networks. *Nature Protocols 2012 7:10*. 2012;7(10):1836-1846. doi:10.1038/nprot.2012.116

68. Beylina A, Langston RG, Rosen D, Reed X, Cookson MR. Generation of fourteen isogenic cell lines for Parkinson's disease-associated leucine-rich repeat kinase (LRRK2). *Stem Cell Res*. 2021;53:102354. doi:10.1016/J.SCR.2021.102354

69. Beccano-Kelly DA, Kuhlmann N, Tatarnikov I, et al. Synaptic function is modulated by LRRK2 and glutamate release is increased in cortical neurons of G2019S LRRK2 knock-in mice. *Front Cell Neurosci*. 2014;8(SEP):1-11. doi:10.3389/FNCEL.2014.00301/BIBTEX

70. Beccano-Kelly DA, Volta M, Lise LN, et al. LRRK2 overexpression alters glutamatergic presynaptic plasticity, striatal dopamine tone, postsynaptic signal transduction, motor activity and memory. *Hum Mol Genet*. 2015;24(5):1336-1349. doi:10.1093/HMG/DDU543

71. Piccoli G, Onofri F, Cirnaru MD, et al. Leucine-Rich Repeat Kinase 2 Binds to Neuronal Vesicles through Protein Interactions Mediated by Its C-Terminal WD40 Domain. *Mol Cell Biol*. 2014;34(12):2147-2161. doi:10.1128/mcb.00914-13

72. Arranz AM, Delbroek L, van Kolen K, et al. LRRK2 functions in synaptic vesicle endocytosis through a kinasedependent mechanism. *J Cell Sci*. 2015;128(3):541-552. doi:10.1242/jcs.158196

73. Hupfeld CJ, Olefsky JM. Regulation of receptor tyrosine kinase signaling by GRKs and beta-arrestins. *Annu Rev Physiol*. 2007;69:561-577. doi:10.1146/ANNUREV.PHYSIOL.69.022405.154626

74. Nazish I, Mamais A, Mallach A, et al. Differential LRRK2 signalling and gene expression in WT-LRRK2 and G2019S-LRRK2 mouse microglia treated with zymosan and MLi2. *bioRxiv*. Published online September 14, 2023. doi:10.1101/2023.09.14.557532

75. Jeremy Nichols R. LRRK2 Phosphorylation. *Adv Neurobiol*. 2017;14:51-70. doi:10.1007/978-3-319-49969-7_3

76. Chen YJ, Xie MR, Zhou SQ, Liu F. Synapses-associated research in Parkinson's disease: an explored trends analysis. *Front Aging Neurosci*. 2025;17. doi:10.3389/FNAGI.2025.1537119

77. (GP2) TGPGP, Leonard HL. Novel Parkinson's Disease Genetic Risk Factors Within and Across European Populations. *medRxiv*. Published online March 17, 2025:2025.03.14.24319455. doi:10.1101/2025.03.14.24319455

78. Uytterhoeven V, Verstreken P, Nachman E. Synaptic sabotage: How Tau and α-Synuclein undermine synaptic health. *J Cell Biol*. 2025;224(2). doi:10.1083/JCB.202409104

79. Gaig C, Martí MJ, Ezquerra M, Rey MJ, Cardozo A, Tolosa E. G2019S LRRK2 mutation causing Parkinson's disease without Lewy bodies. *J Neurol Neurosurg Psychiatry*. 2007;78(6):626-628. doi:10.1136/JNNP.2006.107904

80. Jackson LM, Woodruff BK, Tremblay C, et al. Parkinson's Disease Associated with G2019S LRRK2 Mutations without Lewy Body Pathology. *Mov Disord Clin Pract*. 2024;11(7):874-878. doi:10.1002/MDC3.14068

81. Sekiya H, Franke L, Hashimoto Y, et al. Widespread distribution of α-synuclein oligomers in LRRK2-related Parkinson's disease. *Acta Neuropathol*. 2025;149(1):42. doi:10.1007/S00401-025-02872-9

82. Jensen NM, Vitic Z, Antorini MR, Viftrup TB, Parkkinen L, Jensen PH. Abundant non-inclusion α-synuclein pathology in Lewy body-negative LRRK2-mutant cases. *Acta Neuropathol*. 2025;149(1). doi:10.1007/S00401-025-02871-W

83. Gaertner Z, Azcorra M, Dombeck DA, Awatramani R. Molecular heterogeneity in the substantia nigra: A roadmap for understanding PD motor pathophysiology. *Neurobiol Dis*. 2022;175. doi:10.1016/J.NBD.2022.105925

84. Gaertner Z, Oram C, Schneeweis A, et al. Molecular and spatial transcriptomic classification of midbrain dopamine neurons and their alterations in a LRRK2G2019S

model of Parkinson's disease. *bioRxiv*. Published online June 8, 2024. doi:10.1101/2024.06.06.597807

85. Hedrick NG, Harward SC, Hall CE, Murakoshi H, McNamara JO, Yasuda R. Rho GTPase complementation underlies BDNF-dependent homo- and heterosynaptic plasticity. *Nature*. 2016;538(7623):104-108. doi:10.1038/nature19784

86. Chan D, Citro A, Cordy JM, Shen GC, Wolozin B. Rac1 protein rescues neurite retraction caused by G2019s leucine-rich repeat kinase 2 (LRRK2). *Journal of Biological Chemistry*. 2011;286(18):16140-16149. doi:10.1074/jbc.M111.234005

87. Feng M, Hu X, Li N, et al. Distinctive roles of Rac1 and Rab29 in LRRK2 mediated membrane trafficking and neurite outgrowth. *J Biomed Res*. 2018;32(2):145-156. doi:10.7555/JBR.31.20170039

88. Worth DC, Daly CN, Geraldo S, Oozeer F, Gordon-Weeks PR. Drebrin contains a cryptic F-actin-bundling activity regulated by Cdk5 phosphorylation. *J Cell Biol*. 2013;202(5):793-806. doi:10.1083/JCB.201303005

89. Beaudoin GMJ, Schofield CM, Nuwal T, et al. Afadin, a Ras/Rap effector that controls cadherin function, promotes spine and excitatory synapse density in the hippocampus. *J Neurosci*. 2012;32(1):99-110. doi:10.1523/JNEUROSCI.4565-11.2012

90. Takahashi H, Sekino Y, Tanaka S, Mizui T, Kishi S, Shirao T. Drebrin-dependent actin clustering in dendritic filopodia governs synaptic targeting of postsynaptic density-95 and dendritic spine morphogenesis. *Journal of Neuroscience*. 2003;23(16):6586-6595. doi:10.1523/jneurosci.23-16-06586.2003

91.   Aoki C, Kojima N, Sabaliauskas N, et al. Drebrin a knockout eliminates the rapid form of homeostatic synaptic plasticity at excitatory synapses of intact adult cerebral cortex. *Journal of Comparative Neurology*. 2009;517(1):105-121. doi:10.1002/cne.22137

92.   Sekino Y, Tanaka S, Hanamura K, et al. Activation of N-methyl-d-aspartate receptor induces a shift of drebrin distribution: Disappearance from dendritic spines and appearance in dendritic shafts. *Molecular and Cellular Neuroscience*. 2006;31(3):493-504. doi:10.1016/j.mcn.2005.11.003

93.   Spence EF, Kanak DJ, Carlson BR, Soderling SH. The Arp2/3 complex is essential for distinct stages of spine synapse maturation, including synapse unsilencing. *Journal of Neuroscience*. 2016;36(37):9696-9709. doi:10.1523/JNEUROSCI.0876-16.2016

94.   Kim KS, Marcogliese PC, Yang J, et al. Regulation of myeloid cell phagocytosis by LRRK2 via WAVE2 complex stabilization is altered in Parkinson's disease. *Proc Natl Acad Sci U S A*. 2018;115(22):E5164-E5173. doi:10.1073/pnas.1718946115

95.   Giaime E, Tong Y, Wagner LK, Yuan Y, Huang G, Shen J. Age-Dependent Dopaminergic Neurodegeneration and Impairment of the Autophagy-Lysosomal Pathway in LRRK-Deficient Mice. *Neuron*. 2017;96(4):796-807.e6. doi:10.1016/j.neuron.2017.09.036

96.   Huang G, Bloodgood DW, Kang J, et al. Motor Impairments and Dopaminergic Defects Caused by Loss of Leucine-Rich Repeat Kinase Function in Mice. *J Neurosci*. 2022;42(23):4755-4765. doi:10.1523/JNEUROSCI.0140-22.2022

97.   Kang J, Huang G, Ma L, et al. Cell-autonomous role of leucine-rich repeat kinase in the protection of dopaminergic neuron survival. *Elife*. 2024;12. doi:10.7554/ELIFE.92673

98. Biskup S, Moore DJ, Rea A, et al. Dynamic and redundant regulation of LRRK2 and LRRK1 expression. *BMC Neurosci*. 2007;8. doi:10.1186/1471-2202-8-102

99. Cogo S, Ho FY, Tosoni E, et al. The Roc domain of LRRK2 as a hub for protein-protein interactions: a focus on PAK6 and its impact on RAB phosphorylation. *Brain Res*. 2022;1778. doi:10.1016/j.brainres.2022.147781

100. Narita M, Aoki K, Takagi M, Yajima Y, Suzuki T. Implication of brain-derived neurotrophic factor in the release of dopamine and dopamine-related behaviors induced by methamphetamine. *Neuroscience*. 2003;119(3):767-775. doi:10.1016/S0306-4522(03)00099-X

101. Morella I, Hallum H, Brambilla R. Dopamine D1 and Glutamate Receptors Co-operate With Brain-Derived Neurotrophic Factor (BDNF) and TrkB to Modulate ERK Signaling in Adult Striatal Slices. *Front Cell Neurosci*. 2020;14. doi:10.3389/fncel.2020.564106

102. Hyman C, Hofer M, Barde YA, et al. BDNF is a neurotrophic factor for dopaminergic neurons of the substantia nigra. *Nature*. 1991;350(6315):230-232. doi:10.1038/350230a0

103. Baquet ZC, Bickford PC, Jones KR. Brain-derived neurotrophic factor is required for the establishment of the proper number of dopaminergic neurons in the substantia nigra pars compacta. *Journal of Neuroscience*. 2005;25(26):6251-6259. doi:10.1523/JNEUROSCI.4601-04.2005

104. Rahmani F, Saghazadeh A, Rahmani M, et al. Plasma levels of brain-derived neurotrophic factor in patients with Parkinson disease: A systematic review and meta-analysis. *Brain Res*. 2019;1704:127-136. doi:10.1016/j.brainres.2018.10.006

105. Lin CH, Wu RM, Tai CH, Chen ML, Hu FC. Lrrk2 S1647T and BDNF V66M interact with environmental factors to increase risk of Parkinson's disease. *Parkinsonism Relat Disord*. 2011;17(2):84-88. doi:10.1016/j.parkreldis.2010.11.011

106. Liu J, Zhou Y, Wang C, Wang T, Zheng Z, Chan P. Brain-derived neurotrophic factor (BDNF) genetic polymorphism greatly increases risk of leucine-rich repeat kinase 2 (LRRK2) for Parkinson's disease. *Parkinsonism Relat Disord*. 2012;18(2):140-143. doi:10.1016/j.parkreldis.2011.09.002

107. Siuda J, Patalong-Ogiewa M, Żmuda W, et al. Cognitive impairment and BDNF serum levels. *Neurol Neurochir Pol*. 2017;51(1):24-32. doi:10.1016/j.pjnns.2016.10.001

108. Ziebell M, Khalid U, Klein AB, et al. Striatal dopamine transporter binding correlates with serum BDNF levels in patients with striatal dopaminergic neurodegeneration. *Neurobiol Aging*. 2012;33(2):428.e1-428.e5. doi:10.1016/j.neurobiolaging.2010.11.010

109. Tagliaferro P, Burke RE. Retrograde Axonal Degeneration in Parkinson Disease. *J Parkinsons Dis*. 2016;6(1):1-15. doi:10.3233/JPD-150769

110. Sharma A, Toepfer CN, Ward T, et al. CRISPR/Cas9-Mediated Fluorescent Tagging of Endogenous Proteins in Human Pluripotent Stem Cells. *Curr Protoc Hum Genet*. 2018;96(1):21.11.1-21.11.20. doi:10.1002/CPHG.52

111. Beccano-Kelly DA, Cherubini M, Mousba Y, et al. Calcium dysregulation combined with mitochondrial failure and electrophysiological maturity converge in Parkinson's iPSC-dopamine neurons. *iScience*. 2023;26(7):107044. doi:10.1016/J.ISCI.2023.107044

112. Zaqout S, Kaindl AM. Golgi-cox staining step by step. *Front Neuroanat*. 2016;10:181342. doi:10.3389/FNANA.2016.00038/BIBTEX

113. Risher WC, Ustunkaya T, Alvarado JS, Eroglu C. Rapid Golgi Analysis Method for Efficient and Unbiased Classification of Dendritic Spines. *PLoS One*. 2014;9(9):e107591. doi:10.1371/JOURNAL.PONE.0107591

114. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nature Protocols 2016 11:12*. 2016;11(12):2301-2319. doi:10.1038/nprot.2016.136

115. Aguilan JT, Kulej K, Sidoli S. Guide for protein fold change and: P-value calculation for non-experts in proteomics. *Mol Omics*. 2020;16(6):573-582. doi:10.1039/d0mo00087f

116. Lazar C, Gatto L, Ferro M, Bruley C, Burger T. Accounting for the Multiple Natures of Missing Values in Label-Free Quantitative Proteomics Data Sets to Compare Imputation Strategies. *J Proteome Res*. 2016;15(4):1116-1125. doi:10.1021/acs.jproteome.5b00981

117. Zhao Y, Vavouraki N, Lovering RC, et al. Tissue specific LRRK2 interactomes reveal a distinct striatal functional unit. *PLoS Comput Biol*. 2023;19(1):e1010847. doi:10.1371/JOURNAL.PCBI.1010847

118. Raudvere U, Kolberg L, Kuzmin I, et al. g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res*. 2019;47(W1):W191-W198. doi:10.1093/NAR/GKZ369

Figures

Figure 1

Figure 1. LRRK2 phosphorylation at ser935 rapidly and transiently increases upon BDNF stimulation in primary mouse cortical neurons and differentiated SH-SY5Y cells. **(A)** Schematic representation of the experimental setting of (B). **(B)** Phospho-Ser935 and total Lrrk2 protein levels of primary cortical neurons at DIV14 treated with 100 ng/mL BDNF for 0, 5, 30, 60 180 mins. MLi-2 was used at 500 nM for 90 mins to inhibit Lrrk2 kinase activity. **(C)** Quantification of n=6 independent experiments of (B). One-way ANOVA (***$P < 0.001$), Dunnett's multiple comparison test (t=0 compared with different time points, *$P < 0.05$ t=0 vs. t=5'). **(D)** Schematic representation of the experimental setting of (F) and (G). **(E)** TrkB protein levels in undifferentiated or retinoic acid-differentiated SH-SY5Y cells. **(F)** Phospho Thr202/185 ERK1/2, total ERK1/2 and **(G)** phospho Ser473 AKT, total AKT protein levels of

retinoic acid-differentiated SH-SY5Y cells stimulated with 100 ng/mL BDNF for 0, 5, 30, 60 mins. **(H)** Phospho-Ser935 and total LRRK2 protein levels of retinoic acid-differentiated SH-SY5Y cells stimulated with 100 ng/mL BDNF for 0, 5, 30, 60 mins. MLi-2 was used at 500 nM for 90 mins to inhibit LRRK2 kinase activity. **(I)** Quantification of (H) (n=4 independent experiments). One-way ANOVA (**$P < 0.01$), Dunnett's multiple comparison test (*$P < 0.05$ t=0 vs. t=5'). **(J)** Phospho-T73 and total RAB10 protein levels of retinoic acid-differentiated SH-SY5Y cells stimulated with 100 ng/mL BDNF for 0, 5, 30, 60 mins. MLi-2 was used at 500 nM for 90 mins to inhibit LRRK2 kinase activity. **(K)** Quantification of (**J**) (n=4 independent experiments). One-way ANOVA ($P > 0.5$), Dunnett's multiple comparison test (*$P < 0.05$ t=0 vs. t=30').

Figure 2

Figure 2. BDNF promotes LRRK2 interaction with post-synaptic actin cytoskeleton components.

(A) Volcano plot of GFP-LRRK2 versus GFP enriched interactors from differentiated SH-SY5Y cells (n=6 replicates: n=3 independent experiments each with 2 technical replicates). Interactors that were considered for SynGO analysis have adjusted *P* value < 0.05 and FC > 2. (red dots). **(B)** Donut charts with % of SynGO genes among the total 207 LRRK2 interactors (A) and with % of presynaptic and postsynaptic proteins among SynGO genes. SynGO-CC terms visualized with a sunburst plot showing significant categories (below). **(C)** Volcano plot of GFP-LRRK2 +/-BDNF enriched interactors from differentiated SH-SY5Y cells (n=6 replicates: n=3 independent experiments each with 2 technical replicates). Interactors selected for pathway enrichment analysis fall into two categories: (i) adjusted *P* value < 0.05 and FC > 0.5 or FC < -0.5, or (ii) |FC| > 1 regardless of the *P* value. Proteins with increased interaction upon BDNF stimulation are blue-labeled, proteins with decreased interaction are orange-labeled. Donut chart showing the number of SynGO annotated genes versus non synaptic proteins in BDNF-up versus BDNF-down interactors (top right). **(D)** G:profiler g:GOSt pathway enrichment analysis showing the top 5 enriched biological processes (BP) categories of BDNF-increased interactors (top, blue bar graph) versus BDNF-decreased interactors (bottom, orange bar graph). GO analysis was performed with g:Profiler on 04/14/2025 (version e112_eg59_p19_25aa4782, database updated on 03/02/2025); enriched terms determined using g:SCS correction an size terms set to <500 to increase specificity. **(E)** Protein-Protein Interaction Networks built with STRING (https://string-db.org/) of all BDNF(+) interactors and SynGO BDNF-increased interactors (inset) and **(F)** of SynGO BDNF-decreased interactors. **(G)** Validation of increased drebrin:LRRK2 interaction upon BDNF treatment (15 mins) and quantification (one sample t-test, *P*<0.01). Samples are the same analyzed in the three rounds of MS (A). **(H)** Western blot analysis of brain samples from WT and *Dbn1* (drebrin) KO mice assessing Lrrk2 Ser935 and Rab8 (pan Rabs) phosphorylation Differences between the two genotypes have been evaluated using Student's t-test (significance **P*<0.05 pSer935), *n*=3 animals per genotype per age.

Figure 3

Figure 3. Convergence on actin and synaptic processes in literature-curated LRRK2 interactors and the LRRK2 G2019S striatal phospho-proteome. (A) Network graph showing one of the topological clusters extracted from the LRRK2 protein-protein interaction network. Proteins are represented as nodes while protein-protein interactions are represented as edges. Node size is proportional to the degree centrality: the larger the node, the higher the degree, the more interactions the node has in the network. **(B)** Gene Ontology biological-processes (GO-BPs) enrichment; the analysis was performed using g:Profiler, GO-BPs with term size >500 were discarded as general terms and the remaining terms was clustered based on semantics using the keywords: Postsynaptic (Actin) Cytoskeleton, Lamellipodium, Synaptic Vesicle, Actin Polymerization/Nucleation, DNA Recombination/Repair, Cell/Neuron

Projection. **(C)** Upper network graph as in (A) highlighting actin cytoskeletal proteins in blue and the proportion over the total (pie chart) and lower network graph as in (A) highlighting synaptic proteins in red and the proportion over the total (pie chart). **(D)** Experimental design of phospho-proteomic analysis comparing WT and G2019S knockin mouse striata. Phosphopeptide abundances were normalized to total proteins levels. **(E)** Volcano plot of differentially enriched phosphopeptides, expressed as Log2 fold change (FC) between G2019S-KI versus WT. Highlighted peptides correspond to LRRK2 interactors discovered by MS in Figure 2. **(F)** GO enrichment analysis performed with g:Profiler (accessed on 01/23/2025; version e94_eg41_p11), filtering for GO terms with size <200 to avoid overly general categories. The top 5 most significant terms from the Cellular Component (CC) and Biological Process (BP) domains are shown. **(G)** Sunburst plot showing enriched SynGO cellular component (CC) categories and the percentage of SynGO genes among the differentially phosphorylated proteins. **(H)** Venn diagrams showing common and unique hits across the three datasets: GFP-LRRK2 interactomics, synaptic cluster and phospho-MS.

Figure 4

Figure 4. BDNF signaling is impaired in *Lrrk2* knockout neurons. (A) Six-day RA-differentiated SH-SY5Y naïve and CRISPR-LRRK2 KO cells stimulated with BDNF at different time points (0, 1, 5, 15, 60, 180 mins). Anti-TrkB antibodies were used to confirmed expression of BDNF receptor. To compare BDNF-induced signaling in naïve vs LRRK2-KO cells, phosphorylated Akt (S473) and Erk1/2 (T185/T202) were evaluated. **(B)** Quantification of phosphorylated proteins show from n=3 independent differentiation experiments. Two-way

ANOVA; phospho-Akt: interaction P=0.7079, F (5, 24) = 0.5896; genotype: **P=0.0014, F (1, 24) = 13.07; time: P=0.5610, F (5, 24) = 0.7994. Phospho-Erk1: interaction P=0.0783, F (5, 24) = 2.284; genotype: ***P=0.0003, F (1, 24) = 17.58; time: *P=0.0244, F (5, 24) = 3.174. Phospho-Erk2: interaction: P=0.3725, F (5, 24) = 1.128; genotype: ***P=0.0008 F (1, 24) = 14.59; time: P=0.1356, F (5, 24) = 1.879. **(C)** DIV14 primary cortical neurons from WT vs KO mice stimulated with BDNF at different time points (0, 5, 15, 30, 60 mins) in the presence or absence of LRRK2 inhibitor MLi-2 (90 min, 500 nM). To compare BDNF-induced signaling in WT vs KO neurons, phosphorylated Akt (S473) and Erk1/2 (T185/T202) were evaluated. Phosphorylated LRRK2 was assessed with pS935 antibodies. **(D)** Quantification of phosphorylated proteins show from n=9 independent cultures. Two-way ANOVA; phospho-Akt: interaction P=0.1186, F (3, 63) = 2.031; genotype: **P=0.0037, F (1, 63) = 9.101; time: P=0.0342, F (3, 63) = 3.069. Phospho-Erk1: interaction P=0.3256, F (3, 64) = 1.177; genotype: *P=0.0201, F (1, 64) = 5.680; time: ****P<0.0001, F (3, 64) = 10.04. Phospho-Erk2: interaction P=0.8524, F (3, 62) = 0.2622; genotype: P=0.7528, F (1, 62) = 0.1001; time: ***P=0.0003, F (3, 62) = 7.428. **(E)** Overview of the experimental workflow to induce spine formation/maturation. **(F)** Representative confocal images of primary cortical neurons stimulated with 100 ng/ml of BDNF or vehicle control for 24 hours. GFP has been transfected at DIV4 to fill the neuroplasm and visualize individual dendrites, MAP2 is a neuronal marker and PSD95 is a marker of mature spines. **(G)** Quantification of the number of PSD-positive puncta per unit of length (100 μm). Dots represents individual segments (n≅20 neurites from n=5-6 neurons per replicate) and colors define neuronal cultures prepared in different days from pulled pups (n≅8 pups per culture per genotype, n=3 independent cultures). Two-way ANOVA; interaction **P=0.0038, F (1, 191) = 8.584; genotype: P=0.5833, F (1, 191) = 0.3020; treatment: *P=0.0126, F (1, 191) = 6.346. Šídák's multiple comparisons test: vehicle vs. BDNF (WT) ***P=0.0005; vehicle vs. BDNF (KO) P=0.9440; Vehicle (WT) vs. vehicle (KO) P= 0.3756.

Figure 5

Figure 5. Postsynaptic structural changes in young *Lrrk2* knockout mice.

(A) Overview of experimental design. **(B)** Representative images of neurite segments from Golgi-Cox stained neurons of dorsal striatum. Scale bar: 3 μm. **(C)** Quantification of average spine number (top), width (middle) and length (bottom) of *n*=3 animals per group (same

segments analyzed in B). Statistical significance was determined by two-way ANOVA with Šídák's multiple comparisons test. Number of protrusions: interaction $P=0.3820$, F (2, 12) = 1.044; age: ****$P<0.0001$, F (2, 12) = 27.12; genotype: $P=0.0840$, F (1, 12) = 3.550; WT vs. KO (1 month) $P=0.2013$; WT vs. KO (4 months) $P=0.5080$; WT vs. KO (18 months) $P>0.9999$. Width: interaction $P=0.0815$, F (2, 12) = 3.112; age: ***$P=0.0004$, F (2, 12) = 16.14; genotype: ***$P=0.0007$, F (1, 12) = 20.77; WT vs. KO (1 month) **$P=0.0017$; WT vs. KO (4 months) $P=0.4602$; WT vs. KO (18 months) $P=0.2489$. Length: interaction *$P=0.0345$, F (2, 12) = 4.514; age: *$P=0.0203$, F (2, 12) = 5.488; genotype: ***$P=0.0006$, F (1, 12) = 21.53; WT vs. KO (1 month) ***$P=0.0008$; WT vs. KO (4 months) $P=0.2875$; WT vs. KO (18 months) $P=0.5948$. **(D)** Representative transmission electron microscopy (TEM) micrographs of striatal synapses from 1-month and 18-month old WT vs KO mouse brain slices. Scale bar: 200 nm. **(E)** Quantification of post-synaptic density (PSD) length. Graphs show the length of individual synapses (grey dots) and the average PSD length per animal (colored dots). Statistical significance has been calculated with Student's t-test: 1 month-old mice ($n=4$ mice, 95 synapses WT, 118 synapses KO, ***$P=0.0003$); 18-month old mice ($n=3$ mice, 108 synapses WT, 119 synapses KO, $P=0.8502$). **(F)** Quantitative PCR of *Bdnf, TrkB, Psd95, Shank3* and *Dbn1* mRNA expression in striatum, cortex and midbrain form $n=6$ *Lrrk2* WT and $n=6$ *Lrrk2* KO one-month old mice. Statistical significance has been calculated with Student's t-test. **(G)** Western blot analysis of brain samples from the same Lrrk2 WT and KO mice where Golgi-Cox staining has been performed. The reduction in drebrin content is significant in Lrrk2 KO mice at 1 month of age. Differences between the two genotypes have been evaluated using Student's t-test (significance **$P<0.01$), $n=3$-4 animals per genotype per age.

Figure 6

Figure 6. Effect of BDNF exposure on spontaneous electrical activity recorded in DIV70 LRRK2 WT and KO cortical neurons. (A) Schematic representation of the experimental setup. **(B)** Representative spontaneous traces of hiPSC-derived WT and KO neurons in the presence or absence of 50 ng/ml BDNF for 24 hrs. **(C)** Frequency of miniature postsynaptic excitatory currents (mEPSC) in LRRK2 WT and KO cultures (WT vehicle: 0.16 ± 0.03 Hz (n=15, N=3); WT BDNF: 1.36 ± 0.38 Hz (n=11, N=3); KO vehicle: 1.09 ± 0.21 Hz (n=11, N=3); KO BDNF: 1.11 ± 0.25 Hz (n=12, N=3). Statistical significance was determined using one-way ANOVA (WT vehicle vs. WT BDNF **P<0.01; WT BDNF vs. KO BDNF *P<0.05; WT vehicle vs. KO BDNF *P<0.05; P>0.05 for all the other comparisons). **(D)** Cumulative probability curves of interevent interval (IEIs) in DIV70 WT and KO -/+ BDNF treated cultures. **(E)** Amplitude of miniature postsynaptic excitatory currents (mEPSC) in DIV70 LRRK2 WT and KO cultures. BDNF treatment has no effect on the average peak amplitude (WT vehicle: 11.00 ± 1.85 pA (n=14, N=3); WT BDNF: 7.9 ± 0.84 pA (n=11, N=3); KO vehicle 10.42 ± 1.53 pA (n=11, N=3); KO BDNF 11.12 ± 1.76 pA (n=12, N=3)). Statistical significance was determined using one-way ANOVA (P>0.05 for all comparisons).

Supplementary figures

Supplementary figure 1

Figure S1. (A) Western blot analysis of undifferentiated versus differentiated naïve SH-SY5Y and western blot with anti-phospho-Ser935 LRRK2 and total LRRK2 antibodies. **(B)** Cells stably overexpressing GFP-LRRK2 treated with 100 ng/ml BDNF for 5 mins or with 500 nM MLi-2 for 90 mins. **(C)** On the left, schematic representation of experimental setting. On the right, BDNF stimulation of GFP-LRRK2 SH-SY5Y cells prior to MLi-2 (500 nM) treatment for 90 minutes. Western blot were performed using anti LRRK2 (phospho-Ser935 and total), anti-RAB10 (phospho Thr73 and total) and anti-RAB8 (pan-phopho-RABs and total RAB8).

Supplementary figure 2

Figure S2. Western blot analysis validating the specificity of LRRK2-Drebrin interaction.

(A) Confocal immunofluorescence of DIV14 primary neurons co-stained with MAP2 (neuronal marker), PSD95 (postsynaptic marker) and drebrin. Scale bar: 100 μm; magnifications scale bar: 10 μm. **(B)** To rule out that drebrin does not bind the resin but exclusively GFP-LRRK2 bound to anti-GFP nanobodies, GFP-LRRK2 was purified from cell lysates containing a mix of GFP-LRRK2 OE SH-SY5Y cells and naïve SH-SY5Y cells at different ratios: 4:0, 3:1, 2:2, 1:3, 0:4 (OE:naïve). The levels of drebrin bound to GFP-LRRK2 diminish proportionally with the reduction of LRRK2 purified from the lysate, with a complete lack of signal in the eluate from the resin incubated with naïve cells only (0:4 condition), confirming the specificity of drebrin binding to GFP-LRRK2. **(C)** Co-immunoprecipitation of 3xFlag-LRRK2 with YFP-drebrin domains (full-length, N-terminus and C-terminus;

aminoacid boundaries are indicated in the figure). Drebrin constructs were immunoprecipitated with GFP-trap resin and co-precipitated LRRK2 visualized by western blot using anti Flag antibodies.

Supplementary figure 3

Figure S3. (A) Functional enrichment analysis; the analysis was performed using g:Profiler g:GOSt, proteins related to actin cytoskeleton were manually identified and blue-colored, and the first ten GO-BPs categories with term size <500 were graphed. **(B)** SynGO of enriched biological-processes and cellular components; synaptic proteins were red-colored, and significant terms in SynGO-BP and SynGO-CC categories were graphed.

Supplementary figure 4

Figure S4. Analysis of the predicted aminoacid pathogenicity **in** drebrin using AlphaMissense pathogenicity score. Aminoacid change of S339 or its neighbor S337 is predicted to be benign, suggesting that phosphorylation at this site may play a regulatory function.

Supplementary figure 5

Guide name	sequence
gRNA1	CACACTCGCGACTCTCATAT
gRNA2	TGTGCCTCTGTTGATCGTCT
gRNAB	GAGTCCAAGACGATCAACAG

Figure S5. Western blot of different monoclonal populations of SH-SY5Y cells transfected with pSpCas9(BB)-2A-Puro (PX459) V2.0 vector (Addgene, Watertown, MA, US) for co-expression of Cas9 and gRNAs (gRNA1, gRNA2 and gRNAB) and subsequent single clone isolation and expansion. The band intensities corresponding to the total amount of endogenous LRRK2 were normalized to GAPDH, used as loading control.

Supplementary figure 6

Figure S6. Morphological classification of protrusions into four classes (filopodia, thin, mushroom, branched) and quantified as % of the total number. Each dot represents one segment ($n \geq 20$ segments analyzed per animal; genotype: wild-type vs. *Lrrk2* KO; age: 1, 4, 18 month-old; n=3 mice per group) and error bars represent the mean ± SD of n=3 mice (color coded). Statistical significance was determined by two-way ANOVA with Šídák's multiple comparisons test on mean values. One-month: interaction: **P=0.0017, $F_{(3, 16)}$ = 8.018; genotype: P=0.5731, $F_{(1, 16)}$ = 0.3309; class of protrusion: ****P<0.0001, $F_{(3, 16)}$ = 158.4; filopodia WT vs. KO P=0.0201; thin WT vs. KO *P=0.0102; mushroom WT vs. KO P=0.9585; branched WT vs. KO P=0.8726. Four-months: interaction: P=0.3178, $F_{(3, 16)}$ = 1.271; genotype: P=0.7300, $F_{(1, 16)}$ = 0.1233; class of protrusion: ****P<0.0001, $F_{(3, 16)}$ = 438.9; filopodia WT vs. KO P=0.8478; thin WT vs. KO P=0.4921; mushroom WT vs. KO P=0.8882; branched WT vs. KO P=0.9858. Eighteen-months: interaction: P=0.3237, $F_{(3, 16)}$ = 1.253; genotype: P=0.6880, $F_{(1, 16)}$ = 0.1672; class of protrusion: ****P<0.0001, $F_{(3, 16)}$ = 69.67; filopodia WT vs. KO P=0.9756; thin WT vs. KO P=0.6413; mushroom WT vs. KO P>0.9999; branched WT vs. KO P=0.5290.

Supplementary figure 7

Figure S7. No significant change in the frequency of mEPSC between WT and KO in **(A)** DIV50 and **(B)** DIV90 due to the addition of BDNF (50 mg/ml for 24 hrs). One-way ANOVA with Tukey's post-doc test. $P > 0.05$ for all comparisons (N=2 independent cultures). **(C)** Immunocytochemistry of WT and LRRK2 KO cortical neurons at DIV70 using Bassoon (pre-synaptic marker), Homer (post-synaptic marker), MAP2 (neuronal marker) and DAPI (scale bars 20 μm; zoom 5 μm). **(D)** Quantification of the % of Homer puncta colocalizing with total Bassoon (on MAP2-positive processes). BDNF was applied at 50 mg/ml for 24. One-way ANOVA with Tukey's post-doc test. $P > 0.05$ for all comparisons. N=4 cultures with 5-25 neurons imaged per condition per culture.

Supplementary figure 8

Figure S8. (A) Functional enrichment analysis of the nearest genes to each of the 134 significant loci (obtained from table S3, column F from https://doi.org/10.1101/2025.03.14.24319455) was performed using g:Profiler g:GOSt with term size <500 to increase specificity. (B) SynGO analysis of 32 genes mapped from the input list (from GWAS) revealed significant enrichment of synaptic Cellular Component (5 terms) and Biological Process (6 terms) annotations at 1% FDR. Annotations were based on 31 genes for Cellular Components and 27 for Biological Processes, using a brain-expressed background of 18,035 genes. Experimental-evidence filtering was not applied. SynGO version 20231201 was used.

Table 1. Capacitance and access resistance of LRRK2 KO and WT cortical neurons at 50, 70 and 90 DIV with or without BDNF stimulation.

Capacitance

50 DIV

	Mean (pF)	SEM	n
WT -BDNF	14.14	1.858	12
WT +BDNF	15.1	1.331	12
KO -BDNF	16.21	2.299	9
KO +BDNF	14.66	1.295	11

70 DIV

	Mean (pF)	SEM	n
WT -BDNF	23	2.177	16
WT +BDNF	22.4	2.579	12
KO -BDNF	24.64	2.555	13
KO +BDNF	20.03	2.232	13

90 DIV

	Mean (pF)	SEM	n
WT -BDNF	12.51	1.994	8
WT +BDNF	20.9	6.365	7
KO -BDNF	24.23	3.062	7
KO +BDNF	24.96	6.49	8

Access Resistance

50 DIV

	Mean (MΩ)	SEM	n
WT -BDNF	18.54	1.312	12
WT +BDNF	18.27	1.053	12
KO -BDNF	20.11	1.467	9
KO +BDNF	18.71	1.884	11

70 DIV

	Mean (MΩ)	SEM	n
WT -BDNF	21.18	1.372	16
WT +BDNF	21.6	1.273	12
KO -BDNF	22.36	1.835	13
KO +BDNF	15.1	1.122	13

90 DIV

	Mean (MΩ)	SEM	n
WT -BDNF	17.6	2.221	8
WT +BDNF	18.64	1.68	7
KO -BDNF	16.66	1.609	9
KO +BDNF	14.64	1.591	8

Supplemental file 1

Procedure to check CRISPR-mediated LRRK2 KO in SH-SY5Y cells

Selection of off-targets and primers design

- http://crispor.gi.ucsc.edu/
- Copy and paste the sequence of LRRK2 exon we targeted (exon # XX)
- Select the reference genome: Homo sapiens – Human – USCS Dec. 2013 (GRCh38/hg38) + SNPs: dbSNP148, Kaviar
- Select the Protospacer Adjacent Motif (PAM): 20bp-NGG – Sp Cas9, SpCas9-HF1, eSpCas9 1.1
- Submit
- Select the first 8 results exhibiting a CFD (Cutting Frequency Determination) off-target score > 0.2
- Search for the genomic region containing the off-target sequence reported for each of the selected results
- Design forward and reverse primers, respectively upstream to and downstream of the off-target, for each sequence (off-targets and LRRK2 on-target)
- Select the forward primers for the sequencing

Cell culture procedure and sequencing

- Culture SH-SY5Y cells wild-type and LRRK2 KO until reaching confluence in T25 flasks as described in the material and methods
- Detach and centrifuge the cell populations, keeping the pellets in ice
- Deliver pairs of primers and cellular pellets to BMR Genomics (https://www.bmr-genomics.it/) for PCR amplification and sequencing
- Compare the sequences obtained from wild-type and LRRK2 KO cells with the reference ones using BLAST in order to find potential mismatches
- If the sequence identity is higher than 99% and the mismatch is placed in the final portion of the sequenced region, it is accepted as equal to the reference sequence

Results

Disruption of the sequence was observed only in LRRK2 on-target site in the LRRK2 KO SH-SY5Y cells, while the 8 off-target regions were unchanged across the genotypes as well as when compared to the reference sequence

ontarget_mm0_exon_LRRK2_chr12_40225585
F_TGCTGGTGTTCACGTACTCC; Tm=59.9
R_GCACACTCGCGACTCTCATA; Tm=59.9

`hLRRK2_LRRK2 WT`
CTGTGCTTTTATTTTTGCAAACTTTCTCCCCCTCCTTACATTTGCAAATTTTGTCCTCCTCCCCTTGACCCT
GCTCAAACCCGGACTCTTAAGGAGCCGCAAACTCCCATATCCTTTCCTTAGGGCAGAAAGCAGCTGAGA
ATTTCAGGAAGGTCTTCACCTTTTTGACTTTTCTCCCCGTTTCAGACTAAAAAGGAAAGGGGGTGCTGTG
GATTGTGACTTTGCTTCTTTTCCCCACCCACTTGTTTTCCAGCCTCCAAGTTATTTCAAGGCAAAAATATC
CA`TGTGCCTCTGTTGATCGTCT`TGGACTCCTATATGAGAGTCGCAGCAGGTAAAGGCATTGTTTTCACTTC
AACTCATTCTCCCTTCTGTTTGGAAGGAGACGTTTTACTGGCAATGT

`hLRRK2_LRRK2 KO`
CTGTGCTTTTATTTTTGCAAACTTTCTCCCCCTCCTTACATTTGCAAATTTTGTCCTCCTCCCCTTGACCCT
GCTCAAACCCGGACTCTTAAGGAGCCGCAAACTCCCATATCCTTTCCTTAGGGCAGAAAGCAGCTGAGA
ATTTCAGGAAGGTCTTCACCTTTTTGACTTTTCTCCCCGTTTCAGACTAAAAAGGAAAGGGGGTGCTGTG
GATTGTGACTTTGCTTCTTTTCCCCACCCACTTGTTTTCCAGCCTCCAAGTTATTTCAAGGCAAAAATATC
CA`TGTGCCTCTGTTGATCGTCT`TGGACTCCTATATGAGAGTCGCGGGTTGTGCAAAAGAATATTTTTCTTT
TTCTTTTCTTCCCTTTTTTTTTTTTTTTTTTTTTTTTTTTTTAAAAAGGGGTTTTGGGGCCCCTTTTAAAAAAGCGG
GGAGGGCAAA

1. **mm4_intergenic_APOC1|APOC4/APOC4-APOC2_chr19_44927007**
   F_GCCTCCACTCTGCAAGAACT; Tm=59.9
   R_TCCAACTCCGCAATTCTCCC; Tm=60.0

```
guide:        TGTGCCTCTGTTGATCGTCT  TGG
off-target:   GGTGGTTCTGTCGATCGTCT  TGG
              *    **        *
CFD Off-target score: 0.504871
MIT Off-target score: 0.43
Position: chr19:44927008-44927030:+
```

hAPOC1_LRRK2 WT
CGTGGGAGGGAGGGAGGAGGGTGCCACTGATCCCCTGAACCCCTGCCTCCGCCTCCAGGGTGCCCCT
CCGGCCTCGCCATGAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTGTCGATGGTCTTGGAA
GGTAAAAGTGGGATGGGAGAATTGCGGAGTTGGAA

hAPOC1_LRRK2 KO
CGTGGGAGGGAGGGAGGAGGGTGCCACTGATCCCCTGAACCCCTGCCTCCGCCTCCAGGGTGCCCCT
CCGGCCTCGCCATGAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTGTCGATGGTCTTGGAA
GGTAAAAGTGGGATGGGAGAATTGCGGAGTTGGAA

Score	Expect	Identities	Gaps	Strand
315 bits(170)	4e-91	170/170(100%)	0/170(0%)	Plus/Plus

```
Query  1    CGTGGGAGGGAGGGAGGAGGGTGCCACTGATCCCCTGAACCCCTGCCTCCGCCTCCAGGG  60
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    CGTGGGAGGGAGGGAGGAGGGTGCCACTGATCCCCTGAACCCCTGCCTCCGCCTCCAGGG  60

Query  61   TGCCCCTCCGGCCTCGCCATGAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTG  120
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   TGCCCCTCCGGCCTCGCCATGAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTG  120

Query  121  TCGATGGTCTTGGAAGGTAAAAGTGGGATGGGAGAATTGCGGAGTTGGAA  170
            ||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  121  TCGATGGTCTTGGAAGGTAAAAGTGGGATGGGAGAATTGCGGAGTTGGAA  170
```

2. **mm4_exon_APOE/CTB-129P6.7/APOC1_chr19_44914923**
   F_GACAGGACCTCCCAACCAAG; Tm=59.9
   R_TCCAACTCCGCAATTCTCCC; Tm=60.0

```
guide:       TGTGCCTCTGTTGATCGTCT TGG
off-target:  GGTGGTTCTGTCGATCGTCT TGG
             *    **       *

CFD Off-target score: 0.504871
MIT Off-target score: 0.43
Position: chr19:44914924-44914946:+
```

hAPOE_LRRK2 WT

AAGGGATCGAGGGAGGGAGGGAGGAGGGGAGGGATGATGCCCTGAACCCCTGCCTCCGCCTCCAGG
GTGCCCCTCCGGCCTCCCCATGAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTGTCGATGG
TCTTGGAAGGTAAAAGTGGGATGGGAGAATTGC

hAPOE_LRRK2 KO

GGAGGGATGATGCCCTGAACCCCTGCCTCCGCCTCCAGGGTGCCCCTCCGGCCTCCCCATGAGGCTCT
TCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTGTCGATGGTCTTGGAAGGTAAAAGTGGGATGGGAGAAT
TGCGGAGTTGGAT

Score	Expect	Identities	Gaps	Strand
253 bits(280)	9e-73	140/140(100%)	0/140(0%)	Plus/Plus

```
Query  1    GGAGGGATGATGCCCTGAACCCCTGCCTCCGCCTCCAGGGTGCCCCTCCGGCCTCCCCAT  60
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  28   GGAGGGATGATGCCCTGAACCCCTGCCTCCGCCTCCAGGGTGCCCCTCCGGCCTCCCCAT  87

Query  61   GAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTGTCGATGGTCTTGGAAGGTAA  120
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  88   GAGGCTCTTCCTGTCGCTCCCGGTCCTGGTGGTGGTTCTGTCGATGGTCTTGGAAGGTAA  147

Query  121  AAGTGGGATGGGAGAATTGC  140
            ||||||||||||||||||||
Sbjct  148  AAGTGGGATGGGAGAATTGC  167
```

3. **mm4_intergenic_RP11-329N22.1|RRAGC/RP5-864K19.7_chr1_38822106**
   F_GCCCTCACCTTACCAAAGCT; Tm=59.9
   R_CCGCTTTCATCCTAATGGGGT; Tm=60.1

```
guide:        TGTGCCTCTGTTGATCGTCT  TGG
off-target:   TCTGCCTCTTTCGATCATCT  AGG
               *         * *      *
```

CFD Off-target score: 0.497368
MIT Off-target score: 0.28
Position: chr1:38822107-38822129:+

hRP11-329N22.1_LRRK2 WT
GGCTCGCTTGCTAAAGCTCTCCAGAGAGGCACTTTGAATTAAGACCAGAAATGCTTTTAAAAGACTCTGG
TCTCTGGTCTTCTAATGATGACCTCTTCCTTAATTTTGTGATGGCTTCATCTGCCTCTTTCGATCATCTAGG
TAGCTATAAAAAGCTGAACACCTCCTACTAACCCCATTAGGATGAAAGCGG

hRP11-329N22.1_LRRK2 KO
GGCTCGCTTGCTAAAGCTCTCCAGAGAGGCACTTTGAATTAAGACCAGAAATGCTTTTAAAAGACTCTGG
TCTCTGGTCTTCTAATGATGACCTCTTCCTTAATTTTGTGATGGCTTCATCTGCCTCTTTCGATCATCTAGG
TAGCTATAAAAAGCTGAACACCTCCTACTAACCCCATTAGGATGAAAGCGG

Score	Expect	Identities	Gaps	Strand
357 bits(193)	8e-104	193/193(100%)	0/193(0%)	Plus/Plus

```
Query  1    GGCTCGCTTGCTAAAGCTCTCCAGAGAGGCACTTTGAATTAAGACCAGAAATGCTTTTAA  60
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    GGCTCGCTTGCTAAAGCTCTCCAGAGAGGCACTTTGAATTAAGACCAGAAATGCTTTTAA  60

Query  61   AAGACTCTGGTCTCTGGTCTTCTAATGATGACCTCTTCCTTAATTTTGTGATGGCTTCAT  120
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   AAGACTCTGGTCTCTGGTCTTCTAATGATGACCTCTTCCTTAATTTTGTGATGGCTTCAT  120

Query  121  CTGCCTCTTTCGATCATCTAGGTAGCTATAAAAAGCTGAACACCTCCTACTAACCCCATT  180
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  121  CTGCCTCTTTCGATCATCTAGGTAGCTATAAAAAGCTGAACACCTCCTACTAACCCCATT  180

Query  181  AGGATGAAAGCGG  193
            |||||||||||||
Sbjct  181  AGGATGAAAGCGG  193
```

4. **mm4_intergenic_RP11-295D22.1|AL354931.1_chr9_119731666**
   F_GAACTGGCTGCCCAAGGAG; Tm=60.6
   R_CAAACACTCCCCAAGTAGACTG; Tm=58.5

```
guide:       TGTGCCTCTGTTGATCGTCT TGG
off-target:  GGTGGCTCTGATGATCATCT AGG
                 *     *       *    *

CFD Off-target score: 0.401739
MIT Off-target score: 0.34
Position: chr9:119731667-119731689:-
```

hRP11-295D22.1_LRRK2 WT
TCTTTATTTAGCTCTCAGGCTCTACCTATCCAACTGAAACATTTTCTGAGGTAAGGGCTGAGAATGTGTATT
TTTTTTTTTTTAAAAAAAACAACCCTAGGGGGGGTTTTAAAAAAATATAGGTTTGGGAAACTTTTTCCCGGCC
CCCCCTTGGGCCCCCCCTTAAAA

hRP11-295D22.1_LRRK2 KO
TCTTTATTTAGCTCTCAGGCTCTACCTATCCAACTGAAACATTTTCTGAGGTAAGGGCTGAGAATGTGTATT
TTTTTTTTTTTAAAAAAAACAACCCTAGGGGGGGTTTTAAAAAAATATAGGTTTGGGAGACTTTTTCCCGGCC
CCCCCTTGGGCCCCCCCTTAAAA

Score	Expect	Identities	Gaps	Strand
297 bits(329)	1e-85	166/167(99%)	0/167(0%)	Plus/Plus

```
Query  1    TCTTTATTTAGCTCTCAGGCTCTACCTATCCAACTGAAACATTTTCTGAGGTAAGGGCTG  60
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    TCTTTATTTAGCTCTCAGGCTCTACCTATCCAACTGAAACATTTTCTGAGGTAAGGGCTG  60

Query  61   AGAATGTGTATttttttttttttttaaaaaaaaCAACCCTAGGGGGGGTTTTaaaaaaaTATA  120
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   AGAATGTGTATTTTTTTTTTTTTTTAAAAAAAACAACCCTAGGGGGGGTTTTAAAAAAATATA  120

Query  121  GGTTTGGGAGACTTTTTCCCGGccccccccttgggcccccccCTTAAAA  167
            ||||||||| |||||||||||||||||||||||||||||||||||||||
Sbjct  121  GGTTTGGGAAACTTTTTCCCGGCCCCCCCCTTGGGCCCCCCCCTTAAAA  167
```

5. **mm4_intergenic_RN7SL302P|MIR181A2HG_chr9_124658380**
   F_AGTGACTAGAGGCAGCCAGA; Tm=59.9
   R_TCTGCTCAGCTCGCATTGAA; Tm=60.0

```
guide:      TGTGCCTCTGTTGATCGTCT TGG
off-target: TGTGCTTCTGTCCATCATCT GGG
                 *       **      *

CFD Off-target score: 0.345706
MIT Off-target score: 0.06
Position: chr9:124658381-124658403:+
```

nRN7SL302P_LRRK2 WT

GCCTCTGAATGGTGAGGGCTTCACATTTCACCGATCAGGGCACAGCCCAATCAGAAGACAAACCCCTGC
ACTGCCAGGGACCAGCAAATCCGCTTTCTTCCAGTGTGCACATGACTAAATGTGCTTCTGTCCATCATCT
GGGTCATATATAGTTCAATGCGAGCTGAGCAGAA

nRN7SL302P_LRRK2 KO

GCCTCTGAATGGTGAGGGCTTCACATTTCACCGATCAGGGCACAGCCCAATCAGAAGACAAACCCCTGC
ACTGCCAGGGACCAGCAAATCCGCTTTCTTCCAGTGTGCACATGACTAAATGTGCTTCTGTCCATCATCT
GGGTCATATATAGTTCAATGCGAGCTGAGCAGAA

Score	Expect	Identities	Gaps	Strand
320 bits(173)	8e-93	173/173(100%)	0/173(0%)	Plus/Plus

```
Query  1    GCCTCTGAATGGTGAGGGCTTCACATTTCACCGATCAGGGCACAGCCCAATCAGAAGACA  60
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    GCCTCTGAATGGTGAGGGCTTCACATTTCACCGATCAGGGCACAGCCCAATCAGAAGACA  60

Query  61   AACCCCTGCACTGCCAGGGACCAGCAAATCCGCTTTCTTCCAGTGTGCACATGACTAAAT  120
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   AACCCCTGCACTGCCAGGGACCAGCAAATCCGCTTTCTTCCAGTGTGCACATGACTAAAT  120

Query  121  GTGCTTCTGTCCATCATCTGGGTCATATATAGTTCAATGCGAGCTGAGCAGAA  173
            |||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  121  GTGCTTCTGTCCATCATCTGGGTCATATATAGTTCAATGCGAGCTGAGCAGAA  173
```

6. **mm4_intergenic_RP11-354I10.1|OFCC1_chr6_9603371**
   F_AACTCAGGGCTGACATTGGG; Tm=59.9
   R_GCTGACAGCTGTTCTTTCAGA; Tm=58.5

guide: TGTG**C**CTCTGTTGATC**G**TC**T** *TGG*
off-target: **G**GTG**G**CTCTG<u>TTGATC**A**TC**A**</u> *CGG*
            *   *   * *

CFD Off-target score: 0.301304
MIT Off-target score: 0.27
Position: chr6:9603372-9603394:+

hRP11-354_LRRK2 WT
AATCTTAAGACGTTGAATAGTTTACAACCAAGTAAAGTATACTTTTGCAAAAATGGAAGGACTATGTGCTGT
GGCTTATAATTTTCACAGCCGTGATGATCAACAGAGCCACCACCACCAACACCCAGGATCCCTATATTCCA
GGGAATAAATCCCAATGTCAGCCCTGAGTTA

hRP11-354_LRRK2 KO
TTTAATCTTAAGACGTTGAATAGTTTACAACCAAGTAAAGTATACTTTTGCAAAAATGGAAGGACTATGTGC
TGTGGCTTATAATTTTCACAGCCGTGATGATCAACAGAGCCACCACCACCAACACCCAGGATCCCTATATT
CCAGGGAATAAATCCCAATGTCAGCCCTGAGTTA

Score	Expect	Identities	Gaps	Strand
322 bits(174)	2e-93	174/174(100%)	0/174(0%)	Plus/Plus

```
Query  4    AATCTTAAGACGTTGAATAGTTTACAACCAAGTAAAGTATACTTTTGCAAAAATGGAAGG  63
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    AATCTTAAGACGTTGAATAGTTTACAACCAAGTAAAGTATACTTTTGCAAAAATGGAAGG  60

Query  64   ACTATGTGCTGTGGCTTATAATTTTCACAGCCGTGATGATCAACAGAGCCACCACCACCA  123
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   ACTATGTGCTGTGGCTTATAATTTTCACAGCCGTGATGATCAACAGAGCCACCACCACCA  120

Query  124  ACACCCAGGATCCCTATATTCCAGGGAATAAATCCCAATGTCAGCCCTGAGTTA  177
            |||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  121  ACACCCAGGATCCCTATATTCCAGGGAATAAATCCCAATGTCAGCCCTGAGTTA  174
```

7. **mm4_intergenic_RP11-566H8.3|RP11-566H8.1_chr8_31391168**
   F_TTTGCCACCTCTCTTGCAGT; Tm=59.8
   R_CCACAGCTGCTACTATCTGCA; Tm=59.8

```
guide:       TGTGCCTCTGTTGATCGTCT TGG
off-target:  CGTGCCTTTTTTGATCGTTT GGG
                  *       * *              *

CFD Off-target score: 0.281250
MIT Off-target score: 0.49
Position: chr8:31391169-31391191:-
```

hRP11-566H8.3_LRRK2 WT

CATAAATCGGAATTGCAATTCACATGCAGAAGGGAAGAAGCTATCTTACTTGTCATCTCGATGGAAAGTTG
CTTTTGCTTAGGATTTATTCTTTTATCTTTTTTTCCCCTGACTTTCTGAATAGGTGAGGACACAGATTTCAG
CGTGCCTTTTTTGATCGTTTGGGCTTTTTTGATCTTTTTTGTTGTTGCTTTTTCCCTTTATGTTTTAGGTTAA
CTCACTGCAAGAGAGGTGGGCAAAA

hRP11-566H8.3_LRRK2 KO

TGATTTCTGTCCAGATCATAAATCGGAATTGCAATTCACATGCAGAAGGGAAGAAGCTATCTTACTTGTCAT
CTCGATGGAAAGTTGCTTTTGCTTAGGATTTATTCTTTTATCTTTTTTTCCCCTGACTTTCTGAATAGGTGA
GGACACAGATTTCAGCGTGCCTTTTTTGATCGTTTGGGCTTTTTTGATCTTTTTTGTTGTTGCTTTTTCCCT
TTATGTTTTAGGTTAACTCACTGCAAGAGAGGTGG-CAAAA

Score	Expect	Identities	Gaps	Strand
438 bits(237)	5e-128	240/241(99%)	1/241(0%)	Plus/Plus

```
Query  17   CATAAATCGGAATTGCAATTCACATGCAGAAGGGAAGAAGCTATCTTACTTGTCATCTCG   76
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    CATAAATCGGAATTGCAATTCACATGCAGAAGGGAAGAAGCTATCTTACTTGTCATCTCG   60

Query  77   ATGGAAAGTTGCTTTTGCTTAGGATTTATTCTTTTATCttttttttCCCCTGACTTTCTGA   136
            |||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   ATGGAAAGTTGCTTTTGCTTAGGATTTATTCTTTTATCTTTTTTTCCCCTGACTTTCTGA   120

Query  137  ATAGGTGAGGACACAGATTTCAGCGTGCctttttttgatcgtttgggctttttttgatcttt   196
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  121  ATAGGTGAGGACACAGATTTCAGCGTGCCTTTTTTGATCGTTTGGGCTTTTTTGATCTTT   180

Query  197  tttgttgttgcttttttccctttatgttttAGGTTAACTCACTGCAAGAGAGGTGG-CAAA   255
            |||||||||||||||||||||||||||||||||||||||||||||||||||||||| ||||
Sbjct  181  TTTGTTGTTGCTTTTTCCCTTTATGTTTTAGGTTAACTCACTGCAAGAGAGGTGGGCAAA   240

Query  256  A   256
            |
Sbjct  241  A   241
```

8. **mm4_intron_CDKN3_chr14_54414282**
   F_CTCGGATGTCTGTGTCTTTG; Tm=60.0
   R_CAATAAGACAGGAGAGGCAG; Tm=60.0

```
guide:       TGTGCCTCTGTTGATCGTCT  TGG
off-target:  TGTGCCTCGTTTGGTAGTCT  TGG
                      **    * *

CFD Off-target score: 0.234722
MIT Off-target score: 0.02
Position: chr14:54414283-54414305:-
```

hCDKN3_LRRK2 WT
TAGGATGAGGGCCTGCTCTAATTCGTCTTAACTTGAATATACCTGCAAAGACTCAAATTCTGAATAAGGTC
CCACTCACAGGTACAGGATAAGGACTTCAGCATGTCTTTTGGGGGGACATAATTCAACCAGTAACACCTA
CCGTGTACCACACGCCAAGACTACCAAACGAGGCACAGTCCCCACCTTCGAGGAACCTATAACTAACAC
TACCATTTAACTGCAATAAGACAGGAGAAGGCAGA

hCDKN3_LRRK2 KO
GTAGGAACTAGTCTCTAGGATGAGGGCCTGCTCTAATTCGTCTTAACTTGAATATACCTGCAAAGACTCAA
ATTCTGAATAAGGTCCCACTCACAGGTACAGGATAAGGACTTCAGCATGTCTTTTGGGGGGACATAATTC
AACCAGTAACACCTACCGTGTACCACACGCCAAGACTACCAAACGAGGCACAGTCCCCACCTTCGAGGA
ACCTATAACTAACACTACCATTTAACTGCAATAAGACAGG-GAGGGCAGA

Score	Expect	Identities	Gaps	Strand
440 bits(238)	1e-128	243/245(99%)	1/245(0%)	Plus/Plus

```
Query  16   TAGGATGAGGGCCTGCTCTAATTCGTCTTAACTTGAATATACCTGCAAAGACTCAAATTC   75
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  1    TAGGATGAGGGCCTGCTCTAATTCGTCTTAACTTGAATATACCTGCAAAGACTCAAATTC   60

Query  76   TGAATAAGGTCCCACTCACAGGTACAGGATAAGGACTTCAGCATGTCTTTTGGGGGGACA   135
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  61   TGAATAAGGTCCCACTCACAGGTACAGGATAAGGACTTCAGCATGTCTTTTGGGGGGACA   120

Query  136  TAATTCAACCAGTAACACCTACCGTGTACCACACGCCAAGACTACCAAACGAGGCACAGT   195
            ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
Sbjct  121  TAATTCAACCAGTAACACCTACCGTGTACCACACGCCAAGACTACCAAACGAGGCACAGT   180

Query  196  CCCCACCTTCGAGGAACCTATAACTAACACTACCATTTAACTGCAATAAGACAGG-GAGG   254
            |||||||||||||||||||||||||||||||||||||||||||||||||||||||| ||||
Sbjct  181  CCCCACCTTCGAGGAACCTATAACTAACACTACCATTTAACTGCAATAAGACAGGAGAAG   240

Query  255  GCAGA   259
            |||||
Sbjct  241  GCAGA   245
```