Disparities in spatially variable gene calling highlight the need for benchmarking spatial transcriptomics methods.

Natalie Charitakis[1,2,3], Agus Salim[4,5], Adam T. Piers[1,3,6], Kevin I. Watt[1,3,7,8], Enzo R. Porrello[1,2,3,7,*], David A. Elliott[1,2,3,*], Mirana Ramialison[1,2,3,*]

Murdoch Children's Research Institute, Royal Children's Hospital, Flemington Road, Parkville, Victoria, 3052, Australia
Department of Paediatrics, University of Melbourne, Grattan Street, Parkville, Victoria, 3010, Australia
Novo Nordisk Foundation Center for Stem Cell Medicine, Murdoch Children's Research Institute, Royal Children's Hospital, Flemington Road, Parkville, Victoria, 3052, Australia.
Melbourne School of Population and Global Health, University of Melbourne, Bouverie St, Carlton, Victoria, 3053, Australia
School of Mathematics and Statistics, University of Melbourne, Swanston Street, Parkville, Victoria, 3010, Australia
Melbourne Centre for Cardiovascular Genomics and Regenerative Medicine, The Royal Children's Hospital, Melbourne, Victoria, 3052, Australia.
Department of Anatomy and Physiology, University of Melbourne, Grattan Street, Parkville, Victoria, 3010, Australia
Department of Diabetes, Monash University, Alfred Centre, Commercial Road, Melbourne, Victoria, 3004, Australia

* co-senior & co-corresponding authors

Correspondance to:
mirana.ramialison@mcri.edu.au
Mirana Ramialison

Abstract

Identifying spatially variable genes (SVGs) is a key step in the analysis of spatially resolved

transcriptomics (SRT) data. SVGs provide biological insights by defining transcriptomic

differences within tissues, which was previously unachievable using RNA-sequencing

technologies. However, the increasing number of published tools designed to define SVG sets

currently lack benchmarking methods to accurately assess performance. This study compares

results of 6 purpose-built packages for SVG identification across 9 public and 5 simulated

datasets and highlights discrepancies between results. Additional tools for generation of

simulated data and development of benchmarking methods are required to improve methods

for identifying SVGs.

Background

Spatially resolved transcriptomics (SRT) captures variations in gene expression across tissues

(1–7). Computational methods for analysis of SRT data are being established (8–12). Among

the goals of SRT analysis pipelines is the robust and reliable identification of spatially variable

genes (SVGs) within tissue sections (12–14). SVGs are defined as having expression levels

across a tissue that covary in a location specific manner (13,15). Published methods for SVG

identification employ different mathematical models aiming to capture biological truth

(13,14,16–27). Benchmarking of analysis tools is needed to ensure the reliability of processed

data matches or supersedes that of similar technologies such as single-cell (sc) RNA-Seq

(28,29).

Here, we compared the performance of six packages – SpatialDE (13), SPARK-X (16),

Squidpy (30), Seurat (19), SpaGCN (18), scGCO (31) from healthy and cancerous fresh frozen

(FF) and formalin-fixed paraffin embedded (FFPE) tissues generated with 10X Visium

technology alongside simulated datasets (Additional file 1) (32,33). Our analysis reveals

discrepancies in SVGs identified by each package highlighting the urgent need for further

benchmarking and development of tools to identify SVGs.

Results and Discussion

Comparing SVGs predicted by six packages, we found that the numbers of SVGs identified

differed by one to two orders of magnitude across the same tissue for all nine investigated

datasets (Fig. 1A & Additional files 2-4). For example, in the FF endometrial adenocarcinoma

ovarian tissue dataset, the number of predicted SVGs ranged from 87 to 3707 (Fig. 1A). For

7/9 datasets, SpatialDE identifies the greatest number of SVGs, while SpaGCN identifies the

fewest number of SVGs in 6/9 public datasets (Fig. 1A & Additional files 2-4). Apart from the

FF mouse brain coronal section dataset, packages tested tend to report more SVGs in cancerous

datasets than in healthy tissues, most evident across SpatialDE and SPARK-X (Additional file

5). There is no apparent difference in reported SVGs driven by data generated from FF or FFPE

tissue (Additional file 5).

To understand the disparity in the number of SVGs reported, we hypothesised that methods

used to correct for type I error might influence results. SpatialDE, Squidpy, scGCO and

SPARK-X report a q-value/FDR for all genes. Methods that do not supply associated p-values

report the lowest number of SVGs in half of the datasets (Fig. 1A & Additional files 2-4).

Across all datasets, a minimal number of common SVGs are identified between all 6 packages

that each display an expression pattern that aligns with spatial correlation suggesting that they

are *bona fide* SVGs (Fig. 1B).

Next, to assess the significance of small overlap between predicted SVGs, the ranked lists of

SVGs provided by SpatialDE, Squidpy, scGCO and SPARK-X were compared across datasets

using a Wilcoxon signed rank test. We observed differences in the SVG ranking across the

tested packages, indicating that the lack of overlap is statistically significant (Fig. 1C &

Additional files 6-11, Additional file 12: Table S1). q-values for a single gene differ between

packages (Fig. 1C & Additional files 6-11).

We investigated the downstream impact arising from discordant results by comparing gene

ontology (GO) enrichment analysis using SVGs predicted by each package within the same

tissue. (Fig. 1D & Additional file 13). We observed non-overlapping parent terms between

datasets in the top 10 over-represented GO terms (Fig. 1D & Additional file 13). These results

suggest that gene sets predicted by each package do not overlap functionally. Further,

restricting GO analysis to SVGs can bias downstream interpretation of results, especially if

there is a risk of a large percentage of false positives (FP).

Despite the discrepancies, SVGs identified by all packages within the same dataset display an

expression pattern aligned with a clear spatial correlation. An example is *JCHAIN,* which is

expressed in endometrial macrophages and associated with adenocarcinoma (Fig. 1B) (34–36).

Furthermore, in FF mouse brain coronal section, a dataset with known spatial gene expression

patterns, many of the 368 commonly identified SVGs display expression patterns

corresponding to their known region of expression in the Allen Mouse Brain Atlas (Additional

file 13) (37). These observations indicate that patterns with a stronger signal-to-noise ratio are

identified by all methods.

To gain insight into the inconsistent results between different packages we evaluated their

sensitivity and specificity. To determine true positives and FP rates, we first generated two

simulated datasets with different known SVG patterns with SRTsim (Additional file 15: A-B,

Supplementary methods) (38). Each dataset contained 1500 known SVGs (Supplementary

methods). Using each as the input to four packages (scGCO, SpatialDE, Squidpy and SPARK-

X), all packages identify all true positives across datasets, except for Squidpy when analysing

the dataset with lower signal SVGs. This indicates a potential need for a more relaxed Moran's

I statistic threshold (Additional file 12: Table S2). Additionally, most packages introduce

minimal levels of FPs with simulated datasets, except for scGCO (Additional file 12: Table

S2). This is possibly due to of the type I error correction method or sequential search procedure

employed by scGCO (31).

To further investigate the potential of each package to call FPs, we generated four negative

control datasets from two publicly available datasets. For each, two versions of negative

controls were generated - [1] where spatial coordinates are permutated and [2] where spatial

coordinates and columns of gene expression matrix are permutated. For all negative control

datasets generated from the FF left ventricle, Seurat, scGCO, Squidpy and SPARK-X identified

no SVGs, indicating an ability to distinguish signal from noise. SpaGCN identified <10 SVGs.

SpatialDE identified 4 SVGs when both spatial coordinates and columns of the gene expression

matrix are permutated and 2953 in the other dataset. For negative control datasets generated

from FFPE prostate randomising only spatial coordinates, SpatialDE identified 632 SVGs,

SpaGCN identified 232 and Squidpy identified 1. Seurat, scGCO and SPARK-X did not run

with these inputs. When spatial coordinates and gene expression values were randomised,

SpatialDE was the only package to identify SVGs (713). Patterns of the top SVGs identified

by SpatialDE lack a strong spatial pattern (Additional file 16). This would indicate that

SpatialDE is prone to introducing FPs into a dataset when genes are lowly expressed compared.

However, the original SVGs identified by SpatialDE and the result in Fig. 1B suggest the top

ranked SVGs labelled by SpatialDE in real data display a signal consistent with that of a SVG.

Inconsistencies in the results between different packages may therefore be explained by the

differences in FP rates, rather than the rate of false negatives. To investigate this, expression

of known housekeeping gene *Eef2* (39) was visualised across the FF mouse brain coronal

Section dataset to verify its ubiquitous expression (Additional file 14: D). *Eef2* was found to

be annotated as an SVG by scGCO, SPARK-X, SpatialDE and Squidpy. This indicates genes

with high levels of expression, but that do not display strong spatial expression patterns, may

be a source of confounding.

In addition to differences in FP rates, results generated from different packages applied to the

same dataset may arise due to the different assumptions regarding the underlying distributions

of gene expression and effectiveness of normalisation methods (Additional file 12: Table S3).

Negative binomial distributions can successfully model sequencing-based methods that

employ unique molecular identifiers (UMIs) and most gene expression patterns in Visium

datasets (38,40). While certain packages implement their own normalisation method, there is

no guarantee that data pre-processed using a separate workflow will have been normalised to

fit the assumptions of gene distributions held by a particular package (13,19). Once methods

for generating simulated SRT datasets with different underlying distributions of gene

expression are published, the effect this has on identifying SVGs needs to be investigated

thoroughly. There are many caveats to generating simulated datasets with known patterns of

gene expression with efforts focused on simulating data with defined cell types (41–48).

Conclusions

While different mathematical models for detecting SVGs produce varied results when applied to the same tissue, and across tissues and biological conditions, SVGs with a strong spatial covariance are consistently identified. This indicates that differences between methods do not bias performance in a particular tissue or biological condition. However, more transcriptionally heterogenous tissues appear to affect the performance of certain packages in calling SVGs and genes with unclear spatial patterns but high expression levels may be labelled as FP (14,15). Only four of the tested packages produced q-value ranked gene lists, thus differences in FP rates could confound comparisons of results across packages. Furthermore, due to the resolution of the Visium technology platform the presence of heterogenous cell types could complicate accurate identification of SVGs due to inherent transcriptional variation. Increased resolution may present challenges in terms of computational performance and the mathematical difficulties of dealing with highly sparse datasets. As SVG identification becomes an integral part of the computational analysis of SRT and datasets grow, a subset of SVGs could be used in preference over the entire dataset for downstream analysis (such as highly variable genes in scRNA-Seq analysis). Introducing noise may hamper key downstream steps in understanding spatially restricted disease states as well as novel targets for treatment (49,50). Methods that explore the entire dataset for SVG labelling paired with stricter q-value cut-offs (q-value < 0.01) can be employed when analysing tissues with known transcriptional complexity to decrease the possibility of FP being reported. Future work will include comparison of package performance across datasets generated from different SRT platforms. The development of workflows for simulating datasets and benchmarking current and novel methods for SVG identification will allow for accurate determination of the spatially restricted transcriptional differences that manifest as biological outcomes during development and disease.

Methods

Six packages were selected for comparison: SpatialDE (13), SPARK-X (16), Squidpy (30),
Seurat (19), SpaGCN (18) and scGCO (31), which each employ a different mathematical model
to identify SVGs. Each package was tested on 9 publicly available, V1 Chemistry Visium
datasets generated from healthy and cancerous tissues, most human and one mouse (Additional
file: Table S4). The filtered output files from the Space Ranger v1.0.0 pipeline were
downloaded, and pre-processed using Scanpy v1.8.1 (51). Two reference-free simulated
datasets with features similar to Visium datasets were generated using SRTsim (38). Using the
FF left ventricle and FFPE prostate datasets as inputs, two negative control datasets were
generated in each instance by 1) independently randomising the x and y spot coordinates to
remove spatial correlation and 2) additionally randomising each column in the gene expression
matrix to ensure any overall association was removed.

References

1. Marx V. Method of the Year: spatially resolved transcriptomics. Nat Methods. 2021;18(1):9–14.
2. Asp M, Bergenstråhle J, Lundeberg J. Spatially Resolved Transcriptomes-Next Generation Tools for Tissue Exploration. Bioessays. 2020;42(10).
3. Waylen LN, Nim HT, Martelotto LG, Ramialison M. From whole-mount to single-cell spatial assessment of gene expression in 3D. Communications Biology. 2020;3(1):1–11.
4. Ståhl PL, Salmén F, Vickovic S, Lundmark A, Navarro JF, Magnusson J, et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. Vol. 353, Science. American Association for the Advancement of Science; 2016. p. 78–82.
5. Larsson L, Frisén J, Lundeberg J. Spatially resolved transcriptomics adds a new dimension to genomics. Vol. 18, Nature Methods. Nature Research; 2021. p. 15–8.
6. Moses L, Pachter L. Museum of spatial transcriptomics. Nature Methods. 2022;19(5):534–46.
7. Williams CG, Lee HJ, Asatsuma T, Vento-Tormo R, Haque A. An introduction to spatial transcriptomics for biomedical research. Genome Medicine. 2022;14(1):1–18.
8. Zeng Z, Li Y, Li Y, Luo Y. Statistical and machine learning methods for spatially resolved transcriptomics data analysis. Genome Biology 2022;23(1):1–23.

184	9.	Noel T, Wang QS, Greka A, Marshall JL. Principles of Spatial Transcriptomics
185		Analysis: A Practical Walk-Through in Kidney Tissue. Front Physiol. 2022 Jan
186		6;12:2317.
187	10.	Liu B, Li Y, Zhang L. Analysis and Visualization of Spatial Transcriptomic Data.
188		Front Genet. 2022 Jan 27;0:2852.
189	11.	Dries R, Chen J, Rossi N del, Khan MM, Sistig A, Yuan GC. Advances in spatial
190		transcriptomic data analysis. Genome Res. 2021;31(10):1706–18.
191	12.	Du J, Yang YC, An ZJ, Zhang MH, Fu XH, Huang ZF, et al. Advances in spatial
192		transcriptomics and related data analysis strategies. J Transl Med. 2023;21(1):330.
193	13.	Svensson V, Teichmann SA, Stegle O. SpatialDE: Identification of spatially variable
194		genes. Nat Methods. 2018;15(5).
195	14.	Li K, Yan C, Li C, Chen L, Zhao J, Zhang Z, et al. Computational elucidation of
196		spatial gene expression variation from spatially resolved transcriptomics data. Mol
197		Ther Nucleic Acids. 2022 Mar 8;27:404–11.
198	15.	Walker BL, Cang Z, Ren H, Bourgain-Chang E, Nie Q. Deciphering tissue structure
199		and function using spatial transcriptomics. Communications Biology 2022 5:1. 2022
200		Mar 10;5(1):1–10.
201	16.	Zhu J, Sun S, Zhou X. SPARK-X: non-parametric modeling enables scalable and
202		robust detection of spatial expression patterns for large spatial transcriptomic studies.
203		Genome Biol. 2021;22(1):1–25.
204	17.	Zhang K, Feng W, Wang P. Identification of spatially variable genes with graph cuts
205		[Internet]. bioRxiv. bioRxiv; 2018 [cited 2021 Feb 5]. p. 491472. Available from:
206		https://doi.org/10.1101/491472
207	18.	Hu J, Li X, Coleman K, Schroeder A, Ma N, Irwin DJ, et al. SpaGCN: Integrating
208		gene expression, spatial location and histology to identify spatial domains and spatially
209		variable genes by graph convolutional network. Nature Methods. 2021;18(11):1342–
210		51.
211	19.	Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-
212		cell gene expression data. Nat Biotechnol. 2015; 33(5):495–502.
213	20.	Xu Y, McCord RP. CoSTA: unsupervised convolutional neural network learning for
214		spatial transcriptomics analysis. BMC Bioinformatics. 2021;22(1):1–26.
215	21.	Hao M, Hua K, Zhang X. SOMDE: A scalable method for identifying spatially
216		variable genes with self-organizing map. bioRxiv. 2021
217	22.	Weber LM, Saha A, Datta A, Hansen KD, Hicks SC. nnSVG: scalable identification of
218		spatially variable genes using nearest-neighbor Gaussian processes. bioRxiv. 2022
219	23.	Sun S, Zhu J, Zhou X. Statistical analysis of spatial expression patterns for spatially
220		resolved transcriptomic studies. Nature Methods 2020;17(2):193–200.
221	24.	Jiang X, Xiao G, Li Q. A Bayesian modified Ising model for identifying spatially
222		variable genes from spatial transcriptomics data. Stat Med. 2022.
223	25.	Liu Q, Hsu CY, Shyr Y. Scalable and model-free detection of spatial patterns and
224		colocalization. Genome Res. 2022;32(9):gr.276851.122.
225	26.	Li K, Yan C, Li C, Chen L, Zhao J, Zhang Z, et al. Computational elucidation of
226		spatial gene expression variation from spatially resolved transcriptomics data. Mol
227		Ther Nucleic Acids. 2022;27:404–11.
228	27.	Yu J, Luo X. Identification of cell-type-specific spatially variable genes accounting for
229		excess zeros. Bioinformatics. 2022;38(17):4135–44.
230	28.	Adil A, Kumar V, Jan AT, Asger M. Single-Cell Transcriptomics: Current Methods
231		and Challenges in Data Acquisition and Analysis. Front Neurosci. 2021;15:398.
232	29.	Chen G, Ning B, Shi T. Single-cell RNA-seq technologies and related computational
233		data analysis. Front Genet. 2019;317.

234   30.   Palla G, Spitzer H, Klein M, Fischer D, Schaar AC, Kuemmerle LB, et al. Squidpy: a
235   scalable framework for spatial omics analysis. Nature Methods. 2022;19(2):171–8.
236   31.   Zhang K, Feng W, Wang P. Identification of spatially variable genes with graph cuts.
237   Nature Communications. 2022;13(1):1–15.
238   32.   Palla G, Fischer DS, Regev A, Theis FJ. Spatial components of molecular tissue
239   biology. Nature Biotechnology. 2022;40(3):308–18.
240   33.   Li K, Yan C, Li C, Chen L, Zhao J, Zhang Z, et al. Computational elucidation of
241   spatial gene expression variation from spatially resolved transcriptomics data. Mol
242   Ther Nucleic Acids. 2022;27.
243   34.   Dun EC, Hanley K, Wieser F, Bohman S, Yu J, Taylor RN. Infiltration of tumor-
244   associated macrophages is increased in the epithelial and stromal compartments of
245   endometrial carcinomas. International Journal of Gynecological Pathology.
246   2013;32(6):576–84.
247   35.   The Human Protein Atlas. Available from: https://www.proteinatlas.org/
248   36.   Wang W, Vilella F, Alama P, Moreno I, Mignardi M, Isakova A, et al. Single-cell
249   transcriptomic atlas of the human endometrium during the menstrual cycle. Nature
250   Medicine 2020;26(10):1644–53.
251   37.   Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, Bernard A, et al. Genome-
252   wide atlas of gene expression in the adult mouse brain. Nature. 2006;445(7124):168–
253   76.
254   38.   Zhu J, Shang L, Zhou X. SRTsim: spatial pattern preserving simulations for spatially
255   resolved transcriptomics. 2023,24(1):39.
256   39.   Kouadjo KE, Nishida Y, Cadrin-Girard JF, Yoshioka M, St-Amand J. Housekeeping
257   and tissue-specific genes in mouse tissues. BMC Genomics. 2007;8(1):1–16.
258   40.   Svensson V. Droplet scRNA-seq is not zero-inflated. Nature Biotechnology.
259   2020;38(2):147–50.
260   41.   Ma Y, Zhou X. Spatially informed cell-type deconvolution for spatial transcriptomics.
261   Nat Biotechnol. 2022;
262   42.   Miller BF, Huang F, Atta L, Sahoo A, Fan J. Reference-free cell type deconvolution of
263   multi-cellular pixel-resolution spatially resolved transcriptomics data. Nature
264   Communications 2022;13(1):1–13.
265   43.   Dong R, Yuan GC. SpatialDWLS: accurate deconvolution of spatial transcriptomic
266   data. Genome Biol. 2021;22(1):1–10.
267   44.   Cable DM, Murray E, Zou LS, Goeva A, Macosko EZ, Chen F, et al. Robust
268   decomposition of cell type mixtures in spatial transcriptomics. Nature Biotechnology
269   2021;40(4):517–26.
270   45.   Elosua-Bayes M, Nieto P, Mereu E, Gut I, Heyn H. SPOTlight: seeded NMF
271   regression to deconvolute spatial transcriptomics spots with single-cell transcriptomes.
272   Nucleic Acids Res. 2021;49(9):e50–e50.
273   46.   Kleshchevnikov V, Shmatko A, Dann E, Aivazidis A, King HW, Li T, et al.
274   Cell2location maps fine-grained cell types in spatial transcriptomics. Nature
275   Biotechnology. 2022;40(5):661–71.
276   47.   Bae S, Choi H, Lee DS. spSeudoMap: Cell type mapping of spatial transcriptomics
277   using unmatched single-cell RNA-seq data. bioRxiv. 2022;
278   48.   Song Q, Su J. DSTG: deconvoluting spatial transcriptomics data through graph-based
279   artificial intelligence. Brief Bioinform. 2021;22(5):1–13.
280   49.   Navarro JF, Croteau DL, Jurek A, Andrusivova Z, Yang B, Wang Y, et al. Spatial
281   Transcriptomics Reveals Genes Associated with Dysregulated Mitochondrial
282   Functions and Stress Signaling in Alzheimer Disease. iScience. 2020;23(10):101556.

283   50.   Wang Y, Ma S, Ruzzo WL. Spatial modeling of prostate cancer metabolic gene
284   expression reveals extensive heterogeneity and selective vulnerabilities. Scientific
285   Reports. 2020;10(1):1–14.
286   51.   Wolf FA, Angerer P, Theis FJ. SCANPY: Large-scale single-cell gene expression data
287   analysis. Genome Biol. 2018;19(1):15.
288   52.   Hoffman P, Satija Lab. Analysis, visualization, and integration of spatial datasets with
289   Seurat • Seurat. 2023

Figure Legends

Figure 1. Discrepancies between SVGs in a dataset annotated by six different packages.

A) Upset plots displaying the distinct intersections of SVGs identified in FF endometrial

adenocarcinoma ovarian tissue dataset when analysed with SpaGCN, Squidpy, scGCO, Seurat,

SPARK-X and SpatialDE. **B)** Pattern of expression of *JCHAIN*, a SVG identified by all six

packages in FF endometrial adenocarcinoma ovarian tissue. Spots representing the capture

regions of the Visium slide are overlayed on accompanying histological image. **C)** Sorted q-

values from SpatialDE results across the FF left ventricle datasets plotted against SPARK-X

q-values for the same gene. **D)** Gene ontology enrichment results from FF endometrial

adenocarcinoma ovarian tissue dataset using SVGs identified by each package as inputs.

Figure 2. **Expression patterns of SVGs identified by all tested packages in a single dataset.**

A) Pattern of expression of *ALB*, a SVG identified by all six packages in FF invasive ductal

carcinoma breast tissue. **B)** Pattern of expression of *ACTA1*, a SVG identified by all six

packages in FF left ventricle. **C)** Pattern of expression of *BTG2*, a SVG identified by all six

packages in FFPE prostate. **D)** Pattern of expression of *TACSTD2*, a SVG identified by all six

packages in FFPE invasive ductal carcinoma breast tissue.

Declarations

Ethic approval and consent to participate

NA

Consent for publication

NA

Availability of data and materials

The input files for the Visium data used in this study are available from the [10X Genomics](#)

[website](#). Examples of the data pre-processing and analysis to identify SVGs on a single dataset

for each of the six packages can be found on Github at the repository Ramialison-
Lab/Disparities_in_SVG_calling.

318
Additional file 1: additional_file1.pdf

Overview of the design of benchmarking workflow. First, publicly available datasets are
obtained to be used as inputs for pre-processing in a Scanpy workflow. For each dataset the
processed output is then transformed into a data object most suitable for each of the SVG
analysis packages. Simulated data is directly used as inputs for SVG analysis. Finally, the
results are compared across packages within each dataset.

325
326 Additional file 2: additional_file2.pdf

327 Distinct overlap of SVGs identified by different combinations of the six tested packages. A)

328 FF cerebellum. B) FF lymph node. C) FFPE adenocarcinoma prostate.

329
330 Additional file 3: additional_file3.pdf

331 Distinct overlap of SVGs identified by different combinations of the six tested packages. A)

332 FF invasive ductal carcinoma breast tissue. B) FF left ventricle. C) FFPE prostate.

333
334 Additional file 4: additional_file4.pdf

335 Distinct overlap of SVGs identified by different combinations of the six tested packages. A)

336 FFPE invasive ductal carcinoma breast tissue. B) FF mouse brain coronal section.

337
Additional file 5: additional_file5.pdf

Comparison of the number of SVGs identified by six different packages across different
datasets. FFPE tissues are visualised by a cross and FF tissues are visualised with a square.
High numbers of SVGs are identified in both FF and FFPE tissues, however tissues that are
more transcriptionally complex seem to have more SVGs called across the dataset. Seurat
reports a consistent number of SVGs across datasets as it first identifies highly variable genes
then ranks their expression by how dependent it is on spatial location (52).

Additional file 6: additional_file6.pdf

Comparison of ranked SpatialDE q-values against gene-matched SPARK-X q-values of all

genes generated from each dataset. Order of plots repeats across all datasets. A) FF cerebellum.

B) FF lymph node. C) FFPE adenocarcinoma prostate. D) FF invasive ductal carcinoma breast

tissue. E) FFPE prostate. F) FFPE invasive ductal carcinoma breast tissue. G) FF endometrial

adenocarcinoma ovarian tissue. H) FF mouse brain coronal section.

Additional file 7: additional_file7.pdf

Comparison of ranked SpatialDE q-values against gene-matched scGCO q-values of all genes

generated from each dataset. Order of plots repeats across all datasets. A) FF cerebellum. B)

FF lymph node. C) FFPE adenocarcinoma prostate. D) FF invasive ductal carcinoma breast

tissue. E) FFPE prostate. F) FFPE invasive ductal carcinoma breast tissue. G) FF endometrial

adenocarcinoma ovarian tissue. H) FF left ventricle. I) FF mouse brain coronal section.

Additional file 8: additional_file8.pdf

Comparison of ranked SPARK-X q-values against gene-matched scGCO q-values of all genes

generated from each dataset. Order of plots repeats across all datasets. A) FF cerebellum. B)

FF lymph node. C) FFPE adenocarcinoma prostate. D) FF invasive ductal carcinoma breast

tissue. E) FFPE prostate. F) FFPE invasive ductal carcinoma breast tissue. G) FF endometrial

adenocarcinoma ovarian tissue. H) FF left ventricle. I) FF mouse brain coronal section.

Additional file 9: additional_file9.pdf

Comparison of ranked SpatialDE q-values against gene-matched Squidpy q-values of all genes

generated from each dataset. Order of plots repeats across all datasets. A) FF cerebellum. B)

FF lymph node. C) FFPE adenocarcinoma prostate. D) FF invasive ductal carcinoma breast

tissue. E) FFPE prostate. F) FFPE invasive ductal carcinoma breast tissue. G) FF endometrial

adenocarcinoma ovarian tissue. H) FF left ventricle. I) FF mouse brain coronal section.

Additional file 10: additional_file10.pdf

Comparison of ranked SPARK-X q-values against gene-matched Squidpy q-values of all genes

generated from each dataset. Order of plots repeats across all datasets. A) FF cerebellum. B)

FF lymph node. C) FFPE adenocarcinoma prostate. D) FF invasive ductal carcinoma breast

tissue. E) FFPE prostate. F) FFPE invasive ductal carcinoma breast tissue. G) FF endometrial

adenocarcinoma ovarian tissue. H) FF left ventricle. I) FF mouse brain coronal section.

Additional file 11: additional_file11.pdf

Comparison of ranked scGCO q-values against gene-matched Squidpy q-values of all genes

generated from each dataset. Order of plots repeats across all datasets. A) FF cerebellum. B)

FF lymph node. C) FFPE adenocarcinoma prostate. D) FF invasive ductal carcinoma breast

tissue. E) FFPE prostate. F) FFPE invasive ductal carcinoma breast tissue. G) FF endometrial

adenocarcinoma ovarian tissue. H) FF left ventricle. I) FF mouse brain coronal section.

Additional file 12: additional_file12.pdf

Contains Tables S1-4. Table S1) Reported statistic and associated p-value from each

combination of pairwise comparison of results between SpatialDE, SPARK-X and scGCO

when reported p-values were compared using the Wilcoxon signed rank test. Table S2)

Sensitivity and specificity of scGCO, Squidpy, SPARK-X and SpatialDE when used to analyse

both simulated datasets generated with SRTsim. Sensitivity and specificity are reported over a

range of q-value cut-offs, from 0.01-1.0. Table S3) Packages included for identification of

SVGs in benchmarking study. The packages are ordered by assumptions made on distribution

of gene expression. The rows highlighted in blue indicate grouped packages using graph-based

methods. Table S4) Overview of publicly available 10X Visium datasets to be included in

benchmarking process. FF indicates tissues that are fresh frozen and FFPE indicates tissues

that are formalin-fixed paraffin-embedded. The filtered output files and imaging data from the

Space Ranger v1.0.0, v1.2.0 or V1.3.0 pipeline were downloaded for each dataset.

Additional file 13: additional_file13.pdf

Gene ontology enrichment results using SVGs identified by each package as inputs across

datasets. A) FF cerebellum. B) FF lymph node. C) FFPE adenocarcinoma prostate. D) FF

invasive ductal carcinoma breast tissue. E) FF left ventricle. F) FFPE prostate. G) FFPE

invasive ductal carcinoma breast tissue. H) FF mouse brain coronal Section.

Additional file 14: additional_file14.pdf

Spatial expression patterns of SVGs identified by all packages across the FF mouse brain

coronal section dataset. A) Expression of *Hap1* across the hypothalamus and amygdala, cross-

referenced with the Allen Mouse Brain Reference. B) Expression of *Prkcd* localised to the

thalamus, cross-referenced with the Allen Mouse Brain Reference. C) Expression of *Itpka*, with

highest expression in the isocortex, hippocampal formation (HPF) and cortical subplate

consistent with patterns displayed in the Allen Mouse Brain Reference. D) Expression of *Eef2*,

a known housekeeping gene in mouse (39).

Additional file 15: additional_file15.pdf

Simulated datasets generated with SRT sim. A) Location of simulated SVGs with a hotspot

pattern visualised in blue, while red area indicates expression of noise genes. B) Location of

simulated SVGs in both blue and green corners, while red area indicates expression of noise

genes. C) Distinct overlap of SVGs compared to the control SVG list identified by different

combinations of the four tested packages. 1500 SVGs were present in this dataset. D) Distinct

overlap of SVGs compared to the control SVG list identified by different combinations of the

four tested packages. 750 high signal SVGs and 750 low signal SVGs were present in this

dataset.

Additional file 16: additional_file16.pdf

Top SVG identified by SpatialDE in negative simulated datasets generated by A) Randomising

coordinates of FF Left Ventricle dataset. B) Randomising counts and coordinates of FF left

ventricle dataset. C) Randomising coordinates of FFPE prostate dataset. D) Randomising

counts and coordinates of FFPE prostate dataset.

Additional file 17: additional_file17.pdf

Upset plot of results generated from running SPARK-X and SpatialDE on simulated data with

known pattern of SVGs.

Additional file 18: additional_file18.xlsx

Supplementary methods

Competing Interests

ERP is a cofounder, scientific advisors, and holds equity in Dynomics, a biotechnology

company focused on the development of heart failure therapeutics. The other authors report no

conflicts.

Funding

The authors acknowledge grant and fellowship support from the National Health and Medical

Research Council of Australia (E.R.P., D.A.E.), the Stafford Fox Medical Research Foundation

(E.R.P, D.A.E.), and the Royal Children's Hospital Foundation (E.R.P., D.A.E.). The Novo

Nordisk Foundation Center for Stem Cell Medicine (E.R.P., D.A.E., M.R.) is supported by

Novo Nordisk Foundation grants (NNF21CC0073729). The Murdoch Children's Research

Institute is supported by the Victorian Government's Operational Infrastructure Support

Program.

Authors' contributions

N.C. performed experiments with contribution from A.S. on statistical design; N.C., A.S.,

A.T.P., K.I.W, E.R.P., D.A.E., M.R. designed experiments; N.C., K.I.W, D.A.E., M.R. wrote

the manuscript with input from A.S. and E.R.P.

Acknowledgements

The authors would like to thank Matthew Ritchie, David Powell and Gordon Smyth for their

contributions of ideas to include in the analysis, Hieu Nim for manuscript feedback, Michael

See for use of his code in generating plots and members of Transcriptomics and Bioinformatics

and Heart Regeneration and Disease laboratories for their feedback.