Conditional blastocyst complementation of a defective Foxa2 lineage efficiently promotes generation of the whole lung

generation of the whole lung

Akihiro Miura[1, 2], Hemanta Sarmah[1], Junichi Tanaka[1], Youngmin Hwang[1], Anri Sawada, [1]

Yuko Shimamura[1], Yinshan Fang[1], Dai Shimizu[2], Zurab Ninish[1], Jake Le Suer[3, 4], Nicole C.

Dubois[5], Jennifer Davis[6], Shinichi Toyooka[2], Jun Wu[7], Jianwen Que[1], Finn J. Hawkins[3, 4],

Chyuan-Sheng Lin[8], Munemasa Mori[1] [†]

1 Columbia Center for Human Development and Division of Pulmonary, Allergy, Critical Care,

Department of Medicine, Columbia University Medical Center, New York, NY 10032, USA.

2 Department of Thoracic, Breast and Endocrinological Surgery, Okayama University Graduate

School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, 7008558, Japan.

3 The Pulmonary Center and Department of Medicine, Boston University School of Medicine,

Boston, MA 02118, USA

4 Center for Regenerative Medicine, Boston University and Boston Medical Center, Boston, MA

02118, USA

5 Department of Cell, Developmental and Regenerative Biology, Icahn School of Medicine at

Mount Sinai, New York, NY 10029, USA.

6 Department of Pathology, University of Washington, Seattle, WA 98109, USA.

7 Department of Molecular Biology, University of Texas Southwestern Medical Center, Dallas,

TX, USA

8 Bernard and Shirlee Brown Glaucoma Laboratory, Department of Pathology and Cell Biology,

College of Physicians and Surgeons, Columbia University Irving Medical Center, New York, NY

10032, USA.

† To whom correspondence should be addressed:

Munemasa Mori, M.D., Ph.D.

650W 168th St, New York, NY 10032, USA

E-mail: mm4452@cumc.columbia.edu

Abstract

Millions suffer from incurable lung diseases, and the donor lung shortage hampers organ

transplants. Identifying the crucial lineage and the program for lung organogenesis could facilitate

designing whole-lung bioengineering. Using lineage-tracing mice and human iPSC-derived lung-

directed differentiation, we revealed that gastrulating Foxa2 lineage contributed to both lung

mesenchyme and epithelium formation. Interestingly, Foxa2 lineage-derived cells in the lung

mesenchyme progressively increased and occupied more than half of the mesenchyme niche,

including endothelial cells, during lung development. Foxa2 promoter-driven, conditional Fgfr2

gene depletion caused the lung agenesis phenotype in mice. Importantly, wild-type donor mouse

iPSCs injected into their blastocysts rescued this phenotype by complementing the Fgfr2-defective

niche in the lung epithelium and mesenchyme. Donor cell is shown to replace the entire lung

epithelial and robust mesenchymal niche during early chimeric lung development, resulting in

efficient complementation of the nearly entire lung niche at the late stage of lung development.

These results suggest that lung complementation based on the Foxa2 lineage is a unique model for

the progressive mobilization of donor cells into both epithelial and mesenchymal lung niches and

provides crucial insights for designing new bioengineering strategies to generate whole lungs.

KEYWORDS

Lung generation

Conditional blastocyst complementation

Lung mesenchyme precursors

Mesendoderm

Lung development

Whole organ generation

INTRODUCTION

Tissue regeneration to treat various intractable diseases has long been challenging(Hackett et al., 2010; Kemter et al., 2020; Ott et al., 2010; Petersen et al., 2010; Wang, 2019). Organ bioengineering strategy based on recellularizing tissue-specific progenitors into the decellularized scaffolds, induced pluripotent stem cell (iPSC)-derived organoids, or 3d-bioprinters are the next-generation tissue transplant therapies(Guyette et al., 2014; Kotton and Morrisey, 2014; Petersen et al., 2010; Tian et al., 2021). Even with these techniques, however, the mammalian lung is one of the most difficult organs to replicate because of its anatomical complexity and cellular diversity. It contains hundreds of airway branches and a thin micron-sized alveolar layer of inflated and well-vascularized alveoli composed of billions of cells from more than 50 different cell types(Crapo et al., 1982; Kotton and Morrisey, 2014; Stone et al., 1992; Travaglini et al., 2020). Donor organs for lung transplantation are in short supply worldwide, but the technology does not exist to generate whole lungs composed of tissue-specific epithelial and mesenchymal cells, including endothelial cells. Lungs grow fully only through natural lung development.

During development, lung epithelial and mesenchymal precursors interact to initiate an elaborate developmental program of organogenesis that includes specification, pattern formation, progenitor cell expansion, and differentiation. The lung epithelial cells are derived from the foregut, definitive endoderm (DE) derivatives, classically labeled by Sox17 and Forkhead Box A2 (Foxa2)(Green et al., 2011; Huang et al., 2014). Multiple genetic studies using Sonic Hedgehog (Shh) Cre lineage-tracing mice have also shown that the entire Nkx2-1+ lung and tracheal epithelial primordium arises from Shh+ DE.

The lung mesenchyme primordium is derived from Wnt2$^+$ Isl1$^+$ cardiopulmonary

progenitors (CPP). CPP is the derivative of Osr1$^+$ Nkx6-1$^+$Barx1$^-$ Wnt4^{low} foregut lung mesoderm

that arises from lateral plate mesoderm (LPM)(Han et al., 2020). While DE and LPM arise from

primitive streaks (PS) during gastrulation, the exact lineage origin of LPM has been a complete

mystery.

Mesendoderm is a bipotent transitional state between the PS and nascent mesoderm labeled

by Mixl1, Pdgfrα, and Brachyury (T) during gastrulation that can give rise to both DE and

mesoderm(Hart et al., 2002; Tada et al., 2005). Although it was speculated that mesendoderm

might form LPM and DE, there have been no conclusive genetic studies on whether mesendoderm

gives rise to both lung epithelium and mesenchyme. Pdgfrα is expressed in the epiblast-derived

mesendoderm, the primitive endoderm (PrE), and its extra-embryonic endoderm derivatives, such

as parietal and visceral endoderm, around E5.5~E7.5. Foxa2 plays a pivotal role in alveolarization

and airway goblet cell expansion(Wan et al., 2004), while there was a significant knowledge gap

regarding Foxa2 lineage during lung development.

Blastocyst complementation (BC) has been proposed as a promising option for tissue-

specific niche complementation(Chen et al., 1993). This unique technology has been further

developed into intra- and interspecies organ generations such as kidney, pancreas, and blood

vessels(Hamanaka et al., 2018; Kobayashi et al., 2010; Usui et al., 2012; Yamaguchi et al., 2017).

However, the production of entire organs, including tissue-specific epithelium and mesenchyme,

including endothelium, was still difficult because endothelium significantly impacts the other

organs. Unfortunately, even with BC, the lungs produced were non-functional and very inefficient,

and in addition, the chimeric lungs contained a substantial amount of host-derived tissue(Kitahara

et al., 2020; Li et al., 2021; Wen et al., 2021). Previously, we established the conditional blastocyst

complementation (CBC) approach, which targets specific lineages complemented by donor

pluripotent stem cells(Mori et al., 2019). Using lineage-specific drivers of lung endoderm in CBCs

avoids the effects of genetic manipulation in non-target organs for the generation of empty organ

niches that lead to functional chimeric lung generation(Mori et al., 2019). However, most of the

lung mesenchyme and endothelium were still derived mainly from the host cells, which was the

severe limitation of CBC(Mori et al., 2019). Given that the CBC approach targeted endodermal

lungs, we speculated that this limitation was due to a significant gap in our knowledge of the origin

of all lung cell types, especially pulmonary mesenchyme, including endothelium. In particular, the

complementation of endothelium is a critical issue for overcoming hyperacute rejection after lung

transplantation. To overcome this critical issue, we explored the origin and the program of whole

lung epithelium and mesenchyme, the major components of the lung.

We hypothesized that targeting a single bona fide lung generative lineage (BFL) may

facilitate the designing of the entire lung generation. BFL is based on the strategy for vacating the

desired niche by lineage-specific gene depletion to be complemented by donor cells after injecting

them into the host blastocysts. The lineage is not necessarily lung-specific but rather based on

emptying or mitotically defective in the desired lineage by gene depletion.

We found that the Foxa2 lineage functions as a BFL in CBC. Although the Foxa2 lineage

did not label the entire lung cells during lung development, the mitotically-defective Foxa2 lineage

promoted efficient whole-lung complementation by donor cells.

RESULTS

Pdgfrα⁺ lineage during gastrulation gives rise to the entire lung mesenchyme.

To determine the origin of LPM and pulmonary endothelium for the whole lung generation

via BFL, we performed lung mesenchyme precursor lineage-tracing analysis using $Pdgfra^{CreERT2/+}$;

$Rosa^{tdTomato/+}$ mice. Surprisingly, tamoxifen injection at E5.5 labeled the entire lung mesenchyme

with tdTomato at E14.5 (Figures 1A and 1B). This result suggested that the origin of the whole

lung mesenchyme is the Pdgfrα lineage around early-to-mid-streak-stage embryos. tdTomato

labeled the entire pulmonary mesenchyme, including Sma$^+$ airway smooth muscle cells, Pdgfrβ$^+$

pulmonary mesenchyme, and VE-cadherin$^+$ vascular endothelial cells (Figure 1B). In addition, the

Pdgfrα lineage only partially labels the lung epithelium (Figure 1B, arrows), suggesting that the

contribution of the Pdgfrα lineage to the lung endoderm is low. It suggests that Pdgfrα is difficult

to define as a BFL because of its low contribution to the lung epithelium.

136   Foxa2 lineage labels Pdgfrα$^+$ mesendoderm niche during mouse development

Single-cell RNA-seq (scRNA-seq) analysis using Foxa2-Venus fusion protein reporter

mice indicated that the Foxa2 lineage might give rise to LPM and DE (Scheibner et al., 2021).

Given that Foxa2 and Pdgfrα are expressed during the conversion from mesendoderm to

mesenchyme(Artus et al., 2010; Kopper and Benvenisty, 2012; Scheibner et al., 2021; Tada et al.,

2005), we used Foxa2-lineage tracing mice ($Foxa2^{Cre/+}$; $Rosa^{tdTomato/+}$)(Horn et al., 2012) to

determine whether the Foxa2-lineage would label Mixl1$^+$ or Pdgfrα$^+$ mesendoderm around

E6.25~7.0 during gastrulation. Notably, Foxa2-lineage-driven tdTomato labels a part of the

Pdgfrα$^+$ mesendoderm of the primitive streak (PS) around E6.25~E6.5 relatively early-streak-stage

embryos (Figure 1C, arrow). In the E6.5~E7.0 mid-to-late-streak-stage PS region (Figure 1D,

dotted lines), Foxa2-lineage-derived tdTomato were found in the PS and Pdgfrα$^+$ adjacent nascent

mesoderm, suggesting that Foxa2 lineage-labeled Pdgfrα$^+$ cells ingresses from PS to nascent

mesoderm regions (Figure 1D, arrowheads). These data are reminiscent of the definition of

mesendoderm (Hart et al., 2002; Tada et al., 2005). Our result indicates that the Foxa2 lineage-

labeled Pdgfrα$^+$ cells are the mesendoderm, most likely the derivative of posterior epiblasts, since

Foxa2 is expressed only in the posterior epiblasts before gastrulation (Scheibner et al., 2021).

Further immunostaining analysis of the sequential sections of mid-streak-stage E6.5~6.75 embryos

confirmed that the Foxa2 lineage appeared in the distal portion of Mixl1-weakly-positive

mesendoderm (Figures 1E and 1F, arrows) at anterior primitive streak (APS) besides the Foxa2

protein-expressing DE (Figures 1E and 1F, arrowheads). Based on our Pdgfrα$^+$ lineage tracing

data, we further examined whether Foxa2-lineage would label lung mesenchyme.

Foxa2-lineage labeling increased during lung development, leading to occupy the entire lung

epithelium and half of the lung mesenchyme, including lung endothelium

Foxa2-lineage tracing mice (*Foxa2$^{Cre/+}$; Rosa$^{tdTomato/+}$*) faithfully target Nkx2-5$^+$ cardiac

progenitors associated with the origin of Wnt2$^+$ Isl1$^+$ CPP(Bardot et al., 2017; Peng et al., 2013)

lung mesenchyme. However, there was no conclusive evidence of whether Foxa2-lineage can give

rise to Wnt2$^+$ Isl1$^+$ CPP(Bardot et al., 2017; Peng et al., 2013). Using the Foxa2-lineage tracing

mice, we found from immunostaining that Foxa2 lineage labeling occupies the entire lung

epithelium and most of the lung mesenchyme at E16.5 (Figures 2A-C). Quantitative analyses by

flow cytometry in the E14.5 developing lungs of Foxa2-lineage tracing mice showed that Foxa2-

lineage labeled almost the entire lung epithelium (93.3%± 2.00) and the partial lung mesenchyme

(24.2%± 5.51), including endothelial cells (16.8 %± 5.12) (Figure 2D). Contrary to expectations,

Foxa2-lineage labeled cells increased dramatically throughout lung development (Figure 2E). In

adulthood, the Foxa2-lineage labeling reached about 99.97%± 0.0578 in lung epithelium, 51.27%±

1.72 in lung mesenchyme, and 59.97%± 8.44 in lung vascular endothelial cells, with more than

two-fold change in the lung endothelium, compared with E14.5 (Figure 2E). Morphometric

analysis of immunostaining further confirmed that Foxa2 lineage marked about 30% of E14.5

multiple cell types of lung mesenchyme: Sma$^+$ smooth muscle cells (24.9%± 8.50), VE-cadherin$^+$

(39.3%± 12.5) or Pecam1$^+$ (36.6%± 11.1) endothelial cells, and Pdgfrβ$^+$ pericytes of the pulmonary

arteries and pulmonary veins (41.4%± 10.5) (Figures 1F and 1G). Interestingly, no clear Foxa2

protein level expression was observed in the mesenchyme of the embryonic lungs (Figure S1A),

consistent with previously reported Foxa2 protein expression patterns(Wan et al., 2004). However,

the LungMAP deposited single-cell RNA-seq database analysis showed a sporadic Foxa2

transcriptional expression pattern in developing lung mesenchyme, particularly in proliferating

endothelium, on E15.5 and E17.5 (Figure S1B). We confirmed *Foxa2* transcriptional expression

in lung mesenchyme by in situ hybridization analysis on E18.5 (Figure S1C and S1D). The

expression was detected mostly in tdTomato positive but low frequency in the negative cells,

which indicates that lung mesenchyme spontaneously expressed *Foxa2* at a transcriptional level

and turn on tdTomato during lung development. We also sorted the cell fraction of CD45$^-$ CD31$^-$

EPCAM$^-$ tdTomato$^+$ and CD45$^-$ CD31$^-$ EPCAM$^-$ tdTomato$^-$ from developing lung mesenchyme at

E18.5. We observed a slight increase in the relative expression of *Foxa2* in the tdTomato$^+$ fraction

of embryonic lung mesenchyme, which most likely contributed to the labeling in the Foxa2-lineage

tracing mice (Figure S1E). These results suggest pulmonary mesenchymal progenitor cells turned

on Foxa2 expression slightly at the mRNA level rather than the protein level, which led to a gradual

increase in Foxa2 lineage labeling. Furthermore, tdTomato$^+$ endothelial cells were found to have

a slightly higher proliferative capacity than tdTomato$^-$ cells (Figure S1F). These results suggest

that Foxa2 lineage$^+$ pulmonary mesenchymal appeared during lung development, and lineage

labeling gradually increased compared to Foxa2- lung mesenchyme progenitors throughout lung

development.

Co-development of endodermal and mesodermal lung progenitors derived from MXIL1+

PDGFRα + FOXA2+ mesendoderm in the directed differentiation protocol using hiPSC

To determine whether Foxa2 or Pdgfrα mesoderm is an evolutionarily well-conserved

niche that can give rise to both pulmonary endoderm and mesoderm, we modified a previously

reported protocol to establish a pulmonary endoderm-mesoderm co-developmentally directed

differentiation protocol(Chen et al., 2017; Gotoh et al., 2014; Hawkins et al., 2017; Huang et al.,

2014; Konishi et al., 2016) (Figure 3A). With this optimized protocol, various hiPSC lines were

found to efficiently induce a lung bud-like appearance, indicated by NKX2-1+, in lung epithelial

cells(Gotoh et al., 2014; Konishi et al., 2016). We found that on day 10, TBX4+ lung mesenchyme

emerged and surrounded the NKX2-1+SOX9+ lung epithelium (Figure 3B and 3C). The qPCR

kinetics analyses across the time point further supported the appearance of lung mesenchyme,

represented by LPM marker expression peaked on day6~8; *OSR1, FGF10, BMP4, PDGFRα*, and

smooth muscle cell markers peaked on day8~10; *ACTA2* and *PDGFRβ*, and CPP markers peaked

on day8~12; *ISL1, WNT2, FOXF1,* and *TBX4*, peaked on days 10~14 simultaneously with

pulmonary epithelial markers, *NKX2-1* and *CPM* (Figure 3D).

In this differentiation protocol, NKX2-1+ lung endoderm and WNT2+TBX4+ lung

mesoderm were derived from the anteroventral endoderm and mesoderm at day 15 after Activin-

mediated definitive endoderm and LPM induction, respectively(Chen et al., 2017; Huang et al.,

2014). During primordial streak induction from day0 to day3, cell surface markers of PDGFRα

and EPCAM and intracellular FOXA2 and MIXL1 kinetics were analyzed by flow cytometry

every 12 hours (Figure 3E). Briefly, 12 hours after the Activin induction, more than 60% of the

EPCAM$^+$PDGFRα$^-$ primitive streak first turned on MIXL1, the mesendoderm marker(Hart et al.,

2002; Tada et al., 2005). Subsequently, the epithelial-mesenchymal transition occurred 24 hours

later, as represented by the PDGFRα induction in EPCAM$^+$MIXL1$^+$ mesendoderm. After 36 hours,

more than 90% of MIXL1+EPCAM+ mesoderm cells expressed PDGFRα. At the same time,

expression of FOXA2 appeared in some of those mesoderm cells (Figures 3E and 3F). Thereafter,

PDGFRα expression decreased, and 72 hours later, mutual FOXA2 induction appeared when

EPCAM$^+$FOXA2$^+$ DE and EPCAM$^-$FOXA2$^-$ LPM were presented (Figure 3G). The dynamics of

MIXL1, PDGFRα, and FOXA2 were further revealed by qPCR analysis (Figure 3H). These results

suggest that PDGFRα$^+$ and FOXA2$^+$ lineage are redundant but distinct stages of mesoderm

development. It can give rise to both endoderm and mesoderm lung cells conserved in mouse

development and human-directed differentiation (Figure S4A).

Foxa2-driven Fgfr2 conditional knockout showed a lung agenesis phenotype

The evolutionary-conserved Foxa2-lineage$^+$ mesendoderm forms endodermal and

mesodermal lung niches, suggesting that it may function as a BFL to generate the entire lungs after

donor cells are injected into the vacant niche in the Foxa2 lineage. To explore this possibility, CBC

was performed using Foxa2-driven Fgfr2-conditional knockout mice (*Foxa2$^{Cre/+}$; Fgfr2$^{flox/flox}$*,

hereafter, *Foxa2$^{Cre/+}$; Fgfr2^{cnull}*). Mitotic signaling via Fgfr2 is required for both lung epithelium

and mesenchyme, and systemic knockout mice of Fgf10 or Fgfr2 exhibit a phenotype of lung

agenesis(Arman et al., 1999; De Langhe et al., 2006; De Moerlooze et al., 2000; Sekine et al.,

1999). Based on the results of Foxa2 lineage tracking and the need for Fgfr2 signaling, it was

predicted that *Foxa2$^{Cre/+}$; Fgfr2^{cnull}* mice would be used to generate vacant niches in both lung

epithelium and mesenchyme. Indeed, they exhibited a lung agenesis phenotype (Figures S2A-S2C).

However, we did not observe agenesis phenotype in other major internal organs related to the

Fgfr2 systemic knockout phenotype (Figure S2D) (Arman et al., 1999; De Langhe et al., 2006; De

Moerlooze et al., 2000; Sekine et al., 1999).

Generation of the entire lungs in *Foxa2*-driven *Fgfr2*-deficient mice via CBC

To examine whether donor cells complement the lung agenesis phenotype, we generated

nGFP$^+$iPSCs from Rosa$^{nT\text{-}nG}$ mice (hereafter, nGFP$^+$iPSCs) via Sendai virus-mediated

reprogramming(Huang et al., 2014). nGFP$^+$iPSCs were injected into mouse blastocysts (Figure

4A), and chimerism was analyzed at E17.5. Strikingly, donor nGFP$^+$iPSCs generated whole lungs

in *Foxa2$^{Cre/+}$; Fgfr2^{cnull}* mice, but general chimerism in other organs were diverse (Figure 4B, S3A,

and Table 1). Importantly, almost the entire lung epithelial, mesenchymal, and endothelial cell

population at E17.5 was composed exclusively of nGFP$^+$iPSCs (Figures 4C and S3B). In contrast,

wild-type, Shh-driven heterozygous, or knockout mice showed about 50-70% chimerism in the

mesenchymal and endothelial lineages of the lung (Figures 4C and 4D). Interestingly, although

tdTomato$^+$ Fgfr2 knockout mesenchymal cells remained in early lung development at E14.5

(Figure S3C and Table 2), the percentage of Ki67$^+$ proliferating cells was significantly higher in

GFP$^+$ donor cells compared to tdTomato$^+$ host cells (Figures 4E – 4G). These results suggest that

generating an Fgfr2 deficient niche in the host Foxa2 lineage-derived lung mesenchyme is efficient

for donor iPSCs recruitment in the defective Foxa2 lineage mesenchyme niches.

DISCUSSION

The presence of committed organ precursors capable of contributing to multi-embryonic

layers after pluripotent epiblast formation has been assumed, but the identity and origin of the lung

precursors needed to be better defined. Although the Foxa2 lineage could not give rise to the entire

lung cells that served as a BFL, we have identified a Foxa2 lineage that provides a significant

stepping stone for facilitating whole lung generation during mouse development. Since donor cells

formed about 50~80% chimerism in the complemented lung mesenchyme niches at E14.5 (Table

1) and endogenous Foxa2 lineage forms only 20%~30% in the lung mesenchyme at E14.5, Fgfr2

defects at the Foxa2 lineage in early lung development are important but not sufficient for the

complete lung mesenchyme niche complementation. We showed that losing Fgfr2 expression in

Foxa2 lineage cells resulted in lower proliferative ability than donor cells during the lung

complementation (Figure 4F).

Lung epithelial cell precursors were well-known to be Shh$^+$ DE in the lung development

field(Cardoso and Kotton, 2008; Christodoulou et al., 2011; Harris et al., 2006; Kadzik and

Morrisey, 2012; Tian et al., 2011; Weaver et al., 1999; Xing et al., 2008), and indeed, the Shh-

lineage traces putative DE-derived epithelial lineage but little lung mesenchyme (Figure S3D).

Targeting the endodermal lung lineage driven by Shh was sufficient for lung epithelial

complementation but insufficient to generate whole lungs, and host-derived cells remained

substantially in the mesodermal lung component (Mori et al., 2019).

In contrast to lung epithelium precursors, the orderly commitment and the origin of the

entire pulmonary mesenchyme were not defined well. We showed that gastrulating Pdgfrα-lineage

is the origin of the entire lung mesoderm, including endothelium (Figures 1A and 1B)(Mori et al.,

2019). Furthermore, Foxa2 or Pdgfrα lineage labels a population primarily comprised of the

earliest specified precursors in the Mixl1$^+$ mesendoderm niche, which can be identified in vivo and

in human iPSC cell-derived directed differentiation protocol. Our findings, summarized in Figure

S4, pinpoint the lineage hierarchy of specified lung precursors in gastrulating mesendoderm,

further supported by the scRNA-seq analysis in early embryonic development(Pijuan-Sala et al.,

2019) (Figure S4B).

As previously indicated that nascent mesoderm differentiation into a CPP fate(Bardot et

al., 2017; Devine et al., 2014; Ng et al., 2022; Peng et al., 2013), we clarified the orderly

mesendoderm progression of gene expression, Mixl1, Pdgfrα, and Foxa2, and lung progenitor-

related markers that parallels the commitment of Foxa2+ Pdgfrα+ mesendoderm to an LPM and

DE fate in the lung directed differentiation protocol using human iPSC (Figure S4). Our lineage

tracing analysis also highlighted the unanticipated Foxa2 lineage program, the progressive increase

of lineage labeling by spontaneous expression of *Foxa2 mRNA* that occupies more than half of the

lung mesenchyme during lung development. Foxa2-lineage+ lung mesenchyme is a part of Pdgfrα

lineage+ cells, potentially providing a unique competitive developmental niche during lung

development. Further analyses using Foxa2CreERT2 lineage tracing mice are required to clarify

it. Intriguingly, the Foxa2 lineage+ mesenchyme did not show any discrete fates that would predict

anatomical localization—depleting Fgfr2 in the Foxa2 lineage results in the loss of the Fgfr2

mitogen-mediated function of Foxa2 lineage+ lung mesenchyme, which leads to the loss of

proliferative ability in most host lung mesenchyme. The Foxa2 lineage labeling in lung

mesenchyme is around 20% at E14.5 (Figure. 2D and 2E). Moreover, the donor cell

complementation in lung mesenchyme at E14.5 still holds host-derived lung mesenchyme but

decreases at E17.5. These results indicate that the defective Foxa2 lineage is critically important

for efficient lung complementation.

An unambiguous proof of the developmental origins of patterned organs is critical for

developing regenerative strategies and a better understanding of the genes responsible for

congenital malformations. For the future development of whole lung generation via CBC using

human iPSCs, the human-derived developmental program modification to match the host animals

is the key. More broadly, our studies offer a new paradigm that can apply to modeling various

congenital lung diseases and future autologous transplantation therapies using iPSC in the near

future.

Acknowledgments

We thank Zurab Ninish for his technical assistance. We sincerely appreciate the generous support

from Dr. Hiromitsu Nakauchi at Stanford University and the considerate support and scientific

input from Dr. Wellington Cardoso at the Columbia Center for Human Development (CCHD) and

the members of Cardoso's lab and CCHD. We acknowledge the support from the CCHD Medicine

Microscopy core (MMC), Columbia Stem Cell Initiative (CSCI) Flow Cytometry core (SONY

MA900), and Genetically Modified Mouse Model Shared Resource (GMMMSR) for blastocyst

injection. This work was funded by NIH-NHLBI 1R01 HL148223-01, DoD PR190557, PR191133

to M. M., JSPS202080340, and The Uehara Memorial Foundation to A. M.

Author contributions

A.M. and M.M. designed all experiments; Z.N. and A.M. maintained mutant mice for the injection;

C.S.L. performed blastocyst injection and embryo transfer; J.T., A.S., Y.S., Y.H., H.S., supported

lineage-tracing, chimera analyses, and genotyping; H.S., D.S., and S. T. helped to generate mouse

and human iPSCs, J.S. and F.H. kept human iPSC-directed differentiation, N.D. provided

Foxa2$^{Cre/+}$ mice, A.M. and M.M. wrote the paper; Y.H., H.S., J.W., J.Q., and F.H. gave crucial

insights on the experiments and the manuscripts. There is no competing financial interest.

Declaration of interests

The authors declare no competing interests.

FIGURES

Figure 1. Pdgfrα lineage during gastrulation is the origin of the entire pulmonary mesenchyme. (A) Left: Schematic of Tamoxifen administration (A-B). Representative Immunofluorescence (IF)-confocal imaging of E14.5 *Pdgfrα^{CreERT2/+}; Rosa26^{tdTomato/+}* lineage tracing mouse lungs (A, B). Tamoxifen administration at E5.5 labels Pdgfrα-lineage-driven tdTomato (red) in the entire lung mesenchyme (A), including Sma$^+$ airway smooth muscle cells, Pdgfrα$^+$ mesenchyme, and VE-cadherin$^+$ capillaries (B). Pdgfrα-lineage also labeled a low proportion of epithelial cells (B, white arrows). Enlarged box: dotted box. Cre$^-$ littermate control (A, middle panel) (*n = 3* per group). Scale bars: A, B = 100, 500μm. (C-F) Representative IF-confocal imaging of E6.25 (C), E7.0 (D), and E6.5 (E, F) from *Foxa2^{Cre/+}; Rosa26^{tdTomato/+}* mice. E-cadherin indicates epiblasts (C-F): (C) Foxa2-lineage (red) labeled Pdgfrα (green) expressing mesendoderm (arrow). (D) Foxa2-lineage labeled Pdgfrα$^+$ cells and ingresses from primitive streak (PS) (dotted lines) to nascent mesoderm regions (arrowhead). (E) Mixl1 (red) expression in PS (yellow dotted area). (F) Sequential section of E: Enlarged box: Foxa2-lineage (red) marked the distal portion of the arteriolarizing PS that is a part of the MIxl1^{dim+} mesendoderm (E-F, arrowheads). (*n = 3* per group). Scale bars = 50μm.

Figure 2. Foxa2-lineage gradually increased during lung development and labeled the entire

lung epithelium and half of the mesenchyme. (A-C) IF-confocal imaging of E16.5 *Foxa2[Cre/+];*

Rosa[tdTomato/+] embryonic lung: (A) Foxa2-lineage (red) labeled E-cadherin$^+$ lung epithelium (green)

entirely and Pfgfrβ$^+$ mesenchyme (cyan) partially. (B) Foxa2-lineage partially labeled VE-

cadherin$^+$ capillary (ca) (enlarged box, blue arrowhead). (C) Foxa2-lineage labeled Pdgfrβ$^+$ smooth

muscle cells of the pulmonary artery (pa) (enlarged box, white arrowhead) and pulmonary vein

(pv, yellow arrowhead). Scale bars (A), (B), and (C) = 200μm, 100μm, and 100μm, respectively.

(D and E) Representative histograms and the graphs of FCM quantitative analyses for CD31$^-$

Epcam$^+$ lung epithelium, CD31$^-$Epcam$^-$ mesenchyme, and CD31$^+$Epcam$^-$ endothelium at E14.5,

E18.5, P0, and four weeks adult (n = 4, 6, 7 and 3, independent biological replicates, respectively)

of *Foxa2[Cre/+]; Rosa[tdTomato/+]* mouse lungs. The gradual increase of % tdTomato$^+$ lineage labeling

in both lung mesenchyme and endothelium. Statistical analysis: one-way ANOVA with the Tukey

post hoc test.; statistically significant if *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. (F-

G) IF-confocal imaging of E14.5 *Foxa2[Cre/+]; Rosa[tdTomato/+]* embryonic lungs. tdTomato labeled entirely

with lung epithelial markers Nkx2-1(left) but a relatively low proportion of mesenchyme (Pdgfrβ:

middle, and Sma: right). (n = 3 per group). Scale bars = 50μm. Graphs in (G): The morphometric

analysis: % of Foxa2-lineage labeling in Nkx2-1$^+$epithelial, Pdgfrβ$^+$mesenchyme, Sma$^+$airway

smooth muscle, VE-Cadherin$^+$ capillaries, or PECAM1$^+$ arteries from E14.5 *Foxa2[Cre/+]; Rosa[tdTomato/+]*

lungs. (n = 3 per biological replicates, 5 fields per group).

Figure 3. Co-development of endodermal and mesodermal lung progenitors derived from

MXIL1⁺ PDGFRα⁺ FOXA2⁺ mesendoderm in the directed differentiation protocol using

hiPSC. (A) Schematic culture protocol of hiPSC-derived endodermal and mesodermal lung

progenitor cell co-differentiation. (B) Representative IF-confocal imaging of differentiating

hiPSCs at day 10 culture. Lung epithelium (NKX2-1), distal lung bud epithelium (SOX9),

mesenchyme (TBX4), and nucleus (Hoechst) markers. The budding structures expressed SOX9

and partially NKX2-1 (asterisk), and monolayer cells expressed TBX4. (C) Representative phase-

contrast images of the directed differentiation time course (D) qRT-PCR analyses of lung

mesenchyme and epithelium markers in time course according to the protocol shown in (A). Each

plot showed a different biological experiment (n = 3 independent experiments). (E) FCM-based

protein kinetic analyses during DE and LPM induction; MIXL1 expression preceded compared to

PDGFRα or FOXA2. FOXA2 appearance in the subset of the PDGFRα⁺ population (red asterisk).

(*n = 3* independent experiments) (F) Representative IF imaging of 36 hours-cultured hiPSCs. (G)

Schematic summary of flow cytometry analysis. (H) qRT-PCR analyses further confirmed the

preceded *MIXL1* induction and subsequent expression of *PDGFRα* and *FOXA2*. All graphs: Data

normalized by undifferentiated iPSCs. Each plot showed a different biological experiment (n = 3

independent experiments).. Error bars represent mean $\pm$ SD.

Figure 4. Generation of the entire lungs in *Foxa2*-driven *Fgfr2*-deficient mice via CBC.

(A) Schema of CBC experiment: a2i/VPA/LIF-treated SSEA1^{high} CD31^{high} nGFP$^+$iPSCs were

sorted and injected into WT, *Fgfr2^{hetero}* (heterozygous: *Foxa2$^{cre/+}$; Fgfr2$^{flox/+}$; Rosa26$^{tdTomato/+}$*), and

Fgfr2^{cnull} (homozygous: *Foxa2$^{cre/+}$; Fgfr2$^{flox/flox}$; Rosa26$^{tdTomato/+}$*) blastocysts. (B) Gross morphology,

GFP (green: donor nGFP$^+$iPSCs-derived signals), and tdTomato (host Foxa2-lineage-derived

signals) fluorescence of freshly isolated lungs from E17.5 chimeric *WT*(left), *Fgfr2^{hetero}* (left

middle), and *Fgfr2^{cnull}* (right middle) that were injected with nGFP$^+$iPSCs. Control: littermate

Fgfr2^{cnull} mouse without nGFP$^+$iPSCs injection (right). (C) IF-confocal imaging of E17.5 *Fgfr2^{cnull}*

or *Fgfr2^{hetero}* lungs injected with nGFP$^+$iPSCs. Dotted lines: enlarged images: Compared with

littermate control holding host-derived cells (red), E17.5 *Fgfr2^{cnull}* lungs were entirely composed

of donor-derived nGFP$^+$ cells (green). (D) Graphs: % GFP in CD31$^-$Epcam$^+$ lung epithelium,

CD31$^-$Epcam$^-$ mesenchyme, and CD31$^+$Epcam$^-$ endothelium analyzed by flow cytometry. Each

plot: a different biological animal. *Foxa2$^{Cre/+}$; Fgfr2^{cnull}; Rosa^{tdTomato}* (*n=5*, independent biological

replicates), *Foxa2$^{Cre/+}$; Fgfr2^{hetero}; Rosa$^{tdTomato/+}$*(*n=4*), *Shh$^{Cre/+}$; Fgfr2^{cnull}; Rosa$^{tdTomato/+}$* (*n=2*),

Shh$^{Cre/+}$; Fgfr2^{hetero}; Rosa$^{tdTomato/+}$(*n=3*), and *WT* (*n=4*). (E and F) Representative IF staining of

E14.5 lung of *Fgfr2^{cnull}*. GFP and tdTomato indicate donor and host-derived cells, respectively. (E-

F) Residual tdTomato$^+$ host cells in E14.5 *Foxa2$^{Cre/+}$; Fgfr2^{cnull}* chimeric lungs. (F) Split images of

(E) visualizing GFP$^+$ donor cells and tdTomato$^+$ host cells, co-stained with Ki67. (G) Graphs: %

Ki67 labeling in mesenchymal cells of E14.5 *Foxa2$^{Cre/+}$; Fgfr2^{cnull}* chimeric lungs. Statistical

analyses: paired Student's t-test, significance at **P* < 0.05, ns: non-significant. Scale bars: B, C

(left, right), E, F =1mm, 500μm, 50μm, 20μm, 20μm.

TABLES

Table. 1 E17.5 Chimerism of Foxa2 promoter-driven conditional blastocyst

complementation

		Liver	Lung			
			Whole	**Epithelium**	**Mesenchyme**	**Endothelium**
Foxa2-Cre	**Fgfr2** *cnull*	56.2	91.8	99.8	92.6	78.1
		54.8	90.6	99.2	90	92.8
		47.4	90.3	99.2	89.8	91.9
		38.5	87.7	99.8	87.3	83.5
		14	58.9	97.8	57.6	47.1
	Fgfr2 *hetero*	60.2	65.9	46.2	68.1	33.1
		56.3	88.8	81.8	89.2	90.7
		24.2	42.1	52.9	40	51.9
		9.16	22.7	54.6	19.4	21.1
Shh-Cre	**Fgfr2** *cnull*	35.7	73.6	99.2	71.7	73.9
		13	79.2	99.8	76.5	77.8
	Fgfr2 *hetero*	57.4	83.1	99.3	79.6	86.6
		44.4	84.3	99.2	83.4	78.7
		13.8	35	59.2	33.8	27
	Wild type	56	68.5	79.2	66.2	73.6
		22.3	56.7	18	61.3	68.7
		20.1	53.1	54.9	52.9	53.7
		17	89.5	90.8	80.6	87.7

426 **Table. 2 E14.5 Chimerism of Foxa2 promoter-driven conditional blastocyst**

427 **complementation**

		Liver	Lung			
			Whole	Epithelium	Mesenchyme	Endothelium
Foxa2-Cre	Fgfr2 cnull	59.9	81.9	99.4	81.4	86.2
		51	75.4	99.4	75.4	56.8
		25.3	56.5	97.4	54.1	63.2
	Fgfr2 hetero	37.1	54.4	56.9	54	58.9
		29.8	48.6	83.3	47.7	47.3
		18.7	43.4	18	45	31
		0.54	0.24	0.91	0.1	2.11
		0.4	0.17	1.02	0.058	1.23
	Wild type	39	68.7	56.2	69.2	67.5
		29.4	81.1	62.7	81.8	80.8
		28.6	73.9	39.6	75.6	76.4
		13.8	29.5	5.57	30.4	29.6
		0	2.22	5.52	1.5	16

Methods

Mouse. *Shh*$^{Cre/+}$ mice (cat. 05622), *Rosa26*$^{tdTomato/\ tdTomato}$ mice (cat. 07914), Rosa26$^{nT-nG/nT-nG}$ mice (cat. 023035) and *Pdgfrα*$^{CreERT2/+}$ mice (cat. 032770) were obtained from the Jackson Lab. X. Zhang kindly gifted *Fgfr2*$^{flox/flox}$ mice. We further backcrossed these mice for over three generations with CD-1 mice (cat. 022) from the Charles River. Dr. Nicole C Dubois kindly provided *Foxa2*$^{Cre/Cre}$ mice (129xB6 mixed background). For conditional deletion of *Fgfr2* (*Fgfr2*^{cnull}), we crossed *Fgfr2*$^{flox/flox}$; *Rosa26*$^{tdTomato/tdTomato}$ females with *Foxa2*$^{Cre/Cre}$; *Fgfr2*$^{flox/+}$, *Foxa2*$^{Cre/+}$; *Fgfr2*$^{flox/+}$ or *Shh*$^{Cre/+}$; *Fgfr2*$^{flox/+}$ males, respectively. PCR performed genotyping of the *Shh-Cre*, *Pdgfrα-CreERT2*, *Rosa26-nTnG*, and *Rosa26-tdTomato* alleles according to the protocol provided by the vendor. For the CBC, genotyping of chimeric animals was confirmed by GFP-negative sorted liver cells and lung cells. For detecting the *Fgfr2* floxed allele, we performed PCR using the primer sets: FR2-F1, 5′-ATAGGAGCAACAGGCGG-3′, and FR2-F2, 5′-CAAGAGGCGACCAGTCA-3′(Mori et al., 2019). For lineage tracing with Pdgfrα$^{CreERT2/+}$; Rosa26$^{tdTomato/+}$ mice, 1 dose of 200 μg tamoxifen (MedChem Express, HY-13757A) per g of body weight was given via oral gavage injection. All animal experiments were approved by Columbia University Institutional Animal Care and Use Committee in accordance with US National Institutes of Health guidelines.

Culture of mouse iPSC. We cultured iPSC in a2i/VPA/LIF medium on a feeder, as previously reported(Mori et al., 2019). These PSC cells were passaged at a split ratio of 1:10 every 2–3 d.

Culture of human iPSCs (hiPSCs). All iPSC lines were maintained in feeder-free conditions on laminin iMatrix-511 silk E8 (Amsbio, AMS.892021) in StemFit 04 complete Medium (Amsbio, SFB-504), supplemented with Primocin (Invivogen, ant-pm-1), and passaged with TrypLE Select

(Gibco, A1285901). All human iPSC lines used were characterized for pluripotency and were

found to be karyotypically normal. The BU3NGST cell line was kindly gifted by Dr. Finn Hawkins

and Dr. Darrell Kotton at Boston University, Boston, MA. Dr. Jennifer Davis, the University of

Washington School of Medicine, Seattle, WA, kindly gifted the Rainbow cell line. PD2 and TD1

hiPSC were generated from deidentified commercially available human peripheral blood

mononuclear cell and tracheal epithelial cell lines via the manufacturing protocol of Sendai virus-

mediated reprogramming (CytoTune2.0) (ThermoFisher, A16517). Every other month, all iPSC

lines screened negative for mycoplasma contamination using a MycoAlert PLUS detection kit

(Lonza, LT07-710).

Differentiation of hiPSCs into lung epithelial and mesenchymal cells. The directed

differentiation protocols were modified from previous protocols to maximize lung mesenchymal

cell generation concomitantly with NKX2-1$^+$ lung epithelium. Briefly, DE and LPM precursors

were induced once seeded hiPSC-formed colonies by the Activin induction using the STEMdiff

Definitive Endoderm Kit (StemCell Technologies, 05110) for 72 hours. Differentiated cells were

dissociated and passaged in Laminin511-coated tissue culture plates in a complete serum-free

differentiation medium (cSFDM)(Chen et al., 2017). To induce DE and LPM into the anterior

foregut endoderm and mesoderm, the cSFDM was supplemented with 10μM SB431542

(MedChem Express, HY-10431) and 2μM Dorsomorphin (Tocris, 3093) for 48 hours and 10μM

SB431542 and 2μM IWP2 (Tocris, 3533) for 24 hours. Cells were then cultured for 7-10 additional

days in cSFDM containing 3μM CHIR99021, 10ng/ml recombinant human FGF10 (R&D Systems,

345-FG), 10ng/ml recombinant human KGF (R&D Systems, 251-KG), 10 ng/mL recombinant

human BMP4 (R&D Systems, 314-BP), and 50nM retinoid acid (Sigma-Aldrich, R2625) to induce

NKX2-1 positive lung epithelial cells and WNT2$^+$TBX4$^+$ lung mesenchymal cells.

Immunofluorescence (IF). Before the immunostaining, antigen retrieval was performed using

Unmasking Solution (Vector Laboratories, H-3300) for 10 min at around 100 °C by microwave.

7-μm tissue sections were incubated with primary antibodies in the buffer of M.O.M. kit (Vector

Laboratories, MKB-2213-1) overnight at 4 °C, washed in PBS, and incubated with secondary

antibodies conjugated with Alexa488, 567, or 647 (ThermoScientific, 1:400) with NucBlue Fixed

Cell Ready Probes Reagent (Hoechst) (ThermoScientific, R37605) for 1.5 h, and mounted with

ProLong Gold antifade reagent (Invitrogen, P36930). The images were captured by a Zeiss

confocal 710 microscopy. The antibodies were listed in the Supplemental Table. 1.

Immunocytochemistry. Cells on culture dishes were fixed with 4% Paraformaldehyde (PFA) for

30 min at room temperature (RT), permeabilized, and blocked with staining buffer containing

0.025% Triton X-100 and 1% BSA for 1 hour at RT. Primary antibodies (Supplementary Table 1)

were incubated overnight at 4 °C in the staining buffer. After three washes in PBS, secondary

antibodies (Supplementary Table 1) and NucBlue Fixed Cell Ready Probes Reagent (Hoechst)

were incubated for 1 h. The samples were imaged using DMi8 Leica widefield microscope. The

antibodies were listed in the Supplemental Table. 1.

RNAScope in situ hybridization (ISH). RNA at E18.5 lung sections were stained by RNAScope

probes: Mm-Foxa2-T8 (Advanced Cell Diagnostics, #409111-T8) or Negative control (NC)

(Advanced Cell Diagnostics, #324341) using the RNAScope HiPlex12 Reagent kit v2 (Advanced

Cell Diagnostics, #324419) according to manufacture-provided protocol. Then, sections were

incubated with tdTomato antibody for 2 hours at room temperature, washed in PBS, and incubated

with secondary antibody conjugated with Alexa488 with NucBlue Fixed Cell Ready Probes

Reagent (Hoechst) for 1.5 h, and mounted with ProLong Gold antifade reagent.   The images of

Foxa2 and NC ISH were captured with the same setting by a Zeiss confocal 710 microscopy.

Flow cytometry (FCM) analyses of mouse lung tissue. Lungs from lineage tracing mice at

E14.5, E18.5, P0, four weeks or lungs from CBC chimeric mice at E14.5 and E17.5 were harvested

and prepared for the FCM, as previously described(Mori et al., 2019). Briefly, tissues were minced

with microscissors, and 1 ml of pre-warmed dissociation buffer (1 mg/ml DNase (Sigma, DN25),

5 mg/ml collagen (Roche, 10103578001), and 15 U/ml Dispase II (Stemcell Technologies, 7913)

in HBSS), incubated at 37 °C on the rocker with 50 r.p.m. speed, and neutralized with the

dissociation buffer by FACS buffer containing 2% FBS, Glutamax, 2mM EDTA and 10mM

HEPES in HBSS after the 30 min incubation. After filtrating the cells with a 40-μm filter

(FALCON, 352235), cell pellets were resuspended with 1 ml of cold RBC lysis buffer (Biolegend,

420301) to lyse the remaining erythrocytes for 5 min on ice, and neutralized by 1 ml cold FACS

buffer. After that, it was centrifuged them at 350 rcf, 4 °C, for 3 min to remove the lysed blood

cells. For FCM analysis, one million cells were transferred in 100 μl of FACS buffer supplemented

with 0.5μM Y27632 and then added 2 μl Fc Block (BD Pharmingen, 553141) per sample followed

by 10 min incubation on ice. Cells were incubated with the following antibodies: CD31-APC

(Biolegend, 102510, 1/50), Epcam-BV711 (BioLegend, 118233, 1/50), or Epcam-BV421

(Biolegend, 118225, 1/50), Aqua Zombie (BioLegend, 423101, 1/100), CD45-BV605

(BioLegend, 103104, 1/50) for 30 min on ice. After staining, cells were washed twice with FACS

buffer before resuspending in 500 μl FACS buffer for the subsequent analyses using SONY

MA900 or NovoCyte. Compensation was manually performed to minimize the tdTomato signal

leakage to the GFP channel using FlowJo (ver. 10. 7. 1). 2 samples from E14.5 lineage tracing

mice were removed from analysis based on inaccurate staining of live staining. Total 4 lungs were

calculated from 6 embryos.

nGFP$^+$iPSC establishment and preparation for CBC donor.

E14.5 lung tissues of *Rosa26$^{nTnG/nTnG}$* mice (JAX, cat. 023035, C57BL/6NJ background) were

harvested in a dissociation buffer described above. The dissociated cells were seeded on a 10cm

dish, and only lung fibroblast survived after 1 week in MEF medium(Mori et al., 2019). The

fibroblasts were passaged using Accutase (Innovative Cell Technologies, AT104) and seeded on

gelatin (Millipore-Sigma, ES006B)-coated 6-well plates with a density of 0.1 million cells per

well. Upon cell attachment, Yamanaka reprogramming factors were induced to iPSCs via Sendai

virus using CytoTune2.0 (ThermoFisher, A16517). To establish nGFP$^+$ iPSCs, the Cre plasmid

was transfected using Fugene HD transfection reagent (Promega, E2311), then sorted out

GFP$^+$tdTomato$^-$ live cells by FACS (SONYMA900), and single clones were expanded.

For the CBC donor cell preparation, nGFP$^+$iPSCs cultured in a2i/VPA/LIF(Mori et al., 2019) were

trypsinized and resuspended in 4 ml cold DMEM + 10% FBS immediately and filtering the cells

with a 40-μm filter. Cells were centrifuged at 350 rcf, 4 °C, for 3 min, and the supernatant was

removed. After being washed with flow buffer containing 0.2% BSA, 1% Glutamax, and 1μM

Y27632, the cells were resuspended in 100μl/1 million cells with flow buffer. The following

antibodies were added: Epcam-BV421 (1:50), SSEA1-PE (1:50), CD31-APC (1:50), Zombie

Aqua Fixable Viability Kit (1:100). Epcam^{high}SSEA1^{high}CD31^{high} cells were sorted by FACS

(SONYMA900) and subsequently prepared for the injection.

Blastocyst preparation and embryo transfer. Blastocysts were prepared by mating *Foxa2$^{Cre/Cre}$*;

Fgfr2$^{flox/+}$,*Foxa2$^{Cre/+}$; Fgfr2$^{flox/+}$* or *Shh$^{Cre/+}$; Fgfr2$^{flox/+}$* males (all 129 x B6 x CD-1 background) with

superovulated Fgfr2$^{flox/flox}$; Rosa26$^{tdTomato/tdTomato}$ females (129 x B6 x CD-1 background). Blastocysts

were harvested at E3.5 after superovulation(Mori et al., 2019). 20 sorted nGFP$^+$iPSCs were

injected into each blastocyst. After the iPSC injection, blastocysts were cultured in an M2 medium

(Cosmobio) for a few hours in a 37 °C, 5% CO2 incubator for recovery. Then, blastocysts were

transferred to the uterus of the pseudopregnant foster mother.

Real-time-quantitative RT-PCR (qRT-PCR). Total RNA was extracted using a Direct-zol™

RNA MiniPrep Plus kit (Zymo Research, R2072), and cDNA was synthesized using Primescript™

RT Master Mix (Takara, RR036B). The cDNAs were then used as templates for qRT-PCR analysis

with gene-specific primers. Reactions (10 μl) were performed Luna® Universal qPCR Master Mix

(New England Biolabs, M3003X). mRNA abundance for each gene was determined relative to

GAPDH mRNA using the $2^{-\Delta\Delta Ct}$ method. The primers were listed in the Supplemental Table. 2.

Data were represented as mean ± SD of measurements. The number of animals or cells per group

is provided in the legends. The undetected values in each biological experiment in Fig.3d were

removed from the graphs.

Statistical analysis. Data analysis was performed using Prism 8. Data acquired by performing

biological replicas of two or three independent experiments are presented as the mean ± SD.

Statistical significance was determined using a two-tailed t-test and unpaired one-way or two-way

ANOVA with the Tukey post hoc test. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, ns: non-significant.

Data availability. scRNA-seq data for mouse gastrulation and early organogenesis described in

the manuscript have been analyzed from the deposited database at

https://marionilab.cruk.cam.ac.uk/MouseGastrulation2018/. The authors declare that all data

supporting the results of this study are available within the paper and the Supplementary

Information. Raw data are available from the corresponding author upon reasonable request.

References

Arman E, Haffner-Krausz R, Gorivodsky M, Lonai P. 1999. Fgfr2 is required for limb outgrowth

and lung-branching morphogenesis. *Proceedings of the National Academy of Sciences*

96:11895–11899. doi:10.1073/pnas.96.21.11895

Artus J, Panthier JJ, Hadjantonakis AK. 2010. A role for PDGF signaling in expansion of the

extra-embryonic endoderm lineage of the mouse blastocyst. *Development* **137**:3361–3372.

doi:10.1242/dev.050864

Bardot E, Calderon D, Santoriello F, Han S, Cheung K, Jadhav B, Burtscher I, Artap S, Jain R,

Epstein J, Lickert H, Gouon-Evans V, Sharp AJ, Dubois NC. 2017. Foxa2 identifies a

cardiac progenitor population with ventricular differentiation potential. *Nat Commun* **8**:1–

15. doi:10.1038/ncomms14428

Cardoso W v, Kotton DN. 2008. Specification and patterning of the respiratory system.

Stembook 1–18. doi:10.3824/stembook.1.10.1

Chen J, Lansford R, Stewart V, Young F, Alt FW. 1993. RAG-2-deficient blastocyst

589   complementation: an assay of gene function in lymphocyte development. *Proceedings of*

590   *the National Academy of Sciences* **90**:4528–4532. doi:10.1073/pnas.90.10.4528

591   Chen Y-W, Huang SX, de Carvalho ALRT, Ho S-H, Islam MN, Volpi S, Notarangelo LD,

592   Ciancanelli M, Casanova J-L, Bhattacharya J, Liang AF, Palermo LM, Porotto M, Moscona

593   A, Snoeck H-W. 2017. A three-dimensional model of human lung development and disease

594   from pluripotent stem cells. *Nat Cell Biol* **19**:542–549. doi:10.1038/ncb3510

595   Christodoulou C, Longmire TA, Shen SS, Bourdon A, Sommer CA, Gadue P, Spira A, Gouon-

596   Evans V, Murphy GJ, Mostoslavsky G, Kotton DN. 2011. Mouse ES and iPS cells can form

597   similar definitive endoderm despite differences in imprinted genes. *Journal of Clinical*

598   *Investigation* **121**:2313–2325. doi:10.1172/JCI43853

599   Crapo JD, Barry BE, Gehr P, Bachofen M, Weibel ER. 1982. Cell number and cell

600   characteristics of the normal human lung. *American Review of Respiratory Disease*

601   **126**:332–337. doi:10.1164/arrd.1982.126.2.332

602   De Langhe SP, Carraro G, Warburton D, Hajihosseini MK, Bellusci S. 2006. Levels of

603   mesenchymal FGFR2 signaling modulate smooth muscle progenitor cell commitment in the

604   lung. *Dev Biol* **299**:52–62. doi:10.1016/j.ydbio.2006.07.001

605   De Moerlooze L, Spencer-Dene B, Revest JM, Hajihosseini M, Rosewell I, Dickson C. 2000. An

606   important role for the IIIb isoform of fibroblast growth factor receptor 2 (FGFR2) in

607   mesenchymal-epithelial signalling during mouse organogenesis. *Development* **127**:483–92.

608   Devine WP, Wythe JD, George M, Koshiba-Takeuchi K, Bruneau BG. 2014. Early patterning

609   and specification of cardiac progenitors in gastrulating mesoderm. *Elife* **3**.

610   doi:10.7554/ELIFE.03848

Gotoh S, Ito I, Nagasaki T, Yamamoto Y, Konishi S, Korogi Y, Matsumoto H, Muro S, Hirai T,

Funato M, Mae SI, Toyoda T, Sato-Otsubo A, Ogawa S, Osafune K, Mishima M. 2014.

Generation of alveolar epithelial spheroids via isolated progenitor cells from human

pluripotent stem cells. *Stem Cell Reports* **3**:394–403. doi:10.1016/j.stemcr.2014.07.005

Green MD, Chen A, Nostro M-C, d'Souza SL, Schaniel C, Lemischka IR, Gouon-Evans V,

Keller G, Snoeck H-W. 2011. Generation of anterior foregut endoderm from human

embryonic and induced pluripotent stem cells. *Nat Biotechnol* **29**:267–72.

doi:10.1038/nbt.1788

Guyette JP, Gilpin SE, Charest JM, Tapias LF, Ren X, Ott HC. 2014. Perfusion decellularization

of whole organs. *Nat Protoc* **9**:1451–68. doi:10.1038/nprot.2014.097

Hackett TL, Knight D a, Sin DD. 2010. Potential role of stem cells in management of COPD. *Int*

J Chron Obstruct Pulmon Dis **5**:81–8.

Hamanaka S, Umino A, Sato H, Hayama T, Yanagida A, Mizuno N, Kobayashi T, Kasai M,

Suchy FP, Yamazaki S, Masaki H, Yamaguchi T, Nakauchi H. 2018. Generation of

Vascular Endothelial Cells and Hematopoietic Cells by Blastocyst Complementation. *Stem*

Cell Reports **11**:988–997. doi:10.1016/j.stemcr.2018.08.015

Han L, Chaturvedi P, Kishimoto K, Koike H, Nasr T, Iwasawa K, Giesbrecht K, Witcher PC,

Eicher A, Haines L, Lee Y, Shannon JM, Morimoto M, Wells JM, Takebe T, Zorn AM.

2020. Single cell transcriptomics identifies a signaling network coordinating endoderm and

mesoderm diversification during foregut organogenesis. *Nat Commun* **11**.

doi:10.1038/s41467-020-17968-x

Harris KS, Zhang Z, McManus MT, Harfe BD, Sun X. 2006. Dicer function is essential for lung

epithelium morphogenesis. *Proc Natl Acad Sci U S A* **103**:2208–13.

doi:10.1073/pnas.0510839103

Hart AH, Hartley L, Sourris K, Stadler ES, Li R, Stanley EG, Tam PPL, Elefanty AG, Robb L.

2002. Mixl1 is required for axial mesendoderm morphogenesis and patterning in the murine

embryo. *Development* **129**:3597–608.

Hawkins F, Kramer P, Jacob A, Driver I, Thomas DC, Mccauley KB, Skvir N, Crane AM,

Kurmann AA, Hollenberg AN, Nguyen S, Wong BG, Khalil AS, Huang SXL, Guttentag S,

Rock JR, Shannon JM, Davis BR, Kotton DN. 2017. Prospective isolation of NKX2-1 –

expressing human lung progenitors derived from pluripotent stem cells. *J Clin Invest* **127**:1–

18. doi:10.1172/JCI89950

Horn S, Kobberup S, Jørgensen MC, Kalisz M, Klein T, Kageyama R, Gegg M, Lickert H,

Lindner J, Magnuson MA, Kong YY, Serup P, Ahnfelt-Rønne J, Jensen JN. 2012. Mind

bomb 1 is required for pancreatic β-cell formation. *Proc Natl Acad Sci U S A* **109**:7356–

7361. doi:10.1073/pnas.1203605109

Huang SXL, Islam MN, O'Neill J, Hu Z, Yang Y-G, Chen Y-W, Mumau M, Green MD,

Vunjak-Novakovic G, Bhattacharya J, Snoeck H-W. 2014. Efficient generation of lung and

airway epithelial cells from human pluripotent stem cells. *Nat Biotechnol* **32**:84–91.

doi:10.1038/nbt.2754

Kadzik RS, Morrisey EE. 2012. Directing lung endoderm differentiation in pluripotent stem

cells. *Cell Stem Cell* **10**:355–61. doi:10.1016/j.stem.2012.03.013

Kemter E, Schnieke A, Fischer K, Cowan PJ, Wolf E. 2020. Xeno-organ donor pigs with

multiple genetic modifications - the more the better? *Curr Opin Genet Dev* **64**:60–65.

doi:10.1016/J.GDE.2020.05.034

Kitahara A, Ran Q, Oda K, Yasue A, Abe M, Ye X, Sasaoka T, Tsuchida M, Sakimura K,

Ajioka Y, Saijo Y, Zhou Q. 2020. Generation of Lungs by Blastocyst Complementation in

Apneumic Fgf10-Deficient Mice. *Cell Rep* **31**:107626.

doi:10.1016/J.CELREP.2020.107626

Kobayashi T, Yamaguchi T, Hamanaka S, Kato-Itoh M, Yamazaki Y, Ibata M, Sato H, Lee Y-S,

Usui J-I, Knisely a S, Hirabayashi M, Nakauchi H. 2010. Generation of rat pancreas in

mouse by interspecific blastocyst injection of pluripotent stem cells. *Cell* **142**:787–99.

doi:10.1016/j.cell.2010.07.039

Konishi S, Gotoh S, Tateishi K, Yamamoto Y, Korogi Y, Nagasaki T, Matsumoto H, Muro S,

Hirai T, Ito I, Tsukita S, Mishima M. 2016. Directed Induction of Functional Multi-ciliated

Cells in Proximal Airway Epithelial Spheroids from Human Pluripotent Stem Cells. *Stem*

Cell Reports **6**:18–25. doi:10.1016/j.stemcr.2015.11.010

Kopper O, Benvenisty N. 2012. Stepwise differentiation of human embryonic stem cells into

early endoderm derivatives and their molecular characterization. *Stem Cell Res* **8**:335–345.

doi:10.1016/j.scr.2011.12.006

Kotton DN, Morrisey EE. 2014. Lung regeneration: Mechanisms, applications and emerging

stem cell populations. *Nat Med* **20**:822–832. doi:10.1038/nm.3642

Li E, Ustiyan V, Wen B, Kalin GT, Whitsett JA, Kalin T v., Kalinichenko V v. 2021. Blastocyst

complementation reveals that NKX2-1 establishes the proximal-peripheral boundary of the

airway epithelium. *Dev Dyn* **250**:1001–1020. doi:10.1002/DVDY.298

Mori M, Furuhashi K, Danielsson JA, Hirata Y, Kakiuchi M, Lin CS, Ohta M, Riccio P,

Takahashi Y, Xu X, Emala CW, Lu C, Nakauchi H, Cardoso W v. 2019. Generation of

functional lungs via conditional blastocyst complementation using pluripotent stem cells.

Nat Med **25**:1691–1698. doi:10.1038/s41591-019-0635-8

Ng WH, Johnston EK, Tan JJ, Bliley JM, Feinberg AW, Stolz DB, Sun M, Wijesekara P,

Hawkins F, Kotton DN, Ren X. 2022. Recapitulating Human Cardio-pulmonary Co-

development Using Simultaneous Multilineage Differentiation of Pluripotent Stem Cells.

Elife **11**:1–27. doi:10.7554/eLife.67872

Ott HC, Clippinger B, Conrad C, Schuetz C, Pomerantseva I, Ikonomou L, Kotton D, Vacanti

JP. 2010. Regeneration and orthotopic transplantation of a bioartificial lung. *Nat Med*

16:927–33. doi:10.1038/nm.2193

Peng T, Tian Y, Boogerd CJ, Lu MM, Kadzik RS, Stewart KM, Evans SM, Morrisey EE. 2013.

Coordination of heart and lung co-development by a multipotent cardiopulmonary

progenitor. *Nature* **500**:589–592. doi:10.1038/nature12358

Petersen TH, Calle EA, Zhao L, Lee EJ, Gui L, Raredon MSB, Gavrilov K, Yi T, Zhuang ZW,

Breuer C, Herzog E, Niklason LE. 2010. Tissue-engineered lungs for in vivo implantation.

Science (1979) **329**:538–541. doi:10.1126/science.1189345

Pijuan-Sala B, Griffiths JA, Guibentif C, Hiscock TW, Jawaid W, Calero-Nieto FJ, Mulas C,

Ibarra-Soria X, Tyser RCV, Ho DLL, Reik W, Srinivas S, Simons BD, Nichols J, Marioni

JC, Göttgens B. 2019. A single-cell molecular map of mouse gastrulation and early

organogenesis. *Nature* **566**:490–495. doi:10.1038/s41586-019-0933-9

Scheibner K, Schirge S, Burtscher I, Büttner M, Sterr M, Yang D, Böttcher A, Ansarullah, Irmler

M, Beckers J, Cernilogar FM, Schotta G, Theis FJ, Lickert H. 2021. Epithelial cell plasticity

drives endoderm formation during gastrulation. *Nat Cell Biol* **23**:692–703.

doi:10.1038/s41556-021-00694-x

Sekine K, Ohuchi H, Fujiwara M, Yamasaki M, Yoshizawa T, Sato T, Yagishita N, Matsui D,

Koga Y, Itoh N, Kato S. 1999. Fgf10 is essential for limb and lung formation. *Nat Genet*

21:138–41. doi:10.1038/5096

Stone KC, Mercer RR, Gehr P, Stockstill B, Crapo JD. 1992. Allometric Relationships of Cell

Numbers and Size in the Mammalian Lung. *Am J Respir Cell Mol BioI* **6**:235–243.

Tada S, Era T, Furusawa C, Sakurai H, Nishikawa S, Kinoshita M, Nakao K, Chiba T,

Nishikawa SI. 2005. Characterization of mesendoderm: A diverging point of the definitive

endoderm and mesoderm in embryonic stem cell differentiation culture. *Development*

132:4363–4374. doi:10.1242/dev.02005

Tian L, Gao J, Garcia IM, Chen HJ, Castaldi A, Chen YW. 2021. Human pluripotent stem cell-

derived lung organoids: Potential applications in development and disease modeling. *Wiley*

Interdiscip Rev Dev Biol **10**:1–16. doi:10.1002/wdev.399

Tian Y, Zhang Y, Hurd L, Hannenhalli S, Liu F, Lu MM, Morrisey EE. 2011. Regulation of lung

endoderm progenitor cell behavior by miR302/367. *Development* **138**:1235–45.

doi:10.1242/dev.061762

Travaglini KJ, Nabhan AN, Penland L, Sinha R, Gillich A, Sit R v., Chang S, Conley SD, Mori

Y, Seita J, Berry GJ, Shrager JB, Metzger RJ, Kuo CS, Neff N, Weissman IL, Quake SR,

Krasnow MA. 2020. A molecular cell atlas of the human lung from single-cell RNA

sequencing. *Nature* **587**:619–625. doi:10.1038/s41586-020-2922-4

Usui J, Kobayashi T, Yamaguchi T, Knisely a S, Nishinakamura R, Nakauchi H. 2012.

Generation of kidney from pluripotent stem cells via blastocyst complementation. *Am J*

Pathol **180**:2417–26. doi:10.1016/j.ajpath.2012.03.007

Wan H, Kaestner KH, Ang SL, Ikegami M, Finkelman FD, Stahlman MT, Fulkerson PC,

Rothenberg ME, Whitsett JA. 2004. Foxa2 regulates alveolarization and goblet cell

hyperplasia. *Development* **131**:953–964. doi:10.1242/dev.00966

Wang X. 2019. Bioartificial Organ Manufacturing Technologies. *Cell Transplant* **28**:5–17.

doi:10.1177/0963689718809918

Weaver M, Yingling JM, Dunn NR, Bellusci S, Hogan BL. 1999. Bmp signaling regulates

proximal-distal differentiation of endoderm in mouse lung development. *Development*

126:4005–4015.

Wen B, Li E, Ustiyan V, Wang G, Guo M, Na CL, Kalin GT, Galvan V, Xu Y, Weaver TE,

Kalin T v., Whitsett JA, Kalinichenko V v. 2021. In vivo generation of lung and thyroid

tissues from embryonic stem cells using blastocyst complementation. *Am J Respir Crit Care*

Med **203**:471–483. doi:10.1164/RCCM.201909-

1836OC/SUPPL_FILE/DISCLOSURES.PDF

Xing Y, Li C, Hu L, Tiozzo C, Li M, Chai Y, Bellusci S, Anderson S, Minoo P. 2008.

Mechanisms of TGFbeta inhibition of LUNG endodermal morphogenesis: the role of

TbetaRII, Smads, Nkx2.1 and Pten. *Dev Biol* **320**:340–50. doi:10.1016/j.ydbio.2008.04.044

Yamaguchi T, Sato H, Kato-Itoh M, Goto T, Hara H, Sanbo M, Mizuno N, Kobayashi T,

740   Yanagida A, Umino A, Ota Y, Hamanaka S, Masaki H, Rashid ST, Hirabayashi M,

741   Nakauchi H. 2017. Interspecies organogenesis generates autologous functional islets.

742   *Nature* **542**:191–196. doi:10.1038/nature21070

 SUPPLEMENTAL FIGURES.

Figure S1. Foxa2-lineage gradually increased in the mesenchyme and endothelial cells during

mouse lung development. (A) Representative IF-confocal imaging of E14.5 and E18.5 mouse

embryonic lungs. Foxa2 (green) was expressed in lung epithelial cells labeled by E-cadherin (red)

but not in E-cadherin negative lung mesenchymal area. Scale bar = 50μm (B) Single-cell analysis

of mouse LungMAP. *Foxa2* expression was observed in lung mesenchyme and endothelium in

E15.5 and E17.5 (Arrows). (C), (D) Representative IF and in situ hybridization-confocal imaging

of E18.5 mouse embryonic lungs. Dotted lines: enlarged image. *Foxa2* mRNA was expressed in

lung epithelial cells and lung mesenchyme. Most of the *Foxa2* mRNA expression was colocalized

with tdTomato. Scale bars (left, right) = 50 μm, 20μm (E) Graphs: *Foxa2* qPCR for tdTomato

(tdT)$^+$ or tdTomato$^-$ CD45$^-$ CD31$^-$ EPCAM$^-$ mesenchymal cells and CD45$^-$CD31$^-$EpCAM$^+$

epithelial cells sorted by FACS from E18.5 Foxa2-lineage tracing mice. (n = 3, each biological

replicate) (F) Flow cytometry quantitative analyses: EdU labeling cells in CD31$^+$Epcam$^-$

endothelium at E14.5 and E18.5 (n = 6 each independent biological replicates, respectively) of

Foxa2$^{Cre/+}$; Rosa$^{tdTomato/+}$ lungs. Statistical analysis: paired Student's t-test, significance at *$P<0.05$,

ns: non-significant.

A

Sox2 Nkx2-1 Hoechst

B **C**

tdTomato

D

Figure S2. Lung agenesis phenotype in the *Foxa2$^{Cre/+}$Fgfr2^{cnull}* mice.

(A) Representative IF staining of E10.5 *Foxa2$^{Cre/+}$;Fgfr2^{hetero}* and *Foxa2$^{Cre/+}$;Fgfr2^{cnull}*: *Fgfr2^{cnull}* did

not form Sox2$^+$NKX2-1$^+$proximal airway bifurcation. Scale bar = 200μm. Lu: lung, Eso:

Esophagus. Tr: trachea. Nkx2-1 (blue), Sox2 (green). (B) Representative Gross morphology of

E14.5 *Foxa2$^{Cre/+}$;Fgfr2^{hetero}* and *Foxa2$^{Cre/+}$;Fgfr2^{cnull}*: No difference in the appearance of the embryo

between *Fgfr2^{hetero}* and *Fgfr2^{cnull}* (Top). Limb (arrows) and placenta (pl) are present in mice of all

genotypes. Conversely, the *Fgfr2^{cnull}* embryo showed lung agenesis phenotype (Bottom, asterisk)

(n = 6 per each group). (C) Representative image of *Fgfr2^{cnull}* lungs. Foxa2 lineage$^+$ tdTomato

signal indicates the trachea ends in the middle of the thoracic cavity (asterisk). (D) HE staining:

The internal organs such as the intestine, pancreas, kidney, tooth buds, and hair follicles were

normally formed in *Fgfr2^{cnull}* E16.5 embryos. Scale bar = 500μm.

A

Foxa2$^{Cre/+}$; Fgfr2^{cnull}
+ nGFP$^+$iPSC

Foxa2$^{Cre/+}$; Fgfr2^{hetero}
+ nGFP$^+$iPSC

WT + nGFP$^+$iPSC

C

Epithelium

Mesenchyme

Endothelium

B

Foxa2$^{Cre/+}$; Fgfr2^{cnull} + nGFP$^+$iPSC

Podoplanin Sftpc
GFP Hoechst

Scgb1a
GFP Hoechst

Foxj1 Cgrp
GFP Hoechst

Sma VE-cadherin
GFP Hoechst

D *Shh$^{Cre/+}$; Rosa$^{tdtomato/+}$*

tdTomato Hoechst

Figure S3. The complemented embryos of *Foxa2$^{Cre/+}$; Fgfr2^{cnull} +nGFP+iPSCs showed

normal gross morphology and various chimerism (A) Representative gross morphology of

chimeric embryos of *WT* + nGFP+iPSCs, *Foxa2$^{Cre/+}$; Fgfr2^{hetero}* + nGFP+iPSCs, *Foxa2$^{Cre/+}$;*

Fgfr2^{cnull} +nGFP+iPSCs. The color indicated various GFP+ chimerism on the skin of those

embryos and the residual host Foxa2- lineage+ tdTomato signals. (B) Representative IF-confocal

imaging of E17.5 lungs of chimeric *Fgfr2^{cnull}*+ nGFP+iPSCs. The E17.5 rescued lungs from

Fgfr2^{cnull} +nGFP+iPSCs expressed differentiated cell markers such as Podoplanin (alveolar type1

cells), Sftpc (type 2 cells), Scg1a1 (club cells), Foxj1 (multiciliated cells), Cgrp (neuroendocrine

cells), Sma (smooth muscle cells), and VE-cadherin (endothelial cells). Scale bar = 20μm. (C)

Graphs: % of donor cell chimerism: % of GFP in CD31$^-$Epcam$^+$ lung epithelium fraction, CD31$^-$

Epcam$^-$ mesenchyme, and CD31$^+$Epcam$^-$ endothelium analyzed by flow cytometry. Each plot: a

different biological animal (see table 2). *Foxa2$^{Cre/+}$; Fgfr2^{cnull}; Rosa^{tdTomato}* (*n=3*, independent

biological replicates), *Foxa2$^{Cre/+}$; Fgfr2^{hetero}; Rosa$^{tdTomato/+}$*(*n=5*), and *WT* (*n=5*) at E14.5. (D)

Representative IF-confocal imaging of E14.5 Shh-lineage tracing mouse lungs (*Shh$^{Cre/+}$;*

Rosa$^{tdTomato/+}$). Shh-lineage labeled the entire lung epithelium but rarely mesenchyme, distinct

from Foxa2 lineage mice (see Fig. 2F). Scale bar = 100μm.

A

ME*: mesendoderm (primitive streak ~ nascent mesoderm)

Foxa2 lineage ⇒ All lung epithelium & part of the lung mesenchyme

Pdgfrα lineage ⇒ All lung mesenchyme & part of the lung epithelium

B

Figure S4. Summary of the results and proposed models. (A) Pluripotent stem cell (PSC)

differentiation will be initiated by the Mixl1+ induction and the subsequent temporal expression of

an early mesendoderm (ME) marker, Pdgfrα. After that, the late mesendoderm marker marked by

Foxa2 will turn on. Pdgfrα and Foxa2-lineage partially overlap at the primitive streak stage during

gastrulation. Pdgfrα+ early ME lineage gives rise to the partial lung epithelial cells, most likely by

the overlapping Foxa2+ lineage and entire lung mesenchyme through lateral plate mesoderm

(LPM) induction and cardiopulmonary lineage (CPP). On the other hand, the Foxa2-lineage gives

rise to the whole lung epithelium and about 20% of CPP in early lung development at E12.5~E14.5.

Strikingly, the whole lungs, including lung epithelium, mesenchyme, and endothelial cells, were

produced solely by donor iPSCs via complementing the Foxa2-lineage's mitotic defective lung

niches. It proves that a Foxa2-lineage works as a bona fide lung generative lineage (BFL), enough

to generate entire lungs by donor iPSCs via CBC. (B) Top left panel: Schematic diagram of

mesendoderm lineage trajectory for lung formation based on the single cell RNA-seq (scRNA-

seq) deposited database(Pijuan-Sala et al., 2019). During gastrulation, a bipotent mesendoderm

(Mes: rounded blue area) appears in the transition from PS and NM, labeled by *Mixl1, T, Gsc, and*

Eomes. NM forms lateral plate mesoderm (LPM). Mesendoderm also gives rise to anteriolar PS

(APS) and, subsequently, definitive endoderm (DE). Top right panel: Clustering of scRNA-seq

provided in the deposited database(Pijuan-Sala et al., 2019). Bottom left panel: Timepoint of

scRNA-seq provided in the deposited database(Pijuan-Sala et al., 2019). bottom: *Foxa2* is

expressed in the PS (black arrow) and the part of NM (red arrow). In contrast, *Shh* appeared in DE

but not in PS, NM, or APS. A few cells showed *Pdgfrα* in PS (black arrow) but most expressed

Pdgfrα in NM (red arrow) and LPM.