Differential effects of multiplex and uniplex affiliative relationships on biomarkers of inflammation

Jessica Vandeleest[1]*, Lauren J. Wooddell[2], Amy C. Nathman[1], Brianne A. Beisner[3], Brenda McCowan[1]

[1]California National Primate Research Center, University of California, Davis, CA, United States

[2]Department of Neurosurgery, Emory University, Atlanta, GA, United States

[3]Emory National Primate Research Center Field Station, Lawrenceville, GA, United States

*Corresponding author details: Jessica Vandeleest, PhD

Email: vandelee@ucdavis.edu

Abstract

Social relationships profoundly impact health in social species. Much of what we know regarding the impact of affiliative social relationships on health in nonhuman primates (NHPs) has focused on the structure of connections or the quality of relationships. These relationships are often quantified by comparing different types of affiliative behaviors (e.g., contact sitting, grooming, alliances, proximity) or pooling affiliative behaviors into an overall measure of affiliation. The influence of the breadth of affiliative behaviors (e.g., how many different types or which ones) a dyad engages in on health and fitness outcomes remains unknown. Here we employed a social network approach to explicitly explore whether the integration of different affiliative behaviors within a relationship can point to the potential function of those relationships and their impact on health-related biomarkers (i.e., pro-inflammatory cytokines) in a commonly studied non-human primate model system, the rhesus macaque (*Macaca mulatta*). Being well connected in multiplex grooming networks (networks where individuals both contact sat and groomed), which were more modular and kin biased, was associated with lower inflammation (IL-6, TNF-alpha). In contrast, being well connected in uniplex grooming networks (dyad engaged only in grooming and not in contact sitting), which were more strongly linked with social status, was associated with greater inflammation. Results suggest that multiplex relationships may function as supportive relationships that promote health. In contrast, the function of uniplex grooming relationships may be more transactional and may incur physiological costs. This complexity is important to consider for understanding the mechanisms underlying the association of social relationships on human and animal health.

Keywords: affiliation, *Macaca mulatta*, cytokines, inflammation

Introduction

For decades, research has shown that social relationships impact individual health and fitness in a variety of animal species, with estimates of the magnitude of the association with mortality in humans on par with other well recognized mortality risks (e.g., smoking, alcohol consumption) [1]. While the importance of social factors for health and fitness are widely recognized, the mechanisms by which social life exerts its influence are still not well understood [2,3]. One reason is that social life is complex; it encompasses both agonistic and affiliative interactions that vary across a range of dimensions such as frequency of interaction, symmetry, tenor, predictability, and stability [4–6]. As such, new approaches addressing this complexity are needed to better understand the biological mechanisms by which social relationships influence health.

Traditionally, two approaches taken to unravel this complexity in humans are to examine the structural and/or functional aspects of social relationships. Structural studies frequently concentrate effort on quantifying relationships by calculating the number of social partners, the frequency of interactions, or the higher order structuring of social relationships using social network analysis [1,3,7–13]. Functional studies attempt to examine what purpose or role a relationship serves and whether it meets the needs of the individual. In humans, functional measures of social relationships include surveys of perceived social support, informational support, emotional support, and tangible support [1]. Both have been associated with health outcomes where some studies have found that the quantity of relationships (structural) have health benefits and others have emphasized that the types or quality of relationships (functional) are what matters (see Holt-Lunstad et al., 2010 for a meta-analysis). In comparison, for nonhuman animals, assessing the structural aspects of social relationships is common [3,14–16], but

68   examining the functions of different relationships is far more challenging due to the fact we

69   cannot directly ask animals about the value or perceptions of their relationships. Instead, research

70   on social relationships in animals has more often relied on metrics designed to indirectly assess

71   the quality rather than directly query the function of their relationships [6].

72   As such, studies of animal affiliation have not explicitly attempted to differentiate

between structural and functional aspects of social relationships. Instead they tend to characterize

the quantity and quality of relationships by measuring various affiliative behaviors, and, in doing

so, these behaviors are either analyzed separately (e.g., grooming *or* proximity) or lumped

together giving them roughly equal weight [3,6]. Social network analysis is one technique well

suited to studying the structure of social relationships. Typically, researchers examine the effects

of individual network centrality metrics (e.g., eigenvector, betweenness, and closeness centrality)

and their impact on a variety of health-related metrics. While the general pattern seen across

these measures is that greater connectedness or centrality in specific behavioral networks is

associated with lower risk for gastrointestinal pathogens, increased reproduction and longevity

(Balasubramaniam et al., 2016; Brent et al., 2013; Cheney et al., 2016; Ostner & Schülke, 2018),

there is little consistency across studies in identifying which specific social role or network

metric is important, and some studies find no impact of social network position on fitness at all

(Ellis et al., 2019; Ostner & Schülke, 2018; Snyder-Mackler et al., 2020). Notably, many of

these networks metrics are highly correlated and, in practice, measure similar roles making the

study of the underlying mechanisms challenging [17,18]. Indeed, recent perspectives on network

analysis suggest that certain metrics such as betweenness centrality only become interesting

when they deviate from other network metrics such as degree centrality because if the structure

of the network is a result of a few highly central individuals high betweenness is likely due to

high degree [19–21]. Thus, it is difficult to ascertain which specific metric to use and thus a suite of

metrics is often employed instead [22].

Another commonly used metric to assess the quality of affiliative social relationships in

nonhuman primates is the dyadic sociality index (DSI) which aggregates information on the

quantity (frequency) of multiple, correlated affiliative behaviors (e.g., grooming and proximity).

Relationships with high DSI scores are commonly referred to as strong bonds and tend to be

equitable, stable, involve frequent interaction, and are most common between kin and peers

[3,23,24]. From a health perspective, higher number or quality of these strong bonds has been

associated with acute hormonal responses (e.g., oxytocin or cortisol levels), increased

reproduction, and longer survival [3,9,12,25]. However, these strong affiliative bonds make up only a

small fraction of the affiliative relationships an individual has [12,23,26–28] with weaker affiliative

bonds comprising the remainder. The function of these other, weaker bonds has been

hypothesized to increase social flexibility (e.g., social connections can shift with environmental

demands), allowing general social integration and indirect connections that might provide access

to others who may have resources or information [3,26]. Yet, evidence for an association between

weak bonds and health and fitness is mixed [12,26]. As helpful as these metrics are, such approaches

may overlook key information on the degree to which the diversity or breadth of affiliative

interactions in which a dyad engages are *integrated* [4,6]. For example, a relationship in which a

specific dyad engages in multiple types of affiliation (e.g., contact sitting, grooming, *and*

proximity) may differ from one in which a dyad engages in only one single behavioral domain

(e.g., contact sitting only, grooming only, *or* proximity only), even if the rate of interaction is the

same.

Recent advances in social network analysis and theory, and specifically multilayer or

multiplex networks, ([29]; e.g., multiple types of interactions among the same set of individuals)

may eventually provide a solution with tools to address such issues and, most importantly, to

disentangle the impact of structural and functional social relationships. However, currently,

these analytical methods (e.g., metrics like versatility and multiplex PageRank) are not

sufficiently sophisticated to account for the many structural differences that exist between

behavioral networks of different types (e.g., proximity networks are often dense and undirected,

contact networks may be more sparse and undirected whereas grooming networks are directed)

[29]. Yet, given mounting evidence for the importance of multidimensionality in social

relationships, the integration of diverse affiliative behaviors at this dyadic level may provide

important additional information (1) as to the nature of those relationships, (2) on how their

structure points to their potential function(s), and (3) on their downstream impacts on health and

fitness (Balasubramaniam et al., 2016). Indeed, in many past studies, multidimensional measures

often are the strongest predictors of health and fitness [1,8,13]. Yet, again, few animal studies to date

(except see [27]) have examined whether the integration of that diversity of interactions in a

relationship provide clues as to the nature of those social relationships. One exception, conducted

by Balasubramaniam and colleagues [14] representing one type of integrated approach, found that

highly connected rhesus macaques (i.e., high outdegree or eigenvector) in a grooming network

were less likely to have *Shigella*, a gastrointestinal pathogen, *but only if they were also well*

connected in a huddling network (i.e., high betweenness), suggesting that the presence of

multiple affiliative behaviors across an individual's dyadic connections (e.g., integration) confer

a social buffering effect on individual health (Balasubramaniam et al., 2016).

Here we employed a social network approach to explicitly explore whether the

integration of diverse affiliative behaviors within a relationship can point to the potential

function of those relationships and their impact on health-related outcomes in a commonly

studied non-human primate model system, the rhesus macaque (*Macaca mulatta*). We use

rhesus macaques as a group-living, nonhuman primate (NHP) model because their social

relationships are highly differentiated, exhibit a high degree of complexity and individual

variability, and have been linked to a variety of health and fitness outcomes [30–33]. Affiliation in

rhesus macaques, as in many primate species takes many forms, including grooming, contact

sitting, proximity, embracing, and less commonly coalitionary support [34–36]. In macaques,

grooming is commonly used to indicate the presence of an affiliative relationship [37,38]. Grooming

has been proposed to serve multiple social functions including: to maintain social bonds [37] and

social cohesion [39], and in exchange for tolerance from dominants, for agonistic support, or for

access to resources [40–42]. Although less commonly studied, contact sitting behavior (similar to

huddling) may also be an important indicator of strong affiliative relationships [43], particularly

those that may offer social buffering [14].

Therefore, in this study, an integration of affiliative behaviors was conducted by filtering

these behavioral networks (grooming and contact sitting) into "multiplex" and "uniplex"

networks. We use the term "multiplex" to refer to networks in which edges are represented by

the co-occurrence of multiple affiliative behaviors and the term "uniplex" to refer to networks in

which edges represent only one specific type of interaction (e.g. grooming *or* contact sitting, not

both)[4,8]. Our multiplex networks were defined as the set of dyads who engaged in both grooming

and contact sitting with edge-weights reflecting grooming (multiplex grooming) or contact

sitting (multiplex contact sitting) frequency, while our uniplex networks were defined as the set

of dyads who engaged solely in one behavior with edge weights reflecting the frequency of

interaction (uniplex grooming or uniplex contact sitting). Our rationale for constructing

multiplex and uniplex networks in this manner was inspired by Balasubramaniam [14] which

identified the combination of grooming and huddling behavior to be especially protective against

Shigella infection. Further, this distinction could also be hypothesized to relate to the different

functions of grooming (social bonding/cohesion vs. exchange for tolerance and resources).

Therefore, our primary focus for this study was to examine the differences between multiplex

grooming and uniplex grooming networks. However, we also present results comparing

additional complementary networks (multiplex contact sit vs. uniplex contact sit and grooming

vs. contact sitting) to fully explore the impact of filtering networks in this way. Our analysis had

two main goals: 1) determine if the network structure of related networks differed, and 2) the

effects of individual network centrality on health. To this end, differences in global network

structure of multiplex and uniplex or grooming and contact sitting networks were compared and

individual-level centrality metrics from these networks were examined for their association with

biomarkers of inflammation (i.e., serum pro-inflammatory cytokines), which are common, well-

established indicators of individual health status [33,44]. As such, our prediction was that

individuals exhibiting more central roles in multiplex (socially cohesive) networks would show

lower levels of inflammatory cytokines than those exhibiting more central roles in uniplex

networks.

Materials and Methods

Study system

Rhesus macaques live in large multi-male, multi-female social groups organized by rank and kinship [45]. For females, rank is inherited from their mothers and generally all members of a matriline hold adjacent ranks [46] (although see [47]). In contrast, males generally immigrate into a new social group and may enter at the bottom of the hierarchy, queueing for rank, or attain rank through direct competition [48]. Rhesus macaque females form the core of the social group with affiliation between both kin and non-kin playing a key role in maintaining group stability [45,49]. Although male social bonds have important fitness outcomes in macaques generally [50], male rhesus macaques engage in social affiliation far less frequently [51] and tend to be more socially isolated than females [52]. Therefore, we focused our study on females, which we predict will be more strongly impacted by social bonds than males. We use rhesus macaques as a group-living, nonhuman primate (NHP) model because their social relationships are highly differentiated, exhibit a high degree of complexity and individual variability, and have been linked to a variety of health and fitness outcomes [30–33].

Subjects and housing

Subjects were 248 breeding age (3 years and older) female rhesus macaques (*Macaca mulatta*) that were born at the California National Primate Research Center in Davis, California (Table 1). Subjects lived in one of four large multigenerational and matrilineal social groups containing 100-200 mixed-sex individuals (Table 1), each housed in a 0.2 hectare outdoor enclosure. Subjects were fed commercial monkey chow and foraging enrichment twice daily. Fruits or vegetables were provided weekly. Water was available ad libitum.

Table 1: Group Demographics

Group	Group size (adults)	N (adult females)	# of Matrilines[a]	Mean Matriline[a] size (SD)
Group A	131 (101)	74	13	5.7 (3.6)
Group B	204 (101)	67	33	2.0 (1.0)
Group C	125 (55)	39	6	6.5 (3.9)
Group D	185 (96)	68	13	5.2 (2.3)

[a] Number of matrilines and matriline size statistics include only breeding age females. Individuals were considered part of the same matriline if they could be traced back to the same female common ancestor at the time of group formation.

Behavioral data collection

Subjects were part of a larger study on the associations between social networks and health. Groups A and B were studied for six continuous weeks during the birthing season from March to April 2013 and 2014, respectively. Groups C and D were studied for six continuous weeks during the breeding season from September to October 2013 and 2014, respectively. Behavioral data were collected six hours per day, four days per week from 0900-1200 and 1300-1600 each day by three observers (inter-rater reliability, Krippendorff's alpha ≥0.85). Affiliative behavior was collected by one observer via scan sampling every 20-minutes (maximum 18 scans per day), where identities of all adult female dyads affiliating (i.e. grooming or contact sitting) were recorded[14]. All animals had tattoos and fur markings that allowed accurate individual identification. All observers demonstrated animal ID reliability of > 95%. Grooming was defined as cleaning or manipulating the fur of another animal and contact sitting included ventral contact, embrace, or side by side sitting for at least 3 seconds. During each scan, these behaviors were mutually exclusive for a dyad (an individual grooming another was not contact sitting with that individual). Affiliation scans produced 1637 scans (Group A: N=418, Group B: N=410, Group C: N=378, Group D: N=431) and a median of 38 grooming interactions per female (group range 23 – 49) and 28 contact sitting interactions (group range 13-52). This sampling scheme has

been shown to produce sufficiently sampled grooming and contact sitting networks[38]. Aggression

data (threats, chases, bites) were collected via an event sampling protocol for six hours per day,

four days per week by two other observers (average of 42.5 interactions per individual, group

range 36.2 – 51.9). Because social status has been shown to impact inflammation[31] (although

see[30]), dyadic aggression data was used to calculate dominance ranks and dominance certainty

via the R package *Perc*[30,53]. Dominance rank was expressed as the percent of animals in the

group outranked and therefore ranged from 0 (low) to 1 (high).

Affiliative network analysis

First, weighted networks were constructed from grooming and contact sitting interactions

(Figure 1A). Each of these networks (i.e., grooming or contact sitting) were then separated into

two more networks, a multiplex network where edges between dyads that both groomed and

contact sat were retained (Figure 1C or 1E), and a uniplex network in which edges were retained

for dyads that only groomed (Figure 1D) or only contact sat (Figure 1B). Edge-weights in

contact sitting networks (all contact sitting, uniplex contact sitting, multiplex contact sitting)

reflected the number of unique scans in which a dyad was observed contact sitting over the 6-

week period. Edge-weights in grooming networks reflected the number of unique scans a dyad

was observed grooming (Table S1). For each of the 6 networks (all grooming, multiplex

grooming, uniplex grooming, all contact sitting, multiplex contact sitting, uniplex contact

sitting), centrality and cohesion measures for each individual were calculated in R (ver. 4.0.5)

using iGraph (ver. 1.3.0). The effects of the direct connections for individuals were measured

using degree centrality and strength. The effect of an individual's indirect connections in the

network was evaluated using eigenvector, betweenness, and closeness centralities [3,14,16]. In

addition, the degree to which individuals were part of cohesive local communities was measured

by the local clustering coefficient (i.e., triadic closure). Multiple metrics were chosen to reflect

the different ways social integration can manifest (e.g., bridging, cohesion, embeddedness, etc.

Table 2).

Figure 1: A network example using a subset of animals from Group XX. Edge weight is represented using line thickness and behavior is indicated by color. Network filtering started with networks consisting of (A) all observed grooming (blue) and contact sitting (red) interactions. Edges from these networks were then filtered into multiplex grooming (E) or multiplex contact sitting (C) networks if dyads engaged in both behaviors. Dyads that

270   only groomed were filtered into a uniplex grooming network (D) and dyads only engaged in
271   contact sitting were filtered into a uniplex contact sitting network (B).
272
273   **Table 2: Network Metric Definitions**
274

Measure	Description
Degree/Strength	measures the *number* (unweighted) of partners or *frequency* of interaction (i.e., strength) for each node.[54]
Eigenvector	measures whether individuals are well connected to others that are also well connected, a measure of *social capital*.[54]
Betweenness	measures the number of times a node lies on the shortest path between other nodes, which reflects an individual's role in connecting others in the network or acting as a *social bridge*.[54]
Closeness	measures how close each node is to all other nodes within the network, which reflects how *embedded* an individual is in the network. [55]
Clustering Coefficient	measures the extent to which a node's neighbors are also connected to each other, a measure of *cliquishness or subgrouping*.[56]

Biological sample collection

Blood samples were taken during the fifth week of each group's study period during routine, semi-annual health checks. On a single morning, all animals in a group were lightly sedated with ketamine (10 mg/kg) and given veterinary exams. Blood samples were obtained from the femoral vein and serum was aliquoted and stored at −80 °C for later assay. The order in which animals were processed and samples were collected was recorded to control for any potential impacts of the sampling procedure on the physiological variables examined.

Pro-inflammatory Cytokines

Chronic inflammation is associated with a variety of diseases (e.g., diabetes, cardiovascular disease, cancer) and mortality [57–59], and high levels of pro-inflammatory cytokines, such as IL-6 and TNF-α, have previously been reported to be associated with social variables (e.g., low social status, low social integration, poor quality relationships, loneliness) in both humans and rhesus macaques [30,33,44,60,61]. Therefore, we chose to measure serum levels of IL-6 and TNF-α as a general biomarker of health. Serum levels of IL-6 and TNF-α were measured simultaneously using commercially available, species specific Milliplex multi-analyte profiling (MAP) reagents purchased from EMD/Millipore (Billerica, MA, USA), and utilizing Luminex Xmap technology (Luminex, Austin, TX, USA). Color coded polystyrene microbeads coated with specific antibodies for IL-6 and TNF-α were incubated with the serum samples, washed, and then further reacted with biotinylated detector antibodies followed by Streptavidin-PE to label the immune complexes on the beads. After a final washing to remove all unbound material, the beads were interrogated in a BioPlex dual laser (BioRad, Hercules, CA, USA). The median fluorescent index for each sample was compared to a standard curve to calculate the concentration (IL-6: mean = 12.55 pg/mL, sd = 46.92, range = 0 – 690; TNF-α: mean = 185.0 pg/mL, sd = 442.27, range = 0 – 4052; see Figure S2 for histograms). Samples were tested in

duplicate and had an intra-assay coefficient of variability of 15.3%. Samples were re-analyzed if

the CV was greater than 25% for all analytes measured. Manufacturer provided quality control

samples fell within recommended ranges for all assays. Samples falling below the threshold

sensitivity of the assay (1.6 pg/mL) were assigned a value of zero (IL-6: N = 77, TNF-α: N =

56).

Statistical analysis

Two sets of analyses were done to determine whether 1) multiplex and uniplex grooming

networks (i.e., Fig 1B vs 1C or Fig. 1D vs 1E) or grooming and contact sitting networks (Fig. 1A

red vs. blue) differ in structure and relationships to known social features of rhesus macaques

(e.g., kin bias, hierarchical organization), and 2) whether network metrics from these networks

predicted biomarkers of inflammation, with a specific focus on the relative impact of multiplex

vs uniplex (or grooming vs. contact sitting) network position on inflammation.

First, to validate this method, the structure of the multiplex and uniplex networks, which

were treated as weighted and directed (grooming only) networks, were compared to determine if

they exhibited differences in key structural features of rhesus relationships. For example,

evidence suggests that despotic macaques such as rhesus, particularly in large groups, are likely

to have grooming networks that are modular (i.e., shows subgrouping), expected to be based on

kinship, and have individual network positions (i.e., eigenvector centrality) that are correlated

with rank[62,63]. Therefore, we examined whether these two networks differed in the degree of

clustering (Newman's modularity, clustering coefficient), kin bias (e.g., proportion of kin (kin

unweighted degree/total unweighted degree)), and associations with rank (proportion of

grooming up the hierarchy, rank disparity among grooming dyads) for each of the four groups

studied. Also, because previous research has focused on bond strength, we further examined

reciprocity, strength of relationships (average edge weight), and distribution of grooming

(eigenvector centralization) across these network types. Due to the low number of groups in the

comparison, paired t-tests were used to evaluate if network metrics were consistently different

across groups. Normality of the differences was evaluated using the Shapiro-Wilk test, and if

significant then Wilcoxon signed rank tests were used. Grooming and contact sitting networks

were also compared. As a final structural analysis, we examined the correlations between

individual level network positions from these two network types (Table S2) to evaluate

multicollinearity within networks and associations between networks, including contact sitting

networks.

Next, to determine if an individual's position in the multiplex or uniplex networks was

associated with pro-inflammatory cytokines, we ran generalized linear models using a negative

binomial distribution (R package lme4 v.1.1-34) on each biomarker separately (see [30] for details

on distribution choice and Figure S2 for distributions). For these analyses, networks were treated

as weighted but undirected because of our focus on the qualities of a relationship rather than

focused specifically on grooming behavior and because contact sitting is recorded as an

undirected behavior (i.e., information on who initiated the interaction is unavailable). One

animal was excluded from the IL-6 analyses because it was an outlier with influence (Cook's D

>1); all other outliers had a Cook's D < 0.5 and therefore were included in the analyses. A

second animal was excluded from all analysis due to the fact she was not included in the uniplex

network. Model building proceeded in five steps for each outcome (i.e., IL-6, TNF-α, see Figure

S2). For all steps, ΔAIC > 2 was used to identify potential predictors and candidate models. For

one social group, 6 animals were not present in the uniplex contact sit network (they never

engaged in contact sitting with someone they did not also groom). Therefore to evaluate the

effects of uniplex contact sitting we use a subset of the full dataset for AIC comparisons. First, a

random effect indicating the group ID was evaluated for each outcome, and all subsequent

models were compared to this random effects only model. Second, variables from the literature

(age, dominance rank, dominance certainty, sampling order), although not of direct interest here,

were evaluated to determine if it was necessary to control for their effects on inflammation

before examining social network variables. Third, due to the large number of potential predictors

in our exploratory analysis, a statistical winnowing strategy was used to reduce the number of

social network variables under consideration [64]. This involved running univariate models for

each metric from each network (6 networks with 6 metrics each). Due to our goal of directly

comparing effects of centrality in different networks and predictions for multiplex vs uniplex

networks, if no metric generated improvement in model fit (i.e., ΔAIC > 2) for a given network

all available network metrics for that network were explored in step 4. Fourth, to compare the

effects of uniplex and multiplex network connectivity directly, candidate predictors from step 3

for the uniplex and multiplex grooming (Figure 1D vs E), uniplex and multiplex contact sitting

networks (Figure 1B vs C) were directly compared. Additionally, effects of centrality in

grooming vs contact sitting networks were compared (i.e., Figure 1A red vs blue). Fifth, a final

set of best models was identified by comparing AIC across all models generated in steps 3 and 4.

Metrics from the same network were never included in the same model due to the

interdependence of network metrics. If no single best model emerged, candidate models (i.e.,

those with ΔAIC $\leq$ 2) are discussed. A log of all models tested is available in Tables S3-4.

Ethical Note

All procedures used in this study met all legal requirements of the United States as well

as guidelines set by the American Society of Primatologists regarding the ethical treatment of

non-human primates. This study was approved by the Institutional Care and Use Committee at

the University of California, Davis and was carried out in compliance with the ARRIVE

guidelines.

Results

Multiplex vs. Uniplex Affiliation Networks

For all groups studied, clear differences in network topology, kinship, and associations

with dominance rank were seen between the multiplex and uniplex affiliative networks (Table 3).

Multiplex grooming networks had higher average edge-weight (the average number of

interactions per social partner), clustering coefficient, and modularity (the degree to which the

network can be divided into subgroups) than uniplex grooming networks for all groups. Notably,

although average edge-weights in the multiplex networks were higher than uniplex networks, the

predominant edge weight in all networks was 1-2 interactions (Figure S1). Multiplex grooming

networks also consistently showed more kin bias (proportion kin) and reciprocity than uniplex

grooming networks. In contrast, both multiplex and uniplex grooming networks showed

associations between rank and affiliation (i.e., grooming was directed up the hierarchy and

eigenvector centrality was correlated with rank in both networks) but the disparity in the ranks of

the grooming partners was greater in the uniplex affiliation networks compared to the multiplex

networks. Results for the multiplex vs uniplex contact sitting networks were the same as for

multiplex and uniplex grooming networks with the exception of reciprocity and grooming up the

hierarchy which were not calculated for these undirected networks (Table 3). Individual

centrality metrics generated from the multiplex grooming networks were largely uncorrelated

with metrics from the uniplex grooming networks (mean correlation strength = 0.11, SD = 0.08,

Table S2). In contrast, centrality metrics from grooming and contact sitting (mean correlation

strength = 0.37, SD = 0.07) and multiplex and uniplex contact sitting networks (mean correlation

strength = 0.32, SD = 0.24) were moderately correlated. In contrast, the structure of the all

grooming and all contact sitting networks did not differ on any examined metric with the

exception of average edge-weight which was higher in grooming networks.

Table 3: Whole Network Metric Comparisons

Network	Multi vs. Uni Grooming	Multi vs. Uni Contact Sit	Contact Sit vs Grooming
Density	Multi = Uni	Multi = Uni	CS = GR
Modularity	Multi > Uni[*]	Multi > Uni[*]	CS = GR
Eigenvector Centralization	Multi = Uni	Multi = Uni	CS = GR
Avg Edge Weight	Multi = Uni[a]	Multi = Uni[a]	CS < GR[*]
Clustering Coefficient	Multi > Uni[*]	Multi > Uni[*]	CS = GR
Reciprocity	Multi > Uni[*]	-	-
Proportion Kin	Multi > Uni[*]	Multi > Uni[*]	CS = GR
Proportion Up Rank	Multi < Uni[*]	-	-
Rank Disparity	Multi < Uni[*]	Multi < Uni[*]	CS = GR
Rank/Eigenvector centrality correlation	Multi = Uni	Multi = Uni	CS = GR

Multi: Multiplex network; Uni: Uniplex network; GR: Grooming; CS: Contact Sit. Effect
indicates the overall difference between networks for all groups using a paired t-test. [a] Wilcoxon
test. [*] p < 0.05

Relationship Dimensionality and Biomarkers of Inflammation

IL-6. There were six models that had AIC within 2 of the best fit model, all of which included

metrics from both the multiplex and uniplex grooming networks. For all models, less connected

individuals in the multiplex grooming network (degree, closeness, or clustering coefficient) but

more connected individuals in the uniplex grooming (strength, closeness) network had higher

levels of IL-6, although these effects were not significant in all candidate models (Table 4). Due

to this directional consistency we present the results of the just the best fit model in Figure 3A-B.

Multiplex closeness was not correlated with uniplex closeness (r = -0.04). Contact sitting degree

was weakly negatively correlated with uniplex closeness (-0.13) but more strongly correlated

with multiplex closeness (r = 0.77) yet analysis of VIF (< 2.5) and tolerance (≥ 0.4) did not

indicate any issues with collinearity.

Table 4: IL-6 Candidate Model Outputs

Model Parameters	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
Intercept	0.03	2.8**	1.6***	-0.7	1.8***	1.8**
Uni GR Strength	-	0.05**	0.04**	-	0.04**	0.04**
Uni GR Closeness	7.6**	-	-	6.2**	-	-
Multi GR Degree	-	-	-	-	-	-0.03
Multi GR Closeness	-3.6*	-3.2	-	-	-	-
Multi GR Clustering	-	-	-	-	-0.4	-
ΔAIC	0.00	0.30	0.70	1.40	1.90	1.90

414   * p < 0.05, ** p < 0.01. Uni = Uniplex, Multi = Multiplex, GR = Grooming network. All models
415   were run using a negative binomial distribution and included a random effect of group.
416

417
Figure 2. Effects of affiliative centrality
on cytokines. Effects of multiplex (Multi)
grooming closeness (A) and uniplex (Uni)
grooming closeness (B) on levels of IL-6
with 95% confidence intervals (Model 1).
Effects of multiplex grooming closeness
(C) and uniplex grooming closeness (D)
on levels of TNFα with 95% confidence
intervals (Model 1).

TNF-α. There were five candidate models

identified; the top four of these included

comparisons of multiplex and uniplex

grooming centrality and the last included

centrality metrics from the grooming and contact sitting networks (see Table 5). As with IL-6,

lower centrality in the multiplex grooming network (degree or closeness) but higher centrality in

the uniplex grooming network (strength or closeness) were consistently associated with higher

levels of TNF-α (Table 5, Figure 3C-D). In the fifth candidate model, higher eigenvector

centrality in the grooming network but lower closeness centrality in the contact sitting network

were associated with higher levels of TNF- α.

Table 5: TNF-α Candidate Model Outputs

Model Parameters	Model 1	Model 2	Model 3	Model 4	Model 5
Intercept	2.9*	6.6**	4.9**	0.9	8.3***
Uni GR Strength	-	0.06**	0.06**	-	-
Uni GR Closeness	9.4**	-	-	10.4**	-
Multi GR Degree	-	-	-0.1*	-0.1*	-
Multi GR Closeness	-5.9**	-6.0**	-	-	-
All GR Eigenvector	-	-	-	-	1.7**
All CS Closeness	-	-	-	-	-8.0**
ΔAIC	0.00	0.10	1.30	1.60	1.80

440 * $p < 0.05$, ** $p < 0.01$. Uni = Uniplex, Multi = Multiplex, GR = Grooming network, CS =
441 Contact sitting network. All models were run using a negative binomial distribution and included
442 a random effect of group.
443

Discussion

Social primates have a complex web of differentiated social relationships, which vary in

their structure and function. While strong affiliative social relationships are usually associated

with better health, less is known on how the multidimensionality or integration of different

affiliative behaviors within a social relationship might impact health. We identified affiliative

relationships that were multiplex (animals affiliated using both grooming and contact sitting

behavior) versus uniplex (animals only groomed or only contact sat). Examination of these

networks revealed that they differed in topology, kinship, and associations with rank. Multiplex

networks were more modular, clustered, reciprocal, had higher average edge weights, and were

more strongly associated with kinship. In contrast, dyads in the uniplex networks tended to be of

more disparate ranks. Notably, these differences in kinship and rank between multiplex and

uniplex networks were not apparent when looking at all grooming or contact sitting interactions.

The health impacts of these two networks differed as well, with females that were *less* socially

embedded in multiplex grooming networks exhibiting higher levels of biomarkers of

inflammation (IL-6 and TNF-α), whereas females *more* socially connected in uniplex grooming

networks exhibited higher levels of biomarkers of inflammation. Notably, these effects were

primarily seen in multiplex and uniplex grooming, not in the other networks tested. These

results suggest that grooming which occurs in the context of multiplex affiliative relationships

may result in health benefits (i.e., reduced inflammation) while grooming occurring in uniplex

affiliative relationships may have potential costs.

Networks consisting of dyads with multiplex relationships showed differences from

uniplex relationships in network topology, kinship, and associations with dominance. Multiplex

networks had structural characteristics consistent with strong bonds or supportive affiliative

relationships[23,65,66]. Specifically, interactions in the multiplex networks were more likely to be

reciprocal, frequent (i.e., higher edge-weight), clustered, and associated with kinship, suggesting

they are relationships that are regularly maintained and potentially more stable across time[23,66].

Previous methods demonstrating that strong bonds enhance fitness, particularly those using

sociality indices, have also used multiple behaviors to assess relationship strength (e.g.,

grooming and proximity[12,26,27]). However, these methods rely on total duration or frequency of

affiliation to describe relationships rather than characterizing the breadth or dimensionality of the

relationships (e.g., dyads can have high DSI through grooming, proximity, or both). Similar to

strong bonds, multiplex affiliative relationships may improve health and fitness by buffering

individuals from the negative impacts of stress, improving predator detection, promoting

offspring survival, and improving social stability[14,16,67,68].

Also consistent with the literature on strong affiliative bonds, being well connected to

others was associated with biomarkers of better health. Specifically, the negative association

between multiplex grooming centrality (e.g., degree, closeness centrality, or clustering

coefficient) and biomarkers of inflammation indicated that individuals that were generally well

connected in the network (e.g., at the core of the group) may be at lower risk for inflammation

related diseases[57]. Our results add to the literature suggesting that strong bonds may improve

fitness by altering endocrine and immune function[13,25,69,70]. Consistent with this idea, Yang et

al.[71] found in humans that socially integrated individuals (i.e., those with more social

connections across multiple domains) exhibited lower inflammation, whereas social strain (e.g.,

higher levels of family criticism or demands) was associated with greater inflammation. Given

that familial and friend relationships tend to endure through extended periods, often persisting

over decades (in both humans and NHPs), these relationships may have an important and long-

lasting impact on health.

Uniplex grooming relationships may reflect relationships that are more transactional in

nature [72]. The fact that uniplex grooming relationships are less kin biased but likely to occur

between dyads of more disparate ranks suggests that these relationships may be more related to

grooming being used as a commodity in exchange for tolerance or support from higher ranking

animals. These relationships are likely more transactional in nature, reflecting a desire to

maintain peace/tolerance or used in a biological market exchange[40,41], rather than reflecting a

strong affiliative relationship. The positive association between females' connectedness in

uniplex grooming networks and biomarkers of inflammation suggests that uniplex grooming

relationships may not be supportive on their own and instead are associated with increased

physiological costs, at least in the short term. Specifically, predictors of inflammation in the

uniplex grooming networks included strength or closeness centrality. However, the various

network centrality metrics from the uniplex grooming network were more highly correlated with

each other than the other networks, and therefore it is difficult to identify which specific aspect

of uniplex grooming centrality might be driving these effects. However, collectively this group

of candidate predictors indicates that greater general connectedness (direct and indirect) in

uniplex grooming was associated with increased inflammation. Uniplex grooming relationships

were maintained through generally less frequent interactions that were more likely to occur

between animals of disparate ranks which may result in greater uncertainty regarding the

outcome of any given interaction. This uncertainty may be stressful, and therefore have at least

short-term physiological costs[73]. If these relationships are more transactional in nature, reflecting

a desire to maintain peace/tolerance or used in a biological market exchange[40,41], then

maintaining more of these transactional relationships may result in increased stress, which if

sustained can result in long-term physiological costs[13]. It is possible that these short-term costs

are actually investments that may manifest in future benefits (e.g., tolerance, alliance support)

that would offset these costs, yet this is difficult to test as the "commodities" exchanged may be

heterogeneous and the time-scale for market exchanges is often unclear[74]. However, other work

points to benefits of weak or economically based bonds to survival and reproduction[9,26]

(although see[12]). While these types of connections may have ultimate fitness benefits (e.g.,

alliance support, increased access to food), this research suggests they may also be associated

with proximate costs.

Conclusion

Both humans and many species of NHPs engage in a complex interconnected system of

social interactions. Understanding the mechanisms by which social relationships impact health

and fitness remains a challenge. Decades of research has established that affiliative social

relationships can benefit health, however, the complexity and multidimensionality of

relationships has yet to be explored. By utilizing a network approach, we were able to

characterize two types of affiliative social relationships that differed in their network topology,

kin bias, associations with rank, and importantly their associations with biomarkers of

inflammation. Our research has indicated that features of multiplex affiliative relationships are

consistent with the concept of a strong supportive relationships and may support health and

fitness. In contrast, more transactional affiliative relationships (e.g., uniplex affiliation) may

incur short-term health costs yet may result in ultimate benefits through commodity exchange.

Still unclear is whether these effects are specific to the combination of behaviors used here (i.e.,

contact sitting and grooming), or if other affiliative behaviors (e.g., proximity) might provide

similar information. Further research into the dimensionality of relationships might reflect

different qualities or functions of relationships is needed. However, this complexity is important

to consider for understanding the mechanisms underlying the impact of social relationships on

human and NHP health.

540 **Acknowledgements**

541 We thank the data collection team: A. Barnard, T. Boussina, E. Cano, H. Caparella, C.

542 Carminito, J. Greco, M. Jackson, A. Maness, S. Seil, N. Sharpe, A. Vitale, & S. Winkler. This

543 research was funded by an NIH grant awarded to BM (R01-HD068335) and the California

544 National Primate Research Center base grant (P51-OD01107-53). This is an updated version of

545 a manuscript on the PeerJ preprint server (https://doi.org/10.7287/peerj.preprints.27961v1).

References

1. Holt-Lunstad, J., Smith, T. B. & Layton, J. B. Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine* **7**, (2010).

2. McCowan, B. *et al.* Connections Matter: Social Networks and Lifespan Health in Primate Translational Models. *Front Psychol.* **7**, (2016).

3. Ostner, J. & Schülke, O. Linking Sociality to Fitness in Primates: A Call for Mechanisms. *Advances in the Study of Behavior* **50**, 127–175 (2018).

4. Hinde, R. A. ON DESCRIBING RELATIONSHIPS*. *Journal of Child Psychology and Psychiatry* **17**, 1–19 (1976).

5. Hinde, R. A. Interactions, Relationships and Social-Structure. *Man* **11**, 1–17 (1976).

6. Silk, J. B., Cheney, D. & Seyfarth, R. A practical guide to the study of social relationships. *Evolutionary Anthropology* **22**, 213–225 (2013).

7. Archie, E. A., Tung, J., Clark, M., Altmann, J. & Alberts, S. C. Social affiliation matters: both same-sex and opposite-sex relationships predict survival in wild female baboons. *Proc Biol Sci* **281**, (2014).

8. Berkman, L. F., Glass, T., Brissette, I. & Seeman, T. E. From social integration to health: Durkheim in the new millennium. *Social Science and Medicine* **51**, 843–857 (2000).

9. Ellis, S., Snyder-Mackler, N., Ruiz-Lambides, A., Platt, M. L. & Brent, L. J. N. Deconstructing sociality: the types of social connections that predict longevity in a group-living primate. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20191991 (2019).

10. Lehmann, J., Majolo, B. & McFarland, R. The effects of social network position on the survival of wild Barbary macaques, Macaca sylvanus. *Behavioral Ecology* **27**, 20–28 (2016).

11. Mendes de Leon, C. F. *et al.* Social Networks and Disability Transitions Across Eight

Intervals of Yearly Data in the New Haven EPESE. *The Journals of Gerontology Series B:*

Psychological Sciences and Social Sciences **54B**, S162–S172 (1999).

12. Silk, J. B., Seyfarth, R. M. & Cheney, D. L. Quality versus quantity: do weak bonds enhance

the fitness of female baboons? *Animal Behaviour* **140**, 207–211 (2018).

13. Snyder-Mackler, N. *et al.* Social determinants of health and survival in humans and other

animals. *Science* **368**, (2020).

14. Balasubramaniam, K. N., Beisner, B. A., Vandeleest, J., Atwill, E. & McCowan, B. Social

buffering and contact transmission: network connections have beneficial and detrimental

effects on Shigella infection risk among captive rhesus macaques. *PeerJ* **2016**, e2630 (2016).

15. Brent, L. J. N. *et al.* Genetic origins of social networks in rhesus macaques. *Scientific*

Reports 2013 3:1 **3**, 1–8 (2013).

16. Cheney, D. L., Silk, J. B. & Seyfarth, R. M. Network connections, dyadic bonds and fitness

in wild female baboons. *Royal Society Open Science* **3**, 160255 (2016).

17. Grando, F., Noble, D. & Lamb, L. C. An Analysis of Centrality Measures for Complex and

Social Networks. in *2016 IEEE Global Communications Conference (GLOBECOM)* 1–6

(IEEE, 2016). doi:10.1109/GLOCOM.2016.7841580.

18. Meghanathan, N. Correlation Coefficient Analysis of Centrality Metrics for Complex

Network Graphs. in *Intelligent Systems in Cybernetics and Automation Theory* (eds. Silhavy,

R., Senkerik, R., Oplatkova, Z. K., Prokopova, Z. & Silhavy, P.) 11–20 (Springer

International Publishing, 2015). doi:10.1007/978-3-319-18503-3_2.

19. Makagon, M. M., McCowan, B. & Mench, J. A. How can social network analysis contribute

to social behavior research in applied ethology? *Applied Animal Behaviour Science* **138**,

152–161 (2012).

20. VanderWaal, K. L., Atwill, E. R., Isbell, L. A. & McCowan, B. Quantifying microbe

transmission networks for wild and domestic ungulates in Kenya. *Biological Conservation*

169, 136–146 (2014).

21. Wasserman, S. & Faust, K. *Social Network Analysis: Methods and Applications*.

(Cambbridge University Press, 1994).

22. Rodrigues, F. A. Network Centrality: An Introduction. in *A Mathematical Modeling*

Approach from Nonlinear Dynamics to Complex Systems (ed. Macau, E. E. N.) 177–196

(Springer International Publishing, 2019). doi:10.1007/978-3-319-78512-7_10.

23. Silk, J. B. *et al.* Female chacma baboons form strong, equitable, and enduring social bonds.

Behavioral Ecology and Sociobiology **64**, 1733 (2010).

24. Silk, J. B. Evolutionary perspectives on the links between close social bonds, health, and

fitness. in *Sociality, Hierarchy, Health: Comparative Biodemography: A Collection of*

Papers 121–144 (National Academies Press (US), 2014). doi:10.17226/18822.

25. Crockford, C. *et al.* Urinary oxytocin and social bonding in related and unrelated wild

chimpanzees. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20122765

(2013).

26. McFarland, R. *et al.* The 'strength of weak ties' among female baboons: fitness-related

benefits of social bonds. *Animal Behaviour* **126**, 101–106 (2017).

27. Fischer, J., Farnworth, M. S., Sennhenn-Reulen, H. & Hammerschmidt, K. Quantifying

social complexity. *Animal Behaviour* **130**, 57–66 (2017).

28. Schülke, O., Wenzel, S. & Ostner, J. Paternal Relatedness Predicts the Strength of Social

Bonds among Female Rhesus Macaques. *PLoS ONE* **8**, e59789 (2013).

29. Beisner, B. A. *et al.* A multiplex centrality metric for complex social networks: Sex, social

status, and family structure predict multiplex centrality in rhesus macaques. *PeerJ* **2020**,

(2020).

30. Vandeleest, J. J. *et al.* Decoupling social status and status certainty effects on health in

macaques: a network approach. *PeerJ* (2016) doi:10.7717/peerj.2394.

31. Snyder-Mackler, N. *et al.* Social status alters immune regulation and response to infection in

macaques. *Science* **354**, 1041–1045 (2016).

32. Brent, L. J. N., Ruiz-Lambides, A. & Platt, M. L. Family network size and survival across

the lifespan of female macaques. *Proceedings of the Royal Society B: Biological Sciences*

284, (2017).

33. Capitanio, J. P., Cacioppo, S. & Cole, S. W. Loneliness in monkeys: neuroimmune

mechanisms. *Current Opinion in Behavioral Sciences* **28**, 51–57 (2019).

34. Pallante, V., Ferrari, P. F., Gamba, M. & Palagi, E. Embracing in a Female-Bonded Monkey

Specie (Theropithecus gelada). *Journal of Comparative Psychology* **133**, 442–451 (2019).

35. Aureli, F. & Schaffner, C. M. Aggression and conflict management at fusion in spider

monkeys. *Biology Letters* **3**, 147–149 (2007).

36. Sussman, R. w., Garber, P. A. & Cheverud, J. M. Importance of cooperation and affiliation

in the evolution of primate sociality. *American Journal of Physical Anthropology* **128**, 84–97

(2005).

37. Dunbar, R. I. M. Functional Significance of Social Grooming in Primates. *Folia*

Primatologica **57**, 121–131 (1991).

38. Balasubramaniam, K. N. *et al.* Social network community structure and the contact-mediated

sharing of commensal E. coli among captive rhesus macaques (Macaca mulatta). *PeerJ* **6**,

e4271 (2018).

39. Lehmann, J., Korstjens, A. H. & Dunbar, R. I. M. Group size, grooming and social cohesion

in primates. *Animal Behaviour* **74**, 1617–1629 (2007).

40. Borgeaud, C. & Bshary, R. Wild Vervet Monkeys Trade Tolerance and Specific Coalitionary

Support for Grooming in Experimentally Induced Conflicts. *Current Biology* **25**, 3011–3016

(2015).

41. Schino, G. & Aureli, F. Grooming reciprocation among female primates: A meta-analysis.

Biology Letters **4**, 9–11 (2008).

42. Snyder-Mackler, N. *et al.* Social status drives social relationships in groups of unrelated

female rhesus macaques. *Animal Behaviour* **111**, 307–317 (2016).

43. Campbell, L. A. D., Tkaczynski, P. J., Lehmann, J., Mouna, M. & Majolo, B. Social

thermoregulation as a potential mechanism linking sociality and fitness: Barbary macaques

with more social partners form larger huddles. *Scientific Reports* **8**, 1–8 (2018).

44. Uchino, B. N., Vaughn, A. A., Carlisle, M. & Birmingham, W. Social Support and

Immunity. in *The Oxford Handbook of Psychoneuroimmunology* (Oxford University Press,

2012). doi:10.1093/oxfordhb/9780195394399.013.0012.

45. *Macaque Societies: A Model for the Study of Social Organization*. (Cambridge University

Press, 2004).

46. Berman, C. M. Early agonistic experience and rank acquisition among free-ranging infant

rhesus monkeys. *International Journal of Primatology* **1**, 153–170 (1980).

47. Fushing, H., McAssey, M. P., Beisner, B. A. & McCowan, B. Ranking network of a captive

rhesus macaque society: a sophisticated corporative kingdom. *PLoS ONE* **6**, e17817 (2011).

48. Georgiev, A. V. *et al.* Breaking the succession rule: The costs and benefits of an alpha-status

take-over by an immigrant rhesus macaque on Cayo Santiago. *Behaviour* **153**, 325–351

(2016).

49. Chapais, B., Girard, M. & Primi, G. Non-kin alliances, and the stability of matrilineal

dominance relations in Japanese macaques. *Animal Behaviour* **41**, 481–491 (1991).

50. Schulke, O., Bhagavatula, J., Vigilant, L. & Ostner, J. Social bonds enhance reproductive

success in male macaques. *Curr Biol* **20**, 2207–2210 (2010).

51. Drickamer, L. C. Quantitative observations of grooming behavior in free-ranging Macaca

mulatta. *Primates* **17**, 323–335 (1976).

52. Brent, L. J. N., Ruiz-Lambides, A. & Platt, M. L. Persistent social isolation reflects identity

and social context but not maternal effects or early environment. *Scientific Reports* **7**, 1–11

(2017).

53. Fujii, K. *et al.* Perc: Using Percolation and Conductance to Find Information Flow Certainty

in a Direct Network. (2016).

54. Sosa, S., Sueur, C. & Puga-Gonzalez, I. Network measures in animal social network

analysis: Their strengths, limits, interpretations and uses. *Methods in Ecology and Evolution*

(2020) doi:10.1111/2041-210X.13366.

55. Kim, D.-Y. Closeness Centrality: A Social Network Perspective. *Journal of International &*

Interdisciplinary Business Research **6**, 115–122 (2019).

56. Croft, D. P., James, R. & Krause, J. *Exploring animal social networks*. (Princeton University

Press, 2008).

57. Bruunsgaard, H. *et al.* Predicting death from tumour necrosis factor-alpha and interleukin-6

in 80-year-old people. *Clinical and Experimental Immunology* **132**, 24–31 (2003).

58. Gan, W. Q., Man, S. F. P., Senthilselvan, A. & Sin, D. D. Association between chronic

obstructive pulmonary disease and systemic inflammation: a systematic review and a meta-

analysis. *Thorax* **59**, 574–580 (2004).

59. Holmes, C. *et al.* Systemic inflammation and disease progression in Alzheimer disease.

Neurology **73**, 768–774 (2009).

60. Friedman, E. M. *et al.* Social relationships, sleep quality, and interleukin-6 in aging women.

Proceedings of the National Academy of Sciences of the United States of America **102**,

18757–18762 (2005).

61. Marucha, P. T., Crespin, T. R., Shelby, R. A. & Andersen, B. L. TNF-α levels in cancer

patients relate to social variables. *Brain, Behavior, and Immunity* **19**, 521–525 (2005).

62. Balasubramaniam, K. N. *et al.* The influence of phylogeny, social style, and

sociodemographic factors on macaque social network structure. *American Journal of*

Primatology **80**, e22727 (2018).

63. Sueur, C. *et al.* A comparative network analysis of social style in macaques. *Animal*

Behaviour **82**, 845–852 (2011).

64. Parker, K. J. *et al.* Arginine vasopressin in cerebrospinal fluid is a marker of sociality in

nonhuman primates. *Science Translational Medicine* **10**, 9100 (2018).

65. Silk, J. B. *et al.* The benefits of social capital: close social bonds among female baboons

enhance offspring survival. *Proc Biol Sci* **276**, 3099–3104 (2009).

66. Silk, J. B. Looking inside the black box: Mechanisms linking social behavior to health and

fitness. *Evol Anthropol* **23**, 83–84 (2014).

67. Beisner, B. A., Jackson, M. E., Cameron, A. N. & McCowan, B. Detecting instability in

animal social networks: genetic fragmentation is associated with social instability in rhesus

macaques. *PLoS ONE* **6**, e16365 (2011).

68. Micheletta, J. *et al.* Social bonds affect anti-predator behaviour in a tolerant species of

macaque, Macaca nigra. *Proceedings of the Royal Society B: Biological Sciences* **279**, 4042–

4050 (2012).

69. Boccia, M. L. *et al.* Juvenile friends, behavior, and immune responses to separation in bonnet

macaque infants. *Physiology and Behavior* **61**, 191–198 (1997).

70. Shutt, K., MacLarnon, A., Heistermann, M. & Semple, S. Grooming in Barbary macaques:

Better to give than to receive? *Biology Letters* **3**, 231–233 (2007).

71. Yang, Y. C. *et al.* Social relationships and physiological determinants of longevity across the

human life span. *Proceedings of the National Academy of Sciences of the United States of*

America **113**, 578–583 (2016).

72. Silk, J. B. Using the 'F'-word in primatology. in *Behaviour* vol. 139 421–446 (Brill

Academic Publishers, 2002).

73. Schino, G. & Alessandrini, A. Short-term costs and benefits of grooming in Japanese

macaques. *Primates* **56**, 253–257 (2015).

74. Dunayer, E. S. & Berman, C. M. Biological markets: theory, interpretation, and proximate

perspectives. A response to Sánchez-Amaro and Amici (2015). *Animal Behaviour* **121**, 131–

136 (2016).

725