1 **Heterogeneities in infection outcomes across species: examining sex and tissue differences**
2 **in virus susceptibility**
3

4 Katherine E Roberts[1] and Ben Longdon[1*]
5
6 [1] Centre for Ecology and Conservation, University of Exeter, Penryn Campus, TR10 9FE, UK
7 *corresponding author
8
9 Katherine Roberts bskr13@gmail.com ORCID 0000-0002-8567-3743
10 Ben Longdon b.longdon2@exeter.ac.uk ORCID 0000-0001-6936-1697

11
12
13 **Abstract**
14

Species vary in their susceptibility to pathogens, and this can alter the ability of a pathogen
to infect a novel host. However, many factors can generate heterogeneity in infection
outcomes, obscuring our ability to understand pathogen emergence. Such heterogeneities
can alter the consistency of responses across individuals and host species. For example,
sexual dimorphism in susceptibility means males are often intrinsically more susceptible
than females (although this can vary by host and pathogen). Further, we know little about
whether the tissues infected by a pathogen in one host are the same in another species, and
how this relates to the harm a pathogen does to its host. Here, we first take a comparative
approach to examine sex differences in susceptibility across 31 species of Drosophilidae
infected with Drosophila C Virus (DCV). We found a strong positive inter-specific correlation
in viral load between males and females, with a close to 1:1 relationship, suggesting that
susceptibility to DCV across species is not sex specific. Next, we made comparisons of the
tissue tropism of DCV across seven species of fly. We found differences in viral load between
the tissues of the seven host species, but no evidence of tissues showing different patterns
of susceptibility in different host species. We conclude that, in this system, patterns of viral
infectivity across host species are robust between males and females, and susceptibility in a
given host is general across tissue types.

32
33 **Introduction**
34
Emerging pathogens often arise from a host shift event – where a pathogen jumps into and
establishes in a novel host species. Species vary in their susceptibility to pathogens, but little
is known about the factors underlying this variation, and whether differences between
clades are due to the same or different factors [1, 2]. Understanding this is critical for
determining which hosts pathogens are likely to jump between, and the harm they cause to
their hosts. The host phylogeny has been shown to be an important determinant of host
shifts in a range of systems [3-8] as well as being important for understanding how pathogen
virulence may change when a pathogen finds itself in a new host [9, 10]. For example,
virulence tends to increase, and onward transmission and pathogen load decrease, with
greater evolutionary distance between donor and recipient hosts [10-12] . In addition,

clades of closely related species tend to have similar levels of susceptibility independent
from their distance to the pathogens natural host [4, 9].
47
However, such patterns of susceptibility across species can be affected by heterogeneities in
infection outcomes. These can be viewed at the level of the individual heterogeneities
within species, and by looking at whether these heterogeneities are consistent or different
across host species. Experimental studies typically try and minimise within species or
environmental effects. Often overlooked is variation arising from sexual dimorphism, as
typically only one sex is utilized to remove between sex differences [13]. Sexual dimorphism
is seen across most animal systems in a range of life history traits from body size, growth
rate, reproductive effort and immunity [14-16]. In mammals, males and females often differ
in their pathogen burdens and mortality rates [17]. For example, in SARS-COV-2 infection in
humans, women have a lower risk of morbidity and mortality than men [18]. In HIV infected
individuals, women have up to 40% lower HIV viral RNA in circulation but a greater
likelihood of developing AIDS than men with matched viral loads [19]. Sexually transmitted
infections – which are primarily transmitted between the sexes – are particularly prone to
sex biased infection through either exposure differences or sex biased virulence [20-22]. Sex
biases in parasitism rates in mammals have been suggested to be due to males investing in
traits that favour their reproductive success, which trade off against somatic maintenance,
including immunity. In support of this, sex biased parasitism is positively correlated with
sexual size dimorphism [17]. In insects a comparative analysis found the degree of sex
biased parasitism and mortality is explained by an interaction between the mating system
(polygynous vs non-polygynous) and sexual size dimorphism [23]. This difference is
consistent with parasites having a greater impact on the survival of male insects compared
to females (particularly in polygynous species where males are larger than females). Other
hypotheses suggest that as longevity is a major determinant of female fitness, investment in
costly immune responses are more important than for males, who can maximise their fitness
with shorter term mating success [24]. As such, investment in reproduction may trade-off
with immunity in different ways between males and females. There is some support for this
hypothesis in mammals but a lack of supporting data for insects [15]. However, many
experimental studies of host-parasite interactions do not compare differences between
sexes [13]. Furthermore, for most pathogens we have little understanding of whether sex
differences are consistent across host species, which has important implications for our
understanding of pathogen emergence.
79
Despite phylogenetic patterns of host susceptibility having been observed in a range of
systems [3, 6-8], we know little about why species vary in their susceptibilities. For example,
given equal exposure why do we see high mortality in some species but little in others? One
factor that appears important in determining the severity of disease is the tissue tropism of
a pathogen. In humans, RNA viruses with neural tropism or generalised systemic tropism
tend to result in severe disease [25]. In terms of the patterns of susceptibility across species,
virulence may be a consequence of a virus infecting a sensitive tissue or organ resulting in
damage by the pathogen directly or by autoimmunity. For example, in bacterial meningitis
the pathology is a consequence of bacteria infecting the cerebrospinal fluid and resulting in
inflammatory autoimmune damage to the central nervous system [26]. Alternatively, it may

be due to the pathogen getting into a particularly permissive tissue type and proliferating to
high levels. Virus macroevolutionary change is thought to be driven by cross-species
transmission or codivergence rather than by acquiring new niches – or tissues – within a
host [27]. Likewise, the host specificities of viruses are thought to be more labile than tissue
specificities [28]. As such, heterogeneity in the tissues infected may explain variation in
susceptibility across host species.
96
Here we use a *Drosophila*-virus system to examine the factors underlying susceptibility
across host species. Across *Drosophila* species many physiological traits show sexual
differentiation [14]. Using a comparative approach, we firstly ask if the patterns of infection
seen across the host phylogeny [3, 5, 9, 29] differ between males and females. We infected
both males and females of a panel of 31 species of Drosophilidae with Drosophila C Virus
(DCV), a positive sense RNA virus in the family Dicistroviridae. Males of *Drosophila*
melanogaster have previously been reported to have higher viral loads than females [30]
and show greater rates of shedding, lower clearance and higher transmission potential of
DCV, although these traits can interact with host genotype [31]. Viral load has shown to
have a strong positive correlation with mortality across host species [9, 29]. DCV has been
reported to show tissue tropism in *D. melanogaster,* with high levels of infection in the heart
tissue, fat body, visceral muscle cells around the gut (midgut) and food storage organ (crop)
[32, 33]. To test if the same patterns of tissue infection were observed across species, we
then made comparisons of the tissue tropism of DCV in 7 species of fly.
111
112 **Methods**
113
Viral Infections
Thirty one species of Drosophilidae were used to examine sex differences in viral infection.
Stock populations were reared in the laboratory in multi generation populations, in
Drosophila stock bottles (Fisherbrand) on 50 ml of their respective food medium (Table S1)
at 22˚C and 70% relative humidity with a 12-hour light-dark cycle. Flies were then collected
twice a day in order to try and control for age of maturity in an effort to minimize the
chances that flies would have reached sexual maturity and mated before the sexes were
separated out. Although no effect of mating status on DCV viral load was previously
observed in *D. melanogaster*, this can vary by host genotype and mating status is known to
affect susceptibility to other pathogens [31].
124
To examine differences in viral load between males and females, two vials of 0-1 day old
males flies and two vials of 0-1 day old female flies were collected daily for each species.
Flies were tipped onto fresh vials of food every day to minimise differences in the
microbiomes of flies (Broderick & Lemaitre, 2012; Blum *et al.*, 2013). All vials were kept for
10 days in order to check for larvae, as a sign of successful mating. Only 4 vials from 3
species were found to contain larvae, these were one vial of *D. sturtevanti,* two vials of
Scaptodrosophila lativittata and one vial of *Zaprionus tuberculatus.* After 3 days flies were
experimentally infected with DCV. Three replicate blocks were carried out, with each block
being completed over consecutive days. The order of experimental infection was
randomized each day so that both sex and species were randomised. We carried out three

biological replicates for each species for each sex at time zero and 2 days post infection.
There was a mean of 17 flies per replicate (range across species = 12-20).
137
Viral challenge was carried out by needle inoculation of Drosophila C virus (DCV) strain B6A
[34], derived from an isolate collected from *D. melanogaster* in Charolles, France [35]. The
virus was prepared as described previously [36]. DCV was grown in Schneider's Drosophila
line 2 cells and the Tissue Culture Infective Dose 50 (TCID50) per ml was calculated using the
Reed-Muench end-point method. Flies were anesthetized on CO_2 and inoculated using a
0.0125 mm diameter stainless steel needle bent to a right angle ~0.25mm from the end
(Fine Science Tools, CA, USA). The bent tip of the needle was dipped into the DCV solution
(TCID50 = 6.32×10^9) and pricked into the anepisternal cleft in the thorax of the flies [9, 37].
This mode of infection is used as it creates a more reproducible infection that oral
inoculation, which is found to cause stochastic infection outcomes in *D. melanogaster* [32].
Both methods of infection have been shown to produce systemic infections with the same
tissues ultimately becoming infected [32].
150
To control for relative viral dose between species a time point zero sample of one vial of flies
was immediately snap frozen in liquid nitrogen as soon as they were inoculated. The second
vial of flies were inoculated and placed onto a fresh vial of food, and returned to the
incubator. Two days after challenge (+/- 1 hour) these flies were snap frozen in liquid
nitrogen. This time point is chosen as the sampling time point as previous studies show a
clear increase in viral growth but little mortality at this point in infection [5, 29]. Each
experimental block contained a day 0 and day 2 replicate for each sex and each species (31
species × 2 sexes × 3 experimental blocks).
159
Measuring the change in viral load
Using quantitative Reverse Transcription PCR (qRT- PCR) we measured the change in viral
load in male and female flies from day 0 to day 2 post- infection. Total RNA was extracted
from the snap frozen flies by homogenizing them in Trizol reagent (Invitrogen) using a bead
homogenizer for 2 pulses of 10 seconds (Bead Ruptor 24; Omni international) and stored at
−70°C for later extraction. Samples were defrosted and RNA extracted as described
previously [29]. Briefly, Trizol homogenized flies were processed in a chloroform isopropanol
extraction, eluted in water and reverse- transcribed with Promega GoScript reverse
transcriptase (Sigma) and random hexamer primers. Quantification of the change in viral
RNA load was calculated in relation to a host endogenous control, the housekeeping gene
RpL32. Primers were designed to match the homologous sequence for each of the
experimental species that crossed an intron– exon boundary so will only amplify mRNA. qRT-
PCR was carried out on 1:10 diluted cDNA using Sensifast Hi-Rox Sybr kit (Bioline). Two qRT-
PCR reactions (technical replicates) were carried out per sample with both the viral and
endogenous control primers. All melt curves were checked to verify that the correct
products were being amplified. All experimental plates had experimental replicates
distributed across the plates in a randomized block design to control for between plate
differences. Each qRT- PCR plate contained three standard samples. A linear model which
included plate ID and biological replicate ID was used to correct the cycle threshold (C_t)
values between plates. Any technical replicates that had C_t values more than two cycles

apart after the plate correction were repeated. Change in viral load was calculated as the
mean C_t value of the pairs of technical replicates. We then used these to calculate the ΔC_t as
the difference between the cycle thresholds of the viral DCV qRT- PCR and the RpL32
endogenous control for each sample. The Ct of the day 2 flies relative to day 0 flies was then
calculated as, $2^{-\Delta\Delta Ct}$; where $\Delta\Delta C_t = \Delta C_t$ day0 $- \Delta C_t$ day2.
185
Body Size
We measured wing size of the flies to control for between species and sex differences in
body size. In Drosophilidae wing length has been shown to be a good proxy for body size
(Huey *et al.*, 2006). For measurement wings were removed from a mean of 15 male and
females flies of each species (range 10–18), stored in 80% ethanol, and later photographed
under a dissecting microscope. The length of the IV longitudinal vein from the tip of the
proximal segment to where the distal segment joins vein V was recorded Using ImageJ
software (version 1.48), and the mean taken for each sex of each species.
194
Inferring the host phylogeny
We used a previously inferred phylogenetic tree [5] using seven genes (mitochondrial; *COI,*
COII, ribosomal; *28S* and nuclear; A*dh, SOD, Amyrel, RpL32*). Briefly, we downloaded publicly
available sequences from Genbank and where these were not available they were Sanger
sequenced from our laboratory stocks. For each gene the sequences were aligned in
Geneious (version 9.1.8, www.geneious.com) [38] using the global alignment setting, with
free end gaps and 70% similarity IUB cost matrix. The phylogeny was inferred using these
genes and the BEAST programme (v1.10.4) [39]. Genes were partitioned into three groups;
mitochondria, ribosomal and nuclear, each with separate relaxed uncorrelated lognormal
molecular clock models using random starting trees. Each of the partitions used a HKY
substitution model with a gamma distribution of rate variation with 4 categories and
estimated base frequencies. Additionally, the mitochondrial and nuclear data sets were
partitioned into codon positions 1+2 and 3, with unlinked substitution rates and base
frequencies across codon positions. The tree-shape prior was set to a birth-death process.
We ran the BEAST analysis three times to ensure convergence for 1000 million MCMC
generations sampled every 10000 steps. On completion the MCMC process was examined
by evaluating the model trace files using the program Tracer (version 1.7.1) (Rambaut *et al.*,
2014) to ensure convergence and adequate sampling. The consensus constructed tree was
then visualised using FigTree (v1.4.4) (Rambaut, 2006).
214
Tissue Tropism
In order to examine patterns of tissue infection across species we infected 7 species of flies
used above; *D. melanogaster, D. stuventi, S. lativaitata, D.pseudooscura, D. virilis, D.*
prosaltans and *D.littoralis*). Male flies were infected with DCV using the same inoculation
method as described above. Two days post infection flies were placed on ice to sedate them,
they were then surface sterilized in ice-cold 70% ethanol before being dissected. The head,
crop, gut (all parts), malpighian tubules, sex organs (testis and accessory glands) and
abdominal cuticle including the attached fat body (hereafter referred to as body) were
dissected from each male fly and placed into individual tubes on ice. Six individual flies were
pooled per replicate and then snap frozen in liquid nitrogen for later RNA extraction. For

each species there were six replicate pools of each of the six tissue types. At the same time
as the dissections were carried out whole flies were snap frozen for a "whole fly"
comparative viral load measure. All samples were processed as per the methods for viral
load quantification as described above.
229
230   **Statistical analysis**
231   *Sex differences*
232   Viral load in males and females were analysed using phylogenetic mixed models. We fitted
233   all models using a Bayesian approach in the R package MCMCglmm [40, 41]. We used a
234   multivariate model with viral load of each sex as the response variable.
235
236   The models took the form of:
237   (1) $y_{hit} = \beta_{1:t} + wingsize\beta_{2:t} + u_{p:ht} + u_{s:ht} + e_{hit}$
238   (2) $y_{hit} = \beta_{1:t} + wingsize\beta_{2:t} + u_{p:ht} + e_{hit}$
239
where y is the change in viral load of the *i*th biological replicate of host species *h*, for trait *t*
(male or female). *β* are the fixed effects, with *β₁* being the intercepts for each trait and *β₂*
being the effect of wing size. u_p are the random phylogenetic species effects, and *e* the
model residuals. Models were also run which included the mating status of the species as a
fixed effect. We included this as a binary response for any species that had offspring in at
least one replicate vial (we only found evidence of mating for three species: *D.sturtevanti,*
S.lativittata and *Z.tuberculatus).* Model (1) also includes a species-specific component
independent of the phylogeny $u_{s:ht}$ that allow us to estimate the proportion of variation
that is not explained by the host phylogeny v_s (Longdon et al., 2011). However, this was
removed from model (2) as model (1) failed to separate the phylogenetic and species -
specific effects. The main model therefore assumes a Brownian motion model of evolution
(Felsenstein, 1973). The random effects and the residuals are assumed to follow a
multivariate normal distribution with a zero mean and a covariance structure $V_p \otimes A$ for the
phylogenetic affects and $V_e \otimes I$ for the residuals, $V_s \otimes I$ for species-specific effects, ($\otimes$ here
represents the Kronecker product). **A** is the phylogenetic relatedness matrix, **I** is an identity
matrix and the **V** are 2×2 (co)variance matrices describing the (co)variances between viral
load of the two sexes. The phylogenetic covariance matrix, V_p, describes the phylogenetic
inter-specific variances in each trait and the inter-specific covariances between them, V_s, the
non-phylogenetic between-species variances. The residual covariance matrix, V_e, describes
the within-species variance that can be both due to real within-species effects and
measurement or experimental errors. The off-diagonal elements of V_e (the covariances) are
not estimable because each vial only contains one sex and therefore no vial has multiple
measurements, so were set to zero. The MCMC chain was run for 1,300 million iterations
with a burn- in of 30 million iterations and a thinning interval of 1 million. All the models
were run with different prior structures (as in [5]) in order to test results for sensitivity to
the use of priors, but note they all gave similar results.
266
The proportion of between species variance that can be explained by the phylogeny was
calculated from model (1) using the equation $\frac{V_p}{V_p + V_s}$, where V_p and V_s represent the

phylogenetic and species specific components of between-species variance respectively,
and is equivalent to phylogenetic heritability or Pagel's lambda [42, 43]. The repeatability of
susceptibility measurements was calculated from model (2) as $\frac{V_p}{V_p + V_e}$, where V_e is the
residual variance. Inter-species correlations in viral load between each method were
calculated from model (2) V_p matrix as $\frac{cov_{x,y}}{\sqrt{var_x + var_y}}$ and the slopes (β) of each relationship as
$\frac{cov_{x,y}}{var_x}$. Parameter estimates stated below are means of the posterior density, and 95%
credible intervals reported (CIs) were taken to be the 95% highest posterior density
intervals.

Tissue tropism
278
Viral load data across species and tissues was analysed using a linear mixed effects model
using the lmer function in the lme4 package in R [41, 44] with models compared using the
anova function and the resulting *P* values reported. Tissue type, species and their interaction
were included as fixed effects and experimental replicate as a random effect to account for
the individual pool that each set of tissues came from. With only seven species there is little
power to carry out models controlling for phylogeny which is why species was fitted as a
fixed effect.
286
287 **Data availability**
288
289 All data and scripts are available at dx.doi.org/10.6084/m9.figshare.21437223.
290
291 **Results**
292
Sex differences in viral load
To examine if the sexes respond the same way to viral infection we infected 31 species of
Drosophilidae with DCV and quantified the change in viral load at 2 days post infection using
qRT-PCR. In total we infected 6324 flies across 186 biological replicates (biological replicate
= change in viral load from day 0 to day 2 post-infection), with a mean of 17 flies per
replicate (range across species = 12-20).
299
300 The mean change in viral load across all species was similar between the sexes (females =
301 12.59, 95% CI = 1.16, 23.80; males = 12.93, 95% CI = -0.65, 26.25). We found strong positive
302 interspecific correlation between the viral load of females and males (correlation = 0.92,
303 95% CI = 0.78, 1.00; Figure 1). The estimate of the slope is close to 1 (β = 0.99, 95% CI= 0.58,
304 1.38) suggesting males and females respond similarly to infection.

305
306   **Figure 1. Correlation between viral load in males and females.** Each point represents a
307   species mean, error bars show standard errors and the trend line is estimated from a linear
308   model.
309
The full model including the species-specific random effect independent of the host
phylogeny ($u_{s:ht}$) allowed us to calculate the proportion of the variation between the
species that can be explained by the phylogeny ($\frac{V_p}{V_p + V_s}$), equivalent to phylogenetic
heritability or Pagel's lambda [42, 43]. The host phylogeny explains a large proportion of the
inter-specific variation for both males and females (females = 0.68, 95% CIs: 0.06, 0.99;
males = 0.66, 95% CIs: 0.04, 0.99) consistent with previous findings for males [3, 5, 9, 29].
However, we note these estimates have broad confidence intervals, due to the model
struggling to separate out the phylogenetic and non-phylogenetic components. The
repeatability of viral load across species was relatively high for both sexes (females = 0.63,
95% CIs = 0.41, 0.80; males = 0.52, 95% CIs = 0.31, 0.74). We found no effect of either body
size (-0.05, 95% CI's = -0.28, 0.19) or mating status (0.34, 95% CI's = -5.63, 6.27) on viral load.
321
Tissue Tropism
To look at the how the tissue tropism of DCV varied across host species, we infected seven
species of fly with DCV and dissected them into six tissue types. We found large effects of
species on viral load (χ^2= 320.65, d.f=6, *P*<0.001) with >18 million fold difference in viral load
between the least and most susceptible species. Tissues differed in their viral loads to a
lesser extent, with the maximum difference being seen in *D. pseudobscura* with an
approximately 550 fold difference in viral load between the least and most susceptible
tissues (χ^2= 15.264, d.f=4, *P*=0.009). There was no evidence of tissues showing different
patterns of susceptibility in different hosts i.e. no evidence for a tissue-by-species

interaction (χ^2=41.515, d.f=30, *P*=0.079).

332
333   **Figure 2: Tissue tropism data from DCV infected flies. Male** flies were dissected into the six
334   tissue types 2 days post-infection before undergoing RNA extraction and quantification of
335   viral load. On the right is the phylogeny of the host species. The bars of the individual panels
336   are organised following the order of the *D. melanogaster* ranked from the tissue with lowest
337   to highest viral load. Error bars show standard errors.
338
339   **Discussion**
340
Here, we examined whether heterogeneities in infection outcomes altered patterns of
susceptibility across host species. We first examined whether males and females responded
in consistent or different ways to infection with DCV. We found that viral susceptibility
between females and males of 31 host species showed a strong positive correlation with a
close to 1:1 relationship, suggesting that susceptibility across species is not sex specific. We
next examined whether heterogeneities in the tissues infected across host species altered
the outcome of infection. We found differences in viral load between tissues of seven host

species, but no evidence of tissues showing different patterns of susceptibility in different
host species.
350
A difference between the sexes in immune function and resistance has been in found in a
range of studies [15, 45]. However, this is not universal, and interactions between the host,
pathogen and environmental factors can alter the outcome of infections [46]. Previous
meta-analyses have found mixed results. For example, some studies of arthropods have
found little evidence for consistent sex differences in parasite prevalence or intensity [47]
(although these were largely from natural infections which may have inherently greater
sources of variation). Likewise, a phylogenetically controlled meta-analysis of 105 species
(including 30 insect species) of immune responses found no evidence for sex biases [48].
Other studies have reported a small male bias in parasitism rates for polygynous insects, but
a significant female bias for non-polygynous species, with the extent of sex bias parasitism
increasing with the degree of sexual size dimorphism [23]. However, other studies of insects
(with a smaller number of species) have reported sex differences in some immune traits [15,
23] and sex biased prevalence and impacts are well known for sexually transmitted
infections in insects [20, 21]. In mammals, males have been shown to often have greater
pathogen burdens, with parasitism rates positively correlated to male biased sexual size
dimorphism [15-17, 23].
367
In *Drosophila melanogaster*, some previous studies have reported males have higher DCV
viral loads than females [30]. However, others found no effect of sex on viral load, but did
find effects of sex on viral shedding, clearance, and transmission potential, with these traits
showing interactions with host genotype [31]. Sexual dimorphism in infection avoidance
behaviour has also been reported, when female flies previously exposed to DCV were found
to prefer a clean food source indicating a potentially important dimorphism in infection
avoidance [49]. Here, we used controlled experimental conditions, but in nature sex
differences in behaviour, or how the sexes interact with the environment may lead to
differences in pathogen load. A caveat is that the flies used here were of a fixed age and
largely virgins – future studies should explore if age and mating status can affect these
results.
379
The tissue tropism results here show that susceptibility in a given host is general across
tissue types – for example *D. sturtevanti* has a high viral load across all tissues whereas *D.
virilis* has relatively low viral loads in all tissues (Figure 2). Mortality to DCV infection has
previously been shown to show a strong positive correlation with viral load [9, 29]. The data
presented here show the susceptibility of a given species is general across all tissue types.
This does not exclude the possibility that pathology is due to high viral loads in a given
tissue, but does suggest that the mechanism restricting viral load is general across tissues.
This may be linked to the ability of the virus to bind to or enter hosts cells, utilise the hosts
cellular components for replication or to avoid or suppress the host immune response [50].
Further work should explore this further in a range of conditions including flies of both sexes
and of varying ages. Comparative studies of human viruses have identified the tissue tropism
of viruses to be a significant determinant of virulence; viruses that cause systemic infections
(across multiple organs) or that have neural or renal tropisms are most likely to cause severe

virulence [25]. It has been suggested that high levels of non-adaptive virulence can be the
result of pathogens infecting tissues that do not contribute to onward transmission [26].
Other studies have shown differences in host physiology can be important in determining
the virulence of a novel pathogen [10]. However, further understanding of how infection
results in pathology (i.e. in which tissue the disease tropism occurs [51]) and how virulence
is correlated with transmission potential in infections, is needed to explore this further.
399
In summary, our results show little evidence for sexual dimorphism in susceptibility to viral
infection across species. As such susceptibility in one sex is predictive of that in the other.
We find that susceptibility of a species is general across tissue types, suggesting virulence is
not due to species specific differences in viral tropism. The patterns of susceptibility
observed across species do not appear to be affected by heterogeneities in sex or tissue
tropism. Further work is needed to explore how sex differences can vary with factors such as
host age, mating status, the environment and pathogen type, and the underlying
mechanisms as to why species vary in their susceptibility.
408
409 **Acknowledgments**
410

411 Thanks to group members for discussions of these data. For the purpose of Open Access, the
412 author has applied a CCBY public copyright licence to any Author Accepted Manuscript
413 version arising from this submission. Many thanks for Alison Duncan, Greg Hurst and an
414 anonymous reviewer for useful comments that improved the manuscript.
415
416 **Funding**
417
418 B.L. and K.E.R are supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome
419 Trust and the Royal Society (109356/Z/15/Z).
420

421 **Conflict of interest disclosure**
422
423 The authors declare that they comply have no financial conflicts of interest in relation to the
424 content of the article
425

References
1.   Warren CJ, Meyerson NR, Dirasantha O, Feldman ER, Wilkerson GK, Sawyer SL.
Selective use of primate CD4 receptors by HIV-1. Plos Biology. 2019;17(6). doi:
10.1371/journal.pbio.3000304. PubMed PMID: WOS:000473675900026.
2.   Mollentze N, Keen D, Munkhbayar U, Biek R, Streicker DG. Variation in the ACE2
receptor has limited utility for SARS-CoV-2 host prediction. bioRxiv.
2022:2022.05.16.492068. doi: 10.1101/2022.05.16.492068.
3.   Imrie RM, Roberts KE, Longdon B. Between virus correlations in the outcome of
infection across host species: evidence of virus by host species interactions. Evolution
Letters. 2021;5(5):472-83. doi: 10.1101/2021.02.16.431403.
4.   Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. Host phylogeny
determines viral persistence and replication in novel hosts. PLoS Pathogens.
2011;7((9)):e1002260. doi: 10.1371/journal.ppat.1002260.

5.   Roberts KE, Hadfield JD, Sharma MD, Longdon B. Changes in temperature alter the
potential outcomes of virus host shifts. PLoS Pathog. 2018;14(10):e1007185. Epub
2018/10/20. doi: 10.1371/journal.ppat.1007185. PubMed PMID: 30339695; PubMed Central
PMCID: PMCPMC6209381.
6.   Faria NR, Suchard MA, Rambaut A, Streicker DG, Lemey P. Simultaneously
reconstructing viral cross-species transmission history and identifying the underlying
constraints. Philos Trans R Soc Lond B Biol Sci. 2013;368(1614):20120196. Epub 2013/02/06.
doi: 10.1098/rstb.2012.0196. PubMed PMID: 23382420; PubMed Central PMCID:
PMC3678322.
7.   Martel A, Blooi M, Adriaensen C, Van Rooij P, Beukema W, Fisher MC, et al. Wildlife
disease. Recent introduction of a chytrid fungus endangers Western Palearctic salamanders.
Science. 2014;346(6209):630-1. Epub 2014/11/02. doi: 10.1126/science.1258268. PubMed
PMID: 25359973; PubMed Central PMCID: PMCPMC5769814.
8.   Gilbert GS, Webb CO. Phylogenetic signal in plant pathogen-host range. Proceedings
of the National Academy of Sciences of the United States of America. 2007;104(12):4979-83.
doi: 10.1073/pnas.0607968104. PubMed PMID: WOS:000245256700040.
9.   Longdon B, Hadfield JD, Day JP, Smith SC, McGonigle JE, Cogni R, et al. The Causes
and Consequences of Changes in Virulence following Pathogen Host Shifts. PLoS Pathog.
2015;11(3):e1004728. Epub 2015/03/17. doi: 10.1371/journal.ppat.1004728. PubMed PMID:
25774803.
10.   Mollentze N, Streicker DG, Murcia PR, Hampson K, Biek R. Virulence mismatches in
index hosts shape the outcomes of cross-species transmission. Proc Natl Acad Sci U S A.
2020;117(46):28859-66. Epub 20201029. doi: 10.1073/pnas.2006778117. PubMed PMID:
33122433; PubMed Central PMCID: PMCPMC7682402.
11.   Farrell MJ, Davies TJ. Disease mortality in domesticated animals is predicted by host
evolutionary relationships. Proceedings of the National Academy of Sciences of the United
States of America. 2019;116(16):7911-5. doi: 10.1073/pnas.1817323116. PubMed PMID:
WOS:000464767500050.
12.   Guth S, Visher E, Boots M, Brook CE. Host phylogenetic distance drives trends in
virus virulence and transmissibility across the animal-human interface. Philos Trans R Soc
Lond B Biol Sci. 2019;374(1782):20190296. Epub 20190812. doi: 10.1098/rstb.2019.0296.
PubMed PMID: 31401961; PubMed Central PMCID: PMCPMC6711300.
13.   Belmonte RL, Corbally MK, Duneau DF, Regan JC. Sexual Dimorphisms in Innate
Immunity and Responses to Infection in Drosophila melanogaster. Front Immunol.
2019;10:3075. Epub 20200131. doi: 10.3389/fimmu.2019.03075. PubMed PMID: 32076419;
PubMed Central PMCID: PMCPMC7006818.
14.   Kellermann V, Overgaard J, Sgro CM, Hoffmann AA. Phylogenetic and environmental
patterns of sex differentiation in physiological traits across Drosophila species. J Evol Biol.
2022. Epub 20221005. doi: 10.1111/jeb.14104. PubMed PMID: 36196885.
15.   Nunn CL, Lindenfors P, Pursall ER, Rolff J. On sexual dimorphism in immune function.
Philos Trans R Soc Lond B Biol Sci. 2009;364(1513):61-9. doi: 10.1098/rstb.2008.0148.
PubMed PMID: 18926977; PubMed Central PMCID: PMCPMC2666693.
16.   Regan JC, Partridge L. Gender and longevity: why do men die earlier than women?
Comparative and experimental evidence. Best Pract Res Clin Endocrinol Metab.
2013;27(4):467-79. Epub 20130622. doi: 10.1016/j.beem.2013.05.016. PubMed PMID:
24054925.
17.   Moore SL, Wilson K. Parasites as a viability cost of sexual selection in natural
populations of mammals. Science. 2002;297(5589):2015-8. doi: DOI
10.1126/science.1074196. PubMed PMID: WOS:000178078500036.
18.   Abate BB, Kassie AM, Kassaw MW, Aragie TG, Masresha SA. Sex difference in
coronavirus disease (COVID-19): a systematic review and meta-analysis. BMJ Open.

2020;10(10):e040129. Epub 20201006. doi: 10.1136/bmjopen-2020-040129. PubMed PMID:
33028563; PubMed Central PMCID: PMCPMC7539579.
19.   vom Steeg LG, Klein SL. SeXX Matters in Infectious Disease Pathogenesis. PLoS
Pathog. 2016;12(2):e1005374. Epub 20160218. doi: 10.1371/journal.ppat.1005374. PubMed
PMID: 26891052; PubMed Central PMCID: PMCPMC4759457.
20.   Ashby B, Jones JE, Knell RJ, Hurst GDD. Sexually transmitted infections in natural
populations: what have we learnt from beetles and beyond? In: Fenton A, Tompkins D,
Wilson K, editors. Wildlife Disease Ecology: Linking Theory to Data and Application.
Ecological Reviews. Cambridge: Cambridge University Press; 2019. p. 187-222.
21.   Knell RJ, Webberley KM. Sexually transmitted diseases of insects: distribution,
evolution, ecology and host behaviour. Biol Rev Camb Philos Soc. 2004;79(3):557-81. doi:
10.1017/s1464793103006365. PubMed PMID: 15366763.
22.   Lockhart AB, Thrall PH, Antonovics J. Sexually transmitted diseases in animals:
ecological and evolutionary implications. Biol Rev Camb Philos Soc. 1996;71(3):415-71. doi:
10.1111/j.1469-185x.1996.tb01281.x. PubMed PMID: 8761160.
23.   Wilson K, Cotter SC. Chapter Three - Host–Parasite Interactions and the Evolution of
Immune Defense. In: Brockmann HJ, Roper TJ, Naguib M, Mitani JC, Simmons LW, Barrett L,
editors. Advances in the Study of Behavior. 45: Academic Press; 2013. p. 81-174.
24.   Rolff J. Bateman's principle and immunity. Proc Biol Sci. 2002;269(1493):867-72. doi:
10.1098/rspb.2002.1959. PubMed PMID: 11958720; PubMed Central PMCID:
PMCPMC1690964.
25.   Brierley L, Pedersen AB, Woolhouse MEJ. Tissue tropism and transmission ecology
predict virulence of human RNA viruses. PLoS Biol. 2019;17(11):e3000206. Epub 20191126.
doi: 10.1371/journal.pbio.3000206. PubMed PMID: 31770368; PubMed Central PMCID:
PMCPMC6879112.
26.   Levin BR, Bull JJ. Short-sighted evolution and the virulence of pathogenic
microorganisms. Trends Microbiol. 1994;2(3):76-81. Epub 1994/03/01. doi: 10.1016/0966-
842x(94)90538-x. PubMed PMID: 8156275.
27.   Kitchen A, Shackelton LA, Holmes EC. Family level phylogenies reveal modes of
macroevolution in RNA viruses. Proc Natl Acad Sci U S A. 2011;108(1):238-43. Epub
2010/12/22. doi: 1011090108 [pii]
10.1073/pnas.1011090108. PubMed PMID: 21173251; PubMed Central PMCID:
PMC3017157.
28.   Taber SW, Pease CM. Paramyxovirus Phylogeny - Tissue Tropism Evolves Slower
Than Host Specificity. Evolution. 1990;44(2):435-8. doi: Doi 10.2307/2409419. PubMed
PMID: WOS:A1990CZ91500017.
29.   Roberts KE, Longdon B. Viral susceptibility across host species is largely independent
of dietary protein to carbohydrate ratios. Journal of Evolutionary Biology.
2021;2020.10.21.348987. doi: 10.1111/jeb.13773.
30.   Gupta V, Stewart CO, Rund SSC, Monteith K, Vale PF. Costs and benefits of sublethal
Drosophila C virus infection. J Evol Biol. 2017;30(7):1325-35. Epub 20170524. doi:
10.1111/jeb.13096. PubMed PMID: 28425174.
31.   Siva-Jothy JA, Vale PF. Dissecting genetic and sex-specific sources of host
heterogeneity in pathogen shedding and spread. PLoS Pathog. 2021;17(1):e1009196. Epub
20210119. doi: 10.1371/journal.ppat.1009196. PubMed PMID: 33465160; PubMed Central
PMCID: PMCPMC7846003.
32.   Ferreira AG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. The Toll-Dorsal
Pathway Is Required for Resistance to Viral Oral Infection in Drosophila. Plos Pathogens.
2014;10(12). doi: 10.1371/Journal.Ppat.1004507. PubMed PMID: ISI:000346702400008.

33.   Chtarbanova S, Lamiable O, Lee KZ, Galiana D, Troxler L, Meignin C, et al. Drosophila
C virus systemic infection leads to intestinal obstruction. J Virol. 2014;88(24):14057-69. Epub
2014/09/26. doi: 10.1128/JVI.02320-14. PubMed PMID: 25253354.
34.   Longdon B, Day JP, Alves JM, Smith SCL, Houslay TM, McGonigle JE, et al. Host shifts
result in parallel genetic changes when viruses evolve in closely related species. PLoS
Pathog. 2018;14(4):e1006951. Epub 2018/04/13. doi: 10.1371/journal.ppat.1006951.
PubMed PMID: 29649296; PubMed Central PMCID: PMCPMC5897010.
35.   Jousset FX, Plus N, Croizier G, Thomas M. [Existence in Drosophila of 2 groups of
picornavirus with different biological and serological properties]. C R Acad Sci Hebd Seances
Acad Sci D. 1972;275(25):3043-6. Epub 1972/12/18. PubMed PMID: 4631976.
36.   Longdon B, Cao C, Martinez J, Jiggins FM. Previous Exposure to an RNA Virus Does
Not Protect against Subsequent Infection in Drosophila melanogaster. Plos One.
2013;8(9):e73833. doi: 10.1371/journal.pone.0073833.
37.   CSIRO. Anatomical Atlas of the male Drosophila melanogaster.
38.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious
Basic: an integrated and extendable desktop software platform for the organization and
analysis of sequence data. Bioinformatics. 2012;28(12):1647-9. Epub 20120427. doi:
10.1093/bioinformatics/bts199. PubMed PMID: 22543367; PubMed Central PMCID:
PMCPMC3371832.
39.   Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti
and the BEAST 1.7. Mol Biol Evol. 2012;29(8):1969-73. Epub 2012/03/01. doi:
10.1093/molbev/mss075. PubMed PMID: 22367748; PubMed Central PMCID: PMC3408070.
40.   Hadfield JD. MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. Journal of Statistical Software. 2010;33(2):1-22. PubMed PMID:
WOS:000275203300001.
41.   Team RDC. R: a language and environment for statistical computing. V 2.4. 2006.
42.   Housworth EA, Martins EP, Lynch M. The phylogenetic mixed model. American
Naturalist. 2004;163(1):84-96. doi: 10.1086/380570 PubMed PMID:
WOS:000188954900008.
43.   Pagel M. Inferring the historical patterns of biological evolution. Nature.
1999;401(6756):877-84. Epub 1999/11/30. doi: 10.1038/44766. PubMed PMID: 10553904.
44.   Bates D, Machler M, Bolker BM, Walker SC. Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software. 2015;67(1):1-48. doi: DOI 10.18637/jss.v067.i01.
PubMed PMID: WOS:000365981400001.
45.   Poulin R. Sexual inequalities in helminth infections: A cost of being a male? American
Naturalist. 1996;147(2):287-95. doi: Doi 10.1086/285851. PubMed PMID:
WOS:A1996TR36900007.
46.   Lazzaro BP, Little TJ. Immunity in a variable world. Philosophical Transactions of the
Royal Society B-Biological Sciences. 2009;364(1513):15-26. doi: Doi 10.1098/Rstb.2008.0141.
PubMed PMID: ISI:000261150600002.
47.   Sheridan LAD, Poulin R, Ward DF, Zuk M. Sex Differences in Parasitic Infections
among Arthropod Hosts: Is There a Male Bias? Oikos. 2000;88(2):327-34.
48.   Kelly CD, Stoehr AM, Nunn C, Smyth KN, Prokop ZM. Sexual dimorphism in immunity
across animals: a meta-analysis. Ecol Lett. 2018;21(12):1885-94. Epub 20181004. doi:
10.1111/ele.13164. PubMed PMID: 30288910.
49.   Vale PF, Jardine MD. Infection avoidance behavior: Viral exposure reduces the
motivation to forage in female Drosophila melanogaster. Fly (Austin). 2017;11(1):3-9. Epub
20160630. doi: 10.1080/19336934.2016.1207029. PubMed PMID: 27362557; PubMed
Central PMCID: PMCPMC5354229.
50.   Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. The Evolution and
Genetics of Virus Host Shifts. PLoS Pathog. 2014;10(11):e1004395. Epub 2014/11/07. doi:

10.1371/journal.ppat.1004395. PubMed PMID: 25375777; PubMed Central PMCID:
PMC4223060.
51.   McCall LI. Quo vadis? Central Rules of Pathogen and Disease Tropism. Front Cell
Infect Microbiol. 2021;11:640987. Epub 20210225. doi: 10.3389/fcimb.2021.640987.
PubMed PMID: 33718287; PubMed Central PMCID: PMCPMC7947345.
595
596
597 **Supplementary information**
598
599 **Table S1**: Full list of species used in the sex difference experiment and their rearing food for
600 stock populations. All cornmeal and proprionic medium have dried yeast sprinkled onto the
601 surface of the food, other food types do not unless stated below. The recipes for the food
602 types are described here https://doi.org/10.6084/m9.figshare.21590724.v1
603

Species	Food
D.affinis	Malt
D.americana	Malt
D.ananassae	Cornmeal
D.arizonae	Banana
D.buzzatii	Malt
D.erecta	Malt + yeast
D.flavomontana	Malt + yeast
D.hydei	Cornmeal
D.immigrans	Malt + yeast
D.lacicola	Malt
D.littoralis	Banana
D.mauritiana	Proprionic
D.melanogaster	Cornmeal
D.montana	Malt + yeast
D.novamexicana	Banana
D.obscura	Proprionic
D.persimilis	Malt
D.prosaltans	Proprionic
D.pseudoobscura	Malt
D.putrida	Proprionic

D.santomea	Cornmeal
D.sturtevanti	Cornmeal
D.takahashii	Cornmeal
D.teissieri	Cornmeal
D.virilis	Proprionic
H.duncani	Proprionic
S. lativittata	Banana
S.lebanonensis	Proprionic
Z. inermis	Banana
Z. taronus	Banana
Z. tuberculatus	Banana

604
605