Mitotic lineage adds predictive information beyond cell type in the *C. elegans* connectome

Jordan K. Matelsky[a,b], **Brock Wester**[a], **Konrad P. Körding**[b,c,d]

*For correspondence:
matelsky@upenn.edu (JKM)

[a]Johns Hopkins Applied Physics Laboratory, Laurel, Maryland, United States;
[b]Department of Bioengineering, University of Pennsylvania, Philadelphia, Pennsylvania, United States; [c]Department of Neuroscience, University of Pennsylvania, Philadelphia, Pennsylvania, United States; [d]CIFAR, LMB program

Abstract   During nervous system development, repeated cell divisions of the zygote give rise to a "family tree" of neurons related by mitotic lineage. The developmental process also gives rise to neuronal connections, and neurons phenotypically converge to different cell types. Neural connections are steered by cell type, but they may be driven in part by lineage: We do not know if lineage matters for the developmental neurogenesis process. We thus asked if mitotic lineage predicts neural connections beyond cell type alone. Using three *C. elegans* datasets, we fit models for edge prediction tasks: predicting synaptic targets, and predicting synaptic sources. Adding the mitotic lineage improved these predictions. Our results suggest that the family tree matters for the connections made by neurons, and that developmental lineage is a variable that should be considered more deeply in models of connectome assembly.

Introduction

Brains are built developmentally: cells divide to make new cells, and repeated mitotic branching yields a lineage tree. That tree captures shared developmental ancestry. We ask if access to lineage information improves models of neuronal connectivity beyond what is possible with cell type alone (**Fig. 1**). If lineage matters, then neuroscience should consider this developmental variable much more deeply when modeling how nervous systems are assembled.

Connectome assembly is generally framed as a developmental process in which neurogenesis, neurite outgrowth, and local partner-recognition rules progressively constrain which cells can meet and synapse (*Kerstjens et al., 2022*). In *C. elegans*, this question is especially tractable because neurite neighborhoods remain broadly stable across development even as synaptic connectivity matures (*Moyle et al., 2021*; *Witvliet et al., 2021*). Cell type is a central organizing concept in neuroscience, but it does not fully specify how connectomes are assembled (*Thompson et al., 2014*; *Erö et al., 2018*; *Zeng and Sanes, 2017*; *Barabási and Czégel, 2021*). Physical geometry also matters, yet geometric proximity is likewise an incomplete predictor of connectivity (*Varshney et al., 2011*). The *C. elegans* system enables a direct test because complete lineage resources and synapse-resolved connectomes are available across development and in a mature reference (*Bhatla, 2011*; *Witvliet et al., 2021*; *Cook et al., 2019*). Figure 2 shows this alignment directly for the `Witvliet_8` dataset: the lineage tree, cell-type labels, and connectome adjacency can be coregistered.

Here we focus on the late developmental and mature regime, analyzing `Witvliet_7`, `Witvliet_8`,

Figure 1. Conceptual framing for lineage-connectome integration. A, development yields a mitotic lineage tree. **B**, connectomics yields a synapse-resolved directed graph. **C**, cell-type and lineage labels can be aligned to graph nodes. **D**, pairwise and motif-level analyses relate lineage structure to connectivity structure. **E**, lineage-shuffle null logic asks whether the lineage-to-connectome mapping carries signal beyond the fixed tree and graph topologies. Cell and animal morphology graphics are based on OpenWorm and BossDB resources (*Szigeti et al., 2014*; *Hider et al., 2019*)

Table 1. Held-out directed-edge predictive gains from adding lineage. The main effect is the increase in held-out test R^2 after adding lineage to the cell-type baseline. Raw model R^2 values are shown for context.

Task	Cell Type	Cell Type + lineage	ΔR^2	Relative lift
Target prediction	0.0199	0.0272	0.0073	+37%
Source prediction	0.0177	0.0231	0.0054	+30%

(*Witvliet et al., 2021*) and the Cook/White N2U mosaic (N2U, *Cook et al.* (*2019*)) with held-out linear

models for two directed edge-prediction tasks: predicting a source neuron's targets and predicting

a target neuron's sources (**Fig. 4**). Each ordered row is a candidate directed connection $u \to v$. In

target prediction, we hold the source neuron u fixed and ask which candidate targets v it connects

to; in source prediction, we hold the target neuron v fixed and ask which candidate sources u

connect into it. Here, held-out prediction is used as an assay of information content rather than as

an attempt to maximize predictive accuracy; the question is whether lineage adds out-of-sample

signal beyond cell type.

Results

Cell type is already a strong predictor of directed connectivity

Before asking whether lineage helps, we first establish what a fair non-lineage baseline already

captures. After all, cell type is known to be a strong predictor of connectivity. Using cell-type de-

scriptors alone, the directed models already achieved mean held-out test R^2 values of 0.0199 for

target prediction and 0.0177 for source prediction (Fig. 3). By contrast, a distance-only model sat

essentially on the held-out $R^2 = 0$ line (Appendix). The interpretation of everything that follows

therefore hinges on an incremental question: not whether lineage can predict connectivity from

scratch, but whether it improves prediction after cell type has already explained a substantial part

of the linear signal.

Figure 2. C. elegans lineage aligned to the connectome. Leaves of the two (identical) dendrograms along the perimeter of the plot correspond to neurons shared between the `Witvliet_8` connectome and the lineage tree, colored by composite *cell type* label. The same lineage-derived leaf order is used for the rows and columns of the directed chemical-synapse adjacency matrix, with local branch order chosen to keep nearby connectivity profiles together.

Figure 3. Constituent cell-type baselines. A, target-prediction model summaries using cell type descriptors and lineage. **B**, source-prediction model summaries. Bars show the mean held-out test R^2 across the adult datasets, and error bars show the standard deviation across datasets. This figure establishes the baseline that lineage must beat: source and target type descriptors already explain much of the accessible linear signal, yet adding lineage still improves performance.

Lineage adds held-out signal beyond cell type in both directed tasks

Adding lineage improved held-out performance in both directed edge-prediction tasks (Table 1; Fig. 4). For target prediction (**Fig. 4A**), the mean test R^2 rose from 0.0199 to 0.0272, a gain of 0.0073 (+37%). For source prediction (**Fig. 4B**), it rose from 0.0177 to 0.0231, a gain of 0.0054 (+30%). Crucially, these gains were positive in all three adult datasets.

The lineage-only model performed substantially worse than the combined model in both tasks (0.00860 for target prediction and 0.00677 for source prediction). That pattern is important for interpretation. The result is *not* that lineage is a strong standalone substitute for cell type. (It is not!) Rather, lineage contributes nonredundant information that the cell-type baseline does not already absorb.

The gain is extinguished with a shuffled-null lineage-to-neuron mapping

If lineage is truly contributing task-relevant information, then the gain should depend on the actual mapping between connectome neurons and lineage leaves, not merely on the fixed shapes of the lineage tree and the connectome. We tested that prediction by permuting the lineage-to-neuron mapping within each dataset (**Fig. 1E**), recomputing lineage features, and refitting the same held-out models (**Fig. 4**). For target prediction, the observed mean gain across datasets was 0.00731 test-R^2 units, whereas the 95% leaf-shuffle null interval for the across-dataset mean ranged from −0.00382 to 0.00051. Source prediction also cleared its null, with an observed mean gain of 0.00538 against a 95% null interval from −0.00032 to 0.00339. This null control sharpens the central claim: the advantage of lineage is not a generic consequence of having some hierarchical feature block, but depends on the real alignment between developmental ancestry and connectome identity.

Lineage features work best as an ensemble

Having established that lineage adds signal, we next asked which part of the lineage description carries that gain. To do so, we replaced the full lineage block with one lineage term at a time while keeping the same cell-type baseline. Across both directed tasks, the full three-term lineage block performed best overall (Fig. 5).

In target prediction, the full block improved held-out performance by 0.00710 test-R^2 units over cell type alone, slightly exceeding most recent common ancestor (MRCA) depth alone (0.00621) and raw lineage distance alone (0.00615), while lineage-depth difference alone contributed much less. In source prediction, the same full block improved held-out performance by 0.00318, with raw lineage distance alone (0.00230) slightly ahead of MRCA depth alone (0.00197). The interpretation is that no single scalar lineage summary fully captures the effect. Most of the signal is carried

Figure 4. Lineage-leaf-shuffle null. A. Target prediction. Target prediction asks whether a fixed source neuron u targets a candidate neuron v. In schematics, shaded nodes are held fixed within a split group, and dashed-outline nodes denote the candidate partner being predicted. **B. Source prediction.** Source prediction asks the complementary question of whether a candidate neuron u is an input to a fixed target neuron v. For each dataset, the filled green dot marks the observed held-out gain from adding lineage to the fair baseline, the open circle marks the mean gain after shuffling the mapping between connectome neurons and lineage leaves, and the gray interval shows the central 95% of the shuffled null distribution. The bottom row summarizes the across-dataset mean. Both directed edge-prediction tasks clear the shuffle null.

---

90 by shared developmental ancestry and lineage separation, while absolute depth difference con-
91 tributes comparatively little.

Discussion

In late developmental and adult *C. elegans*, mitotic lineage information adds reproducible improve-
ment in predictive models of connectivity beyond cell type. That effect is consistent across datasets
and can be eliminated with a lineage-leaf-shuffle null, which argues that the real mapping between
lineage and connectome identity carries signal. This is exactly the pattern we expect if developmen-
tal ancestry captures a real part of the residual structure left over after cell type has already done
much of the predictive work.

The lineage decomposition refines that interpretation. MRCA depth and raw lineage distance
carry most of the signal, whereas the absolute difference in lineage depth contributes much less.
This suggests that the relevant information is tied more to shared developmental history and lin-
eage separation than to one neuron simply being developmentally "earlier" or "later" than another.

We should not infer from this work that lineage causally determines the connectome. Stronger
claims would require richer model classes, additional species, or experimental perturbations that
directly test developmental causality. Instead, our results are consistent with the idea that lineage
is one of the variables that reduces the residual wiring entropy that must be addressed by neuro-
genesis.

Genomic-bottleneck arguments suggest that nervous systems cannot specify every synapse
independently and instead must compress wiring instructions into reusable developmental pro-
grams (*Koulakov et al., 2021*). In that framing, cell type removes part of the uncertainty in connec-
tivity, and our results suggest that developmental ancestry removes a bit more.

Because cell type and lineage are correlated, confound control remains essential (*Pearl, 2009*;
Angrist and Pischke, 2014). Here, lineage still provided an improvement after conditioning on type.
That illustrates that lineage captures information not already absorbed by cell type, and it argues
against the lineage effect being merely a proxy for cell-type misclassification noise.

More broadly, lineage has been linked to neuronal and physiological specialization beyond con-
nectomics (*Li et al., 2012*; *Cadwell et al., 2020*; *He et al., 2020*), and emerging lineage-tracing tech-

Figure 5. Lineage features work best as an ensemble. A, target prediction. **B**, source prediction. For each directed edge-prediction task, bars show the mean held-out improvement in test R^2 over the same cell-type baseline across the three adult datasets, and error bars show the standard deviation across datasets, when adding either raw lineage distance alone, MRCA depth alone, lineage depth difference alone, or the full three-term lineage block. The full block performs best overall, indicating that the lineage effect is distributed across multiple correlated summaries rather than being captured cleanly by any single term.

---

nologies should make cross-system tests increasingly feasible (*Masuyama et al., 2019*). We there-
fore view lineage-aware models as part of a larger developmental program for understanding how
wiring constraints are inherited over cell divisions (*Kerstjens et al., 2022*).

As additional matched reconstructions across individuals and stages become available, it should
become possible to estimate lineage effects more precisely and separate reproducible structure
from animal-to-animal variation (*Cook et al., 2019*; *Witvliet et al., 2021*). We therefore view this
analysis as a community resource that should grow more informative as the number of public *C.*
elegans connectomes increases.

Methods
126

Data, normalization, and folding
127

Connectomes were accessed through MotifStudio/BossDB endpoints using host IDs `Witvliet_7`,
`Witvliet_8`, and `N2U` (*Hider et al., 2019*; *Matelsky et al., 2021*, *2025*). These analyses were restricted
to late developmental and mature connectomes. Species/dataset metadata were controlled by
a YAML registry for reproducibility. Lineage was loaded from WormWeb-derived JSON (*Bhatla,*
2011), and soma locations from OpenWorm-derived coordinates with WormBase-supported iden-
tity annotations, with postprocessing performed in Blender (*Szigeti et al., 2014*; *Harris et al., 2010*;
Blender Online Community, 2026). Graph and table analyses were implemented in Python with
NetworkX (*Hagberg et al., 2008*).

Neuron identities were aligned by name with normalization rules from the registry. Bilateral
neurons were folded with right-side preference; edge weights were aggregated by summation
(with mean as an available alternative but unused mode). This folding operation is adopted from
prior work in *C. elegans* connectomics (*Witvliet et al., 2021*).

Prediction tasks
140

The first task was ordered edge-target prediction, $\mathbb{1}[u \to v]$. Each row corresponds to a candidate
source-target pair among neurons shared between the lineage and connectome graphs.

The second task used the same directed edge rows but asked the complementary held-out
question of source prediction: can a target neuron's incoming partners be predicted from candi-
date sources? The outcome variable remained $\mathbb{1}[u \to v]$ but train-test splits were formed within
target-neuron groups rather than source-neuron groups.

Feature blocks

For target prediction and source prediction, the paper's main baseline block contained decomposed `typ` indicators for the source and target. A single physical-distance term was evaluated separately as an appendix negative control rather than included in the paper's main baseline, as it did not substantially improve performance and unnecessarily complicated the models.

The lineage block contained three terms: undirected lineage distance, depth of the most recent common ancestor (MRCA), and the absolute difference between the depths of the two lineage nodes. Additional diagnostic models replaced the full lineage block with one lineage term at a time to determine which component carried the predictive signal.

Model fitting and evaluation

All models were ordinary least squares fits evaluated on held-out data. For target prediction, train-test splits were formed within source-neuron groups so that each source retained both train and test targets. For source prediction, splits were formed within target-neuron groups. The default held-out fraction was 0.25.

As a negative control, we repeated the combined-model fit after 20 random shuffles of the mapping from overlapping connectome neurons to lineage leaves within each dataset. The lineage tree, connectome, baseline features, and train-test split were held fixed; only the lineage-to-neuron assignment used to compute lineage features changed.

We report held-out test R^2 for each model, but in the Results we emphasize the held-out ΔR^2 gained by adding lineage to the cell-type baseline, because that quantity most directly answers the biological question of interest. We then summarize constituent cell-type models and lineage-component models separately.

Reproducibility & Rigor

All analyses produce deterministic figures, summary tables, and run manifests. Supplemental materials, including reproducibility details and generated assets, are provided at https://github.com/KordingLab/LineageConnectomics.

Figure 6. Simple physical-distance terms are weak in this linear setup. A, target prediction. **B**, source prediction. Bars show the mean held-out test R^2 across the three adult datasets for distance alone, cell type alone, and cell type plus a single physical-distance term; error bars show the standard deviation across datasets. The dashed vertical line marks $R^2 = 0$, the constant-predictor reference for this metric. Distance-only models sit at or extremely near that line, while adding distance to cell type changes performance only slightly.

Appendix

Simple physical-distance terms sit near the held-out $R^2 = 0$ line

As a simple negative control, we evaluated models that used only the Euclidean distance between source and target somata, as well as models that added this distance term to the cell-type baseline. Across the adult-only directed tasks, distance-only models achieved mean held-out test R^2 values of -0.000084 for target prediction and 0.000082 for source prediction, i.e. essentially on the $R^2 = 0$ line for this metric. Adding distance to cell type increased mean held-out test R^2 only slightly, from 0.0199 to 0.0217 for target prediction and from 0.0177 to 0.0195 for source prediction (Fig. 6). These checks do not imply that geometry is biologically irrelevant; only that this very simple linear distance term is not a strong standalone predictor here.

Supplemental Materials

Supplemental figures, tables, and reproducibility materials are provided at https://github.com/ KordingLab/LineageConnectomics. The README documents how to reproduce manuscript outputs from the Snakemake workflow or individual analysis entrypoints.

Acknowledgements

The authors thank all contributors to *C. elegans* connectomics, and Nikhil Bhatla for curation of the *C. elegans* lineage tree resources.

Bibliography

References

Angrist JD, Pischke JS. Mastering metrics: The path from cause to effect. Princeton university press; 2014.

Barabási DL, Czégel D. Constructing graphs from genetic encodings. Scientific Reports. 2021; 11(1):1–13.

Bhatla N, C. elegans Interactive Cell Lineage; 2011. http://wormweb.org/celllineage.

Blender Online Community, Blender. Blender Foundation; 2026. http://www.blender.org, blender Institute, Amsterdam.

Cadwell CR, Scala F, Fahey PG, Kobak D, Mulherkar S, Sinz FH, Papadopoulos S, Tan ZH, Johnsson P, Hartmanis L, et al. Cell type composition and circuit organization of clonally related excitatory neurons in the juvenile mouse neocortex. Elife. 2020; 9:e52951.

200 **Cook SJ**, Jarrell TA, Brittin CA, Wang Y, Bloniarz AE, Yakovlev MA, Nguyen KC, Tang LTH, Bayer EA, Duerr JS, et al.
201 Whole-animal connectomes of both Caenorhabditis elegans sexes. Nature. 2019; 571(7763):63–71.

202 **Erö C**, Gewaltig MO, Keller D, Markram H. A cell atlas for the mouse brain. Frontiers in neuroinformatics. 2018;
203 12:84.

204 **Hagberg A**, Swart P, S Chult D. Exploring network structure, dynamics, and function using NetworkX. Los
205 Alamos National Lab.(LANL), Los Alamos, NM (United States); 2008.

206 **Harris TW**, Antoshechkin I, Bieri T, Blasiar D, Chan J, Chen WJ, De La Cruz N, Davis P, Duesbury M, Fang
207 R, et al. WormBase: a comprehensive resource for nematode research. Nucleic acids research. 2010;
208 38(suppl_1):D463–D467.

209 **He S**, Tian Y, Feng S, Wu Y, Shen X, Chen K, He Y, Sun Q, Li X, Xu J, et al. In vivo single-cell lineage tracing in
210 zebrafish using high-resolution infrared laser-mediated gene induction microscopy. Elife. 2020; 9:e52024.

211 **Hider R**, Kleissas DM, Pryor D, Gion T, Rodriguez L, Matelsky J, Gray-Roncal W, Wester B. The block object
212 storage service (bossDB): A cloud-native approach for petascale neuroscience discovery. bioRxiv. 2019; p.
213 217745.

214 **Kerstjens S**, Michel G, Douglas RJ. Constructive Connectomics: how neuronal axons get from here to there
215 using gene-expression maps derived from their family trees. bioRxiv. 2022 Feb; http://dx.doi.org/10.1101/
216 2022.02.26.482112, doi: 10.1101/2022.02.26.482112.

217 **Koulakov A**, Shuvaev S, Zador A. Encoding innate ability through a genomic bottleneck. bioRxiv. 2021 Mar;
218 http://dx.doi.org/10.1101/2021.03.16.435261, doi: 10.1101/2021.03.16.435261.

219 **Li Y**, Lu H, Cheng Pl, Ge S, Xu H, Shi SH, Dan Y. Clonally related visual cortical neurons show similar stimulus
220 feature selectivity. Nature. 2012; 486(7401):118–121.

221 **Masuyama N**, Mori H, Yachie N. DNA barcodes evolve for high-resolution cell lineage tracing. Current Opinion
222 in Chemical Biology. 2019; 52:63–71.

223 **Matelsky JK**, Martinez H, Xenes D, Robinette M, Panigrahi A, Wester B. Analyzing Large Connectome Graphs
224 With BossDB Network Tools. Current Protocols. 2025; 5(12):e70273.

225 **Matelsky JK**, Reilly EP, Johnson EC, Stiso J, Bassett DS, Wester BA, Gray-Roncal W. DotMotif: an open-source
226 tool for connectome subgraph isomorphism search and graph queries. Scientific Reports. 2021 Jun; 11(1).
227 http://dx.doi.org/10.1038/s41598-021-91025-5, doi: 10.1038/s41598-021-91025-5.

228 **Moyle MW**, Barnes KM, Kuchroo M, Gonopolskiy A, Duncan LH, Sengupta T, Shao L, Guo M, Santella A, Chris-
229 tensen R, et al. Structural and developmental principles of neuropil assembly in C. elegans. Nature. 2021;
230 591(7848):99–104.

231 **Pearl J**. Causality. Cambridge university press; 2009.

232 **Szigeti B**, Gleeson P, Vella M, Khayrulin S, Palyanov A, Hokanson J, Currie M, Cantarelli M, Idili G, Larson S.
233 OpenWorm: an open-science approach to modeling Caenorhabditis elegans. Frontiers in computational
234 neuroscience. 2014; 8:137.

235 **Thompson CL**, Ng L, Menon V, Martinez S, Lee CK, Glattfelder K, Sunkin SM, Henry A, Lau C, Dang C, et al.
236 A high-resolution spatiotemporal atlas of gene expression of the developing mouse brain. Neuron. 2014;
237 83(2):309–323.

238 **Varshney LR**, Chen BL, Paniagua E, Hall DH, Chklovskii DB. Structural properties of the Caenorhabditis elegans
239 neuronal network. PLoS computational biology. 2011; 7(2):e1001066.

240 **Witvliet D**, Mulcahy B, Mitchell JK, Meirovitch Y, Berger DR, Wu Y, Liu Y, Koh WX, Parvathala R, Holmyard D, et al.
241 Connectomes across development reveal principles of brain maturation. BioRxiv. 2021; p. 2020–04.

242 **Zeng H**, Sanes JR. Neuronal cell-type classification: challenges, opportunities and the path forward. Nature
243 Reviews Neuroscience. 2017; 18(9):530–546.