FUS regulates RAN translation through modulating the G-quadruplex structure of

GGGGCC repeat RNA in *C9orf72*-linked ALS/FTD

Yuzo Fujino[1,2], Morio Ueyama[1,3,4], Taro Ishiguro[4,5], Daisaku Ozawa[1,3], Hayato Ito[6],

Toshihiko Sugiki[7], Asako Murata[8], Akira Ishiguro[9], Tania F. Gendron[10], Kohji Mori[11],

Eiichi Tokuda[12], Tomoya Taminato[1,3], Takuya Konno[13], Akihide Koyama[13], Yuya

Kawabe[11], Toshihide Takeuchi[3,14], Yoshiaki Furukawa[12], Toshimichi Fujiwara[7], Manabu

Ikeda[11], Toshiki Mizuno[2], Hideki Mochizuki[15], Hidehiro Mizusawa[5], Keiji Wada[4],

Kinya Ishikawa[5], Osamu Onodera[13], Kazuhiko Nakatani[8], Leonard Petrucelli[10], Hideki

Taguchi[6,16], and Yoshitaka Nagai[1,3,4,15]

[1] Department of Neurology, Kindai University Faculty of Medicine, Osaka-Sayama,

Osaka 589-8511, Japan

[2] Department of Neurology, Kyoto Prefectural University of Medicine, Kyoto 602-0841,

Japan

[3] Department of Neurotherapeutics, Osaka University Graduate School of Medicine,

Suita, Osaka 565-0871, Japan

[4] Department of Degenerative Neurological Diseases, National Institute of

Neuroscience, National Center of Neurology and Psychiatry, Kodaira, Tokyo 187-8502,

Japan

[5] Department of Neurology and Neurological Science, Tokyo Medical and Dental

University, Bunkyo-ku, Tokyo 113-8519, Japan

[6] School of Life Science and Technology, Tokyo Institute of Technology, Yokohama,

Kanagawa 226-8503, Japan

[7] Laboratory of Molecular Biophysics, Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

[8] Department of Regulatory Bioorganic Chemistry, The Institute of Scientific and Industrial Research, Osaka University, Ibaraki, Osaka 565-0047, Japan

[9] Research Center for Micro-nano Technology, Hosei University, Koganei, Tokyo 184-0003, Japan

[10] Department of Neuroscience, Mayo Clinic, Jacksonville, FL 32224, USA

[11] Department of Psychiatry, Osaka University Graduate School of Medicine, Suita, Osaka 565-0871, Japan

[12] Department of Chemistry, Keio University, Yokohama, Kanagawa 223-8522, Japan

[13] Department of Neurology, Clinical Neuroscience Branch, Brain Research Institute, Niigata University, Niigata 951-8585, Japan

[14] Life Science Research Institute, Kindai University, Osaka-Sayama, Osaka 589-8511, Japan

[15] Department of Neurology, Osaka University Graduate School of Medicine, Suita, Osaka 565-0871, Japan

[16] Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology, Yokohama, Kanagawa 226-8503, Japan

*Correspondence

Department of Neurology
Kindai University Faculty of Medicine
377-2 Ohnohigashi, Osaka-Sayama, Osaka 589-8511, Japan
Tel.: +81 (72) 366-0246 (ext. 3553); Fax: +81 (72) 368-4846
E-mail: yoshi.nagai@med.kindai.ac.jp

Abstract

Abnormal expansions of GGGGCC repeat sequence in the noncoding region of the *C9orf72* gene is the most common cause of familial amyotrophic lateral sclerosis and frontotemporal dementia (C9-ALS/FTD). The expanded repeat sequence is translated into dipeptide repeat proteins (DPRs) by noncanonical repeat-associated non-AUG (RAN) translation. Since DPRs play central roles in the pathogenesis of C9-ALS/FTD, we here investigate the regulatory mechanisms of RAN translation, focusing on the effects of RNA-binding proteins (RBPs) targeting GGGGCC repeat RNAs. Using C9-ALS/FTD model flies, we demonstrated that the ALS/FTD-linked RBP FUS suppresses RAN translation and neurodegeneration in an RNA-binding activity-dependent manner. Moreover, we found that FUS directly binds to and modulates the G-quadruplex structure of GGGGCC repeat RNA as an RNA chaperone, resulting in the suppression of RAN translation *in vitro*. These results reveal a previously unrecognized regulatory mechanism of RAN translation by G-quadruplex-targeting RBPs, providing therapeutic insights for C9-ALS/FTD and other repeat expansion diseases.

Introduction

Amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) are incurable neurodegenerative diseases with overlapping genetic and neuropathological features. Abnormal expansions of the GGGGCC (G_4C_2) repeat sequence in the noncoding region of the *C9orf72* gene have been found to be the most common genetic mutation responsible for ALS/FTD (DeJesus-Hernandez et al.; 2011, Gijselinck et al., 2012; Renton et al., 2011). Three major pathomechanisms are thought to be involved in the pathogenesis of *C9orf72*-linked ALS/FTD (C9-ALS/FTD): first, expansion of the G_4C_2 repeats results in decreased expression of the *C9orf72* gene, leading to its haploinsufficiency (Boivin et al., 2020; DeJesus-Hernandez et al. 2011; Gijselinck et al., 2012; Shi et al., 2018; Waite et al., 2014; Zhu et al., 2020). Second, the transcribed G_4C_2 repeat-containing RNA accumulates as RNA foci in the affected tissues, sequestering various RNA-binding proteins (RBPs) and altering their function (Conlon et al., 2016; Cooper-Knock et al., 2014; Donnelly et al., 2013; Haeusler et al., 2014; Lee et al., 2013; Mori et al., 2013a). Third, these G_4C_2 repeat RNA is also translated into dipeptide repeat (DPR) proteins, despite the lack of an AUG initiation codon, by noncanonical repeat-associated non-AUG (RAN) translation (Ash et al., 2013; Gendron et al., 2013; Mori et al., 2013b; Mori et al., 2013c; Zu et al, 2011; Zu et al, 2013). Since RAN translation occurs in all reading frames and the expanded G_4C_2 repeat sequence is bidirectionally transcribed, five distinct DPRs, i.e., poly(glycine-arginine) [poly(GR)], poly(glycine-alanine) [poly(GA)], poly(glycine-proline) [poly(GP)], poly(proline-arginine) [poly(PR)], and poly(proline-alanine) [poly(PA)] are produced and observed in patients' brains (Ash et al., 2013; Gendron et al., 2013; Mori et al., 2013b; Mori et al., 2013c; Zu et al., 2013) and cerebrospinal fluid (Gendron et al., 2017; Krishnon et al.,

2022; Lehmer et al., 2017; Su et al., 2014).

DPRs have been shown to exert toxic effects in multiple C9-ALS/FTD models, such

as cultured cells, flies, and mice (Choi et al., 2019; Jovičić et al., 2015; May et al. 2014;

Mizielinska et al., 2014; Rudich et al., 2017; Wen et al., 2014; Zhang et al., 2016; Zhang

et al., 2018). Importantly, the toxicity of DPRs was confirmed in DPR-only flies, which

express DPRs translated from non-G_4C_2 repeat RNAs with alternative codons and show

neurodegeneration, whereas RNA-only flies expressing G_4C_2 repeat RNAs with stop

codon interruptions, which eliminate DPRs production, did not show any obvious

degenerative phenotypes (Mizielinska et al., 2014). In addition, increased DPR

production, but not RNA foci, was reported to correlate with G_4C_2 repeat-induced

toxicity in a C9-ALS/FTD *Drosophila* model (Tran et al., 2015). Taken together, these

studies have strongly suggested that DPRs play a central role in the pathogenesis of C9-

ALS/FTD. Indeed, DPRs have been reported to disrupt various biological pathways,

such as nucleocytoplasmic transport (Hutten et al., 2020; Jovičić et al., 2015; Zhang et

al., 2016) and membraneless organelle dynamics (Kwon et al., 2014; Lee et al., 2016;

Lin et al., 2016). Therefore, elucidating the regulatory mechanism of RAN translation is

a significant challenge toward developing potential therapies for C9-ALS/FTD.

Since the discovery of RAN translation in 2011 (Zu et al., 2011), many studies to

date have focused on its molecular mechanisms, that is, whether it has functional

overlap with canonical AUG-dependent translation. Previous studies on C9-ALS/FTD

using monocistronic reporters containing a G_4C_2 repeat sequence revealed cap-

dependent translation initiation from the upstream near-cognate CUG initiation codon,

requiring the cap-binding eukaryotic translation factor 4F complex (Green et al., 2017;

Tabet et al., 2018). On the other hand, studies using bicistronic reporters with a G_4C_2

repeat sequence in the second cistron also produced DPRs by RAN translation in all

reading frames, suggesting cap-independent translation initiation within the G_4C_2 repeat

sequence (Cheng et al., 2018; Sonobe et al., 2018). This is reminiscent of internal

ribosomal entry site translation initiation, which is another type of noncanonical cap-

independent translation in which specific factors are directly recruited to the highly-

structured mRNA for initiation (Kwan et al., 2019). While such initiation mechanisms

of RAN translation have been explored to date, specific roles of the repeat sequence on

RAN translation remain enigmatic. Considering a repeat length dependency of RAN

translation (Mori et al., 2013c; Zu et al., 2011; Zu et al., 2013), the repeat sequence

itself would also be essential for the initiation or elongation steps of RAN translation.

Based on our previous findings of the protective role of TDP-43 on UGGAA repeat-

induced toxicity in spinocerebellar ataxia type 31 (SCA31) models (Ishiguro et al.,

2017), we hypothesized that RBPs specifically binding to repeat sequences of template

RNA play a role in RAN translation.

Using *Drosophila* models of C9-ALS/FTD, we here demonstrate the regulatory

roles of the ALS/FTD-linked RBP FUS on RAN translation from G_4C_2 repeat RNA,

which lead to the significant modulation of neurodegeneration. We found that FUS

suppresses RNA foci formation and DPR production, resulting in the suppression of

repeat-induced degeneration. This suppressive effect on degeneration was abolished by

mutations in the RNA-recognition motif (RRM) of FUS. In contrast, knockdown of

endogenous *caz*, a *Drosophila* homologue of *FUS*, enhanced DPR aggregation and

RNA foci formation, resulting in the enhancement of repeat-induced degeneration.

Moreover, FUS was found to directly bind to G_4C_2 repeat RNA and modify its G-

quadruplex structure as an RNA chaperone, resulting in the suppression of RAN

translation *in vitro*. In addition, other G-quadruplex-targeting RBPs also suppressed

RAN translation and G_4C_2 repeat-induced toxicity in our C9-ALS/FTD flies. These

results strongly indicate that FUS regulates RAN translation and suppresses DPR

toxicity through modulating the G-quadruplex structure of G_4C_2 repeat RNA. Our

findings shed light on the regulatory mechanisms of RAN translation by G-quadruplex-

targeting RBPs, and propose novel therapeutic strategies for repeat expansion diseases

by regulating RAN translation.

Results

Screening for RBPs that suppress G_4C_2 repeat-induced toxicity in C9-ALS/FTD

flies

We established *Drosophila* models of C9-ALS/FTD that express pathogenic

length 42 or 89 G_4C_2 repeats [$(G_4C_2)_{42}$, or $(G_4C_2)_{89}$ flies, respectively], and confirmed

that expanded G_4C_2 repeat sequences induce eye degeneration and motor dysfunction

accompanied with the formation of RNA foci and the production of three types of DPRs

(Figure 1—figure supplement 1), consistent with previous studies (Freibaum et al.,

2015; Goodman et al., 2019; Mizielinska et al., 2014; Xu et al., 2013). We also

established *Drosophila* expressing normal length 9 G_4C_2 repeats as a control [$(G_4C_2)_9$

flies], and found that they did not show eye degeneration, motor dysfunction, RNA foci

formation, or DPR aggregation (Figure 1—figure supplement 1). We selected 18 RBPs

that have been reported to bind to G_4C_2 repeat RNA (Mori et al., 2013a), as well as

TDP-43, an ALS/FTD-linked RBP that does not bind to G_4C_2 repeat RNA (Xu et al.,

2013) (Figure 1—source data 1), and examined their roles in neurodegeneration in our

C9-ALS/FTD fly models. We found that coexpression of *FUS*, *IGF2BP1*, or

hnRNPA2B1 strongly suppressed the eye degeneration in both flies expressing $(G_4C_2)_{42}$ $_{or\ 89}$, which show decreased eye size and loss of pigmentation (Figures 1A–1D and Figure 1—source data 2). Coexpression of five RBPs, namely, *hnRNPR*, *SAFB2*, *SF3B3*, *hnRNPA1*, and *hnRNPL* also partially suppressed the eye degeneration, whereas coexpression of the other six RBPs had no effect, and two RBPs enhanced the phenotypes (Figures 1A–1D and Figure 1—source data 2). In addition, coexpression of *TDP-43* had no effect on the eye degeneration in $(G_4C_2)_{42}$ flies, and resulted in lethality in $(G_4C_2)_{89}$ flies, likely due to the toxicity of *TDP-43* expression itself (Figures 1A and 1D, and Figure 1—source data 2). The variation in the effects of these G_4C_2 repeat-binding RBPs on G_4C_2 repeat-induced toxicity may be due to their different binding affinities to G_4C_2 repeat RNA, and the different toxicity of overexpressed RBPs themselves. We then analyzed the expression levels of G_4C_2 repeat RNA in flies coexpressing $(G_4C_2)_{89}$ and three RBPs that strongly suppressed eye degeneration. We found that coexpression of *IGF2BP1* or *hnRNPA2B1* significantly decreased G_4C_2 repeat RNA levels, whereas they were not altered upon coexpression of *FUS* (Figure 1E). Although the suppressive effects of IGF2BP1 and hnRNPA2B1 could simply be explained by the decreased levels of G_4C_2 repeat RNA, the molecular mechanisms by which FUS suppresses G_4C_2 repeat-induced toxicity remain to be clarified. The suppressive effects of FUS on G_4C_2 repeat-induced toxicity was confirmed using multiple FUS fly lines, showing the significant suppression of decreased eye size and loss of pigmentation in $(G_4C_2)_{42\ or\ 89}$ flies coexpressing *FUS* (Figure 1—figure supplement 2). Therefore, we decided to further focus on FUS, which is another ALS/FTD-linked RBP, and investigated its mechanism of the suppression of G_4C_2 repeat-induced toxicity.

FUS suppresses G_4C_2 repeat-induced toxicity via its RNA-binding activity

We next investigated whether the suppressive effects of FUS on G_4C_2 repeat-induced toxicity are mediated by its binding to G_4C_2 repeat RNA, using flies expressing FUS with mutations in the RRM (*FUS-RRMmut*), which have been reported to eliminate its RNA-binding activity (Daigle et al., 2013). Western blot analysis confirmed that the *FUS-RRMmut* fly line expresses almost an equivalent level of the FUS proteins to the *FUS* fly line (Figure 2—figure supplement 1). We found that coexpression of *FUS-RRMmut* did not restore the eye degeneration in flies expressing $(G_4C_2)_{89}$, suggesting that the RNA-binding activity of FUS is essential for its suppressive effects on G_4C_2 repeat-induced toxicity (Figures 2A–C). We also evaluated the reduced egg-to-adult viability of $(G_4C_2)_{42}$ flies, and confirmed that this phenotype was rescued by coexpression of *FUS*, but not by coexpression of *FUS-RRMmut* (Figure 2D). Expression of G_4C_2 repeat RNA in the nervous system of flies after eclosion using the *elav-GeneSwitch* driver induces motor dysfunction, and coexpression of *FUS* significantly alleviated this motor dysfunction (Figure 2E), indicating that FUS suppresses the neuronal phenotypes of flies expressing G_4C_2 repeat RNA. It is notable that the motor dysfunction caused by the expression of *FUS* alone was also alleviated by coexpression of $(G_4C_2)_{42}$ (Figure 2E), indicating that the G_4C_2 repeat RNA conversely suppresses FUS toxicity. This result is consistent with our previous observations in SCA31 flies that UGGAA repeat RNA reduced the aggregation and toxicity of TDP-43 (Ishiguro et al., 2017). Moreover, recent studies demonstrated that RNA buffers the phase separation of TDP-43 and FUS, resulting in the suppression of their aggregation (Maharana et al., 2018; Mann et al., 2019). These findings hence

suggest that balancing the crosstalk between repeat RNAs and RBPs neutralizes the

toxicities of each other.

FUS suppresses RNA foci formation and RAN translation from G_4C_2 repeat RNA

We next analyzed the effects of FUS expression on RNA foci and DPR

production in flies expressing G_4C_2 repeat RNA. We performed RNA fluorescence *in*

situ hybridization (FISH) of the salivary glands of fly larvae expressing $(G_4C_2)_{89}$, and

found that coexpression of *FUS* significantly decreased the number of nuclei containing

RNA foci in $(G_4C_2)_{89}$ flies, whereas it was not altered by coexpression of *FUS-RRMmut*

(Figures 3A and 3B). We confirmed that the expression levels of G_4C_2 repeat RNA in

$(G_4C_2)_{89}$ flies were not altered by coexpression of *FUS* or *FUS-RRMmut* (Figure 3C).

These results were in good agreement with our previous study on SCA31 showing the

suppressive effects of FUS and other RBPs on RNA foci formation of UGGAA repeat

RNA through altering RNA structures and preventing aggregation of misfolded repeat

RNA as RNA chaperones (Ishiguro et al., 2017), raising the possibility that FUS has

RNA-chaperoning activity also for G_4C_2 repeat RNA. Immunohistochemistry of the eye

imaginal discs of fly larvae expressing $(G_4C_2)_{89}$ revealed that coexpression of *FUS*

significantly decreased the number of DPR aggregates in $(G_4C_2)_{89}$ flies, whereas

coexpression of *FUS-RRMmut* did not (Figures 3D and 3E). Quantitative analyses of

poly(GP) by immunoassay also demonstrated that poly(GP) levels were greatly

decreased in $(G_4C_2)_{89}$ flies upon coexpression of *FUS*, but not *FUS-RRMmut* (Fig 3F),

indicating that FUS suppresses RAN translation from the G_4C_2 repeat RNA to reduce

DPR production. Considering that the 5′ upstream sequence of the G_4C_2 repeat in the

C9orf72 gene is reported to affect RAN translation activity (Green et al., 2017; Tabet et

al., 2018), we used flies expressing the G_4C_2 repeat sequence with the upstream intronic

sequence of the *C9orf72* gene, namely, *LDS-(G_4C_2)$_{44}$*[GR-GFP] (Goodman et al., 2019).

Since this construct has a 3′-green fluorescent protein (GFP) tag in the GR reading

frame downstream of the G_4C_2 repeat sequence, the GR-GFP fusion protein is produced

by RAN translation (Figure 3—figure supplement 1). We confirmed that coexpression

of *FUS* significantly decreased the expression level of GR-GFP, whereas coexpression

of *FUS-RRMmut* had no effect (Figures 3G–3I).

We further excluded the possibility that FUS directly interacts with DPRs, rather

than with G_4C_2 repeat RNA, to decrease DPR levels and exert its suppressive effects.

Using DPR-only flies expressing DPRs translated from non-G_4C_2 RNAs with

alternative codons (Mizielinska et al., 2014), we confirmed that FUS did not suppress

the eye degeneration in DPR-only flies expressing poly(GR), but rather enhanced their

phenotypes, likely due to the additive effects of FUS toxicity (Figure 3—figure

supplement 2). Together with the finding that FUS decreases not only DPR expression

but also RNA foci formation (Figures 3A and 3B), these results collectively indicate that

FUS indeed interacts with G_4C_2 repeat RNA and regulates RAN translation from G_4C_2

repeat RNA in *Drosophila* models of C9-ALS/FTD.

Reduction of endogenous *caz* expression enhances G_4C_2 repeat-induced toxicity,

RNA foci formation, and DPR aggregation

To elucidate the physiological role of FUS on RAN translation, we also

investigated the role of endogenous *caz*, a *Drosophila* homologue of *FUS*, on G_4C_2

repeat-induced toxicity in flies expressing G_4C_2 repeat RNAs. Coexpression of *caz* as

well as *FUS* suppressed eye degeneration in flies expressing *(G_4C_2)$_{42 \text{ or } 89}$* (Figure 4—

figure supplement 1). These data suggest that *caz* is a functional homologue of *FUS*. In

contrast, knockdown of *caz* by RNA interference or its hemizygous deletion modestly

but significantly enhanced the eye degeneration in $(G_4C_2)_{89}$ flies (Figures 4A–4D),

indicating that reduced *caz* expression enhances G_4C_2 repeat-induced toxicity. We next

analyzed the effects of *caz* knockdown on RNA foci formation and DPR production in

flies expressing $(G_4C_2)_{89}$. FISH analysis of the salivary glands revealed that knockdown

of *caz* significantly increased the number of nuclei containing RNA foci in $(G_4C_2)_{89}$

flies (Figures 4E and 4F). We also confirmed that the expression levels of G_4C_2 repeat

RNA in $(G_4C_2)_{89}$ flies were not altered by the knockdown of *caz* (Figure 4G).

Immunohistochemical analysis showed that knockdown of *caz* significantly increased

the number of DPR aggregates in $(G_4C_2)_{89}$ flies (Figures 4H and 4I). These results

indicate that the reduction of *caz* expression enhances RNA foci formation and DPR

aggregation, compatible with the results of *FUS* coexpression in flies expressing

$(G_4C_2)_{89}$ (Figure 3), and that FUS functions as an endogenous regulator of RAN

translation.

FUS directly binds to and modulates the G-quadruplex structure of G_4C_2 repeat

RNA, resulting in the suppression of RAN translation *in vitro*

We next confirmed the direct interaction of FUS with G_4C_2 repeat RNA by the

filter binding assay. We found that His-tagged FUS binds to the $(G_4C_2)_4$ RNA in a dose-

dependent manner, but not to the control $(AAAAAA)_4$ RNA (Figure 5A), and His-

tagged FUS-RRMmut had almost no binding affinity to the $(G_4C_2)_4$ RNA, consistent

with a previous study (Mori et al., 2013a). We also confirmed the interaction of FUS

with the G_4C_2 repeat RNA in our C9-ALS/FTD flies by showing the colocalization of

FUS with the RNA foci (Figure 5—figure supplement 1), consistent with a recent study

using C9-ALS/FTD patient fibroblasts (Bajc Česnik et al., 2019). Since G_4C_2 repeat

RNA was reported to form both G-quadruplex and hairpin structures (Fratta et al., 2012;

Hauesler et al., 2014; Reddy et al., 2013; Su et al., 2014), we next characterized the

interactions of FUS with G_4C_2 repeat RNA having different structures. G_4C_2 repeat

RNA is known to form G-quadruplex structures in the presence of K^+, whereas they

form hairpin structures in the presence of Na^+ (Su et al., 2014). Surface plasmon

resonance (SPR) analyses demonstrated that FUS preferentially binds to $(G_4C_2)_4$ RNA

with the G-quadruplex structure in KCl buffer (Table 1, dissociation constant (K_D) =

1.5×10^{-8} M), and weakly to $(G_4C_2)_4$ RNA with the hairpin structure in NaCl buffer

(Table 1, $K_D = 1.3 \times 10^{-7}$ μM). We also confirmed that FUS has poor binding affinity to

$(G_4C_2)_4$ RNA in LiCl buffer (Table 1, $K_D = 1.4 \times 10^{-5}$ μM), which destabilizes the G-

quadruplex structure (Herdy et al., 1992), and was an almost similar level to its binding

affinity to the negative control $(A_4C_2)_4$ RNA (not shown). These results suggest the

preferential binding of FUS to G_4C_2 repeat RNA with the G-quadruplex structure,

which is consistent with a previous report showing preferential binding of FUS to G-

quadruplex structured Sc1 and DNMT RNAs (Ozdilek et al., 2017). Considering that

higher-order structures, including G-quadruplex and hairpin structures, are reported to

be involved in RAN translation (Mori et al., 2021; Simone et al., 2018; Wang et al.,

2019; Zu et al., 2011), we next investigated the effects of FUS on the structure of G_4C_2

repeat RNA. The circular dichroism (CD) spectrum of $(G_4C_2)_4$ RNA in KCl buffer was

found to exhibit a positive peak at approximately 260 nm and a negative peak at 240 nm

(Figure 5B, black line), consistent with previous reports (Fratta et al., 2012; Hauesler et

al., 2014; Reddy et al., 2013; Su et al., 2014). Interestingly, upon the addition of FUS,

these two peaks were notably shifted to longer wavelengths with substantial CD

spectrum changes, indicating a significant structural alteration in $(G_4C_2)_4$ RNA (Figure

5B, red line). We confirmed that the CD spectrum of FUS alone in the wavelength range

of 240 to 300 nm was almost negligible (Figure 5—figure supplement 2A, green line),

indicating that this change in CD spectrum is attributed to structural changes in the

$(G_4C_2)_4$ RNA. We also observed CD spectrum changes to some extent in the $(G_4C_2)_4$

RNA upon the addition of FUS in NaCl buffer, but not in LiCl buffer, confirming an

interaction between FUS and hairpin-structured $(G_4C_2)_4$ RNA as well (Figures 5C and

5D). We further analyzed the interaction between FUS and G_4C_2 repeat RNA by imino

proton nuclear magnetic resonance (NMR). In KCl buffer, the NMR signals of the

imino proton for the G-quadruplex structure of $(G_4C_2)_4$ RNA were detected in the region

around 10 to 12 ppm (Figure 5—figure supplement 2D), consistent with previous

studies (Fratta et al., 2012; Su et al., 2014). Upon the addition of FUS, the NMR

intensities of $(G_4C_2)_4$ RNA were decreased in an FUS concentration-dependent manner

(Figure 5—figure supplement 2D), further supporting the possibility that that FUS

interacts with and modulates the G-quadruplex structure of $(G_4C_2)_4$ RNA. Collectively,

these results indicate that FUS directly binds to G_4C_2 repeat RNA, preferentially to its

G-quadruplex form, and modulates its higher-order structures. These structural

alterations of G_4C_2 repeat RNAs by FUS did not require ATP or interactions with other

proteins, suggesting its role as an RNA chaperone for G_4C_2 repeat RNA (Rajkowitsch et

al., 2007).

To further clarify the direct link between the binding of FUS to G_4C_2 repeat

RNA and its effects on RAN translation, we employed a cell-free *in vitro* translation

assay using rabbit reticulocyte lysate. We designed a reporter construct containing the

80 G_4C_2 repeat sequence with the 5′ upstream intronic sequence of the *C9orf72* gene

and the Myc tag sequence in the GA reading frame at the 3′ downstream (Figure 5E).

This upstream sequence contained multiple stop codons in each reading frame and

lacked AUG initiation codons. We confirmed by Western blotting that this reporter

system indeed produces GA-Myc by RAN translation, consistent with previous studies

(Green et al., 2017; Tabet et al., 2018). Notably, upon the addition to this translation

system, FUS suppressed RAN translation efficiently, whereas FUS-RRMmut did not.

FUS decreased the expression levels of GA-Myc at as low as 10nM, and nearly

eliminated RAN translation activity at 100nM. At 400nM, FUS-RRMmut weakly

suppressed the GA-Myc expression levels probably because of the residual RNA-

binding activity (Figures 5F and 5G). Taken together, these results indicate that FUS

suppresses RAN translation *in vitro* through direct interactions with G_4C_2 repeat RNA

as an RNA chaperone.

Identification of G-quadruplex-targeting RBPs that suppress G_4C_2 repeat-induced

toxicity in C9-ALS/FTD flies

Considering that FUS suppresses G_4C_2 repeat-induced toxicity as an RNA

chaperone through its preferential binding to the G-quadruplex structure of G_4C_2 repeat

RNA (Figure 5 and Table 1), we hypothesized that other G-quadruplex-targeting RBPs

might have similar suppressive effects on G_4C_2 repeat-induced toxicity. To investigate

this possibility, we selected 6 representative G-quadruplex-targeting RBPs, all of which

are known to bind to G_4C_2 RNA as well (Cooper-Knock et al., 2014; Haeusler et al.,

2014; Mori et al., 2013a; Xu et al., 2013) (Figure 6—source data 1). Intriguingly,

coexpression of *EWSR1*, *DDX3X*, *DDX5*, or *DDX17* significantly suppressed eye

degeneration in $(G_4C_2)_{89}$ flies without altering G_4C_2 RNA expression (Figures 6A-6D).

As expected, these RBPs also decreased the number of poly(GA) aggregates in the eye

imaginal discs (Figures 6E and 6F). Their effects on G_4C_2 repeat-induced toxicity,

repeat RNA expression, and RAN translation were consistent with those of FUS. In

support of our results, DDX3X was previously reported to suppress RAN translation

and G_4C_2 repeat-induced toxicity in cell culture in a helicase-activity-dependent manner

(Cheng et al., 2019). On the other hand, coexpression of *DHX9* or *DHX36* suppressed

eye degeneration by reducing G_4C_2 repeat RNA levels (Figures 6A-6D). Since G-

quadruplex-targeting RBPs have diverse biological functions, including transcription,

RNA processing, translation, and RNA stabilization (Dumas et al., 2021), these different

effects among G-quadruplex-targeting RBPs on G_4C_2 repeat RNA expression might be

attributed to their different roles in RNA metabolism. Thus, some G-quadruplex-

targeting RBPs regulate RAN translation and G_4C_2 repeat-induced toxicity by binding

to and possibly by modulating the G-quadruplex structure of G_4C_2 repeat RNA.

Discussion

In this study, we revealed a novel regulatory mechanism of RAN translation from

expanded G_4C_2 repeat RNA by the ALS/FTD-linked RBP FUS, which suppresses DPR

production and neurodegeneration in C9-ALS/FTD *Drosophila* models (Figures 1–4).

FUS directly binds to G_4C_2 repeat RNA and modulates its G-quadruplex structure as

evident by CD and NMR analyses (Figure 5, Figure 5—figure supplement 2), and

suppresses RNA foci formation *in vivo* (Figures 3A and 3B), suggesting its functional

role as an RNA chaperone. This is reminiscent of our recent study on SCA31, in which

we demonstrated a novel role of the ALS/FTD-linked RBPs TDP-43, FUS, and

hnRNPA2B1 as RNA chaperones binding to UGGAA repeat RNA and altering its

structure, resulting in the suppression of its neurotoxicity through reducing RNA foci

formation and repeat polypeptide translation (Ishiguro et al., 2017). Considering the

similarities of the effects of FUS on G_4C_2 repeat RNA and UGGAA repeat RNA, we

conclude that FUS functions as an RNA chaperone also for G_4C_2 repeat RNA to

regulate its RAN translation.

The suppressive effects of RBPs in several noncoding repeat expansion diseases by

the amelioration of their sequestration into RNA foci have been reported. For example,

in myotonic dystrophy type 1, MBNL1 was shown to be sequestered into CUG repeat

RNA foci, and overexpression of MBNL1 in a mouse model was found to compensate

its functional loss, resulting in the reversal of myotonia (Kanadia et al., 2006).

Similarly, previous studies reported the suppressive effects of other RBPs on

neurodegeneration, such as hnRNPA2B1 in fragile X ataxia/tremor syndrome (Sofora et

al., 2007), and Pur-α and Zfp106 in C9-ALS/FTD (Xu et al., 2013; Celona et al., 2017).

The suppressive effects of these RBPs have been thought to result from the

supplementation against their loss-of-function due to their sequestration into RNA foci,

although their effects on gain-of-toxic disease pathomechanisms, that is, RAN

translation and repeat RNA expression, remain to be elucidated. In contrast, in this

study we demonstrated that FUS suppresses neurodegeneration in C9-ALS/FTD by

directly targeting G_4C_2 repeat RNA and inhibiting RAN translation as an RNA

chaperone. Similar suppressive effects of RBPs by targeting UGGAA repeat RNA in

SCA31 as RNA chaperones have also been reported (Ishiguro et al., 2017). In addition,

we also showed that the expression of IGF2BP1, hnRNPA2B1, DHX9, and DHX36

decreased G_4C_2 repeat RNA expression and suppressed eye degeneration in our C9-

ALS/FTD *Drosophila* model (Figures 1 and 6), likely via the reduction of DPR levels.

Similarly, we recently reported that hnRNPA3 reduces G_4C_2 repeat RNA expression

levels, leading to the suppression of neurodegeneration in C9-ALS/FTD fly models

(Taminato et al., 2023). Interestingly, these RBPs have been reported to be involved in

RNA decay pathways as components of the P-body or interactors with the RNA

deadenylation machinery (Tran et al., 2004; Katahira et al., 2008; Geissler et al., 2016;

Hubstenberger et al., 2017), possibly contributing to the reduced expression levels of

G_4C_2 repeat RNA. In myotonic dystrophy type 2 models, MBNL1 was also reported to

retain CCUG repeat RNA in the nucleus, resulting in the suppression of RAN

translation (Zu et al., 2017), implying various mechanisms of the effects of RBPs

depending on the combination of RBPs and repeat RNA. Nevertheless, our findings

highlighted the previously unrecognized roles of RBPs directly interacting with repeat

RNA and modulating gain-of-toxic pathomechanisms, including RAN translation in

noncoding repeat expansion diseases.

Several studies have indicated the importance of higher-order structures of repeat

RNA in RAN translation. In SCA8 models, hairpin-forming CAG repeat RNA was

shown to be RAN-translated to produce polyglutamine proteins, but switching the CAG

repeats to non-hairpin-forming CAA repeats abolished the RAN translation (Zu et al.,

2011). In C9-ALS/FTD, G_4C_2 repeat RNA has been reported to form both hairpin and

G-quadruplex structures (Fratta et al., 2012; Hauesler et al., 2014; Reddy et al., 2013;

Su et al., 2014). Although the effect of each structure on RAN translation remains

largely unknown, small molecules binding to the hairpin structure or the G-quadruplex

structure were both reported to inhibit RAN translation from the G_4C_2 repeat RNA,

resulting in reduced DPR levels (Wang et al., 2019, Mori et al., 2021). These findings

are in accordance with our results showing that FUS modifies the G-quadruplex

structure as well as the hairpin structure of G_4C_2 repeat RNA as an RNA chaperone and

reduces DPR production. We further found that G-quadruplex-targeting RNA helicases,

including DDX3X, DDX5, and DDX17, which are known to bind to G_4C_2 repeat RNA

(Cooper-Knock et al., 2014; Haeusler et al., 2014; Mori et al., 2013a; Xu et al., 2013),

also suppress RAN translation and G_4C_2 repeat-induced toxicity without altering the

expression levels of G_4C_2 repeat RNA in our *Drosophila* models. These results suggest

that not only ATP-independent RNA chaperones, but also ATP-dependent RNA

helicases may regulate RAN translation through modifying the higher-order structures

of template repeat RNA. Consistently, a previous study also reported that DDX3X

inhibits RAN translation from G_4C_2 repeat RNA in a helicase activity-dependent

manner (Cheng et al., 2019). Knockdown of another RNA helicase, DHX36, has been

reported to both promote (Cheng et al., 2019) and inhibit (Liu et al., 2021; Tseng et al.,

2021) RAN translation, possibly due to the different effects on repeat RNA structures

depending on the experimental conditions. Unfortunately, most of these studies

reporting the effects of RBPs on RAN translation have limitations of the detailed

structural analyses of repeat RNA. In this study, focusing on FUS, we performed a

series of molecular structural analyses, *in vitro* translation assays, and *in vivo* genetic

analyses to clarify the structure-function relationship of G_4C_2 repeat RNA, and provide

compelling evidence for the modifying effects of FUS on repeat RNA structures leading

to the suppression of RAN translation and repeat-induced toxicity *in vivo*.

FUS has an RRM domain for RNA binding and a low complexity (LC) domain

involved in protein interactions, and exerts multifaceted functions, such as RNA

transcription, RNA splicing, RNA transport, and formation of membraneless organelles,

such as stress granules and nuclear paraspeckles via liquid-liquid phase separation

(Lagier-Tourenne et al., 2010). Recent studies reported that arginine-rich DPRs, such as

poly(GR) and poly(PR), interact with LC domain-containing RBPs, including FUS, and

alter their liquid-liquid phase separation, resulting in the disruption of the dynamics and

functions of membraneless organelles (Kwon et al., 2014; Lee et al., 2016; Lin et al.,

2016). These findings raise the possibility that FUS may exert its suppressive effects by

directly interacting with DPRs. However, we showed that FUS does not suppress eye

degeneration in DPR-only flies (Figure 3—figure supplement 2) indicating that a direct

interaction between FUS and DPRs is unlikely to be the mechanism of the suppression

of DPR toxicity in our C9-ALS/FTD flies. This result supports our conclusion that FUS

suppresses G_4C_2 repeat-induced toxicity through direct binding to G_4C_2 repeat RNA.

In summary, we here provided evidence that FUS modulates the structure of G_4C_2

repeat RNA as an RNA chaperone, and regulates RAN translation, resulting in the

suppression of neurodegeneration in C9-ALS/FTD fly models. Recent advances in

genome sequencing technology unveiled that such expansions of repeat sequences cause

more than 50 monogenic human diseases (Malik et al., 2021), and are also associated

with psychiatric diseases such as autism (Mitra et al., 2021; Trost et al., 2020). Thus,

our findings contribute to the elucidation of the repeat-associated pathogenic

mechanisms underlying not only C9-ALS/FTD, but also a broader range of

neuromuscular and neuropsychiatric diseases than previously thought, and will advance

the development of potential therapies for these diseases.

Materials and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (*D. melanogaster*)	*UAS-(G₄C₂)ₙ, UAS-FUS-2, UAS-FUS-RRMmut*	This paper	N/A	See Materials and Methods, "Generation of constructs and transgenic flies"
strain, strain background (*D. melanogaster*)	*UAS-RBP (FUS-3; IGF2BP1; hnRNPA2B1; hnRNPR; SAFB2; SF3B3; hnRNPA1; hnRNPL; DHX30; SAFB; DHX15; ILF2; DDX21; hnRNPK; SFPQ; ILF3; NONO; ELAVL1; DDX3X; DDX5; DDX17; DHX9; DHX36)*	This paper	N/A	See Materials and Methods, "Generation of constructs and transgenic flies"
strain, strain background (*D. melanogaster*)	*UAS-LDS-(G4C2)₄₄^{GR-GFP}*	Goodman et al., 2019. (PMID: 31110321)	FLYB: FBtp0135960	
strain, strain background (*D. melanogaster*)	*UAS-FUS*	Ishiguro et al., 2017. (PMID: 28343865)	FLYB: FBtp0117594	
strain, strain background (*D. melanogaster*)	*UAS-FUS-4 (UAS-FLAG-FUS)*	Wang et al., 2011 (PMID: 21881207)	FLYB: FBtp0070284	
strain, strain background (*D. melanogaster*)	*UAS-caz (UAS-FLAG-caz)*	Wang et al., 2011 (PMID: 21881207)	FLYB: FBtp0070279	
strain, strain background (*D. melanogaster*)	*caz²*	Frickenhaus et al., 2015 (PMID: 25772687)	FLYB: FBal0323133	

strain, strain background (*D. melanogaster*)	*UAS-TDP-43*	Ishiguro et al., 2017. (PMID: 28343865)	FLYB: FBtp011759 2	
strain, strain background (*D. melanogaster*)	*GMR-GAL4 driver*	Yamaguchi et al., 1999. (PMID: 10597285)	FLYB: FBtp001007 4	
strain, strain background (*D. melanogaster*)	Elav-GAL4 driver*: P{w[+mC]=GAL4-elav.L}2/CyO*	Bloomington Drosophila Stock Center	BDSC: 8765; FLYB: FBst0008765	
strain, strain background (*D. melanogaster*)	Elav-GeneSwitch GAL4 driver*: y[1] w[*]; P{w[+mC]=elav-Switch.O}GSG301*	Bloomington Drosophila Stock Center	BDSC: 43642; FLYB: FBst0043642	
strain, strain background (*D. melanogaster*)	*UAS-EGFP: w[*]; P{w[+mC] = UAS-2xEGFP}AH2*	Bloomington Drosophila Stock Center	BDSC: 6874; FLYB: FBst0006874	
strain, strain background (*D. melanogaster*)	*UAS-DsRed: w[*]; P{w[+mC]=UAS-AUG-DsRed}A*	Bloomington Drosophila Stock Center	BDSC: 6282; FLYB: FBst0006282	
strain, strain background (*D. melanogaster*)	*UAS-EWSR1: w[1118]; P{w[+mC]=UAS-EWSR1.C}26M*	Bloomington Drosophila Stock Center	BDSC: 79592; FLYB: FBst0007959 2	
strain, strain background (*D. melanogaster*)	*UAS-(GR)$_{36}$: w[1118]; P{{y[+t7.7] w[+mC]=UAS-poly-GR.PO-36}attP40*	Bloomington Drosophila Stock Center	BDSC: 58692; FLYB: FBst0005869 2	
strain, strain background (*D. melanogaster*)	*UAS-(GA)$_{36}$: w[1118]; P{{y[+t7.7] w[+mC]=UAS-poly-GA.PO-36}attP40*	Bloomington Drosophila Stock Center	BDSC: 58693; FLYB: FBst0005869 3	
strain, strain background (*D. melanogaster*)	*UAS-(GR)$_{100}$: w[1118]; P{{y[+t7.7] w[+mC]=UAS-poly-GR.PO-100}attP40*	Bloomington Drosophila Stock Center	BDSC: 58696; FLYB: FBst0005869 6	

strain, strain background (*D. melanogaster*)	UAS-(GA)$_{100}$: *w[1118]; P{{y[+t7.7] w[+mC]=UAS-poly-GA.PO-100}attP40*	Bloomington Drosophila Stock Center	BDSC: 58697; FLYB: FBst00058697	
strain, strain background (*D. melanogaster*)	RNAi of *GFP: w[1118]; P{w[+mC] = UAS-GFP.dsRNA.R}142*	Bloomington Drosophila Stock Center	BDSC: 9330; FLYB: FBst0009330	
strain, strain background (*D. melanogaster*)	RNAi of *caz*: P{KK107486}VIE-260B	Vienna Drosophila Resource Center	VDRC: v100291; FLYB: FBst0472165	
antibody	Rat monoclonal anti-poly(GR) antibody (5A2)	Millipore	Car# MABN778; RRID: AB_2728664	IHC(1:1000), WB(1:1000)
antibody	Mouse monoclonal anti-poly(GA) antibody (5E9)	Millipore	Car# MABN889; RRID: AB_2728663	IHC(1:1000)
antibody	Rabbit polyclonal anti-poly(GA) antibody	Cosmo Bio	Cat# CAC-TIP-C9-P01	IHC(1:1000)
antibody	Rabbit polyclonal anti-poly(GP) antibody	Novus Biologicals	Cat# NBP2-25018; RRID: AB_2893239	IHC(1:1000)
antibody	Rabbit polyclonal anti-FUS antibody	Bethyl Laboratories	Cat# A300-302A; RRID: AB_309445	IHC(1:1000), WB(1:1000)
antibody	Mouse monoclonal anti-EGFP antibody	Clontech	Cat# 632569	WB(1:1000)
antibody	Mouse monoclonal anti-actin antibody (AC-40)	Sigma-Aldrich	Cat# A4700; RRID: AB_476730	WB(1:1000)
antibody	Mouse monoclonal anti-c-Myc antibody (9E10)	Wako	Cat# 17-21876	WB(1:3000)
recombinant DNA reagent	pcDNA™5/FRT-*C9orf72* intron1-*(G$_4$C$_2$)$_{80}$* (plasmid)	This paper		See Materials and Methods, "RNA synthesis for in vitro

				translation"
sequence-based reagent	$(G_4C_2)_n$_F(1)	This paper	PCR primers	ATGAATGGGA GCAGTGGTGG
sequence-based reagent	$(G_4C_2)_n$_R(1)	This paper	PCR primers	TGTTGAGAGT CAGCAGTAGC C
sequence-based reagent	$(G_4C_2)_n$_F(2)	This paper	PCR primers	CCCAATCCAT ATGACTAGTA GATCC
sequence-based reagent	$(G_4C_2)_n$_R(2)	This paper	PCR primers	TGTAGGTAGT TTGTCCAATTA TGTCA
sequence-based reagent	gal4_F	Li et al., 2008. (PMID: 18449188)	PCR primers	TTGAAATCGC GTCGAAGGA
sequence-based reagent	gal4_R	Li et al., 2008. (PMID: 18449188)	PCR primers	GGCTCCAATG GCTAATATGC A
peptide, recombinant protein	His-FUS	This paper	N/A	See Materials and Methods, "Filter binding assay"
peptide, recombinant protein	His-FUS-RRMmut	This paper	N/A	See Materials and Methods, "Filter binding assay"
peptide, recombinant protein	FUS (not tagged)	This paper	N/A	See Materials and Methods, "Preparation of recombinant FUS protein"
peptide, recombinant protein	FUS-RRMmut (not tagged)	This paper	N/A	See Materials and Methods, "Preparation of recombinant FUS protein"
commercial assay or kit	In-Fusion Cloning system	TaKaRa Bio	Cat# Z9645N	
commercial assay or kit	EZ-Tn5 <KAN-2> Insertion Kit	Epicentre	Cat# EZI011RK	

commercial assay or kit	QuantiTect Reverse Transcription Kit	QIAGEN	Cat# 205314	
commercial assay or kit	mMESSAGE mMACHINE T7 Transcription Kit	Thermo Fisher Scientific	Cat# AM1344	
commercial assay or kit	Flexi Rabbit Reticulocyte Lysate System	Promega	Cat# L4540	
chemical compound, drug	RU486 (mifepristone)	Wako	M3321; CAS: 84371-65-3	
chemical compound, drug	Formula 4-24 Instant Drosophila medium	Wako	Cat# 534-20571	
software, algorithm	ZEN imaging software	Zeiss	RRID:SCR_013672; https://www.zeiss.com/microscopy/en/products/software/zeiss-zen.html	
software, algorithm	ImageJ	Schneider et al., 2012. (PMID: 22930834)	RRID:SCR_003070; https://imagej.nih.gov/ij/	
software, algorithm	GraphPad Prism Version 8.4.3	GraphPad Software Inc.	RRID:SCR_002798; https://www.graphpad.com	

Flies

All fly stocks were cultured and crossed at 23 °C or 25 °C in standard cornmeal-yeast-

glucose medium. Male adult flies were used for the climbing assay and *GeneSwitch*

experiments. Three to 5-day-old female adult flies were used for the evaluation of eye

phenotype using a stereoscopic microscope model SZX10 (Olympus). Female third

instar larvae were used for quantitative real-time polymerase chain reaction (PCR),

RNA FISH and immunohistochemistry experiments. The transgenic fly line bearing the

GMR-Gal4 transgene has been described previously (Yamaguchi et al., 1999). The

transgenic fly lines bearing *elav-Gal4* (#8765), *elav-GeneSwitch* (#43642), *UAS-EGFP*

(#6874), *UAS-DsRed* (#6282), *UAS-GFP-IR* (inverted repeat) (#9330), *UAS-(GR)$_{36}$*

(#58692), *UAS-(GA)$_{36}$* (#58693), *UAS-(GR)$_{100}$* (#58696), *UAS-(GA)$_{100}$* (#58697), and

UAS-EWSR1 (#79592) were obtained from Bloomington Drosophila Stock Center. The

transgenic fly line bearing *UAS-caz-IR* (#100291) was obtained from Vienna Drosophila

Resource Center. The fly line with the *caz* null allele (*caz^2*), *UAS-LDS-(G$_4$C$_2$)$_{44}$$^{GR-GFP}$*,

and *UAS-caz* (*UAS-FLAG-caz*) and *UAS-FUS*-4 (*UAS-FLAG-FUS*) were kind gifts from

Dr. Erik Storkebaum (Frickenhaus et al., 2015), Dr. Nancy Bonini (Goodman et al.,

2019), and Dr. Brian McCabe (Wang et al., 2011), respectively. Other transgenic fly

lines were generated in this study. Full genotypes of the fly lines used in all figures and

their cultured temperatures are described in Supplementary file 1.

Generation of constructs and transgenic flies

Artificially synthesized (G$_4$C$_2$)$_{50}$ sequences flanked at the 5′ end with an *Eag*I

recognition site and at the 3′ end with a *PspOM*I recognition site were subcloned into T-

vector pMD20 (Takara Bio). To generate a longer repeat size, the pMD20-(G$_4$C$_2$)$_{50}$

vector was digested with *Eag*I and *PspOM*I, followed by ligation into the pMD20-

(G$_4$C$_2$)$_{50}$ vector linearized by digestion with *Eag*I. This vector was digested with *EcoR*I

and *Hind*III, and subcloned into the pcDNA™3.1/*myc*-His(−)A vector (Thermo Fisher

Scientific). We accidentally obtained the pcDNA™3.1/*myc*-His(−)A-(G$_4$C$_2$)$_9$ vector at

this step. The pcDNA™3.1/*myc*-His(−)A-(G$_4$C$_2$)$_n$ vector was digested with *EcoR*I and

*Xba*I, and subcloned into the *Drosophila* pUAST vector. These constructs have no start

codon sequence (ATG) upstream of the G_4C_2 repeat sequence (Figure 1—figure

supplement 1A). These pUAST-$(G_4C_2)_n$ vectors were amplified with a recombinase-

mutated SURE®2 *Escherichia coli* strain (Agilent Technologies) at 28 °C for 72 hours

to prevent repeat length contraction. The number of G_4C_2 repeats in the pUAST-$(G_4C_2)_9$

$_{or\ 50}$ vectors was determined by sequencing. To determine the number of G_4C_2 repeats in

the pUAST-$(G_4C_2)_{89}$ vector, transposable element insertional mutagenesis using EZ-

Tn5™ <KAN-2> Insertion Kit (Epicentre) and sequencing were performed. The entire

sequence of the insert in the pUAST vector is shown in Figure 1—figure supplement

1—source data 1.

To generate pUAST-*FUS* or pUAST-*TDP-43* vectors, the Gateway® Vector

Conversion System (Thermo Fisher Scientific) was used. The human *FUS* or human

TARDBP cDNA was subcloned into the pENTR™/D-TOPO® vector (Thermo Fisher

Scientific). To generate the Gateway destination vector pUAST-DEST, we inserted the

Gateway cassette A sequence (Thermo Fisher Scientific) into the pUAST vector. The

pUAST-*FUS* or pUAST-*TDP-43* vectors were generated using Gateway recombination

reactions (Thermo Fisher Scientific) . The FUS RRM mutant construct (pUAST-*FUS-*

RRMmut), in which leucine residues at positions 305, 341, 359, and 368 in the FUS

protein were substituted to phenylalanine, was generated by PCR and the In-Fusion®

Cloning system (Takara Bio). To generate the other pUASTattB-*RBP* vectors, each

cDNA encoding the RBP shown in Figure 1—source data 1 and Figure 6—source data 1

was subcloned into the pUASTattB vector (VectorBuilder). To establish transgenic flies

harboring *UAS-$(G_4C_2)_n$*, *UAS-FUS*, *UAS-FUS* line 2, *UAS-FUS-RRMmut*, and *UAS-*

TDP-43, the pUAST-$(G_4C_2)_n$, pUAST-*FUS*, pUAST-*FUS* line 2, pUAST-*FUS-RRMmut*,

and pUAST-*TDP-43* vectors, respectively, were injected into fly embryos of the *w^{1118}*

strain. To establish transgenic flies harboring the other *UAS-RBP* constructs including

UAS-FUS line 3, pUASTattB-*RBP* vectors were injected into fly embryos of the attP40

strain. By employing site-specific transgenesis using the pUASTattB vector, each

transgene was inserted into the same locus of the genome, and was expected to be

expressed at the equivalent levels. These transgenic flies were established using

standard methods at BestGene Inc.

The number of repeats in *UAS*-$(G_4C_2)_{9 \text{ or } 42}$ transgenic flies were determined by

genomic PCR using the forward (5′-AACCAGCAACCAAGTAAATCAAC-3′), and

reverse (5′-TGTTGAGAGTCAGCAGTAGCC-3′) primers, which amplifies a part of

the *UAS*-$(G_4C_2)_n$ sequence, including G_4C_2 repeat sequence, followed by sequencing

using the forward (5′-GCCAAGAAGTAATTATTGA-3′) and/or reverse (5′-

TCCAATTATGTCACACC-3′) primers.

Quantitative real-time PCR

Total RNA was extracted from female third instar larvae of each genotype using

TRIzol® reagent (Thermo Fisher Scientific) according to the manufacturer's

instructions. First-strand cDNA was synthesized using QuantiTect Reverse

Transcription Kit (QIAGEN). Real-time PCR was performed using SYBR® Premix Ex

Taq™ II (Takara Bio) and the Mx3000P Real-time quantitative PCR system (Agilent

Technologies) or the CFX96™ Real-Time PCR Detection System (Bio-Rad). For G_4C_2

repeat RNA quantification of flies expressing $(G_4C_2)_n$ in Figure 1—figure supplement

1C, the forward (5′-ATGAATGGGAGCAGTGGTGG-3′) and reverse (5′-

TGTTGAGAGTCAGCAGTAGCC-3′) primers were used. For G_4C_2 repeat RNA

quantification of flies expressing $(G_4C_2)_{89}$(H) and FUS, FUS-RRMmut, other RNA-

binding proteins, or *caz-IR* in Figures 1E, 3C, 4G, and 6D, the forward (5′-

CCCAATCCATATGACTAGTAGATCC-3′) and reverse (5′-

TGTAGGTAGTTTGTCCAATTATGTCA-3′) primers were used. Both of the above-

mentioned primer pairs recognize sequences downstream of the G_4C_2 repeats. For *gal4*

mRNA quantification, the forward (5′-TTGAAATCGCGTCGAAGGA-3′) and reverse

(5′-GGCTCCAATGGCTAATATGCA-3′) primers were used (Li et al., 2008). Data were

analyzed using the standard curve method. The amounts of G_4C_2 repeat transcripts were

normalized to those of *gal4* transcripts expressed in the same tissue to avoid potential

confounding derived from the difference in tissue viability between genotypes. At least

three independent biological replicates per genotype were analyzed. Data were

normalized by setting the values of the samples from flies expressing $(G_4C_2)_{89}$(H)

(Figure 1—figure supplement 1C), both $(G_4C_2)_{89}$(H) and *EGFP* (Figures 1E, 3C, and

6D), or both $(G_4C_2)_{89}$(H) and *GFP-IR* (Figure 4G) as 100.

Imaging and quantification of fly eyes

Light microscopic images of the eyes of 3 to 5-day-old female flies were taken

using a stereoscopic microscope model SZX10 (Olympus) with a CCD camera DP21

(Olympus). Images shown are representative eye phenotypes of the fly crosses. Crosses

were performed three times to validate the specific phenotypes. Eye size and

pigmentation were quantified as previously reported (Saitoh et al., 2015). Five or 10

eyes per genotype were analyzed. Data were normalized by setting the values of

samples from flies expressing one copy of *EGFP* (Figure 4D), those expressing two

copies of *EGFP* (Figures 1B, 1C, 1D, 2B, 2C, 6B, 6C, Figure 1—figure supplements

2B-2D, and Figure 4—figure supplements 1B-1D), or those expressing both *EGFP* and

GFP-IR (Figure 4B), as 100.

Egg-to-adult viability of flies

Mated female flies were placed on grape juice agar with yeast paste for 24

hours. Eggs were collected from the surface of the grape juice agar, and the number of

eggs were counted and placed on new standard fly food. After eclosion, the number of

adult flies was counted. Egg-to-adult viability was calculated by dividing the number of

adult flies by the number of eggs. More than 500 eggs per genotype were used. Data

were normalized by setting the values of samples from flies expressing two copies of

EGFP as 100 (Fig 2D).

Climbing assay

Twenty male flies were gently introduced into a glass vial. After a 5-minute

adaptation period, the bottom of the vial was gently tapped and the height the flies

reached in 10 seconds was recorded using a digital video camera, and scored as follows:

0 (lower than 2 cm), 1 (from 2 to 3.9 cm), 2 (from 4 to 5.9 cm), 3 (from 6 to 7.9 cm), 4

(from 8 to 9.9 cm), and 5 (higher than 10 cm). Five trials were performed in each

experiment at intervals of 20 sec. The assay was performed between 8:00 and 10:00.

Climbing scores were calculated as an average of five trials.

GeneSwitch experiments

Flies were crossed in the absence of RU486 (mifepristone) on standard fly food.

One-day old adult male flies were transferred to Formula 4-24® Instant Drosophila

medium (Wako) with RU486 (100 µg/mL) for the indicated periods. Every 2 or 3 days,

flies were transferred to new medium with RU486. Climbing assays were performed at

0, 7, and 14 days after the start of RU486 treatment (Figure 2E).

RNA fluorescence in situ hybridization

Female third instar larvae were dissected in ice-cold phosphate-buffered saline

(PBS). Salivary glands were fixed with 4% paraformaldehyde (PFA) (pH 7.0) in PBS

for 30 min, and incubated in 100% methanol. Fixed samples were rehydrated in 75%

(v/v), 50%, and 25% ethanol in PBS, and rinsed in PBS and distilled water (DW).

Samples were then treated with 0.2 N HCl/DW for 20 min at room temperature (RT),

and rinsed in DW. Next, the samples were permeabilized with 0.2% Triton X-100 in

PBS for 10 min, rinsed in PBS for 5 min, fixed again in 4% PFA in PBS for 20 min,

then washed twice for 5 min each in PBS, and incubated twice for 15 min each in 2

mg/mL glycine/PBS. After the acetylation treatment, samples were incubated for 1 hour

at 37 °C in hybridization buffer consisting of 50% formamide, $2 \times$ saline sodium

citrate (SSC), 0.2 mg/mL yeast tRNA, and 0.5 mg/mL heparin. For hybridization,

samples were incubated overnight at 80 °C with a 5′ end Alexa594-labeled $(G_2C_4)_4$ or

Alexa488-labeled $(C_2G_4)_4$ locked nucleic acid (LNA) probe (5 nM) in hybridization

buffer. These LNA probes were synthesized by GeneDesign Inc. After the hybridization,

samples were washed once for 5 min in $4 \times$ SSC at 80 °C, three times for 20 min

each in $2 \times$ SSC and 50% formamide at 80 °C, three times for 40 min each in

$0.1 \times$ SSC at 80 °C, and once for 5 min in PBS containing 0.5% Triton X-100 (PBT)

at RT. Nuclei were stained with 4',6-diamidino-2-phenylindole (DAPI) or 2'-(4-

ethoxyphenyl)-5-(4-methyl-1-piperazinyl)-2,5'-bi-1-H-benzimidazole, trihydrochloride

(Hoechst 33342). Stained samples were mounted in SlowFade® Gold antifade reagent

(Thermo Fisher Scientific) and observed under a Zeiss LSM710 or LSM880 confocal

laser-scanning microscope.

After RNA FISH, samples were scanned using a Zeiss LSM710 or LSM880

confocal laser-scanning microscope along the z-axis direction. One z-stack image was

taken per salivary gland using ZEN imaging software (Zeiss). RNA foci-positive nuclei

in more than 30 cells per salivary gland were counted, and the percentage of nuclei

containing RNA foci in the salivary gland was calculated. Ten salivary glands were

analyzed for each genotype.

Immunohistochemistry

Female third instar larvae were dissected in ice-cold PBS. Eye imaginal discs

and salivary glands were fixed with 4% PFA in PBS for 30 min and washed 3 times

with PBT. After blocking with 5% goat serum/PBT, the samples were incubated

overnight at 4 °C with a rat monoclonal anti-poly(GR) antibody (clone 5A2,

MABN778, Millipore), a mouse monoclonal anti-poly(GA) antibody (clone 5E9,

MABN889, Millipore), a rabbit polyclonal anti-poly(GA) antibody (CAC-TIP-C9-P01,

Cosmo Bio), a rabbit polyclonal anti-poly(GP) antibody (NBP2-25018, Novus

Biologicals), or a rabbit polyclonal anti-FUS antibody (A300-302A, Bethyl

Laboratories) at 1:1,000 dilution as the primary antibody. After washing 3 times with

PBT, the samples were incubated with an Alexa 633-conjugated anti-rat antibody (A-

21094, Thermo Fisher Scientific), or an Alexa 488-conjugated or Alexa 555-conjugated

anti-rabbit antibody (A-11008 or A-21428, respectively, Thermo Fisher Scientific) at

1:500 dilution as the secondary antibody. After washing 3 times with PBT, nuclei were

stained with DAPI or Hoechst 33342. Stained samples were mounted in SlowFade®

Gold antifade reagent and observed using confocal laser-scanning microscopes

(LSM710, LSM880 (Zeiss), and FV3000 (Olympus)).

The number of DPR aggregates in the eye discs was quantitatively measured

using ZEN imaging software (Zeiss) and ImageJ (Schneider et al., 2012), as follows: 1)

selection of photoreceptor neurons within the 13 developing ommatidia in rows 2 and 3

at the posterior end of the eye discs (Saitoh et al., 2015) by DAPI or Hoechst 33342

staining, because these ommatidia are at similar stages of development and can be

easily identified; 2) counting of the number of DPR aggregates with a diameter of larger

than 2 µm in the cytoplasm. Ten to 15 eye discs were analyzed for each genotype.

659 Measurement of poly(GP) protein levels

The heads of 5-day-old female flies expressing both $(G_4C_2)_{89}$(H) and either

EGFP, *FUS*, or *FUS-RRMmut* using the *GMR-Gal4* driver were collected and stored at

–80 °C. Samples were prepared using a previously reported method (Tran et al., 2015).

Poly(GP) levels were measured by a sandwich immunoassay that uses Meso Scale

Discovery electrochemiluminescence detection technology, as described previously (Su

et al., 2014). Data were normalized by setting the values of samples from flies

expressing $(G_4C_2)_{89}$(H) and *EGFP* (Figure 3F) as 100.

668 Western blotting

To assess the expression levels of FUS and FUS-RRMmut (Figure 2—figure

supplement 1), or GR-GFP (Figures 3G–3I), 10 heads of 5-day-old female flies

expressing *FUS* or *FUS-RRMmut*, or both *LDS-(G₄C₂)₄₄*$^{GR-GFP}$ and either *FUS* or *FUS-*

RRMmut using the *GMR-Gal4* driver were homogenized in 100 µL of sodium dodecyl

sulfate (SDS) sample buffer using a pestle, boiled for 5 min, and centrifuged at

10,000 $\times$ g for 3 min at 25 °C. Five µL of each supernatant were run on a 5% to 20%

gradient polyacrylamide gel (Wako) and then transferred onto an Immun-Blot[®]

polyvinylidene fluoride membrane (Bio-Rad). Membranes were blocked with 5% skim

milk in PBS containing 0.1% Tween-20 (PBST) or PVDF Blocking Reagent for Can

Get Signal[®] (TOYOBO) for 2 hours at RT, and then incubated overnight at 4 °C with a

rabbit polyclonal anti-FUS antibody (A300-302A, Bethyl Laboratories), a rat

monoclonal anti-poly(GR) antibody (clone 5A2, MABN778, Millipore), a Living

Colors[®] EGFP mouse monoclonal antibody (632569, Clontech), or a mouse monoclonal

anti-actin antibody (clone AC-40, A4700, Sigma-Aldrich) at 1:1,000 dilution as the

primary antibody. After washing 3 times with PBST, membranes were incubated for 2

hours at RT with either HRP-conjugated anti-rat, anti-rabbit, or anti-mouse antibody

(31470, 31460, or 31430, respectively, Invitrogen) at 1:5,000 dilution as the secondary

antibody, washed 3 times with PBST, treated with SuperSignal™ West Dura

chemiluminescent substrate (Thermo Fisher Scientific), and imaged using the

LuminoGraphII imaging system (ATTO). Data were normalized by setting the average

values of samples from flies expressing *FUS* (Figure 2—figure supplement 1B), or

those expressing both *LDS*-$(G_4C_2)_{44}$[GR-GFP] and *DsRed* as 100 (Figures 3H and 3I).

Filter binding assay

For preparation of FUS proteins with an N-terminal His tag (His-FUS and His-

FUS-RRMmut), cDNAs of the human FUS protein (wild type or RRM mutant) from

pUAST-*FUS* or pUAST-*FUS-RRMmut* were cloned into the multiple cloning site (*Xho*I

and *Bam*HI) of the plasmid vector pET-15b (Novagen) (Nomura et al., 2014). After

transfection of the plasmids into *E. coli* BL21 (DE3), the expression of His-FUS

proteins was induced by culturing the transformed cells in the presence of 0.5 mM

isopropyl β-*D*-thiogalactopyranoside (IPTG) at 20 °C for 20 hours. Cells were lysed by

ultrasonication in PBS (pH 7.4) containing 2% (v/v) Triton X-100, 1 M NaCl, DNase I,

$MgSO_4$, and ethylenediaminetetraacetic acid-free Complete Protease Inhibitor Cocktail

(Roche Applied Science). After centrifugation at 20,000 × g for 30 min at 4 °C, the

pellets were redissolved in a buffer (pH 7.0) containing 6 M guanidine hydrochloride

(GdnHCl), 50 mM Tris, and 1 M NaCl. His-FUS proteins in the pellets were purified by

Ni^{2+} affinity chromatography. In brief, the His-FUS proteins were mixed with Profinity

IMAC Ni^{2+}-charged resin (Bio-Rad) for 30 min at 20 °C. Then, FUS proteins bound to

the resin were washed with wash buffer (6 M GdnHCl, 50 mM Tris, and 1 M NaCl, pH

7.0), and eluted with elution buffer (6 M GdnHCl, 50 mM Tris, 100 mM NaCl, and 250

mM imidazole, pH 7.0). For preparation of soluble, refolded FUS, FUS proteins (200

μM) in the elution buffer were diluted 20-fold with a buffer (pH 7.0) containing 50 mM

Tris, 100 mM NaCl, 10% (v/v) glycerol, and 5 mM Tris (2-carboxyethyl) phosphine

(buffer A), which, however, produced significant amounts of precipitate. This insoluble

material was removed by centrifugation at 20,000 × g for 10 min at 4 °C, resulting in

the recovery of soluble FUS proteins in the supernatant fraction. Protein concentrations

were spectroscopically determined from the absorbance at 280 nm using the following

extinction coefficients: 71,630 cm^{-1} M^{-1} for both FUS and FUS-RRMmut.

Biotinylated RNAs were synthesized by GeneDesign Inc. Ten nM biotin-

$(G_4C_2)_4$, 10 nM biotin-$(AAAAAA)_4$, or 10 nM biotin-$(UUAGGG)_4$ (telomeric repeat-

containing RNA: TERRA) were incubated with soluble FUS proteins (5, 10, or 50 nM)

in buffer A with 0.4 U/µL RNase inhibitor (RNasin Plus RNase Inhibitor, Promega).

Biotinylated (AAAAAA)$_4$ and TERRA are negative and positive controls, respectively.

After an hour at RT, the mixture was filtered through a nitrocellulose membrane

(PROTRAN®, 0.2 µm, Amersham Biosciences) overlaid on a nylon membrane

(Hybond-N$^+$, 0.45 µm, Schleicher & Schuell) in a 96-well slot-blot apparatus (ATTO)

(Furukawa et al., 2011). After extensive washing of the membranes with buffer A, the

bound RNAs were crosslinked to the membranes using ultraviolet radiation (254 nm;

UV Stratalinker, Stratagene) at an energy level of 0.12 J. After blocking with 3% (w/v)

BSA in tris-buffered saline with 0.1% Tween-20, the membranes were incubated with

streptavidin-HRP (1:5,000; Nacalai Tesque), and the biotinylated RNAs on the

membranes were detected with ImmunoStar LD reagent (Wako).

Preparation of recombinant FUS protein

For preparation of the FUS proteins, the human *FUS* (WT) and *FUS-RRMmut*

genes flanked at the 5′ end with an *Nde*I recognition site and at the 3′ end with a *Xho*I

recognition site was amplified by PCR from pUAST-*FUS* and pUAST- *FUS-RRMmut*,

respectively. PCR fragments were digested with *Nde*I and *Xho*I. These fragments were

ligated into the cloning sites of the plasmid vector pET-21b (Novagen) between *Nde*I

and *Xho*I. After transfection of the plasmids into *E. coli* BL21 (DE3), expression of the

FUS protein was induced by culturing the transformed cells in the presence of 0.5 mM

IPTG at 37 °C for 6 hours. Cells were harvested by centrifugation and suspended with

buffer B (10% glycerol, 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

[HEPES]-NaOH [pH 7.0], 300 mM NaCl, 1 mM dithiothreitol [DTT], 1 mM

ethylenediaminetetraacetic acid [EDTA], 0.1% Tween-20, and 0.1% benzamidine

hydrochloride) containing 1.5 mg/mL lysozyme, and stored for 30 min on ice. Cell

lysates were sonicated, and insoluble protein was collected by centrifugation. The pellet

was solubilized in buffer C (6 M urea, 10% glycerol, 20 mM HEPES-NaOH [pH 7.0], 1

mM DTT, 1 mM EDTA, and 0.1% benzamidine hydrochloride). After centrifugation,

supernatants were loaded onto a DE52 (GE Healthcare) open column. The flow-through

fraction was loaded onto a CM52 (GE Healthcare) open column. The flow-through

fraction of DE52 was applied to a CAPTO S column (GE Healthcare), and the flow-

through fraction was collected using the ÄKTAexplorer 10S/100 system (GE

Healthcare). The flow-through fraction was applied to a Mono S column (GE

Healthcare). Proteins were fractionated with a 0 to 500 mM linear gradient of NaCl in

buffer D (6 M urea, 10% glycerol, 20 mM HEPES-NaOH [pH 7.0], 1 mM DTT, and 1

mM EDTA) using ÄKTA explorer 10 S/100 system. The FUS fraction was eluted at 150

to 200 mM NaCl. For refolding, the eluted peak fraction was diluted fivefold using

refolding buffer (900 mM arginine, 100 mM N-cyclohexyl-2-hydroxyl-3-

aminopropanesulfonic acid [pH 9.5], 0.3 mM reduced glutathione, 0.03 mM oxidized

glutathione, and 1 mM $ZnCl_2$), and stored overnight at RT. The solution was

concentrated using a centrifugal filter (Vivaspin 6-10 kDa; GE Healthcare) to 1 to 2

mg/mL, and then dialyzed against buffer E (10% glycerol, 20 mM HEPES-NaOH [pH

6.8], 300 mM NaCl, 0.1 mM EDTA, and 10 mM β-cyclodextrin), and stored frozen at

−80 °C.

Surface plasmon resonance analyses

The binding of FUS to $(G_4C_2)_4$ RNA was analyzed using a Biacore T200

instrument (GE Healthcare). $(G_4C_2)_4$ RNAs biotinylated at the 5′ end in 10 mM HEPES

pH 6.8 and 500 mM MCl (M = K, Na, or Li) was injected over the streptavidin-coated

surface of a sensor chip SA (GE Healthcare). The amount of immobilized RNA was as

follows: 240 resonance unit (RU) in KCl, 363 RU in NaCl, or 319 RU in LiCl buffer

condition. Binding experiments were performed using the single cycle kinetics method.

The running buffer used was 20 mM HEPES (pH 6.8), 1 mM $MgCl_2$, 0.05% Tween-20,

and 150 mM KCl, NaCl, or LiCl. FUS was diluted in the running buffer and injected

sequentially over the RNA-immobilized sensor surface in increasing concentrations

(0.016, 0.031, 0.063, 0.13, or 0.25 µM). Sensorgrams were obtained at 25 °C, 30

µL/min flow rate, 60 seconds of contact time, and 120 seconds of dissociation time.

Circular dichroism spectroscopy

CD spectra were measured at 25 °C using a spectropolarimeter model J-820

(JASCO). $(G_4C_2)_4$ RNA was synthesized by GeneDesign Inc. and dissolved in 20 mM

HEPES (pH 6.8), 18.75 mM NaCl, 10 mM $MgCl_2$, 0.625% glycerol, 0.625 mM β-

cyclodextrin, and 0.0625 mM EDTA with 150 mM KCl, NaCl, or LiCl. RNA samples

containing 150 mM KCl were first heated at 95 °C for 5 min and then cooled to RT to

form the G-quadruplex structure. The other samples were not heated. FUS (1 µM) was

added to the RNA sample (4 µM) and mixed before recording the spectrum. CD spectra

were recorded at a speed of 50 nm min^{-1} and a resolution of 1 nm, and 10 scans were

averaged.

Nuclear magnetic resonance spectroscopy

All one-dimensional ^{1}H NMR spectral data were recorded using AVANCE III

800 MHz NMR spectrometers equipped with a TXI cryogenic probe (Bruker BioSpin)

at 25 °C. $(G_4C_2)_4$ RNA dissolved in 20 mM HEPES (pH 6.8), 150 mM KCl, 18.75 mM

NaCl, 10 mM $MgCl_2$, 0.625% glycerol, 0.625 mM β-cyclodextrin, and 0.0625 mM

EDTA was first heated at 95 °C for 5 min and then cooled to room temperature to form

the G-quadruplex structure. The RNA (10 μM) was mixed with FUS at molar ratios

(RNA:FUS) of 1:0, 1:0.2, 1:0.4, and 0:1. The samples were then prepared at a final

concentration of 10% D_2O before recording their spectra. ^1H NMR data were acquired

using simple single 90° hard-pulse excitations following solvent signal suppression with

a jump-and-return pulse scheme. Free induction decay data (1,600 points in total) were

collected by repeating the scans (2,600 times) with an interscan delay of 2.5 seconds.

All NMR data were processed using Topspin 3.6 software (Bruker BioSpin).

RNA synthesis for in vitro translation

For preparation of the C9-RAN reporter plasmid, the pEF6-*C9orf72* intron1-

$(G_4C_2)_{80}$ vector was digested with *HindIII* and *NotI* to obtain the fragment *C9orf72*

intron1-$(G_4C_2)_{80}$, and subcloned into the pcDNA™5/FRT vector (Thermo Fisher

Scientific). To add the T7 promoter upstream of the *C9orf72* intron 1 sequence in this

pcDNA™5/FRT-*C9orf72* intron1-$(G_4C_2)_{80}$ vector, a forward primer including T7

promoter sequences with the 5′-terminal region of *C9orf72* intron 1 flanked at the 5′ end

with an *HindIII* recognition site, and a reverse primer recognizing the 3′-terminal region

of *C9orf72* intron 1 sequences including a *BssHII* recognition site were designed, and

used to amplify a fragment containing *C9orf72* intron 1 with a T7 promoter by PCR.

Then, this fragment was subcloned into the pcDNA™5/FRT-*C9orf72* intron1-$(G_4C_2)_{80}$

vector digested by *HindIII* and *BssHII*. In addition to the T7 promoter, the Myc tag in

the GA frame downstream of $(G_4C_2)_{80}$ was introduced into this vector.

The reporter plasmids were linearized with *XbaI*. Linearized DNA was *in vitro*

transcribed using mMESSAGE mMACHINE™ T7 Transcription Kit (Invitrogen)

according to the manufacturer's instructions. T7 reactions were carried out at 37 °C for

2 hours, treated with TURBO™ DNaseI for 15 min at 37 °C to remove the DNA

template, and then polyadenylated with *E. coli* Poly-A Polymerase (NEB) for 1 hour at

37 °C. Synthesized mRNAs were purified by LiCl precipitation. The size and quality of

the synthesized mRNAs were verified on a denaturing RNA gel.

In vitro translation assay

mRNAs of *C9orf72* intron1-$(G_4C_2)_{80}$ with a Myc tag in the GA frame were *in*

vitro translated with Flexi® Rabbit Reticulocyte Lysate System (Promega) according to

the manufacturer's instructions. Translation reactions were performed with 10 ng/µL

mRNA and contained 30% rabbit reticulocyte lysate, 10 µM amino-acid mix minus

methionine, 10 µM amino-acid mix minus leucine, 0.5 mM MgOAc, 100 mM KCl, and

0.8 U/µL Murine RNAse Inhibitor (NEB). FUS or FUS-RRMmut at each concentration

(10, 100, 200, 400, and 1,000 nM) was preincubated with mRNA for 10 min to facilitate

the interaction between FUS protein and G_4C_2 repeat RNA, and added for translation in

the lysate. Samples were incubated at 30 °C for 90 min before termination by

incubation on ice. Ten µL of samples were analyzed by 13% SDS-polyacrylamide gel

electrophoresis and Western blotting using a mouse monoclonal anti-c-Myc antibody

(clone 9E10, Wako) as the primary antibody.

Quantification and statistical analysis

Statistical parameters including the definitions and exact values of n (e.g.,

number of experiments, number of flies, number of eye imaginal discs, etc.),

distributions, and deviations are stated in the figures and corresponding figure legends.

Multiple comparison tests using one-way ANOVA with Tukey's post hoc analysis were

performed for Figures 1B–1E, 2B–2D, 3B, 3C, 3E, 3F, 3H, 3I, 4B, 4D, 6B, 6C, 6D, 6F,

Figure 1—figure supplements 1C and 1D, Figure 1—figure supplements 2B–2D, and

Figure 4—figure supplements 1B–1D, multiple comparison test using two-way repeated

measures ANOVA with Tukey's post hoc analysis was performed for Figure 2E, and the

unpaired t-test was used for Figures 4F, 4G, 4I, 5G, and Figure 2—figure supplement

1B. Differences in means were considered statistically significant at $P < 0.05$. All

statistical analyses were performed using GraphPad Prism version 8.3.4 (GraphPad

Software, LLC).

As the sample sizes used in the present study were similar to previous

publications (Freibaum et al., 2015; Goodman et al., 2019; Mizielinska et al., 2014; Xu

et al., 2013), statistical analyses were performed afterwards without interim data

analysis. Data were not excluded, and were collected and processed randomly. Sample

collection and analyses for the measurement of poly(GP) protein levels were performed

in a double-blind manner. Data collection and analyses for other experiments were not

performed in a blind manner regarding the conditions of the experiments.

Acknowledgements

We thank Drs E. Storkebaum (Radboud University), N. Bonini (University of Pennsylvania), and B. McCabe (École Polytechnique Fédérale de Lausanne) for kindly providing the *caz* mutant, *UAS-LDS-(G4C2)44*^{GR-GFP}, and *UAS-FLAG-caz* and *UAS-FLAG-FUS* flies, respectively. We acknowledge Bloomington Drosophila Stock Center and Vienna Drosophila Resource Center for providing various fly stocks. We thank Drs H. Imataka and K. Machida (University of Hyogo) for kindly providing human tRNAs for *in vitro* translation. We also thank the members of our laboratory for helpful discussions, K. Yamamoto, E. Wakisaka, T. Yamashita, and A. Sugai for their technical assistance, Dr. H. Akiko Popiel for critical reading and English editing of the manuscript, and the Center for Medical Research and Education, Graduate School of Medicine, Osaka University for supplying experimental equipment.

References

Ash, P. E., Bieniek, K. F., Gendron, T. F., Caulfield, T., Lin, W. L., Dejesus-

Hernandez, M., van Blitterswijk, M. M., Jansen-West, K., Paul, J. W., 3rd, Rademakers,

R., Boylan, K. B., Dickson, D. W., and Petrucelli, L. (2013). Unconventional translation

of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to

c9FTD/ALS. Neuron *77*, 639–646. https://doi.org/10.1016/j.neuron.2013.02.004

Bajc Česnik, A., Darovic, S., Prpar Mihevc, S., Štalekar, M., Malnar, M.,

Motaln, H., Lee, Y. B., Mazej, J., Pohleven, J., Grosch, M., Modic, M., Fonovič, M.,

Turk, B., Drukker, M., Shaw, C. E., and Rogelj, B. (2019). Nuclear RNA foci from

C9ORF72 expansion mutation form paraspeckle-like bodies. J Cell Sci. *132*, jcs224303.

https://doi.org/10.1242/jcs.224303

Boivin, M., Pfister, V., Gaucherot, A., Ruffenach, F., Negroni, L., Sellier, C., and

Charlet-Berguerand, N. (2020). Reduced autophagy upon C9ORF72 loss synergizes

with dipeptide repeat protein toxicity in G4C2 repeat expansion disorders. EMBO J. *39*,

e100574. https://doi.org/10.15252/embj.2018100574

Celona, B., Dollen, J. V., Vatsavayai, S. C., Kashima, R., Johnson, J. R., Tang,

A. A., Hata, A., Miller, B. L., Huang, E. J., Krogan, N. J., Seeley, W. W., and Black, B.

L. (2017). Suppression of *C9orf72* RNA repeat-induced neurotoxicity by the ALS-

associated   RNA-binding   protein   Zfp106.   eLife   *6*,   e19032.

https://doi.org/10.7554/eLife.19032

Cheng, W., Wang, S., Mestre, A. A., Fu, C., Makarem, A., Xian, F., Hayes, L. R.,

Lopez-Gonzalez, R., Drenner, K., Jiang, J., Cleveland, D. W., and Sun, S. (2018).

C9ORF72 GGGGCC repeat-associated non-AUG translation is upregulated by stress

through eIF2α phosphorylation. Nat. Commun. 9, 51. https://doi.org/10.1038/s41467-

017-02495-z

Cheng, W., Wang, S., Zhang, Z., Morgens, D. W., Hayes, L. R., Lee, S., Portz,

B., Xie, Y., Nguyen, B. V., Haney, M. S., Yan, S., Dong, D., Coyne, A. N., Yang, J.,

Xian, F., Cleveland, D. W., Qiu, Z., Rothstein, J. D., Shorter, J., Gao, F. B., Bassik,

M.C., and Sun, S. (2019). CRISPR-Cas9 Screens Identify the RNA Helicase DDX3X as

a Repressor of C9ORF72 (GGGGCC)n Repeat-Associated Non-AUG Translation.

Neuron *104*, 885–898.e8. https://doi.org/10.1016/j.neuron.2019.09.003

Choi, S. Y., Lopez-Gonzalez, R., Krishnan, G., Phillips, H. L., Li, A. N., Seeley,

W. W., Yao, W. D., Almeida, S., and Gao, F. B. (2019). C9ORF72-ALS/FTD-associated

poly(GR) binds Atp5a1 and compromises mitochondrial function in vivo. Nat.

Neurosci. *22*, 851–862. https://doi.org/10.1038/s41593-019-0397-0

Conlon, E. G., Lu, L., Sharma, A., Yamazaki, T., Tang, T., Shneider, N. A., and

Manley, J. L. (2016). The C9ORF72 GGGGCC expansion forms RNA G-quadruplex

inclusions and sequesters hnRNP H to disrupt splicing in ALS brains. eLife *5*, e17820.

https://doi.org/10.7554/eLife.17820

Cooper-Knock, J., Walsh, M. J., Higginbottom, A., Robin Highley, J., Dickman,

M. J., Edbauer, D., Ince, P. G., Wharton, S. B., Wilson, S. A., Kirby, J., Hautbergue, G.

M., and Shaw, P. J. (2014). Sequestration of multiple RNA recognition motif-containing

proteins by C9orf72 repeat expansions. Brain *137*, 2040–2051.

https://doi.org/10.1093/brain/awu120

Daigle, J. G., Lanson, N. A., Jr, Smith, R. B., Casci, I., Maltare, A., Monaghan,

J., Nichols, C. D., Kryndushkin, D., Shewmaker, F., and Pandey, U. B. (2013). RNA-

binding ability of FUS regulates neurodegeneration, cytoplasmic mislocalization and

incorporation into stress granules associated with FUS carrying ALS-linked mutations.

Hum. Mol. Genet. *22*, 1193–1205. https://doi.org/10.1093/hmg/dds526

DeJesus-Hernandez, M., Mackenzie, I. R., Boeve, B. F., Boxer, A. L., Baker, M.,

Rutherford, N. J., Nicholson, A. M., Finch, N. A., Flynn, H., Adamson, J., Kouri, N.,

Wojtas, A., Sengdy, P., Hsiung, G. Y., Karydas, A., Seeley, W. W., Josephs, K. A.,

Coppola, G., Geschwind, D. H., Wszolek, Z. K., Feldman, H., Knopman, D.S.,

Petersen, R.C., Miller, B. L., Dickson, D. W., Boylan, K. B., Graff-Radford, N. R., and

Rademakers, R. (2011). Expanded GGGGCC hexanucleotide repeat in noncoding

region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron*, *72*(2), 245–

256. https://doi.org/10.1016/j.neuron.2011.09.011 (2011). Expanded GGGGCC

hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked

FTD and ALS. Neuron *72*, 245–256. https://doi.org/10.1016/j.neuron.2011.09.011

Donnelly, C. J., Zhang, P. W., Pham, J. T., Haeusler, A. R., Mistry, N. A.,

Vidensky, S., Daley, E. L., Poth, E. M., Hoover, B., Fines, D. M., Maragakis, N.,

Tienari, P. J., Petrucelli, L., Traynor, B. J., Wang, J., Rigo, F., Bennett, C. F., Blackshaw,

S., Sattler, R., and Rothstein, J. D. (2013). RNA toxicity from the ALS/FTD C9ORF72

expansion is mitigated by antisense intervention. Neuron *80*, 415–428.

https://doi.org/10.1016/j.neuron.2013.10.015

Dumas, L., Herviou, P., Dassi, E., Cammas, A., and Millevoi, S. (2021). G-

Quadruplexes in RNA Biology: Recent Advances and Future Directions. Trends

Biochem. Sci. *46*, 270–283. https://doi.org/10.1016/j.tibs.2020.11.001

Fratta, P., Mizielinska, S., Nicoll, A. J., Zloh, M., Fisher, E. M., Parkinson, G.,

and Isaacs, A. M. (2012). C9orf72 hexanucleotide repeat associated with amyotrophic

lateral sclerosis and frontotemporal dementia forms RNA G-quadruplexes. Sci. Rep. *2*,

1016. https://doi.org/10.1038/srep01016

Freibaum, B. D., Lu, Y., Lopez-Gonzalez, R., Kim, N. C., Almeida, S., Lee, K.

H., Badders, N., Valentine, M., Miller, B. L., Wong, P. C., Petrucelli, L., Kim, H. J.,

Gao, F. B., and Taylor, J. P. (2015). GGGGCC repeat expansion in C9orf72

compromises nucleocytoplasmic transport. Nature *525*, 129–133.

https://doi.org/10.1038/nature14974

Frickenhaus, M., Wagner, M., Mallik, M., Catinozzi, M., and Storkebaum, E.

(2015). Highly efficient cell-type-specific gene inactivation reveals a key function for

the Drosophila FUS homolog cabeza in neurons. Sci. Rep. *5*, 9107.

https://doi.org/10.1038/srep09107

Furukawa, Y., Kaneko, K., Watanabe, S., Yamanaka, K., and Nukina, N. (2011).

A seeding reaction recapitulates intracellular formation of Sarkosyl-insoluble

transactivation response element (TAR) DNA-binding protein-43 inclusions. J. Biol.

Chem. *286*, 18664–18672. https://doi.org/10.1074/jbc.M111.231209

Geissler, R., Simkin, A., Floss, D., Patel, R., Fogarty, E. A., Scheller, J., and

Grimson, A. (2016). A widespread sequence-specific mRNA decay pathway mediated

by hnRNPs A1 and A2/B1. Genes Dev. *30*, 1070–1085.

https://doi.org/10.1101/gad.277392.116

Gendron, T. F., Bieniek, K. F., Zhang, Y. J., Jansen-West, K., Ash, P. E.,

Caulfield, T., Daughrity, L., Dunmore, J. H., Castanedes-Casey, M., Chew, J., et al.

(2013). Antisense transcripts of the expanded C9ORF72 hexanucleotide repeat form

nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS.

Acta Neuropathol. *126*, 829–844. https://doi.org/10.1007/s00401-013-1192-8

Gendron, T. F., Bieniek, K. F., Zhang, Y. J., Jansen-West, K., Ash, P. E.,

Caulfield, T., Daughrity, L., Dunmore, J. H., Castanedes-Casey, M., Chew, J., Cosio, D.

M., van Blitterswijk, M., Lee, W. C., Rademakers, R., Boylan, K. B., Dickson, D. W.,

and Petrucelli, L. (2017). Poly(GP) proteins are a useful pharmacodynamic marker for

C9ORF72-associated amyotrophic lateral sclerosis. Sci, Transl. Med. *9*, eaai7866.

https://doi.org/10.1126/scitranslmed.aai7866

Gijselinck, I., Van Langenhove, T., van der Zee, J., Sleegers, K., Philtjens, S.,

Kleinberger, G., Janssens, J., Bettens, K., Van Cauwenberghe, C., Pereson, S.,

Engelborghs, S., Sieben, A., De Jonghe, P., Vandenberghe, R., Santens, P., De Bleecker,

J., Maes, G., Bäumer, V., Dillen, L., Joris, G., Cuijt, I., Corsmit, E., Elinck E., Van

Dongen, J., Vermeulen, S., Van den Broeck, M., Vaerenberg, C., Mattheijssens, M.,

Peeters, K., Robberecht, W., Cras, P., Martin, J. J., De Deyn P. P., Cruts, M., and Van

Broeckhoven, C. (2012). A C9orf72 promoter repeat expansion in a Flanders-Belgian

cohort with disorders of the frontotemporal lobar degeneration-amyotrophic lateral

sclerosis spectrum: a gene identification study. Lancet Neurol. *11*, 54–65.

https://doi.org/10.1016/S1474-4422(11)70261-7

1002

1003   Goodman, L. D., Prudencio, M., Kramer, N. J., Martinez-Ramirez, L. F.,

1004   Srinivasan, A. R., Lan, M., Parisi, M. J., Zhu, Y., Chew, J., Cook, C. N., Berson, A.,

1005   Gitler, A. D., Petrucelli, L., and Bonini, N. M. (2019). Toxic expanded GGGGCC repeat

1006   transcription is mediated by the PAF1 complex in C9orf72-associated FTD. Nat.

1007   Neurosci. *22*, 863–874. https://doi.org/10.1038/s41593-019-0396-1

1008

1009   Green, K. M., Glineburg, M. R., Kearse, M. G., Flores, B. N., Linsalata, A. E.,

1010   Fedak, S. J., Goldstrohm, A. C., Barmada, S. J., and Todd, P. K. (2017). RAN

1011   translation at C9orf72-associated repeat expansions is selectively enhanced by the

1012   integrated stress response. Nat. Commun. *8*, 2005. https://doi.org/10.1038/s41467-017-

1013   02200-0

1014

1015   Haeusler, A. R., Donnelly, C. J., Periz, G., Simko, E. A., Shaw, P. G., Kim, M.

1016   S., Maragakis, N. J., Troncoso, J. C., Pandey, A., Sattler, R., Rothstein, J. D., and Wang,

1017   J. (2014). C9orf72 nucleotide repeat structures initiate molecular cascades of disease.

1018   Nature *507*, 195–200. https://doi.org/10.1038/nature13124

1019

1020   Hardin, C. C., Watson, T., Corregan, M., and Bailey, C. (1992). Cation-

1021   dependent transition between the quadruplex and Watson-Crick hairpin forms of

1022   d(CGCG3GCG). Biochemistry *31*, 833–841. https://doi.org/10.1021/bi00118a028

1023

1024   Hubstenberger, A., Courel, M., Bénard, M., Souquere, S., Ernoult-Lange, M.,

1025   Chouaib, R., Yi, Z., Morlot, J. B., Munier, A., Fradet, M., Daunesse, M., Bertrand, E.,

1026   Pierron, G., Mozziconacci, J., Kress, M., and Weil, D. (2017). P-Body Purification

1027   Reveals the Condensation of Repressed mRNA Regulons. Mol Cell. *68*, 144–157.e5.

1028   https://doi.org/10.1016/j.molcel.2017.09.003

1029

1030   Hutten, S., Usluer, S., Bourgeois, B., Simonetti, F., Odeh, H. M., Fare, C. M.,

1031   Czuppa, M., Hruska-Plochan, M., Hofweber, M., Polymenidou, M., Shorter, J.,

1032   Edbauer, D., Madl, T., and Dormann, D. (2020). Nuclear Import Receptors Directly

1033   Bind to Arginine-Rich Dipeptide Repeat Proteins and Suppress Their Pathological

1034   Interactions. Cell Rep. *33*, 108538. https://doi.org/10.1016/j.celrep.2020.108538

1035

1036   Ishiguro, T., Sato, N., Ueyama, M., Fujikake, N., Sellier, C., Kanegami, A.,

1037   Tokuda, E., Zamiri, B., Gall-Duncan, T., Mirceta, M., Furukawa, Y., Yokota, T., Wada,

1038   K., Taylor, J. P., Pearson, C. E., Charlet-Berguerand, N., Mizusawa, H., Nagai, Y., and

1039   Ishikawa, K. (2017). Regulatory Role of RNA Chaperone TDP-43 for RNA Misfolding

1040   and   Repeat-Associated   Translation   in   SCA31.   Neuron   *94*,   108–124.e7.

1041   https://doi.org/10.1016/j.neuron.2017.02.046

1042

1043   Jovičić, A., Mertens, J., Boeynaems, S., Bogaert, E., Chai, N., Yamada, S. B.,

1044   Paul, J. W., 3rd, Sun, S., Herdy, J. R., Bieri, G., Kramer, N. J., Gage, F. H., Van Den

1045   Bosch, L., Robberecht, W., and Gitler, A. D. (2015). Modifiers of C9orf72 dipeptide

1046   repeat toxicity connect nucleocytoplasmic transport defects to FTD/ALS. Nat. Neurosci.

1047   *18*, 1226–1229. https://doi.org/10.1038/nn.4085

1048

Krishnan, G., Raitcheva, D., Bartlett, D., Prudencio, M., McKenna-Yasek, D.

M., Douthwright, C., Oskarsson, B. E., Ladha, S., King, O. D., Barmada, S. J., Miller, T.

M., Bowser, R., Watts, J. K., Petrucelli, L., Brown, R. H., Kankel, M. W., and Gao, F. B.

(2022). Poly(GR) and poly(GA) in cerebrospinal fluid as potential biomarkers for

C9ORF72-ALS/FTD. Nat. Commun. *13*, 2799. https://doi.org/10.1038/s41467-022-

30387-4

1055

1056   Kanadia, R. N., Shin, J., Yuan, Y., Beattie, S. G., Wheeler, T. M., Thornton, C.

1057   A., and Swanson, M. S. (2006). Reversal of RNA missplicing and myotonia after

1058   muscleblind overexpression in a mouse poly(CUG) model for myotonic dystrophy.

1059   Proc.   Natl.   Acad.   Sci.   USA   *103*,   11748–11753.

1060   https://doi.org/10.1073/pnas.0604970103

1061

1062   Katahira, J., Miki, T., Takano, K., Maruhashi, M., Uchikawa, M., Tachibana, T.,

1063   and Yoneda, Y. (2008). Nuclear RNA export factor 7 is localized in processing bodies

and neuronal RNA granules through interactions with shuttling hnRNPs. Nucleic Acids Res. *36*, 616–628. https://doi.org/10.1093/nar/gkm556

Kwan, T., and Thompson, S. R. (2019). Noncanonical Translation Initiation in Eukaryotes. Cold Spring Harb. Perspect. Biol. *11*, a032672. https://doi.org/10.1101/cshperspect.a032672

Kwon, I., Xiang, S., Kato, M., Wu, L., Theodoropoulos, P., Wang, T., Kim, J., Yun, J., Xie, Y., and McKnight, S. L. (2014). Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede RNA biogenesis, and kill cells. Science *345*, 1139–1145. https://doi.org/10.1126/science.1254917

Lagier-Tourenne, C., Polymenidou, M., and Cleveland, D. W. (2010). TDP-43 and FUS/TLS: emerging roles in RNA processing and neurodegeneration. Hum. Mol. Genet. *19*, R46–R64. https://doi.org/10.1093/hmg/ddq137

Lee, K. H., Zhang, P., Kim, H. J., Mitrea, D. M., Sarkar, M., Freibaum, B. D., Cika, J., Coughlin, M., Messing, J., Molliex, A., Maxwell, B. A., Kim, N. C., Temirov, J., Moore, J., Kolaitis, R. M., Shaw, T. I., Bai, B., Peng, J., Kriwacki, R. W., and Taylor, J. P. (2016). C9orf72 Dipeptide Repeats Impair the Assembly, Dynamics, and Function of Membrane-Less Organelles. Cell *167*, 774–788.e17. https://doi.org/10.1016/j.cell.2016.10.002

Lee, Y. B., Chen, H. J., Peres, J. N., Gomez-Deza, J., Attig, J., Stalekar, M.,

1088   Troakes, C., Nishimura, A. L., Scotter, E. L., Vance, C., Adachi, Y., Sardone, V., Miller,

1089   J. W., Smith, B. N., Gallo, J. M., Ule, J., Hirth, F., Rogelj, B., Houart, C., and Shaw, C.

1090   E. (2013). Hexanucleotide repeats in ALS/FTD form length-dependent RNA foci,

1091   sequester RNA binding proteins, and are neurotoxic. Cell Rep. *5*, 1178–1186.

1092   https://doi.org/10.1016/j.celrep.2013.10.049

1093

Lehmer, C., Oeckl, P., Weishaupt, J. H., Volk, A. E., Diehl-Schmid, J., Schroeter,

M. L., Lauer, M., Kornhuber, J., Levin, J., Fassbender, K., Landwehrmeyer, B., Schludi,

M. H., Arzberger, T., Kremmer, E., Flatley, A., Feederle, R., Steinacker, P., Weydt, P.,

Ludolph, A. C., Edbauer, D., and Otto, M. German Consortium for Frontotemporal

Lobar Degeneration (2017). Poly-GP in cerebrospinal fluid links C9orf72-associated

dipeptide repeat expression to the asymptomatic phase of ALS/FTD. EMBO Mol. Med.

9, 859–868. https://doi.org/10.15252/emmm.201607486

1101

1102   Li, L. B., Yu, Z., Teng, X., and Bonini, N. M. (2008). RNA toxicity is a

1103   component of ataxin-3 degeneration in Drosophila. Nature *453*, 1107–1111.

1104   https://doi.org/10.1038/nature06909

1105

1106   Lin, Y., Mori, E., Kato, M., Xiang, S., Wu, L., Kwon, I., and McKnight, S. L.

1107   (2016). Toxic PR Poly-Dipeptides Encoded by the C9orf72 Repeat Expansion Target

1108   LC Domain Polymers. Cell *167*, 789–802.e12.

1109   https://doi.org/10.1016/j.cell.2016.10.003

1110

1111   Liu, H., Lu, Y. N., Paul, T., Periz, G., Banco, M. T., Ferré-D'Amaré, A. R.,

1112   Rothstein, J. D., Hayes, L. R., Myong, S., and Wang, J. (2021). A Helicase Unwinds

1113   Hexanucleotide Repeat RNA G-Quadruplexes and Facilitates Repeat-Associated Non-

1114   AUG   Translation.   J.   Am.   Chem.   Soc.   *143*,   7368–7379.

1115   https://doi.org/10.1021/jacs.1c00131

1116

1117   Maharana, S., Wang, J., Papadopoulos, D. K., Richter, D., Pozniakovsky, A.,

1118   Poser, I., Bickle, M., Rizk, S., Guillén-Boixet, J., Franzmann, T. M., Jahnel, M.,

1119   Marrone, L., Chang, Y. T., Sterneckert, J., Tomancak, P., Hyman, A. A., and Alberti, S.

1120   (2018). RNA buffers the phase separation behavior of prion-like RNA binding proteins.

1121   Science *360*, 918–921. https://doi.org/10.1126/science.aar7366

1122

1123   Malik, I., Kelley, C. P., Wang, E. T., and Todd, P. K. (2021). Molecular

1124   mechanisms underlying nucleotide repeat expansion disorders. Nat. Rev. Mol. Cell Biol.

1125   *22*, 589–607. https://doi.org/10.1038/s41580-021-00382-6

1126

1127   Mann, J. R., Gleixner, A. M., Mauna, J. C., Gomes, E., DeChellis-Marks, M. R.,

1128   Needham, P. G., Copley, K. E., Hurtle, B., Portz, B., Pyles, N. J., Guo, L., Calder, C. B.,

1129   Wills, Z. P., Pandey, U. B., Kofler, J. K., Brodsky, J. L., Thathiah, A., Shorter, J., and

1130   Donnelly, C. J. (2019). RNA Binding Antagonizes Neurotoxic Phase Transitions of

1131   TDP-43. Neuron *102*, 321–338.e8. https://doi.org/10.1016/j.neuron.2019.01.048

1132

1133   May, S., Hornburg, D., Schludi, M. H., Arzberger, T., Rentzsch, K., Schwenk, B.

1134   M., Grässer, F. A., Mori, K., Kremmer, E., Banzhaf-Strathmann, J., Mann, M.,

1135   Meissner, F., and Edbauer, D. (2014). C9orf72 FTLD/ALS-associated Gly-Ala

1136   dipeptide repeat proteins cause neuronal toxicity and Unc119 sequestration. Acta

1137   Neuropathol. *128*, 485–503. https://doi.org/10.1007/s00401-014-1329-4

1138

1139   Mitra, I., Huang, B., Mousavi, N., Ma, N., Lamkin, M., Yanicky, R., Shleizer-

1140   Burko, S., Lohmueller, K. E., and Gymrek, M. (2021). Patterns of de novo tandem

1141   repeat mutations and their role in autism. Nature *589*, 246–250.

1142   https://doi.org/10.1038/s41586-020-03078-7

1143

Mizielinska, S., Grönke, S., Niccoli, T., Ridler, C. E., Clayton, E. L., Devoy, A.,

Moens, T., Norona, F. E., Woollacott, I., Pietrzyk, J., Cleverley, K., Nicoll, A. J.,

Pickering-Brown, S., Dols, J., Cabecinha, M., Hendrich, O., Fratta, P., Fisher, E.,

Partridge, L., and Isaacs, A. M. (2014). C9orf72 repeat expansions cause

neurodegeneration in Drosophila through arginine-rich proteins. Science *345*, 1192–

1194. https://doi.org/10.1126/science.1256800

1150

Mori, K., Lammich, S., Mackenzie, I. R., Forné, I., Zilow, S., Kretzschmar, H.,

Edbauer, D., Janssens, J., Kleinberger, G., Cruts, M., Herms, J., Neumann, M., Van

Broeckhoven, C., Arzberger, T., and Haass, C. (2013). hnRNP A3 binds to GGGGCC

repeats and is a constituent of p62-positive/TDP43-negative inclusions in the

hippocampus of patients with C9orf72 mutations. Acta Neuropathol. *125*, 413–423.

https://doi.org/10.1007/s00401-013-1088-7. ("Mori et al., 2013a" in the main text)

1157

Mori, K., Arzberger, T., Grässer, F. A., Gijselinck, I., May, S., Rentzsch, K.,

Weng, S. M., Schludi, M. H., van der Zee, J., Cruts, M., Van Broeckhoven, C.,

Kremmer, E., Kretzschmar, H. A., Haass, C., and Edbauer, D. (2013). Bidirectional

transcripts of the expanded C9orf72 hexanucleotide repeat are translated into

aggregating dipeptide repeat proteins. Acta Neuropathol. *126*, 881–893.

https://doi.org/10.1007/s00401-013-1189-3. ("Mori et al., 2013b" in the main text)

1164

1165   Mori, K., Weng, S. M., Arzberger, T., May, S., Rentzsch, K., Kremmer, E.,

1166   Schmid, B., Kretzschmar, H. A., Cruts, M., Van Broeckhoven, C., Haass, C., and

1167   Edbauer, D. (2013). The C9orf72 GGGGCC repeat is translated into aggregating

1168   dipeptide-repeat proteins in FTLD/ALS. Science *339*, 1335–1338.

1169   https://doi.org/10.1126/science.1232927. ("Mori et al., 2013c" in the main text)

1170

1171   Mori, K., Gotoh, S., Yamashita, T., Uozumi, R., Kawabe, Y., Tagami, S., Kamp,

1172   F., Nuscher, B., Edbauer, D., Haass, C., Nagai, Y., and Ikeda, M. (2021). The porphyrin

1173   TMPyP4 inhibits elongation during the noncanonical translation of the FTLD/ALS-

1174   associated GGGGCC repeat in the C9orf72 gene. J. Biol. Chem. *297*, 101120.

1175   https://doi.org/10.1016/j.jbc.2021.101120

1176

1177   Nomura, T., Watanabe, S., Kaneko, K., Yamanaka, K., Nukina, N., and

1178   Furukawa, Y. (2014). Intranuclear aggregation of mutant FUS/TLS as a molecular

1179   pathomechanism of amyotrophic lateral sclerosis. J. Biol. Chem. *289*, 1192–1202.

1180   https://doi.org/10.1074/jbc.M113.516492

1181

1182   Ozdilek, B. A., Thompson, V. F., Ahmed, N. S., White, C. I., Batey, R. T., and

1183   Schwartz, J. C. (2017). Intrinsically disordered RGG/RG domains mediate degenerate

1184   specificity in RNA binding. Nucleic Acids Res. *45*, 7984–7996.

1185   https://doi.org/10.1093/nar/gkx460

1186

1187   Rajkowitsch, L., Chen, D., Stampfl, S., Semrad, K., Waldsich, C., Mayer, O.,

1188   Jantsch, M. F., Konrat, R., Bläsi, U., and Schroeder, R. (2007). RNA chaperones, RNA

1189   annealers and RNA helicases. RNA Biol. *4*, 118–130.

1190   https://doi.org/10.4161/rna.4.3.5445

1191

1192   Reddy, K., Zamiri, B., Stanley, S., Macgregor, R. B., Jr, and Pearson, C. E.

1193   (2013). The disease-associated r(GGGGCC)n repeat from the C9orf72 gene forms tract

1194   length-dependent uni- and multimolecular RNA G-quadruplex structures. J. Biol. Chem.

1195   *288*, 9860–9866. https://doi.org/10.1074/jbc.C113.452532

1196

Renton, A.E., Majounie, E., Waite, A., Simón-Sánchez, J., Rollinson, S., Gibbs,

J. R., Schymick, J. C., Laaksovirta, H., van Swieten, J. C., Myllykangas, L., Kalimo, H.,

Paetau, A., Abramzon, Y., Remes, A. M., Kaganovich, A., Scholz, S. W., Duckworth, J.,

Ding, J., Harmer, D. W., Hernandez, D. G., Johnson. J. O., Mok, K., Ryten, M.,

Trabzuni, D., Guerreiro, R. J., Orrell, R. W., Neal, J., Murray, A., Pearson, J., Jansen, I.

E., Sondervan, D., Seelaar, H., Blake, D., Young, K., Halliwell, N., Callister, J. B.,

Toulson, G., Richardson, A., Gerhard, A., Snowden, J., Mann, D., Neary, D., Nalls, M.

A., Peuralinna, T., Jansson, L., Isoviita, V. M., Kaivorinne, A. L., Höltta-Vuori, M.,

Ikonen, E., Sulkava, R., Benatar, M., Wuu, J., Chiò, A., Restagno, G., Borghero, G.,

Sabatelli, M., Heckerman, D., Rogaeva, E., Zinman, L., Rothstein, J. D., Sendtner, M.,

Drepper, C., Eichler, E. E., Alkan, C., Abdullaev, Z., Pack, S. D., Dutra, A., Pak, E.,

Hardy, J., Singleton, A., Williams, N.M., Heutink, P., Pickering-Brown, S., Morris, H.

R., Tienari, P. J., and Traynor, B. J.. ITALSGEN Consortium (2011). A hexanucleotide

repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD.

Neuron *72*, 257–268. https://doi.org/10.1016/j.neuron.2011.09.010

1212

1213   Rudich, P., Snoznik, C., Watkins, S. C., Monaghan, J., Pandey, U. B., and

1214 Lamitina, S. T. (2017). Nuclear localized C9orf72-associated arginine-containing

1215 dipeptides exhibit age-dependent toxicity in C. elegans. Hum. Mol. Genet. *26*, 4916–

1216 4928. https://doi.org/10.1093/hmg/ddx372

1217

1218   Saitoh, Y., Fujikake, N., Okamoto, Y., Popiel, H. A., Hatanaka, Y., Ueyama, M.,

1219 Suzuki, M., Gaumer, S., Murata, M., Wada, K., and Nagai, Y. (2015). p62 plays a

1220 protective role in the autophagic degradation of polyglutamine protein oligomers in

1221 polyglutamine disease model flies. J. Biol. Chem. *290*, 1442–1453.

1222 https://doi.org/10.1074/jbc.M114.590281

1223

1224   Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to

1225 ImageJ: 25 years of image analysis. Nat. Methods. *9*, 671–675.

1226 https://doi.org/10.1038/nmeth.2089

1227

Simone, R., Balendra, R., Moens, T. G., Preza, E., Wilson, K. M., Heslegrave,

A., Woodling, N. S., Niccoli, T., Gilbert-Jaramillo, J., Abdelkarim, S., Clayton, E. L.,

Clarke, M., Konrad, M. T., Nicoll, A. J., Mitchell, J. S., Calvo, A., Chio, A., Houlden,

H., Polke, J. M., Ismail, M. A., Stephens, C.E., Vo, T., Farahat, A. A., Wilson, W. D.,

Boykin, D. W., Zetterberg, H., Partridge, L., Wray, S., Parkinson, G., Neidle, S., Patani,

R., Fratta, P., and Isaacs, A. M. (2018). G-quadruplex-binding small molecules

ameliorate C9orf72 FTD/ALS pathology in vitro and in vivo. EMBO Mol. Med. *10*,

22–31. https://doi.org/10.15252/emmm.201707850

1236

Shi, Y., Lin, S., Staats, K. A., Li, Y., Chang, W. H., Hung, S. T., Hendricks, E.,

Linares, G. R., Wang, Y., Son, E. Y., Wen, X., Kisler, K., Wilkinson, B., Menendez, L.,

Sugawara, T., Woolwine, P., Huang, M., Cowan, M. J., Ge, B., Koutsodendris, N.,

Sandor K. P., Komberg, J., Vangoor, V. R., Senthilkumar, K., Hennes, V., Seah, C.,

Nelson, A. R., Cheng, T. Y., Lee, S. J., August, P. R., Chen, J. A., Wisniewski, N.,

Hanson-Smith, V., Belgard, T. G., Zhang, A., Coba, M., Grunseich, C., Ward, M. E., van

den Berg, L. H., Pasterkamp, R. J., Trotti, D., Zlokovic, B. V., and Ichida, J. K. (2018).

Haploinsufficiency leads to neurodegeneration in C9ORF72 ALS/FTD human induced

motor neurons. Nat. Med. *24*, 313–325. https://doi.org/10.1038/nm.4490

1246

1247   Sofola, O. A., Jin, P., Qin, Y., Duan, R., Liu, H., de Haro, M., Nelson, D. L., and

1248   Botas, J. (2007). RNA-binding proteins hnRNP A2/B1 and CUGBP1 suppress fragile X

1249   CGG premutation repeat-induced neurodegeneration in a Drosophila model of FXTAS.

1250   Neuron *55*, 565–571. https://doi.org/10.1016/j.neuron.2007.07.021

1251

1252   Sonobe, Y., Ghadge, G., Masaki, K., Sendoel, A., Fuchs, E., and Roos, R. P.

1253   (2018). Translation of dipeptide repeat proteins from the C9ORF72 expanded repeat is

1254   associated with cellular stress. Neurobiol. Dis. *116*, 155–165.

1255   https://doi.org/10.1016/j.nbd.2018.05.009

1256

Su, Z., Zhang, Y., Gendron, T. F., Bauer, P. O., Chew, J., Yang, W. Y., Fostvedt,

E., Jansen-West, K., Belzil, V. V., Desaro, P., Johnston, A., Overstreet, K., Oh, S. Y.,

Todd, P. K., Berry, J. D., Cudkowicz, M. E., Boeve, B. F., Dickson, D., Floeter, M. K.,

Traynor, B. J., Morelli, C., Ratti, A., Silani, V., Rademakers, R., Brown, R. H.,

Rothstein, J. D., Boylan, K. B., Petrucelli, L., and Disney, M. D. (2014). Discovery of a

biomarker and lead small molecules to target r(GGGGCC)-associated defects in

c9FTD/ALS. Neuron *83*, 1043–1050. https://doi.org/10.1016/j.neuron.2014.07.041

1264

1265   Tabet, R., Schaeffer, L., Freyermuth, F., Jambeau, M., Workman, M., Lee, C. Z.,

1266   Lin, C. C., Jiang, J., Jansen-West, K., Abou-Hamdan, H., Désaubry, L., Gendron, T.,

1267   Petrucelli, L., Martin, F., and Lagier-Tourenne, C. (2018). CUG initiation and

1268   frameshifting enable production of dipeptide repeat proteins from ALS/FTD C9ORF72

1269   transcripts. Nat. Commun. *9*, 152. https://doi.org/10.1038/s41467-017-02643-5

1270

1271   Taminato, T., Takeuchi, T., Ueyama, M., Mori, K., Ikeda, M., Mochizuki, H.,

1272   and Nagai, Y. (2023). Therapeutic reduction of GGGGCC repeat RNA levels by

1273   hnRNPA3 suppresses neurodegeneration in drosophila models of C9orf72-linked

1274   ALS/FTD.   Hum.   Mol.   Genet.   ddac298.   Advance   online   publication.

1275   https://doi.org/10.1093/hmg/ddac298

1276

1277   Tran, H., Schilling, M., Wirbelauer, C., Hess, D., and Nagamine, Y. (2004).

1278   Facilitation of mRNA deadenylation and decay by the exosome-bound, DExH protein

1279   RHAU Mol Cell. *13*, 101–111. https://doi.org/10.1016/s1097-2765(03)00481-7

1280

1281   Tran, H., Almeida, S., Moore, J., Gendron, T. F., Chalasani, U., Lu, Y., Du, X.,

1282 Nickerson, J. A., Petrucelli, L., Weng, Z., and Gao, F. B. (2015). Differential Toxicity of

1283 Nuclear RNA Foci versus Dipeptide Repeat Proteins in a Drosophila Model of

1284 C9ORF72 FTD/ALS. Neuron *87,* 1207–1214.

1285 https://doi.org/10.1016/j.neuron.2015.09.015

1286

Trost, B., Engchuan, W., Nguyen, C. M., Thiruvahindrapuram, B., Dolzhenko,

E., Backstrom, I., Mirceta, M., Mojarad, B. A., Yin, Y., Dov, A., Chandrakumar, I.,

Prasolava, T., Shum, N., Hamdan, O., Pellecchia, G., Howe, J. L., Whitney, J., Klee, E.

W., Baheti, S., Amaral, D. G., Anagnostou, E., Elsabbagh, M., Fernandez, B. A., Hoang,

N., Lewis, S., Liu, X., Sjaarda, C., Smith, I. M., Szatmari, P., Zwaigenbaum, L., Glazer,

D., Hartley, D., Stewart, A. K., Eberle, M. A., Sato, N., Pearson, C.E., Scherer, S.W.,

and Yuen, R. (2020). Genome-wide detection of tandem DNA repeats that are expanded

in autism. Nature *586,* 80–86. https://doi.org/10.1038/s41586-020-2579-z

1295

1296   Tseng, Y. J., Sandwith, S. N., Green, K. M., Chambers, A. E., Krans, A., Raimer,

1297 H. M., Sharlow, M. E., Reisinger, M. A., Richardson, A. E., Routh, E. D., Smaldino, M.

1298 A., Wang, Y. H., Vaughn, J. P., Todd, P. K., and Smaldino, P. J. (2021). The RNA

1299 helicase DHX36-G4R1 modulates C9orf72 GGGGCC hexanucleotide repeat-associated

1300 translation. J. Biol. Chem. *297,* 100914. https://doi.org/10.1016/j.jbc.2021.100914

1301

1302   Waite, A. J., Bäumer, D., East, S., Neal, J., Morris, H. R., Ansorge, O., and

1303 Blake, D. J. (2014). Reduced C9orf72 protein levels in frontal cortex of amyotrophic

1304 lateral sclerosis and frontotemporal degeneration brain with the C9ORF72

1305 hexanucleotide repeat expansion. Neurobiol. Aging *35*, 1779.e5–1779.e13.

1306 https://doi.org/10.1016/j.neurobiolaging.2014.01.016

1307

1308 Wang, J. W., Brent, J. R., Tomlinson, A., Shneider, N. A., and McCabe, B. D.

1309 (2011). The ALS-associated proteins FUS and TDP-43 function together to affect

1310 Drosophila locomotion and life span. J. Clin. Invest. *121*, 4118–4126.

1311 https://doi.org/10.1172/JCI57883

1312

Wang, Z. F., Ursu, A., Childs-Disney, J. L., Guertler, R., Yang, W. Y., Bernat, V.,

Rzuczek, S. G., Fuerst, R., Zhang, Y. J., Gendron, T. F., Yildirim, I., Dwyer, B. G., Rice,

J. E., Petrucelli, L., and Disney, M. D. (2019). The Hairpin Form of r(G4C2)^{exp} in

c9ALS/FTD Is Repeat-Associated Non-ATG Translated and a Target for Bioactive

Small Molecules. Cell Chem. Biol. *26*, 179–190.e12.

https://doi.org/10.1016/j.chembiol.2018.10.018

1319

1320 Wen, X., Tan, W., Westergard, T., Krishnamurthy, K., Markandaiah, S. S., Shi,

1321 Y., Lin, S., Shneider, N. A., Monaghan, J., Pandey, U. B., Pasinelli, P., Ichida, J. K., and

1322 Trotti, D. (2014). Antisense proline-arginine RAN dipeptides linked to C9ORF72-

1323 ALS/FTD form toxic nuclear aggregates that initiate in vitro and in vivo neuronal death.

1324 Neuron *84*, 1213–1225. https://doi.org/10.1016/j.neuron.2014.12.010

1325

1326 Xu, Z., Poidevin, M., Li, X., Li, Y., Shu, L., Nelson, D. L., Li, H., Hales, C. M.,

1327 Gearing, M., Wingo, T. S., and Jin, P. (2013). Expanded GGGGCC repeat RNA

1328 associated with amyotrophic lateral sclerosis and frontotemporal dementia causes

1329 neurodegeneration. Proc. Natl. Acad. Sci. USA *110*, 7778–7783.

1330 https://doi.org/10.1073/pnas.1219643110

1331

1332 Yamaguchi, M., Hirose, F., Inoue, Y. H., Shiraki, M., Hayashi, Y., Nishi, Y., and

1333 Matsukage, A. (1999). Ectopic expression of human p53 inhibits entry into S phase and

1334 induces apoptosis in the Drosophila eye imaginal disc. Oncogene *18*, 6767–6775.

1335 https://doi.org/10.1038/sj.onc.1203113

1336

Zhang, Y. J., Gendron, T. F., Grima, J. C., Sasaguri, H., Jansen-West, K., Xu, Y.

F., Katzman, R. B., Gass, J., Murray, M. E., Shinohara, M., Lin, W. L., Garrett, A.,

Stankowski, J. N., Daughrity, L., Tong, J., Perkerson, E. A., Yue, M., Chew, J.,

Castanedes-Casey, M., Kurti, A., Wang, Z. S., Liesinger, A. M., Baker, J. D., Jiang, J.,

Lagier-Tourenne, C., Edbauer, D., Cleveland, D. W., Rademakers, R., Boylan, K. B.,

Bu, G., Link, C. D., Dickey, C. A., Rothstein, J. D., Dickson, D. W., Fryer, J. D., and

Petrucelli, L. (2016). C9ORF72 poly(GA) aggregates sequester and impair HR23 and

nucleocytoplasmic transport proteins. Nat. Neurosci. *19*, 668–677.

https://doi.org/10.1038/nn.4272

1346

1347 Zhang, Y. J., Gendron, T. F., Ebbert, M., O'Raw, A. D., Yue, M., Jansen-West,

1348 K., Zhang, X., Prudencio, M., Chew, J., Cook, C. N., Daughrity, L. M., Tong, J., Song,

1349 Y., Pickles, S. R., Castanedes-Casey, M., Kurti, A., Rademakers, R., Oskarsson, B.,

1350 Dickson, D. W., Hu, W., Gitler, A. D., Fryer, J. D., and Petrucelli, L. (2018). Poly(GR)

1351 impairs protein translation and stress granule dynamics in C9orf72-associated

frontotemporal dementia and amyotrophic lateral sclerosis. Nat. Med. *24*, 1136–1142. https://doi.org/10.1038/s41591-018-0071-1

Zhu, Q., Jiang, J., Gendron, T. F., McAlonis-Downes, M., Jiang, L., Taylor, A., Diaz Garcia, S., Ghosh Dastidar, S., Rodriguez, M. J., King, P., Zhang, Y., La Spada, A. R., Xu, H., Petrucelli, L., Ravits, J., Da Cruz, S., Lagier-Tourenne, C., and Cleveland, D. W. (2020). Reduced C9ORF72 function exacerbates gain of toxicity from ALS/FTD-causing repeat expansion in C9orf72. Nat. Neurosci. *23*, 615–624. https://doi.org/10.1038/s41593-020-0619-5

Zu, T., Gibbens, B., Doty, N. S., Gomes-Pereira, M., Huguet, A., Stone, M. D., Margolis, J., Peterson, M., Markowski, T. W., Ingram, M. A., Nan, Z., Forster, C., Low, W. C., Schoser, B., Somia, N. V., Clark, H. B., Schmechel, S., Bitterman, P. B., Gourdon, G., Swanson, M. S., Moseley, M., and Ranum, L. P. (2011). Non-ATG-initiated translation directed by microsatellite expansions. Proc. Natl. Acad. Sci. USA *108*, 260–265. https://doi.org/10.1073/pnas.1013343108

Zu, T., Liu, Y., Bañez-Coronel, M., Reid, T., Pletnikova, O., Lewis, J., Miller, T. M., Harms, M. B., Falchook, A. E., Subramony, S. H., Ostrow, L. W., Rothstein, J. D., Troncoso, J. C., and Ranum, L. P. (2013). RAN proteins and RNA foci from antisense transcripts in C9ORF72 ALS and frontotemporal dementia. Proc. Natl. Acad. Sci. USA *110*, E4968–E4977. https://doi.org/10.1073/pnas.1315438110

Zu, T., Cleary, J. D., Liu, Y., Bañez-Coronel, M., Bubenik, J. L., Ayhan, F.,

1376   Ashizawa, T., Xia, G., Clark, H. B., Yachnis, A. T., Swanson, M. S., and Ranum, L.

1377   (2017). RAN Translation Regulated by Muscleblind Proteins in Myotonic Dystrophy

1378   Type 2. Neuron *95*, 1292–1305.e5. https://doi.org/10.1016/j.neuron.2017.08.039

1379

 Figure legends

 Figure 1. Screening for RBPs that suppress G_4C_2 repeat-induced toxicity in C9-

 ALS/FTD flies.

 (A) Light microscopic images of the eyes in flies expressing both $(G_4C_2)_{42\ or\ 89}$ and the

 indicated RBPs using the *GMR-Gal4* driver. Coexpression of *FUS*, *IGF2BP1*, or

 hnRNPA2B suppressed eye degeneration in both $(G_4C_2)_{42}$ and $(G_4C_2)_{89}$ flies, indicated by

 "Suppression (strong)". Coexpression of *hnRNPR*, *SAFB2*, *SF3B3*, *hnRNPA1*, or

 hnRNPL suppressed eye degeneration in either $(G_4C_2)_{42}$ or $(G_4C_2)_{89}$ flies, indicated by

 "Suppression (weak)" (see also Figure 1—source data 2). Scale bar: 100 μm. (B)

 Quantification of eye size in $(G_4C_2)_{89}$ flies coexpressing the indicated RBPs (n = 5). (C),

 (D) Quantification of eye pigmentation in $(G_4C_2)_{89}$ flies (C) or $(G_4C_2)_{42}$ flies (D)

 coexpressing the indicated RBPs (n = 5). (E) Expression levels of $(G_4C_2)_{89}$ RNA in flies

 expressing both $(G_4C_2)_{89}$ and the indicated RBPs using the *GMR-Gal4* driver (five

 independent experiments, n = 25 flies per genotype). The $(G_4C_2)_{89}$(H) fly line expresses

 $(G_4C_2)_{89}$ RNA at a high level (see also Figure 1—figure supplement 1). In (B–E), data

 are presented as the mean ± SEM; *P* < 0.0001, as assessed by one-way ANOVA; n.s.:

 not significant, $^*P\Box<\Box0.05$, $^{**}P\Box<\Box0.01$, and $^{***}P\Box<\Box0.001$, as assessed by Tukey's

 post hoc analysis. The detailed statistical information is summarized in Figure 1—

 source data 3.

 Figure 1–figure supplement 1. Characterization of C9-ALS/FTD flies.

 (A) $(G_4C_2)_n$ constructs used in this study. These constructs do not include an ATG start

 codon downstream of the UAS sequence, and were expressed in a tissue-specific

 manner using the *GAL4-UAS* system. (B) Light microscopic images of the eyes in flies

expressing $(G_4C_2)_n$ using the *GMR-Gal4* driver. Scale bar: 100 μm. (C) Expression

levels of $(G_4C_2)_n$ RNAs in flies expressing $(G_4C_2)_n$ using the *GMR-Gal4* driver. Strong

eye degeneration with decreased eye size and loss of pigmentation was observed in

$(G_4C_2)_{42 \text{ or } 89}$ flies, but not in $(G_4C_2)_9$ flies. Eye degeneration was confirmed in $(G_4C_2)_{42}$

and two $(G_4C_2)_{89}$ independent fly lines. Degree of eye degeneration in two $(G_4C_2)_{89}$ fly

lines was expression-level dependent [(L) vs (H) in $(G_4C_2)_{89}$] (three independent

experiments, n = 15 flies per each genotype). Expression of G_4C_2 repeat RNA of the

sense transcripts but not that of the antisense transcripts was confirmed. (D) Climbing

ability at 1 day of age in flies expressing $(G_4C_2)_n$ using the *elav-Gal4* driver. Flies

expressing $(G_4C_2)_{89}$(H) in neurons showed lethality. Decreasing climbing ability was

observed in $(G_4C_2)_{42 \text{ or } 89}$ flies compared with $(G_4C_2)_9$ flies (five independent

experiments, n = 100 flies per each genotype). (E) FISH analyses of G_4C_2 repeat RNA

in the salivary glands of fly larvae with two copies of *GMR-Gal4* and $(G_4C_2)_{9 \text{ or } 89}$ (red:

G_4C_2 RNA; yellow: G_2C_4 RNA; blue [DAPI]: nuclei). RNA foci formation

(arrowheads) of the sense transcripts but not that of the antisense transcripts was

confirmed. RNA foci were observed in $(G_4C_2)_{89}$ flies, but not in $(G_4C_2)_9$ flies. Scale

bars: 100 μm (low magnification) or 20 µm (high magnification). (F)

Immunohistochemical analyses of DPRs stained with anti-DPR antibodies in the eye

imaginal discs of fly larvae with two copies of *GMR-Gal4* and $(G_4C_2)_{9 \text{ or } 89}$ (magenta:

poly(GR); orange: poly(GA); green: poly(GP); blue [DAPI]: nuclei). Expression and

cytoplasmic aggregation (arrowheads) of three DPRs in $(G_4C_2)_{89}$ flies, but not in

$(G_4C_2)_9$ flies, were confirmed. Scale bars: 50 μm (low magnification), 10 μm (middle

magnification), and 5 μm (high magnification). In (B–F), L: low-expression line; H:

high-expression line. In (C) and (D), data are presented as the mean ± SEM; $P < 0.0001$,

as assessed by one-way ANOVA; n.s.: not significant, and $^{***}P < 0.001$, as assessed

by Tukey's post hoc analysis. The detailed statistical information is summarized in

Figure 1—figure supplement 1—source data 2.

1431

Figure 1–figure supplement 2. Coexpression of FUS suppresses G$_4$C$_2$ repeat-

induced toxicity in flies expressing (G$_4$C$_2$)$_{89}$.

(A) Light microscopic images of the eyes in flies expressing both (G$_4$C$_2$)$_{42 \text{ or } 89}$ and *FUS*

using the *GMR-Gal4* driver. *FUS-2* and *FUS-3* are different strains from that in Figure

1. Scale bar: 100 μm. (B) Quantification of the eye size in (G$_4$C$_2$)$_{89}$ flies of the indicated

genotypes (n = 10). (C), (D) Quantification of eye pigmentation in (G$_4$C$_2$)$_{89}$ flies (C) or

(G$_4$C$_2$)$_{42}$ flies (D) of the indicated genotypes (n = 10). In (B–D), data are presented as

the mean ± SEM; $P < 0.0001$, as assessed by one-way ANOVA; $^{***}P < 0.001$, as

assessed by Tukey's post hoc analysis. The detailed statistical information is

summarized in Figure 1—figure supplement 2—source data 1.

1442

Figure 2. FUS suppresses G$_4$C$_2$ repeat-induced toxicity via its RNA-binding

activity.

(A) Light microscopic images of the eyes in flies expressing both (G$_4$C$_2$)$_{89}$ and either

FUS or *FUS-RRMmut* using the *GMR-Gal4* driver. Scale bar: 100 μm. (B)

Quantification of eye size in the flies of the indicated genotypes (n = 10). (C)

Quantification of eye pigmentation in the flies of the indicated genotypes (n = 10). (D)

Egg-to-adult viability in flies expressing both (G$_4$C$_2$)$_{42}$ and either *FUS* or *FUS-RRMmut*

using the *GMR-Gal4* driver (> 500 flies per genotype). (E) Climbing ability in flies

expressing both (G$_4$C$_2$)$_{42}$ and *FUS* using the *elav-GeneSwitch* driver (five independent

experiments, n = 100 flies per each genotype). In (B–E), data are presented as the mean

± SEM. In (B) and (C), $P < 0.0001$, as assessed by one-way ANOVA; n.s.: not

significant, and $^{***}P < 0.001$, as assessed by Tukey's post hoc analysis. In (D), n.s.:

not significant and $^{***}P < 0.001$, as assessed by Tukey's multiple comparison test

using wholly significant difference. In (E), n.s.: not significant, $^{*}P < 0.05$,

$^{**}P < 0.01$ and $^{***}P < 0.001$, as assessed by two-way repeated measures ANOVA

with Tukey's post hoc analysis. The detailed statistical information is summarized in

Figure 2—source data 1.

1460

Figure 2–figure supplement 1. Western blot analysis showing expression levels of

FUS and FUS-RRMmut proteins.

(A) Western blot analysis of the FUS and FUS-RRMmut proteins in the heads of adult

flies expressing *EGFP, FUS* or *FUS-RRMmut* using the *GMR-Gal4* driver, with an anti-

FUS antibody. The arrowhead indicates bands from the FUS and FUS-RRMmut

proteins, whereas the asterisk indicates bands resulting from nonspecific antibody

binding. (B) Quantification of the FUS and FUS-RRMmut proteins from the Western

blot analysis in (A) (n = 3). In (B), data are presented as the mean ± SEM; n.s.: not

significant, as assessed by the unpaired *t*-test. The detailed statistical information is

summarized in Figure 2—figure supplement 1—source data 1.

1471

Figure 3. FUS suppresses RNA foci formation and RAN translation from G_4C_2

repeat RNA.

(A) FISH analyses of G_4C_2 repeat RNA in the salivary glands of fly larvae expressing

both $(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using two copies of the *GMR-Gal4*

driver (red: G_4C_2 RNA; blue [DAPI]: nuclei). Arrowheads indicate RNA foci. Scale bar:

20 µm. (B) Quantification of the number of nuclei containing RNA foci from the FISH

analyses in (A) (n = 10). (C) Expression levels of $(G_4C_2)_{89}$ RNA in fly larvae expressing

both $(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using the *GMR-Gal4* driver (10

independent experiments, n = 50 flies per each genotype). (D) Immunohistochemical

analyses of DPRs stained with anti-DPR antibodies in the eye imaginal discs of fly

larvae expressing both $(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using two copies of

the *GMR-Gal4* driver (magenta: poly(GR); orange: poly(GA); green: poly(GP)).

Arrowheads indicate cytoplasmic aggregates. Scale bars: 20 µm (low magnification) or

5 µm (high magnification). (E) Quantification of the number of DPR aggregates from

the immunohistochemical analyses in (D) [n = 14 or 15 (GR), or 10 (GA or GP)]. (F)

Immunoassay to determine poly(GP) levels in flies expressing both $(G_4C_2)_{89}$ and either

FUS or *FUS-RRMmut* using the *GMR-Gal4* driver (three independent experiments, n =

30 flies per each genotype). (G) Western blot analysis of the heads of adult flies

expressing both LDS-$(G_4C_2)_{44}^{GR-GFP}$ and any of *DsRed*, *FUS* or *FUS-RRMmut* using the

GMR-Gal4 driver, using either an anti-GFP (upper panel) or anti-GR antibody (middle

panel). (H and I) Quantification of GR-GFP protein levels from the Western blot

analysis in (G) (nine independent experiments, n = 90 flies per each genotype). In (B),

(C), (E), (F), (H), and (I), data are presented as the mean ± SEM. In (B), (E), and (F), *P*

< 0.0001, as assessed by one-way ANOVA; n.s.: not significant, $^*P < 0.05$,

$^{**}P < 0.01$, and $^{***}P < 0.001$, as assessed by Tukey's post hoc analysis. In (C), *P* =

0.452, as assessed by one-way ANOVA; n.s.: not significant, as assessed by Tukey's

post hoc analysis. In (H), *P* = 0.0148, as assessed by one-way ANOVA; n.s.: not

significant and $^*P < 0.05$, as assessed by Tukey's post hoc analysis. In (I), *P* =

0.0072, as assessed by one-way ANOVA; n.s.: not significant and $^*P\square<\square0.05$, as

assessed by Tukey's post hoc analysis. The detailed statistical information is

summarized in Figure 3—source data 1.

1503

1504 **Figure 3–figure supplement 1. Schema of the LDS-$(G_4C_2)_{44}^{\text{GR-GFP}}$ construct.**

1505 Schema of the LDS-$(G_4C_2)_{44}^{\text{GR-GFP}}$ construct containing the $(G_4C_2)_{44}$ sequence and 114

1506 nucleotides of the 5′-flanking region of intron 1 of the human *C9orf72* G_4C_2 repeat

1507 sequence. A GFP tag in the GR frame was introduced downstream of the $(G_4C_2)_{44}$ repeat

1508 sequence.

1509

Figure 3–figure supplement 2. Overexpression of *FUS* does not suppress eye

degeneration in DPR-only flies expressing DPRs translated from non-G_4C_2 RNAs.

Light microscopic images of the eyes in DPR-only flies coexpressing either the

poly(GR) or poly(GA) protein, and either *FUS* or *FUS-RRMmut* using the *GMR-Gal4*

driver. Overexpression of *FUS* did not suppress the eye degeneration in flies expressing

either $(GR)_{36}$ or $(GR)_{100}$. Overexpression of *FUS* also caused mild eye degeneration in

flies expressing EGFP, $(GA)_{36}$, or $(GA)_{100}$, likely due to FUS toxicity.

1517

Figure 4. Reduction of endogenous *caz* expression enhances G_4C_2 repeat-induced

toxicity, RNA foci formation, and DPR aggregation.

(A) Light microscopic images of the eyes in flies expressing $(G_4C_2)_{89}$ using the *GMR-*

Gal4 driver, with knockdown of *caz*. Scale bar: 100 μm. (B) Quantification of eye size

in flies of the indicated genotypes shown in (A) (n = 10). (C) Light microscopic images

of the eyes in flies expressing $(G_4C_2)_{89}$ using the *GMR-Gal4* driver, with a hemizygous

deletion of *caz*. Scale bar: 100 μm. (D) Quantification of eye size in the flies of the

indicated genotypes shown in (C) (n = 10). (E) FISH analyses of G_4C_2 repeat RNA in

the salivary glands of fly larvae expressing $(G_4C_2)_{89}$ using the *GMR-Gal4* driver, with

knockdown of *caz* (red: G_4C_2 RNA; blue [DAPI]: nuclei). Arrowheads indicate RNA

foci. Scale bar: 20 μm. (F) Quantification of the number of nuclei containing RNA foci

from the FISH analyses in (E) (n = 10). (G) Expression levels of $(G_4C_2)_{89}$ RNA in fly

larvae expressing $(G_4C_2)_{89}$ using the *GMR-Gal4* driver, with knockdown of *caz* (four

independent experiments, n = 20 flies per each genotype). (H) Immunohistochemical

analyses of DPRs stained with anti-DPR antibodies in the eye imaginal discs of fly

larvae expressing $(G_4C_2)_{89}$ using two copies of the *GMR-Gal4* driver, with the

knockdown of *caz*. (magenta: poly(GR); orange: poly(GA); green: poly(GP)).

Arrowheads indicate cytoplasmic aggregates. Scale bars: 20 μm (low magnification) or

5 μm (high magnification). (I) Quantification of the number of DPR aggregates from

the immunohistochemical analyses in (H) (n = 10). In (B), (D), (F), (G) and (I), data are

presented as the mean ± SEM. In (B) and (D), $P < 0.0001$, as assessed by one-way

ANOVA; [***]$P < 0.001$, as assessed by Tukey's post hoc analysis. In (F), (G), and (I),

n.s.: not significant, [*]$P < 0.05$, [**]$P < 0.01$, and [***]$P < 0.001$, as assessed by the

unpaired *t*-test. The detailed statistical information is summarized in Figure 4—source

data 1.

1543

Figure 4–figure supplement 1. Endogenous *caz* is a functional homologue of *FUS*

for the suppression of G_4C_2 repeat-induced toxicity.

(A) Light microscopic images of the eyes in flies expressing both $(G_4C_2)_{42 \text{ or } 89}$ and

either *caz* (*FLAG-caz*) or *FUS-4* (*FLAG-FUS*) using the *GMR-Gal4* driver. *FUS-4* is a

different strain from those used in Figure 1 and Figure 1–figure supplement 2. Scale

bar: 100 μm. (B) Quantification of the eye size in $(G_4C_2)_{89}$ flies of the indicated

genotypes (n = 10). (C), (D) Quantification of eye pigmentation in $(G_4C_2)_{89}$ flies (C) or

$(G_4C_2)_{42}$ flies (D) of the indicated genotypes (n = 10). In (B–D), data are presented as

the mean ± SEM; $P < 0.0001$, as assessed by one-way ANOVA; n.s.: not significant,

$^*P < 0.05$, $^{**}P < 0.01$,and $^{***}P < 0.001$, as assessed by Tukey's post hoc

analysis. The detailed statistical information is summarized in Figure 4—figure

supplement 1—source data 1.

1556

Figure 5. FUS directly binds to and modulates the G-quadruplex structure of G_4C_2

repeat RNA, resulting in the suppression of RAN translation *in vitro*

(A) Analysis of the binding of His-tagged FUS proteins to biotinylated $(G_4C_2)_4$ RNA by

the filter binding assay. The nitrocellulose membrane (left) traps RNA-bound FUS

proteins, whereas unbound RNAs are recovered on the nylon membrane (right), and

then the RNAs trapped on each of the membranes was probed with streptavidin-

horseradish peroxidase (HRP). Biotinylated $(AAAAAA)_4$ and $(UUAGGG)_4$ were used

as negative and positive controls, respectively. (B–D) CD spectra of $(G_4C_2)_4$ RNA

incubated with or without FUS in the presence of 150 mM KCl (B), NaCl (C), or LiCl

(D). The CD spectrum of FUS alone was subtracted from that of $(G_4C_2)_4$ RNA

incubated with FUS. The original data are shown in Figure 5—figure supplements 2B-

2D. (E) Schema of the template RNA containing the $(G_4C_2)_{80}$ sequence and 113

nucleotides of the 5′-flanking region of intron 1 of the human *C9orf72* G_4C_2 repeat

sequence. A Myc tag in the GA frame was introduced downstream of the $(G_4C_2)_{80}$ repeat

sequence. (F) Western blot analysis of samples from *in vitro* translation using rabbit

reticulocyte lysate in the presence or absence of increasing concentrations of FUS or

FUS-RRMmut. The GA-Myc fusion protein was detected by Western blotting using the

anti-Myc antibody. (G) Quantification of the GA-Myc fusion protein in (F) (n = 3). In

(G), data are presented as the mean ± SEM; $^{*}P \square < \square 0.05$, $^{**}P \square < \square 0.01$,and

$^{***}P \square < \square 0.001$, as assessed by the unpaired *t*-test. The detailed statistical information is

summarized in Figure 5—source data 1.

1578

1579

1580 **Figure 5–figure supplement 1. FUS colocalizes with G_4C_2 RNA foci.**

1581 Combined FISH and immunohistochemical analyses of G_4C_2 repeat RNA and FUS in

1582 the salivary glands of flies expressing both $(G_4C_2)_{89}$ and *FUS* using two copies of the

1583 *GMR-Gal4* driver. Arrowheads indicate colocalization of FUS with RNA foci. Scale

1584 bar: 10 μm (low magnification), and 5 μm (high magnification).

1585

Figure 5–figure supplement 2. FUS modulates the G-quadruplex structure of G_4C_2

repeat RNA.

(A–C) CD spectra of $(G_4C_2)_4$ RNA incubated with or without FUS in the presence of

150 mM KCl (A), NaCl (B), or LiCl (C). CD spectra of $(G_4C_2)_4$ RNA alone (black),

FUS alone (green), sum of $(G_4C_2)_4$ RNA and FUS (blue), and the spectra of their

coincubation (magenta) are shown. FUS interacts with $(G_4C_2)_4$ RNA under KCl or NaCl

buffer conditions. (D) Imino proton NMR spectra of $(G_4C_2)_4$ RNA incubated with

increasing amounts of FUS in the presence of 150 mM KCl.

1594

Figure 6. Identification of G-quadruplex-targeting RBPs that suppress G_4C_2

repeat-induced toxicity in C9-ALS/FTD flies.

(A) Light microscopic images of eyes in flies expressing both $(G_4C_2)_{89}$ and the indicated

G-quadruplex-targeting RBPs using the *GMR-Gal4* driver. Scale bar: 100 μm. (B)

Quantification of eye size in the flies of the indicated genotypes (n = 10). (C)

Quantification of eye pigmentation in the flies of the indicated genotypes (n = 10). (D)

Expression levels of $(G_4C_2)_{89}$ RNA in flies expressing both $(G_4C_2)_{89}$ and the indicated

G-quadruplex-targeting RBPs using the *GMR-Gal4* driver (five independent

experiments, n = 25 flies per each genotype). (E) Immunohistochemical analyses of

poly(GA) stained with anti-GA antibody in the eye imaginal discs of fly larvae

expressing both $(G_4C_2)_{89}$ and the indicated G-quadruplex-targeting RBPs using two

copies of the *GMR-Gal4* driver (orange: poly(GA)). Arrowheads indicate cytoplasmic

aggregates. Scale bars: 20 μm (low magnification) or 5 μm (high magnification). (F)

Quantification of the number of poly(GA) aggregates from the immunohistochemical

analyses in (E) (n=10). In (B), (C), (D), and (F), data are presented as the mean ± SEM;

$P < 0.0001$, as assessed by one-way ANOVA; n.s.: not significant, $^{*}P < 0.05$, and

$^{***}P < 0.001$, as assessed by Tukey's post hoc analysis. The detailed statistical

information is summarized in Figure 6—source data 2.

1613

1614 **Tables**

1615 **Table 1. Association (k_a) and dissociation (k_d) rate and dissociation constants (K_D)**

1616 **between FUS and $(G_4C_2)_4$ RNA in different buffers as assessed by SPR analysis.**

1617

Buffer	k_a ($M^{-1}s^{-1}$) $\times 10^6$	k_d (s^{-1}) $\times 10^{-3}$	K_D (M)
KCl	1.4	22	1.5×10^{-8}
NaCl	0.41	54	1.3×10^{-7}
LiCl	0.0018	25	1.4×10^{-5}

1618
1619

1620 **Source data files and supplementary files list**

1621 Figure 1—source data 1. RNA-binding proteins and their cDNA accession numbers

1622 screened in the genetic analyses in Figure 1.

1623

1624 Figure 1—source data 2. Summary of the genetic analyses in Figure 1.

1625

1626 Figure 1—source data 3. Statistical data related to Figures 1B, 1C, 1D, and 1E.

1627

1628 Figure 1—figure supplement 1‒source data 1. The artificial sequence inserted in the

1629 pUAST vector for generation of $(G_4C_2)_n$ flies.

1630

1631 Figure 1—figure supplement 1‒source data 2. Statistical data related to Figure 1—

1632 figure supplements 1C and 1D.

1633

1634 Figure 1—figure supplement 2‒source data 1. Statistical data related to Figure 1—

1635 figure supplement 2B, 2C and 2D.

1636

1637 Figure 2—source data 1. Statistical data related to Figures 2B, 2C, 2D, and 2E.

1638

1639 Figure 2—figure supplement 1‒source data 1. Statistical data related to Figure 2—

1640 figure supplement 1B.

1641

1642 Figure 2—figure supplement 1‒source data 2. Source data related to Figure 2—figure

1643 supplement 1A.

1644

1645 Figure 3–source data 1. Statistical data related to Figures 3B, 3C, 3E, 3F, 3H, and 3I.

1646

1647 Figure 3–source data 2. Source data related to Figure 3G.

1648

1649 Figure 4—source data 1. Statistical data related to Figures 4B, 4D, 4F, 4G and 4I.

1650

1651 Figure 4—figure supplement 1–source data 1. Statistical data related to Figure 4—

1652 figure supplements 1B, 1C, and 1D.

1653

1654 Figure 5—source data 1. Statistical data related to Figures 5G.

1655

1656 Figure 5—source data 2. Source data related to Figure 5F.

1657

1658 Figure 6—source data 1. RNA-binding proteins and their cDNA accession numbers

1659 screened in the genetic analyses in Figure 6.

1660

1661 Figure 6—source data 2. Statistical data related to Figures 6B, 6C, 6D, and 6F.

1662

1663 Supplementary file 1. Full genotypes of the fly lines and their cultured temperatures.

1664

A

	Suppression (strong)			Suppression (weak)				
	EGFP	*FUS*	*IGF2BP1*	*hnRNPA2B1*	*hnRNPR*	*SAFB2*	*SF3B3*	*hnRNPA1*	*hnRNPL*

$(G_4C_2)_{89}(H)$

$(G_4C_2)_{42}$

EGFP

	No effect						Enhancement		Unknown		
	DHX30	*SAFB*	*DHX15*	*ILF2*	*DDX21*	*TDP-43*	*hnRNPK*	*SFPQ*	*ILF3*	*NONO*	*ELAVL1*

$(G_4C_2)_{89}(H)$ — *TDP-43*: Lethal; *ILF3*: Lethal; *NONO*: Lehtal; *ELAVL1*: Lethal

$(G_4C_2)_{42}$ — *DDX21*: Lethal; *ILF3*: Lethal; *NONO*: Lehtal; *ELAVL1*: Lethal

EGFP

B Eye size — $(G_4C_2)_{89}(H)$

C Eye pigmentation — $(G_4C_2)_{89}(H)$

D Eye pigmentation — $(G_4C_2)_{42}$

E $(G_4C_2)_{89}$ RNA levels — $(G_4C_2)_{89}(H)$

A

	EGFP	*FUS-2*	*FUS-3*
$(G_4C_2)_{89}(H)$			
$(G_4C_2)_{42}$			
EGFP			

B

Eye size

EGFP, EGFP | *EGFP* | *FUS-2* | *FUS-3*
$(G_4C_2)_{89}(H)$

C

Eye pigmentation

EGFP, EGFP | *EGFP* | *FUS-2* | *FUS-3*
$(G_4C_2)_{89}(H)$

D

Eye pigmentation

EGFP, EGFP | *EGFP* | *FUS-2* | *FUS-3*
$(G_4C_2)_{42}$

UAS-LDS-$(G_4C_2)_{44}$^{GR-GFP}

| UAS | 5'flanking | (GGGGCC)₄₄ | GFP (GR frame) |

A

B

C

D

A

B

KCl

- (G$_4$C$_2$)$_4$ RNA
- (G$_4$C$_2$)$_4$ RNA + FUS

C

NaCl

- (G$_4$C$_2$)$_4$ RNA
- (G$_4$C$_2$)$_4$ RNA + FUS

D

LiCl

- (G$_4$C$_2$)$_4$ RNA
- (G$_4$C$_2$)$_4$ RNA + FUS

E

F

G

| | (G₄C₂) RNA | FUS | Merged |

A KCl

CD (mdeg) vs Wavelength (nm)
- (G₄C₂)₄ RNA
- FUS
- sum (G₄C₂)₄ RNA + FUS
- mix (G₄C₂)₄ RNA + FUS

B NaCl

CD (mdeg) vs Wavelength (nm)
- (G₄C₂)₄ RNA
- FUS
- sum (G₄C₂)₄ RNA + FUS
- mix (G₄C₂)₄ RNA + FUS

C LiCl

CD (mdeg) vs Wavelength (nm)
- (G₄C₂)₄ RNA
- FUS
- sum (G₄C₂)₄ RNA + FUS
- mix (G₄C₂)₄ RNA + FUS

D

(G₄C₂)₄ RNA : FUS
- 1:0
- 1:0.2
- 1:0.4
- 0:1

¹H (ppm)

A

	EGFP	*EWSR1*	*DDX3X*	*DDX5*	*DDX17*	*DHX9*	*DHX36*
$(G_4C_2)_{89}$(H)							
EGFP							

B — Eye size

x-axis: *EGFP, EGFP*, *EGFP*, *EWSR1*, *DDX3X*, *DDX5*, *DDX17*, *DHX9*, *DHX36*; $(G_4C_2)_{89}(H)$
*** ***

C — Eye pigmentation

x-axis: *EGFP, EGFP*, *EGFP*, *EWSR1*, *DDX3X*, *DDX5*, *DDX17*, *DHX9*, *DHX36*; $(G_4C_2)_{89}(H)$
*** n.s. ***

D — $(G_4C_2)_{89}$ RNA levels

x-axis: *EGFP*, *EWSR1*, *DDX3X*, *DDX5*, *DDX17*, *DHX9*, *DHX36*; $(G_4C_2)_{89}(H)$
* *** n.s.

E — GMR×2 > $(G_4C_2)_{89}$(H)

GA: *EGFP*, *EWSR1*, *DDX3X*, *DDX5*, *DDX17*

F — Aggregates/13 ommatidia

x-axis: *EGFP*, *EWSR1*, *DDX3X*, *DDX5*, *DDX17*; $(G_4C_2)_{89}(H)$
***