FUS regulates RAN translation through modulating the G-quadruplex structure of

GGGGCC repeat RNA in *C9orf72*-linked ALS/FTD

Yuzo Fujino[1,2], Morio Ueyama[1,3,4], Taro Ishiguro[4,5], Daisaku Ozawa[1,3], Toshihiko

Sugiki[6], Hayato Ito[7], Asako Murata[8], Akira Ishiguro[9], Tania F. Gendron[10], Kohji Mori[11],

Eiichi Tokuda[12], Tomoya Taminato[1,3], Takuya Konno[13], Akihide Koyama[13], Yuya

Kawabe[11], Toshihide Takeuchi[3,14], Yoshiaki Furukawa[12], Toshimichi Fujiwara[6], Manabu

Ikeda[11], Toshiki Mizuno[2], Hideki Mochizuki[15], Hidehiro Mizusawa[5], Keiji Wada[4],

Kinya Ishikawa[5], Osamu Onodera[13], Kazuhiko Nakatani[8], Hideki Taguchi[7,16], Leonard

Petrucelli[10], and Yoshitaka Nagai[1,3,4,14,15]

[1] Department of Neurology, Kindai University Faculty of Medicine, Osaka-Sayama,

Osaka 589-8511, Japan

[2] Department of Neurology, Kyoto Prefectural University of Medicine, Kyoto 602-0841,

Japan

[3] Department of Neurotherapeutics, Osaka University Graduate School of Medicine,

Suita, Osaka 565-0871, Japan

[4] Department of Degenerative Neurological Diseases, National Institute of

Neuroscience, National Center of Neurology and Psychiatry, Kodaira, Tokyo 187-8502,

Japan

[5] Department of Neurology and Neurological Science, Tokyo Medical and Dental

University, Bunkyo-ku, Tokyo 113-8519, Japan

[6] Laboratory of Molecular Biophysics, Institute for Protein Research, Osaka University,

Suita, Osaka 565-0871, Japan

[7] School of Life Science and Technology, Tokyo Institute of Technology, Yokohama,

Kanagawa 226-8503, Japan

[8] Department of Regulatory Bioorganic Chemistry, The Institute of Scientific and

Industrial Research, Osaka University, Ibaraki, Osaka 565-0047, Japan

[9] Research Center for Micro-nano Technology, Hosei University, Koganei, Tokyo 184-

0003, Japan

[10] Department of Neuroscience, Mayo Clinic, Jacksonville, FL 32224, USA

[11] Department of Psychiatry, Osaka University Graduate School of Medicine, Suita,

Osaka 565-0871, Japan

[12] Department of Chemistry, Keio University, Yokohama, Kanagawa 223-8522, Japan

[13] Department of Neurology, Clinical Neuroscience Branch, Brain Research Institute,

Niigata University, Niigata 951-8585, Japan

[14] Life Science Research Institute, Kindai University, Osaka-Sayama, Osaka 589-8511,

Japan

[15] Department of Neurology, Osaka University Graduate School of Medicine, Suita,

Osaka 565-0871, Japan

[16] Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology,

Yokohama, Kanagawa 226-8503, Japan

*Correspondence

Department of Neurology

Kindai University Faculty of Medicine

377-2 Ohnohigashi, Osaka-Sayama, Osaka 589-8511, Japan

Tel.: +81 (72) 366-0246 (ext. 3553); Fax: +81 (72) 368-4846

E-mail: yoshi.nagai@med.kindai.ac.jp

Abstract

Abnormal expansions of GGGGCC repeat sequence in the noncoding region of the *C9orf72* gene is the most common cause of familial amyotrophic lateral sclerosis and frontotemporal dementia (C9-ALS/FTD). The expanded repeat sequence is translated into dipeptide repeat proteins (DPRs) by noncanonical repeat-associated non-AUG (RAN) translation. Since DPRs play central roles in the pathogenesis of C9-ALS/FTD, we here investigate the regulatory mechanisms of RAN translation, focusing on the effects of RNA-binding proteins (RBPs) targeting GGGGCC repeat RNAs. Using C9-ALS/FTD model flies, we demonstrated that the ALS/FTD-linked RBP FUS suppresses RAN translation and neurodegeneration in an RNA-binding activity-dependent manner. Moreover, we found that FUS directly binds to and modulates the G-quadruplex structure of GGGGCC repeat RNA as an RNA chaperone, resulting in the suppression of RAN translation *in vitro*. These results reveal a previously unrecognized regulatory mechanism of RAN translation by G-quadruplex-targeting RBPs, providing therapeutic insights for C9-ALS/FTD and other repeat expansion diseases.

Introduction

Amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) are incurable neurodegenerative diseases with overlapping genetic and neuropathological features. Abnormal expansions of the GGGGCC (G_4C_2) repeat sequence in the noncoding region of the *C9orf72* gene have been found to be the most common genetic mutation responsible for ALS/FTD (DeJesus-Hernandez et al.; 2011, Gijselinck et al., 2012; Renton et al., 2011). Three major pathomechanisms are thought to be involved in the pathogenesis of *C9orf72*-linked ALS/FTD (C9-ALS/FTD): first, expansion of the G_4C_2 repeats results in decreased expression of the *C9orf72* gene, leading to its haploinsufficiency (Boivin et al., 2020; DeJesus-Hernandez et al. 2011; Gijselinck et al., 2012; Shi et al., 2018; Waite et al., 2014; Zhu et al., 2020). Second, the transcribed G_4C_2 repeat-containing RNA accumulates as RNA foci in the affected tissues, sequestering various RNA-binding proteins (RBPs) and altering their function (Conlon et al., 2016; Cooper-Knock et al., 2014; Donnelly et al., 2013; Haeusler et al., 2014; Lee et al., 2013; Mori et al., 2013a). Third, these G_4C_2 repeat RNA is also translated into dipeptide repeat (DPR) proteins, despite the lack of an AUG initiation codon, by noncanonical repeat-associated non-AUG (RAN) translation (Ash et al., 2013; Gendron et al., 2013; Mori et al., 2013b; Mori et al., 2013c; Zu et al, 2011; Zu et al, 2013). Since RAN translation occurs in all reading frames and the expanded G_4C_2 repeat sequence is bidirectionally transcribed, five distinct DPRs, i.e., poly(glycine-arginine) [poly(GR)], poly(glycine-alanine) [poly(GA)], poly(glycine-proline) [poly(GP)], poly(proline-arginine) [poly(PR)], and poly(proline-alanine) [poly(PA)] are produced and observed in patients' brains (Ash et al., 2013; Gendron et al., 2013; Mori et al., 2013b; Mori et al., 2013c; Zu et al., 2013) and cerebrospinal fluid (Gendron et al., 2017; Krishnon et

al., 2022; Lehmer et al., 2017; Su et al., 2014).

DPRs have been shown to exert toxic effects in multiple C9-ALS/FTD models, such

as cultured cells, flies, and mice (Choi et al., 2019; Jovičić et al., 2015; May et al. 2014;

Mizielinska et al., 2014; Rudich et al., 2017; Wen et al., 2014; Zhang et al., 2016;

Zhang et al., 2018). Importantly, the toxicity of DPRs was confirmed in DPR-only flies,

which express DPRs translated from non-G_4C_2 repeat RNAs with alternative codons

and show neurodegeneration, whereas RNA-only flies expressing G_4C_2 repeat RNAs

with stop codon interruptions, which eliminate DPRs production, did not show any

obvious degenerative phenotypes (Mizielinska et al., 2014). In addition, increased DPR

production, but not RNA foci, was reported to correlate with G_4C_2 repeat-induced

toxicity in a C9-ALS/FTD *Drosophila* model (Tran et al., 2015). Taken together, these

studies have strongly suggested that DPRs play a central role in the pathogenesis of C9-

ALS/FTD. Indeed, DPRs have been reported to disrupt various biological pathways,

such as nucleocytoplasmic transport (Hutten et al., 2020; Jovičić et al., 2015; Zhang et

al., 2016) and membraneless organelle dynamics (Kwon et al., 2014; Lee et al., 2016;

Lin et al., 2016). Therefore, elucidating the regulatory mechanism of RAN translation is

a significant challenge toward developing potential therapies for C9-ALS/FTD.

Since the discovery of RAN translation in 2011 (Zu et al., 2011), many studies to

date have focused on its molecular mechanisms, that is, whether it has functional

overlap with canonical AUG-dependent translation. Previous studies on C9-ALS/FTD

using monocistronic reporters containing a G_4C_2 repeat sequence revealed cap-

dependent translation initiation from the upstream near-cognate CUG initiation codon,

requiring the cap-binding eukaryotic translation factor 4F complex (Green et al., 2017;

Tabet et al., 2018). On the other hand, studies using bicistronic reporters with a G_4C_2

repeat sequence in the second cistron also produced DPRs by RAN translation in all reading frames, suggesting cap-independent translation initiation within the G_4C_2 repeat sequence (Cheng et al., 2018; Sonobe et al., 2018). This is reminiscent of internal ribosomal entry site translation initiation, which is another type of noncanonical cap-independent translation in which specific factors are directly recruited to the highly-structured mRNA for initiation (Kwan et al., 2019). While such initiation mechanisms of RAN translation have been explored to date, specific roles of the repeat sequence on RAN translation remain enigmatic. Considering a repeat length dependency of RAN translation (Mori et al., 2013c; Zu et al., 2011; Zu et al., 2013), the repeat sequence itself would also be essential for the initiation or elongation steps of RAN translation. Based on our previous findings of the protective role of TDP-43 on UGGAA repeat-induced toxicity in spinocerebellar ataxia type 31 (SCA31) models (Ishiguro et al., 2017), we hypothesized that RBPs specifically binding to repeat sequences of template RNA play a role in RAN translation.

Using *Drosophila* models of C9-ALS/FTD, we here demonstrate the regulatory roles of the ALS/FTD-linked RBP FUS on RAN translation from G_4C_2 repeat RNA, which lead to the significant modulation of neurodegeneration. We found that FUS suppresses RNA foci formation and DPR production, resulting in the suppression of repeat-induced degeneration. This suppressive effect on degeneration was abolished by mutations in the RNA-recognition motif (RRM) of FUS. In contrast, knockdown of endogenous *caz*, a *Drosophila* homologue of *FUS*, enhanced DPR aggregation and RNA foci formation, resulting in the enhancement of repeat-induced degeneration. Moreover, FUS was found to directly bind to G_4C_2 repeat RNA and modify its G-quadruplex structure as an RNA chaperone, resulting in the suppression of RAN

translation *in vitro*. In addition, other G-quadruplex-targeting RBPs also suppressed

G_4C_2 repeat-induced toxicity in our C9-ALS/FTD flies. These results strongly indicate

that FUS regulates RAN translation and suppresses DPR toxicity through modulating

the G-quadruplex structure of G_4C_2 repeat RNA. Our findings shed light on the

regulatory mechanisms of RAN translation by G-quadruplex-targeting RBPs, and

propose novel therapeutic strategies for repeat expansion diseases by regulating RAN

translation.

Results

Screening for RBPs that suppress G_4C_2 repeat-induced toxicity in C9-ALS/FTD

flies

We established *Drosophila* models of C9-ALS/FTD that express pathogenic

length 42 or 89 G_4C_2 repeats [$(G_4C_2)_{42}$, or $(G_4C_2)_{89}$ flies, respectively], and confirmed

that expanded G_4C_2 repeat sequences induce eye degeneration and motor dysfunction

accompanied with the formation of RNA foci and the production of three types of DPRs

(Figure 1—figure supplement 1), consistent with previous studies (Freibaum et al.,

2015; Goodman et al., 2019; Mizielinska et al., 2014; Xu et al., 2013). We also

established *Drosophila* expressing normal length 9 G_4C_2 repeats as a control [$(G_4C_2)_9$

flies], and found that they did not show eye degeneration, motor dysfunction, RNA foci

formation, or DPR aggregation (Figure 1—figure supplement 1). We selected 18 RBPs

that have been reported to bind to G_4C_2 repeat RNA (Mori et al., 2013a), as well as

TDP-43, an ALS/FTD-linked RBP that does not bind to G_4C_2 repeat RNA (Xu et al.,

2013) (Figure 1—source data 1), and examined their roles in neurodegeneration in our

C9-ALS/FTD fly models. We found that coexpression of *FUS*, *IGF2BP1*, or

hnRNPA2B1 strongly suppressed the eye degeneration in both flies expressing $(G_4C_2)_{42}$

$_{or\ 89}$, which show decreased eye size and loss of pigmentation (Figures 1A–1D and

Figure 1—source data 2). Coexpression of five RBPs, namely, *hnRNPR*, *SAFB2*,

SF3B3, *hnRNPA1*, and *hnRNPL* also partially suppressed the eye degeneration, whereas

coexpression of the other six RBPs had no effect, and two RBPs enhanced the

phenotypes (Figures 1A–1D and Figure 1—source data 2). In addition, coexpression of

TDP-43 had no effect on the eye degeneration in $(G_4C_2)_{42}$ flies, and resulted in lethality

in $(G_4C_2)_{89}$ flies, likely due to the toxicity of *TDP-43* expression itself (Figures 1A and

1D, and Figure 1—source data 2). The variation in the effects of these G_4C_2 repeat-

binding RBPs on G_4C_2 repeat-induced toxicity may be due to their different binding

affinities to G_4C_2 repeat RNA, and their different roles in RNA metabolism. We then

analyzed the expression levels of G_4C_2 repeat RNA in flies coexpressing $(G_4C_2)_{89}$ and

three RBPs that strongly suppressed eye degeneration. We found that coexpression of

IGF2BP1 or *hnRNPA2B1* significantly decreased G_4C_2 repeat RNA levels, whereas they

were not altered upon coexpression of *FUS* (Figure 1E). Although the suppressive

effects of IGF2BP1 and hnRNPA2B1 could simply be explained by the decreased levels

of G_4C_2 repeat RNA, the molecular mechanisms by which FUS suppresses G_4C_2 repeat-

induced toxicity remain to be clarified. The suppressive effects of FUS on G_4C_2 repeat-

induced toxicity was confirmed using multiple FUS fly lines, showing the significant

suppression of decreased eye size and loss of pigmentation in $(G_4C_2)_{42\ or\ 89}$ flies

coexpressing *FUS* (Figure 1—figure supplement 2). Therefore, we decided to further

focus on FUS, which is another ALS/FTD-linked RBP, and investigated its mechanism

of the suppression of G_4C_2 repeat-induced toxicity.

FUS suppresses G_4C_2 repeat-induced toxicity via its RNA-binding activity

We next investigated whether the suppressive effects of FUS on G_4C_2 repeat-

induced toxicity are mediated by its binding to G_4C_2 repeat RNA, using flies expressing

FUS with mutations in the RRM (*FUS-RRMmut*), which have been reported to

eliminate its RNA-binding activity (Daigle et al., 2013). Western blot analysis

confirmed that the *FUS-RRMmut* fly line expresses almost an equivalent level of the

FUS proteins to the *FUS* fly line (Figure 2—figure supplement 1). We found that

coexpression of *FUS-RRMmut* did not restore the eye degeneration in flies expressing

$(G_4C_2)_{89}$, suggesting that the RNA-binding activity of FUS is essential for its

suppressive effects on G_4C_2 repeat-induced toxicity (Figures 2A–C). We also evaluated

the reduced egg-to-adult viability of $(G_4C_2)_{42}$ flies, and confirmed that this phenotype

was rescued by coexpression of *FUS*, but not by coexpression of *FUS-RRMmut* (Figure

2D). Expression of G_4C_2 repeat RNA in the nervous system of flies after eclosion using

the *elav-GeneSwitch* driver induces motor dysfunction, and coexpression of *FUS*

significantly alleviated this motor dysfunction (Figure 2E), indicating that FUS

suppresses the neuronal phenotypes of flies expressing G_4C_2 repeat RNA. It is notable

that the motor dysfunction caused by the expression of *FUS* alone was also alleviated

by coexpression of $(G_4C_2)_{42}$ (Figure 2E), indicating that the G_4C_2 repeat RNA

conversely suppresses FUS toxicity. This result is consistent with our previous

observations in SCA31 flies that UGGAA repeat RNA reduced the aggregation and

toxicity of TDP-43 (Ishiguro et al., 2017). Moreover, recent studies demonstrated that

RNA buffers the phase separation of TDP-43 and FUS, resulting in the suppression of

their aggregation (Maharana et al., 2018; Mann et al., 2019). These findings hence

suggest that balancing the crosstalk between repeat RNAs and RBPs neutralizes the

toxicities of each other.

FUS suppresses RNA foci formation and RAN translation from G_4C_2 repeat RNA

We next analyzed the effects of FUS expression on RNA foci and DPR

production in flies expressing G_4C_2 repeat RNA. We performed RNA fluorescence *in*

situ hybridization (FISH) of the salivary glands of fly larvae expressing $(G_4C_2)_{89}$, and

found that coexpression of *FUS* significantly decreased the number of nuclei containing

RNA foci in $(G_4C_2)_{89}$ flies, whereas it was not altered by coexpression of *FUS-RRMmut*

(Figures 3A and 3B). We confirmed that the expression levels of G_4C_2 repeat RNA in

$(G_4C_2)_{89}$ flies were not altered by coexpression of *FUS* or *FUS-RRMmut* (Figure 3C).

These results were in good agreement with our previous study on SCA31 showing the

suppressive effects of FUS and other RBPs on RNA foci formation of UGGAA repeat

RNA as RNA chaperones (Ishiguro et al., 2017), raising the possibility that FUS has

RNA-chaperoning activity also for G_4C_2 repeat RNA. Immunohistochemistry of the eye

imaginal discs of fly larvae expressing $(G_4C_2)_{89}$ revealed that coexpression of *FUS*

significantly decreased the number of DPR aggregates in $(G_4C_2)_{89}$ flies, whereas

coexpression of *FUS-RRMmut* did not (Figures 3D and 3E). Quantitative analyses of

poly(GP) by immunoassay also demonstrated that poly(GP) levels were greatly

decreased in $(G_4C_2)_{89}$ flies upon coexpression of *FUS*, but not *FUS-RRMmut* (Fig 3F),

indicating that FUS suppresses RAN translation from the G_4C_2 repeat RNA to reduce

DPR production. Considering that the 5′ upstream sequence of the G_4C_2 repeat in the

C9orf72 gene is reported to affect RAN translation activity (Green et al., 2017; Tabet et

al., 2018), we used flies expressing the G_4C_2 repeat sequence with the upstream intronic

sequence of the *C9orf72* gene, namely, *LDS*-$(G_4C_2)_{44}{}^{\text{GR-GFP}}$ (Goodman et al., 2019).

Since this construct has a 3′-green fluorescent protein (GFP) tag in the GR reading

frame downstream of the G_4C_2 repeat sequence, the GR-GFP fusion protein is produced

by RAN translation. We confirmed that coexpression of *FUS* significantly decreased the

expression level of GR-GFP, whereas coexpression of *FUS-RRMmut* had no effect

(Figures 3G–3I).

We further excluded the possibility that FUS directly interacts with DPRs, rather

than with G_4C_2 repeat RNA, to decrease DPR levels and exert its suppressive effects.

Using DPR-only flies expressing DPRs translated from non-G_4C_2 RNAs with

alternative codons (Mizielinska et al., 2014), we confirmed that FUS did not suppress

the eye degeneration in DPR-only flies expressing poly(GR), but rather enhanced their

phenotypes, likely due to the additive effects of FUS toxicity (Figure 3—figure

supplement 1). Together with the finding that FUS decreases not only DPR expression

but also RNA foci formation (Figures 3A and 3B), these results collectively indicate that

FUS indeed interacts with G_4C_2 repeat RNA and regulates RAN translation from G_4C_2

repeat RNA in *Drosophila* models of C9-ALS/FTD.

Reduction of endogenous *caz* expression enhances G_4C_2 repeat-induced toxicity,

RNA foci formation, and DPR aggregation

To elucidate the physiological role of FUS on RAN translation, we also

investigated the role of endogenous *caz*, a *Drosophila* homologue of *FUS*, on G_4C_2

repeat-induced toxicity in flies expressing G_4C_2 repeat RNAs. Coexpression of *caz* as

well as *FUS* suppressed eye degeneration in flies expressing $(G_4C_2)_{42 \text{ or } 89}$ (Figure 4—

figure supplement 1). These data suggest that *caz* is a functional homologue of *FUS*. In

contrast, knockdown of *caz* by RNA interference or its hemizygous deletion modestly

but significantly enhanced the eye degeneration in $(G_4C_2)_{89}$ flies (Figures 4A–4D),

indicating that reduced *caz* expression enhances G_4C_2 repeat-induced toxicity. We next

analyzed the effects of *caz* knockdown on RNA foci formation and DPR production in

flies expressing $(G_4C_2)_{89}$. FISH analysis of the salivary glands revealed that knockdown

of *caz* significantly increased the number of nuclei containing RNA foci in $(G_4C_2)_{89}$

flies (Figures 4E and 4F). We also confirmed that the expression levels of G_4C_2 repeat

RNA in $(G_4C_2)_{89}$ flies were not altered by the knockdown of *caz* (Figure 4G).

Immunohistochemical analysis showed that knockdown of *caz* significantly increased

the number of DPR aggregates in $(G_4C_2)_{89}$ flies (Figures 4H and 4I). These results

indicate that the reduction of *caz* expression enhances RNA foci formation and DPR

aggregation, compatible with the results of *FUS* coexpression in flies expressing

$(G_4C_2)_{89}$ (Figure 3), and that FUS functions as an endogenous regulator of RAN

translation.

FUS directly binds to and modulates the G-quadruplex structure of G_4C_2 repeat

RNA, resulting in the suppression of RAN translation *in vitro*

We next confirmed the direct interaction of FUS with G_4C_2 repeat RNA by the

filter binding assay. We found that His-tagged FUS binds to the $(G_4C_2)_4$ RNA in a dose-

dependent manner, but not to the control $(AAAAAA)_4$ RNA (Figure 5A), and His-

tagged FUS-RRMmut had almost no binding affinity to the $(G_4C_2)_4$ RNA, consistent

with a previous study (Mori et al., 2013a). We also confirmed the interaction of FUS

with the G_4C_2 repeat RNA in our C9-ALS/FTD flies by showing the colocalization of

FUS with the RNA foci (Figure 5—figure supplement 1), consistent with a recent study

using C9-ALS/FTD patient fibroblasts (Bajc Česnik et al., 2019). Since G_4C_2 repeat

RNA was reported to form both G-quadruplex and hairpin structures (Fratta et al., 2012;

Hauesler et al., 2014; Reddy et al., 2013; Su et al., 2014), we next characterized the

interactions of FUS with G_4C_2 repeat RNA having different structures. G_4C_2 repeat

RNA is known to form G-quadruplex structures in the presence of K^+, whereas they

form hairpin structures in the presence of Na^+ (Su et al., 2014). Surface plasmon

resonance (SPR) analyses demonstrated that FUS preferentially binds to $(G_4C_2)_4$ RNA

with the G-quadruplex structure in KCl buffer (Table 1, dissociation constant (K_D) =

1.5×10^{-8} M), and weakly to $(G_4C_2)_4$ RNA with the hairpin structure in NaCl buffer

(Table 1, $K_D = 1.3 \times 10^{-7}$ μM). We also confirmed that FUS has poor binding affinity to

$(G_4C_2)_4$ RNA in LiCl buffer (Table 1, $K_D = 1.4 \times 10^{-5}$ μM), which destabilizes the G-

quadruplex structure (Herdy et al., 1992), and was an almost similar level to its binding

affinity to the negative control $(A_4C_2)_4$ RNA (not shown). These results suggest the

preferential binding of FUS to G_4C_2 repeat RNA with the G-quadruplex structure,

which is consistent with a previous report showing preferential binding of FUS to G-

quadruplex structured Sc1 and DNMT RNAs (Ozdilek et al., 2017). Considering that

higher-order structures, including G-quadruplex and hairpin structures, are reported to

be involved in RAN translation (Mori et al., 2021; Simone et al., 2018; Wang et al.,

2019; Zu et al., 2011), we next investigated the effects of FUS on the structure of G_4C_2

repeat RNA. The circular dichroism (CD) spectrum of $(G_4C_2)_4$ RNA in KCl buffer was

found to exhibit a positive peak at approximately 260 nm and a negative peak at 240 nm

(Figure 5B, black line), consistent with previous reports (Fratta et al., 2012; Hauesler et

al., 2014; Reddy et al., 2013; Su et al., 2014). Interestingly, upon the addition of FUS,

these two peaks were notably shifted to longer wavelengths with substantial CD

spectrum changes, indicating a significant structural alteration in $(G_4C_2)_4$ RNA (Figure

5B, red line). We confirmed that the CD spectrum of FUS alone in the wavelength range

of 240 to 300 nm was almost negligible (Figure 5—figure supplement 2A, green line),

indicating that this change in CD spectrum is attributed to structural changes in the

$(G_4C_2)_4$ RNA. We also observed CD spectrum changes to some extent in the $(G_4C_2)_4$

RNA upon the addition of FUS in NaCl buffer, but not in LiCl buffer, confirming an

interaction between FUS and hairpin-structured $(G_4C_2)_4$ RNA as well (Figures 5C and

5D). We further analyzed the interaction between FUS and G_4C_2 repeat RNA by imino

proton nuclear magnetic resonance (NMR). In KCl buffer, the NMR signals of the

imino proton for the G-quadruplex structure of $(G_4C_2)_4$ RNA were detected in the region

around 10 to 12 ppm (Figure 5—figure supplement 2D), consistent with previous

studies (Fratta et al., 2012; Su et al., 2014). Upon the addition of FUS, the NMR

intensities of $(G_4C_2)_4$ RNA were decreased in an FUS concentration-dependent manner

(Figure 5—figure supplement 2D), further supporting the possibility that that FUS

interacts with and modulates the G-quadruplex structure of $(G_4C_2)_4$ RNA. Collectively,

these results indicate that FUS directly binds to G_4C_2 repeat RNA, preferentially to its

G-quadruplex form, and modulates its higher-order structures. These structural

alterations of G_4C_2 repeat RNAs by FUS did not require ATP or interactions with other

proteins, suggesting its role as an RNA chaperone for G_4C_2 repeat RNA (Rajkowitsch et

al., 2007).

To further clarify the direct link between the binding of FUS to G_4C_2 repeat

RNA and its effects on RAN translation, we employed a cell-free *in vitro* translation

assay using rabbit reticulocyte lysate. We designed a reporter construct containing the

80 G_4C_2 repeat sequence with the 5′ upstream intronic sequence of the *C9orf72* gene

and the Myc tag sequence in the GA reading frame at the 3′ downstream (Figure 5E).

This upstream sequence contained multiple stop codons in each reading frame and

lacked AUG initiation codons. We confirmed by Western blotting that this reporter

system indeed produces GA-Myc by RAN translation, consistent with previous studies

(Green et al., 2017; Tabet et al., 2018). Notably, the addition of FUS to this system

decreased the expression level of GA-Myc in a dose-dependent manner, whereas the

addition of the control bovine serum albumin (BSA) did not (Figure 5F). Taken

together, these results indicate that FUS suppresses RAN translation from G_4C_2 repeat

RNA *in vitro* as an RNA chaperone.

Identification of G-quadruplex-targeting RBPs that suppress G_4C_2 repeat-induced

toxicity in C9-ALS/FTD flies

Considering that FUS suppresses G_4C_2 repeat-induced toxicity as an RNA

chaperone through its preferential binding to the G-quadruplex structure of G_4C_2 repeat

RNA (Figure 5 and Table 1), we hypothesized that other G-quadruplex-targeting RBPs

might have similar suppressive effects on G_4C_2 repeat-induced toxicity. To investigate

this possibility, we selected 6 representative G-quadruplex-targeting RBPs, all of which

are known to bind to G_4C_2 RNA as well (Cooper-Knock et al., 2014; Haeusler et al.,

2014; Mori et al., 2013a; Xu et al., 2013) (Figure 6—source data 1). Intriguingly,

coexpression of *EWSR1*, *DDX3X*, *DDX5*, or *DDX17* significantly suppressed eye

degeneration in $(G_4C_2)_{89}$ flies without altering G_4C_2 RNA expression (Figure 6). Their

effects on G_4C_2 repeat-induced toxicity and repeat RNA expression were consistent with

those of FUS. In support of our results, DDX3X was previously reported to suppress

RAN translation and G_4C_2 repeat-induced toxicity in cell culture in a helicase-activity-

dependent manner (Cheng et al., 2019). On the other hand, coexpression of *DHX9* or

DHX36 suppressed eye degeneration by reducing G_4C_2 repeat RNA levels (Figure 6).

Since G-quadruplex-targeting RBPs have diverse biological functions, including

transcription, RNA processing, translation, and RNA stabilization (Dumas et al., 2021),

these different effects among G-quadruplex-targeting RBPs on G_4C_2 repeat RNA

expression might be attributed to their different roles in RNA metabolism. Thus, some

G-quadruplex-targeting RBPs regulate G_4C_2 repeat-induced toxicity by binding to and

possibly by modulating the G-quadruplex structure of G_4C_2 repeat RNA.

Discussion

In this study, we revealed a novel regulatory mechanism of RAN translation from expanded G_4C_2 repeat RNA by the ALS/FTD-linked RBP FUS, which suppresses DPR production and neurodegeneration in C9-ALS/FTD *Drosophila* models (Figures 1–4). FUS directly binds to G_4C_2 repeat RNA and modulates its G-quadruplex structure, as evident by CD and NMR analyses (Figure 5), suggesting its functional role as an RNA chaperone. This is reminiscent of our recent study on SCA31, in which we demonstrated a novel role of the ALS/FTD-linked RBPs TDP-43, FUS, and hnRNPA2B1 as RNA chaperones binding to UGGAA repeat RNA and altering its structure, resulting in the suppression of its neurotoxicity through reducing RNA foci formation and repeat polypeptide translation (Ishiguro et al., 2017). Considering the similarities of the effects of FUS on G_4C_2 repeat RNA and UGGAA repeat RNA, we conclude that FUS functions as an RNA chaperone also for G_4C_2 repeat RNA to regulate its RAN translation.

The suppressive effects of RBPs in several noncoding repeat expansion diseases by the amelioration of their sequestration into RNA foci have been reported. For example, in myotonic dystrophy type 1, MBNL1 was shown to be sequestered into CUG repeat RNA foci, and overexpression of MBNL1 in a mouse model was found to compensate its functional loss, resulting in the reversal of myotonia (Kanadia et al., 2006). Similarly, previous studies reported the suppressive effects of other RBPs on neurodegeneration, such as hnRNPA2B1 in fragile X ataxia/tremor syndrome (Sofora et al., 2007), and Pur-α and Zfp106 in C9-ALS/FTD (Xu et al., 2013; Celona et al., 2017). The suppressive effects of these RBPs have been thought to result from the supplementation against their loss-of-function due to their sequestration into RNA foci,

although their effects on gain-of-toxic disease pathomechanisms, that is, RAN

translation and repeat RNA expression, remain to be elucidated. In contrast, in this

study we demonstrated that FUS suppresses neurodegeneration in C9-ALS/FTD by

directly targeting G_4C_2 repeat RNA and inhibiting RAN translation as an RNA

chaperone. Similar suppressive effects of RBPs by targeting UGGAA repeat RNA in

SCA31 as RNA chaperones have also been reported (Ishiguro et al., 2017). In addition,

we also showed that the expression of IGF2BP1, hnRNPA2B1, DHX9, and DHX36

decreased G_4C_2 repeat RNA expression and suppressed eye degeneration in our C9-

ALS/FTD *Drosophila* model (Figures 1 and 6), likely via the reduction of DPR levels.

In myotonic dystrophy type 2 models, MBNL1 was also reported to retain CCUG repeat

RNA in the nucleus, resulting in the suppression of RAN translation (Zu et al., 2017),

implying various mechanisms of the effects of RBPs depending on the combination of

RBPs and repeat RNA. Nevertheless, our findings highlighted the previously

unrecognized roles of RBPs directly interacting with repeat RNA and modulating gain-

of-toxic pathomechanisms, including RAN translation in noncoding repeat expansion

diseases.

Several studies have indicated the importance of higher-order structures of repeat

RNA in RAN translation. In SCA8 models, hairpin-forming CAG repeat RNA was

shown to be RAN-translated to produce polyglutamine proteins, but switching the CAG

repeats to non-hairpin-forming CAA repeats abolished the RAN translation (Zu et al.,

2011). In C9-ALS/FTD, G_4C_2 repeat RNA has been reported to form both hairpin and

G-quadruplex structures (Fratta et al., 2012; Hauesler et al., 2014; Reddy et al., 2013;

Su et al., 2014). Although the effect of each structure on RAN translation remains

largely unknown, small molecules binding to the hairpin structure or the G-quadruplex

structure were both reported to inhibit RAN translation from the G_4C_2 repeat RNA,

resulting in reduced DPR levels (Wang et al., 2019, Mori et al., 2021). These findings

are in accordance with our results showing that FUS modifies the G-quadruplex

structure as well as the hairpin structure of G_4C_2 repeat RNA as an RNA chaperone and

reduces DPR production. We further found that G-quadruplex-targeting RNA helicases,

including DDX3X, DDX5, and DDX17, which are known to bind to G_4C_2 repeat RNA

(Cooper-Knock et al., 2014; Haeusler et al., 2014; Mori et al., 2013a; Xu et al., 2013),

also alleviate G_4C_2 repeat-induced toxicity without altering the expression levels of

G_4C_2 repeat RNA in our *Drosophila* models. These results suggest that not only ATP-

independent RNA chaperones, but also ATP-dependent RNA helicases may regulate

RAN translation through modifying the higher-order structures of template repeat RNA.

Consistently, a previous study also reported that DDX3X inhibits RAN translation from

G_4C_2 repeat RNA in a helicase activity-dependent manner (Cheng et al., 2019).

Knockdown of another RNA helicase, DHX36, has been reported to both promote

(Cheng et al., 2019) and inhibit (Liu et al., 2021; Tseng et al., 2021) RAN translation,

possibly due to the different effects on repeat RNA structures depending on the

experimental conditions. Unfortunately, most of these studies reporting the effects of

RBPs on RAN translation have limitations of the detailed structural analyses of repeat

RNA. In this study, focusing on FUS, we performed a series of molecular structural

analyses, *in vitro* translation assays, and *in vivo* genetic analyses to clarify the structure-

function relationship of G_4C_2 repeat RNA, and provide compelling evidence for the

modifying effects of FUS on repeat RNA structures leading to the suppression of RAN

translation and repeat-induced toxicity *in vivo*.

FUS has an RRM domain for RNA binding and a low complexity (LC) domain

involved in protein interactions, and exerts multifaceted functions, such as RNA

transcription, RNA splicing, RNA transport, and formation of membraneless organelles,

such as stress granules and nuclear paraspeckles via liquid-liquid phase separation

(Lagier-Tourenne et al., 2010). Recent studies reported that arginine-rich DPRs, such as

poly(GR) and poly(PR), interact with LC domain-containing RBPs, including FUS, and

alter their liquid-liquid phase separation, resulting in the disruption of the dynamics and

functions of membraneless organelles (Kwon et al., 2014; Lee et al., 2016; Lin et al.,

2016). These findings raise the possibility that FUS may exert its suppressive effects by

directly interacting with DPRs. However, we showed that FUS does not suppress eye

degeneration in DPR-only flies (Figure 3—figure supplement 1) indicating that a direct

interaction between FUS and DPRs is unlikely to be the mechanism of the suppression

of DPR toxicity in our C9-ALS/FTD flies. This result supports our conclusion that FUS

suppresses G_4C_2 repeat-induced toxicity through direct binding to G_4C_2 repeat RNA.

In summary, we here provided evidence that FUS modulates the structure of G_4C_2

repeat RNA as an RNA chaperone, and regulates RAN translation, resulting in the

suppression of neurodegeneration in C9-ALS/FTD fly models. Recent advances in

genome sequencing technology unveiled that such expansions of repeat sequences cause

more than 50 monogenic human diseases (Malik et al., 2021), and are also associated

with psychiatric diseases such as autism (Mitra et al., 2021; Trost et al., 2020). Thus,

our findings contribute not only towards the elucidation of repeat-associated pathogenic

mechanisms underlying a wider range of neuropsychiatric diseases than previously

thought, but also towards the development of potential therapies for these diseases.

Materials and Methods

Flies

All fly stocks were cultured and crossed at 23 °C or 25 °C in standard cornmeal-yeast-glucose medium. Male adult flies were used for the climbing assay and *GeneSwitch* experiments. Three to 5-day-old female adult flies were used for the evaluation of eye phenotype using a stereoscopic microscope model SZX10 (Olympus). Female third instar larvae were used for quantitative real-time polymerase chain reaction (PCR), RNA FISH and immunohistochemistry experiments. The transgenic fly line bearing the *GMR-Gal4* transgene has been described previously (Yamaguchi et al., 1999). The transgenic fly lines bearing *elav-Gal4* (#8765), *elav-GeneSwitch* (#43642), *UAS-EGFP* (#6874), *UAS-DsRed* (#6282), *UAS-GFP-IR* (inverted repeat) (#9330), $UAS\text{-}(GR)_{36}$ (#58692), $UAS\text{-}(GA)_{36}$ (#58693), $UAS\text{-}(GR)_{100}$ (#58696), $UAS\text{-}(GA)_{100}$ (#58697), and *UAS-EWSR1* (#79592) were obtained from Bloomington Drosophila Stock Center. The transgenic fly line bearing *UAS-caz-IR* (#100291) was obtained from Vienna Drosophila Resource Center. The fly line with the *caz* null allele (caz^2), $UAS\text{-}LDS\text{-}(G_4C_2)_{44}^{GR\text{-}GFP}$, and *UAS-caz* (*UAS-FLAG-caz*) and *UAS-FUS*-4 (*UAS-FLAG-FUS*) were kind gifts from Dr. Erik Storkebaum (Frickenhaus et al., 2015), Dr. Nancy Bonini (Goodman et al., 2019), and Dr. Brian McCabe (Wang et al., 2011), respectively. Other transgenic fly lines were generated in this study. Full genotypes of the fly lines used in all figures and their cultured temperatures are described in Supplementary file 1.

Generation of constructs and transgenic flies

Artificially synthesized $(G_4C_2)_{50}$ sequences flanked at the 5′ end with an *Eag*I recognition site and at the 3′ end with a *PspOM*I recognition site were subcloned into T-

vector pMD20 (Takara Bio). To generate a longer repeat size, the pMD20-$(G_4C_2)_{50}$

vector was digested with *Eag*I and *PspOM*I, followed by ligation into the pMD20-

$(G_4C_2)_{50}$ vector linearized by digestion with *Eag*I. This vector was digested with *EcoR*I

and *Hind*III, and subcloned into the pcDNA™3.1/*myc*-His(−)A vector (Thermo Fisher

Scientific). We accidentally obtained the pcDNA™3.1/*myc*-His(−)A-$(G_4C_2)_9$ vector at

this step. The pcDNA™3.1/*myc*-His(−)A-$(G_4C_2)_n$ vector was digested with *EcoR*I and

*Xba*I, and subcloned into the *Drosophila* pUAST vector. These constructs have no start

codon sequence (ATG) upstream of the G_4C_2 repeat sequence (Figure 1—figure

supplement 1A). These pUAST-$(G_4C_2)_n$ vectors were amplified with a recombinase-

mutated SURE$^{®}$2 *Escherichia coli* strain (Agilent Technologies) at 28 °C for 72 hours

to prevent repeat length contraction. The number of G_4C_2 repeats in the pUAST-$(G_4C_2)_9$

$_{or\ 50}$ vectors was determined by sequencing. To determine the number of G_4C_2 repeats in

the pUAST-$(G_4C_2)_{89}$ vector, transposable element insertional mutagenesis using EZ-

Tn5™ <KAN-2> Insertion Kit (Epicentre) and sequencing were performed.

To generate pUAST-*FUS* or pUAST-*TDP-43* vectors, the Gateway$^{®}$ Vector

Conversion System (Thermo Fisher Scientific) was used. The human *FUS* or human

TARDBP cDNA was subcloned into the pENTR™/D-TOPO$^{®}$ vector (Thermo Fisher

Scientific). To generate the Gateway destination vector pUAST-DEST, we inserted the

Gateway cassette A sequence (Thermo Fisher Scientific) into the pUAST vector. The

pUAST-*FUS* or pUAST-*TDP-43* vectors were generated using Gateway recombination

reactions (Thermo Fisher Scientific) . The FUS RRM mutant construct (pUAST-*FUS-

RRMmut*), in which leucine residues at positions 305, 341, 359, and 368 in the FUS

protein were substituted to phenylalanine, was generated by PCR and the In-Fusion$^{®}$

Cloning system (Takara Bio). To generate the other pUASTattB-*RBP* vectors, each

cDNA encoding the RBP shown in Figure 1—source data 1 and Figure 6—source data 1

was subcloned into the pUASTattB vector (VectorBuilder). To establish transgenic flies

harboring *UAS-(G_4C_2)*$_n$, *UAS-FUS*, *UAS-FUS-RRMmut*, and *UAS-TDP-43*, the pUAST-

(G_4C_2)$_n$, pUAST-*FUS*, pUAST-*FUS* line 2, pUAST-*FUS-RRMmut*, and pUAST-*TDP-*

43 vectors, respectively, were injected into fly embryos of the w^{1118} strain. To establish

transgenic flies harboring the other *UAS-RBP* constructs including *UAS-FUS* line 3,

pUASTattB-*RBP* vectors were injected into fly embryos of the attP40 strain. These

transgenic flies were established using standard methods at BestGene Inc.

The number of repeats in *UAS-(G_4C_2)*$_{9 \text{ or } 42}$ transgenic flies were determined by

genomic PCR using the forward (5′-AACCAGCAACCAAGTAAATCAAC-3′), and

reverse (5′-TGTTGAGAGTCAGCAGTAGCC-3′) primers, which amplifies a part of

the *UAS-(G_4C_2)*$_n$ sequence, including G_4C_2 repeat sequence, followed by sequencing

using the forward (5′-GCCAAGAAGTAATTATTGA-3′) and/or reverse (5′-

TCCAATTATGTCACACC-3′) primers.

Quantitative real-time PCR

Total RNA was extracted from female third instar larvae of each genotype using

TRIzol® reagent (Thermo Fisher Scientific) according to the manufacturer's

instructions. First-strand cDNA was synthesized using QuantiTect Reverse

Transcription Kit (QIAGEN). Real-time PCR was performed using SYBR® Premix Ex

Taq™ II (Takara Bio) and the Mx3000P Real-time quantitative PCR system (Agilent

Technologies) or the CFX96™ Real-Time PCR Detection System (Bio-Rad). For G_4C_2

repeat RNA quantification of flies expressing (G_4C_2)$_n$ in Figure 1—figure supplement

1C, the forward (5′-ATGAATGGGAGCAGTGGTGG-3′) and reverse (5′-

TGTTGAGAGTCAGCAGTAGCC-3′) primers were used. For G_4C_2 repeat RNA

quantification of flies expressing $(G_4C_2)_{89}$(H) and FUS, FUS-RRMmut, other RNA-

binding proteins, or *caz-IR* in Figures 1E, 3C, 4G, and 6D, the forward (5′-

CCCAATCCATATGACTAGTAGATCC-3′) and reverse (5′-

TGTAGGTAGTTTGTCCAATTATGTCA-3′) primers were used. Both of the above-

mentioned primer pairs recognize sequences downstream of the G_4C_2 repeats. For *gal4*

mRNA quantification, the forward (5′-TTGAAATCGCGTCGAAGGA-3′) and reverse

(5′-GGCTCCAATGGCTAATATGCA-3′) primers were used (Li et al., 2008). Data were

analyzed using the standard curve method. The amounts of G_4C_2 repeat transcripts were

normalized to those of *gal4* transcripts in the same sample. At least three independent

biological replicates per genotype were analyzed. Data were normalized by setting the

values of the samples from flies expressing $(G_4C_2)_{89}$(H) (Figure 1—figure supplement

1C), both $(G_4C_2)_{89}$(H) and *EGFP* (Figures 1E, 3C, and 6D), or both $(G_4C_2)_{89}$(H) and

GFP-IR (Figure 4G) as 100.

Imaging and quantification of fly eyes

Light microscopic images of the eyes of 3 to 5-day-old female flies were taken

using a stereoscopic microscope model SZX10 (Olympus) with a CCD camera DP21

(Olympus). Images shown are representative eye phenotypes of the fly crosses. Crosses

were performed three times to validate the specific phenotypes. Eye size and

pigmentation were quantified as previously reported (Saitoh et al., 2015). Five or 10

eyes per genotype were analyzed. Data were normalized by setting the values of

samples from flies expressing one copy of *EGFP* (Figure 4D), those expressing two

copies of *EGFP* (Figures 1B, 1C, 1D, 2B, 2C, 6B, 6C, Figure 1—figure supplements

2B-2D, and Figure 4—figure supplements 1B-1D), or those expressing both *EGFP* and

GFP-IR (Figure 4B), as 100.

Egg-to-adult viability of flies

Mated female flies were placed on grape juice agar with yeast paste for 24

hours. Eggs were collected from the surface of the grape juice agar, and the number of

eggs were counted and placed on new standard fly food. After eclosion, the number of

adult flies was counted. Egg-to-adult viability was calculated by dividing the number of

adult flies by the number of eggs. More than 500 eggs per genotype were used. Data

were normalized by setting the values of samples from flies expressing two copies of

EGFP as 100 (Fig 2D).

Climbing assay

Twenty male flies were gently introduced into a glass vial. After a 5-minute

adaptation period, the bottom of the vial was gently tapped and the height the flies

reached in 10 seconds was recorded using a digital video camera, and scored as follows:

0 (lower than 2 cm), 1 (from 2 to 3.9 cm), 2 (from 4 to 5.9 cm), 3 (from 6 to 7.9 cm), 4

(from 8 to 9.9 cm), and 5 (higher than 10 cm). Five trials were performed in each

experiment at intervals of 20 sec. The assay was performed between 8:00 and 10:00.

Climbing scores were calculated as an average of five trials.

GeneSwitch experiments

Flies were crossed in the absence of RU486 (mifepristone) on standard fly food.

One-day old adult male flies were transferred to Formula 4-24® Instant Drosophila

medium (Wako) with RU486 (100 µg/mL) for the indicated periods. Every 2 or 3 days,

flies were transferred to new medium with RU486. Climbing assays were performed at

0, 7, and 14 days after the start of RU486 treatment (Figure 2E).

RNA fluorescence in situ hybridization

Female third instar larvae were dissected in ice-cold phosphate-buffered saline

(PBS). Salivary glands were fixed with 4% paraformaldehyde (PFA) (pH 7.0) in PBS

for 30 min, and incubated in 100% methanol. Fixed samples were rehydrated in 75%

(v/v), 50%, and 25% ethanol in PBS, and rinsed in PBS and distilled water (DW).

Samples were then treated with 0.2 N HCl/DW for 20 min at room temperature (RT),

and rinsed in DW. Next, the samples were permeabilized with 0.2% Triton X-100 in

PBS for 10 min, rinsed in PBS for 5 min, fixed again in 4% PFA in PBS for 20 min,

then washed twice for 5 min each in PBS, and incubated twice for 15 min each in 2

mg/mL glycine/PBS. After the acetylation treatment, samples were incubated for 1 hour

at 37 °C in hybridization buffer consisting of 50% formamide, $2 \times$ saline sodium citrate

(SSC), 0.2 mg/mL yeast tRNA, and 0.5 mg/mL heparin. For hybridization, samples

were incubated overnight at 80 °C with a 5′ end Alexa594-labeled $(G_2C_4)_4$ or Alexa488-

labeled $(C_2G_4)_4$ locked nucleic acid (LNA) probe (5 nM) in hybridization buffer. These

LNA probes were synthesized by GeneDesign Inc. After the hybridization, samples

were washed once for 5 min in $4 \times$ SSC at 80 °C, three times for 20 min each in $2 \times$ SSC

and 50% formamide at 80 °C, three times for 40 min each in $0.1 \times$ SSC at 80 °C, and

once for 5 min in PBS containing 0.5% Triton X-100 (PBT) at RT. Nuclei were stained

with   4',6-diamidino-2-phenylindole   (DAPI)   or   2'-(4-ethoxyphenyl)-5-(4-methyl-1-

piperazinyl)-2,5'-bi-1-*H*-benzimidazole, trihydrochloride (Hoechst 33342). Stained

samples were mounted in SlowFade® Gold antifade reagent (Thermo Fisher Scientific)

and observed under a Zeiss LSM710 or LSM880 confocal laser-scanning microscope.

After RNA FISH, samples were scanned using a Zeiss LSM710 or LSM880

confocal laser-scanning microscope along the z-axis direction. One z-stack image was

taken per salivary gland using ZEN imaging software (Zeiss). RNA foci-positive nuclei

in more than 30 cells per salivary gland were counted, and the percentage of nuclei

containing RNA foci in the salivary gland was calculated. Ten salivary glands were

analyzed for each genotype.

Immunohistochemistry

Female third instar larvae were dissected in ice-cold PBS. Eye imaginal discs

and salivary glands were fixed with 4% PFA in PBS for 30 min and washed 3 times

with PBT. After blocking with 5% goat serum/PBT, the samples were incubated

overnight at 4 °C with a rat monoclonal anti-poly(GR) antibody (clone 5A2,

MABN778, Millipore), a rabbit polyclonal anti-poly(GA) antibody (CAC-TIP-C9-P01,

Cosmo Bio), a rabbit polyclonal anti-poly(GP) antibody (NBP2-25018, Novus

Biologicals), or a rabbit polyclonal anti-FUS antibody (A300-302A, Bethyl

Laboratories) at 1:1,000 dilution as the primary antibody. After washing 3 times with

PBT, the samples were incubated with an Alexa 633-conjugated anti-rat antibody (A-

21094, Thermo Fisher Scientific), or an Alexa 488-conjugated or Alexa 555-conjugated

anti-rabbit antibody (A-11008 or A-21428, respectively, Thermo Fisher Scientific) at

1:500 dilution as the secondary antibody. After washing 3 times with PBT, nuclei were

stained with DAPI or Hoechst 33342. Stained samples were mounted in SlowFade®

Gold antifade reagent and observed using a Zeiss LSM710 or LSM880 confocal laser-

scanning microscope.

The number of DPR aggregates in the eye discs was quantitatively measured

using ZEN imaging software (Zeiss), as follows: 1) selection of photoreceptor neurons

within the 13 developing ommatidia in rows 2 and 3 at the posterior end of the eye discs

(Saitoh et al., 2015) by DAPI or Hoechst 33342 staining, because these ommatidia are

at similar stages of development and can be easily identified; 2) counting of the number

of DPR aggregates with a diameter of larger than 2 μm in the cytoplasm. Ten to 15 eye

discs were analyzed for each genotype.

Measurement of poly(GP) protein levels

The heads of 5-day-old female flies expressing both $(G_4C_2)_{89}$(H) and either

EGFP, *FUS*, or *FUS-RRMmut* using the *GMR-Gal4* driver were collected and stored at

–80 °C. Samples were prepared using a previously reported method (Tran et al., 2015).

Poly(GP) levels were measured by a sandwich immunoassay that uses Meso Scale

Discovery electrochemiluminescence detection technology, as described previously (Su

et al., 2014). Data were normalized by setting the values of samples from flies

expressing $(G_4C_2)_{89}$(H) and *EGFP* (Figure 3F) as 100.

Western blotting

To assess the expression levels of FUS and FUS-RRMmut (Figure 2—figure

supplement 1), or GR-GFP (Figures 3G–3I), 10 heads of 5-day-old female flies

expressing *FUS* or *FUS-RRMmut*, or both *LDS-$(G_4C_2)_{44}$*[GR-GFP] and either *FUS* or *FUS-*

RRMmut using the *GMR-Gal4* driver were homogenized in 100 µL of sodium dodecyl

sulfate (SDS) sample buffer using a pestle, boiled for 5 min, and centrifuged at

$10,000 \times g$ for 3 min at 25 °C. Five µL of each supernatant were run on a 5% to 20%

gradient polyacrylamide gel (Wako) and then transferred onto an Immun-Blot®

polyvinylidene fluoride membrane (Bio-Rad). Membranes were blocked with 5% skim

milk in PBS containing 0.1% Tween-20 (PBST) or PVDF Blocking Reagent for Can

Get Signal® (TOYOBO) for 2 hours at RT, and then incubated overnight at 4 °C with a

rabbit polyclonal anti-FUS antibody (A300-302A, Bethyl Laboratories), a rat

monoclonal anti-poly(GR) antibody (clone 5A2, MABN778, Millipore), a Living

Colors® EGFP mouse monoclonal antibody (632569, Clontech), or a mouse monoclonal

anti-actin antibody (clone AC-40, A4700, Sigma-Aldrich) at 1:1,000 dilution as the

primary antibody. After washing 3 times with PBST, membranes were incubated for 2

hours at RT with either HRP-conjugated anti-rat, anti-rabbit, or anti-mouse antibody

(31470, 31460, or 31430, respectively, Invitrogen) at 1:5,000 dilution as the secondary

antibody, washed 3 times with PBST, treated with SuperSignal™ West Dura

chemiluminescent substrate (Thermo Fisher Scientific), and imaged using the

LuminoGraphII imaging system (ATTO). Data were normalized by setting the average

values of samples from flies expressing *FUS* (Figure 2—figure supplement 1B), or

those expressing both *LDS*-$(G_4C_2)_{44}^{\text{GR-GFP}}$ and *DsRed* as 100 (Figures 3H and 3I).

Filter binding assay

For preparation of FUS proteins with an N-terminal His tag (His-FUS), cDNAs

of the human FUS protein (wild type or RRM mutant) from pUAST-*FUS* or pUAST-

FUS-RRMmut were cloned into the multiple cloning site (*Xho*I and *Bam*HI) of the

plasmid vector pET-15b (Novagen) (Nomura et al., 2014). After transfection of the

plasmids into *E. coli* BL21 (DE3), the expression of His-FUS proteins was induced by

culturing the transformed cells in the presence of 0.5 mM isopropyl β-*D*-

thiogalactopyranoside (IPTG) at 20 °C for 20 hours. Cells were lysed by ultrasonication

in PBS (pH 7.4) containing 2% (v/v) Triton X-100, 1 M NaCl, DNase I, $MgSO_4$, and

ethylenediaminetetraacetic acid-free Complete Protease Inhibitor Cocktail (Roche

Applied Science). After centrifugation at $20,000 \times g$ for 30 min at 4 °C, the pellets were

redissolved in a buffer (pH 7.0) containing 6 M guanidine hydrochloride (GdnHCl), 50

mM Tris, and 1 M NaCl. His-FUS proteins in the pellets were purified by Ni^{2+} affinity

chromatography. In brief, the His-FUS proteins were mixed with Profinity IMAC Ni^{2+}-

charged resin (Bio-Rad) for 30 min at 20 °C. Then, FUS proteins bound to the resin

were washed with wash buffer (6 M GdnHCl, 50 mM Tris, and 1 M NaCl, pH 7.0), and

eluted with elution buffer (6 M GdnHCl, 50 mM Tris, 100 mM NaCl, and 250 mM

imidazole, pH 7.0). For preparation of soluble, refolded FUS, FUS proteins (200 μM) in

the elution buffer were diluted 20-fold with a buffer (pH 7.0) containing 50 mM Tris,

100 mM NaCl, 10% (v/v) glycerol, and 5 mM Tris (2-carboxyethyl) phosphine (buffer

A), which, however, produced significant amounts of precipitate. This insoluble

material was removed by centrifugation at $20,000 \times g$ for 10 min at 4 °C, resulting in the

recovery of soluble FUS proteins in the supernatant fraction. Protein concentrations

were spectroscopically determined from the absorbance at 280 nm using the following

extinction coefficients: 71,630 $cm^{-1} M^{-1}$ for both FUS and FUS-RRMmut.

Biotinylated RNAs were synthesized by GeneDesign Inc. Ten nM biotin-

$(G_4C_2)_4$, 10 nM biotin-$(AAAAAA)_4$, or 10 nM biotin-$(UUAGGG)_4$ (telomeric repeat-

containing RNA: TERRA) were incubated with soluble FUS proteins (5, 10, or 50 nM)

in buffer A with 0.4 U/μL RNase inhibitor (RNasin Plus RNase Inhibitor, Promega).

Biotinylated (AAAAAA)$_4$ and TERRA are negative and positive controls, respectively.

After an hour at RT, the mixture was filtered through a nitrocellulose membrane

(PROTRAN$^®$, 0.2 μm, Amersham Biosciences) overlaid on a nylon membrane

(Hybond-N$^+$, 0.45 μm, Schleicher & Schuell) in a 96-well slot-blot apparatus (ATTO)

(Furukawa et al., 2011). After extensive washing of the membranes with buffer A, the

bound RNAs were crosslinked to the membranes using ultraviolet radiation (254 nm;

UV Stratalinker, Stratagene) at an energy level of 0.12 J. After blocking with 3% (w/v)

BSA in tris-buffered saline with 0.1% Tween-20, the membranes were incubated with

streptavidin-HRP (1:5,000; Nacalai Tesque), and the biotinylated RNAs on the

membranes were detected with ImmunoStar LD reagent (Wako).

Preparation of recombinant FUS protein

For preparation of the FUS protein, the human *FUS* (WT) gene flanked at the 5′

end with an *Nde*I recognition site and at the 3′ end with a *Xho*I recognition site was

amplified by PCR from pUAST-*FUS*. PCR fragments were digested with *Nde*I and

*Xho*I. These fragments were ligated into the cloning sites of the plasmid vector pET-21b

(Novagen) between *Nde*I and *Xho*I. After transfection of the plasmids into *E. coli* BL21

(DE3), expression of the FUS protein was induced by culturing the transformed cells in

the presence of 0.5 mM IPTG at 37 °C for 6 hours. Cells were harvested by

centrifugation and suspended with buffer B (10% glycerol, 20 mM 4-(2-hydroxyethyl)-

1-piperazineethanesulfonic acid [HEPES]-NaOH [pH 7.0], 300 mM NaCl, 1 mM

dithiothreitol [DTT], 1 mM ethylenediaminetetraacetic acid [EDTA], 0.1% Tween-20,

and 0.1% benzamidine hydrochloride) containing 1.5 mg/mL lysozyme, and stored for

30 min on ice. Cell lysates were sonicated, and insoluble protein was collected by

centrifugation. The pellet was solubilized in buffer C (6 M urea, 10% glycerol, 20 mM

HEPES-NaOH [pH 7.0], 1 mM DTT, 1 mM EDTA, and 0.1% benzamidine

hydrochloride). After centrifugation, supernatants were loaded onto a DE52 (GE

Healthcare) open column. The flow-through fraction was loaded onto a CM52 (GE

Healthcare) open column. The flow-through fraction of DE52 was applied to a CAPTO

S column (GE Healthcare), and the flow-through fraction was collected using the

ÄKTAexplorer 10S/100 system (GE Healthcare). The flow-through fraction was applied

to a Mono S column (GE Healthcare). Proteins were fractionated with a 0 to 500 mM

linear gradient of NaCl in buffer D (6 M urea, 10% glycerol, 20 mM HEPES-NaOH

[pH 7.0], 1 mM DTT, and 1 mM EDTA) using ÄKTA explorer 10 S/100 system. The

FUS fraction was eluted at 150 to 200 mM NaCl. For refolding, the eluted peak fraction

was diluted fivefold using refolding buffer (900 mM arginine, 100 mM N-cyclohexyl-2-

hydroxyl-3-aminopropanesulfonic acid [pH 9.5], 0.3 mM reduced glutathione, 0.03 mM

oxidized glutathione, and 1 mM $ZnCl_2$), and stored overnight at RT. The solution was

concentrated using a centrifugal filter (Vivaspin 6-10 kDa; GE Healthcare) to 1 to 2

mg/mL, and then dialyzed against buffer E (10% glycerol, 20 mM HEPES-NaOH [pH

6.8], 300 mM NaCl, 0.1 mM EDTA, and 10 mM β-cyclodextrin), and stored frozen at

−80 °C.

Surface plasmon resonance analyses

The binding of FUS to $(G_4C_2)_4$ RNA was analyzed using a Biacore T200

instrument (GE Healthcare). $(G_4C_2)_4$ RNAs biotinylated at the 5′ end in 10 mM HEPES

pH 6.8 and 500 mM MCl (M = K, Na, or Li) was injected over the streptavidin-coated

surface of a sensor chip SA (GE Healthcare). The amount of immobilized RNA was as

follows: 240 resonance unit (RU) in KCl, 363 RU in NaCl, or 319 RU in LiCl buffer

condition. Binding experiments were performed using the single cycle kinetics method.

The running buffer used was 20 mM HEPES (pH 6.8), 1 mM $MgCl_2$, 0.05% Tween-20,

and 150 mM KCl, NaCl, or LiCl. FUS was diluted in the running buffer and injected

sequentially over the RNA-immobilized sensor surface in increasing concentrations

(0.016, 0.031, 0.063, 0.13, or 0.25 µM). Sensorgrams were obtained at 25 °C, 30

µL/min flow rate, 60 seconds of contact time, and 120 seconds of dissociation time.

Circular dichroism spectroscopy

CD spectra were measured at 25 °C using a spectropolarimeter model J-820

(JASCO). $(G_4C_2)_4$ RNA was synthesized by GeneDesign Inc. and dissolved in 20 mM

HEPES (pH 6.8), 18.75 mM NaCl, 10 mM $MgCl_2$, 0.625% glycerol, 0.625 mM β-

cyclodextrin, and 0.0625 mM EDTA with 150 mM KCl, NaCl, or LiCl. RNA samples

containing 150 mM KCl were first heated at 95 °C for 5 min and then cooled to RT to

form the G-quadruplex structure. The other samples were not heated. FUS (1 µM) was

added to the RNA sample (4 µM) and mixed before recording the spectrum. CD spectra

were recorded at a speed of 50 nm min^{-1} and a resolution of 1 nm, and 10 scans were

averaged.

Nuclear magnetic resonance spectroscopy

All one-dimensional ^1H NMR spectral data were recorded using AVANCE III

800 MHz NMR spectrometers equipped with a TXI cryogenic probe (Bruker BioSpin)

at 25 °C. $(G_4C_2)_4$ RNA dissolved in 20 mM HEPES (pH 6.8), 150 mM KCl, 18.75 mM

NaCl, 10 mM $MgCl_2$, 0.625% glycerol, 0.625 mM β-cyclodextrin, and 0.0625 mM

EDTA was first heated at 95 °C for 5 min and then cooled to room temperature to form

the G-quadruplex structure. The RNA (10 μM) was mixed with FUS at molar ratios

(RNA:FUS) of 1:0, 1:0.2, 1:0.4, and 0:1. The samples were then prepared at a final

concentration of 10% D_2O before recording their spectra. ^1H NMR data were acquired

using simple single 90° hard-pulse excitations following solvent signal suppression with

a jump-and-return pulse scheme. Free induction decay data (1,600 points in total) were

collected by repeating the scans (2,600 times) with an interscan delay of 2.5 seconds.

All NMR data were processed using Topspin 3.6 software (Bruker BioSpin).

RNA synthesis for in vitro translation

For preparation of the C9-RAN reporter plasmid, the pEF6-*C9orf72* intron1-

$(G_4C_2)_{80}$ vector was digested with *HindIII* and *NotI* to obtain the fragment *C9orf72*

intron1-$(G_4C_2)_{80}$, and subcloned into the pcDNA™5/FRT vector (Thermo Fisher

Scientific). To add the T7 promoter upstream of the *C9orf72* intron 1 sequence in this

pcDNA™5/FRT-*C9orf72* intron1-$(G_4C_2)_{80}$ vector, a forward primer including T7

promoter sequences with the 5′-terminal region of *C9orf72* intron 1 flanked at the 5′ end

with an *HindIII* recognition site, and a reverse primer recognizing the 3′-terminal region

of *C9orf72* intron 1 sequences including a *BssHII* recognition site were designed, and

used to amplify a fragment containing *C9orf72* intron 1 with a T7 promoter by PCR.

Then, this fragment was subcloned into the pcDNA™5/FRT-*C9orf72* intron1-$(G_4C_2)_{80}$

vector digested by *HindIII* and *BssHII*. In addition to the T7 promoter, the Myc tag in

the GA frame downstream of $(G_4C_2)_{80}$ was introduced into this vector.

The reporter plasmids were linearized with *XbaI*. Linearized DNA was *in vitro*

transcribed using mMESSAGE mMACHINE™ T7 Transcription Kit (Invitrogen)

according to the manufacturer's instructions. T7 reactions were carried out at 37 °C for

2 hours, treated with TURBO™ DNaseI for 15 min at 37 °C to remove the DNA

template, and then polyadenylated with *E. coli* Poly-A Polymerase (NEB) for 1 hour at

37 °C. Synthesized mRNAs were purified by LiCl precipitation. The size and quality of

the synthesized mRNAs were verified on a denaturing RNA gel.

In vitro translation assay

mRNAs of *C9orf72* intron1-$(G_4C_2)_{80}$ with a Myc tag in the GA frame were *in*

vitro translated with Flexi® Rabbit Reticulocyte Lysate System (Promega) according to

the manufacturer's instructions. Translation reactions were performed with 10 ng/μL

mRNA and contained 30% rabbit reticulocyte lysate, 10 μM amino-acid mix minus

methionine, 10 μM amino-acid mix minus leucine, 0.5 mM MgOAc, 100 mM KCl, 0.8

U/μL Murine RNAse Inhibitor (NEB), and 100 ng/μL human tRNAs purified from

HeLa cells. FUS or BSA at each concentration (10, 100, and 1,000 μM) was added for

translation in the lysate. Samples were incubated at 30 °C for 90 min before termination

by incubation on ice. Ten μL of samples were analyzed by 13% SDS-polyacrylamide

gel electrophoresis and Western blotting using a mouse monoclonal anti-Myc tag

antibody (clone 4A6, Sigma-Aldrich) as the primary antibody.

Quantification and statistical analysis

Statistical parameters including the definitions and exact values of n (e.g.,

number of experiments, number of flies, number of eye imaginal discs, etc.),

distributions, and deviations are stated in the figures and corresponding figure legends.

Multiple comparison tests using one-way ANOVA with Tukey's post hoc analysis were

performed for Figures 1B–1E, 2B–2D, 3B, 3C, 3E, 3F, 3H, 3I, 4B, 4D, 6B, 6C, 6D,

Figure 1—figure supplements 1C and 1D, Figure 1—figure supplements 2B–2D, and

Figure 4—figure supplements 1B–1D, multiple comparison test using two-way repeated

measures ANOVA with Tukey's post hoc analysis was performed for Figure 2E, and the

unpaired t-test was used for Figures 4F, 4G, 4I, and Figure 2—figure supplement 1B.

Differences in means were considered statistically significant at $P < 0.05$. All statistical

analyses were performed using GraphPad Prism 8 software.

As the sample sizes used in the present study were similar to previous

publications (Freibaum et al., 2015; Goodman et al., 2019; Mizielinska et al., 2014; Xu

et al., 2013), statistical analyses were performed afterwards without interim data

analysis. Data were not excluded, and were collected and processed randomly. Sample

collection and analyses for the measurement of poly(GP) protein levels were performed

in a double-blind manner. Data collection and analyses for other experiments were not

performed in a blind manner regarding the conditions of the experiments.

Acknowledgements

We thank Drs E. Storkebaum (Radboud University), N. Bonini (University of Pennsylvania), and B. McCabe (École Polytechnique Fédérale de Lausanne) for kindly providing the *caz* mutant, *UAS-LDS-(G₄C₂)₄₄*$^{GR-GFP}$, and *UAS-FLAG-caz* and *UAS-FLAG-FUS* flies, respectively. We acknowledge Bloomington Drosophila Stock Center and Vienna Drosophila Resource Center for providing various fly stocks. We thank Drs H. Imataka and K. Machida (University of Hyogo) for kindly providing human tRNAs for *in vitro* translation. We also thank the members of our laboratory for helpful discussions, K. Yamamoto, E. Wakisaka, T. Yamashita, and A. Sugai for their technical assistance, Dr. H. Akiko Popiel for critical reading and English editing of the manuscript, and the Center for Medical Research and Education, Graduate School of Medicine, Osaka University for supplying experimental equipment. This work was supported by Grants-in-Aid for Scientific Research on Innovative Areas (Brain Protein Aging and Dementia Control) (17H05699 to Y.N. and 17H05705 to Y. Furukawa), Transformative Research Areas (A) (Multifaceted Proteins) (20H05927 to Y.N. and K.M.), and Strategic Research Program for Brain Sciences (11013026 to Y.N.), from the Ministry of Education, Culture, Sports, Science, and Technology, Japan; by Grants in-Aid for Scientific Research (B) (21H02840 to Y.N. and 20H03602 to K.M.), Scientific Research (C) (15K09331 and 19K07823 to M.U., 17K07291 to A.I.), Young Scientists (A) (17H05091 to K.M.), Young Scientists (B) (25860733 to M.U.), and Challenging Exploratory Research (24659438 to Y.N. and 18K19515 to K.M.) from the Japan Society for the Promotion of Science, Japan; by a Health Labor Sciences Research Grant for Research on Development of New Drugs (H24-Soyaku-Sogo-002 to Y.N.) from the Ministry of Health, Labor and Welfare, Japan; by grants for Strategic Research

Program for Brain Sciences (JP15dm0107026 and JP20dm0107061 to Y.N.), Practical

Research Projects for Rare/Intractable Diseases (JP16ek0109018, JP19ek0109222, and

JP20ek0109316 to Y.N.), and Platform Project for Supporting Drug Discovery and Life

Science Research [Basis for Supporting Innovative Drug Discovery and Life Science

Research] (JP19am0101072 to T.F.) from Japan Agency for Medical Research and

Development, Japan; Intramural Research Grants for Neurological and Psychiatric

Disorders (27-7, 27-9, 30-3, 30-9, and 3-9 to Y.N.) from the National Center of

Neurology and Psychiatry; an IBC Grant (H28 to Y.N.) from the Japan Amyotrophic

Lateral Sclerosis Association; grants from Takeda Science Foundation (2017 to M.U.

and 2016 to K.M.); and a grant from SENSHIN Medical Research Foundation (2018 to

K.M.).

References

869  Ash, P. E., Bieniek, K. F., Gendron, T. F., Caulfield, T., Lin, W. L., Dejesus-
870 Hernandez, M., van Blitterswijk, M. M., Jansen-West, K., Paul, J. W., 3rd, Rademakers,
871 R., Boylan, K. B., Dickson, D. W., and Petrucelli, L. (2013). Unconventional translation
872 of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to
873 c9FTD/ALS. Neuron *77*, 639–646. https://doi.org/10.1016/j.neuron.2013.02.004

875  Bajc Česnik, A., Darovic, S., Prpar Mihevc, S., Štalekar, M., Malnar, M.,
876 Motaln, H., Lee, Y. B., Mazej, J., Pohleven, J., Grosch, M., Modic, M., Fonovič, M.,
877 Turk, B., Drukker, M., Shaw, C. E., and Rogelj, B. (2019). Nuclear RNA foci from
878 C9ORF72 expansion mutation form paraspeckle-like bodies. J Cell Sci. *132*, jcs224303.
879 https://doi.org/10.1242/jcs.224303

881  Boivin, M., Pfister, V., Gaucherot, A., Ruffenach, F., Negroni, L., Sellier, C., and
882 Charlet-Berguerand, N. (2020). Reduced autophagy upon C9ORF72 loss synergizes
883 with dipeptide repeat protein toxicity in G4C2 repeat expansion disorders. EMBO J. *39*,
884 e100574. https://doi.org/10.15252/embj.2018100574

886  Celona, B., Dollen, J. V., Vatsavayai, S. C., Kashima, R., Johnson, J. R., Tang,
A. A., Hata, A., Miller, B. L., Huang, E. J., Krogan, N. J., Seeley, W. W., and Black, B.
L. (2017). Suppression of *C9orf72* RNA repeat-induced neurotoxicity by the ALS-
associated RNA-binding protein Zfp106. eLife *6*, e19032.
https://doi.org/10.7554/eLife.19032

Cheng, W., Wang, S., Mestre, A. A., Fu, C., Makarem, A., Xian, F., Hayes, L. R.,

Lopez-Gonzalez, R., Drenner, K., Jiang, J., Cleveland, D. W., and Sun, S. (2018).

C9ORF72 GGGGCC repeat-associated non-AUG translation is upregulated by stress

through eIF2α phosphorylation. Nat. Commun. *9*, 51. https://doi.org/10.1038/s41467-

017-02495-z

Cheng, W., Wang, S., Zhang, Z., Morgens, D. W., Hayes, L. R., Lee, S., Portz,

B., Xie, Y., Nguyen, B. V., Haney, M. S., Yan, S., Dong, D., Coyne, A. N., Yang, J.,

Xian, F., Cleveland, D. W., Qiu, Z., Rothstein, J. D., Shorter, J., Gao, F. B., Bassik,

M.C., and Sun, S. (2019). CRISPR-Cas9 Screens Identify the RNA Helicase DDX3X as

a Repressor of C9ORF72 (GGGGCC)n Repeat-Associated Non-AUG Translation.

Neuron *104*, 885–898.e8. https://doi.org/10.1016/j.neuron.2019.09.003

Choi, S. Y., Lopez-Gonzalez, R., Krishnan, G., Phillips, H. L., Li, A. N., Seeley,

W. W., Yao, W. D., Almeida, S., and Gao, F. B. (2019). C9ORF72-ALS/FTD-associated

poly(GR) binds Atp5a1 and compromises mitochondrial function in vivo. Nat.

Neurosci. *22*, 851–862. https://doi.org/10.1038/s41593-019-0397-0

Conlon, E. G., Lu, L., Sharma, A., Yamazaki, T., Tang, T., Shneider, N. A., and

Manley, J. L. (2016). The C9ORF72 GGGGCC expansion forms RNA G-quadruplex

inclusions and sequesters hnRNP H to disrupt splicing in ALS brains. eLife *5*, e17820.

https://doi.org/10.7554/eLife.17820

Cooper-Knock, J., Walsh, M. J., Higginbottom, A., Robin Highley, J., Dickman,

M. J., Edbauer, D., Ince, P. G., Wharton, S. B., Wilson, S. A., Kirby, J., Hautbergue, G.

M., and Shaw, P. J. (2014). Sequestration of multiple RNA recognition motif-containing

proteins by C9orf72 repeat expansions. Brain *137*, 2040–2051.

https://doi.org/10.1093/brain/awu120

Daigle, J. G., Lanson, N. A., Jr, Smith, R. B., Casci, I., Maltare, A., Monaghan,

J., Nichols, C. D., Kryndushkin, D., Shewmaker, F., and Pandey, U. B. (2013). RNA-

binding ability of FUS regulates neurodegeneration, cytoplasmic mislocalization and

incorporation into stress granules associated with FUS carrying ALS-linked mutations.

Hum. Mol. Genet. *22*, 1193–1205. https://doi.org/10.1093/hmg/dds526

DeJesus-Hernandez, M., Mackenzie, I. R., Boeve, B. F., Boxer, A. L., Baker, M.,

Rutherford, N. J., Nicholson, A. M., Finch, N. A., Flynn, H., Adamson, J., Kouri, N.,

Wojtas, A., Sengdy, P., Hsiung, G. Y., Karydas, A., Seeley, W. W., Josephs, K. A.,

Coppola, G., Geschwind, D. H., Wszolek, Z. K., Feldman, H., Knopman, D.S.,

Petersen, R.C., Miller, B. L., Dickson, D. W., Boylan, K. B., Graff-Radford, N. R., and

Rademakers, R. (2011). Expanded GGGGCC hexanucleotide repeat in noncoding

region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron*, *72*(2), 245–

256. https://doi.org/10.1016/j.neuron.2011.09.011 (2011). Expanded GGGGCC

hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked

FTD and ALS. Neuron *72*, 245–256.

Donnelly, C. J., Zhang, P. W., Pham, J. T., Haeusler, A. R., Mistry, N. A.,

Vidensky, S., Daley, E. L., Poth, E. M., Hoover, B., Fines, D. M., Maragakis, N.,

Tienari, P. J., Petrucelli, L., Traynor, B. J., Wang, J., Rigo, F., Bennett, C. F., Blackshaw,

S., Sattler, R., and Rothstein, J. D. (2013). RNA toxicity from the ALS/FTD C9ORF72

expansion is mitigated by antisense intervention. Neuron *80*, 415–428.

https://doi.org/10.1016/j.neuron.2013.10.015

Dumas, L., Herviou, P., Dassi, E., Cammas, A., and Millevoi, S. (2021). G-

Quadruplexes in RNA Biology: Recent Advances and Future Directions. Trends

Biochem. Sci. *46*, 270–283. https://doi.org/10.1016/j.tibs.2020.11.001

Fratta, P., Mizielinska, S., Nicoll, A. J., Zloh, M., Fisher, E. M., Parkinson, G.,

and Isaacs, A. M. (2012). C9orf72 hexanucleotide repeat associated with amyotrophic

lateral sclerosis and frontotemporal dementia forms RNA G-quadruplexes. Sci. Rep. *2*,

1016. https://doi.org/10.1038/srep01016

Freibaum, B. D., Lu, Y., Lopez-Gonzalez, R., Kim, N. C., Almeida, S., Lee, K.

H., Badders, N., Valentine, M., Miller, B. L., Wong, P. C., Petrucelli, L., Kim, H. J.,

Gao, F. B., and Taylor, J. P. (2015). GGGGCC repeat expansion in C9orf72

compromises nucleocytoplasmic transport. Nature *525*, 129–133.

https://doi.org/10.1038/nature14974

Frickenhaus, M., Wagner, M., Mallik, M., Catinozzi, M., and Storkebaum, E.

(2015). Highly efficient cell-type-specific gene inactivation reveals a key function for

the Drosophila FUS homolog cabeza in neurons. Sci. Rep. *5*, 9107.

https://doi.org/10.1038/srep09107

Furukawa, Y., Kaneko, K., Watanabe, S., Yamanaka, K., and Nukina, N. (2011).

A seeding reaction recapitulates intracellular formation of Sarkosyl-insoluble

transactivation response element (TAR) DNA-binding protein-43 inclusions. J. Biol.

Chem. *286*, 18664–18672. https://doi.org/10.1074/jbc.M111.231209

Gendron, T. F., Bieniek, K. F., Zhang, Y. J., Jansen-West, K., Ash, P. E.,

Caulfield, T., Daughrity, L., Dunmore, J. H., Castanedes-Casey, M., Chew, J., et al.

(2013). Antisense transcripts of the expanded C9ORF72 hexanucleotide repeat form

nuclear RNA foci and undergo repeat-associated non-ATG translation in c9FTD/ALS.

Acta Neuropathol. *126*, 829–844. https://doi.org/10.1007/s00401-013-1192-8

Gendron, T. F., Bieniek, K. F., Zhang, Y. J., Jansen-West, K., Ash, P. E.,

Caulfield, T., Daughrity, L., Dunmore, J. H., Castanedes-Casey, M., Chew, J., Cosio, D.

M., van Blitterswijk, M., Lee, W. C., Rademakers, R., Boylan, K. B., Dickson, D. W.,

and Petrucelli, L. (2017). Poly(GP) proteins are a useful pharmacodynamic marker for

C9ORF72-associated amyotrophic lateral sclerosis. Sci, Transl. Med. *9*, eaai7866.

https://doi.org/10.1126/scitranslmed.aai7866

Gijselinck, I., Van Langenhove, T., van der Zee, J., Sleegers, K., Philtjens, S.,

Kleinberger, G., Janssens, J., Bettens, K., Van Cauwenberghe, C., Pereson, S.,

Engelborghs, S., Sieben, A., De Jonghe, P., Vandenberghe, R., Santens, P., De Bleecker,

J., Maes, G., Bäumer, V., Dillen, L., Joris, G., Cuijt, I., Corsmit, E., Elinck E., Van

Dongen, J., Vermeulen, S., Van den Broeck, M., Vaerenberg, C., Mattheijssens, M.,

Peeters, K., Robberecht, W., Cras, P., Martin, J. J., De Deyn P. P., Cruts, M., and Van

Broeckhoven, C. (2012). A C9orf72 promoter repeat expansion in a Flanders-Belgian

cohort with disorders of the frontotemporal lobar degeneration-amyotrophic lateral

sclerosis spectrum: a gene identification study. Lancet Neurol. *11*, 54–65.

Goodman, L. D., Prudencio, M., Kramer, N. J., Martinez-Ramirez, L. F.,

Srinivasan, A. R., Lan, M., Parisi, M. J., Zhu, Y., Chew, J., Cook, C. N., Berson, A.,

Gitler, A. D., Petrucelli, L., and Bonini, N. M. (2019). Toxic expanded GGGGCC repeat

transcription is mediated by the PAF1 complex in C9orf72-associated FTD. Nat.

Neurosci. *22*, 863–874. https://doi.org/10.1038/s41593-019-0396-1

Green, K. M., Glineburg, M. R., Kearse, M. G., Flores, B. N., Linsalata, A. E.,

Fedak, S. J., Goldstrohm, A. C., Barmada, S. J., and Todd, P. K. (2017). RAN

translation at C9orf72-associated repeat expansions is selectively enhanced by the

integrated stress response. Nat. Commun. *8*, 2005. https://doi.org/10.1038/s41467-017-

02200-0

1004

1005   Haeusler, A. R., Donnelly, C. J., Periz, G., Simko, E. A., Shaw, P. G., Kim, M.

1006 S., Maragakis, N. J., Troncoso, J. C., Pandey, A., Sattler, R., Rothstein, J. D., and Wang,

1007 J. (2014). C9orf72 nucleotide repeat structures initiate molecular cascades of disease.

1008 Nature *507*, 195–200. https://doi.org/10.1038/nature13124

1009

1010   Hardin, C. C., Watson, T., Corregan, M., and Bailey, C. (1992). Cation-

1011   dependent transition between the quadruplex and Watson-Crick hairpin forms of

1012   d(CGCG3GCG). Biochemistry *31*, 833–841. https://doi.org/10.1021/bi00118a028

1013

1014   Hutten, S., Usluer, S., Bourgeois, B., Simonetti, F., Odeh, H. M., Fare, C. M.,

1015   Czuppa, M., Hruska-Plochan, M., Hofweber, M., Polymenidou, M., Shorter, J.,

1016   Edbauer, D., Madl, T., and Dormann, D. (2020). Nuclear Import Receptors Directly

1017   Bind to Arginine-Rich Dipeptide Repeat Proteins and Suppress Their Pathological

1018   Interactions. Cell Rep. *33*, 108538. https://doi.org/10.1016/j.celrep.2020.108538

1019

Ishiguro, T., Sato, N., Ueyama, M., Fujikake, N., Sellier, C., Kanegami, A.,

Tokuda, E., Zamiri, B., Gall-Duncan, T., Mirceta, M., Furukawa, Y., Yokota, T., Wada,

K., Taylor, J. P., Pearson, C. E., Charlet-Berguerand, N., Mizusawa, H., Nagai, Y., and

Ishikawa, K. (2017). Regulatory Role of RNA Chaperone TDP-43 for RNA Misfolding

and Repeat-Associated Translation in SCA31. Neuron *94*, 108–124.e7.

https://doi.org/10.1016/j.neuron.2017.02.046

1026

1027   Jovičić, A., Mertens, J., Boeynaems, S., Bogaert, E., Chai, N., Yamada, S. B.,

1028   Paul, J. W., 3rd, Sun, S., Herdy, J. R., Bieri, G., Kramer, N. J., Gage, F. H., Van Den

1029   Bosch, L., Robberecht, W., and Gitler, A. D. (2015). Modifiers of C9orf72 dipeptide

1030   repeat toxicity connect nucleocytoplasmic transport defects to FTD/ALS. Nat. Neurosci.

1031   *18*, 1226–1229. https://doi.org/10.1038/nn.4085

1032

1033   Krishnan, G., Raitcheva, D., Bartlett, D., Prudencio, M., McKenna-Yasek, D.

1034   M., Douthwright, C., Oskarsson, B. E., Ladha, S., King, O. D., Barmada, S. J., Miller, T.

M., Bowser, R., Watts, J. K., Petrucelli, L., Brown, R. H., Kankel, M. W., and Gao, F. B. (2022). Poly(GR) and poly(GA) in cerebrospinal fluid as potential biomarkers for C9ORF72-ALS/FTD. Nat. Commun. *13*, 2799. https://doi.org/10.1038/s41467-022-30387-4

Kanadia, R. N., Shin, J., Yuan, Y., Beattie, S. G., Wheeler, T. M., Thornton, C. A., and Swanson, M. S. (2006). Reversal of RNA missplicing and myotonia after muscleblind overexpression in a mouse poly(CUG) model for myotonic dystrophy. Proc. Natl. Acad. Sci. USA *103*, 11748–11753. https://doi.org/10.1073/pnas.0604970103

Kwan, T., and Thompson, S. R. (2019). Noncanonical Translation Initiation in Eukaryotes. Cold Spring Harb. Perspect. Biol. *11*, a032672. https://doi.org/10.1101/cshperspect.a032672

Kwon, I., Xiang, S., Kato, M., Wu, L., Theodoropoulos, P., Wang, T., Kim, J., Yun, J., Xie, Y., and McKnight, S. L. (2014). Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede RNA biogenesis, and kill cells. Science *345*, 1139–1145. https://doi.org/10.1126/science.1254917

Lagier-Tourenne, C., Polymenidou, M., and Cleveland, D. W. (2010). TDP-43 and FUS/TLS: emerging roles in RNA processing and neurodegeneration. Hum. Mol. Genet. *19*, R46–R64. https://doi.org/10.1093/hmg/ddq137

1059   Lee, K. H., Zhang, P., Kim, H. J., Mitrea, D. M., Sarkar, M., Freibaum, B. D.,

1060 Cika, J., Coughlin, M., Messing, J., Molliex, A., Maxwell, B. A., Kim, N. C., Temirov,

1061 J., Moore, J., Kolaitis, R. M., Shaw, T. I., Bai, B., Peng, J., Kriwacki, R. W., and Taylor,

1062 J. P. (2016). C9orf72 Dipeptide Repeats Impair the Assembly, Dynamics, and Function

1063 of  Membrane-Less  Organelles.  Cell *167*,  774–788.e17.

1064 https://doi.org/10.1016/j.cell.2016.10.002

1065

1066   Lee, Y. B., Chen, H. J., Peres, J. N., Gomez-Deza, J., Attig, J., Stalekar, M.,

1067 Troakes, C., Nishimura, A. L., Scotter, E. L., Vance, C., Adachi, Y., Sardone, V., Miller,

1068 J. W., Smith, B. N., Gallo, J. M., Ule, J., Hirth, F., Rogelj, B., Houart, C., and Shaw, C.

1069 E. (2013). Hexanucleotide repeats in ALS/FTD form length-dependent RNA foci,

1070 sequester RNA binding proteins, and are neurotoxic. Cell Rep. *5*, 1178–1186.

1071 https://doi.org/10.1016/j.celrep.2013.10.049

1072

1073   Lehmer, C., Oeckl, P., Weishaupt, J. H., Volk, A. E., Diehl-Schmid, J., Schroeter,

1074 M. L., Lauer, M., Kornhuber, J., Levin, J., Fassbender, K., Landwehrmeyer, B., Schludi,

1075 M. H., Arzberger, T., Kremmer, E., Flatley, A., Feederle, R., Steinacker, P., Weydt, P.,

1076 Ludolph, A. C., Edbauer, D., and Otto, M. German Consortium for Frontotemporal

1077 Lobar Degeneration (2017). Poly-GP in cerebrospinal fluid links C9orf72-associated

1078 dipeptide repeat expression to the asymptomatic phase of ALS/FTD. EMBO Mol. Med.

1079 *9*, 859–868. https://doi.org/10.15252/emmm.201607486

1080

1081   Li, L. B., Yu, Z., Teng, X., and Bonini, N. M. (2008). RNA toxicity is a

1082 component of ataxin-3 degeneration in Drosophila. Nature *453*, 1107–1111.

1083   https://doi.org/10.1038/nature06909

1084

1085   Lin, Y., Mori, E., Kato, M., Xiang, S., Wu, L., Kwon, I., and McKnight, S. L.

1086   (2016). Toxic PR Poly-Dipeptides Encoded by the C9orf72 Repeat Expansion Target

1087   LC   Domain   Polymers.   Cell   *167*,   789–802.e12.

1088   https://doi.org/10.1016/j.cell.2016.10.003

1089

1090   Liu, H., Lu, Y. N., Paul, T., Periz, G., Banco, M. T., Ferré-D'Amaré, A. R.,

1091   Rothstein, J. D., Hayes, L. R., Myong, S., and Wang, J. (2021). A Helicase Unwinds

1092   Hexanucleotide Repeat RNA G-Quadruplexes and Facilitates Repeat-Associated Non-

1093   AUG   Translation.   J.   Am.   Chem.   Soc.   *143*,   7368–7379.

1094   https://doi.org/10.1021/jacs.1c00131

1095

1096   Maharana, S., Wang, J., Papadopoulos, D. K., Richter, D., Pozniakovsky, A.,

1097   Poser, I., Bickle, M., Rizk, S., Guillén-Boixet, J., Franzmann, T. M., Jahnel, M.,

1098   Marrone, L., Chang, Y. T., Sterneckert, J., Tomancak, P., Hyman, A. A., and Alberti, S.

1099   (2018). RNA buffers the phase separation behavior of prion-like RNA binding proteins.

1100   Science *360*, 918–921. https://doi.org/10.1126/science.aar7366

1101

1102   Malik, I., Kelley, C. P., Wang, E. T., and Todd, P. K. (2021). Molecular

1103   mechanisms underlying nucleotide repeat expansion disorders. Nat. Rev. Mol. Cell Biol.

1104   *22*, 589–607. https://doi.org/10.1038/s41580-021-00382-6

1105

1106   Mann, J. R., Gleixner, A. M., Mauna, J. C., Gomes, E., DeChellis-Marks, M. R.,

1107   Needham, P. G., Copley, K. E., Hurtle, B., Portz, B., Pyles, N. J., Guo, L., Calder, C. B.,

1108   Wills, Z. P., Pandey, U. B., Kofler, J. K., Brodsky, J. L., Thathiah, A., Shorter, J., and

1109   Donnelly, C. J. (2019). RNA Binding Antagonizes Neurotoxic Phase Transitions of

1110   TDP-43. Neuron *102*, 321–338.e8. https://doi.org/10.1016/j.neuron.2019.01.048

1111

1112   May, S., Hornburg, D., Schludi, M. H., Arzberger, T., Rentzsch, K., Schwenk, B.

1113   M., Grässer, F. A., Mori, K., Kremmer, E., Banzhaf-Strathmann, J., Mann, M.,

1114   Meissner, F., and Edbauer, D. (2014). C9orf72 FTLD/ALS-associated Gly-Ala

1115   dipeptide repeat proteins cause neuronal toxicity and Unc119 sequestration. Acta

1116   Neuropathol. *128*, 485–503. https://doi.org/10.1007/s00401-014-1329-4

1117

1118   Mitra, I., Huang, B., Mousavi, N., Ma, N., Lamkin, M., Yanicky, R., Shleizer-

1119   Burko, S., Lohmueller, K. E., and Gymrek, M. (2021). Patterns of de novo tandem

1120   repeat mutations and their role in autism. Nature *589*, 246–250.

1121   https://doi.org/10.1038/s41586-020-03078-7

1122

Mizielinska, S., Grönke, S., Niccoli, T., Ridler, C. E., Clayton, E. L., Devoy, A.,

Moens, T., Norona, F. E., Woollacott, I., Pietrzyk, J., Cleverley, K., Nicoll, A. J.,

Pickering-Brown, S., Dols, J., Cabecinha, M., Hendrich, O., Fratta, P., Fisher, E.,

Partridge, L., and Isaacs, A. M. (2014). C9orf72 repeat expansions cause

neurodegeneration in Drosophila through arginine-rich proteins. Science *345*, 1192–

1194. https://doi.org/10.1126/science.1256800

1129

1130   (Mori et al., 2013a)

Mori, K., Lammich, S., Mackenzie, I. R., Forné, I., Zilow, S., Kretzschmar, H., Edbauer, D., Janssens, J., Kleinberger, G., Cruts, M., Herms, J., Neumann, M., Van Broeckhoven, C., Arzberger, T., and Haass, C. (2013). hnRNP A3 binds to GGGGCC repeats and is a constituent of p62-positive/TDP43-negative inclusions in the hippocampus of patients with C9orf72 mutations. Acta Neuropathol. *125*, 413–423. https://doi.org/10.1007/s00401-013-1088-7

(Mori et al., 2013b)

Mori, K., Arzberger, T., Grässer, F. A., Gijselinck, I., May, S., Rentzsch, K., Weng, S. M., Schludi, M. H., van der Zee, J., Cruts, M., Van Broeckhoven, C., Kremmer, E., Kretzschmar, H. A., Haass, C., and Edbauer, D. (2013). Bidirectional transcripts of the expanded C9orf72 hexanucleotide repeat are translated into aggregating dipeptide repeat proteins. Acta Neuropathol. *126*, 881–893. https://doi.org/10.1007/s00401-013-1189-3

(Mori et al., 2013c)

Mori, K., Weng, S. M., Arzberger, T., May, S., Rentzsch, K., Kremmer, E., Schmid, B., Kretzschmar, H. A., Cruts, M., Van Broeckhoven, C., Haass, C., and Edbauer, D. (2013). The C9orf72 GGGGCC repeat is translated into aggregating dipeptide-repeat proteins in FTLD/ALS. Science *339*, 1335–1338. https://doi.org/10.1126/science.1232927

Mori, K., Gotoh, S., Yamashita, T., Uozumi, R., Kawabe, Y., Tagami, S., Kamp, F., Nuscher, B., Edbauer, D., Haass, C., Nagai, Y., and Ikeda, M. (2021). The porphyrin

1155   TMPyP4 inhibits elongation during the noncanonical translation of the FTLD/ALS-

1156   associated GGGGCC repeat in the C9orf72 gene. J. Biol. Chem. *297*, 101120.

1157   https://doi.org/10.1016/j.jbc.2021.101120

1158

1159   Nomura, T., Watanabe, S., Kaneko, K., Yamanaka, K., Nukina, N., and

1160   Furukawa, Y. (2014). Intranuclear aggregation of mutant FUS/TLS as a molecular

1161   pathomechanism of amyotrophic lateral sclerosis. J. Biol. Chem. *289*, 1192–1202.

1162   https://doi.org/10.1074/jbc.M113.516492

1163

1164   Ozdilek, B. A., Thompson, V. F., Ahmed, N. S., White, C. I., Batey, R. T., and

1165   Schwartz, J. C. (2017). Intrinsically disordered RGG/RG domains mediate degenerate

1166   specificity   in   RNA   binding.   Nucleic   Acids   Res.   *45*,   7984–7996.

1167   https://doi.org/10.1093/nar/gkx460

1168

1169   Rajkowitsch, L., Chen, D., Stampfl, S., Semrad, K., Waldsich, C., Mayer, O.,

1170   Jantsch, M. F., Konrat, R., Bläsi, U., and Schroeder, R. (2007). RNA chaperones, RNA

1171   annealers   and   RNA   helicases.   RNA   Biol.   *4*,   118–130.

1172   https://doi.org/10.4161/rna.4.3.5445

1173

1174   Reddy, K., Zamiri, B., Stanley, S., Macgregor, R. B., Jr, and Pearson, C. E.

1175   (2013). The disease-associated r(GGGGCC)n repeat from the C9orf72 gene forms tract

1176   length-dependent uni- and multimolecular RNA G-quadruplex structures. J. Biol. Chem.

1177   *288*, 9860–9866. https://doi.org/10.1074/jbc.C113.452532

1178

Renton, A.E., Majounie, E., Waite, A., Simón-Sánchez, J., Rollinson, S., Gibbs,

J. R., Schymick, J. C., Laaksovirta, H., van Swieten, J. C., Myllykangas, L., Kalimo, H.,

Paetau, A., Abramzon, Y., Remes, A. M., Kaganovich, A., Scholz, S. W., Duckworth, J.,

Ding, J., Harmer, D. W., Hernandez, D. G., Johnson. J. O., Mok, K., Ryten, M.,

Trabzuni, D., Guerreiro, R. J., Orrell, R. W., Neal, J., Murray, A., Pearson, J., Jansen, I.

E., Sondervan, D., Seelaar, H., Blake, D., Young, K., Halliwell, N., Callister, J. B.,

Toulson, G., Richardson, A., Gerhard, A., Snowden, J., Mann, D., Neary, D., Nalls, M.

A., Peuralinna, T., Jansson, L., Isoviita, V. M., Kaivorinne, A. L., Hölttä-Vuori, M.,

Ikonen, E., Sulkava, R., Benatar, M., Wuu, J., Chiò, A., Restagno, G., Borghero, G.,

Sabatelli, M., Heckerman, D., Rogaeva, E., Zinman, L., Rothstein, J. D., Sendtner, M.,

Drepper, C., Eichler, E. E., Alkan, C., Abdullaev, Z., Pack, S. D., Dutra, A., Pak, E.,

Hardy, J., Singleton, A., Williams, N.M., Heutink, P., Pickering-Brown, S., Morris, H.

R., Tienari, P. J., and Traynor, B. J.. ITALSGEN Consortium (2011). A hexanucleotide

repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD.

Neuron *72*, 257–268.

1194

1195   Rudich, P., Snoznik, C., Watkins, S. C., Monaghan, J., Pandey, U. B., and

1196   Lamitina, S. T. (2017). Nuclear localized C9orf72-associated arginine-containing

1197   dipeptides exhibit age-dependent toxicity in C. elegans. Hum. Mol. Genet. *26*, 4916–

1198   4928. https://doi.org/10.1093/hmg/ddx372

1199

1200   Saitoh, Y., Fujikake, N., Okamoto, Y., Popiel, H. A., Hatanaka, Y., Ueyama, M.,

1201   Suzuki, M., Gaumer, S., Murata, M., Wada, K., and Nagai, Y. (2015). p62 plays a

1202   protective role in the autophagic degradation of polyglutamine protein oligomers in

1203 polyglutamine disease model flies. J. Biol. Chem. *290*, 1442–1453.

1204 https://doi.org/10.1074/jbc.M114.590281

1205

Simone, R., Balendra, R., Moens, T. G., Preza, E., Wilson, K. M., Heslegrave,

A., Woodling, N. S., Niccoli, T., Gilbert-Jaramillo, J., Abdelkarim, S., Clayton, E. L.,

Clarke, M., Konrad, M. T., Nicoll, A. J., Mitchell, J. S., Calvo, A., Chio, A., Houlden,

H., Polke, J. M., Ismail, M. A., Stephens, C.E., Vo, T., Farahat, A. A., Wilson, W. D.,

Boykin, D. W., Zetterberg, H., Partridge, L., Wray, S., Parkinson, G., Neidle, S., Patani,

R., Fratta, P., and Isaacs, A. M. (2018). G-quadruplex-binding small molecules

ameliorate C9orf72 FTD/ALS pathology in vitro and in vivo. EMBO Mol. Med. *10*,

22–31. https://doi.org/10.15252/emmm.201707850

1214

Shi, Y., Lin, S., Staats, K. A., Li, Y., Chang, W. H., Hung, S. T., Hendricks, E.,

Linares, G. R., Wang, Y., Son, E. Y., Wen, X., Kisler, K., Wilkinson, B., Menendez, L.,

Sugawara, T., Woolwine, P., Huang, M., Cowan, M. J., Ge, B., Koutsodendris, N.,

Sandor K. P., Komberg, J., Vangoor, V. R., Senthilkumar, K., Hennes, V., Seah, C.,

Nelson, A. R., Cheng, T. Y., Lee, S. J., August, P. R., Chen, J. A., Wisniewski, N.,

Hanson-Smith, V., Belgard, T. G., Zhang, A., Coba, M., Grunseich, C., Ward, M. E., van

den Berg, L. H., Pasterkamp, R. J., Trotti, D., Zlokovic, B. V., and Ichida, J. K. (2018).

Haploinsufficiency leads to neurodegeneration in C9ORF72 ALS/FTD human induced

motor neurons. Nat. Med. *24*, 313–325. https://doi.org/10.1038/nm.4490

1224

1225   Sofola, O. A., Jin, P., Qin, Y., Duan, R., Liu, H., de Haro, M., Nelson, D. L., and

1226 Botas, J. (2007). RNA-binding proteins hnRNP A2/B1 and CUGBP1 suppress fragile X

1227   CGG premutation repeat-induced neurodegeneration in a Drosophila model of FXTAS.

1228   Neuron *55*, 565–571. https://doi.org/10.1016/j.neuron.2007.07.021

1229

1230   Sonobe, Y., Ghadge, G., Masaki, K., Sendoel, A., Fuchs, E., and Roos, R. P.

1231   (2018). Translation of dipeptide repeat proteins from the C9ORF72 expanded repeat is

1232   associated with cellular stress. Neurobiol. Dis. *116*, 155–165.

1233   https://doi.org/10.1016/j.nbd.2018.05.009

1234

Su, Z., Zhang, Y., Gendron, T. F., Bauer, P. O., Chew, J., Yang, W. Y., Fostvedt,

E., Jansen-West, K., Belzil, V. V., Desaro, P., Johnston, A., Overstreet, K., Oh, S. Y.,

Todd, P. K., Berry, J. D., Cudkowicz, M. E., Boeve, B. F., Dickson, D., Floeter, M. K.,

Traynor, B. J., Morelli, C., Ratti, A., Silani, V., Rademakers, R., Brown, R. H.,

Rothstein, J. D., Boylan, K. B., Petrucelli, L., and Disney, M. D. (2014). Discovery of a

biomarker and lead small molecules to target r(GGGGCC)-associated defects in

c9FTD/ALS. Neuron *83*, 1043–1050. https://doi.org/10.1016/j.neuron.2014.07.041

1242

1243   Tabet, R., Schaeffer, L., Freyermuth, F., Jambeau, M., Workman, M., Lee, C. Z.,

1244   Lin, C. C., Jiang, J., Jansen-West, K., Abou-Hamdan, H., Désaubry, L., Gendron, T.,

1245   Petrucelli, L., Martin, F., and Lagier-Tourenne, C. (2018). CUG initiation and

1246   frameshifting enable production of dipeptide repeat proteins from ALS/FTD C9ORF72

1247   transcripts. Nat. Commun. *9*, 152. https://doi.org/10.1038/s41467-017-02643-5

1248

1249   Tran, H., Almeida, S., Moore, J., Gendron, T. F., Chalasani, U., Lu, Y., Du, X.,

1250   Nickerson, J. A., Petrucelli, L., Weng, Z., and Gao, F. B. (2015). Differential Toxicity of

Nuclear RNA Foci versus Dipeptide Repeat Proteins in a Drosophila Model of C9ORF72 FTD/ALS. Neuron *87*, 1207–1214. https://doi.org/10.1016/j.neuron.2015.09.015

Trost, B., Engchuan, W., Nguyen, C. M., Thiruvahindrapuram, B., Dolzhenko, E., Backstrom, I., Mirceta, M., Mojarad, B. A., Yin, Y., Dov, A., Chandrakumar, I., Prasolava, T., Shum, N., Hamdan, O., Pellecchia, G., Howe, J. L., Whitney, J., Klee, E. W., Baheti, S., Amaral, D. G., Anagnostou, E., Elsabbagh, M., Fernandez, B. A., Hoang, N., Lewis, S., Liu, X., Sjaarda, C., Smith, I. M., Szatmari, P., Zwaigenbaum, L., Glazer, D., Hartley, D., Stewart, A. K., Eberle, M. A., Sato, N., Pearson, C.E., Scherer, S.W., and Yuen, R. (2020). Genome-wide detection of tandem DNA repeats that are expanded in autism. Nature *586*, 80–86. https://doi.org/10.1038/s41586-020-2579-z

Tseng, Y. J., Sandwith, S. N., Green, K. M., Chambers, A. E., Krans, A., Raimer, H. M., Sharlow, M. E., Reisinger, M. A., Richardson, A. E., Routh, E. D., Smaldino, M. A., Wang, Y. H., Vaughn, J. P., Todd, P. K., and Smaldino, P. J. (2021). The RNA helicase DHX36-G4R1 modulates C9orf72 GGGGCC hexanucleotide repeat-associated translation. J. Biol. Chem. *297*, 100914. https://doi.org/10.1016/j.jbc.2021.100914

Waite, A. J., Bäumer, D., East, S., Neal, J., Morris, H. R., Ansorge, O., and Blake, D. J. (2014). Reduced C9orf72 protein levels in frontal cortex of amyotrophic lateral sclerosis and frontotemporal degeneration brain with the C9ORF72 hexanucleotide repeat expansion. Neurobiol. Aging *35*, 1779.e5–1779.e13. https://doi.org/10.1016/j.neurobiolaging.2014.01.016

Wang, J. W., Brent, J. R., Tomlinson, A., Shneider, N. A., and McCabe, B. D. (2011). The ALS-associated proteins FUS and TDP-43 function together to affect Drosophila locomotion and life span. J. Clin. Invest. *121*, 4118–4126. https://doi.org/10.1172/JCI57883

Wang, Z. F., Ursu, A., Childs-Disney, J. L., Guertler, R., Yang, W. Y., Bernat, V., Rzuczek, S. G., Fuerst, R., Zhang, Y. J., Gendron, T. F., Yildirim, I., Dwyer, B. G., Rice, J. E., Petrucelli, L., and Disney, M. D. (2019). The Hairpin Form of r(G4C2)^{exp} in c9ALS/FTD Is Repeat-Associated Non-ATG Translated and a Target for Bioactive Small Molecules. Cell Chem. Biol. *26*, 179–190.e12. https://doi.org/10.1016/j.chembiol.2018.10.018

Wen, X., Tan, W., Westergard, T., Krishnamurthy, K., Markandaiah, S. S., Shi, Y., Lin, S., Shneider, N. A., Monaghan, J., Pandey, U. B., Pasinelli, P., Ichida, J. K., and Trotti, D. (2014). Antisense proline-arginine RAN dipeptides linked to C9ORF72-ALS/FTD form toxic nuclear aggregates that initiate in vitro and in vivo neuronal death. Neuron *84*, 1213–1225. https://doi.org/10.1016/j.neuron.2014.12.010

Xu, Z., Poidevin, M., Li, X., Li, Y., Shu, L., Nelson, D. L., Li, H., Hales, C. M., Gearing, M., Wingo, T. S., and Jin, P. (2013). Expanded GGGGCC repeat RNA associated with amyotrophic lateral sclerosis and frontotemporal dementia causes neurodegeneration. Proc. Natl. Acad. Sci. USA *110*, 7778–7783. https://doi.org/10.1073/pnas.1219643110

1299

1300   Yamaguchi, M., Hirose, F., Inoue, Y. H., Shiraki, M., Hayashi, Y., Nishi, Y., and

1301 Matsukage, A. (1999). Ectopic expression of human p53 inhibits entry into S phase and

1302 induces apoptosis in the Drosophila eye imaginal disc. Oncogene *18*, 6767–6775.

1303 https://doi.org/10.1038/sj.onc.1203113

1304

Zhang, Y. J., Gendron, T. F., Grima, J. C., Sasaguri, H., Jansen-West, K., Xu, Y.

F., Katzman, R. B., Gass, J., Murray, M. E., Shinohara, M., Lin, W. L., Garrett, A.,

Stankowski, J. N., Daughrity, L., Tong, J., Perkerson, E. A., Yue, M., Chew, J.,

Castanedes-Casey, M., Kurti, A., Wang, Z. S., Liesinger, A. M., Baker, J. D., Jiang, J.,

Lagier-Tourenne, C., Edbauer, D., Cleveland, D. W., Rademakers, R., Boylan, K. B.,

Bu, G., Link, C. D., Dickey, C. A., Rothstein, J. D., Dickson, D. W., Fryer, J. D., and

Petrucelli, L. (2016). C9ORF72 poly(GA) aggregates sequester and impair HR23 and

nucleocytoplasmic   transport   proteins.   Nat.   Neurosci.   *19*,   668–677.

https://doi.org/10.1038/nn.4272

1314

Zhang, Y. J., Gendron, T. F., Ebbert, M., O'Raw, A. D., Yue, M., Jansen-West,

K., Zhang, X., Prudencio, M., Chew, J., Cook, C. N., Daughrity, L. M., Tong, J., Song,

Y., Pickles, S. R., Castanedes-Casey, M., Kurti, A., Rademakers, R., Oskarsson, B.,

Dickson, D. W., Hu, W., Gitler, A. D., Fryer, J. D., and Petrucelli, L. (2018). Poly(GR)

impairs protein translation and stress granule dynamics in C9orf72-associated

frontotemporal dementia and amyotrophic lateral sclerosis. Nat. Med. *24*, 1136–1142.

https://doi.org/10.1038/s41591-018-0071-1

1322

Zhu, Q., Jiang, J., Gendron, T. F., McAlonis-Downes, M., Jiang, L., Taylor, A.,

Diaz Garcia, S., Ghosh Dastidar, S., Rodriguez, M. J., King, P., Zhang, Y., La Spada, A.

R., Xu, H., Petrucelli, L., Ravits, J., Da Cruz, S., Lagier-Tourenne, C., and Cleveland,

D. W. (2020). Reduced C9ORF72 function exacerbates gain of toxicity from ALS/FTD-

causing repeat expansion in C9orf72. Nat. Neurosci. *23*, 615–624.

https://doi.org/10.1038/s41593-020-0619-5

1329

Zu, T., Gibbens, B., Doty, N. S., Gomes-Pereira, M., Huguet, A., Stone, M. D.,

Margolis, J., Peterson, M., Markowski, T. W., Ingram, M. A., Nan, Z., Forster, C., Low,

W. C., Schoser, B., Somia, N. V., Clark, H. B., Schmechel, S., Bitterman, P. B.,

Gourdon, G., Swanson, M. S., Moseley, M., and Ranum, L. P. (2011). Non-ATG-

initiated translation directed by microsatellite expansions. Proc. Natl. Acad. Sci. USA

108, 260–265. https://doi.org/10.1073/pnas.1013343108

1336

1337   Zu, T., Liu, Y., Bañez-Coronel, M., Reid, T., Pletnikova, O., Lewis, J., Miller, T.

1338   M., Harms, M. B., Falchook, A. E., Subramony, S. H., Ostrow, L. W., Rothstein, J. D.,

1339   Troncoso, J. C., and Ranum, L. P. (2013). RAN proteins and RNA foci from antisense

1340   transcripts in C9ORF72 ALS and frontotemporal dementia. Proc. Natl. Acad. Sci. USA

1341   *110*, E4968–E4977. https://doi.org/10.1073/pnas.1315438110

1342

1343   Zu, T., Cleary, J. D., Liu, Y., Bañez-Coronel, M., Bubenik, J. L., Ayhan, F.,

1344   Ashizawa, T., Xia, G., Clark, H. B., Yachnis, A. T., Swanson, M. S., and Ranum, L.

1345   (2017). RAN Translation Regulated by Muscleblind Proteins in Myotonic Dystrophy

1346   Type 2. Neuron *95*, 1292–1305.e5. https://doi.org/10.1016/j.neuron.2017.08.039

1347

 Figure legends

Figure 1. Screening for RBPs that suppress G_4C_2 repeat-induced toxicity in C9-

ALS/FTD flies.

(A) Light microscopic images of the eyes in flies expressing both $(G_4C_2)_{42 \text{ or } 89}$ and the

indicated RBPs using the *GMR-Gal4* driver. Coexpression of *FUS*, *IGF2BP1*, or

hnRNPA2B suppressed eye degeneration in both $(G_4C_2)_{42}$ and $(G_4C_2)_{89}$ flies, indicated by

"Suppression (strong)". Coexpression of *hnRNPR*, *SAFB2*, *SF3B3*, *hnRNPA1*, or

hnRNPL suppressed eye degeneration in either $(G_4C_2)_{42}$ or $(G_4C_2)_{89}$ flies, indicated by

"Suppression (weak)" (see also Figure 1—source data 2). Scale bar: 100 μm. (B)

Quantification of eye size in $(G_4C_2)_{89}$ flies coexpressing the indicated RBPs (n = 5). (C),

(D) Quantification of eye pigmentation in $(G_4C_2)_{89}$ flies (C) or $(G_4C_2)_{42}$ flies (D)

coexpressing the indicated RBPs (n = 5). (E) Expression levels of $(G_4C_2)_{89}$ RNA in flies

expressing both $(G_4C_2)_{89}$ and the indicated RBPs using the *GMR-Gal4* driver (five

independent experiments, n = 25 flies per genotype). The $(G_4C_2)_{89}$(H) fly line expresses

$(G_4C_2)_{89}$ RNA at a high level (see also Figure 1—figure supplement 1). In (B–E), data

are presented as the mean ± SEM; $P < 0.0001$, as assessed by one-way ANOVA; n.s.:

not significant, $^*P < 0.05$, $^{**}P < 0.01$, and $^{***}P < 0.001$, as assessed by Tukey's post hoc

analysis.

1366

Figure 1–figure supplement 1. Characterization of C9-ALS/FTD flies.

(A) $(G_4C_2)_n$ constructs used in this study. These constructs do not include an ATG start

codon downstream of the UAS sequence, and were expressed in a tissue-specific

manner using the *GAL4-UAS* system. (B) Light microscopic images of the eyes in flies

expressing $(G_4C_2)_n$ using the *GMR-Gal4* driver. Scale bar: 100 μm. (C) Expression

levels of $(G_4C_2)_n$ RNAs in flies expressing $(G_4C_2)_n$ using the *GMR-Gal4* driver. Strong

eye degeneration with decreased eye size and loss of pigmentation was observed in

$(G_4C_2)_{42 \text{ or } 89}$ flies, but not in $(G_4C_2)_9$ flies. Eye degeneration was confirmed in $(G_4C_2)_{42}$

and two $(G_4C_2)_{89}$ independent fly lines. Degree of eye degeneration in two $(G_4C_2)_{89}$ fly

lines was expression-level dependent [(L) vs (H) in $(G_4C_2)_{89}$] (three independent

experiments, n = 15 flies per each genotype). Expression of G_4C_2 repeat RNA of the

sense transcripts but not that of the antisense transcripts was confirmed. (D) Climbing

ability at 1 day of age in flies expressing $(G_4C_2)_n$ using the *elav-Gal4* driver. Flies

expressing $(G_4C_2)_{89}$(H) in neurons showed lethality. Decreasing climbing ability was

observed in $(G_4C_2)_{42 \text{ or } 89}$ flies compared with $(G_4C_2)_9$ flies (five independent

experiments, n = 100 flies per each genotype). (E) FISH analyses of G_4C_2 repeat RNA

in the salivary glands of fly larvae with two copies of *GMR-Gal4* and $(G_4C_2)_{9 \text{ or } 89}$ (red:

G_4C_2 RNA; yellow: G_2C_4 RNA; blue [DAPI]: nuclei). RNA foci formation

(arrowheads) of the sense transcripts but not that of the antisense transcripts was

confirmed. RNA foci were observed in $(G_4C_2)_{89}$ flies, but not in $(G_4C_2)_9$ flies. Scale

bars: 100 μm (low magnification) or 20 μm (high magnification). (F)

Immunohistochemical analyses of DPRs stained with anti-DPR antibodies in the eye

imaginal discs of fly larvae with two copies of *GMR-Gal4* and $(G_4C_2)_{9 \text{ or } 89}$ (magenta:

poly(GR); orange: poly(GA); green: poly(GP); blue [DAPI]: nuclei). Expression and

cytoplasmic aggregation (arrowheads) of three DPRs in $(G_4C_2)_{89}$ flies, but not in

$(G_4C_2)_9$ flies, were confirmed. Scale bars: 50 μm (low magnification), 10 μm (middle

magnification), and 5 μm (high magnification). In (B–F), L: low-expression line; H:

high-expression line. In (C) and (D), data are presented as the mean ± SEM; $P < 0.0001$,

as assessed by one-way ANOVA; n.s.: not significant, and [***]$P < 0.001$, as assessed by

Tukey's post hoc analysis.

1397

Figure 1–figure supplement 2. Coexpression of FUS suppresses G_4C_2 repeat-

induced toxicity in flies expressing $(G_4C_2)_{89}$.

(A) Light microscopic images of the eyes in flies expressing both $(G_4C_2)_{42\ or\ 89}$ and *FUS*

using the *GMR-Gal4* driver. *FUS-2* and *FUS-3* are different strains from that in Figure

1. Scale bar: 100 μm. (B) Quantification of the eye size in $(G_4C_2)_{89}$ flies of the indicated

genotypes (n = 10). (C), (D) Quantification of eye pigmentation in $(G_4C_2)_{89}$ flies (C) or

$(G_4C_2)_{42}$ flies (D) of the indicated genotypes (n = 10). In (B–D), data are presented as

the mean ± SEM; $P < 0.0001$, as assessed by one-way ANOVA; [***]$P < 0.001$, as

assessed by Tukey's post hoc analysis.

1407

Figure 2. FUS suppresses G_4C_2 repeat-induced toxicity via its RNA-binding

activity.

(A) Light microscopic images of the eyes in flies expressing both $(G_4C_2)_{89}$ and either

FUS or *FUS-RRMmut* using the *GMR-Gal4* driver. Scale bar: 100 μm. (B)

Quantification of eye size in the flies of the indicated genotypes (n = 10). (C)

Quantification of eye pigmentation in the flies of the indicated genotypes (n = 10). (D)

Egg-to-adult viability in flies expressing both $(G_4C_2)_{42}$ and either *FUS* or *FUS-RRMmut*

using the *GMR-Gal4* driver (> 500 flies per genotype). (E) Climbing ability in flies

expressing both $(G_4C_2)_{42}$ and *FUS* using the *elav-GeneSwitch* driver (five independent

experiments, n = 100 flies per each genotype). In (B–E), data are presented as the mean

± SEM. In (B) and (C), $P < 0.0001$, as assessed by one-way ANOVA; n.s.: not

significant, and [***]$P < 0.001$, as assessed by Tukey's post hoc analysis. In (D), n.s.: not

significant and $^{***}P < 0.001$, as assessed by Tukey's multiple comparison test using

wholly significant difference. In (E), n.s.: not significant, $^{*}P < 0.05$, $^{**}P < 0.01$ and

$^{***}P < 0.001$, as assessed by two-way repeated measures ANOVA with Tukey's post hoc

analysis.

1424

Figure 2–figure supplement 1. Western blot analysis showing expression levels of

FUS and FUS-RRMmut proteins.

(A) Western blot analysis of the FUS and FUS-RRMmut proteins in the heads of adult

flies expressing *EGFP, FUS* or *FUS-RRMmut* using the *GMR-Gal4* driver, with an anti-

FUS antibody. The arrowhead indicates bands from the FUS and FUS-RRMmut

proteins, whereas the asterisk indicates bands resulting from nonspecific antibody

binding. (B) Quantification of the FUS and FUS-RRMmut proteins from the Western

blot analysis in (A) (n = 3).

In (B), data are presented as the mean ± SEM; n.s.: not significant, as assessed by the

unpaired *t*-test.

1435

Figure 3. FUS suppresses RNA foci formation and RAN translation from G_4C_2

repeat RNA.

(A) FISH analyses of G_4C_2 repeat RNA in the salivary glands of fly larvae expressing

both $(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using two copies of the *GMR-Gal4*

driver (red: G_4C_2 RNA; blue [DAPI]: nuclei). Arrowheads indicate RNA foci. Scale bar:

20 μm. (B) Quantification of the number of nuclei containing RNA foci from the FISH

analyses in (A) (n = 10). (C) Expression levels of $(G_4C_2)_{89}$ RNA in fly larvae expressing

both $(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using the *GMR-Gal4* driver (10

independent experiments, n = 50 flies per each genotype). (D) Immunohistochemical

analyses of DPRs stained with anti-DPR antibodies in the eye imaginal discs of fly

larvae expressing both $(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using two copies of

the *GMR-Gal4* driver (magenta: poly(GR); orange: poly(GA); green: poly(GP); blue

[DAPI]: nuclei). Arrowheads indicate cytoplasmic aggregates. Scale bars: 20 μm (low

magnification) or 5 μm (high magnification). (E) Quantification of the number of DPR

aggregates from the immunohistochemical analyses in (D) [n = 14 or 15 (GR), or 10

(GA or GP)]. (F) Immunoassay to determine poly(GP) levels in flies expressing both

$(G_4C_2)_{89}$ and either *FUS* or *FUS-RRMmut* using the *GMR-Gal4* driver (three

independent experiments, n = 30 flies per each genotype). (G) Western blot analysis of

the heads of adult flies expressing both LDS-$(G_4C_2)_{44}^{\text{GR-GFP}}$ and any of *DsRed*, *FUS* or

FUS-RRMmut using the *GMR-Gal4* driver, using either an anti-GFP (upper panel) or

anti-GR antibody (middle panel). (H and I) Quantification of GR-GFP protein levels

from the Western blot analysis in (G) (nine independent experiments, n = 90 flies per

each genotype). In (B), (C), (E), (F), (H), and (I), data are presented as the mean $\pm$

SEM. In (B), (E), and (F), $P < 0.0001$, as assessed by one-way ANOVA; n.s.: not

significant, $^*P < 0.05$, $^{**}P < 0.01$, and $^{***}P < 0.001$, as assessed by Tukey's post hoc

analysis. In (C), $P = 0.452$, as assessed by one-way ANOVA; n.s.: not significant, as

assessed by Tukey's post hoc analysis. In (H), $P = 0.0148$, as assessed by one-way

ANOVA; n.s.: not significant and $^*P < 0.05$, as assessed by Tukey's post hoc analysis. In

(I), $P = 0.0072$, as assessed by one-way ANOVA; n.s.: not significant and $^*P < 0.05$, as

assessed by Tukey's post hoc analysis.

1466

Figure 3–figure supplement 1. Overexpression of *FUS* does not suppress eye

degeneration in DPR-only flies expressing DPRs translated from non-G_4C_2 RNAs.

Light microscopic images of the eyes in DPR-only flies coexpressing either the

poly(GR) or poly(GA) protein, and either *FUS* or *FUS-RRMmut* using the *GMR-Gal4*

driver. Overexpression of *FUS* did not suppress the eye degeneration in flies expressing

either $(GR)_{36}$ or $(GR)_{100}$. Overexpression of *FUS* also caused mild eye degeneration in

flies expressing EGFP, $(GA)_{36}$, or $(GA)_{100}$, likely due to FUS toxicity.

1474

Figure 4. Reduction of endogenous *caz* expression enhances G_4C_2 repeat-induced

toxicity, RNA foci formation, and DPR aggregation.

(A) Light microscopic images of the eyes in flies expressing $(G_4C_2)_{89}$ using the *GMR-*

Gal4 driver, with knockdown of *caz*. Scale bar: 100 μm. (B) Quantification of eye size

in flies of the indicated genotypes shown in (A) (n = 10). (C) Light microscopic images

of the eyes in flies expressing $(G_4C_2)_{89}$ using the *GMR-Gal4* driver, with a hemizygous

deletion of *caz*. Scale bar: 100 μm. (D) Quantification of eye size in the flies of the

indicated genotypes shown in (C) (n = 10). (E) FISH analyses of G_4C_2 repeat RNA in

the salivary glands of fly larvae expressing $(G_4C_2)_{89}$ using the *GMR-Gal4* driver, with

knockdown of *caz* (red: G_4C_2 RNA; blue [DAPI]: nuclei). Arrowheads indicate RNA

foci. Scale bar: 20 μm. (F) Quantification of the number of nuclei containing RNA foci

from the FISH analyses in (E) (n = 10). (G) Expression levels of $(G_4C_2)_{89}$ RNA in fly

larvae expressing $(G_4C_2)_{89}$ using the *GMR-Gal4* driver, with knockdown of *caz* (four

independent experiments, n = 20 flies per each genotype). (H) Immunohistochemical

analyses of DPRs stained with anti-DPR antibodies in the eye imaginal discs of fly

larvae expressing $(G_4C_2)_{89}$ using two copies of the *GMR-Gal4* driver, with the

knockdown of *caz*. (magenta: poly(GR); orange: poly(GA); green: poly(GP); blue

[DAPI]: nuclei). Arrowheads indicate cytoplasmic aggregates. Scale bars: 20 μm (low

magnification) or 5 μm (high magnification). (I) Quantification of the number of DPR

aggregates from the immunohistochemical analyses in (H) (n = 10). In (B), (D), (F), (G)

and (I), data are presented as the mean ± SEM. In (B) and (D), $P < 0.0001$, as assessed

by one-way ANOVA; [***]$P < 0.001$, as assessed by Tukey's post hoc analysis. In (F), (G),

and (I), n.s.: not significant, [*]$P < 0.05$, [**]$P < 0.01$, and [***]$P < 0.001$, as assessed by the

unpaired t-test.

1499

Figure 4–figure supplement 1. Endogenous *caz* is a functional homologue of *FUS*

for the suppression of G_4C_2 repeat-induced toxicity.

(A) Light microscopic images of the eyes in flies expressing both $(G_4C_2)_{42 \text{ or } 89}$ and

either *caz* (*FLAG-caz*) or *FUS-4* (*FLAG-FUS*) using the *GMR-Gal4* driver. *FUS-4* is a

different strain from those used in Figure 1 and Figure 1–figure supplement 2. Scale

bar: 100 μm. (B) Quantification of the eye size in $(G_4C_2)_{89}$ flies of the indicated

genotypes (n = 10). (C), (D) Quantification of eye pigmentation in $(G_4C_2)_{89}$ flies (C) or

$(G_4C_2)_{42}$ flies (D) of the indicated genotypes (n = 10). In (B–D), data are presented as

the mean ± SEM; $P < 0.0001$, as assessed by one-way ANOVA; n.s.: not significant,

[*]$P < 0.05$, [**]$P < 0.01$,and [***]$P < 0.001$, as assessed by Tukey's post hoc analysis.

1510

Figure 5. FUS directly binds to and modulates the G-quadruplex structure of G_4C_2

repeat RNA, resulting in the suppression of RAN translation *in vitro*

(A) Analysis of the binding of His-tagged FUS proteins to biotinylated $(G_4C_2)_4$ RNA by

the filter binding assay. The nitrocellulose membrane (left) traps RNA-bound FUS

proteins, whereas unbound RNAs are recovered on the nylon membrane (right), and

then the RNAs trapped on each of the membranes was probed with streptavidin-

horseradish peroxidase (HRP). Biotinylated $(AAAAAA)_4$ and $(UUAGGG)_4$ were used

as negative and positive controls, respectively. (B–D) CD spectra of $(G_4C_2)_4$ RNA

incubated with or without FUS in the presence of 150 mM KCl (B), NaCl (C), or LiCl

(D). The CD spectrum of FUS alone was subtracted from that of $(G_4C_2)_4$ RNA

incubated with FUS. The original data are shown in Figure 5—figure supplements 2B-

2D. (E) Schema of the template RNA containing the $(G_4C_2)_{80}$ sequence and 113

nucleotides of the 5′-flanking region of intron 1 of the human *C9orf72* G_4C_2 repeat

sequence. A Myc tag in the GA frame was introduced downstream of the $(G_4C_2)_{80}$ repeat

sequence. (F) Western blot analysis of samples from *in vitro* translation using rabbit

reticulocyte lysate in the presence or absence of increasing concentrations of FUS or

BSA. The GA-Myc fusion protein was detected by Western blotting using the anti-Myc

antibody. Quantifications are shown below each band, using the condition in the

absence of each protein as a reference.

1530

1531   **Figure 5–figure supplement 1. FUS colocalizes with G_4C_2 RNA foci.**

1532   Combined FISH and immunohistochemical analyses of G_4C_2 repeat RNA and FUS in

1533   the salivary glands of flies expressing both $(G_4C_2)_{89}$ and *FUS* using two copies of the

1534   *GMR-Gal4* driver. Arrows indicate colocalization of FUS with RNA foci. Scale bar: 10

1535   μm.

1536

Figure 5–figure supplement 2. FUS modulates the G-quadruplex structure of G_4C_2

repeat RNA.

(A–C) CD spectra of $(G_4C_2)_4$ RNA incubated with or without FUS in the presence of

150 mM KCl (B), NaCl (C), or LiCl (D). CD spectra of $(G_4C_2)_4$ RNA alone (black),

FUS alone (green), sum of $(G_4C_2)_4$ RNA and FUS (blue), and the spectra of their

coincubation (magenta) are shown. FUS interacts with $(G_4C_2)_4$ RNA under KCl or NaCl

buffer conditions. (D) Imino proton NMR spectra of $(G_4C_2)_4$ RNA incubated with

increasing amounts of FUS in the presence of 150 mM KCl.

1545

Figure 6. Identification of G-quadruplex-targeting RBPs that suppress G_4C_2

repeat-induced toxicity in C9-ALS/FTD flies.

(A) Light microscopic images of eyes in flies expressing both $(G_4C_2)_{89}$ and the indicated

G-quadruplex-targeting RBPs using the *GMR-Gal4* driver. Scale bar: 100 μm. (B)

Quantification of eye size in the flies of the indicated genotypes (n = 10). (C)

Quantification of eye pigmentation in the flies of the indicated genotypes (n = 10). (D)

Expression levels of $(G_4C_2)_{89}$ RNA in flies expressing both $(G_4C_2)_{89}$ and the indicated

G-quadruplex-targeting RBPs using the *GMR-Gal4* driver (five independent

experiments, n = 25 flies per each genotype). In (B–D), data are presented as the mean ±

SEM; $P < 0.0001$, as assessed by one-way ANOVA; n.s.: not significant, $^*P < 0.05$, and

$^{***}P < 0.001$, as assessed by Tukey's post hoc analysis.

1557

1558 **Tables**

1559 **Table 1. Association (k_a) and dissociation (k_d) rate and dissociation constants (K_D)**

1560 **between FUS and $(G_4C_2)_4$ RNA in different buffers as assessed by SPR analysis.**

1561

Buffer	k_a (M^{-1}s^{-1}) × 10^6	k_d (s^{-1}) × 10^{-3}	K_D (M)
KCl	1.4	22	1.5×10^{-8}
NaCl	0.41	54	1.3×10^{-7}
LiCl	0.0018	25	1.4×10^{-5}

1562
1563

Source data files and supplementary files list

Figure 1—source data 1. RNA-binding proteins and their cDNA accession numbers screened in the genetic analyses in Figure 1.

Figure 1—source data 2. Summary of the genetic analyses in Figure 1.

Figure 1—source data 3. Statistical data related to Figures 1B, 1C, 1D, and 1E.

Figure 1—figure supplement 1−source data 1. Statistical data related to Figure 1—figure supplements 1C and 1D

Figure 1—figure supplement 2−source data 1. Statistical data related to Figure 1—figure supplement 2B, 2C and 2D

Figure 2—source data 1. Statistical data related to Figures 2B, 2C, 2D, and 2E.

Figure 2—figure supplement 1−source data 1. Statistical data related to Figure 2—figure supplement 1B

Figure 3−source data 1. Statistical data related to Figures 3B, 3C, 3E, 3F, 3H, and 3H

Figure 3−source data 2. Source data related to Figure 3G

1587   Figure 4—source data 1. Statistical data related to Figures 4B, 4D, 4F, 4G and 4I.

1588

1589   Figure 4—figure supplement 1−source data 1. Statistical data related to Figure 4—

1590   figure supplements 1B, 1C, and 1D

1591

1592   Figure 5—source data 1. Source data related to Figure 5F

1593

1594   Figure 6—source data 1. RNA-binding proteins and their cDNA accession numbers

1595   screened in the genetic analyses in Figure 6.

1596

1597   Figure 6—source data 2. Statistical data related to Figures 6B, 6C, and 6D.

1598

1599   Supplementary file 1. Full genotypes of the fly lines and their cultured temperatures.

1600

A

B

Eye size

C

Eye pigmentation

D

Eye pigmentation

	EGFP	*FUS*	*FUS-RRMmut*
(*GR*)$_{36}$			
(*GR*)$_{100}$		Lethal	Lethal
(*GA*)$_{36}$			
(*GA*)$_{100}$			
EGFP			

A

B

C

D

A

B   KCl

C   NaCl

D   LiCl

E

F

(G₄C₂) RNA **FUS** **Merged**

GMR×2 >
(G₄C₂)₈₉(H), FUS