Comparing the effects of host species richness on different metrics of disease

Michael H. Cortez*
Department of Biological Science
Florida State University
Tallahassee, FL 32306
(850) 645-8692
mcortez@fsu.edu

April 19, 2023

*Corresponding author
Synthesis and Review submission
Short title: Comparing responses in disease metrics
Keywords: dilution effect; amplification effect; environmental transmission; direct transmission; density-dependent; frequency-dependent; competence; transmission mode; basic reproduction number
Length: 3 Tables; 5 Color figures; 2 text boxes
Supplementary Material: Online Appendices S1-S4
Statement of Authorship: MHC did all aspects of the study
Data Statement: No new data generated in this study. Calculations and equations are provided in the appendices.

ABSTRACT: Changes in host species richness can alter infection risk and disease

levels in communities. I review theoretical predictions for direct and environmental transmission

pathogens about the effects of host additions (or removals) on three common disease metrics:

the pathogen community reproduction number ($\mathcal{R}_0$) and the proportion (i.e., infection

prevalence) and density of infected individuals in a focal host. To help explain why predictions

differ between studies and between metrics, I analyze an SIR-type model of an environmentally

transmitted pathogen and multiple host species that compete for resources. I use local

sensitivity analysis to explore and compare how variation in an added host's ability to

transmit a pathogen, its density, and the pathogen transmission mechanism affect each

disease metric. I find that if an added host species is a weak interspecific competitor, then

each metric typically increases or decreases, respectively, when the added host has a high

or low ability to transmit the pathogen. However, if the added host species is a strong

interspecific competitor, then the response of each metric can be strengthened, weakened,

or reversed depending on how the competitive effects of the added host propagate through

the community and alter the densities of all host species in the community. The three

metrics often respond in the same direction, but the metrics can respond in different directions

for three reasons: (1) strong density-mediated feedbacks driven by high disease-induced

mortality; (2) host additions or removals cause large changes in focal host density via competition

or disease-induced mortality; and (3) the metrics use different quantities to define a host's

ability to transmit disease: the quantities for infection prevalence and infected density

depend on the instantaneous production rates of infectious propagules whereas the quantities

for $\mathcal{R}_0$ depend on lifetime production of infectious propagules. This study provides a framework

that unifies prior theoretical studies and identifies rules governing the context-dependent

relationships between host species richness and the three metrics of disease.

1 Introduction

The gain or loss of host species from a community can alter levels of disease and disease risk for focal host populations. In empirical systems, increased host species richness can decrease or increase disease risk and levels of disease (Randolph and Dobson, 2012; Salkeld et al., 2013; Wood et al., 2014; Civitello et al., 2015; Ostfeld and Keesing, 2017; Rohr et al., 2019). Theoretical studies have used mathematical models to explore potential mechanisms driving this variation in host species-disease relationships (Dobson, 2004; Joseph et al., 2013; Mihaljevic et al., 2014; Faust et al., 2017). However, because the models make different assumptions about the host and pathogen species and use different metrics of disease, it remains unclear what biological and mathematical assumptions are responsible for the variation and differences in predictions (Buhnerkempe et al., 2015; Johnson et al., 2015; Rohr et al., 2019). Here, I present a framework that helps explain and extend these prior predictions, accounting for the effects of the characteristics of the host and pathogen species and the metric of disease.

Across empirical systems, increased host species richness often decreases levels of disease and disease risk (Civitello et al., 2015), but it can increase disease levels in many systems as well (Wood et al., 2014). The effects of altered host species richness are likely affected by which specific species are already present in the community and which are added to or removed from the community (Wood et al., 2016; Randolph and Dobson, 2012). For example, infection prevalence of the fungus *Batrachochytrium dendrobatidis* in three frog genera depends on which specific genera are present in a community: *Gastrophryne* species decrease, *Bufo* species increase, and *Hyla* species have no effect on infection prevalence in the other genera. Similarly, in laboratory experiments, the prevalence of fugal infections in *Daphnia dentifera* increases when the alternative host *D. lumholtzi* is present but decreases when the alternative host *Ceriodaphnia dubia* is present. In total, empirical studies show that relationships between host species richness and disease are likely to be context-dependent (Salkeld et al., 2013).

Despite the observed variation in how disease levels respond to changes in host species richness, we have a limited understanding of the mechanisms causing that variation and their context-dependent effects. This has led to calls for theory on host richness-disease relationships and, more generally, host biodiversity-disease relationships (Buhnerkempe et al., 2015; Johnson et al., 2015; Halsey, 2019; Rohr et al., 2019). Current theory predicts that responses to host additions and removals are affected by host competence (the ability to transmit the pathogen to susceptible individuals), interspecific host competition, and the pathogen transmission mechanism (Dobson, 2004; Rudolf and Antonovics, 2005; Joseph et al., 2013; Mihaljevic et al., 2014; O'Regan et al., 2015; Strauss et al., 2015; Faust et al., 2017; Cortez and Duffy, 2020; Cortez, 2021); specific predictions are discussed in the next section. While this body of work provides insight into how host species richness can affect levels of disease, general predictions remain unclear because the studies make different assumptions about the characteristics of the host and pathogen species.

Another limitation of current theory is that modeling studies use different metrics of disease and it is unclear how the choice of disease metric influences predictions. Three metrics of disease commonly used in empirical and theoretical studies are: (i) the pathogen's community reproduction number ($\mathcal{R}_0^C$), which is the average number of new infections produced by a single infected individual in an otherwise susceptible community; (ii) the proportion of infected individuals in a focal population (hereafter, focal host infection prevalence; I/N); and (iii) the density of infected individuals in a focal population (hereafter, focal host infected density; I). I note that decreased and increased disease risk in response to increased host species richness are often referred to as amplification and dilution, respectively (Keesing et al., 2006; Rohr et al., 2019), but to avoid confusion about which disease metric is being referenced, I will not use that terminology in this study. The three metrics provide different information about disease dynamics and disease risk. Specifically, $\mathcal{R}_0^C$ provides a measure of the risk of an outbreak and how fast an outbreak will grow. In comparison, infection prevalence and infected density both provide measures of disease burden for a

focal population. In addition, infection prevalence measures disease risk for a focal host population whereas infected density measures the potential risk for a pathogen to spread or spillover into another population or species (Roche et al., 2012; Wood and Lafferty, 2013). Theoretical studies predict that changes in host species richness can have effects of different signs on the three metrics (Roche et al., 2012; Roberts and Heesterbeek, 2018). As an empirical example, infection prevalence and infected density in *Daphnia lumholtzi* both increase when *D. dentifera* is present. In contrast, fungal infection prevalence in *Daphnia dentifera* increases but infected density decreases when *D. lumholtzi* is present (Searle et al., 2016). Thus, the metrics can respond in the same or opposite directions to changes in host species richness, but it is currently unknown when and why the metrics respond similarly versus differently.

In this study, I review the theoretical predictions about the relationships between host species richness and the three metrics of disease, focusing on pathogens with direct transmission (transmission via host-host contact) and environmental transmission (transmission via infectious propagules that are shed into an environment). Then, I present a multihost environmental transmission model and use it to explore the factors influencing the relationships between each metric of disease and host species richness. To do this, I use sensitivity analysis to calculate how each metric responds to variation in components of host competence, host density, and the pathogen transmission mode. This study extends prior work (Roberts and Heesterbeek, 2018; Cortez and Duffy, 2021; Cortez, 2021) using the same approach by computing the relationships for all three metrics and identifying when and why the three metrics can have relationships with the same or different signs. My results provide a framework that unifies prior theoretical studies and explains the biological and mathematical assumptions that are responsible for the variation in prior theoretical predictions. That in turn helps identify the rules and biological mechanisms shaping the context-dependent relationships between host species richness and disease in empirical systems.

2   Current theory on how host species richness affects disease

Many studies have reviewed and debated the empirical evidence for positive and negative effects of host species richness on levels of disease (Keesing et al., 2010; Randolph and Dobson, 2012; Salkeld et al., 2013; Wood et al., 2014; Civitello et al., 2015; Ostfeld and Keesing, 2017; Rohr et al., 2019). Here, I review current theory for directly and environmentally transmitted pathogens.

The foundational work in Keesing et al. (2006) predicts that disease risk in a focal host increases when host additions or removals increase host encounter rates (α), increase per exposure transmission rates (Δ), increase susceptible host densities (S), decrease mortality rates of infected individuals (μ), and decrease host recovery rates (ν). These predictions are derived from an equation describing changes in infected host density,

$$\frac{dI}{dt} = \overbrace{\alpha \Delta SI}^{\text{infection}} - \overbrace{(\mu + m)I}^{\text{mortality}} - \overbrace{\nu I}^{\text{recovery}}$$ (1)

where S and I are susceptible and infected host densities, respectively, and m is the non-disease mortality rate. The predicted responses are based on the idea that increases and decreases in the growth rate of infected individuals (dI/dt) increase and decrease disease risk, respectively. While the five mechanisms are defined for direct transmission pathogens (i.e., transmission via host-host contact), analogues exist for environmental transmission pathogens (i.e., transmission via infectious propagules shed into the environment; Johnson and Thieltges 2010) and vector-borne pathogens (Ostfeld and Keesing, 2000; Keesing et al., 2006). A key limitation of these predictions is that they focus on changes in the growth rate of infected individuals and do not directly address other quantities measured in populations (e.g., infection prevalence and infected density).

Other theoretical studies have explored how the presence and absence of host species

affect the pathogen's community reproduction number ($\mathcal{R}_0^C$) and infection prevalence (I^*/N^*, where N^* is total host density) and infected density (I^*) of a focal host at equilibrium. For density-dependent direct transmission (DDDT) models (host-host contact rates are proportional to the density of infected individuals), host additions often increase $\mathcal{R}_0^C$ and focal host infection prevalence (Dobson, 2004; Rudolf and Antonovics, 2005; Faust et al., 2017), but both metrics can decrease if there is strong interspecific competition (Joseph et al., 2013; Mihaljevic et al., 2014; O'Regan et al., 2015; Cortez, 2021). For frequency-dependent direct transmission (FDDT) models (host-host contact rates are proportional to the frequency of infected individuals), host additions often decrease $\mathcal{R}_0^C$ and focal host infection prevalence (Dobson, 2004; Rudolf and Antonovics, 2005; Joseph et al., 2013; Mihaljevic et al., 2014), but both metrics can increase if added host species have sufficiently high intraspecific and interspecific transmission rates (O'Regan et al., 2015; Faust et al., 2017). Combined, this suggests that host additions increase $\mathcal{R}_0^C$ and focal host infection prevalence more often under DDDT than FDDT (Dobson, 2004; Rudolf and Antonovics, 2005; Mihaljevic et al., 2014; Faust et al., 2017).

For environmental transmission (ET) models (transmission via infectious propagules shed into the environment), responses in $\mathcal{R}_0^C$ and focal host infection prevalence depend on interspecific host competition and whether host species are net producers (sources) or net removers (sinks) of infectious propagules (Cáceres et al., 2014; Cortez and Duffy, 2021; Cortez, 2021; Espira et al., 2022). In particular, added host species are more likely to increase $\mathcal{R}_0^C$ and focal host infection prevalence if they are sources (Cortez, 2021; Espira et al., 2022). Added host species are also more likely to increase focal host infection prevalence if they have stronger competitive effects on sink hosts than on source hosts (Cortez, 2021).

Taken together, current theory points to possible general rules about how host competence, host competition, and the pathogen transmission mechanism affect host species richness-disease relationships. In addition, current theory suggests responses in $\mathcal{R}_0^C$ and infection prevalence are similar. This is useful to modelers because $\mathcal{R}_0^C$ is typically easier to compute

from models than infection prevalence. However, current theory is limited in two key ways. First, studies make different assumptions about the number of host species, the strength of interspecific host interactions, and the rates and mechanisms of disease transmission. For example, some FDDT studies (Dobson, 2004; Joseph et al., 2013; Mihaljevic et al., 2014) assume interspecific competition is absent and interspecific transmission rates are lower than intraspecific transmission rates whereas other studies (O'Regan et al., 2015; Faust et al., 2017) assume interspecific competition is present and intraspecific and interspecific transmission rates are comparable. Similarly, some ET studies (Begon and Bowers, 1994; Fenton et al., 2015) assume interspecific competition is absent and all host species are net producers of infectious propagules, whereas others studies (Strauss et al., 2015; Cáceres et al., 2014; Cortez and Duffy, 2021; Espira et al., 2022) assume interspecific competition is present and host species can be net producers or net removers of infectious propagules. Differences in the assumptions about competition and transmission rates or infectious propagule production rates make it difficult to identify which assumptions are responsible for differences in model behavior. Moreover, the difference in transmission mechanisms complicate efforts to find general rules that apply to all pathogens.

The second limitation of current theory is that prior studies have used different metrics of disease and the three metrics listed above can respond differently to host additions and removal (Keesing et al., 2006; Roche et al., 2012; Roberts and Heesterbeek, 2018). Indeed, Keesing et al. (2006) noted that variation in transmission rates (via α or Δ) could result in infection prevalence and infected density changing in different directions under some conditions (page 488). The differing responses of the three metrics means that predictions for one metric may not directly apply to a different metric. In addition, because we currently do not understand when and why the metrics respond similarly versus differently, it is unclear how to fairly compare the predictions for the different metrics.

These limitations point to a need for theory that can extend and unify the results for each particular metric as well as explain why predictions differ between metrics. The following

presents the model and approach I use as a step toward addressing this need.

3 Models and Methods

3.1 Single-habitat n-host environmental transmission model

I focus on the model in Cortez (2021) of n competing host species and a shared environmentally transmitted pathogen. Infection occurs when susceptible individuals encounter infectious propagules (e.g., spores) in the environment that were shed by infected individuals. All host species shed infectious propagules into a shared habitat and are equally exposed to the infectious propagules. Examples of environmentally transmitted pathogens include the human pathogens *Vibrio cholera* and *Giardia lamblia* (Kaper et al., 1995; Wolfe, 1992), the fungal pathogen *Metschnikowia bicuspidata* of *Daphnia* species (Strauss et al., 2015; Searle et al., 2016), the chytrid fungus *Batrachochytrium dendrobatidis* of amphibians (Daszak et al., 2003; Skerratt et al., 2007), whirling disease in fish (Hedrick et al., 1998; Bartholomew and Reno, 2002), and trematode parasites of snails (Johnson et al., 2012). Without loss of generality, I refer to host 1 is the focal host and the others as alternative hosts; this choice is arbitrary, but in practice, the designation depends on the specific system being studied.

Each host species is divided into subpopulations of susceptible (S_i), infected (I_i), and recovered (R_i) individuals, where the total density for host i is $N_i = S_i + I_i + R_i$. The model describes the changes in each host class and infectious propagule density (P),

$$\frac{dS_i}{dt} = \overbrace{F_i(\cdot)}^{\text{reproduction}} - \overbrace{\beta_i S_i P}^{\text{infection}} - \overbrace{m_i S_i}^{\text{mortality}} + \overbrace{\gamma_i R_i}^{\text{loss of immunity}} \qquad (2a)$$

$$\frac{dI_i}{dt} = \overbrace{\beta_i S_i P}^{\text{infection}} - \overbrace{(m_i + \mu_i)I_i}^{\text{mortality}} - \overbrace{\nu_i I_i}^{\text{recovery}} \qquad (2b)$$

$$\frac{dR_i}{dt} = \overbrace{\nu_i I_i}^{\text{recovery}} - \overbrace{(m_i + \xi_i)R_i}^{\text{mortality}} - \overbrace{\gamma_i R_i}^{\text{loss of immunity}} \qquad (2c)$$

$$\frac{dP}{dt} = \overbrace{\sum_{j=1}^{n} \chi_j I_j}^{\text{shedding}} - \overbrace{\sum_{j=1}^{n}(u_{S_j}S_j + u_{I_j}I_j + u_{R_j}R_j)P}^{\text{uptake}} - \overbrace{\delta P}^{\text{degradation}} \qquad (2d)$$

The parameter values are summarized in Table 1 and additional details are provided in Appendix S1. In equation (2a), host i susceptible density increases due to reproduction (F_i), decreases due to infection ($\beta_i S_i P$) and non-disease mortality ($m_i S_i$), and increases due to waning immunity ($\gamma_i R_i$). In equation (2b), host i infected density increases due to new infections ($\beta_i S_i P$) and decreases due to non-disease mortality ($m_i I_i$), mortality due to disease ($\mu_i I_i$), and recovery ($\nu_i Y_i$). Because infection occurs when infectious propagules are taken up by susceptible individuals, the transmission parameter is the product of the uptake rate of susceptible individuals and the per infectious propagule probability of infection ($\beta_i = \rho_i u_{S_i}$). In equation (2c), host i recovered density increases due to recovery ($\nu_i I_i$) and decreases due to mortality ($m_i R_i + \xi_i R_i$) and waning immunity ($\gamma_i R_i$). I assume susceptible individuals and recovered individuals are identical except that recovered individuals may experience increased mortality due to prior exposure to the pathogen ($\xi_i \geq 0$). In equation (2d), infectious propagule density increases due to shedding by infected individuals (first sum) and decreases due to uptake of infectious propagules by all individuals of all host species (second sum) and degradation (δP).

The reproduction rates, F_i, account for intraspecific and interspecific host competition. For the numerical simulations presented in figures, I use the Lokta-Volterra competition

functions,

$$F_i(\cdot) = \overbrace{r_i(S_i + b_i I_i + R_i)}^{\substack{\text{maximum} \\ \text{reproduction rate}}} \overbrace{\left[1 - \sum_j \alpha_{ij}(S_i + e_{ij}I_j + R_j)\right]}^{\text{intraspecific and interspecific competition}} \tag{3}$$

where for host i, r_i is the maximum reproduction rate, b_i is the reduction in growth rate

for infected individuals, α_{ij} is the per capita competitive effect of host j, and e_{ij} determines

if infected individuals of host j are weaker ($e_{ij} < 1$), equal ($e_{ij} = 1$), or stronger ($e_{ij} >$

1) competitors than susceptible individuals. The latter accounts for the possibility that

infected individuals may be stronger competitors when infection causes increased appetite

or resource acquisition rates (Ponton et al., 2011; Shikano and Cory, 2016; Bernardo and

Singer, 2017).

I note a few important assumptions about model (2); additional mathematical details

about these and other assumptions are provided in Section S1.2.4. First, I assume intraspecific

density dependence and interspecific host competition only affect the reproduction rates

of the host species (F_i) and do not affect the host mortality rates. These assumptions

simplify the mathematical analysis and align with many multi-host-pathogen models (but

see, e.g., Marini et al. (2017)). Second, while I describe and interpret the interspecific

host interactions in terms of competitive interactions (i.e., $-, -$ interactions), my results

also hold for species that have mutualistic $(+, -)$ or contramensalistic $(+, -)$ interactions,

provided the interactions only affect the reproduction rates of the host species. My results

may not apply to species with predator-prey or consumer-resource interactions because

those interspecific interactions necessarily affect the mortality rates of a species.

Third, I assume the n host species stably coexist in the presence and absence of the

pathogen, respectively, at a disease-free equilibrium (with densities $S_i^{\ddagger}$) and an endemic

equilibrium (with densities S_i^*, I_i^*, and R_i^*). Without this assumption, my equilibrium-

based analyses are biologically uninformative. In addition, to explore the effects of host

additions on disease and without loss of generality, I assume host n is the added host species

and assume that host species 1,...,$n - 1$ can stably coexist in presence and absence of

the pathogen. This allows me to compare disease levels when the added host species is

absent and present. However, this assumption means that my results may not apply to

systems where host coexistence is pathogen-mediated or systems where the gain or loss of

the added host causes other host species to go extinct.

3.2   Model Generality

The predictions for the environmental transmission (ET) model (2) apply to direct transmission

models (Cortez, 2021). This is because the ET model (2) can reduce to a model with density-

dependent direct transmission (DDDT), frequency-dependent direct transmission (FDDT),

or direct transmission that is intermediate to DDDT and FDDT; see Box 1 for specific

conditions. This means predictions from model (2) simultaneously apply to ET, DDDT,

and FDDT.

While model (2) can reduce to a direct transmission form, it is not flexible enough to

reduce down to all possible parameterizations of DDDT and FDDT models. The underlying

reason is that model (2) reduces to a direct transmission form where the interspecific (β_{ij})

and intraspecific (β_{ii}) transmissions coefficients are correlated and not independent (mathematical

details provided in Appendix S1.2). To address this issue, in Appendix S2, I extend my

results to an n-habitat version of the model that allows for partially overlapping host habitats.

The key difference between the 1-habitat and n-habitat models is that each host species

can have different shedding rates in their own habitat (χ_{jj}) and the habitats of heterospecifics

(χ_{ij}). This allows for each species to have different rates of transmission to conspecifics

and heterospecifics. That makes the n-habitat model flexible enough to reduce down to all

possible parameterizations of DDDT and FDDT models (mathematical details provided in

Appendix S2.2).

A full analysis of the n-habitat ET model is not tractable. Thus, Appendix S2 primarily

focuses on systems where (i) interspecific host competition is absent and (ii) each host

species has low shedding and uptake rates in the habitats of other host species. When

the n-habitat ET model is reduced down to a direct transmission form, these conditions
result in a direct transmission model where (i) interspecific host competition is absent and
(ii) interspecific direct transmission rates are lower than intraspecific direct transmission
rates ($\hat{\beta}_{ij} > \hat{\beta}_{jj}$, where $\hat{\beta}_{ij}$ is the direct transmission rate parameter for host j transmitting
the pathogento host i). While restrictive, these conditions align with the assumptions in
prior studies on DDDT and FDDT models (Dobson, 2004; Joseph et al., 2013; Mihaljevic
et al., 2014). Thus, the restricted results for the n-habitat model help explain patterns
from these prior studies.

In total, analyzing the 1-habitat model (2) and the n-habitat model provides a way
to extend and unify results for ET, DDDT, and FDDT. The main text focuses on the
simpler 1-habitat model because the results from the 1-habitat and n-habitat models are
qualitatively similar. The only difference is that the predictions for the n-habitat model
depend the host's rates of shedding into the habitats of heterospecifics. These differences
are indicated in the text.

3.3   Defining host competence, source, and sink

My mathematical analysis identifies two characteristics of host species that measure how
well each host species can transmit the pathogen.

First, host competence measures the potential ability of an exposed individual to transmit
the pathogen over the entire length of time the individual is infected. Mathematically, I
define host competence as

$$\mathcal{R}_{0,i}(\infty) = \frac{\beta_i}{u_{s_i}} \frac{\chi_i}{m_i + \mu_i + \nu_i} = \overbrace{\rho_i}^{\substack{\text{per spore probability} \\ \text{of infection}}} \cdot \overbrace{\frac{\chi_i}{m_i + \mu_i + \nu_i}}^{\substack{\text{lifetime production} \\ \text{of infectious propagules}}} \tag{4}$$

where the second equality follows from the transmission rate (β_i) being the product of
the per infectious propagule probability of infection (ρ_i) and the uptake rate of infectious

propagules (u_{S_i}). On the right hand side of the equation, the first term is the probability of infection given exposure and the fraction is the shedding rate (χ_i) multiplied by the average duration of infection ($1/[\mu_i + m_i + \gamma_i]$), which equals the average lifetime production of infectious propagules by an infected individual of host i. Combined, this yields the average lifetime production of infectious propagules by individuals that are exposed to the pathogen. This definition of competence matches that proposed in prior studies (Merrill and Johnson, 2020). I note that my notation for competence comes from equation (4) being mathematically equivalent to the pathogen's basic reproduction number for an arbitrarily large population of host i; see Appendix S1.3 for additional details.

Second, source and sink measure a host species' instantaneous per capita net production rate of infectious propagules. Mathematically, source and sink are defined by,

$$
\begin{aligned}
\varphi_i &= \frac{1}{I_i^*}\left(\overbrace{\chi_i I_i^*}^{\text{shedding}} - \overbrace{[u_{S_i} S_i^* + u_{I_i} I_i^* + u_{R_i} R_i^*]P^*}^{\text{uptake}} \right) \\
&= \overbrace{\chi_i}^{\text{shedding}} - \overbrace{\frac{(m_i + \mu_i + \nu_i)}{\beta_i}u_{S_i} - u_{I_i}P^* - \frac{\nu_i}{\xi_i + m_i + \gamma_i}u_{R_i}P^*}^{\text{uptake}}.
\end{aligned}
\tag{5}
$$

(The second line follows from solving the equilibrium conditions $dI_i/dt = 0$ and $dR_i/dt = 0$ for S_i^* and R_i^*.) Source hosts are net producers of infectious propagules, meaning the host's per capita shedding rate is larger than its per capita uptake rate ($\varphi_i > 0$). Sink hosts are net removers of infectious propagules, meaning the host's per capita shedding rate is smaller than its per capita uptake rate ($\varphi_i < 0$).

While similar, competence and sink/source are distinct because competence measures the lifetime production of infectious propagules per individual whereas sink and source measure the net instantaneous production of infectious propagules per individual. The similarities between competence and sink/source mean that high and low competence often means source and sink, respectively. However, the differences between the definitions

mean that high competence hosts can be sinks and low competence hosts can be sources. First, high competence hosts are sinks when they have low mortality and recovery rates (small μ_i and ν_i, making $\mathcal{R}_{0,i}(\infty)$ large) and small shedding rates (small χ_i, making φ_i negative or small and positive). Such hosts have low instantaneous production rates of infectious propagules, but because individuals remain infected for long periods of time, their lifetime production rates of infectious propagules are large. Second, low competence hosts are large sources when they have high mortality and recovery rates (large $\mu_i + \nu_i$, making $\mathcal{R}_{0,i}(\infty)$ small) and large shedding rates (large χ_i, making φ_i large and positive). Such hosts have high instantaneous production rates of infectious propagules, but because individuals remain infected for short periods of time, their lifetime production rates of infectious propagules are small.

I note two additional things about the definitions of competence, sink, and source. First, in addition to arising naturally in the calculations, an advantage of these definitions is that they provide a density-independent ranking of the host species from highest to lowest competence and from largest source to largest sink. This avoids complications that arise when densities shift as host species are added or removed from a community. However, the rankings can differ across environments because both definitions depend on the non-disease mortality rates, m_i. Second, both definitions apply to direct transmission pathogens because of the connection between direct and environmental transmission models (Box 1). For competence, higher competence hosts have larger intraspecific transmission rates and lower mortality and recovery rates. For source and sink, all hosts are sources ($\varphi_i > 0$) under DDDT because DDDT corresponds to no uptake in the ET model (2). Under FDDT, hosts are sources ($\varphi_i > 0$) when their intraspecific and interspecific transmission rates are sufficiently large and sinks ($\varphi_i < 0$) otherwise; see Appendices S1.3 and S2.2 for mathematical details.

3.4 Computing the sensitivities of the disease metrics

I use model (2) to predict how additions and removals of alternative hosts affect three disease metrics: (i) the parasite's basic reproduction number for the community ($\mathcal{R}_0^C$), (ii) focal host infection prevalence at equilibrium (I_1^*/N_1^*), and (iii) focal host infected density at equilibrium (I_1^*). Because explicit formulas do not exist for the metrics except in special cases, I cannot directly compute how additions and removals of alternative hosts affect each metric. However, it is possible to compute explicit formulas for the slopes of the relationships between each metric of disease and alternative host competence ($\mathcal{R}_{0,n}(\infty)$), alternative host equilibrium density (N_n^*), and the pathogen transmission mode. The signs of the slopes yield insight into the effects of additions/removals on each metric of disease. As in prior studies (Roberts and Heesterbeek, 2018; Cortez and Duffy, 2021; Cortez, 2021), I compute the slopes using local sensitivities (i.e., derivatives). Here, I describe the idea behind the approach. Box 2 and Appendix S2.4 explain how the $\mathcal{R}_0^C$ sensitivities are computed using the theory for stage-structured models. Appendix S1.5 explains how the sensitivities of the other two metrics are computed using the Jacobian-based approach in Cortez and Duffy (2021) and Cortez (2021).

The slopes of the relationships between each metric of disease and the components of alternative host competence are computed using derivatives with respect to the alternative host's transmission (β_n), mortality (μ_n), recovery (ν_n), shedding (χ_n), and uptake ($u_{S_i}, u_{I_i}, u_{R_i}$) rates. For example, the sensitivities for the alternative host's shedding rate are computed using the derivatives,

$$\overbrace{\frac{\partial \mathcal{R}_0^C}{\partial \chi_n}, \quad \frac{\partial}{\partial \chi_n}\frac{I_1^*}{N_1^*}, \quad \frac{\partial I_1^*}{\partial \chi_n}}^{\text{sensitivities to components of alternative host competence}} \quad . \tag{6}$$

Positive and negative values imply that each metric increases and decreases, respectively, as the alternative host's shedding rate increases. These sensitivities are useful because they identify how variation in the epidemiological traits of the alternative host species

influences the disease metrics. In addition, the sensitivities can provide insight about the consequences of intervention strategies that target a specific biological process, e.g., vaccination that lowers infection rates of an alternative host (decreased β_n) or the culling of infected individuals of an alternative host (increased μ_n).

The slopes of the relationships between each metric of disease and alternative host density are computed using the derivatives

$$\overbrace{\frac{d\mathcal{R}_0^C}{dN_n^{\ddagger}}, \quad \frac{d}{dN_n^*}\frac{I_1^*}{N_1^*}, \quad \frac{dI_1^*}{dN_n^*}}^{\text{sensitivities to alternative host density}} \tag{7}$$

where $N_n^{\ddagger}$ and N_n^* are the densities of host n at the disease-free and endemic equilibria, respectively. (The change in derivative notation is mathematically necessary to account for the total effect of variation in alternative host density.) Positive and negative values imply that each metric increases and decreases, respectively, as the alternative host's density increases. This yields insight about the effects of host additions because the effect of adding an alternative host to the community is predicted by the slope of the response as alternative host density is increased from zero to its equilibrium value. In particular, positive and negative values suggest that the addition of the alternative host will increase and decrease, respectively, each metric of disease.

In total, the sensitivity formulas identify the specific conditions under which each metric has positive versus negative relationships with alternative host competence, alternative host density, and the pathogen transmission mechanism. Comparing those conditions yields insight into when the three metrics will have similar or different responses to host additions and removals.

4 Results

I first present the general mechanisms causing the sensitivities for the three metrics to differ in sign. I then compare how the metrics respond to changes in alternative host competence, alternative host density, and the pathogen transmission mode.

4.1 General mechanisms for why sensitivities can have different signs

The three general mechanisms explaining why the sensitivities for the metrics can differ in sign are (1) differences in the definitions of competence and source/sink, (2) density-mediated feedbacks driven by disease-induced mortality, and (3) large changes in focal host density.

For mechanism 1, differences in the definitions of competence and source/sink can cause the sensitivities of $\mathcal{R}_0^C$ to differ in sign from the sensitivities of focal host infection prevalence (I_1^*/N_1^*) and infected density (I_1^*) at equilibrium. The reason is that the sensitivity formulas for $\mathcal{R}_0^C$ depend on host competence $(\mathcal{R}_{0,i}(\infty))$ whereas the sensitivity formulas for focal host infection prevalence and infected density depend on whether hosts are sources or sinks (φ_i). Consequently, predictions for $\mathcal{R}_0^C$ can disagree with predictions for the other two metrics when high and low competence do not imply source and sink, respectively.

For mechanism 2, density-mediated feedbacks driven by disease-induced mortality can cause the sensitivities of $\mathcal{R}_0^C$ to differ in sign from the sensitivities of the other two metrics. The underlying reason is that $\mathcal{R}_0^C$ is computed using the disease-free equilibrium densities $(N_i^\ddagger)$, which are independent of disease-induced mortality rates, whereas focal host infection prevalence and infected density depend on the disease-induced mortality rates. For the latter, as conditions are varied (e.g., a host species is added to the community), changes in infectious propagule densities will lead to changes in infection rates, that will lead to changes in total disease-induced mortality rates for each host $(\mu_i I_i)$, and that will lead to

changes in the total and infected densities of each host. If the feedbacks due to disease-induced mortality are strong, then the changes in total and infected densities will be large. That in turn can cause focal host infection prevalence and infected density to respond in the opposite direction of $\mathcal{R}_0^C$.

For mechanism 3, large changes in focal host density can cause the sensitivities of focal host infection prevalence and focal host infected density to differ in sign. The mathematical reason is that the quotient rule from calculus implies that the sensitivities to alternative host density (equation (7)) are related by

$$
\overbrace{\frac{dI_1^*}{dN_n^*}}^{\substack{\text{Change in} \\ \text{infected density}}} = N_1^* \overbrace{\frac{d}{dN_n^*}\frac{I_1^*}{N_1^*}}^{\substack{\text{Change in} \\ \text{infection prevalence}}} + \frac{I_1^*}{N_1^*} \overbrace{\frac{dN_1^*}{dN_n^*}}^{\substack{\text{Change in} \\ \text{total density}}}. \tag{8}
$$

The same relationship applies to the sensitivities to parameter values (equation (6)). The sensitivities of infected density (left side of equation) and infection prevalence (first term on the right side of the equation) can only differ in sign if the changes in total focal host density (second term on the right side of the equation) are large and in the opposite direction of the changes in infection prevalence. For example, increased density of host n could increase focal infection prevalence ($[d/dN_n^*][I_1^*/N_n^*] > 0$) but also decrease focal host infected density ($dI_1^*/dN_n^* < 0$) because of large decreases in focal host density ($dN_1^*/dN_n^* < 0$). My calculations show that the sensitivities of focal host infection prevalence and infected density have opposite signs only if (i) some host species are more strongly regulated by disease and competition with alternative host species than by competition with the focal host species or (ii) the focal host is a sink or small source and a subset of the alternative hosts are large sources; see Remark S1.6 of Appendix S1.1.5 and Appendix S1.5 for mathematical details. The most likely scenario for condition (i) to be met both biologically and mathematically is when the focal host is strongly regulated by disease or interspecific competition and weakly regulated by intraspecific competition. Because of this, I will only discuss condition (i) in terms of regulation of the focal host.

4.2 Sensitivities of disease metrics to alternative host competence

I now compare the sensitivities of the three metrics to the components that define the competence of the alternative host, $\mathcal{R}_{0,n}(\infty) = \chi_n \beta_n / [u_{S_n}(m_n + \mu_n + \nu_n)]$. These sensitivities are useful because they help identify how variation in the epidemiological traits of the alternative host species influences disease levels and disease risk in the focal host. For instance, the sensitivities can provide insight about the consequences of intervention strategies that target a specific biological process, such as vaccination that lowers infection rates of an alternative host (decreased β_n).

Community reproduction number: Any variation in epidemiological traits that increases the competence of the alternative host always increases $\mathcal{R}_0^C$; mathematical details provided in Appendix S1.4.2. In particular, increased shedding rates (χ_n), decreased uptake rates ($u_{S_n}, u_{I_n}, u_{R_n}$), increased transmission rates (β_n; Figure 1A), decreased mortality rates (μ_n; Figure 1D), and decreased recovery rates (ν_n) of an alternative host always increase $\mathcal{R}_0^C$. These predictions explain why $\mathcal{R}_0^C$ always increased with increased host competence in prior studies of direct (Dobson, 2004; Joseph et al., 2013; O'Regan et al., 2015; Huang et al., 2019) and environmental (Feng et al., 2009; Espira et al., 2022) transmission pathogens; see Appendix S3 for details about each study.

Focal host infection prevalence: Variation in epidemiological traits that increases the competence of the alternative host often increases focal host infection prevalence, but not always; mathematical details provided in Appendix S1.5.3. Specifically, increases in the alternative host's shedding rate (χ_n) or decreases in its uptake rates ($u_{S_n}, u_{I_n}, u_{R_n}$) always increase focal host infection prevalence. In comparison, increases in an alternative host's transmission parameter (β_n) and decreases in its mortality and recovery rates (μ_n, ν_n) often increase focal host infection prevalence (black circles and red triangles in Figure 1BE). However, decreased focal host infection prevalence is also possible; here, I focus on the three most biologically likely scenarios and refer the reader to Cortez (2021) and Appendix S1.5.3 for additional details. First, increases in the alternative host's transmission

parameter (β_n) can decrease focal host infection prevalence when the alternative host is a large source ($\varphi_n > 0$) and interspecific competition is sufficiently strong (blue squares in Figure 1B). Second, decreases in the alternative host's mortality rate (μ_n) can decrease focal host infection prevalence when the alternative host is a large sink ($\varphi_n < 0$; blue squares in Figure 1E). Third, increases in β_n and decreases in μ_n or ν_n can decrease focal host infection prevalence when the alternative host has stronger interspecific competitive effects on sink hosts than source hosts. For all three scenarios, increased competence of the alternative host causes shifts in its densities that reduce the production of infectious propagules, which in turn causes decreases in focal host infection prevalence.

   Focal host infected density: The predictions for focal host infected density are often the same as focal host infection prevalence. However, the predictions can be reversed if (i) the focal host is more strongly regulated by the disease or interspecific competition than by intraspecific competition or (ii) the focal host is a sink or small source and other hosts are large sources. Mathematical details are given in Appendix S1.5.3.

   Comparison of metrics: Intuition suggests that higher competence in the alternative host (larger $\mathcal{R}_{0,n}(\infty)$) will lead to increases in all metrics of disease. This is often the case, resulting in all metrics responding to variation in alternative host competence in the same direction (black circles in Figure 1). However, the intuition can be incorrect for focal host infection prevalence and infected density. This results in variation in the competence of the alternative host having effects of one sign on $\mathcal{R}_0^C$ and effects of the opposite sign on the other focal host infection prevalence and infected density (blue squares in 1). This difference in responses is due to density-mediated feedbacks (mechanism 2) where changes in alternative host competence alter infectious propagule densities, which affects disease-induced mortality in the focal host.

   Variation in the competence of the alternative host also can have effects of opposite sign on focal host infection prevalence and focal host infected density. This difference in responses is due to variation in the competence of the alternative host causing large changes

484 in focal host density (mechanisms 3). This occurs when (i) the focal host is more strongly

485 regulated by the disease or interspecific competition than intraspecific competition (red

486 triangles in Figure 1A-C) or (ii) the focal host is a sink or small source and other hosts are

487 large sources (red triangles in Figure 1D-F).

4.3 Sensitivities of metrics to alternative host density

Here, I compare how each disease metric responds to variation in the density of an alternative

host. These sensitivities yield insight about the effects of host additions because positive

and negative values suggest that the addition of an alternative host will increase and decrease

a metric, respectively. To facilitate that connection and avoid confusion, I vary the density

of host n and refer to host n as the "added host".

The response of each metric depends on how the densities of all host species shift as

the density of the added host increases. In the following formulas, $\alpha_{in}^{total\ddagger}$ and α_{in}^{total*} denote

the total competitive effects (i.e., direct $+$ indirect effect) of the added host on host i at

the disease-free and endemic equilibria, respectively (see Appendix S1.1.5 for details).

The total competitive effects can be positive or negative depending on the strength and

asymmetry of interspecific host competition across the community. If the added host has

a positive (negative) total effect on another host, then increased density of the added host

causes an increase (decrease) in the density of the other host.

Community reproduction number: The sensitivity of $\mathcal{R}_0^C$ to the density of the

added host is

$$\frac{d\mathcal{R}_0^C}{dN_n^{\ddagger}} = C_n^{\ddagger} \overbrace{\left(\mathcal{R}_{0,n}(\infty) - \mathcal{R}_0^C\right)}^{\text{relative competence of host } n} + \sum_{i<n} C_i^{\ddagger} \overbrace{\left(\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C\right)}^{\text{relative competence of host } i} \overbrace{\alpha_{in}^{total\ddagger}}^{\substack{\text{total competitive} \\ \text{effect on host } i}} \tag{9}$$

where the $C_i^{\ddagger}$ are positive constants; see Appendix S1.4.1. Equation (9) shows that the

change in $\mathcal{R}_0^C$ depends on whether the competence of each host species relative to $\mathcal{R}_0^C$ is

high ($\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C > 0$) or low ($\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C < 0$) and whether the added host has

a positive ($\alpha_{in}^{total\ddagger} > 0$) or negative ($\alpha_{in}^{total\ddagger} < 0$) total competitive effect on each species.

In particular, increased density of the added host is more likely to increase $\mathcal{R}_0^C$ when the

added host has higher competence ($\mathcal{R}_{0,n}(\infty) - \mathcal{R}_0^C > 0$), the added host has less negative

total effects on high competence hosts ($\alpha_{in}^{total\ddagger} < 0$ small in magnitude for hosts with

$\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C > 0$), and the added host has more negative total effects on low competence

hosts ($\alpha_{in}^{total\ddagger} < 0$ large in magnitude for hosts with $\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C < 0$). In comparison,

increased density of the added host is more likely to decrease $\mathcal{R}_0^C$ when the added host has

lower competence ($\mathcal{R}_{0,n}(\infty) - \mathcal{R}_0^C < 0$), the added host has less negative total effects on

low competence hosts ($\alpha_{in}^{total\ddagger} < 0$ small in magnitude for hosts with $\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C < 0$),

and the added host has more negative total effects on high competence hosts ($\alpha_{in}^{total\ddagger} < 0$

large in magnitude for hosts with $\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C > 0$).

Combining the above yields the following predictions about how $\mathcal{R}_0^C$ responds to increased

density of the added host. First, if interspecific competition is absent, then the response

is determined solely by the competence of the added host: $\mathcal{R}_0^C$ increases when the added

host has sufficiently high competence ($\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C > 0$; black circles in Figure 2A)

and decreases otherwise ($\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C < 0$; black circles in Figure 2D). Second, these

predictions can be reversed if interspecific competition is sufficiently strong, depending

on the competitive effects of the added host. In particular, increased density of an added

host with high competence can decrease $\mathcal{R}_0^C$ when the added host has more negative (or

less positive) total competitive effects on high competence hosts than low competence

hosts (blue squares in Figure 2A). Similarly, increased density of an added host with low

competence can increase $\mathcal{R}_0^C$ when the added host has more negative (or less positive)

total competitive effects on low competence hosts than high competence hosts (blue squares

in Figure 2D).

The above predictions are only slightly modified when accounting for partially overlapping

habitats. In particular, the conditions depend both on host competence and inter-habitat

shedding rates (i.e., shedding rates into other species' habitats); mathematical details for

the analysis of the n-habitat model are provided in Appendix S2.4.2. Specifically, in the

absence of interspecific competition, $\mathcal{R}_0^C$ increases when the added host has sufficiently

high competence and inter-habitat shedding rates and $\mathcal{R}_0^C$ decreases when the added host

has sufficiently low competence and inter-habitat shedding rates. When interspecific competition

is present, these predictions can be reversed if the added host has more negative total

competitive effects on hosts with high competence and high inter-habitat shedding rates

or hosts with low competence and low inter-habitat shedding rates, respectively.

Focal host infection prevalence: The effect of increased density of the added host

on focal host infection prevalence is

$$
\frac{d}{dN_n^*}\frac{I_1^*}{N_1^*} = \overbrace{C_n^*\varphi_n}^{\substack{\text{host } n \text{ is}\\\text{sink/source}}} + \sum_i C_i^* \overbrace{\varphi_i}^{\substack{\text{host } i \text{ is}\\\text{sink/source}}} \overbrace{\alpha_{in}^{total*}}^{\substack{\text{total competitive}\\\text{effect on host } i}} \tag{10}
$$

where the C_i^* are positive constants; see Appendix S1.5.2. Due to their similar forms, the

interpretation of equation (10) is the same as equation (9) except that high and low competence

are replaced with source and sink, respectively. In particular, increased density of the

added host is more likely to increase focal host infection prevalence when the added host is

a source ($\varphi_n > 0$), the added host has less negative total effects on source hosts ($\alpha_{in}^{total*} < 0$

small in magnitude for hosts with $\varphi_i > 0$), and the added host has more negative total

effects on sink hosts ($\alpha_{in}^{total*} < 0$ large in magnitude for hosts with $\varphi_i < 0$). In comparison,

increased density of the added host is more likely to decrease focal host infection prevalence

when the added host is a sink ($\varphi_n < 0$), the added host has less negative total effects on

sink hosts ($\alpha_{in}^{total*} < 0$ small in magnitude for hosts with $\varphi_i < 0$), and the added host has

more negative total effects on source hosts ($\alpha_{in}^{total*} < 0$ large in magnitude for hosts with

$\varphi_i > 0$).

Combining the above yields the following predictions about how focal host infection

prevalence responds to increased density of the added host. First, if interspecific competition

is absent ($\alpha_{in}^{total*} = 0$), then the response is determined solely by whether the added host is

a source or sink: focal host infection prevalence increases when the added host is a source ($\varphi_n > 0$; black circles in Figure 2BC) and decreases when the added host a sink ($\varphi_n < 0$; black circles in Figure 2EF). Second, these predictions can be reversed if interspecific competition is sufficient strong, depending on how competition affect other source and sink hosts. In particular, increased density of an added source host can decrease focal host infection prevalence when the added host has more negative (or less positive) total competitive effects on source hosts than sink hosts (blue squares in Figure 2BC). Similarly, increased density of an added sink host can decrease focal host infection prevalence when the added host has more negative (or less positive) total competitive effects on sink hosts than source hosts (blue squares in Figure 2EF).

I note that when accounting for partially overlapping habitats, the above predictions are extended to account for whether each host species is a source or sink in each species' habitat; see Appendix S2.5.2 for mathematical details.

Focal host infected density: The predictions for focal host infected density are often the same as focal host infection prevalence (black circles and blue squares in Figure 3ABEF). However, the predictions can be reversed if (i) the focal host is more strongly regulated by the disease or interspecific competition than by intraspecific competition or (ii) the focal host is a sink or small source and other hosts are large sources. Mathematical details are given in Appendix S1.5.2.

Comparison of metrics: Increased alternative host density often has effects of the same sign on all metrics (black circles in Figure 3). However, responses in $\mathcal{R}_0^C$ can differ in sign from the other two metrics due to differences in the definitions of competence and source/sink (mechanism 1). For example, increased density of a high competence sink host increases $\mathcal{R}_0^C$ but decreases focal host infection prevalence and infected density (blue squares in Figure 3A-C). Similarly, increased density of a low competence source host decreases $\mathcal{R}_0^C$ but increases focal host infection prevalence and infected density (blue squares in Figure 3D-F). In addition, while focal host infection prevalence and infected density

often respond in the same direction (black circles and blue squares in Figure 3ABEF), the

two metrics can response in opposite directions due to large changes in focal host density

(mechanism 3) when (i) the focal host is more strongly regulated by disease or interspecific

competition than intraspecific competition (red triangles in Figure 3BCEF) or (ii) the

focal host is a sink or small source and other hosts are large sources (not shown).

4.4 Predictions for host species richness-disease relationships

The sensitivities in the previous section predict how increased density of an added host

affect the three disease metrics. Here, I used those results to predict how interspecific

competition and correlations between host competence and arrival order shape the relationships

between host species richness and each disease metric.

Community reproduction number: The predicted effects of host additions on $\mathcal{R}_0^C$

yield the following kinds of relationships between host species richness and $\mathcal{R}_0^C$. First,

$\mathcal{R}_0^C$ is predicted to increase when added hosts have sufficiently high competence ($\mathcal{R}_0^i(\infty)$

large) and inter-habitat shedding rates (large $\hat{\chi}_{ij}$), unless the added host has sufficiently

strong negative competitive effects on other hosts with high competence and inter-habitat

shedding rates. As an example, consider a system where higher competence hosts are only

present in communities with more species, i.e., there is a positive correlation between host

competence and host arrival order. In such a system, there is typically a positive relationship

between host species richness and $\mathcal{R}_0^C$ (black circle in Figure 4A). However, a negative

relationship can arise if higher competence hosts are also stronger interspecific competitors

because the later arriving hosts suppress host densities (blue squares in Figure 4A).

Second, $\mathcal{R}_0^C$ is predicted to decrease when added hosts have sufficiently low competence

($\mathcal{R}_0^i(\infty)$ small) and inter-habitat shedding rates (small $\hat{\beta}_{ij}$), unless the added host has

sufficiently strong negative competitive effects on other hosts with low competence and

inter-habitat shedding rates. As an example, consider a system where lower competence

hosts are only present in communities with more species, i.e., there is a negative correlation

between host competence and host arrival order. In such a system there is typically a negative relationship between host species richness and $\mathcal{R}_0^C$ (filled black circles in Figure 4D). However, U-shaped relationships between host species richness and $\mathcal{R}_0^C$ are possible if low competence hosts also have strong interspecific competitive effects on other low competence hosts (blue squares in Figure 4D). In this case, the densities of the low competence hosts are suppressed as more species are added to the community, which leads to increases in $\mathcal{R}_0^C$.

I note that if lower competence hosts are only present in communities with more species and the competences of the lower competence hosts are not too low, then $\mathcal{R}_0^C$ can have a positive relationship with host species richness In these counter-intuitive cases, later arriving hosts species are worse at spreading the disease than earlier arriving hosts, but all host species facilitate disease spread across the community. Mathematically, this case corresponds to equation (9) where the later arriving hosts have smaller $\mathcal{R}_{0,i}(\infty)$ values, but all values are large enough that $\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C > 0$.

Focal host infection prevalence: The predicted effects of host additions on focal host infection prevalence yield the following kinds of relationships between host species richness and focal host infection prevalence. First, added source hosts ($\varphi_n > 0$) are predicted to increase focal host infection prevalence, unless the added host has sufficiently strong negative competitive effects on other source hosts. As an example, consider a system where larger source hosts are only present in communities with more species, i.e., there is a positive correlation between net production of infectious propagules and host arrival order. In such a system there is typically a positive relationship between host species richness and focal host infection prevalence. However, a negative relationship can arise if larger source hosts are also stronger interspecific competitors because the later arriving hosts suppress host densities.

Second, added sink hosts ($\varphi_n < 0$) are predicted to decrease focal host infection prevalence, unless the added host has sufficiently strong negative competitive effects on other sink

hosts. As an example, consider a system where large sink hosts are only present in communities with more species, i.e., there is a negative correlation between net production of infectious propagules and host arrival order. In such a system there is typically a negative relationship between host species richness and focal host infection prevalence. However, U-shaped relationships between host species richness and focal host infection prevalence are possible if sink hosts also have strong interspecific competitive effects on other sink hosts. In this case, the densities of the sink hosts are suppressed as more species are added to the community, which leads to increases in infectious propagule density and focal host infection prevalence.

Focal host infected density: Because the predicted effects of host additions on focal host infected density and infected density are often the same, the relationships of those two metrics with host species richness are often the same (compare blue squares in Figure 4B,C and all curves in Figure 4E,F). For example, if large sink hosts are only present in a community with more species (Figure 4,E,F), then focal host infection prevalence and focal host infected density both decrease with increased host richness (black circles), unless sink hosts also have strong interspecific competitive effects on other sink hosts (blue squares). However, the relationships can differ if the focal host is (i) the focal host is more strongly regulated by the disease or interspecific competition than by intraspecific competition or (ii) the focal host is a sink or small source and other hosts are large sources. Both of these conditions explain why adding large source hosts in Figure 4BC causes increased infection prevalence (black circles in Figure 4B) and decreased infected density (black circles in Figure 4C).

Explaining prior results about richness-disease relationships: The sensitivities help explain prior results for environmental transmission models (Table 3); see Appendix S3.2 for mathematical details. First, in prior studies (Rudge et al., 2013; Fenton et al., 2015; Espira et al., 2022) added high competence sources increased $\mathcal{R}_0^C$ and focal host infection prevalence whereas added low competence sinks decreased both metrics because those studies assumed no interspecific competition ($\alpha_{in}^{total} = 0$). Second, other studies of

two-host models (Cáceres et al., 2014; Searle et al., 2016) assumed the focal host was a source ($\varphi_1 > 0$) and there was strong interspecific competition ($\alpha_{1n}^{total} < 0$ where $n = 2$). Consequently, a low competence alternative sink host will necessarily decrease $\mathcal{R}_0^C$ and focal host infection prevalence (Cáceres et al., 2014), but an added alternative source host can increase or decrease focal host infection prevalence (Searle et al., 2016). Third, the added alternative source host in Searle et al. (2016) increased focal host infection prevalence, but decreased focal host infected density. This difference is due to large changes in focal host density (mechanism 3) caused by the focal host being a small source and the alternative host being a very large source; I return this particular example in Section "Application to model of *Daphnia*-fungus system."

Translating the predictions to direct transmission models explains prior results for density-dependent (DDDT) and frequency-dependent (FDDT) direct transmission models (Table 3); mathematical details are provided in Appendix S3.1. Under DDDT, all hosts have high competence ($\mathcal{R}_{0,i}(\infty) - \mathcal{R}_0^C > 0$) and are sources ($\varphi_i > 0$). Thus, equations (9) and (10) are positive in the absence of interspecific competition ($\alpha_{in}^{total} = 0$) and negative only if interspecific competition is sufficiently strong ($\alpha_{in}^{total} < 0$). This explains why host additions always increased $\mathcal{R}_0^C$ and focal host infection prevalence in the absence of interspecific competition (Begon et al., 1992; Dobson, 2004; Rudolf and Antonovics, 2005; McCormack and Allen, 2007; Faust et al., 2017; Huang et al., 2019), but host additions decreased those metrics when interspecific competition was strong (Bowers and Turner, 1997; Peixoto and Abramson, 2006; Joseph et al., 2013; Mihaljevic et al., 2014; O'Regan et al., 2015).

Under FDDT, an alternative host is a source if it has sufficiently high interspecific transmission rates and a sink otherwise (mathematical details given in Special Cases B and C of Appendix S2.4.2 and Special Case 2 of Appendix S2.5.2). Prior studies of FDDT models assume interspecific transmission rates are fractions of intraspecific transmission rates (i.e, $\hat{\beta}_{ij} = c_j \hat{\beta}_{jj}$ where $c_j < 1$), which means an alternative host is source only if it has sufficiently high competence. As a result, additions of lower competence alternative

692 hosts decreased $\mathcal{R}_0^C$ and focal host infection prevalence (Dobson, 2004; Rudolf and Antonovics,

693 2005; Faust et al., 2017; Mihaljevic et al., 2014), but additions of sufficiently high competence

694 alternative hosts increased both metrics (Joseph et al., 2013; O'Regan et al., 2015).

4.5 Sensitivities of metrics to pathogen transmission mechanism

Prior studies (e.g., Dobson 2004; Rudolf and Antonovics 2005; Mihaljevic et al. 2014; Faust

et al. 2017) suggest that host additions decrease all three disease metrics more often under

frequency-dependent direct transmission (FDDT) than density-dependent direct transmission

(DDDT), but not always for focal host infection prevalence (Cortez and Duffy, 2021; Cortez,

2021). I explore this further by comparing how the response of each disease metric to host

additions depends on the pathogen transmission mechanism.

To explain the approach, I use $\mathcal{R}_0^C$ as a concrete example (see Appendix S1.4.3 for

mathematical details). The goal is to determine if larger increases or larger decreases in

$\mathcal{R}_0^C$ occur under FDDT or DDDT when a subcommunity of m resident species ($m < n$)

has species $m + 1, ..., n$ added to it. To do this, I use a new parameter, q, that converts

model (2) from a FDDT form ($\delta = 0$) to a DDDT form ($u_{S_i} = u_{I_i} = u_{R_i} = 0$). To allow

for a fair comparison between the FDDT and DDDT forms of the model, the change of

parameters varies the host uptake and degradation rates while holding constant the value

of $\mathcal{R}_0^C$ for the m-species subcommunity. I then compute how $\mathcal{R}_0^C$ for the full n-species

community changes using the sensitivity $\partial \mathcal{R}_0^C / \partial q$. Positive values mean host additions

cause larger decreases (or smaller increases) in $\mathcal{R}_0^C$ under FDDT than DDDT. Negative

values mean that host additions cause larger increases (or smaller decreases) in $\mathcal{R}_0^C$ under

FDDT than DDDT. The same approach is used for focal host infection prevalence and

infected density, with the only difference being that the change of parameters holds constant

the host and propagule densities at the endemic equilibrium for the m-species subcommunity;

this allows for a fair comparison with those two metrics (see Appendix S1.5.4 for details).

Community reproduction number: For $\mathcal{R}_0^C$, host additions cause larger decreases

(or smaller increases) in $\mathcal{R}_0^C$ under FDDT than DDDT if (i) there is no interspecific competition

(Figure 5A), (ii) added host species are sufficiently weak intraspecific and interspecific

competitors, or (iii) added host species have higher uptake rates than resident host species

(black circles and red triangles in Figure 5D). However, additions can cause larger increases

(or smaller decreases) in $\mathcal{R}_0^C$ under FDDT than DDDT when added host species are stronger

intraspecific and interspecific competitors and have lower uptake rates than resident host

species (blue squares in Figure 5D).

Focal host infection prevalence: Host additions are more likely to cause larger

decreases (or smaller increases) in focal host infection prevalence under FDDT than DDDT

when the added host species (i) are weak intraspecific and interspecific competitors and

(ii) have lower competence than resident host species. For direct transmission models where

interspecific transmission rates are fractions of intraspecific transmission rates (i.e, $\hat{\beta}_{ij} =$

$c_j \hat{\beta}_{jj}$ where $c_j < 1$), condition (ii) implies that added host species have relatively lower

rates of interspecific transmission (smaller c_j) than resident host species; see Appendix

S2.5.4 for details. Host additions are more likely to cause larger increases (or smaller decreases)

in focal host infection prevalence under FDDT than DDDT when the opposite conditions

hold. Consequently, in the absence of interspecific competition, an added low competence

host causes smaller increases in focal host infection prevalence under FDDT than DDDT

(black circles and blue squares in Figure 5B), but an added high competence host can

cause the opposite (red triangles in Figure 5B). When interspecific competition is present,

an added host can cause larger increases in focal host infection prevalence under FDDT

than DDDT when it is a strong interspecific competitor or has high competence (blue

squares and red triangles in Figure 5E, respectively).

Focal host infected density: The predictions for focal host infected density are

often the same as focal host infection prevalence. However, the predictions can be reversed

if host additions cause large changes in focal host density (mechanism 3) because the focal

host is more strongly regulated by disease or interspecific competition than intraspecific

competition.

Comparison of metrics: The pathogen transmission mechanism often has effects of the same sign on all three metrics (black circles in Figure 5). In these cases, host additions cause larger decreases (or smaller increases) in all metrics either under FDDT or under DDDT. However, the pathogen transmission mechanism can have effects of different signs on the metrics under two conditions. First, responses in $\mathcal{R}_0^C$ can differ in sign from the other two metrics when the added host species are large sources (red triangles in Figure 5D-F). This is due to density-mediated feedbacks (mechanism 2) where the added large sources increase infectious propagule density, which increases disease-induced mortality in the focal host. Second, as noted above, the responses in focal host infection prevalence and infected density can have opposite signs. This occurs when there are large changes in focal host density because the focal host is more strongly regulated by disease or interspecific competition than intraspecific competition (blue squares and red triangles in Figure 5BC).

Explaining prior results about richness-disease relationships: These predictions explain patterns in prior studies comparing DDDT and FDDT models. First, the $\mathcal{R}_0^C$ predictions explain why $\mathcal{R}_0^C$ increased more often under DDDT than FDDT in the absence of interspecific competition (Dobson, 2004; Rudolf and Antonovics, 2005; Joseph et al., 2013; Mihaljevic et al., 2014), but not when interspecific competition was stronger (Joseph et al., 2013; Mihaljevic et al., 2014). Second, Rudolf and Antonovics (2005) and Faust et al. (2017) observed that host additions decreased focal host infection prevalence more often under FDDT than DDDT because those studies assumed low interspecific transmission and no interspecific competition, respectively.

4.6 Application to model of *Daphnia*-fungus system

To show how my theory can help explain the differing effects host additions have on the disease metrics, I apply my theory to the parameterized model in Searle et al. (2016). Here, I present a summary of the results and refer the reader to Appendix S3.2.5 for details

about the analysis and to Searle et al. (2016) for details about the empirical system and

model parameterization.

The model describes the dynamics of two host species, *Daphnia dentifera* and *D. lumholtzi*,

and an environmentally transmitted fungal pathogen, *Metschnikowia bicuspidata*. The

model assumes Lotka-Volterra competition and the structure is identical to model (2)

except there are no recovered classes (R_i) because infected individuals do not recover from

infection. To avoid confusion, the host species are indexed using $i = D$ for *D. dentifera*

and $i = L$ for *D. lumholtzi*.

The estimated parameter values from Searle et al. (2016) show that *D. dentifera* has

lower competence than *D. lumholtzi*, i.e., $\mathcal{R}_{0,D}(\infty) = 1.7$ is less than $\mathcal{R}_{0,L}(\infty) = 6$, and is

a smaller source, i.e., $0 < \varphi_D < \varphi_L$. In addition, *D. dentifera* is a weaker intraspecific and

interspecific competitor than *D. lumholtzi*, i.e., $\alpha_{DD} < \alpha_{LL}$ and $\alpha_{DL} < 0 < \alpha_{LD}$ in equation

(3). The small positive direct effect of *D. dentifera* on *D. lumholtzi* ($0 < \alpha_{LD}$) is likely

an artifact due to fitting noisy data and it is expected that the actual value is small and

negative (Searle et al., 2016). In any case, the predicted effects of host additions on each

disease metric are qualitatively unchanged if a small positive or a small negative value is

used for α_{DL}; see Appendix S3.2.5. For the estimated parameter values, the host species

can stably coexist in the absence of the pathogen (at a two-host disease-free equilibrium)

and each host can stably coexist with the pathogen (at a one-host endemic equilibrium).

Community reproduction number: The pathogen's reproduction number is lowest

when *D. dentifera* is alone ($\mathcal{R}_{0,D} = 1.5$), highest when *D. lumholtzi* is alone ($\mathcal{R}_{0,L}^C =$

2.9), and intermediate when both hosts are present ($\mathcal{R}_0^C = 2.8$). This means that $\mathcal{R}_0^C$

increases with additions of *D. lumholtzi* and decreases with additions of *D. dentifera*. The

explanations for the two different effects on $\mathcal{R}_0^C$ are the following.

First, the addition of *D. lumholtzi* to a community with just *D. dentifera* increases

$\mathcal{R}_0^C$. This is because the addition of *D. lumholtzi* causes an increase in the density of a

high competence host (*D. lumholtzi*) and the decrease in the density of a low competence

host (*D. dentifera*) due to competition. Mathematically, we see this in equation (9) where the first and second terms, respectively, are positive because (1) the competence of *D. lumholtzi* is large, making $\mathcal{R}_{0,L}(\infty) - \mathcal{R}_0^C$ large and positive; (2) the competence of *D. dentifera* is small, making $\mathcal{R}_{0,D}(\infty) - \mathcal{R}_0^C$ negative; and (3) *D. lumholtzi* has a negative competitive effect on *D. dentifera*, making $\alpha_{DL}^{total\ddagger}$ negative. Overall, equation (9) shows that increased density of *D. lumholtzi* increases $\mathcal{R}_0^C$, which agrees with $\mathcal{R}_0^C$ being larger when *D. lumholtzi* is added to a community with just *D. dentifera*.

Second, the addition of *D. dentifera* to a community with just *D. lumholtzi* reduces $\mathcal{R}_0^C$. This is because the addition of *D. dentifera* causes a large increase in the density of a lower competence host (*D. dentifera*) and a small increase in the density of a higher competence host (*D. lumholtzi*) due to the positive interspecific effect. Mathematically, we see this in equation (9) where the first term is large and negative and the second term is smaller and positive, respectively, because (1) the competence of *D. dentifera* is small, making $\mathcal{R}_{0,D}(\infty) - \mathcal{R}_0^C$ negative; (2) the competence of *D. lumholtzi* is large, making $\mathcal{R}_{0,L}(\infty) - \mathcal{R}_0^C$ smaller and positive; and (3) *D. dentifera* has a small positive direct effect on *D. lumholtzi*, making $\alpha_{DL}^{total\ddagger}$ positive and small. Overall, equation (9) shows that increased density of *D. dentifera* decreases $\mathcal{R}_0^C$, which agrees with $\mathcal{R}_0^C$ being larger when *D. dentifera* is added to a community with just *D. lumholtzi*.

Focal host infection prevalence: *D. dentifera* and *D. lumholtzi* both cause infection prevalence in the other species to increase. The explanations for the similar responses are the following.

First, the addition of *D. lumholtzi* causes infection prevalence in *D. dentifera* to increase. In this case, the positive effect of adding the large source *D. lumholtzi* is greater than the negative effect of reduced density of the small source *D. dentifera*. Mathematically, we see this in equation (10) where the first term is large and positive and the second term is smaller and negative, respectively, because (1) *D. lumholtzi* is a large source ($\varphi_L > 0$ large); (2) *D. dentifera* is a small source ($\varphi_D > 0$ small); and (3) *D. lumholtzi* has a

negative competitive effect on *D. dentifera* ($\alpha_{DL}^{total*} < 0$). Thus, equation (10) shows that

increased density of *D. lumholtzi* increases infection prevalence in *D. dentifera*, which agrees

with the observed response.

Second, the addition of *D. dentifera* causes infection prevalence in *D. lumholtzi* to

increase. This is caused by the positive effect of adding the small source *D. dentifera* and

the positive effect of increased density of the large source *D. lumholtzi*. Mathematically,

we see this in equation (10) where the first and second terms are positive, respectively,

because (1) *D. dentifera* is a small source ($\varphi_D > 0$ large); (2) *D. lumholtzi* is a large

source ($\varphi_L > 0$ small); and (3) *D. dentifera* has a positive effect on *D. lumholtzi* ($\alpha_{LD}^{total*} >$

0). Thus, equation (10) shows that increased density of *D. dentifera* increases infection

prevalence in *D. lumholtzi*, which agrees with the observed response.

Focal host infected density: The density of infected *D. dentifera* decreases whereas

the density of infected *D. lumholtzi* increases when the other host species is added. The

explanations for the two different effects on infected density are the following.

First, the addition of *D. lumholtzi* causes the density of infected *D. dentifera* to decrease.

Infected density and infection prevalence in *D. dentifera* have responses of opposite signs

because the addition of *D. lumholtzi* causes a large decrease in the total density of *D.*

dentifera. The large decrease in total density of *D. dentifera* is due to *D. lumholtzi* being

a strong interspecific competitor. This aligns with condition (ii) because the focal host

D. dentifera is more strongly regulated by interspecific competition than by intraspecific

competition.

Second, the addition of *D. dentifera* causes the density of infected *D. lumholtzi* to increase.

Infected density and infection prevalence in *D. lumholtzi* have responses of the same sign

because the addition of *D. dentifera* does not cause a large increase or decrease in the

total density of *D. lumholtzi*. Moreover, condition (i) is not satisfied because *D. lumholtzi*

is strongly regulated by intraspecific competition and weakly regulated by disease and

interspecific competition. Condition (ii) is not satisfied because the focal host *D. lumholtzi*

is not a small source.

Comparison of metrics: For both host species, the addition of a second host has

different effects on the three metrics. The addition of *D. lumoltzi* causes $\mathcal{R}_0^C$ and infection

prevalence in *D. dentifera* to increase, but infected density in *D. dentifera* to decrease.

The differing response for infected density is due large changes in the total density of *D.*

dentifera (mechanism 3), driven by *D. dentifera* being more strongly regulated by interspecific

competition with *D. lumholtzi* than by intraspecific competition (condition ii). In comparison,

the addition of *D. dentifera* causes $\mathcal{R}_0^C$ to decrease but infection prevalence and infected

density in *D. lumholtzi* to increase. The differing response for $\mathcal{R}_0^C$ is due to differences in

the definition of competence and source/sink (mechanism 1). In particular, *D. dentifera* is

a lower competence host, which leads to lower $\mathcal{R}_0^C$, but also a small source, which leads to

higher infection prevalence and infected density in *D. lumholtzi*.

5   Discussion

Across empirical studies, increased host species richness can decrease (dilute) or increase

(amplify) disease risk (Keesing et al., 2010; Randolph and Dobson, 2012; Salkeld et al.,

2013; Wood et al., 2014; Civitello et al., 2015; Ostfeld and Keesing, 2017; Rohr et al.,

2019). To understand the variation in responses, theory is needed that identifies the rules

driving context-dependent biodiversity-disease relationships and explains how predicted

outcomes depend on the choice of disease metric (Buhnerkempe et al., 2015; Halsey, 2019;

Rohr et al., 2019). Prior theoretical studies have shown that disease levels and risk can

increase or decreases with increased host species richness, depending on mechanisms related

to host competence, changes in host densities, and transmission (Dobson, 2004; Joseph

et al., 2013; Mihaljevic et al., 2014; Faust et al., 2017; Cortez, 2021). The main goal of

this study was to synthesize this prior theoretical work and clarify how differences between

model predictions were driven by differing assumptions about host competence, interspecific

host competition, the pathogen transmission mechanism, and the metric of disease. My
results (Tables 1 and 2) help explain the variation in predictions across theoretical studies
and identify the rules governing the shapes of host species richness-disease relationships.

Accurate predictions about host richness-disease relationships require measurements of
the ability of each host species to transmit the pathogen, and critically, how that ability
is measured depends on the disease metric of interest. Specifically, predictions about $\mathcal{R}_0^C$
require estimates of life-time production of infectious propagules given exposure (i.e, competence,
$\mathcal{R}_{0,i}(\infty)$, as defined in Merrill and Johnson (2020)). In comparison, predictions about
focal host infection prevalence and infected density require estimates of instantaneous
per capita net production rates of infectious propagules (ϕ_i). In prior studies, a host's
ability to transmit the pathogen has been measured in many ways, including infection
intensity given exposure (Johnson et al., 2013), quantities that include transmission rates
to conspecific and heterospecifics (Gervasi et al., 2015; Downs et al., 2019), and the pathogen's
basic reproduction number for each species when alone (Strauss et al., 2015). These metrics
differ from the current study's definitions of competence and sink/source because they
include additional factors (e.g., host densities) or omit factors (e.g., the average length of
infection, $1/[m_i + \mu_i + \nu_i]$). As a result, the metrics used in these empirical studies could
lead to inaccurate predictions about host species richness-disease relationships in nature.

Accurate predictions about host richness-disease relationships also require knowledge
about density of each host species responses to altered host species richness. One component
of this (only relevant to infection prevalence and infected density) is changes in host densities
due to density-mediated feedbacks driven by disease-induced mortality (mechanism 2).
The other component (relevant to all metrics) is changes in host densities due to interspecific
competition. Prior theoretical studies (Rudolf and Antonovics, 2005; Joseph et al., 2013;
Mihaljevic et al., 2014; O'Regan et al., 2015; Faust et al., 2017) have shown that host
richness-disease relationships can differ depending on whether host additions and removals
have additive effects on host density (corresponding to no interspecific competition) or

substitutive effects on host density (corresponding to strong, symmetric interspecific competition where total host density is held constant); these effects were referred to as subtractive and compensatory effects in Joseph et al. (2013). Because both of these scenarios assume symmetric competitive effects, my results extend this prior work by explaining how asymmetric interspecific competitive effects can be important for shaping host richness-disease relationships. For example, added low competence sink hosts can increase all disease metrics if they have stronger competitive effects on low competence sink hosts than high competence source hosts (Figure 4).

In combination, my results show that accurate predictions about host richness-disease relationships require detailed information about specific measures of each host's ability to transmit the pathogen and changes in host densities in response to altered host species richness. While all of these quantities can be estimated in studies of tractable laboratory systems (Cáceres et al., 2014; Strauss et al., 2015; Searle et al., 2016) or systems where interspecific competition is likely negligible (Fenton et al., 2015), they are likely to be difficult to estimate for many field systems. My analytical results and predictions suggest that transmission abilities (competence and sink/source) are the most important quantities to estimate. The reasoning is that if the transmission ability of an added or removed host correctly predicts the response in a metric, then this suggests the effects of altered host densities are either small in magnitude or in the same direction (i.e., a synergistic effect). For example, Venesky et al. (2014) estimated how the presence and absence of three amphibian species affected the intensity of *Batrachochytrium dendrobatidis* infections in laboratory experiments and found qualitatively similar relationships between presence/absence and infection prevalence in a field study. This agreement suggests that estimates of changes in host densities are not needed to accurately predict the qualitative effect (i.e., increases or decreases in infection prevalence) of altered host species richness. However, an important caveat to the above is that one needs to use the correct quantity when estimating the ability of each host species to transmit the disease; the Venesky et al. (2014) study is not

$_{931}$ ideal in this regard because its metric of transmission ability (infection intensity) does

$_{932}$ not account for the production of new infectious propagules by infected individuals. A

$_{933}$ second important caveat is that without predictions about changes in host densities, it is

$_{934}$ impossible to predict, *a priori*, if infection prevalence and infected density will respond in

$_{935}$ opposite directions.

$_{936}$ My work builds on prior studies that compared the responses of different disease metrics

$_{937}$ (Roche et al., 2012; Roberts and Heesterbeek, 2018) by identifying when and why the

$_{938}$ three metrics of disease change in the same or different directions. Agreement across all

$_{939}$ metrics is an ideal situation for disease management strategies that use culling or host

$_{940}$ removal (Laurenson et al., 2004; Donnelly et al., 2006) because disease risk and disease

$_{941}$ burden can be simultaneously minimized by removing or suppressing specific alternative

$_{942}$ host species in a community. However, cases where the metrics respond differently imply

$_{943}$ the presence of trade-offs. Opposing responses in $\mathcal{R}_0^C$ and focal host infection prevalence

$_{944}$ (mechanism 1 or 2) produce a trade-off between the risk of an outbreak and the risk of a

$_{945}$ focal host individual becoming infected. For example, additions of *D. dentifera* lower risk

$_{946}$ the risk of an outbreak for *D. lumhotzi* by reducing $\mathcal{R}_0^C$, but increase the risk of infection

$_{947}$ for *D. lumhotzi* when an outbreak occurs by increasing infection prevalence in *D. lumhotzi*

$_{948}$ (see "Application to model of *Daphnia*-fungus systems"). In comparison, opposing responses

$_{949}$ in focal host infection prevalence and infected density (mechanism 2) produce a trade-

$_{950}$ off between infection risk for focal host individuals (infection prevalence) and the risk of

$_{951}$ spillover to other species (for which infected density is considered a better metric; Roche

$_{952}$ et al. 2012; Wood and Lafferty 2013). Thus, my work helps identify when disease management

$_{953}$ or conservation strategies need to account for trade-offs between different metrics of disease

$_{954}$ risk.

$_{955}$ My results support prior claims that altered host species richness has context-dependent

$_{956}$ effects on disease risk (Randolph and Dobson, 2012; Salkeld et al., 2013; Wood et al., 2016),

$_{957}$ while also providing testable predictions about the mechanisms driving those context-

dependent effects (Table 2). For example, increases in focal host infection prevalence are predicted to be driven by later added hosts being sources (black curves in Figure 4B) or having stronger competitive effects on sink hosts than source hosts (blue curves in Figure 4E). In comparison, decreases in focal host infection prevalence are predicted to be driven by later added hosts being sinks (black curves in Figure 4E) or having stronger competitive effects on source hosts than sink hosts (blue curves in Figure 4B). Pairing my work with knowledge about how host competence (or sink/source) and competitive effect are correlated with community (dis)assembly also could help explain patterns identified across empirical studies. As one example, based on life-history theory and hypotheses about pathogen adaptation, prior studies (reviewed in Ostfeld and Keesing (2012); Joseph et al. (2013); Huang et al. (2019)) have argued that more competent host species are more likely to be present in low richness communities whereas higher competence host species will only be present in high richness communities. The resulting negative relationship between arrival order and competence can lead to negative relationships between richness and disease (Joseph et al., 2013). However, my results explain why interspecific competition could alter those predictions and cause positive relationships between richness and disease (bottom row of Figure 4) that were also observed in prior theoretical work (Joseph et al., 2013). As a second example, Halliday and Rohr (2019) found that biodiversity gradients driven by mechanisms associated with biodiversity loss (e.g., habitat fragmentation or urbanization) often result in negative host species richness-disease relationships whereas biodiversity gradients driven by other mechanisms (e.g., environmental heterogeneity or elevation) can yield positive or negative host species richness-disease relationships. My theory suggests that these patterns could be driven by different relationships between extinction order and competence or different relationships between extinction order and host densities (due to competition or increased mortality, e.g., due to habitat reduction). More generally, pairing my work with system-specific knowledge about correlations between species traits and extinction/arrival order could explain the different frequencies of unimodal and monotonic

relationships between richness and disease in empirical studies (Halliday and Rohr, 2019).

While this study explains prior theoretical work and has the potential to explain empirical patterns, additional theory is needed to fully explain the observed variation in empirical host species richness-disease relationships. Specific areas directly related model (2) include incorporating the effects of dose-dependent infection and mortality rates (e.g., Clay et al. 2021) and transmission via both direct and environmental pathways (e.g., Eisenberg et al. 2013); these were not addressed in this study because the supplementary material is already at a ridiculous length. More generally, advances in theory are needed in four areas. First, theory is needed for other metrics used in empirical studies, including community prevalence (Suzán et al., 2009; Searle et al., 2011), disease incidence (Haas et al., 2011), infectious propagule density (Roberts and Heesterbeek, 2018; Cortez and Duffy, 2020), and infection intensity (Hechinger and Lafferty, 2005; Becker et al., 2014; Venesky et al., 2014). This is important because the metrics can provide different and complementary information about disease burden and disease risk. Second, more theory is needed to extend our understanding of multi-host systems where host species have non-competitive interspecific interactions (Rohr et al., 2015), including systems where pathogens obligately switch between host species during their life cycle (Hechinger and Lafferty, 2005; Hopkins et al., 2016). Third, additional theory is needed to explain why increased host species richness often, but not always, decreases disease for vector-borne pathogens in empirical studies (Ezenwa et al., 2006; Keesing et al., 2009; Moore and Borer, 2012; Xavier et al., 2012) and theoretical studies (Saul, 2003; Normal et al., 1999; Ogden and Tsao, 2009; Miller and Huppert, 2013). Because vectors are analogous to infectious propagules, some of the variation may be explained by the net production of infected vectors by each host species (analogous to φ_i). However, this analogy is limited because vectors can have complex dynamics and behavior (e.g., biting preferences), which may influence how host additions affect disease metrics (Miller and Huppert, 2013; Marini et al., 2017). Fourth, new theory is needed to explain why host species can have temporally varying effects on disease metrics. For example, empirical

studies show that an additional host species can have effects of the same sign at all points in time (e.g., always decreasing focal host infection prevalence; Strauss et al. 2015) or effects of varying sign (e.g., transition from decreased to increased focal host infection prevalence; Dallas et al. 2016). My results suggest a possible explanation for the temporally varying effects: a low competence source host could decrease infection prevalence early in time because decreases in $\mathcal{R}_0^C$ imply slower rates of spread, but also increase infection prevalence later in time because it is a net producer of infectious propagules. Recent work (Hanthanan Arachchilage et al., 2023) suggests sensitivity-based approaches could be fruitful, however new theory is needed to fully explain when and why host species have temporally variable effects on disease dynamics. Combined, advances in these four areas will help address a current need to understand how the interactions between parasites, pathogens, and their ecosystems shape disease burden and the risk of disease emergence (Cunningham et al., 2017).

This study has made a step towards addressing that need by providing a framework that unifies prior theoretical studies and identifies the rules governing the context-dependent relationships between host species richness and different metrics of disease. By providing testable predictions about underlying mechanisms, this study may help resolve some of the debate about the variation between and frequencies of different patterns of biodiversity and disease.

6   Acknowledgments

I thank MA Duffy, the Duffy lab, the Ecology Reading Group in FSU's Department of Biological Science, and three anonymous reviewers for helpful comments and feedback on the manuscript. MHC was supported by the National Science Foundation under Award DEB-2015280.

References

Bartholomew, J. L., and P. W. Reno. 2002. The history and dissemination of whirling disease. Pages 3–24 *in* American fisheries society symposium. American Fisheries Society.

Becker, C. G., D. Rodriguez, L. F. Toledo, A. V. Longo, C. Lambertini, D. T. Corrêa, D. S. Leite, C. F. Haddad, and K. R. Zamudio. 2014. Partitioning the net effect of host diversity on an emerging amphibian pathogen. Proceedings of the Royal Society B: Biological Sciences 281:20141796.

Begon, M., and R. G. Bowers. 1994. Host-host-pathogen models and microbial pest control: the effect of host self regulation. Journal of Theoretical Biology 169:275–287.

Begon, M., R. G. Bowers, N. Kadianakis, and D. E. Hodgkinson. 1992. Disease and community structure: the importance of host self-regulation in a host-host-pathogen model. American Naturalist 139:1131–1150.

Bernardo, M. A., and M. S. Singer. 2017. Parasite-altered feeding behavior in insects: integrating functional and mechanistic research frontiers. Journal of Experimental Biology 220:2848–2857.

Bowers, R. G., and J. Turner. 1997. Community structure and the interplay between interspecific infection and competition. Journal of Theoretical Biology 187:95–109.

Buhnerkempe, M. G., M. G. Roberts, A. P. Dobson, H. Heesterbeek, P. J. Hudson, and J. O. Lloyd-Smith. 2015. Eight challenges in modelling disease ecology in multi-host, multi-agent systems. Epidemics 10:26–30.

Cáceres, C. E., G. Davis, S. Duple, S. R. Hall, A. Koss, P. Lee, and Z. Rapti. 2014. Complex *Daphnia* interactions with parasites and competitors. Mathematical Biosciences 258:148–161.

Caswell, H. 2019. Sensitivity analysis: matrix methods in demography and ecology. Springer Nature.

Civitello, D. J., J. Cohen, H. Fatima, N. T. Halstead, J. Liriano, T. A. McMahon, C. N. Ortega, E. L. Sauer, T. Sehgal, S. Young, and J. R. Rohr. 2015. Biodiversity inhibits parasites: Broad evidence for the dilution effect. PNAS 112:8667–8671.

Clay, P. A., M. H. Cortez, and M. A. Duffy. 2021. Dose relationships can exacerbate, mute, or reverse the impact of heterospecific host density on infection prevalence. Ecology 102:e03422.

Cortez, M. H. 2021. Using sensitivity analysis to identify factors promoting higher versus lower infection prevalence in multi-host communities. Journal of Theoretical Biology 526:110766.

Cortez, M. H., and M. A. Duffy. 2020. Comparing the indirect effects between exploiters in predator-prey and host-pathogen systems. The American Naturalist 196:to appear.

———. 2021. The context dependent effects of interspecific host competition and the pathogen transmission mode on disease prevalence. The American Naturalist page submitted.

Cortez, M. H., and J. S. Weitz. 2013. Distinguishing between indirect and direct modes of transmission using epidemiological time series. American Naturalist 181:E43–E52.

Cunningham, A. A., P. Daszak, and J. L. Wood. 2017. One health, emerging infectious diseases and wildlife: two decades of progress? Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160167.

Dallas, T., R. J. Hall, and J. M. Drake. 2016. Competition-mediated feedbacks in experimental multispecies epizootics. Ecology 97:661–670.

Daszak, P., A. A. Cunningham, and A. D. Hyatt. 2003. Infectious disease and amphibian population declines. Diversity and Distributions 9:141–150.

Diekmann, O., J. Heesterbeek, and M. G. Roberts. 2010. The construction of next-generation matrices for compartmental epidemic models. Journal of the royal society interface 7:873–885.

Dobson, A. 2004. Population dynamics of pathogens with multiple host species. American Naturalist 164:S64–S78.

Donnelly, C. A., R. Woodroffe, D. Cox, F. J. Bourne, C. Cheeseman, R. S. Clifton-Hadley, G. Wei, G. Gettinby, P. Gilks, H. Jenkins, W. T. Johnston, A. M. Le Fevre, J. P. McInerney, and W. I. Morrison. 2006. Positive and negative effects of widespread badger culling on tuberculosis in cattle. Nature 439:843–846.

Downs, C. J., L. A. Schoenle, B. A. Han, J. F. Harrison, and L. B. Martin. 2019. Scaling of host competence. Trends in parasitology 35:182–192.

Eisenberg, M. C., S. L. Robertson, and J. H. Tien. 2013. Identifiability and estimation of multiple transmission pathways in cholera and waterborne disease. Journal of Theoretical Biology 324:84–102.

Espira, L. M., A. F. Brouwer, B. A. Han, J. Foufopoulos, and J. N. Eisenberg. 2022. Dilution of epidemic potential of environmentally transmitted infectious diseases for species with partially overlapping habitats. The American Naturalist 199:E43–E56.

Ezenwa, V. O., M. S. Godsey, R. J. King, and S. C. Guptill. 2006. Avian diversity and west nile virus: testing associations between biodiversity and infectious disease risk. Proceedings of the Royal Society B: Biological Sciences 273:109–117.

Faust, C. L., A. P. Dobson, N. Gottdenker, L. S. Bloomfield, H. I. McCallum, T. R. Gillespie, M. Diuk-Wasser, and R. K. Plowright. 2017. Null expectations for disease

dynamics in shrinking habitat: dilution or amplification? Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160173.

Feng, Z., R. Liu, D. L. DeAngelis, J. P. Bryant, K. Kielland, F. S. C. III, and R. K. Swihart. 2009. Plant toxicity, adaptive herbivory, and plant community dynamics. Ecosystems 12:534–547.

Fenton, A., D. G. Streicker, O. L. Petchey, and A. B. Pedersen. 2015. Are all hosts created equal? partitioning host species contributions to parasite persistence in multihost communities. The American Naturalist 186:610–622.

Gervasi, S. S., D. J. Civitello, H. J. Kilvitis, and L. B. Martin. 2015. The context of host competence: a role for plasticity in host–parasite dynamics. Trends in parasitology 31:419–425.

Haas, S. E., M. B. Hooten, D. M. Rizzo, and R. K. Meentemeyer. 2011. Forest species diversity reduces disease risk in a generalist plant pathogen invasion. Ecology Letters 14:1108–1116.

Halliday, F. W., and J. R. Rohr. 2019. Measuring the shape of the biodiversity-disease relationship across systems reveals new findings and key gaps. Nature communications 10:1–10.

Halsey, S. 2019. Defuse the dilution effect debate. Nature Ecology & Evolution 3:145.

Hanthanan Arachchilage, K., M. Y. Hussaini, N. G. Cogan, and M. H. Cortez. 2023. Exploring how ecological and epidemiological processes shape multi-host disease dynamics using global sensitivity analysis. Journal of Mathematical Biology page accepted.

Hechinger, R. F., and K. D. Lafferty. 2005. Host diversity begets parasite diversity: bird

final hosts and trematodes in snail intermediate hosts. Proceedings of the Royal Society B: Biological Sciences 272:1059–1066.

Hedrick, R. P., M. A. Adkison, M. El-Matbouli, and E. MacConnell. 1998. Whirling disease: re-emergence among wild trout. Immunological reviews 166:365–376.

Hopkins, S. R., J. M. Ocampo, J. M. Wojdak, and L. K. Belden. 2016. Host community composition and defensive symbionts determine trematode parasite abundance in host communities. Ecosphere 7:e01278.

Huang, J., S. Ruan, Y. Shu, and X. Wu. 2019. Modeling the transmission dynamics of rabies for dog, chinese ferret badger and human interactions in zhejiang province, china. Bulletin of mathematical biology 81:939–962.

Johnson, P., R. S. Ostfeld, and F. Keesing. 2015. Frontiers in research on biodiversity and disease. Ecology Letters 18:1119–1133.

Johnson, P. T. J., D. L. Preston, J. T. Hoverman, J. S. Henderson, S. H. Paull, K. L. D. Richgels, and M. D. Redmond. 2012. Species diversity reduces parasite infection through cross-generational effects on host abundance. Ecology 93:56–64.

Johnson, P. T. J., D. L. Preston, J. T. Hoverman, and K. L. D. Richgels. 2013. Biodiversity decreases disease through predictable changes in host community competence. Nature 494:230–234.

Johnson, P. T. J., and D. W. Thieltges. 2010. Diversity, decoys and the dilution effect: how ecological communities affect disease risk. The Journal of Experimental Biology 213:961–970.

Joseph, M. B., J. R. Mihaljevic, S. A. Orlofske, and S. H. Paull. 2013. Does life history mediate changing disease risk when communities disassemble? Ecology letters 16:1405–1412.

Kaper, J. B., J. G. Morris, Jr., and M. M. Levine. 1995. Cholera. Clinical Microbiological Review 8:48–86.

Keesing, F., L. K. Belden, P. Daszak, A. Dobson, and C. D. Harvell et al. 2010. Impacts of biodiversity on the emergence and transmission of infectious diseases. Nature 468:647–652.

Keesing, F., J. Brunner, S. Duerr, M. Killilea, K. LoGiudice, K. Schmidt, H. Vuong, and R. Ostfeld. 2009. Hosts as ecological traps for the vector of lyme disease. Proceedings of the Royal Society B: Biological Sciences 276:3911–3919.

Keesing, F., R. D. Holt, and R. S. Ostfeld. 2006. Effects of species diversity on disease risk. Ecology Letters 9:485–498.

Laurenson, M. K., R. Norman, L. Gilbert, H. W. Reid, and P. J. Hudson. 2004. Mountain hares, louping-ill, red grouse and harvesting: complex interactions but few data. Journal of Animal Ecology 73:811–813.

Marini, G., R. Rosá, A. Pugliese, and H. Heesterbeek. 2017. Exploring vector-borne infection ecology in multi-host communities: A case study of west nile virus. Journal of Theoretical Biology 415:58–69.

McCormack, R. K., and L. J. Allen. 2007. Disease emergence in multi-host epidemic models. Mathematical Medicine and Biology 24:17–34.

Merrill, T. E. S., and P. T. Johnson. 2020. Towards a mechanistic understanding of competence: a missing link in diversity–disease research. Parasitology 147:1159–1170.

Mihaljevic, J. R., M. B. Joseph, S. A. Orlofske, and S. H. Paull. 2014. The scaling of host density with richness affects the direction, shape, and detectability of diversity-disease relationships. PloS one 9:e97812.

Miller, E., and A. Huppert. 2013. The effects of host diversity on vector-borne disease: the conditions under which diversity will amplify or dilute the disease risk. PLoS One 8:e80279.

Moore, S. M., and E. T. Borer. 2012. The influence of host diversity and composition on epidemiological patterns at multiple spatial scales. Ecology 93:1095–1105.

Normal, R., R. G. Bowers, M. Begon, and P. J. Hudson. 1999. Persistence of tick-borne virus in the presence of multiple host species: Tick reservoirs and parasite mediated competition. Journal of Theoretical Biology 200:111–118.

Ogden, N. H., and J. I. Tsao. 2009. Biodiversity and lyme disease: Dilution or amplification? Epidemics 1:196–206.

O'Regan, S. M., J. E. Vinson, and A. W. Park. 2015. Interspecific contact and competition may affect the strength and direction of disease-diversity relationships for directly transmitted microparasites. The American Naturalist 186:480–494.

Ostfeld, R. S., and F. Keesing. 2000. Biodiversity series: the function of biodiversity in the ecology of vector-borne zoonotic diseases. Canadian Journal of Zoology 78:2061–2078.

———. 2012. Effects of host diversity on infectious disease. Annual review of ecology, evolution, and systematics 43:157–182.

———. 2017. Is biodiversity bad for your health? Ecosphere 8:e01676.

Peixoto, I. D., and G. Abramson. 2006. The effect of biodiversity on the hantavirus epizootic. Ecology 87:873–879.

Ponton, F., F. Lalubin, C. Fromont, K. Wilson, C. Behm, and S. J. Simpson. 2011. Hosts use altered macronutrient intake to circumvent parasite-induced reduction in fecundity. International journal for parasitology 41:43–50.

Randolph, S. E., and A. D. M. Dobson. 2012. Pangloss revisited: a critique of the dilution effect and the biodiversity-buffers-disease paradigm. Parasitology 139:847–863.

Roberts, M., and J. Heesterbeek. 2018. Quantifying the dilution effect for models in ecological epidemiology. Journal of The Royal Society Interface 15:20170791.

Roche, B., A. P. Dobson, J. F. Guegan, and P. Rohani. 2012. Linking community and disease ecology: the impact of biodiversity on pathogen transmission. Philosophical Transactions of the Royal Society B 367:2807–2813.

Rohr, J. R., D. J. Civitello, P. W. Crumrine, N. T. Halstead, A. D. Miller, A. M. Schotthoefer, C. Stenoien, L. B. Johnson, and V. R. Beasley. 2015. Predator diversity, intraguild predation, and indirect effects drive parasite transmission. Proceedings of the National Academy of Sciences 112:3008–3013.

Rohr, J. R., D. J. Civitello, F. W. Halliday, P. J. Hudson, K. D. Lafferty, C. L. Wood, and E. A. Mordecai. 2019. Towards common ground in the biodiversity–disease debate. Nature Ecology & Evolution 4:24–33.

Rudge, J. W., J. P. Webster, D.-B. Lu, T.-P. Wang, G.-R. Fang, and M.-G. Basáñez. 2013. Identifying host species driving transmission of schistosomiasis japonica, a multihost parasite system, in china. Proceedings of the National Academy of Sciences 110:11457–11462.

Rudolf, V. H. W., and J. Antonovics. 2005. Species coexistence and pathogens with frequency-dependent transmission. American Naturalist 166:112–118.

Salkeld, D. J., K. A. Padgett, and J. H. Jones. 2013. A meta-analysis suggesting that the relationship between biodiversity and risk of zoonotic pathogen transmission is idiosyncratic. Ecology Letters 16:679–686.

Saul, A. 2003. Zooprophylaxis or zoopotentiation: the outcome of introducing animals on vector transmission is highly dependent on the mosquito mortality while searching. Malaria Journal 2:1–18.

Searle, C. L., L. M. Biga, J. W. Spatafora, and A. R. Blaustein. 2011. A dilution effect in the emerging amphibian pathogen *Batrachochytrium dendrobatidis*. PNAS 108:16322–16326.

Searle, C. L., M. H. Cortez, K. K. Hunsberger, D. C. Grippi, I. A. Oleksy, C. L. Shaw, S. B. de la Serna, C. L. Lash, K. L. Dhir, and M. A. Duffy. 2016. Population density, not host competence, drives patterns of disease in an invaded community. American Naturalist 188:554–566.

Shikano, I., and J. S. Cory. 2016. Altered nutrient intake by baculovirus-challenged insects: Self-medication or compensatory feeding? Journal of invertebrate pathology 139:25–33.

Skerratt, L. F., L. Berger, R. Speare, S. Cashins, K. R. McDonald, A. D. Phillott, H. B. Hines, and N. Kenyon. 2007. Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. EcoHealth 4:125–134.

Strauss, A. T., D. J. Civitello, C. E. Cáceres, and S. R. Hall. 2015. Success, failure and ambiguity of the dilution effect among competitors. Ecology letters 18:916–926.

Suzán, G., E. Marcé, J. T. Giermakowski, J. N. Mills, G. Ceballos, R. S. Ostfeld, B. Armién, J. M. Pascale, and T. L. Yates. 2009. Experimental evidence for reduced rodent diversity causing increased hantavirus prevalence. PloS one 4.

Van den Driessche, P., and J. Watmough. 2008. Further notes on the basic reproduction number. Mathematical epidemiology pages 159–178.

Venesky, M. D., X. Liu, E. L. Sauer, and J. R. Rohr. 2014. Linking manipulative

experiments to field data to test the dilution effect. Journal of Animal Ecology 83:557–565.

Wolfe, M. S. 1992. Giardiasis. Clinical Microbiology Reviews 5:93–100.

Wood, C. L., and K. D. Lafferty. 2013. Biodiversity and disease: a synthesis of ecological perspectives on lyme disease transmission. Trends in Ecology and Evolution 28:239–247.

Wood, C. L., K. D. Lafferty, G. De Leo, H. S. Young, P. J. Hudson, and A. M. Kuris. 2014. Does biodiversity protect humans against infectious disease? Ecology 95:817–832.

Wood, C. L., K. D. Lafferty, G. DeLeo, H. S. Young, P. J. Hudson, and A. M. Kuris. 2016. Does biodiversity protect humans against infectious disease? Reply. Ecology .

Xavier, S. C. d. C., A. L. R. Roque, V. d. S. Lima, K. J. L. Monteiro, J. C. R. Otaviano, L. F. C. Ferreira da Silva, and A. M. Jansen. 2012. Lower richness of small wild mammal species and chagas disease risk. PLoS neglected tropical diseases 6:e1647.

 Table 1: Model variables, parameters, and important quantities

Notation	Description
Variables	
S_i	host i susceptible density
I_i	host i infected density
R_i	host i recovered density
$N_i = S_i + I_i + R_i$	host i total density
I_i/N_i	host i infection prevalence (i.e., proportion of infected individuals)
P	infectious propagule density
Parameters	
$\beta_i = \rho_i u_{S_i}$	transmission coefficient
ρ_i	per infectious propagule probability of infection m_i m
μ_i	disease-induced mortality rate
ν_i	recovery rate
ξ_i	rate of increased mortality due to prior infection
γ_i	rate of loss of immunity
χ_i	shedding rate of infectious propagules
$u_{S_i}, u_{I_i}, u_{R_i}$	uptake rates of infectious propagules for host classes S_i, I_i, R_i
δ	degradation rate of infectious propagules
$F_i(\cdot)$	density-dependent reproduction rate
	(Lotka-Volterra competition assumed in simulations)
r_i	maximum reproduction rate
b_i	reduction in reproduction rate for infected individuals
α_{ij}	per capita competitive effect of host j on host i
e_{ij}	proportional competitive effect of infected individuals;
	defines if infected individuals are stronger ($e_{ij} > 0$), equal (e_{ij}),
	or weaker ($e_{ij} < 1$) competitors than susceptible individuals
χ_{ij}	shedding rate of host j into the habitat of host i in the n-habitat model
$\hat{\beta}_{ij}$	direction transmission parameter for the transmission rate of host j
	to host i
Quantities	
$\mathcal{R}_0^C$	pathogen community reproduction number
φ_i	per capita net production of infectious propagules by host i;
	defines if host i is a source ($\varphi_i > 0$) or sink ($\varphi_i > 0$)
$\mathcal{R}_{0,i}(\infty) = \beta_i \chi_i / u_{S_i}[m_i + \mu_i + \nu_i]$	competence of host i
$\quad \rho_i \chi_i / [m_i + \mu_i + \nu_i]$	
α_{in}^{total}	total (direct + indirect) competitive effect of host n on host i
$p^{\ddagger}$	n-species disease-free equilibrium with densities $N_i^{\ddagger}$
p^*	n-species endemic equilibrium with densities N_i^*

Table 2: Predicted responses of three metric of disease to variation in components of alternative host competence

Component of alternative host competence	Community Reproduction No. $(\mathcal{R}_0^C)$	Refs.[†]	Focal host infection prevalence (I_1^*/N_1^*)	Refs.[†]	Focal host infection density (I_1^*)
Increased shedding (χ_i)	Always increases	[1]-[6]	Always increases	[7],[8]	Increases unless condition (i)
Increased uptake (u_{S_i})	Always decreases	[6]	Always decreases	[7],[8]	Decreases unless condition (i)
Increased transmission rate (β_i)	Always increases	[4],[6]	Increases unless (1) alternative host is a source and interspecific competition is strong, or (2) alternative host has larger negative competitive effects on sink hosts than source hosts and infected individuals are weaker competitors than susceptible individuals	[7],[8]	Same as infection prevalence unless conditions (i) or (ii)
Increased mortality rate (μ_i)	Always decreases		Decreases unless (1) alternative host is a large sink (2) alternative host has larger negative competitive effects on sinks hosts than source hosts	[7],[8]	Same as infection prevalence unless conditions (i) or (ii)
Increased recovery rate (ν_i)	Always decreases	[6]	Always decreases	[7],[8]	Same as infection prevalence unless conditions (i) or (ii)

Responses in focal host infected density can differ from the responses in focal host infection prevalence when (i) the focal host is more strongly regulated by disease and interspecific competition with alternative hosts than by intraspecific competition or (ii) the focal host is a sink or small source and some alternative hosts are large sources.

† Prior studies whose predictions are explained or generalized are [1] Dobson (2004), [2] (Huang et al., 2019), [3] Fenton et al. (2015), [4] O'Regan et al. (2015), [5] Joseph et al. (2013), [6] Espira et al. (2022), [7] Cortez and Duffy (2021), and [8] Cortez (2021). Mathematical details about each study are provided in Appendix S3.

Table 3: Predicted responses of three metrics of disease to the addition of a host species to a community

Trans. Mode	Interspecific Competition	Community Reproduction Number ($\mathcal{R}_0^C$)	Refs.[†]	Focal host infection prevalence (I_1^*/N_1^*)	Refs.[†]
DDDT	No	Always increases	[1]-[6]	Always increases	[8]-[10]
	Yes	Increases if intersp. competition is sufficiently weak	[4]-[7]	Increases if intersp. competition is sufficiently weak	[9]-[11]
		Decreases if intersp. competition is sufficiently strong	[4]-[7]	Decreases if intersp. competition is sufficiently strong	[7],[10]-[12]
FDDT	No	Increases if added host's competance and intersp. transmission rates are sufficiently high	[1],[4]-[6]	Increases if added host's intrasp. and intersp. transmission rates are sufficiently high	[9],[10]
		Decreases if added host's competance and and intersp. transmission rates are sufficiently low	[4],[6]	Decreases if added host's intrasp. and intersp. transmission rates are sufficiently low	[9]
	Yes	Increases if (i) added host's competance and intersp. transmission rates are sufficiently high, or (ii) hosts with lower competence and intersp. transmission rates decrease more than hosts with lower competence and intersp. transmission rates	[4],[6]	Increases if (i) added host's intrasp. and intersp. transmission rates are sufficiently high, or (ii) hosts with lower intrasp. and intersp. transmission rates decrease more than hosts with lower intrasp. and intersp. transmission rates	
		Decreases if (i) added host's competance and intersp. transmission rates are sufficiently low, or (ii) hosts with higher competence and intersp. transmission rates decrease more than hosts with higher competence and intersp. transmission rates	[4]-[6]	Decreases if (i) added host's intrasp. and intersp. transmission rates are sufficiently low, or (ii) hosts with higher intrasp. and intersp. transmission rates decrease more than hosts with higher intrasp. and intersp. transmission rates	[10]
ET	No	Increases if added host's competence or shedding rates are sufficiently high	[13]-[15]	Increases if added host is a source	[15]-[17]
		Decreases if added host's competence or shedding rates are sufficiently low	[14]	Decreases if added host is a sink	[16],[17]
	Yes	Increases if (i) added host's competence or shedding rates are sufficiently high, or (ii) hosts with lower competence and shedding rates decrease more than hosts with higher competence and shedding rates		Increases (i) if added host is a source, or (ii) sink hosts decrease more than source hosts	[16]-[18]
		Decreases if (i) added host's competence or shedding rates are sufficiently low, or (ii) hosts with higher competence and shedding rates decrease more than hosts with lower competence and shedding rates		Decreases (i) if added host is a sink, or (ii) source hosts decrease more than sink hosts	[16]-[18]
Focal host infected density (I_1^*)		Predictions are the same as focal host infection prevalence unless (i) the focal host is more strongly regulated by disease and intersp. competition than intrasp. competition (ii) the focal host is a sink or small source and some alternative hosts are large sources			[7],[19] [18]

ET = environmental transmission; FDDT = frequency-dependent direct transmission; DDDT = density-dependent direct

transmission; trans. = transmission; intrasp. = intraspecific; intersp. = interspecific.

† Prior studies whose predictions are explained or generalized are [1] Dobson (2004), [2] McCormack and Allen (2007), [3]

Huang et al. (2019), [4] Joseph et al. (2013), [5] Mihaljevic et al. (2014), [6] O'Regan et al. (2015), [7] Roberts and Heesterbeek

(2018), [8] Begon et al. (1992), [9] Faust et al. (2017), [10] Rudolf and Antonovics (2005), [11] Bowers and Turner (1997), [12]

Peixoto and Abramson (2006), [13] Fenton et al. (2015), [14] Espira et al. (2022), [15] Rudge et al. (2013), [16] Cortez and

Duffy (2021), [17] Cortez (2021), [18] Searle et al. (2016), and [19] Cáceres et al. (2014). Mathematical details about each

study are provided in Appendix S3.

 # 7   Figures

Figure 1: Increased competence of an alternative host increases all metrics of disease (black circles), unless there are strong density-mediated feedbacks due to disease-induced mortality (mechanism 2; blue squares) or large changes in focal host density (mechanism 3; red triangles). Note that the x-axes in the bottom row are reversed. (A,D) $\mathcal{R}_0^C$ always increases with increased transmission rates and decreased disease-induced mortality rates of an alternative host. (B,E) Focal host infection prevalence (I_1^*/N_1^*) increases with increased transmission rates and decreased disease-induced mortality rates of an alternative host (black circles and red triangles in B,E), unless there are strong density-mediated feedbacks due to disease-induced mortality (mechanism 2), which occurs when interspecific competition is sufficiently large (blue squares in B) or the alternative host is a large sink (blue squares in E). (C,F) Focal host infected density (I_1^*) responds in the same way as focal host infection prevalence (black circles and blue squares), unless there are large changes in focal host density (mechanism 3), which occurs when the focal host is more strongly regulated by the disease or interspecific competition than intraspecific competition (red triangles in C) or the focal host is a sink and other hosts are large sources (red triangles in F). In each row, curves of the same color are from the same parameterization of a four-host version of model (2); see Appendix S4 for parameter values.

Figure 2: Strong interspecific competition can reverse predictions about the effects of increased aded host density on each disease metric. (A-C) Increased density of an added high competence source host increases all disease metrics in the absence of interspecific competition (black circles), but decreases all disease metrics when the added host has more negative (or less positive) total competitive effects on low competence source hosts than high competence sink hosts (blue squares). (D-F) Increased density of an added low competence sink host decreases all disease metrics in the absence of interspecific competition (black circles), but increases all disease metrics when the added host has more negative (or less positive) total competitive effects on low competence sink hosts than high competence source hosts (blue squares). In each row, curves of the same color are from the same parameterization of a four-host version of model (2); see Appendix S4 for parameter values.

Figure 3: Increased density of an added host causes all metrics to change in the same direction (black circles), unless source and sink do not imply high and low competence (mechanism 1; blue squares) or there are large changes in focal host density (mechanism 3; red triangles). Increased density of the added host (A) always increases $\mathcal{R}_0^C$ when the added host has sufficiently high competence and (D) always decreases $\mathcal{R}_0^C$ when added host has sufficiently low competence. (B,C) Increased density of the added host increases focal host infection prevalence (I_1^*/N_1^*) when the added host is a source (black circles and red triangles in B; blue squares and red triangles in E) and decreases focal host infection prevalence when the added host is a sink (blue squares in B; black circles in E). (C,F) Increased density of the added host often increases focal host infected density (I_1^*) when the added host is a source (black circles in C; blue squares in F) and decreases focal host infected density when the added host is a sink (blue squares in C; black circles in F). However, focal host infected density can decrease when the added host is a source if there are large changes in focal host density (mechanism 3), which occurs when the focal host is more strongly regulated by disease and interspecific competition than by intraspecific competition (red triangles in C,F). In each row, curves of the same color are from the same parameterization of a four-host version of model (2); see Appendix S4 for parameter values.

Figure 4: The shapes of host species richness-disease relationships depend on the characteristics of added host species and levels of interspecific competition. (A) In a system where higher competence hosts are added later, $\mathcal{R}_0^C$ increases with host species richness (black circles), unless interspecific competition is sufficiently strong (blue squares). (B) In a system where larger source hosts are added later, focal host infection prevalence increases with host species richness (black circles), unless interspecific competition is sufficiently strong (blue squares). (D) In a system where lower competence hosts are added later, $\mathcal{R}_0^C$ decreases with host species richness (black circles), unless added low competence hosts have strong negative competitive effects on other sink low competence hosts (blue squares). (E) In a system where larger sink hosts are added later, focal host infection prevalence decreases with host species richness (black circles), unless added sink hosts have strong negative competitive effects on other low competence hosts (blue squares). (C,F) Focal host infected density often responds to increases in host species richness in the same way that focal host infection prevalence responds (blue squares in C, all curves in F). However, differences can arise if there are large changes in focal host density (mechanism 3; black circles in C). Values are from a four-host version of model (2); see Appendix S4 for parameter values.

Figure 5: Host additions cause larger decreases (or smaller increases) in each metric under frequency-dependent direct transmission (FDDT) than density-dependent direct transmission (DDDT), unless the added host species have sufficiently high competence, the added host species are sufficiently strong interspecific competitors, or there are large changes in focal host density. In each panel, a change of parameters is used to convert the environmental transmission model from a FDDT form to a DDDT form; see Appendices S1.4.3 and S1.5.4 for details. Positive slopes mean large decreases or smaller increases in the metric under FDDT than DDDT when a host is added to a community of resident host species; negative slopes mean larger increases or smaller decreases in the metric under FDDT and DDDT. (A,D) Host additions cause larger decreases in $\mathcal{R}_0^C$ under FDDT than DDDT (all curves in A, filled black circles and blue squares in D), unless interspecific competition is sufficiently strong (open black circles in D). (B,E) Host additions cause larger decreases in focal host infection prevalence (I_1^*/N_1^*) under FDDT than DDDT (black circles and filled red triangles in B, filled black circles in E), unless the added host is a sufficiently large source (open red triangles in B) or interspecific competition is sufficiently strong (open black circles and blue squares in E). (C,F) The responses in focal host infected density (I_1^*) are the same as those in focal host infection prevalence (black circles in C, all curves in F), unless there are large changes in focal host density (mechanism 3), which occurs when the focal host is more strongly regulated by disease and interspecific competition than by intraspecific competition (open and filled red triangles in C). In each row, curves of the same color are from the same parameterization of a four-host version of model (2); see Appendix S4 for parameter values.

Box 1: Relationships between environmental and direct transmission models

Environmental transmission (ET) models can reduce down to models with density-dependent direct transmission (DDDT), frequency-dependent direct transmission (FDDT), or direct transmission that is intermediate to DDDT and FDDT (Eisenberg et al., 2013; Cortez and Weitz, 2013; Cortez and Duffy, 2021; Cortez, 2021; Espira et al., 2022). In general, an environmental transmission model reduces to a direct transmission model when the shedding rates and either the uptake rates or degradation rates are very large. Under these conditions infectious propagules only persist in the environment for short periods of time, and as a result a susceptible individual will encounter infectious propagules only if it is in close proximity to an infected individual. Thus, the environmental transmission pathogen effectively behaves like a direct transmission pathogen.

The ET model (2) reduces down to (i) a DDDT model when uptake is negligible compared to degradation ($\sum_k u_{S_k} S_i + u_{I_k} I_i + u_{R_k} R_i + \delta \approx \delta$), (ii) an FDDT model when degradation is negligible compared to uptake ($\delta \approx 0$) and all uptake rates are equal ($u_{S_i} = u_{I_i} = u_{R_i} = u$), and (iii) a model with transmission intermediate to DDDT and FDDT when uptake and degradation are both non-negligible. Here, I provide a brief mathematical explanation; additional details for the 1-habitat and n-habitat models are provided in Appendices S1.2 and S2.2, respectively. When the shedding rates are large and the uptake or degradation rates are large, the infected density equation in the 1-habitat ET model (2) is approximately described by

$$\frac{dI_i}{dt} = \overbrace{\sum_j \frac{\beta_i \chi_j S_i I_j}{\delta + \sum_k (u_{S_k} S_i + u_{I_k} I_i + u_{R_k} R_i)}}^{\text{infection}} - \overbrace{\mu_i I_i - m_i I_i}^{\text{mortality}} - \overbrace{\nu_i I_i}^{\text{recovery}} . \qquad (11)$$

The first term in equation (11) is the sum of the direct transmission rates from each host j to host i; the second and third terms are the mortality rates due to disease and non-disease factors; and the fourth term is the recovery rate. I consider three cases. First, if uptake of infectious propagules is negligible ($\delta + \sum_k u_{S_k} S_i + u_{I_k} I_i + u_{R_k} R_i \approx \delta$), then the transmission rate in equation (11) reduces to $\hat{\beta}_{ij} S_i I_j$ where $\hat{\beta}_{ij} = \beta_i \chi_j / \delta$. This is a DDDT rate because the rate of infection is proportional to the density of infected individuals (I_j). Second, if degradation is negligible ($\delta \approx 0$) and all uptake rates are equal ($u_{S_i} = u_{I_i} = u_{R_i} = u$), then the transmission rate in equation (11) reduces to $\hat{\beta}_{ij} S_i I_j / \sum_k N_k$ where $\hat{\beta}_{ij} = \beta_i \chi_j / u$. This is a FDDT rate because the rate of infection is proportional to the frequency of infected individuals in the community ($I_j / \sum_k N_k$). Third, if uptake and degradation rates are non-negligible, then the direct transmission rate in equation (11) is intermediate to FDDT and DDDT. The rate is more similar to DDDT when degradation is large and uptake small, and the rate is more similar to FDDT under the opposite conditions.

In total, the ET model (2) can be converted from a DDDT form to a FDDT form by varying the magnitudes of the infectious propagule degradation rate and host

uptake rates. Consequently, the results for the ET model 2 apply to DDDT and
FDDT models, after accounting for the restrictions on the parameter values. This
allows one to simultaneously study ET, DDDT, and FDDT pathogens and identify
how the transmission mechanism affects model predictions.

Box 2: Connection between $\mathcal{R}_0^C$ and theory for stage-structured models

The pathogen's basic reproduction number, $\mathcal{R}_0^C$, is the average number of new infections
produced by a single infected individual in an otherwise susceptible community. The
value of $\mathcal{R}_0^C$ determines if a single infected individual will cause an outbreak ($\mathcal{R}_0^C >$
1) or not ($\mathcal{R}_0^C > 1$). Conveniently, $\mathcal{R}_0^C$ for the 1-habitat model (2) has an explicit
formula,

$$
\mathcal{R}_0^C = \sum_i \overbrace{\frac{\chi_i}{(m_i + \mu_i + \nu_i)}}^{\substack{\text{infectious propagules} \\ \text{produced per host}}} \cdot \overbrace{\frac{\beta_i N_i^\ddagger}{U^\ddagger + \delta}}^{\substack{\text{infected hosts} \\ \text{produced per propagule}}} \tag{12}
$$

where $N_i^\ddagger$ is the density of host i and $U^\ddagger = \sum_j u_{S_j} N_j^\ddagger$ is the total uptake rate of
infectious propagules at the disease-free equilibrium; see Appendix S1.4 for mathematical
calculations. Each term in the sum defines the average number of new infections
a single infected individual of host i will produce in its lifetime. In particular, the
first fraction is the shedding rate (χ_j) multiplied by the average duration of infection
($1/[\mu_i + m_i + \gamma_i]$), which equals the average lifetime production of infectious propagules
by an infected individual of host i. The second fraction is the per infectious propagule
infection rate for host i ($\beta_i N_i^\ddagger$) multiplied by the average lifespan of a spore ($1/[U^\ddagger +$
$\delta]$), which equals the average lifetime production of new infected individuals of host
i by an infectious propagule. The sensitivities of $\mathcal{R}_0^C$ (e.g., $\partial \mathcal{R}_0^C / \partial \chi_n$ and $d\mathcal{R}_0^C / dN_n^*$)
can be computed by differentiating equation (12).

For many models, explicit formulas for $\mathcal{R}_0^C$ cannot be computed. This is true for
the n-habitat models analyzed in Appendix S2 and direct transmission models with
three or more host species. Consequently, one cannot compute the sensitivities of
$\mathcal{R}_0^C$ by differentiating a formula analogous to equation (12). However, the matrix
theory developed for discrete-time models of stage-structured populations allows one
to derive formulas for the sensitivities of $\mathcal{R}_0^C$, even when an explicit formula for $\mathcal{R}_0^C$
does not exist.

The intuition for the connection between $\mathcal{R}_0^C$ and stage-structured models is that
infected individuals of each host species represent different life history "stages" for
the pathogen. Matrix models for stage-structured populations describe how each
stage contributes to the abundances of all stages in the next time step. Analogously,
the Next Generation Matrix (Van den Driessche and Watmough, 2008; Diekmann
et al., 2010) describes how the infected class in each host species contributes to infections

in all host species in the next pathogen generation. As a specific example, the Next Generation Matrix for a 3-species version of model (2) is

$$
\mathbf{NGM} = \begin{bmatrix}
\dfrac{\chi_1}{\mu_1 + m_1 + \gamma_1} \dfrac{\beta_1 N_1^{\ddagger}}{U^{\ddagger} + \delta} & \dfrac{\chi_2}{\mu_2 + m_2 + \gamma_2} \dfrac{\beta_1 N_1^{\ddagger}}{U^{\ddagger} + \delta} & \dfrac{\chi_3}{\mu_3 + m_3 + \gamma_3} \dfrac{\beta_1 N_1^{\ddagger}}{U^{\ddagger} + \delta} \\
\dfrac{\chi_1}{\mu_1 + m_1 + \gamma_1} \dfrac{\beta_2 N_2^{\ddagger}}{U^{\ddagger} + \delta} & \dfrac{\chi_2}{\mu_2 + m_2 + \gamma_2} \dfrac{\beta_2 N_2^{\ddagger}}{U^{\ddagger} + \delta} & \dfrac{\chi_3}{\mu_3 + m_3 + \gamma_3} \dfrac{\beta_2 N_2^{\ddagger}}{U^{\ddagger} + \delta} \\
\dfrac{\chi_1}{\mu_1 + m_1 + \gamma_1} \dfrac{\beta_3 N_3^{\ddagger}}{U^{\ddagger} + \delta} & \dfrac{\chi_2}{\mu_2 + m_2 + \gamma_2} \dfrac{\beta_3 N_3^{\ddagger}}{U^{\ddagger} + \delta} & \dfrac{\chi_3}{\mu_3 + m_3 + \gamma_3} \dfrac{\beta_3 N_3^{\ddagger}}{U^{\ddagger} + \delta}
\end{bmatrix}.
\tag{13}
$$

Entry i, j defines the average number of new infections in host j that a single infected individual of host i will produce in its lifetime. In particular, the first fraction in each entry $(\chi_j/[\mu_j + \nu_j + \gamma_j])$ equals the average lifetime production of infectious propagules by an infected individual of host j. The second fraction in each entry $(\beta_i N_i^{\ddagger}/[U^{\ddagger} + \delta])$ equals the average lifetime production of new infected individuals of host i by an infectious propagule. When the fractions are multiplied, they yield the average lifetime production of new infections in host j by a single infected individual of host i.

Importantly, the sensitivity theory for the asymptotic growth rates of stage-structured populations (Caswell, 2019) can be used to compute the sensitivities of $\mathcal{R}_0^C$. The reason is that for matrix models, the leading eigenvalue of the matrix defines the asymptotic growth rate of the species. Analogously, the leading eigenvalue of the Next Generation Matrix (13) defines the pathogen's reproduction number $(\mathcal{R}_0^C)$, which in turn defines the asymptotic growth rate of an outbreak. In total, this means the theory for matrix models can be used to compute the sensitivities of $\mathcal{R}_0^C$ for the n-habitat model and direct transmission models with three or more host species, even though explicit formulas for $\mathcal{R}_0^C$ cannot be computed. Mathematical details are given in Appendix S2.4.