<u>The Circadian Clock is Disrupted in Pancreatic Cancer</u>

Authors: Patrick B. Schwartz, MD[1], Manabu Nukaya, PhD[1], Mark Berres, PhD[2], Clifford D. Rubinstein,

PhD[2], Gang Wu, PhD[3], John B. Hogenesch, PhD[3], Christopher A. Bradfield, PhD[4], and Sean M.

Ronnekleiv-Kelly, MD[1,5]

Affiliations:

[1]Department of Surgery, Division of Surgical Oncology, University of Wisconsin School of Medicine and

Public Health, Madison, WI

[2]Biotechnology Center, University of Wisconsin, Madison, WI

[3]Division of Human Genetics and Immunobiology, Center for Circadian Medicine, Department of

Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

[4]McArdle Laboratory for Cancer Research, Department of Oncology, University of Wisconsin School of

Medicine and Public Health, Madison, WI

[5]University of Wisconsin Carbone Cancer Center, University of Wisconsin School of Medicine and

Public Health, Madison, WI

Corresponding Author: Sean M Ronnekleiv-Kelly, MD ([ronnekleiv-kelly@surgery.wisc.edu)](mailto:ronnekleiv-kelly@surgery.wisc.edu)

Address: 600 Highland Ave. Bx 7375, K3/705 CSC, Madison, WI, 53792

Phone: (608) 262-2025 **Fax:** (608) 252-0913

Running Title: Pancreatic Cancer Clock

Funding: The research reported in this publication was supported by the Department of Defense Peer

Reviewed Cancer Research Program Idea Award Number CA190176 (Grants.gov ID GRANT12935023)

(SRK), by the National Institutes of Health (NIH) National Institute of Environmental Health Sciences

(NIEHS) Award Number R35ES028377 (CAB), by the NIH NIEHS Award Number T32 ES007015

(PBS) and the University of Wisconsin Carbone Cancer Center (SRK). The content is solely the

responsibility of the authors and does not necessarily represent the official views of the Department of

Defense or the National Institutes of Health.

Disclosure: The authors declare no conflict of interest or competing interests

Abstract

Disruption of the circadian clock is inextricably linked to cancer development and progression.

Establishing this connection has proven beneficial for understanding cancer pathogenesis, determining

prognosis, and uncovering novel therapeutic targets. However, barriers to characterizing the circadian

clock in human pancreas and human pancreatic cancer – one of the deadliest malignancies – have

hindered an appreciation of its role in this cancer. Here, we employed normalized coefficient of variation

(nCV) and clock correlation analysis in human population-level data to determine the functioning of the

circadian clock in pancreas cancer and adjacent normal tissue. We found a substantially attenuated clock

in the pancreatic cancer tissue. Then we exploited our existing mouse pancreatic transcriptome data to

perform an analysis of the human normal and pancreas cancer samples using a machine learning method,

cyclic ordering by periodic structure (CYCLOPS). Through CYCLOPS ordering, we confirmed the nCV

and clock correlation findings of an intact circadian clock in normal pancreas with robust cycling of

several core clock genes. However, in pancreas cancer, there was a loss of rhythmicity of many core clock

genes with an inability to effectively order the cancer samples, providing substantive evidence of a

dysregulated clock. The implications of clock disruption were further assessed with a *Bmal1* knock-out

pancreas cancer model, which revealed that an arrhythmic clock caused accelerated cancer growth and

earlier metastatic spread, accompanied by chemoresistance and enrichment of key cancer-related

pathways. These findings provide strong evidence for clock disruption in human pancreas cancer and

demonstrate a link between circadian disruption and pancreas cancer progression.

Introduction

The circadian clock is a conserved molecular feedback loop that regulates many signaling pathways to control metabolism, immunity, apoptosis, and other critical cellular functions in the body.[1] At its core, the positive arm of the clock (i.e. CLOCK and BMAL1 [also known as ARNTL or MOP3]) drives transcription of the negative arm, including PER1-3 and CRY1-2.[2,3] The negative arm represses the transcriptional activation of CLOCK and BMAL1, and a second interlocked loop involves the nuclear receptors RORα/β/λ and NR1D1-2 (also known as REV-ERBα/β), which activate and suppress BMAL1 expression, respectively. These, along with other core clock components, form a tightly regulated series of transcriptional-translational feedback loops that ensure rhythmic expression over 24 hours and function to maintain cellular and organ homeostasis.[2–4]

Environmental cues influence the synchronization of circadian rhythms in various organ systems, and misalignment of external cues with the internal clock (e.g. shift work) can cause clock dysfunction with consequent metabolic derangements and pathologic states.[4–6] For instance, circadian dysregulation has been strongly linked to obesity and diabetes, both risk factors for cancer.[7–11] Concordantly, landmark studies have shown that disruption of the endogenous clock through mutations in or suppression of the core clock genes is intricately linked to tumor growth in several cancers.[12–14] For example, knockout of *Bmal1* in *Kras*- and *p53*-mutant lung cancer causes marked tumor progression *in vivo*,[13] while MYC-induced repression of *BMAL1* in human neuroblastoma drives decreased overall survival in a BMAL1-dependent manner.[15] Importantly, targeting dysfunctional clock components in certain cancers has proven an effective treatment strategy.[15,16] Thus, identifying an aberrantly functioning circadian clock in cancer can lead to key advancements such as understanding pathogenesis, prognosis, and uncovering novel therapeutic targets.

Although indeterminate, there is some evidence that the clock may be dysregulated in pancreatic ductal adenocarcinoma (PDAC), leading to a worse prognosis;[17,18] this is alarming for a deadly malignancy where only 11% of patients survive beyond 5 years.[19] To advance our understanding of how clock

disruption impacts PDAC pathogenesis, and ultimately foster the identification of therapeutic targets, an

essential first step is to establish that clock dysfunction exists in human PDAC. Unfortunately, to date, the

cumulative data does not definitively support this assertion and is inconclusive. Prior studies have relied

on contrasting expression differences between the core clock genes in tumor compared to normal

pancreas as a basis for clock disruption.[17,18,20,21] But differential expression alone is limited, and does not

provide insight into critical components of clock health such as relative amplitude, rhythmicity, or

correlation of expression amongst core clock genes.[22] Moreover, in the pancreas, phase advancement as a

result of chronic jetlag causes differential expression of core clock genes while maintaining a robust and

healthy clock (i.e. strong relative amplitude, rhythmicity, and clock correlation).[23] Thus, much more

substantive data is required to affirm clock disruption in PDAC.

In pre-clinical studies (e.g. mouse or cell-line models), the determinants of clock health can be identified

by obtaining longitudinal data under controlled conditions.[23,24] However, in human studies – particularly

human pancreas – the requisite periodic data by multiple sampling is not feasible (or ethical). Therefore,

alternative means of determining rhythmically expressed genes in humans are necessary. Following the

emergence of several bioinformatics tools, the principal aspects of clock health including relative

amplitude, core clock gene correlation, and statistical determination of rhythmicity can be resolved when

using the appropriate reference data.[22,25–28] We recently generated a robust pancreas dataset demonstrating

diurnally expressed genes over 48 hours.[23] With this foundation, we were able to apply normalized

coefficient of variation (nCV), clock correlation analysis, and cyclic ordering by periodic structure

(CYCLOPS) to test the hypothesis that the circadian clock is disrupted in human PDAC.[22,25,28] While on

the surface this may be construed as a simplistic hypothesis, several limitations have hindered the ability

to evaluate the circadian clock in human pancreas, including the lack of human periodic data and an

absence of a reference transcriptional dataset.

Here, we employed nCV, clock correlation analysis, and CYCLOPS on publicly available human

population-level expression data from The Cancer Genome Atlas (TCGA) and Clinical Proteomic Tumor

Analysis Consortium 3 (CPTAC-3) datasets to determine the health of the circadian clock in PDAC and

adjacent normal tissue.[29,30] For the first time, we identified an intact circadian clock in human normal

pancreas with robust cycling of several core clock genes. We also found a markedly weakened clock in

the cancer tissue, providing substantive evidence of a dysregulated clock in PDAC. These findings

represent significant advancements in evaluating clock function in PDAC, and the potential clinical

implications of clock disruption were further assessed with a pre-clinical PDAC model. We used

CRISPR/Cas9 technology to selectively target *Bmal1* and examined the effects of clock dysfunction *in*

vitro and *in vivo*. This revealed that loss of clock function caused accelerated cancer growth, earlier

metastatic spread, enrichment of key cancer-related pathways, and resistance to commonly used cytotoxic

chemotherapies for PDAC. These findings provide strong evidence for circadian clock disruption in

human PDAC and demonstrate a link between circadian disruption and pancreas cancer progression.

<u>Materials and Methods</u>

Mouse Care

All animal studies were conducted according to an approved protocol (M005959) by the University of

Wisconsin School of Medicine and Public Health (UW SMPH) Institutional Animal Care and Use

Committee (IACUC). Male and female C57Bl/6J mice were housed in an Assessment and Accreditation

of Laboratory Animal Care (AALAC) accredited selective pathogen-free facility (UW Medical Sciences

Center) on corncob bedding with chow diet (Mouse diet 9F 5020; PMI Nutrition International) and water

ad libitum.

Clock Correlation, nCV, and CYCLOPS Pipeline

We assessed the overall clock gene correlation and robustness of the clock with the clock correlation

matrix and normalized coefficient of variation (nCV) – nCV is known to be correlated with the relative

amplitude (rAMP) of oscillating clock genes, indicating the clock robustness.[22,25] The nCV was calculated

for the overall condition with the nCVnet and nCVgene functions.[25] Clock correlation matrices were

created using an available shiny app (https://github.com/gangwug/CCMapp) which compares the

correlation of clock components (17 individual clock genes) to a baseline correlation from the circadian

atlas using the Mantel test.[22,26,27,31,32]

Cyclic ordering by periodic structure (CYCLOPS)[28] was validated for use in the pancreas utilizing our

existing murine normal circadian and chronic jetlag pancreas RNA sequencing (RNA-seq) data (Gene

Expression Omnibus (GEO) Accession number: GSE165198).[23] Specifically, the seed genes for use in

CYCLOPS were selected by cross-referencing genes found to be rhythmically expressed in our dataset

with those genes either rhythmically expressed in the liver (similarly metabolic organ) or those used by

Wu *et al.* when validating CYCLOPS in the skin (**Supplemental Data File 1**).[23,26,31,32] The updated

CYCLOPS pipeline by Wu *et al.* was then used to reorder our murine pancreas datasets with known

sample collection times.[26] Eigengenes were selected with the Oscope package to sharpen CYCLOPS.[33]

Clusters with a $p < 0.05$ and $\text{Met}^{\text{smooth}} < 1$ were considered to be significantly reordered.[28] Rhythmicity of

the reordered genes was determined on cosinor analysis with a $p < 0.01$, rAMP > 0.1, goodness of fit (rsq)

> 0.1, and fitmean > 16.[26] Significantly rhythmic gene phase was then compared to the rhythmic gene

phase detected from the known sample time collection using the meta3d function of Metacyle.[34] Clock

genes were highlighted to demonstrate a correlation between the predicted and actual phase.

The clock was evaluated in human normal and human PDAC RNA-seq datasets from TCGA and

CPTAC-3.[29,30] After batch correction with ComBat and filtering, 50 matched normal and 318 PDAC

samples were obtained for analysis.[35] The pipeline described above was then used to obtain the clock

correlation matrix, nCV, and CYCLOPS reordering. Cosinor analysis was performed to test for

rhythmicity. Given the additional biologic heterogeneity of the human data, a $p < 0.05$, rAMP > 0.1,

goodness of fit (rsq) > 0.1, and fitmean > 16 were used as a rhythmicity cutoff. Rhythmic genes from

normal samples were assessed with phase set enrichment analysis (PSEA).[36] Rhythmic gene sets ordered

by significance were inputted with their calculated phase of expression. Default settings were used for

PSEA, including domain 0-24, min item 10, max sims/test 10,000. The gene set enrichment analysis

(GSEA) gene ontology (GO) (c5.go.bp.v7.5.1.symbols) set was leveraged as the pathway input. The top

15 significant pathways ($q < 0.05$) were selected for representation.

KPC Cell Line Creation and Maintenance

Pancreas cancer cells that harbor $Kras^{G12D}$ and $Trp53^{R172H}$ mutations (KPC cells) were acquired from

Ximbio (Catalog Number 153474; Westfield Stratford City, UK). Cells were cultured in DMEM

supplemented with 10% Fetal Bovine Serum (Cytiva, Marlborough, MA), 1% L-glutamate-Penicillin-

Streptomycin (Gibco, ThermoFisher Scientific, Waltham, MA), and 1% non-essential amino acids

(Gibco) at 37°C at 5.0% CO_2 to their appropriate confluence for use. CRISPR/Cas9 technology was used

to introduce *Bmal1* mutations into the well-established KPC cells, generating the *Bmal1* functional

knock-out line (KPC-BKO).[37] Synthetic tracrRNA and target-specific crRNAs (crRNA:tracrRNA

(ctRNAs)) were annealed as per manufacturer instructions (Integrated DNA Technologies (IDT),

Coralville, IA). Ribonucleoproteins were formed with Cas9 protein (V3, catalog # 1081059, IDT) and

ctRNAs individually (ctRNA 1:AATATGCAGAACACCAAGGA, ctRNA 2:

TTAGAATATGCAGAACACCA) (IDT). Nucleofection was used to introduce RNPs (1.95 µM Cas9, 2

µM ctRNA, 2 µM electroporation enhancer (IDT)) into 2 x 10^5 KPC cells on a Nucleofector 4D (Lonza

Biosciences, Walkersville, MD) with an SF kit (Catalog Number V4CX-2032, Lonza Biosciences) as per

manufacturer instructions using pulse code CM-120. After 48 hours of recovery in DMEM growth media

(described above) at 37°C at 5.0% CO_2, single cells were deposited in 96 well plates using a BD

FACSAria III (BD Biosciences, Franklin Lakes, NJ). Outgrowing clones were condensed to a 96 well

plate in duplicate to propagate clones and generate a genomic DNA source. Genomic DNA was harvested

from KPC cells, nucleofected with ctRNA-1 RNPs and ctRNA-2 RNPs, and the targeted region of *Bmal1*

was PCR amplified using primers: Forward: acactctttccctacacgacgctcttccgatct NNNNNN

CCAAGAATCCTTGTGTGTCTG and Reverse: gtgactggagttcagacgtgtgctcttccgatct

AGAGGACTCCACAGACATGAAC (IDT). PCR products were dual-indexed with indexing PCR,

pooled, sequenced on an Illumina MiSeq instrument (San Diego, CA), and analyzed with CRISPResso2.[38]

Creation of Per2-dLuciferase reporter KPC cell line

KPC cells were stably transfected with a mammalian gene expression vector harboring a destabilized

luciferase reporter driven by the *Per2* promoter fused to intron 2 of *Per2* and a puromycin resistance

cassette (VectorBuilder, Santa Clara, CA).[39] The vector was transfected with lipofectamine 2000

(ThermoFisher Scientific) and incubated for 2 days, and then exposed to media containing 2.5 μg/mL

puromycin for 3 days, and surviving clones were selected. Luciferase activity was measured in the

selected clones with the luciferase assay system (Promega, Madison, WI) on a BMG CLARIOstar (BMG

Labtech, Ortenberg, Germany) plate reader. The two selected clones were then subcloned using a BD

FACSAria III after staining with DAPI. Luciferase activity was again measured in the subclones to

validate expression before use in downstream experiments.

Clock Function Testing

To evaluate for clock function, KPC and KPC-BKO cells were synchronized with 200 nM

dexamethasone for 2 hours in FBS-free DMEM media, followed by RNA isolation 24 hours after the

synchronization using the RNeasy protocol (Qiagen, Hilden, Germany) according to the manufacturer's

recommendations.[40] Samples were collected at 4-hour intervals for 24 hours. Quantitative real-time

polymerase chain reaction (RT-qPCR) was performed for the downstream core clock gene (CCG) *Per1*

(ID: Mm00501813_m1, Life Technologies, Carlsbad, CA) and the housekeeper gene *Hprt* (ID:

Mm03024075_m1) using GoTaq® Probe qPCR and RT-qPCR System (Promega, Madison, WI) and

Quantstudio 7 flex RT-PCR system (ThermoFisher Scientific). Expression was measured and the ΔCT

was calculated. The mean ΔCT values were then tested for rhythmicity using the meta2d function of

Metacycle.[34] For Metacycle settings, the min period and max period were set to 24, and "JTK", "LS", and

"ARS" were selected for cycMethod. An integrated FDR corrected *q* value < 0.05 and rAMP > 0.1 were

taken as rhythmic. To separately evaluate clock function in KPC cells, luciferase activity driven by the

Per2 promoter was measured. Cells from two independent clones were synchronized with 200 nM

dexamethasone for 2 hours in FBS-free DMEM media, and 24 hours later luciferase activity was

measured at 4-hour intervals. To measure activity, 1x cell culture lysis reagent (Promega) was added and

cells were incubated for 5 minutes. Firefly luciferase assay reagent (Promega) was added and

luminescence was measured on a BMG CLARIOstar plate reader.

Western Blotting

Western blotting was performed to determine BMAL1 expression in the KPC cell lines. After

synchronization, protein samples were isolated at 6-hour intervals for 24 hours using CelLytic M lysis

reagent (MilliporeSigma, Burlington, MA) and Halt Protease and Phosphatase Inhibitor Cocktail

(ThermoFisher Scientific). A total of 30 μg of each sample was loaded onto a Mini-PROTEAN TGX

7.5% precast mini-gel (Bio-RAD Laboratories, Hercules, CA). The gel was then transferred using the

semi-dry transfer technique to an Immobilon-P PVDF membrane (MilliporeSigma). The membrane was

then blocked with 5% skim milk and incubated with 1 μg/μL rabbit BMAL1 antibody (NB100-2288,

Novus Biologicals, Littleton, CO) and 1:2000 rabbit β-ACTIN antibody (4967, Cell Signaling

Technologies, Danvers, MA). Finally, the membranes were incubated with alkaline phosphatase-

conjugated goat anti-rabbit IgG (1:10000) (Jackson ImmunoResearch West Grove, PA) and stained using

1-Step NBT/BCIP solution (ThermoFisher).

RNA Isolation, Sequencing, Differential Gene Expression Analysis

To evaluate for transcriptomic differences between wild-type and BKO KPC cells, bulk RNA-seq was

performed on 6 independent samples collected from each condition. Isolation was carried out as above

and quality was tested for an RNA integrity number (RIN) > 7.5 on the Agilent 2100 bioanalyzer (Agilent

Technologies, Santa Clara, CA). A total of 300 ng of mRNA was enriched with poly-A selection and

sequencing on the Illumina HiSeq2500 platform by the University of Wisconsin Biotechnology

Sequencing Core. FASTq files were processed with Skewer and genes were filtered to remove those with

low expression.[41] Samples were normalized by the method of trimmed mean of M-values.[42] Contrasts

were drawn with the edgeR package, with differential expression taken when the FDR $q < 0.05$.[43]

Pathway testing was performed with the KEGG database (Kyoto Encyclopedia of Genes and Genomes)

using previously described methods.[44] The top 500 significant genes were inputted, ordered by q value,

and the top 9 significant pathways (where $p < 0.05$) were plotted for visualization. Pathway dot size is

indicative of the number of genes in each pathway. The RNA-seq data is publicly available through GEO

(Accession number: GSE213680).

Heterotopic Tumor Modeling

To create flank tumors, 1×10^5 KPC or KPC-BKO cells were injected into the right flanks of

immunocompetent C57BL/6J mice obtained from Jackson Laboratory (Bar Harbor, ME). Cells were

mixed in a 1:1 50 μL solution of DMEM media and Matrigel (Corning Inc, Corning, NY). In the first

experiment, a single dose of KPC or KPC-BKO cells was injected into C57BL/6J mice (male: n = 10,

female: n = 10, each group) and tumors were measured twice weekly for four weeks starting on day 6

with the caliper method as previously described.[45] Tumor length and width were measured and tumor

volume was calculated using the formula: tumor volume = length x width^2 x ½. Tumor weight was also

measured (in mg) at the conclusion of the study period. Two independent replicates were performed for

each condition. In the second experiment, the same dose of KPC or KPC-BKO cells was injected into

C57BL/6J mice (n = 7 in each group) and tumors were measured weekly until the mice became moribund

or died. Kaplan Meier log-rank analysis was then performed to compare survival differences between

conditions with the survival package.[46]

Analysis

All analyses were performed in R version 4.2.0 or Julia version 0.3.12 unless otherwise indicated.

<u>Results</u>

The Clock in Human Pancreatic Ductal Adenocarcinoma is Less Robust Than in Normal Pancreatic

Tissue

To test the hypothesis that the circadian clock is disrupted in human pancreatic ductal adenocarcinoma

(PDAC), we used clock correlation, nCV, and CYCLOPS analysis on publicly available TCGA and

CPTAC-3 pancreatic datasets (**Figure 1, Figure 2,** and **Figure 3**).[29,30] However, we first needed to ensure

the validity of the pipeline in the pancreas since these had not been previously applied to pancreatic

tissue. We assessed the clock in our existing murine normal circadian and chronic jetlag pancreas RNA-

seq datasets by examining the clock correlation matrix and nCV.[23] These datasets contained pancreas

samples acquired every 4 hours for 48 hours in male (n = 3 at each timepoint for each condition) and

female (n = 3 at each timepoint for each condition) mice under standard lighting (normal circadian, n =

72) and chronic jetlag (n = 72) conditions. Chronic jetlag is known to affect the phase of gene expression

but not the relative amplitude (rAMP) or the correlation of the core clock genes in the pancreas.[23]

Correspondingly, we found that the core clock relationships remained intact on the correlation matrix

when compared to the baseline clock correlation matrix for both normal and chronic jetlag conditions ($p <$

0.001, z-statistic = 54.96 vs. $p < 0.001$, z-statistic = 54.08) (**Supplemental Figure 1A**).[31,32] Further, the

nCV of 11 clock genes remained unchanged ($p = 0.76$) between the normal circadian (mean nCV ($\pm$

standard error) = 1.78 ($\pm$ 0.27)) and chronic jetlag (mean nCV = 1.9 ($\pm$ 0.28)) conditions (**Supplemental**

Figure 1B) – consistent with the murine pancreatic clock being strongly rhythmic and intact, matching

precisely what we had found in our prior longitudinal analysis.[23]

CYCLOPS was then used to reorder the mouse pancreas samples by their predicted phase, and cluster

reordering of samples by CYCLOPS was analyzed for appropriate phase progression compared to the

known zeitgeber time (ZT) of sample collection. Notably, CYCLOPS accurately (phase appropriately)

reordered both the normal ($p < 0.01$; $\text{Met}^{smooth} = 0.90$) and the chronic jetlag conditions ($p < 0.01$;

$\text{Met}^{smooth} = 0.97$), validating the pipeline for use in the pancreas (**Supplemental Figure 2B-C**).

Collectively, these results confirmed the ability of nCV, correlation matrix, and CYCLOPS to determine

the robustness of the circadian clock in the pancreas.

Considerable data support the tumor suppressor role of the circadian clock and the assertion that circadian

disruption is present in human cancer.[47] In human PDAC, the supposition is that the clock may be altered,

but supporting data is limited to the relative expression of single genes and is consequently

inconclusive.[18,21,48] We therefore applied our pipeline to determine clock health in PDAC using the TCGA

and CPTAC-3 pancreatic datasets.[22,26–30] After processing and batch correction of the 50 available normal

and 318 available PDAC samples (**Supplemental Figure 3**), we examined the correlation matrix between

the core clock genes (CCGs) and found that there was a weaker relationship amongst the CCGs in PDAC

compared to normal (z-statistic PDAC = 13.77 vs. z-statistic normal = 22.42) (**Figure 1A**). Concordantly,

we found that there was a clear decrease ($p = 0.04$) in the nCV between PDAC (mean nCV = 0.69 ($\pm$

0.08)) and normal (mean nCV = 1.03 ($\pm$ 0.13)) (**Figure 1B**), indicating a weaker clock in PDAC

compared to normal. We then applied our pancreatic CYCLOPS analysis (**Supplemental Figure 4**)

which significantly reordered the normal samples ($p < 0.01$; Met^{smooth} = 0.99), indicating an intact clock.

This was further reinforced when evaluating the predicted phase of CCGs and clock-controlled genes

using CYCLOPS. We found that the phase sequence of normal pancreatic samples was similar to the

phase sequence of mouse pancreatic samples (**Figure 2A-B**), with concordant phase in 12 out of 17 genes

and ROR-phased genes largely peaking before E-box phase genes. Based on the CYCLOPS ordering, we

analyzed the proportion of genes that were rhythmically expressed in normal pancreas. We found that

4,034/18,196 (22.17%) of normal pancreatic genes were rhythmic, (**Supplemental Data Files 2**). We

found that several key CCGs and clock-controlled genes were rhythmically expressed in normal samples

on cosinor analysis, including *CLOCK*, *PER1*, *PER3*, *NR1D2*, *RORC*, *NFIL3, TEF*, *NR1D1, BHLHE40,*

and *NPAS2* (**Figure 3A and Supplemental Table 1**). The CCGs *BMAL1* ($p = 0.07$, rAMP = 0.8) and

CRY1 ($p = 0.08$, rAMP = 0.48) were nearly rhythmic in the normal samples based on the pre-defined

cutoff. Finally, phase set enrichment analysis (PSEA) was used to determine phase-dependent gene

enrichment. The top 15 significant pathways were selected for analysis (**Figure 3C**). Consistent with our

prior murine pancreas data, we found that normal pancreas was associated with time-dependent metabolic

gene pathway enrichment.[23,49] In aggregate, this data confirms the health of the circadian clock in normal

pancreatic tissue, which was expected but not previously shown.

Despite the validation steps and success in ordering normal pancreas, CYCLOPS could not reorder PDAC

samples ($p = 0.43$; Met$^{smooth} = 0.99$), revealing a less robust (or disrupted) clock in PDAC compared to

normal. This inability to reorder strongly supports the attenuated circadian clock in PDAC identified with

nCV and clock correlation analysis. We selected the best reordering available based on the eigengenes

(**Supplemental Figure 4**) to enable visual representation of the PDAC samples compared to normal

pancreas samples. This also enabled us to estimate as best as possible the rAMP of the PDAC samples

(**Supplemental Data Files 3**), to coincide with the visual representation. Consistent with the nCV

analysis, we found that rAMP decreased substantially in every core clock gene from normal to tumor

(**Supplemental Table 1**), as evidenced by *PER1* and *PER3* expression (**Figure 3A**). Collectively, the

weaker clock correlation, markedly reduced nCV, and CYCLOPS analysis of human population-level

data convincingly show for the first time that the circadian clock is significantly disrupted in PDAC.

Creation of a Circadian Dysfunction Pancreatic Cancer Model

After demonstrating that the clock in human PDAC was dysfunctional, we sought to understand the

potential clinical implications of clock disruption. Identifying an aberrant clock in other cancers has

consistently demonstrated accelerated cancer progression and worse prognosis, while simultaneously

revealing novel therapeutic targets.[15] Moreover, there appears to be a putative correlation between

suppressed *BMAL1* expression and poor prognosis in patients with PDAC.[17,18] Therefore, we

hypothesized that disruption of the circadian clock in PDAC would cause accelerated cancer progression.

To test this hypothesis, we used a syngeneic *Kras-* and *Trp53*-mutant pancreas cancer cell line (KPC) and

employed CRISPR/Cas9 technology to functionally knock out *Bmal1* (KPC-BKO) so that we could

examine the effects of clock dysfunction *in vitro* and *in vivo*.[37] The reason for mutating *Bmal1* in the

PDAC cells was several-fold: i) KPC cells demonstrated an intact clock (see below) necessitating core

clock gene modulation, ii) *BMAL1* is a central transcriptional regulator of the circadian clock machinery

and suppressed gene expression has been found in several human cancers, correlating with worse

prognosis,[13,15,17,18,20,21,50] iii) *BMAL1* expression is decreased in tumor compared to normal tissue in human

PDAC,[18] iv) we similarly found a substantially dampened nCV (rAMP) of *BMAL1* expression in PDAC

compared to normal tissue (*PER1*, *PER3*, *NR1D1,* and *BMAL1* displayed the greatest decrease in nCV),

and v) segregating patients with PDAC into high and low *BMAL1* expression appears to be prognostic for

survival outcomes.[17,18,21]

Consequently, guide RNAs were designed to target exon 8 of the *Bmal1* gene, which resulted in the

insertion of adenine on one allele and a 2 base pair deletion on the other between nucleotide 81,074 and

81,075 at amino acid 73 (GRCm38) (**Figure 4A**). The result of both mutations was a frameshift just

upstream of the basic helix loop helix (bHLH) domain known to assist the PAS A domain in

heterodimerization with its binding partner CLOCK (**Figure 4B**).[51,52] Confirmatory western blotting

revealed the presence of protein in wild-type (WT) cells and an absence of protein in *Bmal1* knockout

(BKO) KPC cells (**Figure 4C**). To evaluate clock functionality, mRNA was isolated at 4-hour intervals (n

= 3 per condition) for 24 hours after cell synchronization, and we performed RT-qPCR for *Per1*, a core

downstream repressor of the positive arm of the molecular clock and a gene that demonstrated

rhythmicity in our human normal CYCLOPS analysis (**Figure 4D**). Rhythmicity was assessed with the

meta2d function of the Metacycle package.[34] KPC (WT) cells exhibited an intact and robust circadian

clock with a $q = 0.0004$ and rAMP = 0.11 for *Per1*. To further evaluate the functionality of the clock in

KPC cells, luciferase activity driven by the *Per2* promoter was measured. Consistent with our evaluation

of mRNA expression, luciferase activity in two separate clones was highly rhythmic (Clone 1: $q = 5.51E-$

5, rAMP = 0.44; Clone 2: $q = 1.95E-7$, rAMP = 0.39) (**Supplemental Figure 5**). Conversely, KPC-BKO

cells exhibited no detectable clock function with a $q = 0.81$ and rAMP = 0.003 for *Per1* expression

(**Figure 4D**). Consistent with the known cross-talk between the circadian clock and cell cycle, KPC-BKO

cells demonstrated alterations in the cell cycle compared to KPC cells (**Figure 4E**).[20] BKO cells were

found to have a higher mean (± standard error) percentage of cells in S phase (48.66 (± 0.13) vs. 36.96 (±

2.04); $p = 0.029$) and a lower percentage in G2 (18.03 (± 0.27) vs. 27.49 (± 0.89); $p = 0.0053$). There

were no differences between KPC and KPC-BKO cells in G1 (35.56 (± 1.19) vs. 33.31 (± 0.17); $p = 0.2$).

Taken together, these data demonstrate the successful creation of a novel PDAC cell line with an

abolished clock (*Bmal1* gene mutation), which can be used to elucidate the repercussions of clock

disruption in PDAC.

Clock Dysfunction Accelerates Pancreatic Cancer Growth

After establishing the model, we sought to understand how a dysfunctional clock impacted cell growth.

KPC and KPC-BKO cells were grown in culture and injected into the right flanks of syngeneic C57BL/6J

mice. Tumor growth was measured every 3-4 days for a total of 28 days to understand differences in

primary tumor growth. We found that BKO caused an accelerated growth pattern, resulting in higher

mean ($\pm$ standard error) weight tumors at the study conclusion compared to KPC-derived tumors (438.02

± 48.84 mg vs. 280.11 ± 42.73 mg; p = 0.02) (**Figure 5A-B**). These findings were evaluated and

independently confirmed in a second identically created *Bmal1* mutant (functional knock-out) clone

(**Supplemental Figure 6**). Perhaps more relevant to human PDAC – given the aggressiveness of this

cancer – we assessed the effect of clock disruption on survival (i.e. metastatic spread) by implanting KPC

and KPC-BKO heterotopic tumors and observing the mice until moribund status or lethality (**Figure 5C**).

Notably, the terminal endpoint occurred from metastatic spread to the diaphragm, lungs, or abdominal

cavity, or uncommonly from locally advanced tumors. On Kaplan Meier log-rank analysis, mice

harboring tumors derived from KPC-BKO cells experienced worse survival than mice bearing KPC-

derived tumors (median survival: 52 versus 75 days, $p = 0.04$). Thus, clock dysfunction resulted in

accelerated tumor growth and earlier onset of tumor metastases (i.e., PDAC progression) *in vivo*, leading

to worse overall survival.

Loss of Bmal1 Promotes the Enrichment of Cell Growth Pathways

To determine the possible etiologies of clock disruption causing accelerated PDAC progression, we

compared the transcriptomic profiles of KPC and KPC-BKO cells. The principal component analysis

revealed clear transcriptional profile differences between each condition (**Figure 6A**). Differential gene

expression was quantified with edgeR and of the 15,110 genes, 5,235 (34.65%) were upregulated and

5,113 (33.84%) were downregulated in KPC-BKO compared to KPC cells (**Figure 6B, Supplemental**

Data File 4).[43] When we examined the CCGs known to control the circadian cycle, we found that 12/15

examined clock genes were differentially expressed, including *Per1*, *Per2*, *Per3*, *Cry1*, *Cry2*, *Nr1d1*,

Nr1d2, *Bhlhe40*, *Bhlhe41*, *Npas2*, *Arntl2*, and *Dbp* (All q < 0.05).[51] We then performed a KEGG pathway

analysis to examine for enrichment of pathways as a result of *Bmal1* knock-out in the PDAC cells

(**Figure 5C**). We found that there was an enrichment of pathways important for cellular adhesion, such as

ECM-Receptor Interaction, Cell Adhesion, and Focal Adhesion, as well as several cellular growth

pathways including PI3K-AKT Signaling Pathway, MAPK Signaling Pathway, and Rap1 Signaling

Pathway. Collectively, these data indicate that clock disruption in the KPC cells results in significant core

clock gene changes and a transcriptional shift that alters key cancer promotion-related pathways such as

cellular attachment and proliferation.

Cell Survival is Increased with the Loss of Bmal1 through Alterations in Multiple Cell Death Pathways

A hallmark of cancer progression is the inhibition of apoptosis.[53] This phenotype is readily apparent in

response to chemotherapy. In particular, the mechanism of action of gemcitabine (inhibition of DNA

synthesis) and paclitaxel (microtubule stabilization) – backbones in PDAC therapy – is to ultimately

induce apoptosis.[54,55] To understand the potential implications of clock disruption for PDAC patients

undergoing treatment, we subjected KPC and KPC-BKO cells to 72 hours of chemotherapeutic treatment

with either gemcitabine or paclitaxel.[56] We found that the activity of cell death pathways through

apoptosis (measured by Caspase 3/7 activity) was blunted in response to both chemotherapeutic agents as

a result of clock dysfunction (**Figure 7A**). We also assessed cytotoxic cell death, as measured by Dead-

Cell Protease activity, and found that clock disruption promoted resistance to cytotoxic cell death induced

by gemcitabine and paclitaxel (**Figure 7B**). Prior work has implicated alteration of Trp53 signaling to

impact cell survival by suppressed apoptosis,[20] but KPC cells (both WT and BKO) are *Trp53* mutant

(*Trp53*^{R172H}) indicating alternative mechanisms of heightened resistance in the KPC-BKO cells.

Gemcitabine resistance in PDAC often occurs due to the downregulation of the channel protein hENT1

(*SLC29A1* gene), or deoxycytidine kinase (*DCK* gene) which activates gemcitabine once inside the cell.[57]

However, these were only marginally downregulated (1.05-fold, $q = 0.04$ and 1.08-fold, $q = 0.02$) due to

Bmal1 knock-out, and thus unlikely to contribute to the differences seen with clock disruption.

Furthermore, these were non-rhythmic when examining our human samples, suggesting a lack of

circadian control. Meanwhile, paclitaxel drug resistance is thought to occur mostly through upregulation

of drug efflux transporter proteins (P-glycoprotein also known as multidrug-resistance associated protein),

but these ATP-binding cassette transporter proteins (*Abcb1a and Abcb1b* genes) were instead

significantly *downregulated* (8.2-fold, $q < 0.0001$ and 2.1-fold, $q < 0.0001$) in KPC-BKO vs KPC cells.[55]

Concordantly, the *ABCB1* gene was rhythmic with elevated expression in the human normal samples ($p =$

0.015, rAMP = 0.95, mean expression = 73) compared to the human PDAC samples (mean expression =

23). Thus, the commonly described resistance mechanisms for gemcitabine and paclitaxel were not

identified, underscoring the complexity of clock disruption induced chemotherapeutic resistance in

cancer. Regardless, these data demonstrate that clock dysfunction promotes broad resistance in PDAC

including multiple cell death pathways in the setting of two different PDAC backbone agents.

<u>Discussion:</u>

Our work herein used matched normal and tumor samples from TCGA and CPTAC-3 to provide the first

substantive evidence that the circadian clock is disrupted in PDAC while the adjacent normal pancreatic

clock is intact. To evaluate clock functionality in the pancreas, we assessed the clock with three

complementary analyses not previously performed in pancreatic tissues: nCV, clock correlation, and

CYCLOPS. The nCV was pioneered and validated in several tissues by Wu and colleagues.[25] It is a

measurement that is directly related to the relative amplitude (rAMP) of the oscillation, and assesses

overall and individual clock gene robustness.[25] For example, Wu and colleagues demonstrated that the

nCV of core clock genes (CCGs) was consistently and significantly diminished in clock-disrupted tissues

versus wild-type tissues (e.g. *Bmal1* knockout adipocytes vs wild-type adipocytes) and human datasets of

tumor compared to matched (adjacent) normal tissue, where the timing of sample collection was

unspecified.[25] It is therefore extraordinarily beneficial for understanding the rAMP of the circadian clock

genes – a key measure of clock health – in population-level data where sample time acquisition is

unknown. In conjunction with nCV, Shilts *et al.* developed the method of clock correlation to determine

the progression of the clock in time-indeterminate datasets.[22] By capitalizing on the concept of co-

expression of CCGs (intrinsic to the transcriptional translational feedback loop), Shilts and colleagues

were able to computationally discern clock progression in transcriptomic data from numerous human

datasets. Importantly, this technique is not dependent on full coverage of the period by samples, providing

a beneficial approach for our normal data set which included only 50 samples and did not have phase

representation across the 24-hour period. Furthermore, a direct comparison between heatmaps can be

made (i.e. murine vs human or tumor vs normal) because each heatmap has the same corresponding color

to rho correlation value.[22] Thus, when combining nCV and clock correlation analysis in unordered human

samples, the health of the circadian clock in population-level data can be ascertained even when working

with low sample numbers (at least 30).

Wu and colleagues applied nCV analysis to several paired tumor-normal datasets,[25] while Shilts *et al.*

examined clock correlation in normal tissues and paired tumor-normal datasets.[22] However, we are the

first group to analyze the pancreas, possibly because our recent publication was the first to characterize

the diurnal expression of mouse pancreatic genes over 48 hours.[23] Thus, an appropriate mouse reference

group had not been published for comparison. In the study by Shilts *et al.*, the human liver clock

correlation heatmap (a similarly metabolic organ) demonstrated a weaker clock correlation versus the

mouse reference, which they attributed to increased noise in the liver dataset.[22] Therefore, in our human

data, we expected to identify an apparent 'diminished health' of the clock (lower nCV and weaker clock

correlation) in the human pancreatic tissue versus mouse samples due to significant variation in the

normal samples. This variability is contributed by human factors, such as type and timing of diet,[58,59]

underlying genetic differences, and alterations in the 'normal pancreas' that surrounds the tumor

(pancreatic atrophy, fibrosis, inflammation, etc.).[60] These can modulate the relative amplitude of

oscillation or CCG correlation which contributes to significant noise in the data, as compared to the

genetically identical, age-matched, and environmentally matched mouse samples.[22,28] Regardless of these

limitations, the nCV and clock correlation revealed two key components of a healthy clock in human

pancreas, which was an important component of the pipeline.

CYCLOPS is a machine learning method developed by Anafi *et al.*,[28] with subsequent elegant studies by

Wu *et al.*,[26,27] to demonstrate how clock gene relationships can be used to infer and reorder samples by

their predicted phase of expression to understand circadian biology in population-level data. The authors

who developed CYCLOPS recommend roughly 250 samples for a complete phase distribution,[28] which is

dependent on differences in time of surgery (presume specimens acquired between 6 AM and 6 PM) as

well as differences in genetics and environmental factors (e.g. differences in sleep-wake cycle, or shift

worker status).[61] CYCLOPS has been applied to the lung, liver, brain, hepatocellular carcinoma, and skin,

but not the pancreas.[27,28] This is possibly due to known challenges associated with CYCLOPS, which

includes optimizing the seed gene list for appropriate ordering.[28] We used the modified CYCLOPS

approach by Wu *et al.* and were able to leverage our murine pancreatic longitudinal expression data to

generate the seed gene list for use with human data, which increases the signal-to-noise ratio to optimize

ordering capability.[23,27,28] Thus, despite the expected variability in human normal pancreas samples,

CYCLOPS significantly ordered the samples across the period, with several clock genes, including

CLOCK, *PER1*, *PER3*, *NR1D2*, *RORC*, *NFIL3,* T*EF*, *NR1D1, BHLHE40,* and *NPAS2* demonstrating

rhythmic expression. While *BMAL1* and *CRY1* did not pass our predetermined cutoff for rhythmicity in

the normal samples, they demonstrated a robust rAMP with near-significant p values likely reflecting

limitations with our sample number (i.e., distribution of samples across the period) than true lack of

rhythmicity in the normal samples. As evidenced by the apparent gap in phase in the normal data, there

was not uniform distribution of normal samples across the 24-hour period, which signifies a limitation of

our analysis. However, despite these limitations, CYCLOPS statistically ordered the human normal data,

and the phase-set enrichment analysis and predicted phase of expression (of most CCGs) aligned well

with our prior mouse pancreas transcriptomic data (reference dataset). When combined with nCV and

clock correlation, the data clearly demonstrate an intact clock in the human normal pancreas samples,

which was an essential premise for evaluating the clock in human PDAC.

We then proceeded to use nCV, clock correlation, and CYCLOPS to provide compelling evidence of

circadian clock disruption in PDAC. While the concept of circadian disruption in PDAC has been posited

by others, past studies examining global clock function in human PDAC have been limited by detecting

binary differences in overall RNA and protein expression between normal pancreas and PDAC.[17,18,20,21]

Relles *et al.* found that several CCGs demonstrated decreased expression in PDAC compared to benign

tumors or normal pancreas,[18] while Li *et al*. found that low *Bmal1* expression (compared to 'higher

expression') was associated with worse disease-free and overall survival in patients with PDAC.[17] Similar

studies have been repeated with concordant findings.[20,21] However, binary comparisons of expression are

unlikely to capture the complexities of clock health such as rhythmicity, phase changes, changes in

rAMP, or changes in clock progression. With the 318 available PDAC samples, we showed that there was

markedly diminished nCV and a much weaker correlation among clock genes in PDAC compared to

normal pancreas, depicting a loss of clock health in the cancer tissue. Further, although the inability to

reorder PDAC samples may be a limitation of CYCLOPS, it more likely indicates clock dysfunction.[62] In

the PDAC samples, there was a sufficient sample number (n = 318) for phase distribution, but the rAMP

of nearly all CCGs was markedly decreased (as visualized through best reordering), with many

manifesting an arrhythmic pattern. Collectively, our approach to human data (nCV, clock correlation, and

CYCLOPS) provided convincing evidence of clock disruption in PDAC.

The main limitation of our current strategy is the inability to discern why there was a loss of circadian

signatures in human PDAC population data. These observations could be the consequence of i)

dampening of the clock to a significant extent in every patient (i.e., global phenomena), ii) differences in

the extent to which the clock is disrupted (i.e., substantial clock dysfunction in a certain subset of

patients), or iii) biological heterogeneity in the PDAC samples surpassing circadian variability. The last

consideration seems unlikely given the marked decrease in rAMP (nCV) and clock correlation showing

significantly diminished clock health in PDAC. Furthermore, as seen with the CYCLOPS ordered *PER1*

expression data over the 24-hr period (Figure 3A), enhanced variability did not lead to the apparent

arrhythmicity in PDAC. Unfortunately, we were unable to discriminate between the first two possibilities.

The literature in cancer would suggest that cohorts of patients exhibit a differential extent of clock

disruption in their tumors (i.e. some clock intact vs some markedly disrupted),[15,17,18,50] which would result

in an overall loss of circadian signatures seen in the nCV, clock correlation, and CYCLOPS analysis. Our

assessment of clock function in the KPC cells would also support this hypothesis, given that the clock

was intact in the *Kras-* and *Trp53*-mutant pancreas cancer cells and necessitated *Bmal1* mutagenesis to

generate clock disruption. Further, we assessed the cycling of the *Per1* gene and *Per2*-luciferase in the

KPC cells and identified a robust rhythm; by comparison, the human PDAC samples demonstrated

arrhythmic *PER1* expression. Therefore, we generated the novel murine KPC-BKO cell line as a basis to

evaluate clock disruption in PDAC, and intend for this cell line to be a useful approach for investigators

evaluating the clock in PDAC.

Using our KPC and KPC-BKO cells, we found accelerated tumor growth in our syngeneic *in vivo* model

with *Bmal1* functional knock-out. Our preference was to use an immune-competent model given the

known impact of the circadian clock on the immune system (another strength of our model).[63,64]

Interestingly, our primary tumor growth results were similar to results by Jiang and colleagues, who used

implanted human PDAC cells (BxPC-3 and PANC-1) into immunocompromised mice after shRNA

knockdown of *Bmal1*.[20,21] However, patients with PDAC ultimately succumb to distant metastatic spread,

rather than local tumor growth, and so examining the contribution of clock disruption to overall survival

was of greater importance.[65] We found that *Bmal1* mutation promoted earlier metastatic spread of tumors

resulting in earlier lethality, which has not been shown before. Concordant with the aggressive tumor

phenotype, *Bmal1* mutation also caused resistance to chemotherapy. While resistance to gemcitabine has

been shown,[21] we found chemoresistance to two different backbone anti-cancer agents (gemcitabine and

paclitaxel), including suppressed apoptosis and cytotoxicity, indicating a more broad resistance to

standard PDAC chemotherapy. Although suppressed *Bmal1* expression in PDAC has been suggested to

modulate *Trp53* to promote a tumor suppressor effect, this was unlikely the case in our study considering

KPC cells are a *Trp53*-mutant cell line.[20] Other work indicates the transcription factor YY1 is

significantly overexpressed in PDAC and ultimately causes BMAL1 suppression with consequent PDAC

progression and resistance to gemcitabine (unclear mechanism of resistance).[21] Yet, when we examined

our human PDAC and human normal samples, we found that *YY1* expression was equivalent between the

groups (mean expression 101.61 versus 102.87). Turning to other cancers where BMAL1 is suppressed,

MYC amplification in neuroblastoma alters *BMAL1* mRNA expression through induction of *NR1D1*.[50,66]

Notably, this is associated with poor prognosis and is *BMAL1* dependent since ectopic expression of

BMAL1 inhibits tumor growth. This mechanism is unlikely to be occurring in PDAC since there is no

upregulation of *MYC*, *MYC-N,* or *NR1D1* gene expression in human PDAC compared to normal samples.

While common resistance mechanisms of gemcitabine and paclitaxel were not reflected in the human

PDAC versus human normal data or the KPC versus KPC-BKO cells (e.g. channel proteins), RNA

sequencing identified several enriched pathways integral to cancer progression and chemoresistance in the

Bmal1 functional knock-out cells, such as the PI3K-AKT pathway.[67,68] The PI3K-AKT pathway is

inextricably linked to cancer cell proliferation and resistance to apoptosis, indicating a plausible

mechanism for inhibition of programmed cell death to multiple agents seen in our study.[69,70] Further,

resistance to paclitaxel is associated with the activation of the PI3K-AKT pathway,[71] and similar

correlations have been identified with gemcitabine resistance.[72,73] However, the mechanism of

chemoresistance is quite complex and the etiology for suppressed apoptosis and cytotoxicity due to clock

disruption remains unclear. In future work, we can employ a series of clock-manipulated PDAC cell lines

to better understand the contributions of clock disruption to chemoresistance and PDAC progression.

In conclusion, we used a comprehensive approach (nCV, clock correlation, and CYCLOPS) to evaluate

the health of the circadian clock in human normal pancreas and demonstrated clock disruption in human

PDAC. Additionally, we developed novel cell lines to evaluate the repercussions of clock disruption in

PDAC and identified factors associated with poor prognosis (i.e., earlier metastases, resistance to

chemotherapy, and enrichment of cancer-related pathways). Future work will expand on this foundation

to disentangle the clock-dependent effects of PDAC and consequently focus therapeutic efforts.

Acknowledgments:

We would like to acknowledge the Gene Expression Center and the Bioinformatics Resource Center

(BRC) at the University of Wisconsin, Madison for their contributions to this work. We would also like to

thank the Michael W. Oglesby Foundation for their funding support of our work in circadian disruption

and pancreas pathology.

References

1. Bass J, Lazar MA. Circadian time signatures of fitness and disease. *Science*. 2016;354(6315):994-

999. doi:10.1126/science.aah4965

2. Aryal RP, Kwak PB, Tamayo AG, et al. Macromolecular Assemblies of the Mammalian Circadian

Clock. *Mol Cell*. 2017;67(5):770-782.e6. doi:10.1016/j.molcel.2017.07.017

3. Transcriptional architecture of the mammalian circadian clock - PubMed. Accessed October 10,

2022. https://pubmed.ncbi.nlm.nih.gov/27990019/

4. Allada R, Bass J. Circadian Mechanisms in Medicine. *N Engl J Med*. 2021;384(6):550-561.

doi:10.1056/NEJMra1802337

5.   Fishbein AB, Knutson KL, Zee PC. Circadian disruption and human health. *J Clin Invest*.

2021;131(19):e148286. doi:10.1172/JCI148286

6.   West AC, Smith L, Ray DW, Loudon ASI, Brown TM, Bechtold DA. Misalignment with the external

light environment drives metabolic and cardiac dysfunction. *Nat Commun*. 2017;8(1):417.

doi:10.1038/s41467-017-00462-2

7.   Marcheva B, Ramsey KM, Buhr ED, et al. Disruption of the clock components CLOCK and BMAL1

leads to hypoinsulinaemia and diabetes. *Nature*. 2010;466(7306):627-631. doi:10.1038/nature09253

8.   Roenneberg T, Allebrandt KV, Merrow M, Vetter C. Social jetlag and obesity. *Curr Biol CB*.

2012;22(10):939-943. doi:10.1016/j.cub.2012.03.038

9.   Parsons MJ, Moffitt TE, Gregory AM, et al. Social jetlag, obesity and metabolic disorder:

investigation in a cohort study. *Int J Obes 2005*. 2015;39(5):842-848. doi:10.1038/ijo.2014.201

10. Knutson KL, Wu D, Patel SR, et al. Association Between Sleep Timing, Obesity, Diabetes: The

Hispanic Community Health Study/Study of Latinos (HCHS/SOL) Cohort Study. *Sleep*. 2017;40(4).

doi:10.1093/sleep/zsx014

11. Turek FW, Joshu C, Kohsaka A, et al. Obesity and metabolic syndrome in circadian Clock mutant

mice. *Science*. 2005;308(5724):1043-1045. doi:10.1126/science.1108750

12. Dong Z, Zhang G, Qu M, et al. Targeting Glioblastoma Stem Cells through Disruption of the

Circadian Clock. *Cancer Discov*. 2019;9(11):1556-1573. doi:10.1158/2159-8290.CD-19-0215

13. Papagiannakopoulos T, Bauer MR, Davidson SM, et al. Circadian Rhythm Disruption Promotes

Lung Tumorigenesis. *Cell Metab*. 2016;24(2):324-331. doi:10.1016/j.cmet.2016.07.001

14. Kettner NM, Voicu H, Finegold MJ, et al. Circadian Homeostasis of Liver Metabolism Suppresses

Hepatocarcinogenesis. *Cancer Cell*. 2016;30(6):909-924. doi:10.1016/j.ccell.2016.10.007

15. Moreno-Smith M, Milazzo G, Tao L, et al. Restoration of the molecular clock is tumor suppressive in

neuroblastoma. *Nat Commun*. 2021;12(1):4006. doi:10.1038/s41467-021-24196-4

16. Sulli G, Rommel A, Wang X, et al. Pharmacological activation of REV-ERBs is lethal in cancer and

oncogene induced senescence. *Nature*. 2018;553(7688):351-355. doi:10.1038/nature25170

17. Li W, Liu L, Liu D, et al. Decreased circadian component Bmal1 predicts tumor progression and

poor prognosis in human pancreatic ductal adenocarcinoma. *Biochem Biophys Res Commun*.

2016;472(1):156-162. doi:10.1016/j.bbrc.2016.02.087

18. Relles D, Sendecki J, Chipitsyna G, Hyslop T, Yeo CJ, Arafat HA. Circadian gene expression and

clinicopathologic correlates in pancreatic cancer. *J Gastrointest Surg Off J Soc Surg Aliment Tract*.

2013;17(3):443-450. doi:10.1007/s11605-012-2112-2

19. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J Clin*. 2022;72(1):7-

33. doi:10.3322/caac.21708

20. Jiang W, Zhao S, Jiang X, et al. The circadian clock gene Bmal1 acts as a potential anti-oncogene in

pancreatic cancer by activating the p53 tumor suppressor pathway. *Cancer Lett*. 2016;371(2):314-

325. doi:10.1016/j.canlet.2015.12.002

21. Jiang W, Zhao S, Shen J, et al. The MiR-135b-BMAL1-YY1 loop disturbs pancreatic clockwork to

promote tumourigenesis and chemoresistance. *Cell Death Dis*. 2018;9(2):149. doi:10.1038/s41419-

017-0233-y

22. Shilts J, Chen G, Hughey JJ. Evidence for widespread dysregulation of circadian clock progression in

human cancer. *PeerJ*. 2018;6:e4327. doi:10.7717/peerj.4327

23. Schwartz PB, Walcheck MT, Berres ME, et al. Chronic Jetlag-Induced Alterations in Pancreatic

Diurnal Gene Expression. *Physiol Genomics*. Published online May 31, 2021.

doi:10.1152/physiolgenomics.00022.2021

24. Hughes ME, Abruzzi KC, Allada R, et al. Guidelines for Genome-Scale Analysis of Biological

Rhythms. *J Biol Rhythms*. 2017;32(5):380-393. doi:10.1177/0748730417728663

25. Wu G, Francey LJ, Ruben MD, Hogenesch JB. Normalized coefficient of variation (nCV): a method

to evaluate circadian clock robustness in population scale data. *Bioinformatics*. 2021;37(23):4581-

4583. doi:10.1093/bioinformatics/btab731

26. Wu G, Ruben MD, Francey LJ, et al. A population-based gene expression signature of molecular

clock phase from a single epidermal sample. *Genome Med*. 2020;12(1):73. doi:10.1186/s13073-020-

00768-9

27. Wu G, Ruben MD, Schmidt RE, et al. Population-level rhythms in human skin with implications for

circadian medicine. *Proc Natl Acad Sci U S A*. 2018;115(48):12313-12318.

doi:10.1073/pnas.1809442115

28. Anafi RC, Francey LJ, Hogenesch JB, Kim J. CYCLOPS reveals human transcriptional rhythms in

health and disease. *Proc Natl Acad Sci U S A*. 2017;114(20):5312-5317.

doi:10.1073/pnas.1619320114

29. Edwards NJ, Oberti M, Thangudu RR, et al. The CPTAC Data Portal: A Resource for Cancer

Proteomics Research. *J Proteome Res*. 2015;14(6):2707-2713. doi:10.1021/pr501254j

30. Cancer Genome Atlas Research Network, Weinstein JN, Collisson EA, et al. The Cancer Genome

Atlas Pan-Cancer analysis project. *Nat Genet*. 2013;45(10):1113-1120. doi:10.1038/ng.2764

31. Pizarro A, Hayer K, Lahens NF, Hogenesch JB. CircaDB: a database of mammalian circadian gene

expression profiles. *Nucleic Acids Res*. 2013;41(Database issue):D1009-1013.

doi:10.1093/nar/gks1161

32. Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB. A circadian gene expression atlas in

mammals: Implications for biology and medicine. *Proc Natl Acad Sci*. 2014;111(45):16219-16224.

doi:10.1073/pnas.1408886111

33. Leng N, Chu LF, Barry C, et al. Oscope identifies oscillatory genes in unsynchronized single cell

RNA-seq experiments. *Nat Methods*. 2015;12(10):947-950. doi:10.1038/nmeth.3549

34. Wu G, Anafi RC, Hughes ME, Kornacker K, Hogenesch JB. MetaCycle: an integrated R package to

evaluate periodicity in large scale data. *Bioinforma Oxf Engl*. 2016;32(21):3351-3353.

doi:10.1093/bioinformatics/btw405

35. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing batch effects

and other unwanted variation in high-throughput experiments. *Bioinforma Oxf Engl*. 2012;28(6):882-

883. doi:10.1093/bioinformatics/bts034

36. Zhang R, Podtelezhnikov AA, Hogenesch JB, Anafi RC. Discovering Biology in Periodic Data

through Phase Set Enrichment Analysis (PSEA). *J Biol Rhythms*. 2016;31(3):244-257.

doi:10.1177/0748730416631895

37. Hingorani SR, Wang L, Multani AS, et al. Trp53R172H and KrasG12D cooperate to promote

chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. *Cancer

Cell*. 2005;7(5):469-483. doi:10.1016/j.ccr.2005.04.023

38. Clement K, Rees H, Canver MC, et al. CRISPResso2 provides accurate and rapid genome editing

sequence analysis. *Nat Biotechnol*. 2019;37(3):224-226. doi:10.1038/s41587-019-0032-3

39. Mei L, Fan Y, Lv X, Welsh DK, Zhan C, Zhang EE. Long-term in vivo recording of circadian

rhythms in brains of freely moving mice. *Proc Natl Acad Sci U S A*. 2018;115(16):4276-4281.

doi:10.1073/pnas.1717735115

40. Ramanathan C, Khan SK, Kathale ND, Xu H, Liu AC. Monitoring Cell-autonomous Circadian Clock

Rhythms of Gene Expression Using Luciferase Bioluminescence Reporters. *J Vis Exp JoVE*.

2012;(67):4234. doi:10.3791/4234

41. Jiang H, Lei R, Ding SW, Zhu S. Skewer: a fast and accurate adapter trimmer for next-generation

sequencing paired-end reads. *BMC Bioinformatics*. 2014;15:182. doi:10.1186/1471-2105-15-182

42. Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of

RNA-seq data. *Genome Biol*. 2010;11(3):R25. doi:10.1186/gb-2010-11-3-r25

43. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression

analysis of digital gene expression data. *Bioinforma Oxf Engl*. 2010;26(1):139-140.

doi:10.1093/bioinformatics/btp616

44. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters | OMICS: A

Journal of Integrative Biology. Accessed August 28, 2022.

https://www.liebertpub.com/doi/10.1089/omi.2011.0118

45. Euhus DM, Hudd C, Laregina MC, Johnson FE. Tumor measurement in the nude mouse. *J Surg

Oncol*. 1986;31(4):229-234. doi:10.1002/jso.2930310402

46. Therneau T. A Package for Survival Analysis in R. https://CRAN.R-project.org/package=survival

47. Fu L, Kettner NM. The circadian clock in cancer development and therapy. *Prog Mol Biol Transl Sci*.

2013;119:221-282. doi:10.1016/B978-0-12-396971-2.00009-9

48. Pogue-Geile KL, Lyons-Weiler J, Whitcomb DC. Molecular overlap of fly circadian rhythms and

human pancreatic cancer. *Cancer Lett*. 2006;243(1):55-57. doi:10.1016/j.canlet.2005.11.049

49. Schwartz PB, Barrett-Wilt GA, Ronnekleiv-Kelly SM. Chronic Jetlag Alters the Landscape of the

Pancreatic Lipidome. *Pancreas*. 2022;51(1):80-89. doi:10.1097/MPA.0000000000001962

50. Altman BJ, Hsieh AL, Sengupta A, et al. MYC Disrupts the Circadian Clock and Metabolism in

Cancer Cells. *Cell Metab*. 2015;22(6):1009-1019. doi:10.1016/j.cmet.2015.09.003

51. McIntosh BE, Hogenesch JB, Bradfield CA. Mammalian Per-Arnt-Sim proteins in environmental

adaptation. *Annu Rev Physiol*. 2010;72:625-645. doi:10.1146/annurev-physiol-021909-135922

52. Vazquez-Rivera E, Rojas B, Parrott JC, et al. The aryl hydrocarbon receptor as a model PAS sensor.

Toxicol Rep. 2022;9:1-11. doi:10.1016/j.toxrep.2021.11.017

53. Evan GI, Vousden KH. Proliferation, cell cycle and apoptosis in cancer. *Nature*.

2001;411(6835):342-348. doi:10.1038/35077213

54. Mini E, Nobili S, Caciagli B, Landini I, Mazzei T. Cellular pharmacology of gemcitabine. *Ann Oncol

Off J Eur Soc Med Oncol*. 2006;17 Suppl 5:v7-12. doi:10.1093/annonc/mdj941

55. Kampan NC, Madondo MT, McNally OM, Quinn M, Plebanski M. Paclitaxel and Its Evolving Role

in the Management of Ovarian Cancer. *BioMed Res Int*. 2015;2015:413076.

doi:10.1155/2015/413076

56. Goldstein D, El-Maraghi RH, Hammel P, et al. nab-Paclitaxel plus gemcitabine for metastatic

pancreatic cancer: long-term survival from a phase III trial. *J Natl Cancer Inst*. 2015;107(2).

doi:10.1093/jnci/dju413

57. de Sousa Cavalcante L, Monteiro G. Gemcitabine: metabolism and molecular mechanisms of action,

sensitivity and chemoresistance in pancreatic cancer. *Eur J Pharmacol*. 2014;741:8-16.

doi:10.1016/j.ejphar.2014.07.041

58. Brown MR, Matveyenko AV. It's What and When You Eat: An Overview of Transcriptional and

Epigenetic Responses to Dietary Perturbations in Pancreatic Islets. *Front Endocrinol*.

2022;13:842603. doi:10.3389/fendo.2022.842603

59. Fadista J, Vikman P, Laakso EO, et al. Global genomic and transcriptomic analysis of human

pancreatic islets reveals novel genes influencing glucose metabolism. *Proc Natl Acad Sci U S A*.

2014;111(38):13924-13929. doi:10.1073/pnas.1402665111

60. Korpela T, Ristimäki A, Udd M, et al. Pancreatic fibrosis, acinar atrophy and chronic inflammation in

surgical specimens associated with survival in patients with resectable pancreatic ductal

adenocarcinoma. *BMC Cancer*. 2022;22(1):23. doi:10.1186/s12885-021-09080-0

61. Archer SN, Laing EE, Möller-Levet CS, et al. Mistimed sleep disrupts circadian regulation of the

human transcriptome. *Proc Natl Acad Sci U S A*. 2014;111(6):E682-E691.

doi:10.1073/pnas.1316335111

62. Wu G, Ruben MD, Francey LJ, Lee YY, Anafi RC, Hogenesch JB. An in silico genome-wide screen

for circadian clock strength in human samples. Published online May 25, 2022:2022.05.10.491250.

doi:10.1101/2022.05.10.491250

63. Yu X, Rollins D, Ruhn KA, et al. TH17 cell differentiation is regulated by the circadian clock.

Science. 2013;342(6159):727-730. doi:10.1126/science.1243884

64. Druzd D, Matveeva O, Ince L, et al. Lymphocyte Circadian Clocks Control Lymph Node Trafficking

and Adaptive Immune Responses. *Immunity*. 2017;46(1):120-132. doi:10.1016/j.immuni.2016.12.011

65. Conroy T, Hammel P, Hebbar M, et al. FOLFIRINOX or Gemcitabine as Adjuvant Therapy for

Pancreatic Cancer. *N Engl J Med*. 2018;379(25):2395-2406. doi:10.1056/NEJMoa1809775

66. Altman BJ, Hsieh AL, Gouw AM, Dang CV. Correspondence: Oncogenic MYC persistently

upregulates the molecular clock component REV-ERBα. *Nat Commun*. 2017;8:14862.

doi:10.1038/ncomms14862

67. Murthy D, Attri KS, Singh PK. Phosphoinositide 3-Kinase Signaling Pathway in Pancreatic Ductal

Adenocarcinoma Progression, Pathogenesis, and Therapeutics. *Front Physiol*. 2018;9:335.

doi:10.3389/fphys.2018.00335

68. Bryant KL, Stalnecker CA, Zeitouni D, et al. Combination of ERK and autophagy inhibition as a

treatment approach for pancreatic cancer. *Nat Med*. 2019;25(4):628-640. doi:10.1038/s41591-019-

0368-8

69. Hoxhaj G, Manning BD. The PI3K-AKT network at the interface of oncogenic signalling and cancer

metabolism. *Nat Rev Cancer*. 2020;20(2):74-88. doi:10.1038/s41568-019-0216-7

70. Mortazavi M, Moosavi F, Martini M, Giovannetti E, Firuzi O. Prospects of targeting

PI3K/AKT/mTOR pathway in pancreatic cancer. *Crit Rev Oncol Hematol*. 2022;176:103749.

doi:10.1016/j.critrevonc.2022.103749

71. Maloney SM, Hoover CA, Morejon-Lasso LV, Prosperi JR. Mechanisms of Taxane Resistance.

Cancers. 2020;12(11):E3323. doi:10.3390/cancers12113323

72. Yang XL, Lin FJ, Guo YJ, Shao ZM, Ou ZL. Gemcitabine resistance in breast cancer cells regulated

by PI3K/AKT-mediated cellular proliferation exerts negative feedback via the MEK/MAPK and

mTOR pathways. *OncoTargets Ther*. 2014;7:1033-1042. doi:10.2147/OTT.S63145

73. Li W, Zhu Y, Zhang K, et al. PROM2 promotes gemcitabine chemoresistance via activating the Akt

signaling pathway in pancreatic cancer. *Exp Mol Med*. 2020;52(3):409-422. doi:10.1038/s12276-020-

0390-4

Figures:

Figure 1: *The human pancreatic clock robustness is diminished after malignant transformation.* **A**. Clock

correlation matrices were created for normal (left; n = 50) and pancreatic ductal adenocarcinoma (right; n

= 318) samples. **B**. The normalized coefficient of variation (nCV) was calculated to determine the

robustness of the pancreatic clock for the normal (circles) and PDAC (triangles) samples. Each color

indicates a different gene evaluated.

Figure 2: *Clock phase predictions demonstrate conserved clock relationships*: **A.** Graphical

representation and **B.** table of the CCG and clock-controlled gene phase prediction from the cosinor

analysis of CYCLOPS reordered human pancreas samples (n = 50) and Metacycle predicted phases of

mouse pancreas samples (n = 72), relative to *BMAL1* (human) and *Bmal1* (mouse) phase.

Figure 3: *The human pancreatic cancer tumor clock is dysfunctional relative to the normal pancreas.* **A**.

CYCLOPS was used to reorder samples from normal (n = 50) and pancreatic ductal adenocarcinoma (n =

318) TCGA and CPTAC-3 samples. Plots from several core clock genes reordered by CYCLOPS in

normal tissue as compared to best reordering in PDAC (**A**) are shown – ordered from 0 to 2π. Shading

around the blue regression line indicates the 95% confidence interval. **B**. Graphical representation of

Phase Set Enrichment Analysis (PSEA) of normal pancreatic samples, ordered by phase of expression.

Figure 4: *Creation of a mouse clock dysfunction cell line model.* **A**. The core clock gene *Bmal1* was

mutated (functional knock out) in KPC murine pancreatic cells with CRISPR-Cas9 genome editing

technology (BKO) with the frameshift site indicated by the blue line. **B.** The frameshift mutation was

induced upstream of the basic helix loop helix (bHLH) domain (red star). The resulting protein lacked

critical downstream elements for functionality, including the PAS-A, PAS-B, and PAC domains. **C.**

Western blot demonstrating the loss of BMAL1 protein across 24 hours for wild-type (WT) and BKO

cells (M = marker lane). **D.** The mean (± standard error) *Per1* mRNA expression of synchronized KPC (n

= 3 at each time point; black dashed) and KPC-BKO (n = 3 at each time point; grey) cells from ZT0-

ZT24. **E.** Graphs comparing the percent of cells in G1, S, and G2 in WT (n = 3; white) and BKO (n = 3;

grey) KPC cells.

Figure 5: *Mutation of Bmal1 promotes pancreatic cancer progression.* **A.**Comparison of mouse tumor

weight at 28 days for WT (n = 20; white) and BKO (n = 20) KPC tumors.**B.** WT (n = 20; black) and BKO

(n = 20) mean (± standard error) tumor size (in mm^3) over the 28-day growth period. **C.** Kaplan Meier

survival curve for mice implanted with WT (n = 7; black) and BKO (n =7; grey) KPC tumors. The dotted

lines indicate the median survival. [ns = not significant, * = p < 0.05, ** = p < 0.01]

Figure 6: *Functional knockout of Bmal1 promotes widespread transcriptomic alterations and activation*

of multiple oncogenic pathways. **A.** Principal Component Analysis (PCA) demonstrating overall

differences in expression between WT(green) and BKO (purple) samples (n = 6 each) **B.** Volcano plot

showing genes that were significantly upregulated (red) and downregulated (blue) on differential gene

expression (DGE) analysis (q < 0.05). All significant core clock genes are highlighted **C.** KEGG analysis

was then performed, and the top 9 pathways ordered by significance are shown. For each pathway, the

size of each dot corresponds to the number of genes involved in each pathway.

Figure 7: *Clock dysfunction diminishes the chemotherapeutic cell death response.* KPC wildtype (WT)

and KPC-BKO cells were treated with increasing doses of gemcitabine and paclitaxel. Fold change

differences (± standard error) in Caspase 3/7 activity in response to either **A.** gemcitabine or **B.** paclitaxel.

Fold change differences (± standard error) in dead-cell protease activity in response to **C.** gemcitabine or

D. paclitaxel. [ns = not significant, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$]

Supplementary Files:

Supplementary Data Files:

Supplemental Data File 1: CYCLOPS seed gene list.

Supplemental Data File 2: Significant genes on cosinor analysis for normal CYCLOPS reordered

normal pancreas

Supplemental Data File 3: Significant genes on cosinor analysis for normal CYCLOPS reordered PDAC

Supplemental Data File 4: KPC wildtype versus BKO edgeR differential gene expression

Supplementary Figures:

Supplemental Figure 1: *The wildtype mouse pancreatic clock is intact and robust.* **A**. Clock correlation

matrices were created for normal (left, n = 72) and pancreatic ductal adenocarcinoma (right, n = 72)

samples. **B**. The normalized coefficient of variation (nCV) was calculated to determine the robustness of

the pancreatic clock for the normal circadian (circles) and chronic jetlag (triangles) pancreas samples.

Each color indicates a different gene evaluated.

Supplemental Figure 2: *CYCLOPS accurately reorders wildtype mouse pancreas samples.* Eigengenes

identified by Oscope for use in CYCLOPS in **A.** normal circadian (upper) and chronic jetlag (lower)

samples. Shading around the blue regression line indicates the 95% confidence interval for each plot. **B.**

The normal circadian (left) and chronic jetlag (right) clusters ordered by CYCLOPS demonstrate accurate

reordering for both conditions **C.** Normal circadian (left) and chronic jetlag (right) genes found to be

significantly rhythmic on both CYCLOPS reordered cosinor analysis and rhythmicity testing based on the

known sample collection time with the Metacycle meta3d function are ordered by their predicted phase of

expression. Clock genes are shown in orange.

Supplemental Figure 3: *Normal pancreatic and pancreatic ductal adenocarcinoma (PDAC) samples*

were processed and filtered. TCGA and CPTAC-3 samples were processed and batched corrected. An

MDS plot is shown demonstrating differences between the matched normal (green; n = 50) and PDAC

(purple; n = 318) samples.

Supplemental Figure 4: *Eigengenes for the human normal pancreas and pancreatic ductal*

adenocarcinoma (PDAC) samples. Eigengenes were selected by Oscope for normal (upper) and PDAC

(lower) samples for use in CYCLOPS. Shading around the blue regression line indicates the 95%

confidence interval for each plot.

Supplemental Figure 5: *KPC cells expressing dLuciferase driven by the Per2 promoter exhibit circadian*

activity. Two independent clones were examined, and luciferase activity was measured at 4-hour

intervals. Rhythmicity was calculated with Metacycle. Clone 1 (left) was found to be rhythmic with a $q =$

5.51E-5 and rAMP = 0.44. Clone 2 (right) was found to be rhythmic with a $q = 1.95E-7$, rAMP = 0.39.

Supplemental Figure 6: *Cross-validation of mutant Bmal1 (functional knockout) growth in a second*

independent clone. A second independent *Bmal1* knockout (BKO) clone with an identical mutation to the

first was validated with **A.** western blot analysis. We then heterotopically implanted wildtype (WT) and

BKO cells into C57Bl/6J mice and followed **B.** tumor weight and **C.** cell growth. BKO_2 (dark grey; n =

20) had a similar mean (± standard error) tumor weight to BKO_1 (light grey; n = 20) (426.09 (± 40.07)

mg vs 438.02 (± 48.84) mg; p = 0.85), and both BKO_2 (426.09 (± 40.07) mg vs 280.11 (± 42.73) mg; p

= 0.017) and BKO_1 (438.02 (± 48.84) mg vs 280.11 (± 42.73) mg; p = 0.02) were larger than WT

(white; n = 20). BKO_1 and BKO_2 had similar significantly faster growth trajectories. [ns = not

significant, * = p < 0.05, ** = p < 0.01, *** = p < 0.001]

Supplementary Tables

Supplemental Table 1: Cosinor rhythmicity analysis of CYCLOPS ordered normal pancreas and

PDAC

A.

Mouse | Human

B.

Mouse Gene	Phase	Human Gene	Phase
Bmal1	0	BMAL1	0
Clock	23	CLOCK	4.9
Cry1	17.9	CRY1	1.5
Cry2	15	CRY2	13.6
Per1	13	PER1	11.7
Per2	15	PER2	11.6
Per3	12.4	PER3	11.8
Nr1d1	9	NR1D1	11.9
Nr1d2	11	NR1D2	12.5
Rora	9	RORA	18.8
Rorc	19	RORC	11.7
Npas2	0.3	NPAS2	10.6
Nfil3	21	NFIL3	23.7
Bhlhe41	13	BHLHE41	19.1
Dbp	9.9	DBP	10.9
Ciart	12.1	CIART	10
Tef	12.3	TEF	12
Hlf	11	HLF	18.9

A. Frameshift Mutation

5' | 25,000 | 50,000 | 75,000 | 100,000 | 3'

Bmal1
(116,662 bp)

B.

N — NLS — ★ — bHLH — PAS-A — PAS-B — PAC — C

aa36-41   aa79-132   aa333-403

aa150-222   aa408-451

Heterodimerization

DNA Binding

C.

WT   BKO

M   ZT0 ZT6 ZT12 ZT18   ZT0 ZT6 ZT12 ZT18

250 kDa
130 kDa
100 kDa
70 kDa   ← BMAL1
55 kDa
← ACTB
35 kDa
25 kDa
15 kDa

D. *Per1*

● WT
● BKO

Delta CT

Zeitgeber Time

E.

Percent Total G1 (%)   0.2   WT BKO

Percent Total S (%)   0.029   WT BKO

Percent Total G2 (%)   0.0053   WT BKO