<u>The circadian clock is disrupted in pancreatic cancer</u>

Authors: Patrick B. Schwartz, MD PhD[1], Manabu Nukaya, PhD[1], Mark E. Berres, PhD[2], Clifford D.

Rubinstein, PhD[2], Gang Wu, PhD[3], John B. Hogenesch, PhD[3], Christopher A. Bradfield, PhD[4], and Sean

M. Ronnekleiv-Kelly, MD[1,5]

Affiliations:

[1]Department of Surgery, Division of Surgical Oncology, University of Wisconsin School of Medicine and

Public Health, Madison, WI, USA

[2]Biotechnology Center, University of Wisconsin, Madison, WI, USA

[3]Division of Human Genetics and Immunobiology, Center for Circadian Medicine, Department of

Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

[4]McArdle Laboratory for Cancer Research, Department of Oncology, University of Wisconsin School of

Medicine and Public Health, Madison, WI, USA

[5]University of Wisconsin Carbone Cancer Center, University of Wisconsin School of Medicine and

Public Health, Madison, WI, USA

Corresponding Author: Sean M Ronnekleiv-Kelly, MD (ronnekleiv-kelly@surgery.wisc.edu)

Address: 600 Highland Ave. Bx 7375, K3/705 CSC, Madison, WI, 53792

Phone: (608) 262-2025 **Fax:** (608) 252-0913

Running Title: Pancreatic cancer clock

This manuscript has been submitted as a preprint to BioRxiv, last updated on 11/8/22

Funding: The research reported in this publication was supported by the Department of Defense Peer

Reviewed Cancer Research Program Idea Award Number CA190176 (Grants.gov ID GRANT12935023)

(SRK), by the National Institutes of Health (NIH) National Institute of Environmental Health Sciences

(NIEHS) Award Number R35ES028377 (CAB), by the NIH NIEHS Award Number T32 ES007015

(PBS) and the University of Wisconsin Carbone Cancer Center Support Grant P30CA014520 (SRK). The

content is solely the responsibility of the authors and does not necessarily represent the official views of

the Department of Defense or the National Institutes of Health.

Disclosure: The authors declare no conflict of interest or competing interests

<u>Abstract</u>

Disruption of the circadian clock is linked to cancer development and progression. Establishing this

connection has proven beneficial for understanding cancer pathogenesis, determining prognosis, and

uncovering novel therapeutic targets. However, barriers to characterizing the circadian clock in human

pancreas and human pancreatic cancer – one of the deadliest malignancies – have hindered an

appreciation of its role in this cancer. Here, we employed normalized coefficient of variation (nCV) and

clock correlation analysis in human population-level data to determine the functioning of the circadian

clock in pancreas cancer and adjacent normal tissue. We found a substantially attenuated clock in the

pancreatic cancer tissue. Then we exploited our existing mouse pancreatic transcriptome data to perform

an analysis of the human normal and pancreas cancer samples using a machine learning method, cyclic

ordering by periodic structure (CYCLOPS). Through CYCLOPS ordering, we confirmed the nCV and

clock correlation findings of an intact circadian clock in normal pancreas with robust cycling of several

core clock genes. However, in pancreas cancer, there was a loss of rhythmicity of many core clock genes

with an inability to effectively order the cancer samples, providing substantive evidence of a dysregulated

clock. The implications of clock disruption were further assessed with a *Bmal1* knockout pancreas cancer

model, which revealed that an arrhythmic clock caused accelerated cancer growth and worse survival,

accompanied by chemoresistance and enrichment of key cancer-related pathways. These findings provide

strong evidence for clock disruption in human pancreas cancer and demonstrate a link between circadian

disruption and pancreas cancer progression.

<u>Author Summary</u>

The circadian clock is a regulator of human homeostasis. Dysfunction of the clock can lead to the

development of diseases, including cancer. Although several cancers have been shown to have a

dysfunctional clock which may alter prognosis or change treatment, this has been suggested but not

demonstrated in pancreatic cancer. Investigation of this link is important because pancreatic cancer is

highly lethal with few effective treatment options. Here we use recently pioneered bioinformatics

approaches to assess clock functionality in human pancreatic cancer specimens, where we demonstrate

that the clock is dysfunctional relative to normal pancreatic tissue. We then knocked out the core clock

gene, *Bmal1*, in pancreatic cancer cells, which led to faster tumor growth and worse survival in mice and

enhanced chemotherapeutic resistance to standard chemotherapy agents used in the treatment of

pancreatic cancer. Collectively, our findings establish human pancreatic cancer as having clock

dysfunction and clock dysfunction causing a more aggressive cancer.

<u>Introduction</u>

The circadian clock is a conserved molecular feedback loop that regulates many signaling pathways to

control metabolism, immunity, apoptosis, and other critical cellular functions in the body (1). At its core,

the positive arm of the clock (*i.e.,* CLOCK and BMAL1 [also known as ARNTL or MOP3]) drives

transcription of the negative arm, including PER1-3 and CRY1-2 (2,3). The negative arm represses the

transcriptional activation of CLOCK and BMAL1, and a second interlocked loop involves the nuclear

receptors RORα/β/λ and NR1D1-2 (also known as REV-ERBα/β), which activate and suppress BMAL1

expression, respectively. These, along with other core clock components, form a tightly regulated series

of transcriptional-translational feedback loops that ensure rhythmic expression over 24 hours and function

to maintain cellular and organ homeostasis (2–4).

Environmental cues influence the synchronization of circadian rhythms in various organ systems, and

misalignment of external cues with the internal clock (*e.g*., shift work) can cause clock dysfunction with

consequent metabolic derangements and pathologic states (4–6). For instance, circadian dysregulation has

been strongly linked to obesity and diabetes, both risk factors for cancer (7–11). Concordantly, landmark

studies have shown that disruption of the endogenous clock through mutations in or suppression of the

core clock genes is intricately linked to tumor growth in several cancers (12–14). For example, knockout

of *Bmal1* in *Kras-* and *p53*-mutant lung cancer causes marked tumor progression *in vivo* (13), while

MYC-induced repression of *BMAL1* in human neuroblastoma drives decreased overall survival in a

BMAL1-dependent manner (15). Importantly, targeting dysfunctional clock components in certain

cancers has proven an effective treatment strategy (15,16). Thus, identifying an aberrantly functioning

circadian clock in cancer can lead to key advancements such as understanding pathogenesis, prognosis,

and uncovering novel therapeutic targets.

Although indeterminate, there is some evidence that the clock may be dysregulated in pancreatic ductal

adenocarcinoma (PDAC), leading to a worse prognosis (17,18); this is alarming for a deadly malignancy

where only 11% of patients survive beyond 5 years (19). To advance our understanding of how clock

disruption impacts PDAC pathogenesis, and ultimately foster the identification of therapeutic targets, an

essential first step is to establish that clock dysfunction exists in human PDAC. Unfortunately, to date, the

cumulative data does not definitively support this assertion and is inconclusive. Prior studies have relied

on contrasting expression differences between the core clock genes in tumor compared to normal

pancreas as a basis for clock disruption (17,18,20,21). But differential expression alone is limited and

does not provide insight into critical components of clock health such as relative amplitude, rhythmicity,

or correlation of expression amongst core clock genes (22). For example, in the pancreas, phase

advancement as a result of chronic jetlag causes differential expression of core clock genes despite

maintaining a robust and healthy clock (*i.e.,* nearly identical relative amplitude, rhythmicity, and clock

correlation) (23). Furthermore, in population-level data where sample time acquisition is unknown and

the internal clock from individual patient tumors may not be in phase, the interpretation of differential

expression for evaluating clock function becomes challenging. Thus, much more substantive data is

required to affirm clock disruption in PDAC.

In pre-clinical studies (*e.g.*, mouse or cell-line models), the determinants of clock health can be identified

by obtaining longitudinal data under controlled conditions (23,24). However, in human studies –

particularly human pancreas – the requisite periodic data by multiple sampling is not feasible (or ethical).

Therefore, alternative means of determining rhythmically expressed genes in humans are necessary.

Following the emergence of several bioinformatics tools, the principal aspects of clock health including

relative amplitude, core clock gene correlation, and statistical determination of rhythmicity can be

resolved when using the appropriate reference data (22,25–28). We recently generated a robust pancreas

dataset demonstrating diurnally expressed genes over 48 hours (23). With this foundation, we were able

to apply normalized coefficient of variation (nCV), clock correlation analysis, and cyclic ordering by

periodic structure (CYCLOPS) to evaluate the principal aspects of clock function, respectively, and test

the hypothesis that the circadian clock is disrupted in human PDAC (22,25,28). While on the surface this

may be construed as a simplistic hypothesis, several limitations have hindered the ability to evaluate the

circadian clock in human pancreas, including the lack of human periodic data and an absence of a

reference transcriptional dataset.

Here, we employed nCV, clock correlation analysis, and CYCLOPS on publicly available human

population-level expression data from The Cancer Genome Atlas (TCGA) and Clinical Proteomic Tumor

Analysis Consortium 3 (CPTAC-3) datasets to determine the health of the circadian clock in PDAC and

adjacent normal tissue (29,30). We identified an intact circadian clock in human normal pancreas with

robust cycling of several core clock genes. We also found a markedly weakened clock in the cancer

tissue, providing substantive evidence of a dysregulated clock in PDAC. These findings represent

significant advancements in evaluating clock function in PDAC, and the potential clinical implications of

clock disruption were further assessed with a pre-clinical PDAC model. We used CRISPR/Cas9

technology to selectively target *Bmal1* and examined the effects of clock dysfunction *in vitro* and *in vivo*.

This revealed that loss of clock function caused accelerated cancer growth, diminished survival,

enrichment of key cancer-related pathways, and resistance to commonly used cytotoxic chemotherapies

for PDAC. These findings provide strong evidence for circadian clock disruption in human PDAC and

demonstrate a link between circadian disruption and pancreas cancer progression.

Results

The Clock in Human Pancreatic Ductal Adenocarcinoma is Less Robust Than in Normal Pancreatic

Tissue

To test the hypothesis that the circadian clock is disrupted in human pancreatic ductal adenocarcinoma

(PDAC), we used our bioinformatics pipeline that consists of clock correlation, normalized coefficient of

variation (nCV), and cyclic ordering by periodic structure (CYCLOPS) (**Fig 1**). We performed an analysis

on publicly available The Cancer Genome Atlas (TCGA) and Clinical Proteomics Tumor Analysis

Consortium 3 (CPTAC-3) pancreatic datasets (29,30). However, we first needed to ensure the validity of

the pipeline in the pancreas since these had not been previously applied to pancreatic tissue. We assessed

the clock in our existing murine normal circadian and chronic jetlag pancreas RNA-sequencing (RNA-

seq) datasets by examining the clock correlation matrix and nCV (23). These datasets contained pancreas

samples acquired every 4 hours for 48 hours in male (n = 3 at each timepoint for each condition) and

female (n = 3 at each timepoint for each condition) mice under standard lighting (normal circadian, n =

72) and chronic jetlag (n = 72) conditions. Chronic jetlag is known to affect the phase of gene expression

but not the relative amplitude (rAMP) or the correlation of the core clock genes in the pancreas (23).

Correspondingly, we found that the core clock relationships (as measured by the z-statistic or standard

deviations above/below population mean) remained intact on the correlation matrix when compared to the

baseline clock correlation matrix for both normal and chronic jetlag conditions ($p < 0.001$, z-statistic =

54.96 vs. $p < 0.001$, z-statistic = 54.08) (**Fig 2A**) (31,32). Further, the nCV of 11 clock genes remained

unchanged ($p = 0.76$) between the normal circadian (mean nCV ($\pm$ standard error) = 1.78 (± 0.27)) and

chronic jetlag (mean nCV = 1.9 (± 0.28)) conditions (**Fig 2B-C**) – consistent with the murine pancreatic

clock being strongly rhythmic and intact, matching precisely what we had found in our prior longitudinal

analysis (23).

CYCLOPS was then used to reorder the mouse pancreas samples by their predicted phase, and cluster

reordering of samples by CYCLOPS was analyzed for appropriate phase progression compared to the

known zeitgeber time (ZT) of sample collection. CYCLOPS reordering can be assessed by the Met^{smooth}

and $Stat^{err}$ (significant reordering: $Met^{smooth} < 1$ and $Stat^{err} < 0.05$). A full mathematical derivation has

been previously outlined by Anafi *et al* (28). Notably, CYCLOPS accurately (phase appropriately)

reordered both the normal ($p < 0.01$; $Met^{smooth} = 0.90$) and the chronic jetlag conditions ($p < 0.01$;

$Met^{smooth} = 0.97$), validating the pipeline for use in the pancreas (**Supplemental Fig 1A-C**). Collectively,

these results confirmed the ability of nCV, correlation matrix, and CYCLOPS to determine the robustness

of the circadian clock in the pancreas.

Considerable data support the tumor suppressor role of the circadian clock and the assertion that circadian

disruption is present in human cancer (33). In human PDAC, the supposition is that the clock may be

altered, but supporting data is limited to the relative expression of single genes and is consequently

inconclusive (18,21,34). We therefore applied our pipeline to determine clock health in PDAC using the

TCGA and CPTAC-3 pancreatic datasets (22,26–30). After processing and batch correction of the

matched 50 available normal and 318 available PDAC samples (**Supplemental Fig 2**), we examined the

correlation matrix between the core clock genes (CCGs) and found that there was a weaker relationship

amongst the CCGs in PDAC compared to normal (z-statistic PDAC = 13.77 vs. z-statistic normal =

22.42) (**Fig 2D**). Concordantly, we found that there was a clear decrease ($p = 0.04$) in the nCV between

PDAC (mean nCV = 0.69 ($\pm$ 0.08)) and normal (mean nCV = 1.03 ($\pm$ 0.13)) (**Fig 2E-F**), indicating a

weaker clock in PDAC compared to normal pancreas. We then applied our pancreatic CYCLOPS analysis

which significantly reordered the normal samples ($p < 0.01$; Met^{smooth} = 0.99), indicating an intact clock.

This was further reinforced when evaluating the predicted phase of CCGs and clock-controlled genes

using CYCLOPS. We found that the phase sequence of normal pancreatic samples was similar to the

phase sequence of mouse pancreatic samples (**Fig 3A-B**), with the phase of expression in the human

genes equivalent to the mouse genes in 12 out of 17 CCGs and clock-controlled genes and ROR-phased

genes largely peaking before E-box phase genes. Based on the CYCLOPS ordering, we analyzed the

proportion of genes that were rhythmically expressed in normal pancreas. We found that 4,034/18,196

(22.17%) of normal pancreatic genes were rhythmic, (**Supplemental Data File 1**). We found that several

key CCGs and clock-controlled genes were rhythmically expressed in normal samples on cosinor

analysis, including *CLOCK*, *PER1*, *PER3*, *NR1D2*, *RORC*, *NFIL3, TEF*, *NR1D1, BHLHE40,* and *NPAS2*

(**Fig 4A and Supplemental Table 1**). The CCGs *BMAL1* ($p = 0.07$, rAMP = 0.8) and *CRY1* ($p = 0.08$,

rAMP = 0.48) were nearly rhythmic in the normal samples based on the pre-defined cutoff. Finally, phase

set enrichment analysis (PSEA) was used to determine phase-dependent gene enrichment. The top 15

significant pathways were selected for analysis (**Fig 4B**). Consistent with our prior murine pancreas data,

we found that normal pancreas was associated with time-dependent metabolic gene pathway enrichment

(23,35). In aggregate, this data confirms the health of the circadian clock in tumor-adjacent normal

pancreatic tissue, which was expected but not previously shown to our knowledge.

Despite the validation steps and success in ordering normal pancreas, CYCLOPS **could not** reorder

PDAC samples ($p = 0.43$; Met$^{\text{smooth}}$ = 0.99), revealing a less robust (or disrupted) clock in PDAC

compared to normal. This inability to reorder strongly supports the attenuated circadian clock in PDAC

identified with nCV and clock correlation analysis. We selected the best reordering available based on the

eigengenes to enable visual representation of the PDAC samples compared to normal pancreas samples.

This also enabled us to estimate as best as possible the rAMP of the PDAC samples, to coincide with the

visual representation. Consistent with the nCV analysis, we found that rAMP decreased substantially in

every core clock gene from normal to tumor (**Supplemental Table 1**), as evidenced by *PER1* and *PER3*

expression (**Fig 4A**). Collectively, the weaker clock correlation, markedly reduced nCV, and CYCLOPS

analysis of human population-level data show that the circadian clock is significantly disrupted in PDAC.

Creation of a Circadian Dysfunction Pancreatic Cancer Model

After demonstrating that the clock in human PDAC was dysfunctional, we sought to understand the

potential clinical implications of clock disruption. Identifying an aberrant clock in other cancers has

consistently demonstrated accelerated cancer progression and worse prognosis, while simultaneously

revealing novel therapeutic targets (13–16). Moreover, there appears to be a putative correlation between

suppressed *BMAL1* expression and poor prognosis in patients with PDAC (17,18). Therefore, we

hypothesized that disruption of the circadian clock in PDAC would cause accelerated cancer progression.

To test this hypothesis, we used a syngeneic *Kras-* and *Trp53*-mutant pancreas cancer cell line (KPC) and

employed CRISPR/Cas9 technology to functionally knockout *Bmal1* (KPC-BKO) so that we could

examine the effects of clock dysfunction *in vitro* and *in vivo* (36). The reason for mutating *Bmal1* in the

PDAC cells was several-fold: i) KPC cells demonstrated an intact clock (**Fig 5D**) necessitating core clock

gene modulation, ii) *BMAL1* is a central transcriptional regulator of the circadian clock machinery and

functional knockout of this single gene abolishes circadian function (37,38), iii) suppressed *BMAL1*

expression has been found in several human cancers, correlating with worse prognosis

(13,15,17,18,20,21,39), iv) *BMAL1* expression is decreased in tumor compared to normal tissue in human

PDAC (18), v) we similarly found a substantially dampened nCV (rAMP) of *BMAL1* expression in

PDAC compared to normal tissue (*PER1*, *PER3*, *NR1D1,* and *BMAL1* displayed the greatest decrease in

nCV), and vi) segregating patients with PDAC into high and low *BMAL1* expression appears to be

prognostic for survival outcomes (17,18,21).

Consequently, guide RNAs were designed to target exon 8 of the *Bmal1* gene, which resulted in the

insertion of adenine on one allele and a 2 base pair deletion on the other between nucleotide 81,074 and

81,075 at amino acid 73 (GRCm38) (**Fig 5A**). The result of both mutations was a frameshift just upstream

of the basic helix loop helix (bHLH) domain known to assist the PAS A domain in heterodimerization

with its binding partner CLOCK (**Fig 5B**) (40,41). Confirmatory western blotting revealed the presence of

protein in wild-type (WT) cells and an absence of protein in *Bmal1* knockout (BKO) KPC cells (**Fig 5C**).

To evaluate clock functionality, mRNA was isolated at 4-hour intervals (n = 3 per condition) for 24 hours

after cell synchronization, and we performed RT-qPCR for *Per1*, a core downstream repressor of the

positive arm of the molecular clock and a gene that demonstrated rhythmicity in our human normal

CYCLOPS analysis (**Fig 5D**). Rhythmicity was assessed with the meta2d function of the Metacycle

package (42). KPC (WT) cells exhibited an intact and robust circadian clock with a $q = 0.0004$ and rAMP

$= 0.11$ for *Per1*. Although still rhythmic, when compared to normal murine pancreas core clock gene

function from our prior longitudinal transcriptional dataset (normal pancreas *Per1*: $q = 0.0001$, rAMP =

0.394), the relative amplitude was diminished (23). To further evaluate the functionality of the clock in

KPC cells, luciferase activity driven by the *Per2* promoter was measured. Luciferase activity in two

separate clones was highly rhythmic (Clone 1: $q = 5.51E-5$, rAMP = 0.44; Clone 2: $q = 1.95E-7$, rAMP =

0.39) (**Supplemental Fig 3**). Consistent with the comparison of *Per1* expression in KPC versus our prior

normal murine pancreas dataset, the relative amplitude of *Per2* was diminished in the KPC lines

compared to normal pancreas tissue ($q = 1.39E-7$, rAMP = 0.82). Thus, although rAMP was diminished

in the murine pancreas cancer cell line compared to normal pancreas tissue, clock function was still intact.

Conversely, KPC-BKO cells exhibited no detectable clock function with a $q = 0.81$ and rAMP = 0.003 for

Per1 expression (**Fig 5D**). In addition, we looked for differences in the cell cycle consequent to clock

dysfunction as the two pathways are known to interact (43,44). For instance, Bieler *et al.* identified robust

synchrony between cell division and circadian cycle in NIH3T3 fibroblasts that contained a *Rev-Erba-*

YFP reporter (43), while Feillet *et al*. used a similar system (NIH3T3 fibroblasts) to identify a coordinate

entrainment in unsynchronized cells as well as cells that were dexamethasone synchronized (44).

Consistent with this known cross-talk between the circadian clock and cell cycle, KPC-BKO cells

demonstrated alterations in the cell cycle compared to KPC cells (**Fig 5E**) (20). BKO cells were found to

have a higher mean (± standard error) percentage of cells in S phase (48.66 (± 0.13) vs. 36.96 (± 2.04); *p*

= 0.029) and a lower percentage in G2 (18.03 (± 0.27) vs. 27.49 (± 0.89); *p* = 0.0053). There were no

differences between KPC and KPC-BKO cells in G1 (35.56 (± 1.19) vs. 33.31 (± 0.17); *p* = 0.2). Taken

together, these data demonstrate the successful creation of a novel PDAC cell line with an abolished clock

(*Bmal1* gene mutation), which can be used to elucidate the repercussions of clock disruption in PDAC.

Clock Dysfunction Accelerates Pancreatic Cancer Growth

After establishing the model, we sought to understand how a dysfunctional clock impacted cell growth.

KPC and KPC-BKO cells were grown in culture and injected into the right flanks of syngeneic C57BL/6J

mice. Tumor growth was measured every 3-4 days for a total of 28 days to understand differences in

primary tumor growth. We found that BKO caused an accelerated growth pattern, resulting in higher

mean (± standard error) weight tumors at the study conclusion compared to KPC-derived tumors (438.02

± 48.84 mg vs. 280.11 ± 42.73 mg; p = 0.02) (**Fig 6A-B**). These findings were evaluated and

independently confirmed in a second identically created *Bmal1* mutant (functional knockout) clone

(**Supplemental Fig 4A-C**). Perhaps more relevant to human PDAC – given the aggressiveness of this

cancer – we assessed the effect of clock disruption on survival by implanting KPC and KPC-BKO

heterotopic tumors and observing the mice until moribund status or lethality (**Fig 6C**). On Kaplan Meier

log-rank analysis, mice harboring tumors derived from KPC-BKO cells experienced worse survival than

mice bearing KPC-derived tumors (median survival: 52 versus 75 days, $p = 0.04$). Thus, clock

dysfunction resulted in accelerated tumor growth *in vivo*, leading to worse overall survival.

Loss of Bmal1 Promotes the Enrichment of Cell Growth Pathways

To determine the possible etiologies of clock disruption causing accelerated PDAC progression, we

compared the transcriptomic profiles of KPC and KPC-BKO cells. The principal component analysis

revealed clear transcriptional profile differences between each condition (**Fig 7A**). Differential gene

expression was quantified with edgeR and of the 15,110 genes, 5,235 (34.65%) were upregulated and

5,113 (33.84%) were downregulated in KPC-BKO compared to KPC cells (**Fig 7B, Supplemental Data**

File 2) (45). When we examined the CCGs known to control the circadian cycle, we found that 12/15

examined clock genes were differentially expressed, including *Per1*, *Per2*, *Per3*, *Cry1*, *Cry2*, *Nr1d1*,

Nr1d2, *Bhlhe40*, *Bhlhe41*, *Npas2*, *Arntl2*, and *Dbp* (All q < 0.05) (40). We then performed a KEGG

pathway analysis to examine for enrichment of pathways as a result of *Bmal1* knockout in the PDAC cells

(**Fig 7C**). We found that there was an enrichment of pathways important for cellular adhesion, such as

ECM-Receptor Interaction, Cell Adhesion, and Focal Adhesion, as well as several cellular growth

pathways including PI3K-AKT Signaling Pathway, MAPK Signaling Pathway, and Rap1 Signaling

Pathway. Collectively, these data indicate that clock disruption in the KPC cells results in significant core

clock gene changes and a transcriptional shift that alters key cancer promotion-related pathways such as

cellular attachment and proliferation.

Cell Survival is Increased with the Loss of Bmal1 through Alterations in Multiple Cell Death Pathways

A hallmark of cancer progression is the inhibition of apoptosis (46). This phenotype is readily apparent in

response to chemotherapy. In particular, the mechanism of action of gemcitabine (inhibition of DNA

synthesis) and paclitaxel (microtubule stabilization) – backbones in PDAC therapy – is to ultimately

induce apoptosis (47,48). To understand the potential implications of clock disruption for PDAC patients

undergoing treatment, we subjected KPC and KPC-BKO cells to 72 hours of chemotherapeutic treatment

with either gemcitabine or paclitaxel (49). We found that the activity of cell death pathways through

apoptosis (measured by Caspase 3/7 activity) was blunted in response to both chemotherapeutic agents as

a result of clock dysfunction (**Fig 8A-B**). We also assessed cytotoxic cell death, as measured by Dead-

Cell Protease activity, and found that clock disruption promoted resistance to cytotoxic cell death induced

by gemcitabine and paclitaxel (**Fig 8C-D**). These findings were also confirmed in a separately evaluated

independent *Bmal1*-mutant clone (**Supplemental Fig 4 D-G**). Prior work has implicated alteration of

Trp53 signaling to impact cell survival by suppressed apoptosis (20), but KPC cells (both WT and BKO)

are *Trp53* mutant (*Trp53^{R172H}*) indicating alternative mechanisms of heightened resistance in the KPC-

BKO cells. Gemcitabine resistance in PDAC often occurs due to the downregulation of the channel

protein hENT1 (*SLC29A1* gene), or deoxycytidine kinase (*DCK* gene) which activates gemcitabine once

inside the cell (50). However, these were only marginally downregulated (1.05-fold, $q = 0.04$ and 1.08-

fold, $q = 0.02$) due to *Bmal1* knockout, and thus unlikely to contribute to the differences seen with clock

disruption (**Fig 9A**). Furthermore, these were arrhythmic based on rhythmicity cut-off values ($p < 0.05$,

rAMP > 0.1, goodness of fit (rsq) > 0.1, and fitmean > 16) when examining our human samples,

suggesting a lack of circadian control (**Fig 9B, Supplemental Table 1**). Meanwhile, paclitaxel drug

resistance is thought to occur mostly through upregulation of drug efflux transporter proteins (P-

glycoprotein also known as multidrug-resistance associated protein), but these ATP-binding cassette

transporter proteins (*Abcb1a and Abcb1b* genes) were instead significantly *downregulated* (8.2-fold, $q <$

0.0001 and 2.1-fold, $q < 0.0001$) in KPC-BKO vs KPC cells (**Fig 9C**) (48). Concordantly, the *ABCB1*

gene was rhythmic with elevated expression in the human normal samples ($p = 0.015$, rAMP = 0.95,

fitmean = 133, rsq = 0.57) compared to the human PDAC samples that showed low levels of expression

(mean expression = 22) (**Fig 9D, Supplemental Table 1**). Thus, using this combined data analysis, we

found that the commonly described resistance mechanisms for gemcitabine and paclitaxel were not

identified, underscoring the complexity of clock disruption induced chemotherapeutic resistance in

cancer. Regardless, these data demonstrate that clock dysfunction promotes broad resistance in PDAC

including multiple cell death pathways in the setting of two different PDAC backbone agents.

<u>Discussion</u>

Our work herein used matched normal and tumor samples from TCGA and CPTAC-3 to provide the first

substantive evidence, to our knowledge, that the circadian clock is disrupted in PDAC while the adjacent

normal pancreatic clock is intact. Recently published work by Talamanca *et al.* also demonstrated an

intact circadian clock in human pancreas, albeit pancreas from human donors and not tumor-adjacent

pancreas (51). The authors leveraged transcriptional data from the GTEx consortium and used an

algorithm to project phases across tissues in postmortem human donors based on a donor internal phase.

They presumed that relative circadian phases are consistent among individuals to identify sex- and age-

dependent changes in circadian expression. Although the main metabolic organ of focus was the liver, in

the pancreas they identified a relatively strong concordance of rhythmically expressed genes between

males and females. However, when comparing pancreatic tissue from young versus old individuals, there

was a significant decrease in the percentage of conserved rhythmically expressed genes. Meanwhile,

Wucher et al also used the GTEx consortium data and time of death (as well as the season of death) to

determine differentially expressed genes between the day and night and across seasons (52). With this

method, they were able to identify differentially expressed genes between the day and night across

tissues, and they found that among circadian clock genes, *PER1-3* demonstrated the lowest logFC

(night/day) while *BMAL1* and *NPAS2* showed the highest logFC (night/day). Interestingly, *BMAL1* was

the single gene with a day-night pattern across the greatest tissue number (33 tissues). Also of note, the

authors identified a seasonal variation in gene expression with approximately 5% of the pancreatic

transcriptome demonstrating differential expression in at least one season when compared to the other

seasons. These results have implications when evaluating population-level data and indicate sex, age, and

potentially the time of year are important factors depending on the tissue assessed. To evaluate clock

functionality in the pancreas in our study, we examined the clock with three complementary analyses not

previously performed in pancreatic tissues: nCV, clock correlation, and CYCLOPS. The nCV was

pioneered and validated in several tissues by Wu and colleagues (25). It is a measurement that is directly

related to the relative amplitude (rAMP) of the oscillation and assesses overall and individual clock gene

robustness (25). For example, Wu and colleagues demonstrated that the nCV of core clock genes (CCGs)

was consistently and significantly diminished in clock-disrupted tissues versus wild-type tissues (*e.g*.,

Bmal1 knockout adipocytes vs wild-type adipocytes) and human datasets of tumor compared to matched

(adjacent) normal tissue, where the timing of sample collection was unspecified (25). It is therefore

extraordinarily beneficial for understanding the rAMP of the circadian clock genes – a key measure of

clock health – in population-level data where sample time acquisition is unknown. In conjunction with

nCV, Shilts *et al*. developed the method of clock correlation to determine the progression of the clock in

time-indeterminate datasets (22). By capitalizing on the concept of co-expression of CCGs (intrinsic to

the transcriptional translational feedback loop), Shilts and colleagues were able to computationally

discern clock progression in transcriptomic data from numerous human datasets. Importantly, this

technique is not dependent on full coverage of the period by samples, providing a beneficial approach for

our normal data set which included only 50 samples and did not have phase representation across the 24-

hour period. Furthermore, a direct comparison between heatmaps can be made (*i.e*., murine vs human or

tumor vs normal) because each heatmap has the same corresponding color to rho correlation value (22).

Thus, when combining nCV and clock correlation analysis in unordered human samples, the health of the

circadian clock in population-level data can be ascertained even when working with low sample numbers

(at least 30).

Wu and colleagues applied nCV analysis to several paired tumor-normal datasets (25), while Shilts *et al.*

examined clock correlation in normal tissues and paired tumor-normal datasets (22). However, we are the

first group, to our knowledge, to analyze the pancreas, possibly because our recent publication was the

first, to our knowledge, to characterize the diurnal expression of mouse pancreatic genes over 48 hours

(23). Thus, an appropriate mouse reference group had not been published for comparison. In the study by

Shilts *et al*., the human liver clock correlation heatmap (a similarly metabolic organ) demonstrated a

weaker clock correlation versus the mouse reference, which they attributed to increased noise in the liver

dataset (22). Therefore, in our human data, we expected to identify an apparent 'diminished health' of the

clock (lower nCV and weaker clock correlation) in the human pancreatic tissue versus mouse samples due

to significant variation in the normal samples. This variability is contributed by human factors, such as

type and timing of diet (53,54), underlying genetic differences, and alterations in the 'normal pancreas'

that surrounds the tumor (pancreatic atrophy, fibrosis, inflammation, etc.) (55). These can modulate the

relative amplitude of oscillation or CCG correlation which contributes to significant noise in the data, as

compared to the genetically identical, age-matched, and environmentally matched mouse samples (22,28).

Regardless of these limitations, the nCV and clock correlation revealed two key components of a healthy

clock in human pancreas, which was an important component of the pipeline.

CYCLOPS is a machine learning method developed by Anafi *et al* (28), with subsequent elegant studies

by Wu *et al* (26,27)., to demonstrate how clock gene relationships can be used to infer and reorder

samples by their predicted phase of expression to understand circadian biology in population-level data.

The authors who developed CYCLOPS recommend roughly 250 samples for a complete phase

distribution (28), which is dependent on differences in time of surgery (presume specimens acquired

between 6 AM and 6 PM) as well as differences in genetics and environmental factors (*e.g*., differences in

sleep-wake cycle, or shift worker status) (56). CYCLOPS has been applied to the lung, liver, brain,

hepatocellular carcinoma, and skin, but not the pancreas (27,28). This is possibly due to known

challenges associated with CYCLOPS, which includes optimizing the seed gene list for appropriate

ordering (28). We used the modified CYCLOPS approach by Wu *et al*. and were able to leverage our

murine pancreatic longitudinal expression data to generate the seed gene list for use with human data,

which increases the signal-to-noise ratio to optimize ordering capability (23,27,28). Thus, despite the

expected variability in human normal pancreas samples, CYCLOPS significantly ordered the samples

across the period, with several clock genes, including *CLOCK*, *PER1*, *PER3*, *NR1D2*, *RORC*, *NFIL3,*

T*EF*, *NR1D1, BHLHE40,* and *NPAS2* demonstrating rhythmic expression. While *BMAL1* and *CRY1* did

not pass our predetermined cutoff for rhythmicity in the normal samples, they demonstrated a robust

rAMP with near-significant p values likely reflecting limitations with our sample number (*i.e.*,

distribution of samples across the period) than true lack of rhythmicity in the normal samples. As

evidenced by the apparent gap in phase in the normal data, there was not uniform distribution of normal

samples across the 24-hour period, which signifies a limitation of our analysis. Incorporating time-

stamped normal samples (the hybrid design) can significantly increase the accuracy of CYCLOPS

ordering results, as suggested in Wu et al (26,27). Currently, we cannot implement a hybrid experimental

design in this study. However, despite these limitations, CYCLOPS statistically ordered the human

normal data, and the phase-set enrichment analysis and predicted phase of expression (of most CCGs)

aligned well with our prior mouse pancreas transcriptomic data (reference dataset). When combined with

nCV and clock correlation, the data clearly demonstrate an intact clock in the human normal pancreas

samples, which was an essential premise for evaluating the clock in human PDAC.

We then proceeded to use nCV, clock correlation, and CYCLOPS to provide compelling evidence of

circadian clock disruption in human PDAC. While the concept of circadian disruption in PDAC has been

posited by others, past studies examining global clock function in human PDAC have been limited by

detecting binary differences in overall RNA and protein expression between normal pancreas and PDAC

(17,18,20,21). Relles *et al.* found that several CCGs demonstrated decreased expression in PDAC

compared to benign tumors or normal pancreas (18), while Li *et al.* found that low *Bmal1* expression

(compared to 'higher expression') was associated with worse disease-free and overall survival in patients

with PDAC (17). Similar studies have been repeated with concordant findings (20,21). However, binary

comparisons of expression are unlikely to capture the complexities of clock health such as rhythmicity,

phase changes, changes in rAMP, or changes in clock progression. With the 318 available PDAC

samples, we showed that there was markedly diminished nCV and a much weaker correlation among

clock genes in PDAC compared to normal pancreas, depicting a loss of clock health in the cancer tissue.

Further, although the inability to reorder PDAC samples may be a limitation of CYCLOPS, it more likely

indicates clock dysfunction (57). In the PDAC samples, there was a sufficient sample number (n = 318)

for phase distribution, but the rAMP of nearly all CCGs was markedly decreased (as visualized through

best reordering), with many manifesting an arrhythmic pattern. Collectively, our approach to human data

(nCV, clock correlation, and CYCLOPS) provided convincing evidence of clock disruption in PDAC.

The main limitation of our current strategy is the inability to discern why there was a loss of circadian

signatures in human PDAC population data. While we have shown compelling evidence from the human

sample transcriptomic data that circadian disruption is present in PDAC, the population-level data doesn't

accurately represent individual patient tumors. The literature in cancer would suggest that cohorts of

patients exhibit a differential extent of clock disruption in their tumors (*i.e.*, some patient tumors have an

intact clock while others are markedly disrupted) (15,17,18,39). This is seen in other cancers. For

example, in human neuroblastoma, MYC amplification distorts circadian repressors (NR1D1) and

circadian activators (RORA and BMAL1) to cause metabolic rewiring that fuels cancer cell growth and

promotes poor patient outcomes (15,39). Consequently, clock function in neuroblastoma can be

subcategorized based on low/normal MYC expression (intact clock) versus MYC amplification (clock

disrupted). Similarly, in human PDAC, the overall loss of circadian signatures seen in the nCV, clock

correlation, and CYCLOPS analysis could be the consequence of i) dampening of the clock to a

significant extent in every patient (*i.e.*, global phenomena), ii) differences in the extent to which the clock

is disrupted (*i.e.*, clock disrupted vs intact clock), or iii) biological heterogeneity in the PDAC samples

surpassing circadian variability. The last consideration seems unlikely given the marked decrease in

rAMP (nCV) and clock correlation showing significantly diminished clock health in PDAC. Furthermore,

as seen with the CYCLOPS ordered *PER1* expression data over the 24-hr period (**Fig 3A**), enhanced

variability did not lead to the apparent arrhythmicity in PDAC. Unfortunately, we are currently unable to

discriminate between the first two possibilities and test the hypothesis that patient tumor subgroups

harbor differences in clock function. Ideally, we would be able to subcategorize the population of patients

instead of evaluating the entire cohort of PDAC patients, given the known molecular heterogeneity in

PDAC (58). For example, there are alterations in different gene programs (*e.g.*, *Kras*-mutant, *TGF-β*

signaling, G1/S checkpoint signaling, etc.) that determine the molecular subtypes in PDAC such as

squamous (including MYC pathway activation), pancreatic progenitor (including programs controlling

fatty acid metabolism), immunogenic, and ADEX (58). Therefore, separating the cohort into subtypes to

evaluate clock function could help elucidate whether circadian disruption in human PDAC co-segregates

with specific molecular subtypes or if there are alterations in specific gene programs that promote clock

disruption. However, the strength in our current evaluation derives from the number of patients included

in the analysis – and this is requisite for CYCLOPS analysis (minimum of ~200 patients needed for

optimal phase distribution and functioning of the algorithm) (28). Thus, despite batch correction

algorithms, the significant heterogeneity in transcriptomic data (*e.g.,* institutional differences in

experimental conditions, tumor purity, etc. (59–62)) results in challenges in interpreting differences in

clock function and drawing conclusions from smaller subsets of patients, particularly with algorithms like

CYCLOPS (28). Therefore, the focus of the current work was to establish that clock dysfunction is

present in human PDAC. In future work, we will build on this foundation and integrate whole genome

sequencing with RNA sequencing from our institutional samples, combined with uniform pathologic

evaluation and consistent sequencing conditions to minimize the variability/batch effect and optimize the

signal-to-noise ratio. This strategy will enable us to investigate whether specific molecular subtypes of

pancreas cancer co-segregate with clock dysfunction.

Our assessment of clock function in the KPC cells would also suggest that clock function is

heterogeneous in PDAC, given that the clock was intact in the *Kras-* and *Trp53*-mutant pancreas cancer

cells and necessitated *Bmal1* mutagenesis to generate clock disruption. Further, we assessed the cycling of

the *Per1* gene and *Per2*-luciferase in the KPC cells and identified a robust rhythm; by comparison, the

human PDAC samples demonstrated arrhythmic *PER1* expression. This discrepancy highlights the

challenge associated with the use of immortalized cancer cell lines for extrapolating to circadian clock

function and cancer-specific distortion of the clock in human patients. These *in vitro* and *in vivo* models

can be used to evaluate the impact of specific clock alterations, which can also yield information on

potential novel therapeutic approaches (*e.g.,* uncovering the mechanism through which *Bmal1* knockout

causes accelerated tumor growth could lead to therapeutic vulnerabilities). Yet, this type of model system

is uniform with respect to underlying PDAC-specific mutations (all lines in our study are derived from

the same *Kras* and *Trp53* mutations) and does not recapitulate the unique nature of human PDAC *in situ*,

where individual patient tumors exhibit mutational heterogeneity (58). Therefore, we did not have a

'representative model' in our immortalized studies of each human PDAC subtype to determine which

subtypes may harbor a disrupted clock. Unfortunately, there is currently no methodology to obtain

longitudinal data from each individual patient tumor; such an approach would enable an understanding of

the circadian clock at the individual level and would help to determine the circumstances under which

tumors manifest clock disruption, such as with specific gene program alterations or in the setting of

metastatic disease. As noted previously, the strategy of future work to uniformly evaluate subsets of

PDAC to determine clock disrupted versus clock intact subtypes can help to address this significant

discrepancy.

We generated the novel murine KPC-BKO cell line as a basis to evaluate clock disruption in PDAC.

Using our KPC and KPC-BKO cells, we found accelerated tumor growth in our syngeneic *in vivo* model

with *Bmal1* functional knockout. Our preference was to use an immune-competent model given the

known impact of the circadian clock on the immune system (another strength of our model) (63,64).

Interestingly, our primary tumor growth results were similar to results by Jiang and colleagues, who used

implanted human PDAC cells (BxPC-3 and PANC-1) into immunocompromised mice after shRNA

knockdown of *Bmal1* (20,21). However, patients with PDAC ultimately succumb to distant metastatic

spread, rather than local tumor growth, so examining the contribution of clock disruption to overall

survival was of greater importance (65). We found that *Bmal1* mutation promoted earlier lethality, which

has not been shown before. Concordant with the aggressive tumor phenotype, *Bmal1* mutation also

caused resistance to chemotherapy. While resistance to gemcitabine has been shown (21), we found

chemoresistance to two different backbone anti-cancer agents (gemcitabine and paclitaxel), including

suppressed apoptosis and cytotoxicity, indicating a more broad resistance to standard PDAC

chemotherapy. Although suppressed *Bmal1* expression in PDAC has been suggested to modulate *Trp53*

to promote a tumor suppressor effect, this was unlikely the case in our study considering KPC cells are a

Trp53-mutant cell line (20). Other work indicates the transcription factor YY1 is significantly

overexpressed in PDAC and ultimately causes BMAL1 suppression with consequent PDAC progression

and resistance to gemcitabine (unclear mechanism of resistance) (21). Yet, when we examined our human

PDAC and human normal samples, we found that *YY1* expression was equivalent between the groups

(mean expression 101.61 versus 102.87). Turning to other cancers where BMAL1 is suppressed, *MYC*

amplification in neuroblastoma alters *BMAL1* mRNA expression through the induction of *NR1D1*

(39,66). Notably, this is associated with poor prognosis and is *BMAL1* dependent since ectopic expression

of *BMAL1* inhibits tumor growth. Given the known interaction between MYC and the circadian clock,

and the known presence of MYC pathway activation in PDAC (squamous subtype), this will be a focus in

future work to evaluate the molecular subtypes of PDAC (15,39,67–69). However, this mechanism was

not apparent at the global population-level since we did not identify upregulation of *MYC*, *MYC-N,* or

NR1D1 gene expression in human PDAC compared to normal samples. While common resistance

mechanisms of gemcitabine and paclitaxel were not reflected in the human PDAC versus human normal

data or the KPC versus KPC-BKO cells (*e.g*., channel proteins), RNA sequencing identified several

enriched pathways integral to cancer progression and chemoresistance in the *Bmal1* functional knockout

cells, such as the PI3K-AKT pathway (70,71). The PI3K-AKT pathway is inextricably linked to cancer

cell proliferation and resistance to apoptosis, indicating a plausible mechanism for inhibition of

programmed cell death to multiple agents seen in our study (72,73). Further, resistance to paclitaxel is

associated with the activation of the PI3K-AKT pathway (74), and similar correlations have been

identified with gemcitabine resistance (75,76). However, the mechanism of chemoresistance is quite

complex and the etiology for suppressed apoptosis and cytotoxicity in the KPC-BKO line versus the KPC

line remains unclear. To add additional complexity (and highlight another limitation), it is currently

unresolved whether the effects we have seen with the loss of functional BMAL1 protein are due to clock

disruption or an independent role. Furthermore, clock-dependent effects can depend on which arm is

perturbed, the positive arm (circadian activators) or negative arm (circadian repressors) of the clock. For

example, in recent seminal work by Liu and colleagues, abolishing different arms of the circadian clock

resulted in reciprocal effects on the expression of *c-Myc* and downstream target genes (68), with the

knockout of *Bmal1* causing increased expression and *Cry1/2* knockout causing suppressed expression of

c-Myc and *c-Myc* regulated genes, respectively. Therefore, in future work, we can employ a series of

clock-manipulated PDAC cell lines (*e.g., Per1/2* double knockout to disrupt the negative arm) to better

understand the contributions of clock disruption and various components of the clock to chemoresistance

and PDAC progression.

In conclusion, we used a comprehensive approach (nCV, clock correlation, and CYCLOPS) to evaluate

the health of the circadian clock in human normal pancreas and demonstrated clock disruption in human

PDAC. Additionally, we developed novel cell lines to evaluate the repercussions of clock disruption in

PDAC and identified factors associated with poor prognosis (*i.e.*, worse survival, resistance to

chemotherapy, and enrichment of cancer-related pathways). While we acknowledge that significant work

needs to be done to improve our understanding of the circadian clock in PDAC, we can use the

foundation from this study as a basis for future work, where we will disentangle the clock-dependent

effects of PDAC and consequently focus therapeutic efforts.

Materials and Methods

Mouse Care

All animal studies were conducted according to an approved protocol (M005959) by the University of

Wisconsin School of Medicine and Public Health (UW SMPH) Institutional Animal Care and Use

Committee (IACUC). Male and female C57BL/6J mice were housed in an Assessment and Accreditation

of Laboratory Animal Care (AALAC) accredited selective pathogen-free facility (UW Medical Sciences

Center) on corncob bedding with chow diet (Mouse diet 9F 5020; PMI Nutrition International) and water

ad libitum. All *in vivo* experiments were performed with age-, cage-, and sex-matched mice under

identical lighting conditions. All experiments were performed with mice 6-8 weeks of age, and cages

were selected at random for each experiment.

Clock Correlation, nCV, and CYCLOPS Pipeline

We assessed the overall clock gene correlation and robustness of the clock with the clock correlation

matrix and normalized coefficient of variation (nCV) – nCV is known to be correlated with the relative

amplitude (rAMP) of oscillating clock genes, indicating the clock robustness as previously described

(22,25). The nCV was calculated for the overall condition with the nCVnet and nCVgene functions (25).

Clock correlation matrices were created using an available shiny app

(https://github.com/gangwug/CCMapp) which compares the correlation of clock components (17

individual clock genes) to a baseline correlation from the circadian atlas using the Mantel test

(22,26,27,31,32).

Cyclic ordering by periodic structure (CYCLOPS) (28) was validated for use in the pancreas utilizing our

existing murine normal circadian and chronic jetlag pancreas RNA sequencing (RNA-seq) data (Gene

Expression Omnibus (GEO) Accession number: GSE165198) (23). Specifically, the seed genes for use in

CYCLOPS were selected by cross-referencing genes found to be rhythmically expressed in our dataset

with those genes either rhythmically expressed in the liver (similarly metabolic organ) or those used by

Wu *et al.* when validating CYCLOPS in the skin (**Supplemental Data File 3**) (23,26,31,32). The updated

CYCLOPS pipeline by Wu *et al*. (https://github.com/gangwug/Oslops) was then used to reorder our

murine pancreas datasets with known sample collection times (26). Eigengenes were selected with the

Oscope package to sharpen CYCLOPS (77). Clusters with a $p < 0.05$ (also known as $Stat^{err}$) and Met^{smooth}

< 1 were considered to be significantly reordered (28). Rhythmicity of the reordered genes was

determined on cosinor analysis with a $p < 0.01$, rAMP > 0.1, goodness of fit (rsq) > 0.1, and fitmean > 16

(26). Significantly rhythmic gene phase was then compared to the rhythmic gene phase detected from the

known sample time collection using the meta3d function of Metacyle (42). Clock genes were highlighted

to demonstrate a correlation between the predicted and actual phase.

The clock was evaluated in human normal and human PDAC RNA-seq datasets from TCGA and

CPTAC-3 (29,30). After batch correction with ComBat and filtering, 50 matched normal and 318 PDAC

samples were obtained for analysis (78). The pipeline described above was then used to obtain the clock

correlation matrix, nCV, and CYCLOPS reordering. Cosinor analysis was performed to test for

rhythmicity. Given the additional biologic heterogeneity of the human data, a $p < 0.05$, rAMP > 0.1,

goodness of fit (rsq) > 0.1, and fitmean > 16 were used as a rhythmicity cutoff. Rhythmic genes from

normal samples were assessed with phase set enrichment analysis (PSEA) (79). Rhythmic gene sets

ordered by significance were inputted with their calculated phase of expression. Default settings were

used for PSEA, including domain 0-24, min item 10, max sims/test 10,000. The gene set enrichment

analysis (GSEA) gene ontology (GO) (c5.go.bp.v7.5.1.symbols) set was leveraged as the pathway input.

The top 15 significant pathways ($q < 0.05$) were selected for representation.

KPC Cell Line Creation and Maintenance

Pancreas cancer cells that harbor $Kras^{G12D}$ and $Trp53^{R172H}$ mutations (KPC cells) were acquired from

Ximbio (Catalog: 153474; Westfield Stratford City, UK). Cells were cultured in DMEM supplemented

with 10% Fetal Bovine Serum (Cytiva, Marlborough, MA), 1% L-glutamate-Penicillin-Streptomycin

(Gibco, ThermoFisher Scientific, Waltham, MA), and 1% non-essential amino acids (Gibco) at 37°C at

5.0% CO_2 to their appropriate confluence for use. CRISPR/Cas9 technology was used to introduce *Bmal1*

mutations into the well-established KPC cells, generating the *Bmal1* functional knockout line (KPC-

BKO) (36). Synthetic tracrRNA and target-specific crRNAs (crRNA:tracrRNA (ctRNAs)) were annealed

as per manufacturer instructions (Integrated DNA Technologies (IDT), Coralville, IA).

Ribonucleoproteins were formed with Cas9 protein (V3, Catalog: 1081059, IDT) and ctRNAs

individually (ctRNA 1:AATATGCAGAACACCAAGGA, ctRNA 2: TTAGAATATGCAGAACACCA)

(IDT). Nucleofection was used to introduce RNPs (1.95 µM Cas9, 2 µM ctRNA, 2 µM electroporation

enhancer (IDT)) into 2 x 10^5 KPC cells on a Nucleofector 4D (Lonza Biosciences, Walkersville, MD)

with an SF kit (Catalog: V4CX-2032, Lonza Biosciences) as per manufacturer instructions using pulse

code CM-120. After 48 hours of recovery in DMEM growth media (described above) at 37°C at 5.0%

CO_2, single cells were deposited in 96 well plates using a BD FACSAria III (BD Biosciences, Franklin

Lakes, NJ). Outgrowing clones were condensed to a 96 well plate in duplicate to propagate clones and

generate a genomic DNA source. Genomic DNA was harvested from KPC cells, nucleofected with

ctRNA-1 RNPs and ctRNA-2 RNPs, and the targeted region of *Bmal1* was PCR amplified using primers:

Forward: acactctttccctacacgacgctcttccgatct NNNNNN CCAAGAATCCTTGTGTGTCTG and Reverse:

gtgactggagttcagacgtgtgctcttccgatct AGAGGACTCCACAGACATGAAC (IDT). PCR products were

dual-indexed with indexing PCR, pooled, sequenced on an Illumina MiSeq instrument (San Diego, CA),

and analyzed with CRISPResso2 (80).

Creation of Per2-dLuciferase reporter KPC cell line

KPC cells were stably transfected with a mammalian gene expression vector harboring a destabilized

luciferase reporter driven by the *Per2* promoter fused to intron 2 of *Per2* and a puromycin resistance

cassette (VectorBuilder, Santa Clara, CA) (81). The vector was transfected with lipofectamine 2000

(ThermoFisher Scientific) and incubated for 2 days, and then exposed to media containing 2.5 µg/mL

puromycin for 3 days, and surviving clones were selected. Luciferase activity was measured in the

selected clones with the luciferase assay system (Promega, Madison, WI) on a BMG CLARIOstar (BMG

Labtech, Ortenberg, Germany) plate reader. The two selected clones were then subcloned using a BD

FACSAria III after staining with DAPI. Luciferase activity was again measured in the subclones to

validate expression before use in downstream experiments.

Clock Function Testing

To evaluate for clock function, KPC and KPC-BKO cells were synchronized with 200 nM

dexamethasone for 2 hours in FBS-free DMEM media, followed by RNA isolation 24 hours after the

synchronization using the RNeasy protocol (Qiagen, Hilden, Germany) according to the manufacturer's

recommendations (82). Samples (n = 3 biologic replicates) were collected at 4-hour intervals for 24 hours.

Quantitative real-time polymerase chain reaction (RT-qPCR) was performed (in technical triplicate) for

the downstream core clock gene (CCG) *Per1* (ID: Mm00501813_m1, Life Technologies, Carlsbad, CA)

and the housekeeper gene *Hprt* (ID: Mm03024075_m1) using GoTaq Probe qPCR and RT-qPCR System

(Promega, Madison, WI) and Quantstudio 7 flex RT-PCR system (ThermoFisher Scientific). Expression

was measured and the ΔCT was calculated. The mean ΔCT values were then tested for rhythmicity using

the meta2d function of Metacycle (42). For Metacycle settings, the min period and max period were set to

24, and "JTK", "LS", and "ARS" were selected for cycMethod. An integrated FDR corrected q value <

0.05 and rAMP > 0.1 were taken as rhythmic as previously described (26). To separately evaluate clock

function in KPC cells, luciferase activity driven by the *Per2* promoter was measured. Cells from two

independent clones were synchronized with 200 nM dexamethasone for 2 hours in FBS-free DMEM

media, and 24 hours later luciferase activity was measured at 4-hour intervals (n = 6 technical replicates

per time point). To measure activity, 1x cell culture lysis reagent (Promega) was added and cells were

incubated for 5 minutes. Firefly luciferase assay reagent (Promega) was added and luminescence was

measured on a BMG CLARIOstar plate reader and tested for rhythmicity using Metacycle.

Western Blotting

Western blotting was performed to determine BMAL1 expression in two independent KPC cell lines.

After synchronization, protein samples were isolated at 6-hour intervals for 24 hours using CelLytic M

lysis reagent (MilliporeSigma, Burlington, MA) and Halt Protease and Phosphatase Inhibitor Cocktail

(ThermoFisher Scientific). A total of 30 μg of each sample was loaded onto a Mini-PROTEAN TGX

7.5% precast mini-gel (Bio-RAD Laboratories, Hercules, CA). The gel was then transferred using the

semi-dry transfer technique to an Immobilon-P PVDF membrane (MilliporeSigma). The membrane was

then blocked with 5% skim milk and incubated with 1 μg/μL rabbit BMAL1 antibody (Catalog: NB100-

2288, RRID: AB_10000794, Novus Biologicals, Littleton, CO) and 1:2000 rabbit β-ACTIN antibody

(Catalog: 4967, RRID: AB_330288, Cell Signaling Technologies, Danvers, MA). Finally, the membranes

were incubated with alkaline phosphatase-conjugated goat anti-rabbit IgG (1:10000) (Catalog: 111-055-

144, RRID: AB_2337953, Jackson ImmunoResearch West Grove, PA) and stained using 1-Step

NBT/BCIP solution (ThermoFisher).

RNA Isolation, Sequencing, Differential Gene Expression Analysis

To evaluate for transcriptomic differences between wild-type and BKO KPC cells, bulk RNA-seq was

performed on 6 independent samples collected from each condition. Isolation was carried out as above

and quality was tested for an RNA integrity number (RIN) > 7.5 on the Agilent 2100 bioanalyzer (Agilent

Technologies, Santa Clara, CA). A total of 300 ng of mRNA was enriched with poly-A selection and

sequencing on the Illumina HiSeq2500 platform by the University of Wisconsin Biotechnology

Sequencing Core. FASTq files were processed with Skewer and genes were filtered to remove those with

low expression (83). Samples were normalized by the method of trimmed mean of M-values (84).

Contrasts were drawn with the edgeR package, with differential expression taken when the FDR $q < 0.05$

(45). Pathway testing was performed with the KEGG database (Kyoto Encyclopedia of Genes and

Genomes) using previously described methods (85). The top 500 significant genes were inputted, ordered

by q value, and the top 9 significant pathways (where $p < 0.05$) were plotted for visualization. Pathway

dot size is indicative of the number of genes in each pathway. The RNA-seq data is publicly available

through GEO (Accession number: GSE213680).

Heterotopic Tumor Modeling

To create flank tumors, 1×10^5 KPC or KPC-BKO cells were injected into the right flanks of

immunocompetent C57BL/6J mice obtained from Jackson Laboratory (RRID: IMSR_JAX:000664, Bar

Harbor, ME). Cells were mixed in a 1:1 50 μL solution of DMEM media and Matrigel (Corning Inc,

Corning, NY). In the tumor growth experiment, a single dose of KPC or KPC-BKO cells was injected

into C57BL/6J mice (male: n = 5, female: n = 5, each group) and tumors were measured twice weekly for

four weeks starting on day 6 with the caliper method as previously described (86). Tumor length and

width were measured and tumor volume was calculated using the formula: tumor volume = length x

width^2 x ½. Tumor weight was also measured (in mg) at the conclusion of the study period. Two

independent replicate experiments were performed for tumors derived from KPC cells and KPC-BKO

cells (total n = 20 mice per group). Additionally, two separate KPC-BKO clones were tested. Our power

analysis, based on prior literature, indicated that using 10 mice per group would have 80% power to

detect a 1.5-fold difference (alpha = 0.05) in tumor size (21 days after heterotopic injection) between

KPC and KPC-BKO cells (87). Mean differences between groups were computed using a t-test. In the

tumor metastasis/survival experiment, the same dose of KPC or KPC-BKO cells was injected into

C57BL/6J mice (n = 7 in each group) and tumors were measured weekly until the mice became moribund

or died. Kaplan Meier log-rank analysis was then performed to compare survival differences between

conditions with the survival package (88). A $p < 0.05$ is taken as significant for both experiments.

Cell cycle analysis

To evaluate the cell cycle of KPC and KPC-BKO cells, cells were grown to confluence, fixed with 70%

ethanol, and stained with 50 μg/mL propidium iodide (Catalog: 421301, BioLegend, San Diego, CA) and

100 μg/mL RNAse A (Catalog: FEREN0531, Thermo Fisher). Samples (n = 3 technical replicates, each

condition) were sorted with an Attune NxT flow cytometer (Thermo Fisher) and the DNA content was

analyzed for the percent of cells in G1, S, and G2 phase with ModFit LT 6.0 software at the University of

Wisconsin Carbone Cancer Center Flow Cytometry Laboratory. The average coefficient of variation (CV)

of each sample was < 6%. Differences between conditions were tested with a t-test and a $p < 0.05$ was

taken as statistically significant. This experiment was performed independently twice.

Chemotherapeutic Resistance Evaluation

To evaluate for chemotherapeutic resistance, an equal number of KPC or KPC-BKO cells (3000 each)

were grown in 96-well plates for 24 hours. Cells were washed and media containing either vehicle

709 (DMSO, Catalog: D2650-100ML, MilliporeSigma) or increasing concentrations of gemcitabine (dose

710 range = 10 nM to 1μM) (Catalog: G6423-50MG, MilliporeSigma) or paclitaxel (dose range = 10 nM to

711 1μM) (Catalog: AAJ62734MC, Thermo Fisher) was added to each well. Apoptosis was measured by

712 detecting Caspase 3/7 activity after 24 hours of treatment using the Caspase-Glo 3/7 Assay System

713 (Promega) on a BMG CLARIOstar plate reader. Dead-cell protease activity was similarly measured in

714 cells after 20 hours of treatment using the CytoTox-Glo Cytotoxicity Assay (Promega) on a BMG

715 CLARIOstar plate reader. Mean differences in the fold change between chemotherapy- and vehicle-

716 treated cells were made with a Student's t-test and a $p < 0.05$ was taken as statistically significant. Each

717 experiment was completed with technical triplicates and independently replicated twice. A representative

718 result is shown for each experiment.

719 *Analysis*

720 All analyses were performed in R version 4.2.0 or Julia version 0.3.12 unless otherwise indicated.

721 Samples were not excluded from the analysis unless explicitly stated. Standard comparisons of the mean

722 were performed with a t-test and survival was assessed with the log-rank test unless explicitly stated.

723 **Acknowledgments**

724 We would like to acknowledge the Gene Expression Center and the Bioinformatics Resource Center

725 (BRC) at the University of Wisconsin, Madison for their contributions to this work. The authors thank the

726 University of Wisconsin Carbone Cancer Center Flow Cytometry Laboratory, supported by P30

727 CA014520, for use of its facilities and services. We would also like to thank the Michael W. Oglesby

728 Foundation for their funding support of our work in circadian disruption and pancreas pathology.

729 **Materials Availability Statement**

730 The modified KPC cell lines (KPC-BKO and KPC Per2-dLuc) created for use in this manuscript are

731 available upon reasonable request from the corresponding author, Sean Ronnekleiv-Kelly.

Data Availability Statement

The TCGA and CPTAC-3 data were requested from the following URL: https://portal.gdc.cancer.gov/.

All RNAseq data can be found on the Gene Expression Omnibus (GEO) at accession numbers

GSE213680 and GSE165198.

References

1.   Bass J, Lazar MA. Circadian time signatures of fitness and disease. Science. 2016 25;354(6315):994–

9.

2.   Aryal RP, Kwak PB, Tamayo AG, Gebert M, Chiu PL, Walz T, et al. Macromolecular Assemblies of

the Mammalian Circadian Clock. Mol Cell. 2017 Sep 7;67(5):770-782.e6.

3.   Transcriptional architecture of the mammalian circadian clock - PubMed [Internet]. [cited 2022 Oct

10]. Available from: https://pubmed.ncbi.nlm.nih.gov/27990019/

4.   Allada R, Bass J. Circadian Mechanisms in Medicine. N Engl J Med. 2021 Feb 11;384(6):550–61.

5.   Fishbein AB, Knutson KL, Zee PC. Circadian disruption and human health. J Clin Invest. 2021 Oct

1;131(19):e148286.

6.   West AC, Smith L, Ray DW, Loudon ASI, Brown TM, Bechtold DA. Misalignment with the external

light environment drives metabolic and cardiac dysfunction. Nat Commun. 2017 12;8(1):417.

7.   Marcheva B, Ramsey KM, Buhr ED, Kobayashi Y, Su H, Ko CH, et al. Disruption of the clock

components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes. Nature. 2010 Jul

29;466(7306):627–31.

8.   Roenneberg T, Allebrandt KV, Merrow M, Vetter C. Social jetlag and obesity. Curr Biol CB. 2012

May 22;22(10):939–43.

9.   Parsons MJ, Moffitt TE, Gregory AM, Goldman-Mellor S, Nolan PM, Poulton R, et al. Social jetlag,

obesity and metabolic disorder: investigation in a cohort study. Int J Obes 2005. 2015

May;39(5):842–8.

10. Knutson KL, Wu D, Patel SR, Loredo JS, Redline S, Cai J, et al. Association Between Sleep Timing,

Obesity, Diabetes: The Hispanic Community Health Study/Study of Latinos (HCHS/SOL) Cohort

Study. Sleep. 2017 Apr 1;40(4).

11. Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, et al. Obesity and metabolic

syndrome in circadian Clock mutant mice. Science. 2005 May 13;308(5724):1043–5.

12. Dong Z, Zhang G, Qu M, Gimple RC, Wu Q, Qiu Z, et al. Targeting Glioblastoma Stem Cells

through Disruption of the Circadian Clock. Cancer Discov. 2019 Nov 1;9(11):1556–73.

13. Papagiannakopoulos T, Bauer MR, Davidson SM, Heimann M, Subbaraj L, Bhutkar A, et al.

Circadian Rhythm Disruption Promotes Lung Tumorigenesis. Cell Metab. 2016 09;24(2):324–31.

14. Kettner NM, Voicu H, Finegold MJ, Coarfa C, Sreekumar A, Putluri N, et al. Circadian Homeostasis

of Liver Metabolism Suppresses Hepatocarcinogenesis. Cancer Cell. 2016 Dec 12;30(6):909–24.

15. Moreno-Smith M, Milazzo G, Tao L, Fekry B, Zhu B, Mohammad MA, et al. Restoration of the

molecular clock is tumor suppressive in neuroblastoma. Nat Commun. 2021 Jun 28;12(1):4006.

16. Sulli G, Rommel A, Wang X, Kolar MJ, Puca F, Saghatelian A, et al. Pharmacological activation of

REV-ERBs is lethal in cancer and oncogene induced senescence. Nature. 2018 Jan

18;553(7688):351–5.

17. Li W, Liu L, Liu D, Jin S, Yang Y, Tang W, et al. Decreased circadian component Bmal1 predicts

tumor progression and poor prognosis in human pancreatic ductal adenocarcinoma. Biochem Biophys

Res Commun. 2016 Mar 25;472(1):156–62.

18. Relles D, Sendecki J, Chipitsyna G, Hyslop T, Yeo CJ, Arafat HA. Circadian gene expression and

clinicopathologic correlates in pancreatic cancer. J Gastrointest Surg Off J Soc Surg Aliment Tract.

2013 Mar;17(3):443–50.

19. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA Cancer J Clin. 2022

Jan;72(1):7–33.

20. Jiang W, Zhao S, Jiang X, Zhang E, Hu G, Hu B, et al. The circadian clock gene Bmal1 acts as a

potential anti-oncogene in pancreatic cancer by activating the p53 tumor suppressor pathway. Cancer

Lett. 2016 Feb 28;371(2):314–25.

21. Jiang W, Zhao S, Shen J, Guo L, Sun Y, Zhu Y, et al. The MiR-135b-BMAL1-YY1 loop disturbs

pancreatic clockwork to promote tumourigenesis and chemoresistance. Cell Death Dis. 2018 Feb

2;9(2):149.

22. Shilts J, Chen G, Hughey JJ. Evidence for widespread dysregulation of circadian clock progression in

human cancer. PeerJ. 2018;6:e4327.

23. Schwartz PB, Walcheck MT, Berres ME, Nakaya M, Wu G, Carrillo ND, et al. Chronic Jetlag-

Induced Alterations in Pancreatic Diurnal Gene Expression. Physiol Genomics. 2021 May 31;

24. Hughes ME, Abruzzi KC, Allada R, Anafi R, Arpat AB, Asher G, et al. Guidelines for Genome-Scale

Analysis of Biological Rhythms. J Biol Rhythms. 2017 Oct;32(5):380–93.

25. Wu G, Francey LJ, Ruben MD, Hogenesch JB. Normalized coefficient of variation (nCV): a method

to evaluate circadian clock robustness in population scale data. Bioinformatics. 2021 Dec

1;37(23):4581–3.

26. Wu G, Ruben MD, Francey LJ, Smith DF, Sherrill JD, Oblong JE, et al. A population-based gene

expression signature of molecular clock phase from a single epidermal sample. Genome Med. 2020

Aug 21;12(1):73.

27. Wu G, Ruben MD, Schmidt RE, Francey LJ, Smith DF, Anafi RC, et al. Population-level rhythms in

human skin with implications for circadian medicine. Proc Natl Acad Sci U S A. 2018 Nov

27;115(48):12313–8.

28. Anafi RC, Francey LJ, Hogenesch JB, Kim J. CYCLOPS reveals human transcriptional rhythms in

health and disease. Proc Natl Acad Sci U S A. 2017 May 16;114(20):5312–7.

29. Edwards NJ, Oberti M, Thangudu RR, Cai S, McGarvey PB, Jacob S, et al. The CPTAC Data Portal:

A Resource for Cancer Proteomics Research. J Proteome Res. 2015 Jun 5;14(6):2707–13.

30. Cancer Genome Atlas Research Network, Weinstein JN, Collisson EA, Mills GB, Shaw KRM,

Ozenberger BA, et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet. 2013

Oct;45(10):1113–20.

31. Pizarro A, Hayer K, Lahens NF, Hogenesch JB. CircaDB: a database of mammalian circadian gene

expression profiles. Nucleic Acids Res. 2013 Jan;41(Database issue):D1009-1013.

32. Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB. A circadian gene expression atlas in

mammals: Implications for biology and medicine. Proc Natl Acad Sci. 2014 Nov 11;111(45):16219–

24.

33. Fu L, Kettner NM. The circadian clock in cancer development and therapy. Prog Mol Biol Transl Sci.

2013;119:221–82.

34. Pogue-Geile KL, Lyons-Weiler J, Whitcomb DC. Molecular overlap of fly circadian rhythms and

human pancreatic cancer. Cancer Lett. 2006 Nov 8;243(1):55–7.

35. Schwartz PB, Barrett-Wilt GA, Ronnekleiv-Kelly SM. Chronic Jetlag Alters the Landscape of the
Pancreatic Lipidome. Pancreas. 2022 Jan 1;51(1):80–9.

36. Hingorani SR, Wang L, Multani AS, Combs C, Deramaudt TB, Hruban RH, et al. Trp53R172H and
KrasG12D cooperate to promote chromosomal instability and widely metastatic pancreatic ductal
adenocarcinoma in mice. Cancer Cell. 2005 May;7(5):469–83.

37. Bunger MK, Wilsbacher LD, Moran SM, Clendenin C, Radcliffe LA, Hogenesch JB, et al. Mop3 is
an essential component of the master circadian pacemaker in mammals. Cell. 2000 Dec
22;103(7):1009–17.

38. Ness-Cohn E, Allada R, Braun R. Comment on "Circadian rhythms in the absence of the clock gene
Bmal1." Science. 2021 Apr 16;372(6539):eabe9230.

39. Altman BJ, Hsieh AL, Sengupta A, Krishnanaiah SY, Stine ZE, Walton ZE, et al. MYC Disrupts the
Circadian Clock and Metabolism in Cancer Cells. Cell Metab. 2015 Dec 1;22(6):1009–19.

40. McIntosh BE, Hogenesch JB, Bradfield CA. Mammalian Per-Arnt-Sim proteins in environmental
adaptation. Annu Rev Physiol. 2010;72:625–45.

41. Vazquez-Rivera E, Rojas B, Parrott JC, Shen AL, Xing Y, Carney PR, et al. The aryl hydrocarbon
receptor as a model PAS sensor. Toxicol Rep. 2022;9:1–11.

42. Wu G, Anafi RC, Hughes ME, Kornacker K, Hogenesch JB. MetaCycle: an integrated R package to
evaluate periodicity in large scale data. Bioinforma Oxf Engl. 2016 01;32(21):3351–3.

43. Bieler J, Cannavo R, Gustafson K, Gobet C, Gatfield D, Naef F. Robust synchronization of coupled
circadian and cell cycle oscillators in single mammalian cells. Mol Syst Biol. 2014 Jul 15;10(7):739.

44. Feillet C, Krusche P, Tamanini F, Janssens RC, Downey MJ, Martin P, et al. Phase locking and

multiple oscillating attractors for the coupled mammalian clock and cell cycle. Proc Natl Acad Sci U

S A. 2014 Jul 8;111(27):9828–33.

45. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression

analysis of digital gene expression data. Bioinforma Oxf Engl. 2010 Jan 1;26(1):139–40.

46. Evan GI, Vousden KH. Proliferation, cell cycle and apoptosis in cancer. Nature. 2001 May

17;411(6835):342–8.

47. Mini E, Nobili S, Caciagli B, Landini I, Mazzei T. Cellular pharmacology of gemcitabine. Ann Oncol

Off J Eur Soc Med Oncol. 2006 May;17 Suppl 5:v7-12.

48. Kampan NC, Madondo MT, McNally OM, Quinn M, Plebanski M. Paclitaxel and Its Evolving Role

in the Management of Ovarian Cancer. BioMed Res Int. 2015;2015:413076.

49. Goldstein D, El-Maraghi RH, Hammel P, Heinemann V, Kunzmann V, Sastre J, et al. nab-Paclitaxel

plus gemcitabine for metastatic pancreatic cancer: long-term survival from a phase III trial. J Natl

Cancer Inst. 2015 Feb;107(2).

50. de Sousa Cavalcante L, Monteiro G. Gemcitabine: metabolism and molecular mechanisms of action,

sensitivity and chemoresistance in pancreatic cancer. Eur J Pharmacol. 2014 Oct 15;741:8–16.

51. Talamanca L, Gobet C, Naef F. Sex-dimorphic and age-dependent organization of 24-hour gene

expression rhythms in humans. Science. 2023 Feb 3;379(6631):478–83.

52. Wucher V, Sodaei R, Amador R, Irimia M, Guigó R. Day-night and seasonal variation of human

gene expression across tissues. PLoS Biol. 2023 Feb;21(2):e3001986.

53. Brown MR, Matveyenko AV. It's What and When You Eat: An Overview of Transcriptional and

Epigenetic Responses to Dietary Perturbations in Pancreatic Islets. Front Endocrinol.

2022;13:842603.

54. Fadista J, Vikman P, Laakso EO, Mollet IG, Esguerra JL, Taneera J, et al. Global genomic and

transcriptomic analysis of human pancreatic islets reveals novel genes influencing glucose

metabolism. Proc Natl Acad Sci U S A. 2014 Sep 23;111(38):13924–9.

55. Korpela T, Ristimäki A, Udd M, Vuorela T, Mustonen H, Haglund C, et al. Pancreatic fibrosis, acinar

atrophy and chronic inflammation in surgical specimens associated with survival in patients with

resectable pancreatic ductal adenocarcinoma. BMC Cancer. 2022 Jan 3;22(1):23.

56. Archer SN, Laing EE, Möller-Levet CS, van der Veen DR, Bucca G, Lazar AS, et al. Mistimed sleep

disrupts circadian regulation of the human transcriptome. Proc Natl Acad Sci U S A. 2014 Feb

11;111(6):E682–91.

57. Wu G, Ruben MD, Francey LJ, Lee YY, Anafi RC, Hogenesch JB. An in silico genome-wide screen

for circadian clock strength in human samples [Internet]. bioRxiv; 2022 [cited 2022 Sep 8]. p.

2022.05.10.491250. Available from: https://www.biorxiv.org/content/10.1101/2022.05.10.491250v2

58. Bailey P, Chang DK, Nones K, Johns AL, Patch AM, Gingras MC, et al. Genomic analyses identify

molecular subtypes of pancreatic cancer. Nature. 2016 Mar 3;531(7592):47–52.

59. Molania R, Foroutan M, Gagnon-Bartsch JA, Gandolfo LC, Jain A, Sinha A, et al. Removing

unwanted variation from large-scale RNA sequencing data with PRPS. Nat Biotechnol. 2023

Jan;41(1):82–95.

60. Peixoto L, Risso D, Poplawski SG, Wimmer ME, Speed TP, Wood MA, et al. How data analysis

affects power, reproducibility and biological insight of RNA-seq studies in complex datasets. Nucleic

Acids Res. 2015 Sep 18;43(16):7664–74.

61. Zhang C, Cheng W, Ren X, Wang Z, Liu X, Li G, et al. Tumor Purity as an Underlying Key Factor in

Glioma. Clin Cancer Res Off J Am Assoc Cancer Res. 2017 Oct 15;23(20):6279–91.

62. Vieth B, Parekh S, Ziegenhain C, Enard W, Hellmann I. A systematic evaluation of single cell RNA-

seq analysis pipelines. Nat Commun. 2019 Oct 11;10(1):4667.

63. Yu X, Rollins D, Ruhn KA, Stubblefield JJ, Green CB, Kashiwada M, et al. TH17 cell differentiation

is regulated by the circadian clock. Science. 2013 Nov 8;342(6159):727–30.

64. Druzd D, Matveeva O, Ince L, Harrison U, He W, Schmal C, et al. Lymphocyte Circadian Clocks

Control Lymph Node Trafficking and Adaptive Immune Responses. Immunity. 2017 Jan

17;46(1):120–32.

65. Conroy T, Hammel P, Hebbar M, Ben Abdelghani M, Wei AC, Raoul JL, et al. FOLFIRINOX or

Gemcitabine as Adjuvant Therapy for Pancreatic Cancer. N Engl J Med. 2018 20;379(25):2395–406.

66. Altman BJ, Hsieh AL, Gouw AM, Dang CV. Correspondence: Oncogenic MYC persistently

upregulates the molecular clock component REV-ERBα. Nat Commun. 2017 Mar 23;8:14862.

67. Shostak A, Ruppert B, Ha N, Bruns P, Toprak UH, ICGC MMML-Seq Project, et al. MYC/MIZ1-

dependent gene repression inversely coordinates the circadian clock with cell cycle and proliferation.

Nat Commun. 2016 Jun 24;7:11807.

68. Liu Z, Selby CP, Yang Y, Lindsey-Boltz LA, Cao X, Eynullazada K, et al. Circadian regulation of c-

MYC in mice. Proc Natl Acad Sci U S A. 2020 Sep 1;117(35):21609–17.

69. Maddipati R, Norgard RJ, Baslan T, Rathi KS, Zhang A, Saeid A, et al. MYC Levels Regulate

Metastatic Heterogeneity in Pancreatic Adenocarcinoma. Cancer Discov. 2022 Feb;12(2):542–61.

70. Murthy D, Attri KS, Singh PK. Phosphoinositide 3-Kinase Signaling Pathway in Pancreatic Ductal

Adenocarcinoma Progression, Pathogenesis, and Therapeutics. Front Physiol. 2018;9:335.

71. Bryant KL, Stalnecker CA, Zeitouni D, Klomp JE, Peng S, Tikunov AP, et al. Combination of ERK

and autophagy inhibition as a treatment approach for pancreatic cancer. Nat Med. 2019

Apr;25(4):628–40.

72. Hoxhaj G, Manning BD. The PI3K-AKT network at the interface of oncogenic signalling and cancer

metabolism. Nat Rev Cancer. 2020 Feb;20(2):74–88.

73. Mortazavi M, Moosavi F, Martini M, Giovannetti E, Firuzi O. Prospects of targeting

PI3K/AKT/mTOR pathway in pancreatic cancer. Crit Rev Oncol Hematol. 2022 Aug;176:103749.

74. Maloney SM, Hoover CA, Morejon-Lasso LV, Prosperi JR. Mechanisms of Taxane Resistance.

Cancers. 2020 Nov 10;12(11):E3323.

75. Yang XL, Lin FJ, Guo YJ, Shao ZM, Ou ZL. Gemcitabine resistance in breast cancer cells regulated

by PI3K/AKT-mediated cellular proliferation exerts negative feedback via the MEK/MAPK and

mTOR pathways. OncoTargets Ther. 2014;7:1033–42.

76. Li W, Zhu Y, Zhang K, Yu X, Lin H, Wu W, et al. PROM2 promotes gemcitabine chemoresistance

via activating the Akt signaling pathway in pancreatic cancer. Exp Mol Med. 2020 Mar;52(3):409–

22.

77. Leng N, Chu LF, Barry C, Li Y, Choi J, Li X, et al. Oscope identifies oscillatory genes in

unsynchronized single cell RNA-seq experiments. Nat Methods. 2015 Oct;12(10):947–50.

78. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing batch effects
and other unwanted variation in high-throughput experiments. Bioinforma Oxf Engl. 2012 Mar
15;28(6):882–3.

79. Zhang R, Podtelezhnikov AA, Hogenesch JB, Anafi RC. Discovering Biology in Periodic Data
through Phase Set Enrichment Analysis (PSEA). J Biol Rhythms. 2016;31(3):244–57.

80. Clement K, Rees H, Canver MC, Gehrke JM, Farouni R, Hsu JY, et al. CRISPResso2 provides
accurate and rapid genome editing sequence analysis. Nat Biotechnol. 2019 Mar;37(3):224–6.

81. Mei L, Fan Y, Lv X, Welsh DK, Zhan C, Zhang EE. Long-term in vivo recording of circadian
rhythms in brains of freely moving mice. Proc Natl Acad Sci U S A. 2018 Apr 17;115(16):4276–81.

82. Ramanathan C, Khan SK, Kathale ND, Xu H, Liu AC. Monitoring Cell-autonomous Circadian Clock
Rhythms of Gene Expression Using Luciferase Bioluminescence Reporters. J Vis Exp JoVE. 2012
Sep 27;(67):4234.

83. Jiang H, Lei R, Ding SW, Zhu S. Skewer: a fast and accurate adapter trimmer for next-generation
sequencing paired-end reads. BMC Bioinformatics. 2014 Jun 12;15:182.

84. Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of
RNA-seq data. Genome Biol. 2010 Mar 2;11(3):R25.

85. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters | OMICS: A
Journal of Integrative Biology [Internet]. [cited 2022 Aug 28]. Available from:
https://www.liebertpub.com/doi/10.1089/omi.2011.0118

86. Euhus DM, Hudd C, Laregina MC, Johnson FE. Tumor measurement in the nude mouse. J Surg
Oncol. 1986;31(4):229–34.

87. Alexander RK, Liou YH, Knudsen NH, Starost KA, Xu C, Hyde AL, et al. Bmal1 integrates

mitochondrial metabolism and macrophage activation. eLife. 2020 May 12;9:e54090.

88. Therneau T. A Package for Survival Analysis in R [Internet]. Available from: https://CRAN.R-

project.org/package=survival

<u>Figures</u>

Fig 1: *Bioinformatics pipeline for circadian function*. Recently, there have been three separate

bioinformatics tools that have substantially improved the ability to determine circadian clock function in

population-level transcriptional data (22,25–28). **<u>Normalized coefficient of variation (nCV)</u>**: Clock gene

expression produces robust oscillations with the amplitude of the oscillation defined by the difference

between peak and trough, and relative amplitude (rAMP) determined by the ratio between amplitude and

baseline level of expression (<u>upper left</u>). Clock disruption (such as with clock gene knockout) causes

suppressed rAMP, and thus rAMP can be considered a strong determinant of clock function (<u>lower left</u>).

Because rAMP is calculated from time-labeled samples, it can only be measured in human population

data when sample time acquisition is known (*e.g.,* MetaCycle) or when using programs to predict

longitudinal time-course data (*e.g.,* CYCLOPS). To overcome this limitation and assess core clock gene

'robustness' of oscillation in human data (without labeled time), normalized coefficient of variation

(nCV) is used. The nCV is the coefficient of variation (CV) of the core clock gene (in the population data)

divided by the mean CV of all genes in the dataset. And the CV is determined by the standard deviation

of the gene expression divided by its average expression. Thus, when the circadian clock is intact,

circadian genes in large population data sets would be expected to have a relatively high nCV due to

oscillating gene expression (<u>upper middle</u>). Conversely, with a disrupted clock, the variation of the core

clock genes would be diminished (oscillation is suppressed), resulting in a relatively low nCV (<u>lower

middle</u>). Importantly, the nCV was found – across multiple cancer types and tissues – to correspond

directly to the normalized rAMP (<u>upper right</u>) (25); it can thus be used confidently as a surrogate for

rAMP, which is a key determinant of clock health. Of note, the rAMP (and thus nCV) of core clock genes

differs due to differences in amplitude, and so the between tissue difference (*e.g.,* normal vs cancer) for

each core clock gene is the value of importance. **<u>Clock Correlation</u>**: When the circadian clock is intact,

there is an expected progression of core clock gene expression, where the positive arm of the clock (*e.g.*,

BMAL1, *CLOCK*) drives transcription of the negative arm of the clock (*e.g., PER*, *CRY*). Thus, when

BMAL1 expression peaks, *CLOCK* expression should also be near its peak, and this should be evident in

population-level data (*i.e.,* can look at the spread of individual expression points across the population). In

turn, when *BMAL1* expression peaks, negative arm members (e.g., *PER1-2*) should be at a trough.

Concordantly, strong positive correlations ($0.5 < \rho < 1$, red) should be apparent among transcriptional

activators (e.g., *BMAL1* and *CLOCK*) and among transcriptional repressors (e.g., *NR1D1* and *PER2*), and

a strong negative correlation ($-0.5 > \rho > -1$, blue) should be present amongst activators and repressor

targets (e.g., *BMAL1* and *PER2*). If this is the case, then the clock is intact; if these correlations are not

preserved (*i.e.,* $\rho \sim 0$), this indicates the clock is disrupted. Using the set of core clock genes and clock-

associated genes, the correlation of each gene against the others is determined by Spearman's rho (ρ) and

mapped in matrix form. This experimental matrix is then compared to a baseline correlation matrix from

the mouse circadian gene atlas using the Mantel test, which compares the correlation between the two

matrices to produce a z-statistic (z-stat). A higher z-stat value corresponds to a correlation matrix that is

closer to the circadian gene atlas baseline (*i.e.,* highly preserved clock correlation). **<u>CYCLOPS:</u>** Intrinsic

to circadian clock genes is rhythmic expression. CYCLOPS (cyclic ordering by periodic structure) is an

algorithm that can identify rhythmic (longitudinal) data from population data where sample time

acquisition is unknown. The basis for this type of program is that each sample has a different 'clock time'

due to differences in time of sample acquisition and environmental factors such as sleep/wake cycle or

shift worker status (*e.g.,* 15-20% of patients are night shift workers). Seed genes that are known to be

rhythmic in the tissue of interest (*i.e.,* pancreas) are inputted into CYCLOPS, and the population level

data is reduced to two vectors (Eigengene 1 and Eigengene 2) derived from the seed genes. When plotted,

the optimal Eigengene pair will demonstrate an ellipse, which indicates that the two Eigengenes are

rhythmic and anti-phasic – this pair can then be used for each patient sample to determine the sample

'time', and thus the order of that sample relative to the 24-hr period (phase). By incorporating enough

patients, population-level data can be transformed into longitudinal data. Subsequently, with the patient

sample dataset ordered by CYCLOPS (by the Eigengene pair), individual genes (*e.g.,* oscillating genes or

core clock genes) can be evaluated for rhythmicity. An intact circadian clock is indicated by the ability of

CYCLOPS to order the data (statistically) and the identification of rhythmically expressed core clock

genes. Meanwhile, a disrupted clock is designated as arrhythmic core clock gene expression or an

inability of CYCLOPS to order the data (statistically). Statistically significant rhythmic gene expression

is determined by $p < 0.05$, rAMP > 0.1, fitmean > 16 and goodness of fit (rsq) > 0.1. The fitmean value

can be conceptualized as a mean level of expression of that gene across the dataset, and therefore a

minimum level of expression (fitmean > 16) is required for rhythmicity cutoff. The goodness of fit of the

experimental values by cosinor regression is calculated by R squared (rsq). CYCLOPS reordering is

assessed by the Met^{smooth} and $Stat^{err}$ (significant reordering: $Met^{smooth} < 1$ and $Stat^{err} < 0.05$). The Met^{smooth}

compares the smoothness of the reconstructed circular trajectory versus a linear ordering based on the

first principal component, while the $Stat^{err}$ (we designate as *p*-value) is analogous to the F statistic of a

typical nested regression and compares the model fit when a circular rather than linear bottleneck node is

used. A full mathematical derivation has been previously outlined by Anafi *et al.* (28) Figure created with

BioRender.com.

1008

Fig 2: *The human pancreatic clock robustness is diminished after malignant transformation.* **A**. Clock

correlation matrices were created for normal circadian (left, n = 72) and chronic jetlag (right, n = 72)

mouse pancreas samples. **B, C**. The normalized coefficient of variation (nCV) was calculated to

determine the robustness of the pancreatic clock for the normal circadian (circles) and chronic jetlag

(triangles) pancreas samples, with the mean nCV in each group also depicted. Each color indicates a

different gene evaluated. **D.** Clock correlation matrices were created for normal (left; n = 50) and

pancreatic ductal adenocarcinoma (right; n = 318) samples. **E, F**. The normalized coefficient of variation

(nCV) was calculated to determine the robustness of the pancreatic clock for the normal (circles) and

PDAC (triangles) samples, with the mean nCV in each group also depicted. Each color indicates a

different gene evaluated.

Fig 3: *Clock phase predictions demonstrate conserved clock relationships*: **A.** Graphical representation

and **B.** table of the CCG and clock-controlled gene phase prediction from the cosinor analysis of

CYCLOPS reordered human pancreas samples (0 to 2π; n = 50) and Metacycle predicted phases of mouse

pancreas samples (0 to 24; n = 72), relative to *BMAL1* (human) and *Bmal1* (mouse) phase.

Fig 4: *The human pancreatic cancer tumor clock is dysfunctional relative to the normal pancreas.* **A**.

CYCLOPS was used to reorder samples from normal (n = 50) and pancreatic ductal adenocarcinoma (n =

318) TCGA and CPTAC-3 samples. Each point represents an individual patient with order determined by

paired eigengenes. Plots from several core clock genes reordered by CYCLOPS in normal tissue as

compared to best reordering in PDAC (**A**) are shown – ordered from 0 to 2π. Shading around the blue

regression line indicates the 95% confidence interval. Relative amplitude (rAMP) for each gene is

determined by dividing the amplitude by the fitted expression baseline (fitmean). The fitmean value can

be conceptualized as a mean level of expression of that gene across the dataset, and therefore a minimum

level of expression (fitmean > 16) is required for rhythmicity cutoff. The goodness of fit of the

experimental values by cosinor regression was calculated by R^2(rsq). Using these collective parameters, a

$p < 0.05$, rAMP > 0.1, goodness of fit (rsq) > 0.1, and fitmean > 16 were considered rhythmic (27). **B**.

Graphical representation of Phase Set Enrichment Analysis (PSEA) of normal pancreatic samples,

ordered by phase of expression.

Fig 5: *Creation of a mouse clock dysfunction cell line model.* **A**. The core clock gene *Bmal1* was mutated

(functional knockout) in KPC murine pancreatic cells with CRISPR-Cas9 genome editing technology

(BKO) with the frameshift site indicated by the blue line. **B.** The frameshift mutation was induced

upstream of the basic helix loop helix (bHLH) domain (red star). The resulting protein lacked critical

downstream elements for functionality, including the PAS-A, PAS-B, and PAC domains. **C.** Western blot

demonstrating the loss of BMAL1 protein across 24 hours for wild-type (WT) and BKO cells (M =

marker lane). **D.** The mean (± standard error) *Per1* mRNA expression of synchronized KPC (n = 3 at each

time point; black dashed) and KPC-BKO (n = 3 at each time point; grey) cells from ZT0-ZT24. **E.** Graphs

comparing the percent of cells in G1, S, and G2 in WT (n = 3; white) and BKO (n = 3; grey) KPC cells.

Fig 6: *Mutation of Bmal1 promotes pancreatic cancer progression.* **A.** Comparison of mouse tumor

weight at 28 days for WT (n = 20; white) and BKO (n = 20) KPC tumors. **B.** WT (n = 20; black) and

BKO (n = 20; grey) mean (± standard error) tumor size (in mm^3) over the 28-day growth period. **C.**

Kaplan Meier survival curve for mice implanted with WT (n = 7; black) and BKO (n =7; grey) KPC

tumors. The dotted lines indicate the median survival. [ns = not significant, * = p < 0.05, ** = p < 0.01]

Fig 7: *Functional knockout of Bmal1 promotes widespread transcriptomic alterations and activation of*

multiple oncogenic pathways. **A.** Principal Component Analysis (PCA) demonstrating overall differences

in expression between WT (green) and BKO (purple) samples (n = 6 each) **B.** Volcano plot showing

genes that were significantly upregulated (red) and downregulated (blue) on differential gene expression

(DGE) analysis (q < 0.05). All significant core clock genes are highlighted **C.** KEGG analysis was then

performed, and the top 9 pathways ordered by significance are shown. For each pathway, the size of each

dot corresponds to the number of genes involved in each pathway.

Fig 8: *Clock dysfunction diminishes the chemotherapeutic cell death response.* KPC wildtype (WT) and

KPC-BKO cells (n = 3) were treated with increasing doses of gemcitabine and paclitaxel. Fold change

differences (± standard error) in Caspase 3/7 activity in response to either **A.** gemcitabine or **B.** paclitaxel.

Fold change differences (± standard error) in dead-cell protease activity in response to **C.** gemcitabine or

D. paclitaxel. Comparisons between conditions at each concentration were made with t-test. [ns = not

significant, * = *p* < 0.05, ** = *p* < 0.01, *** = *p* < 0.001]

Fig 9: *Evaluating for mechanisms of chemotherapy resistance due to clock dysfunction.* Gemcitabine

resistance mechanism: **A**. Fold change in expression between KPC-BKO and KPC for channel protein

hENT1 (*SLC29A1* gene) and deoxycytidine kinase (*DCK* gene) is shown, with negative values

representing downregulation in the KPC-BKO (clock disrupted) cells. **B.** CYCLOPS reordered samples

(0 to 2π) from normal (n = 50) and pancreatic ductal adenocarcinoma (n = 318) is depicted for each of the

respective genes from (**A**), demonstrating arrhythmic gene expression in human normal and pancreas

cancer tissue. Shading around the blue regression line indicates the 95% confidence interval. Paclitaxel

resistance mechanism: **C.** Fold change in expression between KPC-BKO and KPC for ATP-binding

cassette drug efflux transporters (*Abcb1a and Abcb1b* genes) are shown, with negative values

representing downregulation in the KPC-BKO (clock disrupted) cells. **D.** CYCLOPS reordered samples

(0 to 2π) depict the human *ABCB1* gene expression in normal (rhythmic) and PDAC (arrhythmic and

suppressed) tissue. Shading around the blue regression line indicates the 95% confidence interval. *p* <

0.05, rAMP > 0.1, goodness of fit (rsq) > 0.1, and fitmean > 16 were considered rhythmic for CYCLOPS

analysis (27).

Supplementary Files

Supplementary Figures:

Supplemental Fig 1: *CYCLOPS accurately reorders wildtype mouse pancreas samples.* Eigengenes

identified by Oscope for use in CYCLOPS in **A.** normal circadian (upper) and chronic jetlag (lower)

samples. Shading around the blue regression line indicates the 95% confidence interval for each plot. **B.**

The normal circadian (left) and chronic jetlag (right) clusters ordered by CYCLOPS demonstrate accurate

reordering for both conditions **C.** Normal circadian (left) and chronic jetlag (right) genes found to be

significantly rhythmic on both CYCLOPS reordered cosinor analysis and rhythmicity testing based on the

known sample collection time with the Metacycle meta3d function are ordered by their predicted phase of

expression. Clock genes are shown in orange.

Supplemental Fig 2: *Normal pancreatic and pancreatic ductal adenocarcinoma (PDAC) samples were*

processed and filtered. TCGA and CPTAC-3 samples were processed and batched corrected. An MDS

plot is shown demonstrating differences between the matched normal (green; n = 50) and PDAC (purple;

n = 318) samples.

Supplemental Fig 3: *KPC cells expressing dLuciferase driven by the Per2 promoter exhibit circadian*

activity. Two independent clones were examined, and luciferase activity was measured at 4-hour intervals

(n = 6 per time point). Rhythmicity was calculated with Metacycle. Clone 1 (left) was found to be

rhythmic with a q = 5.51E-5 and rAMP = 0.44. Clone 2 (right) was found to be rhythmic with a q =

1.95E-7, rAMP = 0.39.

Supplemental Fig 4: *Cross-validation of mutant Bmal1 (functional knockout) growth in a second*

independent clone. A second independent *Bmal1* knockout (BKO) clone with an identical mutation to the

first was validated with **A.** western blot analysis. We then heterotopically implanted wildtype (WT) and

BKO cells into C57BL/6J mice and followed **B.** tumor weight and **C.** cell growth. BKO_2 (dark grey; n =

20) had a similar mean (± standard error) tumor weight to BKO_1 (light grey; n = 20) (426.09 (± 40.07)

mg vs 438.02 (± 48.84) mg; p = 0.85), and both BKO_2 (426.09 (± 40.07) mg vs 280.11 (± 42.73) mg; p

= 0.017) and BKO_1 (438.02 (± 48.84) mg vs 280.11 (± 42.73) mg; p = 0.02) were larger than WT

(white; n = 20). BKO_1 and BKO_2 had similar significantly faster growth trajectories. **D-G.** KPC

wildtype (WT) and BKO_2 cells (n = 3) were treated with increasing doses of gemcitabine and paclitaxel.

Fold change differences (± standard error) in Caspase 3/7 activity in response to either **D.** gemcitabine or

E. paclitaxel. Fold change differences (± standard error) in dead-cell protease activity in response to **F.**

gemcitabine or **G.** paclitaxel. Comparisons between conditions at each concentration were made with t-

test. [ns = not significant, * = p < 0.05, ** = p < 0.01, *** = p < 0.001]

Supplementary Data Files:

Supplemental Data File 1: Significant genes on cosinor analysis for CYCLOPS reordered normal

pancreas

Supplemental Data File 2: KPC wildtype versus BKO edgeR differential gene expression

Supplemental Data File 3: CYCLOPS seed gene list.

Supplemental Data File 4: Data for Reconstruction of Figures

Supplementary Tables

Supplemental Table 1: Cosinor rhythmicity analysis of CYCLOPS ordered normal pancreas and

PDAC

A.

Mouse

Human

B.

Mouse Gene	Phase	Human Gene	Phase
Bmal1	0	BMAL1	0
Clock	23	CLOCK	1.3
Cry1	17.9	CRY1	0.4
Cry2	15	CRY2	3.6
Per1	13	PER1	3.1
Per2	15	PER2	3
Per3	12.4	PER3	3.1
Nr1d1	9	NR1D1	3.1
Nr1d2	11	NR1D2	3.2
Rora	9	RORA	4.9
Rorc	19	RORC	3.1
Npas2	0.3	NPAS2	2.8
Nfil3	21	NFIL3	6.2
Bhlhe41	13	BHLHE41	5
Dbp	9.9	DBP	2.9
Ciart	12.1	CIART	2.6
Tef	12.3	TEF	3.1
Hlf	11	HLF	4.9

A. Frameshift Mutation

Bmal1 (116,662 bp)

B.
NLS aa36-41
bHLH aa79-132
PAS-A aa150-222
PAS-B aa333-403
PAC aa408-451
Heterodimerization
DNA Binding

C.

BMAL1
ACTB

D. *Per1*

WT [q = 0.0004; rAMP = 0.11]
BKO [q = 0.81; rAMP = 0.003]

E.