Title: Examining the Reproductive Success of Bull Kelp (*Nereocystis luetkeana*) in Climate Change Conditions

List of Authors:

Korabik, Angela R.* Department of Environmental Science and Policy, University of California, Davis, Davis, CA, USA 95616

Winquist, Tallulah. Department of Biology, San Diego State University, San Diego, CA, USA 95616

Grosholz, Edwin D. Department of Environmental Science and Policy, University of California, Davis, CA USA 95616

Hollarsmith, Jordan A. Alaska Fisheries Science Center, Juneau, AK, USA 99801

Contact Information: * arkorabik@ucdavis.edu

Keywords: kelp forests, climate change, marine heat waves, ocean acidification, bull kelp, reproduction

Abstract:

Climate change is affecting marine ecosystems in many ways including rising temperatures and ocean acidification. From 2014-2016, an extensive marine heatwave extended along the west coast of North America and had devastating effects on numerous species during this period, including bull kelp (*Nereocystis luetkeana*). Bull kelp is an important foundation species in coastal ecosystems that can be affected by marine heat waves and ocean acidification, however these impacts have not been investigated on sensitive early life stages. To determine the effects of changing temperatures and carbonate levels on Northern California's bull kelp populations, we collected sporophylls from mature bull kelp individuals in Point Arena, CA. At the Bodega Marine Laboratory, we released spores from field-collected bull kelp, and cultured microscopic

gametophytes in a common garden experiment with a fully factorial design crossing modern conditions (11.63±0.54°C and pH 7.93±0.26) with observed extreme climate conditions (15.56±0.83°C and 7.64±0.32pH). Our results found that both increased temperature and decreased pH influenced growth-and egg production of bull kelp microscopic stages. Increased temperature generally resulted in decreased gametophyte survival and offspring production. In contrast, decreased pH had less of an effect, but generally resulted in increased gametophyte survival and offspring production. Additionally, we found that increased temperature significantly impacted reproductive timing by causing female gametophytes to produce offspring earlier than under ambient temperature conditions. Our findings inform better predictions of the impacts of climate change on coastal ecosystems as well as provide key insight into environmental dynamics regulating the bull kelp lifecycle.

Introduction:

Globally, marine systems are under pervasive threats from climate change. Chief among these threats are marine heat waves and ocean acidification (OA) (Cooley et al. 2022). Changing temperature and OA have negative impacts on the critical structure-forming foundational species of the world's oceans, namely kelps and corals, especially in terms of reduced reproduction (Straub et al. 2019, Smith et al. 2022) and juvenile mortality (Kroeker et al. 2013, Harvey et al. 2013, Przeslawksi et al. 2014). In the ocean, early life stages are already subject to high mortality rates due to a number of environmental bottlenecks, and increased temperature and decreased pH can further increase juvenile mortality through reduced recruitment and growth of microscopic stage canopy kelps (Gaitan-Espitia et al. 2014, Shukla and Edwards 2017, Hollarsmith et al. 2020, Lind and Konar 2017), reduced calcification and increased disease in juvenile invertebrates (Kroeker et al. 2013, Ban et al. 2013, Small et al. 2016, Miner et al. 2018), and altered larval fish behavior (Munday 2010, Ferrari 2011).

Kelp forests are critical to temperate, nearshore subtidal and intertidal marine systems worldwide, and sustain numerous economically important recreational and commercial fisheries (Bennett et al. 2016, Blamey and Bolton 2018, Carr and Reed 2016). In addition, kelp forests provide numerous ecosystem functions and services such as shelter of structural habitat and food sources to surrounding ecosystems, buffering coastlines from wave energy, ameliorating the effects of ocean acidification, reduction of current speeds and larval delivery to the shore, and modification of seawater chemistry (Hamilton et al. 2022, Malone et al. 2022).

Globally, the effects of marine heatwaves are already having extreme effects on kelp forests (Arafeh-Dalmau et al. 2019, Straub et al. 2019, Filbee-Dexter et al. 2020). From 2014-2017, Northern California lost 90% of its bull kelp (*Nereocystis luetkeana*) kelp canopy cover

over an area of roughly 350 km (Rogers-Bennett and Catton 2019). This loss of kelp forest cover has been attributed to a dramatic increase in purple urchin (*Strongylocentrotus purpuratus*) density due to loss of keystone predators, coupled with a pervasive system of marine heatwaves (McPhearson et al. 2021). The results of such widespread canopy loss were drastic changes in community structure and composition (Beas-Luna et al. 2020) and the collapse of the several fisheries in the area, such as that of the red sea urchin (*Mesocentrotus franciscanus*) (Rogers-Bennett and Okamoto 2020) and the closure of the world's largest recreational abalone fishery (*Haliotis rufescens*) (Rogers-Bennett and Catton 2019, Reid et al. 2016).

Numerous studies in recent years have documented the effects of increased temperature on bull kelp canopies (Rogers-Bennett and Catton 2019, Hamilton et al. 2020, Helen Berry's work in PS?), and have found that decreases in adult bull kelp canopy abundance have been related to local and large scale processes associated with warm water (Schiel et al. 2004, Pfister et al. 2017). Bull kelp exposure to warm temperatures also reduces adult blade morphological plasticity to changes in hydrodynamic flow regimes (Suprataya et al. 2020), but the physiological impacts of warm waters on bull kelp need to be further studied. Studies of bull kelp microscopic developmental stages in British Columbia and Alaska have found that increased temperatures have resulted in reductions in settlement and reduced germination and growth (Schiltroth 2021, Lind and Konar 2017, Muth et al. 2019), but the impact of rising temperatures on microscopic bull kelp stages in the southern portion of their range in northern California remains unclear. California bull kelp populations represent the range extreme of bull kelp, existing in low-latitude areas that are the most exposed to ENSO warm water events compared to more northern populations. As a result, California bull kelp populations could either be more warm-water adapted than the higher latitude populations previously studied, or they could be existing much closer to the thermal maxima and therefore very vulnerable. As the bulk of kelp die offs during the 2014 to

2017 marine heatwave occurred near the lower-latitude portion of kelp species' ranges (Arafeh-Dalmau et al. 2019, Beas-Luna et al. 2020, Cavanaugh et al. 2019, Finger et al. 2021, Rogers-Bennett and Catton 2019), it is necessary to further study how future marine heatwaves may affect the ability of these foundation species to remain in their lower latitude ranges.

In addition to the increasing threat of marine heat waves, coastal temperate ecosystems are also subject to stress from ocean acidification (OA), which, on average, has already caused a global lowering of surface water pH by 0.11 pH units (Feely et al., 2004, Feely et al., 2009, Gattuso et al., 2015a). Variability of pH levels in nearshore systems is normal to a degree as seasonal oceanographic shifts like upwelling bring deep offshore waters to the surface and expose nearshore ecosystems to reduced pH levels. This exposure varies with local bathymetry and coastal topography, which often changes the intensity of upwelling events along the coast (Feely et al. 2008). While pH variation in the California Current System generally stays between 7.720 and 8.413 (Feely et al. 2018), climate change projections predict an increasing frequency and duration of low-pH extremes (Bakun et al. 2015, García-Reyes et al. 2015), which may result in an average decrease of up to 0.4 pH units (Feely et al. 2008). Low pH may impact physiological functions among a variety of organisms. Studies have shown that OA will more proportionately impact organisms that form calcium carbonate skeletons (Kroeker et al. 2013), but it is critical that we also understand how the compounding stress of these combined threats will impact our critical temperate nearshore systems.

Kelps are very efficient at processing multiple carbon species in the water column. Kelps utilize CO_2 via photosynthesis, and also have the ability to concentrate calcium carbonate into their tissues. There is some evidence to suggest that the excess of carbon predicted for future ocean conditions may increase kelp growth in climate change conditions. For example, increased

pCO$_2$ has been shown to have beneficial impacts on mature bull kelp net apparent productivity (Thom 1996) and growth (Swanson and Fox 2007). At the microscopic stage, however, the effects of pCO2 and pH on kelp can be variable, ranging from having no effect (Fernández et al. 2015, Hollarsmith et al 2020) to positive effects on growth and photosynthesis.

Understanding how different life stages respond to environmental stress is critical when trying to predict population resilience to disturbance events. Laminariales, or the large canopy forming kelps, have a multistage process of development that presents numerous areas for the imposition of bottlenecks from climate stress. However, to our knowledge, no studies have yet investigated the role that pH may play in juvenile bull kelp development, nor the combined threats of increased temperature and ocean acidification on any bull kelp life stage.

In this study, we ask how increased temperatures and decreasing pH will impact: 1) bull kelp gametophyte development, 2) egg and sporeling production, and 3) juvenile growth. Based on the observed negative effects of the 2014-17 marine heatwave on bull kelp adults, we hypothesized that increased temperature will generally result in decreased growth, survival and reproduction. In contrast, we hypothesized that decreased pH has less of an effect than temperature on growth and egg production, but generally resulted in increased growth, survival and reproduction.

Materials and Methods:

Bull Kelp Life Cycle

In California, one of the most dominant canopy forming kelp species is bull kelp (*Nereocystis luetkeana*). The range of bull kelp extends from the eastern Aleutian Islands, Alaska in the north to Point Conception, CA in the south. Within its California range, it is considered to be the dominant canopy-forming kelp species in Northern California, between San Francisco and

the California-Oregon border. Bull kelp is an annual species, and is thought to be a more opportunistic, resilient colonizer, especially in areas too turbulent for the persistence of giant kelp (*Macrocystis pyrifera*) (Foster and Schiel 1985, Graham 1997, Graham et al. 2007).

Bull kelp have a heteromorphic life cycle consisting of a large diploid sporophyte, and a microscopic haploid gametophyte. Adult sporophytes develop sporophyll patches on their blades at the ocean surface, and at maturity, begin to release spores. The released zoospores then settle on hard substrate at the benthos, where they grow into microscopic male and female gametophytes. The female gametophytes begin to develop eggs, and then release a pheromone to trigger sperm release from nearby males. Once the sperm fertilizes the egg, a new sporophyte begins to develop (Reed 1990).

Collection

Sporophylls from approximately 10 individuals were collected at the surface by boat from a single kelp bed in Point Arena, California (38.916271°N, 123.725644°W) in October 2017. Sporophylls were cleaned in iodine and freshwater, layered in a cooler with wet paper towels separating individual sporophylls, and transported to the Bodega Marine Laboratory (BML, 38.318164°N, 123.072019°W) for sporulation. Spore densities were determined using a hemocytometer (model number CTL-HEMM-GLDR, LW Scientific, Lawrenceville, U.S.A.), and were introduced into the experimental petri dishes at densities of approximately 8 spores/mm^2.

Ex-situ culturing experiment

We conducted a fully factorial common garden experiment that consisted of four treatments representing ambient and high temperature and ambient and low pH, with ten replicates per treatment, for a total of forty experimental petri dishes (Fisher Brand 100mm

x15mm). Temperature was maintained at 15.6±0.8°C and 11.6±0.5°C using incubators at BML, and pH was maintained at 7.93±0.26 pH and 7.64±0.32 pH using chemical additions of equal parts 1M HCl and 1M NaHCO3 (NaHCO3 + HCl → NaCl + H2CO3) (Riebesell et al., 2011). Temperatures were chosen to represent ambient sea surface temperatures for our ambient temperature treatment, whereas our high temperature treatment represented the 4°C increase in SST observed during the 2014-17 marine heatwave (Gentemann et al. 2017). Ambient and low pH were chosen to represent the pH of incoming seawater at BML and pH during an extreme upwelling event (Feely et al., 2008), respectively. Light was set at 14:10 photoperiod and 30-45 umol m^{-2} s^{-1}. The pH of incoming, manipulated, and outgoing seawater was measured to 0.01 pH units immediately after collection using a spectrophotometer. Total alkalinity (T$_{alk}$, µmol kg^{-1}) was measured using potentiometric acid titration. Water was changed in all experimental dishes every 2 to 3 days for the duration of the 27 day experiment. We added standard 20 mL L^{-1} Provasoli nutrient mix to all treatment water (Provasoli 1968).

Photo Analysis

Beginning one week after spore inoculation, petri dishes were photographed weekly with a Micropublisher 5.0 RTV digital camera (QImaging, Surrey, Canada) mounted on an inverted microscope at 40x magnification, resulting in 4 weeks of photos documenting population growth and reproduction. Within each dish, 3 points were randomly selected to be photographed, with different points being photographed each week. Each photo encompassed 1.08mm^2 of the petri dish (15,708mm^2 bottom surface area).

After the growth experiment was completed, each photo was analyzed using ImageJ (Rasband 2019). Week 1 and 2 photos did not contain any gametophytes large enough to identify by sex, so only Weeks 3 and 4 were used for analysis. Count data was obtained from each photo

for female gametophytes, male gametophytes, eggs, and juvenile sporelings (Figure 1). Females

were also categorized as "productive" (having produced at least one egg or juvenile) or "non-

productive" (have no eggs or juveniles). The proportion of productive females in each photo was

calculated by dividing the number of productive females by the total number of females counted.

Count data was also used to calculate three additional variables: average number of eggs

per female, average number of juveniles per female, and average number of offspring per female

((# eggs + # juveniles)/# females). These three ratios were then used to approximate fertilization

and maturity rates between Weeks 3 and 4. Juvenile sizes were also obtained by using the

freehand trace tool and measuring the number of pixels encapsulated. Sizes were then converted

to $\lceil m^2$ using a conversion factor of 71330 pixels per 62,500$\lceil m^2$, which was calculated by

measuring the area of a photo of a 0.0625mm^2 hemocytometer cell at 40x magnification.

Statistical Analysis

All count outcome variables were analyzed using linear mixed models with temperature,

pH, and their interaction as fixed effects and Dish ID as a random effect. In order to meet the

parametric assumptions of normality of residuals and homogeneity of variances, all count data

was subjected to a square-root transformation as needed before being analyzed. We tested the

significance of our fixed effects by conducting log-likelihood tests via model comparison using

the anova function, where one model included the effect of interest while the other model

excluded it.

We analyzed the proportion of productive females using a generalized linear mixed

model with a beta distribution. Size data was also analyzed with a generalized linear mixed

model using a gamma distribution. GLMMs included temperature, pH, and their interactions as

fixed effects, and petri dish ID as a random effect. Average number of gametophytes per photo in

a given dish was also calculated and included in the size model as a covariate to account for possible density dependence. We also separately analyzed the relationship between average size of juveniles per photo and the covariate (average number of gametophytes per photo) using a linear regression model that included only the covariate as a fixed effect.

All count and size data were only analyzed for Week 4 of our experiment, but calculated ratios of eggs per female (eggs/fem), juveniles per female (juvs/fem), and offspring per female (offspring/fem) were analyzed for both Weeks 3 and 4 in order to draw conclusions about differences in rates of fertilization or maturation. Specifically, we used the ratio of offspring/fem to ask whether females, regardless of treatment, were making the same effort to reproduce, and the ratios of eggs/fem and juvs/fem were calculated to inform us about which stage reproduction was within each treatment. We tested the significance of our fixed effects via model comparison using the anova function, where one model included the effect of interest while the other model excluded it. Hypothesis testing was conducted via log-likelihood tests for count and offspring ratio LMMs and chi-squared tests for size GLMMs. All analyses were performed using R version 4.1.2 (R Core Team 2021) and the packages *nlme* (Pinheiro et al. 2022), *lme4* (Bates et al. 2015), and *glmmTMB* (Brooks et al. 2017).

Results:

Female and Male Gametophyte Development

High temperatures resulted in a significant decline in the number of females present in Week 4 (Log-Likelihood=51.1283, DF=36 , p<0.0001) (Figure 2, Table 1) . Neither pH nor the interaction between pH and temperature had a significant effect on female gametophyte numbers. The random effect of Dish had a significant effect on female gametophyte numbers in

Week 4 (Log-Likelihood=7.8795, p=0.005), indicating that there was some variation between petri dishes for this test.

Males were similarly negatively affected by high temperatures in both Week 3 (Log-Likelihood=18.9288, DF=36 , p<0.0001) and Week 4 (Log-Likelihood=45.393, DF=36 , p<0.0001). However, they varied from females in that their numbers were significantly higher under lower pH conditions in Week 4 (Log-Likelihood=8.6378, DF=36 , p=0.0033). Neither the pH:temperature interaction term nor the random effect Dish had any significant effect on male gametophyte numbers. In summary, these results indicate that temperature caused a significant decrease in female and male gametophyte numbers, whereas low pH only caused a significant increase in male gametophyte numbers.

Egg and Sporeling Counts

After 4 weeks, high temperatures were correlated with significantly lower numbers of both eggs (Log-Likelihood=33.73, DF=36 , p<0.0001) and juveniles (Log-Likelihood=36.6391, DF=36 , p<0.0001) (Figure 3). Low pH was associated with significantly larger numbers of eggs (Log-Likelihood=4.3958, DF=36, p=0.036) whereas there were no significant effects on juvenile counts (Log-Likelihood=1.0702, DF=36 , p=0.3009). The interaction term for pH:temperature was insignificant for counts of both eggs and juveniles. The random effect of Dish was insignificant for juveniles in Week 4 (Log-Likelihood=3.0544, p=0.0805), but was significant for eggs in Week 4 (Log-Likelihood=5.2080, p=0.0225). Overall, temperature caused the greatest decreases in both egg and juvenile numbers, whereas low pH caused a significant increase in eggs only.

Proportion of Productive Females

The proportion of productive females was uniformly high across all treatments, but the high temperature treatments consistently resulted in nearly 100% of females reaching

productivity by Week 4 (Figure 4). We found that the proportion of productive females was not significantly affected by the interaction between temperature and pH (Chi-Sq=0.5117, p=0.4744) nor the individual effect of pH (Chi-Sq=1.1619, p=0.2811). High temperature was the only variable to result in a significant increase in the proportion of productive females (Chi-Sq=28.187, p<0.0001).

Ratios of Offspring per Female

For mean number of eggs per female (egg/fem), we found a marginally significant effect of the interaction between temperature and pH in Week 3 (Log-Likelihood=3.7737, DF=36 , p=0.0521) but not Week 4 (Log-Likelihood=0.0976, DF=36 , p=0.7547) (Table 2, Figure 5). Investigating temperature and pH individually in Week 3, we found that low pH (Log-Likelihood=3.7345, DF=36 , p=0.0533) was marginally significantly associated with a decreased egg/fem ratio under ambient temperature treatments, but an increased egg/fem ratio under high temperature treatments. Temperature was found to be insignificant in Week 3 (Log-Likelihood=0.1406, DF=36 , p=0.7077). In Week 4, low pH was found to be significantly associated with a higher egg/fem (Log-Likelihood=9.3663, DF=36 , p=0.0022) whereas low temperature resulted in lower egg/fem (Log-Likelihood=13.114, DF=36 , p=0.0003). The effect of Dish was insignificant in both Week 3 (Log-Likelihood=0.2318, p=0.6302) and Week 4 (Log-Likelihood=0.5643, p=0.4525).

For mean number of juveniles per female (juv/fem), we found that high temperatures resulted in larger juv/fem in both Week 3 (Log-Likelihood=45.2639, DF=36 , p<0.0001) and Week 4 (Log-Likelihood=9.1867, DF=36 , p=0.0024). Low pH treatments had significantly lower juv/fem ratio than ambient pH treatments in Week 3 (Log-Likelihood=16.7485, DF=36 ,

p<0.0001) but not in Week 4 (Log-Likelihood=0.1262, DF=36 , p=0.7225). Neither the interaction term pH:temperature nor the random effect Dish were significant in either week. Increased ratios of total offspring per female (offspring/fem) were significantly correlated with high temperatures in Week 3 (Log-Likelihood=30.4186, DF=36 , p<0.0001) but not Week 4 (Log-Likelihood=0.2279, DF=36 , p=0.6331), whereas more offspring/fem were significantly correlated with low pH in Week 4 (Log-Likelihood=5.2345, DF=36 , p=0.0221) but not Week 3 (Log-Likelihood=1.3622, DF=36 , p=0.2432). Neither the interaction between temperature and pH nor the random effect Dish were significant in either week.

Across all ratios, high temperature had the greatest impacts in Week 3, resulting in lower ratios of juvs/fem and offspring/fem, whereas low pH was most significant in Week 4, resulting in high eggs/fem and offspring/fem.

   Growth of Juvenile Bull Kelp

   When analyzing the global trend across all treatments, we found that juvenile size was significantly influenced by the covariate (average number of gametophytes within each dish) in Week 4 (R^2 =0.639, p<0.0001), indicating possible density dependence where increased number of gametophytes resulted in significantly smaller juveniles (Figure 6). When included in the GLMM, the 3-way interaction between pH, temperature, and the covariate was significant (Chi-Sq=6.3387, p=0.0118), but all 2-way interactions were insignificant.

   In order to examine the role of each fixed factor (pH, temperature, and the covariate) in the 3-way interaction, we subset our model within the two pH and two temperature levels to elucidate the significance of the covariate and the other non-subset factor. When low pH and ambient pH treatments were separately analyzed, the covariate was never independently significant, but high temperatures resulted in a significant increase in size by itself under low pH

conditions (Chi-Sq=10.051 , p=0.0015) and the temperature by covariate interaction was significant under ambient pH conditions (Chi-Sq=5.3001 , p=0.02132). When the two temperature treatments were analyzed separately, the covariate was again never significant by itself, but pH resulted in a significant decrease in juvenile sizes under ambient temperature conditions (Chi-Sq=3.955 , p=0.04673), and the pH by covariate interaction had a significant effect on size under high temperature conditions (Chi-Sq=6.0391 , p=0.01399) (Figure 6). In summary, juvenile sizes were most significantly impacted by high temperatures, but pH and the number of gametophytes present greatly influenced this effect.

Discussion:

Our results demonstrated that both temperature and pH do significantly impact bull kelp reproduction and development, but the effects were more variable and nuanced than we predicted. Our most consistent finding was that high temperatures resulted in significant reduction in the number of gametophytes that survived and/or settled and the number of eggs and juveniles produced. The number of male gametophytes, female gametophytes, eggs, and juveniles were always lower in high temperature treatments than ambient temperature treatments, regardless of pH. These results align with previous findings that bull kelp exposed to increased temperature conditions had reduced spore germination rates and reduced gametophyte growth (Lind and Konar 2017; Muth et al 2019, Schiltroth 2021). Increased temperatures also result in decreased gametophyte survival and sporophyte recruitment in numerous other kelp species including giant kelp (*Macrocystis pyrifera)* (Hollarsmith et al. 2020*)*, stalked kelp (*Pterygophora californica*) (Howard 2014), spiny kelp (*Ecklonia radiat*a) (Alsuwaiyan 2021), sugar kelp (*Saccharina latissima)*, and dragon kelp (*Eualaria fistulosa*) (Lind and Konar 2017).

Low pH had a non-significant and lower effect size on the number of female gametophytes or juveniles than temperature in our study, but there was a significant increase in male gametophytes and eggs. Other studies have found varying impacts of pH on kelp gametophyte growth and survival. Several studies have found overall positive effects of low pH on *M. pyrifera* gametophyte growth, survival, and size (Roleda et al. 2012, Leal et al. 2017), where as other studies found that elevated pCO2 had little effect on rates of growth and photosynthesis (Fernández et al. 2015) or reproduction (Hollarsmith et al 2020). The variation in kelp organismal responses across studies, species, and location (Roleda and Hurd 2012; Hollarsmith et al 2020) indicates that there is much more to be understood about the impacts of ocean acidification on kelp reproduction.

Our results also support a hypothesis that high temperatures may result in bull kelp gametophytes potentially allocating more effort into reproduction. While the overall number of gametophytes and juveniles declined under high temperature conditions, the female gametophytes that survived were more productive on average, and produced more juveniles earlier than female gametophytes under ambient temperature conditions. This result suggests that female gametophytes possessed similar capacities for offspring production regardless of treatment, but that females become reproductive sooner under high temperature conditions. This idea is also supported by the separate ratios for eggs per female and juveniles per female in each week.

We also saw that our results align with this hypothesis via slower reproduction and development under ambient temperature conditions. In Week 3, high temperature treatments had higher ratios of both eggs/fem and juvs/fem than ambient temperature treatments. By Week 4, however, the eggs/fem ratio in ambient temperature treatments exceeded that of high temperature

treatments, and the juvs/fem ratio was similar regardless of temperature treatment. The later

increases in egg/fem and juvs/fem ratios in ambient temperatures seem to indicate that female

gametophytes have equal reproductive capacity under both our temperature treatments, but

females growing under high temperature treatments were progressing through reproduction

earlier. These results are consistent with those of Howard (2014), who found that ambient

temperatures lead to slower microstage reproduction of bull kelp as well as giant kelp

(*Macrocystis pyrifera*) and stalked kelp (*Pterygophora californica*)..

In contrast, low pH conditions seemed to impact reproductive efforts differently based on

temperature conditions. Under ambient temperatures, low pH appeared to slow reproduction by

producing the lowest egg/fem, juv/fem, and offspring/fem ratios of all treatments in Week 3, but

then increased enough to match the ratios of other treatments in Week 4 (Figure 5). Under high

temperature conditions on the other hand, low pH treatments produced the highest egg/fem and

offspring/fem ratios and second highest juv/fem ratios of all treatments in Week 3, with only

nominal increases in these ratios in Week 4. Additionally, the lowest proportion of productive

females was observed in low pH treatments under ambient temperatures (Figure 4), and these

females seemed to produce more offspring per female, later in the experiment. The greatest

increase in productivity between Week 3 and Week 4 under any pH treatments can be found in

the large increase in egg/fem ratios under ambient temperature conditions. This late increase in

production of eggs may potentially signal that a delay in reproduction occurs under low pH and

ambient temperature conditions.

Recent advances in kelp reproduction studies have given needed attention to delayed

development of microscopic stages and the resulting "bank of microscopic forms" (Hoffman and

Santelices 1991, Carney and Edwards 2006, Schoenrock et al. 2021), but less focus has been

placed on the factors that may accelerate microscopic kelp development. In terrestrial plants, increased temperatures have been found to result in an acceleration of pollen tube growth and stigma and ovule development, which correspond to an overall reduction of the length of time females are receptive to pollination (Hedhly et al. 2009). Reviews of other marine organisms, specifically benthic invertebrates, have shown that increased sea surface temperatures may increase the rate and timing of development and spawning (Przeslawski et al. 2008). In order to better understand the ability of populations to recover from extreme climate disturbance events, more research is needed to better understand the effect of climate stressors on survival, time to development, and propagule production.

For all variables other than egg/fem in Week 3, we found no significant temperature and pH interactions on bull kelp microscopic stages, which contrasts with studies of giant kelp, that did find interactive effects of pH or pCO2 and temperature on spore, gametophyte, and sporeling production (Gaitan-Espitia et al. 2014, Shukla and Edwards 2017, Hollarsmith et al. 2020). Our results interestingly reflect natural seasonal fluctuations in northern California's coastal waters, where conditions of low pH and high temperature rarely overlap. Oceanographic conditions in California's coastal waters are typically determined by three seasons: upwelling season (April to June), relaxation season (July to October), and storm season (October to March) (García-Reyes and Largier 2012). Upwelling season is characterized by the upwelling of cold, dense, nutrient rich water from the deep ocean to the coastline, causing coastal waters to be colder and more acidic. During non-upwelling season, on the other hand, the intensity of upwelling processes decreases, and coastal waters become relatively warmer, more basic, and exhibit less primary productivity and chlorophyll-a content due to shortened day lengths and decreased nutrient concentrations (García-Reyes and Largier 2012). The bull kelp lifecycle has a relatively

consistent, yet still flexible, phenology. The majority of visible juveniles appear in the spring (upwelling season) and most adults become reproductive by the end of July (relaxation season), but these two events of visible recruitment and spore release have been observed to occur in all seasons, albeit at much lower rates (Maxell and Miller 1996, Dobkowski et al. 2018).

As a result, the need for gametophytes to endure high temperature or low pH conditions is dependent on the life cycle of the parent sporophyte. Gametophytes and sporelings that develop in the spring will likely be most exposed to low temperatures and low pH as a result of upwelling, but the vast majority of gametophytes and sporelings that develop in the fall will be exposed to non-upwelling season late summer conditions. As such, it is conceivable that high temperatures in September and October (relaxation season) would affect the first month of juvenile development, whereas upwelling season would likely be more important for visible juveniles than the microscopic ones.

However, our results potentially contrast with those of Dobkowski et al. 2018 in that we found that low pH (most often seen in the Spring upwelling season) resulted in slower reproduction and growth whereas high temperature (most often seen in late summer and fall) accelerated it. In their study, Dobkowski et al. (2018) witnessed the quickest recruitment of visible juveniles (indicating faster microscopic development times) in the spring (upwelling season), and slowest recruitment (implying slower microscopic development times) in the late summer and fall (relaxation season. A potential explanation for the different observed reproductive rates is that Dobkowski et al. conducted their experiments in the field, where they were exposed to a full of array of abiotic conditions, whereas our experiments were conducted in a laboratory setting where only temperature and pH were manipulated, and all other variables were held constant, including nutrients. Previous studies have shown that delayed development

of microscopic kelp stages is often closely tied to insufficient nutrient and light regimes (Carney and Edwards 2010), both of which are present between September to March due to dampened upwelling conditions and reduced daylength. As such, the slow development over winter in natural populations suggests that nutrients from upwelling and daylength could be more important than temperature and pH fluctuations in promoting the development of microscopic kelp stages.

Our results suggest that there may be some density dependence effects on juvenile growth at these microscopic stages. The difference in juvenile sizes between treatments was most significantly correlated with temperature, but also showed at least a marginally significant correlation with the number of gametophytes present in both weeks (Figure 5). However, due to the fact that high temperatures consistently resulted in significant decreases in gametophyte numbers, the relationships of both temperature and number of gametophytes to gametophyte size are confounded, and direct causation cannot be determined. As a result, more research is needed to see whether these increased sizes were really a result of high temperatures or whether they were a result of lowered density of individuals.

In natural populations, there are numerous density-dependent effects that impact kelp reproduction and recruitment. At initial spore settlement, high densities of gametophytes are needed for fertilization between male and female gametophytes to occur, so Allee effects may occur if spores settle at a density of less than 1 spore/mm^2 (Reed 1990). The direction of density-dependence then reverses somewhere between the gametophyte stage and the point where a juvenile becomes easily detectable to the naked eye, and numerous kelps, including bull kelp, exhibit subsequent increases in mortality until they reach the adult life stage (Schiel and Foster 2006, Reed et al. 1991, Dobkowski et al. 2018). Our petri dishes, each containing thousands of

spores and gametophytes, contained a bottom surface area of approximately 157.8cm^2, whereas an adult bull kelp holdfast can grow up to an estimated 40cm in diameter and 1256cm^2 surface area (Abbot and Hollenberg 1976). As a result, less than one juvenile sporophyte in each petri dish would be expected to survive to adulthood and reproductive maturity in natural environments.

The results of this research indicate that climate change will significantly affect bull kelp reproduction via increased temperatures, and, to a lesser extent, ocean acidification. Increasing frequency and intensity of extreme temperature events such as marine heatwaves will likely lead to a massive decrease in the survival of gametophytes and decreased, but accelerated, production of juvenile sporophytes. Lowered pH, mimicking ocean acidification, resulted in an increase in numbers of male gametophytes and juvenile sporelings, as well as a slower reproduction rate. The ability of bull kelp to recover from extreme climate events depends on the ability of all lifestages to withstand abiotic stress. Currently, managers and scientists are focused on understanding the natural processes that guide kelp recovery and whether those may require some degree of management intervention. In order to better understand the ability of species to recover from climate change, more research is needed to understand the ability of intermediate stages between propagules and adults to withstand abiotic stress. In order to intervene successfully through restoration, an understanding of potential bottlenecks and challenges present at each life stage is necessary. This study lends a little insight into the otherwise largely unknown world of microscopic life stages of kelp

Acknowledgments:

We would like to thank Rob Coyan for assistance collecting sporophylls, and Carol Vines for help with microscopy methods. A. Blandino of the UC Davis Statistical Consulting Group and

References:

Abbott, I. A., A. Isabella, and G. J. Hollenberg. 1992. Marine Algae of California. Stanford University Press.

Arafeh-Dalmau, N., G. Montaño-Moctezuma, J. A. Martínez, R. Beas-Luna, D. S. Schoeman, and G. Torres-Moye. 2019. Extreme Marine Heatwaves Alter Kelp Forest Community Near Its Equatorward Distribution Limit. Frontiers in Marine Science 6.

Alsuwaiyan, N. A., S. Vranken, K. Filbee-Dexter, M. Cambridge, M. A. Coleman, and T. Wernberg. 2021. Genotypic variation in response to extreme events may facilitate kelp adaptation under future climates. Marine Ecology Progress Series 672:111–121.

Bakun, A., B. A. Black, S. J. Bograd, M. García-Reyes, A. J. Miller, R. R. Rykaczewski, and W. J. Sydeman. 2015. Anticipated Effects of Climate Change on Coastal Upwelling Ecosystems. Current Climate Change Reports 1:85–93.

Ban, S. S., N. A. J. Graham, and S. R. Connolly. 2014. Evidence for multiple stressor interactions and effects on coral reefs. Global Change Biology 20:681–697.

Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects Models Using lme4." *Journal of Statistical Software*, **67**(1), 1–48. doi:10.18637/jss.v067.i01.

Beas-Luna, R., F. Micheli, C. B. Woodson, M. Carr, D. Malone, J. Torre, C. Boch, J. E. Caselle, M. Edwards, J. Freiwald, S. L. Hamilton, A. Hernandez, B. Konar, K. J. Kroeker, J. Lorda, G. Montaño-Moctezuma, and G. Torres-Moye. 2020. Geographic variation in responses of kelp forest communities of the California Current to recent climatic changes. Global Change Biology 26:6457–6473.

Bennett, S., T. Wernberg, S. D. Connell, A. J. Hobday, C. R. Johnson, and E. S. Poloczanska. 2016. The "Great Southern Reef": social, ecological and economic value of Australia's neglected kelp forests. Marine and Freshwater Research 67:47.

Blamey, L. K., and J. J. Bolton. 2018. The economic value of South African kelp forests and temperate reefs: Past, present and future. Journal of Marine Systems 188:172–181.

Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Maechler M, Bolker BM (2017). "glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling." *The R Journal*, **9**(2), 378–400. doi:10.32614/RJ-2017-066.

Carney, L. T., and M. S. Edwards. 2006. Cryptic Processes in the Sea: A Review of Delayed Development in theMicroscopic Life Stages of Marine Macroalgae. ALGAE 21:161–168.

Carney, L. T., and M. S. Edwards. 2010. Role of Nutrient Fluctuations and Delayed Development in Gametophyte Reproduction by Macrocystis Pyrifera (phaeophyceae) in Southern California1. Journal of Phycology 46:987–996.

Carr, M. H., and D. C. Reed. 2015. Shallow Rocky Reefs and Kelp Forests. Pages 311–336 in H. A. Mooney and E. S. Zavaleta, editors. Ecosystems of California. University of California Press, Berkeley.

Cavanaugh, K. C., D. C. Reed, T. W. Bell, M. C. N. Castorani, and R. Beas-Luna. 2019. Spatial Variability in the Resistance and Resilience of Giant Kelp in Southern and Baja California to a Multiyear Heatwave. Frontiers in Marine Science 6.

Cooley, S., D. Schoeman, L. Bopp, P. Boyd, S. Donner, S.-I. Ito, W. Kiessling, P. Martinetto, E. Ojea, M.-F. Racault, B. Rost, M. Skern-Mauritzen, and D. Y. Ghebrehiwet. 2022. Ocean and coastal ecosystems and their services. Page in H.-O. Pörtner, D. C. Roberts, M. M. B.

Tignor, E. S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, and B. Rama, editors. Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.

Dickson, A. G., and F. J. Millero. 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. Deep Sea Research Part A. Oceanographic Research Papers 34:1733–1743.

Dickson, A. G., C. L. Sabine, J. R. Christian, C. P. Bargeron, and North Pacific Marine Science Organization, editors. 2007. Guide to best practices for ocean CO2 measurements. North Pacific Marine Science Organization, Sidney, BC.

Dobkowski, K. A., K. D. Flanagan, and J. R. Nordstrom. 2019. Factors influencing recruitment and appearance of bull kelp, Nereocystis luetkeana (phylum Ochrophyta). Journal of Phycology 55:236–244.

Feely, R. A., C. L. Sabine, K. Lee, W. Berelson, J. Kleypas, V. J. Fabry, and F. J. Millero. 2004. Impact of Anthropogenic CO2 on the CaCO3 System in the Oceans. Science 305:362–366.

Feely, R. A., C. L. Sabine, J. M. Hernandez-Ayon, D. Ianson, and B. Hales. 2008. Evidence for Upwelling of Corrosive "Acidified" Water onto the Continental Shelf. Science 320:1490–1492.

Feely, R. A., S. C. Doney, and S. R. Cooley. 2009. Ocean Acidification: Present Conditions and Future Changes in a High-CO₂ World. Oceanography 22:36–47.

Feely, R. A., R. R. Okazaki, W.-J. Cai, N. Bednaršek, S. R. Alin, R. H. Byrne, and A. Fassbender. 2018. The combined effects of acidification and hypoxia on pH and aragonite

saturation in the coastal waters of the California current ecosystem and the northern Gulf of
Mexico. Continental Shelf Research 152:50–60.

Fernández, P. A., M. Y. Roleda, and C. L. Hurd. 2015. Effects of ocean acidification on the
photosynthetic performance, carbonic anhydrase activity and growth of the giant kelp
Macrocystis pyrifera. Photosynthesis Research 124:293–304.

Ferrari, M. C. O., M. I. McCormick, P. L. Munday, M. G. Meekan, D. L. Dixson, Ö. Lonnstedt,
and D. P. Chivers. 2011. Putting prey and predator into the CO2 equation – qualitative and
quantitative effects of ocean acidification on predator–prey interactions. Ecology Letters
14:1143–1148.

Filbee-Dexter, K., T. Wernberg, S. P. Grace, J. Thormar, S. Fredriksen, C. N. Narvaez, C. J.
Feehan, and K. M. Norderhaug. 2020. Marine heatwaves and the collapse of marginal
North Atlantic kelp forests. Scientific Reports 10:13388.

Finger, D. J. I., M. L. McPherson, H. F. Houskeeper, and R. M. Kudela. 2021. Mapping bull
kelp canopy in northern California using Landsat to enable long-term monitoring. Remote
Sensing of Environment 254:112243.

Foster, M. S., and D. R. Schiel. 1985. The Ecology of Giant Kelp Forests in California: A
Community Profile. U.S. Fish and Wildlife Service.

Gaitán-Espitia, J. D., J. R. Hancock, J. L. Padilla-Gamiño, E. B. Rivest, C. A. Blanchette, D. C.
Reed, and G. E. Hofmann. 2014. Interactive effects of elevated temperature and pCO2 on
early-life-history stages of the giant kelp Macrocystis pyrifera. Journal of Experimental
Marine Biology and Ecology 457:51–58.

García-Reyes, M., and J. L. Largier. 2012. Seasonality of coastal upwelling off central and northern California: New insights, including temporal and spatial variability. Journal of Geophysical Research: Oceans 117.

García-Reyes, M., W. J. Sydeman, D. S. Schoeman, R. R. Rykaczewski, B. A. Black, A. J. Smit, and S. J. Bograd. 2015. Under Pressure: Climate Change, Upwelling, and Eastern Boundary Upwelling Ecosystems. Frontiers in Marine Science 2.

Gattuso, J.-P., A. Magnan, R. Billé, W. W. L. Cheung, E. L. Howes, F. Joos, D. Allemand, L. Bopp, S. R. Cooley, C. M. Eakin, O. Hoegh-Guldberg, R. P. Kelly, H.-O. Pörtner, A. D. Rogers, J. M. Baxter, D. Laffoley, D. Osborn, A. Rankovic, J. Rochette, U. R. Sumaila, S. Treyer, and C. Turley. 2015. Contrasting futures for ocean and society from different anthropogenic CO2 emissions scenarios. Science 349:aac4722.

Gattuso, J.-P., J.-M. Epitalon, H. Lavigne, J. Orr, B. Gentili, M. Hagens, A. Hofmann, J.-D. Mueller, A. Proye, J. Rae, and K. Soetaert. 2022, July 4. seacarb: Seawater Carbonate Chemistry.

Gentemann, C. L., M. R. Fewings, and M. García-Reyes. 2017. Satellite sea surface temperatures along the West Coast of the United States during the 2014–2016 northeast Pacific marine heat wave. Geophysical Research Letters 44:312–319.

Graham, M., C. Harrold, S. Lisin, K. Light, J. Watanabe, and M. Foster. 1997. Population dynamics of giant kelp Macrocystis pyrifera along a wave exposure gradient. Marine Ecology Progress Series 148:269–279.

Graham, M., J. Vásquez, and A. Buschmann. 2007. Global Ecology of the Giant Kelp Macrocystis. Pages 39–88.

Hamilton, S. L., T. W. Bell, J. R. Watson, K. A. Grorud-Colvert, and B. A. Menge. 2020. Remote sensing: generation of long-term kelp bed data sets for evaluation of impacts of climatic variation. Ecology 101:e03031.

Hamilton, S. L., M. G. Gleason, N. Godoy, N. Eddy, and K. Grorud-Colvert. 2022. Ecosystem-based management for kelp forest ecosystems. Marine Policy 136:104919.

Harvey, B. P., D. Gwynn-Jones, and P. J. Moore. 2013. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. Ecology and Evolution 3:1016–1030.

Hedhly, A. 2011. Sensitivity of flowering plant gametophytes to temperature fluctuations. Environmental and Experimental Botany 74:9–16.

Hoffmann, A., and B. Santelices. 1991. Banks of algal microscopic forms: hypotheses on their functioning and comparisons with seed banks. Marine Ecology Progress Series 79:185–194.

Hollarsmith, J. A., A. H. Buschmann, C. Camus, and E. D. Grosholz. 2020. Varying reproductive success under ocean warming and acidification across giant kelp (Macrocystis pyrifera) populations. Journal of Experimental Marine Biology and Ecology 522:151247.

Howard, A. 2014. Effects of Temperature on Sexual Competition in Kelps: Implications for Range Shifts in Foundation Species. Master's Theses.

Kroeker, K. J., R. L. Kordas, R. Crim, I. E. Hendriks, L. Ramajo, G. S. Singh, C. M. Duarte, and J.-P. Gattuso. 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. Global Change Biology 19:1884–1896.

Leal, P. P., C. L. Hurd, P. A. Fernández, and M. Y. Roleda. 2017. Ocean acidification and kelp development: Reduced pH has no negative effects on meiospore germination and

gametophyte development of Macrocystis pyrifera and Undaria pinnatifida. Journal of Phycology 53:557–566.

Lind, A. C., and B. Konar. 2017. Effects of abiotic stressors on kelp early life-history stages. ALGAE 32:223–233.

Malone, D. P., K. Davis, S. I. Lonhart, A. Parsons-Field, J. E. Caselle, and M. H. Carr. 2022. Large-scale, multidecade monitoring data from kelp forest ecosystems in California and Oregon (USA). Ecology 103:e3630.

Maxell, B. A., and K. A. Miller. 1996. Demographic Studies of the Annual Kelps Nereocystis luetkeana and Costaria costata (Laminariales, Phaeophyta) in Puget Sound, Washington 39:479–490.

McPherson, M. L., D. J. I. Finger, H. F. Houskeeper, T. W. Bell, M. H. Carr, L. Rogers-Bennett, and R. M. Kudela. 2021. Large-scale shift in the structure of a kelp forest ecosystem co-occurs with an epizootic and marine heatwave. Communications Biology 4:1–9.

Mehrbach, C., C. H. Culberson, J. E. Hawley, and R. M. Pytkowicx. 1973. Measurement of the Apparent Dissociation Constants of Carbonic Acid in Seawater at Atmospheric Pressure1. Limnology and Oceanography 18:897–907.

Miner, C. M., J. L. Burnaford, R. F. Ambrose, L. Antrim, H. Bohlmann, C. A. Blanchette, J. M. Engle, S. C. Fradkin, R. Gaddam, C. D. G. Harley, B. G. Miner, S. N. Murray, J. R. Smith, S. G. Whitaker, and P. T. Raimondi. 2018. Large-scale impacts of sea star wasting disease (SSWD) on intertidal sea stars and implications for recovery. PLOS ONE 13:e0192870.

Munday, P. L., D. L. Dixson, M. I. McCormick, M. Meekan, M. C. O. Ferrari, and D. P. Chivers. 2010. Replenishment of fish populations is threatened by ocean acidification. Proceedings of the National Academy of Sciences 107:12930–12934.

Muth, A. F., M. H. Graham, C. E. Lane, and C. D. G. Harley. 2019. Recruitment tolerance to increased temperature present across multiple kelp clades. Ecology 100:e02594.

Pfister, C. A., M. A. Altabet, and B. L. Weigel. 2019. Kelp beds and their local effects on seawater chemistry, productivity, and microbial communities. Ecology 100:e02798.

Pfister, C. A., H. D. Berry, and T. Mumford. 2018. The dynamics of Kelp Forests in the Northeast Pacific Ocean and the relationship with environmental drivers. Journal of Ecology 106:1520–1533.

Przeslawski, R., S. Ahyong, M. Byrne, G. Wörheide, and P. Hutchings. 2008. Beyond corals and fish: the effects of climate change on noncoral benthic invertebrates of tropical reefs. Global Change Biology 14:2773–2795.

Przeslawski, R., M. Byrne, and C. Mellin. 2015. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. Global Change Biology 21:2122–2140.

Rasband, W.S. 2019. *ImageJ*. U. S. National Institutes of Health, Bethesda, Maryland, USA. https://imagej.nih.gov/ij/, 1997-2018.

Reed, D. C. 1990. The Effects of Variable Settlement and Early Competition on Patterns of Kelp Recruitment. Ecology 71:776–787.

Reed, D. C., M. Neushul, and A. W. Ebeling. 1991. Role of Settlement Density on Gametophyte Growth and Reproduction in the Kelps Pterygophora Californica and Macrocystis Pyrifera (phaeophyceae)1. Journal of Phycology 27:361–366.

Reid, J., L. Rogers-Bennett, F. Lavín, M. Pace, C. Catton, and I. Taniguchi. 2016. The economic value of the recreational red abalone fishery in northern California. California Fish and Game 102:119–130.

Riebesell, U., V. J. Fabry, L. Hansson, and J.-P. Gattuso. 2011. Guide to best practices for ocean acidification research and data reporting. Page (U. Riebesell, V. J. Fabry, L. Hansson, and J.-P. Gattuso, Eds.). Office for Official Publications of the European Communities, Luxembourg.

Rogers-Bennett, L., and C. A. Catton. 2019. Marine heat wave and multiple stressors tip bull kelp forest to sea urchin barrens. Scientific Reports 9:1–9.

Rogers-Bennett, L., and D. Okamoto. 2007. Chapter 32 - Mesocentrotus franciscanus and Strongylocentrotus purpuratus. Pages 593–608 in J. M. Lawrence, editor. Developments in Aquaculture and Fisheries Science. Elsevier.

Roleda, M. Y., and C. L. Hurd. 2012. Seaweed Responses to Ocean Acidification. Pages 407–431 in C. Wiencke and K. Bischof, editors. Seaweed Biology: Novel Insights into Ecophysiology, Ecology and Utilization. Springer Berlin Heidelberg, Berlin, Heidelberg.

Roleda, M. Y., J. N. Morris, C. M. McGraw, and C. L. Hurd. 2012. Ocean acidification and seaweed reproduction: increased CO2 ameliorates the negative effect of lowered pH on meiospore germination in the giant kelp Macrocystis pyrifera (Laminariales, Phaeophyceae). Global Change Biology 18:854–864.

Santelices, B. 2002. Recent Advances in Fertilization Ecology of Macroalgae1. Journal of Phycology 38:4–10.

Schiel, D. R., and M. S. Foster. 2006. The Population Biology of Large Brown Seaweeds: Ecological Consequences of Multiphase Life Histories in Dynamic Coastal Environments. Annual Review of Ecology, Evolution, and Systematics 37:343–372.

Schiel, D. R., J. R. Steinbeck, and M. S. Foster. 2004. Ten Years of Induced Ocean Warming Causes Comprehensive Changes in Marine Benthic Communities. Ecology 85:1833–1839.

Schiltroth, B. 2021, January 22. Effects of climate change on two species of foundational brown algae, Nereocystis luetkeana and Fucus gardneri, within the Salish Sea. Thesis, Science: Biological Sciences Department.

Schoenrock, K. M., T. A. McHugh, and S. A. Krueger-Hadfield. 2021. Revisiting the 'bank of microscopic forms' in macroalgal-dominated ecosystems. Journal of Phycology 57:14–29.

Shukla, P., and M. S. Edwards. 2017. Elevated pCO_2 is less detrimental than increased temperature to early development of the giant kelp, *Macrocystis pyrifera* (Phaeophyceae, Laminariales). Phycologia 56:638–648.

Small, D. P., P. Calosi, D. Boothroyd, S. Widdicombe, and J. I. Spicer. 2016. The sensitivity of the early benthic juvenile stage of the European lobster Homarus gammarus (L.) to elevated pCO2 and temperature. Marine Biology 163:53.

Smith, J. G., and M. T. Tinker. 2022. Alternations in the foraging behaviour of a primary consumer drive patch transition dynamics in a temperate rocky reef ecosystem. Ecology Letters 25:1827–1838.

Smith, K. E., M. T. Burrows, A. J. Hobday, N. G. King, P. J. Moore, A. Sen Gupta, M. S. Thomsen, T. Wernberg, and D. A. Smale. 2023. Biological Impacts of Marine Heatwaves. Annual Review of Marine Science 15:null.

Steneck, R. S., M. H. Graham, B. J. Bourque, D. Corbett, J. M. Erlandson, J. A. Estes, and M. J. Tegner. 2002. Kelp forest ecosystems: biodiversity, stability, resilience and future. Environmental Conservation 29:436–459.

Strain, E. M. A., R. J. Thomson, F. Micheli, F. P. Mancuso, and L. Airoldi. 2014. Identifying the interacting roles of stressors in driving the global loss of canopy-forming to mat-forming algae in marine ecosystems. Global Change Biology 20:3300–3312.

Straub, S. C., T. Wernberg, M. S. Thomsen, P. J. Moore, M. T. Burrows, B. P. Harvey, and D. A. Smale. 2019. Resistance, Extinction, and Everything in Between – The Diverse Responses of Seaweeds to Marine Heatwaves. Frontiers in Marine Science 6.

Supratya, V. P., L. J. M. Coleman, and P. T. Martone. 2020. Elevated Temperature Affects Phenotypic Plasticity in the Bull Kelp (Nereocystis luetkeana, Phaeophyceae). Journal of Phycology 56:1534–1541.

Swanson, A. K., and C. H. Fox. 2007. Altered kelp (Laminariales) phlorotannins and growth under elevated carbon dioxide and ultraviolet-B treatments can influence associated intertidal food webs. Global Change Biology 13:1696–1709.

Thom, R. M. 1996. CO2-Enrichment effects on eelgrass (Zostera marina L.) and bull kelp (Nereocystis luetkeana (mert.) P & R.). Water, Air, and Soil Pollution 88:383–391.

Figure 1: The microscopic stages of bull kelp: A) Female gametophyte (image area =0.065 mm^2); B) Male gametophyte (image area = 0.065 mm^2); C) Female gametophyte producing an egg (image area = 0.077 mm^2); D) Female gametophytes with juvenile sporelings (image area = 0.065 mm^2).

Figure 2: Female and male gametophytes present in each photo after 4 weeks of growth. The box plots summarize the mean (diamond) and median (box midline) for each treatment, the first and third quartiles (upper and lower box limits), outliers within 1.5 times the inter-quartile range (vertical lines), and outliers beyond that range (dots).

Figure 3: Eggs and juveniles present in each photo after 4 weeks of growth. The box plots summarize the mean (diamond) and median (box midline) for each treatment, the first and third quartiles (upper and lower box limits), outliers within 1.5 times the inter-quartile range (vertical lines), and outliers beyond that range (dots).

Figure 4: Proportion of productive female gametophytes after 4 weeks of growth. The box plots summarize the mean (diamond) and median (box midline) for each treatment, the first and third quartiles (upper and lower box limits), outliers within 1.5 times the inter-quartile range (vertical lines), and outliers beyond that range (dots).

Figure 5: Eggs, juveniles, and total offspring (eggs + juveniles) per female after 3 and 4 weeks of growth. The box plots summarize the mean (diamond) and median (box midline) for each treatment, the first and third quartiles (upper and lower box limits), outliers within 1.5 times the inter-quartile range (vertical lines), and outliers beyond that range (dots).

Figure 6: Left panel shows the average size of juveniles after 4 weeks of growth. The left panel shows box plots that summarize the mean (diamond) and median (box midline) for each treatment, the first and third quartiles (upper and lower box limits), outliers within 1.5 times the inter-quartile range (vertical lines), and outliers beyond that range (dots). The right panel shows the relationship of the covariate (mean number of gametophytes) to the response variable (mean juvenile size). Trends within groups are represented by colored lines, whereas the trend across all groups is represented by the dotted black line. Heterogeneous slopes and different ranges of values for each treatment indicate that the different treatments are confounded with differences in the covariate.

Week 4

Treatment	Eggs	Females	Juveniles	Males
Ambient Temp, Low pH	15.6 ± 10.48	15.83 ± 4.05	20.03 ± 6.99	12.83 ± 3.79
Ambient Temp, Ambient pH	12.24 ± 6.65	17.55 ± 5.13	21.72 ± 6.14	10.17 ± 4.26
High Temp, Low pH	6.03 ± 5.17	8.5 ± 3.88	12 ± 5.26	6.1 ± 3.17
High Temp, Ambient pH	3.17 ± 2.74	8.13 ± 2.21	12.47 ± 4.29	4.17 ± 2.29

Table 1: Means and SDs of Count data. Counts represent the mean value of three photos per 10 dishes in each treatment

Week 3

Treatment	Proportion Productive Females	Eggs per Female	Juveniles per Female	Total Offspring per Female
Ambient Temp, Low pH	0.68 ± 0.2	0.76 ± 0.4	0.43 ± 0.27	1.2 ± 0.48
Ambient Temp, Ambient pH	0.8 ± 0.15	0.76 ±0.4	0.66 ± 0.29	1.41 ± 0.41
High Temp, Low pH	0.89 ± 0.13	0.96 ± 0.55	0.94 ± 0.35	1.9 ± 0.49
High Temp, Ambient pH	0.95 ± 0.08	0.63 ± 0.42	1.26 ± 0.42	1.89 ± 0.53

Week 4

Treatment	Proportion Productive Females	Eggs per Female	Juveniles per Female	Total Offspring per Female

Ambient Temp, Low pH	0.89 ± 0.11	0.97 ± 0.53	1.3 ± 0.4	2.26 ± 0.65
Ambient Temp, Ambient pH	0.92 ± 0.1	0.68 ± 0.3	1.29 ± 0.37	1.97 ± 0.42
High Temp, Low pH	0.99 ±0.03	0.67 ± 0.48	1.49 ± 0.38	2.16 ± 0.49
High Temp, Ambient pH	1 ± 0	0.4 ± 0.36	1.53 ± 0.33	1.93 ± 0.41

Table 2: Means and SDs of Productivity Data. The mean productivity is the number of offspring per female per three photos per 10 dishes in each treatment

Week 4

Treatment	Juvenile blade area (um^2)	Covariate: Mean # gametophytes per dish
Ambient Temp, Low pH	1561.11 ± 1101.82	28.17 ± 3.97
Ambient Temp, Ambient pH	1841.37 ± 1288.6	28.22 ± 5.47
High Temp, Low pH	3088.05 ± 3079.67	15.06 ± 4.16
High Temp, Ambient pH	3304.37 ± 2726.05	11.97 ± 1.31

Table 3: Means and SDs of sizes and covariates. Size represents the area of juvenile blades in um2, where as the covariate refers to the mean number of gametophytes per dish.

SUPPLEMENTARY TABLES:

Week 4 Females

Model: lmer(sqrt(Females)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.08195				7.879457	**0.005**
Residual	0.21277					

Interactions	Estimate	SE	Df	T-value	LLR	P-value
Temperature : pH	-0.213035	0.2478791	36	-0.859432	0.81337	0.3671

Fixed Effects	Estimate	SE	Df	T-value	LLR	P-value
Intercept	2.82775	0.12364	79	22.8702		
Temperature	1.31545	0.1757	36	7.4871	51.1283	**< 0.0001**

| pH | 0.02026 | 0.17486 | 36 | 0.11587 | 0.51791 | 0.4717 |

Table 1: Statistical outcomes for models of female count data

Week 4 Males

Model: lmer(sqrt(Males)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.04678				1.082005	0.2982
Residual	0.41635					
Interactions	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Temperature : pH	0.032674	0.2733936	36	0.119513	0.01556	0.9007
Fixed Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Intercept	1.94558	0.13622	79	14.2825		
Temperature	1.17496	0.19399	36	6.05686	45.393	**< 0.0001**
pH	0.38634	0.19265	36	2.00546	8.63776	**0.0033**

Table 2: Statistical outcomes for models of male count data

Week 4 Eggs

Model: lmer(sqrt(Eggs)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.28648				5.207977	**0.0225**
Residual	0.98171					
Interactions	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**

	Estimate	SE	Df	T-value	LLR	P-value
Temperature : pH	-0.204592	0.496794	36	-0.411825	0.18959	0.6633
Fixed Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Intercept	1.52398	0.24773	79	6.15174		
Temperature	1.82336	0.35222	36	5.17672	33.73	**<0.0001**
pH	0.61102	0.35035	36	1.74406	4.39583	**0.036**

Table 3: Statistical outcomes for models of egg count data

Week 4 Juveniles

Model: lmer(sqrt(Juveniles)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
TrtRep (Dish)	0.08057				3.054386	0.0805
Residual	0.38809					
Interactions	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Temperature : pH	-0.122581	0.290655	36	-0.42174	0.19862	0.6558
Fixed Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Intercept	3.48191	0.14489	79	24.0311		
Temperature	1.13264	0.20614	36	5.49452	36.6391	**<0.0001**
pH	-0.0829	0.20491	36	-0.4047	1.07023	0.3009

Table 4: Statistical outcomes for models of juvenile count data

Week 3 Eggs per Female

Model: lmer(EPF~ Temp + pH + Temp:pH + (1|Trt_Rep), data=wk3dt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.00925				0.231782	0.6302
Residual	0.19129					
Interactions	Estimate	SE	Df	T-value	LLR	P-value
Temperature : pH	-0.322532	0.1709	36	-1.887257	3.77374	**0.0521**
Fixed Effects	Estimate	SE	Df	T-value	LLR	P-value
Intercept	0.62774	0.08545	80	7.34626		
Temperature	0.12938	0.12084	36	1.0706	0.14057	0.7077
pH	0.3294	0.12084	36	2.72582	3.73457	**0.0533**

Week 4 Eggs per Female

Model: lmer(log(EPF+1)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.00455				0.564319	0.4525
Residual	0.05809					
Interactions	Estimate	SE	Df	T-value	LLR	P-value
Temperature : pH	-0.028905	0.098165	36	-0.29445	0.09764	0.7547
Fixed Effects	Estimate	SE	Df	T-value	LLR	P-value
Intercept	0.30672	0.0489	79	6.27227		
Temperature	0.19842	0.06967	36	2.84811	13.114	**0.0003**
pH	0.16569	0.06916	36	2.39591	9.36634	**0.0022**

Table 5: Statistical outcomes for models of egg/female ratios.

Week 3 Juveniles per Female

Model: lmer(sqrt(JPF)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=wk3dt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.00127				0.11794	0.7313
Residual	0.03731					

Interactions	Estimate	SE	Df	T-value	LLR	P-value
Temperature : pH	-0.01826	0.074029	36	-0.2466	0.06719	0.7955

Fixed Effects	Estimate	SE	Df	T-value	LLR	P-value
Intercept	1.10355	0.03701	80	29.814		
Temperature	-0.313	0.05235	36	-5.9799	45.2639	**<0.0001**
pH	-0.1512	0.05235	36	-2.8877	16.7485	**<0.0001**

Week 4 Juveniles per Female

Model: lmer(sqrt(JPF)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.00259				1.14462	0.2847
Residual	0.22314					

Interactions	Estimate	SE	Df	T-value	LLR	P-value

	Estimate	SE	Df	T-value	LLR	P-value
Temperature : pH	0.019201	0.063551	36	0.30213	0.10041	0.7513
Fixed Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Intercept	1.23074	0.03167	79	38.8668		
Temperature	-0.1066	0.04509	36	-2.364	9.18669	**0.0024**
pH	-0.0192	0.04478	36	-0.4516	0.12616	0.7225

Table 6: Statistical outcomes for models of juvenile/female ratios.

Week 3 Offspring per Female

Model: lmer(OPF~ Temp + pH + Temp:pH + (1|Trt_Rep), data=wk3dt)

Random Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
TrtRep (Dish)	0.00557				0.065118	0.7986
Residual	0.22368					
Interactions	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Temperature : pH	-0.2254192	0.179037	36	-1.2591	1.70902	0.1911
Fixed Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Intercept	1.88519	0.08952	80	21.0593		
Temperature	-0.4731	0.1266	36	-3.7368	30.4186	**<0.0001**
pH	0.01056	0.1266	36	0.08344	1.36223	0.2432

Week 4 Offspring per Female

Model: lmer(log(OPF)~ Temp + pH + Temp:pH + (1|Trt_Rep), data=pdt)

Random Effects	Estimate	SE	Df	T-value	LLR	P-value
TrtRep (Dish)	0.00612				1.053258	0.3048
Residual	0.05503					
Interactions	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Temperature : pH	0.007708	0.099264	36	0.07765	0.00619	0.9373
Fixed Effects	**Estimate**	**SE**	**Df**	**T-value**	**LLR**	**P-value**
Intercept	0.63724	0.04946	79	12.8842		
Temperature	0.01849	0.07043	36	0.2625	0.22787	0.6331
pH	0.10749	0.06995	36	1.53682	5.23446	**0.0221**

Table 7: Statistical outcomes for models of offspring/female ratios.

Week 4 Proportion of Productive Females

Model: glmmTMB((Prop_prod-0.00001) ~ Temp*pH +(1|FullTrt), family = beta_family(link="logit"), data = pdt)

Random Effects	Estimate	SE	z-value	ChiSq	P-value
TrtRep (Dish)	2.24E-01				
Interactions	**Estimate**	**SE**	**z-value**	**ChiSq**	**P-value**
Temperature : pH	-0.34443	0.47961	-0.718	0.5117	0.4744
Fixed Effects	**Estimate**	**SE**	**z-value**	**ChiSq**	**P-value**
Intercept	4.05985	0.35058	11.58		

	Estimate	SE	T-value	ChiSq	P-value
Temperature	-1.25286	0.37167	-3.371	28.187	**<0.0001**
pH	-0.08916	0.339	-0.263	1.1619	0.2811

Table 8: Statistical outcomes for models of ratios of proportion productive females.

Week 4 Juvenile Sizes

Full Model: glmer(I(area_um2/100) ~ Temp*pH*gammean +(1|FullTrt), family = Gamma(link="inverse"), data = wk4dt)

Interactions	Estimate	SE	T-value	ChiSq	P-value (nested)
gammean:Temp:pH	0.48066	0.14353	3.349	6.3387	**0.01181**

Subset Model for Low pH Treatments

Model: glmer(I(area_um2/100) ~ Temp*gammean +(1|FullTrt), family = Gamma(link="inverse"), data = wk4lph)

Variables	Estimate	SE	T-value	ChiSq	P-value
TrtRep (Dish)	2.93E-03				
Residual	6.53E+01				
Intercept	7.56E+00	1.86766	4.048		
temp:gammean	0.11974	0.08914	1.343	1.6106	0.2044
temp	-5.40444	20.7699	-2.602	10.051	**0.001523**
gammean	-0.03965	0.06543	-0.606	0.3192	0.5721

Subset Model for Ambient pH Treatments

Model: glmer(I(area_um2/100) ~ Temp*gammean +(1|FullTrt), family = Gamma(link="inverse"), data = wk4hph)

Variables	Estimate	SE	T-value	ChiSq	P-value
TrtRep (Dish)	1.33E-03				
Residual	5.21E+01				
Intercept	2.57E+00	1.03042	2.49		
temp:gammean	-0.36249	0.13335	-2.718	5.3001	**0.02132**
temp	3.63827	1.87179	1.944	2.0815	0.1491
gammean	0.10426	0.03703	2.816	3.1812	0.07449

Subset Model for Ambient Temperature Treatments

Model: glmer(I(area_um2/100) ~ pH*gammean +(1|FullTrt), family = Gamma(link="inverse"), data = wk4ltemp)

Variables	Estimate	SE	T-value	ChiSq	P-value
TrtRep (Dish)	4.03E-03				
Residual	4.58E+01				
Intercept	2.62E+00	1.57501	1.665		
pH:gammean	-1.46E-01	0.11731	-1.246	2.2666	0.1322
pH	5.05E+00	3.32134	1.521	3.955	**0.04673**
gammean	1.04E-01	0.05653	1.837	1.1728	0.2788

Subset Model for High Temperature Treatments

Model: glmer(I(area_um2/100) ~ pH*gammean +(1|FullTrt), family = Gamma(link="inverse"), data = wk4htemp)

Variables	Estimate	SE	T-value	ChiSq	P-value

TrtRep (Dish)	4.86E-04				
Residual	8.07E+01				
Intercept	5.95E+00	1.2176	4.887		
pH:gammean	3.22E-01	0.10705	3.011	6.0391	**0.01399**
pH	-3.90E+00	1.35655	-2.874	0.0811	0.7758
gammean	-2.40E-01	0.09895	-2.427	0.988	0.3202

Table 9: Statistical outcomes for models of juvenile size data (area in um2).