A polymerase ribozyme increases copying fidelity through pyrophosphate-mediated RNA repair

Alexandra D. Kent,[1,2] Lucy L. Yang,[1] Kyle H. Cole,[3] Ivee A. Lee,[2] and Andrej Lupták[1,2,3]*

[1] Department of Chemistry, University of California at Irvine; Irvine, CA 92617

[2] Department of Pharmaceutical Sciences, University of California at Irvine; Irvine, CA 92617

[3] Department of Molecular Biology and Biochemistry, University of California at Irvine; Irvine CA 92617

Abstract: Prior to the emergence of the contemporary biosphere, the first replicating systems are thought to have progressed through an RNA-based stage. Such an evolving world would likely have transferred heritable information during replication using RNA polymerase ribozymes. Though substantial effort has been put forth towards evolving RNA polymerases, many variants suffer from premature termination and low fidelity, resulting in low yields of full-length or active sequences. Replication of longer sequences requires a sufficiently high fidelity to lend an evolutionary advantage to an evolvable system. Here we demonstrate ribozyme-mediated repair of mismatched and damaged RNA sequences. Under conditions of saturating pyrophosphate concentrations, we show that a polymerase ribozyme can repair RNA sequences terminated in a mismatch, a non-extendable 2′-3′ cyclic phosphate, or both, to generate a triphosphorylated nucleotide. This repair step increases the fidelity and allows polymerization along an extended template, including the ribozyme itself. This increase of copying fidelity advances the longstanding goal of developing a self-replicating polymerase ribozyme.

 Main Text

Because the cellular functions of ribonucleic acids (RNAs) include critical roles in catalysis,

translation, and metabolite-dependent regulation of gene expression, RNAs may have been the

first biopolymers to store information and catalyze chemical reactions on an early Earth.[1–5] The

replication process is fundamental to life and is the mechanism by which heritable information is

stored and propagated. In the extant biosphere, this is achieved through protein-based

polymerase enzymes, but in a hypothetical RNA World—prior to the advent of genetically

encoded proteins—replication would likely have been carried out by catalytic RNAs termed

polymerase ribozymes.[6,7] These ribozymes could serve as the central molecule linking the

phenotype of an early protocell to its genotype, but sufficient activity and fidelity would be critical

to build up an information storage system.[8,9]

The central reaction catalyzed by enzymes replicating phosphodiester-based informational

polymers is the transfer of the α-phosphate of a triphosphorylated nucleoside to a 3′-hydroxyl of

another nucleotide, with concomitant release of pyrophosphate (Fig. 1). The reverse reaction is

pyrophosphorolysis of polynucleotides, yielding a nucleoside triphosphate and a one-residue

shorter strand. Beginning with the first *in vitro* selection of a ligase ribozyme, a large body of work

using molecular evolution and engineering has sought to isolate ribozymes that catalyze

phosphodiester formation.[10–14] The ligase ribozyme has been evolved into progressively more

sophisticated polymerases capable of promoter-based polymerization, production of other

functional nucleic acids, reverse transcription, and transcription of non-canonical (xeno-) nucleic

acids, among other functions.[15–21] However, most of the advanced polymerase ribozymes suffer

from both mutation burdens that reduce the activity of produced transcripts and low yields of full-

length products due to premature termination and degradation.[22–24] Recent reports demonstrate

that evolving the polymerase ribozymes further under selection pressure to produce active

ribozyme sequences increased the fidelity enough to enable successive rounds of amplification

of an active, short self-cleaving ribozyme.[25] Even with these advances, replicating active

polymerase ribozymes remains an unsolved challenge, highlighting a need for novel approaches

to increasing replication fidelity. An early process for RNA repair could serve to increase

replication fidelity, but very few mechanisms of RNA repair have been reported to date and none

have been catalyzed by ribozymes. A mechanism for repairing both degraded and mismatched

sequences would lend an evolutionary advantage to an evolving system and is an unexplored

approach to increasing polymerase ribozyme fidelity.[26–28]

We demonstrate that ribozyme-catalyzed pyrophosphorolysis of mismatched RNA sequences

yields a nucleoside triphosphate, repairs the mutation, and increases the fidelity of the subsequent

extension. Using sequencing analysis of mismatched and repaired RNAs, we show that repair of

terminal mismatches improves downstream replication fidelity. Sequencing of RNAs generated

from the polymerization of long templates reveal an increase in overall length and fidelity in the

presence of pyrophosphate, and even include copies of the polymerase ribozyme. We further

demonstrate that addition of pyrophosphate and the polymerase ribozyme to 2′-3′ cyclic

phosphate-terminated (damaged) RNA sequences also facilitate damage repair and subsequent

RNA polymerization. Thus, we present a mechanism for increasing the copying fidelity of long

RNAs by a single polymerase ribozyme bringing us one step closer to a fully self-replicating RNA

system.

Results

Pyrophosphorolysis   Extension

Figure 1. Reversible RNA synthesis by the polymerase ribozyme. The RNA polymerase ribozyme,
tC19Z, catalyzes templated addition of a nucleotide to the 3′-end of a primer–template RNA with
pyrophosphate (PPi) as a byproduct of the reaction. The reverse reaction, pyrophosphorolysis of RNA,
removes the last nucleotide, yielding a nucleoside triphosphate. Dashed line indicates a linkage
between the primer and the template, forming a single-molecule RNA hairpin used in this study.

One of the higher-fidelity RNA polymerases ribozymes, tC19Z, was selected *in vitro* from an

earlier polymerase (R18), which in turn originated from the class I RNA ligase [16,18]. The tC19Z

sequence was evolved to have improved sequence generality and polymerase activity, although

the template sequence also has a significant influence on replication efficiency. To extend a

sequence, the 5′ end of the ribozyme base-pairs with the 5′ end of the template sequence and

extends a primer (Fig. 1). Extension of the primer generates pyrophosphate, a byproduct of

polymerization that has also been suggested as a source of available phosphate on an early

Earth, which would have been critical due to the role phosphate plays in energy metabolism and

genome replication.[29–33] Inspired by a previous study that proposed pyrophosphorolysis was

possible with a class I ligase derived ribozyme, we hypothesized that addition of pyrophosphate

to the polymerization reaction would shift the equilibrium of the reaction and induce a low level of

pyrophosphorolysis (Fig. 1) that could increase replication fidelity by selectively removing

mismatched and damaged RNA.[34]

Pyrophosphate-mediated mismatch repair

To maintain fidelity of genome copying at non-deleterious levels, modern replication machinery

uses distinct active sites or entire protein complexes to correct mismatched base-pairs, but at the

onset of evolvable systems, it is unlikely that mismatch repair was available as an independent

biochemical activity.[35,36] Thus, a simpler process for increasing replication fidelity would have

been a desirable feature of an early replicating polymerase. To test whether ribozyme-mediated

pyrophosphorolysis removes a mismatched nucleotide, we prepared hairpin sequences with 3′-

terminal mismatches. We fused the primer and template to form a hairpin (Fig. 1) that serves both

as a substrate for the tC19Z ribozyme at exact stoichiometry and as a practical construct for

downstream analysis using high-throughput sequencing. We generated the hairpins *via* an *in vitro*

transcription reaction of a self-cleaving HDV-like ribozyme (drz-Mbac-1) fused to the 3′ terminus

of the target sequence.[37] The ribozyme self-cleaves, yielding a 2′-3′-cyclic phosphate (2′-3′-cP)

on the 3′ end of the primer segment. The cyclic phosphate can be removed enzymatically by T4

polynucleotide kinase (PNK) (Extended Data Fig. 1).[38,39]

We began by characterizing the reactivity of a terminal A/C mismatch in the primer/template

sequence with the cyclic phosphate removed from the 3′ end. RNA polymerization by the

ribozyme was not observed upon addition of nucleoside triphosphates (NTPs), indicating that an

A/C mismatch acts as a chain-terminator (Fig. 2a,b). After addition of saturating concentrations

of pyrophosphate (10 mM, with clearly visible Mg-PPi precipitate) and NTPs, we observed

ribozyme-mediated extension of the sequence (Fig. 2), but we did not observe the -1 product by

denaturing polyacrylamide gel electrophoresis (PAGE), likely because the equilibrium highly

favors polymerization (SI Note 1, Fig. 2b, and Extended Data Fig. 2). When removed through

pyrophosphorolysis, the terminal nucleotide is converted into a nucleoside triphosphate, and an

A/C-terminated hairpin would produce a molecule of ATP. To obtain independent evidence of

pyrophosphorolysis, we detected this critical metabolite using an ultrasensitive firefly-luciferase-

based assay, which yields a photon for each ATP molecule present in the reaction.[40] Because

pyrophosphate inhibits the luciferase reaction, insoluble pyrophosphate was first removed from

the samples by centrifugation, and the remaining pyrophosphate was subsequently hydrolyzed

by addition of pyrophosphatase enzyme. The resulting samples were incubated with the firefly

luciferase and D-luciferin and were then imaged using an ultracooled EM-CCD camera. The

integrated luminescence was measured for both the pyrophosphorolysis reaction and control

reactions lacking one of the components. In the absence of either pyrophosphate, the ribozyme,

or the hairpin substrate, we observed a baseline level of luminescence. In reactions where all

components were present, the luminescence was significantly greater, indicating pyrophosphate-

mediated synthesis of ATP (Fig. 2c and Extended Data Fig. 3). These experiments provide

evidence that the mechanism of repair occurs by a pyrophosphorolysis reaction and demonstrate

the first instance of ribozyme-catalyzed production of ATP.

To further analyze the repair reaction, we used high-throughput sequencing (HTS) of the

mismatched-terminated hairpin reactions (Fig. 2d,e). HTS analysis revealed that

pyrophosphorolysis of the substrate yields approximately 100-times higher concentrations of a

hairpin lacking the terminal nucleotide than the no-pyrophosphate controls (Extended Data Fig.

4), providing experimental confirmation for the pyrophosphorolysis of the terminal nucleotide.

We repeated this analysis for each of the eleven other possible combinations of terminal

mismatches (Fig. 2 and Extended Data Figs. 2, 4, and 5). Sequence analysis of the extension

reactions correlated well with the gel electrophoresis analysis. In addition, the HTS data provided

a deeper insight into the reactions, revealing the fractions of the sequences that were repaired

and subsequently extended. PAGE analysis was consistent over replicate experiments. We found

that the efficiency of repair and extension depends on the identity of the mismatch:

purine/pyrimidine mismatches were repaired the most efficiently, with the exception of the U/G

wobble. Purine/purine mismatches were repaired and extended at a low frequency upon the

addition of pyrophosphate but were not extendable by the ribozyme in the absence of

pyrophosphate (Fig. 2f,g and Extended Data Fig. 5). In contrast, pyrimidine/pyrimidine

mismatches were extended well by the ribozyme, but most of the extension products retained the

mismatch (Fig. 2f,g and Extended Data Fig. 5). Sequences containing terminal C/A or G/U

mismatches behave like the A/C mismatch and only showed extension (with varying efficiencies)

after pyrophosphorolysis of the mismatch.

In addition to PAGE analyses, we obtained quantitative results for the fidelity of extension

products from the sequencing data. We measured the fidelity at each position of the extension

and found that for all mismatches, repair of the mismatched base led to an increase in the fidelity

of downstream extensions (Fig. 2f,g). In some cases, such as the pyrimidine/pyrimidine

mismatches, the effect is strongly pronounced: although the frequency of repair is low, when a

hairpin is repaired, the extension occurs with high fidelity. Specifically, in the case of the U/C

mismatch, extension of the terminal mismatch preferentially adds a second mismatch (C/A) which

then acts as a chain terminator (Extended Data Fig. 6). Therefore, we conclude that mismatched

sequences that do not terminate the extension are more likely to acquire additional mutations, as

has been observed in enzyme-free polymerization of nucleic acid sequences, implying a

mechanism for error propagation that must involve an allosteric modulation of the fidelity of

incoming nucleotide incorporation by upstream mismatches.[41]

We observed that addition of pyrophosphate inhibited the forward polymerization reaction in the

positive control (Supplementary Fig. 1). To quantify the pyrophosphate reaction inhibition, we

used a non-hairpin, primer-template complex to measure reaction kinetics of the forward

extension reaction. Initial rates were plotted as a function of substrate concentration and revealed

an approximate K_M of 4 mM with respect to NTPs (Extended Data Fig. 7A,B,C). We then

measured the reaction kinetics in the presence of pyrophosphate and plotted the initial rates of

tC19Z-mediated polymerization as a function of pyrophosphate concentration (Extended Data

Fig. 7D,E). At millimolar PPi concentrations, the forward reaction was inhibited approximately five-

fold, compared to when no PPi is present.

Figure 2. Pyrophosphorolysis-mediated mismatch repair and ATP generation by a polymerase ribozyme. (a) Graphical representation of a 3′ terminal mismatch (shown in red) undergoing ribozyme-catalyzed pyrophosphorolysis and subsequent repair. (b) High-resolution PAGE analysis of a ^{32}P-labeled A/C mismatched hairpin. The left two lanes show no extension of the hairpin in the absence of nucleotides or pyrophosphate. Upon addition of pyrophosphate, the sequence can be extended (right lane). (c) ATP generated by the removal of the terminal AMP of an A/C-mismatched hairpin. Reactions lacking either the polymerase ribozyme or pyrophosphate show only background levels of luminescence by firefly luciferase. Open and filled circles represent two independent RNA preparations. Gray lines link experiments performed side-by-side from the same component stocks. Representative luminescence image is shown below. (d) Distribution of sequences extended by the ribozyme in the presence of pyrophosphate (mismatched and repaired), determined by HTS. (e) HTS analysis of sequences of the A/C-mismatched hairpin after incubation with the polymerase ribozyme and PPi. The sequences show the starting A/C mismatch (red A), the removal of the mismatched nucleotide by phosphorolysis (second row sequence, comprising 5.5% of total hairpin reads), the addition of the correct nucleotide (blue G at position 0), and the extension of the corrected sequence (light-blue boxes at positions 1–5). Total percentage of sequences extended are shown in d. (f) Fraction of each mismatch repaired through ribozyme-catalyzed pyrophosphorolysis, grouped by base-pair type. (g) Fidelity of extensions, comparing the fractions of the correctly extended sequences in sequences containing a mismatch (black bars) and the correctly extended sequences that are extended after mismatch repair (gray bars), indicating the fidelity of all sequences that follow the mismatch-repaired position. In all cases, pyrophosphate-mediated mismatch repair leads to an increase in the fidelity of extended sequences. The large differences between the dark and light bars imply that unrepaired mismatches lead to further errors in RNA polymerization.

Addition of pyrophosphate to extension reactions of a long template results in increased length and fidelity

Based on our demonstration that a polymerase ribozyme can repair hairpin sequences terminating in a mismatch and the mechanism for that repair, we hypothesized that pyrophosphate-mediated repair would suppress mutations in long extension reactions of primers, leading to increased fidelity in the polymerization of long templates. To do this, we used a primer–template system that can result in a templated polymerization of 118 nucleotides (Fig. 3a). We performed the ribozyme–catalyzed primer extension reactions for either three hours or four days at varying concentrations of pyrophosphate, and the RNAs were analyzed using HTS. As the concentration of pyrophosphate increased, we observed an increase in the average length of the primer extensions, although the total number of sequences decreased overall (Fig. 3b, Supplementary Table 3), consistent with pyrophosphate inhibition of polymerization. As hypothesized, the sequencing data showed an increase in fidelity at saturating concentrations of

pyrophosphate (Fig. 3c,d). Further analysis revealed that many sequences resulting from early

termination had mismatches at the 3′-termini, suggesting that the mismatches caused the

polymerization to abort (Supplementary Fig. 2). We observed a negative correlation between the

number of sequences and their length in the three-hour experiment, suggesting a trade-off

between copying fidelity and the number of extension products. We also observed complements

of tC19Z sequences in the HTS data with strand-specific ligation products from sequencing library

preparation, indicating that the sequences originated from off-target priming and extension of the

polymerase ribozyme itself. Nearly full-length (98%) ribozyme sequences were present in

samples containing pyrophosphate and exhibited higher fidelity, whereas no long-read RNA

segments antisense to the ribozyme were detected in the no-pyrophosphate HTS dataset

(Supplementary Fig. 3).

Figure 3. Effect of pyrophosphate on RNA polymerization along a long template. (a) The primer–
template construct used for long extension experiments. (b) Plots showing the length of sequences
resulting from polymerase-mediated extension of the primer at varying concentrations of
pyrophosphate at either three hours or four days. Black horizontal bar indicates average length of
sequences. FL dashed line indicates full length sequences. "% Ext" shows the percentage of
sequences that were extended from primers, and % FL indicates the percentage of extended
sequences that are full length. (c,d) Graphs showing the overall fidelity of extension products at varying
concentrations of pyrophosphate at either three hours or four days. All analyses are based on HTS
data. At ~10 mM, Mg-PPi is above saturating concentration and is partly precipitated.

Pyrophosphate-mediated repair of 2′-3′-cyclic-phosphate-terminated RNA sequences

When the phosphodiester of the RNA backbone undergoes nucleophilic attack by an adjacent 2′

oxyanion, the RNA strand is cleaved, generating sequences terminated with a 2′-3′-cyclic

phosphate (2′-3′-cP); this damage is the most common type of RNA degradation. This cyclization

reaction also occurs in self-cleaving ribozymes, many RNases, and base-catalyzed RNA

degradation,[42–44] and is responsible for the relatively low stability of RNA, when compared with

DNA or other nucleic acids that lack a phosphate-adjacent hydroxyl group. Because the 2′-3′-cP

acts as a chain terminator and prevents the 3′ hydroxyl from participating in extension during

polymerization, it represents a replication dead end (Fig. 4a,b). Thus, we tested the polymerase

ribozyme for pyrophosphate-mediated damage repair of an RNA hairpin terminated with a 2′-3′-

cP. The cyclic phosphate can be removed enzymatically by T4 polynucleotide kinase (PNK),

which we used as a positive control for testing the extension efficiencies of the hairpin substrates

(Extended Data Fig. 1).[38,39]

We prepared RNA hairpins terminated with each of the four canonical nucleotides base-paired to

the matching nucleotide on the template strand and terminated with a 2′-3′-cP. As expected, no

polymerization by the ribozyme was observed when extension of the 2′-3′-cP-damaged sequence

was attempted. However, upon addition of pyrophosphate, the terminal damaged nucleotide was

removed, and each of the sequences could be extended by the polymerase ribozyme (Fig. 4c-g,

Extended Data Fig. 8–9, and Supplementary Fig. 4). The sequences could also be extended after

removal of the 2′-3′-cP by PNK. Both extension and repair were largely dependent on the

sequence, with the U/A hairpin undergoing repair the most efficiently.

We hypothesized that the mechanism of damage repair proceeds through pyrophosphorolysis of

the 3′ terminal 2′-3′-cP-damaged nucleotide, as described earlier in the context of mismatch

repair. Therefore, we analyzed the pyrophosphate-mediated removal of the terminal nucleotide

by high-resolution PAGE but did not observe a gel mobility change that could be unequivocally

attributed to removal of a single nucleotide from the 2′-3′-cP-terminated hairpin (see SI note 1).

To better understand pyrophosphate-mediated repair of damaged RNAs, we began by performing

repair experiments that could indirectly indicate the removal of the terminal nucleotide. First, we

omitted the rNTP that corresponded to the damaged nucleotide in the extension reaction. For

example, in the case of an A/U hairpin, we omitted ATP from the rNTPs mixture, because following

excision of the terminal AMP-2′-3′-cP, the ribozyme would need to replace it with undamaged

AMP to continue the extension reaction. As expected, we found that if ATP was not present in the

reaction, the sequence was not extended (Supplementary Fig. 5). To gather additional evidence

of damage repair, we transcribed hairpins without ^{32}P and performed the polymerization reaction

with α-^{32}P-NTPs to incorporate the ^{32}P tracer. We observed that the α-^{32}P-NTP is only utilized in

sequences that have been repaired, and although the ribozyme can incorporate it as a

mismatched nucleotide during the extension, this occurs with lower efficiency (Supplementary

Fig. 6). Finally, we observed no reaction in the presence of only uridine and pyrophosphate,

indicating that a non-canonical reaction, such as opening of the 2′-3′-cP by pyrophosphate to form

a triphosphate that can subsequently react with unphosphorylated uridine, did not take place

(Supplementary Fig. 7).

Figure 4. Pyrophosphate-facilitated repair of an RNA damage site by a polymerase ribozyme. (a) A hairpin substrate terminated with a 2′-3′-cyclic phosphate (2′-3′-cP; red nucleotide) cannot be extended by the polymerase ribozyme. However, upon removal of the terminal adenosine by ribozyme-mediated pyrophosphorolysis, the revealed 3′-OH becomes available for further addition of a nucleoside (blue) from a "correct" NTP. (b) Graphical representation of a hairpin with 2′-3′-cP damage and subsequent repair *via* pyrophosphorolysis. (c) Full sequence of an A/U hairpin and aligned reads from high-throughput sequencing analysis showing removal of the terminal adenosine (dark blue), and subsequent extension up to five bases (light blue). Percentages on the right indicate the fraction of each sequence in the dataset. The numbering above the alignment represents terminal nucleotides defined in a. (d-g) Analysis of ^{32}P-ATP-labeled 2′-3′-cP-damaged hairpins terminated in all four base-pairs; terminal base-pairs are indicated alongside each PAGE analysis (left-hand image of each set). Graphs on the right of each panel show the percentage of each sequence as determined from HTS

281 data, starting with sequences in which the damaged terminal nucleotide has been removed (fraction of
282 sequences ending at position -1) and proceeding to extensions of the repaired RNA (0–5). The starting
283 RNA containing the 2′-3′-cP cannot be ligated to the primer used for sequencing and is therefore not
284 included in this analysis.

285 Repair of both cyclic-phosphate damage and mismatches

For certain hairpins, we observed both damage and mismatch repair upon addition of

pyrophosphate. Specifically, the purine/purine mismatches (A/A, G/A, and A/G) and the G·U

wobble pairs (G/U and U/G) could be repaired by the ribozyme when pyrophosphate was added

to the extension reaction of these 2′-3′-cP-terminated mismatches (Extended Data Fig. 10). These

reactions exemplify a repair and copying of RNA strands that were mismatched, cleaved, or

degraded and formed imperfect primer-template pairs either in *cis* or in *trans*, enabling replication

of novel segments or duplication of existing segments as a pathway in genetic innovation.

293 Discussion

Modern biological systems possess distinct complexes and pathways dedicated to repairing

different types of genome damage that could lead to degradation of the genetic information. Early

replicating systems in an RNA world would not have had those functionalities or enzymatic

diversity available to them, but damage and mismatch repair would likely have been necessary

to replicate long genomes. It has been widely hypothesized that for known polymerase ribozymes

to truly achieve efficient self-replication capability, an increase in the current standard of fidelity

of replication is necessary. Though early studies on the first polymerase ribozymes hypothesized

that pyrophosphorolysis could improve fidelity, this reactivity was not detected.[34] We address this

gap by demonstrating repair activity in the presence of pyrophosphate by an evolved polymerase

ribozyme originally selected to extend a primer along a template in an effort to generate a system

of self-sustained replication.[18] We show that the tC19Z polymerase ribozyme can repair

mismatched sequences, which presents a possible solution to the fidelity problems faced by early

replication ribozymes: in a reaction with a long extension product, we observed an increase in the

average length of sequences and an overall higher fidelity in the presence of millimolar

pyrophosphate. In addition, tC19Z can use pyrophosphate to repair 2′-3′-cyclic-phosphate-

damaged RNA sequences that are formed *via* base-catalyzed RNA cleavage. Although many

ribozymes exhibit varying levels of replication efficiency, under saturating pyrophosphate

conditions, tC19Z is the first ribozyme with a demonstrated capability of repairing RNA.

Although our findings established that tC19Z can repair terminal mismatched sequences, the

extent of ribozyme-mediated repair is highly dependent on mismatch identity, and distinct trends

are present. Previously published findings suggested that less-evolved polymerase ribozymes

(more closely resembling the class I ligase) also exhibit nucleotide preferences and add

mismatches or extend mismatches with varying efficiencies.[15] Our observations on single

nucleotide incorporation and patterns of extended sequences show that tC19Z has preferences

for some nucleotides and sequences over others. For example, the ribozyme will add to uridine

nucleotides in preference to other nucleotides, and similarly will repair damaged uridine

nucleotides with greater frequency. Based on the mismatch repair trends, we hypothesize that

there is a structural foundation for the trends in ribozyme-mediated extension of mismatched

sequences and the effect of pyrophosphate on extension. Specifically, the width of the terminal

base-pair appears to affect the recognition and polymerization capabilities of the ribozyme [45–49].

Although a recent cryo-EM structure has provided new insight into the structural basis for the

function of polymerase ribozymes, additional structural data for this ribozyme-hairpin RNA

complex are needed to definitively conclude what the ramifications of mismatches are on the

ribozyme's polymerization reactivity.[50]

The relationship of sequence length and replication fidelity was originally outlined by Eigen,

stating that as conserved sequence length increases so must replication fidelity to maintain

functionality over successive rounds of copying, and hypothesizing that a mechanism of repair

would be critical in an evolving world.[51,52] Our sequencing analysis reveals a connection between

downstream fidelity and mismatch repair: a substantial increase the downstream fidelity of

extended sequences is observed when mismatches are corrected. Some mismatches negatively

affect the fidelity of downstream polymerization more prominently than others, indicating that if

the ribozyme inserts one incorrect base, it is more likely to add additional mismatches. For

example, if the U/C mismatched hairpin is not repaired, over 60% of subsequently extended

sequences have a second mismatch in the +1 position, and the majority involve addition of a

cytosine to make a C/A mismatch. Furthermore, in the long extension experiments, shorter

sequences present in the data exhibited a varying number of mismatches at their 3′-end.

Ultimately, compounding downstream mismatches leads to termination of replication, which

would result in genome replication catastrophe in an evolving world. By improving fidelity through

a ribozyme-mediated repair mechanism, we show a new functionality accessible to an evolving

system that can directly address the theoretical limits on replicating longer sequences that

maintain their function.

Our study establishes that a single ribozyme can perform several distinct biochemical functions,

likely with the same active site.[16,19,23] Such economy of function may be a hallmark of emerging

genome-encoded functional macromolecules before dedicated synthesis and repair machinery

evolved. These results bring us one step closer to generating truly self-replicating systems that

could have evolved in an RNA world.

Materials and Methods

Cloning tC19Z into a pUC19 vector.

The insert, tC19Z, was synthesized and obtained from Integrated DNA Technologies and then was subsequently amplified with primers containing the restriction enzyme sites BamHI and EcoRI (all enzymes were obtained from New England BioLabs) (**Supplementary Table 1**; sequence #1 is the ribozyme construct).[18] The tC19Z insert and pUC19 vector were double digested with BamHI and EcoRI restriction enzymes for 4 hours. Additionally, the pUC19 vector was further treated with calf-intestinal, alkaline phosphatase (CIP) for 1 hour following the double digestion. The insert was ligated into the vector using T4 DNA ligase and incubating at 16 ºC for 16 hours. The resulting ligation was transformed into DH5α competent cells and subsequently plated on ampicillin-containing agar plates which were grown at 37 ºC for 14 hours. Single colonies were picked and grown in 3 mL of ampicillin containing LB broth for 14 hours. Cultures were subjected to alkaline lysis and the plasmids were purified. The presence of tC19Z in the vector was confirmed by Sanger sequencing. Future polymerase chain reactions (PCR) of tC19Z were performed fresh from the plasmid for each batch of ribozyme.

PCR and transcription of tC19Z and generation of hairpin constructs.

PCR was performed to isolate tC19Z from the plasmid prepared above. The plasmid (5 ng), each primer (0.5 µM) (**Supplementary Table 1**; primers # 4 and 5), and *Taq* DNA polymerase were used to amplify the sequence at an annealing temperature of 49 ºC. PCR products of tC19Z from the plasmid prepared above were used for *in vitro* transcription. Transcription was performed under the following conditions: 50 mM Tris HCl, 2 mM spermidine, 12 mM MgCl$_2$, 8 mM DTT, 2 mM each rNTP, and 0.01% Triton X-100 in a total volume of 100 µL. T7 RNA polymerase was added to the reaction and allowed to react at 37 ºC for 3 hours. The transcriptions were desalted with G15 Sephadex and purified using PAGE [18% w/v acrylamide/bisacrylamide 19:1 solution,

7 M urea, and 0.5x TBE Buffer (50 mM tris(hydroxymethyl)aminomethane hydrochloride, 45 mM

boric acid, and 0.5 mM EDTA, pH 8.4)]. RNA was detected by UV shadowing and then was (i)

eluted in 300 µL of 300 mM KCl from gel pieces, (ii) precipitated by adding 700 µL of ethanol, and

(iii) incubated at –20 ºC overnight. The samples were then centrifuged to pellet the precipitate,

which was subsequently dried in air and resuspended in water.

To prepare the RNA hairpin substrates for the tC19Z polymerase ribozyme

(**Supplementary Table 1**), the hairpin sequences (0.5 µM primers #7–#22; **Supplementary**

Table 1) were generated as fusions with the drz-Mbac-1 HDV-like ribozyme (0.5 µM; #6 construct;

Supplementary Table 1) using primer extension.[37] Following amplification, the sequence were

transcribed in the presence of α-[32]P[ATP] (PerkinElmer) by adding 50 mM Tris HCl, 2 mM

spermidine, 12 mM $MgCl_2$, 8 mM DTT, 2 mM GTP, 2 mM CTP, 2 mM UTP, 200 µM ATP, and

0.01% Triton X-100 to a 50 µL reaction containing T7 RNA polymerase, and the mixture was

allowed to react at 37 ºC for 2 hours. The HDV-like ribozyme self-cleaves co-transcriptionally to

yield cyclic-phosphate-terminated hairpins. The transcriptions were desalted with Sephadex G10

spin columns (10 mM Tris, 1 mM EDTA, pH 7.2) purified using PAGE, and worked up as described

above to generate internally [32]P-labeled polymerase substrates.

Adding [32]P-phosphate to an RNA primer.

An RNA primer 10 nucleotides in length was obtained from Dharmacon (primer #2;

Supplementary Table 1). A [32]P radiolabel was added to the 5′-end of the sequence by adding

the RNA primer at a concentration of 5 µM to a solution containing γ-[32]P[ATP], 1x T4

polynucleotide kinase buffer, and 5 units T4 polynucleotide kinase. The reaction was incubated

at 37 °C for 2 hours, purified using PAGE, and worked up as described above.

Kinetic analysis of pyrophosphate inhibition of RNA primer extension.

Kinetic data for pyrophosphate inhibition were obtained by monitoring the extension of a

^{32}P-labeled primer by tC19Z over time and varying concentrations of pyrophosphate. Equimolar

concentrations (0.5 µM) of tC19Z, primer, and template were mixed and heated to 82 ºC for 2

minutes, then cooled slowly to 17 ºC for 10 minutes. Varying concentrations of pyrophosphate

were then added to the reactions followed by buffer containing 200 mM MgCl$_2$, 50 mM Tris HCl

buffer (pH 8.3), and 4 mM rNTPs, and the solution was then allowed to incubate at 17 ºC. Aliquots

(5 µL) were removed at various time points and were quenched with 250 mM EDTA and 7 M urea

gel-loading buffer. The reactions were analyzed using an 18% denaturing PAGE to obtain single-

nucleotide-resolution images. The gels were imaged on a Typhoon 9410 Variable Mode Imager,

and the pixel density of the bands was analyzed using ImageJ to generate the kinetic data. Initial

rates were calculated using the first four time points and used to demonstrate pyrophosphate

inhibition.

<u>Repair of cyclic phosphate damaged RNA hairpins.</u>

Hairpin substrate repair was visualized and quantified by imaging the extension of an

internally ^{32}P-labeled hairpin by tC19Z. For negative controls, equimolar concentrations (0.5 µM)

of tC19Z and 2′-3′-cyclic-phosphate-terminated hairpins were mixed and heated to 82 ºC for 2

minutes, and then cooled slowly to 17 ºC for 10 minutes. At this point, one of two buffers was

added (1st buffer: 200 mM MgCl$_2$ and 50 mM Tris buffer; 2nd buffer: 200 mM MgCl$_2$, 50 mM Tris

buffer, and 4 mM rNTPs), and the reactions were incubated at 17 ºC for 24 hours. The reactions

were quenched with 250 mM EDTA and 7 M urea gel-loading buffer, and then were analyzed on

a 15% PAGE gel. For the repair reaction, equimolar concentrations (0.5 µM) of tC19Z and cyclic-

phosphate-terminated hairpin were mixed and heated to 82 ºC for 2 minutes, followed by slow

cooling to 17 ºC for 10 minutes and subsequent addition of 10 mM pyrophosphate. A buffer

containing 200 mM MgCl$_2$, 50 mM Tris buffer, and 4 mM rNTPs was added, and the reaction was

allowed to incubate at 17 ºC for 24 hours. The reactions were quenched with 250 mM EDTA and

7 M urea gel-loading buffer and were analyzed on a 15% denaturing polyacrylamide gel. To

generate a positive control sample, the cyclic phosphate was removed from the hairpin by treating

with polynucleotide kinase in 1x PNK buffer (NEB) for 1 hour at 37 °C then inactivated at 65 °C

for 10 minutes. The PNK-repaired hairpin substrate was then added to the tC19Z polymerase

ribozyme, both at concentrations of 0.5 µM, and the solution was heated to 82 °C for 2 minutes

and then cooled slowly to 17 °C for 10 minutes. Next, one of two buffers was added (1st buffer:

200 mM $MgCl_2$ and 50 mM Tris buffer; 2nd buffer: 200 mM $MgCl_2$, 50 mM Tris buffer, and 4 mM

rNTPs), and the reactions were allowed to incubate at 17 °C for 24 hours. The reactions were

quenched with 250 mM EDTA and 7 M urea gel-loading buffer and were analyzed on a 15%

denaturing polyacrylamide gel. The gels were imaged on a Typhoon 9410 Variable Mode Imager.

<u>Kinetic analysis of cyclic-phosphate damage repair.</u>

The kinetics of tC19Z-mediated extension of pyrophosphate-repaired hairpin, PNK-

repaired hairpin, and pyrophosphate inhibited PNK-repaired hairpin were obtained under similar

conditions as those described above. Either hairpin or PNK-repaired hairpin was incubated with

tC19Z at 82 °C for 2 minutes and then cooled slowly to 17 °C for 10 minutes. For the reactions

containing pyrophosphate-repaired hairpin and pyrophosphate-inhibited hairpin, pyrophosphate

was added at a concentration of 10 mM. A buffer containing 200 mM $MgCl_2$, 50 mM Tris buffer,

and 4 mM rNTPs was added to each reaction, and the solutions were allowed to incubate at 17

°C. Time points were removed from the reactions in 5 µL aliquots and quenched with 250 mM

EDTA and 7 M urea gel-loading buffer. The reactions were analyzed on a 15% denaturing

polyacrylamide gel that was imaged on a Typhoon 9410 Variable Mode Imager. ImageJ was used

to quantify band density and generate the kinetic data for tC19Z extension of hairpins under

varying pyrophosphate and repair conditions.

<u>Hairpin migration and extension under various conditions.</u>

To better understand hairpin migration and reactivity, an A/U hairpin was either left

untreated or allowed to react with an alkaline phosphatase [5 µM hairpin, 1x Cutsmart buffer

(NEB), and shrimp alkaline phosphatase (NEB)] for 30 mins at 37 °C. The two conditions were

then either left untreated, treated with PNK as described above, or incubated with acid (25 mM

HCl) followed by quenching. Each solution was then analyzed on a 15% denaturing

polyacrylamide gel. Furthermore, each treated hairpin (0.5 µM) was heated with tC19Z (0.5 µM)

for 2 minutes at 85 °C and then cooled to 17 °C over 10 minutes. A buffer containing 200 mM

$MgCl_2$ and 50 mM Tris HCl (and, in some reactions, 4 mM rNTPs) was added. Additionally, 4 mM

uridine and/or 10 mM pyrophosphate were added to some reactions. The mixtures were incubated

at 17 °C for 24 hours followed by quenching with 250 mM EDTA and 7 M urea gel-loading buffer.

The reactions were analyzed on a 15% denaturing polyacrylamide gel that was imaged on a

Typhoon 9410 Variable Mode Imager.

<u>Incorporation of ^{32}P-rNMPs into non-labeled, repaired hairpins.</u>

To further demonstrate repair, an A/U hairpin was transcribed with an α-^{32}P[ATP] or

without (cold). Some of the hairpins were then treated with PNK as described above. The ^{32}P-

labeled hairpin (0.5 µM) was incubated with tC19Z (0.5 µM) at 82 °C for 2 mins and cooled to 17

°C over 10 minutes. A buffer containing 200 mM $MgCl_2$ and 50 mM Tris HCl (and, in some

reactions, 4 mM rNTPs) was added. To some reactions was added either 10 mM pyrophosphate

or α-^{32}P[ATP] and 10 mM pyrophosphate. The cold hairpin was treated in the same way. The

mixtures were incubated at 17 °C for 24 hours followed by quenching with 250 mM EDTA and 7

M urea gel-loading buffer. The reactions were analyzed on a 15% denaturing polyacrylamide gel

that was imaged on a Typhoon 9410 Variable Mode Imager.

A U/A hairpin substrate was transcribed without α-^{32}P-ATP and either left untreated or

allowed to react with PNK as described above. The hairpin (0.5 µM) was heated with tC19Z (0.5

µM) for 2 minutes at 85 °C and then cooled to 17 °C over 10 minutes. A buffer containing 200 mM

MgCl$_2$, 50 mM Tris HCl and, in some reactions, 10 mM pyrophosphate was added. Additionally,

either α -^{32}P[ATP], α -^{32}P[UTP], or α -^{32}P[CTP] was added. The reactions were incubated at 17 ºC

for 24 hours followed by quenching with 250 mM EDTA and 7 M urea gel-loading buffer. The

reactions were analyzed on a 15% denaturing polyacrylamide gel that was imaged on a Typhoon

9410 Variable Mode Imager.

<u>Extension of mismatched RNA hairpin sequences.</u>

Mismatched RNA hairpins were extended by the tC19Z polymerase ribozyme under

conditions of PNK repair or pyrophosphate. Pyrophosphate facilitated extension of both a cyclic-

phosphate-damaged hairpin and a mismatched hairpin when 0.5 µM hairpin was added to 0.5 µM

tC19Z. The solution was incubated at 82 ºC for 2 minutes and then cooled slowly to 17 ºC for 10

minutes. Pyrophosphate was added at a final concentration of 10 mM, followed by a buffer

containing 200 mM MgCl$_2$, 50 mM Tris buffer, and 4 mM rNTPs, and the solution was incubated

at 17 ºC for 24 hours. A reaction containing a PNK-repaired mismatched hairpin and tC19Z at

concentrations of 0.5 µM each was incubated at 82 ºC for 2 minutes and then cooled slowly to 17

ºC for 10 minutes. Pyrophosphate was added at a concentration of 10 mM, followed by a buffer

containing 200 mM MgCl$_2$, 50 mM Tris buffer, and 4 mM rNTPs, and the reaction was allowed to

proceed at 17 ºC for 24 hours. All reactions were quenched with 250 mM EDTA and 7 M urea

gel-loading buffer and then analyzed on a 15% denaturing polyacrylamide gel that was imaged

on a Typhoon 9410 Variable Mode Imager. ImageJ was used to quantify band density to generate

fraction extended graphs.

<u>High-throughput sequencing of RNA hairpins.</u>

RNA hairpins were analyzed by Illumina Sequencing to determine sequence identity. Cyclic

phosphate repair and mismatch extension reactions, as described above, were performed and

fractionated by PAGE. The images obtained were analyzed for extension products, and the

corresponding bands were excised from the gel, eluted, and precipitated with ethanol. The RNA

sequences were resuspended in water and then ligated to a DNA sequence containing an Illumina

forward adapter (#24 and 25, **Supplementary Table 1**). The DNA adapter sequence was

designed with a 5′-phosphate, followed by four random nucleotides to reduce ligation bias at the

3′-RNA terminus. The adapter also contained a 3′-amine to prevent self-ligation. The oligo (10

µM) was added to a mixture containing PEG 8000, 1x T4 RNA ligase buffer, 15 µM ATP, and T4

RNA ligase 1 to a total volume of 100 µL to adenylylate the 5′-end of the DNA sequence (5′-App

oligo). G15 Sephadex was used to desalt and remove residual ATP from the 5′-App oligo.

To anneal the sequences, the 5′-App oligo was added in approximately three times excess

of the RNA hairpin, and the resulting solution was heated to 95 ºC for 3 mins and then cooled to

25 ºC at a rate of 1 ºC/min. To ligate, the annealed oligos were added to 1x T4 RNA ligase buffer,

PEG 8000, and T4 RNA ligase 2, truncated KQ to a total volume of 20 µL, and allowed to react

at 25 ºC for 18 hours and then at 4 ºC for 2 hours.

To a 5 µL aliquot of the ligation reaction was added 2.5 µM of the RT primer (#29 in

Supplementary Table 1), which was annealed at 75 ºC for 3 mins, 37 ºC for 10 mins, and 25 ºC

for 10 mins. Following annealing, 1x ProtoScript II buffer, 0.5 mM each dNTP, 5 mM $MgCl_2$, 10

mM DTT, and ProtoScript II were added and the reverse transcription reaction was heated to 50

ºC for 2 hours to generate cDNA.

The Illumina forward adapter was added to the cDNA through a PCR reaction (using

primer #31 in **Supplementary Table 1**). Forward and reverse primers (#31 and #30; 1 µM) were

added to the cDNA, as well as 1x Q5 reaction buffer, 5 mM of each dNTP, and Q5 DNA

polymerase. The reaction was annealed at 52 ºC and extended at 72 ºC. The Illumina barcodes

were added in a second PCR reaction containing 1 µM of each primer, 1x Q5 buffer, 5 mM each

dNTP, and Q5 DNA polymerase. The reaction to install the Illumina barcodes was annealed at 65

ºC and extended at 72 ºC, and the reaction was purified via agarose gel. The libraries were

submitted for Illumina Sequencing. PhiX was removed from the resulting reads using bowtie2,

and the remaining reads were trimmed using CutAdapt to remove adapter regions. The 5′-adapter

(TCTACACTCTTTCCCTACACGACGCTCTTCCGATCT) was removed first, then the 3′-adapter

(GATCGGAAGAGCACACGTCTGAACTCCAGTCAC), with at most 10% error in single-end

mode. Sequences were counted in FastAptamer using fastaptamer_count. Starting and extended

sequences were extracted from data for analysis in FastAptamer using fastaptamer_search for

each sequence with unique identity and length.[53] Specifically, the two-nucleotide tag (AC or GT)

incorporated from sequences #24 or #25, followed by four random N nucleotides, and ending with

the sequence of interest were searched for and compiled. These sequences were then curated

visually in Jalview. Sequence identity was analyzed at each position and the number of

sequences in each case were used to calculate extension efficiency, pyrophosphorolysis, and

fidelity (SI Note 2). The sequencing was performed on one representative reaction for each

condition, as is convention in the field when determining polymerase ribozyme fidelity.

<u>Long-read primer extension and sequencing.</u>

For sequencing the long-read experiments, we prepared an RNA template corresponding

to a sequence previously used with the tC19Z rRNA polymerase ribozyme .[18] The sequence was

transcribed from the long-read DNA template (sequence #23 in **Supplementary Table 1**)

prepared as a double-stranded DNA using the template strand and a T7 RNAP primer-extended

as described above, and PAGE-purified, as described above. The template RNA was combined

with the RNA primer (sequence #2 in **Supplementary Table 1**) and the tC19Z ribozyme at 1 µM

concentration (each) and this mixture was incubated at various concentrations of pyrophosphate

for 4 days at 17 ºC as described above. The reactions were desalted using Sephadex G10. The

desalted samples were not further purified. The samples were then ligated to the ligation primer

for sequencing (sequences #24 and #25 in **Supplementary Table 1**) and reverse-transcribed as

described above. The cDNAs were then ligated to the cDNA ligation hairpin (sequence #26 in

Supplementary Table 1), which contains a 7-nt random sequence to allow splint-ligation by a

DNA ligase independent of the 3′ terminal sequence of the cDNA and allow amplification of all

RNAs present in the samples, as described previously.[54] To suppress self-ligation, the cDNA

ligation hairpin DNAs contained 3′ terminal carbon spacer. The samples were incubated in T4

DNA ligase buffer (NEB) with 1 mM ATP, 0.5 μL of T4 DNA ligase, 0.5 M betaine, 20% PEG 8000,

and 0.5 μL of RNaseH (NEB). To preferentially amplify the ribozyme-extended RNA primers, a

DNA primer that overlapped with the RNA primer sequence (CUGCCAACCG) extended by the

ligated hairpin sequence was designed and used for PCR amplification (sequence #27 in

Supplementary Table 1) together with the RT primer for sequencing (#29 in **Supplementary**

Table 1). A nested PCR reaction was performed to further enrich for sequences of the RNA

primer–template ribozyme extension reaction and introduced the Illumina reverse primer

sequence (#28 in **Supplementary Table 1**) and the reactions were sequenced as described

above. Whereas reactions were designed to bias towards amplification of the RNA primer-

extensions, we observed many other amplicons, including many sequences that corresponded to

products of primer or hairpin self-ligations.

The libraries were sequenced by Illumina Sequencing, and the resulting reads were

trimmed using CutAdapt to remove primer regions. Sequences were counted in FastAptamer

using fastaptamer_count. Sequences containing the primer sequence were extracted from the

data for analysis in FastAptamer using fastaptamer_search for each sequence and subsequently

visually aligned and curated using Jalview. Sequence lengths and fidelity were obtained using

Jalview (SI Note 2). The sequencing was performed on one representative reaction for each

condition, as is convention in the field when determining polymerase ribozyme fidelity.

Measuring ATP concentrations using a firefly luciferase assay.

A solution containing a 2-μM PNK-repaired A/C hairpin and 0.5-μM tC19Z polymerase

ribozyme was incubated at 82 ºC for 2 minutes and then cooled slowly to 17 ºC for 10 minutes.

Pyrophosphate was added at a concentration of 10 mM followed by a buffer containing 200 mM

$MgCl_2$ and 50 mM Tris buffer, and the reaction was allowed to proceed at 17 ºC for 3 days. Control

reactions were incubated as above, but in the absence of individual components, as indicated.

After 3 days, the reactions were centrifuged, and 0.5 µL of the supernatant was added to 4.5 µL

of a solution containing inorganic pyrophosphatase and firefly luciferase (preincubated for 3 h in

presence of D-luciferin to deplete any ATP co-purified with the enzyme). Luminescence was

generated using the StayBrite Highly Stable ATP Bioluminescence Assay Kit (BioVision), as

described previously.[40] The reaction contained 50 µM D-luciferin, 1 µL of 4x StayBrite reaction

buffer, and 1 µL StayBrite luciferase enzyme, in addition to 0.5 µL of inorganic pyrophosphatase

(NEB). Each reaction was loaded in a well of a black 384-well plate and imaged using a high

numerical aperture lens (Voigtländer 29 mm F/0.8) mounted on an EMCCD camera (iXon Ultra

888, Andor). The temperature of the camera electronics was maintained at –100 ºC during

detection using a thermoelectric recirculating chiller (Thermocube 300, Solid State Cooling

Systems). During imaging, 10 one-minute exposures were collected. The images were then

stacked, a Z-projection was generated, and relative luminescence was measured using ImageJ.

To generate a standard curve, different ATP concentrations in nM range were incubated in the

same buffer system, including pyrophosphate. For the no-pyrophosphate control, pyrophosphate

was added to the samples at the end of the experiment and the samples were treated so that the

same amount of soluble pyrophosphate was present in each sample when incubated with the

pyrophosphatase and luciferase enzymes.

References

1.   Leslie E., O. Prebiotic Chemistry and the Origin of the RNA World. *Crit. Rev. Biochem.*
604   *Mol. Biol.* **39**, 99–123 (2004).

605   2.   Robertson, M. P. & Joyce, G. F. The origins of the RNA world. *Cold Spring Harb.*
606   *Perspect. Biol.* **4**, a003608 (2012).

607   3.   Schrum, J. P., Zhu, T. F. & Szostak, J. W. The Origins of Cellular Life. *Cold Spring Harb.*
608   *Perspect. Biol.* **2**, (2010).

4.   Bartel, D. P. & Unrau, P. J. Constructing an RNA world. *Trends Cell Biol.* **9**, 9–13 (1999).

5.   Joyce, G. F. The antiquity of RNA-based evolution. *Nature* **418**, 214–221 (2002).

6. Ma, W. & Yu, C. Intramolecular RNA replicase: Possibly the first self-replicating molecule
in the RNA world. *Orig. Life Evol. Biosph.* **36**, 413–420 (2006).

7. Chen, X., Li, N. & Ellington, A. D. Ribozyme Catalysis of Metabolism in the RNA World.
Chem. Biodivers. **4**, 633–655 (2007).

8. Benner, S. A., Ellington, A. D. & Tauer, A. Modern metabolism as a palimpsest of the
RNA world. *Proc. Natl. Acad. Sci. U. S. A.* **86**, 7054–7058 (1989).

9. Neveu, M., Kim, H.-J. & Benner, S. A. The "Strong" RNA World Hypothesis: Fifty Years
Old. *Astrobiology* **13**, 391–403 (2013).

10. Ekland, E. H., Szostak, J. W. & Bartel, D. P. Structurally complex and highly active RNA
ligases derived from random RNA sequences. *Science* **269**, 364–370 (1995).

11. Bartel, D. P. & Szostak, J. W. Isolation of new ribozymes from a large pool of random
sequences [see comment]. *Science* **261**, 1411–1418 (1993).

12. Ekland, E. H. & Bartel, D. P. The secondary structure and sequence optimization of an
RNA ligase ribozyme. *Nucleic Acids Res.* **23**, 3231–3238 (1995).

13. Jaeger, L., Wright, M. C. & Joyce, G. F. A complex ligase ribozyme evolved in vitro from
a group I ribozyme domain. *Proc. Natl. Acad. Sci.* **96**, 14712–14717 (1999).

14. Shechner, D. M. *et al.* Crystal Structure of the Catalytic Core of an RNA-Polymerase
Ribozyme. *Science* **326**, 1271–1275 (2009).

15. Ekland, E. H. & Bartel, D. P. RNA-catalysed RNA polymerization using nucleoside
triphosphates. *Nature* **382**, 373–376 (1996).

16. Johnston, W. K., Unrau, P. J., Lawrence, M. S., Glasner, M. E. & Bartel, D. P. RNA-
Catalyzed RNA Polymerization: Accurate and General RNA-Templated Primer Extension.
Science **292**, 1319–1325 (2001).

17. Samanta, B. & Joyce, G. F. A reverse transcriptase ribozyme. *Elife* **6**, e31153 (2017).

18. Wochner, A., Attwater, J., Coulson, A. & Holliger, P. Ribozyme-Catalyzed Transcription of
an Active Ribozyme. *Science* **332**, 209–212 (2011).

19. Horning, D. P. & Joyce, G. F. Amplification of RNA by an RNA polymerase ribozyme.
Proc. Natl. Acad. Sci. **113**, 9786–9791 (2016).

20. Horning, D. P., Bala, S., Chaput, J. C. & Joyce, G. F. RNA-Catalyzed Polymerization of
Deoxyribose, Threose, and Arabinose Nucleic Acids. *ACS Synth. Biol.* **8**, 955–961
(2019).

21. Razvan, C. & J., U. P. Processive RNA polymerization and promoter recognition in an
RNA World. *Science* **371**, 1225–1232 (2021).

22. Tjhung, K. F., Shokhirev, M. N., Horning, D. P. & Joyce, G. F. An RNA polymerase
ribozyme that synthesizes its own ancestor. *Proc. Natl. Acad. Sci.* **117**, 2906–2913
(2020).

23. Zaher, H. S. & Unrau, P. J. Selection of an improved RNA polymerase ribozyme with
superior extension and fidelity. *RNA* **13**, 1017–1026 (2007).

24. Attwater, J. *et al.* Chemical fidelity of an RNA polymerase ribozyme. *Chem. Sci.* **4**, 2804–
2814 (2013).

25. Papastavrou, N., Horning, D. P. & Joyce, G. F. RNA-catalyzed evolution of catalytic RNA.
Proc. Natl. Acad. Sci. **121**, e2321592121 (2024).

26. Beate, S., Rana, S., Kiong, H. C. & Stewart, S. Portability and fidelity of RNA-repair
systems. *Proc. Natl. Acad. Sci.* **101**, 2788–2793 (2004).

27. Burroughs, A. M. & Aravind, L. RNA damage in biological conflicts and the diversity of
responding RNA repair systems. *Nucleic Acids Res.* **44**, 8525–8555 (2016).

28. Nandakumar, J., Schwer, B., Schaffrath, R. & Shuman, S. RNA Repair: An Antidote to
Cytotoxic Eukaryal RNA Damage. *Mol. Cell* **31**, 278–286 (2008).

29. Kamerlin, S. C. L., Sharma, P. K., Prasad, R. B. & Warshel, A. Why nature really chose
phosphate. *Q. Rev. Biophys.* **46**, 1–132 (2013).

30. Pasek, M. A. Rethinking early Earth phosphorus geochemistry. *Proc. Natl. Acad. Sci.*
105, 853–858 (2008).

31. Miller, S. L. & Parris, M. Synthesis of pyrophosphate under primitive earth conditions.
Nature **204**, 1248–1250 (1964).

32. Hørder, M. Complex formation of inorganic pyrophosphate with magnesium: The
influence of ionic strength, supporting medium and temperature. *Biochim. Biophys. Acta -*
Enzymol. **358**, 319–328 (1974).

33. Holm, N. G. & Baltscheffsky, H. Links Between Hydrothermal Environments,
Pyrophosphate, Na+, and Early Evolution. *Orig. Life Evol. Biosph.* **41**, 483–493 (2011).

34. Glasner, M. E., Yen, C. C., Ekland, E. H. & Bartel, D. P. Recognition of
Nucleoside Triphosphates during RNA-Catalyzed Primer Extension. *Biochemistry* **39**,
15556–15562 (2000).

35. Li, G.-M. Mechanisms and functions of DNA mismatch repair. *Cell Res.* **18**, 85–98 (2008).

36. Iyer, R. R., Pluciennik, A., Burdett, V. & Modrich, P. L. DNA mismatch repair: functions
and mechanisms. *Chem. Rev.* **106**, 302–323 (2006).

37. Riccitelli, N. J., Delwart, E. & Lupták, A. Identification of Minimal HDV-Like Ribozymes
with Unique Divalent Metal Ion Dependence in the Human Microbiome. *Biochemistry* **53**,
1616–1626 (2014).

38. Wang, L. K., Lima, C. D. & Shuman, S. Structure and mechanism of T4 polynucleotide
kinase: an RNA repair enzyme. *EMBO J.* **21**, 3873–3880 (2002).

39. Das, U. & Shuman, S. Mechanism of RNA 2',3'-cyclic phosphate end healing by T4
polynucleotide kinase-phosphatase. *Nucleic Acids Res.* **41**, 355–365 (2013).

40. Chizzolini, F., Kent, A. D., Passalacqua, L. F. M. & Lupták, A. Enzymatic RNA Production
from NTPs Synthesized from Nucleosides and Trimetaphosphate. *ChemBioChem* **22**,
2098–2101 (2021).

41. Leu, K. *et al.* Cascade of Reduced Speed and Accuracy after Errors in Enzyme-Free
Copying of Nucleic Acid Sequences. *J. Am. Chem. Soc.* **135**, 354–366 (2013).

42. Jimenez, R. M., Polanco, J. A. & Lupták, A. Chemistry and Biology of Self-Cleaving
Ribozymes. *Trends Biochem. Sci.* **40**, 648–661 (2015).

43. Li, Y. & Breaker, R. R. Kinetics of RNA Degradation by Specific Base Catalysis of
Transesterification Involving the 2'-Hydroxyl Group. *J. Am. Chem. Soc.* **121**, 5364–5372
(1999).

44. Seith, D. D., Bingaman, J. L., Veenis, A. J., Button, A. C. & Bevilacqua, P. C. Elucidation
of Catalytic Strategies of Small Nucleolytic Ribozymes from Comparative Analysis of

Active Sites. *ACS Catal.* **8**, 314–327 (2018).

45.   Cruse, W. B. *et al.* Structure of a mispaired RNA double helix at 1.6-A resolution and
implications for the prediction of RNA secondary structure. *Proc. Natl. Acad. Sci.* **91**,
4160–4164 (1994).

46.   Pan, B., Mitra, S. N. & Sundaralingam, M. Crystal Structure of an RNA 16-mer Duplex
R(GCAGAGUUAAAUCUGC)2 with Nonadjacent G(Syn)·A+(Anti) Mispairs. *Biochemistry*
38, 2826–2831 (1999).

47.   Sugimoto, N., Kierzek, R. & Turner, D. H. Sequence dependence for the energetics of
terminal mismatches in ribooligonucleotides. *Biochemistry* **26**, 4559–4562 (1987).

48.   Sydow, J. F. *et al.* Structural basis of transcription: mismatch-specific fidelity mechanisms
and paused RNA polymerase II with frayed RNA. *Mol. Cell* **34**, 710–721 (2009).

49.   Kahn, J. D. & Hearst, J. E. Reversibility of nucleotide incorporation by Escherichia coli
RNA polymerase, and its effect on fidelity. *J. Mol. Biol.* **205**, 291–314 (1989).

50.   McRae, E. K. S. *et al.* Cryo-EM structure and functional landscape of an RNA polymerase
ribozyme. *Proc. Natl. Acad. Sci.* **121**, e2313332121 (2024).

51.   Eigen, M. Selforganization of matter and the evolution of biological macromolecules.
Naturwissenschaften **58**, 465–523 (1971).

52.   Eigen, M., McCaskill, J. & Schuster, P. Molecular quasi-species. *J. Phys. Chem.* **92**,
6881–6891 (1988).

53.   Alam, K. K., Chang J. L. & Burke, D. H. FASTAptamer: A bioinformatic toolkit for high-
throughput sequence analysis of combinatorial selections. *Mol. Ther. Nucleic Acids* **4**,
e230 (2015).

54.   Latifi, B., Cole, K. H., Vu, M. M. K., Lupták, A. Rapid discovery of functional RNA
domains. *Nucleic Acids Res.* **53**, gkaf307 (2025).

Acknowledgements

We thank the staff at the Genomics High-Throughput Facility (GHTF) Shared Resource of the
Cancer Center, Support Grant (P30CA-062203) at the University of California, Irvine and NIH
shared instrumentation grants 1S10RR025496-01, 1S10OD010794-01, and 1S10OD021718-
01; and S. Flynn for assistance with sequencing data. We thank C. Langeberg and D. Treco for
comments on the manuscript, and J. H. D. Cate for his support. This work was supported by the
John Templeton Foundation through a grant to the Foundation for Applied Molecular Evolution
(to A. L.), NSF CBET Biophotonics 1804220 (to A. L.), NASA Exobiology 80NSSC21K0488 (to
A. L.), NASA ICAR 80NSSC21K0596 (to A.L.), NSF GRFP (to A. K.), Goldwater Scholarship (to
L. Y.), and UCI Undergraduate Research Opportunities Program awards (to L. Y. and I. L.).
730

731   **Affiliations**

732
733   *Department of Chemistry, University of California, Irvine, Irvine, CA, USA*
734   Alexandra D. Kent, Lucy L. Yang, and Andrej Lupták
735
736   *Department of Pharmaceutical Sciences, University of California, Irvine, Irvine, CA, USA*
737   Alexandra D. Kent and Andrej Lupták

738
739 *Department of Molecular Biology and Biochemistry, University of California, Irvine, Irvine, CA,*
740 *USA*
741 Kyle H. Cole and Andrej Lupták
742
743 **Current Affiliations**
744
California Institute for Quantitative Biosciences, University of California at Berkeley, Berkeley,
CA, USA
Alexandra D. Kent
Department of Chemistry, Harvard University; Cambridge, MA, USA
Lucy L. Yang
Amgen; Thousand Oaks, CA, USA
Kyle H. Cole
752
753

754 **Author contributions**
755
756 A.D.K. and A.L. conceived and designed the study. A.D.K. performed the experiments with
757 assistance from L.L.Y., K.H.C. and A. L. The data were analyzed by A.D.K. and A.L. The
758 manuscript was prepared by A.D.K and A.L.

759 **Data and materials availability:** All data are available in the main text, the supplementary
760 materials, or deposited in a sequence repository.
761

Extended Data Figures

A/U Hairpin   A/U Hairpin + PNK   A/U Hairpin + Acid

Extended Data Fig. 1. Gel mobility of A/U hairpin. An A/U hairpin has a 2′-3′-cyclic phosphate resulting from drz-Mbac-1 ribozyme self-scission (red and left lane on PAGE gel). After repair with PNK, the cyclic phosphate is removed (blue), and the corresponding band migrates faster on the PAGE gel (middle lane). When the 2′-3′-cyclic-phosphate-terminated hairpin is treated with hydrochloric acid, the cyclic phosphate is partially hydrolyzed, yielding a 3′ monophosphate (green) or 2′ monophosphate (not shown), and the mobility of the corresponding PAGE band is retarded (right lane).

Extended Data Fig. 2. Ribozyme-catalyzed, pyrophosphate-facilitated repair reactions for

all mismatch combinations. All mismatches were ^{32}P-labeled and reacted under standard

conditions with variations indicated with + and - symbols. The reactions were then analyzed by

high-resolution PAGE.

Extended Data Fig. 3. Luciferase-based assay measures ATP produced during ribozyme-

catalyzed pyrophosphorolysis. (A) Firefly luciferase assay measuring the concentration of ATP

in nM generated by the removal of the terminal adenosine monophosphate of an A/C mismatched

hairpin, as calculated by use of a standard curve. Reactions lacking the polymerase ribozyme,

hairpin, or pyrophosphate show only background levels of luminescence. Higher concentrations

of ATP are present in reactions containing all three components. Open and filled circles represent

two independent preparations of the polymerase ribozyme and the hairpin substrate, with filled

circles showing higher background ATP levels apparently co-purified with the polymerase

ribozyme. Lines connect individual experiments performed in parallel on the same day and from

the same stocks. A representative luminescence image is shown below the graph. (B) Additional

luminescence image used to generate the values shown in A and showing additional control

reactions lacking both the ribozyme and pyrophosphate.

Extended Data Fig. 4. Removal of terminal nucleotides in the presence and absence of

pyrophosphate. Addition of pyrophosphate increases the fraction of shorter sequences as

determined by high-throughput sequencing. (A) Matched U/A and C/G hairpins terminated with a

2′-3′-cyclic phosphate and incubated in absence of pyrophosphate show very low fractions of

shorter sequences, indicating low background of these sequences in the samples. (B) Matched

hairpins terminated with a 2′-3′-cyclic phosphate yield a fraction of shorter sequences that is about

2 orders of magnitude higher, particularly in the −1 position, after treatment with ribozyme and

pyrophosphate. These results provide direct evidence for ribozyme-catalyzed pyrophosphorolysis

of RNA. (C) All mismatched sequences following incubation with pyrophosphate and ribozyme.

Some sequences exhibit higher percentages of pyrophosphorolysis than others, largely following

the trends described in Figure 4. To ease comparison, all graphs are plotted on the same scale,

with the horizontal axis placed at 0.01% and a gray dashed line placed at 0.1%.

Extended Data Fig. 5. Extension and repair of mismatches. Mismatched hairpins repaired and extended with distinct efficiencies, as determined by high-throughput sequencing. The left graph of each panel shows the fraction of sequences terminated at each position along the template relative to the starting length (0) after incubation with the polymerase ribozyme, pyrophosphate, and rNTPs. All hairpins were treated with PNK to remove the 2′-3′-cyclic phosphate. "−1" corresponds to the fraction of sequences terminated at nucleotides corresponding to the products of pyrophosphorolysis of the mismatch-terminated hairpin. Hairpins terminated in A/G (B), A/C (D), C/A (E), and U/U (I) show particularly high fractions of sequences with the terminal mismatches removed. The right graph of each panel shows the percentage of each extended segment with the starting mismatch repaired. (A-C) Purine/purine mismatches are repaired and extended at low efficiency, but longer products contain high fraction of repaired sequences. (D–F) Purine/pyrimidine mismatches are extended well and have high percentages of repair, except the U/G wobble pair. For example, a hairpin that started with an A/C mismatch (D), showed more than 15% of sequences extended by one nucleotide (+1, left graph) and ~20 % of these 1-nt extensions had been repaired (+1, right graph). (G–I) Pyrimidine/pyrimidine mismatches are

822   extended with high efficiencies but are not repaired well.

Extended Data Fig. 6. One mismatch can induce addition of more mutations. When a hairpin having a terminal U/C mismatch is incubated with a polymerase ribozyme and tC19Z it can either undergo pyrophosphorolysis or be extended by one nucleotide. In only a low percentage is the mismatch removed, but if it is, the subsequent extension proceeds with high fidelity. If the mismatch is not corrected, the ribozyme will add a second mismatch preferentially which results in chain termination.

Extended Data Fig. 7. Reaction kinetics of primer extension and pyrophosphate inhibition.
(A) The fraction extended of the [32]P-labeled primer is quantified as a function of time at varying NTP concentrations by high resolution PAGE. (B) High-resolution PAGE of the primer extension at 4 mM rNTPs which is plotted on the graph in A. (C) The fraction of primer extended per minute is plotted as a function of NTP concentration and shows an estimated K_M of ~4 mM. (D) The fraction of the [32]P-labeled primer extended quantified as a function of time at varying pyrophosphate conditions. (E) The fraction extended per minute is plotted as a function of the log of pyrophosphate concentration.

Extended Data Fig. 8. Damage repair reactions for all matched hairpins. The general sequence of the hairpins is shown above the PAGE gels, with terminal base-pair shown as N-N in blue and specific terminal base-pairs indicated above each set of experiments. The site of RNA damage, corresponding to the 2′-3′-cyclic phosphate is indicated as >P in red. All hairpins were allowed to react under standard conditions with reaction conditions variations shown by the + and - symbols. Hairpins were analyzed by high-resolution PAGE. The A/U, U/A, and C/G hairpins were all resolved on the same gel, and the G/C hairpin was resolved on a separate PAGE gel. PNK indicates removal of the terminal 2′-3′-cyclic phosphate by the polynucleotide kinase enzyme to yield canonical 2′ and 3′ hydroxyls, which results in faster migration. All hairpins were extendable by the tC19Z polymerase ribozyme when either treated by the PNK enzyme or pyrophosphate (right two lanes of each set of experiments).

Extended Data Fig. 9. Kinetics of RNA polymerization of damage-repaired substrates. A
^{32}P-labeled hairpin terminated with 2′-3′-cyclic phosphate was either treated with PNK or
pyrophosphate and then subjected to standard polymerase-ribozyme extension conditions. Nine
time points were collected and analyzed by high-resolution PAGE. (A) For an A/U hairpin,
triangles indicate the fraction extended of the PNK-repaired hairpin, crosses indicate the fraction
extended of the PNK repaired hairpin in the presence of 10 mM PPi to determine the inhibition of
the reaction by pyrophosphate, and open circles indicate the fraction extended of the 2′-3′-cyclic-
phosphate-terminated hairpin that was repaired by the ribozyme in presence of 10 mM PPi. For
hairpins terminated in G/C (B), U/A (C), or C/G (D), triangles indicate the fraction extended of the
PNK-repaired hairpin, and open circles indicate the fraction extended of the 2′-3′-cyclic-
phosphate-terminated hairpin that was repaired by addition of 10 mM PPi.

Extended Data Fig. 10. Trends in mismatch repair and damage and mismatch repair. (A)

Grid showing the correlation between mismatch repair and overall mismatch extension. The box

colors indicate what percentage of each mismatch is extended when repaired by the ribozyme in

the presence of pyrophosphate, compared to a no-pyrophosphate control, as determined by

PAGE analysis. The gray color indicates no significant difference observed in extension reactions

in the presence/absence of pyrophosphate. The numbers correspond to the total fraction of

mismatch repair derived from HTS data. (B) Grid showing mismatches in which both 2′-3′-cP

phosphate damage and the mismatch could be repaired *via* ribozyme-catalyzed

pyrophosphorolysis of the terminal nucleotide.

Supplementary Information

SI Note 1. Gel migration of cyclic-phosphate–terminated hairpins

Previous reports showed that treatment of 2′-3′-cP-terminated oligonucleotide sequences

with PNK to remove the cyclic phosphate leads to a gel mobility retardation, and treatment of a

2′-3′-cyclic-phosphate-terminated sequence with acid to generate a linear monophosphate leads

to a faster-migrating species.[52–54] We consistently observed the opposite effect and have

concluded that the anomalous migration is a property of the hairpin sequence, which differs

structurally from the linear RNA sequences used in previously reported experiments. Following

treatment with PNK, all 2′-3′-cP-terminated hairpin sequences exhibit faster gel migration. The

relative gel mobility of the 2′-3′-cP-terminated hairpins treated with acid (which initially hydrolyzes

just one of the phosphodiesters of the cyclic phosphate) shows a gel mobility retardation. This

data confirmed that PNK treated hairpin sequences indeed migrate faster upon removal of the 2′-

3′-cP (Extended Data Fig. 1). Many of the hairpin sequences when visualized by PAGE show two

bands depending on the resolution of the experiment. Based on the gel migration of the acid

treated 2′-3′-cP, we determined that these two bands arise from the difference between closed

and open states of the terminal cyclic phosphate.

SI Note 2. Analysis of HTS sequencing data for each relevant figure

Additional details on the HTS analysis for each figure:

All libraries were submitted for Illumina sequencing. See Supplementary Table 2 for explanation

of datasets.

Illumina MiSeq Single Read 150 sequencing was used for tC19Z_MM_1, tC19Z_MM_2,

tC19Z_MM_3, tC19Z_MM_4, tC19Z_MM_5, tC10Z_MM_6, tC19Z_MM_7, tC19Z_MM_8,

tC19Z_MM_9, tC19Z_MM_10, tC19Z_MM_11, tC19Z_MM_12, tC19Z_MA_1, tC19Z_MA_2,

tC19Z_MA_3, tC19Z_MA_4, tC19Z_MA_NoPPi_1, tC19Z_MA_NoPPi_2

Illumina MiSeq Paired End sequencing was used for tC19Z_LT_3_0, tC19Z_LT_3_2.5,

tC19Z_LT_3_10, tC19Z_LT_4_0, tC19Z_LT_4_3, tC19Z_LT_4_10, tC19Z_LT_5_10,

tC19Z_LT_5_2.5, tC19Z_LT_5_1, tC19Z_LT_5_0

A PhiX library was added into all samples and its sequences were first removed from all

sequencing files using Bowtie2.

Analysis for Figure 2d:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the full hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated. The

extension percentages shown in Figure 2d were calculated by generating percentages of the total

sequences for each extension length regardless of extended nucleotide identity.

Analysis for Figure 2e:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated. The

extension percentages shown in Figure 2e were calculated by generating percentages of just

sequences, in which the nucleotides were extended correctly as shown in the figure. The same

normalization factor as used in Figure 2e was applied.

Figure 2f:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated. The

percentages shown in Figure 2f were calculated by tabulating all sequences that had the

mismatched nucleotide repaired and extended from zero to five nucleotides for each mismatched

sequence.

Figure 2g:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated. The

percentages shown in the black bars in Figure 2g were calculated by tabulating the sequences

that were not repaired and the percentage that were subsequently extended correctly was

calculated. The grey bars in Figure 2g were calculated by tabulating the sequences that were

repaired and the percentage that were subsequently extended correctly was calculated.

Figure 3b and 3c:

CutAdapt was used to remove the adapters and primer regions on first the 5′ and then the 3′-end

of the sequences. Sequences were counted in FastAptamer using fastaptamer_count.

Sequences containing the primer were extracted from data for analysis in FastAptamer using

fastaptamer_search for each sequence and subsequently visually aligned and curated using

Jalview. Sequences that were not extended, ligation artifacts, and sequencing artifacts were

removed. Jalview was used to view the aligned sequences. The sequence lengths were verified

visually. This procedure was repeated for each condition shown in the figure.

Figure 3d and 3e:

CutAdapt was used to remove the adapters and primer regions on first the 5′ and then the 3′-end

of the sequences. Sequences were counted in FastAptamer using fastaptamer_count.

Sequences containing the primer were extracted from data for analysis in FastAptamer using

fastaptamer_search for each sequence and subsequently visually aligned and curated using

Jalview. Sequences that were not extended, ligation artifacts, and sequencing artifacts were

removed. The fidelity was determined by visually inspecting the sequences in Jalview and

tabulating the number of incorrectly incorporated nucleotides. The percentage fidelity was

determined as the average of the fidelity at each position.

Figure 4c-g:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated. The

extension percentages shown in Figure 4c-g were calculated by generating percentages of the

total sequences for each extension length.

1000

Extended Data Fig. 4:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-end of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present including sequences where the 3′-end was either

extended or shortened up to 5 nucleotides. The total counts of these sequences were tabulated.

FastAptamer search was then used to find sequences that corresponded to sequences that were

-1, -2, -3, and -4. These sequences were then visually inspected and manually curated. The

counts of the remaining sequences were tabulated. The percentages shown in Extended Data

Fig. 4 were calculated by generating percentages of sequences that had 1-4 nucleotides

removed.

1011

Extended Data Fig. 5:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated. The

extension percentages shown in the left graph of each panel in Extended Data Fig. 5 were

calculated by generating percentages of the total sequences for each extension length regardless

of extended nucleotide identity. The percentages were then normalized to add to 100%. The right

graph of each panel in Extended Data Fig. 5 shows the percentage of the sequences in the left

panel at each length where the mismatch was corrected.

1028

Extended Data Fig. 6:

CutAdapt was used to remove the adapters on first the 5′ and then the 3′-termini of the sequences.

FastAptamer was used to count the sequences. FastAptamer search was then used to find all

sequences that had the hairpin present, including sequences in which the 3′-ends were either

extended or shortened by up to 5 nucleotides. The total counts of these sequences were

tabulated. FastAptamer search was then used to find sequences that corresponded to the starting

hairpin, sequences that were -1, sequences that were extended by one nucleotide correctly,

sequences that were extended by one nucleotide incorrectly, sequences that were extended by

two nucleotides correctly, sequences that were extended by one nucleotide correctly and the

second nucleotide incorrectly, etc., up to five nucleotides. These sequences were then visually

inspected and manually curated. The counts of the remaining sequences were tabulated.

Extension percentages shown in Extended Data Fig. 6 were calculated for each sequence shown

as a fraction of the total hairpin containing sequences.

1042

Supplementary Fig. 2:

CutAdapt was used to remove the adapters and primer regions on first the 5′ and then the 3′-end

of the sequences, including the seven random nucleotides. Sequences were counted in

FastAptamer using fastaptamer_count. Sequences containing the primer were extracted from

data for analysis in FastAptamer using fastaptamer_search for each sequence and subsequently

visually aligned and curated using Jalview. Sequences that were not extended, ligation artifacts,

and sequencing artifacts were removed. The sequences were visually inspected, manually

aligned, and sequences that did not align to the start of the sequence were removed. The

sequences were exported to Adobe Illustrator, the reference sequence was colored green and

mismatches were colored blue.

1053

G/C Primer/Template

PolRz	-	+	+	+	-	+
NTPs	+	+	+	+	-	-
PPi (mM)	0	0	3	10	10	10

1055

Supplementary Fig. 1. Pyrophosphate inhibits ribozyme mediated extension. A primer/template sequence is extended by the ribozyme under standard conditions. The extension is inhibited when pyrophosphate is also added to the reaction (see also Extended Data Fig. 3). However, addition of pyrophosphate either with or without the tC19Z polymerase ribozyme does not result in any additional bands corresponding to RNA degradation or significant observable pyrophosphorolysis.

1062

Supplementary Fig. 2. Alignment of long extension sequences from HTS. Sequences corresponding to extensions present in the HTS data generated from the three-hour polymerization reaction and subsequent sequencing described in Figure 3B. Many of the sequences that are not full length terminate in one or more mismatches. The primer region is shown in the bracket, and the expected full-length sequence the HTS data was aligned to is highlighted in green. Mismatches are highlighted in blue.

1069

1070

1071

Supplementary Fig. 3. tC19Z sequences present in HTS analysis of long extension

reactions. (Top) Sequences of cDNAs generated by 3′–terminal ligation of the RNA strand and,

after reverse transcription, 3′–terminal ligation of the cDNA strand. The high-throughput

sequencing forward adapter was ligated to the 3′–terminus of the RNA. The cDNAs shown

matching the tC19Z ribozyme (top sequence) therefore report generation of RNAs antisense to

the ribozyme during the extension reactions (data shown are for analysis of 5-day experiments).

RNAs corresponding to long segments of tC19Z were observed only in experiments containing

pyrophosphate. The pyrophosphate concentrations are annotated on the left. The longest copies

of the ribozyme lack just 4 nucleotides (2%) from the ribozyme termini. (Bottom) All cDNA

sequences matching the sense strand of the tC19Z ribozyme generated under 0-, 1-, 2.5-, and

10-mM pyrophosphate conditions. No long sequences were observed in samples lacking

pyrophosphate. Thick lines represent the ribozyme matching segments, whereas the thin lines

represent segments of other sequences, such as ligation and sequencing adapters.

1085

1086

1087 **Supplementary Fig. 4. Damage repair reactions with a full set of controls for the U/A-**

1088 **terminated hairpin substrate.** A U/A hairpin was allowed to react under standard extension

1089 conditions with variations shown by + and - symbols below the gel image.

1090

1091

1092

Supplementary Fig. 5. Omission of a single nucleotide prevents extension of a repaired A/U hairpin. A [32]P-labeled A/U hairpin was extended under standard reaction conditions with variations indicated by the + and − symbols below each reaction resolved by PAGE. The left 5 lanes are negative controls, that lack any NTP; lane 6 is a positive control (PNK-repaired hairpin extension); and lane 7 is the pyrophosphate-facilitated repair reaction at 10 mM pyrophosphate. Lane 8 indicates that upon addition of all nucleotides except ATP (indicated by (B)TP (all, except ATP)), no extension is observed, strongly suggesting that the pyrophosphate-facilitated repair removes adenosine 5′ monophosphate-2′-3′-cyclic phosphate from the damaged terminus and requires ATP to incorporate the first nucleotide. Lanes 9–12 show that the repair does not occur at subsaturating concentrations of pyrophosphate.

A

UCUGCCAACCA>P 3'
C
|||||||||
C
G GACGGUUGGUACGCUUCGCACAGUAACUG 5'

	A/U α-³²P hairpin						A/U non-radiolabeled hairpin				
PolRz	+	+	+	+	+	+	+	+	+	+	+
PNK	-	-	-	+	+	+	-	-	-	+	+
α-³²P ATP	-	+	-	-	+	-	+	+	+	+	+
NTPs	+	-	+	-	-	+	-	-	+	-	+
PPi (10 mM)	-	+	+	-	-	-	-	+	+	-	-

B

UCUGCCAACCU>P 3'
C
|||||||||
C
G GACGGUUGGAACGCUUCGCACAGUAACUG 5'

U/A non-radiolabeled hairpin

tC19Z	+	+	+	+	+	+
PNK	-	-	-	+	+	+
α-³²P ATP	+	-	-	+	-	-
α-³²P UTP	-	+	-	-	+	-
α-³²P CTP	-	-	+	-	-	+
PPi (10 mM)	+	+	+	-	-	-

Supplementary Fig. 6. Incorporation of α-^{32}P-labeled nucleotides into repaired, non-labeled hairpins. (A) An A/U hairpin was prepared with or without internal ^{32}P, as indicated above the PAGE gel image. The hairpin was then subjected to repair and extension under standard reaction conditions, except that α-^{32}P-ATP was added to some reactions. In the case of the non-labeled hairpin, the ^{32}P-AMP is only incorporated if the sequence has been repaired either by pyrophosphate-facilitated reaction or by PNK. (B) A U/A hairpin was transcribed without α-^{32}P-ATP and was subjected to standard extension conditions but using α-^{32}P-rNTP (indicated) in place of the standard NTPs. Different α-^{32}P-rNTPs are utilized by the tC19Z polymerase ribozyme with different efficiencies, as shown by the varying intensities of the bands. ^{32}P-UMP is incorporated most efficiently because the first nucleotide on the template sequence is an adenosine. However, misincorporation is seen for the other nucleotides, and a second nucleotide appears to be added in presence of UTP. No ^{32}P labeling is observed in control reactions that lack PNK or pyrophosphate (as shown in panel A).

A/U Hairpin

	Hairpin				PNK			Acid			
Uridine (4 mM)	-	-	+	-	-	+	-	-	-	+	-
NTP	-	-	-	+	-	-	+	-	+	-	+
PPi (10 mM)	-	+	+	+	-	-	-	-	+	-	-

1120

Supplementary Fig. 7. The mechanism of repair does not involve cyclic-phosphate opening

by pyrophosphate. Pyrophosphate and uridine were added to a damaged A/U hairpin to

determine whether the cyclic phosphate could be opened by pyrophosphate to form a

triphosphate and allow for subsequent addition of uridine (lane 3). No additional bands were

observed on the PAGE gel, indicating that this is not the mechanism of repair. Uridine was added

to PNK-repaired or acid-treated hairpins as positive and negative controls, respectively.

1127

1128

Hairpin Workflow

1129

Supplementary Fig. 8. Workflow for HTS analysis of hairpin substrates of the polymerase

ribozyme. The hairpins were first prepared by primer extension of an drz-Mbac-1 HDV-like

ribozyme sequence (#6) and a hairpin sequence (#7–#22). This DNA construct with the ribozyme

on the 3′-end of the hairpin was transcribed into RNA where the HDV-like ribozyme co-

transcriptionally cleaves, leaving a hairpin with a 2′-3′-cyclic phosphate. For mismatch

experiments, that cyclic phosphate can be removed during incubation with PNK. The hairpins are

then extended by the tC19Z polymerase ribozyme. Following this reaction, the RNA is ligated to

a DNA oligo with a randomized 5′-end (#24/#25) and the resulting construct is reverse-transcribed

with RT primer #29. The Illumina adapters (#30 and #31) are then added via PCR prior to

sequencing.

Supplementary Table 1. DNA and RNA sequences.

T7 RNA polymerase promoter sequences are underlined. HDV overlap sequences in the hairpin

constructs are italicized. The terminal match or mismatch in the hairpin sequences is bolded.

1	tC19Z Ribozyme with T7 RNAP promoter sequence (underlined)	GCTAATACGACTCACTATA GTCATTGAAAAAAAAAGACAAATCTGCCCTCAGAGCTTGAGAACATC TTCGGATGCAGAGGAGGCAGCCTTCGGTGGCGCGATAGCGCCAAC GTTCTCAACAGACACCCAATACTCCCGCTTCGGCGGGTGGGGATAA CACCTGACGAAAAGGCGATGTTAGACACGCCCAGGTCATAATCCCC GGAGCTTCGGCTCC
2	RNA Primer	CUGCCAACCG
3	Short RNA Template	GUCAAUGACACGCUUCGCACGGUUGGCAG
4	tC19Z forward primer	GCTAATACGACTCACTATAGTCATTG
5	tC19Z reverse primer	GGAGCCGAAGCTCCGGGGATTATGAC
6	drz-Mbac-1 HDV-like ribozyme template (italicized segment overlaps with individual hairpin RNA constructs)	ATACGTATAACTATTAGCATGACCCATCGTCAACCACGAGGA*GGTTG TTATACTATGACGAC*
7	G/C Hairpin with drz-Mbac-1 HDV-like ribozyme overlap (italicized)	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**C**GGTTGGCAGGCCTCTGCCAACC**G** *GTCGTCATAGTATAACAACC*
8	C/G Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**G**GGTTGGCAGGCCTCTGCCAACC**C***GTC GTCATAGTATAACAACC*
9	A/U Hairpinwith drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**T**GGTTGGCAGGCCTCTGCCAACC**A***GTC GTCATAGTATAACAACC*
10	U/A Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**A**GGTTGGCAGGCCTCTGCCAACC**T***GTC GTCATAGTATAACAACC*
11	G/G Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**G**GGTTGGCAGGCCTCTGCCAACC**G***GTC GTCATAGTATAACAACC*
12	C/C Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**C**GGTTGGCAGGCCTCTGCCAACC**C***GTC GTCATAGTATAACAACC*
13	A/A Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**A**GGTTGGCAGGCCTCTGCCAACC**A***GTC GTCATAGTATAACAACC*
14	U/U Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**T**GGTTGGCAGGCCTCTGCCAACC**T***GTC GTCATAGTATAACAACC*
15	G/A Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**A**GGTTGGCAGGCCTCTGCCAACC**G***GTC GTCATAGTATAACAACC*
16	G/U Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**T**GGTTGGCAGGCCTCTGCCAACC**G***GTC GTCATAGTATAACAACC*
17	C/A Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**A**GGTTGGCAGGCCTCTGCCAACC**C***GTC GTCATAGTATAACAACC*
18	C/U Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**T**GGTTGGCAGGCCTCTGCCAACC**C***GTC GTCATAGTATAACAACC*
19	A/G Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**G**GGTTGGCAGGCCTCTGCCAACC**A***GTC GTCATAGTATAACAACC*
20	A/C Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	GCTAATACGACTCACTATA GTCAATGACACGCTTCGCA**C**GGTTGGCAGGCCTCTGCCAACC**A***GTC GTCATAGTATAACAACC*

21	U/C Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	<u>GCTAATACGACTCACTATA</u>GTCAATGACACGCTTCGCA**C**GGTTGGCAGGCCTCTGCCAACC**T***GTCGTCATAGTATAACAACC*
22	U/G Hairpin with drz-Mbac-1 HDV-like ribozyme overlap	<u>GCTAATACGACTCACTATA</u>GTCAATGACACGCTTCGCA**G**GGTTGGCAGGCCTCTGCCAACC**T***GTCGTCATAGTATAACAACC*
23	Long read template with T7 RNAP promoter sequence (underlined)	CTGCCAACCGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTGCGAAGCGTGTCATTGAC<u>TATAGTGAGTCGTATTAGC</u>
24	Ligation primer for sequencing	5Phos/NNNNACAGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTAGATC/3AmMO
25	Ligation primer for sequencing	5Phos/NNNNGTAGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGTAGATC/3AmMO
26	cDNA Ligation Primer for Sequencing of long reads	5Phos/CCTATAGTGAGTCGTATTAAAAAAAATATAGGNNNNNNN/3SpC3
27	Primer for Long Read Sequencing overlapping sequences #26 and #2	CGACTCACTATAGG CTGCCAA
28	Illumina reverse primer to Extend from primer 27 and overlap primer #2	GTGACTGGAGTTCAGACGTGTGCTCTTC CGATCT CACTATA GGCTGCCAACCG
29	RT primer for used after ligation of RNA with #24 and #25	GATCTACACTCTTTCCCTACACG
30	Illumina (R) adapter for hairpin sequencing overlapping the 5' terminus of the hairpin template strand (green)	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATC GTCAATGACACGCTTCGC
31	TrueSeq Universal Fwd to amplify after RT with #29 from ligated oligos #24&25	AATGATACGGCGACCACCGA GATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT

1143

1144 **Supplementary Table 2. Correlation of figures to deposited sequencing data.**

1145

Dataset	Main Figures	Extended Data and SI Figures
tC19Z_MM_1	2f (U/G), 2g (U/G)	Extended Data Fig. 4C (U/G), Extended Data Fig. 5F
tC19Z_MM_2	2f (G/U), 2g (G/U)	Extended Data Fig. 4C (G/U)
tC19Z_MM_3	2d, 2e, 2f (A/C), 2g (A/C)	Extended Data Fig. 4C (A/C), Extended Data Fig. 5D
tC19Z_MM_4	2f (C/A), 2g (C/A)	Extended Data Fig. 4C (C/A), Extended Data Fig. 5E
tC19Z_MM_5	2f (A/A), 2g (A/A)	Extended Data Fig. 4C (A/A)
tC19Z_MM_6	2f (A/G), 2g (A/A)	Extended Data Fig. 4C (A/G), Extended Data Fig. 5B
tC19Z_MM_7	2f (G/G), 2g (G/G)	Extended Data Fig. 4C (G/G), Extended Data Fig. 5A
tC19Z_MM_8	2f (G/A), 2g (G/A)	Extended Data Fig. 4C (G/G), Extended Data Fig. 5C
tC19Z_MM_9	2f (U/U), 2g (U/U)	Extended Data Fig. 4C (U/U)
tC19Z_MM_10	2f (U/C), 2g (U/C)	Extended Data Fig. 4C (U/C), Extended Data Fig. 6
tC19Z_MM_11	2f (C/C), 2g (C/C)	Extended Data Fig. 4C (C/C), Extended Data Fig. 5H
tC19Z_MM_12	2f (C/U), 2g (C/U)	Extended Data Fig. 4C (C/U), Extended Data Fig. 5G
tC19Z_MA_1	4e	Extended Data Fig. 4B (U/A)
tC19Z_MA_2	4c, 4d	
tC19Z_MA_3	4f	
tC19Z_MA_4	4g	Extended Data Fig. 4B (C/G)
tC19Z_MA_NoPPi_1		Extended Data Fig. 4A (U/A)
tC19Z_MA_NoPPi_2		Extended Data Fig. 4A (C/G)
tC19Z_LT_3_0	3b, 3c (0 mM PPi)	Supplementary Fig. 2
tC19Z_LT_3_2.5	3b, 3c (2.5 mM PPi)	
tC19Z_LT_3_10	3b, 3c (10 mM PPi)	Supplementary Fig. 2
tC19Z_LT_4_0	3b, 3d (0 mM PPi)	
tC19Z_LT_4_3	3b, 3d (3 mM PPi)	
tC19Z_LT_4_10	3b, 3d (10 mM PPi)	
tC19Z_LT_0_0	3b (0 mM PPi)	

1146 These datasets are available in NLM under SRA PRJNA1174777.
1147

1148   **Supplementary Table 3: Sequencing data in Figure 3.**

1149

Dataset	Total seq.	% Seq. with primer	% Extended
0 hours, 0 mM PPi	50825	14.7	0.08
3 hours, 0 mM PPi	101862	30.9	0.83
3 hours, 2.5 mM PPi	56588	37.9	0.77
3 hours, 10 mM PPi	42930	35.6	0.52
4 days, 0 mM PPi	140423	25.2	1.11
4 days, 3 mM PPi	65769	17.1	1.39
4 days, 10 mM PPi	26457	22.3	1.13

1150

1151
Supplementary Information: Full Length Gels
Boxes have been drawn around relevant lanes.
1154

1155
Figure 2b
1157
1158
1159
1160

1161
1162   Figures 4d, 4e, 4g and Extended Data Fig. 8 (A/U, U/A, C/G)

1163
1164   Extended Data Fig. 8 (G/C)

1165
1166   Extended Data Fig. 1

1167
1168   Extended Data Fig. 2: A/A left, G/G right

1169
1170   Extended Data Fig. 2: C/C left, U/C middle, U/U left

1171
1172   Extended Data Fig. 2: U/G left, G/A left middle, C/U right middle, G/U right
1173

1174
1175   Supplementary Fig. 1

1176
1177   Extended Data Fig. 7
1178

1179
1180   Supplementary Fig. 4

1181
1182   Supplementary Fig. 5

1183
1184   Supplementary Fig. 6A

1185
1186   Supplementary Fig. 6B

1187
1188   Supplementary Fig. 7
1189

1190
1191
1192