Cross-species activation of hydrogen cyanide production by a promiscuous quorum-sensing receptor promotes *Chromobacterium subtsugae* competition in a dual-species model

Cheyenne Loo[a, b, §], Pratik Koirala[a§], Nathan C. Smith[a], Anna Oller[a, c], Kara Evans[a, d], Saida Benomar[a, e], Isabelle R. Parisi[a], and Josephine Chandler[a, #]

[§]Contributed equally

[a]Stowers Institute for Medical Research, Kansas City, MO 64110 USA

[b]Department of Molecular Biosciences, University of Kansas, Lawrence, KS 66045 USA

[c]Current location: Department of Biological and Clinical Sciences, University of Central Missouri, Warrensburg, MO 64093 USA

[d]Current location: International Flavors and Fragrances Health and Biosciences, Madison WI, 53716 USA

[e]Current location: Department of Microbiology & Immunology, Emory University School of Medicine, Atlanta, GA, 30322 USA.

Running title: *C. subtsugae* eavesdropping and interspecies competition

[#]Address correspondence to Josephine R. Chandler, jrchandler@ku.edu.

ABSTRACT:

Many saprophytic bacteria have LuxR-I-type acyl-homoserine lactone (AHL) quorum-sensing systems that may be important for competing with other bacteria in complex soil communities. LuxR AHL receptors specifically interact with cognate AHLs to cause changes in gene expression. Some LuxR-type AHL receptors have relaxed specificity and are responsive to non-cognate AHLs. These promiscuous receptors might be used to sense and respond to AHLs produced by other bacteria by eavesdropping. We are interested in understanding the role of eavesdropping during interspecies competition. The soil saprophyte *Chromobacterium subtsugae* has a single AHL circuit, CviR-I, which produces and responds to *N*-hexanoyl-HSL (C6-HSL). The AHL receptor CviR can respond to a variety of AHLs in addition to C6-HSL. In prior studies we have utilized a coculture model with *C. subtsugae* and another soil saprophyte *Burkholderia thailandensis*. Using this model we previously showed that promiscuous activation of CviR by *B. thailandensis* AHLs provides a competitive advantage to *C. subtsugae.* Here, we show that *B. thailandensis* AHLs activate transcription of dozens of genes in *C. subtsugae,* including the *hcnABC* genes coding for production of hydrogen cyanide. We show that hydrogen cyanide production is cell density-dependent and demonstrate that the cross-induction of hydrogen cyanide by *B. thailandensis* AHLs provides a competitive advantage to *C. subtsugae.* Our results provide new information on *C. subtsugae* quorum sensing and are the basis for future studies aimed at understanding the role of eavesdropping in interspecies competition.

IMPACT STATEMENT:

In quorum sensing, population density-dependent changes in gene regulation are the result of a cytoplasmic transcription regulator binding to a quorum sensing signal. The signal-receptor interaction is considered to be specific to ensure fidelity of the system. However, some quorum-sensing receptor proteins have relaxed specificity and can recognize and respond to a range of signals. These promiscuous receptors might provide some benefit by enabling interspecies activation of quorum sensing by "eavesdropping," although the potential benefits of eavesdropping are not well studied. The current study utilizes a dual-species laboratory competition model, where one species has a promiscuous signal receptor and can respond to signals produced by the other species. In our study, we identify the signals that enable quorum sensing cross talk and show that cross talk promotes competition by inducing hydrogen cyanide production. Our results highlight how quorum sensing-enabled interspecies cross talk might provide an advantage during competition and provide a new basis for understanding how receptor-signal pairs might evolve in natural environments.

DATA SUMMARY

The authors confirm all supporting data, code and protocols have been provided within the article or through supplementary data files.

INTRODUCTION:

Quorum sensing is a cell density-dependent signaling system that relies on small diffusible signaling molecules. Once a critical population threshold is reached, the molecules interact with cytoplasmic transcription factors to cause activation of specific genes and coordinate gene expression across the population. In *Proteobacteria*, one type of quorum sensing is carried out by acyl-homoserine lactones (AHLs) signals. These molecules have a lactone ring and a fatty acid tail that varies in length from 4 to 20 carbons and can also have modifications at the third carbon position, which may be unsubstituted, hydroxylated, or have an oxo-substitution. Typical AHL quorum-sensing gene circuits consist of a LuxR-family signal receptor and a LuxI-family signal synthase, which produces a cognate signal. A bacterium may possess multiple LuxR-I circuits that typically produce and respond to different AHLs. Although the AHLs generally have similar structures, the receptors are thought to interact with AHLs in a highly specific and selective manner (1). However, there are exceptions such as the CviR receptor of *Chromobacterium subtsugae* (formerly *C. violaceum*), which can promiscuously detect and respond to a range of AHLs (2, 3). The existence of such receptors suggests receptor promiscuity might have certain advantages in some conditions.

Quorum sensing systems control different behaviors in different bacteria, such as the production of toxins, exoproducts, and biofilm matrix components. Many of these behaviors might provide a benefit to individuals within groups or complex communities. For example, toxin production might be important for competition with other strains or species of bacteria. There is an increasing interest in understanding the specific benefits of quorum sensing in dynamic polymicrobial communities. A major barrier to understanding quorum sensing in this context is the need for laboratory models that enable studies of mixed-strain and mixed-species interactions in a controlled setting. Such models are increasingly being utilized, and are

providing significant new insights into the role and mechanisms of quorum sensing in mixed microbial communities (for a review, see (4)).

This study utilizes a laboratory dual-species model that was previously developed where two soil bacteria, *Burkholderia thailandensis* and *Chromobacterium subtsugae,* use AHL-dependent quorum sensing to compete with the other species (5) (Fig. 1). *B. thailandensis* produces a ribosome-targeting antibiotic called bactobolin (6). Bactobolin is regulated by a LuxR-I system called BtaR2-I2, which senses and responds to 3OHC8-HSL and 3OHC10-HSL (7). Both bactobolin and the BtaR2-I2 system are important for *B. thailandensis* to compete with *C. subtsugae* (5). In addition to BtaR2-I2, *B. thailandensis* has two other complete quorum-sensing circuits, BtaR1-I1 and BtaR3-I3, which produce and respond to C8-HSL and 3OHC10-HSL, respectively (8). *C. subtsugae* has a single LuxR-I-type system, CviR-I, which produces and responds to C6-HSL (2). CviR-I is important for competition with *B. thailandensis* (5). CviR is known to activate production of a purple pigmented antibiotic, violacein, however mutational disruption of violacein had no effect on competition (5). We also showed that CviR can be activated by *B. thailandensis* AHLs by "eavesdropping" to promote the competitive ability of *C. subtsugae* (5).

In this study, we identify C8-HSL as the *B. thailandensis* AHL best able to activate *C. subtsugae* production of quorum-dependent antimicrobials. We use RNAseq to identify the full set of *C. subtsugae* genes that can be activated by either C8-HSL or the native signal, C6-HSL. Among the genes most highly activated by both signals are those coding for hydrogen cyanide biosynthesis. We use our coculture model to validate the importance of hydrogen cyanide for *C. subtsugae* competition. We also demonstrate that the hydrogen cyanide genes can be transcriptionally activated by CviR and both native and non-native signals. The results provide

new information on promiscuous AHL responses in *C. subtsugae* and further support the idea

that AHLs drive important interactions through interspecies crosstalk.

METHODS:

Bacterial culture conditions and reagents. All strains were grown in Luria-Bertani (LB) broth (10g tryptone, 5g yeast extract, and 10g NaCl in 1L water), LB with morpholine-propanesulfonic acid (LB-MOPS, 50mM; pH 7), or in LB with 1.5% (wt/vol) Bacto-Agar. All broth cultures were incubated at 30 °C (*C. subtsugae* or *C. subtsugae-B. thailandensis* cocultures) or 37 °C (*B. thailandensis* or *E. coli*). Synthetic acyl-homoserine lactone signals were purchased from Cayman Chemicals (MI, USA) and stored in ethyl acetate acidified with 0.1% of glacial acetic acid. In all cases, synthetic signals were added to the culture flask and dried down completely using a stream of nitrogen gas prior to adding the culture media. For strain constructions, we used gentamicin at 50 μg ml^{-1} (*C. subtsugae*) or 15 μg ml^{-1} (*E. coli*). For selection from cocultures, we used gentamicin at 100 μg ml^{-1} (*B. thailandensis*) and trimethoprim at 100 μg ml^{-1} (*C. subtsugae*).

Genomic DNA, PCR and DNA fragments, and plasmid DNA were purified by using a Puregene Core A kit, plasmid purification miniprep kit, or PCR cleanup/gel extraction kits (Qiagen or IBI-MidSci) according to the manufacturer's protocol. RNA was isolated using the Qiagen RNEasy Minikit.

Bacterial strains and plasmids. Strains are summarized in Table 1. For cocultures, we used *B. thailandensis* strains JBT125 and BD20. Both are derived from strain E264 (9); BD20 is an AHL-deficient *btaK* mutant (7), and JBT125 is an AHL- and bactobolin-deficient *bta1-3, btaK* quadruple mutant (8). *C. subtsugae* strain CV017 (referred to as wild type, previously known as *C. violaceum* CV017) is a derivative of strain ATCC 31532 with a transposon insertion in gene CV_RS05185 causing overexpression of violacein (2). All *C. subtsugae* mutant strains were

constructed from CV017 using allelic exchange and methods described below. For plasmid construction, we used *E. coli* strain DH10B (Invitrogen).

Unmarked, in-frame deletions of *cviR, cviI,* and *hcnB* (Cv017_06925) were constructed using the following method. DNA fragments were generated by PCR or DNA synthesis (Genscript, New Jersey) containing ~500-1000 bp DNA flanking each gene and fused together, creating an unmarked, non-polar deletion of each gene with incorporated XbaI and SacI restriction enzymes sites. These fragments were digested with XbaI and SacI and cloned into XbaI, SacI-digested pEXG2 (10). The relevant pEXG2 gene deletion plasmid was subsequently used to make the mutant *C. subtsugae* strain using methods described previously (5). Briefly, the relevant pEXG2 plasmid was transformed into *C. subtsugae* by electroporation or conjugation. Merodiploids were selected on LB agar containing gentamicin, and deletion mutants were counterselected on LB + 15% sucrose. Mutant strains were verified by testing for gentamicin sensitivity and by PCR-amplifying the deletion region and sequencing the PCR product.

For transcription reporter assays, plasmid pMP220 *hcnA-lacZ* was made by amplifying the *hcnA* promoter from *C. subtsugae* with primers hcn220R (5'-TATATCTAGAGATTAGCGGTTTGCCGTTCA-3') and hcn220F (5'-TATAGGTACCGCCACTGAATGCCAATGTTCTT-3'), which incorporated the XbaI and KpnI sites. The amplified product was digested with XbaI and KpnI and ligated to XbaI- and KpnI-digested pMP220 (11). pMP220 *hcnA-lacZ* was introduced to *C. subtsugae* Δ*cviI* and Δ*cviR* by electroporation.

RNA isolation. To isolate RNA for RNAseq, cultures were grown to an OD_{600} of 4.0 in 10 ml LB-MOPS in 125 ml non-baffled flasks. Bacteria (~1 × 10^9 cells) were removed from the

cultures and suspended in RNAprotect bacteria reagent (Qiagen), pelleted by centrifugation, and stored at −80°C. Thawed cells were suspended in 1 ml RLT buffer (Qiagen) containing 2-mercaptoethanol and lysed by bead beating. RNA was purified using the RNAeasy minikit (Qiagen). Contaminating DNA was removed with Turbo DNase (Ambion), and RNA was obtained by using RNeasy MinElute cleanup kit (Qiagen).

RNA-seq library construction, sequencing, mapping and analysis. Libraries for RNA-seq were prepared by Novogene using the Novogene mRNA-seq services project workflow. cDNA libraries were sequenced by Novogene using an Illumina platform and PE-150 reads. Sequences were aligned and mapped to the *C. subtsugae* CV017 genome (accession NZ_JAHDTB000000000) using Featurecounts and the Rsubread package in R (12). We determined differentially regulated genes for biological replicates using function edgeR (13) and using a false discovery rate [FDR] cutoff of 0.05. We proceeded with genes with >15 aligned reads and 2-fold or more regulation by C6- or C8-HSL relative to no AHL. Functions and their sources can be found at https://github.com/ChandlerLabKU/Cs_rnaseq. The data have been deposited in the NCBI sequence read archive (SRA database) (BioProject identification [ID] PRJNA894552). For Fig. 6, promoter sites and DNA recognition motifs were predicted using RSAT (14) and the transcription start site in the DNA region upstream of the *C. subtsugae hcnA* gene was predicted from the RNAseq data generated in this study using the TSSr program in R (15).

Coculture experiments. Coculture experiments were conducted in 20 mL LB-MOPS medium in 125 mL non-baffled flasks. The inoculum was from logarithmic-phase pure cultures of *C.*

subtsugae and *B. thailandensis*. The initial OD_{600} in the coculture was 0.05 for *B. thailandensis* $(2—4x10^7$ cells per ml) and 0.005 for *C. subtsugae* $(2 – 4x10^6$ cells per mL). After inoculating, cocultures were incubated at 30 ºC with shaking at 250 rpm for 24 hours. Colony forming units of each species were determined by using differential antibiotic selection on LB agar plates. *B. thailandensis* was selected with gentamicin and *C. subtsugae* was selected with trimethoprim.

Transcription reporter assays. To assess activation of the *hcnA* promoter in *C. subtsugae,* overnight cultures of *C. subtsugae* strains carrying the pMP220_*hcn-lacZ* reporter plasmid were used as starters by diluting to an OD_{600} of 0.05. When experimental cultures reached an OD_{600} of 0.5 they were distributed to 2 ml Eppendorf tubes containing different concentrations of dried C6-HSL, C8-HSL, 3OHC8-HSL, and 3OHC10-HSL. The volume in each tube was 0.5 ml. After 5 h with shaking at 30°C, β-galactosidase activity was measured using the Tropix Galacto-Light Plus™ chemiluminescent kit according to the manufacturer's protocol (Applied Biosystems, Foster City, CA).

RESULTS:

C. subtsugae **response to *B. thailandensis* AHLs in coculture.** Previously, we showed that extracted fluid from *B. thailandensis* cultures containing all three *B. thailandensis* AHLs (C8-HSL, 3OHC8-HSL and 3OHC10-HSL) increase *C. subtsugae* competitiveness in a manner dependent on the *C. subtsugae* quorum-sensing receptor CviR (5). These results suggest that *C. subtsugae* CviR is responsive to one or several of the *B. thailandensis* AHLs. Therefore, we carried out coculture experiments to determine the individual contribution of each of the *B. thailandensis* AHL(s) to *C. subtsugae* competitiveness. We used signal synthase mutants of *C.*

subtsugae and *B. thailandensis* for the competition experiment and added each of the *B. thailandensis* synthetic AHLs individually at 1 uM AHLs, which is a concentration in the range of that reported for similarly-grown pure cultures of *B. thailandensis* (16). The *B. thailandensis* strain JBT125 was also bactobolin-defective (Δ*btaK*) so that *C. subtsugae* would not be killed by AHL-induced bactobolin in the experiment. In our cocultures experiment we observed that the *C. subtsugae* native signal C6-HSL improved *C. subtsugae* competitiveness by ~100-fold compared with the no signal condition (Fig. 2), similar to our prior study (5). Adding C8-HSL and 3OHC8-HSL also significantly improved competitiveness of *C. subtsugae* although to a lesser degree than the native C6-HSL. 3OHC10-HSL had a more variable effect than the other AHLs and was not significantly different from no AHLs, although this signal also appeared to improve competitiveness. Together, these results show that *C. subtsugae* is responsive to the *B. thailandensis* AHLs C8-HSL and 3OHC8-HSL, and possibly also to 3OHC10-HSL.

Production of hydrogen cyanide is induced by quorum sensing and *hcnB*. We used RNAseq transcriptomic analysis to identify AHL-activated *C. subtsugae* genes as a first step to identify the factor important for competition with *B. thailandensis*. We compared transcripts in our Δ*cviI* strain grown with or without exogenously added synthetic AHLs. We grew cultures with either C6-HSL or C8-HSL, with the goal of identifying a set of genes with the ability to respond to both signals. We anticipated that the gene(s) required for *C. subtsugae* to compete via eavesdropping would be among this gene set. We collected cells for our analysis during the transition from logarithmic growth to stationary phase, when cells are at an optical density at 600 nm (OD$_{600}$) of 4.0. At this cell density, the culture just begins turning purple due to production of

the quorum sensing-activated purple pigment violacein. AHLs were added at 2 µM to ensure robust responses for detecting differences in gene expression.

Overall, 348 *C. subtsugae* genes were activated by C6-HSL, and 97 of these genes were also activated by C8-HSL (Table S1). In the set of genes most highly activated by both C6-HSL and C8-HSL (Table 2), we identified *hcnA*, *hcnB* and *hcnC,* which are predicted to code for biosynthesis of hydrogen cyanide. Hydrogen cyanide is important for *C. subtsugae* to protect itself from predation by other bacteria and kill mosquito larvae (17, 18), and is also important for *P. aeruginosa* to compete with *B. multivorans* in cocultures (19). Therefore, we focused our attention on hydrogen cyanide. To determine whether *C. subtsugae* produces hydrogen cyanide, we measured cyanide ion (CN$^-$) in cell-free fluid from wild-type *C. subtsugae* cultures grown to various growth stages (early logarithmic, late logarithmic, stationary phase) (Fig. 3A). The production of CN$^-$ was population-density dependent, consistent with quorum sensing activation of these genes. We detected CN$^-$ in fluid from stationary-phase cultures at a maximum concentration of 3.3 ± 1.1 mM at an OD$_{600}$ ~7.

To determine if either the *hcn* genes or quorum sensing was required for hydrogen cyanide production, we introduced a $\Delta hcnB$ mutation to the wild type genome and measured CN$^-$ in culture fluid of the $\Delta hcnB$ and $\Delta cviR$ mutants and wild type. Fluid from stationary-phase (OD$_{600}$ of 7) cultures of both $\Delta hcnB$ and $\Delta cviR$ were similarly low compared with identically grown wild type (Fig. 3B). These results support the conclusion that quorum sensing and *hcnB* are important for hydrogen cyanide production in *C. subtsugae.*

Cyanide is important for *C. subtsugae* to compete with *B. thailandensis.* Because expression of the hydrogen cyanide biosynthesis genes is responsive to both C6-HSL and C8-HSL, we

hypothesized that *B. thailandensis* AHLs increase competitiveness of *C. substugae* by activating hydrogen cyanide production through eavesdropping. As a first test of this hypothesis, we determined the susceptibility of *B. thailandensis* strain BD20 to a range of concentrations of potassium cyanide (0-750 uM), which mimics cyanide ions (20). We determined the 50% lethal dose (LD_{50}) of potassium cyanide for *B. thailandensis* BD20 to be 119 µM (Fig. 4A). Thus, the concentration of cyanide produced by *C. subtsugae* (~3 mM, Fig. 2A) is more than sufficient to kill *B. thailandensis*.

Next, we investigated the effects of hydrogen cyanide in cocultures (Fig. 4B). We competed the bactobolin-deficient *B. thailandensis* (BD20) with different genetic mutants of *C. subtsugae*. In these conditions, *B. thailandensis* BD20 produces AHLs that can activate the quorum-sensing system of *C. subtsugae* (5). Consistent with prior results (5), in these conditions the *C. subtsugae* Δ*cviI* muant and wild type are similarly competitive, because the Δ*cviI* mutant has CviR that can be activated by *B. thailandensis* AHLs. The Δ*cviR* mutant cannot respond to any AHLs and, as expected, was significantly less competitive than the wild-type and Δ*cviI* mutant. To determine the role of hydrogen cyanide in competition, we tested the competitive ability of the Δ*hcnB* mutant. The Δ*hcnB* single mutant competed poorly with BD20 and was similar to the Δ*cviR* mutant. These results support that hydrogen cyanide is important for the competitive ability of *C. subtsugae*. We also tested the Δ*cviI*, Δ*hcnB* double mutant. In this strain, the competitive ability of *C. subtsugae* relies on AHLs produced by *B. thailandensis*. The Δ*cviI*, Δ*hcnB* double mutant also competed poorly, similar to the Δ*cviR* and Δ*hcnB* single mutants. These results support that hydrogen cyanide is important for *C. subtsugae* to compete in response to *B. thailandensis* AHLs. Together, the results support the idea that hydrogen cyanide

is required for the competitive advantage provided to *C. subtsugae* in response to native AHLs and non-native AHLs.

Sensitivity of *hcnA* promoter to AHLs. The hydrogen cyanide biosynthesis genes *hcnA, hcnB* and *hcnC* (the *hcn* genes) are adjacently encoded in the genome facing the same direction. There is a single putative promoter upstream of *hcnA*. We hypothesized that this promoter is activated either directly or indirectly by CviR in response to the *C. subtsugae*-produced signal C6-HSL, as well as the *B. thailandensis* signals C8-HSL, 3OHC8-HSL, and possibly 3OHC10-HSL. To test our hypothesis, we created a plasmid with a DNA fragment containing the putative promoter region upstream of *hcnA* fused to a promoterless *lacZ* gene. This plasmid was introduced into the *C. subtsugae* Δ*cviI* and Δ*cviR* mutants. The *hcnA-lacZ* reporter was activated by 5 μM of each of the four AHLs (C6-, C8-, 3OHC8- and 3OHC10-HSL) in the Δ*cviI* mutant, but not in the Δ*cviR* mutant (Fig. 5A). Thus, activation of *hcnA-lacZ* required CviR and AHLs. We also determined the signal sensitivity of the *hcnA-lacZ* reporter by testing each AHL at a range of concentrations (Fig. 5B). The *hcnA-lacZ* reporter responded most sensitively to the native signal C6-HSL. There was also a response to C8-HSL. The other two AHLs, 3OHC8-HSL and 3OHC10-HSL, activated the reporter only at the highest signal concentrations (1-5 μM). These results were consistent with the coculture results (Fig. 2), and demonstrate that C8-HSL and, to a lesser extent 3OHC8-HSL and 3OHC10-HSL, can activate the *hcn* gene promoter in a manner dependent on CviR.

DISCUSSION:

The *B. thailandensis-C. subtsugae* model system was developed previously to examine the effects of quorum sensing and AHL-dependent eavesdropping in competition (5). In those studies, we elucidated the importance of *B. thailandensis* bactobolin for competition, however, the *C. subtsugae* toxin remained elusive. Here, we identified hydrogen cyanide as the *C. substugae* toxin, filling in this gap in knowledge. We have also shown that noncognate AHLs can induce dozens of genes in addition to those coding for hydrogen cyanide. Results with our coculture model support the idea that promiscuous receptors might enable interspecies interactions by coordinating broad changes in gene expression in response to AHLs from other species. Results of our studies also expand the list of genes known to be activated by the CviR-I system in *C. subtsugae.*

The *C. substugae* CviR receptor can activate certain genes in response to a wide range of AHLs (2). Although detailed structure-function studies of CviR and different AHLs have been carried out (21, 22), few studies have addressed the potential benefits of AHL receptor promiscuity for the cell. Receptor activation by non-self AHLs could enable detection of other bacteria and might provide certain benefits for interspecies competition (5) or cooperation (23). Indeed, there is evidence of eavesdropping in natural polymicrobial communities in plants (23-25), and eavesdropping has also been shown to occur in communities relevant to infections (26). Recent studies also support the idea that promiscuity might be a feature of a wider range of receptors than previously recognized (27). These studies support the idea that AHL-dependent cross talk might be an important but underrecognized facet of quorum sensing communication and that it warrants further study.

We find it interesting that in our study, the majority of genes most strongly activated by both C6-HSL and C8-HSL in *C. subtsugae* are predicted to code for antibiotic production (Table

2). The receptor BtaR2 in *B. thailandensis* is also highly promiscuous and regulates production of a potent and broad-spectrum antibiotic, bactobolin (6, 7). The role of these receptors in regulating antibiotic production supports the idea that promiscuous activation of receptors could play an important role in competition. The promiscuous activation of CviR by C8-HSL leads to the induction of a subset of genes that overlaps with that controlled by C6-HSL. It may be that these genes are more highly sensitive to CviR, enabling response to CviR activation by non-native signals. An alternative possibility is that C8-HSL changes CviR recognition of gene promoters, possibly through allosteric changes in the DNA binding motif. Future studies focused on the regulation of particular genes by different AHLs could help to elucidate the mechanism of CviR activation of gene promoters by non-native AHLs.

Prior studies show that receptor selectivity falls on a continuum from highly selective (e.g. *P. aeruginosa* RhlR) to intermediate (e.g. *P. aeruginosa* LasR) to highly promiscuous (e.g. CviR, and *B. thailandensis* BtaR2) (27). The affinity of receptors for a particular AHL can be influenced by changing specific residues within the AHL binding pocket of the receptor (21, 28-30). For example, although CviR is normally inhibited by the noncognate signal C10-HSL, replacing methionine at position 89 with serine converts this receptor to become C10-HSL-responsive (21). These results suggest that a particular receptor could easily adapt to become more or less promiscuous dependent on the benefits of promiscuity. In addition, receptor selectivity is influenced by receptor expression levels (27). Although very little is known about the environmental conditions that regulate quorum sensing receptors, one can imagine that conditions that increase receptor levels in the cell could increase AHL promiscuity. For example, it was recently demonstrated that the CviR receptor is activated by certain antibiotics (31), although the effects of antibiotics on receptor selectivity remain to be tested.

The results of this study are consistent with prior studies (17, 19, 32) supporting a role for hydrogen cyanide in competition. In *P. aeruginosa,* the hydrogen cyanide biosynthesis genes are part of an operon driven by a promoter upstream of *hcnA* (see Fig. 6). The *P. aeruginosa hcnA* promoter is quorum-sensing controlled, and has an experimentally validated binding site for the quorum sensing receptor LasR at position -101 relative to the translation start site (33). This region also has a binding site for the anaerobic regulator (Anr) at position -72 (33, 34) (Fig. 6). The intergenic region upstream of *hcnA* in *P. aeruginosa* and *C. substugae* share 45% identity; however, we were unable to find evidence of a *lux* box or Anr binding site anywhere in the *C. substugae hcnA* promoter. It is possible that CviR can recognize the *hcnA* promoter in the absence of a recognizable *lux* box, which has been reported in some cases for LasR (33). It is also possible that CviR controls the *hcn* genes indirectly. Either way, our results provide additional support for the idea that CviR can activate production of hydrogen cyanide in response to AHL from other species, supporting the idea that some quorum sensing-dependent competitive behaviors might be induced by interspecies cross talk.

AUTHOR STATEMENTS

The authors declare no conflicts of interest.

This work was supported by the National Institutes of Health (NIH) R35GM133572 to J.R.C. S.B. and C.L. were supported by an NIH K-INBRE fellowship (P20GM103418), and K.C.E. was supported by the NIH Chemical Biology Training Program (T32 GM08545). P.K. was supported by the KU Weaver Fellowship. We also thank the KU Genomics Core, which is supported by KU and NIH grants P20GM103638 (COBRE CMADP) and P20GM103418 (K-INBRE).

We acknowledge Drs. Ajai Dandekar and Amy Schaefer for helpful discussions as well as Nicole Smalley for technical input.

TABLES AND FIGURES

Table 1. Bacterial strains and plasmids used in this study.

Strain or plasmid	Relevant properties[a]	Reference or source
Strain		
B. thailandensis		
JBT125	*bta1-3, btaK* quadruple mutant of strain E264	(8)
BD20	*btaK* mutant of strain E264	(7)
C. substugae		
CV017	Mini-Tn5 mutant of ATCC 31532, 'wild type'	(2)
CV017 Δ*cviI*	CV017 with an in frame, unmarked *cviI* deletion	This study
CV017 Δ*cviR*	CV017 with an in frame, unmarked *cviR* deletion	This study
CV017 Δ*hcnB*	CV017 with an in frame, umarked *hcnB* deletion	This study
CV017 Δ*cviI*, Δ*hcnB*	CV017 Δ*cviI* with the Δ*hcnB* mutation	This study
Escherichia coli		
DH5α	[F- φ80 lacZΔM15] Δ(lacZYA-argF)U169 recA1 endA1 hsdR17, [rK−mK+] supE44 thi-1 gyrA relA1	Invitrogen
Rho-3	thi-1 thr-1 leuB26 tonA21 lacY1 supE44 recA integrated RP4-2 Tcr::Mu (λpir+)Δasd::FRT ΔaphA::FRT	(35)
Plasmids		
pEXG2	Suicide vector, Gm^r	(10)
pEXG2 Δ*cviI*	pEXG2 with Δ*cviI* extending from XX to XX with regard to the *cviI* translation start site	This study
pEXG2 Δ*hcnB*	pEXG2 with Δ*hcnB* extending from XX to XX	This study

with regard to the *cviI* translation start site

pMP220   lacZ transcriptional reporter vector, IncP, Tc^R (11)

pMP220_hcn-lacZ   pMP220 with the *hcnA* promoter extending from XX to XX, Tc^R This study

---

^{a}Tc^R, tetracycline resistant; Gm^R, gentamicin resistant.

Table 2. Genes most highly activated by C6-HSL and C8-HSL

Locus tag[a]	Description	Fold change[b]	
		C6-HSL	C8-HSL
KIF53_RS20405	amino acid adenylation domain-containing protein	7.4	4.8
KIF53_RS06810	cyanide-forming glycine dehydrogenase subunit HcnA.	6.3	6.4
KIF53_RS14555	DUF1842 domain-containing protein	6.2.	3.3
KIF53_RS20400	hybrid non-ribosomal peptide synthetase/type I polyketide synthase	6.5	5.0
KIF53_RS01860	long-chain-fatty-acyl-CoA reductase	5.9	2.1
KIF53_RS13810	activator protein	7.3	4.4
KIF53_RS05635	FAD-dependent oxidoreductase	5.4	4.1
KIF53_RS15685	M9 family metallopeptidase N-terminal domain-containing protein	5.0	3.7
KIF53_RS15785	methyl-accepting chemotaxis protein	4.4	2.6
KIF53_RS20410	non-ribosomal peptide synthetase	6.8	4.0
KIF53_RS16340	ornithine carbamoyltransferase	5.2	5.2
KIF53_RS06800	cyanide-forming glycine dehydrogenase subunit HcnC	5.4	5.6
KIF53_RS06795	ABC transporter substrate-binding protein	4.7	4.6
KIF53_RS02825	hydrogenase maturation protein	4.5	3.2
KIF53_RS06805	NAD(P)/FAD-dependent oxidoreductase HcnB	6.9	6.9

[a]Locus tags correspond to the *C. subtsugae* CV017 genome (accession

NZ_JAHDTB000000000.1)

[b]Fold change value of induction by AHLs (C6-HSL or C8-HSL) when added to strain CV017

Δ*cviI*.

Fig. 1. *Burkholderia thailandensis-Chromobacterium subtsugae* **coculture model.** The *B. thailandensis* quorum sensing system BtaR2-I2 produces and responds to 3OHC8-HSL and 3OHC10-HSL, and activates production of bactobolin antibiotic. BtaR2-I2 and bactobolin are both important for *B. thailandensis* to compete with *C. subtsugae*. The *C. subtsugae* quorum sensing system CviR-I produces and responds to C6-HSL. CviR-I is important for *C. subtsugae* to compete with *B. thailandensis,* through an unknown mechanism.

Fig. 2. Contribution of *B. thailandensis* AHLs to *C. subtsugae* competitiveness. Cocultures were with *C. subtsugae* ΔcviI and *B. thailandensis* ΔbtaI1, ΔbtaI2, ΔbtaI3, ΔbtaK (JBT125). After 24 h of coculture growth, the ratio of *C. subtsugae* to *B. thailandensis* was determined by selective plating and colony counts. The dashed line indicates the initial ratio of *C. subtsugae* to *B. thailandensis*. The solid line represents the means and the vertical bars shows the standard error of the mean for each group. Synthetic AHLs were added at the beginning of the coculture to a final concentration of 1 μM. Statistical comparisons of each condition with no signal coculture were by one-way ANOVA; ***, $p<0.001$; **, $p<0.005$ and ns, not significant.

Fig. 3. Cyanide measurements in *C. subtsugae* culture fluid. Cyanide (CN⁻) was measured in culture fluid of *C. subtsugae*. A. Wild type *C. subtsugae* (CV017) was grown to different cell densities prior to measuring CN⁻. Data are shown as growth adjusted CN⁻. B. The indicated *C. subtsugae* strain was grown for 12 hr prior to measuring cell density and CN⁻. Data are shown as the growth-adjusted CN⁻ as a percentage of that of wild type. In all cases, data represent the average of 3 biological replicates and the vertical bars represents the standard error of the mean.

Fig. 4. Cyanide toxicity and role in cocultures. A. Survival of *B. thailandensis* BD20 grown in the presence of different concentrations of cyanide for 8 hr from an initial OD_{600} of 0.005. The cell density was measured and *B. thailandensis* (Bt) survival was determined as a percent of untreated culture grown under identical conditions. B. Cocultures were with *B. thailandensis* Δ*btaK* (BD20) and the *C. subtsugae* strain indicated. After 24 h of coculture growth, the ratio of *C. subtsugae* to *B. thailandensis* was determined by selective plating and colony counts. The initial ratio of *C. subtsugae* to *B. thailandensis* was 10^{-1}. The solid line represents the means and the vertical bars shows the standard error for each group. Statistical comparisons of each condition compared with no signal were by one-way ANOVA; **, $p<0.005$; * = $p < 0.05$ and ns, not significant.

Fig. 5. Sensitivity of the *hcnA* promoter to native and non-native AHLs. A. *C. subtsugae*

Δ*cviI* or Δ*cviR* mutant harboring the *hcnA-lacZ* reporter was treated with 5 µM of AHL (C6-

HSL, C8-HSL, 3OHC8-HSL or 3OHC10-HSL). B. *C. subtsugae* Δ*cviI* harboring the *hcnA-lacZ*

reporter was treated with AHLs at the indicated concentrations. Results show the average of four

independent experiments from two separate days, and the vertical bars represent the standard

error.

Fig. 6. Hydrogen cyanide biosynthesis genes (*hcnABC*) in *C. subtsugae* and *P. aeruginosa*.
Red arrows and lettering indicate the transcription start site. The arrow is dashed for *C. subtsugae* as it was predicted to be between nucleotides -79 to -81 (see Methods). Green boxes and lettering indicate the binding site for the anaerobic regulator (Anr) and the blue box and lettering indicate the *lux* box of *P. aeruginosa,* and the canonical lux box. Bolding in the *P.aeruginosa lux* box sequence indicate nucleotides that are conserved with the canonical *lux* box. No *lux* box or Fnr/Anr like sequences were identified in the *C. subtsugae* sequence (see Discussion text).

CITATIONS

1. Hawver LA, Jung SA, Ng WL. 2016. Specificity and complexity in bacterial quorum-sensing systems. FEMS Microbiol Rev 40:738-52.

2. McClean KH, Winson MK, Fish L, Taylor A, Chhabra SR, Camara M, Daykin M, Lamb JH, Swift S, Bycroft BW, Stewart GS, Williams P. 1997. Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones. Microbiology 143 (Pt 12):3703-11.

3. Swem LR, Swem DL, O'Loughlin CT, Gatmaitan R, Zhao B, Ulrich SM, Bassler BL. 2009. A quorum-sensing antagonist targets both membrane-bound and cytoplasmic receptors and controls bacterial pathogenicity. Mol Cell 35:143-53.

4. Abisado RG, Benomar S, Klaus JR, Dandekar AA, Chandler JR. 2018. Bacterial quorum sensing and microbial community interactions. mBio 9:e02331-17.

5. Chandler JR, Heilmann S, Mittler JE, Greenberg EP. 2012. Acyl-homoserine lactone-dependent eavesdropping promotes competition in a laboratory co-culture model. ISME J 6:2219-28.

6. Seyedsayamdost MR, Chandler JR, Blodgett JA, Lima PS, Duerkop BA, Oinuma K, Greenberg EP, Clardy J. 2010. Quorum-sensing-regulated bactobolin production by *Burkholderia thailandensis* E264. Org Lett 12:716-9.

7. Duerkop BA, Varga J, Chandler JR, Peterson SB, Herman JP, Churchill ME, Parsek MR, Nierman WC, Greenberg EP. 2009. Quorum-sensing control of antibiotic synthesis in *Burkholderia thailandensis*. J Bacteriol 191:3909-18.

8. Chandler JR, Duerkop BA, Hinz A, West TE, Herman JP, Churchill ME, Skerrett SJ, Greenberg EP. 2009. Mutational analysis of *Burkholderia thailandensis* quorum sensing and self-aggregation. J Bacteriol 191:5901-9.

9. Brett PJ, DeShazer D, Woods DE. 1998. *Burkholderia thailandensis* sp. nov., a *Burkholderia pseudomallei*-like species. Int J Syst Bacteriol 48 Pt 1:317-20.

10. Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. 2005. ExsE, a secreted regulator of type III secretion genes in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 102:8006-11.

11. Spaink HP, Okker RJ, Wijffelman CA, Pees E, Lugtenberg BJ. 1987. Promoters in the nodulation region of the Rhizobium leguminosarum Sym plasmid pRL1JI. Plant Mol Biol 9:27-39.

12. Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Res 47:e47.

13. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26:139-40.

14. Santana-Garcia W, Castro-Mondragon JA, Padilla-Galvez M, Nguyen NTT, Elizondo-Salas A, Ksouri N, Gerbes F, Thieffry D, Vincens P, Contreras-Moreira B, van Helden J, Thomas-Chollier M, Medina-Rivera A. 2022. RSAT 2022: regulatory sequence analysis tools. Nucleic Acids Res doi:10.1093/nar/gkac312.

15. Lu Z, Berry K, Hu Z, Zhan Y, Ahn TH, Lin Z. 2021. TSSr: an R package for comprehensive analyses of TSS sequencing data. NAR Genom Bioinform 3:lqab108.

16. Majerczyk C, Brittnacher M, Jacobs M, Armour CD, Radey M, Schneider E, Phattarasokul S, Bunt R, Greenberg EP. 2014. Global analysis of the *Burkholderia thailandensis* quorum sensing-controlled regulon. J Bacteriol 196:1412-24.

17. Short SM, van Tol S, MacLeod HJ, Dimopoulos G. 2018. Hydrogen cyanide produced by the soil bacterium *Chromobacterium* sp. Panama contributes to mortality in *Anopheles gambiae* mosquito larvae. Sci Rep 8:8358.

18. Mun W, Kwon H, Im H, Choi SY, Monnappa AK, Mitchell RJ. 2017. Cyanide production by *Chromobacterium piscinae* shields It from *Bdellovibrio bacteriovorus* HD100 predation. mBio 8.

19. Smalley NE, An D, Parsek MR, Chandler JR, Dandekar AA. 2015. Quorum sensing protects *Pseudomonas aeruginosa* against cheating by other species in a laboratory coculture model. J Bacteriol 197:3154-9.

20. Gilchrist FJ, Alcock A, Belcher J, Brady M, Jones A, Smith D, Spanel P, Webb K, Lenney W. 2011. Variation in hydrogen cyanide production between different strains of *Pseudomonas aeruginosa*. Eur Respir J 38:409-14.

21. Chen G, Swem LR, Swem DL, Stauff DL, O'Loughlin CT, Jeffrey PD, Bassler BL, Hughson FM. 2011. A strategy for antagonizing quorum sensing. Mol Cell 42:199-209.

22. Stauff DL, Bassler BL. 2011. Quorum sensing in *Chromobacterium violaceum*: DNA recognition and gene regulation by the CviR receptor. J Bacteriol 193:3871-8.

23. Valente RS, Nadal-Jimenez P, Carvalho AFP, Vieira FJD, Xavier KB. 2017. Signal integration in quorum sensing enables cross-species induction of virulence in *Pectobacterium wasabiae*. mBio 8.

24. Hosni T, Moretti C, Devescovi G, Suarez-Moreno ZR, Fatmi MB, Guarnaccia C, Pongor S, Onofri A, Buonaurio R, Venturi V. 2011. Sharing of quorum-sensing signals and role of interspecies communities in a bacterial plant disease. ISME J 5:1857-70.

25. Dulla GF, Lindow SE. 2009. Acyl-homoserine lactone-mediated cross talk among epiphytic bacteria modulates behavior of *Pseudomonas syringae* on leaves. ISME J 3:825-34.

26. Riedel K, Hentzer M, Geisenberger O, Huber B, Steidle A, Wu H, Hoiby N, Givskov M, Molin S, Eberl L. 2001. N-acylhomoserine-lactone-mediated communication between *Pseudomonas aeruginosa* and *Burkholderia cepacia* in mixed biofilms. Microbiology (Reading) 147:3249-62.

27. Wellington S, Greenberg EP. 2019. Quorum sensing signal selectivity and the potential for interspecies cross talk. mBio 10.

28. Collins CH, Arnold FH, Leadbetter JR. 2005. Directed evolution of *Vibrio fischeri* LuxR for increased sensitivity to a broad spectrum of acyl-homoserine lactones. Mol Microbiol 55:712-23.

29. Hawkins AC, Arnold FH, Stuermer R, Hauer B, Leadbetter JR. 2007. Directed evolution of *Vibrio fischeri* LuxR for improved response to butanoyl-homoserine lactone. Appl Environ Microbiol 73:5775-81.

30. Lintz MJ, Oinuma K, Wysoczynski CL, Greenberg EP, Churchill ME. 2011. Crystal structure of QscR, a *Pseudomonas aeruginosa* quorum sensing signal receptor. Proc Natl Acad Sci U S A 108:15763-8.

31. Lozano GL, Guan C, Cao Y, Borlee BR, Broderick NA, Stabb EV, Handelsman J. 2020. A chemical counterpunch: *Chromobacterium violaceum* ATCC 31532 produces violacein in response to translation-inhibiting antibiotics. mBio 11.

32. Wang M, Schaefer AL, Dandekar AA, Greenberg EP. 2015. Quorum sensing and policing of *Pseudomonas aeruginosa* social cheaters. Proc Natl Acad Sci U S A 112:2187-91.

33. Schuster M, Urbanowski ML, Greenberg EP. 2004. Promoter specificity in *Pseudomonas aeruginosa* quorum sensing revealed by DNA binding of purified LasR. Proc Natl Acad Sci U S A 101:15833-9.

34. Pessi G, Haas D. 2000. Transcriptional control of the hydrogen cyanide biosynthetic genes *hcnABC* by the anaerobic regulator ANR and the quorum-sensing regulators LasR and RhlR in *Pseudomonas aeruginosa*. J Bacteriol 182:6940-9.

35. Lopez CM, Rholl DA, Trunck LA, Schweizer HP. 2009. Versatile dual-technology system for markerless allele replacement in *Burkholderia pseudomallei*. Appl Environ Microbiol 75:6496-503.