1
2
3
4 Cross-species activation of hydrogen cyanide production by a promiscuous quorum-sensing
5 receptor promotes *Chromobacterium subtsugae* competition in a dual-species model
6
7

8 Cheyenne Loo[a, b, §], Pratik Koirala[a§], Nathan C. Smith[a], Kara C. Evans[a, c], Saida Benomar[a, d],
9 Isabelle R. Parisi[a], and Anna Oller[e] and Josephine R. Chandler[a, #]
10
11 [§]Contributed equally
12
13 [a]Department of Molecular Biosciences, University of Kansas, Lawrence, KS 66045 USA
14
15 [b]Current location: Stowers Institute for Medical Research, Kansas City, MO 64110 USA
16
17 [c]Current location: International Flavors and Fragrances Health and Biosciences, Madison WI,
18 53716 USA
19
20 [d]Current location: Department of Microbiology & Immunology, Emory University School of
21 Medicine, Atlanta, GA, 30322 USA.
22
23 [e]Department of Biological and Clinical Sciences, University of Central Missouri, Warrensburg,
24 MO 64093 USA
25
26 Running title: *C. subtsugae* eavesdropping and interspecies competition
27
28 [#]Address correspondence to Josephine R. Chandler, jrchandler@ku.edu.

ABSTRACT:

Many saprophytic bacteria have LuxR-I-type acyl-homoserine lactone (AHL) quorum-sensing systems that may be important for competing with other bacteria in complex soil communities. LuxR AHL receptors specifically interact with cognate AHLs to cause changes in expression of target genes. Some LuxR-type AHL receptors have relaxed specificity and are responsive to non-cognate AHLs. These promiscuous receptors might be used to sense and respond to AHLs produced by other bacteria by eavesdropping. We are interested in understanding the role of eavesdropping during interspecies competition. The soil saprophyte *Chromobacterium subtsugae* has a single AHL circuit, CviR-I, which produces and responds to *N*-hexanoyl-HSL (C6-HSL). The AHL receptor CviR can respond to a variety of AHLs in addition to C6-HSL. In prior studies we have utilized a coculture model with *C. subtsugae* and another soil saprophyte *Burkholderia thailandensis*. Using this model, we previously showed that promiscuous activation of CviR by *B. thailandensis* AHLs provides a competitive advantage to *C. subtsugae.* Here, we show that *B. thailandensis* AHLs activate transcription of dozens of genes in *C. subtsugae,* including the *hcnABC* genes coding for production of hydrogen cyanide. We show that hydrogen cyanide production is population density-dependent and demonstrate that the cross-induction of hydrogen cyanide by *B. thailandensis* AHLs provides a competitive advantage to *C. subtsugae.* Our results provide new information on *C. subtsugae* quorum sensing and are the basis for future studies aimed at understanding the role of eavesdropping in interspecies competition.

IMPACT STATEMENT:

In quorum sensing, population density-dependent changes in gene regulation are the result of a

cytoplasmic transcription regulator binding to a quorum sensing signal. The signal-receptor

interaction is considered to be specific to ensure fidelity of the system. However, some quorum-

sensing receptor proteins have relaxed specificity and can recognize and respond to a range of

signals. These promiscuous receptors might provide some benefit by enabling interspecies

activation of quorum sensing by "eavesdropping," although the potential benefits of

eavesdropping are not well studied. The current study utilizes a dual-species laboratory

competition model, where one species has a promiscuous signal receptor and can respond to

signals produced by the other species. In our study, we identify the signals that enable quorum

sensing cross talk and show that cross talk promotes competition by inducing hydrogen cyanide

production. Our results highlight how quorum sensing-enabled interspecies cross talk might

provide an advantage during competition and provide a new basis for understanding how

receptor-signal pairs might evolve in natural environments.

82 **DATA SUMMARY**

83 The authors confirm all supporting data, code and protocols have been provided within the

84 article or through supplementary data files.

INTRODUCTION:

Quorum sensing is a population density-dependent signaling system that relies on small diffusible signaling molecules. Once a critical population threshold is reached, the molecules interact with cytoplasmic transcription factors to cause activation of specific genes and coordinate gene expression across the population. In *Proteobacteria*, one type of quorum sensing is carried out by acyl-homoserine lactones (AHLs) signals. These molecules have a lactone ring and a fatty acid tail that varies in length from 4 to 20 carbons and can also have modifications at the third carbon position, which may be unsubstituted, hydroxylated, or have an oxo-substitution. Typical AHL quorum-sensing gene circuits consist of a LuxR-family signal receptor and a LuxI-family signal synthase, which produces a cognate signal. A bacterium may possess multiple LuxR-I circuits that typically produce and respond to different AHLs. Although the AHLs generally have similar structures, the receptors are thought to interact with AHLs in a highly specific and selective manner (1). However, there are exceptions such as the CviR receptor of *Chromobacterium subtsugae* (formerly *C. violaceum*), which can promiscuously detect and respond to a range of AHLs (2, 3). The existence of such receptors suggests receptor promiscuity might have certain advantages in some conditions.

Quorum sensing systems control different behaviors in different bacteria, such as the production of toxins, exoproducts, and biofilm matrix components. Many of these behaviors might provide a benefit to individuals within groups or complex communities. For example, toxin production might be important for competition with other strains or species of bacteria. There is an increasing interest in understanding the specific benefits of quorum sensing in dynamic polymicrobial communities. A major barrier to understanding quorum sensing in this context is the need for laboratory models that enable studies of mixed-strain and mixed-species

interactions in a controlled setting. Such models are increasingly being utilized, and are

providing significant new insights into the role and mechanisms of quorum sensing in mixed

microbial communities (for a review, see (4)).

This study utilizes a laboratory dual-species model that was previously developed where

two soil bacteria, *Burkholderia thailandensis* and *Chromobacterium subtsugae,* use AHL-

dependent quorum sensing to compete with the other species (5) (Fig. 1). *B. thailandensis*

produces a ribosome-targeting antibiotic called bactobolin (6). Bactobolin is regulated by a

LuxR-I system called BtaR2-I2, which senses and responds to 3OHC8-HSL and 3OHC10-HSL

(7). Both bactobolin and the BtaR2-I2 system are important for *B. thailandensis* to compete with

C. subtsugae (5). In addition to BtaR2-I2, *B. thailandensis* has two other complete quorum-

sensing circuits, BtaR1-I1 and BtaR3-I3, which produce and respond to C8-HSL and 3OHC10-

HSL, respectively (8). *C. subtsugae* has a single LuxR-I-type system, CviR-I, which produces

and responds to C6-HSL (2). CviR-I is important for competition with *B. thailandensis* (5). CviR

is known to activate production of a purple pigmented antibiotic, violacein, however mutational

disruption of violacein had no effect on competition (5). We also showed that CviR can be

activated by *B. thailandensis* AHLs by "eavesdropping" to promote the competitive ability of *C.*

subtsugae (5).

In this study, we identify C8-HSL as the *B. thailandensis* AHL best able to activate *C.*

subtsugae production of quorum-dependent antimicrobials. We use RNAseq to identify the full

set of *C. subtsugae* genes that can be activated by either C8-HSL or the native signal, C6-HSL.

Among the genes most highly activated by both signals are those coding for hydrogen cyanide

biosynthesis. We use our coculture model to validate the importance of hydrogen cyanide for *C.*

subtsugae competition. We also demonstrate that the hydrogen cyanide genes can be

transcriptionally activated by CviR and both native and non-native signals. The results provide

new information on promiscuous AHL responses in *C. subtsugae* and further support the idea

that AHLs drive important interactions through interspecies crosstalk.

METHODS:

Bacterial culture conditions and reagents. All strains were grown in Lysogeny Broth

(LB) broth (10g tryptone, 5g yeast extract, and 10g NaCl in 1L water), LB with morpholine-

propanesulfonic acid (LB-MOPS, 50mM; pH 7), or in LB with 1.5% (wt/vol) Bacto-Agar. All

broth cultures were incubated at 30 °C (*C. subtsugae* or *C. subtsugae-B. thailandensis*

cocultures) or 37 °C (*B. thailandensis* or *E. coli*). Synthetic acyl-homoserine lactone signals were

purchased from Cayman Chemicals (MI, USA) and stored in ethyl acetate acidified with 0.1% of

glacial acetic acid. In all cases, synthetic signals were added to the culture flask and dried down

completely using a stream of nitrogen gas prior to adding the culture media. For strain

constructions, we used gentamicin at 50 μg ml^{-1} (*C. subtsugae*) or 15 μg ml^{-1} (*E. coli*). For

selection from cocultures, we used gentamicin at 100 μg ml^{-1} (*B. thailandensis*) and

trimethoprim at 100 μg ml^{-1} (*C. subtsugae*).

Genomic DNA, PCR and DNA fragments, and plasmid DNA were purified by using a

Puregene Core A kit, plasmid purification miniprep kit, or PCR cleanup/gel extraction kits

(Qiagen or IBI-MidSci) according to the manufacturer's protocol. RNA was isolated using the

Qiagen RNEasy Minikit.

Bacterial strains and plasmids. Strains are summarized in Table 1. For cocultures, we used *B.*

thailandensis strains JBT125 and BD20. Both are derived from strain E264 (9); BD20 is an

bactobolin-deficient *btaK* mutant (7), and JBT125 is an AHL- and bactobolin-deficient *bta1-3,*

btaK quadruple mutant (8). *C. subtsugae* strain CV017 (referred to as wild type, previously

known as *C. violaceum* CV017) is a derivative of strain ATCC 31532 with a transposon insertion

in gene CV_RS05185 causing overexpression of violacein (2). All *C. subtsugae* mutant strains

were constructed from CV017 using allelic exchange and methods described below. For plasmid

construction, we used *E. coli* strain DH10B (Invitrogen).

Unmarked, in-frame deletions of *cviR, cviI,* and *hcnB* (Cv017_06925) were constructed

using the following method. DNA fragments were generated by PCR or DNA synthesis

(Genscript, New Jersey) containing ~500-1000 bp DNA flanking each gene and fused together,

creating an unmarked, non-polar deletion of each gene with incorporated XbaI and SacI

restriction enzymes sites. These fragments were digested with XbaI and SacI and cloned into

XbaI, SacI-digested pEXG2 (10). The relevant pEXG2 gene deletion plasmid was subsequently

used to make the mutant *C. subtsugae* strain using methods described previously (5). Briefly, the

relevant pEXG2 plasmid was transformed into *C. subtsugae* by electroporation or conjugation.

Merodiploids were selected on LB agar containing gentamicin, and deletion mutants were

counterselected on LB + 15% sucrose. Mutant strains were verified by testing for gentamicin

sensitivity and by PCR-amplifying the deletion region and sequencing the PCR product.

For transcription reporter assays, plasmid pMP220 *hcnA-lacZ* was made by amplifying the *hcnA*

promoter from *C. subtsugae* with primers hcn220R (5'-

TATATCTAGAGATTAGCGGTTTGCCGTTCA-3') and hcn220F (5'-

TATATCTAGAGAAGCACCTTGCTGTGGTGAATG -3'), which incorporated the XbaI and

KpnI sites. The amplified product was digested with XbaI and KpnI and ligated to XbaI- and

KpnI-digested pMP220 (11). pMP220 *hcnA-lacZ* was introduced to *C. subtsugae* Δ*cviI* and

Δ*cviR* by electroporation.

RNA isolation. To isolate RNA for RNAseq, cultures were grown to an OD_{600} of 4.0 in 10 ml

LB-MOPS in 125 ml non-baffled flasks. Bacteria (~1 × 10^9 cells) were removed from the

cultures and suspended in RNAprotect bacteria reagent (Qiagen), pelleted by centrifugation, and

stored at −80°C. Thawed cells were suspended in 1 ml RLT buffer (Qiagen) containing 2-

mercaptoethanol and lysed by bead beating. RNA was purified using the RNAeasy minikit

(Qiagen). Contaminating DNA was removed with Turbo DNase (Ambion), and RNA was

obtained by using RNeasy MinElute cleanup kit (Qiagen).

RNA-seq library construction, sequencing, mapping and analysis. Libraries for RNA-seq

were prepared by Novogene using the Novogene mRNA-seq services project workflow. cDNA

libraries were sequenced by Novogene using an Illumina platform and PE-150 reads. Sequences

were aligned and mapped to the *C. subtsugae* CV017 genome (accession

NZ_JAHDTB000000000) using Featurecounts and the Rsubread package in R (12). We

determined differentially regulated genes for biological replicates using function edgeR (13) and

using a false discovery rate [FDR] cutoff of 0.05. We proceeded with genes with >15 aligned

reads and 2-fold or more regulation by C6- or C8-HSL relative to no AHL. Functions and their

sources can be found at https://github.com/ChandlerLabKU/Cs_rnaseq. The data have been

deposited in the NCBI sequence read archive (SRA database) (BioProject identification [ID]

PRJNA894552). For Fig. 6, promoter sites and DNA recognition motifs were predicted using

RSAT (14) and the transcription start site in the DNA region upstream of the *C. subtsugae hcnA*

gene was predicted from the RNAseq data generated in this study using the TSSr program in R

(15).

Coculture experiments. Coculture experiments were conducted in 20 mL LB-MOPS medium in

125 mL non-baffled flasks. The inoculum was from logarithmic-phase pure cultures of *C.*

subtsugae and *B. thailandensis*. The initial OD_{600} in the coculture was 0.05 for *B. thailandensis*

(2—4x10^7 cells per ml) and 0.005 for *C. subtsugae* (2 – 4x10^6 cells per mL). After inoculating,

cocultures were incubated at 30 ºC with shaking at 250 rpm for 24 hours. Colony forming units

of each species were determined by using differential antibiotic selection on LB agar plates. *B.*

thailandensis was selected with gentamicin and *C. subtsugae* was selected with trimethoprim.

Transcription reporter assays. To assess activation of the *hcnA* promoter in *C. subtsugae,*

overnight cultures of *C. subtsugae* strains carrying the pMP220_*hcn-lacZ* reporter plasmid were

used as starters by diluting to an OD_{600} of 0.05. When experimental cultures reached an OD_{600} of

0.5 they were distributed to 2 ml Eppendorf tubes containing different concentrations of dried

C6-HSL, C8-HSL, 3OHC8-HSL, and 3OHC10-HSL. The volume in each tube was 0.5 ml. After

5 h with shaking at 30°C, β-galactosidase activity was measured using the Tropix Galacto-Light

Plus™ chemiluminescent kit according to the manufacturer's protocol (Applied Biosystems,

Foster City, CA).

Cyanide measurements. Cyanide concentrations were measured using a CN^- ion-selective

electrode (Cole-Parmer, USA). Cells were clarified from culture fluid using centrifugation

followed by filter sterilization and the cell-free fluid was brought to a pH of 12 using NaOH

prior to measugin the CN^- ion by conductivity. A standard curve using KCN was similarly

measured and used to calculate the CN^- concentration in samples.

RESULTS:

C. subtsugae **response to** *B. thailandensis* **AHLs in coculture.** Previously, we showed that

extracted fluid from *B. thailandensis* cultures containing all three *B. thailandensis* AHLs (C8-

HSL, 3OHC8-HSL and 3OHC10-HSL) increase *C. subtsugae* competitiveness in a manner

dependent on the *C. subtsugae* quorum-sensing receptor CviR (5). These results suggest that *C.*

subtsugae CviR is responsive to one or several of the *B. thailandensis* AHLs. Therefore, we

carried out coculture experiments to determine the individual contribution of each of the *B.*

thailandensis AHL(s) to *C. subtsugae* competitiveness. We used signal synthase mutants of *C.*

subtsugae and *B. thailandensis* for the competition experiment and added each of the *B.*

thailandensis synthetic AHLs individually at 1 uM AHLs, which is a concentration in the range

of that reported for similarly-grown pure cultures of *B. thailandensis* (16). The *B. thailandensis*

strain JBT125 was also bactobolin-defective (Δ*btaK*) so that *C. subtsugae* would not be killed by

AHL-induced bactobolin in the experiment. In our cocultures experiment we observed that the *C.*

subtsugae native signal C6-HSL improved *C. subtsugae* competitiveness by ~100-fold compared

with the no signal condition (Fig. 2), similar to our prior study (5). Adding C8-HSL and 3OHC8-

HSL also significantly improved competitiveness of *C. subtsugae* although to a lesser degree

than the native C6-HSL. 3OHC10-HSL had a more variable effect than the other AHLs and was

not significantly different from no AHLs, although this signal also appeared to improve

competitiveness. Together, these results show that *C. subtsugae* is responsive to the *B.*

thailandensis AHLs C8-HSL and 3OHC8-HSL, and possibly also to 3OHC10-HSL.

Production of hydrogen cyanide is induced by quorum sensing and *hcnB*. We used RNAseq

transcriptomic analysis to identify AHL-activated *C. subtsugae* genes as a first step to identify

the factor important for competition with *B. thailandensis*. We compared transcripts in our Δ*cviI*

strain grown with or without exogenously added synthetic AHLs. We grew cultures with either

C6-HSL or C8-HSL, with the goal of identifying a set of genes with the ability to respond to

both signals. We anticipated that the gene(s) required for *C. subtsugae* to compete via

eavesdropping would be among this gene set. We collected cells for our analysis during the

transition from logarithmic growth to stationary phase, which corresponds with an optical

density at 600 nm (OD_{600}) of 4.0. AHLs were added at 2 μM to ensure robust responses for

detecting differences in gene expression.

Overall, 348 *C. subtsugae* genes were activated by C6-HSL, and 97 of these genes were also

activated by C8-HSL (Table S1). In the set of genes most highly activated by both C6-HSL and

C8-HSL (Table 2), we identified *hcnA*, *hcnB* and *hcnC,* which are predicted to code for

biosynthesis of hydrogen cyanide. Hydrogen cyanide is important for *C. subtsugae* to protect

itself from predation by other bacteria and kill mosquito larvae (17, 18), and is also important for

P. aeruginosa to compete with *B. multivorans* in cocultures (19). Therefore, we focused our

attention on hydrogen cyanide. To determine whether *C. subtsugae* produces hydrogen cyanide,

we measured cyanide ion (CN^-) in cell-free fluid from wild-type *C. subtsugae* cultures grown to

various growth stages (early logarithmic, late logarithmic, stationary phase). The maximum

concentration of CN^- was 3.3 ± 1.1 mM at an OD_{600} ~8 (Fig. 3A). When adjusted for growth, the

production of CN^- was population density-dependent (Fig. 3A), consistent with quorum sensing

activation of these genes. To determine if either the *hcn* genes or quorum sensing was required

for hydrogen cyanide production, we introduced a Δ*hcnB* mutation to the wild type genome and

measured CN^- in culture fluid of the Δ*hcnB* and Δ*cviR* mutants and wild type. Growth-adjusted

CN^- levels were low in both Δ*hcnB* and Δ*cviR* culture fluid compared with that of wild type (Fig.

3B). These results support the conclusion that *hcnB* and quorum sensing are important for

hydrogen cyanide production in *C. subtsugae.*

Cyanide is important for *C. subtsugae* to compete with *B. thailandensis.* Because expression

of the hydrogen cyanide biosynthesis genes is responsive to both C6-HSL and C8-HSL, we

hypothesized that *B. thailandensis* AHLs increase competitiveness of *C. substugae* by activating

hydrogen cyanide production through eavesdropping. As a first test of this hypothesis, we

determined the susceptibility of *B. thailandensis* strain BD20 to a range of concentrations of

potassium cyanide (0-750 uM), which mimics cyanide ions (20). We determined the 50% lethal

dose (LD_{50}) of potassium cyanide for *B. thailandensis* BD20 to be 119 μM (Fig. 4A). Thus, the

concentration of cyanide produced by *C. subtsugae* (~3 mM, Fig. 2A) is more than sufficient to

kill *B. thailandensis.*

Next, we investigated the effects of hydrogen cyanide in cocultures (Fig. 4B). We

competed the bactobolin-deficient *B. thailandensis* (BD20) with different genetic mutants of *C.*

subtsugae. In these conditions, *B. thailandensis* BD20 produces AHLs that can activate the

quorum-sensing system of *C. subtsugae* (5). Consistent with prior results (5), in these conditions

the *C. subtsugae* ΔcviI mutant and wild type are similarly competitive, because the ΔcviI mutant

has CviR that can be activated by *B. thailandensis* AHLs. The ΔcviR mutant cannot respond to

any AHLs and, as expected, was significantly less competitive than the wild-type and ΔcviI

mutant. To determine the role of hydrogen cyanide in competition, we tested the competitive

ability of the ΔhcnB mutant. The ΔhcnB single mutant competed poorly with BD20 and was

similar to the ΔcviR mutant. These results support that hydrogen cyanide is important for the

competitive ability of *C. subtsugae.* We also tested the ΔcviI, ΔhcnB double mutant. In this

strain, the competitive ability of *C. subtsugae* relies on AHLs produced by *B. thailandensis*. The

Δ*cviI*, Δ*hcnB* double mutant also competed poorly, similar to the Δ*cviR* and Δ*hcnB* single

mutants. These results support that hydrogen cyanide is important for *C. subtsugae* to compete in

response to *B. thailandensis* AHLs. Together, the results support the idea that hydrogen cyanide

is required for the competitive advantage provided to *C. subtsugae* in response to native AHLs

and non-native AHLs.

Sensitivity of *hcnA* promoter to AHLs. The hydrogen cyanide biosynthesis genes *hcnA, hcnB*

and *hcnC* (the *hcn* genes) are adjacently encoded in the genome facing the same direction. There

is a single putative promoter upstream of *hcnA*. We hypothesized that this promoter is activated

either directly or indirectly by CviR in response to the *C. subtsugae*-produced signal C6-HSL, as

well as the *B. thailandensis* signals C8-HSL, 3OHC8-HSL, and possibly 3OHC10-HSL. To test

our hypothesis, we created a plasmid with a DNA fragment containing the putative promoter

region upstream of *hcnA* fused to a promoterless *lacZ* gene. This plasmid was introduced into the

C. subtsugae Δ*cviI* and Δ*cviR* mutants. The *hcnA-lacZ* reporter was activated by 5 µM of each of

the four AHLs (C6-, C8-, 3OHC8- and 3OHC10-HSL) in the Δ*cviI* mutant, but not in the Δ*cviR*

mutant (Fig. 5A). Thus, activation of *hcnA-lacZ* required CviR and AHLs. We also determined

the signal sensitivity of the *hcnA-lacZ* reporter by testing each AHL at a range of concentrations

(Fig. 5B). The *hcnA-lacZ* reporter responded most sensitively to the native signal C6-HSL. There

was also a response to C8-HSL. The other two AHLs, 3OHC8-HSL and 3OHC10-HSL,

activated the reporter only at the highest signal concentrations (1-5 µM). These results were

consistent with the coculture results (Fig. 2), and demonstrate that C8-HSL and, to a lesser extent

3OHC8-HSL and 3OHC10-HSL, can activate the *hcn* gene promoter in a manner dependent on

CviR.

DISCUSSION:

The *B. thailandensis-C. subtsugae* model system was developed previously to examine

the effects of quorum sensing and AHL-dependent eavesdropping in competition (5). In those

studies, we elucidated the importance of *B. thailandensis* bactobolin for competition, however,

the *C. subtsugae* toxin remained elusive. Here, we identified hydrogen cyanide as the *C.*

substugae toxin, filling in this gap in knowledge. We have also shown that noncognate AHLs

can induce dozens of genes in addition to those coding for hydrogen cyanide. Results with our

coculture model support the idea that promiscuous receptors might enable interspecies

interactions by coordinating broad changes in gene expression in response to AHLs from other

species. Results of our studies also expand the list of genes known to be activated by the CviR-I

system in *C. subtsugae.*

The *C. substugae* CviR receptor can activate certain genes in response to a wide range of

AHLs (2). Although detailed structure-function studies of CviR and different AHLs have been

carried out (21, 22), few studies have addressed the potential benefits of AHL receptor

promiscuity for the cell. Receptor activation by non-self AHLs could enable detection of other

bacteria and might provide certain benefits for interspecies competition (5) or cooperation (23).

Indeed, there is evidence of eavesdropping in natural polymicrobial communities in plants (23-

25), and eavesdropping has also been shown to occur in communities relevant to infections (26).

Recent studies also support the idea that promiscuity might be a feature of a wider range of

receptors than previously recognized (27). These studies support the idea that AHL-dependent

cross talk might be an important but underrecognized facet of quorum sensing communication

and that it warrants further study.

We find it interesting that in our study, the majority of genes most strongly activated by

both C6-HSL and C8-HSL in *C. subtsugae* are predicted to code for antibiotic production (Table

2). The receptor BtaR2 in *B. thailandensis* is also highly promiscuous and regulates production

of a potent and broad-spectrum antibiotic, bactobolin (6, 7). The role of these receptors in

regulating antibiotic production supports the idea that promiscuous activation of receptors could

play an important role in competition. The promiscuous activation of CviR by C8-HSL leads to

the induction of a subset of genes that overlaps with that controlled by C6-HSL. It may be that

these genes are more highly sensitive to CviR, enabling response to CviR activation by non-

native signals. An alternative possibility is that C8-HSL changes CviR recognition of gene

promoters, possibly through allosteric changes in the DNA binding motif. Future studies focused

on the regulation of particular genes by different AHLs could help to elucidate the mechanism of

CviR activation of gene promoters by non-native AHLs.

Prior studies show that receptor selectivity falls on a continuum from highly selective

(e.g. *P. aeruginosa* RhlR) to intermediate (e.g. *P. aeruginosa* LasR) to highly promiscuous (e.g.

CviR, and *B. thailandensis* BtaR2) (27). The affinity of receptors for a particular AHL can be

influenced by changing specific residues within the AHL binding pocket of the receptor (21, 28-

30). For example, although CviR is normally inhibited by the noncognate signal C10-HSL,

replacing methionine at position 89 with serine converts this receptor to become C10-HSL-

responsive (21). These results suggest that a particular receptor could easily adapt to become

more or less promiscuous dependent on the benefits of promiscuity. In addition, receptor

selectivity is influenced by receptor expression levels (27). Although very little is known about

the environmental conditions that regulate quorum sensing receptors, one can imagine that

conditions that increase receptor levels in the cell could increase AHL promiscuity. For example,

it was recently demonstrated that the CviR receptor is activated by certain antibiotics (31),

although the effects of antibiotics on receptor selectivity remain to be tested.

The results of this study are consistent with prior studies (17, 19, 32) supporting a role for

hydrogen cyanide in competition. In *P. aeruginosa,* the hydrogen cyanide biosynthesis genes are

part of an operon driven by a promoter upstream of *hcnA* (see Fig. 6). The *P. aeruginosa hcnA*

promoter is quorum-sensing controlled, and has an experimentally validated binding site for the

quorum sensing receptor LasR at position -101 relative to the translation start site (33). This

region also has a binding site for the anaerobic regulator (Anr) at position -72 (33, 34) (Fig. 6).

The intergenic region upstream of *hcnA* in *P. aeruginosa* and *C. substugae* share 45% identity;

however, we were unable to find evidence of a *lux* box or Anr binding site anywhere in the *C.*

substugae hcnA promoter. It is possible that CviR can recognize the *hcnA* promoter in the

absence of a recognizable *lux* box, which has been reported in some cases for LasR (33). It is

also possible that CviR controls the *hcn* genes indirectly. Either way, our results provide

additional support for the idea that CviR can activate production of hydrogen cyanide in response

to AHL from other species, supporting the idea that some quorum sensing-dependent

competitive behaviors might be induced by interspecies cross talk.

AUTHOR STATEMENTS

Conflict of interest

The authors declare no conflicts of interest.

Funding information

This work was supported by the National Institutes of Health (NIH) R35GM133572 to J.R.C.

S.B. and C.L. were supported by NIH K-INBRE fellowships (P20GM103418), K.C.E. was

supported by the NIH Chemical Biology Training Program (T32 GM08545), and P.K. was

supported by the KU Weaver Fellowship. We also thank the KU Genomics Core, which is

supported by NIH grants P20GM103638 (COBRE CMADP) and P20GM103418 (K-INBRE).

Acknowledgements

We acknowledge Drs. Ajai Dandekar and Amy Schaefer for helpful discussions and Nicole

Smalley for technical input. BioRender was used for the creation of Fig. 1.

426
427 **TABLES AND FIGURES**

428
Table 1. Bacterial strains and plasmids used in this study.

───
Strain or plasmid   Relevant properties[a]   Reference or source
───
Strain

B. thailandensis

JBT125   *bta1-3, btaK* quadruple mutant of strain E264   (8)

BD20   *btaK* mutant of strain E264   (7)

C. substugae

CV017   Mini-Tn5 mutant of ATCC 31532, 'wild type'   (2)

CV017 Δ*cviI*   CV017 with an in frame, unmarked *cviI* deletion   This study

CV017 Δ*cviR*   CV017 with an in frame, unmarked *cviR* deletion   This study

CV017 Δ*hcnB*   CV017 with an in frame, umarked *hcnB* deletion   This study

CV017 Δ*cviI*, Δ*hcnB* CV017 Δ*cviI* with the Δ*hcnB* mutation   This study

Escherichia coli

DH5α   [F- φ80 lacZΔM15] Δ(lacZYA-argF)U169 recA1   Invitrogen

445   endA1 hsdR17, [rK–mK+] supE44 thi-1 gyrA relA1

446 Rho-3   thi-1 thr-1 leuB26 tonA21 lacY1 supE44 recA   (35)

447   integrated RP4-2 Tcr::Mu (λpir+)Δasd::FRT ΔaphA::FRT

449 Plasmids

450	pEXG2	Suicide vector, Gm^R	(10)
451	pEXG2 Δ*cviI*	pEXG2 containing Δ*cviI* with flanking sequences	This study
452	pEXG2 Δ*hcnB*	pEXG2 containing Δ*hcnB* with flanking sequences	This study
453		with regard to the *cviI* translation start site	
454	pMP220	lacZ transcriptional reporter vector, IncP, Tc^R	(11)
455	pMP220_hcn-lacZ	pMP220 with the *hcnA* promoter extending from	This study
456		-1 to -500 relative to the translation start site, Tc^R	

457 ^{a}Tc^R, tetracycline resistant; Gm^R, gentamicin resistant.

Table 2. Genes most highly activated by C6-HSL and C8-HSL

Locus tag[a]	Description	Fold change[b]	
		C6-HSL	C8-HSL
KIF53_RS20405	amino acid adenylation domain-containing protein	7.4	4.8
KIF53_RS13810	activator protein	7.3	4.4
KIF53_RS06805	NAD(P)/FAD-dependent oxidoreductase HcnB	6.9	6.9
KIF53_RS20400	hybrid non-ribosomal peptide synthetase/type I polyketide synthase	6.5	5.0
KIF53_RS06810	cyanide-forming glycine dehydrogenase subunit HcnA	6.3	6.4
KIF53_RS21020	PfaD family polyunsaturated fatty acid/polyketide biosynthesis protein	6.2	3.0
KIF53_RS14555	DUF1842 domain-containing protein	6.2.	3.3
KIF53_RS01860	long-chain-fatty-acyl-CoA reductase	5.9	2.1
KIF53_RS14680	U32 family peptidase	5.5	2.3
KIF53_RS06800	cyanide-forming glycine dehydrogenase subunit HcnC	5.4	5.6
KIF53_RS05635	FAD-dependent oxidoreductase	5.4	4.1
KIF53_RS15685	M9 family metallopeptidase N-terminal domain-containing protein	5.0	3.7

482	KIF53_RS16340	ornithine carbamoyltransferase	5.2	5.2
483	KIF53_RS06795	ABC transporter substrate-binding protein	4.7	4.6
484	KIF53_RS02825	hydrogenase maturation protein	4.5	3.2
485	KIF53_RS15785	methyl-accepting chemotaxis protein	4.4	2.6

486 [a]Locus tags correspond to the *C. subtsugae* CV017 genome (accession

487 NZ_JAHDTB000000000.1)

488 [b]Fold change value of induction by AHLs (C6-HSL or C8-HSL) when added to strain CV017

489 Δ*cviI*.

490
491
492

Fig. 1. *Burkholderia thailandensis-Chromobacterium subtsugae* **coculture model.** The *B.*

thailandensis quorum sensing system BtaR2-I2 produces and responds to 3OHC8-HSL and

3OHC10-HSL, and activates production of bactobolin antibiotic. *B. thailandensis* also has two

other quorum sensing systems, BtaR1-I1 and BtaR3-I3, which produce C8-HSL and 3OHC8-

HSL, respectively. BtaR2-I2 is important for *B. thailandensis* to compete with *C. subtsugae* due

to activation of bactobolin prodution. The *C. subtsugae* quorum sensing system CviR-I produces

and responds to C6-HSL. CviR-I is important for *C. subtsugae* to compete with *B. thailandensis,*

through an unknown mechanism.

501
502
503

Fig. 2. Contribution of *B. thailandensis* AHLs to *C. subtsugae* competitiveness. Cocultures

were with *C. subtsugae* ΔcviI and *B. thailandensis* ΔbtaI1, ΔbtaI2, ΔbtaI3, ΔbtaK (JBT125).

After 24 h of coculture growth, the ratio of *C. subtsugae* to *B. thailandensis* was determined by

selective plating and colony counts. The dashed line indicates the initial ratio of *C. subtsugae* to

B. thailandensis. The solid line represents the means and the vertical bars shows the standard

error of the mean for each group. Synthetic AHLs were added at the beginning of the coculture

to a final concentration of 1 μM. Statistical comparisons of each condition with no signal

coculture were by one-way ANOVA; ***, $p<0.001$; **, $p<0.005$ and ns, not significant.

Fig. 3. Cyanide measurements in *C. subtsugae* culture fluid. Cyanide (CN⁻) was measured in

culture fluid of *C. subtsugae*. A. Wild type *C. subtsugae* (CV017) was grown to different

population densities as indicated by the optical density at 600 nm (OD_{600}) prior to measuring CN^-

. Data are shown as growth-adjusted CN^-. The total CN^- concentration at OD_{600} of ~8

corresponds with 3.3 ± 1.1 mM. B. The indicated *C. subtsugae* strain was grown for 12 h, which

corresponds with OD_{600} of ~8, prior to measuring population density and CN^-. Data are shown as

the growth-adjusted CN^- as a percentage of that of wild type. In all cases, data represent the

average of 3 biological replicates and the vertical bars represents the standard error of the mean.

527
528
529
530

Fig. 4. Cyanide toxicity and role in cocultures. A. Survival of *B. thailandensis* BD20 grown in

the presence of different concentrations of cyanide for 8 hr from an initial OD_{600} of 0.005. The

population density was measured and *B. thailandensis* (Bt) survival was determined as a percent

of untreated culture grown under identical conditions. B. Cocultures were with *B. thailandensis*

Δ*btaK* (BD20) and the *C. subtsugae* strain indicated. After 24 h of coculture growth, the ratio of

C. subtsugae to *B. thailandensis* was determined by selective plating and colony counts. The

initial ratio of *C. subtsugae* to *B. thailandensis* was 10^{-1}. The solid line represents the means and

the vertical bars shows the standard error for each group. Statistical comparisons of each

condition compared with no signal were by one-way ANOVA; **, $p<0.005$; * = $p < 0.05$ and ns,

not significant.

560
561
562
563

Fig. 5. Sensitivity of the *hcnA* promoter to native and non-native AHLs. A. *C. subtsugae*

Δ*cviI* or Δ*cviR* mutant harboring the *hcnA-lacZ* reporter was treated with 5 µM of AHL (C6-

HSL, C8-HSL, 3OHC8-HSL or 3OHC10-HSL). B. *C. subtsugae* Δ*cviI* harboring the *hcnA-lacZ*

reporter was treated with AHLs at the indicated concentrations. Results show the average of four

independent experiments from two separate days, and the vertical bars represent the standard

error.

580
581
Fig. 6. Hydrogen cyanide biosynthesis genes (*hcnABC*) in *C. subtsugae* and *P. aeruginosa*.

Red arrows and lettering indicate the transcription start site. The arrow is dashed for *C.*

subtsugae as it was predicted to be between nucleotides -79 to -81 (see Methods). Green boxes

and lettering indicate the binding site for the anaerobic regulator (Anr) and the blue box and

lettering indicate the *lux* box of *P. aeruginosa,* and the canonical lux box. Bolding in the

P.aeruginosa lux box sequence indicate nucleotides that are conserved with the canonical *lux*

box. No *lux* box or Fnr/Anr like sequences were identified in the *C. subtsugae* sequence (see

Discussion text).

CITATIONS

1. Hawver LA, Jung SA, Ng WL. 2016. Specificity and complexity in bacterial quorum-sensing systems. FEMS Microbiol Rev 40:738-52.

2. McClean KH, Winson MK, Fish L, Taylor A, Chhabra SR, Camara M, Daykin M, Lamb JH, Swift S, Bycroft BW, Stewart GS, Williams P. 1997. Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones. Microbiology 143 (Pt 12):3703-11.

3. Swem LR, Swem DL, O'Loughlin CT, Gatmaitan R, Zhao B, Ulrich SM, Bassler BL. 2009. A quorum-sensing antagonist targets both membrane-bound and cytoplasmic receptors and controls bacterial pathogenicity. Mol Cell 35:143-53.

4. Abisado RG, Benomar S, Klaus JR, Dandekar AA, Chandler JR. 2018. Bacterial quorum sensing and microbial community interactions. mBio 9:e02331-17.

5. Chandler JR, Heilmann S, Mittler JE, Greenberg EP. 2012. Acyl-homoserine lactone-dependent eavesdropping promotes competition in a laboratory co-culture model. ISME J 6:2219-28.

6. Seyedsayamdost MR, Chandler JR, Blodgett JA, Lima PS, Duerkop BA, Oinuma K, Greenberg EP, Clardy J. 2010. Quorum-sensing-regulated bactobolin production by *Burkholderia thailandensis* E264. Org Lett 12:716-9.

7. Duerkop BA, Varga J, Chandler JR, Peterson SB, Herman JP, Churchill ME, Parsek MR, Nierman WC, Greenberg EP. 2009. Quorum-sensing control of antibiotic synthesis in *Burkholderia thailandensis*. J Bacteriol 191:3909-18.

8. Chandler JR, Duerkop BA, Hinz A, West TE, Herman JP, Churchill ME, Skerrett SJ, Greenberg EP. 2009. Mutational analysis of *Burkholderia thailandensis* quorum sensing and self-aggregation. J Bacteriol 191:5901-9.

9. Brett PJ, DeShazer D, Woods DE. 1998. *Burkholderia thailandensis* sp. nov., a *Burkholderia pseudomallei*-like species. Int J Syst Bacteriol 48 Pt 1:317-20.

10. Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. 2005. ExsE, a secreted regulator of type III secretion genes in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 102:8006-11.

11. Spaink HP, Okker RJ, Wijffelman CA, Pees E, Lugtenberg BJ. 1987. Promoters in the nodulation region of the Rhizobium leguminosarum Sym plasmid pRL1JI. Plant Mol Biol 9:27-39.

12. Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Res 47:e47.

13. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26:139-40.

14. Santana-Garcia W, Castro-Mondragon JA, Padilla-Galvez M, Nguyen NTT, Elizondo-Salas A, Ksouri N, Gerbes F, Thieffry D, Vincens P, Contreras-Moreira B, van Helden J, Thomas-Chollier M, Medina-Rivera A. 2022. RSAT 2022: regulatory sequence analysis tools. Nucleic Acids Res doi:10.1093/nar/gkac312.

15. Lu Z, Berry K, Hu Z, Zhan Y, Ahn TH, Lin Z. 2021. TSSr: an R package for
comprehensive analyses of TSS sequencing data. NAR Genom Bioinform 3:lqab108.

16. Majerczyk C, Brittnacher M, Jacobs M, Armour CD, Radey M, Schneider E,
Phattarasokul S, Bunt R, Greenberg EP. 2014. Global analysis of the *Burkholderia*
thailandensis quorum sensing-controlled regulon. J Bacteriol 196:1412-24.

17. Short SM, van Tol S, MacLeod HJ, Dimopoulos G. 2018. Hydrogen cyanide produced by
the soil bacterium *Chromobacterium* sp. Panama contributes to mortality in *Anopheles*
gambiae mosquito larvae. Sci Rep 8:8358.

18. Mun W, Kwon H, Im H, Choi SY, Monnappa AK, Mitchell RJ. 2017. Cyanide
production by *Chromobacterium piscinae* shields It from *Bdellovibrio bacteriovorus*
HD100 predation. mBio 8:e01370-17.

19. Smalley NE, An D, Parsek MR, Chandler JR, Dandekar AA. 2015. Quorum sensing
protects *Pseudomonas aeruginosa* against cheating by other species in a laboratory
coculture model. J Bacteriol 197:3154-9.

20. Gilchrist FJ, Alcock A, Belcher J, Brady M, Jones A, Smith D, Spanel P, Webb K,
Lenney W. 2011. Variation in hydrogen cyanide production between different strains of
Pseudomonas aeruginosa. Eur Respir J 38:409-14.

21. Chen G, Swem LR, Swem DL, Stauff DL, O'Loughlin CT, Jeffrey PD, Bassler BL,
Hughson FM. 2011. A strategy for antagonizing quorum sensing. Mol Cell 42:199-209.

22. Stauff DL, Bassler BL. 2011. Quorum sensing in *Chromobacterium violaceum*: DNA
recognition and gene regulation by the CviR receptor. J Bacteriol 193:3871-8.

23. Valente RS, Nadal-Jimenez P, Carvalho AFP, Vieira FJD, Xavier KB. 2017. Signal
integration in quorum sensing enables cross-species induction of virulence in
Pectobacterium wasabiae. mBio 8: e00398-17.

24. Hosni T, Moretti C, Devescovi G, Suarez-Moreno ZR, Fatmi MB, Guarnaccia C, Pongor
S, Onofri A, Buonaurio R, Venturi V. 2011. Sharing of quorum-sensing signals and role
of interspecies communities in a bacterial plant disease. ISME J 5:1857-70.

25. Dulla GF, Lindow SE. 2009. Acyl-homoserine lactone-mediated cross talk among
epiphytic bacteria modulates behavior of *Pseudomonas syringae* on leaves. ISME J
3:825-34.

26. Riedel K, Hentzer M, Geisenberger O, Huber B, Steidle A, Wu H, Hoiby N, Givskov M,
Molin S, Eberl L. 2001. N-acylhomoserine-lactone-mediated communication between
Pseudomonas aeruginosa and *Burkholderia cepacia* in mixed biofilms. Microbiology
(Reading) 147:3249-62.

27. Wellington S, Greenberg EP. 2019. Quorum sensing signal selectivity and the potential
for interspecies cross talk. mBio 10: e00146-19.

28. Collins CH, Arnold FH, Leadbetter JR. 2005. Directed evolution of *Vibrio fischeri* LuxR
for increased sensitivity to a broad spectrum of acyl-homoserine lactones. Mol Microbiol
55:712-23.

29. Hawkins AC, Arnold FH, Stuermer R, Hauer B, Leadbetter JR. 2007. Directed evolution
of *Vibrio fischeri* LuxR for improved response to butanoyl-homoserine lactone. Appl
Environ Microbiol 73:5775-81.

30. Lintz MJ, Oinuma K, Wysoczynski CL, Greenberg EP, Churchill ME. 2011. Crystal
structure of QscR, a *Pseudomonas aeruginosa* quorum sensing signal receptor. Proc Natl
Acad Sci U S A 108:15763-8.

31. Lozano GL, Guan C, Cao Y, Borlee BR, Broderick NA, Stabb EV, Handelsman J. 2020.
A chemical counterpunch: *Chromobacterium violaceum* ATCC 31532 produces violacein
in response to translation-inhibiting antibiotics. mBio 11:e00948-20.

32. Wang M, Schaefer AL, Dandekar AA, Greenberg EP. 2015. Quorum sensing and
policing of *Pseudomonas aeruginosa* social cheaters. Proc Natl Acad Sci U S A
112:2187-91.

33. Schuster M, Urbanowski ML, Greenberg EP. 2004. Promoter specificity in *Pseudomonas*
aeruginosa quorum sensing revealed by DNA binding of purified LasR. Proc Natl Acad
Sci U S A 101:15833-9.

34. Pessi G, Haas D. 2000. Transcriptional control of the hydrogen cyanide biosynthetic
genes *hcnABC* by the anaerobic regulator ANR and the quorum-sensing regulators LasR
and RhlR in *Pseudomonas aeruginosa*. J Bacteriol 182:6940-9.

35. Lopez CM, Rholl DA, Trunck LA, Schweizer HP. 2009. Versatile dual-technology
system for markerless allele replacement in *Burkholderia pseudomallei*. Appl Environ
Microbiol 75:6496-503.