Podocyte lineage marker expression is preserved across Wilms tumor subtypes and enhanced in tumors harboring the SIX1/2-Q177R mutation

Matthew J. Stevenson, Urvi Patel, Lori L. O'Brien

Affiliations:
Department of Cell Biology and Physiology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

Corresponding author:
Lori L. O'Brien
lori_obrien@med.unc.edu

SUMMARY

Wilms tumors present as an amalgam of varying proportions of three tissues normally located within the developing kidney, one being the multipotent nephron progenitor population. While incomplete differentiation of the nephron progenitors is widely-considered the underlying cause of tumor formation, where this barrier occurs along the differentiation trajectory and how this might promote therapeutic resistance in high-risk blastemal-predominant tumors is unclear. Comprehensive integrated analysis of genomic datasets from normal human fetal kidney and high-risk Wilms tumors has revealed conserved expression of genes indicative of podocyte lineage differentiation in tumors of all subtypes. Comparatively upregulated expression of several of these markers, including the non-canonical WNT ligand *WNT5A*, was identified in tumors with the relapse-associated mutation SIX1/2-Q177R. These findings highlight the shared progression of cellular differentiation towards the podocyte lineage within Wilms tumors and enhancement of this differentiation program through promotion of non-canonical WNT/planar cell polarity signaling in association with SIX1/2-Q177R.

INTRODUCTION

Wilms tumor, the most common childhood kidney cancer is typically diagnosed between 2-5 years of age and accounts for an estimated 5% of all cancers in patients under the age of 14 (Steliarova-Foucher et al 2017, Breslow et al 2006, Hol et al 2018). The histological composition of Wilms tumors resembles that of the normal developing kidney, being comprised of varying degrees of blastemal, epithelial, and stromal tissues. Each of these tissues is thought to represent distinct compartments within the developing kidney: 1) nephron progenitor cells (NPCs), the multipotent progenitor population that gives rise to all epithelial cell types of the nephron, 2) differentiated tubules, and 3) interstitial cells (ICs), respectively (reviewed in Rivera and Huber, 2005). Beyond the morphological similarities, numerous microarray and transcriptomic analyses have revealed gene expression signatures in Wilms tumors resembling those of normal pre- and post-induction NPCs with varying degrees of differentiation, altogether implying stalled nephrogenesis underlies formation of these tumors (Li et al 2002, Fukuzawa et al 2017, Wegert et al 2015, Walz et al 2015, Gadd et al 2012, Gadd et al 2017, Young et al 2018, Trink et al 2018). Despite the apparent morphological and molecular similarities to normal human fetal kidney (hFK) and the low mutational burden characteristic of Wilms tumors and pediatric malignancies in general compared to that of adult cancers, the genetic drivers and molecular mechanisms underlying the genesis of these tumors remain elusive (Wegert et al 2015, Grobner et al 2018, Kandoth et al 2013).

Current treatment of Wilms tumor is generally considered a success with a five-year relative survival over 90% (SEER cancer stats 1975-2017), yet high-risk blastemal predominant tumors continue to present a challenge to therapeutic intervention. From 2001-2012, 20% of patients with blastemal-predominant tumors treated according to International

Society of Paediatric Oncology protocols relapsed within five years of diagnosis and 95% of relapses were distant metastases (Van den Heuvel-Eibrink et al 2015). In patients treated according to Children's Oncology Group guidelines, 88% of relapses of stage III tumors occurred within the first two years after diagnosis, including five of seven blastemal tumors analyzed in that study (Fernandez et al 2018). Five-year overall survival after relapse of favorable histology Wilms tumor is estimated at 60%-70% (Mullen et al 2018), highlighting the need for improved therapies to limit the risk of relapse for all Wilms tumors, but particularly in the case of high-risk blastemal-predominant tumors.

Mutations in NPC-associated transcription factors *SIX1* and *SIX2* (~7%) are most associated with blastemal-predominant tumors (Gadd et al 2017, Walz et al 2015, Wegert et al 2015). Of the variants detected in *SIX1* across several studies, the overwhelming majority were a glutamine to arginine substitution, Q177R. This mutation is significantly associated with relapse having recently been identified in ~13% of relapsed tumors (Gadd et al 2022). Wegert et al provided the first mechanistic investigation of the SIX1-Q177R mutation, demonstrating a unique DNA binding motif for SIX1-Q177R in primary Wilms tumor tissue compared to that of wild type SIX1 in a similar tumor. Increased binding of SIX1-Q177R was observed near the *TGFA* gene with corresponding increases in expression of *TGFA* in SIX1-Q177R mutant tumors compared to wild type SIX1-expressing tumors (Wegert et al 2015). More broadly however, the consequences of this altered DNA binding on the fidelity of downstream transcriptional regulatory networks and its potential role in Wilms tumorigenesis and relapse remain unexplored.

Six1 is required for kidney development in the mouse as knockout results in kidney agenesis and Six1 acts upstream of canonical NPC markers including Six2 and Pax2, placing it near the top of the NPC gene regulatory network hierarchy (Xu et al 2003, Li et al 2003). Comparative investigations of mouse and human kidney development demonstrated prolonged temporal expression of *SIX1*/SIX1 in NPCs through later stages of development in addition to novel regulatory interactions between SIX1 and SIX2 not observed in the mouse, suggestive of an expanded regulatory role for SIX1 in human nephrogenesis that may underlie its frequent mutation and association with high-risk Wilms tumors (O'Brien et al 2016).

Over the last several years, pioneering studies of human kidney development through morphological and transcriptomic analyses have characterized the morphogenesis and molecular signatures intrinsic to the progression from proliferative, multipotent NPCs to the complex multifaceted structure of the nephron. These investigations provided the spatial resolution necessary to more precisely distinguish the transcriptomes of cells in the renal vesicle (RV), the earliest epithelial descendent of the NPCs, from polarized constituents of the S-shaped body (SSB) destined to differentiate to tubules or podocytes (Lindstrom et al 2018a-d, Lindstrom et al 2021, Hochane et al 2019, Menon et al 2018, Tran et al

2019). Consequently, these enhanced molecular profiles can be repurposed to refine the molecular identity of cells in Wilms tumors.

The goal of this study is to expand upon previous works to attempt to define the direct regulatory role of SIX1-Q177R in Wilms tumor within the context of the normal human nephrogenic niche utilizing genomic data available from large-scale studies of Wilms tumors, in tandem with SIX1 ChIP-seq data from week 17 hFK (wk17hFK). Furthermore, we sought to classify the Wilms tumor transcriptomes within the framework of the developing hFK by merging single cell RNA-seq datasets from week 14 hFK (wk14hFK) and wk17hFKs and integrating the findings from our genomic analyses of Wilms tumor (Lindstrom et al 2018d, Lindstrom et al 2021, Tran et al 2019). In doing so, we identify upregulation of genes associated with promotion of the non-canonical WNT/planar cell polarity (PCP) pathway and/or inhibition of the canonical WNT/β-catenin-mediated pathway in SIX1-Q177R-expressing Wilms tumors. Furthermore, we provide a mechanistic link to *WNT5A* upregulation attributable to enhanced binding affinity of SIX1-Q177R for putative cis-regulatory elements (CREs) and demonstrate conserved expression of podocyte lineage markers in all Wilms tumors analyzed, a subset of which are specifically upregulated in SIX1-Q177R-expressing tumors.

RESULTS

Putative SIX1-Q177R target genes are associated with distinct biological processes compared to those of SIX1

Analysis of Wilms tumor SIX1 ChIP-seq peaks containing the previously identified SIX1/SIX1-Q177R DNA binding motifs revealed marked differences in putative regulatory target genes of the two proteins (Figure 1A, Supplemental Figure 1A) (Data kindly provided by Dr. Manfred Gessler, Wegert et al 2015). Comparison with wk17hFK SIX1 ChIP-seq showed modest overlap in peak locations, approximately 20% of peaks from each tumor (Figure 1B). Genomic Regions Enrichment of Annotations Tool (GREAT) (McLean et al 2010) was used to identify putative target genes in each tumor dataset. Gene ontology analysis using GREAT revealed enrichment in kidney development-related biological processes in those putative target genes shared between the SIX1 and SIX1-Q177R tumors. Despite only modest overlap in physical peak locations between SIX1 tumor and wk17hFK, the SIX1 tumor targets are nonetheless enriched for kidney developmental processes. Putative target genes exclusive to SIX1-Q177R, however, are associated with distinct biological processes not explicitly related to kidney development. (Figure 1C).

Intriguingly, the MEIS1 DNA binding motif is significantly enriched in all peak sets analyzed (Supplemental Figure 1B). Expression of *Meis1* is restricted to the interstitial lineage in the developing mouse kidney. However, comparative

FIGURE 1: SIX1-Q177R tumor-only putative target genes minimally overlap with wk17hFK SIX1 peaks and are associated with biological processes unrelated to kidney development. A) SIX1 and SIX1-Q177R DNA binding motifs discovered using the STREME tool from Wilms tumor datasets (Wegert et al 2015, motifs published previously in that study) and week 17 human fetal kidney (wk17hFK) (O'Brien et al 2016). **B)** Grid displaying the pairwise fraction of SIX1 ChIP-seq peak location overlap between SIX1 Wilms tumor, SIX1-Q177R Wilms tumor, and wk17hFK. **C)** Venn diagram displaying overlaps of putative target genes identified using motif-enriched peaks in GREAT between SIX1-Q177R Wilms tumor and SIX1 Wilms tumor. Bar charts displaying top 10 most significantly enriched biological processes for shared tumor, SIX1-Q177R tumor-only, and SIX1 tumor-only putative target genes generated using GREAT.

analysis of marker expression by immunofluorescence and *in situ* hybridization in developing hFK revealed overlap of NPC marker SIX2/*SIX2* with MEIS1/*MEIS1* as well as FOXD1/*FOXD1*, another marker used to distinguish interstitial progenitors from NPCs in the mouse (Lindstrom et al 2018a). Overexpression of either SIX1 or SIX1-Q177R, alongside MEIS1-3xFLAG *in vitro* followed by immunoprecipitation (IP) with a FLAG tag antibody resulted in co-IP of both SIX1 and SIX1-Q177R (Supplemental Figure 1C). To our knowledge this is the first biochemical evidence of an interaction between SIX1 and MEIS1 in any context. Further investigation is needed to characterize potential regulatory functions of complexes containing these proteins in NPCs.

WNT5A and other WNT signaling effectors are upregulated in SIX1/2-Q177R Wilms tumors

To further refine the general peak/gene associations, we utilized RNA-seq data from 114 chemotherapy-naïve high-risk favorable histology Wilms tumors (FHWT) and diffuse anaplastic Wilms tumors (DAWT) of varying histological

classifications that subsequently relapsed, generated as part of the Therapeutically Applicable Research to Generate Effective Treatments (TARGET) program, to pinpoint differentially expressed genes unique to SIX1/2-Q177R tumors. The same Q177R mutation has been identified in SIX2 in blastemal-predominant Wilms tumors but occurs almost half as frequently as SIX1-Q177R (Wegert et al 2015, Walz et al 2015, Gadd et al 2017). SIX2 is a closely related Six family transcription factor to SIX1, acts downstream of SIX1 in kidney development (Li et al 2003), and binds to most of the same genomic sites as SIX1 in hFK (O'Brien et al 2016). Accounting for these similarities between SIX1/SIX2 and to increase the statistical power of our analyses, we grouped SIX1-Q177R and SIX2-Q177R tumors.

Nearly 30% of SIX1/2-Q177R tumors also harbor inactivating mutations in the miRNA processing genes *DROSHA* or *DGCR8* (Wegert et al 2015, Walz et al 2015, Gadd et al 2017). Strikingly, co-occurrence of SIX1/2-Q177R and DROSHA/DGCR8 mutations significantly increased the rates of relapse and death in a synergistic manner compared to tumors with other mutations and tumors with SIX1/2-Q177R or DROSHA/DGCR8 mutations alone (Walz et al 2015). Notably, Wegert et al observed no strong effects on gene expression in DROSHA/DGCR8 mutant tumors, only a significant reduction in miRNA levels (Wegert et al 2015). Due to the scarcity of SIX1/2-Q177R tumors without mutations in miRNA processing genes in this dataset and to best isolate the transcriptional effects attributable to SIX1/2-Q177R, tumors with both SIX1/2-Q177R and miRNA mutations were used in differential expression analysis (SIX1/2miRNA, n=9). Specific histology classification of other tumors used in this analysis was based on prior classification if available (Walz et al 2015, TARGET data matrix https://portal.gdc.cancer.gov/projects), otherwise tumors were included in the group containing mixed histology tumors, generating three additional tumor groups: wild type SIX1-expressing blastemal tumors (Blastemal, n=22), tumors of mixed/epithelial/stromal histology (MIXED/ES, n=56), and DAWT (n=27) (Supplemental Table 1).

Limma-voom was utilized for differential gene expression analysis (Supplemental Figure 2) (Smyth 2005, Law et al 2014, Liu et al 2015). Due to the overwhelming evidence supporting the nephrogenic origin of Wilms tumors (reviewed in Li et al 2021 and Hohenstein et al 2015, Coorens et al 2019), we reasoned an approach focusing on transcripts expressed in all tumor samples could elucidate important deviations along the nephrogenic trajectory between tumors, while also highlighting critical shared intrinsic characteristics of these tumors. Therefore, weakly-expressed genes were stringently filtered to exclude potential false positives due to heterogeneity in tumor microenvironments or sample-to-sample processing variability (genes with counts per million (CPM) < 2 in any sample were removed). Unsupervised hierarchical clustering by Pearson's correlation coefficient using normalized and scaled expression values illustrated a high degree of transcriptional similarity between the majority of tumors in the dataset, regardless of histological classification (Figure 2A). Moreover, tumors with Chromosome 1 q21-q23 gain, recently identified in 75% of relapsed tumors (Gadd et al 2022), did not noticeably cluster together suggesting transcriptional changes associated with this particular copy number gain are minimal or

heterogeneous. As has been noted previously, most SIX1/2miRNA tumors clustered together (Wegert et al 2015, Walz et al 2015).

Few genes were identified as differentially expressed between Blastemal and MIXED/ES groups using a less stringent fold change cutoff, eight genes upregulated and 20 genes downregulated (log2 fold change > 1, adj. p < 0.05) (Supplemental Figure 3A, Supplemental File 1). Compared to Blastemal tumors, 28 genes were significantly upregulated in SIX1/2miRNA tumors, 14 of which were also significantly upregulated compared to MIXED/ES tumors (log2 fold change > 1.5, adj. p <0.05) (Figure 2B, Supplemental Figure 3A, Supplemental File 1). Remarkably, the expression levels of well-characterized nephron progenitor markers including *SIX1*, *SIX2*, *SALL1*, *MEOX1*, *PAX2*, *LYPD1*, and *CRABP2* were not significantly different between the four groups (Supplemental File 1). Thus, blastemal-predominant tumors do not appear to represent an augmentation of the pre-induction NPC transcriptional regulatory identity compared to tumors of other histologies. *CCND2*, encoding cyclin D2, was significantly upregulated in SIX1/2miRNA tumors compared to both Blastemal and DAWT groups and was upregulated 1.46 log2 fold compared to MIXED/ES tumors, similar to findings of other studies

FIGURE 2: Several genes upregulated in SIX1/2miRNA tumors have functions related to enhancement of non-canonical WNT/PCP signaling and/or inhibition of canonical WNT/β-catenin-mediated signaling. A) Heatmap displaying results of unsupervised hierarchical clustering of all tumors used in RNA-seq differential gene expression analysis by Pearson's correlation coefficient. **B)** Violin plots showing the distributions of log2 counts per million (CPM) of genes significantly upregulated in SIX1/2miRNA tumors compared to Blastemal tumors. Dot within each group plot represents the mean. Unless indicated by n.s. (not significant), the log2 fold change of the SIX1/2miRNA group was > 1.5 and adjusted p-value < 0.05 with respect to that tumor group.

(Wegert et al 2015, Walz et al 2015). Nine genes were significantly upregulated in SIX1/2miRNA tumors compared to all groups, including known NPC markers *TMEM100* and *ITGA8.* This suggests the sustained proliferation reflected by enhanced *CCND2* expression in SIX1/2miRNA tumors may be governed by non-transcriptional mechanisms including cell adhesion. Several genes were significantly downregulated in SIX1/2miRNA tumors as well, however we chose to focus on upregulated genes for initial follow-up analyses (Supplemental Figure 3B, Supplemental File 1).

Functional similarities among the upregulated SIX1/2miRNA genes were discovered through literature searches, revealing an enrichment in protein functions either promoting non-canonical WNT/PCP signaling and/or antagonizing canonical WNT/β-catenin signaling. Positive regulators of non-canonical WNT/PCP signaling include non-canonical WNT ligand WNT5A (Qian et al 2007, Nishita et al 2010), IQGAP2 through interaction with Cdc42/Rac1 (Logue et al 2011, Ozdemir et al 2018, Fukata et al 2002), ARHGEF3 through activation of RhoA (D'Amato et al 2015, You et al 2021), and FOXC1 through direct regulation of *WNT5A* expression (Han et al 2018). Negative regulators of canonical WNT/β-catenin signaling include WNT5A through interaction with the receptor Ror2 (Mikels and Nusse 2006)*,* KDM2B through independent demethylation of β-catenin and transcriptional repression of β-catenin target genes (Lu et al 2015, Ladinovich et al 2020)*,* as well as SLIT3 and IQGAP2, as the expression of both has been associated with decreased β-catenin target gene activation or decreased nuclear β-catenin, respectively. (Kim et al 2018, Ng et al 2018, Deng et al 2016). These findings suggest imbalance in WNT/β-catenin signaling in favor of non-canonical WNT/PCP may contribute to the aggressive nature of SIX1/2miRNA tumors. Of particular interest as it relates to the increased rates of relapse of SIX1/2-Q177R tumors is *WNT5A* as it has been associated with increased resistance to chemotherapeutics in ovarian and breast cancer cells (Peng et al 2011, Hung et al 2014).

SIX1 and SIX1-Q177R enhance transcription from proximal and distal WNT5A CREs *in vitro* and SIX1-Q177R binds the *WNT5A* proximal CRE with higher affinity than SIX1

Again, leveraging the available SIX1 ChIP-seq data we identified shared peaks in both SIX1-expressing and SIX1-Q177R-expressing tumors, which did not appear in the wk17hFK dataset, within both proximal and distal regions near the *WNT5A* gene at sites of high DNA sequence conservation suggesting these regions may represent CREs (Figure 3A). Apart from *WNT5A*, proximal and/or distal SIX1-Q177R or shared tumor peaks were identified within candidate CREs as predicted by ENCODE or GeneHancer for other upregulated SIX1/2miRNA genes including *KDM2B, CDKN1C, TMEM100, ARHGEF3,* and *FOXC1* (Supplemental Figure 4A) (Abascal et al 2020, Fishilevich et al 2017).

To explore the potential regulation of gene expression by SIX1-Q177R through the *WNT5A* CREs, luciferase activity was measured from minimal promoter constructs containing each putative regulatory element in MCF-7 cells while overexpressing either SIX1 or SIX1-Q177R alongside the cofactor EYA1. As shown in Figure 3B, both wild type and mutant

FIGURE 3: **SIX1 and SIX1-Q177R bound highly-conserved putative proximal and distal cis-regulatory elements (CREs) for *WNT5A* in Wilms tumors and both proteins enhanced expression from each CRE in *in vitro* luciferase enhancer assays. A)** IGV genome browser snapshots displaying SIX1-Q177R tumor, SIX1 tumor, and wk17hFK SIX1 ChIP-seq tracks, along with UCSC genome browser basewise conservation track. Shaded regions indicate positions of shared SIX1 and SIX1-Q177R peaks in Wilms tumors. **B)** Bar graphs displaying results from the luciferase enhancer assays for each of the indicated DNA elements tested, * = $p < 0.05$.

SIX1 drove higher expression of luciferase from the proximal DNA element and both distal DNA elements. These data support the regulation of *WNT5A* expression by both wild type SIX1 and SIX1-Q177R. Nevertheless, overexpression of the proteins in these assays could mask subtle, yet biologically significant differences in their regulatory activities when expressed at normal physiological levels.

The SIX1-Q177R mutation is almost exclusively heterozygous in Wilms tumors and in the heterozygous context, both wild type SIX1 and SIX1-Q177R alleles were found to be expressed at similar levels (Walz et al 2015, Wegert et al 2015). Accordingly, differences in binding affinity at peaks shared between the two proteins could account for aberrant gene expression *in vivo*. To interrogate this possibility, we carried out electrophoretic mobility shift assays (EMSAs) using purified wild type SIX1 or SIX1-Q177R protein expressed in *E. Coli* and biotinylated oligonucleotide probes containing the putative DNA binding sequence found within the *WNT5A* promoter peak. Shown in Figure 4, SIX1-Q177R binds this sequence with an estimated 10-fold higher affinity compared to SIX1. Furthermore, changing the guanine to an adenine at the nucleotide position that appears to be preferred by SIX1-Q177R (Wegert et al 2015, Figure 1A) resulted in a loss of affinity of SIX1-Q177R with a concomitant, albeit modest increase in affinity of SIX1. Therefore, under conditions of equal expression of SIX1 and SIX1-Q177R alleles, SIX1-Q177R likely outcompetes its wild type counterpart for binding to this CRE, promoting aberrant expression of *WNT5A*.

FIGURE 4: SIX1-Q177R binds core DNA motif sequence from *WNT5A* proximal CRE with higher affinity than SIX1. Chemiluminescence images of blots from Electrophoretic Mobility Shift Assays (EMSA) using varying concentrations of purified SIX1 or SIX1-Q177R protein and biotin-labeled oligonucleotide probes with core DNA motif sequence indicated above. Below, quantification of EMSA binding derived from signal intensities of bound and unbound probe measured using ImageJ software.

Several upregulated genes in SIX1/2miRNA tumors are characteristic of the podocyte lineage in the developing human kidney

To relate our findings thus far to the normal human nephrogenic niche, wk14hFK and wk17hFK single cell RNA-seq datasets (GSE112570, GSE139280, GSE124472 (only sample GSM3534656)) were integrated using Seurat to generate a powerful reference with which to assess the expression and localization of the differentially expressed Wilms tumor genes (Supplemental Figure 5A) (Hao and Hao et al 2021, Stuart and Butler et al 2019, Butler et al 2018, Satija and Farrell et al 2015). These datasets were derived from specific cortical isolation procedures of hFK, ensuring enrichment of predominantly nephrogenic tissue (Lindstrom et al 2018d, Lindstrom et al 2021, Tran et al 2019). Consisting of more than 30,000 cells, UMAP dimensionality generated 20 clusters. Using the same sets of markers for cluster annotation as used in the data source publications, consistent clusters were identified in the integrated dataset (Supplemental Figure 5B, Supplemental File 2). In addition, unsupervised hierarchical clustering by Pearson's correlation coefficient using average

FIGURE 5: Several genes significantly upregulated in SIX1/2miRNA Wilms tumors show podocyte-enriched expression pattern in developing human kidney. A) Left, UMAP dimensional reduction plot showing clustering of cells from LindstromTranWk17 and LindstromWk14 single cell RNA-seq datasets after integrated analysis using Seurat. Right, UMAP dimensional reduction plot showing cell type classification of cell clusters. **B)** Violin plots displaying normalized expression levels of the significantly upregulated genes in SIX1/2miRNA tumors compared to Blastemal tumors in each cell cluster as shown in panel A.

z-scores for each gene across clusters identified cluster similarities which, in combination with the known cluster marker genes, was used to guide the grouping of individual clusters to generate the broad cluster identities shown in Figure 5A (Supplemental Figure 5C).

Probing this dataset for the SIX1/2miRNA upregulated genes shown in Figure 2B reveals abundant and podocyte precursor-enriched expression of *CDKN1C*, *ARHGEF3*, *FOXC1*, and *IQGAP2* (Figure 5B). Expression of *NOTCH2* also appears in podocyte precursors as well as PTA/RV, alongside *PAX8*, a canonical marker of NPC differentiation induction. Notably, *WNT5A* is also highly expressed in a small population of podocyte precursors (Supplemental Figure 6). Although *KDM2B* does not appear strongly in this dataset, *in situ* hybridization in hFK has illustrated high expression in RV/SSB structures (Lindstrom et al 2018d). Interrogation of the genes significantly downregulated in SIX1/2miRNA tumors compared to both Blastemal and MIXED/ES tumors reveal enrichment of genes with IC-specific expression patterns, likely indicative of smaller proportions of stromal/ICs in the SIX1/2miRNA tumors (Supplemental Figure 6). The expression pattern of the upregulated genes in differentiating structures of the developing human kidney is indicative of a farther progression along the podocyte differentiation trajectory in SIX1/2miRNA tumors than has been characterized previously.

Numerous genes exhibiting podocyte-enriched expression pattern in the developing kidney are expressed in all Wilms tumors analyzed

Due to our stringent filtering of weakly-expressed genes described earlier, we can infer with high confidence that the genes passing this filtering step are expressed at appreciable levels in all tumor samples. As such, we examined these genes more closely, regardless of significant differential expression, to gain additional insight into the correlation of the Wilms tumor transcriptome with that of the developing hFK. Removal of human housekeeping genes (Eisenberg and

FIGURE 6: **Numerous genes expressed in all Wilms tumor groups show podocyte-enriched expression pattern in developing human kidney, recapitulating both early and late signatures of podocyte lineage differentiation.** A) Heatmap displaying z-scores of average expression of all cells within each cluster for the 1,149 genes expressed in all Wilms tumor groups that were also identified as a marker in any of the single cell RNA-seq clusters. B) Bar chart showing the number of genes from panel A annotated in each of the shown Gene Ontology Biological Process terms identified using the Database for Annotation, Visualization and Integrated Discovery (DAVID) . C) Dot plots showing scaled average expression of all cells within the indicated cell clusters and the percent of cells in each cluster expressing the indicated genes derived from Tran et al early podocyte (top) and late podocyte (bottom) gene signatures. D) Density ridge plots showing the log2 CPM expression of the indicated genes from panel C in all Wilms tumor subgroups.

Levanon 2013) and intersection of the remaining genes with markers of all clusters in the integrated single cell RNA-seq dataset generated a list of 1,149 genes (Supplemental File 2). Analysis of average z-scores for each gene across all cell clusters shows the largest proportions of genes are expressed predominantly in endothelial cells, immune cells, and podocyte precursors (Figure 6A). Gene Ontology analysis using the Database for Annotation, Visualization, and Integrated Discovery (DAVID) revealed significant enrichment of biological processes related to angiogenesis, cell migration, and cell adhesion (Figure 6B) (Sherman et al 2022, Huang et al 2009). That large fractions of these genes are associated with endothelial and immune cells is unsurprising, as there is evidence for the presence of both endothelium and tumor-associated macrophages in Wilms tumors (Skoldenberg et al 2001, Ghanem et al 2003, Liou 2013, Vakkila 2016, Tian 2020). The prevalence of podocyte-precursor specific genes is consistent with the observation within the SIX1/2miRNA upregulated gene set and implies the acquisition of podocyte-like attributes may provide a selective advantage to Wilms tumor cells.

To characterize the podocyte relationship further, we examined the expression of genes identified by Tran et al as signatures of cells at early (EP) and late (LP) stages along the podocyte differentiation trajectory. The hFK expression patterns of those genes that are expressed in all Wilms tumors analyzed and also identified as EP and LP genes by Tran et al are shown in Figure 6C (Tran et al 2019). The expression of selected genes from that list in each tumor group are shown in Figure 6D, including canonical podocyte markers *MAFB*, *PODXL*, and *SYNPO*. These data illustrate that Wilms tumors of all histological subtypes recapitulate the continuum of podocyte specification at the transcriptional level.

DISCUSSION

Here we have integrated genomic datasets from pre-chemotherapy high-risk Wilms tumors and developing hFK to demonstrate the commonality of a podocyte-like gene expression signature in Wilms tumors of all histological subtypes and the augmentation of podocyte-specific gene expression in SIX1/2miRNA tumors. Wilms tumors have been characterized generally as arising from aberrant kidney development, yet until the availability of single cell RNA sequencing the field has lacked the necessary resolution to place Wilms tumors along the trajectory of normal human kidney development. Our results appear to contradict those of Gadd et al, in which the authors observed enrichment of pre-induction gene expression with accompanying low post-induction gene expression in most of the tumors included in their dataset using microarray expression data from a subset of the same FHWTs utilized for our study (Gadd et al 2017). However, that work preceded the use of single cell RNA sequencing in the context of human fetal kidney development beginning in 2018.

Subsequently, genes once characterized almost exclusively as pre-induction NPC genes are now known to be expressed in differentiating epithelialized structures of the RV including *SIX2, TMEM100,* and *MEOX1*, and even in the SSB including *SIX1* which is localized to the podocyte precursors of the proximal SSB (Lindstrom et al 2018b). This raises the

likelihood that SIX1 plays a meaningful functional role in the gene regulatory networks governing podocyte lineage specification. Thus, the persistent expression of these genes in Wilms tumors does not necessarily reflect a proliferative, naïve NPC-like state. In fact, a study using serial transplantation of Wilms tumor xenografts suggested cells resembling induced epithelium were the primary culprit of propagation in these tumors (Shukrun et al 2014).

Podocytes, highly specialized epithelial derivatives of NPCs, form extensive mesh-like networks with one another in which cellular protrusions termed "foot processes" interdigitate and surround the glomerular capillaries to filter the incoming blood. While it is well-understood that podocytes derive from the NPCs, the temporal relationship and transcriptional similarity between these cell populations was only recently uncovered. Lindstrom et al used time-lapse imaging of the developing mouse kidney to demonstrate that the last committed NPCs integrated into the RV contribute to the podocyte lineage. Pseudotime temporal analysis of single-cell RNA-seq data from nephrogenic hFK tissue reinforced those findings, indicating podocytes are the most closely related descendent of NPCs and arise from a distinct differentiation timeline to that of the tubule precursors, so much so that the authors concluded podocytes differentiate directly form committed NPCs (Lindstrom et al 2018d). Tran et al characterized this intimate relationship in detail using single-cell RNA-seq to distinguish mature podocytes located in the inner cortex of the developing kidney to the NPCs and early podocyte precursors located in the outer cortex of the kidney. This analysis generated a panel of marker genes distinguishing the EP lineage from the LP lineage. In agreement with Lindstrom et al's evidence, approximately half of the EP genes are expressed in NPCs including *SIX1, SIX2, CRABP2, MEOX1,* and *TMEM100* (Tran et al 2019).

Our analyses suggest this podocyte lineage trajectory is broadly conserved in Wilms tumors, with certain cytoskeletal and cell adhesion attributes potentially enhanced by the SIX1-Q177R mutation. Imbalance in WNT pathway signaling in favor of non-canonical WNT/PCP in the context of SIX1/2-Q177R would also favor podocyte lineage progression, as canonical WNT/β-catenin signaling has been demonstrated to inhibit podocyte differentiation in mouse and chick model systems, as well as in human induced pluripotent stem cell (hiPSC)-derived podocytes (Lindstrom et al 2015, Grinstein et al 2013, Yoshimura et al 2019). Supported by the findings of Lindstrom et al and Tran et al, acquisition of podocyte-like characteristics represents the differentiation path of least resistance. Moreover, the specialized traits of podocytes would likely be advantageous for tumor survival and propagation. Podocytes require a complex cytoskeletal architecture for foot-process formation and maintenance of slit-diaphragm integrity to ensure proper glomerular filtration (reviewed in Welsh and Saleem 2011). Developing podocytes also produce the angiogenic growth factor VEGF-A, as do Wilms tumors (Figure 6D), thought to attract endothelial progenitor cells to promote vascularization of the developing glomerulus and could feasibly enhance vascularization of Wilms tumors (Eremina et al 2003, Kitamoto and Tomita 1997).

Altogether, the evidence presented here supports a more podocyte-restricted developmental path of SIX1/2-Q177R-expressing cells in Wilms tumor, the characteristics of which may aid in resistance to chemotherapeutic intervention. However, whether this mutation is sufficient for oncogenic transformation in isolation remains unanswered and unlikely. The most common genomic anomaly in Wilms tumor is the loss of imprinting (LOI)/loss of heterozygosity (LOH) at chromosome 11p15 resulting in overexpression of *IGF2*, occurring in 75-80% of tumors (Gadd et al 2017, Wegert et al 2015). *IGF2* was the most highly expressed gene in our differential expression analysis (log2 CPM ~15). Therefore, synergism between SIX1/2-Q177R and *IGF2* overexpression or miRNA processing mutations may be required for tumorigenesis.

Interrogation of the mechanism through which the podocyte lineage might contribute to Wilms tumor and be further promoted by SIX1-Q177R will require an *in vitro* model system that recapitulates this developmental progression. Tran et al extended their findings to hiPSC-derived kidney organoid differentiations and illustrated podocyte-like cells in kidney organoids followed a developmental trajectory resembling that of *in vivo* hFK podocytes characterized by a high degree of conservation of most of the identified EP and LP genes. Additionally, xenotransplantation of these avascular organoids in immunocompromised mice resulted in vascularization of the organoids by the host, as has been demonstrated similarly by several groups (Sharmin et al 2016, Van den Berg et al 2018, Ryan et al 2021, Koning et al 2022). Hence, hiPSC-derived kidney organoids represent an appropriate model system for future mechanistic studies of podocyte lineage contribution to tumorigenesis.

Our study and the findings herein are not without limitations. Wilms tumors are notoriously heterogenous and histological classification is fundamentally subjective. However, tumors classified as mixed histology imply the presence of some identifiable blastemal components, supporting our observation of the overall high transcriptional correlation between tumors of different histological subtypes. Due to the limited availability of SIX1/2-Q177R tumor tissue, the nine samples included in this analysis undoubtedly fail to capture the full extent of variability among tumors with this mutation. Nonetheless, our use of stringent filtering, expression, and statistical thresholds increases the likelihood of biologically significant findings resulting from of our differential gene expression analysis. Conversely, these same stringent thresholds likely obscured some differentially expressed genes of biological significance. For example, NPC and podocyte marker *WT1*, as well as *CDH1* encoding E-cadherin, were excluded from this analysis due to several tumors not meeting the expression threshold. Additionally, the SIX1-Q177R ChIP-seq data utilized in this study was derived from a chemotherapy-treated tumor specimen (Wegert et al 2015). As such, this sample does not serve as a direct comparison to the SIX1-Q177R tumors used in the RNA-seq analysis. Therefore, inference of putative target gene regulation by SIX1-Q177R in chemotherapy-naïve tumors is only correlative at this time.

Despite these limitations, our data support a reassessment of the differentiation status of Wilms tumors in which progression through podocyte lineage specification is a common feature of all Wilms tumor subtypes, is augmented in high-risk blastemal-predominant-associated SIX1/2-Q177R-expressing tumors, and is advantageous for oncogenesis in the developing hFK.

EXPERIMENTAL PROCEDURES

Cell culture and transfections

HEK293T and MCF-7 (gift from Dr. Richard Cheney) cell lines were cultured in DMEM/F12 w/ L-glutamine and HEPES (Gibco) + 10% Fetal Bovine Serum (Omega Scientific) + 1X Penicillin/Streptomycin (Gibco), unless indicated elsewhere. Media was exchanged every 3-4 days and cells were passaged at confluence. Transfections for all assays were carried out using Lipofectamine 3000 kit (Invitrogen) following manufacturer's protocol, scaled for tissue culture vessel used and cells would be assayed ~48 hours post-transfection.

Cloning

All restriction digest reactions were carried out at 37C using CutSmart Buffer (New England Biolabs). Restriction digest products were subjected to gel electrophoresis in 1% agarose gel + 1X SYBR Safe DNA Gel Stain (Invitrogen). DNA bands of digested DNA were excised and purified using NucleoSpin Gel and PCR Clean-Up kit (Machery-Nagel) following manufacturer's protocol. All ligations were carried out at RT using T4 DNA ligase (New England Biolabs) following manufacturer's protocol. Unless indicated elsewhere, plasmids were transformed in 5-alpha Competent *E. coli* (New England Biolabs) cells following manufacturer's protocol. LB agar plates containing 1X ampicillin (Sigma Aldrich) were then streaked with transformed bacteria and incubated at 37C overnight. Individual bacteria colonies were picked using P1000 pipet tips and cultured in LB broth + 1X ampicillin at 37C on orbital shaker from several hours to overnight. Plasmids were purified from cultures using NucleoSpin Plasmid EasyPure plasmid purification kit (Machery-Nagel) following manufacturer's protocol. Plasmids were submitted to GENEWIZ (Azenta Life Sciences) for Sanger sequencing to verify DNA sequence. Purified plasmids verified by DNA sequencing were then either used directly in applicable assay or re-transformed in 5-alpha cells and an individual bacteria colony would be used to inoculate a 200mL liquid culture in LB broth + 1X ampicillin and culture would incubate overnight at 37C on orbital shaker. Plasmid would then be purified using NucleoBond Xtra Midi Plus EF Kit (Machery-Nagel), following manufacturer's protocol including use of NucleoBond finalizers.

SDS-PAGE and Western Blots

All samples were prepared using NuPAGE LDS Sample Buffer (Invitrogen) and incubated at 70C for 10 minutes prior to electrophoresis. Samples were run on Novex 4-20% Tris-Glycine gels (Invitrogen) along with Precision Plus Protein All Blue Prestained Protein Standards (Biorad) using 1X Tris-Glycine running buffer (25mM Tris Base (Fisher BioReagents), 190mM Glycine (Fisher Chemical), and 3.5 mM SDS (Fisher Chemical)) in a Mini Gel Tank (Invitrogen). Proteins were then transferred to a nitrocellulose membrane (GE Water and Process Technologies) using a wet transfer protocol and a Mini Trans-Blot Cell (Bio-Rad) transfer apparatus at 4C, 100V for 1 hour in transfer buffer (192 mM Glycine, 25mM Tris Base, and 20% methanol). Membranes were then washed several times with ddH20. Membranes were blocked in PBS + 0.1% TWEEN-20 (Fisher BioReagents) + 5% dry milk at RT for 20-40 minutes and residual milk was washed from membrane with water. Primary antibody solutions were made in PBS + 0.1% TWEEN-20 + 3% BSA (Fisher BioReagents).   Unless indicated elsewhere, membranes were incubated in primary antibody solution with gentle rocking at either RT for 1 hour or overnight at 4C. Membranes were washed at least three times with PBS + 0.1% TWEEN-20, 5 minutes per wash. Secondary antibody solutions were made in PBS + 0.1% TWEEN-20 + 1% dry milk. Membranes were incubated in secondary antibody solution for 1 hour at RT. Membranes were washed at least three times with PBS + 0.1% TWEEN-20 and then incubated in Pierce ECL Western Blotting Subtrate (Thermo Scientific) for 1-2 minutes at RT. Blots were imaged using iBright FL1500 Imaging System (Invitrogen).

ChIP-seq data analysis

SIX1 ChIP-seq data from Wilms tumors was kindly shared by Dr. Manfred Gessler (Wegert et al. 2015) in the form of bigWig files from two SIX1 wild type ChIP-seq samples and two SIX1-Q177R ChIP-seq samples (50 bp bins, normalized to pooled input by signal extraction scaling (SES)). BigWig files were converted to bedGraph format using UCSC tools bigWigToBedGraph (Kent et al 2010). Duplicates were merged and $\log_2$ fold enrichments over input were averaged between duplicates, generating a single bedGraph file for each set of duplicate samples. Coordinates were transformed from hg19 genome assembly to hg38 using the UCSC LiftOver tool (Kent et al 2002). Peaks were called using MACS2 bdgpeakcall (Zhang et al 2008) through the Galaxy web platform usegalaxy.org (Afgan et al 2016) using a $\log_2$ fold cutoff of +/- 2. Week 17 human fetal kidney SIX1 ChIPseq peak BED file was obtained from Dr. Lori O'Brien (O'Brien et al, 2016). Peak overlaps between samples were identified using bedtools (Quinlan and Hall 2010) intersect intervals in Galaxy. Peak sequences were obtained using bedtools GetFastaBed in Galaxy and these FASTA files were used as input for motif discovery using STREME (MEME Suite 5.4.1) (Bailey 2021) and the default settings, or the SEA tool in MEME Suite 5.4.1 to assess enrichment of motifs of potential SIX1 co-factor MEIS1 (Bailey and Grant 2021 bioRxiv). The identified SIX1 or SIX1-Q177R motifs from STREME were also used as input into FIMO (Grant 2011) using p-value cutoffs of either <1E-4 or <1E-5. The peak coordinates containing the desired motifs were converted to BED format and duplicate coordinates within each file were removed using bedtools MergeBED in Galaxy. These BED files were then used as input in the Genomic Regions Enrichment of Annotations Tool (GREAT) (McLean et al 2010) using whole genome background and basal plus extension association rules, changing distal up to 500kb. Putative target genes were identified using a binomial false discovery rate cutoff of 0.2.

Wilms tumor RNA-seq data analysis

Wilms tumor RNAseq data was obtained through the National Cancer Institute TARGET Data Matrix (https://ocg.cancer.gov/programs/target/data-matrix) in the form of gene quantification text files. From each gene quantification file, raw counts were transferred to an excel spreadsheet to create a count matrix. This count matrix was then used in Galaxy for differential gene expression analysis using limma-voom with sample quality weights, filtering out lowly expressed genes with CPM < 2 if threshold was not met in all samples, and using a log2 fold change cutoff of +/- 1.5, and all other default settings (Smyth 2005, Law et al 2014, Liu et al 2015).

Single cell RNA-seq analysis

Raw and processed data was obtained from three studies: GSE112570, GSE139280, GSE124472 (only sample GSM3534656). All subsequent data processing and analyses was performed using the Seurat package and following the vignettes "PBMC 3K guided tutorial" and "Introduction to scRNA-seq integration" (https://satijalab.org/seurat/index.html, Hao and Hao et al 2021, Stuart and Butler et al 2019, Butler et al 2018, Satija and Farrell et al 2015). Each dataset was filtered using the following parameters: LindstromWk14 – nFeature_RNA = 1500-4000, mitochondrial counts < 5%; LindstromWk17 – nFeature_RNA = 1000-3500, mitochondrial counts < 5%; TranWk17zone1 – nFeature_RNA = 1000-5000, mitochondrial counts < 5%. Each dataset was normalized independently and variable features were identified independently. Integration features were selected and integration anchors were identified. An integrated assay was then created, data was scaled, PCA and UMAP dimensional reduction were performed using n=20 principle components/dimensions. Neighbors were found and clusters were found using resolution = 0.5. Cluster markers were found using FindAllMarkers() function, min.pct = 0.15, logfc.threshold = 0.25. AverageExpression() function was used to calculate average expression value for each cluster, used return.seurat = TRUE to return SeuratObject with scaled and centered expression values generated from ScaleData() function. Dot plots were generated using DotPlot() function, heatmaps were generated using DoHeatmap() function, and violin plots were generated using VlnPlot() function.

Luciferase assays

- ### *Cloning and plasmids*

pBV-Luc was a gift from Bert Vogelstein (Addgene plasmid # 16539; http://n2t.net/addgene:16539; RRID:Addgene_16539). This plasmid encodes firefly luciferase driven by a minimal promoter element and was digested with NheI and HindIII, however this digestion removed the minimal promoter from the pBV-Luc vector. SIX1_enhancer gBlock (Supplemental Table 2) was PCR amplified and digested with NheI and HindII restriction enzymes and annealed with digested pBV-Luc. To re-insert the minimal promoter sequence, single-stranded DNA oligos containing the minimal promoter sequence flanked by 5'-HindIII and 3'-NcoI restriction sites were annealed in 1X annealing buffer (10mM Tris Base, 50mM NaCl (Fisher Chemical), 1mM EDTA (Invitrogen)) and incubated on thermocycler 95C for 2 minutes followed by cooling to 25C at a rate of -0.1C/second. Annealed minimal promoter oligo and SIX1_enhancer-pBV-Luc were then

digested with HindIII and NcoI restriction enzymes and annealed. All subsequent cloning of DNA elements was performed using NheI/HindIII sites from this plasmid.

pRL-SV40P was a gift from Ron Prywes (Addgene plasmid # 27163; http://n2t.net/addgene:27163; RRID:Addgene_27163) and was used as renilla luciferase expression control. Empty pCIG and empty pCS2+ plasmids were used as empty vectors for total DNA transfection normalization. pCS2+-EYA1-2xHA plasmid was generated by PCR amplifying EYA1 coding sequence from pCS2+-EYA1-FLAG plasmid and swapping out FLAG tag for 2xHA tag (Supplemental Table 2). pCIG-SIX1 and pCIG-SIX1-Q177R plasmids were generated using gBlocks synthesized by Integrated DNA Technologies (IDT) containing the coding sequences for the respective proteins (Supplemental Table 2). The gBlocks were PCR amplified and flanking ClaI and XhoI restriction sites were added and used for subsequent digestion and ligation into pCIG vector.

- *Luciferase assay*

MCF-7 cells were cultured in 6-well plates to ~90% confluence and media was exchanged prior to transfection. For each biological replicate of each DNA element assayed, three wells would be transfected with 5 µg total DNA/well: one no protein control condition, one SIX1/EYA1 condition, and one SIX1-Q177R/EYA1 condition. Control transfections consisted of 500 ng firefly luciferase vector, 10 ng renilla luciferase vector, 1.5 µg empty pCIG, and 3 µg empty pCS2+. Luciferase assays were performed using the Dual Luciferase Reporter Assay System (Promega). Each well was harvested using a cell scraper and 100 µL 1X Passive Lysis Buffer. After trituration with a P200 pipet tip, 20 µL lysate was transferred to each of 3x wells of a clear, flat-bottom 96-well plate for technical triplicate per condition. Luminescence was measured using a BioTek Synergy HT plate reader with the following settings: 10 second integration time, 135 gain, 1 mm read height. 100 µL/well Luciferase Assay Reagent II was dispensed into all wells containing cell lysate using P200 multichannel pipet and plate was immediately placed in plate reader. After firefly luciferase luminescence was measured, 100 µL/well Stop & Glo reagent was dispensed using P200 multichannel pipet and plate was immediately placed in plate reader. Fold change relative to no protein control was calculated by comparing ratios of firefly/renilla luminescence from SIX1 or SIX1-Q177R conditions to that of no protein control.

Protein expression and purification

Same SIX1 and SIX1-Q177R gBlocks used for pCIG cloning were PCR amplified to add flanking BamHI and XhoI restriction sites. Digested gBlocks were ligated to digested pGEX-6p1-N-HA (gift from Andrew Jackson & Martin Reijns, Addgene plasmid # 119756; http://n2t.net/addgene:119756; RRID:Addgene_119756). BL21 (DE3) Competent *E. Coli* (New England Biolabs) were transformed following manufacturer's protocol and streaked LB agar + 1X ampicillin plates were incubated overnight at 37C. Picked one colony from each transformation and cultured in 14 mL tubes each containing 7 mL LB broth + 1X ampicillin in orbital shaker at 37C overnight. Made glycerol stocks from each overnight liquid culture by mixing 50% glycerol solution and liquid culture at a 1:1 ratio, stored at -80C. Used pipet tip to transfer a small amount of each glycerol stock to flasks containing 50 mL LB broth + 1X ampicillin and incubated cultures on orbital shaker overnight at 37C. Transferred each 50 mL culture to 2 L flasks each containing 950 mL LB broth + 1X ampicillin and cultured at 37C on orbital shaker until OD600 = 0.55-0.56. Added 5 mL 100mM IPTG (Sigma-Aldrich) solution to each culture, incubated flasks on orbital shaker at 25C for 19 hours. For each culture, distributed entire culture volume into four 250 mL centrifuge bottles and centrifuged at 5,500 RPM for 20 minutes at 4C. Resuspended and pooled all pellets for each culture in 35 mL supernatant and transferred each final suspension to 50 mL tubes. Centrifuged tubes at 4,000 x g for 18 minutes at 4C. Discarded supernatants and stored pellets at -80C.

Thawed frozen bacteria pellets on ice, loosened each pellet in 25 mL lysis buffer (20 mM Tris-HCl, 150 mM NaCl, 1% Triton X100, 10 mM DTT, 1X Protease Inhibitor Cocktail (Roche)). Sonicated samples on ice in 4C cold room using Branson Sonifier 250, 10 cycles of 10 sec ON/OFF at 50% amplitude/duty cycle. Incubated tubes on ice for 15 minutes, centrifuged at 13,000 x g for 15 minutes at 4C. Decanted H20 from glutathione-agarose bead mixture (0.84 g. glutathione-agarose beads in 168 mL water, incubated at 4C overnight). Added 120 mL lysis buffer to beads, mixed, incubated at 4C for 20 minutes. Decanted most of lysis buffer from beads, transferred 8 mL bead slurry to each of 2x 50 mL tubes. Added bacterial supernatants to tubes containing beads, put tubes on tube rotator at 4C for 1.5 hours. Wet and washed EconoColumns (BioRad) with 1x column volume PBS + 1% Triton-X100, emptied. Added supernatant/bead mixtures to columns and opened column spigots full to allow gravity flow. As flow was near stopping, washed beads with full column volumes four times with cleavage buffer (50 mM Tris-HCl, 1 mM EDTA, 1 mM DTT, 1% Triton-X100 (Fisher BioReagents). Mixed 200 µL PreCission Protease (Cytiva) with 9.8 mL cleavage buffer, added 5 mL to each column, inverted to mix, incubated at 4C for 2 hours, inverted to mix, and incubated at 4C for 2 hours. Collected flow-through in 15

mL conical tubes. Cut 12-14,000 MWCO dialysis tubing (Spectrum Laboratories) and incubated at RT in 1.5L H2O + 5 mM EDTA for 2-3 hours, rinsed thoroughly with H2O.

Transferred entire protein volumes to dialysis tubing, clipped ends shut, incubated overnight at 4C submerged in 1 L dialysis buffer (50 mM Tris-HCl, 1 mM EDTA, 0.8 mM DTT) with gentle stirring. Discarded used dialysis buffer, replaced with fresh 1 L dialysis buffer and continued incubation at 4C for 2.5 hours. Transferred contents of each dialysis tubing to 15 mL conical tubes. Pre-washed membranes of 4x Amicon Ultra-4 30,000 MWCO Centrifugal Filter Units (Millipore) with water, removed. For each protein solution, transferred 2.5 mL to each of 2x filter units. Centrifuged tubes at 3,000 x g for 25 minutes at 4C. For each protein concentrate, pooled volumes from filter units. Measured protein concentration using Pierce BCA Assay Kit (Thermo Scentrific) and following manufacturer's protocol with the following changes: in 96-well plate loaded BSA controls and 10 µL 1:10 sample dilution with 190 µL working reagent in duplicate, incubated at 37C for 30 minutes. Measured absorbance at 562 nm on BioTek Synergy HT plate reader. Using BSA controls to generate standard curve, calculated protein concentration of each purified protein sample. Added dialysis buffer to each protein solution to bring concentrations to 2 mg/mL, made aliquots and stored at -80C. Protein purification was validated by SDS-PAGE of dilution series of each protein solution, followed by western blot using anti-SIX1 antibody (Cell Signaling Technology #12891) (Supplemental Figure 4B).

Electrophoretic Mobility Shift Assays

Single-stranded DNA oligonucleotide probes were synthesized by IDT and biotin end-labeled using Pierce 3' Biotin end-labeling DNA kit (Thermo Scientific) following manufacturer's protocol with the following changes: labeled 25 pmol oligo per reaction, stopped reactions with 1 µL 0.5M EDTA, mixed complementary oligo labeling reactions prior to centrifugation at 13,000 x g for 2 minutes. For unlabeled oligos, mixed 50 µL H2O with 25 µL of each complementary 100 µM oligo. Added annealing buffer to 1X and annealed following same procedure as described in luciferase assay, stored annealed oligos at -20C.

Purified SIX1 and SIX1-Q177R protein was diluted in H20 to the following molarities (nM): 1, 2, 4, 10, 20, 40, 100, 150, 200, and 300. EMSAs were performed using either Gelshift Chemiluminescent EMSA Kit (Active Motif) or LightShift Chemiluminescent EMSA Kit (Thermo Scientific) following manufacturer's protocol for setting up binding reactions, with the following changes: did not include glycerol or poly d(I-C) in reactions, used 3 µL/reaction of 1:10 diluted biotin end-labeled probe, used 12 µL/reaction H2O (13µL for no protein control reaction), 1µL appropriate protein dilution per reaction incubated reactions at RT for 25 minutes. Pre-ran 6% DNA Retardation Gel (Invitrogen) at 100V for 30 minutes in 0.5X TBE buffer (45 mM Tris Base, 45 mM boric acid, 1 mM EDTA) in a Mini Gel Tank. Added 5 µL/reaction 5X Loading Dye, loaded 20 µL/reaction in gel and ran gel at 100V for one hour. Transferred using a wet transfer protocol similar to that used for western blot described previously but using 0.5X TBE buffer as transfer buffer and transferring to Immobilin Ny+ nylon membrane (Millipore). Put membrane on paper towel to dry briefly. Placed membrane face-down in BioDoc-IT gel imager (UVP), turned UV lamp on and incubated for 15 minutes. Crosslinked membrane was then either stored at -20C, or proceeded directly to staining following kit manufacturer's protocol. Stained membranes were imaged using iBright FL1500 Imaging System.

Co-immunoprecipitation and Western Blot

pCS2-MEIS1-3xFLAG construct was generated by PCR amplification of MEIS1 from human induced pluripotent stem cell-derived cDNA (Supplemental Table 2). HEK293 cells were grown in 6cm dishes and transfected as described above, but using 4 ug total DNA per dish, 2µg/plasmid (Supplemental Table 2). One dish transfected with only pCS2-MEIS1-3xFLAG, one dish with pCS2-MEIS1-3xFLAG and pCIG-SIX1, one dish with pCS2-MEIS1-3xFLAG and pCIG-SIX1-Q177R, and one dish with only pCIG-SIX1. Nuclear protein complexes were extracted ~48 hours post-transfection using Active Motif Nuclear Complex Co-IP kit following manufacturer's protocol for preparation of nuclear extract but doubled digestion buffer volume used for each sample and incubated in digestion buffer at 37C for 25 minutes. Saved 1uL/sample as input and used 75uL/sample for immunoprecipitations. For each sample, added 25 µL Protein G Dynabeads (Invitrogen) to new microcentrifuge tube, washed 3x with 1 mL/wash PBS + 0.5% BSA. Resuspended beads in 250 µL PBS + 0.5% BSA, added 5 µL anti-SIX1 antibody (Cell Signaling Technology, SIX1 (D4A8K) Rabbit mAb #12891), incubated with gentle rotation at 4C for at least one hour. Washed beads 5x with 500uL/wash PBS + 0.25% Triton X-100. Removed final wash, added 30uL/tube LDS sample buffer (Invitrogen) + 2.5% 2-Mercaptoethanol (Sigma) to each immunoprecipitation tube and input tubes, incubated at 70C for 10 minutes. Ran samples on 4-20% Tris-Glycine Novex gel and transferred to nitrocellulose membrane same as detailed above, but transferred at 30V at 4C overnight.

Washed membrane and stained with primary and secondary antibodies following procedure described above, but using the following antibodies: ms anti-FLAG (Sigma #F3165, 1:500) and donkey anti-mouse HRP (Invitrogen #A16017, 1:10,000).

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Manfred Gessler for generously sharing the Wilms tumor ChIP-seq data, as well as the Therapeutically Applicable Research to Generate Effective Treatments (https://ocg.cancer.gov/programs/target) initiative (TARGET) study phs000218.v24.p8 for generating the Wilms tumor RNA-seq data used in this analysis (data available at https://portal.gdc.cancer.gov/projects).

FUNDING

Work in this study was supported by the Maren Endowment and UNC Lineberger Tier 1 Pilot/Development Award (L.L.O.), as well as NCI F31CA257443 and NIGMS Training Grant 5T32 GM007092 (M.J.S).

REFERENCES

Abascal, F., Acosta, R., Addleman, N. J., Adrian, J., Afzal, V., Ai, R., Aken, B., Akiyama, J. A., Jammal, O. A., Amrhein, H., Anderson, S. M., Andrews, G. R., Antoshechkin, I., Ardlie, K. G., Armstrong, J., Astley, M., Banerjee, B., Barkal, A. A., Barnes, I. H. A., . . . Weng, Z. (2020). Expanded encyclopaedias of DNA elements in the human and mouse genomes. Nature, 583(7818), 699–710. https://doi.org/10.1038/s41586-020-2493-4

Beckwith, J. B. (1993). Precursor lesions of Wilms tumor: Clinical and biological implications. *Medical and Pediatric Oncology*, *21*(3), 158–168. https://doi.org/10.1002/mpo.2950210303

Beckwith, J. B., Kiviat, N. B., & Bonadio, J. F. (1990). Nephrogenic Rests, Nephroblastomatosis, and the Pathogenesis of Wilms' Tumor. *Pediatric Pathology*, *10*(1–2), 1–36. https://doi.org/10.3109/15513819009067094

Breslow, N. E., Beckwith, J. B., Perlman, E. J., & Reeve, A. E. (2006). Age distributions, birth weights, nephrogenic rests, and heterogeneity in the pathogenesis of Wilms tumor. *Pediatric Blood & Cancer*, *47*(3), 260–267. https://doi.org/10.1002/pbc.20891

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., & Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nature Biotechnology, 36(5), 411–420. https://doi.org/10.1038/nbt.4096

Coorens, T. H. H., Treger, T. D., Al-Saadi, R., Moore, L., Tran, M. G. B., Mitchell, T. J., Tugnait, S., Thevanesan, C., Young, M. D., Oliver, T. R. W., Oostveen, M., Collord, G., Tarpey, P. S., Cagan, A., Hooks, Y., Brougham, M., Reynolds, B. C., Barone, G., Anderson, J., . . . Behjati, S. (2019). Embryonal precursors of Wilms tumor. Science, 366(6470), 1247–1251. https://doi.org/10.1126/science.aax1323

D'Amato, L., Dell'Aversana, C., Conte, M., Ciotta, A., Scisciola, L., Carissimo, A., Nebbioso, A., & Altucci, L. (2015). ARHGEF3 controls HDACi-induced differentiation via RhoA-dependent pathways in acute myeloid leukemias. *Epigenetics*, *10*(1), 6–18. https://doi.org/10.4161/15592294.2014.988035

DENG, Z., WANG, L., HOU, H., ZHOU, J., & LI, X. (2015). Epigenetic regulation of IQGAP2 promotes ovarian cancer progression via activating Wnt/β-catenin signaling. *International Journal of Oncology*, *48*(1), 153–160. https://doi.org/10.3892/ijo.2015.3228

Eisenberg, E., & Levanon, E. Y. (2013). Human housekeeping genes, revisited. Trends in Genetics, 29(10), 569–574. https://doi.org/10.1016/j.tig.2013.05.010

Eremina, V., Sood, M., Haigh, J., Nagy, A., Lajoie, G., Ferrara, N., Gerber, H. P., Kikkawa, Y., Miner, J. H., & Quaggin, S. E. (2003). Glomerular-specific alterations of VEGF-A expression lead to distinct congenital and acquired renal diseases. Journal of Clinical Investigation, 111(5), 707–716. https://doi.org/10.1172/jci17423

Farrell, J. A., Gennert, D., Schier, A. F., & Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. Nature Biotechnology, 33(5), 495–502. https://doi.org/10.1038/nbt.3192

Fernandez, C. V., Mullen, E. A., Chi, Y. Y., Ehrlich, P. F., Perlman, E. J., Kalapurakal, J. A., Khanna, G., Paulino, A. C., Hamilton, T. E., Gow, K. W., Tochner, Z., Hoffer, F. A., Withycombe, J. S., Shamberger, R. C., Kim, Y., Geller, J. I., Anderson, J. R., Grundy, P. E., & Dome, J. S. (2018). Outcome and Prognostic Factors in Stage III Favorable-Histology Wilms Tumor: A Report From the Children's Oncology Group Study AREN0532. *Journal of Clinical Oncology*, *36*(3), 254–261. https://doi.org/10.1200/jco.2017.73.7999

Fishilevich, S., Nudel, R., Rappaport, N., Hadar, R., Plaschkes, I., Iny Stein, T., Rosen, N., Kohn, A., Twik, M., Safran, M., Lancet, D., & Cohen, D. (2017). GeneHancer: genome-wide integration of enhancers and target genes in GeneCards. Database, 2017. https://doi.org/10.1093/database/bax028

Fukata, M., Watanabe, T., Noritake, J., Nakagawa, M., Yamaga, M., Kuroda, S., Matsuura, Y., Iwamatsu, A., Perez, F., & Kaibuchi, K. (2002). Rac1 and Cdc42 Capture Microtubules through IQGAP1 and CLIP-170. *Cell*, *109*(7), 873–885. https://doi.org/10.1016/s0092-8674(02)00800-0

Fukuzawa, R., Anaka, M. R., Morison, I. M., & Reeve, A. E. (2017). The developmental programme for genesis of the entire kidney is recapitulated in Wilms tumour. *PLOS ONE*, *12*(10), e0186333. https://doi.org/10.1371/journal.pone.0186333

Gadd, S., Huff, V., Huang, C. C., Ruteshouser, E. C., Dome, J. S., Grundy, P. E., Breslow, N., Jennings, L., Green, D. M., Beckwith, J. B., & Perlman, E. J. (2012). Clinically Relevant Subsets Identified by Gene Expression Patterns Support a Revised Ontogenic Model of Wilms Tumor: A Children's Oncology Group Study. *Neoplasia*, *14*(8), 742-IN21. https://doi.org/10.1593/neo.12714

Gadd, S., Huff, V., Walz, A. L., Ooms, A. H. A. G., Armstrong, A. E., Gerhard, D. S., Smith, M. A., Auvil, J. M. G., Meerzaman, D., Chen, Q. R., Hsu, C. H., Yan, C., Nguyen, C., Hu, Y., Hermida, L. C., Davidsen, T., Gesuwan, P., Ma, Y., Zong, Z., . . . Perlman, E. J. (2017). A Children's Oncology Group and TARGET initiative exploring the genetic landscape of Wilms tumor. *Nature Genetics*, *49*(10), 1487–1494. https://doi.org/10.1038/ng.3940

Gadd, S., Huff, V., Skol, A. D., Renfro, L. A., Fernandez, C. V., Mullen, E. A., Jones, C. D., Hoadley, K. A., Yap, K. L., Ramirez, N. C., Aris, S., Phung, Q. H., & Perlman, E. J. (2022). Genetic changes associated with relapse in favorable histology Wilms tumor: A Children's Oncology Group AREN03B2 study. Cell Reports Medicine, 3(6), 100644. https://doi.org/10.1016/j.xcrm.2022.100644

Ghanem, M. A. (2003). Expression and prognostic relevance of vascular endothelial growth factor (VEGF) and its receptor (FLT-1) in nephroblastoma. Journal of Clinical Pathology, 56(2), 107–113. https://doi.org/10.1136/jcp.56.2.107

Grinstein, M., Yelin, R., Herzlinger, D., & Schultheiss, T. M. (2013). Generation of the podocyte and tubular components of an amniote kidney: timing of specification and a role for Wnt signaling. Development, 140(22), 4565–4573. https://doi.org/10.1242/dev.097063

Gröbner, S. N., Worst, B. C., Weischenfeldt, J., Buchhalter, I., Kleinheinz, K., Rudneva, V. A., Johann, P. D., Balasubramanian, G. P., Segura-Wang, M., Brabetz, S., Bender, S., Hutter, B., Sturm, D., Pfaff, E., Hübschmann, D., Zipprich, G., Heinold, M., Eils, J., Lawerenz, C., . . . Pfister, S. M. (2018). The landscape of genomic alterations across childhood cancers. *Nature*, *555*(7696), 321–327. https://doi.org/10.1038/nature25480

Han, B., Zhou, B., Qu, Y., Gao, B., Xu, Y., Chung, S., Tanaka, H., Yang, W., Giuliano, A. E., & Cui, X. (2017). FOXC1-induced non-canonical WNT5A-MMP7 signaling regulates invasiveness in triple-negative breast cancer. Oncogene, 37(10), 1399–1408. https://doi.org/10.1038/s41388-017-0021-2

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J., Wilk, A. J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J., Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., . . .

Satija, R. (2021). Integrated analysis of multimodal single-cell data. Cell, 184(13), 3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048

Hikasa, H., Shibata, M., Hiratani, I., & Taira, M. (2002). The Xenopus receptor tyrosine kinase Xror2 modulates morphogenetic movements of the axial mesoderm and neuroectoderm via Wnt signaling. *Development*, *129*(22), 5227–5239. https://doi.org/10.1242/dev.129.22.5227

Hochane, M., van den Berg, P. R., Fan, X., Bérenger-Currias, N., Adegeest, E., Bialecka, M., Nieveen, M., Menschaart, M., Chuva de Sousa Lopes, S. M., & Semrau, S. (2019). Single-cell transcriptomics reveals gene expression dynamics of human fetal kidney development. PLOS Biology, 17(2), e3000152. https://doi.org/10.1371/journal.pbio.3000152

Hohenstein, P., Pritchard-Jones, K., & Charlton, J. (2015). The yin and yang of kidney development and Wilms' tumors. Genes &Amp; Development, 29(5), 467–482. https://doi.org/10.1101/gad.256396.114

Hol, J. A., Lopez-Yurda, M. I., van Tinteren, H., van Grotel, M., Godzinski, J., Vujanic, G., Oldenburger, F., de Camargo, B., Ramírez-Villar, G. L., Bergeron, C., Pritchard-Jones, K., Graf, N., & van den Heuvel-Eibrink, M. M. (2019). Prognostic significance of age in 5631 patients with Wilms tumour prospectively registered in International Society of Paediatric Oncology (SIOP) 93–01 and 2001. *PLOS ONE*, *14*(8), e0221373. https://doi.org/10.1371/journal.pone.0221373

Howlader N, Noone AM, Krapcho M, Miller D, Brest A, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). SEER Cancer Statistics Review, 1975-2017, National Cancer Institute. Bethesda, MD, https://seer.cancer.gov/csr/1975_2017/, based on November 2019 SEER data submission, posted to the SEER web site, April 2020.

Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2008). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nature Protocols, 4(1), 44–57. https://doi.org/10.1038/nprot.2008.211

Hung, T. H., Hsu, S. C., Cheng, C. Y., Choo, K. B., Tseng, C. P., Chen, T. C., Lan, Y. W., Huang, T. T., Lai, H. C., Chen, C. M., & Chong, K. Y. (2014). Wnt5A regulates ABCB1 expression in multidrug-resistant cancer cells through activation of the non-canonical PKA/β-catenin pathway. Oncotarget, 5(23), 12273–12290. https://doi.org/10.18632/oncotarget.2631

Kandoth, C., McLellan, M. D., Vandin, F., Ye, K., Niu, B., Lu, C., Xie, M., Zhang, Q., McMichael, J. F., Wyczalkowski, M. A., Leiserson, M. D. M., Miller, C. A., Welch, J. S., Walter, M. J., Wendl, M. C., Ley, T. J., Wilson, R. K., Raphael, B. J., & Ding, L. (2013). Mutational landscape and significance across 12 major cancer types. *Nature*, *502*(7471), 333–339. https://doi.org/10.1038/nature12634

Kawasaki, A., Torii, K., Yamashita, Y., Nishizawa, K., Kanekura, K., Katada, M., Ito, M., Nishimoto, I., Terashita, K., Aiso, S., & Matsuoka, M. (2007). Wnt5a promotes adhesion of human dermal fibroblasts by triggering a phosphatidylinositol-3 kinase/Akt signal. *Cellular Signalling*, *19*(12), 2498–2506. https://doi.org/10.1016/j.cellsig.2007.07.023

Kim, H., Choi, Y. J., Lee, Y. S., Park, S. Y., Baek, J. E., Kim, H. K., Kim, B. J., Lee, S. H., & Koh, J. M. (2018). SLIT3 regulates endochondral ossification by β-catenin suppression in chondrocytes. *Biochemical and Biophysical Research Communications*, *506*(4), 847–853. https://doi.org/10.1016/j.bbrc.2018.10.167

Kitamoto, Y., Tokunaga, H., & Tomita, K. (1997). Vascular endothelial growth factor is an essential molecule for mouse kidney development: glomerulogenesis and nephrogenesis. Journal of Clinical Investigation, 99(10), 2351–2357. https://doi.org/10.1172/jci119416

Koning, M., Dumas, S. J., Avramut, M. C., Koning, R. I., Meta, E., Lievers, E., Wiersma, L. E., Borri, M., Liang, X., Xie, L., Liu, P., Chen, F., Lin, L., Luo, Y., Mulder, J., Spijker, H. S., Jaffredo, T., van den Berg, B. M., Carmeliet, P., . . . Rabelink, T. J. (2022). Vasculogenesis in kidney organoids upon transplantation. Npj Regenerative Medicine, 7(1). https://doi.org/10.1038/s41536-022-00237-4

Lađinović, D., Pinkas, D., ŠOpin, T., Raška, O., Liška, F., Raška, I., & Vacík, T. (2020). Alternative isoforms of KDM2A and KDM2B lysine demethylases negatively regulate canonical Wnt signaling. *PLOS ONE*, *15*(10), e0236612. https://doi.org/10.1371/journal.pone.0236612

Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology*, *15*(2), R29. https://doi.org/10.1186/gb-2014-15-2-r29

Li, C. M., Guo, M., Borczuk, A., Powell, C. A., Wei, M., Thaker, H. M., Friedman, R., Klein, U., & Tycko, B. (2002). Gene Expression in Wilms' Tumor Mimics the Earliest Committed Stage in the Metanephric Mesenchymal-Epithelial Transition. *The American Journal of Pathology*, *160*(6), 2181–2190. https://doi.org/10.1016/s0002-9440(10)61166-2

Li, H., Hohenstein, P., & Kuure, S. (2021). Embryonic Kidney Development, Stem Cells and the Origin of Wilms Tumor. Genes, 12(2), 318. https://doi.org/10.3390/genes12020318

Li, X., Ohgi, K. A., Zhang, J., Krones, A., Bush, K. T., Glass, C. K., Nigam, S. K., Aggarwal, A. K., Maas, R., Rose, D. W., & Rosenfeld, M. G. (2003). Eya protein phosphatase activity regulates Six1–Dach–Eya transcriptional effects in mammalian organogenesis. *Nature*, *426*(6964), 247–254. https://doi.org/10.1038/nature02083

Lindström, N. O., Guo, J., Kim, A. D., Tran, T., Guo, Q., De Sena Brandine, G., Ransick, A., Parvez, R. K., Thornton, M. E., Basking, L., Grubbs, B., McMahon, J. A., Smith, A. D., & McMahon, A. P. (2018). Conserved and Divergent Features of Mesenchymal Progenitor Cell Types within the Cortical Nephrogenic Niche of the Human and Mouse Kidney. Journal of the American Society of Nephrology, 29(3), 806–824. https://doi.org/10.1681/asn.2017080890

Lindström, N. O., McMahon, J. A., Guo, J., Tran, T., Guo, Q., Rutledge, E., Parvez, R. K., Saribekyan, G., Schuler, R. E., Liao, C., Kim, A. D., Abdelhalim, A., Ruffins, S. W., Thornton, M. E., Basking, L., Grubbs, B., Kesselman, C., & McMahon, A. P. (2018). Conserved and Divergent Features of Human and Mouse Kidney Organogenesis. Journal of the American Society of Nephrology, 29(3), 785–805. https://doi.org/10.1681/asn.2017080887

Lindström, N. O., Tran, T., Guo, J., Rutledge, E., Parvez, R. K., Thornton, M. E., Grubbs, B., McMahon, J. A., & McMahon, A. P. (2018). Conserved and Divergent Molecular and Anatomic Features of Human and Mouse Nephron Patterning. Journal of the American Society of Nephrology, 29(3), 825–840. https://doi.org/10.1681/asn.2017091036

Lindström, N. O., de Sena Brandine, G., Tran, T., Ransick, A., Suh, G., Guo, J., Kim, A. D., Parvez, R. K., Ruffins, S. W., Rutledge, E. A., Thornton, M. E., Grubbs, B., McMahon, J. A., Smith, A. D., & McMahon, A. P. (2018). Progressive Recruitment of Mesenchymal Progenitors Reveals a Time-Dependent Process of Cell Fate Acquisition in Mouse and Human Nephrogenesis. *Developmental Cell*, *45*(5), 651–660.e4. https://doi.org/10.1016/j.devcel.2018.05.010

Lindström, N. O., Sealfon, R., Chen, X., Parvez, R. K., Ransick, A., de Sena Brandine, G., Guo, J., Hill, B., Tran, T., Kim, A. D., Zhou, J., Tadych, A., Watters, A., Wong, A., Lovero, E., Grubbs, B. H., Thornton, M. E., McMahon, J. A., Smith, A. D., . . . McMahon, A. P. (2021). Spatial transcriptional mapping of the human nephrogenic program. *Developmental Cell*, *56*(16), 2381–2398.e6. https://doi.org/10.1016/j.devcel.2021.07.017

Lindström, N. O., Lawrence, M. L., Burn, S. F., Johansson, J. A., Bakker, E. R., Ridgway, R. A., Chang, C. H., Karolak, M. J., Oxburgh, L., Headon, D. J., Sansom, O. J., Smits, R., Davies, J. A., & Hohenstein, P. (2015). Integrated β-catenin, BMP, PTEN, and Notch signalling patterns the nephron. ELife, 4. https://doi.org/10.7554/elife.04000

Liou, P., Bader, L., Wang, A., Yamashiro, D., & Kandel, J. J. (2013). Correlation of tumor-associated macrophages and clinicopathological factors in Wilms tumor. Vascular Cell, 5(1), 5. https://doi.org/10.1186/2045-824x-5-5 Menon, R., Otto, E. A., Kokoruda, A., Zhou, J., Zhang, Z., Yoon, E., Chen, Y. C., Troyanskaya, O., Spence, J. R., Kretzler, M., & Cebrián, C. (2018). Single-cell analysis of progenitor cell dynamics and lineage specification in the human fetal kidney. Development, 145(16). https://doi.org/10.1242/dev.164038

Liu, R., Holik, A. Z., Su, S., Jansz, N., Chen, K., Leong, H. S., Blewitt, M. E., Asselin-Labat, M. L., Smyth, G. K., & Ritchie, M. E. (2015). Why weight? Modelling sample and observational level variability improves power in RNA-seq analyses. *Nucleic Acids Research*, *43*(15), e97. https://doi.org/10.1093/nar/gkv412

Liu, Y., Shi, J., Lu, C. C., Wang, Z. B., Lyuksyutova, A. I., Song, X. J., & Zou, Y. (2005). Ryk-mediated Wnt repulsion regulates posterior-directed growth of corticospinal tract. *Nature Neuroscience*, *8*(9), 1151–1159. https://doi.org/10.1038/nn1520

Logue, J. S., Whiting, J. L., Tunquist, B., Langeberg, L. K., & Scott, J. D. (2011). Anchored Protein Kinase A Recruitment of Active Rac GTPase. *Journal of Biological Chemistry*, *286*(25), 22113–22121. https://doi.org/10.1074/jbc.m111.232660

Lu, L., Gao, Y., Zhang, Z., Cao, Q., Zhang, X., Zou, J., & Cao, Y. (2015). Kdm2a/b Lysine Demethylases Regulate Canonical Wnt Signaling by Modulating the Stability of Nuclear β-Catenin. *Developmental Cell*, *33*(6), 660–674. https://doi.org/10.1016/j.devcel.2015.04.006

McLean, C. Y., Bristor, D., Hiller, M., Clarke, S. L., Schaar, B. T., Lowe, C. B., Wenger, A. M., & Bejerano, G. (2010). GREAT improves functional interpretation of cis-regulatory regions. *Nature Biotechnology*, *28*(5), 495–501. https://doi.org/10.1038/nbt.1630

Mikels, A. J., & Nusse, R. (2006). Purified Wnt5a Protein Activates or Inhibits β-Catenin–TCF Signaling Depending on Receptor Context. *PLoS Biology*, *4*(4), e115. https://doi.org/10.1371/journal.pbio.0040115

Mikels, A., Minami, Y., & Nusse, R. (2009). Ror2 Receptor Requires Tyrosine Kinase Activity to Mediate Wnt5A Signaling. *Journal of Biological Chemistry*, *284*(44), 30167–30176. https://doi.org/10.1074/jbc.m109.041715

Moon, R., Campbell, R., Christian, J., McGrew, L., Shih, J., & Fraser, S. (1993). Xwnt-5A: a maternal Wnt that affects morphogenetic movements after overexpression in embryos of Xenopus laevis. *Development*, *119*(1), 97–111. https://doi.org/10.1242/dev.119.1.97

Mullen, E. A., Chi, Y. Y., Hibbitts, E., Anderson, J. R., Steacy, K. J., Geller, J. I., Green, D. M., Khanna, G., Malogolowkin, M. H., Grundy, P. E., Fernandez, C. V., & Dome, J. S. (2018). Impact of Surveillance Imaging Modality on Survival After Recurrence in Patients With Favorable-Histology Wilms Tumor: A Report From the Children's Oncology Group. *Journal of Clinical Oncology*, *36*(34), 3396–3403. https://doi.org/10.1200/jco.18.00076

Ng, L., Chow, A. K. M., Man, J. H. W., Yau, T. C. C., Wan, T. M. H., Iyer, D. N., Kwan, V. H. T., Poon, R. T. P., Pang, R. W. C., & Law, W. L. (2018). Suppression of Slit3 induces tumor proliferation and chemoresistance in hepatocellular carcinoma through activation of GSK3β/β-catenin pathway. *BMC Cancer*, *18*(1). https://doi.org/10.1186/s12885-018-4326-5

Nishita, M., Itsukushima, S., Nomachi, A., Endo, M., Wang, Z., Inaba, D., Qiao, S., Takada, S., Kikuchi, A., & Minami, Y. (2010). Ror2/Frizzled Complex Mediates Wnt5a-Induced AP-1 Activation by Regulating Dishevelled Polymerization. *Molecular and Cellular Biology*, *30*(14), 3610–3619. https://doi.org/10.1128/mcb.00177-10

O'Brien, L. L., Guo, Q., Lee, Y., Tran, T., Benazet, J. D., Whitney, P. H., Valouev, A., & McMahon, A. P. (2016). Differential regulation of mouse and human nephron progenitors by the Six family of transcriptional regulators. *Development*, *143*(4), 595–608. https://doi.org/10.1242/dev.127175

Oishi, I., Suzuki, H., Onishi, N., Takada, R., Kani, S., Ohkawara, B., Koshida, I., Suzuki, K., Yamada, G., Schwabe, G. C., Mundlos, S., Shibuya, H., Takada, S., & Minami, Y. (2003). The receptor tyrosine kinase Ror2 is involved in non-canonical Wnt5a/JNK signalling pathway. *Genes to Cells*, *8*(7), 645–654. https://doi.org/10.1046/j.1365-2443.2003.00662.x

Ozdemir, E. S., Jang, H., Gursoy, A., Keskin, O., Li, Z., Sacks, D. B., & Nussinov, R. (2018). Unraveling the molecular mechanism of interactions of the Rho GTPases Cdc42 and Rac1 with the scaffolding protein IQGAP2. *Journal of Biological Chemistry*, *293*(10), 3685–3699. https://doi.org/10.1074/jbc.ra117.001596

Peng, C., Zhang, X., Yu, H., Wu, D., & Zheng, J. (2011). Wnt5a as a Predictor in Poor Clinical Outcome of Patients and a Mediator in Chemoresistance of Ovarian Cancer. International Journal of Gynecologic Cancer, 21(2), 280–288. https://doi.org/10.1097/igc.0b013e31820aaadb

Rivera, M. N., & Haber, D. A. (2005). Wilms' tumour: connecting tumorigenesis and organ development in the kidney. *Nature Reviews Cancer*, *5*(9), 699–712. https://doi.org/10.1038/nrc1696

Ryan, A. R., England, A. R., Chaney, C. P., Cowdin, M. A., Hiltabidle, M., Daniel, E., Gupta, A. K., Oxburgh, L., Carroll, T. J., & Cleaver, O. (2021). Vascular deficiencies in renal organoids and ex vivo kidney organogenesis. Developmental Biology, 477, 98–116. https://doi.org/10.1016/j.ydbio.2021.04.009

Schambony, A., & Wedlich, D. (2007). Wnt-5A/Ror2 Regulate Expression of XPAPC through an Alternative Noncanonical Signaling Pathway. *Developmental Cell*, *12*(5), 779–792. https://doi.org/10.1016/j.devcel.2007.02.016

Sharmin, S., Taguchi, A., Kaku, Y., Yoshimura, Y., Ohmori, T., Sakuma, T., Mukoyama, M., Yamamoto, T., Kurihara, H., & Nishinakamura, R. (2015). Human Induced Pluripotent Stem Cell–Derived Podocytes Mature into Vascularized Glomeruli upon Experimental Transplantation. Journal of the American Society of Nephrology, 27(6), 1778–1791. https://doi.org/10.1681/asn.2015010096

Sherman, B. T., Hao, M., Qiu, J., Jiao, X., Baseler, M. W., Lane, H. C., Imamichi, T., & Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic Acids Research, 50(W1), W216–W221. https://doi.org/10.1093/nar/gkac194

Shukrun, R., Pode-Shakked, N., Pleniceanu, O., Omer, D., Vax, E., Peer, E., Pri-Chen, S., Jacob, J., Hu, Q., Harari-Steinberg, O., Huff, V., & Dekel, B. (2014). Wilms' Tumor Blastemal Stem Cells Dedifferentiate to Propagate the Tumor Bulk. Stem Cell Reports, 3(1), 24–33. https://doi.org/10.1016/j.stemcr.2014.05.013

SKÖLDENBERG, E. G., CHRISTIANSSON, J., SANDSTEDT, B., LARSSON, A., LÄCKGREN, G., & CHRISTOFFERSON, R. (2001). ANGIOGENESIS AND ANGIOGENIC GROWTH FACTORS IN WILMS TUMOR. Journal of Urology, 165(6 Part 2), 2274–2279. https://doi.org/10.1016/s0022-5347(05)66183-6

Smyth, G. K. (2005). limma: Linear Models for Microarray Data. *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*, 397–420. https://doi.org/10.1007/0-387-29362-0_23

Steliarova-Foucher, E., Colombet, M., Ries, L. A. G., Moreno, F., Dolya, A., Bray, F., Hesseling, P., Shin, H. Y., Stiller, C. A., Bouzbid, S., Hamdi-Cherif, M., Hablas, A., Chirpaz, E., Buziba, N., Chesumbai, G., Manraj, S., Reynders, D., Wabinga, H., Chokunonga, E., . . . Steliarova-Foucher, E. (2017). International incidence of childhood cancer, 2001–10: a population-based registry study. *The Lancet Oncology*, *18*(6), 719–731. https://doi.org/10.1016/S1470-2045(17)30186-9

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y., Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell, 177(7), 1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031

Tian, K., Wang, X., Wu, Y., Wu, X., Du, G., Liu, W., & Wu, R. (2020). Relationship of tumour-associated macrophages with poor prognosis in Wilms' tumour. Journal of Pediatric Urology, 16(3), 376.e1-376.e8. https://doi.org/10.1016/j.jpurol.2020.03.016

Tran, T., Lindström, N. O., Ransick, A., De Sena Brandine, G., Guo, Q., Kim, A. D., Der, B., Peti-Peterdi, J., Smith, A. D., Thornton, M., Grubbs, B., McMahon, J. A., & McMahon, A. P. (2019). In Vivo Developmental Trajectories of Human Podocyte Inform In Vitro Differentiation of Pluripotent Stem Cell-Derived Podocytes. Developmental Cell, 50(1), 102-116.e6. https://doi.org/10.1016/j.devcel.2019.06.001

Trink, A., Kanter, I., Pode-Shakked, N., Urbach, A., Dekel, B., & Kalisky, T. (2018). Geometry of Gene Expression Space of Wilms' Tumors From Human Patients. Neoplasia, 20(8), 871–881. https://doi.org/10.1016/j.neo.2018.06.006

Uehara, S., Udagawa, N., Mukai, H., Ishihara, A., Maeda, K., Yamashita, T., Murakami, K., Nishita, M., Nakamura, T., Kato, S., Minami, Y., Takahashi, N., & Kobayashi, Y. (2017). Protein kinase N3 promotes bone resorption by osteoclasts in response to Wnt5a-Ror2 signaling. *Science Signaling*, *10*(494). https://doi.org/10.1126/scisignal.aan0023

Vakkila, J., Jaffe, R., Michelow, M., & Lotze, M. T. (2006). Pediatric Cancers Are Infiltrated Predominantly by Macrophages and Contain a Paucity of Dendritic Cells: a Major Nosologic Difference with Adult Tumors. Clinical Cancer Research, 12(7), 2049–2054. https://doi.org/10.1158/1078-0432.ccr-05-1824

van den Berg, C. W., Ritsma, L., Avramut, M. C., Wiersma, L. E., van den Berg, B. M., Leuning, D. G., Lievers, E., Koning, M., Vanslambrouck, J. M., Koster, A. J., Howden, S. E., Takasato, M., Little, M. H., & Rabelink, T. J. (2018). Renal Subcapsular Transplantation of PSC-Derived Kidney Organoids Induces Neo-vasculogenesis and Significant Glomerular and Tubular Maturation In Vivo. Stem Cell Reports, 10(3), 751–765. https://doi.org/10.1016/j.stemcr.2018.01.041

van den Heuvel-Eibrink, M., van Tinteren, H., Bergeron, C., Coulomb-L'Hermine, A., de Camargo, B., Leuschner, I., Sandstedt, B., Acha, T., Godzinski, J., Oldenburger, F., Gooskens, S., de Kraker, J., Vujanic, G., Pritchard-Jones, K., & Graf, N. (2015). Outcome of localised blastemal-type Wilms tumour patients treated according to intensified treatment in the SIOP WT 2001 protocol, a report of the SIOP Renal Tumour Study Group (SIOP-RTSG). *European Journal of Cancer*, *51*(4), 498–506. https://doi.org/10.1016/j.ejca.2014.12.011

Walz, A., Ooms, A., Gadd, S., Gerhard, D., Smith, M., Guidry Auvil, J. M., Meerzaman, D., Chen, Q. R., Hsu, C., Yan, C., Nguyen, C., Hu, Y., Bowlby, R., Brooks, D., Ma, Y., Mungall, A., Moore, R., Schein, J., Marra, M., . . . Perlman, E. (2015). Recurrent DGCR8, DROSHA, and SIX Homeodomain Mutations in Favorable Histology Wilms Tumors. *Cancer Cell*, *27*(2), 286–297. https://doi.org/10.1016/j.ccell.2015.01.003

Wegert, J., Ishaque, N., Vardapour, R., Geörg, C., Gu, Z., Bieg, M., Ziegler, B., Bausenwein, S., Nourkami, N., Ludwig, N., Keller, A., Grimm, C., Kneitz, S., Williams, R., Chagtai, T., Pritchard-Jones, K., Van Sluis, P., Volckmann, R., Koster, J., . . . Gessler, M. (2015). Mutations in the SIX1/2 Pathway and the DROSHA/DGCR8 miRNA Microprocessor Complex Underlie High-Risk Blastemal Type Wilms Tumors. *Cancer Cell*, *27*(2), 298–311. https://doi.org/10.1016/j.ccell.2015.01.002

Welsh, G. I., & Saleem, M. A. (2011). The podocyte cytoskeleton—key to a functioning glomerulus in health and disease. Nature Reviews Nephrology, 8(1), 14–21. https://doi.org/10.1038/nrneph.2011.151

Xu, P. X., Zheng, W., Huang, L., Maire, P., Laclef, C., & Silvius, D. (2003). Six1 is required for the early organogenesis of mammalian kidney. *Development*, *130*(14), 3085–3094. https://doi.org/10.1242/dev.00536

Yoshimura, Y., Taguchi, A., Tanigawa, S., Yatsuda, J., Kamba, T., Takahashi, S., Kurihara, H., Mukoyama, M., & Nishinakamura, R. (2019). Manipulation of Nephron-Patterning Signals Enables Selective Induction of Podocytes from Human Pluripotent Stem Cells. Journal of the American Society of Nephrology, 30(2), 304–321. https://doi.org/10.1681/asn.2018070747

You, J. S., Singh, N., Reyes-Ordonez, A., Khanna, N., Bao, Z., Zhao, H., & Chen, J. (2021). ARHGEF3 Regulates Skeletal Muscle Regeneration and Strength through Autophagy. *Cell Reports*, *34*(1), 108594. https://doi.org/10.1016/j.celrep.2020.108594

Young, M. D., Mitchell, T. J., Vieira Braga, F. A., Tran, M. G. B., Stewart, B. J., Ferdinand, J. R., Collord, G., Botting, R. A., Popescu, D. M., Loudon, K. W., Vento-Tormo, R., Stephenson, E., Cagan, A., Farndon, S. J., del Castillo Velasco-Herrera, M., Guzzo, C., Richoz, N., Mamanova, L., Aho, T., . . . Behjati, S. (2018). Single-cell transcriptomes from human kidneys reveal the cellular identity of renal tumors. *Science*, *361*(6402), 594–599. https://doi.org/10.1126/science.aat1699

Yun, K., Ajima, R., Sharma, N., Costantini, F., Mackem, S., Lewandoski, M., Yamaguchi, T. P., & Perantoni, A. O. (2014). Non-canonical Wnt5a/Ror2 signaling regulates kidney morphogenesis by controlling intermediate mesoderm extension. *Human Molecular Genetics*, *23*(25), 6807–6814. https://doi.org/10.1093/hmg/ddu397