PARK15/FBXO7 is dispensable for PINK1/Parkin mitophagy in iNeurons and HeLa cell systems

Felix Kraus[1,2], Ellen A. Goodall[1,2], Ian R. Smith[1], Yizhi Jiang[1], Julia C. Paoli[1], Jiuchun Zhang[1], Joao A. Paulo[1], and J. Wade Harper[1,2*]

[1]Department of Cell Biology, Blavatnik Institute, Harvard Medical School, 240 Longwood Ave, Boston MA 02115, USA
[2]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA
*Correspondence: wade_harper@hms.harvard.edu (J.W.H.)

ABSTRACT

The protein kinase PINK1 and ubiquitin ligase Parkin promote removal of damaged mitochondria via a feed-forward mechanism involving ubiquitin (Ub) phosphorylation, Parkin activation, and ubiquitylation of mitochondrial outer membrane proteins to support recruitment of mitophagy receptors. The ubiquitin ligase substrate receptor FBXO7/PARK15 is mutated in an early-onset parkinsonian-pyramidal syndrome. Previous studies have proposed a role for FBXO7 in promoting Parkin-dependent mitophagy. Here, we systematically examine the involvement of FBXO7 in depolarization-dependent mitophagy in the well-established HeLa and induced-neurons cell systems. We find that FBXO7$^{-/-}$ cells have no demonstrable defect in: 1) kinetics of pUb accumulation, 2) pUb puncta on mitochondria by super-resolution imaging, 3) recruitment of Parkin and autophagy machinery to damaged mitochondria, 4) mitophagic flux, and 5) mitochondrial clearance as quantified by global proteomics. Moreover, global proteomics of neurogenesis in the absence of FBXO7 reveals no obvious alterations in mitochondria or other organelles. These results argue against a general role for FBXO7 in Parkin-dependent mitophagy and point to the need for additional studies to define how FBXO7 mutations promote parkinsonian-pyramidal syndrome.

INTRODUCTION

Organelle quality control underlies cellular health and is often defective in disease and pathological states. Arguably the best understood such quality control pathway is mitophagy, wherein damaged or supernumerary mitochondria are targeted for removal from the cell via selective autophagy (Goodall *et al*, 2022; Harper *et al*, 2018; McWilliams & Muqit, 2017; Pickrell & Youle, 2015). In this process, individual organelles are marked for sequestration within a double-membrane vesicle called an autophagosome, which then fuses with a lysosome to facilitate degradation of the organelle by resident lysosomal hydrolases (Dikic & Elazar, 2018; Stolz *et al*, 2014). Multiple forms of mitophagy have been reported, which differ in the types of regulatory mechanisms involved in marking the organelle for degradation, but these fall into two primary types – ubiquitin-dependent and ubiquitin-independent (Harper *et al.*, 2018; Pickrell & Youle, 2015). Our understanding of ubiquitin-dependent mitophagy has been advanced by the discovery of a signalling pathway composed of the PINK1 protein kinase and the Parkin (also called PRKN) ubiquitin (Ub) E3 ligase, which marks damaged mitochondria for elimination (Narendra *et al*, 2008). Parkin and PINK1 are both mutated in early-onset recessive forms of Parkinson's disease and understanding how these enzymes work has been a major focus of the field (Ng *et al*, 2021; Pickrell & Youle, 2015).

In healthy mitochondria, PINK1 is imported into the mitochondrial translocon and
rapidly processed for degradation (Jin *et al*, 2010; Yamano & Youle, 2013). In response to
mitochondrial damage – such as depolarization, accumulation of mitochondrial misfolded
proteins, or defects in mitochondrial fusion – PINK1 is stabilized on the mitochondrial outer
membrane (MOM) in association with the translocon (Lazarou *et al*, 2012), where it can
phosphorylate Ser65 on Ub already conjugated to proteins on the MOM (Kane *et al*, 2014;
Kazlauskaite *et al*, 2015; Koyano *et al*, 2014; Ordureau *et al*, 2014; Wauer *et al*, 2015).
Accumulation of phospho S65-Ub (referred to henceforth as pUb) on the MOM promotes
recruitment of Parkin – a pUb-binding protein – thereby facilitating phosphorylation of Parkin
on S65 of its N-terminal Ub-like (UBL) domain and activation of its Ub ligase activity
(Gladkova *et al*, 2018; Kane *et al.*, 2014; Kazlauskaite *et al.*, 2015; Ordureau *et al.*, 2014;
Sauve *et al*, 2018; Wauer *et al.*, 2015). Parkin then ubiquitylates numerous MOM proteins,
resulting in both the accumulation of additional pUb and further MOM protein ubiquitylation
(Antico *et al*, 2021; Bingol *et al*, 2014; Ordureau *et al*, 2020; Sarraf *et al*, 2013). MOM
ubiquitylation then promotes recruitment of selective autophagy cargo receptors including
OPTN, CALCOCO2 (also called NDP52), and SQSTM1 (also called p62) to facilitate the
assembly of an autophagosome around the ubiquitylated organelle (Evans & Holzbaur,
2020; Heo *et al*, 2015; Lazarou *et al*, 2015; Wong & Holzbaur, 2014). Thus, PINK1, Parkin,
and pUb function in a positive feedback loop to promote selective ubiquitylation and
elimination of the damaged organelle (Goodall *et al.*, 2022).
Previous work (Burchell *et al*, 2013) concluded that FBXO7, the product of an early-
onset parkinsonian-pyramidal syndrome risk gene also designated as *PARK15* (Di Fonzo *et*
al, 2009; Houlden & Singleton, 2012; Paisan-Ruiz *et al*, 2010), is a positive modulator of
Parkin-dependent mitophagy (**Supplemental Figure 1A**). FBXO7 is a member of the F-box
family of proteins, which assemble with SKP1, CUL1, and RBX1 to form a modular SCF Ub
ligase complex wherein the F-box protein to binds substrates (Jin *et al*, 2004). FBXO7 is
characterized by an N-terminal UBL domain, a central FP domain with structural similarity to
a domain in the proteasome inhibitory factor PI31/PSMF1, followed by the F-box motif, and
a C-terminal proline rich region (Kirk *et al*, 2008). Patient mutations in FBXO7 are found in
multiple regions of the protein (Di Fonzo *et al.*, 2009; Houlden & Singleton, 2012; Paisan-
Ruiz *et al.*, 2010). Initial studies on FBXO7 proposed a functional interaction with the
Parkin/PINK1 pathway and concluded that FBXO7 functions as a positive regulator of
mitophagy through direct interaction with Parkin and with PINK1 (Burchell *et al.*, 2013).
Depletion of FBXO7 by siRNA in cell lines was reported to reduce depolarization-dependent
loss in mitochondria based on immunoblotting of matrix proteins and likewise to reduce
depolarization-dependent ubiquitylation of MFN1, a substrate of Parkin (Burchell *et al.*,
2013). However, studies that further substantiate a functional link between FBXO7 and

mitophagy are lacking, although FBXO7 depletion has recently been reported to alter

mitochondrial dynamics (Al Rawi *et al*, 2022).

Given the strong genetic association of FBOX7/PARK15 with Parkinson's Disease

(Houlden & Singleton, 2012) and major advances in our understanding of Parkin-dependent

mitophagy (Goodall *et al.*, 2022) since initial links between FBXO7 and mitophagy were

reported, we set out to elucidate how FBXO7 might function within the positive feedback

loop to promote mitophagy (**Supplemental Figure 1A**). Relevant advances include the

identification of pUb as a sentinel response to PINK1/Parkin activation (Kane *et al.*, 2014;

Kazlauskaite *et al.*, 2015; Koyano *et al.*, 2014; Ordureau *et al.*, 2014; Wauer *et al.*, 2015),

the development of highly quantitative mitophagic flux assays using a mitochondrially

localized Keima fluorescent reporter protein (Katayama *et al*, 2011; Lazarou *et al.*, 2015),

and the development of genetically tractable iNeuron systems for functional analysis of the

Parkin pathway in neuronal cell types (Ordureau *et al.*, 2020; Ordureau *et al*, 2018).

Unexpectedly, we find no evidence of a role for FBXO7 in the kinetics of depolarization-

dependent pUb accumulation, or mitophagic flux in either iNeurons or the conventional HeLa

cell used extensively to examine Parkin and PINK1-dependent mitophagy (Narendra *et al.*,

2008). Moreover, global proteomics of FBXO7$^{-/-}$ HeLa cells undergoing mitophagy revealed

no defect in mitochondrial elimination by autophagy. Taken together with previous studies

that failed to verify Parkin-FBXO7 interactions during mitophagy (Sarraf *et al.*, 2013), these

data indicate that FBXO7 does not play a general role as a positive modulator of Parkin and

PINK1-dependent mitophagy in HeLa and iNeuron systems. This work suggests that

elucidation of FBXO7 biochemical functions is needed to understand how it might contribute

to suppression of early-onset parkinsonian-pyramidal syndrome.

RESULTS

A tool kit for functional analysis of cells lacking FBXO7

To test if FBXO7 is a pivotal amplifier of the PINK1/Parkin pathway, we employed

CRISPR-Cas9-based gene editing to generated knockouts for FBXO7 in HeLa HFT/TO-

PRKN cells (Ordureau *et al*, 2015) and confirmed insertion of a frameshift mutation by

sequencing and immunoblotting (**Supplemental Figure 1 B-D**). HeLa HFT/TO-PRKN cells

contain a DOX-inducible PRKN cassette allowing regulated expression of Parkin to facilitate

an analysis of mitophagic flux in response to mitochondrial damage (Ordureau *et al.*, 2015).

Three independent FBXO7$^{-/-}$ clones were identified (clones 27, 41 and 45) (**Supplemental**

Figure 1C,D). In parallel, we generated two independent human ES cell clones lacking

FBXO7 via CRISPR-Cas9 (clones 52 and 89) (**Supplemental Figure 1B,E,F**). Here, an ES

cell line harbouring a DOX inducible NGN2 gene was used, allowing for conversion of these

cells to cortical-like iNeurons over a two week or greater time course (Ordureau *et al.*, 2020;

Zhang *et al*, 2013). We have previously demonstrated that iNeurons derived from these cells
exhibit Parkin and PINK1-dependent mitophagic flux in response to mitochondrial damage
(Ordureau *et al.*, 2020). Both clones displayed normal karyotypes and the absence of
FBXO7 was verified using MiSeq sequencing (**Supplemental Figure 1F,G**).

Robust depolarization-dependent Ub phosphorylation by PINK1 in cells lacking
FBXO7

The earliest known event in Parkin-dependent mitophagy is accumulation of PINK1
on the mitochondrial outer membrane followed by rapid S65 phosphorylation of pre-existing
Ub linked with mitochondrial proteins (**Supplemental Figure 1A**), as has been worked out in
detail using the well-validated HeLa cell system with ectopically expressed Parkin (Kane *et*
al., 2014; Lazarou *et al.*, 2012; Narendra *et al*, 2010; Yamano & Youle, 2013). In the
presence of Parkin, Ub phosphorylation is enhanced as a result of increased Ub conjugation
to mitochondrial proteins, thereby providing additional substrate molecules for
phosphorylation by PINK1 (Kane *et al.*, 2014; Koyano *et al.*, 2014; Ordureau *et al.*, 2014). To
examine pUb accumulation in the context of HeLa cells lacking FBXO7, we induced Parkin
expression (16 h) and then depolarized mitochondria for 1 or 6 h with antimycin
A/Oligomycin A (AO), which inhibit Complex III and Complex IV activity in the electron
transport chain, respectively. As expected, control HeLa cells display time-dependent
accumulation of pUb as measured by immunoblotting, but the absence of FBXO7 had no
impact on the extent of pUb accumulation, as assessed in all three FBXO7$^{-/-}$ cell lines in
biological triplicate analyses (**Figure 1A,B**).

Similarly, we assessed depolarization-dependent pUb accumulation in the iNeuron
system. control or FBXO7$^{-/-}$ ES cells were converted to day 12 iNeurons and depolarized
with AO in duplicate (**Figure 1C**). pUb was detected by immunoblotting at 6 and 24 h and
was indistinguishable between WT and two independent FBXO7$^{-/-}$ clones (**Figure 1D**).
Taken together, these data do not support a universal role for FBXO7 in promoting the
earliest step in mitophagy signalling – PINK1-dependent Ub phosphorylation – in the HeLa
cell system or iNeurons.

Super-resolution imaging of pUb formation independent of FBXO7

While depolarization-dependent Ub phosphorylation was not impaired, we
considered the possibility that the organization of pUb on damaged mitochondria might be
altered in cells lacking FBXO7. To our knowledge, super-resolution microscopy has not been
used to examine the kinetics and spatial properties of pUb accumulation in response to
depolarization. We therefore optimized conditions to monitor pUb via volumetric 3D-SIM
super-resolution microscopy in both HeLa and iNeuron systems (see **EXPERIMENTAL**

MATERIALS AND METHODS). After AO treatment of HeLa cells for 1, 3 or 6 h, the

mitochondria matrix protein HSP60 and pUb were examined by super-resolution imaging. In

control cells at 1 h, fragmented mitochondria were often proximal to pUb foci with an

average volume of 5.71 µm^3, which were not observed in cells lacking PINK1 (**Figure 2A-C,**

Supplemental Figure 2A,B). The volume of pUb increased ~8-fold over 6 h (47.88 µm^3),

consistent with the known feed-forward response (**Figure 2B, Supplemental Figure 2A**). At

6 h of AO treatment, we observed a near complete coating of mitochondrial fragments with

pUb, as quantified by the significant increase in pUb object volume (**Figure 2B,C,**

Supplemental Figure 2A). Consistent with analysis of pUb by immunoblotting, two FBXO7$^{-/-}$

clones did not show any significant differences in pUb recruitment, volume or overall

morphological differences compared to control cells, indicating that FBXO7 is not required

for the feed-forward amplification of PINK1/Parkin mitophagy after AO-treatment (**Figure 2A-**

C). Analogous super-resolution experiments in day 12 iNeurons with 0.5, 1, and 3 h of AO

revealed similar results, with no obvious effect of FBXO7 deletion on pUb recruitment,

volume or overall morphology when compared to WT iNeurons (**Figure 2D-F,**

Supplemental Figure 3A). As expected (Ordureau *et al*., 2020; Ordureau *et al*., 2018),

iNeurons lacking PINK1 did not accumulate pUb (**Figure 2D,E**). These results were also

confirmed in iNeurons treated with AO for 1 or 6 h using conventional immunofluorescence

and image analysis of HSP60 and pUb co-localization (**Figure 2G,H**). To ensure that kinetic

effects were not obscured by overt depolarization, we reduced the concentrations of AO

used by 10-fold, but again no defect in pUb recruitment to mitochondria was observed

(**Supplemental Figure 3B,C**). These data indicate that the absence of FBXO7 does not

alter the ability of PINK1-dependent pUb foci to accumulate on damaged mitochondria in

either HeLa or iNeuron systems.

Parkin and autophagy regulator recruitment in FBXO7$^{-/-}$ cells

Parkin recruitment depends upon the accumulation of pUb on damaged mitochondria

(**Supplemental Figure 1A**) (Okatsu *et al*, 2015). Given that the existing model for FBXO7 in

Parkin-dependent mitophagy posited that Parkin and FBXO7 physically interact and that

FBXO7 promotes Parkin function (Burchell *et al*., 2013), we directly tested whether FBXO7

is required for Parkin recruitment to mitochondria using the HeLa cell system. Control or

FBXO7$^{-/-}$ cells were stably transduced with a lentivirus expressing GFP-Parkin and subjected

to imaging with or without depolarization with AO (**Supplemental Figure 4A**). We used

CellProfiler (Stirling *et al*, 2021) to analyse the characteristics of GFP-Parkin and

mitochondria and their relationship to each other. By segmenting out cytosolic and

mitochondrially-localized Parkin, we could quantify alterations in Parkin abundance within

the mitochondrial mask as a measure of Parkin-translocation (**Supplemental Figure 4B**).

Our expectation was that a requirement for FBXO7 in Parkin recruitment would result in
reduced Parkin intensity within the mitochondrial mask in response to depolarization. First,
we tested whether mitochondrial morphology (mitochondrial integrated density [==mtIntDen]
as proxy, stained with HSP60) is changing after AO-treatment on a single-cell level. Under
fed conditions, control and three FBXO7$^{-/-}$ clones displayed comparable mtIntDen.
Depolarization by AO led to the expected mitochondrial fragmentation and clustering, thus
causing an increase in local HSP60 signal and a right-shift in mtIntDen (**Supplemental**
Figure 4C, top panel). This shift was also observed to a similar extend in FBXO7$^{-/-}$ cells.
Thus, both control and FBXO7$^{-/-}$ cells robustly respond to AO-induced mitophagy. Next, we
tested if Parkin intensities are increasing on mitochondria on an "organelle object" level. As
expected, mtIntDen did shift to the right after AO-treatment, in line with previous
experiments. Parkin signal was comparable between the genotypes under fed conditions,
and increased in intensity after AO-treatment, consistent with Parkin accumulation inside the
mitochondrial area (**Supplemental Figure 4C, lower panel**). This Parkin shift was observed
in both control and FBXO7$^{-/-}$ cell lines, arguing that FBXO7 is not universally required for
Parkin recruitment to damaged mitochondria.

Mitochondrial ubiquitylation by Parkin promotes recruitment of downstream
autophagy machinery, including the FIP200-ULK1 complex and Ub-bind autophagy
adaptors, ultimately leading to the formation of an autophagosome (Heo *et al.*, 2015;
Lazarou *et al.*, 2015; Ravenhill *et al*, 2019; Vargas *et al*, 2019). Lipidation of LC3 is
associated with autophagosome formation and is thought to contribute to recruitment of
autophagy receptors. As expected, depolarization of Parkin-expressing HeLa cells resulted
in recruitment of both p62 and FIP200 to mitochondria in a manner that dependent on
PINK1, as revealed by immunostaining and confocal imaging (**Supplemental Figure 4D**).
However, FBXO7$^{-/-}$ cells were as proficient as control cells in recruitment of p62 or FIP200
after 3 h or 16 h or AO-induced mitophagy (**Supplemental Figure 4D-G**). Previous studies
using siRNAs targeting FBXO7 also reported that LC3 lipidation in response to mitochondrial
depolarization is defective compared with non-depleted cells (Burchell *et al.*, 2013).
However, all three FBXO7$^{-/-}$ HeLa cell clones underwent depolarization-dependent LC3
lipidation to similar extents, consistent with a functional Parkin pathway (**Supplemental**
Figure 4H,I). Taken together, these data indicate that FBXO7 is not universally required to
promote either Parkin-dependent recruitment to damaged mitochondria or subsequent steps
that depend upon Ub chain assembly on mitochondria such as recruitment of autophagy
machinery and initiation of LC3 lipidation.

Additionally, we have not able to reproduce a previously reported association
between overexpressed FBXO7 and PINK1 (Huang *et al*, 2020) in either HEK293T or
HCT116 cells. Briefly, as part of our BioPlex project (Huttlin *et al*, 2021), we ectopically

expressed PINK1-HA and performed interaction proteomics. While we identified the known
protein kinase chaperone HSP90 subunits in association with PINK1, we did not detect
FBXO7 (**Supplemental Figure 4J, left panel; Supplemental Table S1**). Likewise, FBXO7
associated reciprocally with members of the CUL1-SKP1-RBX1, the COP9/Signalosome,
and both core particle and regulatory subunits of the proteasome in HEK293T and/or
HCT116 cells, as expected (Bader *et al*, 2011; Liu *et al*, 2019; Vingill *et al*, 2016), but neither
PINK or Parkin were detected (**Supplemental Figure 4J,K; Supplemental Table S2**)
(Huttlin *et al.*, 2021) (see **METHODS**). These results are consistent with our previous Parkin
interaction proteomics experiments that failed to demonstrate an interaction with FBXO7
during mitophagy induction (Sarraf *et al.*, 2013).
250
Mitophagic flux in FBXO7$^{-/-}$ cells
A significant advance has been the development of mitophagy reporters, which
cumulatively measure flux of mitochondrial turnover through to the final stages of mitophagy:
the fusion of mitochondria-laden autophagosomes with lysosomes (Lazarou *et al.*, 2015).
mKeima is a fluorescent protein that undergoes a pH-dependent Stokes shift, and delivery of
a Keima-tagged cargo from the cytoplasm to the acidic interior of the lysosome can be
monitored by measuring the ratio of emission at 620 nm with maximal excitation at 440 nm
or 586 nm under neutral or acidic conditions, respectively, by flow cytometry or microscopy
(**Figure 3A**). To monitor differences in mitophagic flux upon deletion of FBXO7, we
employed mtKeima, an mKeima protein targeted to the mitochondrial with a COX8-matrix
targeting sequence (Ordureau *et al.*, 2020).
In Parkin-expressing HeLa cells, the ratio of acidic-Keima (measured with excitation
at 561 nm and emission 620 nm) to neutral Keima (ex. 405 nm, em. 603 nm) increases as
cells are treated with AO (**Figure 3B, Supplemental Figure 4L**). This mitophagic flux is
PINK1-dependent as deletion of PINK1 abolishes this shift (**Figure 3B, Supplemental
Figure 4L**). As expected, there is little flux without depolarization with AO, and inhibition of
the lysosomal V-ATPase with BafilomycinA1 (BafA) restores the mtKeima acidic:neutral ratio
to levels similar to that seen fed cells (**Supplemental Figure 4L**). In three FBXO7$^{-/-}$ clones,
AO-dependent mitophagic flux is similar to control cells with increasing flux observed with
increasing length of AO treatment (**Figures 3B; Supplemental Figure 4L**). We validated the
mtKeima flux results using the mitochondrial DNA clearance assay (Heo *et al.*, 2015;
Lazarou *et al.*, 2015). Staining for mtDNA foci per cell after 24 h AO treatment in control and
FBXO7$^{-/-}$ HeLa cells showed a significant decrease in mtDNA number in all samples (**Figure
3G**). The remaining mtDNA levels were indistinguishable between control and FBXO7$^{-/-}$ cells
(**Figure 3G**).

We also measured mtKeima foci and flux in control and FBXO7$^{-/-}$ iNeurons at day 12
of differentiation (**Figure 3C-E**). As with HeLa cells, red-shifted mtKeima foci increased ~2-
fold after 6 h AO treatment in control and FBXO7$^{-/-}$ cells (**Figure 3C,D**). Using flow
cytometry, mitophagic flux (normalized to BafA) after 6 and 24 h AO treatment was
comparable between control and FBXO7$^{-/-}$ iNeurons, and was absent in PINK1$^{-/-}$ cells
(**Figure 3E**). Additionally, we investigated mtDNA turnover in d12 control and two
independent FBXO7$^{-/-}$ iNeuron clones. Within 6 or 24 h of depolarization of iNeurons, we
observed a reduction in the number of mitochondrial DNA/cell, and the effect was
independent of the presence or absence of FBXO7 (**Figure 3F,G**). Taken together, these
data indicate that FBXO7 is not universally required for depolarization-dependent mitophagic
flux in either the HeLa or iNeuron systems.
287
Proteomic analysis of HeLa cells lacking FBXO7 during mitophagy
As an alternative to mtKeima for measuring mitophagic flux, we examined the total
proteome of depolarized HeLa cells with the expectation that in control HeLa cells,
mitochondrial turnover would result in bulk loss of the mitochondrial proteome in depolarized
cells. Thus, control or FBXO7$^{-/-}$ HeLa cells expressing Parkin were depolarized for 16 h in
triplicate and total cell extracts subjected to 18-plex Tandem Mass Tagging (TMT)-based
proteomics (**Figure 4A, Supplemental Table S3,S4**) (Li *et al*, 2021). Replicates were highly
correlated, with correlation coefficients greater that 0.95 and Principal Component Analysis
(PCA) revealed tight clustering of replicates, with the major feature separating the samples
being AO-treatment (**Supplemental Figure 5A,B**). Through hierarchical clustering
(**Supplemental Figure 5C**), we identified major alterations in the abundance of mitochondria
in response to depolarization (**Figure 4B, Supplemental Figure 5D-I**). Indeed, the majority
of proteins annotated as mitochondrial were found to have reduced levels in control cells in
response to depolarization, as indicated in the overlap of mitochondrial proteins on the
volcano plots for the ~8000 proteins quantified (leftward skew of red dots in **Figure 4B**, left
panel). Importantly, the patterns of protein abundance were indistinguishable from control for
two independent FBXO7$^{-/-}$ cell lines, with leftward skew of mitochondrial proteins in the
volcano plot (**Figure 4B,C**). Proteins with altered abundance were enriched for mitochondrial
matrix, inner membrane, outer membrane and oxidative phosphorylation categories,
consistent with autophagy of entire organelles (**Figure 4C**). The behaviour of other
organelles and large protein complexes were similar for both control and FBXO7$^{-/-}$ cells
(**Figure 4C**). Moreover, both FBXO7 clones displayed similar behaviour to each other and to
the control, as indicated by correlation analysis (Pearson's R=0.82, 0.86 and 0.74,
respectively) (**Figure 4D,E**). These results are consistent with the absence of a discernible
alteration in depolarization-dependent mitophagic flux in cells lacking FBXO7. A previous

report proposed that depletion of FBXO7 resulted in accumulation of PINK1 proteins levels
under basal conditions (Huang *et al.*, 2020), but in our experiments, PINK1 levels in the two
FBXO7$^{-/-}$ clones were unaffected based on TMT intensities (**Figure 4F**).
316

Proteomic analysis of human ES cells lacking FBXO7 during neurogenesis

As an initial unbiased approach to examining FBXO7 function in iNeurons, we
performed 18-plex TMT proteomics on control and two clones of FBXO7$^{-/-}$ cells at day 0, 4
and d12 of differentiation (**Figure 5A, Supplemental Table S5-S8**). We have previously
demonstrated dramatic remodelling of mitochondria around day 4 of differentiation to
support a switch from glycolysis to oxidative phosphorylation, which is accompanied by
BNIP3L-dependent mitophagy of a fraction of mitochondria (Ordureau *et al*, 2021). From the
~6000 proteins quantified across all replicates and conditions, we found strong
correspondence of control and FBXO7$^{-/-}$ cells, with PC1 being drive by differentiation
(**Supplemental Figure 6A,B**). FBXO7 itself was expressed at comparable levels across the
time course but was not detected in FBXO7$^{-/-}$ iNeurons, as expected (**Supplemental Figure
6C**). Remarkably, the abundance of the proteome was largely immune to the loss of FBXO7
at both day 4 and day 12 of differentiation (**Figure 5B-D, Supplemental Figure 6D**).
Moreover, the abundance of mitochondrial proteins (including inner and out membrane and
OXPHOS components) were unchanged relative to control cells at day 4 or 12 of
differentiation (**Figure 5B, Supplemental Figure 6D**).
Recent reports have directly linked L250P mutations in the dimerization domain of
FBXO7 to MiD49/MiD51, a core adapter protein involved in mitochondrial fission (Al Rawi *et
al.*, 2022). Indeed, DNM1L-mediated mitochondrial fission has been previously indicated in
the efficient scission of damaged parts of mitochondria and protection of the remaining
healthy mitochondrial network (Burman *et al*, 2017). Cells expressing FBXO7^{L250P} were
found to have reduced levels of proteasome subunits as well (Al Rawi *et al.*, 2022). Our
proteomic data from iNeurons indicates no alterations in the abundance of proteasome
subunits or mitochondrial fission and fusion machinery (DNM1L, MFF, and MiD51) in
iNeurons lacking FBXO7 (**Figure 5C,E**). To test if loss of FBXO7 causes in morphological
changes in the mitochondrial network, we stained day 12 iNeurons with for the matrix marker
HSP60 prior to image analysis (**Figure 5F,G**). However, we observed no significant
alterations in the number of mitochondria/cell or the mean mitochondrial length in control or
FBXO7$^{-/-}$ iNeurons. The mean mitochondrial circularity was slightly reduced in FBXO7$^{-/-}$
iNeurons treated with AO for 3h (**Figure 5F**). Analogous analyses of mitochondria and
fission/fusion proteins also did not reveal significant alterations in response to loss of FBXO7
in two independent clones of HeLa cells (**Figure 5H,I**). However, there was a ~10%
decrease in the abundance of proteasome subunits in HeLa cells lacking FBXO7 in the

absence of depolarization, but an increase after mitophagy induction (**Supplemental Figure**
5E, Figure 4C). If this response is a compensatory mechanism of the knockout cell line to
cope with cytotoxic insults is an intriguing idea. Last, we examined whether FBXO7$^{-/-}$ is
required for the maintenance of the mitochondrial organelle pool during differentiation.
Previous work has shown significant metabolic rewiring during the differentiation from hESC
to iNeurons, including a switch from glycolysis to oxidative phosphorylation as the cells main
energy source (Ordureau *et al.*, 2021). Since this switch is accompanied by increased
mitophagy, we set out to see if the absence of FBXO7 would lead to accumulation of
mitochondria and/or impair differentiation. FBXO7$^{-/-}$ cells did not display obvious differences
in the abundance of mitochondrial proteins, when normalized to day 0 (**Figure 5B,**
Supplemental Figure 6D). Likewise, other abundance of other organelles was unchanged
(**Figure 5C**). Finally, neither differentiation markers or autophagy proteins displayed obvious
changes in control versus the two FBXO7$^{-/-}$ clones as measured by TMT-proteomics and
displayed using a T^2-statstic (Ordureau *et al.*, 2021) (**Supplemental Figure 7A-F**).
364
DISCUSSION
Despite strong genetic evidence linking mutations in FBXO7 with parkinsonian-
pyramidal syndrome (Di Fonzo *et al.*, 2009; Houlden & Singleton, 2012; Paisan-Ruiz *et al.*,
2010), our understanding of the cellular roles of FBXO7 are limited. An early study reported
biochemical links between FBXO7 and Parkin/PINK1-dependent mitophagy in fibroblasts
and SH-SY5Y cells, including interaction of overexpressed FBXO7 with both PINK1 and
Parkin (Burchell *et al.*, 2013). Using siRNA to deplete FBXO7, it was also concluded that
FBXO7 promotes Parkin's ability to ubiquitylate the outer mitochondrial membrane substrate
MFN1 and also promotes clearance of mitochondria by autophagy. These biochemical and
physiological results led to the conclusion that FBXO7 functions as a biochemical amplifier
of Parkin/PINK1-deepndent mitophagy (Burchell *et al.*, 2013). However, follow-up studies
have been limited, and the field has experienced dramatic advances in the understanding of
the biochemical roles of Parkin and PINK1 in promoting mitochondrial clearance by
autophagy, thereby providing the tools to examine potential roles of FBXO7 in the pathway.
With these tools, we set out to define where in the pathway FBXO7 might operate to
promote mitophagy. However, in both the HeLa cells with overexpressed Parkin and
iNeurons with a fully endogenous Parkin/PINK1 pathway, we failed to validate a role for
FBXO7 in any of several steps in the pathway, including pUb accumulation, Parkin
recruitment to the mitochondrial outer membrane, recruitment of autophagy machinery,
mitophagic flux as measured by mtKeima, and mitochondrial proteome degradation. The
finding that deletion of FBXO7 has no obvious effect on the pathway with either endogenous
or overexpressed Parkin in iNeurons and HeLa cells indicates that FBXO7 does not play a

general or universally required role in the pathway. A previous study largely based on
overexpression concluded that FBXO7 may bind and regulate PINK1 levels, with <2-fold
increases in PINK1 by immunoblotting in FBXO7-depleted cells (Huang *et al.*, 2020).
However, our quantitative proteomics experiments (**Figure 4**) indicated that deletion of
FBXO7 has no effect on PINK1 abundance, consistent with the finding that pUb
accumulation in response to depolarization is unaffected in FBXO7$^{-/-}$ cells. Moreover, our
previous interaction proteomics experiments failed to identify either PINK1 in association
with overexpressed FBXO7 (Huttlin *et al.*, 2021) or FBXO7 in association with
overexpressed Parkin (Sarraf *et al.*, 2013). Thus, our functional and biochemical results do
not support a biochemical linkage between FBXO7 and the PINK1-Parkin pathway.

In order to broadly examine FBXO7$^{-/-}$ iNeurons for pathways that are affected, we
performed global proteomic analysis during neurogenesis *in vitro*. However, we did not
observe any alterations in pathways linked with mitochondria, or other quality control
pathways, although a reduction in the abundance of alcohol dehydrogenase enzymes
among a small number of other proteins, was observed. Whether these signatures are
related to FBXO7 function remains to be examined. While a recent study linked the L250P
mutation in the dimerization domain of FBXO7 with alterations in proteins associated with
mitochondrial fusion and fission (Al Rawi *et al.*, 2022), we did not observe alterations in
either the abundance of these proteins (MiD51, MFF, FIS1) nor did we observe changes in
mitochondrial morphology in either HeLa cells or iNeurons lacking FBXO7.

An apparent ortholog of FBXO7 was identified in *Drosophila* – referred to as
nutcracker, *ntc* – and shown to associate with components of the core particle of the
proteasome, as well as with DmPI31 (Bader *et al.*, 2011). Association of human FBXO7 with
proteasome components and PI31(also called PSMF1) have also been observed in both
focused studies and in our previous interaction proteomic studies (Al Rawi *et al.*, 2022;
Huttlin *et al.*, 2021; Vingill *et al.*, 2016). In this context, PI31/DmPI31 and FBXO7/*ntc* have
been proposed to functionally link proteasome trafficking via dynein motors in neuronal
processes (Liu *et al.*, 2019). Our interaction proteomics analysis has also confirmed
association of FBXO7 with PSMF1, CRL, COP9/Signalosome, and proteasome core
subunits (**Supplemental Figure 4J,K**). Further studies are required to elucidate the
relationship between FBXO7 and parkinsonian-pyramidal syndrome, to understand any
functional relationships between FBXO7 and other Parkinson's disease risk alleles, and to
examine whether FBXO7's physical association with the proteasome is linked with disease.
420

ACKNOWLEDGMENTS
This work was supported by Aligning Science Across Parkinson's (ASAP) (J.W.H.), the NIH
(R01 NS083524 to J.W.H. and K01DK098285 to J.A.P.), and the Harvard Medical School

Cell Biology Initiative for Molecular Trafficking and Neurodegeneration. Michael J Fox

Foundation administers the grant ASAP-000282 on behalf of ASAP and itself. For the

purpose of open access, the author has applied a CC-BY public copyright license to the

Author Accepted Manuscript (AAM) version arising from this submission. We thank Laura

Pontano Vaites for help with the BioPlex 3.0 interactome analysis. We thank Jennifer

Waters, Talley Lambert, Federico Gasparoli and Rylie Walsh in the Nikon Imaging Center

and the Cell Biology Microscopy Facility (CBMF) at Harvard Medical School for microscopy

support and technical support.

KEYWORDS

FBXO7, mitophagy, proteomics, iNeurons

DATA & CODE AVAILABILITY

Proteomic data and analysis files part of this study are deposited at ProteomeXchange

Consortium by the PRIDE partner (Deutsch *et al*, 2020; Perez-Riverol *et al*, 2022). The

PRIDE project identification number is PXD037797 and can be accessed for reviewers

under username reviewer_pxd037797@ebi.ac.uk; password 65wg8SiI. Macros and

pipelines used in this work can be found on GitHub

(https://github.com/harperlaboratory/FBXO7.git) or Zenodo

(https://doi.org/10.5281/zenodo.7258918). Raw files associated with this work are deposited

on Zenodo (https://doi.org/10.5281/zenodo.7268030).

AUTHOR CONTRIBUTIONS

Study design: J.W.H., F.K., E.A.G.; Data collection: F.K., E.A.G., J.A.P., J.Z., Y.Z.; Data

analysis and interpretation: F.K., E.G., I.R.S., J.A.P., J.Z., Y.Z. L.P.V.; Manuscript

preparation: J.W.H., F.K. with input from all authors.

DECLARATION OF INTERESTS

J.W.H. is a consultant and founder of Caraway Therapeutics and a founding scientific

advisory board member of Interline Therapeutics. No other authors declare a conflict of

interest.

 FIGURES

Figure 1: Robust depolarization-dependent Ub phosphorylation by PINK1 cells
lacking FBXO7. (**A**) Immunoblot for pUb on HeLa control and FBXO7$^{-/-}$ cells after treatment
with AO for indicated times. (**B**) Quantification of pUb signal relative to loading. Immunoblots
run in triplicates. Error bars depict S.D.. (**C**) Scheme depicting production of iNeurons for
biochemical analysis. (**D**) Immunoblot for pUb on extracts from day 12 iNeuron control and
FBXO7$^{-/-}$ cells after treatment with BafA or AO for indicated times.
466
467

Figure 2. Super-resolution imaging of pUb formation independent of FBXO7. (**A**) 3D-SIM images of HeLa control, PINK1$^{-/-}$ and FBXO7$^{-/-}$ cell lines after AO-induced mitophagy. Cells were stained for nuclear DNA (DAPI), mitochondria (HSP60) and pUb. Zoom-ins of regions of interested are enlarged in the middle panel. 3D-surface renderings of insets are shown on the right. Scale bar = 5 µm or 1 µm. (**B,C**) Evaluation of 3D-SIM images from HeLa datasets. The changes in pUb volume and minimal distances between mitochondria and pUb after mitophagy-induction are plotted. Error bars depict S.D. from 8-14 measured cells per condition. (**D**) 3D-SIM images of iN day 12 control, PINK1$^{-/-}$ and FBXO7$^{-/-}$ cell lines after AO-induced mitophagy. Cells were stained for nuclear DNA (DAPI), mitochondria (HSP60) and pUb. Zoom-ins of regions of interested are enlarged in the middle panel. 3D-surface renderings of insets are shown on the right. Scale bar = 5 µm or 1 µm. (**E,F**) Evaluation of 3D-SIM images from iNeuron datasets. The changes in pUb volume and minimal distances between mitochondria and pUb after mitophagy-induction are plotted. Error bars depict S.D. from 7-14 measured cells per condition. (**G**) Confocal images of iNeuron d12 Control, PINK1$^{-/-}$ and FBXO7$^{-/-}$ cell lines after AO-induced mitophagy. Cells were stained for nuclear DNA (Hoechst33342), mitochondria (HSP60) and pUb. Scale bar = 10µm and 5µm. (**H**) Evaluation of pUb volume after mitophagy induction. Error bars depict S.D. from 3 replicates.

Figure 3. Mitophagic flux in iNeurons and HeLa cells lacking FBXO7. (**A**) Scheme depicting mitophagic flux assay using mtKeima. See text for details. (**B**) Mean Acidic:Neutral mtKeima per-cell ratios measured by flow cytometry for HeLa cells expressing Parkin ndicating the number of hours treated with AO (Antimycin A (0.5 µM) and Oligomycin (0.5 µM)) or three hours with 25 nM BafilomycinA (BafA). Error bars depict S.D. from triplicate measurements. (**C**) Indicated hESCs expressing mtKeima were differentiated to iNeurons (day 12) and were subjected to image analysis. Scale bar, 5 µm. (**D**) The number of red-shifted Keima foci per cell is plotted, where each dot represents the average foci-number per image stack, originating from 7-12 image stacks of iNeuron differentiations. Error bars represent S.D.. (**E**) mtKeima mitophagy flux readout of indicated iNeuron genotypes. Cells were treated with AO for either 1 or 6h and mitophagic flux measured by flow cytometry (>10,000 cells). Pooled data from 2 biological replicates is shown, normalized to 6h BafA treated cells. Error bars depict S.D.. (**F**) Bar graph showing the number of mtDNA puncta/cell with or without treatment with AO (6 h) in control or FBXO7$^{-/-}$ day 12 iNeurons. Error bars depict S.D. from triplicate differentiations. (**G**) Bar graph showing the number of mtDNA puncta/cell with or without treatment with AO (6 h) in HeLa control or FBXO7$^{-/-}$ cells. Error bars depict S.D. from triplicate replicates.

509
Figure 4. Proteomic analysis of HeLa cells lacking FBXO7 during mitophagy. (**A**)
Workflow for analysis of total protein abundance in HeLa cells expressing Parkin with and
without depolarization with AO (16 h). Cell extracts were digested with trypsin prior to 18-
plex TMT labelling and analysis by mass spectrometry. (**B**) Volcano plots [Log_2 FC (16 h AO
/ Fed) versus -Log_{10}(q-value)] for control or one of two FBXO7$^{-/-}$ cell lines with or without
treated with AO (16 h). Red dots, mitochondrial proteins; blue dots, autophagy proteins; grey
dots, remainder of the proteome. (**C**) Violin plots [Log_2 (16 h AO/Fed)] of control or FBXO7$^{-/-}$
cells depicting alterations in the abundance of mitochondrial protein (left panel) or specific

organelles or protein complexes (right panel). (**D**) Correlation plot of c27 and c41 FBXO7$^{-/-}$
clones. Log$_2$FC (16 h AO for each clone relative to control cells) is plotted. (**E**) Correlation
plots [Log$_2$ FC (16 h AO / Fed)] of the proteome of FBXO7$^{-/-}$c27 or c41 clones against
control. Red dots, mitochondrial proteins; blue dots, autophagy proteins; grey dots,
remainder of the proteome. (**F**) PINK1 levels in control cells and in two FBXO7$^{-/-}$ clones were
measured by TMT-proteomics in fed cells (n=3).
524
525

526
Figure 5. Proteomic analysis of human ES cells lacking FBXO7 during neurogenesis.
(A) Workflow for analysis of total protein abundance in ES cells undergoing NGN2-driven
neurogenesis with or without FBXO7. Cell extracts were digested with trypsin prior to 18-
plex TMT labelling and analysis by mass spectrometry. (**B**) Log_2 FC for the indicated
mitochondrial protein-groups in control or FBXO7$^{-/-}$ cells at day 0, 4 or 12 during
neurogenesis is shown in violin plots. (**C**) Log_2 FC for the indicated cellular organelle
proteins in control or FBXO7$^{-/-}$ cells at day 0, 4 or 12 during neurogenesis is shown in violin
plots. (**D**) Volcano plots [Log_2 FC (FBXO7$^{-/-}$/Control) versus -Log_{10}(q-value)] for FBXO7$^{-/-}$
(c52 and c89) and control cells at day 4 (left panel) or day 12 (right panel) of differentiation.
Proteins showing decreased or increased abundance are shown as blue or red dots. (**E**)
Relative abundance of DNM1L (Drp1), MFF, and FIS1 in control or FBXO7$^{-/-}$ cells at day 0, 4
or 12 during neurogenesis. (**F,G**) Mitochondrial morphology in iNeurons (iN) was assessed
using confocal imaging after staining cells with HSP60 to detect mitochondria and DAPI to
identify nuclei, in either fed cells or cells treated with AO (3 h) (left panel). The number of
median # of mitochondria/cell, median mitochondrial length, and median mitochondrial
circularity is shown. Error bars depict S.D. from biological triplicate experiments (9-12 image

stacks each), as shown in panel I left, center, and right, respectively. (**H**) Western Blot
analysis on HeLa control and FBXO7$^{-/-}$ whole cell lysate, probed for the mitochondrial fission
adapter MiD51. (**I**) Relative abundance of DNM1L, MFF, MiD51 and FIS1 in WT (Control) or
FBXO7$^{-/-}$ HeLa cells either in the fed state or after 16 h AO as determined from the
proteomics data in **Figure 5A,B**.
548
549

550
551
Supplemental Figure 1: Tool kit for analysis of HeLa cells and iNeurons lacking
FBXO7. (**A**) Working model summarizing the suggested modes of action of FBXO7 in
PINK1/Parkin mitophagy. (**B**) Targeting of gRNA sequence within FBXO7 gene along with
the sequences of alleles A and B. MiSeq analysis of the three clones used for this study are
shown. (**B**) Schematic for generation of FBXO7$^{-/-}$ in hESC cell lines. (**C**) Western Blot
analysis on HeLa control and FBXO7$^{-/-}$ whole cell lysate. (**D**) Targeting of gRNA sequence
within FBXO7 gene along with the sequences of alleles A and B. MiSeq analysis of the three
HeLa clones (c27, c41, c45) used for this study are shown. (**E**) Western Blot analysis on
hESC control and FBXO7$^{-/-}$ whole cell lysate. Asterisk indicates non-specific band. (**F**)
Targeting of gRNA sequence within FBXO7 gene along with the sequences of alleles A and
B. MiSeq analysis of the two clones (c52, c89) used for this study are shown. (**G**) Karyotype
analysis of FBXO7$^{-/-}$ c52 and c89 hESCs.
564
565

566
567
Supplemental Figure 2. Super-resolution pUb detection in HeLa cells in response to
mitochondrial depolarization. 3D-SIM images of HeLa control, PINK1$^{-/-}$ and FBXO7$^{-/-}$ cell
lines after 1h, 3h or 6h, AO-induced mitophagy (related to **Figure 2A,B**). Cells were stained
for nuclear DNA (DAPI), mitochondria (HSP60) and pUb. (**B**) Zoom-ins of regions of
interested after 1h AO-induced mitophagy are depicted and 3D-surface renderings of insets
are shown on the right. Scale bar = 5 µm or 1 µm.
574
575

576
577
Supplemental Figure 3. Super-resolution pUb detection in iNeurons in response to
mitochondrial depolarization. (**A**) 3D-SIM images of iNeurons day 12 control, PINK1[-/-] and
FBXO7[-/-] cell lines after AO-induced mitophagy. Cells were stained for nuclear DNA (DAPI),
mitochondria (HSP60) and pUb. Related to **Figure 2D**. Scale bar = 5 μm or 1μm. (**B**)
Confocal images of iNeurons day 12 control, PINK1[-/-] and FBXO7[-/-] cell lines after AO-
induced mitophagy with lower AO concentrations. Cells were stained for nuclear DNA
(DAPI), mitochondria (HSP60) and pUb. (**C**) Evaluation of images depicted in **B.** Error bars
depict S.D. from 12 image stacks measured per condition. Scale bar = 10 μm and 5 μm.
586
587

588
589
Supplemental Figure 4. Parkin and autophagy regulator recruitment in FBXO7$^{-/-}$ cells.
(**A**) Overview of methods for analysis of Parkin recruitment to damaged mitochondria. (**B**)
Cells were imaged for GFP-Parkin, nuclei, and mitochondria and segmented to facilitate
analysis of Parkin recruitment to mitochondria in single cells / mitochondrial organelles. (**C**)
Std mitochondrial signal vs Parkin intensity and Std Parkin vs mitochondria with or without
1h AO-induced mitophagy are shown. Data from three replicates, including a total of 238
image stacks, containing 16888 cells and 237848 mitochondrial objects. (**D**) Recruitment of
p62 to mitochondria (not stained) in control or FBXO7$^{-/-}$ HeLa cells with or without treatment
with AO (16 h) was examined by confocal imaging. Scale bar = 10 and 5 μm. (**E,F**)
Quantification of cells in panel D. Assays were performed in biological triplicate with 4 image

stacks taken per repeat. n = 6712 cells. Error bars depict S.D.. (**G**) Quantification of #p62
foci normalized per cell in control, PINK1[-/-] and FBXO7[-/-] cells with or without 3h AO
treatment. Assays were performed in biological triplicate with 4 image stacks taken per
repeat . n = 12646 cells. Error bars depict S.D.. (**H**) Control or FBXO7[-/-] HeLa cells
expressing Parkin were either left untreated or incubated with AO for 1 or 6 h and extracts
subjected to immunoblotting with α-LC3B and α-Actin as a loading control. (**I**) The ratio of
LC3B lipidation (LC3B-II/LC3B-I) was quantified in 3 biological triplicate experiments. (**J**)
Interaction network analysis of PINK1 (left) and FBXO7 (right) based on interaction
proteomics data from our BioPlex Interactome (Huttlin *et al.*, 2021) (see METHODS for
details). (**K**) Summary of major interactions observed for FBXO7 in the context of the
relevant protein complexes. (**L**) Ridgeline plots of mtKeima-shift analysis in HeLa control,
PINK1[-/-] and FBXO7[-/-] cell lines treated with AO at the indicated times. All lines are
normalized to the BafA sample.
613
614

Supplemental Figure 5. Proteomic analysis of HeLa cells lacking FBXO7 during
mitophagy. (**A,B**) Correlation plots (panel A) and PCA analysis (panel B) for ~8000 proteins
quantified by TMT proteomics in individual replicates for the experiment outlined in **Figure**
4A. (**C**) Hierarchical clustering of 8000 proteins quantified in the experiment outlined in
Figure 4A. (**D,E**) Violin plots [Log$_2$ (control / FBXO7$^{-/-}$)] for FBXO7$^{-/-}$ c27 and c41 cells
depicting alterations in the abundance of mitochondrial proteins (left panel) or specific
organelles or protein complexes (right panel). (**F**) Venn diagrams depicting the overlap in the
number of proteins whose levels are reduced (left panel) or increased (right panel) for the

data shown in **Figure 4B** are shown. Proteins whose abundance is significantly reduced
upon AO treatment are enriched in proteins limited with mitochondrial function (right panel),
as expected for cells undergoing mitophagy. (**G**) GO-term enrichment analysis of genes
significantly down in both Control and FBXO7$^{-/-}$ after 16h AO mitophagy. (**H,I**) GO-term
enrichment analysis of unique genes significantly down (panel H) or up (panel I) in FBXO7$^{-/-}$
after 16h AO mitophagy.
631
632

633
634
Supplemental Figure 6. Proteomic analysis of human ES cells lacking FBXO7 during
neurogenesis. (**A**) Hierarchical clustering and correlation plots of control and FBXO7[−/−] c52
(top) or c89 (bottom) at indicated days of differentiation in the experiment outlined in **Figure**
5A. (**B**) PCA analysis of whole cell proteomic samples from control and FBXO7[−/−] cells.
Triplicates for each timepoint and genotype were used. (**C**) FBXO7 peptide abundance in
control and FBXO7[−/−] cells at indicated days of differentiation. (**D**) Violin plots [Log_2 (control /
FBXO7[−/−])] for FBXO7[−/−] c89 cells depicting alterations in the abundance of mitochondrial
proteins.
643
644
645

646
647
Supplemental Figure 7. Comparison of human ES cell neurogenesis with or without
FBXO7. (**A**) Hotelling plots (Log$_{10}$ T^2 statistic versus Log$_2$ FC day 12/day 0) of control (left
panel) or FBXO7$^{-/-}$ c52 hES cells undergoing differentiation. Selected neurogenesis factors
are indicated. (**B**) The relative abundance of selected proteins is shown in the lower
histograms at day 0, 4 and 12 of differentiation. (**C,D**) Hotelling plots similar to **A,B** for
control and FBXO7$^{-/-}$ c89. (**E**) Patterns of changes in the abundance of autophagy proteins in
control (left panel) and FBXO7$^{-/-}$ (right panel) day 12 iNeurons relative to day 0 cells
displayed on a Hotelling plot. (**F**) Cluster analysis of control and FBXO7$^{-/-}$ cells comparing
day 12 of differentiation to day 0. Individual clusters are displayed on Hotelling plots for each
genotype.
658

EXPERIMENTAL MATERIALS AND METHODS

All details and catalogue numbers can be found in the **Materials Table (Supplemental Table S9).** Protocols associated with this work can be found on protocols.io (dx.doi.org/10.17504/protocols.io.kxygx99pwg8j/v1)

Cloning and plasmid generation

The construction of lentiviral expression constructs used in this study have been previously described: pHAGE-mt-mKeima in (Heo *et al.*, 2015), PB-mt-mKeimaXL (Ordureau *et al.*, 2021) and pHAGE-GFP-Parkin in (Ordureau *et al.*, 2018).

Cell culture, generation of lentiviral stable cell lines.

HeLa Flip-In T-Rex (HFT) expressing doxycycline-inducible WT-Parkin (HeLa HFT/TO-PRKN-WT) cells (Heo *et al.*, 2015) and HEK293T were maintained in Dulbecco's modified Eagles medium (DMEM), supplemented with 10% vol/vol fetal bovine serum (FBS), 5% vol/vol penicillin-streptomycin (P/S), 5% vol/vol GlutaMAX and 5% vol/vol Non-essential amino acids (NEAA) at 37°C, 5% O_2.

   Stable cell lines were generated using lentivirus generated from HEK293T cells. pHAGE-mt-mKeima or pHAGE-eGFP-Parkin were co-transfected in HEK293T cells together with the lentiviral vectors pSpax2 and pMD2.1 using Lipofectaime LTX reagent (Thermo Fisher Scientific, 15338100), according to manufacturer's instructions. Virus was harvested, filtered (0.45 µm) and added 8µg/ml polybrene and target cells infected for 18 h. Antibiotic selection was performed using 1 µg/ml puromycin or FACS sorting and cells verified using immunoblotting or fluorescence microscopy.

If not stated otherwise, Parkin expression in HeLa HFT/TO-PRKN was induced using 2 µg/ml doxycycline o/n, and subsequent mitophagy induced using Antimycin A / Oligomycin A (==AO) for the indicated times.

   H9 ES cells harboring the mitochondrial mt-mKeima flux reporter (Ordureau *et al.*, 2020) were generated by electroporation of $1x10^6$ cells with 2.5 µg of pAC150-PiggyBac-matrix-mKeima^{XL} along with 2.5 µg of pCMV-HypBAC-PiggyBac-Helper, as described (Ordureau *et al.*, 2020). The cells were selected and maintained in E8 medium supplemented with 50 µg/ml Hygromycin and Hygromycin was kept in the medium during differentiation to iNeurons.

Gene-Editing and iNeuron differentiation

Generation of HeLa HFT cells lacking FBXO7$^{-/-}$ was facilitated using CRISPR/Cas9 with target sites determined using CHOPCHOP (Labun *et al*, 2019). Guide RNAs were ligated into the px459 plasmid (Addgene plasmid # 62988) and cells transfected using Lipofectaime LTX reagent (Thermo Fisher Scientific, 15338100), according to manufacturer's instructions. Two days post-transfection, single GFP positive cells were sorted into 96-well dishes containing 300µl full growth medium (composition as described above). Single cells were allowed to grow into colonies, then duplicated for MiSeq analysis and maintenance. Knockout candidates were confirmed by Western blot on whole cell lysates. The sgRNAs were generated using *GeneArt Precision gRNA Synthesis Kit* (Thermo Fisher Scientific) according to the manufacturer's instruction and purified using RNeasy Mini Kit (Qiagen). The sgRNA target sequence: ACCGATTCACTACAGAGCAT.

   The PINK1$^{-/-}$ H9 cells used here, wherein sequences in exon 1 were deleted using CRISPR-Cas9 to create a null allele, were reported previously (Ordureau *et al.*, 2018). To generate FBXO7$^{-/-}$ H9 ES or HeLa cells, 0.6 µg sgRNA was incubated with 3 µg SpCas9 protein for 10 minutes at room temperature and electroporated into $2x10^5$ WT H9 cells using Neon transfection system (Thermo Fisher Scientific). Mutants were identified by Illumina MiSeq and further confirmed by Western blotting. For introduction of TRE3G-NGN2 into the AAVS1 site, a donor plasmid pAAVS1-TRE3G-NGN2 was generated by replacing the EGFP sequence with N-terminal flag-tagged human NGN2 cDNA sequence in plasmid pAAVS1-TRE3G-EGFP (Addgene plasmid # 52343). Five micrograms of pAAVS1-TRE3G-NGN2, 2.5 µg hCas9 (Addgene plasmid # 41815), and 2.5 µg gRNA_AAVS1-T2 (Addgene plasmid #

41818) were electroporated into 1×10^6 H9 cells. The cells were treated with 0.25 µg/ml Puromycin for 7 days and surviving colonies were expanded and subjected to genotyping.

Human ES cells (H9, WiCell Institute) were cultured in E8 medium (Chen *et al*, 2011) on Matrigel-coated tissue culture plates with daily medium change. Cells were passaged every 4-5 days with 0.5 mM EDTA in 1× DPBS (Thermo Fisher Scientific). SpCas9 and AsCas12a/AsCpf1 expression plasmids pET-NLS-Cas9-6xHis (Addgene plasmid # 62934) and modified pDEST-his-AsCpf1-EC, generated by deleting the MBP sequence from plasmid pDEST-hisMBP-AsCpf1-EC (Addgene plasmid # 79007), were transformed into *Rosetta™(DE3)pLysS* Competent Cells (Novagen), respectively, for expression. SpCas9 and AsCas12a/AsCpf1 proteins were purified as described elsewhere (Hur *et al*, 2016; Zuris *et al*, 2015). Briefly, cells expressing SpCas9 [0.5 mM isopropylthio-β-galactoside, 14-hour induction] were lysed in FastBreak buffer (Promega, Inc) and the NaCl concentration adjusted to 500 mM. Extracts were centrifuged at 38,000g for 10 min at 4°C and the supernatant incubated with Ni-NTA resin for 1 hour. The resin was washed extensively with 50 mM Tris pH8.0, 500 mM NaCl, 10% glycerol, 20 mM imidazole, and 2 mM TCEP prior to elution with this buffer supplemented with 400 mM imidazole. Proteins were diluted two volumes/volume in PBS and fractionated on a Heparin-Sepharose column using a 0.1 to 1.0 M NaCl gradient. Cas9-containing fractions were stored in PBS, 20% glycerol, 2 mM TCEP at -80°C. AsCpf1 expression was induced similarly, and cells pelleted by centrifugation. Cells were lysed by sonication in 50 mM HEPES pH7, 200 mM NaCl, 5 mM $MgCl_2$, 1 mM DTT, and 10 mM imidazole supplemented with lysozyme (1 mg/ml) and protease inhibitors (Roche complete, EDTA-free). After centrifugation (16,000g for 30 min), the supernatant was incubated with Ni-NTA resin, the resin washed with 2M NaCl, and bound proteins eluted with 250 mM imidazole, and buffer exchanged into lysis buffer lacking $MgCl_2$ and imidazole prior to storage at -80°C.

For human ES cell conversion to iNeurons, cells were expanded and plated at $2 \times 10^4/cm^2$ on Matrigel-coated tissue plates in DMEM/F12 supplemented with 1x N2, 1x NEAA (Thermo Fisher Scientific), human Brain-derived neurotrophic factor (BDNF, 10 ng/ml, PeproTech), human Neurotrophin-3 (NT-3, 10 ng/ml, PeproTech), mouse laminin (0.2 µg/ml, Cultrex), Y-27632 (10 µM, PeproTech) and Doxycycline (2 µg/ml, Alfa Aesar) on Day 0. On Day 1, Y-27632 was withdrawn. On Day 2, medium was replaced with Neurobasal medium supplemented with 1x B27 and 1x Glutamax (Thermo Fisher Scientific) containing BDNF, NT-3 and 1 µg/ml Doxycycline. Starting on Day 4, half of the medium was replaced every other day thereafter. On Day 7, the cells were treated with Accutase (Thermo Fisher Scientific) and plated at $3-4 \times 10^4/cm^2$ on Matrigel-coated tissue plates. Doxycycline was withdrawn on Day 10.

Immunoblotting
At the indicated times, ES cells, iNeurons or HeLa cells were washed on ice in 1x PBS, harvested and pellet wash with 1x PBS and resuspended in 8 M urea buffer (8 M urea, 150 mM TRIS pH 7.4, 50 mM NaCl, PhosSTOP phosphatase inhibitor cocktail). Resuspended cell lysates were sonicated for 10 seconds and debris pelleted at 13,000 rpm for 10 min. Protein concentration was determined by BCA assay according to manufacturer's instructions (Thermo Fisher Scientific, 23227). Indicated amounts of proteins were resuspended in 1xLDS + 100 mM DTT and boiled for 10 minutes at 85°C. Equal amount of protein and volume were loaded run on 4%-20% Bis-Tris, 8% Tris NuPAGE gels for 5 minutes at 100V, 5 min at 150 V and then run at 200 V for the required time. Gels were transferred via wet transfer system onto PDVF membranes for immunoblotting. Chemiluminescence and colorimetric images were acquired using a BioRad ChemiDoc MP imaging system. Images from Western Blots were exported and analysed using Image Lab and ImageJ/FiJi (Schindelin *et al*, 2012).

PROTEOMICS
Proteomics – general sample preparation
Sample preparation of proteomic analysis of whole-cell extract from HeLa, hESC, NPC and iNeurons was performed according to previously published studies (Ordureau *et al.*, 2021; Ordureau *et al.*, 2020). Cells were harvested on ice and plates were washed twice with 1x PBS and detached in 1x PBS using cell scraper. After pelleting at 2000 rpm for 5 min at 4°C, cells were washed 2x with 1x PBS and resuspended in 8 M urea buffer (composition stated above). After sonification for 10 seconds, resuspended cells were pelleted for 10 min at 13000 rpm. Protein concentration was determined using BCA kit (Thermo Fisher Scientific, 23227).

Unless otherwise noted, proteomics and data analysis was performed as described (Ordureau *et al.*, 2021; Ordureau *et al.*, 2020). Briefly, protein extracts (100 μg) were subjected to disulfide bond reduction with 5 mM TCEP (room temperature, 10 min) and alkylation with 25 mM chloroacetamide (room temperature, 20 min). Methanol–chloroform precipitation was performed prior to protease digestion. In brief, four parts of neat methanol were added to each sample and vortexed, one part chloroform was then added to the sample and vortexed, and finally three parts water was added to the sample and vortexed. The sample was centrifuged at 6,000 rpm for 2 min at room temperature and subsequently washed twice with 100% methanol. Samples were resuspended in 100 mM EPPS pH8.5 containing 0.1% RapiGest and digested at 37 °C for 8h with Trypsin at a 100:1 protein-to-protease ratio. Samples were acidified with 1% Formic Acid for 15 min and subjected to C18 solid-phase extraction (SPE) (Sep-Pak, Waters).

Proteomics – quantitative proteomics using TMT
Tandem mass tag labeling of each sample (50 mg peptide input) was performed by adding 5 uL of the 25 ng/mL stock of TMTpro reagent along with acetonitrile to achieve a final acetonitrile concentration of approximately 30% (v/v). Following incubation at room temperature for 1 h, the reaction was quenched with hydroxylamine to a final concentration of 0.5% (v/v) for 15 min. The TMTpro- labeled samples were pooled together at a 1:1 ratio. The sample was vacuum centrifuged to near dryness, and following reconstitution in 1% FA, samples were desalted using C18 solid-phase extraction (SPE) (50 mg, Sep-Pak, Waters), according to manufacturer protocol.

Dried TMTpro-labeled peptides (~300 ug) were resuspended in 10 mM NH_4HCO_3 pH 8.0 and fractionated using basic pH reverse phase HPLC (Wang *et al*, 2011). Briefly, samples were offline fractionated over a 90 min run, into 96 fractions by high pH reverse-phase HPLC (Agilent LC1260) through an Aeris peptide xb-c18 column (Phenomenex; 250 mm x 3.6 mm) with mobile phase A containing 5 % acetonitrile and 10 mM NH_4HCO_3 in LC-MS grade H_2O, and mobile phase B containing 90 % acetonitrile and 10 mM NH_4HCO_3 in LC-MS grade H_2O (both pH 8.0). The 96 resulting fractions were then pooled in a non-continuous manner into 24 fractions (as outlined in Figure S5 of (Paulo *et al*, 2016)) and sets of 12 fractions (even or odd numbers) were used for subsequent mass spectrometry analysis. Fractions were vacuum centrifuged to near dryness. Each consolidated fraction was desalted via StageTip, dried again via vacuum centrifugation, and reconstituted in 5 % acetonitrile, 1 % formic acid for LC-MS/MS processing.

TMT proteomics samples were subjected to analysis using a Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Fisher Scientific, San Jose, CA) online with Proxeon EASY-nLC1200 liquid chromatography (Thermo Scientific). Peptides were resuspended in 5 % ACN/5 % FA and 10 % of the samples were loaded on a 35 cm analytical column (100mm inner diameter) packed in-house with Accurcore150 resin (150 Å, 2.6 mm, Thermo Fisher Scientific, San Jose, CA) for LC-MS analysis. Peptide separation was performed with a gradient of acetonitrile (ACN, 0.1% FA) from 3-13 % (0-83 min) and 13-28 % (80-83 min) during a 90 min run.

For the HeLa TMTpro proteomic samples, LC-hrMS/MS was combined with 3
optimized CV parameters on the FAIMS Pro Interface to reduced precursor ion interference
(Schweppe *et al*, 2019). Data-dependent acquisition (DDA) was performed by selecting the
most abundant precursors from each CV's (-40/-60/-80) MS^1 scans for hrMS/MS over a 1-
1.5s duty cycle (1s/1.5s/1s respectively). The MS^1 scan parameters include a 400-1,600 m/z
mass range at 60,000 resolution (at 200 Th) with 4 x 10^5 automated gain control (AGC) (100
%), and a maximum injection time (max IT) of 50 ms. Precursors (z=2-5) were isolated with
0.7 Th (quadrupole), fragmented with high energy C-trap dissociation (HCD) at 36
normalized collision energy (NCE), and subjected to hrMS/MS on the Orbitrap at 50,000
resolution (at 200 Th), 120 ms max IT, fixed first mass 110 Th, and 1.0 x 10^5 AGC (200%).
Precursors were placed on 90 s dynamic exclusion (+/- 10 ppm) to prevent redundant
sampling.
In the iNeuron TMTpro experiments, the same FAIMS and MS^1 parameters were
implemented (with 1.25 s duty cycle/CV for DDA) with the Multi-Notch SPS-MS3 acquisition
method (McAlister *et al*, 2014), to further reduce ion interference in TMT reporter
quantification (Paulo *et al.*, 2016). Most abundant precursors (with 120 s dynamic exclusion
+/- 10 ppm) were selected from MS^1 scans, isolated using the quadrupole (0.6 Th isolation),
fragmented with collision-induced dissociation (CID) at 35 NCE, and subjected to MS/MS in
the ion trap (turbo scan speed, 35 ms max IT, 1.0 x 10^4 AGC). Using Real Time Search
analysis software (Erickson *et al*, 2019; Schweppe *et al*, 2020), a synchronous-precursor-
selection (SPS) API-MS^3 scan was collected on the top 10 most intense b- or y-ions from the
matched peptide identification (determined by an online search of its respective MS/MS
scan). MS^3 scans were performed on the Orbitrap (AGC 2.0 x 10^5; NCE 55; max IT 250 ms,
50,000 resolution at 200 Th). To increase quantitative sampling (MS^3 scans) of proteins
during each mass spectrometry injection, a 2 peptide per protein per sample closeout was
set. This ensures no more than two peptide-spectrum matches per protein (that pass quality
filters) are subjected to MS^3 scans, reducing redundant protein MS sampling and potentially
increasing proteome depth (Schweppe *et al.*, 2020).
851
Proteomics Data Analysis
Mass spectrometry raw data were converted to mzXML and monoisotopic peaks were
reassigned using Monocle (Rad *et al*, 2021). Mass spectra were database searched using
Sequest algorithm (2019.01 rev. 5; (Eng *et al*, 1994)) against the Human Reference
Proteome Uniprot database (2019-01 SwissProt entries only; UniProt Constortium, 2015)
appended with sequences of common contaminates and reverse sequences of proteins as
decoys, for target-decoy competition (Elias & Gygi, 2007). Sequest search parameters
include: 50 ppm precursor tolerance, 0.9 Da product ion tolerance, trypsin endopeptidase
specificity (C-terminal to [KR], 2 max missed cleavages), static modifications on peptide N-
terminus and lysines with TMTpro tags (+304.207 Da) and carbamidomethylation on
cysteines (57.021 Da), and variable modification of oxidation on methionines (+15.995 Da).
Peptide-spectrum matches were filtered at 2% false discovery rate (FDR) using linear
discriminant analysis (Huttlin *et al*, 2010), based on XCorr, DeltaCn, missed cleavages,
peptide length, precursor mass accurracy, fraction of matched product ions, charge state,
and number of modifications per peptide (additionally restricting PSM Xcorr >1 and peptide
length>6). Following a 2% protein FDR filter (Savitski *et al*, 2015), PSMs reporter ion
intensities were quantified (most intense centroid within 0.003 Da of theoretical TMT reporter
mass), filtered based on a precursor isolation specificity > 0.5, and filtered by a summed
signal-to-noise across TMT channels > 100.
Protein quantification was performed via the summation (weighted average) of its
constituent PSMs' reporter intensities and TMT channels were normalized for protein input
to total TMT channel intensity across all quantified PSMs (adjusted to median total TMT
intensity for the TMT channels) (Plubell *et al*, 2017). Log$_2$ normalized summed protein
reporter intensities were compared using a Student's t-test and p-values were corrected for
multiple hypotheses using the Benjamini-Hochberg adjustment (Benjamini & Hochberg,
1995). Resultant q-values and mean log$_2$ fold changes between conditions were used to

generate volcano plots. Hotelling T^2 analysis was performed using the normalized summed
protein reporter ion intensities and the timecourse (v. 1.66.0; (Tai, 2022)) package in R.
The annotation list for the subcellular localization of organellar protein markers was
derived from previously published high confidence HeLa dataset ((Itzhak *et al*, 2016); "high"
& "very high" confidence) and additional manual entries (Ordureau *et al.*, 2021). MitoCharta
3.0 (Rath *et al*, 2021) was used for mitochondrial annotation, whereas annotations for
ribosome and autophagy components were used from previous studies (An *et al*, 2020).
Transcription factor and neuronal development markers were based on previously published
databases and publications (Lambert *et al*, 2018; Ordureau *et al.*, 2021). Figures were
generated using a combination of Excel, R (v.4.2.0) in RStudio (2022.07.01 Build 554),
Perseus (v1.6.5 (Tyanova & Cox, 2018)), GraphPad Prism (v9.1.0), and Adobe Illustrator
(26.2.1).
890
891 The **Supplemental Data Table S3-8** contains the quantified proteins as well as associated
892 TMT reporter ratio to control channels used for quantitative analysis.
893
Interaction Proteomic Analysis
For interaction analysis of FBXO7 and PINK1, interaction proteomics data from the Bioplex
Interactome (Huttlin *et al.*, 2021) was extracted for HEK293T and HCT116 samples
(**Supplemental Table S1,2**). The number of replicates in which a peptide-spectral match
(PSM) mapped to PINK1's or FBXO7's the Bioplex 3.0 interactomes was plotted via a
network diagram in R. (packages to add from R include (igraph 1.3.1, tidygraph 1.2.1, and
ggraph 2.0.5).
901
902 *MICROSCOPY*
903 Macros and pipelines used in this work can be found on GitHub
904 (https://github.com/harperlaboratory/FBXO7.git) or
905 Zenodo (https://doi.org/10.5281/zenodo.7258918).
906
Live-cell confocal microscopy for mitophagic flux analysis over differentiation (mt-mKeima[XL])
For quantitative Keima-flux analysis (Ordureau *et al.*, 2020), hESC were seeded on day 4 of
differentiation into 6-well 1.5 high performance glass bottom plates (Cellvis, P06-1.5H-N)
and further differentiated in the vessel for the indicated times until reaching appropriate
confluency and imaged. iNeurons were imaged using a Yokogawa CSU-W1 spinning disk
confocal on a Nikon Eclipse Ti-E motorized microscope. The system is equipped with a
Tokai Hit stage top incubator and imaging was performed at 37°C, 5% CO_2 and 95%
humidity under a Nikon Plan Apo 60×/1.40 N.A immersion oil objective lens. For ratiometic
imaging, mtKeimaXL were excited in sequential manner with a Nikon LUN-F XL solid state
laser combiner ([laser line – laser power]: 445 - 80mW, 561 - 65mW]) using a
Semrock Di01-T445/515/561 dichroic mirror. Fluorescence emissions were collected
through a Chroma ET605/52m [for 445 nm] and a 568 Chroma ET605/52m [for 561 nm],
filters, respectively (Chroma Technologies). Confocal images were acquired with a
Hamamatsu ORCA-Fusion BT CMOS camera (6.5 μm^2 photodiode, 16-bit) camera and NIS-
Elements image acquisition software. Consistent laser intensity and exposure time were
applied to all the samples, and brightness and contrast were adjusted equally by applying
the same minimum and maximum display values in ImageJ/FiJi (Schindelin *et al.*, 2012).
Image quantification was performed in ImageJ/FiJi using custom-written batch-macros.
In brief, raw confocal images of mitochondrial targeted mt-mKeima[XL] were divided
[ex:561/ex:445] resulting in a ratiometic image of only acidic Keima-puncta. These signals
were subjected to background subtraction (rolling kernel size 25, sliding paraboloid) and
converted to binary objects. The "Analyze Particles..." command (pixel size exclusion: 0.5-∞,
exclude edge objects) was used to measure foci-abundance and other morphological
parameters. Results for each image-stack saved as .csv files, together with the original
ratiometic .tiff file for QC purposes. Unless stated otherwise all images represent z-

projections. Statistical analysis and plotting of microscopy data was performed in Prism
(v9.1.0, GraphPad).

934
935
936
937
Immunocytochemical analysis
iNeurons or HeLa cells were fixed with warm 4% paraformaldehyde (Electron Microscopy
Science, #15710, purified, EM grade) in PBS at 37°C for 30 min and permeabilized with
0.5% Triton X-100 in PBS for 15 minutes at room temperature. After three washes with
0.02% Tween20 in PBS (PBST), cells were blocked for 10 min in 3% BSA-1xPBS at room
temperature and washed again three times in PBST. Cells were incubated for 3h in primary
antibodies in 3% BSA-1xPBS and washed three times with PBST. Secondary antibodies
(Thermo Scientific, 1:400 in 3% BSA-1xPBS) where applied for 1h at room temperature.
Alexa Fluor 633 Phalloidin (Thermo Fisher, 1:200, A22284) was added with secondary
antibodies to label F-actin. To stain nuclei, Hoechst33342 (1:10000) was added for 5 min to
cells in PBST and final three washes performed before mounting in Vectashield (Vector
Laboratories, H-1000-10). Primary and secondary antibodies used in this study can be found
in the **Materials Table (Supplemental Table S9).**

951
Fixed-cell microscopy – general acquisition parameters
Immunofluorescently labelled Hela or iNeurons (antibodies indicated in figures and figure
legends and details in **Materials Table, (Supplemental Table S9)**) were imaged at room
temperature using a Yokogawa CSU-W1 spinning disk confocal on a Nikon Eclipse Ti-E
motorized microscope equipped with a Nikon Plan Apochromat 40×/0.40 N.A air-objective
lens, Nikon Apochromat 60×/1.42 N.A oil-objective lens and a Plan Apochromat 100×/1.45
N.A oil-objective lens. Signals of 405/488/568/647 fluorophores were excited in sequential
manner with a Nikon LUN-F XL solid state laser combiner ([laser line – laser power]: 405 -
80mW, 488 - 80mW, 561 - 65mW, 640nm - 60mW]) using a Semrock Di01-
T405/488/568/647 dichroic mirror. Fluorescence emissions were collected with
Chroma ET455/50m [405 nm], 488 Chroma ET525/50m [488 nm], 568 Chroma ET605/52m
[561 nm], 633 Chroma ET705/72m [640 nm] filters, respectively (Chroma Technologies).
Confocal images were acquired with a Hamamatsu ORCA-Fusion BT CMOS camera (6.5
μm^2 photodiode, 16-bit) camera and NIS-Elements image acquisition software. Consistent
laser intensity and exposure time were applied to all the samples, and brightness and
contrast were adjusted equally by applying the same minimum and maximum display values
in ImageJ/FiJi (Schindelin *et al.*, 2012).

969
Microscopy-based mitochondrial morphology measurements in iNeurons
For quantitative measurement of mitochondrial morphology in d12 iNeurons under fed or
mitophagy conditions, HeLa cells were seeded on 6-well 1.5 high performance glass bottom
plates (Cellvis, P06-1.5H-N) and mitophagy induced for the indicated time durations.
Alternatively, hESC were seeded on day 4 of differentiation into 6-well 1.5 high performance
glass bottom plates (Cellvis, P06-1.5H-N) and differentiated in the vessel into iNeurons and
mitophagy induced for the indicated time durations. Cells were fixed and stained as
described above. Z-stacks were acquired with a Nikon Plan Apo 100×/1.45 N.A oil-objective
lens and with the parameters stated above. Image quantification was performed in
ImageJ/FiJi using custom-written batch-macros.
Statistical analysis and plotting of microscopy data was performed in Prism (v9.1.0,
GraphPad). Primary and secondary antibodies used in this study can be found in the
Materials Table (Supplemental Table 9).

983
984 *Microscopy-based mtDNA turnover measurements in HeLa and iNeurons*
985 For quantitative measurement of mtDNA turnover after AO-induced mitophagy, HeLa cells
986 were seeded on 6-well 1.5 high performance glass bottom plates (Cellvis, P06-1.5H-N) and

987	mitophagy induced for the indicated time durations. Alternatively, hESC were seeded on day
988	4 of differentiation into 6-well 1.5 high performance glass bottom plates (Cellvis, P06-1.5H-
989	N) and differentiated in the vessel into iNeurons and mitophagy induced for the indicated
990	time durations. Cells were fixed and stained as described above, but primary antibody
991	incubation with aDNA (1:200) was performed overnight. Z-stacks were acquired with a Nikon
992	Plan Apo 100×/1.45 N.A oil-objective lens and with the parameters stated above. Image
993	quantification was performed in ImageJ/FiJi using custom-written batch-macros. In brief,
994	both aDNA and nuclear signals were converted to binary files and the nuclear signal
995	subtracted from the aDNA signal, resulting in an image stack containing only the mtDNA
996	intensities. The "Analyze Particles..." command (pixel size exclusion: 0.05-3, exclude edge
997	objects) was used to measure morphological features and results for each image-stack
998	saved as .csv files, together with the analyzed binary-mask overlay .tiff file for QC purposes.
999	Number of mtDNA signals were normalized to cell number found in the same image stack.
1000	Statistical analysis and plotting of microscopy data was performed in Prism (v9.1.0,
1001	GraphPad). Primary and secondary antibodies used in this study can be found in the
1002	**Materials Table (Supplemental Table S9)**.

Microscopy-based measurements of p62 recruitment in HeLa

For quantitative measurement of p62 recruitment to mitochondria, HeLa control and knockout cells were seed in 1.5 high performance glass bottom plates and doxycycline added over night to induce Parkin expression. Mitophagy was induced using Oligomycin A / Antimycin A for 16h in full DMEM in presence of doxycycline. Cells were fixed and stained as stated above and imaged using a Nikon Plan Apo 60×/1.42 N.A air-objective lens and with the parameters stated above. 8 µM z-stacks were taken for each selected field of view.

Image quantification was performed in ImageJ/FiJi using custom-written batch-macros. The p62 channel was filtered (Gaussian Blue, sigma =2) and converted into binary files using the Intermodes thresholding method. p62 spots were counted using the "Analyze Particles..." function (pizel size exclusion: 0.1-30, exclude edge objects) and results for each image-stack saved as .csv files, together with the analyzed binary-mask overlay .tiff file for QC purposes. The DAPI channel was used to count nuclei for per cell normalization. Statistical analysis and plotting of microscopy data was performed in Prism (v9.1.0, GraphPad). Primary and secondary antibodies used in this study can be found in the **Materials Table (Supplemental Table S9).**

Microscopy-based pUb-coverage measurements of mitochondria in iNeurons

For quantitative measurement of pUb coverage over mitochondria AO-induced mitophagy, hESC were seeded on day 4 of differentiation into 6-well 1.5 high performance glass bottom plates (Cellvis, P06-1.5H-N) and differentiated in the vessel into iNeurons and mitophagy induced for the indicated time durations. Cells were fixed and stained as described above, using anti-HSP60 to label mitochondria and anti-pUb (Ser65) to label pUb. 3-5 10 µm thick z-stacks per replicate per sample were acquired using a Nikon Plan Apo 60×/1.42 N.A air-objective lens and with the parameters stated above.

Image quantification was performed in ImageJ/FiJi using custom-written batch-macros. In brief, mitochondrial signal was filtered (Gaussian Blur, sigma=2), converted into binary files and holes in the resulting mask filled. pUb channel was thresholded into a binary file (Triangle method); these masks were measured using the "Analyze Particles..." command (pixel size exclusion: 0.5-∞, exclude edge objects) and results for each image-stack saved as .csv files, together with the analyzed binary-mask overlay .tiff file for QC purposes. % of mitochondrial pUb coverage was calculated and normalized to [t]=6h AO. Statistical analysis and plotting of microscopy data was performed in Prism (v9.1.0, GraphPad). Primary and secondary antibodies used in this study can be found in the **Materials Table (Supplemental Table S9).**

Microscopy-based evaluation of Parkin translocation and mitophagy in FBXO7[-/-] cell lines

For quantitative measurement of Parkin translocation kinetics in FBXO7$^{-/-}$ cell lines, stable
HeLa cell lines expressing GFP-Parkin were created. WT and knockout cell lines were
seeded into 12-well 1.5 high performance glass bottom plates (Cellvis, P12-1.5H-N) two
days prior to experimental manipulation. Mitophagy was induced as described above using
Antimycin A / Oligomycin A for 1h. Treated and control (fed) cells were fixed as described
above, stained for mitochondria (HSP60), Parkin (GFP) and DNA (SpyDNA-555) and 8µm z-
stacks were acquired with a Nikon Plan Apo 60×/1.42 N.A oil-objective lens and with the
parameters stated above. For quantitate single-cell analysis, a total of ~240 randomly xy-
marked stacks were acquired, and MIPs used for subsequent analysis. CellProfiler (Stirling
et al., 2021) was used for the quantitative analysis of single-cells and mitochondrial objects
(segmentation and analysis pipelines can be found on GitHub). Plotting of microscopy data
was performed in Prism (v9.1.0, GraphPad) and R using the following libraries: tidyverse,
dplyr, tibble, viridis, ggplot2, ggridges, ggsci. Primary and secondary antibodies used in this
study can be found in the **Materials Table (Supplemental Table S9)**.
1055
Immunocytochemical sample preparation for 3D-SIM
Sample preparation and SIM acquisition guidelines from (Kraus *et al*, 2017) were used as a
basis for the super-resolution analysis of pUb spreading after mitophagy induction. iNeurons
or HeLa cells were seeded on 18x18 mm Marienfeld Precision cover glasses thickness No.
1.5H (tol. ± 5 µm). Cover glasses were coated, if necessary, for hESC / iNeuron culture as
described above. After experimental manipulation, cells were fixed with warm
paraformaldehyde 3% Glutaraldehyde 0.35% in 0.1M Sodium Cacodylate, pH 7.4 (Electron
Microscopy Science) at 37°C for 30 min and permeabilized with 0.5% Triton X-100 in PBS
for 15 minutes at room temperature. After three washes with 0.02% Tween20 in PBS
(PBST), cells were blocked for 10 min in 3% BSA-1xPBS at room temperature and washed
again three times in PBST. iNeurons were incubated overnight in primary antibodies in 3%
BSA-1xPBS and washed three times with PBST. Secondary antibodies (Thermo Scientific,
1:400 in 3% BSA-1xPBS) where applied for 1 h at room temperature. To stain nuclei, DAPI
was added for 5 min to cells in PBST, washed three times for 5 min in 1xPBST before a
post-fixation in 4% paraformaldehyde was performed. After 2 washes in PBST, coverslips
were washed once in 1xPBS and mounted in Vectashield (Vector Laboratories, H-1000-10)
on glass slides. Primary and secondary antibodies used in this study can be found in the
Materials Table (Supplemental Table S9).
1074
3D-SIM microscopy – acquisition parameters
3D-SIM microscopy was performed on a DeltaVision OMX v4 using an Olympus 60x / 1.42
Plan Apo oil objective (Olympus, Japan). The instrument is equipped with 405 nm, 445 nm,
488 nm, 514 nm, 568 nm and 642 nm laser lines (all >= 100 mW) and images were recorded
on a front-illuminated sCMOS (PCO Photonics, USA) in 512x512px image size mode, 1x
binning, 125 nm z-stepping and with 15 raw images taken per z-plane (5 phase-shifts, 3
angles). Raw image data was computationally reconstructed using CUDA-accelerated 3D-
SIM reconstruction code (https://github.com/scopetools/cudasirecon) based on (Gustafsson
et al, 2008). Optimal optical transfer function (OTF) was determined via an in-house build
software, developed by Talley Lambert from the NIC / CBMF (GitHub:
https://github.com/tlambert03/otfsearch, all channels were registered to the 528nm output
channel, Wiener filter: 0.002, background: 90).
1087
3D-SIM microscopy – pUb – mitochondria 3D renderings & analysis
3D renderings of 3D-SIM images were performed in UCSF Chimera X
(https://www.cgl.ucsf.edu/chimerax/) using 32-bit .tiff image stacks. Image channels were
sequentially imported and visualized as surfaces using the "Volume Viewer" tool. Surface
representations were cleaned up using the "High dust" command, based on size filtering
thresholding.
3D-analysis of HeLa and iNeuron datasets from 3D-SIM datasets was performed
using Imaris (Oxford Instruments, v9.7). After converting all multi-color .tiff to native .ims

files, import into Imaris Arena and global background subtraction, mitochondrial and pUb objects were segmented from seeds (XY starting diameter: 0.08 µm == pixel size of images), segmented based on automatic thresholding with local background subtraction and splitting of touching objects (0.4µm). Objects were piped into Imaris Vantage module for further analysis. In Vantage, nearest neighbour distances of pUb to pUb and between pUb and mitochondria, as well as volume of segmented objects were computed. This pipeline was tested on WT control cells and then applied for batch processing on all other genotype to allow for unbiased segmentation and analysis. On average, 10 cells were analysed per genotype per condition (121 total) for HeLa cells, and 12 cells (139 total) for iNeurons.

Unless stated otherwise, all images depicted in figures are maximum-intensity projections.

Flow cytometry-based measurement of mitophagic flux

HeLa HFT/TO-PRKN cells of the indicated genotype (control, PINK1$^{-/-,}$ FBXO7 $^{-/-}$) expressing mtKeima (Heo *et al.*, 2015) were seeded into 12-well dishes. Upon reaching 60% confluency, 2 µg/ml doxycycline was added to induce Parkin expression for at least eight hours before depolarization of mitochondria. Following Parkin induction, cells were treated with Antimycin A (0.5 µM) and Oligomycin (0.5 µM) for the indicated number of hours. Bafilomycin A (25 nM, Sigma-Aldrich, B1793)-treated samples were treated for three hours. Control cells were fed three hours before experimental manipulation. To harvest cells for analysis, each well was washed with 1 mL PBS then treated with 100 µL of 0.25% trypsin for 3 min at room temperature, resuspended in 300 µL of DMEM with 10% FBS. 200 µL of each sample was transferred to a flat-bottom 96-well plate for analysis by flow cytometry.

hESCs were seeded at day 4 into coated 6-well dishes and differentiated. At day 12, mitophagy was induced with Antimycin A (0.5 µM) and Oligomycin (0.5 µM) or Bafilomycin A (25 nM, Sigma-Aldrich, B1793) for the indicated number of hours. Control cells were fed 2h before experimental manipulation. Cells were dissociated from the wells using Accutase and resuspended in 600 µL ND2 medium + 10 µM Y27632 and filtered through a cell strainer cap tube (Corning, 352235).

Analysis of a population of at least 10,000 cells was performed on an Attune NxT (Thermo Fisher Scientific) detecting neutral mtKeima signal with excitation at 445 nm and emission 603 nm with a 48 nm bandpass and acidic mtKeima with 561 nm excitation and emission 620 nm and a 15 nm band pass. mtKeima ratio was analysed as previously described (An & Harper, 2018). Briefly, acidic:neutral Keima ratios were measured by gating a population of well-behaved single cells using forward and side scatter, followed by calculation of acidic:neutral mtKeima ratio on a per-cell basis in FlowJo Software (FlowJo, LLC).

QUANTIFICATION AND STATISTICAL ANALYSIS

Unless stated otherwise all quantitative experiments were performed in triplicate and average with S.E.M. or S.D. as indicated in legends reported.

REFERENCES

Al Rawi S, Simpson L, McDonald NQ, Chernuha V, Elpeleg O, Zeviani M, Barker RA, Spiegel R, Laman H (2022) Study of an FBXO7 patient mutation reveals Fbxo7 and PI31 co-regulate proteasomes and mitochondria. *BioRxiv*

An H, Harper JW (2018) Systematic analysis of ribophagy in human cells reveals bystander flux during selective autophagy. *Nat Cell Biol* 20: 135-143

An H, Ordureau A, Korner M, Paulo JA, Harper JW (2020) Systematic quantitative analysis of ribosome inventory during nutrient stress. *Nature* 583: 303-309

Antico O, Ordureau A, Stevens M, Singh F, Nirujogi RS, Gierlinski M, Barini E, Rickwood ML, Prescott A, Toth R *et al* (2021) Global ubiquitylation analysis of mitochondria in primary

neurons identifies endogenous Parkin targets following activation of PINK1. *Sci Adv* 7:
eabj0722
Bader M, Benjamin S, Wapinski OL, Smith DM, Goldberg AL, Steller H (2011) A conserved F
box regulatory complex controls proteasome activity in Drosophila. *Cell* 145: 371-382
Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society, Series B* 57: 289-300
Bingol B, Tea JS, Phu L, Reichelt M, Bakalarski CE, Song Q, Foreman O, Kirkpatrick DS, Sheng
M (2014) The mitochondrial deubiquitinase USP30 opposes parkin-mediated mitophagy.
Nature 510: 370-375
Burchell VS, Nelson DE, Sanchez-Martinez A, Delgado-Camprubi M, Ivatt RM, Pogson JH,
Randle SJ, Wray S, Lewis PA, Houlden H *et al* (2013) The Parkinson's disease-linked proteins
Fbxo7 and Parkin interact to mediate mitophagy. *Nat Neurosci* 16: 1257-1265
Burman JL, Pickles S, Wang C, Sekine S, Vargas JNS, Zhang Z, Youle AM, Nezich CL, Wu X,
Hammer JA *et al* (2017) Mitochondrial fission facilitates the selective mitophagy of protein
aggregates. *J Cell Biol* 216: 3231-3247
Chen G, Gulbranson DR, Hou Z, Bolin JM, Ruotti V, Probasco MD, Smuga-Otto K, Howden SE,
Diol NR, Propson NE *et al* (2011) Chemically defined conditions for human iPSC derivation
and culture. *Nat Methods* 8: 424-429
Deutsch EW, Bandeira N, Sharma V, Perez-Riverol Y, Carver JJ, Kundu DJ, Garcia-Seisdedos
D, Jarnuczak AF, Hewapathirana S, Pullman BS *et al* (2020) The ProteomeXchange
consortium in 2020: enabling 'big data' approaches in proteomics. *Nucleic Acids Res* 48:
D1145-D1152
Di Fonzo A, Dekker MC, Montagna P, Baruzzi A, Yonova EH, Correia Guedes L, Szczerbinska
A, Zhao T, Dubbel-Hulsman LO, Wouters CH *et al* (2009) FBXO7 mutations cause autosomal
recessive, early-onset parkinsonian-pyramidal syndrome. *Neurology* 72: 240-245
Dikic I, Elazar Z (2018) Mechanism and medical implications of mammalian autophagy. *Nat*
Rev Mol Cell Biol 19: 349-364
Elias JE, Gygi SP (2007) Target-decoy search strategy for increased confidence in large-scale
protein identifications by mass spectrometry. *Nat Methods* 4: 207-214
Eng JK, McCormack AL, Yates JR (1994) An approach to correlate tandem mass spectral data
of peptides with amino acid sequences in a protein database. *J Am Soc Mass Spectrom* 5:
976-989
Erickson BK, Mintseris J, Schweppe DK, Navarrete-Perea J, Erickson AR, Nusinow DP, Paulo
JA, Gygi SP (2019) Active Instrument Engagement Combined with a Real-Time Database
Search for Improved Performance of Sample Multiplexing Workflows. *J Proteome Res* 18:
1299-1306
Evans CS, Holzbaur EL (2020) Degradation of engulfed mitochondria is rate-limiting in
Optineurin-mediated mitophagy in neurons. *Elife* 9
Gladkova C, Maslen SL, Skehel JM, Komander D (2018) Mechanism of parkin activation by
PINK1. *Nature* 559: 410-414
Goodall EA, Kraus F, Harper JW (2022) Mechanisms underlying ubiquitin-driven selective
mitochondrial and bacterial autophagy. *Mol Cell* 82: 1501-1513
Gustafsson MG, Shao L, Carlton PM, Wang CJ, Golubovskaya IN, Cande WZ, Agard DA, Sedat
JW (2008) Three-dimensional resolution doubling in wide-field fluorescence microscopy by
structured illumination. *Biophys J* 94: 4957-4970
Harper JW, Ordureau A, Heo JM (2018) Building and decoding ubiquitin chains for
mitophagy. *Nat Rev Mol Cell Biol* 19: 93-108

Heo JM, Ordureau A, Paulo JA, Rinehart J, Harper JW (2015) The PINK1-PARKIN

Mitochondrial Ubiquitylation Pathway Drives a Program of OPTN/NDP52 Recruitment and

TBK1 Activation to Promote Mitophagy. *Mol Cell* 60: 7-20

Houlden H, Singleton AB (2012) The genetics and neuropathology of Parkinson's disease.

Acta Neuropathol 124: 325-338

Huang T, Fang L, He R, Weng H, Chen X, Ye Q, Qu D (2020) Fbxo7 and Pink1 play a reciprocal

role in regulating their protein levels. *Aging (Albany NY)* 13: 77-88

Hur JK, Kim K, Been KW, Baek G, Ye S, Hur JW, Ryu SM, Lee YS, Kim JS (2016) Targeted

mutagenesis in mice by electroporation of Cpf1 ribonucleoproteins. *Nat Biotechnol* 34: 807-

808

Huttlin EL, Bruckner RJ, Navarrete-Perea J, Cannon JR, Baltier K, Gebreab F, Gygi MP,

Thornock A, Zarraga G, Tam S *et al* (2021) Dual proteome-scale networks reveal cell-specific

remodeling of the human interactome. *Cell* 184: 3022-3040 e3028

Huttlin EL, Jedrychowski MP, Elias JE, Goswami T, Rad R, Beausoleil SA, Villen J, Haas W,

Sowa ME, Gygi SP (2010) A tissue-specific atlas of mouse protein phosphorylation and

expression. *Cell* 143: 1174-1189

Itzhak DN, Tyanova S, Cox J, Borner GH (2016) Global, quantitative and dynamic mapping of

protein subcellular localization. *Elife* 5

Jin J, Cardozo T, Lovering RC, Elledge SJ, Pagano M, Harper JW (2004) Systematic analysis

and nomenclature of mammalian F-box proteins. *Genes Dev* 18: 2573-2580

Jin SM, Lazarou M, Wang C, Kane LA, Narendra DP, Youle RJ (2010) Mitochondrial

membrane potential regulates PINK1 import and proteolytic destabilization by PARL. *J Cell*

Biol 191: 933-942

Kane LA, Lazarou M, Fogel AI, Li Y, Yamano K, Sarraf SA, Banerjee S, Youle RJ (2014) PINK1

phosphorylates ubiquitin to activate Parkin E3 ubiquitin ligase activity. *J Cell Biol* 205: 143-

153

Katayama H, Kogure T, Mizushima N, Yoshimori T, Miyawaki A (2011) A sensitive and

quantitative technique for detecting autophagic events based on lysosomal delivery. *Chem*

Biol 18: 1042-1052

Kazlauskaite A, Martinez-Torres RJ, Wilkie S, Kumar A, Peltier J, Gonzalez A, Johnson C,

Zhang J, Hope AG, Peggie M *et al* (2015) Binding to serine 65-phosphorylated ubiquitin

primes Parkin for optimal PINK1-dependent phosphorylation and activation. *EMBO Rep* 16:

939-954

Kirk R, Laman H, Knowles PP, Murray-Rust J, Lomonosov M, Meziane el K, McDonald NQ

(2008) Structure of a conserved dimerization domain within the F-box protein Fbxo7 and the

PI31 proteasome inhibitor. *J Biol Chem* 283: 22325-22335

Koyano F, Okatsu K, Kosako H, Tamura Y, Go E, Kimura M, Kimura Y, Tsuchiya H, Yoshihara

H, Hirokawa T *et al* (2014) Ubiquitin is phosphorylated by PINK1 to activate parkin. *Nature*

510: 162-166

Kraus F, Miron E, Demmerle J, Chitiashvili T, Budco A, Alle Q, Matsuda A, Leonhardt H,

Schermelleh L, Markaki Y (2017) Quantitative 3D structured illumination microscopy of

nuclear structures. *Nat Protoc* 12: 1011-1028

Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E (2019) CHOPCHOP

v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids Res* 47: W171-

W174

Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, Chen X, Taipale J, Hughes TR,

Weirauch MT (2018) The Human Transcription Factors. *Cell* 172: 650-665

Lazarou M, Jin SM, Kane LA, Youle RJ (2012) Role of PINK1 binding to the TOM complex and alternate intracellular membranes in recruitment and activation of the E3 ligase Parkin. *Dev Cell* 22: 320-333

Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, Sideris DP, Fogel AI, Youle RJ (2015) The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. *Nature* 524: 309-314

Li J, Cai Z, Bomgarden RD, Pike I, Kuhn K, Rogers JC, Roberts TM, Gygi SP, Paulo JA (2021) TMTpro-18plex: The Expanded and Complete Set of TMTpro Reagents for Sample Multiplexing. *J Proteome Res* 20: 2964-2972

Liu K, Jones S, Minis A, Rodriguez J, Molina H, Steller H (2019) PI31 Is an Adaptor Protein for Proteasome Transport in Axons and Required for Synaptic Development. *Dev Cell* 50: 509-524 e510

McAlister GC, Nusinow DP, Jedrychowski MP, Wuhr M, Huttlin EL, Erickson BK, Rad R, Haas W, Gygi SP (2014) MultiNotch MS3 enables accurate, sensitive, and multiplexed detection of differential expression across cancer cell line proteomes. *Anal Chem* 86: 7150-7158

McWilliams TG, Muqit MM (2017) PINK1 and Parkin: emerging themes in mitochondrial homeostasis. *Curr Opin Cell Biol* 45: 83-91

Narendra D, Tanaka A, Suen DF, Youle RJ (2008) Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *J Cell Biol* 183: 795-803

Narendra DP, Jin SM, Tanaka A, Suen DF, Gautier CA, Shen J, Cookson MR, Youle RJ (2010) PINK1 is selectively stabilized on impaired mitochondria to activate Parkin. *PLoS Biol* 8: e1000298

Ng MYW, Wai T, Simonsen A (2021) Quality control of the mitochondrion. *Dev Cell*

Okatsu K, Koyano F, Kimura M, Kosako H, Saeki Y, Tanaka K, Matsuda N (2015) Phosphorylated ubiquitin chain is the genuine Parkin receptor. *J Cell Biol* 209: 111-128

Ordureau A, Heo JM, Duda DM, Paulo JA, Olszewski JL, Yanishevski D, Rinehart J, Schulman BA, Harper JW (2015) Defining roles of PARKIN and ubiquitin phosphorylation by PINK1 in mitochondrial quality control using a ubiquitin replacement strategy. *Proc Natl Acad Sci U S A* 112: 6637-6642

Ordureau A, Kraus F, Zhang J, An H, Park S, Ahfeldt T, Paulo JA, Harper JW (2021) Temporal proteomics during neurogenesis reveals large-scale proteome and organelle remodeling via selective autophagy. *Mol Cell*

Ordureau A, Paulo JA, Zhang J, An H, Swatek KN, Cannon JR, Wan Q, Komander D, Harper JW (2020) Global Landscape and Dynamics of Parkin and USP30-Dependent Ubiquitylomes in iNeurons during Mitophagic Signaling. *Mol Cell* 77: 1124-1142 e1110

Ordureau A, Paulo JA, Zhang W, Ahfeldt T, Zhang J, Cohn EF, Hou Z, Heo JM, Rubin LL, Sidhu SS *et al* (2018) Dynamics of PARKIN-Dependent Mitochondrial Ubiquitylation in Induced Neurons and Model Systems Revealed by Digital Snapshot Proteomics. *Mol Cell* 70: 211-227 e218

Ordureau A, Sarraf SA, Duda DM, Heo JM, Jedrychowski MP, Sviderskiy VO, Olszewski JL, Koerber JT, Xie T, Beausoleil SA *et al* (2014) Quantitative proteomics reveal a feedforward mechanism for mitochondrial PARKIN translocation and ubiquitin chain synthesis. *Mol Cell* 56: 360-375

Paisan-Ruiz C, Guevara R, Federoff M, Hanagasi H, Sina F, Elahi E, Schneider SA, Schwingenschuh P, Bajaj N, Emre M *et al* (2010) Early-onset L-dopa-responsive parkinsonism with pyramidal signs due to ATP13A2, PLA2G6, FBXO7 and spatacsin mutations. *Mov Disord* 25: 1791-1800

Paulo JA, O'Connell JD, Gygi SP (2016) A Triple Knockout (TKO) Proteomics Standard for
Diagnosing Ion Interference in Isobaric Labeling Experiments. *J Am Soc Mass Spectrom* 27:
1620-1625
Perez-Riverol Y, Bai J, Bandla C, Garcia-Seisdedos D, Hewapathirana S, Kamatchinathan S,
Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M *et al* (2022) The PRIDE database
resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids*
Res 50: D543-D552
Pickrell AM, Youle RJ (2015) The roles of PINK1, parkin, and mitochondrial fidelity in
Parkinson's disease. *Neuron* 85: 257-273
Plubell DL, Wilmarth PA, Zhao Y, Fenton AM, Minnier J, Reddy AP, Klimek J, Yang X, David LL,
Pamir N (2017) Extended Multiplexing of Tandem Mass Tags (TMT) Labeling Reveals Age and
High Fat Diet Specific Proteome Changes in Mouse Epididymal Adipose Tissue. *Mol Cell*
Proteomics 16: 873-890
Rad R, Li J, Mintseris J, O'Connell J, Gygi SP, Schweppe DK (2021) Improved Monoisotopic
Mass Estimation for Deeper Proteome Coverage. *J Proteome Res* 20: 591-598
Rath S, Sharma R, Gupta R, Ast T, Chan C, Durham TJ, Goodman RP, Grabarek Z, Haas ME,
Hung WHW *et al* (2021) MitoCarta3.0: an updated mitochondrial proteome now with sub-
organelle localization and pathway annotations. *Nucleic Acids Res* 49: D1541-D1547
Ravenhill BJ, Boyle KB, von Muhlinen N, Ellison CJ, Masson GR, Otten EG, Foeglein A,
Williams R, Randow F (2019) The Cargo Receptor NDP52 Initiates Selective Autophagy by
Recruiting the ULK Complex to Cytosol-Invading Bacteria. *Mol Cell* 74: 320-329 e326
Sarraf SA, Raman M, Guarani-Pereira V, Sowa ME, Huttlin EL, Gygi SP, Harper JW (2013)
Landscape of the PARKIN-dependent ubiquitylome in response to mitochondrial
depolarization. *Nature* 496: 372-376
Sauve V, Sung G, Soya N, Kozlov G, Blaimschein N, Miotto LS, Trempe JF, Lukacs GL, Gehring
K (2018) Mechanism of parkin activation by phosphorylation. *Nat Struct Mol Biol* 25: 623-
630
Savitski MM, Wilhelm M, Hahne H, Kuster B, Bantscheff M (2015) A Scalable Approach for
Protein False Discovery Rate Estimation in Large Proteomic Data Sets. *Mol Cell Proteomics*
14: 2394-2404
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for biological-image
analysis. *Nat Methods* 9: 676-682
Schweppe DK, Eng JK, Yu Q, Bailey D, Rad R, Navarrete-Perea J, Huttlin EL, Erickson BK, Paulo
JA, Gygi SP (2020) Full-Featured, Real-Time Database Searching Platform Enables Fast and
Accurate Multiplexed Quantitative Proteomics. *J Proteome Res* 19: 2026-2034
Schweppe DK, Prasad S, Belford MW, Navarrete-Perea J, Bailey DJ, Huguet R, Jedrychowski
MP, Rad R, McAlister G, Abbatiello SE *et al* (2019) Characterization and Optimization of
Multiplexed Quantitative Analyses Using High-Field Asymmetric-Waveform Ion Mobility
Mass Spectrometry. *Anal Chem* 91: 4010-4016
Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A (2021)
CellProfiler 4: improvements in speed, utility and usability. *BMC Bioinformatics* 22: 433
Stolz A, Ernst A, Dikic I (2014) Cargo recognition and trafficking in selective autophagy. *Nat*
Cell Biol 16: 495-501
Tai YC (2022) timecourse: Statistical Analysis for Developmental Microarray Time Course
Data. R package version 1.68.0. *http://wwwbioconductororg*

Tyanova S, Cox J (2018) Perseus: A Bioinformatics Platform for Integrative Analysis of
Proteomics Data in Cancer Research. *Methods Mol Biol* 1711: 133-148
Vargas JNS, Wang C, Bunker E, Hao L, Maric D, Schiavo G, Randow F, Youle RJ (2019)
Spatiotemporal Control of ULK1 Activation by NDP52 and TBK1 during Selective Autophagy.
Mol Cell 74: 347-362 e346
Vingill S, Brockelt D, Lancelin C, Tatenhorst L, Dontcheva G, Preisinger C, Schwedhelm-
Domeyer N, Joseph S, Mitkovski M, Goebbels S *et al* (2016) Loss of FBXO7 (PARK15) results
in reduced proteasome activity and models a parkinsonism-like phenotype in mice. *EMBO J*
35: 2008-2025
Wang Y, Yang F, Gritsenko MA, Wang Y, Clauss T, Liu T, Shen Y, Monroe ME, Lopez-Ferrer D,
Reno T *et al* (2011) Reversed-phase chromatography with multiple fraction concatenation
strategy for proteome profiling of human MCF10A cells. *Proteomics* 11: 2019-2026
Wauer T, Simicek M, Schubert A, Komander D (2015) Mechanism of phospho-ubiquitin-
induced PARKIN activation. *Nature* 524: 370-374
Wong YC, Holzbaur EL (2014) Optineurin is an autophagy receptor for damaged
mitochondria in parkin-mediated mitophagy that is disrupted by an ALS-linked mutation.
Proc Natl Acad Sci U S A 111: E4439-4448
Yamano K, Youle RJ (2013) PINK1 is degraded through the N-end rule pathway. *Autophagy*
9: 1758-1769
Zhang Y, Pak C, Han Y, Ahlenius H, Zhang Z, Chanda S, Marro S, Patzke C, Acuna C, Covy J *et*
al (2013) Rapid single-step induction of functional neurons from human pluripotent stem
cells. *Neuron* 78: 785-798
Zuris JA, Thompson DB, Shu Y, Guilinger JP, Bessen JL, Hu JH, Maeder ML, Joung JK, Chen ZY,
Liu DR (2015) Cationic lipid-mediated delivery of proteins enables efficient protein-based
genome editing in vitro and in vivo. *Nat Biotechnol* 33: 73-80
1360