Transcriptome profiles of eggplant (*Solanum melongena*) and its wild relative *S.*

***dasyphyllum* under different levels of osmotic stress provide insights into response**

mechanisms to drought

Gloria Villanueva[a], Santiago Vilanova[a], Mariola Plazas[a], Jaime Prohens[a], Pietro

Gramazio[b].

Affiliations

[a]*Instituto de Conservación y Mejora de la Agrodiversidad Valenciana, Universitat*

Politècnica de València, Camino de Vera 14, 46022 Valencia, Spain

[b]*Instituto de Biología Molecular y Celular de Plantas, Consejo Superior de*

Investigaciones Científicas-Universitat Politècnica de València, Camino de Vera 14,

46022 Valencia, Spain

*Corresponding author:

Gloria Villanueva: glovilpa@etsiamn.upv.es

Abstract

Defence mechanisms to abiotic stresses, like drought, are very broad and RNA sequencing (RNA-Seq) can help in understanding the complex responses triggered. In this study, we performed RNA-Seq of the transcriptomes of eggplant (*Solanum melongena*) and its related wild species (*S. dasyphyllum*) under two PEG concentrations (20% and 30%), two different times (after 0.5 h and 2 h of osmotic stress) and at two plant phenological stages (three and five true fully developed leaves). *Solanum dasyphyllum* was more tolerant to osmotic stress, and a differential expression pattern of drought-related genes was identified between the two species. Plants subjected to a higher osmotic potential, at a more adult stage and at a higher stress exposure time displayed a higher number of DEGs (differential expressed genes). Gene ontology (GO) enrichment analysis revealed that, compared to *S. melongena*, *S. dasyphyllum* triggered the regulation of a wide range of transcription factors (*AP2/ERF*, DREB, bZIP, WRKY and bHLH). In both species, the abscisic acid (ABA) signaling response pathway played a crucial role leading to stomatal closure. Other important pathways involved in abiotic stresses tolerance including flavonoid, carotenoid and phenylpropanoid biosynthesis, chlorophyll metabolism and photosynthesis pathway among others were found to have a relevant role under both moderate and severe osmotic stresses. Our results reveal that *S. dasyphyllum* is a potential source of genes for breeding resilient eggplant varieties.

Keywords: transcriptome, RNA Seq, osmotic stress, drought, *Solanum melongena*, *S. dasyphyllum.*

1. Introduction

Drought spells occur naturally in many areas of the world, but climate change has accelerated and intensified them, with dramatic consequences on agriculture (Mukherjee et al., 2018). Projections indicate that the risk and severity of drought episodes will increase across the subtropics and mid-latitudes in both hemispheres as a consequence of global warming and decreased regional precipitation (Cook et al., 2018; IPCC, 2022). Drought stress triggers morphological, physiological, biochemical, cellular and molecular response mechanisms in plants with a potentially severe reduction in plant growth and crop production as a major consequence. Therefore, determining plant response and tolerance mechanisms against drought stress is fundamental to mitigating its effects (Hossain et al., 2016; Ilyas et al., 2021).

The development of new molecular and bioinformatics tools has allowed the expansion of applied knowledge in breeding programs. In this way, transcriptomics has provided new potential resources for studying the molecular response of abiotic stress in crops (Zhuang et al., 2014), being RNA sequencing (RNA-Seq) the general method of choice. This method allows a broad coverage of the transcriptome, providing a significant characterization of mRNA transcripts of specific tissue and time and, in addition, is a quantitative method that yields a digital gene expression atlas at a genomic scale (Kido et al., 2016).

Drought tolerance is a complex trait involving different components at the physiological, biochemical and genetic levels (Samarah, 2016). Osmotic stress, resulting in an increased difficulty for water uptake by the roots, is one of the most important factors in drought (Haswell and Verslues, 2015). To unravel the effects of water deficit in genetic networks, the use of a solution containing polyethylene glycol (PEG) in hydroponic culture is a common practice to induce osmotic stress and reduce

the water potential of tissues in plants (Claeys and Inzé, 2013; Osmolovskaya et al.,

2018). In this way, the transcriptome of PEG-treated plants provides information

regarding drought-related genes, which can be primarily classified in protective and

regulatory genes (Kido et al., 2016). Regarding the former, these are genes that encode

LEA proteins, chaperones, osmoprotectants, water channels, ion exchangers, and

enzymes involved in the osmolyte biosynthesis and the reactive oxygen species (ROS),

among others (Chen et al., 2019; Riyazuddin et al., 2021). On the other hand, genes

encoding regulatory proteins act on the expression of stress-responsive, including

transcription factors, protein kinases and phosphatases, enzymes involved in

phospholipid metabolism and abscisic acid (ABA) biosynthesis and epigenetic-related

genes (Shinozaki and Yamaguchi-Shinozaki, 2007; Wang et al., 2021).

Crop wild relatives (CWRs) are an increasingly fundamental resource for plant breeding

to improve the adaptative capacity of agricultural systems to climate change-related

stresses (Prohens et al., 2017). Among vegetable crops, eggplant (*Solanum melongena*

L.) can be highly benefited by introgression breeding, as many eggplant CWRs thrive in

areas affected by moderate to severe drought (Vorontsova and Knapp, 2016). Eggplant

is an important crop, being the eighth vegetable crop in terms of cultivated area in the

world, being widely grown in Asia, Africa and Europe (FAO, 2020). It has been

described as a relatively drought-tolerant crop and different degrees of drought

tolerance have been observed in cultivated accessions and CRWs (Díaz-Pérez and

Eaton, 2015; Kouassi et al., 2020; Plazas et al., 2019a). Among these CWRs, *S.*

dasyphyllum Schumach. and Thonn. grows naturally in areas where drought spells are

frequent and it has been reported to exhibit significant drought tolerance both under

field and experimental conditions (Kouassi et al., 2020; Plazas et al., 2022). It is

considered the wild ancestor of the gboma eggplant (*S. macrocarpon* L.) (Bukenya and

Carasco, 1994; Plazas et al., 2014) and is classified in the Anguivi clade, which includes

several African and Southeast Asian "prickly" species (Vorontsova et al., 2013;

Vorontsova and Knapp, 2016). *Solanum dasyphyllum* is a member of the secondary

genepool of eggplant (Syfert et al., 2016), and interspecific hybrids and advanced

backcross materials of *S. dasyphyllum* with *S. melongena* have been obtained (Kouassi

et al., 2016; Plazas et al., 2016).

In the present study, we analyzed the transcriptomes of a cultivated *S. melongena* and a

drought-tolerant *S. dasyphyllum* accessions under PEG-induced osmotic stress in two

different plant phenological stages and at two times for each phenological stage. By

evaluating its physiological responses in conjunction with the analysis of the gene

expression we aimed at a better comprehensive understanding of the different response

mechanisms against osmotic stress in these materials. The results are of great interest

for a better understanding of drought tolerance and to foster introgression breeding of

drought-tolerant resilient cultivars in eggplant.

2. Material and Methods

2.1. Plant materials and growth conditions

Solanum melongena MEL1 and *S. dasyphyllum* DAS1 accessions were used for the

present study. Seeds were germinated according to Ranil et al. (2015) protocol for

uniform eggplant CWRs germination and plants were grown in hydroponic culture

according to Renau-Morata et al. (2016) with Hoagland solution (Hoagland and Arnon,

1950) in a growth chamber with a 16/8 h light/dark photoperiod, 25ºC temperature and

60-65% of humidity. The nutrient solution was resupplied every four days and an air

compressor was used to supply aeration.

2.2. PEG-induced osmotic stress

To evaluate the effect of the plant phenological stage and the stress response, two

osmotic stress experiments were conducted using PEG 6000 (Bio Basic Inc., Ontario,

Canada). One experiment was performed with 20% PEG at a phenological stage of three

fully developed true leaves (Ex_1), while the other with 30% PEG at the five fully

developed true leaves stage (Ex_2). In each experiment, leaves of three biological

replicates (i.e., three different plants uniformly developed, each one constituting a

replicate and for each one a library was developed) were taken for each species at three

times: 0 h (control; T0), 0.5 h (T0.5) and 2 h (T2) after initiation of the stress treatment.

Immediately, leaf samples were frozen with liquid nitrogen and stored at -80ºC for RNA

extraction. Plant symptoms were registered at different times of the treatments.

2.3. RNA extraction, sequencing and data processing

Total RNA was extracted from leaves samples of each biological replicate using

TRIzol^{TM} Reagent (Invitrogen, Carlsbad, CA, USA). For each of the 36 replicates, the

RNA library was performed by Novogene Co., LTD (Beijing, China) and sequenced on

an Illumina NovaSeq 6000 (paired-end 150 bp). Raw data in FASTQ format were

filtered by removing reads with adaptor contamination, reads containing N > 10% and

low-quality reads (Qscore of over 50% bases below 5). Error rate (%), Q20 (%), Q30

(%) and GC content (%) were calculated for data quality control of clean data. Gene

expression levels were estimated by calculating fragments per kilobase of transcript

sequence per millions of base pairs sequenced (FPKM).

2.4. Transcriptomic analysis

Differentially expressed genes (DEGs) analysis was performed using the DEseq2 R

package (Anders and Huber, 2010), and the resulting *p*-values were adjusted using

Benjamini and Hochberg's correction for controlling the false discovery rate (FDR)

(Benjamini and Hochberg, 1995). Genes with adjusted p-value < 0.05 and |log$_2$(fold

change)| > 1 were considered as differentially expressed. DEGs were annotated based

on the functional annotation information of genes of the eggplant reference genome

"67/3" V3 (Barchi et al., 2019). Venn diagrams of DEGs were displayed using jvenn, a

plug-in for the jQuery JavaScript library (Bardou et al., 2014).

Hierarchical clustering analysis was carried out of log$_2$ (FPKM+1) of union differential

expression genes, within all comparison groups. Heatmaps were performed selecting

drought-related DEGs, based on the scientific literature, which were classified

according to their function into four groups: osmoprotectants, phytohormones, protein

kinases and transcription factors using the web tool ClustVis (Metsalu and Vilo, 2015).

Gene ontology (GO, http://www.geneontology.org/) and Kyoto Encyclopedia of Genes

and Genomes (KEGG, http://www.genome.jp/kegg/) enrichment analyses of the DEGs

were performed. The tomato (*S. lycopersicum* L.) KEGG pathways annotated database

was used for the analysis, being the closest species with more comprehensive and

reliable information. GO and KEGG terms with an adjusted p-value < 0.05 were

considered significantly enriched for the DEGs.

3. Results

3.1. Physiological responses to osmotic stress

As a general trend, in both experiments, *S. dasyphyllum* (DAS) displayed a better water

stress tolerance than *S. melongena* (MEL). In Ex_1, DAS presented visual symptoms

only at T2 while MEL started to show symptoms of stress at T0.5 (Figure 1). In Ex_2,

manifestations of water stress in plants were observed at T0.5 and T2 in both species in

a faster way with more severe symptoms compared with Ex_1, although DAS, again,

exhibit more tolerance, with fewer symptoms of wilting (Figure 1).

Figure 1. Representative phenotypes of *S. melongena* (MEL) and *S. dasyphyllum*

(DAS) after 0, 0.5 and 2h of PEG stress in hydroponic culture in both experiments

(Ex_1 and Ex_2).

3.2. Differential gene expression over time in response to PEG treatment

After filtering raw sequencing data, clean reads showed an error rate between 0.02 and

0.03%, an average Q30 of 93.85% and GC content of 43.17% (Table S1). For each

experiment, DEGs with an adjusted *p*-value < 0.05 and a |$\log_2$ (fold change)| > 1 were

selected by performing pairwise comparisons at each time of PEG treatment (T0.5 and

T2) with the non-stressed control (T0).

In Ex_1 (20% PEG and three fully developed true leaves stage), a total of 894 and 433

DEGs were detected for DAS and MEL, respectively. For DAS a total of 114 (74 up-

regulated [UR], 40 down-regulated [DR] and 33 related to drought stress) and 840

DEGs (475 UR, 365 DR and 171 related to drought stress) were detected at T0.5 and

T2, respectively (Table 1). For MEL, a total of 327 (273 UR, 54 DR and 89 related to

drought stress) and 117 DEGs (76 UR, 41 DR and 24 related to drought stress) were

detected at T0.5 and T2, respectively (Table 1). Venn diagram analysis showed that in

DAS 52 DEGs were commonly regulated at T0.5 and T2 while 49 and 707 DEGs were

specific at 0.5 and T2, respectively (Figure 2A). In MEL, seven DEGs were commonly

regulated after both times of treatment, 273 and 67 DEGs at T0.5 and T2 respectively

(Figure 2A).

In Ex_2 (30% PEG and five fully developed true leaves) a total of 2,037 and 4,375

DEGs were detected for DAS and MEL, respectively. For DAS, a total of 147 (109 UR

38 DR and 53 related to drought stress) and 1,999 DEGs (1,040 UR, 959 DR and 363

related to drought stress) were detected at T0.5 and T2, respectively (Table 1). For

MEL, a total of 198 (134 UR, 64 DR and 62 related to drought stress) and 4,360 DEGs

(2,252 UR, 2,108 DR and 774 related to drought stress) were detected at T0.5 and T2,

respectively (Table 1). Venn diagram analysis showed that 31 and 80 DEGs were

commonly regulated at T0.5 and T2 exclusively in DAS and MEL respectively (Figure

2B). A total of 20 and 1,065 DEGs were detected only in DAS at T0.5 and T2

respectively. In MEL, 11 and 3,363 DEGs were detected exclusively at T0.5 and T2

respectively. A total of 38 common DEGs were detected for both times and both

accessions. (Figure 2B).

Table 1. Differentially expressed genes that were up-regulated or down-regulated after

0.5 h (T0.5) and 2 h (T2) of PEG stress in *S. dasyphyllum* (DAS) and *S. melongena*

(MEL) in experiments 1 and 2 (Ex_1 and Ex_2).

Experiment	Time comparison	Number of DEGs		Up-Regulated DEGs		Down-Regulated DEGs		Number of DEGs related to drought	
		DAS	MEL	DAS	MEL	DAS	MEL	DAS	MEL
Ex_1	T0 vs T0.5	114	327	74	273	40	54	33	89
	T0 vs T2	840	117	475	76	365	41	171	24
Ex_2	T0 vs T0.5	147	198	109	134	38	64	53	62
	T0 vs T2	1,999	4,360	1,040	2,252	959	2,108	363	774

Figure 2. Venn diagram of DEGs under 0.5 and 2h of PEG stress of *S. dasyphyllum*

(DAS) and *S. melongena* (MEL) in experiment 1 (Ex_1; **A**) and experiment 2 (Ex_2;

B).

Drought-responsive DEGs were classified according to their function into four groups:

osmoprotectants, phytohormones, protein kinases and transcription factors related to the

drought stress response. A total of 264 DEGs related to drought were observed in Ex_1,

of which 38 of them were genes related to osmoprotectants, 46 were related to the

synthesis of phytohormones, 67 were protein kinases genes and 113 were transcription

factors. In Ex_2 a total of 953 DEGs were detected, of which 150 were genes that

encode for proteins related to osmoprotectants, 180 were related to phytohormones, 296

for protein kinases and 327 transcription factors genes (Figure 3).

In both experiments, in general, the expression pattern of drought-responsive genes

changed over time for both accessions, allowing clear differentiation between

accessions and time of exposure to stress. In all cases, up-regulated and down-regulated

genes from the different groups of the classification were observed (Figure 3).

Figure 3. Heatmap of DEGs related to drought stress, osmoprotectants (OP), phytohormones (PHY), protein kinases (PK) and transcription factor (TF) after 0, 0.5 and 2 h of PEG stress of *S. dasyphyllum* (DAS) and *S. melongena* (MEL) in experiments 1 (Ex_1; **A**) and 2 (Ex_2; **B**).

3.3. GO and KEGG enrichment in DEGs according to phenological stage and stress

conditions

A gene ontology (GO) analysis was performed with DEGs being annotated as a

biological process (BP), cellular components (CC) and molecular function (MF). In

Ex_1, for DAS at T0.5, 37 DEGs were annotated as MF, 13 of them as DNA-binding

transcription factor activity (10 UR and three DR), three as xyloglucan:xyloglucosyl

transferase activity (UR), six as sequence-specific DNA binding (UR), eight as

transferase activity (transferring glycosyl groups; six UR and two DR), seven as

transferase activity (transferring hexosyl groups; five UR and two DR). Regarding CC,

three were annotated as apoplast and cell wall (Figure 4A). After 2 h of osmotic stress

(T2), in DAS, all significant DEGs were annotated as MF, 45 as DNA-binding

transcription factor activity (24 UR and 21 DR), 25 as sequence-specific DNA binding

(16 UR and nine DR) and eight as terpene synthase activity (DR; Figure 4B). For MEL,

at T0.5, a total of 45 DEGs were annotated as MF, 22 of them as DNA-binding

transcription factor activity (19 UR and three DR), nine as calcium ion binding (UR)

and 10 as sequence-specific DNA binding (seven UR and three DR) and four as G

protein-coupled receptor signaling pathway as biological process (UR; Figure 4C). In

MEL at T2, three CC DEGs were annotated as apoplast and cell wall (DR) and three BP

DEGs as phosphorelay signal transduction system (two UR and one DR; Figure 4D).

249
Figure 4. Gene ontology (GO) terms enrichment scatter plot in DEGs of *S. dasyphyllum*

(DAS) after 0.5 (T0.5) **(A)** and 2 h (T2) **(B)** versus 0 h of PEG stress and *S. melongena*

(MEL) after 0.5 (T0.5) **(C)** and 2 h (T2) **(D)** compared with 0 h of PEG stress in

experiment 1 (Ex_1).

In Ex_2 for DAS at T0.5, a total of 44 DEGs were annotated, 17 of them as BP, three as

CC and 24 as MF. Within the biological process category, seven DEGs were annotated

as a response to chemical (one UR and six DR), six as response to auxin (DR) and four

as cellular glucan metabolic process (UR), while as CC, three as apoplast and cell wall

(UR). As molecular function, 11 were annotated as DNA-binding transcription factor

activity (10 UR and one DR), three as xyloglucan:xyloglucosyl transferase activity

(UR), four as glucosyltransferase activity (UR) and six as sequence-specific DNA

binding (UR; Figure 5A). At T2, in DAS, a total of 82 DEGs were annotated as BP, 37

as response to chemical (10 UR and 27 DR), 24 as response to hormone (one UR and 23

DR) and 21 as response to auxin (DR). As MF, 49 DEGs as protein dimerization

activity (29 UR and 20 DR) and 57 as DNA-binding transcription factor activity (35 UR

and 22 DR) (Figure 5B). For MEL, significant GO terms annotated for BP at T0.5 were

seven to response to hormone (one UR and six DR), six to response to auxin (DR) and

eight to response to chemical (two UR and six DR). Also, 16 DEGs were annotated as

DNA-binding transcription factor activity (15 UR and one DR) in MF classification

(Figure 5C). At T2, 161 DEGs were classified as BP, 51 as response to hormone (12 UR

and 39 DR), 44 as response to auxin (10 UR and 34 DR) and 66 as response to chemical

(24 UR and 42 DR). As molecular function, 125 were annotated as DNA-binding

transcription factor activity (102 UR and 23 DR), 72 as sequence-specific DNA binding

(52 UR and 15 DR), 41 as enzyme inhibitor activity (27 UR and 14 DR), 52 as

transferase activity (transferring acyl groups other than amino-acyl groups; 31 UR and

21 DR), 62 as transferase activity (transferring acyl groups; 37 UR and 25 DR), 11

oxidoreductase activity (acting on the aldehyde or oxo group of donors as; five UR and

six DR), seven as chitinase activity (six UR and one DR), seven as calcium-dependent

phospholipid binding (UR) and 19 as endopeptidase inhibitor activity (17 UR and two

DR) (Figure 5D). Enriched genes annotated in each GO term classification were

included in Table S2.

Figure 5. Gene ontology (GO) terms enrichment scatter plot in DEGs of *S. dasyphyllum*

(D) after 0.5 h (T0.5) **(A)** and 2 h (T2) **(B)** versus 0 h of PEG stress and *S. melongena*

(M) after 0.5 h (T0.5) **(C)** and 2 h (T2) **(D)** compared with 0h of PEG stress experiment

2 (Ex_2).

A pathway enrichment analysis using the Kyoto Encyclopedia of Genes and Genomes

(KEGG) was performed to identify significant (padj < 0.05) enriched metabolic or

signal transduction pathways associated with differentially expressed genes (DEGs)

comparing the whole genome background. In Ex_1 more DEGs were assigned to

KEGG pathways in DAS than in MEL. For DAS, at T0.5 and T2, plant hormone signal

transduction and MAPK (mitogen activated protein kinase) signaling pathway were

identified as enriched pathway (five UR and five DR DEGs). For T2 were also

determined circadian rhythm (seven UR and two DR DEGs), sesquiterpenoid and

triterpenoid biosynthesis (seven DR DEGs), galactose metabolism (seven UR DEGs)

and zeatin biosynthesis (one UR and eight DR) as enriched pathways. For MEL at T0.5,

plant-pathogen interaction (10 UR DEGs) and, at T2, circadian rhythm (eight UR and

two DR DEGs) were enriched pathways detected (Table 2). In Ex_2, more expressed

genes were assigned to metabolic pathways in MEL than for DAS. For DAS at T0.5,

DEGs were assigned to plant hormone signal transduction (six UR and eight DR) and

also to MAPK signaling pathway (three UR and four DR). At T2, plant hormone signal

transduction (16 UR and 29 DR DEGs) and the phenylpropanoid biosynthesis (17 UR

and 11 DR) were determined as enriched pathways. For MEL at T0.5, plant hormone

signal transduction (nine UR and six DR DEGs), MAPK signaling pathway (three UR

and four DR), fatty acid elongation (three DR), and carotenoid biosynthesis (three UR)

were found to be enriched pathways. At T2, DEGs were linked to porphyrin and

chlorophyll metabolism (seven UR and 21 DR), plant hormone signal transduction (58

UR and 37 DR), photosynthesis and antenna proteins (10 DR), α-linolenic acid

metabolism (24 UR and two DR MAPK signaling pathway (53 UR and 16 DR),

flavonoid biosynthesis (nine UR and nine DR and glutathione metabolism (21 UR and

seven DR) (Table 2). Enriched genes annotated in each KEGG pathway classification

were included in Table S3.

Table 2. Significant Kyoto Encyclopedia of Genes and Genomes (KEGG) enriched

pathways and its ID of tomato database in *S. dasyphyllum* (DAS) and *S. melongena*

(MEL) after 0.5 h (T0.5) and 2 h (T2) of PEG stress in experiments 1 (Ex_1) and 2

(Ex_2).

Experiment	Time comparison	Accession	Pathway Terms	ID	Count	Up-Regulated	Down-Regulated
Ex_1	T0 vs T0.5	DAS	Plant hormone signal transduction	sly04075	10	5	5
			MAPK signaling	sly04016	8	3	5

			pathway - plant				
		MEL	Plant-pathogen interaction	sly04626	10	10	0
	T0 vs T2	DAS	Plant hormone signal transduction	sly04075	32	12	20
			MAPK signaling pathway - plant	sly04016	27	10	17
			Circadian rhythm - plant	sly04712	9	7	2
			Sesquiterpenoid and triterpenoid biosynthesis	sly00909	7	0	7
			Galactose metabolism	sly00052	7	7	0
			Zeatin biosynthesis	sly00908	8	1	7
		MEL	Circadian rhythm - plant	sly04712	10	8	2
Ex_2	T0 vs T0.5	DAS	Plant hormone signal transduction	sly04075	14	6	8
			MAPK signaling pathway - plant	sly04016	7	3	4
		MEL	Plant hormone signal transduction	sly04075	15	9	6
			MAPK signaling pathway - plant	sly04016	10	7	3
			Fatty acid elongation	sly00062	3	0	3
			Carotenoid biosynthesis	sly00906	3	3	0
	T0 vs T2	DAS	Plant hormone signal transduction	sly04075	45	16	29
			Phenylpropanoid biosynthesis	sly00940	28	17	11
		MEL	Porphyrin and chlorophyll metabolism	sly00860	28	7	21
			Plant hormone signal transduction	sly04075	95	58	37
			Photosynthesis - antenna proteins	sly00196	10	0	10
			α-Linolenic acid metabolism	sly00592	26	24	2
			MAPK signaling pathway - plant	sly04016	69	53	16
			Flavonoid biosynthesis	sly00941	18	9	9
			Glutathione metabolism	sly00480	28	21	7

4. Discussion

Eggplant has been considered a relatively drought-tolerant crop since a long time ago (Behboudian, 1977) and several studies to evaluate the physiological and biochemical responses to water stress of different eggplant cultivars and wild relatives have been performed (Delfin et al., 2021; Fu et al., 2013; Plazas et al., 2019). However, detailed molecular mechanisms in response to drought stress in eggplant are not well known and, to our knowledge, transcriptional analysis by RNA-Seq method has not been

reported so far. In the current study, we evaluated plants of the cultivated eggplant *S.*

melongena and its wild relative *S. dasyphyllum* under two concentrations of PEG (20%

and 30%) at two different phenological stages (three and five fully developed true

leaves) in hydroponic conditions in order to obtain a general overview of their response

to osmotic stress and get insight in the gene expression involved in response and

tolerance to drought. *Solanum dasyphyllum* displayed a better water deficit tolerance

than *S. melongena*, confirming its already recently reported drought tolerance in field

and experimental conditions (Kouassi et al., 2020; Plazas et al., 2022). PEG

concentration had a visually significant effect in physiological response, with more

symptoms in Ex_2, in which plants were subjected to a higher PEG concentration,

resulting in a higher osmotic potential (Michel and Kaufmann, 1973).

RNA sequencing is a tool for transcriptome analysis that has allowed a better

understanding of the functions of the genome (Stark et al., 2019). In this research, the

analysis of differential gene expression has enabled the study of the response to osmotic

stress in both species at the genomic level. One of the most important components of

drought stress is osmotic stress and it has been widely used to study drought tolerance in

many species (Haswell and Verslues, 2015). In our study, in general, osmotic stress

treatments mainly triggered an activation response, as more significantly up-regulated

than down-regulated DEGs were observed. The number of DEGs increased as PEG

concentration was higher and longer in time, as was previously reported in potato

(*Solanum tuberosum* L.) (Moon et al., 2018; Yang et al., 2019). The expression pattern

of drought-responsive genes displayed large differences between *S. dasyphyllum* and *S.*

melongena, revealing very divergent response mechanisms under an osmotic stress

according to plant physiological observations.

This study has disclosed the main functions and pathways expressed of two related

species with large differences in osmotic stress response. GO enrichment of the

identified DEGs has allowed establishing the biological functions associated to those

genes. *Solanum dasyphyllum* expressed genes were involved in diverse functions related

to osmotic stress response. On one side, genes involved in the modification of cell wall

and apoplast structure, such as xyloglucan:xyloglucosyl transferases (Stratilová et al.,

2020; Tenhaken, 2015), were enriched in the wild species. Other up-regulated genes in

S. dasyphyllum in response to osmotic stress were classified in the DNA-binding

transcription factor activity and sequence-specific DNA binding GO terms, including a

wide range of transcription factors (TFs), which exert crucial roles in diverse signaling

pathways in different abiotic stress response as *AP2/ERF* (APETALA2/Ethylene

Response Factor) family (D. Li et al., 2021) and two of its major subfamilies such as

dehydration-responsive element binding proteins (DREBs) and ethylene-responsive

element (ERE) binding factors (Islam and Wang, 2009; Sun et al., 2008). The same

occurs with TFs, from homeobox-leucine zipper family (Gao et al., 2015; Jiao et al.,

2022; Perotti et al., 2017), basic leucine zipper (bZIP) (Zhu et al., 2018) and WRKY

family (Chen et al., 2017). Meanwhile, in the case of *S. melongena*, the expression of

AP2/ERF, WRKY and bZIP TFs was also observed, however, in general, the number of

differential genes expressed under the stress treatments was fewer. When plants were

subjected to the higher osmotic potential, the overall gene expression was also higher

and included DEGs classified in response to chemical and hormones and also down-

regulated genes related to auxin response. Auxins are involved in the regulation of plant

growth and development and auxin response factors (ARFs) gene family play an

essential role in the regulation of auxin-relative genes in abiotic stress responses in

tomato (*S. lycopersicum*) (Bouzroud et al., 2018). Basic helix-loop-helix (bHLH)

transcription factors were overexpressed in *S. dasyphyllum*, which they have been

reported to be involved in the response to abiotic stresses in potato (*S. tuberosum*)

(Wang et al., 2018) and pepper (*Capsicum annuum* L.) (Zhang et al., 2020). For *S.*

melongena, exposure to a higher osmotic stress resulted in the differential expression of

genes related to enzyme inhibitor activity, transferases, chitinase and oxidoreductase

activities, among others. The overall response observed was very broad, with the wild

species (*S. dasyphyllum*) showing a greater and more diverse expression of genes

involved in drought response, which could be related to its increased tolerance.

KEGG analysis revealed significant enriched pathways related to osmotic stress such as

plant hormone signal transduction and MAPK signaling. In these pathways, genes

encoding for the three main components of the core Abscisic Acid (ABA) signaling

response were up-regulated, a pathway that has been widely reported as a key drought

stress response (Daszkowska-Golec, 2016). Among those genes, protein phosphatases

type-2C (PP2Cs), ABA receptors PYRPYR/PYL/RCAR (PYRABACTIN-

RESISTANCE 1/PYRABACTIN RESISTANCE LIKE/REGULATORY

COMPONENT OF ABA RECEPTOR) and SNF1-Related Protein Kinases type 2

(SnRK2s) were identified as DEGs (de Zelicourt et al., 2016). Although PP2Cs are

negative regulators of ABA signalling, an increased relative expression under drought

stress conditions has been reported in other similar studies (Gong et al., 2015; He et al.,

2019; Yang et al., 2018), suggesting that these apparent contrasting effects need to be

further investigated. AREB/ABF transcription factors and MAPKKs (mitogen activated

protein kinase kinase) were also activated as a response to ABA signaling, which leads

to stomatal closure, one of the most important drought responses (de Zelicourt et al.,

2016; Wu et al., 2014). *Solanum dasyphyllum* displayed a wide variety of response

mechanisms along with the ABA pathway. These included galactinol synthase and

transferases related genes, which have been reported to improve drought tolerance

(Selvaraj et al., 2017). Also, zeatin biosynthesis was down-regulated, in particular the

cytokinin signaling repressors A-type ARABIDOPSIS RESPONSE REGULATORS

(ARRs), which have been reported to negatively regulate by drought stress, promoting

cell division in meristems (Le et al., 2012; S. M. Li et al., 2021; Nishiyama et al., 2011).

In addition, GIGANTEA (GI) protein synthesis was activated, which is a regulator in

the circadian rhythm plant pathway and improves drought tolerance (Baek et al., 2020).

Finally, phenylpropanoid biosynthesis pathway was detected, which exhibits different

important roles in the regulation under abiotic stress conditions (Sharma et al., 2019).

On the other hand, *S. melongena* showed different drought response pathways,

including the carotenoid biosynthesis, which has been reported to have a similar

regulation in *S. tuberosum* (Gong et al., 2015), the inactivation of porphyrin,

chlorophyll metabolism and photosynthesis pathways as a consequence of the osmotic

stress (Reddy et al., 2004). Furthermore, the regulation of flavonoid biosynthesis, which

has an important role in coping with environmental stress (Sharma et al., 2019), the

expression of plant glutathione transferases (GSTs), which has been reported to be

involved in responses to biotic and abiotic stress (Dixon et al., 2010), and the synthesis

of the stress signaling molecule, such as jasmonic acid (JA) by the metabolism of α-

Linolenic acid (Zi et al., 2022) were linked to osmotic stress. When the plants are more

adult and under a more intense osmotic stress, ABA signaling response leads to

stomatal closure and to the down regulation of small auxin up-regulated RNA (SAUR)

genes, which induce plant growth (Stortenbeker and Bemer, 2019). In our study, a

common response as stress adaptation has been observed, including ABA signaling

response and inhibition of plant growth.

5. Conclusions

The present work provides an overview of the osmotic stress response at the transcriptomic level of cultivated eggplant (*S. melongena*) and its drought-tolerant wild relative *S. dasyphyllum*. We have found that osmotic potential and plant phenological stage play a crucial role in the response, which is increased when the exposure time was longer and osmotic stress was more intense. Our data showed that response mechanisms at the gene expression level were very wide-ranging, including transcription factors, phytohormones, osmoprotectants and protein kinases, being ABA response signaling an important pathway. Clear differences observed between the two species in the response to osmotic stress and overall gene expression pattern confirmed that *S. dasyphyllum* is a potential source for breeding to drought tolerance in eggplant. Overall, our work provided insights into the gene expression mechanisms of tolerance to osmotic stress in eggplant and its wild relative *S. dasyphyllum*, which is of great relevance in the improvement of drought tolerance of cultivated eggplant.

Supplementary information

Supplementary Table S1. Data quality summary of samples of experiment 1 and 2, after 0, 0.5 and 2 h of PEG stress of *S. dasyphyllum* (DAS) and *S. melongena* (MEL).

Supplementary Table S2. Gene ontology (GO) terms enrichment and regulation, ID, description and transcription factor family of DEGs in *S. dasyphyllum* after 0.5 (T0.5) (A) and 2 h (T2) versus 0h of PEG stress and *S. melongena* (M) after 0.5 (T0.5) and 2 h (T2) compared with 0h of PEG stress experiment 2 (Ex_2).

Supplementary Table S3. Significant Kyoto Encyclopedia of Genes and Genomes (KEGG) enriched pathways, its ID of tomato database, and regulation ID, description

and transcription factor family of DEGs in *S. dasyphyllum* (D) and *S. melongena* (M)

after 0.5 h (T0.5) and 2 h (T2) of PEG stress in experiments 1 (Ex_1) and 2 (Ex_2).

Acknowledgements

This work was funded by MCIN/AEI/ 10.13039/501100011033/ and by "ERDF A way

of making Europe" through grant RTI-2018–094592-B-I00. The Spanish Ministerio de

Ciencia e Innovación, Agencia Estatal de Investigación and Fondo Social Europeo,

funded a predoctoral fellowship to Gloria Villanueva (PRE2019-103375). Pietro

Gramazio is grateful to Spanish Ministerio de Ciencia e Innovación for a post-doctoral

grant (FJC2019-038921-I) funded by (MCIN/AEI /10.13039/501100011033).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

References

Anders, S., Huber, W., 2010. Differential expression analysis for sequence count data.

Nature Precedings. 11, 1-1. https://doi.org/10.1186/gb-2010-11-10-r106

Baek, D., Kim, W.Y., Cha, J.Y., Park, H.J., Shin, G., Park, J., Lim, C.J., Chun, H.J., Li,

N., Kim, D.H., Lee, S.Y., Pardo, J.M., Kim, M.C., Yun, D.J., 2020. The

GIGANTEA-ENHANCED em LEVEL complex enhances drought tolerance via

regulation of abscisic acid synthesis. Plant Physiol. 184, 443–458.

https://doi.org/10.1104/PP.20.00779

Barchi, L., Pietrella, M., Venturini, L., Minio, A., Toppino, L., Acquadro, A., Andolfo,

G., Aprea, G., Avanzato, C., Bassolino, L., Comino, C., Molin, A.D., Ferrarini, A.,

Maor, L.C., Portis, E., Reyes-Chin-Wo, S., Rinaldi, R., Sala, T., Scaglione, D.,

Sonawane, P., Tononi, P., Almekias-Siegl, E., Zago, E., Ercolano, M.R., Aharoni,

A., Delledonne, M., Giuliano, G., Lanteri, S., Rotino, G.L., 2019. A chromosome-

anchored eggplant genome sequence reveals key events in Solanaceae evolution.

Sci. Rep. 9, 11769. https://doi.org/10.1038/s41598-019-47985-w

Bardou, P., Mariette, J., Escudié, F., Djemiel, C., Klopp, C., 2014. jvenn: an interactive

Venn diagram viewer. BMC Bioinformatics 15, 293. https://doi:10.1186/1471-

2105-15-293

Behboudian, M.H., 1977. Responses of eggplant to drought. I. Plant water balance. Sci.

Hortic. (Amsterdam). 7, 303–310. https://doi.org/10.1016/0304-4238(77)90002-4

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J. R. Stat. Soc. Ser. B 57, 289–300.

https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bouzroud, S., Gouiaa, S., Hu, N., Bernadac, A., Mila, I., Bendaou, N., Smouni, A.A.,

Bouzayen, M., Zouine, M., 2018. Auxin response factors (ARFs) are potential

mediators of auxin action in tomato response to biotic and abiotic stress (*Solanum

lycopersicum*). PLOS ONE 13, e0193517.

https://doi.org/10.1371/journal.pone.0193517

Bukenya, Z.R., Carasco, J.F., 1994. Biosystematic study of *Solanum macrocarpon—S.

dasyphyllum* complex in Uganda and relations with *Solanum linnaeanum*. East

African Agric. For. J. 59, 187–204.

https://doi.org/10.1080/00128325.1994.11663195

Chen, F., Hu, Y., Vannozzi, A., Wu, K., Cai, H., Qin, Y., Mullis, A., Lin, Z., Zhang, L.,

2017. The WRKY transcription factor family in model plants and crops. CRC.

Crit. Rev. Plant Sci. 36, 311–335. https://doi.org/10.1080/07352689.2018.1441103

Chen, L., Meng, J., Luan, Y., 2019. miR1916 plays a role as a negative regulator in

drought stress resistance in tomato and tobacco. Biochem. Biophys. Res. Commun.

508, 597–602. https://doi.org/10.1016/j.bbrc.2018.11.165

Claeys, H., Inzé, D., 2013. The agony of choice: How plants balance growth and

survival under water-limiting conditions. Plant Physiol. 162, 1768–1779.

https://doi.org/10.1104/pp.113.220921

Cook, B.I., Mankin, J.S., Anchukaitis, K.J., 2018. Climate change and drought: from

past to future. Curr. Clim. Chang. Reports 4, 164–179.

https://doi.org/10.1007/s40641-018-0093-2

Daszkowska-Golec, A., 2016. The role of abscisic acid in drought stress: How ABA

helps plants to cope with drought stress, in: Drought Stress Tolerance in Plants,

Vol 2. Springer International Publishing, pp. 123–186.

de Zelicourt, A., Colcombet, J., Hirt, H., 2016. The role of MAPK modules and ABA

during abiotic stress signaling. Trends Plant Sci. 21, 677–685.

https://doi.org/10.1016/j.tplants.2016.04.004

Delfin, E.F., Drobnitch, S.T., Comas, L.H., 2021. Plant strategies for maximizing

growth during water stress and subsequent recovery in *Solanum melongena* L.

(eggplant). PLOS ONE 16, e0256342.

https://doi.org/10.1371/journal.pone.0256342

Díaz-Pérez, J.C., Eaton, T.E., 2015. Eggplant (*Solanum melongena* L.) Plant growth and

fruit yield as affected by drip irrigation rate. HortScience 50, 1709–1714.

https://doi.org/10.21273/hortsci.50.11.1709

Dixon, D.P., Skipsey, M., Edwards, R., 2010. Roles for glutathione transferases in plant

secondary metabolism. Phytochemistry 71, 338–350.

https://doi.org/10.1016/j.phytochem.2009.12.012

FAO, 2020. FAOSTAT database collections. Available online at:

http://www.fao.org/faostat/en/#home, Accessed date: 11 April 2022.

Fu, Q.S., Yang, R.C., Wang, H.S., Zhao, B., Zhou, C.L., Ren, S.X., Guo, Y.D., 2013.

Leaf morphological and ultrastructural performance of eggplant (*Solanum*

melongena L.) in response to water stress. Photosynthetica 51, 109–114.

https://doi.org/10.1007/s11099-013-0005-6

Gao, Y., Gao, S., Xiong, C., Yu, G., Chang, J., Ye, Z., Yang, C., 2015. Comprehensive

analysis and expression profile of the homeodomain leucine zipper IV transcription

factor family in tomato. Plant Physiol. Biochem. 96, 141–153.

https://doi.org/10.1016/j.plaphy.2015.07.025

Gong, L., Zhang, H., Gan, X., Zhang, L., Chen, Y., Nie, F., Shi, L., Li, M., Guo, Z.,

Zhang, G., Song, Y., 2015. Transcriptome profiling of the potato (*Solanum*

tuberosum L.) plant under drought stress and water-stimulus conditions. PLOS

ONE 10, e0128041. https://doi.org/10.1371/journal.pone.0128041

Haswell, E.S., Verslues, P.E., 2015. The ongoing search for the molecular basis of plant

osmosensing. J. Gen. Physiol. 145, 389–394.

https://doi.org/10.1085/jgp.201411295

He, Z., Wu, J., Sun, X., Dai, M., 2019. The Maize Clade A PP2C Phosphatases play

critical roles in multiple abiotic stress responses. Int. J. Mol. Sci. 20, 3573.

https://doi.org/10.3390/ijms20143573

Hoagland, D.R., Arnon, D.I., 1950. The water-culture method for growing plants

without soil. Circ. Calif. Agric. Exp. Stn. 347, 1–32.

Hossain, M.A., Wani, S.H., Bhattacharjee, S., Burritt, D.J., Tran, L.-S.P., 2016. Drought

stress tolerance in plants, Vol 2. Springer International Publishing.

https://doi.org/10.1007/978-3-319-32423-4

Ilyas, M., Nisar, M., Khan, N., Hazrat, A., Khan, A.H., Hayat, K., Fahad, S., Khan, A.,

Ullah, A., 2021. Drought tolerance strategies in plants: a mechanistic approach. J.

Plant Growth Regul. 40, 926–944. https://doi.org/10.1007/s00344-020-10174-5

IPCC, 2022. Intergovernmental panel on climate change. Proceedings of the Sixth

Assesment Report, WGII, Climate change 2022: impacts, adaptation and

vulnerability.

Islam, M.S., Wang, M.H., 2009. Expression of dehydration responsive element-binding

protein-3 (DREB3) under different abiotic stresses in tomato. BMB Rep. 42, 611–

616. https://doi.org/10.5483/BMBRep.2009.42.9.611

Jiao, P., Jiang, Z., Wei, X., Liu, S., Qu, J., Guan, S., Ma, Y., 2022. Overexpression of

the homeobox-leucine zipper protein ATHB-6 improves the drought tolerance of

maize (*Zea mays* L.). Plant Sci. 316, 111159.

https://doi.org/10.1016/j.plantsci.2021.111159

Kido, É.A., Ferreira-Neto, J.R.C., Pandolfi, V., de Melo Souza, A.C., Benko-Iseppon,

A.M., 2016. Drought stress tolerance in plants: insights from transcriptomic

studies, in: Drought Stress Tolerance in Plants, Vol 2. Springer International

Publishing, pp. 153–185.

Kouassi, A.B., Kouassi, K.B.A., Sylla, Z., Plazas, M., Fonseka, R.M., Kouassi, A.,

Fonseka, H., N'guetta, A.S.P., Prohens, J., 2020. Genetic parameters of drought

tolerance for agromorphological traits in eggplant, wild relatives, and interspecific

hybrids. Crop Sci. 61, 55–68. https://doi.org/10.1002/csc2.20250

Kouassi, B., Prohens, J., Gramazio, P., Kouassi, A.B., Vilanova, S., Galán-Ávila, A.,

Herraiz, F.J., Kouassi, A., Seguí-Simarro, J.M., Plazas, M., 2016. Development of

backcross generations and new interspecific hybrid combinations for introgression

breeding in eggplant (*Solanum melongena*). Sci. Hortic. 213, 199–207.

https://doi.org/10.1016/j.scienta.2016.10.039

Le, D.T., Nishiyama, R., Watanabe, Y., Vankova, R., Tanaka, M., Seki, M., Ham, L.H.,

Yamaguchi-Shinozaki, K., Shinozaki, K., Tran, L.S.P., 2012. Identification and

expression analysis of cytokinin metabolic genes in soybean under normal and

drought conditions in relation to cytokinin levels. PLOS ONE 7, e42411.

https://doi.org/10.1371/journal.pone.0042411

Li, D., He, Y.J., Li, S., Shi, S., Li, L., Liu, Y., Chen, H., 2021. Genome-wide

characterization and expression analysis of AP2/ERF genes in eggplant (*Solanum

melongena* L.). Plant Physiol. Biochem. 167, 492–503.

https://doi.org/10.1016/j.plaphy.2021.08.006

Li, S.M., Zheng, H.X., Zhang, X.S., Sui, N., 2021. Cytokinins as central regulators

during plant growth and stress response. Plant Cell Rep. 40, 271–282.

https://doi.org/10.1007/s00299-020-02612-1

Metsalu, T., Vilo, J., 2015. ClustVis: a web tool for visualizing clustering of

multivariate data using Principal Component Analysis and heatmap. Nucleic Acids
Res. 43, 566–570. https://doi.org/10.1093/nar/gkv468

Michel, B.E., Kaufmann, M.R., 1973. The osmotic potential of polyethylene glycol
6000. Plant Physiol. 51, 914–916. https://doi.org/10.1104/pp.51.5.914

Moon, K.B., Ahn, D.J., Park, J.S., Jung, W.Y., Cho, H.S., Kim, H.R., Jeon, J.H., Park,
Y. Il, Kim, H.S., 2018. Transcriptome profiling and characterization of drought-
tolerant potato plant (*Solanum tuberosum* L.). Mol. Cells 41, 979–992.
https://doi.org/10.14348/molcells.2018.0312

Mukherjee, S., Mishra, A., Trenberth, K.E., 2018. Climate change and drought: a
perspective on drought indices. Curr. Clim. Chang. Reports 4, 145–163.
https://doi.org/10.1007/s40641-018-0098-x

Nishiyama, R., Watanabe, Y., Fujita, Y., Le, D.T., Kojima, M., Werner, T., Vankova,
R., Yamaguchi-Shinozaki, K., Shinozaki, K., Kakimoto, T., Sakakibara, H.,
Schmülling, T., Tran, L.S.P., 2011. Analysis of cytokinin mutants and regulation
of cytokinin metabolic genes reveals important regulatory roles of cytokinins in
drought, salt and abscisic acid responses, and abscisic acid biosynthesis. Plant Cell
23, 2169–2183. https://doi.org/10.1105/tpc.111.087395

Osmolovskaya, N., Shumilina, J., Kim, A., Didio, A., Grishina, T., Bilova, T.,
Keltsieva, O.A., Zhukov, V., Tikhonovich, I., Tarakhovskaya, E., Frolov, A.,
Wessjohann, L.A., 2018. Methodology of drought stress research: Experimental
setup and physiological characterization. Int. J. Mol. Sci. 19, 4089.
https://doi.org/10.3390/ijms19124089

Perotti, M.F., Ribone, P.A., Chan, R.L., 2017. Plant transcription factors from the
homeodomain-leucine zipper family I. Role in development and stress responses.

IUBMB Life 69, 280–289. https://doi.org/10.1002/iub.1619

Plazas, M., Andújar, I., Vilanova, S., Gramazio, P., Javier Herraiz, F., Prohens, J., 2014.

Conventional and phenomics characterization provides insight into the diversity

and relationships of hypervariable scarlet (*Solanum aethiopicum* L.) and gboma (*S.*

*macrocarpo*n L.) eggplant complexes. Front. Plant Sci. 5, 318.

https://doi.org/10.3389/fpls.2014.00318

Plazas, M., González-Orenga, S., Nguyen, H.T., Morar, I.M., Fita, A., Boscaiu, M.,

Prohens, J., Vicente, O., 2022. Growth and antioxidant responses triggered by

water stress in wild relatives of eggplant. Sci. Hortic. 293, 110685.

https://doi.org/10.1016/j.scienta.2021.110685

Plazas, M., Nguyen, H.T., González-Orenga, S., Fita, A., Vicente, O., Prohens, J.,

Boscaiu, M., 2019. Comparative analysis of the responses to water stress in

eggplant (*Solanum melongena*) cultivars. Plant Physiol. Biochem. 143, 72–82.

https://doi.org/10.1016/j.plaphy.2019.08.031

Plazas, M., Vilanova, S., Gramazio, P., Rodríguez-Burruezo, A., Fita, A., Herraiz, F.J.,

Ranil, R., Fonseka, R., Niran, L., Fonseka, H., Kouassi, B., Kouassi, A., Kouassi,

A., Prohens, J., 2016. Interspecific hybridization between eggplant and wild

relatives from different genepools. J. Am. Soc. Hortic. Sci. 141, 34–44.

https://doi.org/10.21273/jashs.141.1.34

Prohens, J., Gramazio, P., Plazas, M., Dempewolf, H., Kilian, B., Díez, M.J., Fita, A.,

Herraiz, F.J., Rodríguez-Burruezo, A., Soler, S., Knapp, S., Vilanova, S., 2017.

Introgressiomics: a new approach for using crop wild relatives in breeding for

adaptation to climate change. Euphytica 213, 158. https://doi.org/10.1007/s10681-

017-1938-9

638 Ranil, R.H.G., Niran, H.M.L., Plazas, M., Fonseka, R.M., Fonseka, H.H., Vilanova, S.,

639   Andújar, I., Gramazio, P., Fita, A., Prohens, J., 2015. Improving seed germination

640   of the eggplant rootstock *Solanum torvum* by testing multiple factors using an

641   orthogonal array design. Sci. Hortic. 193, 174–181.

642   https://doi.org/10.1016/j.scienta.2015.07.030

643 Reddy, A.R., Chaitanya, K.V., Vivekanandan, M., 2004. Drought-induced responses of

644   photosynthesis and antioxidant metabolism in higher plants. J. Plant Physiol. 161,

645   1189–1202. https://doi.org/10.1016/j.jplph.2004.01.013

646 Renau-Morata, B., Sánchez-Perales, M., Medina, J., Molina, R., Corrales, R., Carrillo,

647   L., Fernández-Nohales, P., Marqués, J., Pollmann, S., Vicente-Carbajosa, J.,

648   Granell, A., Nebauer, S., 2016. Salinity assay in tomato. Bio-protocol. 4, e1215.

649   https://doi.org/10.21769/bioprotoc.1215

650 Riyazuddin, R., Nisha, N., Singh, K., Verma, R., Gupta, R., 2021. Involvement of

651   dehydrin proteins in mitigating the negative effects of drought stress in plants.

652   Plant Cell Rep. https://doi.org/10.1007/s00299-021-02720-6

653 Samarah, N.H., 2016. Understanding How Plants Respond to Drought Stress at the

654   Molecular and Whole Plant Levels, in: Drought Stress Tolerance in Plants, Vol 2.

655   Springer International Publishing, pp. 1–38.

656 Selvaraj, M.G., Ishizaki, T., Valencia, M., Ogawa, S., Dedicova, B., Ogata, T.,

657   Yoshiwara, K., Maruyama, K., Kusano, M., Saito, K., Takahashi, F., Shinozaki,

658   K., Nakashima, K., Ishitani, M., 2017. Overexpression of an Arabidopsis thaliana

659   galactinol synthase gene improves drought tolerance in transgenic rice and

660   increased grain yield in the field. Plant Biotechnol. J. 15, 1465–1477.

661   https://doi.org/10.1111/pbi.12731

Sharma, A., Shahzad, B., Rehman, A., Bhardwaj, R., Landi, M., Zheng, B., 2019.

Response of phenylpropanoid pathway and the role of polyphenols in plants under

abiotic stress. Molecules 24, 1–22. https://doi.org/10.3390/molecules24132452

Shinozaki, K., Yamaguchi-Shinozaki, K., 2007. Gene networks involved in drought

stress response and tolerance. J. Exp. Bot. 58, 221–227.

https://doi.org/10.1093/jxb/erl164

Stark, R., Grzelak, M., Hadfield, J., 2019. RNA sequencing: the teenage years. Nat.

Rev. Genet. 20, 631–656. https://doi.org/10.1038/s41576-019-0150-2

Stortenbeker, N., Bemer, M., 2019. The SAUR gene family: The plant's toolbox for

adaptation of growth and development. J. Exp. Bot. 70, 17–27.

https://doi.org/10.1093/jxb/ery332

Stratilová, B., Kozmon, S., Stratilová, E., Hrmova, M., 2020. Plant xyloglucan

xyloglucosyl transferases and the cell wall structure: Subtle but significant.

Molecules 25, 1–25. https://doi.org/10.3390/molecules25235619

Sun, S., Yu, J.P., Chen, F., Zhao, T.J., Fang, X.H., Li, Y.Q., Sui, S.F., 2008. TINY, a

dehydration-responsive element (DRE)-binding protein-like transcription factor

connecting the DRE- and ethylene-responsive element-mediated signaling

pathways in Arabidopsis. J. Biol. Chem. 283, 6261–6271.

https://doi.org/10.1074/jbc.M706800200

Syfert, M.M., Castañeda-Álvarez, N.P., Khoury, C.K., Särkinen, T., Sosa, C.C.,

Achicanoy, H.A., Bernau, V., Prohens, J., Daunay, M.-C., Knapp, S., 2016. Crop

wild relatives of the brinjal eggplant (*Solanum melongena*): Poorly represented in

genebanks and many species at risk of extinction. Am. J. Bot. 103, 635–651.

https://doi.org/10.3732/ajb.1500539

Tenhaken, R., 2015. Cell wall remodeling under abiotic stress. Front. Plant Sci. 5, 1–9.

https://doi.org/10.3389/fpls.2014.00771

Vorontsova, M.S., Knapp, S., 2016. A revision of the spiny solanums, *Solanum*

subgenus *Leptostemonum* (Solanaceae) in Africa and Madagascar. Syst. Bot.

Monogr. 99, 1–436. https://doi.org/https://doi.org/10.5519/0055154

Vorontsova, M.S., Stern, S., Bohs, L., Knapp, S., 2013. African spiny *Solanum*

(subgenus *Leptostemonum*, solanaceae): A thorny phylogenetic tangle. Bot. J.

Linn. Soc. 173, 176–193. https://doi.org/10.1111/boj.12053

Wang, J., Li, C., Li, L., Reynolds, M., Mao, X., Jing, R., 2021. Exploitation of drought

tolerance-related genes for crop improvement. Int. J. Mol. Sci. 22.

https://doi.org/10.3390/ijms221910265

Wang, R., Zhao, P., Kong, N., Lu, R., Pei, Y., Huang, C., Ma, H., Chen, Q., 2018.

Genome-wide identification and characterization of the Potato bHLH Transcription

factor family. Genes (Basel). 9, 54. https://doi.org/10.3390/genes9010054

Wu, J., Wang, J., Pan, C., Guan, X., Wang, Y., Liu, S., He, Y., Chen, J., Chen, L., Lu,

G., 2014. Genome-wide identification of MAPKK and MAPKKK gene families in

tomato and transcriptional profiling analysis during development and stress

response. PLOS ONE 9, e103032. https://doi.org/10.1371/journal.pone.0103032

Yang, Q., Liu, K., Niu, X., Wang, Q., Wan, Y., Yang, F., Li, G., Wang, Y., Wang, R.,

2018. Genome-wide identification of PP2C genes and their expression profiling in

response to drought and cold stresses in *Medicago truncatula*. Sci. Rep. 8, 12841.

https://doi.org/10.1038/s41598-018-29627-9

Yang, X., Liu, J., Xu, J., Duan, S., Wang, Q., Li, G., Jin, L., 2019. Transcriptome

profiling reveals effects of drought stress on gene expression in diploid potato

genotype P3-198. Int. J. Mol. Sci. 20, 852. https://doi.org/10.3390/ijms20040852

Zhang, Z., Chen, J., Liang, C., Liu, F., Hou, X., Zou, X., 2020. Genome-wide

identification and characterization of the bHLH transcription factor family in

pepper (*Capsicum annuum* L.). Front. Genet. 11, 570156.

https://doi.org/10.3389/fgene.2020.570156

Zhu, M., Meng, X., Cai, J., Li, G., Dong, T., Li, Z., 2018. Basic leucine zipper

transcription factor SlbZIP1 mediates salt and drought stress tolerance in tomato.

BMC Plant Biol. 18, 83. https://doi.org/10.1186/s12870-018-1299-0

Zhuang, J., Zhang, J., Hou, X.L., Wang, F., Xiong, A.S., 2014. Transcriptomic,

proteomic, metabolomic and functional genomic approaches for the study of

abiotic stress in vegetable crops. CRC. Crit. Rev. Plant Sci. 33, 225–237.

https://doi.org/10.1080/07352689.2014.870420

Zi, X., Zhou, S., Wu, B., 2022. Alpha-Linolenic Acid Mediates Diverse Drought

Responses in Maize (*Zea mays* L.) at Seedling and Flowering Stages. Molecules

27, 771. https://doi.org/10.3390/molecules27030771

Ex_1

MEL

DAS

Ex_2

MEL

DAS

T0   T0.5   T2