**CineFinch: An animated female zebra finch for studying courtship interactions**

Nikhil Phaniraj[1], Sanjana Joshi[1], Pradeepkumar Trimbake[1], Aditya Pujari[1,3], Samyuktha Ramadurai[1,3],

Shikha Kalra[1,3], Nikhil Ratnaparkhi[2] and Raghav Rajan[1]*

1.  Division of Biology, Indian Institute of Science Education and Research Pune, Pune,

Maharashtra, 411008 INDIA.

2.  Birla Institute of Science and Technology, Pilani, Rajasthan, 333031, INDIA.

3.  These authors contributed equally

* Correspondence should be addressed to Raghav Rajan. Email: raghav@iiserpune.ac.in

**Running Head:** Animated female zebra finch

**Key words:** Bird song, courtship displays, video interactions, animations

## 11 SUMMARY STATEMENT

We develop and test an animation of a female zebra finch to study song and courtship interactions in

the male zebra finch.

**ABSTRACT**

Dummies, videos and computer animations have been used extensively in animal behaviour to study
simple social interactions. These methods allow complete control of one interacting animal, making it
possible to test hypotheses about the significance and relevance of different elements of animal
displays. Recent studies have demonstrated the potential of videos and interactive displays for studying
more complex courtship interactions in the zebra finch, a well-studied songbird. Here, we extended
these techniques by developing an animated female zebra finch and showed that ~40% of male zebra
finches (n=5/12) sing to this animation. To study real-time social interactions, we developed two
possible methods for closed loop control of animations; (1) an arduino based system to initiate
videos/animations based on perch hops and (2) a video game engine based system to change
animations. Overall, our results provide an important tool for understanding the dynamics of complex
social interactions during courtship.

## INTRODUCTION

Complex behavioural displays are used by animals to communicate with each other (Bradbury and Vehrencamp, 2011). These behavioural displays carry information necessary for successful communication and often, have multiple different components that are produced together. For example, the courtship dance of the blue-capped cordon bleu, a songbird, involves multiple rapid foot and head movements that occur just before song, a vocal signal (Ota et al., 2015). Are all of these different components (foot, head movements and vocal signal) important for successful courtship? Do individual components of the display carry information about courtship potential or is courtship potential signalled only by presence of the entire display? Answering these questions requires complete control over one of the interacting animals to ensure that different components can be produced independent of each other. Such control is provided by robotic dummies, videos and computer animations and animal behaviour studies have a rich tradition of using such stimuli (Tinbergen, 1948; Woo and Rieucau, 2011). For example, Patricelli and colleagues used robotic female satin bowerbirds in artificially staged courtship interactions to demonstrate that male satin bowerbirds regulate the intensity of their courtship displays based on female responses (Patricelli et al., 2006). Van Dyk and Evans used computer animations of Jacky dragon lizards to understand the dynamics of aggressive encounters and showed that lizards use multiple signals to assess the level of aggression (Van Dyk and Evans, 2008).

Recent studies have extended the use of dummies and videos to understand more complex interactions like courtship in songbirds. In the zebra finch, a well-studied songbird native to Australia, courtship involves a song and a dance by the male (Sossinka and Böhner, 1980; Ullrich et al., 2016; Zann, 1996). Females also respond with vocalizations and tail-quivering displays (Zann, 1996). Male zebra finches sing to taxidermically stuffed female zebra finches and to videos of female zebra finches and the characteristics of these songs are highly similar to courtship song directed at a live female bird (Bischof et al., 1981; Galoch and Bischof, 2007; James et al., 2019). In addition to courtship interactions, male zebra finches also interact with juvenile zebra finches during tutoring sessions. Juveniles learn songs more accurately from a live tutor than from song playbacks from a speaker (Derégnaucourt et al., 2013). This suggests that the visual stimulus of the tutor and possibly social interactions with a tutor are also important for accurate learning. While the mere presence of a visual tutor, in the form of a video, is not sufficient to enhance learning (Varkevisser et al., 2022a), a recent study showed that a robotic zebra finch that vocally interacts with juvenile birds does enhance learning (Araguas et al., 2022).

Importantly, the robotic zebra finch provided closed-loop vocal interactions, i.e. vocal interactions were

provided only when juveniles interacted with the robot and the timing of these vocal interactions were

comparable to the timing of interactions with live tutors. The robotic zebra finch highlighted the

potential of appropriately manipulatable, artificial stimuli for probing complex social interactions, but

such stimuli have not been tested in the context of courtship.

In addition to robots, animations provide an attractive method to provide closed-loop social

interactions. The advent of fast computer hardware and user-friendly open-source graphics software

has made it much easier to generate and control animations (Stowers et al., 2017). They provide a

complementary approach to robots with considerable flexibility for studying social interactions. Here,

we developed an animation of a female zebra finch and showed that ~40% of male zebra finches tested,

sang to this animation. We also demonstrated two possible ways to interactively control these

animations.

**MATERIALS AND METHODS**

All experimental procedures conducted were approved by IISER Pune's Institute Animal Ethical

Committee (IAEC) and were in accordance with the guidelines of the Committee for the Purpose of

Control and Supervision of Experiments on Animals (CPSCEA), New Delhi. Zebra finches (n=25

males and n=5 females) were procured from a local vendor (n=7) or bred in our colony at IISER Pune

(n=23). Birds bought from an outside vendor were used for experiments only after they had been in our

colony at IISER Pune for more than 30 days. Birds were raised in individual cages (120 cm x 50 cm x

50 cm cage, up to 6 birds per cage) along with other birds of the same gender. Light conditions were

regulated to maintain a 14/10 hour day/night cycle. A*d libitum* access to food and water were provided

at all times, unless otherwise mentioned. All birds were >100 dph at the time of the experiment (males

> 100dph and females > 300dph).

*Experimental Apparatus*

The apparatus consisted of a metal double-cage (46 cm x 23 cm x 23 cm) with each half separated by a

glass slab (23 cm x 23 cm x 0.3 cm). One half housed the subject male bird. During live female trials,

the other half housed a stimulus female bird. During video playback or animation trials, the other half

had a Samsung galaxy tab S4 placed upright lengthwise with its screen touching the glass slab. To

record song produced by the male, we placed a microphone (AKG Acoustics C417PP omnidirectional
condenser microphone) on top of the cage above the male. A camera (GoPro Hero7) was placed outside
the apparatus to record videos of the sessions.

*Experimental subjects*
For the experiments where males were presented with videos of females (video trials), we used 10 male
zebra finches for the 30s trials and 10 males for the 4-minute trials. For the animation experiments, we
used 12 males. 4 males were common across the 30s and 4-minute trials, 1 male was common across
both 4 minute trials and animation trials and 1 bird was common for 30s and 4-minute video trials and
animation trials. For the video experiments, we used 3 female zebra finches for the 30s trials and 2
female zebra finches for the 4-minute trials. For the animation experiments, we used 2 females. One
female was common across 4-minute and 30s video trials and one female was common across 30s
video and animation trials.

*Stimulus videos*
4 minute long videos (2560 x 1440 pixels @60 fps) of a female bird were recorded while the female
interacted with a male bird (see Movie 1 for part of a video). The same experimental setup that was
used for video trials with the male was used for this and the male was positioned just beside the video
camera, on the other side of the glass slab (Fig. 1A). The aspect ratio of the video was fixed at 15.5 cm
x 8.7 cm to ensure that the female bird in the video appeared life-sized.

*Construction of Animations*
Initially, a 13.67s long animation of a female zebra finch was made using Blender 2.90.1
(https://www.blender.org/). This initial animation consisted of 820 frames which were played at 60fps.
To make this, first, a 3D model of a female zebra finch was sculpted (Fig. 2A). Next, colours from a
photograph of a female zebra finch were extracted and used to paint the skin of the 3D model (Fig. 2B).
A feathery texture was added. 2 core bones (grey in Fig. 2C) and 9 structural bones (yellow in Fig. 2C)
were rigged into the body and 5 control bones (blue in Fig. 2C) were placed outside that connected to a
single bone or multiple distant bones for synchronized body movements and provided for puppeteer-
like control of the zebra finch (Fig. 2A-2D). One of the control bones was placed in front of the beak
and the beak bone programmed to always point towards the beak control bone. The beak-head joint

was kept rigid whereas the head bone was free to rotate around the neck bone giving the head-neck
joint 3 degrees of freedom. The second control bone was placed behind the tail and the tail bone
programmed to always point to the tail control bone. The tail bone could rotate around the joint in both
horizontal and vertical plane but not around its long axis, giving it 2 degrees of freedom. 2 of the
control bones were placed in front of the claw and the claw bones were programmed to always point to
the claw control bone. 2 additional bones were placed behind the knee, outside the body, to mark
endpoints for inverse kinematics of the knee (red in Fig. 2C). These two external bones prevented the
knee joints from displacing beyond the marked endpoints. The ankle joint was kept rigid whereas the
knee and the pelvis were allowed restricted rotation in only the vertical plane giving them 1 degree of
freedom. The 5th control bone was the master control bone (deep blue in Fig. 2C) placed directly
below the centre of mass of the bird and this bone connected the other 4 control bones and allowed for
coordinated movements of the two legs, tail, and head.

The initial animation started with the female zebra finch hopping in from the left, performing 2 head
turns (towards its right and its left), hopping to the centre, rotating towards the viewer, bending, and tail
quivering (see Fig. 2E-2H for frames from the animation). After the tail quivering, the female was
made to go back to the initial posture, hop to the right, perform 2 head turns (towards its right and its
left), and hop to the right again – out of view. The velocity of hops, head turns and tail quivering were
roughly similar to those of the female in one of the 30s responsive videos. We only matched the
velocity qualitatively and did not reproduce exact statistics of movements. This initial animation was
duplicated, laterally inverted and stitched to the end of the original animation to make the female enter
from the left, exit from the right, then enter from the right, and exit from the left. This entire animation
sequence, now 1640 frames (27.33s) long was again duplicated thrice and the 4 parts stitched together.
A 10s clip showing an image of an empty cage was added to the beginning and the entire animation
video was ~ 2mins (119.33s) long. A 5s long audio clip with 3 female calls was added to the first 5s of
the animation (see Movie 2). The final animation video was played with the audio on (Call animation)
or muted (No call animation). A color control animation was made where the bird in the animation was
entirely painted with deep blue and the feathery texture removed (see Movie 2). Everything else was
kept the same.

*General experimental procedures*
All birds used for experiments were housed singly in small cages (23 cm x 23 cm x 23 cm) starting a
few days before the start of the experiment and for the entire duration of the experiment, unless

otherwise mentioned. For 30s trials, pairs of male subject birds were acclimatized to the setup by
housing them in the apparatus – one bird on each half of the setup for 48 hours – 2 weeks before the
start of the experimental trials, and again for 48 hours, before the time of the experimental trial. All
experimental trials were conducted in sound-attenuation enclosures (Newtech Engineering Systems,
Bengaluru or custom-made enclosures). For all experiments, songs were recorded with a microphone
placed at the top of the cage. Signals from the microphone were amplified using a mixer (Behringer
XENYX 802) and then digitized on a computer system at a sampling rate of 44,100 Hz using a custom-
written Python based software. Undirected songs were recorded over a period of 24 hours, with birds
housed singly in a cage (23 cm x 23 cm x 23 cm) in the same sound isolation box. 1 of the 7 birds
never sang undirected songs.

*Video presentation procedures*
Across all trial types, videos were played manually by the experimenter, without the audio, as a
window on the screen. The video window overlapped an image of an empty cage in the background.
The tablet was placed in its position as soon as the video was started. Once the video ended, the video
window disappeared showing only the image of the empty cage on the screen. Animations were also
converted into videos as described above and played using the same procedure.

*Short duration experimental trials (30s Video trials)*
For the 30s video trials, experimental trials were conducted over a period of 10 days with one bird
going through the experiment each day. Two, 30s long, silent, videos of each female were chosen as
stimuli; (1) a "responsive" video where the female showed tail quivering and (2) an "unresponsive"
video where the female did not show tail quivering. These videos were cut out of the 4 minute long
videos of female birds that were recorded as described above. Along with the "responsive" and
"unresponsive" videos, we also presented a live female as a third stimulus. Each of the three stimuli
was repeated thrice and the 9 stimuli were presented to the male in random order separated by 5
minutes (Fig. 1B).

*Long duration experimental trials (4-minute video trials)*
For the 4-minute video trials, we used the 4 minute long videos recorded as described above.
Experimental trials were conducted over multiple days. On a given day, male birds were briefly

checked with ~10s exposure to the tablet to see if they sang to the video. Birds went through the trial
only if they sang to the video during the brief exposure (n=3/10 birds did not sing to the video on their
first exposure and were not used further). During the actual trial, the subject bird was exposed to either
the live female or the video for 4 minutes. For the live female condition, one of the two females was
randomly chosen as the stimulus. For the video playback condition, one of the two videos (one of each
female stimulus) was chosen to be played. For each male, only one type of trial was conducted on a
given day. This procedure was repeated for each male on separate days until we obtained atleast 6 song
bouts for each trial type (median of 7 days per male; range: 6-18).

*Animation trials*
For the animation trials, birds were tested with a single exposure to the 2 minute animation of the
female zebra finch without audio. Only 5/12 birds tested sang to this animation and these 5 birds were
chosen for further experiments. Experimental trials consisted of 3 days of experiments with each day
consisting of 5 types of trials (live female, video of female, animation with call, animation without call
and colour control animation) in a random sequence. Each trial lasted 2 minutes and trials were
separated by 5 minutes.

*Song Analysis*
Songs were analysed as described earlier (Kalra et al., 2021; Rajan and Doupe, 2013; Rao et al., 2019),
using custom-written scripts on MATLAB (MathWorks). Briefly, songs were first segmented into
syllables based on an amplitude threshold. Syllables with inter-syllable intervals < 5 ms were merged
into one syllable and syllables shorter than 10s were discarded. Then, syllables were manually labelled
based on their appearance on a spectrogram; similar looking syllables were given the same label. The
sequence of syllables that repeated in a bird's song was identified as a motif (Sossinka and Böhner,
1980; Zann, 1996). Short syllables that were repeated a variable number of times at the beginning of a
bout were considered as introductory notes (Kalra et al., 2021; Rajan, 2018; Rajan and Doupe, 2013;
Rao et al., 2019). The number of introductory notes at the beginning of a bout was quantified by
starting with the introductory note immediately preceding the first syllable of the bout and counting
backwards until either (1) the previous note was not an introductory note or (2) there was a silence >
500 ms (Kalra et al., 2021; Kao et al., 2005; Rajan and Doupe, 2013; Rao et al., 2019; Sossinka and
Böhner, 1980). Song bouts were defined as groups of vocalizations that contained at least one motif

syllable and were separated from other such groups by more than 2 s of silence. All song bouts sung by

birds towards live females and video/animation were considered for analysis as independent song bouts

and were combined with all other song bouts across sessions/days. Of all the undirected bouts sung by

a bird, 15 were randomly chosen for analysis for the 4 minute trials (median: 15; range: 14-16) and 4

were chosen randomly for analysis for the 30s trials (median: 4; range: 2-6). The number of undirected

bouts to analyse was chosen to be comparable to the number of song bouts produced by the male in

response to the live female and video trials.

*Tutoring of birds with videos*

Juveniles used for tutoring were isolated as described earlier (Kalra et al., 2021). Briefly, fathers were

removed from the nest around 12 days post hatching (range: 9-13) and juveniles were housed along

with their mother until they could feed on their own (range: ~35-40 days post hatch). Following this,

individual juveniles were housed singly and tutoring began on ~37dph (range: ~ 36-46 dph). Tutoring

was done in a cage that contained two perches; one was active and associated with playback of the tutor

song as described below and one perch was a dummy. The active perch had an IR beam running across

it, which was monitored by an Arduino board (https://www.arduino.cc/). When the bird jumped onto

the perch a break in the IR beam was detected by the Arduino-MATLAB interface. This triggered the

tablet screen to play the video of the male bird along with audio playback of the tutor song.

Specifically, an IR-beam break initiated a command to virtually "tap" on a specific coordinate on the

tablet screen. This "tap" opened the video in a VLC window of whose size was chosen to ensure that

the bird in the video was of a realistic size. The latency of the perch-hop to display loop was generally

<1 sec but we noticed that it was longer for the first few hops in a session. After the video ended, the

display was reset to an image of the far end of the cage as it would have appeared if the tablet were not

present. This was also used as the initial image before tutoring began. Tutoring was done for 1-2

sessions per day (median number of tutoring sessions across birds = 45; range = 24-51; some days, we

did not conduct tutoring). In each session, we limited the number of possible song playbacks to 20 as

too much exposure to song has been shown to result in poor imitation (Tchernichovski et al., 1999).

Each session lasted until the bird had exhausted the quota of 20 playbacks. If the bird did not trigger 20

playbacks, then the session was terminated after 1 hour. Passive playbacks were used occassionally to

prompt the bird to hop on the perch. Playback songs consisted of introductory notes (INs) followed by

2 motifs (n=3 were tutored with 2 INs and n=3 were tutored with 7 INs). At ~70 dph (range: 63-79

dph), we switched the number of INs for the tutor song and tutoring with the changed song continued
till 90 dph. These were done to determine if the number of INs can be changed during the course of
learning and are part of another ongoing study. However, for the purpose of this study, we only
considered the accuracy of imitation of the motif and this motif was not changed during the entire
duration of tutoring.

Song similarity was calculated using 5 example motifs recorded from these birds at ~90 dph (range: 90-
100 dph). Similarity calculations were done using Sound Analysis Pro (SAP) (Tchernichovski et al.,
2000) as described earlier (Kalra et al., 2021). Briefly, asymmetric time-course similarity
measurements were made between tutor song and 5 example motifs and the average of these similarity
values was taken as a measure of similarity to the tutor song. For comparison with birds tutored without
videos, we used birds tutored at IISER Pune for a different study (n=10) (Kalra et al., 2021). These
birds were tutored with 2 motifs without INs using active playback methods.

*Closed-loop animations*

The animated female zebra finch described earlier was used in conjunction with a video game engine to
create interactive animations. First, animations were added to the female model using keyframing, a
technique used in blender. Animations included movements like hops, neck movements, body rotation,
tail quivering, and panting. The animation also included female calls (3 calls per stimulus). The
animated female was then exported in FBX (filmbox) format. FBX format was chosen for
compatibility with the game development software, Unreal Game engine 4
(https://www.unrealengine.com/en-US). This model was imported into Unreal engine 4 (version
4.27.2). Unreal engine 4 was used to create a simple game for generating potential stimuli. First, we
created a 1D blend space, i.e. an axis that allowed us to gradually blend a stationary animated female
zebra finch into a hopping, tail-quivering zebra finch. Then, key-presses were associated with different
positions along this axis, thereby allowing us to control the speed of hopping and tail-quivering. The
key-press associations were made within the code of the game engine (Fig. S3) and such stimuli
generated from the game would allow interactive-control of the animation in real-time.

*Statistical analysis*

We did not perform any apriori sample size calculations. All statistical analyses was done with R.
Sample sizes used were comparable to other studies (Ikebuchi and Okanoya, 1999; James et al., 2019;
Takahasi et al., 2005) Generalized Linear Mixed Models (GLMMs) were used to statistically analyse
the data. We considered each song bout as a single unit and considered all song bouts across trial types
for statistical analyses. Response variables were fit using the appropriate distributions; bouts per
presentation and number of INs were count variables and were fit using a Poisson distribution, bout
durations were skewed towards lower duration values, so we log transformed these values and a log-
linked Gaussian distribution was used. Bird identity was a random effect and stimulus type was a fixed
effect for all analyses. Day Number, Presentation number and bout number (all ordinals) were included
as fixed terms while analyzing the number of INs and bout durations in the animation trials. Only day
number and presentation number were included as fixed terms (along with stimulus type) for analyzing
the number of bouts per stimulus for animation trials. Only presentation number and stimulus type were
fixed terms for analyzing the number of bouts per stimulus for the 30 s video trials. For the rest,
stimulus type was the only fixed term (day, presentation, and bout numbers could not be included when
comparisons with undirected songs were made). Type II Wald Chi Sq. tests were used to determine
statistical significance followed by pairwise comparisons between stimulus type using estimated
marginal means with Holm correction. To compare the log ratio of time spent singing across different
conditions, and to analyze the number of bouts per stimulus in the 4-minute video trials, we used a
Kruskal-Wallis test. An alpha level of 0.05 was considered for all statistical analyses.

**RESULTS**

*Male zebra finches sang short song bouts to 30s videos of female zebra finches*

Our goal was to make an animated female zebra finch that could be used for understanding social
interactions. We first tested our display systems by examining responses of male zebra finches to
videos of female zebra finches. We used two different videos (see Fig. 1A and Methods for video
recording procedures) of the same female: (1) a "responsive" video where the female made tail-
quivering movements and (2) an "unresponsive" video without tail-quivering movements (Movie 1).
Individual males were presented with 3 presentations each of the two types of videos and 3
presentations of a live female in random order (Fig. 1B; see Methods for details). Male birds sang
comparable number of songs to videos of female birds and to live females (Fig. 1C, 1D, Movie 1, Fig.
S1A, Fig. S2A). As reported earlier (James et al., 2019), song bouts directed to either of the videos

were considerably shorter than song bouts directed to a live female (Fig. 1E). Songs directed towards live females and songs directed towards videos of females were preceded by greater number of introductory notes as compared to "undirected" songs produced when the bird was alone (Fig. 1F), indicating the "directed" nature of songs sung to the videos (Sossinka and Böhner, 1980). Overall, these results showed that males sing "directed" songs to videos of females and singing behaviour was not influenced by responsiveness of the female.

*Longer duration videos elicited longer song bouts*

To test whether longer stimulus durations elicit longer song bouts, we next presented males with 4 minute videos consisting of both "responsive" and "unresponsive" segments. Each day, individual males were presented with either a video or a live female each day (Fig. 1G) and trials were conducted for multiple days in a random order until we obtained atleast 6 song bouts in both conditions (see Methods for details). Birds sang comparable number of song bouts to the live female and to the videos (Fig. 1H, 1I, S1C). Song bouts to the live female were still longer (Fig. 1J), but the difference in length of song bouts to videos and live females was smaller with longer video durations (Fig. S1C, S2C). This was largely a result of more singing after the first 30s (Fig. S1C). Songs directed to the 4 minute videos were also preceded by higher number of introductory notes compared to undirected songs (Fig. 1K). Overall, these results suggest that video duration plays a role in determining song bout duration possibly by giving the bird more time to detect and respond to the presence of a female on the screen.

*Building an animation of a female zebra finch*

To obtain greater control of the behaviour, we next constructed an animation of a female zebra finch. The animation was constructed in Blender by sculpting a 3D model of a female zebra finch, painting the skin with realistic colours and adding a feather texture (Fig. 2A, 2B). Bones were rigged into the body to allow for puppeteer-like control of movement of specific body parts (Fig. 2C; see Methods for details). Finally, the bird was placed in a background of a cage (Fig. 2D). We used this to construct a simple animation that involved the female hopping in, performing a few head turns and quivering its tail. This sequence was repeated multiple times to construct a 2 minute video that began with the image of an empty cage (Fig. 2E – 2H; first 2 minutes of Movie 2; see Methods for full details). This video was played with the 5s of female calls on appearance of the female (Call animation) or without audio (No call animation). As a control, we used a colour-modified version where the entire bird was painted

with deep blue and the feathery texture removed (last 2 minutes of Movie 2). Everything else including all the movements were kept the same.

*Male zebra finches produced courtship songs directed at the animations*

To test the ability of this animation to elicit song, we tested male birds across 3 sessions (1 session/day) using 5 stimuli presented in random order in each session. The five stimuli used were (1) a live female, (2) a video of a female, (3) video of the animated female zebra finch with and without audio (3 and 4) and the (5) colour-modified control animation without audio (Fig. 2I; see Methods for details). Stimulus duration was kept at 2 minutes. We selected birds based on response to a single 2 minute exposure to the animation without calls. 5/12 males sang to this single exposure and were chosen for the 3 sessions. Male birds sang to all stimuli with comparable numbers of song bouts (Fig. 2J, 2K, S1B, S2B). Song bouts directed at the animation (Movie 3) were comparable in duration to song bouts directed at videos (Fig. S2D), but significantly shorter than song bouts directed at live females (Fig. 2L). Songs directed at animations began with a large number of introductory notes confirming the "directed" nature of these songs (Fig. 2M). Interestingly, 3/5 birds also sang to the colour-modified control animations (Fig. 2J, 2K, 2L). These bouts were also preceded by a large number of introductory notes and were comparable in length to bouts directed towards the animation (Fig. 2M). These results demonstrate the potential of using animations to elicit song and suggest that such animations could be used to further study courtship interactions in the zebra finch.

*Methods for closed loop control of the animation*

To use these animations to study courtship interactions, we needed methods to change these animations in real-time, i.e. we needed a closed-loop system. Here, we used two different approaches to produce interactive animations; (1) using an Arduino to control the appearance of a video and (2) using the animation in conjunction with a gaming engine.

In the first approach, we used an Arduino (a single-board microcontroller) based approach to control the appearance of a video for tutoring young birds. Since young birds learn better from a live tutor than from song playback from a speaker (Derégnaucourt et al., 2013), we used active tutoring methods to test whether including a video of a male bird along with song playback increased copying accuracy. Juveniles could hop on a perch to elicit song playback and video playback of a male bird (Fig. 3A; see

Methods for details of procedure). Juveniles tutored by this method (n=6) copied tutor songs with
accuracies similar to juveniles tutored by song playback alone (Fig. 3B shows spectrograms of tutor
song and songs of two juveniles; Fig. 3C; n=6 birds with video and n=10 birds with only audio
playbacks from Kalra et al. (Kalra et al., 2021)). These results suggested that the visual presence of an
adult bird, albeit on a screen, does not improve copying accuracy and is supported by similar results
from a recent study (Varkevisser et al., 2022a).

While, the Arduino based approach was simple and easy to use, real-time control of the animation was
not possible, as latencies from perch hop to video playback were long (mean +/- std. was 0.74s +/-
0.16s). In our second approach, we used a video game engine, Unreal Engine 4 to construct interactive
animations. We imported the female zebra finch from Blender into Unreal Engine 4 and created a 1D
blend space, i.e. an axis that allowed us to gradually blend a stationary animated female zebra finch
into a tail-quivering zebra finch (Fig. 3D, S3; see Methods for details). Different speeds of movement
along the axis were connected to different durations of a "key press" on a regular keyboard. While, we
connected "key-presses" to different states of the animation, it should be possible to use input from a
microphone to control movement of the animation allowing for song-contingent control. Such methods
have been used to make birds change the fundamental frequency of their vocalizations (Tumer and
Brainard, 2007). Overall, these two methods provide the potential for generating closed-loop,
interactive, animations for studying courtship dynamics.

**DISCUSSION**
Here, we demonstrated potential use of animations to study courtship interactions in a songbird, the
zebra finch. Specifically, we first showed that male zebra finches sang to videos of female zebra
finches and song bouts were longer in response to longer videos. Second, we built an animation of a
female zebra finch and showed that 42% of males sang to the animation. Finally, we showed the
potential for real-time control with tools for interactive control of the animation; a feature that is
essential for analysing courtship interactions in real-time. Overall, our results highlight the power of
animations for studying courtship interactions in songbirds and provide a new tool for the same.

**Duration of the video is a factor that influences the amount of song produced**

Artificial stimuli like animations, need to capture important details of the natural stimulus to be
effective. What aspects of a video/animation stimulus are important for eliciting song from a male
zebra finch? Our results show that stimulus duration is an important factor to consider. Male birds sang
longer song bouts when presented with 4 minute videos as compared to 30s videos of females (Fig. 1).
Both in earlier studies (James et al., 2019) and our experiments, videos were played without sounds.
Live females call and so, when presented with a live female, male birds would experience both visual
and auditory stimuli. Such multimodality is known to be essential for learning from a tutor (Varkevisser
et al., 2022b) and multimodal signals are common in the courtship displays of other songbird species
(Mitoyen et al., 2019; Ota et al., 2015). Given the absence of auditory cues in the videos, longer
duration videos may give birds more time to notice the video and decide to sing. Integration of auditory
cues may increase responses to videos. In line with this, a recent study also used audio and visual cues
to successfully allow two birds to interact with each other virtually (Larsen et al., 2022).

**Improving responses to animations**
Our animation successfully elicited songs from only 42% of the birds tested. What can be done to make
more birds respond to animations? One possibility is to make animations more realistic by
incorporating realistic movement parameters by using pose-estimation software like DeepLabCut to
track, quantify movements of live females (Lauer et al., 2022; Mathis et al., 2018). Another possibility
is to use colour corrections based on the bird's visual system. Such colour-corrected videos have been
shown to increase the attention of juvenile birds to videos of tutors (Varkevisser et al., 2022b). In our
own experiments, the response of some birds to colour control animations suggests the possibility that
the colour scale was not realistic enough.

**Outlook for using animations for closed-loop studies of social interactions**
Our animations and the use of these animations in conjunction with video game engines (like Unreal
Engine) provide a novel way to study social interactions in birds. Recent studies have used a robotic
zebra finch for studying social interactions between juvenile birds and a tutor (the robotic bird)
(Araguas et al., 2022). While a robot provides physical interactions that animations cannot provide,
animations can be built and manipulated more easily. Animations, like the one we have developed
would provide a complementary approach to robotic zebra finches to study social interactions using
virtual-reality setups. A recent study used a virtual reality arrangement to show that two zebra finches

can successfully interact with each other virtually (Larsen et al., 2022). Our animations and video game

engines, combined with this setup could facilitate testing and analyses of social interactions in much

greater detail.

**Broader applications of animated animals**

Here, we focused on developing and controlling an animation for studying courtship interactions in

zebra finches. Similar animations with other animal species can be useful for a diverse range of socio-

cognitive and behavioural experiments. For instance, such animations could be used in social learning

experiments to study the process of social learning. Such experiments typically require one animal in a

group to learn a novel task asocially, following which social transmission of this learning can be

studied (Heyes, 1994; Rendell et al., 2010; Van Schaik and Burkart, 2011). However, such asocial

learning can take time. This can be speeded up using videos to teach the first animal a novel task.

Animations could also be used to study interspecies encounters and animals' responses towards

heterospecifics (Gröning and Hochkirch, 2008; Peiman and Robinson, 2010; Sridhar and Guttal, 2018).

In some species, response to heterospecifics is plastic and is observed to vary with heterospecific

individuals (Lehtonen et al., 2010; Magellan, 2020). However whether physical features (size, body

shape, etc.) or behavioural features (aggression, submission, etc.) are used by animals to identify

heterospecifics remains unclear. Animations, such as ours, offer independent control over physical and

behavioural features and would understand interspecies interactions. Overall, our methods will not only

aid the study of animal courtship interactions but will also be extendable to other fields from animal

cognition to ecology.

**SUPPLEMENTARY INFORMATION**

Supplementary information includes 3 supplemental figures and 3 movies.

**DATA ACCESSIBILITY**

All data, videos, animations and files related to animation are available on reasonable request from the

corresponding author (RR – raghav@iiserpune.ac.in).

**COMPETING INTERESTS**

We declare we have no competing interests.

**ACKNOWLEDGMENTS**

We would like to thank Prakash Raut for bird colony maintenance and Vrushali Rao Gumnur and Aboli Ektare for preliminary tests with different monitors and video stimuli. We would like to thank Hamish Mehaffey, Mimi Kao and Girish Deshpande for helpful comments on the manuscript and the Rajan lab for useful discussion related to the project.

**AUTHOR CONTRIBUTIONS**

NP and RR designed the study. NP did the experiments with videos and created the animated female zebra finch in Blender. SJ carried out the animation related experiments. NR did some initial work on interactive animations using Unity and PT used Unreal engine 4 to construct interactive animations. SK, AP and SR constructed the tutoring setup and tutored juveniles with video and audio playbacks. NP and RR wrote the manuscript in consultation with SJ, PT, SK, AP, SR and NR.

**FUNDING**

This work was supported by grants from the Department of Science and Technology, India (DST/CSRI/2017/163), the Department of Biotechnology (DBT), India, Ramalingaswami Fellowship (BT/HRD/35/02/2006), the Science and Engineering Research Board, India (SERB EMR/2015/000829), the DBT/Wellcome Trust India Alliance Senior Fellowship (IA/S/21/1/505621) and IISER Pune to RR. SK is a graduate student supported by funding from IISER Pune and a senior research fellowship from CSIR, India (9/936(0159)/2016 EMR-1). NP was a recipient of the INSPIRE Scholarship for Higher Education.

479 **REFERENCES**

**Araguas, A., Guellaï, B., Gauthier, P., Richer, F., Montone, G., Chopin, A. and Derégnaucourt, S.** (2022). Design of a robotic zebra finch for experimental studies on developmental song learning. *Journal of Experimental Biology* **225**, jeb242949.

**Bischof, H.-J., Böhner, J. and Sossinka, R.** (1981). Influence of external stimuli on the quality of the song of the zebra finch (Taeniopygia guttata castanotis Gould). *Ethology* **57**, 261–267.

**Bradbury, J. W. and Vehrencamp, S. L.** (2011). Principles of animal communication.

**Derégnaucourt, S., Poirier, C., Van der Kant, A., Van der Linden, A. and Gahr, M.** (2013). Comparisons of different methods to train a young zebra finch (Taeniopygia guttata) to learn a song. *Journal of Physiology-Paris* **107**, 210–218.

**Galoch, Z. and Bischof, H.-J.** (2007). Behavioural responses to video playbacks by zebra finch males. *Behavioural Processes* **74**, 21–26.

**Gröning, J. and Hochkirch, A.** (2008). Reproductive interference between animal species. *The Quarterly review of biology* **83**, 257–282.

**Heyes, C. M.** (1994). Social learning in animals: categories and mechanisms. *Biological Reviews* **69**, 207–231.

**Ikebuchi, M. and Okanoya, K.** (1999). Male zebra finches and Bengalese finches emit directed songs to the video images of conspecific females projected onto a TFT display. *Zoological Science* **16**, 63–70.

**James, L. S., Fan, R. and Sakata, J. T.** (2019). Behavioural responses to video and live presentations of females reveal a dissociation between performance and motivational aspects of birdsong. *Journal of Experimental Biology* **222**, jeb206318.

**Kalra, S., Yawatkar, V., James, L. S., Sakata, J. T. and Rajan, R.** (2021). Introductory gestures before songbird vocal displays are shaped by learning and biological predispositions. *Proc Biol Sci* **288**, 20202796.

**Kao, M. H., Doupe, A. J. and Brainard, M. S.** (2005). Contributions of an avian basal ganglia-forebrain circuit to real-time modulation of song. *Nature* **433**, 638–643.

**Larsen, L. B., Adam, I., Berman, G. J., Hallam, J. and Elemans, C. P. H.** (2022). Driving singing behaviour in songbirds using a multi-modal, multi-agent virtual environment. *Sci Rep* **12**, 13414.

**Lauer, J., Zhou, M., Ye, S., Menegas, W., Schneider, S., Nath, T., Rahman, M. M., Di Santo, V., Soberanes, D., Feng, G., et al.** (2022). Multi-animal pose estimation, identification and tracking with DeepLabCut. *Nat Methods* **19**, 496–504.

**Lehtonen, T. K., McCrary, J. K. and Meyer, A.** (2010). Territorial aggression can be sensitive to the status of heterospecific intruders. *Behavioural Processes* **84**, 598–601.

**Magellan, K.** (2020). Behavioural crypsis in a South African galaxiid fish induced by predatory and non-predatory heterospecifics. *Journal of Fish Biology* **96**, 1278–1283.

**Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W. and Bethge, M.** (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* **21**, 1281–1289.

**Mitoyen, C., Quigley, C. and Fusani, L.** (2019). Evolution and function of multimodal courtship displays. *Ethology* **125**, 503–515.

**Ota, N., Gahr, M. and Soma, M.** (2015). Tap dancing birds: the multimodal mutual courtship display of males and females in a socially monogamous songbird. *Sci Rep* **5**, 16614.

**Patricelli, G. L., Coleman, S. W. and Borgia, G.** (2006). Male satin bowerbirds, Ptilonorhynchus violaceus, adjust their display intensity in response to female startling: an experiment with robotic females. *Animal Behaviour* **71**, 49–59.

**Peiman, K. and Robinson, B.** (2010). Ecology and evolution of resource-related heterospecific aggression. *The Quarterly Review of Biology* **85**, 133–158.

**Rajan, R.** (2018). Pre-bout neural activity changes in premotor nucleus HVC correlate with successful initiation of learned song sequence. *Journal of Neuroscience* **38**, 5925–5938.

**Rajan, R. and Doupe, A. J.** (2013). Behavioral and neural signatures of readiness to initiate a learned motor sequence. *Curr. Biol.* **23**, 87–93.

**Rao, D., Kojima, S. and Rajan, R.** (2019). Sensory feedback independent pre-song vocalizations correlate with time to song initiation. *J. Exp. Biol.* **222**,.

**Rendell, L., Boyd, R., Cownden, D., Enquist, M., Eriksson, K., Feldman, M. W., Fogarty, L., Ghirlanda, S., Lillicrap, T. and Laland, K. N.** (2010). Why copy others? Insights from the social learning strategies tournament. *Science* **328**, 208–213.

**Sossinka, R. and Böhner, J.** (1980). Song Types in the Zebra Finch Poephila guttata castanotis1. *Zeitschrift für Tierpsychologie* **53**, 123–132.

**Sridhar, H. and Guttal, V.** (2018). Friendship across species borders: factors that facilitate and constrain heterospecific sociality. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**, 20170014.

**Stowers, J. R., Hofbauer, M., Bastien, R., Griessner, J., Higgins, P., Farooqui, S., Fischer, R. M., Nowikovsky, K., Haubensak, W. and Couzin, I. D.** (2017). Virtual reality for freely moving animals. *Nature methods* **14**, 995–1002.

**Takahasi, M., Ikebuchi, M. and Okanoya, K.** (2005). Spatiotemporal properties of visual stimuli for song induction in Bengalese finches. *Neuroreport* **16**, 1339–1343.

**Tchernichovski, O., Lints, T., Mitra, P. P. and Nottebohm, F.** (1999). Vocal imitation in zebra finches is inversely related to model abundance. *Proceedings of the National Academy of Sciences* **96**, 12901–12904.

**Tchernichovski, Nottebohm, Ho, Pesaran and Mitra** (2000). A procedure for an automated measurement of song similarity. *Anim Behav* **59**, 1167–1176.

**Tinbergen, N.** (1948). Social releasers and the experimental method required for their study. *The Wilson Bulletin* 6–51.

**Tumer, E. C. and Brainard, M. S.** (2007). Performance variability enables adaptive plasticity of "crystallized" adult birdsong. *Nature* **450**, 1240–1244.

**Ullrich, R., Norton, P. and Scharff, C.** (2016). Waltzing Taeniopygia: integration of courtship song and dance in the domesticated Australian zebra finch. *Animal Behaviour* **112**, 285–300.

**Van Dyk, D. A. and Evans, C. S.** (2008). Opponent assessment in lizards: examining the effect of aggressive and submissive signals. *Behavioral Ecology* **19**, 895–901.

**Van Schaik, C. P. and Burkart, J. M.** (2011). Social learning and evolution: the cultural intelligence hypothesis. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 1008–1016.

**Varkevisser, J. M., Simon, R., Mendoza, E., How, M., van Hijlkema, I., Jin, R., Liang, Q., Scharff, C., Halfwerk, W. H. and Riebel, K.** (2022a). Adding colour-realistic video images to audio playbacks increases stimulus engagement but does not enhance vocal learning in zebra finches. *Animal Cognition* **25**, 249–274.

**Varkevisser, J. M., Mendoza, E., Simon, R., Manet, M., Halfwerk, W., Scharff, C. and Riebel, K.** (2022b). Multimodality during live tutoring is relevant for vocal learning in zebra finches. *Animal Behaviour* **187**, 263–280.

**Woo, K. L. and Rieucau, G.** (2011). From dummies to animations: a review of computer-animated stimuli used in animal behavior studies. *Behav Ecol Sociobiol* **65**, 1671–1685.

**Zann, R. A.** (1996). *Zebra Finch*. Oxford.

 **FIGURE LEGENDS**

**FIGURE 1 Male zebra finches sing longer song bouts to 4 minute videos as compared to 30s**
**videos**

A) Schematic depicting the way in which videos of females were recorded. The male is on the left side
and the female is on the right side, separated by a glass partition. A camera was placed near the male to
record videos of the female.

(B) Order of 30s trials in a session for one representative bird. LF represents "live female", RV
represents "responsive" video and UV represents "unresponsive" video. Inter-trial-time of 5 minutes
was used.

(C) Example spectrograms showing songs produced by one male for the 3 different kinds of stimuli and
undirected singing (UD) when the bird was alone.

(D), (E), (F) mean number of song bouts per session (D), mean song bout duration (E) and mean
number of introductory notes (INs) (F) for all birds for the different trial types. Grey circles represent
data for individual birds and grey lines connect data from the same bird. Black circles and whiskers
represent mean and s.e.m across all birds (n=9 birds).

(G) Schematic depicting the experimental protocol for long duration experimental trials (4 minute
trials). LF represents live female and VP represents video presentation.

(H) Example spectrograms showing the songs produced by one male for the 2 different kinds of stimuli
and undirected singing (UD) when the bird was alone.

(I), (J), (K) mean number of song bouts per session (I), mean song bout duration (J) and mean number
of introductory notes (INs) (K) for all birds for the different trial types. Grey circles represent data for
single birds and grey lines connect data from the same bird. Black circles and whiskers represent mean
and s.e.m across all birds (n=7 birds).

* represents $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$ - Type II Wald Chi Square test followed by
pairwise comparisons using estimated marginal means with Holm correction.

**FIGURE 2 Construction and validation of an animated female zebra finch**

(A), (B), (C), (D) Different stages in the construction of an animated female zebra finch in Blender; 3D
model (A), with colour and feather texture (B). Positions of control bones with yellow representing
structural bones, grey representing core bones, blue representing control bones, deep blue representing

the master control bone and red representing the knee stop bones (C) (see Methods for details) and final image with empty cage (D).

(E), (F), (G), (H) represent different animation frames with the female hopping into view, looking to either side and quivering its tail.

I) Order of 2 minute experimental trials across 3 sessions for one representative bird. LF represents live female, CA represents call with animation, NCA represents no call animation, VP represents video presentation and CC represents colour control animation. Inter-trial-time of 5 minutes was used.

(J) Example spectrograms showing the songs produced by one male for the 5 different kinds of stimuli.

(K), (L), (M) mean number of song bouts per session (K), mean song bout duration (L) and mean number of introductory notes (INs) (M) are shown for all birds for the different trial types. Grey circles represent data for individual birds and grey lines connect data from the same bird. Black circles and whiskers represent mean and s.e.m across all birds (n=5 birds).

* represents $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$ - Type II Wald Chi Square test followed by pairwise comparisons using estimated marginal means with Holm correction.

**FIGURE 3 Potential methods for interactive control of the animation**

(A) Schematic of experimental setup and Arduino based approach to control song audio and video playback based on perch hopping behaviour of a juvenile zebra finch. Briefly, hopping on a perch triggers playback of song audio and video through an Arduino-MATLAB connection.

(B) Example spectrograms of tutor song and the songs of two juveniles tutored using this paradigm.

(C) Boxplots showing the similarity to tutor song for juveniles tutored using only audio playbacks (subset of birds from an earlier study (Kalra et al., 2021)) and audio-visual playbacks. Circles represent data from individual birds.

(D) A view of the 1D blendspace in Unreal Engine 4 with the axis extremes representing "An idle animation" and "Tail quivering animation". Different positions along the axis represent different speeds of the tail quivering.

 **SUPPLEMENTARY INFORMATION**

Supplementary information consists of 3 supplementary figures and 3 movies.

**SUPPLEMENTARY FIGURE LEGENDS**

**FIGURE S1 Raster plots showing song bouts produced by all birds during short-duration, long-**

**duration and animation experimental trials**

(A), (B), (C) Raster plots showing song bouts produced by all birds during short-duration (A),

animation (B) and long-duration experimental trials (C). Each bar represents a song bout and the length

of the bar represents the duration of the song bout. Different colours represent different birds. Bird

indices are indices given to unique birds. Birds that were common across different experimental trials

can be tracked using this bird index (for eg. bird #7 was used for all types of experimental trials).

**FIGURE S2 Probability and amount of song produced towards different stimulus types**

(A), (B) Probability of singing atleast 1 bout of song for 30s video trials (A) and animation trials (B).

(C) Log ratio of time spent singing to video presentations relative to live females.

(D) Log ratio of time spent singing to animation relative to video presentation.

Grey circles represent data from individual birds and grey lines connect data from the same bird. Black

circles and whiskers represent mean and s.e.m across all birds.

**FIGURE S3 Construction of interactive animations using video game engine Unreal Engine 4**

Flowchart depicting the process used to create interactive animations using the video game engine

Unreal Engine 4.

**MOVIE CAPTIONS**

**MOVIE 1 Examples of responsive and unresponsive videos and song bouts of a male zebra finch**

Movie shows an example of a responsive video of a female (30s), an unresponsive video of a female

(30s) followed by examples of a male singing to a live female and to a responsive video.

**MOVIE 2 Male zebra finches sing to animations of female zebra finches**

Movie shows one cycle of the animation and one cycle of the colour control animation. This is

followed by examples of males singing to the animation and the colour control animation.

**MOVIE 3 1D blend space in Unreal Engine 4**

Movie shows a 1D blend space in Unreal Engine 4 with one end of the axis corresponding to an idle

female zebra finch and the other end corresponding to a tail quivering zebra finch. Speed of tail

quivering varies along the axis and an intermediate speed is also shown in the video.

**Figure 1 Male zebra finches sing longer song bouts to 4 minute videos as compared to 30s videos**

**<u>Figure 1 Male zebra finches sing longer song bouts to 4 minute videos as compared to 30s videos</u>**

A) Schematic depicting the way in which videos of females were recorded. The male is on the left side and the female is on the right side, separated by a glass partition. A camera was placed near the male to record videos of the female.

(B) Order of 30s trials in a session for one representative bird. LF represents "live female", RV represents "responsive" video and UV represents "unresponsive" video. Inter-trial-time of 5 minutes was used.

(C) Example spectrograms showing songs produced by one male for the 3 different kinds of stimuli and undirected singing (UD) when the bird was alone.

(D), (E), (F) mean number of song bouts per session (D), mean song bout duration (E) and mean number of introductory notes (INs) (F) for all birds for the different trial types. Grey circles represent data for individual birds and grey lines connect data from the same bird. Black circles and whiskers represent mean and s.e.m across all birds (n=9 birds).

(G) Schematic depicting the experimental protocol for long duration experimental trials (4 minute trials). LF represents live female and VP represents video presentation.

(H) Example spectrograms showing the songs produced by one male for the 2 different kinds of stimuli and undirected singing (UD) when the bird was alone.

(I), (J), (K) mean number of song bouts per session (I), mean song bout duration (J) and mean number of introductory notes (INs) (K) for all birds for the different trial types. Grey circles represent data for single birds and grey lines connect data from the same bird. Black circles and whiskers represent mean and s.e.m across all birds (n=7 birds).

* represents $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$ - Type II Wald Chi Square test followed by pairwise comparisons using estimated marginal means with Holm correction.

**Figure 2 Construction and validation of an animated female zebra finch**

**Figure 2 Construction and validation of an animated female zebra finch**

(A), (B), (C), (D) Different stages in the construction of an animated female zebra finch in Blender; 3D model (A), with colour and feather texture (B). Positions of control bones with yellow representing structural bones, grey representing core bones and blue representing control bones (C) (see Methods for details) and final image with empty cage (D).

(E), (F), (G), (H) represent different animation frames with the female hopping into view, looking to either side and quivering its tail.

I) Order of 2 minute experimental trials across 3 sessions for one representative bird. LF represents live female, CA represents call with animation, NCA represents no call animation, VP represents video presentation and CC represents colour control animation. Inter-trial-time of 5 minutes was used.

(J) Example spectrograms showing the songs produced by one male for the 5 different kinds of stimuli.

(K), (L), (M) mean number of song bouts per session (K), mean song bout duration (L) and mean number of introductory notes (INs) (M) are shown for all birds for the different trial types. Grey circles represent data for individual birds and grey lines connect data from the same bird. Black circles and whiskers represent mean and s.e.m across all birds (n=5 birds).

* represents $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$ - Type II Wald Chi Square test followed by pairwise comparisons using estimated marginal means with Holm correction.

**Figure 3 Potential methods for interactive control of the animation**

(A) Schematic of experimental setup and Arduino based approach to control song audio and video playback based on perch hopping behaviour of a juvenile zebra finch. Briefly, hopping on a perch triggers playback of song audio and video through an Arduino-MATLAB connection.

(B) Example spectrograms of tutor song and the songs of two juveniles tutored using this paradigm.

(C) Boxplots showing the similarity to tutor song for juveniles tutored using only audio playbacks (subset of birds from Kalra et al. (Kalra et al., 2021)) and audio-visual playbacks. Circles represent data from individual birds.

(D) A view of the 1D blendspace in Unreal Engine 4 with the axis extremes representing "An idle animation" and "Tail quivering animation". Different positions along the axis represent different speeds of the tail quivering.

**FIGURE S1 Raster plots showing song bouts produced by all birds during short-duration, long-duration and animation experimental trials**

**FIGURE S1 Raster plots showing song bouts produced by all birds during short-duration, long-duration and animation experimental trials**

(A), (B), (C) Raster plots showing song bouts produced by all birds during short-duration (A), animation (B) and long-duration experimental trials (C). Each bar represents a song bout and the length of the bar represents the duration of the song bout. Different colours represent different birds. Bird indices are indices given to unique birds. Birds that were common across different experimental trials can be tracked using this bird index (for eg. bird #7 was used for all types of experimental trials).

**<u>Figure S2 Probability and amount of song produced towards different stimulus types</u>**

(A), (B) Probability of singing atleast 1 bout of song for 30s video trials (A) and animation trials (B).

(C) Log ratio of time spent singing to video presentations relative to live females.

(D) Log ratio of time spent singing to animation relative to video presentation.

Grey circles represent data from individual birds and grey lines connect data from the same bird. Black circles and whiskers represent mean and s.e.m across all birds.

1) Animations are made in Blender and exported to work with Unreal Engine 4.

2) The animations are imported into Unreal Engine 4.

Idle animation

Tail quivering animation

4) Make animation blueprint to convert the movements that are given via the keyboard key presses into forward speed and blend the animation from idle state to tail quivering state. The weight of the key press determines the magnitude of blending (a heavy key press transitions completely to the tail quivering animation and conversely for a light key press the transition is incomplete and the animated bird undertakes an intermediate state between idle and tail quiver).

3) Create 1D blendspace in the game engine. A blendspace is simply a 1D axis that blends (merges) two animations into a continuous single animation. Here, on the extreme left of the axis is the idle animation whereas on the right is the tail quivering animation.

5) Finally, the controls for the animated model in the game engine are added with character blueprint class. The character blueprint class specifies the keys, timing and sequence of the events. For example the red highlighted box shows two events: 1) AddLocalOffset event associated with an object called plane. It adds a 1000 points offset in the z direction of the plane. After the execution of the AddLocalOffset, the execution of Play Animation event starts and it is associated with the Zebra finch model imported from blender. The whole blueprint shows how the "Z" key press translates into the execution of animation and playback of sound when pressed and released respectively.

**Figure S3 Construction of interactive animations using video game engine Unreal Engine 4**

Flowchart depicting the process used to create interactive animations using the video game engine Unreal Engine 4.