**Towards universal synthetic heterotrophy using a metabolic coordinator**

Sean F. Sullivan[a], Anuj Shetty[b], Tharun Bharadwaj[b], Naveen Krishna[b], Vikas D. Trivedi[a,c], Venkatesh

Endalur Gopinarayanan[a], Todd C. Chappell[a], Daniel M. Sellers[a], Pravin Kumar R.[b], and Nikhil U. Nair[a]*

[a] Department of Chemical & Biological Engineering, Tufts University, Medford, MA 02155

[b] Kcat Enzymatic Private Limited, Bengaluru, Karnataka, India 560005

[c] Department of Structural Biology and Center for Data Driven Discovery, St. Jude Children's Research

Hospital, Memphis, TN

* Corresponding author: nikhil.nair@tufts.edu; @nair_lab

**ABSTRACT**

Engineering the utilization of non-native substrates, or synthetic heterotrophy, in proven industrial

microbes such as *Saccharomyces cerevisiae* represents an opportunity to valorize plentiful and renewable

sources of carbon and energy as potential inputs to biotechnological processes. We previously

demonstrated that activation of the galactose (GAL) regulon, a regulatory structure used by this yeast to

coordinate substrate utilization with biomass formation during growth on galactose, during growth on the

non-native substrate xylose results in a vastly altered gene expression profile and faster growth compared

with constitutive overexpression of the same heterologous catabolic pathway. However, this effort

involved the creation of a xylose-inducible variant of Gal3p (Gal3p$^{Syn4.1}$), the sensor protein of the GAL

regulon, preventing this semi-synthetic regulon approach from being easily adapted to additional non-

native substrates. Here, we report the construction of a variant Gal3p$^{MC}$ (metabolic coordinator) that

exhibits robust GAL regulon activation in the presence of structurally diverse substrates and recapitulates

the dynamics of the native system. Multiple molecular modeling studies confirm that Gal3p$^{MC}$ occupies

conformational states corresponding to galactose-bound Gal3p in an inducer-independent manner. Using

Gal3p$^{MC}$ to test a regulon approach to the assimilation of the non-native lignocellulosic sugars xylose,

arabinose, and cellobiose yields higher growth rates and final cell densities when compared with a

constitutive overexpression of the same set of catabolic genes. The subsequent demonstration of rapid and

complete co-utilization of all three non-native substrates suggests that Gal3p$^{MC}$-mediated dynamic global

gene expression changes by GAL regulon activation may be universally beneficial for engineering

synthetic heterotrophy.

**KEYWORDS:** Gene regulation, sustainability, synthetic biology, metabolic engineering, substrate

utilization, molecular dynamics

**INTRODUCTION**

The adoption of abundant and renewable substrates as inputs for biotechnology will be essential for

creating a sustainable, circular bioeconomy [1,2]. But the inability of industrially important microbes, like

*Saccharomyces cerevisiae* (henceforth, yeast)*,* to utilize many potential substrates poses a major hurdle to

realizing this goal. The existing paradigm for engineering the assimilation of non-natives substrates (i.e.,

synthetic heterotrophy) in yeast begins with the identification and constitutive overexpression of catabolic

genes that enable the substrate to enter central carbon metabolism (CCM) where it is expected to be

transformed into the key primary metabolites used for growth and biosynthesis. However, this method

ignores the tight regulation of CCM and how these resources are distributed to accomplish cellular

objectives [3]. The consequences of this oversight are reflected in the need for subsequent interventions,

including flux balancing [4–6], functional genomics [7–11], and adaptive lab evolution [12–14], undertaken to

improve growth and/or substrate utilization rate by resolving conflicts with cellular processes. Moreover,

existing efforts are often focused on a specific substrate or small set of substrates – a siloed approach that

makes it hard to translate findings to other substrates and even disincentivizes holistic thinking about the

limits of metabolic plasticity/adaptability in this yeast. This has, at least in part, motivated domestication

of other yeasts with broader substrate ranges (e.g., *Scheffersomyces stipitis, Kluyveromyces marxianus*)

for biomanufacturing applications.

An alternate approach that we advocate here, is that instead of combating natural regulation we

should leverage existing regulatory structures that have evolved to coordinate complex phenotypes like

substrate utilization with biomass formation and metabolite synthesis. One such system is the galactose

(GAL) regulon that yeast uses to coordinate substrate catabolism with global metabolism during growth

on the native substrate galactose. In this system, the interaction of galactose with sensor protein Gal3p

enables it to relieve the repression of Gal80p on the transcription factor Gal4p via protein-protein

interactions [15]. Once freed from repression, Gal4p – the master activator of the GAL regulon – binds to its

cognate Upstream Activating Sequences (UAS) to directly induce transcription of galactose catabolic

genes (Leloir pathway) and indirectly modulates the transcript and/or protein abundance of numerous

growth-associated genes (**Figure 1A**) [16,17]. We previously reported the construction of a variant of sensor

protein Gal3p (Gal3p$^{Syn4.1}$) that strongly activates the GAL regulon by the pentose sugars xylose and

arabinose [18,19]. By placing heterologous genes for either xylose or arabinose catabolism under the control

of GAL-responsive (Leloir pathway) promoters, the GAL regulon could be adapted for growth on these

non-native substrates. Upon comparing this regulon-coordinated (REG) approach to growth with simple

constitutive overexpression of the same catabolic genes, which we term a constitutive (CONS) approach,

we found that our semi-synthetic regulon yielded superior growth rates on both xylose ($\mu = 0.24$ h$^{-1}$ vs.

0.11 h$^{-1}$) and arabinose ($\mu = 0.27$ h$^{-1}$ vs. 0.06 h$^{-1}$), respectively with minimal metabolic engineering [19].

Significantly, a transcriptomic comparison showed relative upregulation of genes associated with cell

division and mitochondrial biogenesis in the REG strain while the CONS strain showed upregulation of

stress response and starvation-associated genes [18]. This suggests that the GAL regulon dynamically

reshapes the cellular response for growth through direct and/or indirect action of Gal3p-Gal80p-Gal4p to

potentiate the cells for rapid growth [17,20], irrespective of the identity of the available substrate.

In this study, we demonstrate that, in addition to the known genome-wide changes in gene

expression that occur upon GAL regulon activation, the expression from GAL-inducible promoters is

dynamic and appears to be coordinated with growth whereas expression from constitutive promoters is

constant. Hypothesizing that both aspects of this metabolic coordination may be universally beneficial for

the assimilation of non-native substrates, we sought to develop this semi-synthetic regulon system into a

platform approach that enables the rapid engineering of efficient synthetic heterotrophy without having to

continually re-engineer substrate-specific induction. To do so, we created a variant of sensor protein that

we term Gal3p$^{MC}$ (i.e., metabolic coordinator) and demonstrate that it, unlike the wild-type Gal3p

(Gal3p$^{WT}$), can strongly activate the GAL regulon on numerous structurally diverse substrates. We show

that Gal3p$^{MC}$ can recapitulate the dynamic activation of the native regulon and, through the use of

molecular dynamics and metadynamics simulations, that the mutations it carries enable it to do so in an

inducer-independent manner. We found that this substrate-agnostic system retains the benefits of

substrate-specific activation without any undue burden when engineering synthetic heterotrophy. We also

show that using Gal3p$^{MC}$ to implement a REG approach to growth on three non-native substrates –

xylose, arabinose, and cellobiose – yields superior performance when compared to a CONS approach.

Finally, we provide a first demonstration that a single strain expressing Gal3p$^{MC}$ is capable of rapid,

simultaneous, and complete utilization of all three non-native substrates concurrent with rapid growth –

paving the way for a universal synthetic heterotrophy platform.

**RESULTS**

**Expression under the GAL regulon is dynamic and recapitulates growth phase.**

Existing efforts to engineer synthetic heterotrophy generally utilize strong, constitutive promoters to

control the expression of the necessary catabolic genes. Constitutive promoters are natively able to recruit

cellular transcriptional machinery to yield a roughly consistent expression level independent of cellular

state and environmental context. This 'always on' phenotype contrasts with the control of catabolic gene

expression observed in many regulatory structures that coordinate native substrate utilization in response

to nutrient availability, including the GAL regulon of *S. cerevisiae*. To understand how these different

approaches could impact the efficiency of substrate utilization, we used a fluorescent reporter (*EGFP*) to

quantify the expression dynamics of several GAL-responsive (*GAL1p*, *GAL7p*, *GAL10p*, *GAL3p,* and

*GAL80p*) and

**Figure 1: Expression from GAL-inducible promoters is coordinated with growth. (A)** Simplified regulatory overview of the GAL regulon. The binding of galactose to sensor protein Gal3p enables it to relieve the repression of Gal80p on Gal4p via protein-protein interactions. Gal4p binds to its cognate Upstream Activating Sequences (UAS) to directly transcribe galactose catabolic genes as well as to indirectly reshape global metabolism for rapid growth. The regulon is also repressed in the presence of excess glucose, a regulatory logic that ensures the regulon is activated only in the presence of its target carbon source but not when a preferred carbon source is present. Expression of EGFP in wild-type strain W303-1a from **(B)** strong constitutive promoters (*TEF1p*, *TPI1p*, *TDH3p*, *GPM1p*) and **(C)** GAL-inducible promoters (*GAL1p*, *GAL7p*, *GAL10p*, *GAL3p*, *GAL80p*) at different time points during cultivation on galactose overlaid with cell density (OD$_{600}$). Each data point represents the average of three (for OD$_{600}$ values) or four (for fluorescence) biological replicates ± sd.

104
constitutive (*TPI1p*, *TEF1p*, *TDH3p*, and *GPM1p*) yeast promoters in a wild-type strain (W303-1a) that

contains the native GAL regulon. During growth on galactose, we found that expression from the

constitutive promoters was relatively stable across a 24-hour period (varying less than two-fold) (**Figure**

**1B**). In contrast, expression from GAL-responsive promoters starts lower (~50% of constitutive

promoters) but increases by ~4-6 fold for the promoters that control the expression of the galactose

catabolic (Leloir pathway) genes (*GAL1p*, *GAL7p*, and *GAL10p*) while remaining relatively constant for

the promoters controlling regulatory genes (*GAL3p* and *GAL80p*) (**Figure 1C**). When we overlaid the

$OD_{600}$ profile, we observed that expression from Leloir pathway promoters tracks with growth phase, with

the lowest expression levels occurring when the cells are in lag phase (t ~ 0 h), maximum expression

during the exponential growth phase, and finally diminishing expression as galactose is depleted and the

cells enter stationary phase (t > 20 h). During growth on glucose, the expression profile of the constitutive

promoters was largely unchanged (**Figure S1A**) while expression from GAL-inducible promoters is

strongly repressed because of carbon catabolite repression (**Figure S1B**). Overall, GAL-responsive

promoters are more dynamic and stronger (upon full activation) relative to constitutive promoters [21,22].

**Gal3p$^{MC}$ activates the GAL regulon in the presence of structurally diverse carbon sources.**

While the creation of a xylose-inducible sensor protein variant Gal3p$^{Syn4.1}$ enabled us to adapt the GAL

regulon for growth on xylose [18], we wanted to develop a method for rapidly applying a regulon approach

to other non-native substrates without having to repeatedly re-engineer substrate-specific sensing. Blank

et al. previously reported various autoactivating Gal3p variants but we found that the best among those

(*viz.* F237Y, S509P) could only partially activate GAL promoters without galactose (**Figure S2A**) [23]. To

enable complete activation, equivalent to the WT GAL regulon with galactose, we combined the F237Y

mutation to our Gal3p$^{Syn4.1}$ variant [24] to yield a mutant we term Gal3p$^{MC}$ (metabolic coordinator) (**Figure**

**S2B**). To compare the ability of Gal3p$^{WT}$ and Gal3p$^{MC}$ to activate the regulon in the presence of a given

carbon source, we monitored the fluorescence resulting from a *GAL1p-yEGFP* reporter construct (**Figure**

**2A**) in the presence of numerous structurally distinct carbon sources substrates – galactose, arabinose,

xylose, cellobiose, raffinose, ethanol/glycerol, sucrose, and glucose – in a strain (VEG16) that lacks the

galactose sensing and catabolic genes (*ΔGAL3; ΔGAL1; ΔGAL10; ΔGAL7; ΔGRE3)* but retains *GAL80*

and *GAL4*. We performed this experiment using two native carbon sources, raffinose (**Figure 2B**) or

sucrose (**Figure S3**), to support cell growth. We observed that Gal3p$^{WT}$ strongly activates the regulon in

the presence of its native inducer galactose and to a lesser extent xylose and arabinose, likely due to the

structural similarity between these substrates. The indistinguishable signal between Gal3p$^{WT}$ and the

*ΔGAL3* control in presence of glucose reflects carbon catabolite repression while the small increase in

fluorescence on raffinose and cellobiose likely reflects the weak activation known to result from

overexpression of Gal3p$^{WT}$ in the absence of glucose [25].

**Figure 2: Sensor protein variant Gal3p$^{MC}$ enables robust GAL regulon activation in an inducer-independent manner.** (**A**) Schematic of fluorescent reporter assay used to quantify GAL regulon activation by different *GAL3* variants. Cells expressing a given variant (Δ*GAL3*, Gal3p$^{WT}$, or Gal3p$^{MC}$) are grown in the presence of both native and non-native substrates. Regulon activation drives EGFP expression from the GAL-inducible promoter, *GAL1p*. (**B**) Comparison of normalized fluorescence resulting from interaction between *GAL3* variant and carbon source (2% w/v) using RAF to support growth on substrates GAL, ARA, XYL, and CEL. SUC and GLU are repressing substrates. (**C**) Expression of EGFP from *TEF1p* and *GAL1p* during growth on galactose in a strain (SFS6) expressing Gal3p$^{MC}$ to activate the regulon. Each data point represents the average of 5–6 biological replicates ± sd.

In contrast, Gal3p$^{MC}$ activates the GAL regulon in the presence of all the non-native substrates to a similar

extent as Gal3p$^{WT}$ on galactose while remaining repressible by glucose and sucrose (**Figure 2B**), a result

that suggests Gal3p$^{MC}$ may be capable of recapitulating the dynamic catabolic gene expression profile and

genome-wide expression changes known to support optimal growth on the native substrate galactose

during growth on non-native substrates [26]. To determine if regulon activation using Gal3p$^{MC}$ remains

dynamic, we constructed a strain (SFS6) that contains copies of *GAL3$^{MC}$* integrated into the chromosome

under the control of both *GAL1p* and *GAL3p*. This dual-feedback loop configuration mimics the

organization of the native GAL regulon and has been demonstrated to increase both the overall magnitude

and the homogeneity of activation to better effect a switch-like behavior among cells in the population

[18,27]. As SFS6 lacks the Leloir pathway (△*GAL1/7/10*), these genes were supplied under the control of

their native promoters on a plasmid (pRS423-GAL-REG) to facilitate growth on galactose. Using strain

SFS6, we assessed the expression profile of the constitutive *TEF1p-EGFP* and GAL-responsive *GAL1p-*

*EGFP* constructs during growth on galactose (**Figure 2C**). While expression from *TEF1p* was observed

to increase by 2–3-fold at one time point, it generally stayed within a two-fold range whereas expression

from *GAL1p* appeared to mimic the profile observed upon activation by the wild-type regulon (**Figure**

**1C**), exhibiting a >5-fold increase in fluorescence.

**Molecular simulations support inducer-independent regulon activation by Gal3p$^{MC}$**

Experiments have shown that the binding of Gal3p$^{WT}$ to its ligand galactose induces and stabilizes a

conformational change from an 'open' to a 'closed' state that facilitates interaction with Gal80p and

subsequent de-repression of Gal4p [15]. However, mapping the mutations carried by Gal3p$^{MC}$ onto a

structure of Gal3p$^{WT}$ in complex with Gal80p (PDB: 3V2U) reveals that the mutations are distributed

throughout the protein (**Figure S4**). Combined with the structural diversity of substrates (pentoses,

hexose, disaccharide, and trisaccharide) on which Gal3p$^{MC}$ exhibits robust regulon activation,

observations suggest that Gal3p$^{MC}$ may adopt a 'closed' conformation that enables it to bind Gal80p and

relieve repression on Gal4p without requiring interaction with a ligand/inducer. As the transition between

open and closed states is associated with the relative movement of two domains in Gal3p$^{WT}$, while a third

'hinge' region remains relatively static, we decided to explore this inducer-independent activation

hypothesis using molecular dynamics (MD) simulations. To do so, we monitored the distance between

two residues (D104 and L370) found on opposing moving domains (**Figure S5A**) over the course of 1000

ns molecular dynamics (MD) simulations of Gal3p$^{WT}$ in complex with ATP and Mg$^{2+}$ without galactose

(WT$^{Gal-}$), Gal3p$^{WT}$ in complex with ATP, Mg$^{2+}$, and galactose (WT$^{Gal+}$), and Gal3p$^{MC}$ in complex ATP

and Mg$^{2+}$ without galactose (MC$^{Gal-}$). Simulations revealed that distances between the residues were more

similar between MC$^{Gal-}$ and WT$^{Gal+}$ (**Figure S5B**). Calculations of active site cavities at the end of the

1000 ns simulation also revealed a similar trend (**Figure S5C–E**). To understand which residues most

contributed to the dynamics observed in each of the three systems during these simulations, we performed

cross-correlational studies of the relative movement between the residues of WT$^{Gal-}$, WT$^{Gal+}$, and MC$^{Gal-}$

(**Figure S5F–H**). The lower intensities of residue-residue correlation in WT$^{Gal+}$ and MC$^{Gal-}$ maps indicate

relatively stable dynamics relative to those of WT$^{Gal-}$. To visualize the domain movements in detail, we

used principal component analysis (PCA) for all three systems across the 1000 ns MD trajectories.

Porcupine plots generated based on the extreme projections of the PCA show the intensity and direction

of the motion of the C$_\alpha$ atoms in each structure. WT$^{Gal-}$ showed highly dynamic conformational states

while MC$^{Gal-}$ and WT$^{Gal+}$ showed relatively limited dynamics across the structure (**Figure S5I–K**).

Initializing the molecular dynamics simulations of the three conditions in the 'closed'

conformation introduces a bias that prevents each system from adopting a range of conformational states.

Metadynamics-based simulations use a potential bias to move a system out of a local energy minimum

enabling the calculation of the free energy surface (FES) across a range of conformations. To perform

these complementary metadynamics analyses, we began to define the conformational landscape to be

explored by identifying centers of mass (COM) for each of the three Gal3p domains (C1–C3). Next, we

defined the two collective variables (CV) to be manipulated during the simulation as (i) the distance

between the COM of the two moving domains (C1 and C2) and (ii) the angle formed by the three COM at

the hinge region (C3) (**Figure 3A**). Examining the resulting FES for each system reveals differences in

the CV values that yield energetically favorable structural conformations. The FES of $WT^{Gal-}$ (**Figure 3B**)

showed two Gaussian wells, one for a short-lived partially open state and one for an exceptionally

prominent minimum corresponding to an irreversibly collapsed closed conformation catalyzed by strong

interaction networks between the two moving domains (**Figure S6A**) not observed between the moving

domains in the $WT^{Gal+}$ (**Figure S6B**) and $MC^{Gal-}$ (**Figure S6C**) systems. Therefore, it can be called a

misfolded state ($W_m$) of the

**Figure 3: Metadynamics simulations identify energetically favorable conformations. (A)** Two collective variables (CV) were defined to enable energetic exploration of the Gal3p conformational landscape: **CV1** as the distance between C1 and C2, the centers of mass of the two moving domains, and **CV2** as the angle formed between C1, C3 (the hinge region center of mass), and C2. Calculating the energy associated with different values of CV1 and CV2 enabled construction of the free energy surfaces (FES) for the **(B)** $WT^{Gal-}$ (W), **(C)** $WT^{Gal+}$ (G), and **(D)** $MC^{Gal-}$ (M) systems. The subscripts "m", "c", and "o", corresponding to misfolded (collapsed), closed, and open states, respectively, are used to associate prominent energetic minima with distinct conformations of Gal3p variants. Significantly, the most favorable conformations in the $WT^{Gal+}$ and $MC^{Gal-}$ systems correspond to a closed state that likely favors interaction with Gal80p and thus regulon activation.

201
protein as the recovery of an open conformation would be highly energy-consuming, and therefore,

unlikely. The FES of $WT^{Gal+}$ showed a path of conformational transitions that leads from a closed ($G_c$),

but not collapsed, state to a fully open state ($G_o$) of the protein (**Figure 3C**). Significantly, the closed state

of $WT^{Gal+}$ ($G_c$) is structurally distinct from the closed, collapsed/misfolded state ($W_m$) adopted by $WT^{Gal-}$

suggesting that the latter is unlikely to interact with Gal80p in a productive manner. As the metadynamics

simulation progresses, $WT^{Gal+}$ eventually adopts an open state ($G_o$) that visualization studies reveal to be a

consequence of substate egress. The FES of $MC^{Gal-}$ (**Figure 3D**) showed a single predominant minimum

corresponding to a closed state ($M_c$) that closely resembles the closed state ($G_c$) of $WT^{Gal+}$. These results

suggest that the mutant $Gal3p^{MC}$ forms a stable closed conformation closely resembling the closed $WT^{Gal+}$

conformation, with a similar interaction network observed between the moving domains. When combined

with the previously observed robust activation on structurally diverse substrates, the similarity between

the closed state conformations adopted by $WT^{Gal+}$ and the $MC^{Gal-}$ systems suggests that $Gal3p^{MC}$ can bind

and sequester Gal80p in an inducer-independent manner. Recognizing that this substrate-agnostic

activation phenotype of $Gal3p^{MC}$ represents a departure from the inducer-dependent regulatory logic of

the native regulon, we decided to test the fitness consequences of this transition when utilizing a regulon

approach to engineer synthetic heterotrophy.

**Interactions between $Gal3p^{MC}$ with Gal80p are energetically favorable.**

We compared the energetics of Gal3p-Gal80p binding using the minimum energy conformations ($W_m$, $G_c$,

$M_c$) obtained from metadynamics-based simulations. Docking simulations between Gal3p and Gal80p

were run for 50 ns and binding energy was calculated every 0.02 ns (**Figure 4A**). Among the complexes

formed, the WT$^{Gal+}$–Gal80p was energetically the most like the crystal structure complex (**Figure 4B**).

The results also revealed that the complex formed by WT$^{Gal-}$ with Gal80p was significantly less

energetically favorable than the complexes formed by WT$^{Gal+}$ and MC$^{Gal-}$. This data further supports the

idea that Gal3p$^{MC}$ may be able to activate the GAL regulon in a substrate independent manner.

**Figure 4: Docking interactions between Gal3p and Gal80p.** (**A**) Detailed representation of the interface between Gal3p (purple) and Gal80p (pink) complex. The residues involved in the interface are highlighted in the call-out. (**B**) The average binding energy between Gal3p and Gal80p was calculated by sampling every 0.02 ns from the docked Gal3p-Gal80p complexes obtained through protein-protein docking study. A statistical analysis was performed on the average of the four binding energy values using ANOVA to determine whether there was a significant difference in the four sets of calculated binding energies. The results of the analysis showed an F-value of 2137 with 3 degrees of freedom and p-value of <0.0001, indicating a significant difference between all four.

**Substrate-agnostic and substrate-specific GAL regulon activation are equivalent.**

During our previous effort to adapt the GAL regulon to growth on xylose, regulon activation was

accomplished using the xylose-inducible sensor-protein variant Gal3p$^{Syn4.1}$ [18]. To compare the

performance of inducer-dependent (substrate-specific) and inducer-independent (substrate-agnostic)

approaches for engineering synthetic heterotrophy, we transformed a plasmid (pRS426-XYL-REG)

encoding the isomerase pathway genes for xylose utilization (*XYLA*3* [28]*, XKS1*) under the control of

GAL-responsive promoters into strains engineered for (i) xylose-inducible regulon activation using

Gal3p$^{Syn4.1}$ (strain VDT13), (ii) substrate-agnostic regulon activation using Gal3p$^{MC}$ (strain SFS6), and

(iii) substrate-agnostic activation via deletion of *GAL80*, the master repressor of the GAL regulon (strain

SFS11), as disruption of this regulatory gene has been demonstrated to yield constitutive expression of

galactose catabolic genes and enhance galactose utilization [29,30] (**Figure 5A**). While regulon activation in

both SFS6 and VDT13 requires an interaction between Gal80p and their respective Gal3p variants

(Gal3p$^{MC}$ and Gal3p$^{Syn4.1}$),

**Figure 5: Comparing the fitness of substrate-specific and substrate-agnostic approaches to GAL regulon activation. (A)** Schematic illustrating different ways of activating the GAL regulon in cells carrying xylose catabolic genes under the control of GAL-responsive promoters. Gal3p$^{Syn4.1}$ requires induction by xylose, Gal3p$^{MC}$ does not require an inducer but maintains the native regulatory architecture, while $\Delta GAL80$ effects inducer-independent activation by removing native regulatory elements. Growth of the three systems in defined (SC) medium containing 2% of **(B)** sucrose, **(C)** glucose, or **(D)** raffinose reveals that a de-regulated approach ($\Delta GAL80$) reduces fitness on native carbon sources. **(E)** Comparing growth of the three GAL regulon activation systems on the non-native sugar xylose after being subcultured from sucrose. Comparison of constitutive (CONS) and regulon (REG) approaches to growth in complex media containing 2% of **(F)** arabinose or **(G)** cellobiose as sole carbon sources. All data points represent the average of three biological replicates ± sd.

activation in SFS11 is totally de-regulated due to the absence of Gal80p. In addition, all three strains

contain two accessory genes found to improve pentose utilization – the transporter *GAL2*$^{2.1}$ [31] and the

transaldolase *TAL1* – integrated into the chromosome under the control of GAL-responsive promoters.

The growth of these three strains in defined (SC) media containing the native carbon sources sucrose

(**Figure 5B**), glucose (**Figure 5C**), or raffinose (**Figure 5D**) was monitored to identify the potential

fitness defects of each approach during strain handling. While all three approaches reached similar final

cell concentrations on each of the three substrates, the $\Delta GAL80$ condition took roughly twice the time to

do so compared with Gal3p$^{Syn4.1}$ and Gal3p$^{MC}$ (>30 h vs. ~18 h) suggesting that de-regulation of the

regulon results in reduced fitness relative to strains in which the native regulatory organization is

maintained. As sucrose and glucose are both capable of exerting repression on the GAL regulon [19],

subculturing all three strains from these two native substrates into complex (YP) media containing the

non-native substrate xylose should therefore result in growth-coupled expression of xylose catabolic

genes and rapid growth. Upon subculturing from sucrose, we observed that all three approaches enabled

growth. However, the lower growth rate seen for the $\Delta GAL80$ strain compared with cells carrying

*GAL3*$^{Syn4.1}$ or *GAL3*$^{MC}$ ($\mu$ = 0.14 ± 0.001 h$^{-1}$, 0.21 ± 0.008 h$^{-1}$, and 0.21 ± 0.001 h$^{-1}$, respectively)

combined with the drastically lower final cell density supports maintenance of native regulatory structure

over abolished control (**Figure 5E**). Subculturing from glucose as an overnight carbon source yielded

similar results (**Figure S7**). Significantly, the equivalent performance of *GAL3*$^{Syn4.1}$ and *GAL3*$^{MC}$

backgrounds on both native and non-native carbon sources indicates that inducer-independent activation

enables the metabolic coordination of a regulon approach to be applied for engineering synthetic

heterotrophy in a substrate-agnostic manner provided that the native regulatory organization is

maintained.

To confirm that these results were not xylose-specific, we proceeded to use Gal3p$^{MC}$ to compare

REG and CONS approaches to growth on two additional non-native, lignocellulosic substrates –

cellobiose and arabinose. To do so, we placed catabolic genes previously identified to enable the efficient

utilization of arabinose (*araA*, *araB*, and *araD* from *Lactobacillus plantarum* [32]) and cellobiose (*cdt-1* and

*gh1-1* from *Neurospora crassa* [33]) under the control of the GAL-responsive promoters *GAL1p*, *GAL10p*,

and *GAL7p* and the strong, constitutive promoters *TEF1p*, *TPI1p*, and *GPM1p* to yield plasmids pRS426-

ARA-REG, pRS426-ARA-CONS, pRS426-CEL-REG, and pRS426-CEL-CONS, respectively.

Transforming pRS426-ARA-REG into *GAL3$^{MC}$* integrant strain SFS6 (containing GAL-inducible copies

of *GAL2$^{2.1}$* and *TAL1*) yielded strain ARA-REG while transforming pRS426-ARA-CONS into strain

VDT27 (that contains *GAL2$^{2.1}$* and *TAL1* under the control of strong constitutive promoters) yielded strain

ARA-CONS. Similarly, transforming pRS426-CEL-REG and pRS426-CEL-CONS into strains SFS3 and

VEG16, respectively, that are comparable to SFS6 and VDT27 except that they lack accessory genes

*GAL2$^{2.1}$* and *TAL1*, yielded strains CEL-REG and CEL-CONS. The REG strain exhibited a higher growth

rate than the CONS strain on both defined (0.16 ± 0.005 h$^{-1}$ vs. 0.05 ± 0.005 h$^{-1}$) (**Figure S8A**) and

complex (0.19 ± 0.002 h$^{-1}$ vs. 0.17 ± 0.003 h$^{-1}$) (**Figure 5F**) media containing 2% arabinose as a sole

carbon source. Similarly, the REG strain grew faster than its CONS counterpart on both defined (0.14 ±

0.001 h$^{-1}$ vs. 0.09 ± 0.005 h$^{-1}$) (**Figure S8B**) and complex (0.25 ± 0.002 h$^{-1}$ vs. 0.17 ± 0.003 h$^{-1}$) (**Figure

5G**) media containing 2% cellobiose as a sole carbon source. In general, we found that the strength of the

promoters driving the catabolic gene expression corresponded well with growth phenotype (**Figure S9**).

Taken together, these data demonstrate that the metabolic coordination effected by Gal3p$^{MC}$ is broadly

beneficial to growth on non-native carbon sources compared with the existing constitutive paradigm with

minimal additional engineering.

**Gal3p$^{MC}$ enables rapid and complete co-utilization of multiple non-native substrates.**

Having demonstrated that a regulon approach is beneficial to growth on all three non-native, lignocellulosic substrates individually, we sought to utilize the inducer-independent regulon activation of Gal3p$^{MC}$ to test a REG approach to their co-utilization. As previous studies have observed that the division of catabolic labor among a consortium of strains can enable faster and/or more efficient co-utilization when compared with a single strain engineered to utilize all substrates simultaneously [34–36], we decided to test both approaches. To accommodate all three plasmids into a single strain, the catabolic genes for each substrate were moved to plasmids containing unique selectable markers to create pRS423-XYL-REG, pRS425-ARA-REG, and pRS424-CEL-REG. Transforming these plasmids into strains SFS6, SFS6, and SFS3, respectively, represented a three-strain (Consortium) approach, while transforming all three plasmids into

**Figure 6**: **Gal3p$^{MC}$ enables rapid and complete co-utilization of multiple non-native substrates.** **(A)** Placing REG plasmids for the catabolism of cellobiose, xylose, and arabinose into separate strains represents a Consortium approach to co-utilization while placing all three plasmids into a single strain represents a Consolidated approach. **(B)** Cell density (black) and percent residual substrate of cellobiose (yellow), xylose (light blue), and arabinose (pink) of Consortium (dotted lines) and Consolidated (solid lines) approaches to co-utilization in complex media containing 1% of each substrate. Incomplete pentose utilization motivated the exploration of all six possible pairings between GAL-inducible promoters (*GAL1p*, *GAL10p*, and *GAL7p* in order) and the catabolic genes for **(C)** arabinose (A = L-arabinose isomerase, B = L-ribulokinase, D = L-ribulose-5-phosphate-4-epimerase) and **(D)** xylose utilization (XI = xylose isomerase, XKS = D-xylulokinase, □ = no gene). Promoter-gene pairings used in subsequent co-utilization experiments are indicated by (**\***). **(E)** Re-testing a Consolidated approach to co-utilization using expression-balanced pentose plasmids results in complete utilization of all three substrates. Data points represent the average of three (panel **B**) or two (panel **E**) biological replicates ± sd.

SFS6 represented a single strain (Consolidated) approach (**Figure 6A**). While both Consortium and

Consolidated approaches exhibited simultaneous co-utilization of all carbon sources and achieved similar

final cell densities during growth on YP media containing 1% of each substrate, we were surprised to find

that the Consolidated approach resulted in a faster maximum growth rate than the Consortium approach

($0.20 \pm 0.003$ h$^{-1}$ vs. $0.17 \pm 0.003$ h$^{-1}$; two-tailed, two-sample t-test, t = 8.41, df = 3, p < 0.01) (**Figure**

**6B**). However, neither approach resulted in complete pentose consumption even after cellobiose was fully

depleted. This led us to hypothesize that there may be flux imbalances in the pentose utilization pathways.

The success of combinatorial pathway design in optimizing both catabolic [4,37,38] and anabolic

pathways [39,40] motivated us to test the performance of each of the six possible GAL-responsive promoter-

catabolic gene pairings for arabinose (**Figure 6C**) and xylose (**Figure 6D**) in Gal3p$^{MC}$ integrant strain

SFS6. The arabinose constructs all performed similarly, with growth rates clustered in a small range of

values ($0.20 - 0.22$ h$^{-1}$). A similar range of growth rates was observed for the xylose constructs ($0.20 -$

$0.21$ h$^{-1}$) but the substantially reduced final cell density of two of the arrangements (□-XK-XI and XK-□-

XI) indicates that there can be fitness consequences to expression imbalances. As no clearly superior

combination could be identified to the initial arabinose and xylose designs, we decided to re-test a

Consolidated approach to co-utilization on 1% of each substrate (**Figure 6E**) using designs (ABD for

arabinose, XI-□-XK for xylose) identified as optimal in a related investigation by our lab into efficient

pentose assimilation [19]. Hypothesizing that incomplete pentose utilization may also reflect limitations in

the carrying capacity of the medium itself, we also tested different substrate ratios (cellobiose-xylose-

arabinose: 0.7%–0.7%–0.7%, 1%–0.5%–0.5%, and 0.4%–0.8%–0.8%) (**Figure S10A-C**) at a lower

(~2%) total substrate concentration. Encouragingly, all substrates were utilized to undetectable levels

(<0.05%, limit of detection) in approximately 50 h of cultivation for all conditions. For the 3% total sugar

condition, we observed an 81% improvement in final cell density when using the optimized pentose

plasmids compared with the non-optimized ($OD_{600}$ = 17.2 vs. 9.5, respectively). For the three 2% total

substrate conditions, the final cell density was 41% greater on average using the same comparison ($OD_{600}$

~13.5 vs. 9.5, respectively). Despite their similar performance during growth on a single sugar, the higher

final $OD_{600}$ values and complete substrate utilization observed upon switching from the initial to the

optimized designs suggests that the relative expression level of catabolic genes optimal for co-utilization

may be different than for the single substrate. Considering that the highest single-sugar growth rates

obtained in this study are either the best (for cellobiose) or competitive with the best (xylose, arabinose)

aerobic growth rates observed in the literature (**Table S1**) without the need for extensive genome editing

or adaptive laboratory evolution (ALE), we believe that together these results demonstrate the flexibility

and effectiveness of utilizing a regulon approach to engineer efficient synthetic heterotrophy on diverse

substrates.

**DISCUSSION**

The existing paradigm for engineering synthetic heterotrophy consists of constitutive expression of

catabolic genes that channel the carbon source into central carbon metabolism (CCM) and largely ignores

cellular regulation. In contrast, the assimilation of native carbon sources is often controlled by regulatory

structures called regulons that coordinate substrate catabolism with broader metabolism to optimize

biomass formation. Activation of the galactose (GAL) regulon, for instance, results not only in the

expression of the genes necessary for the catabolism of galactose but global metabolic remodeling to

support efficient growth [17,20]. In previous work, we established that adapting the GAL regulon to growth

on the non-native substrate xylose improved growth rate and substrate utilization when compared with a

constitutive approach [18]. This surprising result motivated us to explore the potential of a semi-synthetic

regulon system as a general approach to engineering synthetic heterotrophy in the important industrial

microbe *S. cerevisiae* so that it can be easily and rapidly applied to other attractive potential substrates for

yeast biotechnology.

In this work, we demonstrate that expression from the GAL-responsive promoters that natively

control the expression of galactose catabolic genes (Leloir pathway) is dynamic and mirrors growth

phases whereas expression from constitutive promoters is constant throughout cultivation. Thus, the

regulatory organization of the GAL regulon functions to avoid the burden of unproductive protein

synthesis by only expressing the genes for galactose assimilation only when it is present. The

conservation of cellular resources effected by this inducible phenotype represents a design principle that

has also been learned by metabolic engineers. Early observations of growth inhibition resulting from the

overproduction of heterologous protein [41] led to the use of separate culturing phases for growth and

production and strategies for dynamically switching between them [42]. This concept of dynamic flux

control has subsequently been applied at finer resolution to biosynthetic pathways. Engineered circuits

that feature a biosensor enable a cell to continually monitor some aspect of its internal state or external

environment and automatically adjust their metabolism in response [43], resulting in improvements in titer,

rate, and yield (TRY) [44–47]. While the use of sugar-responsive biosensors has been proposed to engineer

cells to respond to the carbon sources present in a given lignocellulosic hydrolysate for consolidated

bioprocessing, the suggested implementations decouple sensing from the broader regulatory and

metabolic context in which the sensors evolved [48]. To our knowledge, our semi-synthetic regulon

approach is the first time that this dynamic activation concept has been applied to engineering catabolism

in a way that interfaces with native metabolism for the purpose of efficient biomass formation. This

coordination is significant as the regulation of CCM has evolved to ensure that resource uptake and

utilization are balanced [3], with the transcription level of a large fraction of the yeast genome (~15 – 27%)

[49,50] known to be correlated with growth rate. We, therefore, posit that regulon control of catabolic gene

expression, where the amount of carbon fed into CCM is innately connected to the utilization of those

resources to match cellular needs, is likely beneficial to fast growth regardless of the carbon source being

assimilated.

375            While we were previously able to engineer a sensor protein variant (Gal3p$^{Syn4.1}$) capable of strong

regulon activation upon induction with xylose, that effort involved an extensive directed evolution

campaign that would have to be repeated for each new substrate [18]. Here, we report the construction of a

sensor protein variant, Gal3p$^{MC}$ (metabolic coordinator), which robustly activates the GAL regulon on

structurally diverse, non-native carbon sources to a similar degree as Gal3p$^{WT}$ upon induction with

galactose. Using metadynamics simulations, we show that the most energetically favorable conformation

for Gal3p$^{MC}$ without galactose (MC$^{Gal-}$) is similar to the closed conformational state of Gal3p$^{WT}$ in

complex with galactose (WT$^{Gal+}$). In contrast, the free energy surface of Gal3p$^{WT}$ in the absence of

galactose (WT$^{Gal-}$) is characterized by a deep energy well reflective of the active site

collapsing/misfolding, a conformation that is unproductive for signal transduction. Molecular dynamics

(MD) simulations of these three systems reveal large-scale domain movements in the WT$^{Gal-}$ system

throughout the course of the simulation whereas the WT$^{Gal+}$ and MC$^{Gal-}$ systems exhibit much reduced

and more localized movement indicating that the mutations carried by Gal3p$^{MC}$ stabilize the protein in a

closed state that enables interaction with Gal80p without requiring an inducer. The inducer-independent

mechanism of Gal3p$^{MC}$ would normally ablate the dynamic expression profile exhibited by the native,

inducer-dependent regulon as it would constantly be turned on. However, we also observed repression of

the Gal3p$^{MC}$-controlled GAL regulon on repressing substrates (glucose, sucrose), which allows for

dynamic activation. Gal3p$^{MC}$ thus enables both the dynamic catabolic gene expression and global

metabolic remodeling of the semi-synthetic regulon approach to be applied in a substrate-agnostic

manner.

395            Although the inducer-independent activation effected by Gal3p$^{MC}$ represents a departure from the

inducer-dependent regulatory logic of the native GAL regulon that served as our inspiration, we,

surprisingly, did not observe any major growth defects on either native or non-native substrates compared

with a substrate-specific activation approach using Gal3p$^{Syn4.1}$. In contrast, a Δ*GAL80* strain, which

represents an alternative approach of achieving substrate-agnostic regulon activation, showed reduced

fitness in both conditions, emphasizing the importance of maintaining the native regulatory organization

when utilizing a regulon approach. During growth on the native carbon sources glucose, sucrose, and

raffinose, the lower fitness likely results from the burden of constitutive GAL regulon activation.

As we previously observed with xylose [18], using Gal3p$^{MC}$ to implement a regulon approach for

growth on arabinose and cellobiose resulted in faster growth rates and higher final cell densities compared

to the traditional constitutive overexpression approach despite expressing the same set of catabolic genes.

Significantly, the growth rates achieved using this rational approach are comparable or superior to the

highest aerobic growth rates for these substrates identified in the literature (**Table S1**) without the need

for time-consuming and/or expensive experiments like ALE (adaptive laboratory evolution), functional

genomics, and reverse genetics, or even deletion of genes previously found to mitigate stress of growth on

pentoses (e.g., *PHO13, ALD6, ASK10, ADH6, COX4, CYC8*, etc. [51]). The success of a semi-synthetic

regulon approach in all the single-sugar conditions motivated us to explore their co-utilization as the

simultaneous and complete co-utilization of the constituent sugars in lignocellulosic hydrolysates has

been identified as essential to their valorization [52]. One of the major challenges of engineering the co-

utilization of lignocellulosic sugars is that the presence of glucose exerts carbon catabolite repression

(CCR) in many organisms, hindering their ability to metabolize other sugars until glucose is depleted and

repression is relieved [53,54]. However, Jin & Cate et al. found that CCR could be avoided by implementing

a heterologous pathway for cellobiose, a glucose disaccharide that does not trigger CCR [33,55]. We

demonstrate that using Gal3p$^{MC}$ to implement a regulon approach enabled rapid and complete co-

utilization of three non-native carbon sources – xylose, arabinose, and cellobiose – in a single strain in

only ~48 h. Despite being evolved for the catabolism of galactose, these results reinforce the universality

and robustness of the metabolic coordination effected by the GAL regulon for engineering synthetic

heterotrophy as well as the plasticity of yeast metabolism to diverse substrates. However, our findings do

not discount the importance of existing metabolic engineering tools and strategies. For example, complete

utilization of pentoses during co-utilization required balancing the expression levels of pathway genes.

Additionally, while Gal3p$^{MC}$ enables a regulon approach to be rapidly adapted to structurally diverse,

non-native carbon sources, it is dependent on the development and importation of an efficient catabolic

module. This requirement limits its immediate application to attractive renewable feedstocks like C1

compounds (i.e., $CO_2$ [56], formate, methanol [57], methane [58]) as efforts to engineer their assimilation in *S.*

*cerevisiae* remain nascent. Researchers have noted the potential of platform strains that maximize the

concentration of a key precursor to speed the construction and optimization of biosynthetic pathways [59].

We envision that the integration of substrate catabolism with broader metabolism characteristic of a

regulon approach could similarly accelerate the process of engineering catabolism by reducing the

number of iterative, semi-rational genetic interventions currently needed to engineer efficient growth on a

given non-native substrate. Together, these results establish the metabolic coordination of a regulon

approach as a fast, flexible, and effective strategy for engineering synthetic heterotrophy and provide a

tool (Gal3p$^{MC}$) for adapting it to growth on structurally diverse non-native substrates.

## MATERIALS AND METHODS

### Strains and Plasmids

A full list of strains and plasmids used in this study can be found in **Table S2** and **Table S3**, respectively.

### Media and Transformation

Yeast strains were grown at 30 °C in either defined Synthetic Complete (SC) (Yeast nitrogen base (1.67 g/L), ammonium sulfate (5 g/L), complete supplement mixture without His, Leu, Ura and Trp (0.6 g/L) (Sunrise Science Products, Inc.) or complex Yeast Peptone (YP) medium (20 g/L Yeast Extract, 40 g/L Casein Peptone, 200 mg adenine) medium. SC media was augmented with appropriate nutrients to select for maintenance of plasmids carrying a given auxotrophy. Luria Bertani (LB) broth and LB agar plates with 100 μg/mL of ampicillin when required were used for all *E. coli* propagation and transformation experiments. Commercial *E. coli* strain NEB-5α (New England Biolabs) was used for MES transformation when constructing plasmids as well as for plasmid propagation. Upon isolation, plasmids were sequenced before being transformed into yeast strains using the protocol of Gietz [60] and recovered upon agar plates with the appropriate auxotrophy.

### Strain Construction

The yeast strain W303-1a was used for constructing all the strains used in the study. The generation of knockout strain VEG16 from W303-1a was described previously [18]. All subsequent strains were derived from VEG16 via chromosomal integrations using a previously described set of disintegrator plasmids and associated protocols [61].

### Plasmid Construction

*S. cerevisiae* promoter, gene, and terminator sequences were amplified from the genome using Phusion Polymerase (ThermoFisher Scientific). *XYLA*3* DNA sequence was provided by Prof. Hal S. Alper (University of Texas at Austin). *GAL2^{2.1}* DNA sequence was provided by Prof. Bernard Hauer

(University of Stuttgart, Stuttgart, Germany). Cellobiose utilization genes *cdt-1* and *gh1-1* were provided

by Prof. Huimin Zhao (University of Illinois at Urbana-Champaign). Synthesis of DNA primers and

Sanger sequencing of plasmid DNA was outsourced to Genewiz. Substrate utilization plasmids were

created by combining promoter, catabolic gene, and terminator sequences into previously described yeast

shuttle vectors containing either *CEN6* [62] or *2μ* [63] origins. All plasmids were constructed using either

NEBuilder HiFi DNA Assembly or restriction enzyme digest and ligation using reagents purchased from

New England Biolabs.

**Quantifying Promoter Dynamics via Fluorescence**

468    Overnight inoculums were grown in 200 µL in 96-well plates containing the required dropout SC medium

469    with glucose (2%) for 24 hours. Cultures were then subcultured (5 μL) into 200 µL SC medium

470    containing the appropriate carbon source (2%) and returned to the incubator. Fluorescence (excitation at

471    488 nm and emission at 525 nm) and $OD_{600}$ were measured at numerous time points in a Spectramax M3

472    spectrophotometer to obtain Fluorescence/$OD_{600}$.

**Molecular Dynamics (MD) Simulations**

Extended molecular dynamics and metadynamics simulation were conducted on three systems: $Gal3p^{WT}$

complexed with ATP and $Mg^{2+}$ ($WT^{Gal-}$), $Gal3p^{WT}$ complexed with galactose, ATP, and $Mg^{2+}$ ($WT^{Gal+}$),

and $Gal3p^{MC}$ complexed with ATP & $Mg^{2+}$ ($MC^{Gal-}$). To establish a common simulation starting point for

the three systems, each of the protein structures was modeled into the closed state via homology modeling

using Swiss Model [64] based on the experimentally determined closed state structure (PDB ID: 3V2U) as a

template (**Figure S11**). Side chain conformations of the template were retained when generating models.

Ramachandran plot was used to confirm that none of the side chains were in disallowed regions (**Figure**

**S12**). A comparison of the crystal structure (PDB ID: 3V2U) with the homology model revealed low Root

Mean Square Deviation (RMSD) of 0.09 (**Figure S13**). Molecular dynamic simulation of the three

different experimental setups was performed for 1000 ns using GROMACS–2019.4 [65]. The molecules

were parameterized using antechamber package from AMBER [66]. Quantum polarized charges were

derived using the GAMESS package [67] with 6-31G basis set and B3LYP level of theory AMBER99SB [66]

force field was used to perform a simulation with the TIP3p water model. To solvate the complex, the

TIP3P water model with a triclinic box volume of 1000 nm$^3$ was used to mimic or replace the natural

environment around the protein. The system was neutralized by adding Na$^+$ ions to the system, 16574

water molecules and ions overall. Energy minimization of the system was performed using the steepest

descent algorithm with convergence energy cut-off of 1000 kJ mol$^{-1}$ with 1000 steps. Equilibration of the

system was performed using NVT and NPT, where NVT is for constant volume and temperature and NPT

is for constant pressure. The NVT and NPT were carried out for 500 PS with 300 K temperature using a

Berendsen thermostat and the system was prepared for Molecular dynamic simulation for 1 µs at 300 K

temperature. The system was made to run for 1000 ns, for every 100 ns the trajectory was saved for

analysis. The first 100 ns were excluded for every analysis. The distance between C$_\alpha$ atoms of L370 and

D104 has been calculated across the simulation 1000 ns long trajectory for all three proteins to know the

variation in the movement of two domain regions across the time. RMSD (Root mean square Deviation)

and RMSF (Root mean square fluctuation) of the backbone atoms were calculated over time. Using

WORDOM tools [68], cross-correlation studies of residue-residue displacements along the trajectory were

calculated for each system to quantify dynamic changes of across the moving domains. Similarly,

principal component analysis (PCA) was calculated for each 100-1000 ns trajectory and porcupine plots

were generated to analyze the intensity and movement of C$_\alpha$ atom. Visualization of structures was done

using UCSF Chimera [69] and Visual Molecular Dynamics (VMD) [70].

**Metadynamics Simulations**

Metadynamics-based simulations use a potential bias to move the system out of a local minimum, which

is then used to explore the resulting free energy landscape (FES). Metadynamics-based simulations were

conducted using the PLUMED software package [71] and GROMACS toolkit for all three experimental

setups (WT$^{Gal-}$, WT$^{Gal+}$, and MC$^{Gal-}$) after thorough equilibration obtained from NVT, NPT ensembles. C1,

C2, and C3 are Centers of Mass (COMs) determined for the three regions of Gal3p that fall within the

residues regions as shown in **Figure S5A**. One of the Collective Variables (CV) for metadynamics

simulation was defined as the distance between C1 and C2. The other CV was defined as the angle

between the intersection point of the vectors that pass through C1, C3 and C2, C3, respectively. Different

simulations in this study were performed for 50 ns with a bias factor set to 8.0, the temperature set to 300

K, and the results were calculated accordingly. The free energy landscape was explored as the distance

between the two domains and the change in the angle formed by the three COMs were varied. A

superposition of modeled closed form and apo form was used to visualize the structural changes (**Figure**

**S14**).

**Gal3p-Gal80p Docking Studies**

Crystal structure of Gal80p complexed with the closed Gal3p structure (PDB ID 3V2U) was used as the

reference for protein-protein docking studies. Protein-protein docking studies were conducted using PatchDock

[72,73] and then refined using Fiberdock [74,75]. The minimum energy conformations of $WT^{Gal+}$, $WT^{Gal-}$, and $MC^{Gal-}$

attained from the metadynamics simulations were docked with Gal80p and the complexes obtained from

protein-protein docking studies were simulated using GROMACS–2019.4 [76] for 50 ns ×2 to better equilibrate

the complexes that were obtained. The coordinates of the complexes were sampled every 0.02 ns and the

binding affinity in kcal/mol between the Gal3p and Gal80p structures for all three complexes was calculated

using the molecular mechanics/Poisson-Boltzmann surface area (MM/PBSA) method [77,78].

**Growth Studies**

Overnight inoculums were grown in the required dropout SC medium with 2% sucrose (unless explicitly

stated to be glucose/raffinose) for 24 h. The culture was washed twice in the growth medium (either SC or

YP containing 2% of the target sugar) and resuspended at an initial $OD_{600}$ of 0.1 in 250 mL shake flasks

containing 20 mL of media. Cell growth $OD_{600}$ measurement was checked at frequent time intervals (3–6

h) on SpectraMax M3 spectrophotometer (Molecular Devices). Growth rates (μ) for each biological

replicate were quantified in Microsoft Excel using an exponential curve equation fit to at least three time

points during logarithmic growth phase. These values were used to calculate the average growth rate and

the standard error for a given condition.

**Residual Substrate Quantification**

Residual substrate during co-utilization studies was measured using an Agilent HPLC system equipped

with a Hi-Plex H-column and detected using 1260 Agilent ELSD detector. Mobile phase was 0.1%

trifluoroacetic acid (TFA) with a flow rate of 0.6 mL/min. The ELSD detector's nebulizer and

evaporation temperature were set at 30 °C and nitrogen flow rate at 1.6 SLM (standard liter per minute).

Using these settings, we observed characteristic retention times for cellobiose (9.3 min), arabinose (12.6

min), and xylose (11.8 min).

**DATA AVAILABILITY**

All plasmids and strains used in this study can be obtained from the corresponding author under a

material transfer agreement. Source data are provided with this paper.

**ACKNOWLEDGEMENTS**

The authors would like to thank current and former Nair lab members for helpful discussions. This work

was supported by NIH grant #DP2HD91798, #R21HD105934, and NSF grant #1935354, and Tufts

Launchpad | Accelerator to N.U.N.

**AUTHOR CONTRIBUTIONS**

N.U.N., S.F.S., V.D.T., and V.E.G. conceived and designed the research project. S.F.S., A.S., and N.U.N.

co-wrote the manuscript. S.F.S., A.S., V.D.T., T.C.C., V.E.G., D.M.S., and T.B. performed the

experiments. S.F.S., A.S., T.B., P.K.R., and N.U.N. analyzed the data. All the authors have reviewed the

manuscript and approved it for submission.

## COMPETING INTERESTS

None.

**REFERENCES**

1. Langholtz, M., Stokes, B. & Eaton, L. 2016 billion-ton report: Advancing domestic resources for a thriving bioeconomy (Executive Summary). *Industrial Biotechnology* **12**, 282–289 (2016).

2. Rogers, J. N. *et al.* An assessment of the potential products and economic and environmental impacts resulting from a billion ton bioeconomy. *Biofuels, Bioproducts and Biorefining* **11**, 110–128 (2017).

3. Nielsen, J. & Keasling, J. D. Engineering Cellular Metabolism. *Cell* **164**, 1185–1197 (2016).

4. Latimer, L. N. *et al.* Employing a combinatorial expression approach to characterize xylose utilization in Saccharomyces cerevisiae. *Metab Eng* **25**, 20–29 (2014).

5. Kobayashi, Y., Sahara, T., Ohgiya, S., Kamagata, Y. & Fujimori, K. E. Systematic optimization of gene expression of pentose phosphate pathway enhances ethanol production from a glucose/xylose mixed medium in a recombinant Saccharomyces cerevisiae. *AMB Express* **8**, 1–11 (2018).

6. Kim, S. R. *et al.* Rational and evolutionary engineering approaches uncover a small set of genetic changes efficient for rapid xylose fermentation in Saccharomyces cerevisiae. *PLoS One* **8**, e57048 (2013).

7. Ni, H., Laplaza, J. M. & Jeffries, T. W. Transposon mutagenesis to improve the growth of recombinant Saccharomyces cerevisiae on D-xylose. *Appl Environ Microbiol* **73**, 2061–6 (2007).

8. Mukherjee, V., Lind, U., St. Onge, R. P., Blomberg, A. & Nygård, Y. A CRISPR Interference Screen of Essential Genes Reveals that Proteasome Regulation Dictates Acetic Acid Tolerance in Saccharomyces cerevisiae. *mSystems* **6**, (2021).

9. Unrean, P., Gätgens, J., Klein, B., Noack, S. & Champreda, V. Elucidating cellular mechanisms of Saccharomyces cerevisiae tolerant to combined lignocellulosic-derived inhibitors using high-Throughput phenotyping and multiomics analyses. *FEMS Yeast Res* **18**, 1–10 (2018).

10. HamediRad, M., Lian, J., Li, H. & Zhao, H. RNAi Assisted Genome Evolution Unveils Yeast Mutants with Improved Xylose Utilization. *Biotechnol Bioeng* (2018) doi:10.1002/bit.26570.

11. Chen, Y., Stabryla, L. & Wei, N. Improved acetic acid resistance in Saccharomyces cerevisiae by overexpression of the WHI2 gene identified through inverse metabolic engineering. *Appl Environ Microbiol* **82**, 2156–2166 (2016).

12. Ha, S. J. *et al.* Engineered Saccharomyces cerevisiae capable of simultaneous cellobiose and xylose fermentation. *Proc Natl Acad Sci U S A* **108**, 504–509 (2011).

13. Zhou, H., Cheng, J.-S., Wang, B., Fink, G. R. & Stephanopoulos, G. Xylose isomerase overexpression along with engineering of the pentose phosphate pathway and evolutionary engineering enable rapid xylose utilization and ethanol production by Saccharomyces cerevisiae. *Metab Eng* **14**, 611–622 (2012).

14.  Sanchez, R. G. *et al.* Improved xylose and arabinose utilization by an industrial recombinant.
*Biotechnol Biofuels* **3**, (2010).

15.  Lavy, T., Kumar, P. R., He, H. & Joshua-Tor, L. The Gal3p transducer of the GAL regulon interacts
with the Gal80p repressor in its ligand-induced closed conformation. *Genes Dev* **26**, 294–303
(2012).

16.  Griffin, T. J. *et al.* Complementary profiling of gene expression at the transcriptome and
proteome levels in Saccharomyces cerevisiae. *Mol Cell Proteomics* **1**, 323–333 (2002).

17.  Ren, B. *et al.* Genome-wide location and function of DNA binding proteins. *Science (1979)* **290**,
2306–2309 (2000).

18.  Endalur Gopinarayanan, V. & Nair, N. U. A semi-synthetic regulon enables rapid growth of yeast
on xylose. *Nat Commun* **9**, (2018).

19.  Trivedi, V. D., Sullivan, S. F., Choudhury, D., Endalur Gopinarayanan, V. & Nair, N. U. Integration
of metabolism and regulation reveals rapid adaptability to growth on non-native substrates.
*bioRxiv* (2022) doi:10.1101/2022.05.21.492926.

20.  Reimand, J., Vaquerizas, J. M., Todd, A. E., Vilo, J. & Luscombe, N. M. Comprehensive reanalysis
of transcription factor knockout expression data in Saccharomyces cerevisiae reveals many new
targets. *Nucleic Acids Res* **38**, 4768–4777 (2010).

21.  Peng, B., Williams, T. C., Henry, M., Nielsen, L. K. & Vickers, C. E. Controlling heterologous gene
expression in yeast cell factories on different carbon substrates and across the diauxic shift: A
comparison of yeast promoter activities. *Microb Cell Fact* **14**, 1–11 (2015).

22.  Deng, J. *et al.* A synthetic promoter system for well-controlled protein expression with different
carbon sources in Saccharomyces cerevisiae. *Microb Cell Fact* **20**, 1–10 (2021).

23.  Blank, T. E., Woods, M. P., Lebo, C. M., Xin, P. & Hopper, J. E. Novel Gal3 proteins showing altered
Gal80p binding cause constitutive transcription of Gal4p-activated genes in Saccharomyces
cerevisiae. *Mol Cell Biol* **17**, 2566–2575 (1997).

24.  Gopinarayanan, V. E. & Nair, N. U. A semi-synthetic regulon enables rapid growth of yeast on
xylose. *Nat Commun* **9**, (2018).

25.  Bhat, P. J. & Hopper, J. E. Overproduction of the GAL1 or GAL3 protein causes galactose-
independent activation of the GAL4 protein: evidence for a new model of induction for the yeast
GAL/MEL regulon. *Mol Cell Biol* **12**, 2701–2707 (1992).

26.  Malakar, P. & Venkatesh, K. V. GAL regulon of Saccharomyces cerevisiae performs optimally to
maximize growth on galactose. *FEMS Yeast Res* **14**, 346–356 (2014).

27.  Venturelli, O. S., El-Samad, H. & Murray, R. M. Synergistic dual positive feedback loops
established by molecular sequestration generate robust bimodal response. *Proc Natl Acad Sci U S*
*A* **109**, (2012).

28. Lee, S. M., Jellison, T. & Alper, H. S. Directed evolution of xylose isomerase for improved xylose
catabolism and fermentation in the yeast Saccharomyces cerevisiae. *Appl Environ Microbiol* **78**,
5708–5716 (2012).

29. Quarterman, J. *et al.* Rapid and efficient galactose fermentation by engineered Saccharomyces
cerevisiae. *J Biotechnol* **229**, 13–21 (2016).

30. Torchia, T. E., Hamilton, R. W., Cano, C. L. & Hopper, J. E. Disruption of regulatory gene GAL80 in
Saccharomyces cerevisiae: effects on carbon-controlled regulation of the galactose/melibiose
pathway genes. *Mol Cell Biol* **4**, 1521–1527 (1984).

31. Reznicek, O. *et al.* Improved xylose uptake in Saccharomyces cerevisiae due to directed evolution
of galactose permease Gal2 for sugar co-consumption. *J Appl Microbiol* **119**, 99–111 (2015).

32. Wisselink, H. W. *et al.* Engineering of Saccharomyces cerevisiae for efficient anaerobic alcoholic
fermentation of L-arabinose. *Appl Environ Microbiol* **73**, 4881–91 (2007).

33. Galazka, J. M. *et al.* Cellodextrin Transport in Yeast for Improved Biofuel Production. *Science
(1979)* **330**, 84–86 (2010).

34. Eiteman, M. A., Lee, S. A. & Altman, E. A co-fermentation strategy to consume sugar mixtures
effectively. *J Biol Eng* **2**, 1–8 (2008).

35. Verhoeven, M. D., de Valk, S. C., Daran, J. M. G., van Maris, A. J. A. & Pronk, J. T. Fermentation of
glucose-xylose-Arabinose mixtures by a synthetic consortium of single-sugar-fermenting
Saccharomyces cerevisiae strains. *FEMS Yeast Res* **18**, 1–12 (2018).

36. Flores, A. D. *et al.* Catabolic Division of Labor Enhances Production of D-Lactate and Succinate
From Glucose-Xylose Mixtures in Engineered Escherichia coli Co-culture Systems. *Front Bioeng
Biotechnol* **8**, 1–14 (2020).

37. Kim, B., Du, J., Eriksen, D. T. & Zhao, H. Combinatorial design of a highly efficient xylose-utilizing
pathway in Saccharomyces cerevisiae for the production of cellulosic biofuels. *Appl Environ
Microbiol* **79**, 931–941 (2013).

38. Protzko, R. J. *et al.* Engineering Saccharomyces cerevisiae for co-utilization of d-galacturonic acid
and d-glucose from citrus peel waste. *Nat Commun* **9**, 5059 (2018).

39. Ajikumar, P. K. *et al.* Isoprenoid pathway optimization for Taxol precursor overproduction in
Escherichia coli. *Science (1979)* **330**, 70–74 (2010).

40. Smanski, M. J. *et al.* Functional optimization of gene clusters by combinatorial design and
assembly. *Nat Biotechnol* **32**, 1241–1249 (2014).

41. Kurland, C. G. & Dong, H. Bacterial growth inhibition by overproduction of protein. *Mol Microbiol*
**21**, 1–4 (1996).

42. Burg, J. M. *et al.* Large-scale bioprocess competitiveness: the potential of dynamic metabolic
control in two-stage fermentations. *Curr Opin Chem Eng* **14**, 121–136 (2016).

43.  Hartline, C. J., Schmitz, A. C., Han, Y. & Zhang, F. Dynamic control in metabolic engineering:
Theories, tools, and applications. *Metab Eng* **63**, 126–140 (2021).

44.  Zhang, F., Carothers, J. M. & Keasling, J. D. Design of a dynamic sensor-regulator system for
production of chemicals and fuels derived from fatty acids. *Nature Biotechnology 2012 30:4* **30**,
354–359 (2012).

45.  Xiao, Y., Bowen, C. H., Liu, D. & Zhang, F. Exploiting nongenetic cell-to-cell variation for enhanced
biosynthesis. *Nature Chemical Biology 2016 12:5* **12**, 339–344 (2016).

46.  Xu, P., Li, L., Zhang, F., Stephanopoulos, G. & Koffas, M. Improving fatty acids production by
engineering dynamic pathway regulation and metabolic control. *Proc Natl Acad Sci U S A* **111**,
11299–11304 (2014).

47.  Dahl, R. H. *et al.* Engineering dynamic pathway regulation using stress-response promoters.
*Nature Biotechnology 2013 31:11* **31**, 1039–1046 (2013).

48.  Alvarez-Gonzalez, G. & Dixon, N. Genetically encoded biosensors for lignocellulose valorization.
*Biotechnology for Biofuels 2019 12:1* **12**, 1–14 (2019).

49.  Castrillo, J. I. *et al.* Growth control of the eukaryote cell: A systems biology study in yeast. *J Biol* **6**,
(2007).

50.  Brauer, M. J. *et al.* Coordination of growth rate, cell cycle, stress response, and metabolic activity
in yeast. *Mol Biol Cell* **19**, 352–67 (2008).

51.  Li, X. *et al.* Metabolic network remodelling enhances yeast's fitness on xylose using aerobic
glycolysis. *Nat Catal* **4**, (2021).

52.  Gao, M., Ploessl, D. & Shao, Z. Enhancing the Co-utilization of Biomass-Derived Mixed Sugars by
Yeasts. *Front Microbiol* **9**, 1–21 (2019).

53.  Boles, E. & Hollenberg, C. P. The molecular genetics of hexose transport in yeasts. *FEMS*
*Microbiol Rev* **21**, 85–111 (1997).

54.  Roca, C., Haack, M. B. & Olsson, L. Engineering of carbon catabolite repression in recombinant
xylose fermenting Saccharomyces cerevisiae. *Appl Microbiol Biotechnol* **63**, 578–583 (2004).

55.  Li, S. *et al.* Overcoming glucose repression in mixed sugar fermentation by co-expressing a
cellobiose transporter and a β-glucosidase in Saccharomyces cerevisiae. *Mol Biosyst* **6**, 2129–
2132 (2010).

56.  Liu, Z., Wang, K., Chen, Y., Tan, T. & Nielsen, J. Third-generation biorefineries as the means to
produce fuels and chemicals from CO2. *Nat Catal* **3**, 274–288 (2020).

57.  Antoniewicz, M. R. Synthetic methylotrophy: Strategies to assimilate methanol for growth and
chemicals production. *Curr Opin Biotechnol* **59**, 165–174 (2019).

58.  Clomburg, J. M., Crumbley, A. M. & Gonzalez, R. Industrial biomanufacturing: The future of
chemical production. *Science (1979)* **355**, (2017).

59.    Nielsen, J. Yeast cell factories on the horizon: Metabolic engineering in yeast gets increasingly
702           more versatile. *Science (1979)* **349**, 1050–1051 (2015).

703    60.    Gietz, R. D. Yeast Transformation by the LiAc/SS Carrier DNA/PEG Method. in *Yeast Protocols* vol.
704           1163 33–44 (2014).

705    61.    Sadowski, I., Su, T.-C. & Parent, J. Disintegrator vectors for single-copy yeast chromosomal
706           integration. *Yeast* **24**, 447–455 (2007).

707    62.    Frazer, L. N. & O'Keefe, R. T. A new series of yeast shuttle vectors for the recovery and
708           identification of multiple plasmids from Saccharomyces cerevisiae. *Yeast* **24**, 777–789 (2007).

709    63.    Christianson, T. W., Sikorski, R. S., Dante, M., Shero, J. H. & Heiter, P. Multifunctional yeast high-
710           copy shuttle vectors. *Gene* **110**, 119–122 (1992).

711    64.    Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and complexes.
712           *Nucleic Acids Res* **46**, W296–W303 (2018).

713    65.    Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-level
714           parallelism from laptops to supercomputers. *SoftwareX* **1–2**, 19–25 (2015).

715    66.    Lindorff-Larsen, K. *et al.* Improved side-chain torsion potentials for the Amber ff99SB protein
716           force field. *Proteins: Structure, Function, and Bioinformatics* **78**, 1950–1958 (2010).

717    67.    Guest, M. F. *et al.* The GAMESS-UK electronic structure package: algorithms, developments and
718           applications. *https://doi.org/10.1080/00268970512331340592* **103**, 719–747 (2007).

719    68.    Seeber, M. *et al.* Wordom: A user-friendly program for the analysis of molecular structures,
720           trajectories, and free energy surfaces. *J Comput Chem* **32**, 1183–1194 (2011).

721    69.    Pettersen, E. F. *et al.* UCSF Chimera—A visualization system for exploratory research and analysis.
722           *J Comput Chem* **25**, 1605–1612 (2004).

723    70.    Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. *J Mol Graph* **14**, 33–38
724           (1996).

725    71.    Abrams, J. B. & Tuckerman, M. E. Efficient and direct generation of multidimensional free energy
726           surfaces via adiabatic dynamics without coordinate transformations. *Journal of Physical*
727           *Chemistry B* **112**, 15742–15757 (2008).

728    72.    Duhovny, D., Nussinov, R. & Wolfson, H. J. Efficient unbound docking of rigid molecules. *Lecture*
729           *Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture*
730           *Notes in Bioinformatics)* **2452**, 185–200 (2002).

731    73.    Schneidman-Duhovny, D., Inbar, Y., Nussinov, R. & Wolfson, H. J. PatchDock and SymmDock:
732           servers for rigid and symmetric docking. *Nucleic Acids Res* **33**, W363–W367 (2005).

733    74.    Mashiach, E., Nussinov, R. & Wolfson, H. J. FiberDock: Flexible induced-fit backbone refinement
734           in molecular docking. *Proteins* **78**, 1503–19 (2010).

75. Mashiach, E., Nussinov, R. & Wolfson, H. J. FiberDock: a web server for flexible induced-fit
backbone refinement in molecular docking. *Nucleic Acids Res* **38**, W457-61 (2010).

76. Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-level
parallelism from laptops to supercomputers. *SoftwareX* **1–2**, 19–25 (2015).

77. Kumari, R., Kumar, R. & Lynn, A. *g_mmpbsa* —A GROMACS Tool for High-Throughput MM-PBSA
Calculations. *J Chem Inf Model* **54**, 1951–1962 (2014).

78. Baker, N. A., Sept, D., Joseph, S., Holst, M. J. & McCammon, J. A. Electrostatics of nanosystems:
Application to microtubules and the ribosome. *Proceedings of the National Academy of Sciences*
**98**, 10037–10041 (2001).

744