Towards universal synthetic heterotrophy using a metabolic coordinator

Sean F. Sullivan[a], Anuj Shetty[b], Tharun Bharadwaj[b], Naveen Krishna[b], Vikas D. Trivedi[a,c], Venkatesh Endalur Gopinarayanan[a], Daniel M. Sellers[a], Pravin Kumar R.[b], and Nikhil U. Nair[a]*

[a] Department of Chemical & Biological Engineering, Tufts University, Medford, MA 02155

[b] Kcat Enzymatic Private Limited, Bengaluru, Karnataka, India 560005

[c] Department of Structural Biology and Center for Data Driven Discovery, St. Jude Children's Research Hospital, Memphis, TN

* Corresponding author: nikhil.nair@tufts.edu; @nair_lab

ABSTRACT

Engineering the utilization of non-native substrates, or synthetic heterotrophy, in proven industrial microbes such as *Saccharomyces cerevisiae* represents an opportunity to valorize plentiful and renewable sources of carbon and energy as potential inputs to biotechnological processes. We previously demonstrated that activation of the galactose (GAL) regulon, a regulatory structure used by this yeast to coordinate substrate utilization with biomass formation during growth on galactose, during growth on the non-native substrate xylose results in a vastly altered gene expression profile and faster growth compared with constitutive overexpression of the same heterologous catabolic pathway. However, this effort involved the creation of a xylose-inducible variant of Gal3p (Gal3p$^{\text{Syn4.1}}$), the sensor protein of the GAL regulon, preventing this semi-synthetic regulon approach from being easily adapted to additional non-native substrates. Here, we report the construction of a variant Gal3p$^{\text{MC}}$ (metabolic coordinator) that exhibits robust GAL regulon activation in the presence of structurally diverse substrates and recapitulates the dynamics of the native system. Multiple molecular modeling studies confirm that Gal3p$^{\text{MC}}$ occupies conformational states corresponding to galactose-bound Gal3p in an inducer-independent manner. Using Gal3p$^{\text{MC}}$ to test a regulon approach to the assimilation of the non-native lignocellulosic sugars xylose, arabinose, and cellobiose yields higher growth rates and final cell densities when compared with a constitutive

overexpression of the same set of catabolic genes. The subsequent demonstration of rapid and complete co-

utilization of all three non-native substrates suggests that Gal3p^{MC}-mediated dynamic global gene

expression changes by GAL regulon activation may be universally beneficial for engineering synthetic

heterotrophy.

KEYWORDS: Gene regulation, sustainability, synthetic biology, metabolic engineering, substrate

utilization, molecular dynamics

INTRODUCTION

The adoption of abundant and renewable substrates as inputs for biotechnology will be essential for creating

a sustainable, circular bioeconomy [1,2]. But the inability of industrially important microbes, like

Saccharomyces cerevisiae (henceforth, yeast)*, to utilize many potential substrates poses a major hurdle to

realizing this goal. The existing paradigm for engineering the assimilation of non-natives substrates (i.e.,

synthetic heterotrophy) in yeast begins with the identification and constitutive overexpression of catabolic

genes that enable the substrate to enter central carbon metabolism (CCM) where it is expected to be

transformed into the key primary metabolites used for growth and biosynthesis. However, this method

ignores the tight regulation of CCM and how these resources are distributed to accomplish cellular

objectives [3]. The consequences of this oversight are reflected in the need for subsequent interventions,

including flux balancing [4–6], functional genomics [7–11], and adaptive lab evolution [12–14], undertaken to

improve growth and/or substrate utilization rate by resolving conflicts with cellular processes. Moreover,

existing efforts are often focused on a specific substrate or small set of substrates – a siloed approach that

makes it hard to translate findings to other substrates and even disincentivizes holistic thinking about the

limits of metabolic plasticity/adaptability in this yeast. This has, at least in part, motivated domestication

of other yeasts with broader substrate ranges (e.g., *Scheffersomyces stipitis, Kluyveromyces marxianus*) for

biomanufacturing applications.

An alternate approach that we advocate here, is that instead of combating natural regulation we

should leverage existing regulatory structures that have evolved to coordinate complex phenotypes like

substrate utilization with biomass formation and metabolite synthesis. One such system is the galactose

(GAL) regulon that yeast uses to coordinate substrate catabolism with global metabolism during growth on

the native substrate galactose. In this system, the interaction of galactose with sensor protein Gal3p enables

it to relieve the repression of Gal80p on the transcription factor Gal4p via protein-protein interactions [15].

Once freed from repression, Gal4p – the master activator of the GAL regulon – binds to its cognate

Upstream Activating Sequences (UAS) to directly induce transcription of galactose catabolic genes (Leloir

pathway) and indirectly modulates the transcript and/or protein abundance of numerous growth-associated

genes (**Figure 1A**) [16,17]. We previously reported the construction of a variant of sensor protein Gal3p

(Gal3p$^{Syn4.1}$) that strongly activates the GAL regulon by the pentose sugars xylose and arabinose [18,19]. By

placing heterologous genes for either xylose or arabinose catabolism under the control of GAL-responsive

(Leloir pathway) promoters, the GAL regulon could be adapted for growth on these non-native substrates.

Upon comparing this regulon-coordinated (REG) approach to growth with simple constitutive

overexpression of the same catabolic genes, which we term a constitutive (CONS) approach, we found that

our semi-synthetic regulon yielded superior growth rates on both xylose ($\mu = 0.24$ h^{-1} vs. 0.11 h^{-1}) and

arabinose ($\mu = 0.27$ h^{-1} vs. 0.06 h^{-1}), respectively with minimal metabolic engineering [19]. Significantly, a

transcriptomic comparison showed relative upregulation of genes associated with cell division and

mitochondrial biogenesis in the REG strain while the CONS strain showed upregulation of stress response

and starvation-associated genes [18]. This suggests that the GAL regulon dynamically reshapes the cellular

response for growth through direct and/or indirect action of Gal3p-Gal80p-Gal4p to potentiate the cells for

rapid growth [17,20], irrespective of the identity of the available substrate.

In this study, we demonstrate that, in addition to the known genome-wide changes in gene

expression that occur upon GAL regulon activation, the expression from GAL-inducible promoters is

dynamic and appears to be coordinated with growth whereas expression from constitutive promoters is

constant. Hypothesizing that both aspects of this metabolic coordination may be universally beneficial for

the assimilation of non-native substrates, we sought to develop this semi-synthetic regulon system into a

platform approach that enables the rapid engineering of efficient synthetic heterotrophy without having to

continually re-engineer substrate-specific induction. To do so, we created a variant of sensor protein that

we term Gal3p^{MC} (i.e., metabolic coordinator) and demonstrate that it, unlike the wild-type Gal3p

(Gal3p^{WT}), can strongly activate the GAL regulon on numerous structurally diverse substrates. We show

that Gal3p^{MC} can recapitulate the dynamic activation of the native regulon and, through the use of molecular

dynamics and metadynamics simulations, that the mutations it carries enable it to do so in an inducer-

independent manner. We found that this substrate-agnostic system retains the benefits of substrate-specific

activation without any undue burden when engineering synthetic heterotrophy. We also show that using

Gal3p^{MC} to implement a REG approach to growth on three non-native substrates – xylose, arabinose, and

cellobiose – yields superior performance when compared to a CONS approach. Finally, we provide a first

demonstration that a single strain expressing Gal3p^{MC} is capable of rapid, simultaneous, and complete

utilization of all three non-native substrates concurrent with rapid growth – paving the way for a universal

synthetic heterotrophy platform.

RESULTS

Expression under the GAL regulon is dynamic and recapitulates growth phase.

Existing efforts to engineer synthetic heterotrophy generally utilize strong, constitutive promoters to control

the expression of the necessary catabolic genes. Constitutive promoters are natively able to recruit cellular

transcriptional machinery to yield a roughly consistent expression level independent of cellular state and

environmental context. This 'always on' phenotype contrasts with the control of catabolic gene expression

observed in many regulatory structures that coordinate native substrate utilization in response to nutrient

availability, including the GAL regulon of *S. cerevisiae*. To understand how these different approaches

could impact the efficiency of substrate utilization, we used a fluorescent reporter (*EGFP*) to quantify the

expression dynamics of several GAL-responsive (*GAL1p*, *GAL7p*, *GAL10p*, *GAL3p,* and *GAL80p*) and

Figure 1: Expression from GAL-inducible promoters is coordinated with growth. (A) Simplified regulatory overview of the GAL regulon. The binding of galactose to sensor protein Gal3p enables it to relieve the repression of Gal80p on Gal4p via protein-protein interactions. Gal4p binds to its cognate Upstream Activating Sequences (UAS) to directly transcribe galactose catabolic genes as well as to indirectly reshape global metabolism for rapid growth. The regulon is also repressed in the presence of excess glucose, a regulatory logic that ensures the regulon is activated only in the presence of its target carbon source but not when a preferred carbon source is present. Expression of EGFP in wild-type strain W303-1a from **(B)** strong constitutive promoters (*TEF1p*, *TPI1p*, *TDH3p*, *GPM1p*) and **(C)** GAL-inducible promoters (*GAL1p*, *GAL7p*, *GAL10p*, *GAL3p*, *GAL80p*) at different time points during cultivation on galactose overlaid with cell density (OD$_{600}$). Each data point represents the average of three (for OD$_{600}$ values) or four (for fluorescence) biological replicates ± sd.

constitutive (*TPI1p*, *TEF1p*, *TDH3p*, and *GPM1p*) yeast promoters in a wild-type strain (W303-1a) that

contains the native GAL regulon. During growth on galactose, we found that expression from the

constitutive promoters was relatively stable across a 24-hour period (varying less than two-fold) (**Figure**

1B). In contrast, expression from GAL-responsive promoters starts lower (~50 % of constitutive promoters)

but increases by ~4-6 fold for the promoters that control the expression of the galactose catabolic (Leloir

pathway) genes (*GAL1p*, *GAL7p*, and *GAL10p*) while remaining relatively constant for the promoters

controlling regulatory genes (*GAL3p* and *GAL80p*) (**Figure 1C**). When we overlaid the OD_{600} profile, we

observed that expression from Leloir pathway promoters tracks with growth phase, with the lowest

expression levels occurring when the cells are in lag phase (t ~ 0 h), maximum expression during the

exponential growth phase, and finally diminishing expression as galactose is depleted and the cells enter

stationary phase (t > 20 h). During growth on glucose, the expression profile of the constitutive promoters

was largely unchanged (**Figure S1A**) while expression from GAL-inducible promoters is strongly repressed

because of carbon catabolite repression (**Figure S1B**).

Gal3p^{MC} activates the GAL regulon in the presence of structurally diverse carbon sources.

While the creation of a xylose-inducible sensor protein variant Gal3p$^{Syn4.1}$ enabled us to adapt the GAL

regulon for growth on xylose [18], we wanted to develop a method for rapidly applying a regulon approach

to other non-native substrates without having to repeatedly re-engineer substrate-specific sensing. To this

end, we added mutation F237Y [21] to Gal3p$^{Syn4.1}$ variant to yield a variant we term Gal3p^{MC} (<u>m</u>etabolic

<u>c</u>oordinator). To compare the ability of Gal3p^{WT} and Gal3p^{MC} to activate the regulon in the presence of a

given carbon source, we monitored the fluorescence resulting from a *GAL1p-yEGFP* reporter construct

(**Figure 2A**) in the presence of numerous structurally distinct carbon sources substrates – galactose,

arabinose, cellobiose, and xylose – in a strain (VEG16) which lacks the galactose sensing and catabolic

genes (*ΔGAL3; ΔGAL1; ΔGAL10; ΔGAL7; ΔGRE3)* but retains *GAL80* and *GAL4*. We performed this

experiment using two native carbon sources, raffinose (**Figure 2B**) or sucrose (**Figure S2**), to support cell

growth. We observed that Gal3p^{WT} strongly activates the regulon in the presence of its native inducer

galactose and to a lesser extent xylose and arabinose, likely due to the structural similarity between these

substrates. The indistinguishable signal between Gal3p^{WT} and the *ΔGAL3* control in presence of glucose

reflects carbon catabolite repression while the small increase in fluorescence on raffinose and cellobiose

likely reflects the weak activation known to result from overexpression of Gal3p^{WT} in the absence of glucose

[22].

Figure 2: Sensor protein variant Gal3pᴹᶜ enables robust GAL regulon activation in an inducer-independent manner. (A) Schematic of fluorescent reporter assay used to quantify GAL regulon activation by different *GAL3* variants. Cells expressing a given variant (ΔGAL3, Gal3pᵂᵀ, or Gal3pᴹᶜ) are grown in the presence of both native and non-native substrates. Regulon activation drives EGFP expression from the GAL-inducible promoter, *GAL1p*. **(B)** Comparison of normalized fluorescence resulting from interaction between *GAL3* variant and carbon source (2 % w/v). **(C)** Expression of EGFP from *TEF1p* and *GAL1p* during growth on galactose in a strain (SFS6) expressing Gal3pᴹᶜ to activate the regulon. Each data point represents the average of 5–6 biological replicates ± sd.

In contrast, Gal3pᴹᶜ activates the GAL regulon in the presence of all the non-native substrates to a similar

extent as Gal3pᵂᵀ on galactose while remaining repressible by glucose (**Figure 2B**), a result that suggests

Gal3pᴹᶜ may be capable of recapitulating the dynamic catabolic gene expression profile and genome-wide

expression changes known to support optimal growth on the native substrate galactose during growth on

non-native substrates [23]. To determine if regulon activation using Gal3pᴹᶜ remains dynamic, we constructed

a strain (SFS6) that contains copies of *GAL3ᴹᶜ* integrated into the chromosome under the control of both

GAL1p and *GAL3p*. This dual-feedback loop configuration mimics the organization of the native GAL

regulon and has been demonstrated to increase both the overall magnitude and the homogeneity of

activation to better effect a switch-like behavior among cells in the population [18,24]. As SFS6 lacks the Leloir

pathway (ΔGAL1/7/10), these genes were supplied under the control of their native promoters on a plasmid

(pRS423-GAL-REG) to facilitate growth on galactose. Using strain SFS6, we assessed the expression

profile of the constitutive *TEF1p-EGFP* and GAL-responsive *GAL1p-EGFP* constructs during growth on

galactose (**Figure 2C**). While expression from *TEF1p* was observed to increase by 2–3-fold at one time

point, it generally stayed within a two-fold range whereas expression from *GAL1p* appeared to mimic the

profile observed upon activation by the wild-type regulon (**Figure 1C**), exhibiting a >5-fold increase in

fluorescence.

Molecular dynamics (MD) and metadynamics simulations support inducer-independent regulon

activation by Gal3p^{MC}

Experiments have shown that the binding of Gal3p^{WT} to its ligand galactose induces and stabilizes a

conformational change from an 'open' to a 'closed' state that facilitates interaction with Gal80p and

subsequent de-repression of Gal4p [15]. However, mapping the mutations carried by Gal3p^{MC} onto a structure

of Gal3p^{WT} in complex with Gal80p (PDB: 3V2U) reveals that none are found within the ligand binding

pocket but rather in the Gal80p interface and hinge regions (**Figure S3**). Combined with the structural

diversity of substrates (pentoses, hexose, disaccharide, and trisaccharide) on which Gal3p^{MC} exhibits robust

regulon activation, observations suggest that Gal3p^{MC} may adopt a 'closed' conformation that enables it to

bind Gal80p and relieve repression on Gal4p without requiring interaction with a ligand/inducer. As the

transition between open and closed states is associated with the relative movement of two domains in

Gal3p^{WT}, while a third 'hinge' region remains relatively static, we decided to explore this inducer-

independent activation hypothesis using molecular dynamics simulations. To do so, we monitored the

distance between two residues (D104 and L370) found on opposing moving domains (**Figure 3A**) over the

course of 1000 ns molecular dynamics (MD) simulations of Gal3p^{WT} in complex with ATP and Mg^{2+}

without galactose (WT^{Gal-}), Gal3p^{WT} in complex with ATP, Mg^{2+}, and galactose (WT^{Gal+}), and Gal3p^{MC} in

complex ATP and Mg^{2+} without galactose (MC^{Gal-}). The resulting distance graph showed a dip in the

conformation for WT^{Gal-} around 320 ns into the simulation, reflecting a compaction of the active site not

Figure 3: Observations of 1000 ns scale molecular dynamics (MD) simulations. (A) Structural representation of Gal3p in the closed conformation with dotted lines representing the distance between residues L370 and D104, found on the opposing 'moving' domains (cyan- and pink-colored regions), that is monitored during the simulation. **(B)** Time series of the distance measured between residues L370 and D104 records domain movement across 1000 ns of simulations. Volume of active site pocket (depicted as blue spheres) calculated for the last frame of the simulation for the **(D)** WT^{Gal-}, **(E)** WT^{Gal+}, and **(F)** MC^{Gal-} systems. Cross correlation studies were used to derive residue-residue correlation heatmaps for **(G)** WT^{Gal-}, **(H)** WT^{Gal+}, and **(I)** MC^{Gal-}. Cross-correlation maps showed an intense residue-residue correlation between residues 90–250 and residues 250–350 in WT^{Gal-} that was not observed in the other two systems. Porcupine plots constructed from a principal component analysis (PCA) of the dynamics from the MD trajectories of **(D)** WT^{Gal-}, **(E)** WT^{Gal+} and **(F)** MC^{Gal-} use spikes to show the intensity and direction of the principal component of motion for each residue. WT^{Gal-} showed approximately three-fold and four-fold greater motion relative to WT^{Gal+} and MC^{Gal-}, respectively. Regions showing increased dynamics in the PCA of WT^{Gal-} correspond to the same regions showing defined residue-residue correlation from the cross-correlation studies.

observed in the other two systems (**Figure 3B**). By the end of the simulation (1000 ns), the active site

cavities of the WT^{Gal+} and MC^{Gal-} structures contained pocket volumes of around 1034 Å^3 (**Figure 3D**) and

1056 Å^3 (**Figure 3E**), respectively, the pocket volume of WT^{Gal-} was only 828 Å^3 (**Figure 3C**) due to the

strong interaction network between the interface of the two moving domains. The root mean square

deviations (RMSD) of all three systems were mostly distributed between 2 Å and 3 Å across the 1000 ns

of the MD trajectories, though WT^{Gal-} showed an increase in deviation to about 4 Å before settling into its

compacted state near 320 ns (**Figure S4A**). In contrast, the WT^{Gal+} and MC^{Gal-} systems behaved similarly

throughout. The root mean square fluctuations (RMSF) calculated for the α-carbon (C$_\alpha$) atoms of the

backbone, averaged for each residue obtained throughout the simulation, showed that the fluctuations are

muted for the majority of the protein in both WT^{Gal+} and MC^{Gal-} while in WT^{Gal-} fluctuations are more

apparent across the protein (**Figure S4B**). Overall, these results indicate backbone structural fluctuations

are reduced in both WT^{Gal+} and MC^{Gal-}. Principal component analysis of the same trajectories revealed,

more evidently, that the backbone fluctuations of WT^{Gal-} were greatly increased when compared to WT^{Gal+}

and MC^{Gal-} (**Figure S4C**).

To understand which residues most contributed to the dynamics observed in each of the three

systems during these simulations, we performed cross-correlational studies of the relative movement

between the residues of WT^{Gal-} (**Figure 3F**), WT^{Gal+} (**Figure 3G**), and MC^{Gal-} (**Figure 3H**). The intensity

of correlation and magnitude of the movement observed in the WT^{Gal-} map indicate large-scale movements

in the two 'moving' domains defined by regions containing residues 90–250 and 250–350. The lower

intensity of residue-residue correlation in WT^{Gal+} map indicates relatively stable dynamics that may be a

consequence of the galactose bound in the active site. MC^{Gal-} displayed residue-residue correlations that

were limited to the local secondary structures, and we did not observe large domain movements, which

suggests a stable closed conformation similar to that of WT^{Gal+}. To visualize the domain movements in

detail, we used principal component analysis (PCA) for all three systems across the 1000 ns MD trajectories.

Porcupine plots generated based on the extreme projections of the PCA show the intensity and direction of

the motion of the C_α atoms in each structure. WT^{Gal-} showed highly dynamic conformational states, with

the direction of major motion towards a compact state observable among moving domain residues (**Figure**

3I). While MC^{Gal-} showed conformational transitions in certain dynamic loop regions, in general both

WT^{Gal+} (**Figure 3J**) and MC^{Gal-} showed relatively limited dynamics across the structure (**Figure 3K**).

Initializing the molecular dynamics simulations of the three conditions in the 'closed' conformation

introduces a bias that prevents each system from adopting a range of conformational states. Metadynamics-

based simulations use a potential bias to move a system out of a local energy minimum enabling the

calculation of the free energy surface (FES) across a range of conformations. To perform these

complementary metadynamics analyses, we began to define the conformational landscape to be explored

by identifying centers of mass (COM) for each of the three Gal3p domains (C1–C3). Next, we defined the

two collective variables (CV) to be manipulated during the simulation as (i) the distance between the COM

of the two moving domains (C1 and C2) and (ii) the angle formed by the three COM at the hinge region

(C3) (**Figure 4A**). Examining the resulting FES for each system reveals differences in the CV values that

yield energetically favorable structural conformations. The FES of WT^{Gal-} (**Figure 4B**) showed two

Gaussian wells, one for a short-lived partially open state and one for an exceptionally prominent minimum

corresponding to an irreversibly collapsed closed conformation catalyzed by strong interaction networks

between the two moving domains (**Figure S5A**) not observed between the moving domains in the WT^{Gal+}

(**Figure S5B**) and MC^{Gal-} (**Figure S5C**) systems. Therefore, it can be called a misfolded state (W_m) of the

Figure 4: Metadynamics simulations identify energetically favorable conformations. (A) Two collective variables (CV) were defined to enable energetic exploration of the Gal3p conformational landscape: **CV1** as the distance between C1 and C2, the centers of mass of the two moving domains, and **CV2** as the angle formed between C1, C3 (the hinge region center of mass), and C2. Calculating the energy associated with different values of CV1 and CV2 enabled construction of the free energy surfaces (FES) for the **(B)** WT^{Gal-} (W), **(C)** WT^{Gal+} (G), and **(D)** MC^{Gal-} (M) systems. The subscripts "m", "c", and "o", corresponding to misfolded (collapsed), closed, and open states, respectively, are used to associate prominent energetic minima with distinct conformations of Gal3p variants. Significantly, the most favorable conformations in the WT^{Gal+} and MC^{Gal-} systems correspond to a closed state that likely favors interaction with Gal80p and thus regulon activation.

protein as the recovery of an open conformation would be highly energy-consuming, and therefore,

unlikely. The FES of WT^{Gal+} showed a path of conformational transitions that leads from a closed (G$_c$), but

not collapsed, state to a fully open state (G$_o$) of the protein (**Figure 4C**). Significantly, the closed state of

WT^{Gal+} (G$_c$) is structurally distinct from the closed, collapsed/misfolded state (W$_m$) adopted by WT^{Gal-}

suggesting that the latter is unlikely to interact with Gal80p in a productive manner. As the metadynamics

simulation progresses, WT^{Gal+} eventually adopts an open state (G$_o$) that visualization studies reveal to be a

consequence of substate egress. The FES of MC^{Gal-} (**Figure 4D**) showed a single predominant minimum

corresponding to a closed state (M_c) that closely resembles the closed state (G_c) of WT[Gal+]. These results

suggest that the mutant Gal3p[MC] forms a stable closed conformation closely resembling the closed WT[Gal+]

conformation, with a similar interaction network observed between the moving domains. When combined

with the previously observed robust activation on structurally diverse substrates, the similarity between the

closed state conformations adopted by WT[Gal+] and the MC[Gal−] systems suggests that Gal3p[MC] can bind and

sequester Gal80p in an inducer-independent manner. Recognizing that this substrate-agnostic activation

phenotype of Gal3p[MC] represents a departure from the inducer-dependent regulatory logic of the native

regulon, we decided to test the fitness consequences of this transition when utilizing a regulon approach to

engineer synthetic heterotrophy.

Substrate-agnostic and substrate-specific GAL regulon activation are equivalent.

During our previous effort to adapt the GAL regulon to growth on xylose, regulon activation was

accomplished using the xylose-inducible sensor-protein variant Gal3p[Syn4.1] [18]. To compare the performance

of inducer-dependent (substrate-specific) and inducer-independent (substrate-agnostic) approaches for

engineering synthetic heterotrophy, we transformed a plasmid (pRS426-XYL-REG) encoding the

isomerase pathway genes for xylose utilization (*XYLA*3* [25], *XKS1*) under the control of GAL-responsive

promoters into strains engineered for (i) xylose-inducible regulon activation using Gal3p[Syn4.1] (strain

VDT13), (ii) substrate-agnostic regulon activation using Gal3p[MC] (strain SFS6), and (iii) substrate-agnostic

activation via deletion of *GAL80*, the master repressor of the GAL regulon (strain SFS11), as disruption of

this regulatory gene has been demonstrated to yield constitutive expression of galactose catabolic genes

and enhance galactose utilization [26,27] (**Figure 5A**). While regulon activation in both SFS6 and VDT13

requires an interaction between Gal80p and their respective Gal3p variants (Gal3p[MC] and Gal3p[Syn4.1]),

activation in SFS11 is totally de-regulated due to the absence of Gal80p. In addition, all three strains contain

two accessory genes found to improve pentose utilization – the transporter *GAL2^{2.1}* [28] and the transaldolase

TAL1 – integrated into the chromosome under the control of GAL-responsive promoters.

The growth of these three strains in defined (SC) media containing the native carbon sources

sucrose (**Figure 5B**), glucose (**Figure 5C**), or raffinose (**Figure 5D**) was monitored to identify the potential

fitness defects of each approach during strain handling. While all three approaches reached similar final

Figure 5: Comparing the fitness of substrate-specific and substrate-agnostic approaches to GAL regulon activation. (A) Schematic illustrating different ways of activating the GAL regulon in cells

carrying xylose catabolic genes under the control of GAL-responsive promoters. Gal3p$^{Syn4.1}$ requires induction by xylose, Gal3p^{MC} does not require an inducer but maintains the native regulatory

architecture, while $\Delta GAL80$ effects inducer-independent activation by removing native regulatory elements. Growth of the three systems in defined (SC) medium containing 2 % of **(B)** sucrose, **(C)**

glucose, or **(D)** raffinose reveals that a de-regulated approach ($\Delta GAL80$) reduces fitness on native carbon sources. **(E)** Comparing growth of the three GAL regulon activation systems on the non-native sugar

xylose after being subcultured from sucrose. Comparison of constitutive (CONS) and regulon (REG)

approaches to growth in complex media containing 2 % of **(F)** arabinose or **(G)** cellobiose as sole carbon sources. All data points represent the average of three biological replicates ± sd.

cell concentrations on each of the three substrates, the $\Delta GAL80$ condition took roughly twice the time to do

so compared with Gal3p$^{Syn4.1}$ and Gal3p^{MC} (>30 h vs. ~18 h) suggesting that de-regulation of the regulon

results in reduced fitness relative to strains in which the native regulatory organization is maintained. As

sucrose and glucose are both capable of exerting repression on the GAL regulon [19], subculturing all three

strains from these two native substrates into complex (YP) media containing the non-native substrate xylose

should therefore result in growth-coupled expression of xylose catabolic genes and rapid growth. Upon

subculturing from sucrose, we observed that all three approaches enabled growth. However, the lower

growth rate seen for the $\Delta GAL80$ strain compared with cells carrying $GAL3^{Syn4.1}$ or $GAL3^{MC}$ ($\mu = 0.14 \pm$

0.001 h^{-1}, 0.21 ± 0.008 h^{-1}, and 0.21 ± 0.001 h^{-1}, respectively) combined with the drastically lower final

cell density supports maintenance of native regulatory structure over abolished control (**Figure 5E**).

Subculturing from glucose as an overnight carbon source yielded similar results (**Figure S6**). Significantly,

the equivalent performance of $GAL3^{Syn4.1}$ and $GAL3^{MC}$ backgrounds on both native and non-native carbon

sources indicates that inducer-independent activation enables the metabolic coordination of a regulon

approach to be applied for engineering synthetic heterotrophy in a substrate-agnostic manner provided that

the native regulatory organization is maintained.

To confirm that these results were not xylose-specific, we proceeded to use Gal3p^{MC} to compare

REG and CONS approaches to growth on two additional non-native, lignocellulosic substrates – cellobiose

and arabinose. To do so, we placed catabolic genes previously identified to enable the efficient utilization

of arabinose (*araA*, *araB*, and *araD* from *Lactobacillus plantarum* [29]) and cellobiose (*cdt-1* and *gh1-1* from

Neurospora crassa [30]) under the control of the GAL-responsive promoters *GAL1p*, *GAL10p*, and *GAL7p*

and the strong, constitutive promoters *TEF1p*, *TPI1p*, and *GPM1p* to yield plasmids pRS426-ARA-REG,

pRS426-ARA-CONS, pRS426-CEL-REG, and pRS426-CEL-CONS, respectively. Transforming pRS426-

ARA-REG into $GAL3^{MC}$ integrant strain SFS6 (containing GAL-inducible copies of $GAL2^{2.1}$ and *TAL1*)

yielded strain ARA-REG while transforming pRS426-ARA-CONS into strain VDT27 (that contains

$GAL2^{2.1}$ and *TAL1* under the control of strong constitutive promoters) yielded strain ARA-CONS. Similarly,

transforming pRS426-CEL-REG and pRS426-CEL-CONS into strains SFS3 and VEG16, respectively, that

are comparable to SFS6 and VDT27 except that they lack accessory genes *GAL2²·¹* and *TAL1*, yielded

strains CEL-REG and CEL-CONS. The REG strain exhibited a higher growth rate than the CONS strain

on both defined (0.16 ± 0.005 h^{-1} vs. 0.05 ± 0.005 h^{-1}) (**Figure S7A**) and complex (0.19 ± 0.002 h^{-1} vs.

0.17 ± 0.003 h^{-1}) (**Figure 5F**) media containing 2 % arabinose as a sole carbon source. Similarly, the REG

strain grew faster than its CONS counterpart on both defined (0.14 ± 0.001 h^{-1} vs. 0.09 ± 0.005 h^{-1}) (**Figure**

S7B) and complex (0.25 ± 0.002 h^{-1} vs. 0.17 ± 0.003 h^{-1}) (**Figure 5G**) media containing 2 % cellobiose as

a sole carbon source. Taken together, these data demonstrate that the metabolic coordination effected by

Gal3p^{MC} is broadly beneficial to growth on non-native carbon sources compared with the existing

constitutive paradigm with minimal additional engineering.

Gal3p^{MC} enables rapid and complete co-utilization of multiple non-native substrates

Having demonstrated that a regulon approach is beneficial to growth on all three non-native, lignocellulosic

substrates individually, we sought to utilize the inducer-independent regulon activation of Gal3p^{MC} to test

a REG approach to their co-utilization. As previous studies have observed that the division of catabolic

labor among a consortium of strains can enable faster and/or more efficient co-utilization when compared

with a single strain engineered to utilize all substrates simultaneously [31–33], we decided to test both

approaches. To accommodate all three plasmids into a single strain, the catabolic genes for each substrate

were moved to plasmids containing unique selectable markers to create pRS423-XYL-REG, pRS425-ARA-

REG, and pRS424-CEL-REG. Transforming these plasmids into strains SFS6, SFS6, and SFS3,

respectively, represented a three-strain (Consortium) approach, while transforming all three plasmids into

SFS6 represented a single strain (Consolidated) approach (**Figure 6A**). While both Consortium and

Consolidated approaches exhibited simultaneous co-utilization of all carbon sources and achieved similar

final cell densities during growth on YP media containing 1 % of each substrate, we were surprised to find

that the Consolidated approach resulted in a faster maximum growth rate than the Consortium approach

(0.20 ± 0.003 h^{-1} vs. 0.17 ± 0.003 h^{-1}) (**Figure 6B**). However, neither approach resulted in complete pentose

Figure 6: **Gal3p^{MC} enables rapid and complete co-utilization of multiple non-native substrates.** **(A)** Placing REG plasmids for the catabolism of cellobiose, xylose, and arabinose into separate strains represents a Consortium approach to co-utilization while placing all three plasmids into a single strain represents a Consolidated approach. **(B)** Cell density (black) and percent residual substrate of cellobiose (yellow), xylose (light blue), and arabinose (pink) of Consortium (dotted lines) and Consolidated (solid lines) approaches to co-utilization in complex media containing 1 % of each substrate. Incomplete pentose utilization motivated the exploration of all six possible pairings between GAL-inducible promoters (*GAL1p*, *GAL10p*, and *GAL7p* in order) and the catabolic genes for **(C)** arabinose and **(D)** xylose utilization. Promoter-gene pairings used in subsequent co-utilization experiments are indicated by (*). **(E)** Re-testing a Consolidated approach to co-utilization using expression-balanced pentose plasmids results in complete utilization of all three substrates. Data points represent the average of three (panel **A**) or two (panel **E**) biological replicates ± sd.

consumption even after cellobiose was fully depleted. This led us to hypothesize that there may be flux

imbalances in the pentose utilization pathways.

The success of combinatorial pathway design in optimizing both catabolic [4,34,35] and anabolic

pathways [36,37] motivated us to test the performance of each of the six possible GAL-responsive promoter-

catabolic gene pairings for arabinose (**Figure 6C**) and xylose (**Figure 6D**) in Gal3p^{MC} integrant strain SFS6.

The arabinose constructs all performed similarly, with growth rates clustered in a small range of values

$(0.20 - 0.22 \text{ h}^{-1})$. A similar range of growth rates was observed for the xylose constructs $(0.20 - 0.21 \text{ h}^{-1})$

but the substantially reduced final cell density of two of the arrangements (ϕ-XK-XI and XK-ϕ-XI) indicates

that there can be fitness consequences to expression imbalances. As no clearly superior combination could

be identified to the initial arabinose and xylose designs, we decided to re-test a Consolidated approach to

co-utilization on 1 % of each substrate (**Figure 6E**) using designs (ABD for arabinose, XI-ϕ-XK for xylose)

identified as optimal in a related investigation by our lab into efficient pentose assimilation [19].

Hypothesizing that incomplete pentose utilization may also reflect limitations in the carrying capacity of

the medium itself, we also tested different substrate ratios (cellobiose-xylose-arabinose: 0.7 %-0.7 %-0.7

%, 1 %-0.5 %-0.5 %, and 0.4 %-0.8 %-0.8 %) (**Figure S8A-C**) at a lower (~2 %) total substrate

concentration. Encouragingly, all substrates were utilized to undetectable levels (< 0.05 %, limit of

detection) in approximately 50 h of cultivation for all conditions. For the 3 % total sugar condition, we

observed an 81 % improvement in final cell density when using the optimized pentose plasmids compared

with the non-optimized (OD_{600} = 17.2 vs. 9.5, respectively). For the three 2 % total substrate conditions,

the final cell density was 41 % greater on average using the same comparison (OD_{600} ~13.5 vs. 9.5,

respectively). Despite their similar performance during growth on a single sugar, the higher final OD_{600}

values and complete substrate utilization observed upon switching from the initial to the optimized designs

suggests that the relative expression level of catabolic genes optimal for co-utilization may be different than

for the single substrate. Considering that the highest single-sugar growth rates obtained in this study are

either the best (for cellobiose) or competitive with the best (xylose, arabinose) aerobic growth rates

observed in the literature (**Table S1**) without the need for extensive genome editing or adaptive laboratory

evolution (ALE), we believe that together these results demonstrate the flexibility and effectiveness of

utilizing a regulon approach to engineer efficient synthetic heterotrophy on diverse substrates.

DISCUSSION

The existing paradigm for engineering synthetic heterotrophy consists of constitutive expression of

catabolic genes that channel the carbon source into central carbon metabolism (CCM) and largely ignores

cellular regulation. In contrast, the assimilation of native carbon sources is often controlled by regulatory

structures called regulons that coordinate substrate catabolism with broader metabolism to optimize

biomass formation. Activation of the galactose (GAL) regulon, for instance, results not only in the

expression of the genes necessary for the catabolism of galactose but global metabolic remodeling to

support efficient growth [17,20]. In previous work, we established that adapting the GAL regulon to growth

on the non-native substrate xylose improved growth rate and substrate utilization when compared with a

constitutive approach [18]. This surprising result motivated us to explore the potential of a semi-synthetic

regulon system as a general approach to engineering synthetic heterotrophy in the important industrial

microbe *S. cerevisiae* so that it can be easily and rapidly applied to other attractive potential substrates for

yeast biotechnology.

In this work, we demonstrate that expression from the GAL-responsive promoters that natively

control the expression of galactose catabolic genes (Leloir pathway) is dynamic and mirrors growth phases

whereas expression from constitutive promoters is constant throughout cultivation. Thus, the regulatory

organization of the GAL regulon functions to avoid the burden of unproductive protein synthesis by only

expressing the genes for galactose assimilation only when it is present. The conservation of cellular

resources effected by this inducible phenotype represents a design principle that has also been learned by

metabolic engineers. Early observations of growth inhibition resulting from the overproduction of

heterologous protein [38] led to the use of separate culturing phases for growth and production and strategies

for dynamically switching between them [39]. This concept of dynamic flux control has subsequently been

applied at finer resolution to biosynthetic pathways. Engineered circuits that feature a biosensor enable a

cell to continually monitor some aspect of its internal state or external environment and automatically adjust

their metabolism in response [40], resulting in improvements in titer, rate, and yield (TRY) [41–44]. While the

use of sugar-responsive biosensors has been proposed to engineer cells to respond to the carbon sources

present in a given lignocellulosic hydrolysate for consolidated bioprocessing, the suggested

implementations decouple sensing from the broader regulatory and metabolic context in which the sensors

evolved [45] . To our knowledge, our semi-synthetic regulon approach is the first time that this dynamic

activation concept has been applied to engineering catabolism in a way that interfaces with native

metabolism for the purpose of efficient biomass formation. This coordination is significant as the regulation

of CCM has evolved to ensure that resource uptake and utilization are balanced [3], with the transcription

level of a large fraction of the yeast genome (~15 – 27 %) [46,47] known to be correlated with growth rate.

We, therefore, posit that regulon control of catabolic gene expression, where the amount of carbon fed into

CCM is innately connected to the utilization of those resources to match cellular needs, is likely beneficial

to fast growth regardless of the carbon source being assimilated.

While we were previously able to engineer a sensor protein variant (Gal3p$^{\text{Syn4.1}}$) capable of strong

regulon activation upon induction with xylose, that effort involved an extensive directed evolution

campaign that would have to be repeated for each new substrate [18]. Here, we report the construction of a

sensor protein variant, Gal3p$^{\text{MC}}$ (<u>m</u>etabolic <u>c</u>oordinator), which robustly activates the GAL regulon on

structurally diverse, non-native carbon sources to a similar degree as $Gal3p^{WT}$ upon induction with

galactose. Using metadynamics simulations, we show that the most energetically favorable conformation

for $Gal3p^{MC}$ without galactose (MC^{Gal-}) is similar to the closed conformational state of $Gal3p^{WT}$ in complex

with galactose (WT^{Gal+}). In contrast, the free energy surface of $Gal3p^{WT}$ in the absence of galactose (WT^{Gal-}

) is characterized by a deep energy well reflective of the active site collapsing/misfolding, a conformation

that is unproductive for signal transduction. Molecular dynamics (MD) simulations of these three systems

reveal large-scale domain movements in the WT^{Gal-} system throughout the course of the simulation whereas

the WT^{Gal+} and MC^{Gal-} systems exhibit much reduced and more localized movement indicating that the

mutations carried by $Gal3p^{MC}$ stabilize the protein in a closed state that enables interaction with Gal80p

without requiring an inducer. The inducer-independent mechanism of $Gal3p^{MC}$ would normally ablate the

dynamic expression profile exhibited by the native, inducer-dependent regulon as it would constantly be

turned on. However, we also observed repression of the $Gal3p^{MC}$-controlled GAL regulon on repressing

substrates (glucose, sucrose), which allows for dynamic activation. $Gal3p^{MC}$ thus enables both the dynamic

catabolic gene expression and global metabolic remodeling of the semi-synthetic regulon approach to be

applied in a substrate-agnostic manner.

Although the inducer-independent activation effected by $Gal3p^{MC}$ represents a departure from the

inducer-dependent regulatory logic of the native GAL regulon that served as our inspiration, we,

surprisingly, did not observe any major growth defects on either native or non-native substrates compared

with a substrate-specific activation approach using $Gal3p^{Syn4.1}$. In contrast, a $\Delta GAL80$ strain, which

represents an alternative approach of achieving substrate-agnostic regulon activation, showed reduced

fitness in both conditions, emphasizing the importance of maintaining the native regulatory organization

when utilizing a regulon approach. During growth on the native carbon sources glucose, sucrose, and

raffinose, the lower fitness likely results from the burden of constitutive GAL regulon activation.

As we previously observed with xylose [18], using $Gal3p^{MC}$ to implement a regulon approach for

growth on arabinose and cellobiose resulted in faster growth rates and higher final cell densities compared

to the traditional constitutive overexpression approach despite expressing the same set of catabolic genes.

Significantly, the growth rates achieved using this rational approach are comparable or superior to the

highest aerobic growth rates for these substrates identified in the literature (**Table S1**) without the need for

time-consuming and/or expensive experiments like ALE (adaptive laboratory evolution), functional

genomics, and reverse genetics, or even deletion of genes previously found to mitigate stress of growth on

pentoses (e.g., *PHO13, ALD6, ASK10, ADH6, COX4, CYC8*, etc. [48]). The success of a semi-synthetic

regulon approach in all the single-sugar conditions motivated us to explore their co-utilization as the

simultaneous and complete co-utilization of the constituent sugars in lignocellulosic hydrolysates has been

identified as essential to their valorization [49]. One of the major challenges of engineering the co-utilization

of lignocellulosic sugars is that the presence of glucose exerts carbon catabolite repression (CCR) in many

organisms, hindering their ability to metabolize other sugars until glucose is depleted and repression is

relieved [50,51]. However, Jin & Cate et al. found that CCR could be avoided by implementing a heterologous

pathway for cellobiose, a glucose disaccharide that does not trigger CCR [30,52]. We demonstrate that using

Gal3p^{MC} to implement a regulon approach enabled rapid and complete co-utilization of three non-native

carbon sources – xylose, arabinose, and cellobiose – in a single strain in only ~48 h. Despite being evolved

for the catabolism of galactose, these results reinforce the universality and robustness of the metabolic

coordination effected by the GAL regulon for engineering synthetic heterotrophy as well as the plasticity

of yeast metabolism to diverse substrates. However, our findings do not discount the importance of existing

metabolic engineering tools and strategies. For example, complete utilization of pentoses during co-

utilization required balancing the expression levels of pathway genes. Additionally, while Gal3p^{MC} enables

a regulon approach to be rapidly adapted to structurally diverse, non-native carbon sources, it is dependent

on the development and importation of an efficient catabolic module. This requirement limits its immediate

application to attractive renewable feedstocks like C1 compounds (i.e., CO_2 [53], formate, methanol [54],

methane [55]) as efforts to engineer their assimilation in *S. cerevisiae* remain nascent. Researchers have noted

the potential of platform strains that maximize the concentration of a key precursor to speed the construction

and optimization of biosynthetic pathways [56]. We envision that the integration of substrate catabolism with

broader metabolism characteristic of a regulon approach could similarly accelerate the process of

engineering catabolism by reducing the number of iterative, semi-rational genetic interventions currently

needed to engineer efficient growth on a given non-native substrate. Together, these results establish the

metabolic coordination of a regulon approach as a fast, flexible, and effective strategy for engineering

synthetic heterotrophy and provide a tool (Gal3p^{MC}) for adapting it to growth on structurally diverse non-

native substrates.

MATERIALS AND METHODS

Strains and Plasmids

A full list of strains and plasmids used in this study can be found in **Table S2** and **Table S3**, respectively.

Media and Transformation

Yeast strains were grown at 30 °C in either defined Synthetic Complete (SC) (Yeast nitrogen base (1.67

g/L), ammonium sulfate (5 g/L), complete supplement mixture without His, Leu, Ura and Trp (0.6 g/L)

(Sunrise Science Products, Inc.) or complex Yeast Peptone (YP) medium (20 g/L Yeast Extract, 40 g/L

Casein Peptone, 200 mg adenine) medium. SC media was augmented with appropriate nutrients to select

for maintenance of plasmids carrying a given auxotrophy. Luria Bertani (LB) broth and LB agar plates with

100 μg/mL of ampicillin when required were used for all *E. coli* propagation and transformation

experiments. Commercial *E. coli* strain NEB-5α (New England Biolabs) was used for MES transformation

when constructing plasmids as well as for plasmid propagation. Upon isolation, plasmids were sequenced

before being transformed into yeast strains using the protocol of Gietz [57] and recovered upon agar plates

with the appropriate auxotrophy.

Strain Construction

The yeast strain W303-1a was used for constructing all the strains used in the study. The generation of

knockout strain VEG16 from W303-1a was described previously [18]. All subsequent strains were derived

from VEG16 via chromosomal integrations using a previously described set of disintegrator plasmids and

associated protocols [58].

Plasmid Construction

S. cerevisiae promoter, gene, and terminator sequences were amplified from the genome using Phusion

Polymerase (ThermoFisher Scientific). *XYLA*3* DNA sequence was provided by Prof. Hal S. Alper

(University of Texas at Austin). *GAL2^{2.1}* DNA sequence was provided by Prof. Bernard Hauer (University

of Stuttgart, Stuttgart, Germany). Cellobiose utilization genes *cdt-1* and *gh1-1* were provided by Prof.

Huimin Zhao (University of Illinois at Urbana-Champaign). Synthesis of DNA primers and Sanger

sequencing of plasmid DNA was outsourced to Genewiz. Substrate utilization plasmids were created by

combining promoter, catabolic gene, and terminator sequences into previously described yeast shuttle

vectors containing either *CEN6* [59] or *2μ* [60] origins. All plasmids were constructed using either NEBuilder

HiFi DNA Assembly or restriction enzyme digest and ligation using reagents purchased from New England

Biolabs.

Quantifying Promoter Dynamics via Fluorescence

473   Overnight inoculums were grown in 200 µL in 96-well plates containing the required dropout SC medium

474   with glucose (2 %) for 24 hours. Cultures were then subcultured (5 µL) into 200 µL SC medium containing

475   the appropriate carbon source (2 %) and returned to the incubator. Fluorescence (excitation at 488 nm and

476   emission at 525 nm) and OD_{600} were measured at numerous time points in a Spectramax M3

477   spectrophotometer to obtain Fluorescence/OD_{600}.

Molecular dynamics simulations

Extended molecular dynamics and metadynamics simulation were conducted on three systems: $Gal3p^{WT}$

complexed with ATP and Mg^{2+} (WT^{Gal-}), $Gal3p^{WT}$ complexed with galactose, ATP, and Mg^{2+} (WT^{Gal+}),

and $Gal3p^{MC}$ complexed with ATP & Mg^{2+} (MC^{Gal-}). To establish a common simulation starting point for

the three systems, each of the protein structures was modeled into the closed state via homology modeling

using Swiss Model [61] based on the experimentally determined closed state structure (PDB ID: 3V2U) as a

template (**Figure S9**). Side chain conformations of the template were retained when generating models.

Molecular dynamic simulation of the three different experimental setups was performed for 1000 ns using

GROMACS–2019.4 [62] . The molecules were parameterized using antechamber package from AMBER [63] .

Quantum polarized charges were derived using the GAMESS package [64] with 6-31G basis set and B3LYP

level of theory AMBER99SB [63] force field was used to perform a simulation with the TIP3p water model.

To solvate the complex, the TIP3P water model with a triclinic box volume of 1000 nm^3 was used to mimic

or replace the natural environment around the protein. The system was neutralized by adding Na^+ ions to

the system, 16574 water molecules and ions overall. Energy minimization of the system was performed

using the steepest descent algorithm with convergence energy cut-off of 1000 kJ mol^{-1} with 1000 steps.

Equilibration of the system was performed using NVT and NPT, where NVT is for constant volume and

temperature and NPT is for constant pressure. The NVT and NPT were carried out for 500 PS with 300 K

temperature using a Berendsen thermostat and the system was prepared for Molecular dynamic simulation

for 1 μs at 300 K temperature. The system was made to run for 1000 ns, for every 100 ns the trajectory was

saved for analysis. The first 100 ns were excluded for every analysis. The distance between C_α atoms of

L370 and D104 has been calculated across the simulation 1000 ns long trajectory for all three proteins to

know the variation in the movement of two domain regions across the time. RMSD (Root mean square

Deviation) and RMSF (Root mean square fluctuation) of the backbone atoms were calculated over time.

Using WORDOM tools [65] , cross-correlation studies of residue-residue displacements along the trajectory

were calculated for each system to quantify dynamic changes of across the moving domains. Similarly,

principal component analysis (PCA) was calculated for each 100-1000 ns trajectory and porcupine plots

were generated to analyze the intensity and movement of C_α atom. Visualization of structures was done

using UCSF Chimera [66] and Visual Molecular Dynamics (VMD) [67] .

Metadynamics simulations

Metadynamics-based simulations use a potential bias to move the system out of a local minimum, which is

then used to explore the resulting free energy landscape (FES). Metadynamics-based simulations were

conducted using the PLUMED software package [68] and GROMACS toolkit for all three experimental setups

(WT^{Gal-}, WT^{Gal+}, and MC^{Gal-}) after thorough equilibration obtained from NVT, NPT ensembles. C1, C2,

and C3 are Centers of Mass (COMs) determined for the three regions of Gal3p that fall within the residues

regions as shown in **Figure 3A**. One of the Collective Variables (CV) for metadynamics simulation was

defined as the distance between C1 and C2. The other CV was defined as the angle between the intersection

point of the vectors that pass through C1, C3 and C2, C3, respectively. Different simulations in this study

were performed for 50 ns with a bias factor set to 8.0, the temperature set to 300 K, and the results were

calculated accordingly. The free energy landscape was explored as the distance between the two domains

and the change in the angle formed by the three COMs were varied.

Growth Studies

Overnight inoculums were grown in the required dropout SC medium with 2 % sucrose (unless explicitly

stated to be glucose/raffinose) for 24 h. The culture was washed twice in the growth medium (either SC or

YP containing 2 % of the target sugar) and resuspended at an initial OD$_{600}$ of 0.1 in 250 mL shake flasks

containing 20 mL of media. Cell growth OD$_{600}$ measurement was checked at frequent time intervals (3–6

h) on SpectraMax M3 spectrophotometer (Molecular Devices). Growth rates (μ) for each biological

replicate were quantified in Microsoft Excel using an exponential curve equation fit to at least three time

points during logarithmic growth phase. These values were used to calculate the average growth rate and

the standard error for a given condition.

Residual Substrate Quantification

Residual substrate during co-utilization studies was measured using an Agilent HPLC system equipped

with a Hi-Plex H-column and detected using 1260 Agilent ELSD detector. Mobile phase was 0.1 %

trifluoroacetic acid (TFA) with a flow rate of 0.6 mL/min. The ELSD detector's nebulizer and evaporation temperature were set at 30 °C and nitrogen flow rate at 1.6 SLM (standard liter per minute). Using these settings, we observed characteristic retention times for cellobiose (9.3 min), arabinose (12.6 min), and xylose (11.8 min).

DATA AVAILABILITY

All plasmids and strains used in this study can be obtained from the corresponding author under a material transfer agreement. Source data are provided with this paper.

ACKNOWLEDGEMENTS

The authors would like to thank current and former Nair lab members and Dr. Todd C. Chappell for helpful discussions. This work was supported by NIH grant #DP2HD91798 and NSF grant #1935354, and Tufts Launchpad | Accelerator to N.U.N.

AUTHOR CONTRIBUTIONS

N.U.N., S.F.S., V.D.T., and V.E.G. conceived and designed the research project. S.F.S., A.S., and N.U.N. co-wrote the manuscript. S.F.S., A.S., V.D.T., D.S. and T.B. performed the experiments. S.F.S., A.S., T.B., P.K.R., and N.U.N. analyzed the data. All the authors have reviewed the manuscript and approved it for submission.

COMPETING INTERESTS

None.

REFERENCES

1. Langholtz M, Stokes B, Eaton L. 2016 billion-ton report: Advancing domestic resources for a thriving bioeconomy (Executive Summary). *Industrial Biotechnology*. 2016;12(5):282-289. doi:10.1089/ind.2016.29051.doe

2. Rogers JN, Stokes B, Dunn J, et al. An assessment of the potential products and economic and environmental impacts resulting from a billion ton bioeconomy. *Biofuels, Bioproducts and Biorefining*. 2017;11(1):110-128. doi:10.1002/bbb.1728

3. Nielsen J, Keasling JD. Engineering Cellular Metabolism. *Cell*. 2016;164(6):1185-1197. doi:10.1016/j.cell.2016.02.004

4. Latimer LN, Lee ME, Medina-Cleghorn D, Kohnz RA, Nomura DK, Dueber JE. Employing a combinatorial expression approach to characterize xylose utilization in Saccharomyces cerevisiae. *Metab Eng*. 2014;25:20-29. doi:10.1016/j.ymben.2014.06.002

5. Kobayashi Y, Sahara T, Ohgiya S, Kamagata Y, Fujimori KE. Systematic optimization of gene expression of pentose phosphate pathway enhances ethanol production from a glucose/xylose mixed medium in a recombinant Saccharomyces cerevisiae. *AMB Express*. 2018;8(1):1-11. doi:10.1186/s13568-018-0670-8

6. Kim SR, Skerker JM, Kang W, et al. Rational and evolutionary engineering approaches uncover a small set of genetic changes efficient for rapid xylose fermentation in Saccharomyces cerevisiae. *PLoS One*. 2013;8(2):e57048. doi:10.1371/journal.pone.0057048

7. Ni H, Laplaza JM, Jeffries TW. Transposon mutagenesis to improve the growth of recombinant Saccharomyces cerevisiae on D-xylose. *Appl Environ Microbiol*. 2007;73(7):2061-2066. doi:10.1128/AEM.02564-06

8. Mukherjee V, Lind U, St. Onge RP, Blomberg A, Nygård Y. A CRISPR Interference Screen of Essential Genes Reveals that Proteasome Regulation Dictates Acetic Acid Tolerance in Saccharomyces cerevisiae. Vickers C, ed. *mSystems*. 2021;6(4). doi:10.1128/mSystems.00418-21

9. Unrean P, Gätgens J, Klein B, Noack S, Champreda V. Elucidating cellular mechanisms of Saccharomyces cerevisiae tolerant to combined lignocellulosic-derived inhibitors using high-Throughput phenotyping and multiomics analyses. *FEMS Yeast Res*. 2018;18(8):1-10. doi:10.1093/femsyr/foy106

10. HamediRad M, Lian J, Li H, Zhao H. RNAi Assisted Genome Evolution Unveils Yeast Mutants with Improved Xylose Utilization. *Biotechnol Bioeng*. Published online 2018. doi:10.1002/bit.26570

11. Chen Y, Stabryla L, Wei N. Improved acetic acid resistance in Saccharomyces cerevisiae by overexpression of the WHI2 gene identified through inverse metabolic engineering. *Appl Environ Microbiol*. 2016;82(7):2156-2166. doi:10.1128/AEM.03718-15

12. Ha SJ, Galazka JM, Kim SR, et al. Engineered Saccharomyces cerevisiae capable of simultaneous cellobiose and xylose fermentation. *Proc Natl Acad Sci U S A*. 2011;108(2):504-509. doi:10.1073/pnas.1010456108

13.   Zhou H, Cheng JS, Wang B, Fink GR, Stephanopoulos G. Xylose isomerase overexpression along
with engineering of the pentose phosphate pathway and evolutionary engineering enable rapid
xylose utilization and ethanol production by Saccharomyces cerevisiae. *Metab Eng*.
2012;14(6):611-622. doi:10.1016/j.ymben.2012.07.011

14.   Sanchez RG, Karhurmaa K, Fonseca C, et al. Improved xylose and arabinose utilization by an
industrial recombinant. *Biotechnol Biofuels*. 2010;3(13).

15.   Lavy T, Kumar PR, He H, Joshua-Tor L. The Gal3p transducer of the GAL regulon interacts with the
Gal80p repressor in its ligand-induced closed conformation. *Genes Dev*. 2012;26(3):294-303.
doi:10.1101/gad.182691.111

16.   Griffin TJ, Gygi SP, Ideker T, et al. Complementary profiling of gene expression at the
transcriptome and proteome levels in Saccharomyces cerevisiae. *Mol Cell Proteomics*.
2002;1(4):323-333. doi:10.1074/mcp.M200001-MCP200

17.   Ren B, Robert F, Wyrick JJ, et al. Genome-wide location and function of DNA binding proteins.
Science (1979). 2000;290(5500):2306-2309. doi:10.1126/science.290.5500.2306

18.   Endalur Gopinarayanan V, Nair NU. A semi-synthetic regulon enables rapid growth of yeast on
xylose. *Nat Commun*. 2018;9(1). doi:10.1038/s41467-018-03645-7

19.   Trivedi VD, Sullivan SF, Choudhury D, Endalur Gopinarayanan V, Nair NU. Integration of
metabolism and regulation reveals rapid adaptability to growth on non-native substrates.
bioRxiv. Published online 2022. doi:10.1101/2022.05.21.492926

20.   Reimand J, Vaquerizas JM, Todd AE, Vilo J, Luscombe NM. Comprehensive reanalysis of
transcription factor knockout expression data in Saccharomyces cerevisiae reveals many new
targets. *Nucleic Acids Res*. 2010;38(14):4768-4777. doi:10.1093/nar/gkq232

21.   Blank TE, Woods MP, Lebo CM, Xin P, Hopper JE. Novel Gal3 proteins showing altered Gal80p
binding cause constitutive transcription of Gal4p-activated genes in Saccharomyces cerevisiae.
Mol Cell Biol. 1997;17(5):2566-2575. doi:10.1128/mcb.17.5.2566

22.   Bhat PJ, Hopper JE. Overproduction of the GAL1 or GAL3 protein causes galactose-independent
activation of the GAL4 protein: evidence for a new model of induction for the yeast GAL/MEL
regulon. *Mol Cell Biol*. 1992;12(6):2701-2707. doi:10.1128/mcb.12.6.2701

23.   Malakar P, Venkatesh K V. GAL regulon of Saccharomyces cerevisiae performs optimally to
maximize growth on galactose. *FEMS Yeast Res*. 2014;14(2):346-356. doi:10.1111/1567-
1364.12109

24.   Venturelli OS, El-Samad H, Murray RM. Synergistic dual positive feedback loops established by
molecular sequestration generate robust bimodal response. *Proc Natl Acad Sci U S A*.
2012;109(48). doi:10.1073/pnas.1211902109

25.   Lee SM, Jellison T, Alper HS. Directed evolution of xylose isomerase for improved xylose
catabolism and fermentation in the yeast Saccharomyces cerevisiae. *Appl Environ Microbiol*.
2012;78(16):5708-5716. doi:10.1128/AEM.01419-12

26.   Quarterman J, Skerker JM, Feng X, et al. Rapid and efficient galactose fermentation by
629   engineered Saccharomyces cerevisiae. *J Biotechnol*. 2016;229:13-21.
630   doi:10.1016/j.jbiotec.2016.04.041

631   27.   Torchia TE, Hamilton RW, Cano CL, Hopper JE. Disruption of regulatory gene GAL80 in
632   Saccharomyces cerevisiae: effects on carbon-controlled regulation of the galactose/melibiose
633   pathway genes. *Mol Cell Biol*. 1984;4(8):1521-1527. doi:10.1128/mcb.4.8.1521

634   28.   Reznicek O, Facey SJ, de Waal PP, et al. Improved xylose uptake in Saccharomyces cerevisiae due
635   to directed evolution of galactose permease Gal2 for sugar co-consumption. *J Appl Microbiol*.
636   2015;119(1):99-111. doi:10.1111/jam.12825

637   29.   Wisselink HW, Toirkens MJ, del Rosario Franco Berriel M, et al. Engineering of Saccharomyces
638   cerevisiae for efficient anaerobic alcoholic fermentation of L-arabinose. *Appl Environ Microbiol*.
639   2007;73(15):4881-4891. doi:10.1128/AEM.00177-07

640   30.   Galazka JM, Tian C, Beeson WT, Martinez B, Glass NL, Cate JHD. Cellodextrin Transport in Yeast
641   for Improved Biofuel Production. *Science (1979)*. 2010;330(6000):84-86.
642   doi:10.1126/science.1192838

643   31.   Eiteman MA, Lee SA, Altman E. A co-fermentation strategy to consume sugar mixtures
644   effectively. *J Biol Eng*. 2008;2:1-8. doi:10.1186/1754-1611-2-3

645   32.   Verhoeven MD, de Valk SC, Daran JMG, van Maris AJA, Pronk JT. Fermentation of glucose-xylose-
646   Arabinose mixtures by a synthetic consortium of single-sugar-fermenting Saccharomyces
647   cerevisiae strains. *FEMS Yeast Res*. 2018;18(8):1-12. doi:10.1093/femsyr/foy075

648   33.   Flores AD, Choi HG, Martinez R, et al. Catabolic Division of Labor Enhances Production of D-
649   Lactate and Succinate From Glucose-Xylose Mixtures in Engineered Escherichia coli Co-culture
650   Systems. *Front Bioeng Biotechnol*. 2020;8(May):1-14. doi:10.3389/fbioe.2020.00329

651   34.   Kim B, Du J, Eriksen DT, Zhao H. Combinatorial design of a highly efficient xylose-utilizing pathway
652   in Saccharomyces cerevisiae for the production of cellulosic biofuels. *Appl Environ Microbiol*.
653   2013;79(3):931-941. doi:10.1128/AEM.02736-12

654   35.   Protzko RJ, Latimer LN, Martinho Z, et al. Engineering Saccharomyces cerevisiae for co-utilization
655   of d-galacturonic acid and d-glucose from citrus peel waste. *Nat Commun*. 2018;9(1):5059.
656   doi:10.1038/s41467-018-07589-w

657   36.   Ajikumar PK, Xiao WH, Tyo KEJ, et al. Isoprenoid pathway optimization for Taxol precursor
658   overproduction in Escherichia coli. *Science (1979)*. 2010;330(6000):70-74.
659   doi:10.1126/science.1191652

660   37.   Smanski MJ, Bhatia S, Zhao D, et al. Functional optimization of gene clusters by combinatorial
661   design and assembly. *Nat Biotechnol*. 2014;32(12):1241-1249. doi:10.1038/nbt.3063

662   38.   Kurland CG, Dong H. Bacterial growth inhibition by overproduction of protein. *Mol Microbiol*.
663   1996;21(1):1-4. doi:10.1046/J.1365-2958.1996.5901313.X

39.   Burg JM, Cooper CB, Ye Z, Reed BR, Moreb EA, Lynch MD. Large-scale bioprocess
competitiveness: the potential of dynamic metabolic control in two-stage fermentations. *Curr*
Opin Chem Eng. 2016;14:121-136. doi:10.1016/J.COCHE.2016.09.008

40.   Hartline CJ, Schmitz AC, Han Y, Zhang F. Dynamic control in metabolic engineering: Theories,
tools, and applications. *Metab Eng*. 2021;63:126-140. doi:10.1016/J.YMBEN.2020.08.015

41.   Zhang F, Carothers JM, Keasling JD. Design of a dynamic sensor-regulator system for production
of chemicals and fuels derived from fatty acids. *Nature Biotechnology 2012 30:4*. 2012;30(4):354-
359. doi:10.1038/nbt.2149

42.   Xiao Y, Bowen CH, Liu D, Zhang F. Exploiting nongenetic cell-to-cell variation for enhanced
biosynthesis. *Nature Chemical Biology 2016 12:5*. 2016;12(5):339-344.
doi:10.1038/nchembio.2046

43.   Xu P, Li L, Zhang F, Stephanopoulos G, Koffas M. Improving fatty acids production by engineering
dynamic pathway regulation and metabolic control. *Proc Natl Acad Sci U S A*.
2014;111(31):11299-11304.
doi:10.1073/PNAS.1406401111/SUPPL_FILE/PNAS.1406401111.SAPP.PDF

44.   Dahl RH, Zhang F, Alonso-Gutierrez J, et al. Engineering dynamic pathway regulation using stress-
response promoters. *Nature Biotechnology 2013 31:11*. 2013;31(11):1039-1046.
doi:10.1038/nbt.2689

45.   Alvarez-Gonzalez G, Dixon N. Genetically encoded biosensors for lignocellulose valorization.
Biotechnology for Biofuels 2019 12:1. 2019;12(1):1-14. doi:10.1186/S13068-019-1585-6

46.   Castrillo JI, Zeef LA, Hoyle DC, et al. Growth control of the eukaryote cell: A systems biology study
in yeast. *J Biol*. 2007;6(2). doi:10.1186/jbiol54

47.   Brauer MJ, Huttenhower C, Airoldi EM, et al. Coordination of growth rate, cell cycle, stress
response, and metabolic activity in yeast. *Mol Biol Cell*. 2008;19(1):352-367.
doi:10.1091/mbc.e07-08-0779

48.   Li X, Wang Y, Li G, et al. Metabolic network remodelling enhances yeast's fitness on xylose using
aerobic glycolysis. *Nat Catal*. 2021;4(September). doi:10.1038/s41929-021-00670-6

49.   Gao M, Ploessl D, Shao Z. Enhancing the Co-utilization of Biomass-Derived Mixed Sugars by
Yeasts. *Front Microbiol*. 2019;9(January):1-21. doi:10.3389/fmicb.2018.03264

50.   Boles E, Hollenberg CP. The molecular genetics of hexose transport in yeasts. *FEMS Microbiol*
Rev. 1997;21(1):85-111. doi:10.1111/J.1574-6976.1997.TB00346.X

51.   Roca C, Haack MB, Olsson L. Engineering of carbon catabolite repression in recombinant xylose
fermenting Saccharomyces cerevisiae. *Appl Microbiol Biotechnol*. 2004;63(5):578-583.
doi:10.1007/S00253-003-1408-2/TABLES/3

52.   Li S, Du J, Sun J, et al. Overcoming glucose repression in mixed sugar fermentation by co-
expressing a cellobiose transporter and a β-glucosidase in Saccharomyces cerevisiae. *Mol Biosyst*.
2010;6(11):2129-2132. doi:10.1039/C0MB00063A

53. Liu Z, Wang K, Chen Y, Tan T, Nielsen J. Third-generation biorefineries as the means to produce
fuels and chemicals from CO2. *Nat Catal*. 2020;3(3):274-288. doi:10.1038/s41929-019-0421-5

54. Antoniewicz MR. Synthetic methylotrophy: Strategies to assimilate methanol for growth and
chemicals production. *Curr Opin Biotechnol*. 2019;59:165-174. doi:10.1016/j.copbio.2019.07.001

55. Clomburg JM, Crumbley AM, Gonzalez R. Industrial biomanufacturing: The future of chemical
production. *Science (1979)*. 2017;355(6320). doi:10.1126/science.aag0804

56. Nielsen J. Yeast cell factories on the horizon: Metabolic engineering in yeast gets increasingly
more versatile. *Science (1979)*. 2015;349(6252):1050-1051. doi:10.1126/SCIENCE.AAD2081

57. Gietz RD. Yeast Transformation by the LiAc/SS Carrier DNA/PEG Method. In: *Yeast Protocols*. Vol
1163. ; 2014:33-44. doi:10.1007/978-1-4939-0799-1_4

58. Sadowski I, Su TC, Parent J. Disintegrator vectors for single-copy yeast chromosomal integration.
Yeast. 2007;24(5):447-455. doi:10.1002/yea.1469

59. Frazer LN, O'Keefe RT. A new series of yeast shuttle vectors for the recovery and identification of
multiple plasmids from Saccharomyces cerevisiae. *Yeast*. 2007;24(9):777-789.
doi:10.1002/yea.1509

60. Christianson TW, Sikorski RS, Dante M, Shero JH, Heiter P. Multifunctional yeast high-copy shuttle
vectors. *Gene*. 1992;110:119-122.

61. Waterhouse A, Bertoni M, Bienert S, et al. SWISS-MODEL: homology modelling of protein
structures and complexes. *Nucleic Acids Res*. 2018;46(W1):W296-W303.
doi:10.1093/NAR/GKY427

62. Abraham MJ, Murtola T, Schulz R, et al. GROMACS: High performance molecular simulations
through multi-level parallelism from laptops to supercomputers. *SoftwareX*. 2015;1-2:19-25.
doi:10.1016/J.SOFTX.2015.06.001

63. Lindorff-Larsen K, Piana S, Palmo K, et al. Improved side-chain torsion potentials for the Amber
ff99SB protein force field. *Proteins: Structure, Function, and Bioinformatics*. 2010;78(8):1950-
1958. doi:10.1002/PROT.22711

64. Guest MF, Bush IJ, J Van Dam HJ, et al. The GAMESS-UK electronic structure package: algorithms,
developments and applications. *https://doi.org/101080/00268970512331340592*. 2007;103(6-
8):719-747. doi:10.1080/00268970512331340592

65. Seeber M, Felline A, Raimondi F, et al. Wordom: A user-friendly program for the analysis of
molecular structures, trajectories, and free energy surfaces. *J Comput Chem*. 2011;32(6):1183-
1194. doi:10.1002/JCC.21688

66. Pettersen EF, Goddard TD, Huang CC, et al. UCSF Chimera—A visualization system for exploratory
research and analysis. *J Comput Chem*. 2004;25(13):1605-1612. doi:10.1002/JCC.20084

67. Humphrey W, Dalke A, Schulten K. VMD: Visual molecular dynamics. *J Mol Graph*. 1996;14(1):33-
38. doi:10.1016/0263-7855(96)00018-5

68.   Abrams JB, Tuckerman ME. Efficient and direct generation of multidimensional free energy
surfaces via adiabatic dynamics without coordinate transformations. *Journal of Physical*
Chemistry B. 2008;112(49):15742-15757. doi:10.1021/JP805039U/ASSET/IMAGES/LARGE/JP-
2008-05039U_0015.JPEG

741