Infant low-frequency EEG cortical power, cortical tracking and phase-amplitude coupling predicts language a year later

Adam Attaheri[1]*, Áine Ní Choisdealbha[1], Sinead Rocha[1,2,3], Perrine Brusini[1,4], Giovanni M. Di Liberto[5,6], Natasha Mead[1], Helen Olawole-Scott[1], Panagiotis Boutris[1], Samuel Gibbon[1], Isabel Williams[1], Christina Grey[1], Maria Alfaro e Oliveira[1], Carmel Brough[1], Sheila Flanagan[1] and Usha Goswami[1].

Department of Psychology, Centre for Neuroscience in Education, University of Cambridge, Cambridge, United Kingdom

Psychology and Sports Science, Anglia Ruskin University, Cambridge, United Kingdom

Department of Psychology, Goldsmiths, University of London, London, UK

Institute of Population Health, University of Liverpool, Liverpool, United Kingdom

School of Computer Science and Statistics, Trinity College Dublin, Dublin, Ireland

Laboratoire des Systèmes Perceptifs, UMR 8248, CNRS, Ecole Normale Supérieure, PSL Research University, Paris, France

* Corresponding author, E-mail: aa974@cam.ac.uk

Abstract

Cortical signals have been shown to track acoustic and linguistic properties of continuous speech. This phenomenon has been measured in both children and adults, reflecting speech understanding by adults as well as cognitive functions such as attention and prediction. Furthermore, atypical low-frequency cortical tracking of speech is found in children with phonological difficulties (developmental dyslexia). Accordingly, low-frequency cortical signals may play a critical role in language acquisition. A recent investigation with infants Attaheri et al., 2022 (1) probed cortical tracking mechanisms at the ages of 4, 7 and 11 months as participants listened to sung speech. Results from temporal response function (TRF), phase-amplitude coupling (PAC) and dynamic theta-delta power (PSD) analyses indicated speech

envelope tracking and stimulus-related power (PSD) for delta and theta neural signals.

Furthermore, delta- and theta-driven PAC was found at all ages, with theta phases displaying

stronger PAC with high-frequency amplitudes than delta. The present study tests whether these

previous findings replicate in the second half of the full cohort of infants (N = 122) who were

participating in this longitudinal study (first half: N=61, (1); second half: N=61). In addition to

demonstrating good replication, we investigate whether cortical tracking in the first year of life

predicts later language acquisition for the full cohort (122 infants recruited, 113 retained) using

both infant-led and parent-estimated measures and multivariate and univariate analyses.

Increased delta cortical tracking in the univariate analyses, increased ~2Hz PSD power and

stronger theta-gamma PAC in both multivariate and univariate analyses were related to better

language outcomes using both infant-led and parent-estimated measures. By contrast, increased

~4Hz PSD power in the multi-variate analyses, increased delta-beta PAC and a higher theta/delta

power ratio in the multi-variate analyses were related to worse language outcomes. The data are

interpreted within a "Temporal Sampling" framework for developmental language trajectories.

40   1 Introduction

Infant speech perception develops rapidly during the first year of life, yet our

understanding of the neural factors underpinning language acquisition is still incomplete. Whilst

infant behavioral experiments have highlighted perceptual factors contributing to early

phonological and morphological development, for example perceptual 'magnet' effects whereby

native language phonetic category prototypes pull neighbouring speech sounds towards them (2),

the neural mechanisms supporting infant language acquisition are poorly understood.

Information about neural mechanisms could eventually identify early neural markers of

developmental disorders of language, as neural processing is automatic and key neural

mechanisms may differ in their efficiency between infants. To explore this possibility, here we

contribute novel longitudinal neural data from a relatively large sample of 113 infants, utilising

EEG recordings taken during the first year of life while infants listened to sung or rhythmically

spoken (chanted) nursery rhymes. Nursery rhymes were chosen because of their rhythmic

properties, as infant behavioural research has suggested that a sensitivity to acoustic rhythm is a

universal precursor of language acquisition (3). *A priori*, it was expected that individual

differences in infants' cortical tracking of sung and chanted speech may predict later language

outcomes. Accordingly, a range of neural measures are used to explore relations with subsequent

language acquisition using a selection of standardised and experimental phonological, gestural

and lexical measures administered between 12 to 24 months of age (4).

The auditory neuroscience of adult speech processing has revealed that accurate encoding

via neuronal oscillations of amplitude modulations (AMs) nested in the envelope of the speech

signal is fundamental to many different aspects of language processing (5–12). Regarding earlier

development, and building on these adult insights, Temporal Sampling (TS) theory describes

how the low-frequency neural oscillations that encode speech rhythm patterns may be key to

learning and extracting phonological information from the speech signal (13,14). According to

TS theory, auditory cortical networks operating at delta, theta, beta and gamma frequencies

'sample' the envelope of the speech signal at matched frequencies, thereby underpinning the

phonological (linguistic) encoding of speech by infants and children. Cortical signals have been

shown to track various key properties of speech, using methodological advances such as

Temporal Response Functions (TRFs), an analysis technique that explores how neural signals

encode continuous sensory stimuli. Adult TRF studies have revealed how cortical tracking of the

speech envelope reflects both bottom-up and top-down cortical processes in adult listeners,

linking to selective attention, prediction, signal parsing and speech intelligibility (7–9,15–18).

Recent infant EEG data have revealed robust cortical tracking of the speech envelope (19–21),

particularly in the delta and theta frequency bands (1). Indeed, patterns of cortical tracking in

delta and theta frequency bands make distinct contributions to speech encoding by adults (7), and

there is evidence for preferential tracking in the delta band during the first year of life ($\sim$0.5Hz–4

Hz, see 1,22). Consequently, TS theory proposes that language acquisition by infants may

depend in part on successful low-frequency cortical tracking of speech envelope information.

This hypothesis regarding low-frequency cortical tracking is tested here by using TRFs.

Further, low-frequency cortical oscillations do not work in isolation from each other but

show dynamic interactions during speech processing. There is a nested hierarchy of cortical

oscillations that matches the frequency of multiple components of the speech signal (22). Low-

frequency (delta, theta) neural phase dynamics temporally organize the amplitude of high-

frequency signals in a process known as phase-amplitude coupling (PAC; (23–25)). Adult

electrophysiological and MEG recordings have shown that PAC between delta-gamma, delta-

beta and theta-gamma is induced in response to continuous speech (22,26,27), as well as to

rhythmic speech stimuli (sung and chanted nursery rhymes, (28)). PAC is thought to group

speech information into linguistic units (6,29), and accordingly could constitute another source

of individual differences in language acquisition. Like natural speech, rhythmic speech induces

PAC in infants and adults (1). Interestingly, PAC emerges as early as 4 months of age and

remains stable, while both delta- and theta-band cortical tracking show developmental changes

over the first year of life (1). Whilst the exact mechanisms linking PAC and language are still

being explored in the adult literature, it is clear that PAC plays a key role in the initial encoding

processes that are necessary for phonological learning and speech comprehension (18,22,26).

Therefore, it is likely that successful PAC in infants may relate to later language outcomes.

Studies relating infant cortical tracking to later language outcomes are rare (30).

However, infants at risk for dyslexia differ from not-at-risk infants in measures of acoustic

processing that relate to cortical tracking, with implications for later language development (31).

Meanwhile, research on the role of cortical dynamics in language development has progressed by

studying older children with language difficulties. For example, recent EEG modelling studies

(32) based on TS theory have shown that delta-theta PAC during passive story listening is

atypical in children with developmental language disorder (DLD) while the theta/delta power

ratio during passive story listening is atypical in children with developmental dyslexia. Children

with dyslexia have poor phonological processing skills while children with DLD have poor

syntactic skills. Based on (32), we can expect that individual differences in the theta/delta power

ratio for infants may also predict subsequent language development. Furthermore, previous

electrophysiological and magnetoencephalography studies testing TS theory with children with

dyslexia have revealed a significant role for delta band encoding in both phonological and

vocabulary development (33–36). Filtered speech interventions that enhance delta-band

information also improve the theta-delta oscillatory power ratio during speech listening for

dyslexic children (34). These neural data from children suggest that cortical tracking within

specific low-frequency bands (i.e. delta, 0.5–4 Hz and theta, 4–8 Hz) and their interactive

dynamics may be central to language acquisition by infants (13,37,38).

In the current study, we explore the potential role of individual differences in low-frequency

cortical tracking, PAC and theta-delta power relations in predicting individual differences in later

language acquisition. Participating infants were drawn from the Cambridge UK BabyRhythm

project, a longitudinal study of 122 infants (122 recruited, 113 retained) from two- to 42-months-of-age, investigating early neural and motor tracking of auditory, visual and audiovisual rhythms in relation to language outcomes. Here we examine whether increases in low-frequency EEG power spectral density (PSD) in the delta and theta range, low-frequency cortical tracking (delta, theta and alpha; alpha band data are used as a control band, as TS theory does not accord a role to alpha-band cortical tracking regarding initial language acquisition) and PAC in response to rhythmic audiovisual speech predict performance on multiple language and communication measures administered in later infancy and toddlerhood. EEG data were collected longitudinally at 4-, 7- and 11-months in response to both nursery rhymes and during a silent state control period. The neural outcome measures were then used to predict subsequent performance in phonological, gestural and lexical measures taken from 12 to 24 months of age. It was expected that stronger cortical tracking would be predictive of better language outcomes (30). As PAC did not show development with age in (1), no strong predictions were made. However, given the DLD modelling data noted earlier (32), poorer theta-driven PAC may be linked to worse language outcomes. Further, given the dyslexia modelling data, a higher theta/delta power ratio may be linked to worse language outcomes (32). In Araújo et al., preprint, higher ratios in the dyslexic group were not driven by group differences in either delta or theta power per se, but by their joint dynamics.

2 Materials and Methods

2.1 Infant participants and ethics

All the data reported here are from the longitudinal Cambridge UK BabyRhythm project (122 infants recruited, 113 retained). The cohort was split into two halves based on recruitment

date, to allow for confirmatory analyses of neural factors by comparing EEG data from the first

and second halves of the cohort. The present results primarily address findings from the second

half of the sample. Although results from the first half of the sample are reported in (1); it's

important to highlight that minor discrepancies exist between the values presented here and those

in (1). These variations stem from nuanced adjustments made to the analytical methods,

elaborated upon within each respective section. Due to participant withdrawals, missed

appointments and data exclusion the final number of data points for the second half of the sample

was, ~4-months (N=56), ~7-months (N=51), ~11-months (N=52). Missing data were due to

missed appointments or technical issues with the data files, ~4-months (N=4), ~7-months (N=10)

and ~11-months (N=7). Other infants were excluded due to having fewer than 42 data epochs

available after pre-processing, ~4-months (N=1), ~7-months (N=0) and ~11-months (N=2).

The two halves of the cohort were then combined for the novel language analyses

reported here (language outcome data were still being collected when Attaheri et al., 2022 (1)

was being prepared). The full cohort provided EEG data for analysis as follows, ~4-months

(N=110, aged 115.5 ± 5.0 days), ~7-months (N=104, aged 213.5 ± 6.1 days) and ~11-months

(N=107, aged 333.6, ± 5.0 days) [mean ± standard deviation (*SD*)].

Infants were recruited from a medium sized city in the United Kingdom and surrounding

areas via multiple means including flyers in hospitals, schools and antenatal classes, research

presentations at maternity classes and online advertising. All infants were born full term (37-42

gestational weeks) and had no diagnosed developmental disorder. The study was reviewed by the

Psychology Research Ethics Committee of the University of Cambridge. Parents gave written

informed consent after a detailed explanation of the study and families were reminded that they

could withdraw from the study at any point during the repeated appointments (8 EEG recordings

at 2-, 4-, 5-, 6-, 7-, 8-, 9- and 11-months; 6 language follow-ups at 12-, 15-, 18-, 24-, 30- and 42-

months). For the current report, the EEG data collected at 4, 7 and 11 months are utilized, as the

nursery rhyme stimuli were only presented to the infant during these recording visits. The

language outcome data are analysed using EEG measures computed from the full Cambridge UK

BabyRhythm cohort (N=113) rather than separating the first and second halves of the cohort.

167 2.2 Stimuli

A selection of 18 common English language nursery rhymes were chosen as the stimuli.

Audio-visual stimuli of a singing female (head only) were recorded using a Canon XA20 video

camera at 1080p, 50fps and with audio at 4800 Hz. A native female speaker of British English

used infant directed speech to melodically sing (for example "Mary Mary Quite Contrary") or

rhythmically chant (for nursery rhymes like "There was an old woman who lived in a shoe") the

nursery rhymes whilst listening to a 120 bpm metronome played to one ear. Although the nursery

rhymes had a range of beat rates and indeed some utilized a 1 Hz rate, the metronome was used

to keep the singer on time. The beat was not present on the stimulus videos, but it ensured that a

consistent quasi-rhythmic production was maintained throughout the 18 nursery rhymes. To

ensure natural vocalisations the nursery rhyme videos were recorded while being sung or

rhythmically chanted to an alert infant.

179 2.3 EEG data collection

Infants were seated in a highchair (with neck pillow support if required) approximately

one meter in front of their primary care giver, within a sound-proof acoustic chamber (on

occasion it was necessary for the infant to sit on the lap of the primary care giver). EEG data

were recorded at a sampling rate of 1000 Hz using a GES 300 amplifier connected to an

appropriately sized 64 channel infant net (Geodesic Sensor Net, Electrical Geodesics Inc.,

Eugene, OR, USA). The infant was seated ~650mm away from the presentation screen and

sounds were presented at 60dB (checked by sound level meter) from speakers (Q acoustics 2020i

driven by a Cambridge Audio Topaz AM5 stereo amplifier) placed either side of the screen.

Whilst the infant attended to the screen, 18 nursery rhyme videos were randomized and then

played sequentially in a block, each block was then repeated 3 times (54 videos, with a

presentation time of 20' 33'' in total). If the infant lost attention to the screen an attention

grabber video was played at the end of that nursery rhyme. If the infant became too fussy a short

break was allowed before continuing, otherwise the session was ended. All infants included for

analysis listened to at least 2 repetitions of each nursery rhyme (minimum of 36 nursery rhymes,

lasting 13' 42''). This stimulus period was followed by 5 minutes of silent recording (hereafter

silent state). The silent state was recorded after the nursery rhymes, and commenced after a 2-

minute break to check on the welfare of the participant and parent. The nursery rhymes were

given first in order to maximise infant attention. Note that this may risk stimulus after-effects

during the silent state (39), however the time-course of cortical tracking of the EEG signal is not

comparable to fNIRs (5,40), and such after-effects would work against our hypothesis by making

it more difficult to demonstrate cortical tracking when comparing the nursery rhyme data to the

silent state data, making the PSD and TRF analyses more stringent. To ensure compliance from

the infant, and therefore a good EEG signal during the silent state, it was sometimes necessary

for a researcher to sit alongside the infant. To ensure consistency across participants, the

researcher performed the same action of silently blowing bubbles and showing the same picture

book to the infant during this period.

2.4 EEG preprocessing

All analyses were conducted with custom-made scripts in Matlab 2017a (The MathWorks, Inc., Natick, MA) incorporating the EEGLab toolbox (41). The EEG data recorded in response to the nursery rhymes and the silent state were treated as one continual data set for the initial preprocessing steps. Therefore, they underwent the same cleaning and filtering procedure prior to epoching.

The four (channels 61, 62, 63 and 64) facial electrodes (present on the larger nets) were excluded. The EEG data, from the remaining 60 channels, was filtered (*pop_eegfiltnew* function of EEGLab toolbox) into a broadband signal (0.5-45 Hz for PSD and PAC, methods for acquiring PAC bands of interest further detailed in (1)) or the frequency band of interest (delta, 0.5-4 Hz, theta, 4-8 Hz or alpha [control band] 8-12 Hz for the mTRF analyses). A zero-phase bandpass Hamming windowed FIR filter (transition band widths of 2 Hz with cutoff frequencies at -6 dB, 0-5 Hz, 3-9 Hz and 7-13 Hz respectively) was implemented. The EEG data were down sampled to 100 Hz to reduce the computational load. Next, Artifact Subspace Reconstruction (ASR; *clean_asr* EEGLab function (41,42)) was used to clean noise artifacts from the data by identifying and removing bad principal components via a modified PCA procedure (see (1)). ASR effectively removed the transient or large-amplitude artifacts common in infant EEG data. Further bad channels were identified via probability and kurtosis and were interpolated (via spherical interpolation). A channel was rejected and interpolated if it was 3*SD* away from the mean (average number of interpolated channels; 4mo = 6.9, 7mo = 6.5, 11mo = 6.6). Finally, all channels were re-referenced to a 60-channel average reference.

EEG responses to the 18 nursery rhymes were epoched into trials aligned to the start of a nursery rhyme phrase (e.g. "Mary had a little lamb"). Each nursery rhyme contained

approximately 5 phrases (*M number of phrases per nursery rhyme ± SD*: *4.61 phrases ±4.22,*

range =1:20*) producing EEG responses to 83 phrases (*M length ± SD*: 4.23sec ±0.88) which

were repeated a maximum of 3 times in the experiment (249 epochs in total). This epoching

procedure was selected as infant EEG typically contains short, irregular, movement artifacts and

using shorter epochs increases the likelihood of generating a whole artifact-free epoch whilst

maintaining long enough epochs for optimising the model fit with the mTRF toolbox (43). The

infants occasionally rested their head or touched the EEG cap in such a way that specific

channels showed short bursts of noise whilst the rest of the channels remained clean. To account

for this, and retain a larger volume of data, a second stage of channel rejection and interpolation

was conducted epoch by epoch. Per epoch, probability and kurtosis were used to identify bad

channels and were interpolated (via spherical interpolation) if they were 3*SD* away from the

mean. Finally, bad epochs were rejected with the *pop_autorej* function (EEGLab), removing

epochs with fluctuations above 1000uV and values outside a 3*SD* of the probability threshold.

Due to the interpolation and ASR procedure, less than 1% of the epochs were rejected via

pop_autorej. This step was maintained as a sanity check to ensure all extreme fluctuations had

been removed. After preprocessing, the EEG responses to each of the nursery rhymes were

averaged (data across a maximum of three repeats of the 83 nursery rhyme phrases were

averaged), thereby creating 83 "averaged trials". This averaging was conducted to improve the

signal to noise ratio of the data for the mTRF analysis. Due to the increased susceptibility of PSD

and PAC analyses to noise, all the remaining epochs were concatenated back into a continual

signal and further manual rejection was conducted to remove noisy periods of data from the

signal prior to running the PSD and PAC analyses.

2.5 Multivariate Temporal Response Function (mTRF)

TRFs describe the linear relationship between an input and an output signal, taking into account that such a relationship may not be instantaneous and could extend over a certain window of time (43). TRFs can be estimated with system identification methodologies, such as the multivariate temporal response function (mTRF), which is a de-convolution method based on multiple lagged linear regression. Here, we applied the mTRF in the backward direction to reconstruct the stimulus envelope from the EEG signals. In turn, this informs us on the strength of the stimulus-EEG relationship, which we take as an index of cortical tracking of the stimulus envelope. The backwards 'stimulus reconstruction' mTRF model has important advantages compared to the forward linear encoding model which has previously been used in infant studies (19,20). Specifically, backward TRFs combine data from all the EEG channels simultaneously in a multivariate manner to reconstruct the univariate stimulus envelope, while forward TRF models only predict one EEG channel at a time. Pearson's correlation (r) was used to validate how well the reconstructed stimulus envelope correlated to the original (a metric for cortical tracking). This process was repeated in a cross-validated manner for the M-1 trials (maximum of 83 rotations if the data set had the full 83 trials remaining after pre-processing). The average r value represented how well the model reconstructed the stimulus envelope, when trained with either the delta (0.5-4Hz), theta (4-8Hz) or alpha (8-12Hz) EEG band data. Chance r scores were created for each participant to measure the average stimulus reconstruction (r) that could be obtained by chance. The chance TRF procedure was conducted per participant for each frequency band. To obtain a chance permutation of the data, whilst maintaining phase integrity, each of the stimulus envelopes were first reversed and a random circular shift was applied. Next, the mTRF cross-validation was run in the same way as the real data (see above for details), to

give a stimulus reconstruction (r) value. The r value using the shuffled model was used as the

chance r score per participant. Please note that the chance r values in the tables, for the first half

of the sample, differ slightly from (1) as a single random model per participant rather than an

average random model was implemented here. A linear mixed effects model (LMEM) was next

constructed using the same parameters as in Attaheri et al., 2022 (1), to discover if the mean of

the "real" model was larger than the mean of the "shuffled" model, across participants.

280   2.6 mTRF auditory stimuli preprocessing

The envelope of the auditory signal was extracted by taking the absolute value of the

analytic signal generated by the Hilbert transform (Matlab). As the lower frequencies of the

envelope are linearly relatable to the EEG signal (10,11,44) the envelope of the stimuli was

filtered between 0.5 Hz and 15 Hz (lowpass; 6th order Butterworth filter. Highpass; 9th order

Butterworth filter). The resultant envelopes were normalised using *nt_normcol* (Noisetools).

Finally, the stimulus envelopes were down sampled to 100 Hz to match the EEG signal. The

preprocessed and filtered envelopes, from now on will be referred to as envelopes.

288   2.7 Phase amplitude coupling (PAC)

To assess PAC, a modified version of the WinPACT plugin (EEGLab) (41) was used to

acquire normalised modulation index (nMI) values (45) a measure adapted from Canolty and

colleagues' modulation index (MI) (23). This measure is a widely validated metric of the

coupling strength and preferred phase between two frequencies. For each infant's EEG data,

low-frequency phase (LFP) and high-frequency amplitude (HFA) were extracted with a zero-

phase FIR filter (Hamming window), separately for all 60 electrodes. LFP data were filtered in

1Hz steps from 2 to 8 Hz. Each centre frequency had a 2 Hz filter bandwidth around the centre

frequency. HFA centre frequencies were extracted from 17.5 Hz to 42.5 Hz, in 5 Hz steps with a

5 Hz filter bandwidth around the centre frequency, Next, a normalised modulation index value

was calculated for each channel using a 5 second sliding analysis window (with 2.5 second

overlaps)(23,45). To ensure only windows displaying significant PAC were considered, 200

surrogate statistical iterations were created for each window. A statistically normalised MI

estimate was obtained for each analysis window by subtracting the mean and dividing by the

standard deviation obtained from a Gaussian fit of surrogate MI estimates (nMI = (Canolty's MI

– surrogate MI Mean) / surrogate MI Std). This statistical procedure is based on the work of

Özkurt and Schnitzler, 2011 (45) and implemented via the winPACT EEGLab plugin. Each

iteration of the surrogate data was created by shuffling the high-frequency amplitude time series

via circular rotation. A MI estimate was obtained for each of the 200 surrogate data iterations,

from which a 95% confidence interval (CI) was calculated. This step accounted for the mean and

standard deviation of the surrogate data set, thus creating an appropriate threshold for the

frequency band analysed. Finally, the 95% CIs were averaged across all windows to generate a

statistical threshold for each channel per participant. All the nMI windows were averaged per

channel for each of the PAC pairs (i.e. each LFP and HFA step) separately for each participant.

The channel exhibiting the strongest MI, within predefined phase and amplitude band groupings

(delta/beta, delta/gamma, theta/beta, theta/gamma, with delta=2-4Hz, theta=4-8Hz, beta=15-

30Hz and gamma=30-45Hz), was taken forward for the LMEM and for the group level grand

average plots. Due to the Nyquist frequency required for our 5 second window size, the first

frequency band analysed in the PAC was 2Hz. See (1) for a detailed description of the PAC

method and analysis.

2.8 Spectral analysis (periodogram PSD estimate)

A one-sided PSD estimate was conducted separately for each channel using the periodogram function (Matlab). The grand average PSD across channels was next calculated by averaging across channels, recording type and ages. Visual inspection revealed three frequency peaks centered around 1.92 Hz, 4.05 Hz and 4.35 Hz (Fig. S1). The spectra for each individual age group per condition were next calculated and taken forward for further analysis (Fig. 1). To allow for individual participant variation, the maximum value within a 0.25Hz window centered on the peak of interest was taken per participant for further analysis. A smaller window was used around the peaks of interest than used in Attaheri et al., 2022 (1), this was necessary to avoid overlap between the 4.05 Hz and 4.35 Hz windows. See (1) for a detailed description of the PSD method and analysis.

2.8.1 Theta / delta (PSD) ratio

The PSD peak per participant at 4.35 Hz and 1.92 Hz, taken from the spectral decomposition of the EEG signal described above, was used to create a theta / delta ratio calculation. Per participant, the 4.35 Hz PSD value was divided by the 1.92 Hz PSD value (4.35/1.92) to give a theta / delta ratio.

2.9 Language measures

A full description of the language measures used here can be found in Rocha et al., (4). The parent-estimated measures were taken from a carer-completed questionnaire, the UK-CDI. The UK-CDI estimates both receptive and productive vocabulary (46), which we refer to here as language comprehension or language production. In the current report the parent-estimated 24-month CDI measures were selected for analysis, as by the age of 2 years both word comprehension and word production data showed a wide range of scores, enabling more

sensitive measurement of individual differences (see (4), for detailed rationale). Only measures

up to the age of 24 months are included here, as the 30- and 42-month measures are still being

processed. Five measures were selected as the infant-led (in that the infant made an active

response) language measures. These were (i) gesture (adapted from (47)), in which pointing

towards a present and absent target was used as a pre-verbal measure of communicative intent at

12 months, and (ii) the computerized comprehension task (CCT), a touch screen-based measure

of infant vocabulary administered at 18 months, in which the infant is asked to select a picture of

a named item (48). Thirdly a non-word repetition (NWR) task based on (49), was used to

measure the child's productive phonology at 24 months and generate three different measures:

(iii) accurate reproduction of consonants, (iv) number of syllables reproduced, and (v) stress

patterns produced (canonical or non-canonical, see Rocha et al., (4)). The NWR task measured

toddlers' phonological production of both nonsense words (e.g. "punky") and real words (e.g.

"puppy") via a game about naming a series of toys.

354   2.10 Language outcomes: statistics

Our statistical approach was decided *a priori* and was established when exploring the

language outcome measures (4). We ran a multivariate model for each of the infant-led and

parent-estimated groups of measures. Given that missing cases vary across the multiple tasks,

and that the multivariate models employed use listwise deletion, additional follow-up univariate

models for each of the predictors were run separately, to utilise the breadth of data collected and

maximise statistical power. This approach was applied across multiple BabyRhythm studies

using different neural measures as language predictors (see (50), using phase measures, and (51),

which uses neural responses to visual-only speech), accordingly an alpha threshold of 0.05 was

applied. As the follow-up univariate analyses required multiple comparisons, one per language

measure, they were Bonferroni-corrected for the number of language measures in the group (i.e.

5 for infant-led, and 2 for parent-estimated). Respective alpha levels were therefore 0.01 and

0.025. Although the main multivariate tests may show that some predictors had a significant

effect and others did not, we did not remove any of the original covariates in the univariate

models, to ensure comparability between the multivariate and univariate tests.

Due to the inherent differences between the infant-led and parent-estimated measures,

separate multivariate linear models were conducted for each, providing a global measure of

language abilities taken from either infant-led or parent-estimated language measures.

Multivariate linear models (using Rstudio *lm* function) were run separately for the PSD, mTRF,

theta-delta ratio and PAC measures. Secondly, Bonferroni-corrected individual univariate linear

models were run, which typically included more infants, to explore the individual contribution of

each language measure regarding these different neural variables.

3 Results

3.1 PSD, mTRF, PAC and theta/delta ratio

First, we present the power spectral density (using a periodogram; PSD), cortical tracking

(using multivariate temporal response functions; mTRFs), phase-amplitude coupling (normalised

modulation index, nMI; PAC) and theta/delta PSD ratio analyses using EEG recorded from the

second half of our cohort of infants. We compare these data to the analyses from the first half of

the cohort (1), and also merge the neural data to explore these neural factors in the combined full

sample of 113 infants. Due to missed recording sessions, technical issues with data files and not

enough trials after preprocessing, less than 113 data points are included in each analysis.

Furthermore, outlier analysis was also conducted to remove extreme data points that would

compromise each of the LMEMs. The number of data points included are given separately for

each analysis.

3.1.1 Power spectral density (PSD) analyses

Figure 1, Spectral decomposition of the EEG signal averaged across the a) first half, b) second
half and c) full infant cohort (1–12 Hz plotted, from 0.5 to 45 Hz calculated), in response to

nursery rhyme stimulation. A periodogram was used to obtain a power spectral density (PSD)
estimate separately for 4- (red), 7- (green) and 11- (blue) months data. First the PSD estimate
was averaged across channels per participant. Bold lines indicate the grand average mean values
and pale shading plots the standard deviation of the data across the participants. Outlier analysis
was also conducted to remove extreme data points that would compromise the LMEM. Panel A
is reproduced from Attaheri et al. 2022 (1), with permission.

The distribution of low-frequency oscillations in the EEG recorded from the second half

of the sample and the full cohort were obtained using the power spectral density (PSD) estimate

(Fig. 1). The results from the first half of the cohort (1) are reproduced for comparison. Given the

differences in the location of the PSD peaks for the first and second halves of the cohort, the

distribution of PSD peaks in low-frequency power for the full sample was first established by

creating a grand average PSD estimate across channels, recording type (nursery rhymes, silent

state) and ages. Visual inspection revealed three frequency peaks centered around 1.92 Hz, 4.05

Hz and 4.35 Hz (Fig. S1, and in the second half of the sample, Fig. S2). The distribution of the

PSD peaks were similar to those observed in the first half of the sample (see (1) and Fig. S3). It

is important to note that the PSD peaks matched the most prominent modulation spectrum peaks

that were present in the nursery rhyme stimuli (see Fig. S4–S6), which were respectively within

the delta and theta band definitions used for the EEG analyses (delta, 0.5–4 Hz; theta, 4–8 Hz).

Whilst infant frequency bandings appear to change throughout infancy (52–55), there are no

prior infant data regarding center frequencies in studies using speech stimuli. Therefore, we

selected the delta and theta boundaries based on the prior adult speech processing literature, and

this decision is supported by the modulation bands identified by modelling the temporal

modulation architecture of the 18 nursery rhymes used in our study (56)). Further, when the EEG

from the full cohort was analysed, the ~4Hz peak split into two locations, peaking at 4.05 Hz and

4.35 Hz. Both theta peaks were therefore retained for further investigation. Whilst the relevant

data are provided in the results text, the mean and standard error of the PSD peak amplitudes for

both silent state and nursery rhyme recording conditions, are also provided in Table S1. Plots of

the silent state PSD are also provided to aid comparison (Fig. S7). To investigate whether the

PSD was significantly higher in response to the nursery rhymes compared to the silent state, and

whether this relationship was affected by the age of the participants, a linear mixed effects model

(LMEM) was conducted separately for the 1.92 Hz, 4.05 Hz and 4.35 Hz peaks. A random

intercept (AR1) per participant was included to account for individual differences across the 3

recording sessions (4-, 7- and 11-months). The full sample submitted to the LMEM was as

follows, 4-months (N=96), 7-months (N=88) and 11-months (N=92). This process was

conducted separately for the first half of the sample, the second half of the sample and the full

cohort (Table 1). To allow for individual variation across participants, a maximum peak value

per participant was taken from the participants' 60 channel averaged data, within 0.25 Hz

windows centered around the peak of interest (1.92 Hz, 4.05 Hz and 4.35 Hz). Note that a

smaller window was used around the peaks of interest than used in Attaheri et al., 2022 (1), this

was necessary to avoid overlap between the 4.05 Hz and 4.35 Hz windows. Fixed effects of

recording type (nursery rhymes, silent state), age (4-, 7- or 11-months) and a recording type by

age interaction were included in the model. The ~4.35 Hz peak is reported first, as this was the

most robust.

~4.35 Hz

	1^{st} half of sample		2^{nd} half of sample	
	F	p	F	p
Recording type	F(1, 124.16 = 12.51)	*0.001*	F(1, 107.77 = 7.67)	*0.007*
Age	F(2, 153.41 = 10.01)	*$7.7x10^{-5}$*	F(2, 114.77 = 4.86)	*0.009*
Recording type * Age	F(2, 123.93 = 3.30)	*0.04*	F(2, 107.90 = 3.54)	*0.032*

	Full sample	
	F	p
Recording type	F(1, 236.60 = 19.25)	*$1.7x10^{-5}$*
Age	F(2, 281.12 = 14.13)	*$1.0x10^{-6}$*

Recording type * Age	F(2, 236.48 = 6.95)	*0.001*

Table 1. Tests of fixed effects for LMEM conducted on PSD power peak at 4.35Hz. Results reported for separate LMEMs run on the first half, second half and full sample of data. Significant results (p<0.05) denoted by **bold italic text**. The Satterthwaite approximation was applied to approximate the degrees of freedom, due to the missing cases.

The 4.35 Hz peak showed the largest stimulus-related increase in PSD of the three peaks analysed. The LMEM's revealed a significant effect of recording type, with the nursery rhyme (NR) condition producing significantly greater PSD power than the silent state (SS; Table 1). A significant effect of age was observed, with ~4.35 Hz power increasing parametrically with age from 4- months to 7- months to 11- months (estimates of fixed effects provided in Table S2). The observed recording type by age interaction was due to a developmental increase in NR induced power above the SS. Bonferroni-corrected pairwise comparisons showed that PSD increased significantly between 4- and 11-months (M increase = 18.41, SE ± 7.43, $p = 0.043$) as well as between 4- and 7- months (M increase = 30.56, SE ± 5.77, $p = 6.7 \times 10^{-7}$). Hence there was a developmental increase in 4.35 Hz PSD for both the nursery rhymes and the silent state. This may indicate age-related maturation of theta activity in general. Importantly, we also observed a stimulus-driven interaction with age for the 4.35 Hz PSD peaks, with the largest increase observed at 11-months (4-months, NR, *M* = 38.9 *SE* ± 4.9; SS, *M* = 37.9 *SE* ± 5.5. 7-months, NR, *M* = 63.0 *SE* ± 6.9; SS, *M* = 50.6 *SE* ± 7.2. 11-months, NR, *M* = 84.8 *SE* ± 5.8; SS, *M* = 53.1 *SE* ± 6.0). This showed that the interaction was driven by a greater PSD in response to the nursery rhymes at 11-months. This pattern replicated the theta PSD findings observed in the first half of the sample (see (1)). In summary, the peak at 4.35 Hz showed robust replication across the first and second half of the sample, with consistent increases in NR related power displaying a developmental increase in PSD magnitudes, above SS, indicating a highly reliable peak.

~4.05 Hz

	1st half of sample		2nd half of sample	
	F	p	F	p
Recording type	F(1, 223.47 = 2.72)	0.100	F(1, 101.13 = 5.65)	*0.019*
Age	F(2, 160.05 = 4.74)	*0.010*	F(2, 111.74 = 2.39)	0.097
Recording type * Age	F(2, 223.50= 1.07)	0.345	F(2, 101.07 = 1.50)	0.228

	Full sample	
	F	p
Recording type	F(1,424.40 = 7.33)	*0.007*
Age	F(2,298.70 = 3.54)	*0.030*
Recording type * Age	F(2, 424.16 = 1.77)	0.172

Table 2. Tests of fixed effects for LMEM on PSD power peak at 4.05Hz. Results reported for
separate LMEMs run on the first half, second half and full sample of data. Significant results
(p<0.05) denoted by ***bold italic text***. The Satterthwaite approximation was applied to
approximate the degrees of freedom, due to the missing cases.

For the 4.05 HZ peak, the LMEM results replicated less well across the first half of the

sample, the second half of the sample and the full cohort. Significant effects of both recording

type and age were only observed when analysing the full cohort. The significant effect of age

was driven by the 7-month ($M = 63.1$ *SE* ± 6.0) PSD power being larger than the 11-month ($M =$

52.5 *SE* ± 3.2), however this result is in part due to the 0.25 Hz window not fully encapsulating

the 11-month peak for all participants (Fig. 1). A significant effect of recording type was

observed in the second half of the sample, with a trend observed in the first half of the sample

(Table 2). A significant effect of age was observed in the first half of the sample with a trend

towards significance observed in the second half of the sample. As the data were more robust for

the 4.35 HZ peak, and as the 0.25 Hz window did not fully capture the 4.05 Hz peak for all

participants, the 4.35 Hz PSD peak was taken forward as the 'theta' peak for analyses with the

language data.

~1.92 Hz

	1st half of sample		2nd half of sample	
	F	p	F	p
Recording type	F(1, 96.59 = 9.71)	*0.002*	F(1, 87.49 = 0.25)	0.616

Age	F(2, 127.50 = 1.00)	0.371	F(2, 109.00 = 3.34)	*0.039*
Recording type * Age	F(2, 96.74 = 0.61)	0.544	F(2, 87.50 = 7.39)	*0.001*

	Full sample	
	F	p
Recording type	F(1,188.40 = 8.47)	*0.004*
Age	F(2,232.10 = 0.36)	0.698
Recording type * Age	F(2,96.74 = 0.61)	*0.021*

Table 3. Tests of fixed effects for LMEM on PSD power peak at 1.92Hz. Results reported for separate LMEMs run on the first half, second half and full sample of data. Significant results (p<0.05) denoted by **bold italic text**. The Satterthwaite approximation was applied to approximate the degrees of freedom, due to the missing cases.

For the 1.92 Hz peak, the LMEM conducted on the full cohort revealed a significant effect of recording type (Table 3). However, there were differences in this effect across the first and second halves of the sample. Upon further inspection, the non-significant level of recording type in the second half of the sample was due to the large amount of PSD power in the silent state (SS) data at 4-months (4-month SS, $M = 116.3$ *SE ± 10.6,* NR, M= 97.3 *SE ± 10.2*). Posthoc paired sample t-tests on the 7- and 11-month data, for the second half of the sample, both showed a consistent stimulus-induced increase in PSD power, matching the first half of the sample (7-month SS, *M= 116.3 SE ±13.5,* NR, *M*= 124.2 *SE ±13.1, p = 0.007;* 11-month SS, *M=76.7 SE $\pm$ 11.9 ,* NR, *M*= 94.0 *SE ± 11.6 , p =* 1.90×10^{-27}). A significant recording type by age interaction was observed in both the second half of the sample and the full cohort. In conclusion, the 1.92Hz stimulus-related increase in PSD power is broadly replicated across the first and second halves of the sample once the anomalous SS power, present in the 4-month data in the second half of the sample, was taken into account.

3.1.2 Cortical tracking (mTRF) analyses

Figure 2, mTRF replication. Grand averaged stimulus reconstruction values (Pearson's r) per frequency band in the a) first half, b) second half and c) full infant cohort. Each bar represents the grand average r value across the participants at the 3 age points: 4-months (red), 7-months

(green) and 11-months (blue). Age responses are grouped along the X axis by EEG frequency
band, delta (0.5–4 Hz) theta (4–8 Hz) and alpha (8–12 Hz). Light grey boxes signify the average
chance *r* value across subjects, calculated separately for each band. Outlier analysis was
conducted to remove extreme values which would compromise the LMEMs. Panel a is
reproduced from Attaheri et al. (2022), with permission.

A linear mixed effects model (LMEM) was constructed using the same parameters as

described in Attaheri et al., 2022 (1) to examine potential developmental changes in cortical

tracking in the delta, theta and alpha (control) bands. The full sample submitted to the mTRF

LMEM was as follows, 4-months (N=108), 7-months (N=104) and 11-months (N=106). Main

effects of data type (real *r* values, chance *r* values), frequency band (delta, theta or alpha) and

age (4-, 7- or 11-months) were investigated along with interactions between data type by

frequency band, data type by age and age by frequency band. A random intercept (AR1) per

participant was included to account for individual differences across the 3 recording sessions (4-,

7- and 11-months).

The main pattern of results replicated between the first and second halves of the sample,

with significant fixed effects of data type and frequency band, as well as significant interactions

between data type by frequency band and age by frequency band (Table 4). There was no

significant effect of age when analysing the second half of the sample in isolation, although this

effect was present in the first half of the sample (Table 4). When using the full cohort, tests of

fixed effects showed significant effects of data type, frequency band and age. These results

indicate that the real *r* values (cortical tracking) in response to nursery rhymes were significantly

above randomly permuted *r* values and that the cortical tracking values differed by frequency

band and age. The significant interaction of frequency band by data type indicated that cortical

tracking in the delta and theta bands was driven by their values being significantly above the

randomly permuted *r* values (as compared to the alpha base case; Table S3). The marginal means

across the three ages, revealed that cortical tracking estimates (r values) were significantly higher

in both delta (random permuted r values, $M = 0.015$, $SE \pm 0.001$; real r values, $M = 0.03$, $SE \pm$

0.001), and theta bands (random permuted r values, $M = 0.011$, $SE \pm 0.001$; real r values, $M =$

0.016, $SE \pm 0.001$), compared to the alpha base case (random permuted r values, $M = 0.007$, SE

± 0.001; real r values, $M = 0.007$, $SE \pm 0.001$). Furthermore, alpha tracking was again not

significantly above chance (Table S3) when using the full sample, mimicking the results from

the first half of the sample (1). The significant interaction between age and frequency band

showed that the relative difference in r values between the three age groups were different in the

three frequency bands. This was predominantly driven by 4-month delta ($M = 0.25$, $SE \pm 0.001$)

producing larger r values than the 11-month delta ($M = 0.20$, $SE \pm 0.001$) and the 7- month delta

($M = 0.23$, $SE \pm 0.001$).

In summary, the main effects reported by Attaheri et al. 2022 (1) were replicated in the

second half of the sample, apart from some main effects of age being inconsistent when

comparing the first and second halves of the cohort for the ~4.05Hz power peak. Complete

consistency regarding age was only achieved for the 4.35Hz power peak, which is the measure

taken forwards below. The replication of PSD and cortical tracking results in the lower

frequencies strengthens the conclusion that delta and theta EEG responses track the acoustic

envelope of sung and chanted speech from the age of 4-months and that this infant neural

response is robust and reliable.

Cortical tracking

	1st half of sample		2nd half of sample	
	F	*p*	*F*	*p*
Data type	F(1,851.69) = 157.53	*2.82 x10⁻³³*	F(1,766.02) = 83.62	*5.29x10⁻¹⁹*
Age	F(2, 519.14) = 3.15	*0.044*	F(2, 198.42) = 1.61	0.202

Frequency band	F(2, 852.32) = 217.39	*5.17x10^{-77}*	F(2, 767.80) = 189.87	*1.01x10^{-67}*
Data type * age	F(2, 851.68) = 2.39	0.093	F(2, 766.00) = 0.65	0.524
Data type * freq. band	F(2, 851.67) = 71.47	*2.02x10^{-29}*	F(2, 765.96) = 32.63	*2.51x10^{-14}*
Age *freq. band	F(4, 852. 31) = 3.06	*0.016*	F(4, 768.17) = 3.52	*0.007*

	Full sample	
	F	*p*
Data type	F(1, 1625.31) = 228.71	*1.92x10^{-48}*
Age	F(2, 603.11) = 3.77	*0.024*
Frequency band	F(2, 1627.67) = 401.38	*1.99x10^{-142}*
Data type * age	F(2, 1625.30) = 2.53	0.080
Data type * freq. band	F(2, 1625.29) = 97.11	*1.45x10^{-40}*
Age *freq. band	F(4, 1627.89) = 5.74	*1.38x10^{-4}*

Table 4. Tests of fixed effects for LMEM of cortical tracking across data type (real r values, random permutation *r* values), frequency band (delta, theta or alpha) and age (4-, 7- or 11-months). Results reported for separate LMEMs run on the first half, second half and full sample of data. Significant results (p<0.05) denoted by ***bold italic text***. The Satterthwaite approximation was applied to approximate the degrees of freedom, due to the missing cases.

3.1.2 Phase amplitude coupling (PAC) analysis

Figure 3, PAC replication. Violin plots of phase amplitude coupling (normalised modulation index, nMI) at 4-months (red), 7-months (green) and 11-months (blue) for the a) first half, b) second half and c) full infant cohort. The nMIs, from all significant analysis windows, were averaged for each individual infant. The PAC pairing with the maximum nMI per infant, from within the pre-defined frequency bands of interest (delta/beta, delta/gamma, theta/beta and theta/gamma; beta 2–4 Hz, theta 4–8 Hz, beta 15–30 Hz and gamma 30–45 Hz), were included in the grand average violin plot. Horizontal black and red lines represent the group mean and median respectively. Outlier analysis was conducted to remove extreme data points that would compromise the LMEM. Panel A is reproduced from Attaheri et al. 2022 (1), with permission.

The PAC analyses (Fig. 3) were conducted for the second half of the sample and the full

cohort, replicating the LMEM conducted with the first half of the sample (1). The full sample

submitted to the PAC LMEM was as follows, 4-months (N=108), 7-months (N=103) and 11-

months (N=107). This LMEM included fixed effect factors of data type (real nMI values or

chance nMI values), low-frequency phase (delta or theta), high-frequency amplitude (beta or

gamma) and age (4-, 7- or 11-months). Interactions of data type by age, by low-frequency phase,

data type by high-frequency amplitude, data type by age, low-frequency phase by age, high-

frequency amplitude by age and low-frequency phase by high-frequency amplitude were also

included in the model, with a random intercept (AR1) per participant (see methods section and

Attaheri et al., 2022 (1) for detailed methods). Tests of fixed effects are reported in Table 5 and

estimates of fixed effects are provided in Table S4. The LMEM intended to explore whether the

nMI values were significantly above chance and if there were any significant differences when

either delta or theta was the low-frequency carrier phase or when either beta or gamma were the

high-frequency amplitudes, and if this relationship was affected by age.

The main pattern of results replicated between the first and second halves of the sample,

with significant fixed effects of data type and low-frequency phase, as also reflected in the full

cohort. The main effect of LFP arose because the nMI values where larger when theta (M

=2.568, SE ± 0.013) was the low-frequency phase coupling with high-frequency amplitudes

rather than delta (M = 2.405, SE ± 0.013, Table 5). A non-significant effect of HFA suggested

that the beta and gamma amplitudes coupled equally as strongly with delta and theta phases

(Table 5). A significant interaction between data type and LFP was also observed. Whilst both

delta and theta displayed significant above-chance PAC, the significant interaction of LFP by

data type reflected the fact that the relative difference between chance and real nMI values was

significantly greater in the theta band than in the delta band, delta (chance nMI, M=1.770, SE ±

0.016; real nMI, M = 3.041, SE ± 0.016), Theta (chance nMI, M=1.772, SE ± 0.016; real nMI, M

= 3.364, SE ± 0.016). Whilst the main pattern of results replicated between the first and second

halves of the sample, one inconsistency was that the second half of the sample also showed a

significant interaction between data type and HFA and between data type and age. However, as

no main effects of age nor HFA were observed in the second half of the sample, these results are

difficult to interpret.

PAC

	1^{st} half of sample		2^{nd} half of sample	
	F	p	F	p
Data type (rand vs real nMI)	F(1, 1178.08) = 4921.51	*<0.001*	F(1, 1113.52) = 4409.87	*<0.001*
Low freq. phase (LFP)	F(1, 1178.08) = 55.79	*<0.001*	F(1, 1113.52) = 64.77	*<0.001*
Age	F(2,686.71) = .06	0.941	F(2, 202.36) = 1.81	0.167
High freq. amp (HFA)	F(1, 1178.08) = .05	0.824	F(1, 1113.52) = 1.68	0.196
Data type * LFP	F(1, 1178.08) = 54.34	*<0.001*	F(1, 1113.52) = 62.86	*<0.001*
Data type * HFA	F(1, 1178.08) = .00	0.949	F(1, 1113.52) = 4.00	*0.046*
Data type * Age	F(2, 1178.08) = 1.68	0.187	F(2, 1113.52) = 3.26	*0.039*
LFP * Age	F(2, 1178.08) = .47	0.626	F(2, 1113.52) = 2.10	0.123
HFA * Age	F(2, 1178.08) = 2.01	0.134	F(2, 1113.52) = .28	0.757
LFP * HFA	F(1, 1178.08) = .11	0.738	F(1, 1113.52) = 2.48	0.115
Age * LFP * HFA	F(2, 1178.08) = .64	0.526	F(2, 1113.52) = .69	0.504

	Full sample	
	F	p
Data type (rand vs real nMI)	F(1,2269.31) = 9217.24	*<0.001*
Low freq. phase (LFP)	F(1,2269.31) = 118.88	*<0.001*
Age	F(2,790.87) = 1.40	0.247
High freq. amp (HFA)	F(1,2269.31) = 1.10	0.294
Data type * LFP	F(1, 2269.31) = 115.42	*<0.001*
Data type * HFA	F(1, 2269.31) = 1.78	0.182
Data type * Age	F(2, 2269.31) = 2.31	0.100
LFP * Age	F(2, 2269.31) = 2.21	0.110
HFA * Age	F(2, 2269.31) = 1.79	0.167
LFP * HFA	F(1, 2269.31) = 0.71	0.399
Age * LFP * HFA	F(1, 2269.31) = 0.44	0.642

Table 5, Tests of fixed effects generated from LMEM, assess how PAC (nMI) varied across data
types (real or random nMI values), low-frequency phase (LFP), high-frequency amplitude (HFA)
and age (4-, 7- or 11-months). Results reported for separate LMEMs run on the first half, second
*half and full sample of data. Significant results (p<0.05) denoted by **bold italic text**. The*
Satterthwaite approximation was applied to approximate the degrees of freedom, due to the
missing cases.

3.1.4 Theta/delta PSD power ratio

A significant difference between the theta/delta PSD power ratio across the three ages

was also found (Fig. 4). Ratio values significantly increased with age (4 months, $M = 0.546$, $SE \pm$

0.027 ; 7 months, $M = 0.78$, $SE \pm 0.073$; 11 months, $M = 1.017$, $SE \pm 0.110$), indicated by a one-

way ANOVA ($F(2,273) = 9.54$, $p = 9.9 \times 10^{-5}$). The ratio increased because the amount of theta

PSD power increased relative to the amount of delta PSD power as our sample aged from 4- to

11- months (see Figure 1). For comparison, the child modelling using EEG recorded from

children with and without dyslexia aged around 9 years showed that a higher theta/delta ratio

was associated with poorer phonological awareness (32). In the filtered speech intervention study

with dyslexic children (34), in which delta-band modulations in natural speech were amplified

via filtering, the theta/delta ratio improved when the children were listening to the filtered speech

(compared to natural speech) because delta power increased. Accordingly, the theta/delta ratio

appears to be a sensitive index of speech-related neural dynamics in infants and children.

Figure 4, Theta / delta PSD ratio. The PSD peak per individual at 4.35 Hz and 1.92 Hz, taken
from the spectral decomposition of the EEG signal, was used to create a theta / delta ratio
calculation (4.35/1.92). Data provided at 4-months (red), 7-months (green) and 11-months (blue)
with standard error bars in black.

3.2 Neural measures as predictors of language acquisition

Multivariate linear models were used to test whether the neural findings reported in 3.1

(for the full sample) were related to the different language measures using the analysis strategy

described at 2.1 above. As explained in Rocha et al., (4), these language measures were selected

as most reliable in previous analyses of the Cambridge UK BabyRhythm cohort, as they had the

largest number of infants contributing data and scores were neither at floor nor ceiling (4). As

noted, the parent-estimated measures were receptive and productive vocabulary scores from the

UK-CDI at 24 months (46) and the infant-led measures were (i) The ability to point at 12-

months, (ii) Word recognition at 18-months on a computerised infant-operated comprehension

test (CCT), and (iii-v) Phonological processing at 24-months, as measured by non-word

repetition of consonants, syllables and stress patterns (NWR). Due to the inherent difference in

parent-estimated and infant-led measures of language, separate multivariate and univariate linear

models were run for each set of scores. The results of these analyses for both parent-estimated

and infant-led outcomes are reported in Tables 6-11. The univariate analyses often included more

cases, as the multivariate analyses only included infants who contributed data points to all the

language measures as well as to the neural measure being analysed.

3.2.1. PSD and language outcomes

The 11-months EEG data were selected for comparison to both the parent-estimated and

infant-led language measures, due to the observed developmental maturation of the ~4 Hz peak

(see Figure 1). As described above, the 4.35 Hz peak was the more robust and consistent of the

two theta peaks (see Tables 1 and 2). It was therefore used for predicting the language outcomes.

The analyses for the 4-month and 7-month EEG are also provided as Tables S5 to S11 (with 4.05

Hz peak results reported in Tables S7-S9). The analyses for the ~1.92 Hz peak are also reported

here. 77 infants provided complete data for all the parent-estimated measures and PSD power

results, whilst 70 infants provided data for all the infant-led language measures and PSD power.

Regarding global language outcomes, the multivariate models indicated that increased

~1.92 Hz PSD power in response to nursery rhymes did not lead to significant prediction of

either infant-led ($F(5,65) = 1.257$, $p= 0.285$) nor parent-estimated ($F(2,75) = 0.560$, $p= 0.574$)

outcomes (Table 6). However, the univariate linear models revealed a trend for pointing at 12-

months to be correlated with increases in ~1.92 Hz power (Table 6a; $p= 0.014$, Bonferroni

corrected trend α values, $p = <0.02$ for infant-led univariate models).

The multivariate analyses for ~4.35 Hz PSD power in response to nursery rhymes

showed no effect on language outcomes for the infant-led measures ($F(5,65) = 1.758$, $p= 0.134$,

Table 7a). This was also the case in the univariate analyses. However, for the parent-estimated

measures, increased theta power led to a significant *decrease* (F(2,75) = 3.18, *p*= 0.047) in the

global parent-estimated language outcomes (Table 7b). It is important to note that whilst the

univariate linear model analyses indicated that comprehension and production did not

individually make significant contributions to this prediction (Table 7b), the direction of the beta

estimates suggest that higher power was associated with fewer known words later in

development. Accordingly there is a global effect of theta power on parent-estimated (UK-CDI)

language outcomes, but this is not driven by either single parent-estimated measure.

~1.92 Hz PSD power at 11 months

a, Infant-led language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Infant-led language measures	5	65	1.257	0.29
Univariate linear models	*Df*	$\beta_{estimate}$	*SE*	p
NWR Consonants, 24 months	1,79	5.23×10^{-5}	8.50×10^{-5}	0.95
NWR Syllables, 24 months	1,79	5.54×10^{-4}	8.56×10^{-4}	0.53
NWR Stress, 24 months	1,79	5.51×10^{-4}	7.83×10^{-4}	0.48
CCT, 18 months	1,79	7.16×10^{-4}	6.32×10^{-4}	0.26
Pointing, 12 months	1,90	0.003	0.001	**0.014**

b, Parent-estimated language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Parent-estimated language measures	2	75	0.560	0.57
Univariate linear models	*Df*	$\beta_{estimate}$	*SE*	p
CDI comprehension, 24 months	1,76	0.373	0.445	0.40
CDI production, 24 months	1,76	0.179	0.465	0.70

Table 6, Multivariate and univariate linear models, investigating whether ~1.92 Hz PSD power at
11 months predicted either infant-led or parent-estimated language outcomes. The table details
the multivariate linear models followed by the univariate linear models describing whether ~1.92
Hz PSD power at 11 months predicted, a) infant-led language measures or b) the parent-
estimated language measures. Bonferroni correction for multiple comparisons in the univariate
models led to modified significant alpha levels of, p = <0.01 for infant-led and p = <0.025 for
parent-estimated (denoted by ***bold italic text***). Modified alpha level for trends are p = <0.02 for
infant-led and p = <0.05 for parent-estimated (denoted by **bold text**).

~4.35 Hz PSD power at 11 months

a, Infant-led language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Infant-led language measures	5	65	1.76	0.13
Univariate linear models	Df	$\beta_{estimate}$	SE	p
NWR Consonants, 24 months	1,79	-4.93×10^{-4}	3.69×10^{-4}	0.19
NWR Syllables, 24 months	1,79	-8.36×10^{-4}	3.65×10^{-4}	0.025
NWR Stress, 24 months	1,79	-5.31×10^{-4}	3.40×10^{-4}	0.12
CCT, 18 months	1,79	-1.62×10^{-4}	2.76×10^{-4}	0.56
Pointing, 12 months	1,90	-2.95×10^{-4}	-5.98×10^{-4}	0.62

b, Parent-estimated language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Parent-estimated language measures	2	75	3.18	***0.047***
Univariate linear models	Df	$\beta_{estimate}$	SE	p
CDI comprehension, 24 months	1,76	-0.086	0.209	0.68
CDI production, 24 months	1,76	-0.346	0.215	0.11

Table 7, Multivariate and univariate linear models, investigating whether ~4.35 Hz PSD power at
11 months predicted either infant-led or parent-estimated language outcomes. The table details
the multivariate linear models followed by the univariate linear models describing whether ~4.35
Hz PSD power at 11 months predicted a) infant-led language measures or b) the parent-estimated
language measures. Bonferroni correction for multiple comparisons in the **univariate models** led
to modified significant alpha levels of, p = <0.01 for infant-led and p = <0.025 for parent-
estimated (denoted by ***bold italic text***). Modified alpha level for trends are p = <0.02 for infant-
led and p = <0.05 for parent-estimated (denoted by **bold text**).

A further pair of multivariate linear model analyses showed that a larger theta/delta PSD

power ratio (4.35 Hz / 1.92 Hz) led to a decrease in performance for both infant-led (F(5,65) =

2.684, *p*= 0.029, Table 8a) and parent-estimated (F(2,75) = 3.65, *p*= 0.031, Table 8b) global

outcomes. The more theta power that was present relative to delta power, the worse the infants'

performance on either type of outcome. Again, the univariate linear models suggested that these

global effects were not driven by any single individual measure. However, it was measures of

language *production* that showed *p* values <.10 (NWR of syllables, *p*= .07, CDI production, *p*=

.09).

Theta / delta ratio of PSD power

a, Infant-led language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Infant-led language measures	5	65	2.68	***0.029***
Univariate linear models	**Df**	**$\beta_{estimate}$**	**SE**	**p**
NWR Consonants, 24 months	1,79	-0.019	0.032	0.56
NWR Syllables, 24 months	1,79	-0.058	0.032	0.07
NWR Stress, 24 months	1,79	-0.044	0.029	0.14
CCT, 18 months	1,79	-0.023	0.023	0.32
Pointing, 12 months	1,90	-0.065	0.049	0.19

b, Parent-estimated language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Parent-estimated language measures	2	75	3.65	***0.031***
Univariate linear models	**Df**	**$\beta_{estimate}$**	**SE**	**P**
CDI comprehension, 24 months	1,76	-6.760	16.44	0.68
CDI production, 24 months	1,76	-28.55	16.82	0.094

Table 8, Multivariate and univariate linear models, investigating whether delta/theta ratio of PSD
power (~1.92 Hz/4.35 Hz) at 11 months predicted either infant-led or parent-estimated language
outcomes. The table details the multivariate linear models followed by the univariate linear
models describing whether delta/theta ratio of PSD power (~1.92 Hz/4.35 Hz) at 11 months
predicted a) infant-led language measures or b) the parent-estimated language measures.
Bonferroni correction for multiple comparisons in the **univariate models** led to modified
significant alpha levels of, p = <0.01 for infant-led and p = <0.025 for parent-estimated (denoted
by ***bold italic text***). Modified alpha level for trends are p = <0.02 for infant-led and p = <0.05 for
parent-estimated (denoted by **bold text).**

3.2.3 mTRF and language outcomes

The 11-months EEG data were selected for analyses with both the parent-estimated and

infant-led language measures, as these were recorded the closest in time to the language

measures (please see Tables S12-15 for matching data for the earlier age groupings; these did not

reach significance). 89 infants provided complete data for all the parent-estimated measures and

a cortical tracking value, whilst 82 infants provided data for all the infant-led language measures

and a cortical tracking value.

The multivariate linear models regarding the parent-estimated language measures (Table

9a) displayed a trend (not significant at an alpha value of 0.05) towards better language

outcomes, with increased accuracy of delta-band cortical tracking in response to nursery rhymes

predicting better outcomes ($F_{(2,87)} = 3.00$, $p = 0.055$). No multivariate effect was present for the

infant-led measures, ($F_{(5,77)} = 0.83$, $p = 0.350$). However, the univariate linear model analyses

showed that both CDI comprehension and CDI production at 24 months were significantly

predicted by delta band cortical tracking (p's = 0.02, Table 9b). By contrast, the accuracy of theta

band cortical tracking did not predict either infant-led ($F_{(5,77)} = 0.886$, $p = 0.495$) or parent-

estimated ($F_{(2,87)} = 0.250$, $p = 0.780$) language outcomes (Table 10). This was unexpected given

the adult literature. As alpha cortical oscillations did not show above chance cortical tracking,

comparison to language measures were not considered (however the data are provided in Tables

S16-S18 for completeness)

Delta cortical tracking

a, Infant-led language measures

Multivariate linear model	Df_{num}	Df_{denom}	F	p
Infant-led language measures	5	77	0.83	0.35
Univariate linear models	Df	$\beta_{estimate}$	SE	p
NWR Consonants, 24 months	1,96	1.65	1.85	0.37
NWR Syllables, 24 months	1,96	1.43	1.85	0.44
NWR Stress, 24 months	1,96	2.89	1.67	0.09
CCT, 18 months	1,93	1.04	1.46	0.48
Pointing, 12 months	1,105	2.91	2.95	0.33

b, Parent-estimated language measures

Multivariate linear model	Df_{num}	Df_{demom}	F	p
Parent-estimated language measures	2	87	3.00	**0.055**
Univariate linear models	Df	$\beta_{estimate}$	SE	p
CDI comprehension, 24 months	1,90	2043.00	877.43	*0.02*
CDI production, 24 months	1,90	2235.55	948.65	*0.02*

Table 9, Multivariate and univariate linear models, investigating whether delta cortical tracking
at 11 months predicted either infant-led or parent-estimated language outcomes. The table details
the multivariate linear models followed by the univariate linear models describing whether delta
cortical tracking at 11 months predicted a) infant-led language measures or b) the parent-
estimated language measures. Bonferroni correction for multiple comparisons in the **univariate**
models led to modified significant alpha levels of, p = <0.01 for infant-led and p = <0.025 for
parent-estimated (denoted by ***bold italic text***). Modified alpha level for trends are p = <0.02 for
infant-led and p = <0.05 for parent-estimated (denoted by **bold text**).

Theta cortical tracking

a, Infant-led language measures

Multivariate linear model	Df_{num}	Df_{demom}	F	p
Infant-led language measures	5	77	0.89	0.50
Univariate linear models	Df	$\beta_{estimate}$	SE	p
NWR Consonants, 24 months	1,96	-4.32	2.91	0.14
NWR Syllables, 24 months	1,96	-3.51	2.92	0.23
NWR Stress, 24 months	1,96	-2.24	2.67	0.40
CCT, 18 months	1,93	-2.72	2.28	0.24
Pointing, 12 months	1,105	5.72	4.61	0.22

b, Parent-estimated language measures

Multivariate linear model	Df_{num}	Df_{demom}	F	p
Parent-estimated language measures	2	87	0.25	0.78
Univariate linear models	Df	$\beta_{estimate}$	SE	p
CDI comprehension, 24 months	1,90	-828.31	1452.9	0.57
CDI production, 24 months	1,90	-1114.12	1570.3	0.48

Table 10, Multivariate and univariate linear models, investigating whether theta cortical tracking
at 11 months predicted either infant-led or parent-estimated language outcomes. The table details
the multivariate linear models followed by the univariate linear models describing whether theta
cortical tracking at 11 months predicted a) infant-led language measures or b) the parent-
estimated language measures. Bonferroni correction for multiple comparisons in the **univariate**
models led to modified significant alpha levels of, p = <0.01 for infant-led and p = <0.025 for
parent-estimated (denoted by ***bold italic text***). Modified alpha level for trends are p = <0.02 for
infant-led and p = <0.05 for parent-estimated (denoted by **bold text**).

3.2.4 PAC and language outcomes

Phase amplitude coupling (PAC) at 4-months was selected for the language outcome

analyses as PAC did not show any developmental changes indicating that the early presence of

stronger PAC may be informative (please see Tables S19-20 for matching data for the earlier age

groupings). 87 infants provided complete data for all the parent-estimated measures and a PAC

value, whilst 81 infants provided data for all the infant-led language measures and a PAC value.

Multivariate linear models were run separately to investigate whether delta-beta, delta-gamma,

theta-beta or theta-gamma PAC respectively predicted the language outcome measures.

Considering delta as the LFP first, increased delta-beta PAC predicted a global *decrease*

in the parent-estimated ($F(2,85) = 3.38$, $p= 0.039$) language measures (Table 11b). The

univariate linear models indicated that CDI production at 24 months drove this overall effect

(Table 11b). Increased delta-beta PAC did not predict the infant-led language outcome measures

in the multivariate analysis ($F(5,76) = 0.447$, $p= 0.815$) nor in the univariate analyses (Table

11a). By contrast, stronger delta-gamma PAC showed a trend to increase parent-estimated

language measures ($F(2,85) = 2.68$, $p= 0.075$), with the significant univariate linear model

indicating that CDI comprehension at 24 months drove this trend (*p=0.02*). Delta-gamma PAC

did not predict the infant-led language outcome measures ($F(5,76) = 0.447$, $p= 0.815$, Table

11a), and the delta-gamma univariate models showed similar results.

When theta was the LFP, the multivariate linear models indicated that increased theta-

gamma PAC predicted a significant global increase in the infant-led language measures ($F(5,76)$

$= 4.126$, $p= 0.002$, Table 11a) and the parent-estimated ($F(2,85) = 4.86$, $p= 0.010$) language

measures (Table 11b). Bonferroni-corrected univariate linear models highlighted that better non-

word repetition of syllables at 24-months ($p= .002$) was the strongest contributing factor. Theta-

beta PAC showed no effect on later language outcomes in the multivariate analyses, both for the

infant-led measures ($F(5,76) = 0.897$, $p= 0.488$) and the parent-estimated measures ($F(2,85) =$

146, $p= 0.240$, Table 11b). The univariate analyses showed similar results.

In summary, theta-gamma PAC was associated with a significant increase in both the

infant-led and parent-estimated language outcomes, in both the multivariate and univariate

analyses. Trend effects suggested that delta-gamma PAC also contributed to increases in parent-

estimated language outcomes. Conversely, delta-beta PAC predicted a significant global

decrease in parent-estimated language measures, and theta-beta PAC exerted no effects. The data

suggest that the coupling of LFPs to gamma-band HFAs could be a key factor in language

acquisition.

PAC

a, Infant-led language

	Delta-beta		Delta-gamma		Theta-beta		Theta-gamma	
Multivariate linear model	F	p	F	p	F	p	F	p
Infant-led language measures	0.81	0.54	0.45	0.81	0.90	0.49	4.13	*0.002*
	Delta-beta		Delta-gamma		Theta-beta		Theta-gamma	
Univariate linear models	$\beta_{est.}$	P	$\beta_{est.}$	P	$\beta_{est.}$	p	$\beta_{est.}$	p
NWR Consonants, 24 months	-0.11	0.17	0.05	0.51	-0.04	0.52	0.08	0.09
NWR Syllables, 24 months	-0.09	0.27	0.03	0.70	-0.04	0.49	0.14	*0.002*
NWR Stress, 24 months	-0.07	0.37	0.02	0.79	-0.08	0.05	0.0.	0.65
CCT, 18 months	-0.13	0.03	0.03	0.54	0.04	0.39	0.06	0.05
Pointing, 12 months	-0.1	0.41	-0.03	0.78	-0.08	0.39	0.02	0.77

b, Parent-estimated language

	Delta-beta		Delta-gamma		Theta-beta		Theta-gamma	
Multivariate linear model	F	p	F	p	F	p	F	p
Parent-est. language measures	3.38	*.039*	2.68	**0.07**	1.46	0.24	4.86	*0.01*
	Delta-beta		Delta-gamma		Theta-beta		Theta-gamma	
Univariate linear models	$\beta_{est.}$	p	$\beta_{est.}$	p	$\beta_{est.}$	p	$\beta_{est.}$	p
CDI comprehension, 24 mo	-53.4	0.20	85.0	*0.02*	-26.5	0.38	24.2	0.28
CDI production, 24 mo	-103.3	*0.02*	77.1	0.06	3.35	0.92	60.1	*0.01*

Table 11, Multivariate and univariate linear models, investigating whether delta-beta, delta-
gamma, theta-beta or theta-gamma PAC at 4 months predicted either infant-led or parent-
estimated language outcomes. The table details the multivariate linear models followed by the
univariate linear models describing whether PAC at 4 months predicted a) infant-led language
measures or b) the parent-estimated language measures. Bonferroni correction for multiple
comparisons in the **univariate models** led to modified significant alpha levels of, p = <0.01 for

infant-led and p = <0.025 for parent-estimated (denoted by ***bold italic text***). Modified alpha level
for trends are p = <0.02 for infant-led and p = <0.05 for parent-estimated (denoted by **bold text**).

4 Discussion

According to recent extensions of Temporal Sampling theory (13,37) cortical tracking of

infant directed speech (IDS) by delta- and theta-band neural signals and delta-theta neural

dynamics may be key factors in explaining individual differences in the developmental

trajectories for language acquisition. In the Cambridge UK BabyRhythm infant sample assessed

here (N=113), individual differences in delta-band cortical tracking (in the univariate analyses),

theta-gamma PAC (in both univariate and multivariate analyses), theta power (in the multivariate

analyses) and theta/delta power ratio (multivariate analyses) were all significant predictors of

language outcomes in infants during the first year of life. These infant data are consistent with

neural oscillatory factors that have been found to explain individual differences in language

development in children, for example the fidelity and synchronicity of delta-band encoding of

speech envelope information for phonological development (35,57). They are also consistent

with a recent EEG modelling study based on child recordings during passive speech listening,

which showed that a higher theta/delta power ratio was related to phonological difficulties in

dyslexia (32), and a child MEG study which showed that amplifying speech information in the

delta band reduced this power ratio when dyslexic children were listening to filtered speech (34).

Further, the infant neural oscillatory factors identified here were broadly consistent in terms of

replicating our previous findings with the first half of the infant cohort (1). The presence and

maturation of low-frequency (<12 Hz) cortical oscillations (PSD), cortical speech tracking

(mTRF) and phase amplitude coupling (PAC) in response to sung IDS was largely similar across

both halves of the longitudinal Cambridge UK BabyRhythm cohort.

The replication of stronger delta band tracking than theta band tracking in the youngest

(4-month-old) infants suggests that delta band cortical tracking may be a key early building

block for language acquisition. Efficient delta-band cortical tracking would enable infants to lock

on to the information-rich strong (or stressed) syllables in the speech signal, acoustic landmarks

that occur on average every 500ms across languages (58) . This mechanism would enable the

infant brain to begin to represent the acoustic structure of human speech (56,59). Dynamic

interactions with theta-band cortical tracking may then support further linguistic learning (see

below). The importance of delta-band cortical tracking for language outcomes in our univariate

analyses is consistent with child MEG studies, which show that typically-developing children

exhibit efficient delta-band tracking of speech-in-noise, but not efficient theta-band tracking

(60). The finding that delta band cortical tracking predicted early language comprehension and

production in the univariate analyses is also congruent with research in the adult brain, where

delta-band tracking is associated with key language-related skills such as perceptual grouping

(61,62), discourse-level parsing related to phrasing (63,64) and comprehending speech-in-noise

(65).

In many adult studies, the theta band is considered the key linguistic oscillator (6). In our

infant cohort, better theta-gamma PAC at 4-months was associated with better outcomes in both

the infant-led and parent-estimated multivariate analyses (assessing global language outcomes)

and the univariate analyses. Adult work shows how PAC could operate as a neural mechanism

for grouping fine-grained phonetic information into syllabic units, that in turn could be

concatenated to build up words and phrases (22,26,29). The univariate analyses for theta-gamma

PAC revealed that nonword repetition, a phonological output measure (specifically syllable

production at 24-months) and CDI production (parental estimation of vocabulary output) were

strong drivers of these effects. By contrast, delta-based PAC predicted increases in the language

outcome measures when gamma was the HFA (parent-estimated univariate analyses), but

decreases in the language outcome measures when beta was the HFA (parent-estimated

multivariate and univariate analyses). Further, not all the effects found for theta-band cortical

tracking were positive in nature. Theta-band PSD increased significantly during the first year of

life, and greater theta-band PSD was associated with poorer language outcomes as measured by

parental estimates of comprehension and production at 24-months in the multivariate analyses.

Greater theta-band PSD also showed a trend to be associated with poorer infant-led nonword

repetition in the univariate analyses (p= .025). A larger theta/delta PSD power ratio was also

associated with poorer language outcomes, as measured by multivariate analyses of both parent-

estimated and infant-led measures. This latter finding underscores the importance of considering

neural dynamics when assessing developmental trajectories. In the previous EEG modelling

work with 9-year-old children with and without dyslexia that investigated theta/delta ratios, a

dyslexia-specific pattern of delta and theta power dynamics during natural speech processing was

seen to become progressively more atypical the worse the child's phonological awareness, and

this was not related to age (32). Further, this ratio could be improved for children with dyslexia

by increasing delta power via a filtered speech manipulation (34).

The infant literature is sparse regarding PAC dynamics. For adults, theta-gamma PAC

has been suggested as a neural mechanism for processing phonetic information in the speech

signal (6,66,67). When the same rhythmic speech stimuli used in the current study were

presented to adult participants, theta/gamma coupling was stronger than delta/gamma coupling

(28). For our infant participants presented with the same nursery rhyme stimuli, theta-gamma

PAC was the consistent predictor of language outcomes in both the multivariate and the

univariate analyses. Indeed, it is the phase of theta band rather than delta band oscillations that

adjust to speech rate variations for adults, coupling with gamma amplitudes (26). Whilst both

delta and theta phases couple with higher-frequency amplitudes in the infant brain, it seems that

theta-based PAC is more strongly predictive of early language outcomes. The univariate analyses

suggested that measures involving speech production, namely nonword repetition and productive

vocabulary, were important regarding these relations. The adult PAC literature suggests that

delta-beta PAC is particularly associated with measures of speech production via audio-motor

dynamics (27,68). In our data, stronger delta-beta PAC before 1 year of age was associated with

poorer language outcomes at 24 months. In the univariate analyses, the negative association was

for CDI production ($p=$ .02). This may be a transitory developmental effect. As the infants in our

cohort continue to develop and produce more speech, it may be that delta-beta PAC will begin to

play a positive role in predicting language outcomes.

The current study has several limitations. Recording EEG with infants is an inherently

noisy process, and this affected our data, as different infants had to be excluded from different

analyses, reducing sample size for the longitudinal language outcome assessments, particularly

regarding the multivariate analyses. A key source of noisy data are infant fussing and movement,

and noise was particularly apparent in the 7-month-old recordings for our sample. This could be

age-related, as by 7 months of age infants have a much larger range of movement. A second

limitation is that even though our study followed a relatively large cohort of infants (122

recruited and 113 retained), a study with a larger participant pool may have had increased

chances of uncovering developmentally important time-lagged associations. Notwithstanding the

limitations of the study, here we present a large longitudinal sample demonstrating that

individual differences in neural dynamics in response to speech stimuli predict some aspects of

early infant language acquisition.

5 Conclusion

Estimates of the efficiency of low-frequency cortical tracking, PSD, phase-amplitude

coupling and theta-delta power dynamics in an infant's first year of life predict their early

language outcomes. Importantly, the neural mechanisms measured here are automatic processes

that are inherent to the human brain, and as such may represent physiological priors for language

learning in a Bayesian sense (69). Accordingly, individual differences in the efficiency of these

mechanisms in the pre-verbal stage of language learning assessed here are unlikely to reflect top-

down linguistic processes such as language comprehension (18). Delta-band cortical tracking at

11 months in the univariate analyses and theta–gamma PAC at 4 months of age in both the

univariate and multivariate analyses significantly predicted language comprehension and

production at 24 months, with the latter predicting both word production and nonword repetition

in the univariate analyses. Conversely, increased theta PSD power and a larger theta/delta ratio

were associated with decreased performance on the language outcome measures in our

multivariate analyses. While other studies have also indicated a predictive relationship between

cortical tracking efficiency in infancy and language development (30), the current study is the

first to show that individual differences in PAC and theta/delta power dynamics also predict

infant language outcomes.

A Declaration of Competing Interest

The authors do not report any conflicts of interest.

Author Contributions

Adam Attaheri: EEG Paradigm Development, EEG Preprocessing, Investigation - Data Curation, Formal Analysis - design, creation and implementation of analysis, Writing - Original Draft.

Áine Ní Choisdealbha: EEG Preprocessing - Investigation - Data Curation - Writing - Review and Editing

Sinead Rocha: Investigation - Data Curation, Writing - Review and Editing

Perrine Brusini: EEG Paradigm Development, Investigation - Data Curation

Giovanni Di Liberto: Formal analysis, Writing - Review and Editing

Natasha Mead: Investigation - Data Curation

Helen Olawole-Scott: Investigation - Data Curation

Panagiotis Boutris: Investigation - Data Curation

Samuel Gibbon: Investigation - Data Curation

Isabel Williams: Investigation - Data Curation

Christina Grey: Investigation - Data Curation

Maria Alfaro e Oliveira - Data Curation

Carmel Brough - Data Curation

Shelia Flanagan – Data curation

887   *Usha Goswami*: Conceptualization - Methodology, Funding Acquisition, Supervision, Project

888   Administration, Writing - Original Draft

Data Sharing agreement

The analyses were conducted by using publicly available MATLAB toolboxes that can be

downloaded at http://audition.ens.fr/adc/NoiseTools/,

https://sccn.ucsd.edu/eeglab/download.php, https://github.com/mickcrosse/mTRF-Toolbox.

The custom code used for the mTRF, PSD and PAC analysis can be downloaded from

https://osf.io/q9s5k/. The final data included in the manuscript figures and statistics can also be

downloaded from https://osf.io/q9s5k/.

Funding sources

897   This project has received funding from the European Research Council (ERC) under the

898   European Union's Horizon 2020 research and innovation programme (grant agreement No.

899   694786).

Acknowledgements

901   We would like to thank all the families and infants who kindly donated their time to this project.

902   We would also like to thank Henna Ahmed for their time spent as an RA on the Cambridge UK

903   BabyRhythm project. Finally, we would like to acknowledge Mike X Cohen's statistical

904   textbooks and online materials for guiding our statistical analysis.

References

1.   Attaheri A, Choisdealbha ÁN, Di Liberto GM, Rocha S, Brusini P, Mead N, et al. Delta- and
theta-band cortical tracking and phase-amplitude coupling to sung speech by infants.
NeuroImage. 2022 Feb 15;118698.

2. Kuhl PK. Early language acquisition: cracking the speech code. Nat Rev Neurosci. 2004
Nov;5(11):831–43.

3. Mehler J, Jusczyk P, Lambertz G, Halsted N, Bertoncini J, Amiel-Tison C. A precursor of
language acquisition in young infants. Cognition. 1988;29(2):143–78.

4. Rocha, S., Ní Choisdealbha, Á., Attaheri, A., Mead, N., Olawole-Scott, H., Grey, C.,
Williams, I., Gibbon, S., Boutris, P., Brusini, P., Brough, C., Alfaro e Oliveira, M., &
Goswami, U. 2024. Language Acquisition in the Longitudinal Cambridge UK BabyRhythm
Cohort. Collabra: Psychology, 10(1). https://doi.org/10.1525/collabra.92998

5. Obleser J, Kayser C. Neural Entrainment and Attentional Selection in the Listening Brain.
Trends Cogn Sci. 2019 Nov 1;23(11):913–26.

6. Giraud AL, Poeppel D. Cortical oscillations and speech processing: emerging computational
principles and operations. Nat Neurosci. 2012 Mar 18;15(4):511–7.

7. Ding N, Simon JZ. Cortical entrainment to continuous speech: functional roles and
interpretations. Front Hum Neurosci [Internet]. 2014 [cited 2020 Jul 3];8. Available from:
https://www.frontiersin.org/articles/10.3389/fnhum.2014.00311/full

8. Di Liberto GM, O'Sullivan JA, Lalor EC. Low-Frequency Cortical Entrainment to Speech
Reflects Phoneme-Level Processing. Curr Biol CB. 2015 Oct 5;25(19):2457–65.

9. Luo H, Poeppel D. Phase patterns of neuronal responses reliably discriminate speech in
human auditory cortex. Neuron. 2007 Jun 21;54(6):1001–10.

10. O'Sullivan JA, Power AJ, Mesgarani N, Rajaram S, Foxe JJ, Shinn-Cunningham BG, et al.
Attentional Selection in a Cocktail Party Environment Can Be Decoded from Single-Trial
EEG. Cereb Cortex N Y NY. 2015 Jul;25(7):1697–706.

11. Zion Golumbic EM, Ding N, Bickel S, Lakatos P, Schevon CA, McKhann GM, et al.
Mechanisms Underlying Selective Neuronal Tracking of Attended Speech at a 'Cocktail
Party.' Neuron. 2013 Mar 6;77(5):980–91.

12. Peelle J, Davis M. Neural Oscillations Carry Speech Rhythm through to Comprehension.
Front Psychol. 2012;3:320.

13. Goswami U. Speech rhythm and language acquisition: an amplitude modulation phase
hierarchy perspective. Ann N Y Acad Sci. 2019;1453(1):67–78.

14. Goswami U. A Neural Basis for Phonological Awareness? An Oscillatory Temporal-
Sampling Perspective. Curr Dir Psychol Sci. 2018 Feb 1;27(1):56–63.

15. Baltzell LS, Srinivasan R, Richards VM. The effect of prior knowledge and intelligibility on
the cortical entrainment response to speech. J Neurophysiol. 2017 Sep 6;118(6):3144–51.

16. Di Liberto GM, Crosse MJ, Lalor EC. Cortical Measures of Phoneme-Level Speech
Encoding Correlate with the Perceived Clarity of Natural Speech. eNeuro [Internet]. 2018
Apr 16 [cited 2020 Jul 3];5(2). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5900464/

17. Millman RE, Johnson SR, Prendergast G. The Role of Phase-locking to the Temporal
Envelope of Speech in Auditory Perception and Speech Intelligibility. J Cogn Neurosci.
2015 Mar 1;27(3):533–45.

18. Peelle JE, Gross J, Davis MH. Phase-locked responses to speech in human auditory cortex
are enhanced during comprehension. Cereb Cortex N Y N 1991. 2013 Jun;23(6):1378–87.

19. Jessen S, Fiedler L, Münte TF, Obleser J. Quantifying the individual auditory and visual
brain response in 7- month-old infants watching a brief cartoon movie. bioRxiv. 2019 Apr
17;610709.

20. Kalashnikova M, Peter V, Di Liberto GM, Lalor EC, Burnham D. Infant-directed speech
facilitates seven-month-old infants' cortical tracking of speech. Sci Rep [Internet]. 2018 Sep
13 [cited 2020 Jul 3];8. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6137049/

21. Ortiz Barajas MC, Guevara R, Gervain J. The origins and development of speech envelope
tracking during the first months of life. Dev Cogn Neurosci. 2021 Jan 20;48:100915–
100915.

22. Gross J, Hoogenboom N, Thut G, Schyns P, Panzeri S, Belin P, et al. Speech Rhythms and
Multiplexed Oscillatory Sensory Coding in the Human Brain. PLOS Biol. 2013 Dec
31;11(12):e1001752.

23. Canolty RT, Edwards E, Dalal SS, Soltani M, Nagarajan SS, Kirsch HE, et al. High Gamma
Power Is Phase-Locked to Theta Oscillations in Human Neocortex. Science. 2006 Sep
15;313(5793):1626–8.

24. Canolty RT, Knight RT. The functional role of cross-frequency coupling. Trends Cogn Sci.
2010 Nov;14(11):506–15.

25. Lakatos P, Shah AS, Knuth KH, Ulbert I, Karmos G, Schroeder CE. An oscillatory hierarchy
controlling neuronal excitability and stimulus processing in the auditory cortex. J
Neurophysiol. 2005 Sep;94(3):1904–11.

26. Lizarazu M, Lallier M, Molinaro N. Phase–amplitude coupling between theta and gamma
oscillations adapts to speech rate. Ann N Y Acad Sci. 2019 Oct;1453(1):140–52.

27. Arnal LH, Doelling KB, Poeppel D. Delta-Beta Coupled Oscillations Underlie Temporal
Prediction Accuracy. Cereb Cortex N Y N 1991. 2014/05/20 ed. 2015 Sep;25(9):3077–85.

28. Attaheri A, Panayiotou D, Phillips A, Ní Choisdealbha Á, Di Liberto GM, Rocha S, et al.
Cortical Tracking of Sung Speech in Adults vs Infants: A Developmental Analysis. Front

Neurosci [Internet]. 2022;16. Available from: https://www.frontiersin.org/article/10.3389/fnins.2022.842447

29. Hyafil A, Fontolan L, Kabdebon C, Gutkin B, Giraud AL. Speech encoding by coupled cortical theta and gamma oscillations. Brownell H, editor. eLife. 2015 May 29;4:e06213.

30. Menn KH, Ward EK, Braukmann R, van den Boomen C, Buitelaar J, Hunnius S, et al. Neural Tracking in Infancy Predicts Language Development in Children With and Without Family History of Autism. Neurobiol Lang. 2022 Aug 17;3(3):495–514.

31. Kalashnikova M, Goswami U, Burnham D. Sensitivity to amplitude envelope rise time in infancy and vocabulary development at 3 years: A significant relationship. Dev Sci. 2019 Nov 1;22(6):e12836.

32. Araújo J, Simons BD, Varghese P, Mandke K, Kalashnikova M, Macfarlane A, et al. Atypical encoding of speech identifies children with Dyslexia versus Developmental Language Disorder [Internet]. BioRxiv; Preprint. Available from: https://www.biorxiv.org/content/10.1101/2022.10.26.513864v1

33. Keshavarzi M, Di Liberto G, Gabrielczyk F, Wilson A, Macfarlane A, Goswami U. Atypical speech production of multisyllabic words by children with developmental dyslexia. 2022.

34. Mandke K, Flanagan S, Macfarlane A, Feltham G, Gabrielczyk F, Wilson AM, et al. Neural responses to natural and enhanced speech edges in children with and without dyslexia. Front Hum Neurosci [Internet]. 2023;17. Available from: https://www.frontiersin.org/articles/10.3389/fnhum.2023.1200950

35. Power AJ, Colling LJ, Mead N, Barnes L, Goswami U. Neural encoding of the speech envelope by children with developmental dyslexia. Brain Lang. 2016 Sep;160:1–10.

36. Power AJ, Mead N, Barnes L, Goswami U. Neural entrainment to rhythmic speech in children with developmental dyslexia. Front Hum Neurosci [Internet]. 2013 [cited 2020 Jul 3];7. Available from: https://www.frontiersin.org/articles/10.3389/fnhum.2013.00777/full

37. Goswami U. Language acquisition and speech rhythm patterns: an auditory neuroscience perspective. R Soc Open Sci. 2022 Jul 27;9.

38. Menn KH, Michel C, Meyer L, Hoehl S, Männel C. Natural infant-directed speech facilitates neural tracking of prosody. NeuroImage. 2022 May 1;251:118991.

39. Homae F, Watanabe H, Nakano T, Taga G. Large-Scale Brain Networks Underlying Language Acquisition in Early Infancy. Front Psychol [Internet]. 2011;2. Available from: https://www.frontiersin.org/articles/10.3389/fpsyg.2011.00093

40. Zoefel B, Oever ST, Sack AT. The Involvement of Endogenous Neural Oscillations in the Processing of Rhythmic Input: More Than a Regular Repetition of Evoked Neural Responses. 2018 Jun 6 [cited 2020 Jul 6]; Available from: https://www.repository.cam.ac.uk/handle/1810/276669

41. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. J Neurosci Methods. 2004 Mar
15;134(1):9–21.

42. Kothe CA, Makeig S. BCILAB: a platform for brain–computer interface development. J
Neural Eng. 2013 Aug 28;10(5):056014.

43. Crosse MJ, Di Liberto GM, Bednar A, Lalor EC. The Multivariate Temporal Response
Function (mTRF) Toolbox: A MATLAB Toolbox for Relating Neural Signals to Continuous
Stimuli. Front Hum Neurosci [Internet]. 2016 [cited 2020 Jul 3];10. Available from:
https://www.frontiersin.org/articles/10.3389/fnhum.2016.00604/full

44. Pasley BN, David SV, Mesgarani N, Flinker A, Shamma SA, Crone NE, et al.
Reconstructing Speech from Human Auditory Cortex. PLOS Biol. 2012 Jan
31;10(1):e1001251.

45. Özkurt TE, Schnitzler A. A critical note on the definition of phase–amplitude cross-
frequency coupling. J Neurosci Methods. 2011 Oct 15;201(2):438–43.

46. Alcock K, Meints K, Rowland CF, Brelsford V, Christopher A, Just J. The UK
Communicative Development Inventories: Words and Gestures - Manual and Norms.
Emsworth, UK: J & R Press; 2020.

47. Liszkowski U, Carpenter M, Tomasello M. Reference and attitude in infant pointing. J Child
Lang. 2007/01/25 ed. 2007;34(1):1–20.

48. Friend M, Keplinger M. An Infant-Based Assessment of Early Lexicon Acquisition. Behav
Res Methods Instrum Comput J Psychon Soc Inc. 2003 Jun 1;35:302–9.

49. Hoff E, Core C, Bridges K. Non-word repetition assesses phonological memory and is
related to vocabulary development in 20- to 24-month-olds. J Child Lang. 2008/10/06 ed.
2008;35(4):903–16.

50. Ní Choisdealbha Á, Attaheri A, Rocha S, Mead N, Olawole-Scott H, Brusini P, et al. Neural
phase angle from two months when tracking speech and non-speech rhythm linked to
language performance from 12 to 24 months. Brain Lang. 2023 Aug 1;243:105301.

51. Ní Choisdealbha Á, Attaheri A, Rocha S, Mead N, Olawole-Scott H, Alfaro e Oliveira M, et
al. Cortical tracking of visual rhythmic speech by 5- and 8-month-old infants: Individual
differences in phase angle relate to language outcomes up to 2 years. Dev Sci. 2024 Mar
14;n/a(n/a):e13502.

52. Marshall PJ, Bar-Haim Y, Fox NA. Development of the EEG from 5 months to 4 years of
age. Clin Neurophysiol. 2002;113(8):1199–208.

53. Orekhova EV, Stroganova TA, Posikera IN, Elam M. EEG theta rhythm in infants and
preschool children. Clin Neurophysiol Off J Int Fed Clin Neurophysiol. 2006
May;117(5):1047–62.

54. Stroganova TA, V. Orekhova E, Posikera IN. Externally and internally controlled attention
in infants: an EEG study. Int J Psychophysiol. 1998 Nov 1;30(3):339–51.

55. Cuevas K, Bell MA. 293C13EEG Frequency Development Across Infancy and Childhood.
In: Gable PA, Miller MW, Bernat EM, editors. The Oxford Handbook of EEG Frequency
[Internet]. Oxford University Press; 2022 [cited 2023 Feb 26]. p. 0. Available from:
https://doi.org/10.1093/oxfordhb/9780192898340.013.13

56. Leong V, Kalashnikova M, Burnham D, Goswami U. The Temporal Modulation Structure of
Infant-Directed Speech. 2017 Mar 27 [cited 2020 Jul 3]; Available from:
https://www.repository.cam.ac.uk/handle/1810/263724

57. Molinaro N, Lizarazu M, Lallier M, Bourguignon M, Carreiras M. Out-of-synchrony speech
entrainment in developmental dyslexia. Hum Brain Mapp. 2016;37(8):2767–83.

58. Dauer RM. Stress-timing and syllable-timing reanalyzed. J Phon. 1983 Jan 1;11(1):51–62.

59. Leong V, Goswami U. Acoustic-Emergent Phonology in the Amplitude Envelope of Child-
Directed Speech. PLOS ONE. 2015 Dec 7;10(12):e0144411.

60. Ghinst MV, Bourguignon M, Niesen M, Wens V, Hassid S, Choufani G, et al. Cortical
Tracking of Speech-in-Noise Develops from Childhood to Adulthood. J Neurosci. 2019 Apr
10;39(15):2938–50.

61. Ding N, Melloni L, Zhang H, Tian X, Poeppel D. Cortical tracking of hierarchical linguistic
structures in connected speech. Nat Neurosci. 2016 Jan;19(1):158–64.

62. Kösem A, van Wassenhove V. Distinct contributions of low- and high-frequency neural
oscillations to speech comprehension. Lang Cogn Neurosci. 2017 May 28;32(5):536–44.

63. Boucher VJ, Gilbert AC, Jemel B. The role of low-frequency neural oscillations in speech
processing: Revisiting delta entrainment. J Cogn Neurosci. 2019;31(8):1205–15.

64. Doelling KB, Arnal LH, Ghitza O, Poeppel D. Acoustic landmarks drive delta-theta
oscillations to enable speech comprehension by facilitating perceptual parsing. NeuroImage.
2014 Jan 15;85 Pt 2:761–8.

65. Vanthornhout J, Decruy L, Wouters J, Simon JZ, Francart T. Speech Intelligibility Predicted
from Neural Entrainment of the Speech Envelope. J Assoc Res Otolaryngol. 2018 Apr
1;19(2):181–91.

66. Fries P. The Model- and the Data-Gamma. Neuron. 2009 Dec 10;64(5):601–2.

67. Schroeder CE, Lakatos P. Low-frequency neuronal oscillations as instruments of sensory
selection. Trends Neurosci. 2009 Jan 1;32(1):9–18.

68. Poeppel D, Assaneo MF. Speech rhythms and their neural foundations. Nat Rev Neurosci.
2020 Jun 1;21(6):322–34.

69. Doelling KB, Arnal LH, Assaneo MF. Adaptive oscillators provide a hard-coded Bayesian
mechanism for rhythmic inference. bioRxiv. 2022 Jan 1;2022.06.18.496664.

1086