Antiviral action of aqueous extracts of propolis from *Scaptotrigona aff. postica* against Zica, Chikungunya, and Mayaro virus

Mendonça RZ[1]*, Nascimento RM[1], Fernandes ACO[1], da Silva Junior PI[2]

1) Laboratory of Parasitology, Butantan Institute;
2) Special Laboratory for Applied Toxinology (LETA), Center of Toxins, Immune-response and cell signaling (CeTICS/CEPID), Butantan Institute,

*Corresponding author. Laboratório de Parasitologia, Instituto Butantan, Av. Vital Brasil 1500, CEP 05503-900, São Paulo, Brasil. Tel.: phone 55 1126279420, fax: +55 011 3726 1505.
E-mail address: **zucatelli@uol.com.br**

ABSTRACT

The limited availability of antivirals for new highly pathogenic strains of virus has become a serious public health problem that kills thousands of people annually. For this reason, the search for new products against these agents has become an urgent necessity. Many studies have been carried out with this aim. Among the multiple sources of research for new antibiotics and antivirals, bioprospecting from insect exudates or their products has become an increasingly frequent option. Insects appeared on the planet about 350 million years ago and have been one of the beings with greater adaptability and resistance to the most varied biomes. Insects have been found in all known ecosystems. Their survival for so long, in such different environments, is an indication that they have a very efficient protection system against environmental infections, despite not having a developed immune system like mammals. Historically, since the ancient civilizations of Egypt and Rome, the products obtained from the bee, such as honey and propolis, have been of great pharmacological importance, being used as antimicrobial, anti-inflammatory, antitumor, healing several other functions. Investigations on the chemical composition and biological activity of propolis have been carried out, mainly in the species Apis mellifera, and this product has shown activity against some important viruses, such as influenza, poliovirus, hepatitis, HIV, and others. However, for the Meliponini species, known as stingless bees, there are few studies, either on their chemical composition or on their biological activities. The importance of studying these bees is because they come from regions with native forests, and therefore with many species of plants not yet studied, in addition to which they are regions still free of pesticides, which guarantees a greater fidelity of the obtained data. Previous studies by our group with crude hydroalcoholic extract of propolis demonstrated an intense antiviral activity against Herpes, influenza, and rubella viruses. All studies carried out with propolis are done with alcoholic extracts. In this work, we chose to use aqueous extracts, which eliminates the presence of other compounds besides those originally present in propolis, in addition to extracting substances different from those obtained in alcoholic extracts, which makes this work unprecedented. Therefore, this study aimed to identify, isolate and characterize compounds with antiviral effects from aqueous propolis extracts from Scaptotrigona aff postica, in emerging viruses such as zicavirus, chikungunya, and Mayaro. The evaluation of the antiviral activity of the crude and purified material was performed by reducing infectious foci in cultures of infected cells treated with propolis extracts in infected cultures and treated or not treated with propolis. The chemical characterization of the elements present in the extracts was performed by liquid chromatography. The results obtained indicate a high reduction of Zicavirus, chikungunya, and Mayaro virus replication when compared to infected-only cultures. Likewise, it was observed that the antiviral response was dose-dependent, being more intense when propolis was added 2 hours after the viral infection.

Keywords – Scaptotrigona aff postica, propolis, antiviral activity, Zicavirus, Chikungunya, Mayaro

1. INTRODUCTION

Arboviruses are viruses transmitted by the bite of arthropods, such as Aedes aegypti. Aedes aegypti is a vector of great importance for public health in the tropics and subtropics and practically the entire American continent, as well as in Southeast Asia and throughout India. More than 200 arbovirus species of arbovirus have been isolated in Brazil, being around them related to diseases in humans (Rosa et al., 1998). Arboviruses are emergent viruses by nature. None of them are originally humans disease. They are originally zoonoses, diseases of animals that only occasionally affect humans, and only become important when some significant ecological modification alters their natural habitat and leads to changes in reservoirs, vectors, and even virulence (Casseb et al., 2013). The process of economic development, population growth and urbanization verified in Brazil from the second half of the 20th century placed arboviruses among the viruses of great impact, real or potential, on public health. In some situations, the adaptation to humans, of the virus or the arthropod vector, is sufficient to make the animal reservoir unnecessary for the maintenance of the virus cycle in nature. The main arthropod vectors are mosquitoes, flies, and ticks. The most common vertebrate hosts are rodents, but it also occurs in small mammals, primates, birds, and ungulates (Lopes et al., 2014). In Brazil, the best-known and most widely circulated arboviruses with epidemiological importance today are Dengue, Zika, and Chikungunya.

Among the known arboviruses, Zika has left the world on alert since the recent outbreaks associated with it, especially in Brazil between 2013 and 2014, mainly in the Northeast region, with a high number of cases of microcephaly in newborns and other complications, as is the case of Guillain-Barré syndrome, which had not been reported in the same epidemiological proportion before. Although the Zika virus has been identified in humans in Africa in 1952 (Dick et al., 1952), the first major outbreak occurred only in 2007, in Micronesia (Duffy et al., 2009). 2013 and 2014, outbreaks occurred in French Polynesia in 2013 (Cao-Lormeau et al., 2013 and New Caledonia in 2014 (Dupont-Rouzeyrol et al 2015). A few times after this, still in 2014, reported cases of an illness characterized by rash and fever in northeastern Brazil. In May 2015, local transmission of the Zika virus is confirmed. In 2016, the Brazilian government reported 214,193 probable cases and 128,266 confirmed cases of Zika virus (SVS, 2017) although many cases were probably not reported. In this time is estimated that there were between 500,000 and 1.5 million cases from 2015 to early 2016 (Heukelbach et al., 2016). As of May 2017, 85 countries and territories have reported documented cases of Zika virus transmission.

Recently, the Mayaro virus has become a potential public health risk for which there is no specific treatment. Less common in humans than Zica and Chikungunya, the Mayaro virus also causes outbreaks of febrile illness in the North and Midwest regions of Brazil, as well as in other South American countries (Bolivia, Venezuela, and Peru) (Alves et al, 2017). It belongs to the genus Alphavirus, family Togaviridae, along with eastern equine encephalitis, venezuelan equine encephalitis, and Chikungunya viruses. Illness caused by the Mayaro virus is usually benign but can be temporarily disabling, often causing rash and arthralgia or even arthritis, myalgia, and fever lasting three to five days. During convalescence, arthralgia often persists for a few weeks. The sylvatic cycle of the Mayaro virus is similar to that of sylvatic yellow fever, a disease of monkeys, transmitted by mosquitoes of the genus Haemagogus. In addition to monkeys, the Mayaro virus can also be found in birds. All of them have a short life in the body, causing acute infections. In 2016, there was an explosion of cases of Chikungunya (63.810 confirmed cases) (Silva et al., 2018). There is a potential risk of the Mayaro virus becoming a virus

with a greater public health impact than the previous two. The co-circulation of arboviruses in Brazil has made clinical diagnosis difficult, increasing cases of complications such as neurological fever syndromes, hemorrhages, and autoimmune diseases. The association of prevention and control of vectors in the development of vaccines to prevent the population against these diseases, can help the control and reduction of cases of arboviruses in Brazil and the world. Another important aspect is the development of drugs that can inhibit the replication of the virus in the body and thus reduce the severity of clinical conditions.

However, despite the great importance in public health, few drugs are available for the treatment of these viruses after they are installed in the organism. Most treatments are only to overcome the effects of the virus and not to fight it. Seeking to combat these infections, the population has used natural products obtained from plants with antiviral actions for millennia. One of the products frequently used is propolis, a bee product that has a very wide range of pharmacological actions. Among the best-known activities of propolis, its antiviral action is one of the most outstanding, as it demonstrates better efficacy against viruses than standard drugs available on the market (Cornara, et al., 2017). An example of this is the comparative study between an ointment prepared with Canadian propolis, rich in flavonoids, and acyclovir ointment (a drug indicated for the treatment of Herpes simplex, Varicella zoster, Epstein-Barr and Cytomegalovirus viruses). In this study, the propolis-based ointment was more effective than acyclovir in healing genital herpes lesions and in reducing local symptoms in the treatment of genital herpes simplex virus (Vynograd, et al., 2000). An in vitro study with Canadian propolis proved to be efficient against Herpes simplex, types 1 and 2 during the viral adsorption phase or when incubated directly with the virus. These results indicate that propolis can directly interfere with the virus, preventing the penetration of the virus into cells (Bankova et al., 2014). Propolis harvested in the city of Moravia, the Czech Republic, when tested against HVS-1, showed a protective effect when the virus was previously incubated with the extracts (Schnitzler, et al., 2010). The composition of the word propolis comes from the Greek, pro means "in defense", and polis means "city", that is, "in defense of the city". Propolis is a resinous material that bees collect from plants, mix with wax, and use to build their nests. Bees apply propolis in thin layers to the inner walls to strengthen and block holes and crevices, as well as use it as an "embalming" substance on dead invaders or even dead bees, thus inhibiting the growth of fungi and bacteria inside the hive. (Bankova et al., 2000; Ghisalberti, 1979). The medicinal properties of propolis have been reported since ancient times. In Egypt it was used in the mummification process to prevent bodies from rotting; the Greeks and Romans used it as an antiseptic and healing agent, and the Incas as an antipyretic (Ghisalberti, 1979, Sforcin Bankova, 2011) Due to its great pharmacological potential, propolis became the target of several researches, arousing the interest of the scientific society that has sought to investigate its compounds and biological activities. This interest has intensified in recent decades (Bankova et al., 2000; Araújo et al., 2010). Countries such as Switzerland and Germany already legally recognize propolis as a medicine (Bogdanov, 2014). In Brazil, the main substances described are those derived from prenylates of p-coumaric acid and acetophenone, such as diterpenes and lignans (Bogdanov, 2014). Studies with propolis have already demonstrated biological activities, such as immunomodulation, antitumor, antioxidant, antibacterial, antiviral, and analgesic (Búfalo et al., 2009; Dobrowolski et al., 1991; Paulino et al., 2003).

One of the components of hives is resins. Resins are pre-formed substances produced by secondary metabolites of plants, which have specific functions, such as pigmentation of flowers to attract pollinating insects, repellent action, and defense mechanisms against pathogenic microorganisms (Garcia and Carril, 2009). In propolis, the resin is used in prophylactic protection and combat against microorganisms. Studies show that epidemics in nests decrease both productivity (Evans and Pettis, 2005) and the number of bees (Moret and Schmidt, 2000). Therefore, the chemical compounds present in these products are essential for the protection and survival of bees, being considered an evolutionary mechanism for the benefit of the colony (Simone-Finstrom et al, 2017). The composition of propolis is extremely variable, both in terms of geographic location and season (Negri et al, 2022). In Europe, North America, and temperate zones, flavonoid compounds, phenolic acids, and esters predominate in propolis (Falcao et al, 2010). In the Mediterranean terpenoids have been identified in high concentrations (Popova, et al., 2010; Popova, et al., 2011). In Africa, the main compounds are triterpenoids (Zabaiou, et al., 2017).

In Brazil, due to the great biodiversity, several types of propolis have been reported, such as red, green, and brown. The vast majority of studies are carried out with propolis from bees of commercial interest, mainly Apis mellifera. The propolis of the species Apis mellifera was classified into 12 distinct groups, based on the physicochemical characteristics and place of origin. Among the main compounds isolated are coumaric acid, ferulic acid, pinobanksin, kaempfero, apigenin, and caffeic acid (Park, et al., 2002). The components of Brazilian red and green propolis are commonly extracted from Dalbergia ecastaphyllum, Baccharis spp, Betula pendula, Betula verrucosa, and Betula pubescens species and are composed of phenolic acids, phenylpropanoids, diterpenic acids p-coumaric acids, (Zabaiou, et al., 2017). Studies on the glandular secretions of bees have demonstrated antibacterial activity, as is the case with the hypopharyngeal secretion of the species Apis mellifera (White, et al., 1963). The same hypopharyngeal glands were found in the species *Scaptotrigona postica*, presuming they have the same functions (Costa and Cruz-Landim, 1999). The vast majority of studies on the pharmacological effects of propolis are related to propolis obtained from Apis mellifera bees (Sforcin and Bankova, 2011). The chemical composition of propolis is influenced not only by the vegetation of the environment but also by the seasons and the bees that produce it (Bankova et al., 2000). These variations in chemical components affect their biological properties (Nakamura et al., 2010). For this reason, it is so important to study the propolis of different bee species and regions. Among the different species of bees, studies dealing with propolis from stingless bees are rare (Bankova et al., 1998). In Brazil, one of the species of stingless bees belongs to the Apidae family, of the Meliponinae subfamily. These bees mix plant resin with wax and clay, thus forming a geopropolis. This denomination is precisely due to the use of land for the final composition of the material produced (Bankova et al., 1998; Nogueira-neto, 1997). Recently, compounds such as phenolic acids and water-soluble tannins were isolated from the geopropolis of Melipona fasciculata, (gallotannins and ellagitannins). These compounds, from the aqueous and alcoholic fractions of geopropolis from Melipona scutellaris (Franchin et al., 2012, Franchin et al., 2013), demonstrated antioxidant activity (Dutra et al., 2014), anti-inflammatory and antinociceptive activities. Within the Meliponinae subfamily, there is the genus Scaptotrigna, which is distributed throughout the Neotropical region (Silveira et al., 2002). This genus, is found the Scaptotrigona aff postica, popularly known in Maranhão as the "tubi" bee. However, despite being in the family of geopropolis producers, they are an exception to the rule, since they do not use land in the composition of propolis (Araújo et al., 2010).

There are few studies with propolis from Scaptotrigona aff postica, and there is no consensus in the literature on how to refer to it, some groups use the name geopropolis (Coelho et al., 2015), while others use propolis (Farias, et al. 2014; Araújo et al., 2010, 2011). For this work, we will use the term propolis, as we understand it to be the most appropriate. This propolis has been used by the population of the Maranhão region in the treatment of cancer and wound healing (Araújo et al., 2010). Earlier, we have shown that propolis from Scaptotrigona aff postica presents a potent antiviral action against Herpes Simplex Virus (HSV-1) (Coelho et al., 2015) as well against Rubella virus (Coelho et al., 2018). Scientific studies have corroborated the possible medicinal effects of this propolis. Recently, we have shown (Carmona et al., 2020) that propolis and its components can help reduce the physiopathological consequences of covid-19 infection, and that, pyrrolizidine alkaloids and flavonoids glycosides from propolis of Scaptotrigona aff. postica has a potent antimicrobial action (Cantero et al., 2022). Martin et al., 2013, treated rats with corneal injuries caused by burns with an emulsion of the crude extract and observed both healing and anti-inflammatory effects. Antitumor activity against Ehrlich tumor and reduction in pathology associated with asthma due to an inhibition of migration of inflammatory cells into the alveolar space are also described. In addition to those mentioned, there are no reports on other actions of this product. Therefore, due to the importance of these arboviruses in public health, the limited availability of effective drugs on the market, and the abundant information on the antiviral activity of propolis from several bee species, this work aims to study the antiviral activities of Scaptotrigona. aff postica propolis against arboviruses.

2. Material and methods

2.1. Própolis

The propolis used in this work was obtained from a colony of Scaptotrigona aff postica, located in the State of Maranhão, in the Barra do Corda region, Brazil. The city is located in the geographic center of Maranhão, at the confluence of the Rio da Corda and Rio Mearim.

2.2 Preparation of propolis extract by aqueous extraction

In this work, the aqueous extraction of propolis was used as a source of substances for studies of antiviral action. Aqueous extraction of propolis was carried out as described below. The propolis, obtained by scraping the meliponiculture box, was gathered, forming a "paste". This material was transported to the laboratory and frozen at -20ºC for 24 hours to make the material rigid, forming a frozen propolis "stone" to facilitate the spraying process. This "stone" of propolis was then manually macerated until it became a very fine granular material. This crushed material was passed through stainless granulometric sieves, fine mesh of 0.15 mm. The sieved material was ground again by one degree until the material above became a very fine powder which was passed again through ABNT granulometric (75 to 100 mm Astn) sieves with a porosity of 1 to 5 microns. After this procedure, ultrapure water was added to the powder (100 ml of water for every 10 grams of propolis powders) and homogenized under rotation at 1,000 rpm for 24 hours at room temperature, protected from light. To separate the precipitate (wax) from the liquid medium (extract), the supernatant was centrifuged at 4°C for 30 minutes at 15,000 rpm (Eppendorf® centrifuge 5804R). The supernatant was then filtered through a 0.22μm millipore membrane, aliquoted, and stored in a sterile glass refrigerator until use. This material was considered to be 100%.

2.3 . Cell line, culture medium, and cell culture

VERO cells (African green monkey kidney fibroblast cells - Cercopithecus aethiops) were used in this study to determine the antiviral action of propolis. These cells were cultured at 37°C in T-flasks or 96-well microplates containing Leibovitz medium (L-15), supplemented with 10% fetal bovine serum. Cells were maintained at 37°C in a cell culture oven. Cell growth and morphology were monitored daily using an inverted optical microscope Olympus CK2 at 200x magnification.

2.4 . Cytotoxicity assay of samples in cells

The cytotoxicity test of all samples and the purified fractions of propolis was performed in VERO cells to determine the maximum concentration of propolis or its fractions that is not toxic. For this, cells were seeded at a concentration of 10^4 cells/well in 96-well plates and cultured at 37°C for 24 hours. After this period, the cells were treated with different concentrations of the samples to be tested. PBS was used as a negative control and DMSO 10% as a positive control. After 24, 48, and 72 hours of treatment, cell viability was determined by visualizing cell morphology and treating the cells with a vital dye, trypan blue. In each experiment, an aliquot of the samples, at the concentration to be used in the tests, was added to a plate excavation containing VERO cells, as a way of verifying the non-toxicity of the sample, serving as a basis for comparing the morphology of the cells in the test excavations.

2.5 Vírus and infection

2.5.1 Virus

The antiviral action of Scaptotrigona aff postica propolis was tested in 3 different arbovirus strains of interest in public health; (Zicavirus, Chikungunya, and Mayaro). These viral strains are part of the collection of the Parasitology laboratory at Instituto Butantan

2.5.2 Viral titulation

Initial viral stocks were titrated at the beginning of the work by the viral extinction technique as described by (Reed and Muench., 1938) and detailed in the following item. The viral titer was expressed as TCID50/mL (50% Tissue culture infectious dose), which is the highest dilution of the virus capable of infecting 50% of the cells. To determine the viral titer, aliquots of the virus samples under study were placed on a semi-confluency mat of VERO cells (2×10^5 cells/mL), grown in 96 microplates. The viruses were added to the cell cultures in serial dilutions 2. The microplates were incubated at 37°C and the viral cytopathic effect was observed daily and the titer of infectious viral particles was calculated as described by Reed and Muench. For this study, an aliquot of the virus was thawed immediately before experimenting.

2.5.3 Test of antiviral activity

2.5.3.1. Test of antiviral activity in cell culture by the technique of determination of the minimum infective dose (TCID/50)

In the first phase of tests, VERO cells, an exponential growth phase, grown in a 96-well plate, were treated with crude samples of the test substances, 1 hour before infection. After this treatment period, the cells were infected with the different viral strains in serial dilutions ratio 2, with the first dilution varying from 100 to 500 TCI/50 of virus. The cultures were kept at 37°C and observed daily under an optical microscope, seeking to determine the inhibitory effect of the samples on virus replication by determining the cytopathic effect. After the appearance of the cytopathic effect, the culture supernatant was removed and the cells were stained with crystal violet to reveal the cytopathic effect. The antiviral effect was determined by the difference in the highest dilution of the virus capable of inducing a cytopathic effect observed in the infected-only culture and the infected but treated culture. Negative controls for infection and cytotoxicity of the samples were performed in all experiments

2.5.3.2 Dose/effect determination

To define the best amount of fraction to be added to infected cultures, fractions showing antiviral activity were added 1 hour before infection at concentrations of 2, 5, or 10% (v/v). The cultures were kept at 37°C and observed daily under an optical microscope, seeking to determine the inhibitory effect of the samples on virus replication through the determination of the cytopathic effect and the determination of the effect performed as described above.

2.5.3.3 Determination of the optimal time of infection (TOI)

To define the best time of infection (TOI), the fractions that showed antiviral activity were tested for the best time to add these fractions to the infected cultures. For this, samples of the fractions, at the optimal concentration described above, were added 2 hours before infection, and 2 hours after infection, or the cultures were treated with a mixture of fraction + virus and kept in contact for 2 hours. before being added to crops. The cultures were kept at 37ºC and observed daily under an optical microscope, seeking to determine the inhibitory effect of the samples on virus replication through the determination of the cytopathic effect and the determination of the effect performed as described above. Six days after the start of the experiment, the supernatant was removed, 0.25% Crystal Violet was added and washed with PBS

2.6. Fractionation of samples by Reverse-Phase HPLC DAD-ESI-MS/MS and antiviral activity test of purified fractions

2.6.1 Aqueous Propolis Extract HPLC Purification

Scapotrigona aff, postica propolis (1,2g) was suspended in 180 mL of ultrapure water and homogenized for 2 hours at 4ºC. The supernatant (20 mL) was fractionated by a reverse-phase high-performance liquid chromatography (RP-HPLC) at room temperature on a Shimadzu LC-10 HPLC system (using a semi-preparative C18 Jupiter column (30 mm; 300A; 10μ 250 mm - Phenomenex International, Torrance, CA, USA) The colunm was equilibrated at room temperature with 0.1% TFA in ultrapure water. The elution was done under a linear ACN gradient and the flow rate was 8 mL/min during 60 min (0-80% of B). Each eluted fraction (under absorbance monitored at 225 nm) was collected and   concentrated in a vacuum centrifuge SpeedVac Savant (Thermo Fisher Scientific, Waltham, MA, USA). The fractions obtanined were stored in the freezer at - 20 °C until its usage. The concentrated fractions were reconstituted in 1 mL ultrapure water and used in antiviral activity assays. The purified fractions were tested for their antiviral capacity in the same way as performed with the crude propolis sample

2.6.2 Characterization of propolis fractions with activity by mass spectrometry

The peptides with activity were resuspended in 20 μL in ACN/H2O 1:1 with 0.1% formic acid and analyzed by direct infusion in a Thermo Scientific LTQ-Orbitrap LC/MS (LTQ – Linear Trap Quadrupole), under 0.5 μL/min in a 50 μL syringe (Hamilton, Reno, NV, USA). Fragmentation of the mono or doubly-charged ions was performed under different collision energies (Ecol) against inert gas (argon) so that the best MS/MS spectra can be used in the sequencing. DIC spectra against inert gas molecules (Ar) were acquired every 0.5 second. The final composition of spectra was obtained after 1.5 min of acquisition

2.7 Protein dosage

The concentration of the fractions was determined by the Lambert-Beer law, (Grimsley and Pace, 2004). The active fractions concentration was determined at 205 nm using a NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific).

3 RESULTS

3.1. Determination of the cytotoxicity of the agents to be tested

At the beginning of the work, the propolis cytotoxicity of Scaptotrigona aff. postica was determined in VERO cells. The concentrations used were the same used in the tests. For this, VERO cells were grown in 96-well microplates. When the cell reached confluently, 1, 2, 5, or 10% v/v propolis. The cultures were observed daily and for 72 hours to determine the morphological alteration in the cultures. None of the concentrations used showed deleterious effect on the growth of Vero cells. **(Figure 01)** As a negative control of cytotoxicity, in all the tests performed, an aliquot of the propolis to be tested was always added to the normal VERO cells, at the same concentration as the test. The results obtained as control of propolis cytotoxicity, performed in all experiments, were similar to the results presented below.

FIGURE 1

Figure 1: Cell viability of vero cells treated with different concentrations propolis. The culture was treated com 1, 2, 5 or 10% v/v of propolis. The culture was maintained at 37 ºC for 72 hours and the culture was observed daily to determine the cytotoxic effect. The results above represents the morphology observed in cells in all experiments

3.2 Chromatography of the crude extract of propolis and identification of fractions with viral activity

To identify the molecules present in the sample with antiviral action, phytochemical analysis of the propolis extract was performed using a reversed-phase HPLC- Thermo Scientific LTQ-Orbitrap LC/MS (LTQ – Linear Trap Quadrupole). As can be observed in **Figure 02,** various fractions were obtained. These fractions were then tested against viruses to determine the chromatographic fraction with antiviral activity. The purified fractions were tested for their antiviral capacity in the same way as performed with the crude propolis sample. Of all the fractions tested, the one that showed antiviral activity was fraction 1. This fraction showed a size of 370DA and a concentration of 107μg/ml**.** The results are presented below.

Figure 2: Phytochemical analysis of the propolis extract. The propolis samples were resuspended in 20 μL in ACN/H2O 1:1 with 0.1% formic acid and analyzed by direct infusion in a Thermo Scientific LTQ-Orbitrap LC/MS (LTQ – Linear Trap Quadrupole), under 0.5 μL/min in a 50 μL syringe (Hamilton, Reno, NV, USA). Fragmentation of the mono or doubly-charged ions was performed under different collision energies (Ecol) against inert gas (argon) so that the best MS/MS spectra can be used in the sequencing. DIC spectra against inert gas molecules (Ar) were acquired every 0.5 seconds. The final composition of spectra will be obtained after 1.5 min of acquisition. The fractions obtained were tested by their antiviral activity

3.3 Determination of antiviral action

3.3.1. Determination of the antiviral action of propolis on different arboviruses

In this work, the antiviral action of propolis was tested against Zica virus, Chikungunya, and Mayaro virus. For this experiment, plates from 96 wells with VERO cells were treated with propolis being maintained in contact with the cells for 1 hour. After this period, serial dilutions (ratio 2) of the virus, with an initial titer of 100 to 500 TCID/50, were added to the treated cultures and observed for 72 hours to determine the cytopathic effect.

3.3.1.1. Antiviral action of propolis and its purified fraction against Zica virus

The test was performed as described above. As can be seen in **Figure 3**, there is an intense reduction in the viral effect of 256 TCID/50 of Zicavirus after the treatment of cells with 10% v/v propolis. Under these conditions, the viral reduction ranged from 256 to 4 TCID/50, or 64 fold (98,4%), with crude propolis. With the treatment of cells with purified fraction 1, the antiviral action was 16-fold (256 to 16 TCID/50).

Figure 3: Antiviral action of propolis from Scaptotrigona aff. postica (10% v/v) against the Zica virus. The propolis extract or the purified fractions were added to the cultures of VERO cells and 1 hour later, serial dilutions (r:2) of Zica virus (initial titer of 256 TCID 50/0.1ml) were added to the culture. The culture was maintained at 37 °C for 72 hours and the culture was observed daily to determine the cytopathic effect. The final titer was determined by the highest dilution of the virus to cause a cytopathic effect in the culture. The result represents 4 individual experiments (performed in duplicate each).

In the photos in **Figure 4**, can be seen that there was a clear reduction in the cytopathic effect caused by the Zicavirus 72 hours after infection, in cultures treated with propolis (10% v/v)

Figure 04. Photomicrography of normal VERO cell cultures (A), infected with Zica virus (B) and infected and treated with 10% v/v propolis from Scaptotrigona aff. postica (C). The cultures were kept for 3 days at 37ºC. Cells were observed daily under an inverted microscope at 10x magnification. After this period, the culture supernatant was removed and the cells stained with crystal violet (0,25%)

3.2.1.2 Antiviral action of propolis against Mayaro virus

The antiviral action of the aqueous extract of propolis was also tested against the Mayaro virus. The experimental procedures were the same as mentioned above but using 512 TCID/50 of virus and 2 % propolis. Under these conditions, the viral reduction was similar to that obtained in the tests with the Zica virus, using crude or purified propolis fraction. In this case, the reduction was on average 128-fold for crude propolis (512 to 4 TCID50) and 32-fold for the fraction (512 to 16 TCID/50) **(Figure 5).**
The result represents 4 individual experiments (performed in duplicate each).

Figure 5: Antiviral action of propolis from Scaptotrigona aff. postica against the Mayaro virus. Aqueous propolis extract (2% v/v) was added to the VERO cell culture. One hour after this procedure, serial dilutions (r:2) of the Mayaro virus (512 TCID 50/0.1ml) were added to the culture. The culture was maintained at 37 ºC for 72 hours and the culture was observed daily to determine the cytopathic effect. The final titer was determined by the highest dilution of the virus to cause a cytopathic effect in the culture.

In the photos in **Figure 6**, can be seen that there was a clear reduction in the cytopathic effect caused by the Mayaro virus 72 hours after infection, in cultures treated with propolis (2 % v/v)

Figure 06. Photomicrography of normal VERO cell cultures (A), infected cells with Mayaro virus (B), and infected cells treated with 2 % v/v propolis from Scaptotrigona aff. Post (C)
The cultures were kept for 3 days at 37ºC. Cells were observed daily under an inverted microscope at 10x magnification. After this period, the culture supernatant was removed and the cells were stained with crystal violet 025%.

3.2.1.3. Antiviral action of propolis on Chikungunya virus

The antiviral action of the aqueous extract of propolis was tested against Chikungunya. The experimental procedures were the same as mentioned above, only with a greater amount of virus (1000 TCID/50%) and 5% propolis. As can be seen in **Figure 7**, a 256-fold reduction in viral replication was observed when crude propolis was added or a 512-fold reduction when the purified fraction was used.

Figure 7: Antiviral activity of the aqueous extract of propolis from Scaptotrigona aff. postica (5 % v/v) against the Chikungunya virus. The propolis extract or the purified fractions were added to the VERO cell cultures. 1 hour later, serial dilutions (r:2) of the Chikungunya virus (initial titer of 1000 TCID 50/0.1ml) were added to the culture. The culture was maintained at 37 ℃ for 72 hours and the culture was observed daily to determine the cytopathic effect. The final titer was determined by the highest dilution of the virus to cause a cytopathic effect in the culture. The data in the figure is representative of 4 experiments (performed in duplicate each)

In the photos in **Figure 8**, it can be seen that there was a clear reduction in the cytopathic effect caused by 5% (v/v) of propolis or its purified fraction against the Chikungunya virus (72 hours after infection).

Figure 08. Photomicrography of normal VERO cell cultures (A), infected with Chikungunya virus (B), and infected cells treated with 5% v/v propolis from Scaptotrigona aff. postica (C) or with infected and treated with 5% of purified fraction of propolis (D). The cultures were kept for 3 days at 37℃. Cells were observed daily under an inverted microscope at 10x magnification (A, B, and C) and 20x in photo D. After this period, the culture supernatant was removed and the cells stained with crystal violet 025%.

3.2.2. Effect of the concentration of propolis aqueous extract on viral replication

The action of concentration of propolis aqueous extract on antiviral activity was tested with Zicavirus or with Mayaro. For this, the VERO cell cultures were treated with 2, 5, or 10% propolis and after 1 hour the cells were infected with 128 or 512 TCID/50% of Zica virus or Mayaro virus. The cultures were observed for 72 hours and the viral titer was defined as the highest dilution of the virus capable of inducing a cytopathic effect in the cells. As can be seen in **Figure 9,** the antiviral action of propolis is dose-dependent, and the viral reduction with the concentration of 10% of propolis extract was 64-fold to Zicavirus and 128-fold to the Mayaro virus (99,6% of reduction).

Figure 9: Antiviral action of the concentration of the aqueous propolis extract of Scaptotrigona aff. postica against Mayaro virus (A) or Zicavirus (B). Aqueous propolis extract (2, 5, or 10% v/v) was added to the VERO cell culture. 1 hour after this procedure, serial dilutions (r:2) of Zicavirus or Mayaro virus (initial titer of 128-512 TCID 50/0.1ml) were added to the culture. The culture was maintained at 37 ℃ for 72 hours and the culture was observed daily to determine the cytopathic effect. The final titer was determined by the highest dilution of the virus to cause a cytopathic effect in the culture

3.2.3. Effect of time of addition of aqueous propolis extract on viral replication

The effect of the time of addition of the aqueous extract of propolis on the antiviral activity was tested against different dilutions of Zicavirus (from 0 to 2048 TCID/50/0.1 ml). For this, VERO cell cultures were treated with 5% propolis 2 hours before infection, 2 hours after infection, or treated with a mixture of virus + propolis kept together for 2 hours. The culture was observed for 6 days and the viral titer was defined as the highest dilution of the virus capable of inducing a cytopathic effect in the cells. As can be seen in **Figure 10,** the antiviral action of propolis was more effective when propolis was added 2 hours after infection. The experiments were performed in triplicate.

Figure 10: Action of propolis aqueous extract (TOI) addition time on antiviral activity against different Zicavirus dilutions (from 0 to 2048 TCID/50/0.1 ml). For this, VERO cell cultures were treated with 5% propolis 2 hours before infection, 2 hours after infection, or treated with a mixture of virus + propolis kept together for 2 hours. The culture was observed for 6 days and the viral titer was defined as the highest dilution of the virus capable of inducing a cytopathic effect in the cells.

DISCUSSION

Arboviruses are viruses transmitted by arthropods (Arthropod-borne virus) They are transmitted to humans and animals by the bite of blood-sucking arthropods (mosquitoes). Of the more than 545 known arbovirus species, about 150 cause disease in humans. (Lopes et al. 2014). Arboviruses have represented a major challenge to public health, due to environmental and climate changes and deforestation that favor amplification, and viral transmission, in addition to the transposition of the barrier between species. Among the arboviruses that cause diseases in humans and other warm-blooded animals are Dengue, Zica, Chikungunya, and Mayaro, among many others. These viruses are a major problem for which there is no specific treatment Treatment is usually focused only on relieving symptoms because there are no drugs that specifically attack the virus (Donalisio et al. 2017). In general, medications are indicated to control fever, headache, skin spots, and body pain, the same used in patients with dengue, zikavirus, or other arboviruses. That is why it is extremely important to develop a drug that acts directly on the virus and not just on the symptoms. Various studies have reported the antiviral activity obtained of products from invertebrates. Among the various animal sources of antiviral studies, propolis has gained great attention for obtaining new drugs (Bastos et al. 2011; Bankova et al. 2014), mainly because of its extensive and varied chemical composition (Bankova et al. 2005; Bankova et al., 2014, Negri et al, 2022) that generate huge amounts of molecules of high pharmacological value. Peter et al, 2017, describe the antiviral effect of three hydroalcoholic extracts of propolis (green, brown, and jataí bees (Tetragonisca angustula), against bovine viral diarrhea Virus (BVDV and bovine herpesvirus type-1 (BoHV-1). This propolis showed high efficacy in
treatments pre ad post aexposure against BoHV1. Extracts from Jataí showed activity against BoHV-1 asd well BVDV infection, when in the pre-test, meantime, brown propolis demonstrated action only against the BoHV-1 in the pre-infection assay. Is this study, green propolis led to a reduction in viral titer of 4.33log, while was observed a reduction of 3.5log to brown propolis and of the 3.24log to jataí propolis. Like this one, to date, most of the scientific work on the therapeutic activities of propolis has been carried out with the bee species Apis mellifera. However, some studies using propolis from stingless bees have been described, demonstrating its therapeutic potential against viruses of importance in human and veterinary health such as human herpes virus type 1 (Amoros et al. 1992, Coelho et al. 2015, Yildirim et al. 2016), Parvovírus Suíno (PPV) (Ma et al. 2015, Herpes Vírus Bovino (BoHV-5) (Fischer et al. 2007), influenza virus (Kujumgiev 1999) and vírus da imunodeficiência human (HIV) (Gekker et al. 2005).

One of the advantages of propolis as a therapeutic agent is, in general, the absence of toxicity to humans and animals (Vynograd et al. 2000). In the literature, studies have been observed in which the concentrations of propolis used in antiviral assays vary from 0.0003% to 5% (Amoros et al. 1992), with no morphological alterations being observed in the cells (Vynograd et al. 2000). However, Peter et al. 2017, observed cytotoxicity of 0.097µg/mL for green and brown propolis and 0.39µg/mL for jataí propolis. In a similar study, Bankova et al. (2014) evaluated the antiviral activity of a hydroalcoholic extract of Canadian propolis, where it was observed that the maximum tolerated non-cytotoxic concentration of propolis in MDBK cells was 0.032mg/mL. This relative toxicity may be related to the extraction method used, which was a hydroalcoholic extraction, which removes a good amount of bioactive substances present in propolis. Contrary to what is normally used, our experiments were carried out with aqueous extraction. With this procedure, substances that are not isolated in alcoholic extracts are

obtained, which leads to the discovery of new molecules with therapeutic potential, in addition to avoiding the presence of other substances that may be toxic in addition to those present in the material under study. In our study, with the aqueous extraction, we did not observe any toxic effect on VERO cells, when we used final concentrations of up to 10% of propolis, which allows for obtaining a higher concentration of bioactive products in the formulation.

There are several study about antiviral activity of propolis. However, the mechanisms of action of propolis on viruses are not yet fully elucidated, but some studies seek to identify where this activity occurs. Bankova et al. (2014) evaluated an alcoholic extract of propolis against Herpes simplex type 2 (HSV2) before contact with the cell, and suggest that propolis can be act in the viral envelope, affecting the adsorption or entry of the virus into the cell. The same study was carried out against Herpes simplex type 1 (HSV1), and the data suggest that when propolis is added before or at the time of infection, a greater virus-inhibiting effect is observed. In the study by Peter et al (2017) with extracts of brown propolis and jataí, a greater effect of propolis was observed when previously inoculated the cells, before viral infection, suggesting the destruction of the virus inside the cell. However, the same result was not observed when this propolis sample was inoculated previously to BVDV, possibly showing different mechanisms of viral neutralization. There appears to be a wide variation in the activity of different types of propolis in antiviral activity, as when Peter et al (2017) used green propolis extract, contrary to what was observed with brown and Jatai propolis, they did not observe significant antiviral activity in any of the experiments used. These data with green propolis differ from those obtained by Fischer et al. (2007), who evaluated the effect of the antiviral activity of a Brazilian green propolis sample against bovine herpesvirus type 1 and BVDV. These differences may be related to the chemical composition of propolis from different crops. Recently, in a study carried out by us with 12-month samples of propolis from Scaptotrigona aff. Postica, we have observed a great variation in the chemical composition of propolis according to the harvest month. This variation in chemical composition is related to the variation of flowering plants in each month, which bees use for the production of honey and propolis. Unlike other studies carried out with propolis from other sources, which derives mostly from a single botanical source, the propolis from Barra do Corda used in our studies is composed of several sources, as it was obtained in a region of native forest, with a very large variety of plants (Negri et al, 2022).

One of the positive aspects of this propolis is that its origin is in a native region, without the use of pesticides, which prevents its contamination by other unnatural products. Cueto et al. (2011) also reported the antiviral action of two ethanolic extracts of propolis from Rio Grande do Sul (Brazil), against some animal viruses, such as Bovine Viral Diarrhea Virus, Adenovirus type 2 and feline Calicivirus, Canine. In this study, the presence of flavonoids such as gallic acid, quercetin, and rutin, was identified. In this work, antiviral activity against BVDV and CAV-2 was observed when propolis was added before viral infection. In a study carried out by us, (Coelho et al. 2015) we determined the antiviral action of a hydroethanolic extract of Scaptotrigona aff postica from the region of Barra de Corda, state of Maranhão, Brazil against Herpes Simplex Virus (HSV-1). The chemical analysis of propolis showed higher amounts of C-glycosyl flavones and pyrrolizidine alkaloids, which is a compound detected for the first time in a sample of origin of meliponian bees. Apparently, the viral type also influences the results of antiviral activity. Peter et al (2017), observed that jataí propolis - T. angustula) obtained better antiviral results against RNA virus (BVDV) than when using DNA virus (BoHV-1). The BVDV genome consists of a single strand of RNA, surrounded by a lipid

envelope, which constitutes a structural difference concerning BoHV-1, which is a DNA virus (Cueto et al. 2011). Eventually, this may explain the greater action of the propolis obtained from the jataí bee. Yildirim et al. (2016) observed a significant reduction in HSV-1 and HSV-2 replication with Jatay propolis. However, toxic action was observed at concentrations of 200 and 400 μg/mL. The time of action of propolis also varied according to the virus used. While the replication of HSV-1 started 24 h after infection, for HSV-2 the start of replication was after 48 h of incubation. In our study, propolis of Scaptotrigona aff postica was tested, for potential antiviral activity against arboviruses. Experiments with total propolis led to an average 256-fold reduction in replication of the three tested viruses, while the semi-purified protein led to a 512-fold reduction in Chikungunya virus production, a 64-fold reduction in Mayaro replication, and a 16-fold reduction in Zica virus production. The action of propolis was higher when the cell culture was infected 2 hours before the addition of propolis, which suggests an intracellular mechanism of action. Maybe the substance with antiviral action present in propolis may act as an agent that affects the innate antiviral immune response. IFNs, cytokines and chemokines are essential for host defense against infection. Melchjorsen et al., 2009 showed that the initial response of HSV-infected cells is the secretion of antiviral substances such as nitric oxide, defensins and cytokine production, including chemokines and IFN. Earlier we have reported IFN-I induce to NO synthesis in *T. cruzi* infection, which act in the control of the parasitemia in resistant mice. (Costa et al., 2006). As occur with trypanosomes, we believe that the mechanism of action of some antivirals is to stimulate cells to produce interferon, nitric oxide, or other cytokines. Nevertheless, its mechanism of action needs further investigation. Concluding, this study showed antiviral activity against various arbovirus. It is can be an important target for the development of new compounds as an alternative to commercial antivirals. However, in addition to the antiviral activity, it is expected to obtain many other pharmacological activities in the propolis of Scaptotrigona aff postica, given the great complexity in the composition of this product and the great variability in the seasonal chemical composition that we observed in this propolis (Negri et al., 2022)

Acknowledgment: Research support by Butantan Foundation

Conflict of interest. The authors of the article "Antiviral action of aqueous extracts of propolis from *Scaptotrigona aff. Postica* against Zica, Chikungunya and Mayaro virus" declare that there are no known conflicts of interest associated with this publication and that there was no significant financial support for this work that could have influenced its outcome.

4. References

Alves, D.L.; Benedito, E. and Fonseca, A.L., 2017. Will the Mayaro virus be responsible for the next outbreak of an arthropod-borne virus in Brazil? The Brazilian Journal of Infectious Diseases Volume 21, Issue 5, 540-544

Amoros M., Sauvager F. and Girre L., 1992. In vitro antiviral activity of propolis. Apidologie 23(3), 231-240.

Araújo, M.J.A.M., Dutra, R.P., Costa, G.C., Reis, A.S., Assunção, K.M., Libério, S.A., Silva L., Guerra, R.N.M., Ribeiro,M.N.S., and Nascimento, F.R.F., 2010. Efeito do tratamento com própolisde Scaptotrigona aff. postica sobre o desenvolvimento do tumor de Ehrlich em camundongos. Rev. Bras. farmacogn.20(4), 580-587.

Araújo, M.J.A.M., Dutra, R.P., Costa, G.C., Reis, A.S., Assunção, K.M., Libério, S.A., Silva L., Guerra, R.N.M., Ribeiro,M.N.S., and Nascimento, F.R.F., 2011. Pharmacognostic and acute toxicological evaluation of *Scaptotrigona aff. postica* propolis extract in pre-clinical assays. Natural Product Research, 1037-1046

Bankova, V., Christov, R., Marcucci, M.C., and Popov, S. 1998. Constituents of Brazilian geopropolis. Z. Naturforschung 53c: 402-406.

Bankova, V.S., Castro, S.L., Marcucci, M.C., 2000. Propolis: recent advances in chemistry and plant origin. Apidologie. 31, 3-15.

Bankova, V. 2005. Chemical diversity of propolis and the problem of standardization. J. Ethnopharmacol. 100, 114–117

Bankova, V., Galabov, A.S., Antonova, D., Vilhelmova, N., Di Perri B., 2014. Chemical composition of Propolis Extract ACF® and activity against herpes simplex virus. Phytomedicine. 21, 1432-1438.

Bastos, E.M.A.F., Galbiati, C., Loureire, M.and Scoaris, D.O. 2011. Indicadores físico-químicos e atividade antibacteriana de própolis marrom frente à Escherichia coli. Arq. Bras. Med. Vet. Zootec. 63(5), 1255-1259.

Bogdanov, S. 2014. Propolis: Composition, Health, Medicine: A Review. Bee Product Science, 1-40 http://www.beehexagon.net/files/file/fileE/Health/PropolisBookReview.pdf

Buffalo, M.C., Figueiredo, A.S., de Sousa, J.P., Candeias, J.M., Bastos, J.K., and Sforcin, J.M. 2009. The anti-poliovirus activity of *Baccharis dracunculifolia* and propolis by cellviability determination and real-time PCR. J. Appl. Microbiol. 107, 1669-1680

Cantero, T.M, da Silva Junior, P.I, Negri, Negri., Nascimento, R.M., and Mendonça, R.Z. 2022, Antimicrobial activity of flavonoids glycosides and pyrrolizidine alkaloids from propolis of Scaptotrigona aff. postica. Toxin Reviews, (in press)

CaoLormeau, V.M., Roche, C., Teissier, A., Robin, E., Berry, A.L., Mallet, H.P., 2014. Zika virus, French Polynesia, South Pacific, Emerg Infect Dis. 20, 1085–6

Carmona,G.S., Sousa,I.L.M., Laurentino,R.V., Junior,D.L., Pantoja,A.V., Mendonça,R.Z., Nascimento,R.M and Valentin, F.N. 2020. Propolis and its components can help reduce the physiopathological consequences of covid-19 infection International Journal of Development Research Vol. 10, Issue, 10, 41067-41069

Casseb, A.R., Casseb, L.M.N., Silva, S.P. and Vasconcelos, P.F.C. 2013. Arbovírus: importante zoonose na Amazônia brasileira. Vet. e Zootec.20(3), 9-21

Coelho, G.R., Mendonça, R.Z., Villar, K.S., Figueiredo, C.A., Badari, J.C., Taniwaki,N., Namiyama, G., Oliveira, M.I., Curti, S.P., Silva, P.E., Negri G., 2015. Antiviral action of hydroethanolic extract of geopropolis from *Scaptotrigona postica* against the anti-herpes simplex virus (HSV-1). Evid. Complement Altern Med. 2015, 1-10

Coelho, G.R., Figueiredo, C.A., Negri, G., Fernandes-Silva, C.C., Senna Villar, K., Badari,J.C., Oliveira,M.I., Barbosa,T.F., Taniwaki,N.N., Namiyama, G.M and Mendonça, R.M. 2018. Antiviral Activity of Geopropolis Extract from Scaptotrigona Aff. Postica against Rubella Virus Journal of Food Research; Vol. 7, No. 6, 91-106.

Cornara, L., Biagi, M., Xiao, J., Burlando B. 2017. Therapeutic Properties of Bioactive Compounds from Different Honeybee Products. Front Pharmacol. 28, 8:412.

Costa,V.M A ., Karen, C.L., Mendonça, R.Z., Gresser,I., Gollob, K.J., Abrahamsohn, I.A. 2006. Type I IFNs stimulate nitric oxide production and resistance to Trypanosoma cruzi infection. J Immunol Sep 1;177(5), 3193-200.

Costa, R.A.C., and Cruz-Landim, C. 1999. Electrophoretic polypeptide pattern of hypopharyngeal glands extracts from workers and males of *Scaptotrigona postica* Latr. (Hymenoptera, Apidae, Meliponinae). Netherlands journal of zoology. v. 49, n. 3, 157-166.

Cueto, A.P., Alves, S.H. and Pilau, M. 2011. Atividade antiviral do extrato deprópolis contra o Calicivírus felino, adenovírus canino 2 e vírus da Diarréia viral bovina. Ciência Rural 41(10), 1800-1806.

Dick, G.W., Kitchen, S.F., and Haddow, A.J. 1952. Zika virus. I. Isolations and serological specificity. Trans R Soc Trop Med Hyg. 46, 509–20

Dobrowolski, J.W., Vohoraq, S.B., Sharma, K., Shah, S.A., Naqvi, S.A.H. and Dandiya, P.C. 1991. Antibacterial, Antifungal, Antiamoebic, Antiinflammatory, and Antipyretic Studies on Propolis Bee Products. Journal of Ethnopharmacology, 35, 77-82.

Donalisio, M.R., Freitas, A.R.R and Von Zuben, A.P.B. 2017 . Arboviruses emerging in Brazil: challenges for clinic and implications for public health Rev. Saúde Pública 51, https://doi.org/10.1590/S1518-8787.2017051006889

Duffy, MR., Chen, T.H., Hancock, W.T., and Powers, A.M. 2009. Zika virus outbreak on Yap Island, Federated States of Micronesia. N Engl J Med. 360, 2536–43.

Dupont-Rouzeyrol, M., O'Connor, O., Calvez, E., Daures, M., John, M., and Grangeon, J.P. 2015. Co-infection with zika and dengue viruses in 2 patients, New Caledonia, Emerg Infect Dis. 2015;21, 381–2

Dutra, R.P., Abreu, B.V.B., Cunha, M.S., Batista, M.C.A., Torres, L.M.B., Nascimento, F.R.F., Ribeiro, M.N.S., and Guerra, R.N.M. 2014. Phenolic acids, hydrolyzable tannins, and antioxidant activity of geopropolis from the stingless bee *Melipona fasciculata* Smith. Journal of Agricultural and Food Chemistry, 62, 2549−2557.

deFarias, J.H.C., Reis, A.S., Araújo, M.A.R., Araújo, M.J.A., Assunção, A.K.M., Farias,J.C., Fialho, E.M.S., Silva, L.A., Costa, G.C., Guerra, R.N.M., Ribeiro, M.N.S., Nascimento, R.F., 2014. Effects of Stingless Bee Propolis on experimental asthma. Evid Based Complement Alternat Med 2014, 951478.

Evans, J.D., and Pettis, JS. 2005. Colony-level impacts of immune responsiveness in honey bees, *Apis mellifera*. Evolution. *59*, 2270–2274

Falcão, S.I., Vilas-Boas, M., Estevinho, L.M., Barros, C., Domingues, M.R.M., and Cardoso, S.M. 2010. Phenolic characterization of Northeast Portuguese propolis: usual andunusual compounds. Anal. Bioanal. Chem. 396, 887–897.

Fischer, G., Cleff, M.B., Dummer, L.A., Paulino, N., Paulino, A.S., Vilela, C.O., Campos ,F.S., Storch, T., Vargas G.D.A., Hübner, S.O. and Vidor, T. 2007. Adjuvant effect of green propolis on the humoral immune response of cattle immunized with bovine herpesvirus type 5. Vet. Immunol. Immunopathol.116:79-84.

Franchin, M., Da Cunha, M.G., Denny, C., Napimoga, M.H., Cunha, T.M., Bueno-Silva, B., and Rosalen, P. 2012. Geopropolis from Melipona scutellaris decreases the mechanical inflammatory hypernociception by inhibiting the production of IL-1β and TNF-α.Journal of Ethnopharmacology. 143(2), 709–715

Franchin, M., Da Cunha, M.G., Denny, C., Napimoga, M.H., Cunha, T.M., Bueno-Silva, B., and Rosalen, P. 2013. Bioactive fraction of geopropolis from Melipona scutellaris decreases neutrophils migration in the inflammatory process: Involvement of nitric oxide pathway. Evid Based Complement Alternat Med. 2013, 907041.

García, A. A., and Carril, E. P.U. 2009. Metabolismo secundario de plantas. Reduca (Biología). Serie Fisiología Vegetal. 2 (3): 119-145.

Gekker, G., Hu, S., Spivak, M., Lokensgard, J.R.and Peterson P.K. 2005. The anti-HIV-1 activity of propolis in CD4+ lymphocyte and microglial cell cultures. J Ethnopharmacol. 102:158-163

Ghisalberti, EL. 1979. Propolis: a review. Bee World. 60,59–84

Grimsley, G.R. and Pace, N.C. 2004. Spectrophotometric determination of protein concentration. Curr Protoc Protein Sci., Nov; Chapter 3:Unit 3.1.

Heukelbach, J., C.H., Kelvin, A.A., Oliveira, W.K., and Cavalcanti, L.P.G. 2016. Zika virus outbreak in Brazil. J Infect Dev Ctries, Feb 28;10(2):116-20.

Huang, S., Zhang, C.P., Wang, K., Li, G.Q., and Hu, F.L. 2014. Recent Advances in the Chemical Composition of Propolis. Molecules, 19, 19610-19632.

Kujumgiev, A. 1999. Antibacterial, antifungal, and antiviral activity of propolis of different geographic origins. J. Ethnopharmacol. 64(3), 235-240.

Lopes, N., Nozawa, C. and Linhares, R.E.C. 2014. General features and epidemiology of emerging arboviruses in Brazil. Rev Pan-Amaz Saude, 5(3), 55-64.

Ma, X., Guo, Z., Shen, Z., Liu, Y., Wang, J. and Fan Y. 2015. The anti-porcine parvovirus activity of nanometer propolis flavone and propolis flavone in vitro and in vivo. J. Evid. Based Complement. Altern. Med. 2015:1-11.

Martin, L.F.T., Rocha, E.M., Garcia, SB., and Paula, J.S. 2013. Tropical Brazilian propolis improves corneal wound healing and inflammation in rats following alkali burns. BMC Complement Altern Med. 2013; 13, 337

Melchjorsen, J., Matikainen, S. and Paludan, S.R. 2009Activation and Evasion of Innate Antiviral Immunity by Herpes Simplex Virus. Viruses. Dec; 1(3), 737–759.

Moret, Y. and Schmid-Hempel, P. 2000, Survival for immunity: The price of immune system activation for bumblebee workers. Science, 290, 1166–1168.

Nakamura, R., Watanabe, K., Oka, K., Ohta, S., Mishima, S. And Teshima, R. 2010. Effects of propolis from different areas on mast cell degranulation and identification of theeffective components in propolis. Int Immunopharmacol. Sep;10(9), 1107-12.

Negri, G., Salatino, A., Regnier, L.L.P., Salatino, M.L.F., Nascimento, R.M. and Mendonça, R.Z. 2022. A highly complex stingless bee propolis: composition and seasonal variation JSFA Reports, 2, 64-80

Nogueira-Neto, Paulo. 1997. Vida e Criação de Abelhas indígenas sem ferrão. (Meliponinae), São Paulo: Editora Nogueirapis, 445pg

Park, Y.K., Alencar, S.M., and Aguiar, C.L. 2002. Botanical origin and chemical composition of Brazilian propolis. J. Agric. Food Chem. 50, 2502–2506

Paulino, N., Dantas, AP., Bankova, V., Ionghi, DT., Scremin, A., De Castro, S.L. and Calixto, J.B. 2003. Bulgarian propolis induces analgesic and anti-inflammatory effects in mice and inhibits in vitro contraction of airway smooth muscle. Journal of Pharmacological Sciences, 93 (3), 307-313.

Peter, C.M., Picoli, T., Zani, J.L., Latosinski, G.S., de Lima, M., Vargas, G.D., Hübner, S.O. and Fischer, G. 2017. Atividade antiviral e virucida de extratos hidroalcoólicos de própolis marrom, verde e de abelhas Jataí (Tetragonisca angustula) frente ao herpersvírus bovino tipo 1 (BoHV-1) e ao vírus da diarreia viral bovina (BVDV) Pesq. Vet. Bras. 37(7), 667-675.

Popova, M.P., Graikou, K., Chinou, I. and Bankova, V.S. 2010. GC–MS profiling of diterpene compounds in Mediterranean propolis from Greece. J. Agric. Food Chem. 58, 3167–3176.

Popova, M.P., Trusheva, B., Antonova, D., Cutajar, S., Mifsud, D., Farrugia, C., Tsvetkova, I., Najdenski, H.and Bankova, V. 2011. The specific chemical profile of Mediterranean propolis from Malta. Food Chem. 1431–1435

Reed, L.J. and Muench, H., 1938. A simple Method of estimating Fifty percentendpoints. *Am J Epidemiol*. 27, 493-97.

Rosa, J..FS.T., Rosa ,A.P.A.T., Vasconcelos, .PF.C., Pinheiro, F.P., Rodrigues, S.G., Rosa, E.S.T., Dias, L.B and Cruz, A.C.R. 1998. Arboviruses isolated in the Evandro Chagas Institute, including some described for the first time in the Brazilian Amazon region, their known hosts, and their pathology for man. *In*: Rosa APAT, Vasconcelos PFC, Rosa JFST (eds). An overview of arbovirology in Brazil and neighboring countries. Instituto Evandro Chagas, Belém, p. 19-31.

Schnitzler, P., Neuner, A., Nolkemper, S., Zundel, C., Nowack, H., Sensch, K.H., Reichling J., 2010. Antiviral Activity and Mode of Action of Propolis Extracts and Selected Compounds. Phytotherapy Res. 24, 20-28.

Silveira, F. S., Melo, G. A. R. And Almeida, E. A. B. 2002. Abelhas Brasileiras, Sistemática e Identificação. Belo Horizonte, Fernando A. Silveira. 253p.

Simone-Finstrom, M., Borba, R.S., Wilson, M. and Spivak, M. 2017. Propolis Counteracts Some Threats to Honey Bee Health. Insects. Apr 29, 8(2).

Sforcin, J.M., Bankova, V., 2011. Propolis: is there a potential for the development of new drugs? J. Ethnopharmacol.133, 253-260.

SVS - Secretaria de Vigilância em Saúde, Ministério da Saúde, "Boletim Epidemiológico: Monitoramento de casos de dengue, febre de chikungunya e febre pelo vírus Zika até semana epidemiológica 51, 2017, p. 8,

Vynograd, N., Vynograd, I.and Sosnowski, Z. 2000. A comparative multi-center study of the efficacy of propolis, acyclovir, and placebo in the treatment of genital herpes (HSV). Phytomedicine. Mar;7(1), 1-6.

White, J.W., Subers, M. H. and Schepartz, A.I. 1963. Identification of inhibin, the antibacterialfactor in the honey glucose oxidase system. Biochim. Biophys. Acta. v. 73, 57-70.

Zabaiou, N., Fouache, A., Trousson, A., Baron, S., Zellagui, A., Lahouel, M.and Lobaccaro, J.A. 2017. Biological properties of propolis extracts: Something new from an ancient product. Chem Phys Lipids. Oct; 207(Pt B), 214-222.

Yildirim, A., Duran, G.G., Duran, N., Jenedi, K., Bolgul, B.S., Miraloglu, M. and Muz, M. 2016. Antiviral activity of Hatay Propolis against replication of Herpes Simplex Virus Type 1 and Type 2. Med. Sci. Monit. 22:422-430

Antiviral action of aqueous extracts of propolis from *Scaptotrigona aff. postica* against
Zica, Chikungunya, and Mayaro virus

Mendonça RZ, Nascimento RM, Fernandes ACO, da Silva Junior PI

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7

FIGURE 8

FIGURE 9

FIGURE 10

FIGURE 11

Author agreement

The authors of the article "Antiviral action of aqueous extracts of propolis from *Scaptotrigona aff. Postica* against Zica, Chikungunya and Mayaro virus" declare that there are no known conflicts of interest associated with this publication and that there was no significant financial support for this work that could have influenced its outcome and that everyone agrees with this publication