Catching more air:

A method to spatially quantify aerial triazole resistance in *Aspergillus fumigatus*

Hylke H. Kortenbosch [1]*, Fabienne van Leuven [1], Cathy van den Heuvel [1],

Sijmen S. Schoustra [1], Bas J. Zwaan [1], Eveline Snelders [1]*

[1] Laboratory for Genetics, Wageningen University & Research, Wageningen, Gelderland, The Netherlands

*Address correspondence to Hylke H. Kortenbosch, hylke.kortenbosch@wur.nl, and Eveline Snelders, eveline.snelders@wur.nl.

ABSTRACT

Airborne triazole-resistant spores of the human fungal pathogen *Aspergillus fumigatus* are a significant human health problem as the agricultural use of triazoles has selected for cross-resistance to life-saving clinical triazoles. However, how to measure the health risk posed by these inhaled spores remains unclear. Here, we describe a method for cost-effective wide-scale outdoor air sampling to measure both spore abundance as well as antifungal resistance fractions. We show that prolonged outdoor exposure of sticky seals placed in delta traps, when combined with a two-layered cultivation approach, can consistently yield sufficient colony-forming units (CFUs) for the quantitative assessment of aerial resistance levels at a spatial scale that was up to now unfeasible. When testing our method in a European pilot sampling of 12 regions, we demonstrate that the triazole-resistant fraction of airborne spores is widespread and varies between 0 and 0.1 for itraconazole ($\sim$4 mg/L) and voriconazole ($\sim$2 mg/L). This method facilitates the

assessment of health risks by pinpointing potential hot- and coldspots of environmental resistance selection. This efficient and accessible air sampling protocol opens up extensive options for fine-spatial sampling and surveillance studies.

IMPORTANCE

Aspergillus fumigatus is an opportunistic fungal pathogen which humans and other animals are primarily exposed to through inhalation. Due to the limited availability of antifungals, resistance to the first choice class of antifungals, the triazoles, in *A. fumigatus* can make infections by this fungus untreatable and uncurable. Here we describe and validate a method that allows for the quantification of airborne resistance fractions and quick genotyping of *A. fumigatus* TR-types. Our pilot study provides proof of concept of the suitability of the method for use by citizen scientists for large-scale spatial air sampling. Spatial air sampling opens up extensive options for surveillance, health-risk assessment, and studying the ecology of *A. fumigatus* and the development of triazole resistance.

INTRODUCTION

Aspergillus fumigatus is a saprophytic fungus that is ubiquitous in soils rich in organic matter [1]. *A. fumigatus* is especially abundant in dead plant material such as plant waste heaps, where its high thermotolerance allows it to grow under the high temperatures generated during the composting process and form vast numbers of asexual conidiospores [2, 3, 4]. Due to their small size and hydrophobic nature, these spores readily disperse through the air when plant waste heaps are disturbed [5, 6]. The plant waste heaps are considered the optimal habitat for *A. fumigatus*, while this fungus also is an opportunistic pathogen that can infect or colonise the lungs of birds, humans and other mammals [7, 1, 8]. Humans are believed to inhale roughly 100 spores per day on average [9, 1], and although the immune systems of most humans efficiently remove these spores, they can cause symptoms ranging from mild allergies to often-lethal acute invasive aspergillosis infections, depending on the health status of an individual's lungs and their

immune system status [10, 1]. Unfortunately, only a limited pool of antifungals is

available to treat *A. fumigatus* infections, with triazoles being the most widely used class.

Quantifying growth and antifungal resistance fractions in plant waste material is useful

for the identification of sources of environmental resistance [11], but the direct health risk

is posed by inhaled airborne spores. Efforts have been made to assess aerial resistance,

predominantly in hospital environments [12, 13, 14]. The methods range from the air

compaction of 250 L of air on agar plates [12], collection by homemade dust vanes [14]

and the overnight exposure of agar plates [13]. All of these methods are useful for the

qualitative detection of *A. fumigatus* conidiospores and resistance genotypes and have

even been used to quantitatively estimate colony forming units (CFU) per m^3 [13].

However, the CFUs reported per sample are generally quite low (<10) in these hospital

building studies, probably due to HEPA filtration of the air. When estimating a proportion

of a natural population, in our case the fraction of triazole-resistant CFUs, an adequate

sample size is key for accuracy. Specifically because at low sample sizes, sampling error

increases rapidly to the point where the sample estimates become completely

uninformative on quantitative differences in proportions, such as for antifungal resistance

fractions [15, 16]). However, quantitative assessment of aerial resistance fractions is key

for identifying hotspots and coldspots of triazole resistance, possible transmission routes

to patients, and the health risks they pose in different geographical regions. This leads to

the question; how can we increase *A. fumigatus* CFUs in air samples to accurately estimate

triazole resistance fractions?

A limitation of air compaction or other types of active air sampling devices capable of

sampling large volumes of air is that they are costly [17]. This is not a hurdle in

longitudinal studies at set locations such as hospitals, however, this makes these devices

impractical for more extensive environmental surveys that aim to estimate resistance

fractions at various geographic scales. Shelton et al. 2023 [18] therefore proposed a

sampling method that at low cost collects airborne fungal spores and also at a wide

geographical range by involving citizen-scientists. In short, this method involves

attaching two halves of a MicroAmp^{TM} clear adhesive film (Applied Biosystems^{TM}, UK)

to a window sill with poster putties and exposing them to outside air for six to eight hours on a predetermined sampling day. Subsequently, the film was recovered and returned to the lab by Freepost. In the lab, the film was then pressed onto an agar plate and cultured at 43°C. This method proved effective at capturing *A. fumigatus* spores on a nationwide level, but on average yield only 1.25 *A.fumigatus* CFUs per sample [18]. Thus, the number of recovered CFUs was too low to accurately estimate antifungal resistance fractions per individual sample.

Building on the method of Shelton et al. 2023, we also used sticky seals to capture spores, now with exposure times of four weeks. Compared to agar plates, these seals are less prone to drying out, weathering, or unwanted microbial growth. As prolonged exposure also means potentially increased exposure to other microbes, we tested a culturing method in which we poured Flamingo medium [19] selective for *A. fumigatus*, directly on top of exposed seals in two distinct layers. The first layer allows all *A. fumigatus* spores to germinate and grow, while the second layer is selective for resistance against the triazole compound of choice. In addition to developing this culturing approach, we tested it in an international pilot study, including 12 sampled regions, for which we asked fellow scientists to follow a provided protocol supported with instructions via a video. With this, we assessed whether the use of foldable delta traps, protecting the seals from UV light and rain, could be combined with the use of sticky seals in a flexible, accessible and effective protocol for affordable and large-scale outdoor spore capture.

RESULTS AND DISCUSSION

Validation layer culturing approach

We tested whether culturing from seals inoculated with *A. fumigatus* conidia by adding Flamingo media with different triazole treatments directly on top would, relative to plating conidia on the agar surface, affect the assay's utility to visually screen for triazole-resistant *A. fumigatus* colonies. We found that an interaction between the seals and itraconazole strongly diminished suppression of itraconazole-sensitive strains,

compared to using itraconazole in Flamingo agar plates without seals. (see Supplement -

Effect of seals on triazole selectivity). To address this technical issue, we validated a

layered culturing approach where a triazole-negative layer of medium acts as a buffer

between the triazole-containing layer and the seal. For the two clinical triazoles

itraconazole and voriconazole, we found that culturing with the described two-layer

culturing method only allowed colonies of *A. fumigatus* strains with a $MIC_{95} \geq 2$ mg/L to

breach the agar surface of the upper azole-containing layer. The rate at which different *A.*

fumigatus strains did so by the end of incubation seemed largely tied to their MIC_{95} value

(Fig. 1). Similarly, we tested whether this two-layer culturing approach could also be used

to detect resistance to the agricultural triazoles difenoconazole (4 mg/L) and tebuconazole

(6 mg/L). However, these triazoles were found to be insufficiently selective when added

to the triazole-containing layer during the layered culturing approach (see Tables S1 and

S2). We also conducted limited tests with the agricultural triazoles bromuconazole,

epoxiconazole, and propiconazole (data not shown), and these compounds also appeared

to be ineffective for resistance screening with the layered culturing method.

Strain	Treatment			MIC$_{95}$	
	_	ITR 4	VOR 4	ITR	VOR
AfIR974				0,5	1
AfIR964				0,5	1
11-1				16	2
78-2				16	4
39-3				16	1
19-2				1	16
8-3				2	16
40-1				2	16

FIG 1 Growth from seals innoculated with *A. fumigatus* conidia of strains with known minimal inhibitory concentration (MIC) values through the top layer of Flamingo medium after eight days of incubation at 48 °C. The included treatments in the upper agar layer are triazole-negative, 4 m/L itraconazole (ITR 4), and 4 mg/L voriconazole (VOR 4). MIC$_{95}$ values in mg/L for itraconazole (ITR) and voriconazole (VOR) of the strains are given in the two right-most columns.

We do not have a full explanation for why some triazoles are applicable in this

culturing method and not others. Table 1 lists key characteristics for several medical and

agricultural triazole compounds, indicating whether they were tested to be effective with
the described layered culturing method to discriminate between azole-susceptible and
azole-resistant *A. fumigatus*. Ideally, the applied triazole concentration in the upper agar
layer should be higher than the mean MIC_{95} of wild-type strains for discrete selection for
resistant strains. The higher mean wild-type MIC_{95} values in agricultural fungicides(Table
1) indicate that they should be present in higher concentrations in the agar to be effective
at screening for resistance. Unfortunately, technical aspects of layered culturing directly
from sticky seals can make achieving such concentrations difficult. We hypothesize that
low solubility in water (Table 1) may cause some triazoles to dissolve in the adhesive of
the seal preferentially when applied directly onto the seal in water-based agar, lowering
the effective concentration in the agar and reducing the effectiveness of the assay. In the
case of itraconazole, its low solubility in water limits its diffusion into the lower
triazole-negative layer of the medium. We believe the lower agar layer functions as a
buffer here, limiting interaction with the seal. However, in the case of difenoconazole,
tebuconazole, and bromuconazole, their intermediate solubilities in water may render this
buffering layer less effective. These triazoles may still diffuse into the triazole-negative
layer and subsequently bind to the seal, allowing sensitive colonies to grow uninhibited.
Voriconazole has a much higher water solubility than itraconazole (Table 1) and likely
diffuses into the triazole-negative layer but has seemingly no interaction with the seal (see
supplement - Effect of seals on triazole selectivity).

TABLE 1   Water solubilities of different triazoles, the mean MIC_{95} values[a] for *A. fumigatus* strains with different *cyp*51A genotypes (wild type, TR34/L98H,

TR46/Y121F/T289A), and whether the double-layer culturing approach can be used to screen for resistance to these triazoles.

Triazole	Solubility[a]	Mean WT[b] MIC$_{95}$	Mean TR34[b] MIC$_{95}$	Mean TR46[b] MIC$_{95}$	Applicable culturing method this study
Itraconazole	10 [20]	0.125	>16	2	Yes
Voriconazole	97.8 [21]	0.5	4	>16	Yes
Difenoconazole	15 [22]	1	>16	>16	No
Tebuconazole	36 [23]	1	16	16	No
Bromuconazole	50 [24]	1	16	>16	No
Epoxiconazole	6.63 [23]	2	>16	>16	No
Propioconazole	150 [25]	2	>16	>16	No

[a]Solubility is given at mg/L in water of 25 °C

[b]Mean MIC$_{95}$ values are given in mg/L and were adapted from Snelders et al. 2012 [26].

Generally, MIC tests are intended to predict the clinical outcome of treatment of antifungal drugs on isolates under standardised conditions based on established breakpoints [27]. For our purpose of large-scale resistance screening of environmental air samples, we apply the triazoles in Flamingo medium, at 48 ° C, in a non-homogenous (layered) agar, to already germinated colonies growing on the same plate. These culturing conditions are all linked to factors which are known to impact MIC values [27, 28, 29, 30]. The fact that there are inherently no established clinical breakpoints for agricultural fungicides makes finding appropriate concentrations to assay for cross-resistance, and interpreting the output of any such non-standard assay, complex. Therefore, before using this two-layered cultured approach to screen for fungicide resistance it is important to consider the chemical properties of a fungicide. Specifically, its solubility in water, the

162 MIC values of the WT *A. fumigatus* population to it, and its inhibitory effect on *A.*

163 *fumigatus* when used in the described culturing method.

International air sampling pilot

Of the 70 air sampling packages that were distributed for the international pilot study,

54 were returned in good order and selected for analysis. These samples were grouped

into 12 different regions, in which samples were taken within a circular area with a 50 km

radius (Fig. 2). Five of these 54 samples were excluded because of low CFUs (>3x lower

than the regional average), three of them were likely taken indoors rather than outdoors.

Of the other two samples this was reported to be the case by the participants. Three

samples were excluded because of high CFUs (>2x higher than the regional average): one

of these traps had fallen from the tree it was hung up in and had been on the ground for

an unknown time, and the remaining two were likely deployed close to an *A. fumigatus*

growth environment such as a heap of degrading plant waste material. This left 46

representative samples for the analysis of aerial baseline triazole resistance in *A. fumigatus*.

To assess regional differences in CFU counts, we also excluded samples deployed later

than December 19th 2022 to mitigate the known effect of seasonality on the prevalence of

airborne *A. fumigatus*. This left 39 samples for analysis of regional differences.

FIG 2   The European regions sampled for the air sampling pilot between
November 2022 and January 2023. The green circles designate the regions in
which the delta traps were deployed (within a 50 km radius).

Regional differences in CFU counts

Among the sampled regions, we found significant differences in spore counts between

regions (see Fig.3 and Table S3). Samples from The Netherlands, Belgium, and the

bordering region of Düsseldorf (Germany) contained 3-fold higher levels of $\sim$ CFUs than

most other regions. We assessed whether weather conditions during the sampling

window were significantly correlated with the median CFU counts per trap. We tested for

regional averaged UV light, total precipitation, wind speed, and temperature and only

found a significant correlation with wind speed (Estimate = 0.5370, SE = 0.1407, z value =
3.818, Pr(>|z|) = 0.000135). This is an expected positive correlation, given that our
passive spore capture method requires airflow over the stickers to capture spores.
However, in an earlier longitudinal study where more controlled volumes of air were
sampled, this correlation was also observed [31]. This suggests that increased CFU counts
at greater wind speeds may not simply be a result of more air flowing over the sticker but
also reflect an absolute increase in amounts of spores in the air, possibly due to increased
disturbance of, and spore release from, *A. fumigatus* sources [1, 32]. Furthermore, the
relationship with wind speed is not linear. The regions of Copenhagen, Verton, and
Nantes with the highest average wind speed have the lowest counts. These are all coastal
regions, which could also have an effect, given that *A. fumigatus* is a terrestrial mould.
Moreover, given the limited size of our pilot data and that the regions with the highest
counts are adjacent, we cannot rule out spatial auto-correlation in the data. More
systematic spatial sampling will be needed under variable weather conditions to separate
the factor space from the effect of weather conditions.

FIG 3   The median CFU count per trap per region (mean ± SE). The
Netherlands, Belgium, and the bordering region of Düsseldorf have noticeably
higher CFU counts than most other sampled regions.

Our analysis shows regional differences in CFU capture by the traps. Pilot studies,
similar to the one described here should precede systematic sampling for aerial triazole
resistance in new regions to assess what number of CFUs can be expected. Due to the
passive nature of our air sampling method and the possibility of seasonality in *A.*
fumigatus abundance [18], CFU counts may, where necessary be increased by 1) increasing
the length of the sampling window and/or 2) adjusting the sampling season. This raises
the question: how are these CFU counts per sample relevant for the assessment of
resistance fractions?

211 **How accuracy matters**

212   When estimating a true value in a population, in our case the resistance fraction to
213 triazoles, power calculations are key to determining what number of colonies will provide
214 the desired precision of the population estimate. To determine the number of CFUs (S)
215 needed to estimate a proportion of a population with a binomial distribution the
216 following equation can be used,

217
$$S = Z^2 \times P \times \frac{(1-P)}{E^2} \tag{1}$$

218 which we can rework to calculate the error size:

219
$$E = \frac{Z \times \sqrt{P(1-P)}}{\sqrt{S}} \tag{2}$$

The key factors here are the expected population resistance fraction (P), the sampling error
(E) and the critical value (Z) for the desired confidence level of your sample estimate,
which is 1.96 for the 95% confidence intervals commonly applied in biology [33]. Based on
data from previous studies [34, 35, 18], we expected a resistance fraction of ∼5%, so will
fill in 0.05 as our P. Combining this value with the common 95% confidence interval (Z =
1.96), we can simplify the equation to:

226
$$E = \frac{0.43}{\sqrt{S}} \tag{3}$$

On the one hand, plotting this equation shows that the sampling error, or how much a
sample resistance fraction may differ from the true population resistance fraction, rapidly
increases at counts <25 (Fig. 4). Previous estimates of baseline resistance based on aerial
surveys reported resistance fractions of ∼0.05 [18, 17]. Therefore, a detection limit of >0.04
is undesirable for quantitative assessment of aerial resistance fractions. Hence, we only
included samples that contained ≥25 CFUs in our quantitative assessment of regional
differences in triazole resistance fractions. On the other hand, having over 250 CFUs
makes counts less accurate due to overcrowding, and only provides marginal statistical
benefits (Fig. 4). This gives us our desired range of 25-250 CFUs per sample. Based on this,
only regions with two or more samples in this range were included in our quantitative
analysis of resistance fractions. This further restricted our analysis to 22 samples for
itraconazole resistance, and 18 for voriconazole resistance.

FIG 4   A plot of equation 3, showing the effect of CFU counts or sample size on
the estimated resistance fraction's 95% confidence intervals, or precision. Where
we are under the conservative assumption that the true population resistance
fraction is 0.05. The desirable range of 25-250 CFUs per sample for resistance
fraction estimations is highlighted.

Validation resistance selection environmental air samples

All isolates scored as resistant on itraconazole double-layered flamingo medium plates grew when subcultured on 4 mg/L itraconazole supplemented standard medium, while the isolates scored as resistant on the voriconazole double-layered flamingo medium plates grew when supplemented on 2 mg/L voriconazole supplemented standard medium. The two *cyp*51A gene haplotypes known to be most common in the environment, tandem repeats of either 34 (TR_{34}) or 46 (TR_{46}) basepairs [36], were dominant among the resistant isolates (51/58). TR_{34} was more abundant than TR_{46} (42/51) (see table S3). TR_{34} haplotypes are known to confer high-level resistance to itraconazole and often have lower-level cross-resistance to voriconazole [26, 37]. The reverse is true for TR_{46} haplotypes, which confers high-level resistance to voriconazole, and often lower-level cross-resistance to itraconazole [26, 37]. This higher abundance of TR_{34} relative to TR_{46} haplotypes in air samples is in line with what has been observed among airborne spores in the UK [18]. The ratio between the two haplotypes seems to more closely resemble that of European clinical isolates [37, 35, 38, 39] than the ratio observed in known environmental resistance hotspots sampled in the Netherlands, where TR_{46} haplotypes were more common [40]. These phenotypic and genotypic data, combined with our earlier described validation experiment, demonstrate that the described culturing method is reliable for resistance fraction estimations in environmental air samples.

Regional baseline resistance

Figure 5 shows the estimated resistance fractions to itraconazole [A] and voriconazole [B] per sampled region. For both triazoles baseline resistance fractions ranged from 0-0.1. We found no significant regional differences for either screened triazole. Triazole resistant *A. fumigatus* is known to commonly be cross-resistant to multiple 14α-demethylase inhibitors (DMIs) [26]. In light of this, our findings are comparable to those of multiple European studies assessing resistance in both soil and air which reported similar resistance fractions ranging between 0 and 0.1 [41, 42, 17, 43, 18]. Given that neither air nor soil are prolific growth environments for *A. fumigatus*, we hypothesise that such

268 values represent local baseline resistance levels present in the environment rather than
269 resistance hotspots such as agricultural plant waste piles containing residues of triazoles.

270 [A] [B]

FIG 5 Itraconazole ($\sim$4 mg/L) [A] and voriconazole ($\sim$2 mg/L) [B] resistance fractions per region (mean $\pm$ SE). Only regions where multiple samples used for resistance fraction estimations grew $\geq$25 CFUs, were included.

The fact that air sampling efforts have so far yielded similar antifungal resistance
fractions [17, 18] seems to indicate that spatial differences in airborne triazole resistance
are absent. However, the air is not a growth environment of *A. fumigatus*, but rather the
medium through which it disperses. The airborne *A. fumigatus* population represents a
product of many growth environments under the influence of atmospheric mixing [18].
The effects of resistance hotspots on aerial resistance fractions may therefore be subtle or
relatively local. To detect such spatial differences a method that can both provide precise
point estimations of antifungal resistance and practically be used for large-scale spatial
sampling is needed. Our pilot study provides proof of concept that the presented method
is suitable for such large-scale air sampling efforts.

281 Hotspot or coldspot?

With this new tool for large-scale environmental air sampling, we argue for a sharper
definition of triazole resistance hotspots [44, 36]. That is, resistance hotspots are *A.*
fumigatus growth environments in which the resistance fraction to triazole(s) increases
significantly above the aerial baseline resistance. Such hotspots may be considered

potential health hazards if they, in turn, significantly increase the local aerial triazole
resistance fraction and the prevalence of triazole resistant *A. fumigatus* spores in their
immediate surroundings above the regional baseline resistance level. Peaks in aerial
antifungal resistance may be linked to known and unknown hotspots by combining
spatial data on aerial resistance with spatial data on land use (type of agriculture) and
economic activity.

Outlook

Here we provide proof concept of a combination of air sampling and a culturing
method that holds the potential to assess aerial resistance fractions through space. The
effectiveness of the method, its relatively low cost, simplicity of use in the field and thus
suitability for use by citizen scientists [45], make this method suitable for large-scale air
sampling efforts. This may prove key to bridging the knowledge gap in understanding
the transmission of triazole-resistant *A. fumigatus* from the environment to patients.
Numbers of airborne *A. fumigatus* appear to vary through space. Therefore, pilot studies
will need to precede large-scale spatial sampling. In global regions with lower (<25) CFU
yields from the traps adjustments in exposure time and sampling window will have to be
made to provide higher yields and reliable resistance fraction estimations. Furthermore,
due to the greater technical complexity of culturing in layered agar directly from a sticky
seal, the described method is not suitable for all triazoles. These unsuitable triazoles
include tebuconazole which has been used for screening in previous air sampling studies
[17, 18]. These mixed results highlight the need for further validation preceding the
application of other fungicides in the here-described culturing approach. Nonetheless,
having an efficient and accessible air sampling and resistance screening protocol opens up
extensive options for surveillance and research. The impact of known resistance hotspots
[44, 40] on local aerial resistance levels can now be quantified. This is key for evaluating
preventive measures where they matter most, in the air.

MATERIALS AND METHODS

Validation layered selective culturing

To test whether the selection for triazole-resistant colonies was effective when culturing with BIORAD^{TM} microseal seals 'B' (MSB1001), hereafter called 'seals', we selected eight *A. fumigatus* strains with known, contrasting MIC$_{95}$ values for itraconazole and voriconazole (Fig. 1). We selected an additional set of four strains with contrasting MIC$_{95}$ values for difenoconazole and tebuconazole for a similar test with these compounds (See Table S1). Spore suspensions were made with 0.05% saline-Tween. This solution was prepared by adding 500 µl tween-80 to 1 l 0.9% saline solution. We diluted the spore suspension such that they contained 1-40 spores per 75 µl. We cut seals into small fragments (4 cm x 6,85 cm), such that they could be placed in 9 cm Petri dishes. Within a levelled laminal flow hood, we then placed the sticky seals inside of the Petri dishes, the sticky side facing up and exposed. 75 µl of spore suspension was pipetted in droplets across the seal to ensure colony spread. We left the droplets to dry on the seal for an hour under a laminar flow hood. We prepared Flamingo medium as described in (Zhang et al. 2021) except that, after the autoclaving step, we let the medium cool to 60 °C before adding the antimicrobial supplements and 12.5 ml of 2M sucrose as a carbon source. Subsequently, 24 ml of Flamingo medium was poured into the plate so that the seal and bottom of the plate were covered with an 8mm layer of medium. This ensured that the agar was liquid but still within the temperature tolerance range (Kozakiewicz and Smith 1994) of *A. fumigatus*. We left the agar to solidify at room temperature and incubated the plates at 48 °C for 30 hours, then kept them at 4 °C overnight, and cultured them for 8 more hours at 48 °C. Following this intermittent incubation, we marked all visible colonies and added the second layer of Flamingo medium of 12 ml (4 mm thickness) containing either no triazoles, 4 mg/L itraconazole or 4 mg/L voriconazole to plates from a set of plates for each of the reference strains. The plates were then incubated for six more days at 48 °C and the growth was scored on the agar surface.

International Delta trap air sampling

During November 2022 a total of 70 air sampling packages were distributed or posted across Western Europe to the participants, who had no prior experience using the air sampling method. The sampling packages contained a foldable delta trap (Biogrow, Leuven, BE), a zip-lock bag, containing three sticky seals (of which the length was cut down to 11 cm by removing one of the non-sticky ends), a piece of paper to note down postal code, country and sampling start and end dates, a 30 cm piece of rope to tie up the delta trap, a strip of six poster putties, and sampling instructions. All of these materials were packaged in a 22.9 by x 33.4 cm envelope. Participants were given instructions to place the sticky seals in the delta trap and expose them for four weeks in an outdoor location near their homes. After exposure, the participants re-covered the seals and returned them via post. For a detailed description of the sampling package and the sampling instructions, including a link to the instruction video, see Supplement - Delta trap air sampling.

Resistance screening protocol environmental air samples

To test the applicability of the layered culturing approach in environmental samples, we modified the layered culturing approach to selectively culture *A. fumigatus* from sticky seals that had been exposed to outdoor air. The volumes of Flamingo medium were adjusted to 60 ml for the initial triazole-negative flamingo medium, and 30 ml for the triazole-treatment layer. We did so to retain 8 mm and 4mm thickness, respectively, of the agar layers in larger square Petri dishes with vents (120 mm X 120 mm X 17mm) that can accommodate a sticky seal of which one non-sticky strip has been cut such that all seals are 11 cm long. Furthermore, the seals are attached to the bottom of the agar plate with Pritt^{TM} Compact Adhesive Rollers to keep the larger sticky seals flat. We adjusted the initial incubation times to 16h at 48 °C, six hours at 4 °C, and 20 hours at 48 °C. After adding the secondary triazole layer, the plates were incubated for 24 more hours at 48 °C. At this time, the total number of CFUs on the plates was counted. Finally, plates were incubated for seven more days at 48 °C after which the sporulating colonies were counted

367 as resistant. The number of resistant colonies per seal (R) and the total number of CFUs
368 (Tot) per seal can be used together to estimate a resistance fraction (RF) by RF = R/Tot.
369 For a detailed day-by-day description including pictures of the environmental air sample
370 resistance screening protocol and its rationale, see Supplement - Layered culturing
371 protocol.

Validation resistance screening environmental samples

To validate the selectivity of our culturing approach, we selected colonies from
environmental air samples which visually sporulated with the unaided eye within nine
days following the start of incubation. We selected 28 and 30 colonies from voriconazole
and itraconazole plates, respectively. Colonies from voriconazole and itraconazole plates
were transferred to a set of triazole-negative growth control slants, and slants containing 2
mg/L of voriconazole or 4 mg/L of itraconazole, respectively. We used 2 mg/L
voriconazole to validate the resistance of colonies from the voriconazole plates rather than
the 4 mg/L used in the selective layer because in earlier validation experiments the rapid
(within 20 hours) suppression of growth of colonies of voriconazole-susceptible strains in
the permissive layer indicated more diffusion through permissive agar layer than in the
itraconazole plates. This observed greater diffusion corresponds to the 10x higher
solubility in water of voriconazole than itraconazole [20, 21]. Due to this diffusion, the
effective concentration in the double-layered voriconazole plates is lower than 4 mg/L,
likely closer to 2 mg/L. To validate clinical resistance, the concentrations of itraconazole
and voriconazole in the slants were chosen based on the EUCAST clinical breakpoints for
these compounds [29]. The slants were subsequently incubated at 37 °C and scored for
growth after four days. The isolates were then genotyped by harvesting spores from the
growth control slants and extracting DNA using the protocol described in Fraczek et al.
2019 [46]. To detect known tandem repeat haplotypes we designed a new primer
combination which allows for genotyping different TR types on agarose gel without the
need for Sanger sequencing. We used this primer combination to amplify the promoter
region of *cyp*51A as a resistance marker. For the PCR protocol, see Supplement - PCR
protocol *cyp*51A genotyping.

Data analysis

For the analysis of the CFU yield per seal, we used a negative binomial GLM with region as a fixed effect. Weather data was extracted from "ERA5 monthly averaged data on single levels from 1940 to present" database of the Copernicus Institute. To model weather effects on the CFU yield we included Total precipitation, 2m temperature, 10m windspeed and Downward UV radiation at the surface as fixed effects. A post hoc Tukey test was performed to test the significance of pairwise differences. For the analysis of the resistance fraction data per region, we used binomial GLMs. Statistical analysis and data visualisation were performed in R version 4.3.1 [47], using the following R packages: AER 1.2-10 [48], AICcmodavg 2.3-2 [49], cmsafops 1.3.0 [50], dplyr 1.1.2 [51], emmeans 1.8.8 [52], ggplot2 3.4.3 [53], ggpubr 0.6.0 [54], lme4 1.1-34 [55], MASS 7.3-60 [56], multcomp 1.4-25 [57], multcompview 0.1-9 [58], ncdf4 1.21 [59], rgdal 1.6-7 [60], tidyr 1.3.0 [61], rnaturalearth 0.3.4 [62], rnaturalearthdata 0.1.0 [63], raster 3.6-23 [64], sf 1.0-14 [65, 66].

ACKNOWLEDGMENTS

We want to thank Bart Pannebakker, Frank Becker, Ben Auxier, Eric Bastiaans, Jianhua Zhang from the Laboratory of Genetics and the other members of the Groen III One Health Consequences of Circularity team for their helpful suggestions and critical discussions. We thank Bo Briggeman for filming and editing the revised Delta trap instruction videos. We also want to thank all our collaborators and fellow scientists who contributed to the international air sampling pilot. We finally thank the BSc students who helped conceptualise and test the air sampling method during the MEE course at Wageningen University & Research during spring 2022 and spring 2023.

DATA AVAILABILITY STATEMENT

The R code and data are made available at:

https://git.wur.nl/korte058/catching_more_air.

FUNDING

This research was funded by The Netherlands Organization for Scientific Research (NWO) under the Groen III program (GROEN.2019.002): One health consequences of circularity—What lessons to learn from the saprophytic and human pathogenic fungus *Aspergillus fumigatus*?

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

References

[1] **Kwon-Chung KJ, Sugui JA**. 12 2013. Aspergillus fumigatus—What Makes the Species a Ubiquitous Human Fungal Pathogen? *PLoS Pathogens* 9 (12):e1003743. doi: 10.1371/journal.ppat.1003743.

[2] **Altenburger A, Bender M, Ekelund F, Elmholt S, Jacobsen CS**. 2014. Steam-treatment-based soil remediation promotes heat-tolerant , potentially pathogenic microbiota. *Environmental Technology* 35 (6):773–780. doi: http://dx.doi.org/10.1080/09593330.2013.850520.

[3] **Zhang J, Snelders E, Zwaan BJ, Schoustra SE, Meis JF, van Dijk K, Hagen F, van der Beek MT, Kampinga GA, Zoll J, Melchers WJG, Verweij PE, Debets AJM**. 7 2017. A Novel Environmental Azole Resistance Mutation in Aspergillus fumigatus and a Possible Role of Sexual Reproduction in Its Emergence. *mBio* 8 (3):1–13. doi: 10.1128/mBio.00791-17.

[4] **Schoustra SE, Zhang J, Zwaan BJ, Debets AJF, Verweij PE, Buijtenhuijs D, Rietveld AG**. 2018. New insights in the development of azole-resistance in Aspergillus fumigatus. National Institute for Public Health and the Environment, Bilthoven, The Netherlands.

[5] **Sewell TR, Zhu J, Rhodes J, Hagen F, Meis JF, Fisher MC, Jombart T**. 5 2019.

Nonrandom Distribution of Azole Resistance across the Global Population of

Aspergillus fumigatus. *mBio* 10 (3):1–11. doi:10.1128/mBio.00392-19.

[6] **Williams B, Douglas P, Roca Barcelo A, Hansell AL, Hayes E**. 2 2019. Estimating

Aspergillus fumigatus exposure from outdoor composting activities in England

between 2005 and 14. *Waste Management* 84:235–244. doi:

10.1016/j.wasman.2018.11.044.

[7] **Hohl TM, Feldmesser M**. 11 2007. Aspergillus fumigatus: principles of pathogenesis

and host defense. *Eukaryotic cell* 6 (11):1953–63. doi:10.1128/EC.00274-07.

[8] **Debergh H, Becker P, Vercammen F, Lagrou K, Haesendonck R, Saegerman C,**

Packeu A. 3 2023. Pulmonary Aspergillosis in Humboldt Penguins—Susceptibility

Patterns and Molecular Epidemiology of Clinical and Environmental Aspergillus

fumigatus Isolates from a Belgian Zoo, 2017–2022. *Antibiotics* 12 (3):584. doi:

10.3390/antibiotics12030584.

[9] **Ascioglu S, Rex JH, de Pauw B, Bennett JE, Bille J, Crokaert F, Denning DW,**

Donnelly JP, Edwards JE, Erjavec Z, Fiere D, Lortholary O, Maertens J, Meis JF,

Patterson TF, Ritter J, Selleslag D, Shah PM, Stevens DA, Walsh TJ, Invasive Fungal

Infections Cooperative Group of the European Organization for Research and

Treatment of Cancer, Mycoses Study Group of the National Institute of Allergy and

Infectious Diseases. 1 2002. Defining opportunistic invasive fungal infections in

immunocompromised patients with cancer and hematopoietic stem cell transplants:

an international consensus. *Clinical infectious diseases : an official publication of the*

Infectious Diseases Society of America 34 (1):7–14. doi:10.1086/323335.

[10] **Verweij PE, Snelders E, Kema GH, Mellado E, Melchers WJ**. 12 2009. Azole

resistance in Aspergillus fumigatus: a side-effect of environmental fungicide use? *The*

Lancet Infectious Diseases 9 (12):789–795. doi:10.1016/S1473-3099(09)70265-8.

[11] **Snelders E, Huis in 't Veld RAG, Rijs AJMM, Kema GHJ, Melchers WJG, Verweij**

PE. 6 2009. Possible Environmental Origin of Resistance of Aspergillus fumigatus to

Medical Triazoles. *Applied and Environmental Microbiology* 75 (12):4053–4057. doi:
10.1128/AEM.00231-09.

[12] **Reboux G, Gbaguidi-Haore H, Bellanger AP, Demonmerot F, Houdrouge K,**
Deconinck E, Bertrand X, Millon L. 2014. A 10-year survey of fungal
aerocontamination in hospital corridors: A reliable sentinel to predict fungal exposure
risk? *Journal of Hospital Infection* 87 (1):34–40. doi:10.1016/j.jhin.2014.02.008.

[13] **Demuyser T, De Cock E, Sermijn E**. 11 2019. Airborne Aspergillus fumigatus
contamination in an intensive care unit: Detection, management and control. *Journal of*
Infection and Public Health 12 (6):904–906. doi:10.1016/j.jiph.2019.04.016.

[14] **Godeau C, Reboux G, Scherer E, Laboissiere A, Lechenault-Bergerot C, Millon L,**
Rocchi S. 6 2020. Azole-resistant Aspergillus fumigatus in the hospital: Surveillance
from flower beds to corridors. *American Journal of Infection Control* 48 (6):702–704. doi:
10.1016/j.ajic.2019.10.003.

[15] **Davis C**. 8 2014. Enumeration of probiotic strains: Review of culture-dependent and
alternative techniques to quantify viable bacteria. *Journal of Microbiological Methods*
103:9–17. doi:10.1016/j.mimet.2014.04.012.

[16] **Baveja CP, Aggarwal P**. 4 2017. Statistical Analysis of Microbiological Diagnostic
Tests. *Indian Journal of Medical Microbiology* 35 (2):184–193. doi:
10.4103/ijmm.IJMM{_}16{_}215.

[17] **Fraaije B, Atkins S, Hanley S, Macdonald A, Lucas J**. 12 2020. The Multi-Fungicide
Resistance Status of Aspergillus fumigatus Populations in Arable Soils and the Wider
European Environment. *Frontiers in Microbiology* 11 (December):1–17. doi:
10.3389/fmicb.2020.599233.

[18] **Shelton JMG, Rhodes J, Uzzell CB, Hemmings S, Brackin AP, Sewell TR,**
Alghamdi A, Dyer PS, Fraser M, Borman AM, Johnson EM, Piel FB, Singer AC,
Fisher MC. 7 2023. Citizen science reveals landscape-scale exposures to
multiazole-resistant Aspergillus fumigatus bioaerosols. *Science Advances* 9 (29).
doi:10.1126/sciadv.adh8839.

[19] **Zhang J, Debets AJM, Verweij PE, Schoustra SE**. 5 2021. Selective Flamingo
Medium for the Isolation of Aspergillus fumigatus. *Microorganisms* 9 (6):1155. doi:
10.3390/microorganisms9061155.

[20] **Van Peer A, Woestenborghs R, Heykants J, Gasparini R, Gauwenbergh G**. 7 1989.
The effects of food and dose on the oral systemic availability of itraconazole in healthy
subjects. *European Journal of Clinical Pharmacology* 36 (4):423–426. doi:
10.1007/BF00558308.

[21] **Reber E, Neyer P, Schönenberger KA, Saxer C, Bernasconi L, Stanga Z, Huber A,**
Hammerer-Lercher A, Mühlebach S. 9 2021. Physicochemical Stability and
Compatibility Testing of Voriconazole in All-in-One Parenteral Nutrition Admixtures.
Pharmaceutics 13 (9):1447. doi:10.3390/pharmaceutics13091447.

[22] **Khalaf B, Hamed O, Jodeh S, Bol R, Hanbali G, Safi Z, Dagdag O, Berisha A,**
Samhan S. 2021. Cellulose-based hectocycle nanopolymers: Synthesis, molecular
docking and adsorption of difenoconazole from aqueous medium. *International Journal*
of Molecular Sciences 22 (11). doi:10.3390/ijms22116090.

[23] **Li H, Li Y, Wang W, Wan Q, Yu X, Sun W**. 2022. Uptake, translocation, and
subcellular distribution of three triazole pesticides in rice. *Environmental Science and*
Pollution Research 29 (17). doi:10.1007/s11356-021-17467-6.

[24] **Hariyadi HR**. 2018. The disappearance of bromuconazole fungicide in domestic
waste anaerobic fermentation. *In* IOP Conference Series: Earth and Environmental
Science, vol 160. doi:10.1088/1755-1315/160/1/012013.

[25] **Battaglin WA, Sandstrom MW, Kuivila KM, Kolpin DW, Meyer MT**. 2011.
Occurrence of azoxystrobin, propiconazole, and selected other fungicides in US
streams, 2005-2006. *Water, Air, and Soil Pollution* 218 (1-4). doi:
10.1007/s11270-010-0643-2.

[26] **Snelders E, Camps SMT, Karawajczyk A, Schaftenaar G, Kema GHJ, van der Lee**
HA, Klaassen CH, Melchers WJG, Verweij PE. 3 2012. Triazole Fungicides Can
Induce Cross-Resistance to Medical Triazoles in Aspergillus fumigatus. *PLoS ONE* 7
(3):e31801. doi:10.1371/journal.pone.0031801.

[27] **Berkow EL, Lockhart SR, Ostrosky-Zeichner L**. 6 2020. Antifungal Susceptibility Testing: Current Approaches. *Clinical Microbiology Reviews* 33 (3). doi: 10.1128/CMR.00069-19.

[28] **Rex JH, Pfaller MA, Galgiani JN, Bartlett MS, Espinel-Ingroff A, Ghannoum MA, Lancaster M, Odds FC, Rinaldi MG, Walsh TJ, Barry AL**. 2 1997. Development of Interpretive Breakpoints for Antifungal Susceptibility Testing: Conceptual Framework and Analysis of In Vitro-In Vivo Correlation Data for Fluconazole, Itraconazole, and Candida Infections. *Clinical Infectious Diseases* 24 (2):235–247. doi: 10.1093/clinids/24.2.235.

[29] **Arendrup MC, Verweij PE, Mouton JW, Lagrou K, Meletiadis J**. 12 2017. Multicentre validation of 4-well azole agar plates as a screening method for detection of clinically relevant azole-resistant Aspergillus fumigatus. *The Journal of antimicrobial chemotherapy* 72 (12):3325–3333. doi:10.1093/jac/dkx319.

[30] **Vahedi-Shahandashti R, Stubenböck MM, Lass-Flörl C**. 9 2023. The Influence of Medium Composition on EUCAST and Etest Antifungal Susceptibility Testing. *Journal of Fungi* 9 (10):973. doi:10.3390/jof9100973.

[31] **Priyamvada H, Singh RK, Akila M, Ravikrishna R, Verma RS, Gunthe SS**. 9 2017. Seasonal variation of the dominant allergenic fungal aerosols – One year study from southern Indian region. *Scientific Reports* 7 (1):11171. doi:10.1038/s41598-017-11727-7.

[32] **Elbert W, Taylor PE, Andreae MO, Pöschl U**. 9 2007. Contribution of fungi to primary biogenic aerosols in the atmosphere: wet and dry discharged spores, carbohydrates, and inorganic ions. *Atmospheric Chemistry and Physics* 7 (17):4569–4588. doi:10.5194/acp-7-4569-2007.

[33] **Nakagawa S, Cuthill IC**. 11 2007. Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews* 82 (4):591–605. doi: 10.1111/j.1469-185X.2007.00027.x.

[34] **Lestrade PP, Bentvelsen RG, Schauwvlieghe AFAD, Schalekamp S, van der Velden WJFM, Kuiper EJ, van Paassen J, van der Hoven B, van der Lee HA, Melchers WJG, de Haan AF, van der Hoeven HL, Rijnders BJA, van der Beek MT, Verweij PE**. 4

2019. Voriconazole Resistance and Mortality in Invasive Aspergillosis: A Multicenter

Retrospective Cohort Study. *Clinical Infectious Diseases* 68 (9):1463–1471. doi:

10.1093/cid/ciy859.

[35] **Risum M, Hare RK, Gertsen JB, Kristensen L, Rosenvinge FS, Sulim S,**

Abou-Chakra N, Bangsborg J, Røder BL, Marmolin ES, Astvad KMT, Pedersen M,

Dzajic E, Andersen SL, Arendrup MC. 4 2022. Azole resistance in <i>Aspergillus

fumigatus</i> . The first 2-year's Data from the Danish National Surveillance Study,

2018–2020. *Mycoses* 65 (4):419–428. doi:10.1111/myc.13426.

[36] **Verweij PE, Lucas JA, Arendrup MC, Bowyer P, Brinkmann AJF, Denning DW,**

Dyer PS, Fisher MC, Geenen PL, Gisi U, Hermann D, Hoogendijk A, Kiers E,

Lagrou K, Melchers WJG, Rhodes J, Rietveld AG, Schoustra SE, Stenzel K, Zwaan

BJ, Fraaije BA. 12 2020. The one health problem of azole resistance in Aspergillus

fumigatus: current insights and future research agenda. *Fungal Biology Reviews* 34

(4):202–214. doi:10.1016/j.fbr.2020.10.003.

[37] **Lestrade PPA, Buil JB, van der Beek MT, Kuijper EJ, van Dijk K, Kampinga GA,**

Rijnders BJA, Vonk AG, de Greeff SC, Schoffelen AF, van Dissel J, Meis JF,

Melchers WJG, Verweij PE. 7 2020. Paradoxal Trends in Azole-Resistant

<i>Aspergillus fumigatus</i> in a National Multicenter Surveillance Program, the

Netherlands, 2013–2018. *Emerging Infectious Diseases* 26 (7):1447–1455. doi:

10.3201/eid2607.200088.

[38] **Morio F, Aubin GG, Danner-Boucher I, Haloun A, Sacchetto E, Garcia-Hermoso D,**

Bretagne S, Miegeville M, Le Pape P. 8 2012. High prevalence of triazole resistance in

Aspergillus fumigatus, especially mediated by TR/L98H, in a French cohort of

patients with cystic fibrosis. *Journal of Antimicrobial Chemotherapy* 67 (8):1870–1873. doi:

10.1093/jac/dks160.

[39] **Steinmann J, Hamprecht A, Vehreschild MJGT, Cornely OA, Buchheidt D, Spiess**

B, Koldehoff M, Buer J, Meis JF, Rath PM. 5 2015. Emergence of azole-resistant

invasive aspergillosis in HSCT recipients in Germany. *Journal of Antimicrobial*

Chemotherapy 70 (5):1522–1526. doi:10.1093/jac/dku566.

[40] **Zhang J, Jimenez LL, Snelders E, Debets AJM, Rietveld AG, Zwaan BJ, Verweij PE,**
Schoustra SE. 1 2020. Dynamics of Aspergillus fumigatus in Azole Fungicide-
Containing Plant Waste in the Netherlands (2016-2017). *Applied and Environmental*
Microbiology 87 (2):1–12. doi:10.1128/AEM.02295-20.

[41] **Barber AE, Riedel J, Sae-Ong T, Kang K, Brabetz W, Panagiotou G, Deising HB,**
Kurzai O. 12 2020. Effects of Agricultural Fungicide Use on Aspergillus fumigatus
Abundance, Antifungal Susceptibility, and Population Structure. *mBio* 11 (6). doi:
10.1128/mBio.02213-20.

[42] **Tsitsopoulou A, Posso R, Vale L, Bebb S, Johnson E, White PL**. 6 2018.
Determination of the Prevalence of Triazole Resistance in Environmental Aspergillus
fumigatus Strains Isolated in South Wales, UK. *Frontiers in Microbiology* 9 (JUN):1–8.
doi:10.3389/fmicb.2018.01395.

[43] **Bromley MJ, van Muijlwijk G, Fraczek MG, Robson G, Verweij PE, Denning DW,**
Bowyer P. 12 2014. Occurrence of azole-resistant species of Aspergillus in the UK
environment. *Journal of Global Antimicrobial Resistance* 2 (4):276–279. doi:
10.1016/j.jgar.2014.05.004.

[44] **Schoustra SE, Debets AJM, Rijs AJMM, Zhang J, Snelders E, Leendertse PC,**
Melchers WJG, Rietveld AG, Zwaan BJ, Verweij PE. 7 2019. Environmental Hotspots
for Azole Resistance Selection of Aspergillus fumigatus, the Netherlands. *Emerging*
Infectious Diseases 25 (7):1347–1353. doi:10.3201/eid2507.181625.

[45] **Shelton JMG, Fisher MC, Singer AC**. 2020. Campaign-Based Citizen Science for
Environmental Mycology: The Science Solstice and Summer Soil-Stice Projects to
Assess Drug Resistance in Air- and Soil-Borne Aspergillus fumigatus. *Citizen Science:*
Theory and Practice 5 (1):1–13. doi:10.1101/2020.06.11.146241.

[46] **Fraczek MG, Zhao C, Dineen L, Lebedinec R, Bowyer P, Bromley M, Delneri D**. 9
2019. Fast and Reliable PCR Amplification from <i>Aspergillus fumigatus</i> Spore
Suspension Without Traditional DNA Extraction. *Current Protocols in Microbiology* 54
(1). doi:10.1002/cpmc.89.

[47] **R Core Team**. 2023. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria .

[48] **Kleiber C, Zeileis A**. 2008. Applied Econometrics with R. Springer New York, New York, NY. doi:10.1007/978-0-387-77318-6.

[49] **Mazerolle MJ**. 2023. AICcmodavg: Model selection and multimodel inference based on (Q)AIC(c) https://cran.r-project.org/package=AICcmodavg.

[50] **Kothe S**. 2023. cmsafops: Tools for CM SAF NetCDF Data https://CRAN.R-project.org/package=cmsafops.

[51] **Wickham H, François R, Henry L, Müller K**. 2018. dplyr: A Grammar of Data Manipulation https://cran.r-project.org/package=dplyr.

[52] **Lenth R**. 2018. emmeans: Estimated Marginal Means, aka Least-Squares Means. https://cran.r-project.org/package=emmeans.

[53] **Villanueva RAM, Chen ZJ**. 7 2019. ggplot2: Elegant Graphics for Data Analysis (2nd ed.). *Measurement: Interdisciplinary Research and Perspectives* 17 (3):160–167. doi:10.1080/15366367.2019.1565254.

[54] **Kassambara A**. 2017. ggpubr:"ggplot2" based publication ready plots. R package version 0.1.6 .

[55] **Bates D, Mächler M, Bolker B, Walker S**. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67 (1). doi:10.18637/jss.v067.i01.

[56] **Venables BD W N Ripley**. 2002. Modern Applied Statistics with S. Fourth ed Springer, New York.

[57] **Hothorn T, Bretz F, Westfall P**. 6 2008. Simultaneous Inference in General Parametric Models. *Biometrical Journal* 50 (3):346–363. doi:10.1002/bimj.200810425.

[58] **Graves S SLGSDRS Piepho H**. 2023. multcompView: Visualizations of Paired Comparisons https://CRAN.R-project.org/package=multcompView.

[59] **Pierce D**. 2023. ncdf4: Interface to Unidata netCDF (Version 4 or Earlier) Format Data Files https://CRAN.R-project.org/package=ncdf4.

[60] **Bivand R RB Keitt T**. 2023. rgdal: Bindings for the 'Geospatial' Data Abstraction
Library. https://CRAN.R-project.org/package=rgdal .

[61] **Wickham H GM Vaughan D**. 2023. tidyr: Tidy Messy Data
https://CRAN.R-project.org/package=tidyr.

[62] **Massicotte P SA**. 2023. rnaturalearth: World Map Data from Natural Earth
https://CRAN.R-project.org/package=rnaturalearth.

[63] **South A**. 2017. rnaturalearthdata: World Vector Map Data from Natural Earth Used
in 'rnaturalearth' https://CRAN.R-project.org/package=rnaturalearthdata.

[64] **Hijmans R**. 2023. raster: Geographic Data Analysis and Modeling
https://CRAN.R-project.org/package=raster.

[65] **Pebesma E**. 2018. Simple Features for R: Standardized Support for Spatial Vector
Data. *The R Journal* 10 (1):439. doi:10.32614/RJ-2018-009.

[66] **Pebesma E, Bivand R**. 5 2023. Spatial Data Science. Chapman and Hall/CRC, New
York. doi:10.1201/9780429459016.