Transition to transversion ratio estimation by accounting codon degeneracy

and pretermination codons in coding sequences

Pratyush Kumar Beura[1], Ruksana Aziz[1], Piyali Sen[2], Saurav Das[1], Nima Dondu Namsa[1,3],

Edward J Feil[4], Siddharatha Sankar Satapathy*[2,3], Suvendra Kumar Ray[*1,3]

Department of [1]Molecular Biology and Biotechnology, [2]Computer Science and Engineering,

[3]Centre for Multidisciplinary Research, Tezpur University, Tezpur, Assam, India, 784028, [4]The

Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath

BA2 7AY, United Kingdom

For correspondence*

Siddhartha Shankar Satapathy: ssankar@tezu.ernet.in

Suvendra Kumar Ray: suven@tezu.ernet.in

ABSTRACT

Degeneracy in the genetic code table influences the transition to transversion ratio in the coding

sequences because a *two-fold degenerate* codon undergoes synonymous changes only by

transition while a *four-fold degenerate* codon undergoes the same by transition as well as

transversion. In addition, there are eighteen pretermination codons in the table where non-sense

substitution is rare to observe. Therefore, the simple estimation of transition to transversion ratio

$(\frac{ti}{tv})$ is inappropriate to compare among coding sequences in a genome. Here, we developed a

method to estimate transition to transversion ratio in coding sequences by accounting codon

degeneracy as well as pretermination codons. In an analysis of 2516 genes in *Escherichia coli* a

distinct observation of $\frac{ti}{tv}$ difference across the coding sequences and the influence of codon

degeneracy and pretermination codon was observed. Interestingly, coding sequences were found

to be variable regarding synonymous $\frac{ti}{tv}$ values, which might be attributed to their intrinsic nature.

The proposed method can be used to estimate different $\frac{ti}{tv}$ ratios in genes.

INTRODUCTION

The genetic code table is a beautiful gift of nature to study molecular evolution (1,2). The

different degenerate codons influence the transition (*ti*) to transversion (*tv*) ratio ($\frac{ti}{tv}$) as follows:

synonymous variation in a *two-fold degenerate* (TFD) codon occurs *via* only one transition

whereas the same in a *four-fold degenerate* (FFD) codon occurs via one transition and two

transversions (Supplementary Figure S1A). Therefore, in any organism, the $\frac{ti}{tv}$ ratio is likely to

vary across the coding sequences having differences in degenerate codon composition. In

addition to the above, compositional variation of pretermination codons (PTC) (Supplementary

Figure S1B and S1C) (3) among the coding sequences is also likely to influence the $\frac{ti}{tv}$ ratio

because non-synonymous variations leading to termination codons are rapidly purged out of the

population by purifying selection (4-6). Considering in general, a *ti* being more frequent than a *tv*

(7-13) which is known as Kimura's two parameter model (14), and a stronger purifying selection

on non-synonymous variation than a synonymous variation (15), the $\frac{ti}{tv}$ ratios of TFD codons and

FFD codons are expected to be different. Accordingly, it has been recently reported that the

synonymous variation in TFD codons is relatively higher than the FFD codons in *Escherichia*

coli (16). Therefore, a comparative study of $\frac{ti}{tv}$ among coding sequences in a genome will be

always insightful.

Figure 1 accurately elucidates the two types of base variations in genomic DNA (17).

The ratio of *ti* to *tv* is theoretically expected to be 0.5. However, the ratio is generally observed

around 2.0 in genomes (7,8). Factors such as geometry of DNA double helix (18), modification

of DNA bases such as deamination of cytosine and adenine (19-21), secondary structure in the

RNA (9) and codon degeneracy are known to elevate *ti* phenomenon in a genome over *tv* (16,22).

In a standard genetic code table out of 549 total single nucleotide variations (SNVs) involving 61

sense codons (9 SNVs per codon), only 134 SNVs are synonymous (24.4%). Among

synonymous (S*ti* and S*tv*) and non-synonymous (N*ti* and N*tv*) variations, the theoretically

estimated ($_e$) numbers of S*ti*, S*tv*, N*ti* and N*tv* in a genetic codon table are 62, 72, 121 and 294,

respectively (see Supplementary Table S1). Yet the observed synonymous polymorphisms are

always higher than the expected proportion across genes (23-26) as purifying selection is

stronger on non-synonymous polymorphisms than the synonymous polymorphism (15). Hence,

by accounting for the polymorphism patterns affected by codon degeneracy as well as PTC, now

the theoretically estimated polymorphisms can be re-calculated in the form of S*ti*, S*tv*, N*ti* and

N*tv* for each codon (Table 1). Also, we have summarized the total numbers of theoretically

estimated synonymous *ti* and *tv* and non-synonymous *ti* and *tv* in the genetic codon table (Table

2).

The codon substitution model suggests that sites within codons evolve at different rates

and consequently they should not be equally treated (22,27). In the codon substitution model,

using likelihood approach the $\frac{ti}{tv}$ ratio was estimated in accordance with codon degeneracy class

which was implemented in estimating *dN/dS* in coding sequences (22,28,29). Recently, by

accounting $\frac{ti}{tv}$ substitution rate difference in codons as well as nonsense variations in PTC, the

dN/dS values have been improved (16). This finding implies that factors such as codon

degeneracy and PTC composition should be considered while analyzing the evolution of protein-

coding genes. The ratio of *ti* to *tv* has been studied for reliable estimation of sequence distance

and phylogeny reconstruction (30). Using maximum likelihood approach, the $\frac{ti}{tv}$ ratio (commonly

known as kappa *κ*) has been used to mainly compare the $\frac{ti}{tv}$ values among different species (30).

The parameter *κ* has not been implemented to compare coding sequences within a genome in the

context of their compositional differences of different degenerate classes before. As the selective

pressure on synonymous and non-synonymous variations are heterogeneous (31), it is important

to analyze the $\frac{ti}{tv}$ values among different genes within an organism by accounting synonymous

sites and non-synonymous sites regarding their compositional differences of different degenerate

codons.

Given that there is a tremendous variation of substitution rates among protein coding

sequences within a genome (32), an appropriate methodology for the estimation of *ti* to *tv* ratio is

required. We are presenting an improved equation that builds upon the research conducted by Ina

in 1998 (33). Considering the differential selective constraint in synonymous and non-

synonymous variations it is crucial to separately analyze the synonymous $\frac{ti}{tv}$ and non-

synonymous $\frac{ti}{tv}$ to gain a comprehensive understanding of the overall $\frac{ti}{tv}$. The improved equations

used in this study are helpful to estimate the *ti* and *tv* rate differences in coding sequences by

accounting the compositional difference in terms of codon degeneracy and PTC. Though it is

known that TFD codons only undergo synonymous *ti*, the $\frac{Sti}{Stv}$ ratio among the coding sequences

in relation to their compositional differences (TFD, FFD and PTC) has not been reported

previously. Further, we observed a correlation between TFD: FFD and $\frac{Sti}{Stv}$ values and the PTC%

and $\frac{Nti}{Ntv}$ values. We tried to establish the biological significance of codon composition and $\frac{ti}{tv}$ ratio

across 2516 coding sequences of *E. coli*. The improved method developed in the present study

may be applied to study different $\frac{ti}{tv}$ ($\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$) across coding sequences.

MATERIALS AND METHODS

Coding sequences of *E. coli* genome

In this study, we carried out a SNVs based computational analysis of 2516 coding sequences

across 157 strains of *E. coli* (34). The information regarding the dataset selected for the present

study is provided in Supplementary Table S2. Although the public database has ~4500 coding

sequences, we excluded coding sequences showing length variation and improper

alignment/annotation. Codons having variations in more than one position or variations leading

to ambiguous nucleotides were not considered in the analyses of the variations in this study. A

detailed procedure for estimating variation is represented in (see Supplementary Table S3) using

example of a hypothetical sequence. This procedure is based on intra-species genome sequence

comparison to find out single nucleotide variations in bacterial genomes. A reference sequence

has been generated considering the most frequent nucleotide occurrence at a site in the gene

sequence. The logic is that the most abundant nucleotide in a species is the ancestral nucleotide

in the species (9,16,35,36).The observed SNVs (SNV$_o$) at different positions of each codon were

categorized into *Sti$_o$*, *Stv$_o$*, *Nti$_o$* and *Ntv$_o$*, where S for synonymous, N for non-synonymous, *ti$_o$* for

observed transition and *tv$_o$* for observed transversion, and subsequently *ti$_o$* (*Sti$_o$* $_+$ *Nti$_o$*) and *tv$_o$*

(*Stv$_{o+}$* *Ntv$_o$*) for each gene were calculated (see Supplementary Table S4). After finding observed

variation values for a gene, we calculated the theoretically estimated variation values *Sti$_e$*, *Stv$_e$*

Nti$_e$, *Ntv$_e$*, *ti$_e$* and *tv$_e$*. We derived the reference sequence for each gene based on the methodology

explained (see Supplementary Table S3). The reference sequences were further used to find the

codon counts of each gene. Then we multiplied the count of each codon by the theoretically

estimated variation values for each codon. For example, if the codon frequency of GGG is

observed to be 10 in a gene, then the total theoretically estimated value for *Sti*, *Stv*, *Nti* and *Ntv*

are 10, 20, 20 and 40 (as estimated *Sti*, *Stv*, *Nti* and *Ntv* for each GGG is 1, 2, 2 and 4,

respectively). We then summed up the individual variation values for 61 sense codons to get the

total theoretically estimated individual substitution values (Sti_e, Stv_e, Nti_e and Ntv_e) for the gene

(see Supplementary Table S5). The values of Sti_o, Stv_o, Nti_o, Ntv_o, ti_o, and tv_o and the values of

Sti_e, Stv_e Nti_e, Ntv_e, ti_e, and tv_e of a coding sequence are then applied in the modified equation to

find out different $\dfrac{ti}{tv}$ ratio (see Supplementary Table S6). A detailed workflow is provided in

Figure 2.

Improved equations

We proposed an improved mathematical equation for the calculation of different $\dfrac{ti}{tv}$ ratio as

described below.

$$\frac{ti'}{tv'} = \frac{\left(ti_o/ti_e\right)}{\left(tv_o/tv_e\right)}$$ **Equation 1**

$$\frac{Sti'}{Stv'} = \frac{\left(Sti_o/Sti_e\right)}{\left(Stv_o/Stv_e\right)}$$ **Equation 2**

$$\frac{Nti'}{Ntv'} = \frac{\left(Nti_o/Nti_e\right)}{\left(Ntv_o/Ntv_e\right)}$$ **Equation 3**

Where, $\dfrac{ti'}{tv'}$ is improved $\dfrac{ti}{tv}$ ratio, $\dfrac{Sti'}{Stv'}$ is improved $\dfrac{Sti}{Stv}$ ratio, $\dfrac{Nti'}{Ntv'}$ is improved $\dfrac{Nti}{Ntv}$ ratio, ti_o is transition

observed, ti_e is the theoretically estimated transition, tv_o is transversion observed, tv_e is

transversion estimated, Sti_o is synonymous transition observed, Sti_e is the theoretically estimated

synonymous transition, Stv_o is synonymous transversion observed, Stv_e is the theoretically

estimated synonymous transversion, Nti_o is non-synonymous transition observed, Nti_e is the

theoretically estimated non-synonymous transition, Ntv_o is non-synonymous transversion observed,

Ntv_e is the theoretically estimated non-synonymous transversion. We calculated $\frac{ti}{tv}$ ratios for all the

genes using a program written in the C language. (to be modified later)

134   It is noteworthy that in a genome the number of potential replacements is greater for non-

synonymous alterations, whereas the frequency of actual variations is higher for synonymous

alterations among coding sequences. The improved estimators $\frac{ti'}{tv'}$, $\frac{Sti'}{Stv'}$, and $\frac{Nti'}{Ntv'}$ account for the

observed variations out of the total possible variations, on the contrary, the conventional method

only takes an account of the *ti_o* and *tv_o*. The possible sites for *Sti* and *Stv* are also different among

genes, which is dependent upon the codon composition belonging to different degeneracy classes.

The conventional $\frac{ti}{tv}$ ratio does not adequately account for the limitations of variable selection

against synonymous and non-synonymous variations. Hence, all these parameters make the

requirement of an improved estimator instead of the conventional estimator.

Comparative representation of different $\frac{ti}{tv}$ through conventional and improved method

We have done a comparative analysis of $\frac{ti}{tv}$ calculated using the existing and the improved

method to support the validation of method proposed in this study. Also, we have calculated the

TFD: FFD and the percentage of PTC (PTC%) of each gene and performed statistical analysis

between different parameters. A summary of the different parameters observed throughout the

dataset is represented in Table 3. We also calculated the percentage (%) change in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$

values obtained from conventional to improved method as follows.

(a) $$\frac{Sti}{Stv} \text{ \% change} = \frac{\frac{Sti}{Stv} - \frac{Sti'}{Stv'}}{\frac{Sti}{Stv}} \times 100$$

(b) $$\frac{Nti}{Ntv} \text{ \% change} = \frac{\frac{Nti}{Ntv} - \frac{Nti'}{Ntv'}}{\frac{Nti}{Ntv}} \times 100$$

We found out $\frac{ti}{tv}$ in all these 2516 genes using maximum likelihood (ML) approach

available in MEGA-X (37).

Statistical analyses

Comparison among the conventional, ML approach and the improved $\frac{ti}{tv}$ values were done using

different statistical methods to demonstrate the performance and accuracy of the improved

method. OriginPro, Version 2022, OriginLab Corporation, Northampton, MA, USA was used to

draw the Box-plot as well as to perform the Mann-Whitney test (38) to find out the *p*-value. The

correlation plot was drawn for R^2 value and Pearson's correlation coefficient (39) was also

calculated between the $\frac{ti}{tv}$ values obtained using the conventional and improved method.

RESULTS

Compositional analysis of TFD, FFD, and PTC codons in 2516 coding sequences of *E. coli*

We found out the total number of TFD, FFD, and PTC codons across 2516 coding sequences of *E.*

coli. The percentage of TFD, FFD, and PTC was calculated (see Supplementary Table S2). The

percentage (%) compositional differences in the 2516 coding sequences represented as TFD%,

FFD%, and PTC% is shown in a box-plot (Figure 3). A variable range of TFD: FFD ratio in the

coding sequences of *E. coli* was observed. The minimum to maximum TFD:FFD ratio was observed

as 0.229 and 3.333 in *sugE* and *mntS*, respectively. It suggests that FFD codons are present

approximately four times more frequent than TFD codons in certain coding sequences. Similarly,

TFD codons are present approximately three times more frequent than FFD codons in the selected

coding sequences. Among 2516 coding sequences, only 35 coding sequences possessed the

TFD:FFD ratio of 1.000. Hence, this observation suggests that there is a compositional asymmetry

of TFD and FFD codons in the coding sequences. The minimum to maximum PTC% was observed

in the gene *atpE* as 15.00% and in the gene *chaB* as 57.14% (Figure 3). The compositional variation

(TFD, FFD and PTC) observed in coding sequences were considered to validate the accuracy of

different $\frac{ti}{tv}$ ratio using the conventional and improved method.

Impact of codon degeneracy and PTC on $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ values

A gene entirely composed of FFD codons will result into a twice $\frac{Sti'}{Stv'}$ value than $\frac{Sti}{Stv}$ value, similarly

a gene having equal proportion of FFD codons and TFD codons will result into equal values of $\frac{Sti'}{Stv'}$

and $\frac{Sti}{Stv}$. The increase in TFD codon proportion will directly impact the difference between $\frac{Sti}{Stv}$ values

and $\frac{Sti'}{Stv'}$ values. We have observed a similar pattern as well (Supplementary Figure S2). The

complete set of observed and estimated values of different variations is enlisted (see Supplementary

Table S2). These values were used to calculate the $\frac{ti}{tv}, \frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$. Further the improved equation

was used to calculate the $\frac{ti'}{tv'}, \frac{Sti'}{Stv'}$ and $\frac{Nti'}{Ntv'}$. We compared the values derived using the conventional

method ($\frac{ti}{tv}, \frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$) and improved method ($\frac{ti'}{tv'}, \frac{Sti'}{Stv'}$ and $\frac{Nti'}{Ntv'}$) (see Supplementary Table S2). A

box-plot showing the comparison between the values of different $\frac{ti}{tv}$ ratio obtained through the

conventional and improved method is represented in Figure 4. The minimum % change in $\frac{Sti}{Stv}$

between conventional and improved method was observed in the gene *garP* as -30.99% and

maximum in the gene *yfhL* as 30.76%. Similarly, the minimum % change in $\frac{Nti}{Ntv}$ between

conventional and improved method was observed in the gene *thiS* as -194.675% and maximum in

the gene *bhsA* as -129.600%. The detailed % change in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ is represented (see

Supplementary Table S2). While it is evident that, TFD codons and FFD codons have different

probabilities of obtaining synonymous variations, the composition of TFD: FFD was cross-checked

against the % change in $\frac{Sti}{Stv}$ for each gene. Likewise, the impact of PTC% in estimating the $\frac{Nti}{Ntv}$

values have been considered in this study.

The synonymous variation in the genes is likely to be different considering the differential

composition of TFD codons and FFD codons, because when FFD codon proportion is higher in a

gene, $\frac{ti}{tv}$ value is going to be lower. Similarly, when the FFD codon proportion is lower, $\frac{ti}{tv}$ value is

going to be higher. So, this observation is distinct, and the improved estimator is taking care of the

compositional differences (estimated numbers) of different degenerate codons. To show that the

compositional difference is indeed influencing the $\frac{ti}{tv}$ ratio, we have performed the correlation

analyses between TFD: FFD and % change in $\frac{Sti}{Stv}$ as well as between PTC% and % change in $\frac{Nti}{Ntv}$.

We have also observed this similar result in our simulation study.　The Pearson correlation

coefficient value between TFD: FFD and % change in $\frac{Sti}{Stv}$ (conventional to improved) was obtained

to be 0.891, signifying a strong correlation coefficient. Similarly, the Pearson correlation coefficient

value between PTC% and % change in $\frac{Nti}{Ntv}$ was obtained to be -0.488, a moderate negative

correlation between the two variables. The strong correlation between TFD: FFD and % change in

$\frac{Sti}{Stv}$ suggested that there is an influence of the TFD: FFD codons in overall synonymous $\frac{ti}{tv}$. Similarly,

the moderate negative correlation also suggested the contribution of PTC codons in overall non-

synonymous $\frac{ti}{tv}$. The results were found to be statistically different at $p<0.01$, for $\frac{ti}{tv}$ & $\frac{ti'}{tv'}$, $\frac{Sti}{Stv}$ & $\frac{Sti'}{Stv'}$

and $\frac{Nti}{Ntv}$ & $\frac{Nti'}{Ntv'}$. Further the correlation graph between % change in $\frac{Sti}{Stv}$ and TFD: FFD and % change

in $\frac{Nti}{Ntv}$ and PTC% also substantiate the role of codon composition and $\frac{ti}{tv}$ ratio (Figure 5A & 5B). In

the trendline analyses the % change in $\frac{Sti}{Stv}$ was perceived to be directly influenced by the TFD: FFD

with a R^2 value of 0.795. A weak negative association was obtained between % change in $\frac{Nti}{Ntv}$ and

PTC% with a R^2 value of 0.216. Hence, it is pertinent to note that the overall % change in $\frac{Sti}{Stv}$ and

$\frac{Nti}{Ntv}$ were directly influenced by the TFD and PTC codon composition across the coding sequences.

Therefore, the findings of correlation analysis further support the contribution of TFD: FFD and

PTC% in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ respectively. We showed that the role of codon composition in estimating the

$\frac{ti}{tv}$ ratio using the improved method and this parameter was not accounted in the conventional

method. We performed a simulation study to substantiate the impact of the compositional difference

of degeneracy class in estimating $\frac{Sti}{Stv}$ by using our improved method in coding sequences. For

detailed methodology and results (see Supplementary Table S7, Supplementary Figure S3). The

results of comparative analysis between ML based method and improved method are presented in

supplementary files for 2516 genes (see Supplementary Table S8, Supplementary Figure S4).

DISCUSSION

Various methods have been used in the estimation of the *ti* to *tv* ratio by molecular evolutionary

biologists to understand the mutation as well as selection bias. The phylogeny-based method and

Bayesian methods have been implemented by researchers in the inter-species study of $\frac{ti}{tv}$ (40,41).

The coding sequences within a genome are highly variable in terms of compositions of different

degenerate codons and PTC. It suggests that while estimating $\frac{ti}{tv}$ in coding sequences within a

genome, compositional difference of different degenerate class as well as PTC should be

considered. A species-specific dependence of the deleterious nature of *Nti* and *Ntv* in coding

sequences has been reported recently and the plausible contribution of PTC was not accounted

for in the study (42). However, our observation in the form of codon degeneracy and PTC in

estimating the $\frac{ti}{tv}$ ratio is in concordance with the recent findings on the role of codon degeneracy

and PTC in coding sequences to estimate *dN/dS* values (16). Considering synonymous variations

are more abundant than the non-synonymous variations, comparison of coding sequences

regarding $\frac{ti}{tv}$ separately for synonymous and non-synonymous sites is important. In the present

work, we have extended the work of Ina 1998 (33) and developed modified estimators to

separately calculate $\frac{ti}{tv}$ at synonymous and non-synonymous sites. The new estimators consider

the compositional differences among the coding sequences in terms of codon degeneracy and

PTC.

The interesting observation came out from our study is the large variation across the

coding sequences regarding their $\frac{Sti}{Stv}$ even after accounting for the codon degeneracy and PTC. It

might be the result of the non-identical 12 substitution patterns as follows. The C→T is known as

the most frequent change that occur among the *ti* variation. Among the *tv*, G→T or C→A change

are the most frequently observed variation in the genome. So, in the future, the contribution of

different mutation rates among the nucleotides will be more insightful in this context. There has

been recent description of context-dependent mutation in organism which might induce different

ti and *tv* in coding sequences (43-46). We did a correlation study between the gene expression

study using the CAI value against the $\frac{Sti}{Stv}$ values, but we could not observe any strong correlation.

Therefore, our study agrees that the role of intrinsic contributing factors across the coding

sequences might attribute to the difference in $\frac{ti}{tv}$ values.

Overall, our method is simple, succinct, easy to follow for researchers in the field with

limitations in mathematics and provides an insight to understand the evolution of protein coding

sequences. We also did a comparison with the ML method. There are several genes where we

could observe the $\frac{ti}{tv}$ values estimated by the improved method is more in concordance with the

observed *ti* and *tv* values than the ML method. It has been said that the likelihood methods are

computer intensive and difficult to apply to large datasets and can fall into local traps (47).

Furthermore, the extrapolation of our method in various inter-species studies, as well as

phylogenetic tools can enhance the horizon of our understanding regarding mutational bias in

coding sequences. However, the differential usage of amino acids, gene expression, and the

evolutionary significance of codon degeneracy and PTC composition is yet to be investigated.

DATA AVAILABILITY

265 The authors confirm that the data supporting the findings of this study are available within the

266 article and Supplementary Data.

ACKNOWLEDGMENTS

PKB is grateful to Tezpur University for the institutional fellowship. RA is thankful for the

JRF fellowship from the DBT grant (BT/511/NE/TBP/2013) and (BT/403/NE/U-Excel/2013). PS is

grateful to UGC, GoI New Delhi, for the JRF. SD is thankful to DBT for MSc. Scholarship. SSS and

SKR are thankful to DBT, GoI for the twinning grant (BT/PR16361/NER/95/192/2015 date 18-10-

2016) to them. NDN, SSS & SKR thankfully acknowledge the DBT funded Bioinformatics and

Computational Biology Centre at Tezpur University.

Authors contributions

P.K.B, S.S.S. and S.K.R: Idea, design, executed, data analysis, writing, and discussion; P.S

and R.A.: executed, data analysis, and writing; S.D: writing and discussion. N.D.N: Writing

and Discussion; E.F; Writing, Discussion, and critical analysis.

Author Disclosure Statement

The authors declare no potential conflict of interest. The authors also declare no competing

financial interests.

FUNDING

The authors received no specific funding for this work.

REFERENCE

1. Crick, F.H.C., Barnett, L., Brenner, S. and Watts-Tobin, R.J. (1961) General nature of

the genetic code for proteins. *Nature*, **30**;192:1227-32. https://doi.org/ 10.1038/1921227a0.

2. Khorana, H.G., Büchi, H., Ghosh, H.,Gupta, N., Jacob, T. M., Kössel, H., Morgan, R.,

Narang, S.A., Ohtsuka, E. and Wells, R.D. (1966) Polynucleotide synthesis and the genetic code. Cold Spring Harbor Symp Quant Biol. **31**:39-49. https://doi.org/10.1101/sqb.1966.031.01.010.

3. Modiano, G., Battistuzzi, G. and Motulsky, A.G. (1981) Nonrandom patterns of codon usage and of nucleotide substitutions in human alpha-and beta-globin genes: an evolutionary strategy reducing the rate of mutations with drastic effects? *Proc Natl Acad Sci,* **78**:1110–1114. https://doi.org/ 10.1073/pnas.78.2.1110

4. van den Eynden, J., Basu, S. and Larsson, E. (2016) Somatic mutation patterns in hemizygous genomic regions unveil purifying selection during tumor evolution. *PLoS Genet,* **12**: e1006506. https://doi.org/10.1371/journal.pgen.1006506

5. Morales, A.C., Rice, A.M., Ho, A.T.l., Mordstein, C., Mühlhausen, S., Watson, S., Cano, L., Young, B., Kudla, G. and Hurst, L.D. (2021) Causes and consequences of purifying selection on SARS-CoV-2. *Genome Biol Evol,* **13**: evab196. https://doi.org/10.1093/gbe/evab196

6. Li, W-H., Gojobori, T. and Nei, M. (1981) Pseudogenes as a paradigm of neutral evolution. *Nature,* **292**:237–239. https://doi.org/10.1038/292237a0

7. Duchêne, S., Ho, S.Y.W. and Holmes, E.C. (2015) Declining transition/transversion ratios through time reveal limitations to the accuracy of nucleotide substitution models. *BMC Evol Biol,* **15**:1–10. https://doi.org/10.1186/s12862-015-0312-6

8. Stoltzfus, A. and Norris, R.W. (2016) On the causes of evolutionary transition: transversion bias. *Mol Biol Evol,* **33**:595–602. https://doi.org/10.1093/molbev/msv274

9. Sen, P., Aziz, R., Deka, R.C., Feil. E. J., Ray, S.K. and Satapathy S.S. (2022) Stem region

of tRNA genes favors transition substitution towards keto bases in bacteria. *J Mol Evol,*

90:114–123. https://doi.org/10.1007/s00239-021-10045-x

10. Seplyarskiy, V.B., Kharchenko, P., Kondrashov, A.S. and Bazykin, G.A. (2012)

Heterogeneity of the transition/transversion ratio in Drosophila and Hominidae genomes.

Mol Biol Evol, **29**:1943–1955. https://doi.org/10.1093/molbev/mss071

11. Lewis Jr, C.A., Crayle, J., Zhou, S. Swanstrom, R. and Wolfenden, R. (2016) Cytosine

deamination and the precipitous decline of spontaneous mutation during Earth's history.

Proc Natl Acad Sci, **113**:8194–8199. https://doi.org/ 10.1073/pnas.1607580113.

12. Lyons, D.M. and Lauring, A.S. (2017) Evidence for the selective basis of transition-to-

transversion substitution bias in two RNA viruses. *Mol Biol Evol,* **34**:3205–3215.

https://doi.org/ 10.1093/molbev/msx251

13. Schroeder, J.W., Randall, J.R., Hirst, W.G., O'Donnel, M.E. and Simmons, L.A. (2017)

Mutagenic cost of ribonucleotides in bacterial DNA. *Proc Natl Acad Sci,* **114**:11733–

11738. https://doi.org/10.1073/pnas.1710995114

14. Kimura, M. (1980) A simple method for estimating evolutionary rates of base substitutions

through comparative studies of nucleotide sequences. *J Mol Evol,* **16**:111–120.

https://doi.org/ 10.1007/BF01731581

15. Hurst, L.D. and Pál, C. (2001) Evidence for purifying selection acting on silent sites in

BRCA1. *TRENDS in Genetics,* **17**:62–65. https://doi.org/ 10.1016/s0168-9525(00)02173-9

16. Aziz, R., Sen, P., Beura, P.K., Das, S., Tula, D., Dash, M., Namsa, N.D.N., Deka, R.C.,

Feil, E.J. Satapathy, S.S. *et al* (2022) Incorporation of transition to transversion ratio and

nonsense mutations, improves the estimation of the number of synonymous and non-

synonymous sites in codons. *DNA Research,* **29**: dsac023. https://doi.org/10.1093/dnares/dsac023

17. Fersht, A.R. and Knill-Jones, J.W. (1981) DNA polymerase accuracy and spontaneous mutation rates: frequencies of purine. purine, purine. pyrimidine, and pyrimidine. pyrimidine mismatches during DNA replication. *Proc Natl Acad Sci,* **78**(7):4251–4255.https://doi.org/10.1073/pnas.78.7.4251

18. Topal, M.D. and Fresco, J.R. (1976) Complementary base pairing and the origin of substitution mutations. *Nature,* **263**(5575):285-289. https://doi.org/10.1038/263285a0.

19. Kino K, Sugiyama H (2001) Possible cause of G. C> C. G transversion mutation by guanine oxidation product, imidazolone. *Chem. Biol*, 8, 369–378. https://doi.org/10.1016/S1074-5521(01)00019-9

20. Rocha, E.P.C., Touchon, M. and Feil. E. J. (2006) Similar compositional biases are caused by very different mutational effects. *Genome Res,* **16**, 1537–1547. https://doi.org/10.1101/gr.5525106

21. Bhagwat, A. S., Hao, W., Townes, J.P., Lee, H., Tang, H. and Foster, P.L. (2016) Strand-biased cytosine deamination at the replication fork causes cytosine to thymine mutations in Escherichia coli. *Proc Natl Acad Sci,* **113**, 2176–2181. https://doi.org/10.1073/pnas.1522325113

22. Muse, S.V. and Gaut, B.S. (1994) A Likelihood Approach for Comparing Synonymous and Nonsynonymous Nucleotide Substitution Rates, with Application to the Chloroplast Genome. *Mol Bio Evol,* **1** **1(5)**:7 15-724. https://doi.org/10.1093/oxfordjournals.molbev.a040152

23. Tamura, K. (1992) The rate and pattern of nucleotide substitution in Drosophila
mitochondrial DNA. *Mol Biol Evol,* **9**:814–825.
https://doi.org/10.1093/oxfordjournals.molbev.a040763

24. Wolfe, K.H., Sharp, P.M. and Li, W-H. (1989) Rates of synonymous substitution in plant
nuclear genes. *J Mol Evol,* **29**:208–211. https://doi.org/10.1007/BF02100204

25. Moriyama, E.N. and Powell, J.R. (1997) Synonymous substitution rates in Drosophila:
mitochondrial versus nuclear genes. *J Mol Evol,* **45**:378–391. https://doi.org/
10.1007/pl00006243

26. Ngandu, N.K., Scheffler, K., Moore, P., Woodman, Z., Martin, D. and Seoighe, C. (2008)
Extensive purifying selection acting on synonymous sites in HIV-1 Group M sequences.
Virol J, **5**:1–11. https://doi.org/ 10.1186/1743-422X-5-160

27. Shapiro, B., Rambaut, A. and Drummond, A.J. (2006) Choosing appropriate substitution
models for the phylogenetic analysis of protein-coding sequences. *Mol Biol Evol,* **23**, 7–9.
https://doi.org/10.1093/molbev/msj021

28. Goldman, N. and Yang, Z. (1994) A codon-based model of nucleotide substitution for
protein-coding DNA sequences. *Mol Biol Evol,* **11**(5), 725-736. https://doi.org/
10.1093/oxfordjournals.molbev.a040153.

29. Arenas, M. (2015) Trends in substitution models of molecular evolution. *Front Genet,*
6:319. https://doi.org/10.3389/fgene.2015.00319

30. Yang, Z. and Yoder, A.D. (1999) Estimation of the Transition/Transversion Rate Bias and
Species Sampling. *J Mol Evol* **48**:274–283. https://doi.org/10.1007/PL00006470

31. Bartolomé. C., Maside. X., Yi. S., Grant. A.L. and Charlesworth, B. (2005) Patterns of

377   selection on synonymous and nonsynonymous variants in Drosophila miranda. *Genetics,*

378   **169**(3):1495-507. https://doi.org/10.1534/genetics.104.033068

379   32. Wu, C-I. and Li, W.H. (1985) Evidence for higher rate of nucleotide substitution in rodents

380   than in man. *Proc Natl Acad Sci,* **82**(6):1741-1745. https://doi.org/ 10.1073/pnas.82.6.1741

381   33. Ina, Y. (1998) Estimation of the transition/transversion ratio. *J Mol Evol,* **46**:521–533.

382   https://doi.org/10.1007/PL00006333

383   34. Thorpe, H.A., Bayliss, S.C. and Hurst, L.D. and Feil, E.J. (2017) Comparative analyses of

384   selection operating on nontranslated intergenic regions of diverse bacterial species.

385   *Genetics,* **206**:363–376. https://doi.org/10.1534/genetics.116.195784

386   35. Sen, P. (2022) Computational Analysis of Codon Usage Bias Single Nucleotide

387   Polymorphism and RNA Secondary Structures in Microbial Genome Sequences. Ph.D.

388   Thesis, Tezpur University http://hdl.handle.net/10603/473607

389   36. Beura, P.K., Sen, P., Aziz, R., Satapathy, S.S. and Ray, S.K. (2023) Transcribed intergenic

390   regions exhibit a lower frequency of nucleotide polymorphism than the untranscribed

391   intergenic regions in the genomes of Escherichia coli and Salmonella enterica. *J Genet,*

392   **102**:22. https://doi.org/10.1007/s12041-023-01418-w

393   37. Kumar, S., Stecher, G., and Li, M., Knyaz, C. and Tamura, K. (2018) MEGAX: Molecular

394   evolutionary genetics analysis across computing platforms. *Mol Biol Evol,* **35**(6):1547–

395   1549 https://doi.org/ 10.1093/molbev/msy096

396   38. Mann, H.B. and Whitney, D.R. (1947) On a test of whether one of two random variables is

397   stochastically larger than the other. *Ann Math Statist,* **18**(1), 50-60. https://doi.org/

398   10.1214/aoms/1177730491

39. Pearson, K. (1896) Mathematical contributions to the theory of evolution. On a form of

spurious correlation which may arise when indices are used in the measurement of organs.

Proc R Soc Lond, **60**(359-367), 489-498. https://doi.org/10.1098/rspl.1896.0076

40. Purvis, A. and Bromham, L. (1997) Estimating the transition/transversion ratio from

independent comparisons with an assumed phylogeny. *J Mol Evol,* **44**, 112–119.

https://doi.org/ 10.1007/pl00006117

41. Huelsenbeck, J. and Ronquist, F. (2001) MrBayes: Bayesian inference of phylogenetic

trees. *Bioinform* **17**, 754–755. https://doi.org/ 10.1093/bioinformatics/17.8.754.

42. Zou, Z. and Zhang, J. (2021) Are Nonsynonymous Transversions Generally More

Deleterious than Nonsynonymous Transitions? *Mol Biol Evol,* **38**(1):181–191.

https://doi.org/ 10.1093/molbev/msaa200

43. Sung, W., Ackerman, M.S., Gout, J.F., Miller, S.F., Williams, Emily., Foster, P.L.

and Lynch, M. (2015) Asymmetric context-dependent mutation patterns revealed through

mutation-accumulation experiments. *Mol Biol Evol.* 32:1672–1683.

doi: 10.1093/molbev/msv055.

44. Zhu, Y., Neeman, T., Yap, V.B. and Huttley. G.A. Statistical methods for identifying

sequence motifs affecting point mutations. *Genetics.* 2017

doi: 10.1534/genetics.116.195677.

45. Aikens. R.C., Johnson. K.E. and Voight. B.F. (2019) Signals of variation in human

mutation rate at multiple levels of sequence context. *Mol Biol Evol.* 36:955–965.

doi: 10.1093/molbev/msz023.

46. Ling. G., Miller, D., Nielsen, R. and Stern. A. A. (2020) Bayesian framework for inferring

the influence of sequence context on point mutations. *Mol Biol Evol.* 37:893–903.

doi: 10.1093/molbev/msz248.

47. Golding, B. and Felsenstein, J. (1990) A maximum likelihood approach to the detection of

selection from a phylogeny. *J Mol Evol,* **31**:511-523. https://doi.org/10.1007/bf02102078

428   **Figure_legend_Table**

Figure 1. Schematic representation of the different substitutions occurring in the DNA. The

transition *(ti)* is marked with dark double headed arrows and *(tv)* is marked with light double headed

arrows. R symbolizes purines (A, G) and Y symbolizes pyrimidines (T, C). Out of the twelve

possible base substitutions, four are *ti* in which a purine (or pyrimidine) nucleotide is replaced by

another purine (or pyrimidine) nucleotide, and eight are *tv* in which a purine (or pyrimidine)

nucleotide is replaced by another pyrimidine (or purine) nucleotide.

Figure 2. Schematic representation of the workflow showing the steps to execute our improved

method. By considering each codon as j, ti_e^j and tv_e^j can be calculated for each codon as presented

in Table 1. The reference sequence needs to be calculated from the coding region (gene/query).

From the reference sequence the observed ($_o$) substitutions are calculated for the gene. The other

usage of the reference sequence is to find out the codon count of the gene. Then, the respective

codon count is multiplied by estimated transition and transversion for each codon. The next step is to

find out the sum of ti and sum of tv in all codons to represent the estimated ($_e$) ti and tv for the gene.

After getting the observed and estimated values for ti and tv, now the improved equation is applied

to find out the $\frac{ti'}{tv'}$ of the gene. This scenario is only showing the $\frac{ti'}{tv'}$ ratio, whereas in similar way the

$\frac{Sti'}{Stv'}$ & $\frac{Nti'}{Ntv'}$ can be calculated.

Figure 3. The box-plot presents the percentage (%) of TFD, FFD and PTC codons in the coding

sequences considered to validate the improved method proposed in this study. The minimum and

maximum TFD% in the dataset was observed as 10.38% and 53.25%, similarly the minimum and

maximum FFD% in the dataset was observed as 13.95% and 52.05%. The minimum and maximum

PTC% in the dataset was observed as 15.00% and 57.14%.

Figure 4. The box-plot presents the different $\frac{ti}{tv}$ ratio values. Here we have compared the different $\frac{ti}{tv}$

ratios of the entire set of genes through conventional and improved method. $\frac{ti}{tv}$ shows the

conventional method and $\frac{ti'}{tv'}$ shows the result of improved method. Similarly, $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ represent

the conventional values and $\frac{Sti'}{Stv'}$ and $\frac{Nti'}{Ntv'}$ represent the improved values. This illustration shows that

the median values between $\frac{ti}{tv}$ & $\frac{ti'}{tv'}$, $\frac{Sti}{Stv}$ & $\frac{Sti'}{Stv'}$ and $\frac{Nti}{Ntv}$ & $\frac{Nti'}{Ntv'}$ are statistically different (p-value <

0.01).

A

B

467 **Figure 5.** Regression plot of coding sequence considered to validate the improved method. The x-

468 axis shows the codon composition (TFD: FFD and PTC%) and y-axis shows the % change in $\frac{Sti}{Stv}$

469 and $\frac{Nti}{Ntv}$. (A) % change in $\frac{Sti}{Stv}$ is showing a strong correlation with the TFD: FFD with a R^2 value of

470 0.795. (B) % change in $\frac{Nti}{Ntv}$ is showing a weak negative correlation with the PTC% with a a R^2 value

471 of 0.216.

Table 1. Theoretically estimated ($_e$) polymorphism for each codon by excluding the non-synonymous

substitutions (*Nti* & *Ntv*) leading to stop codons through PTC

Codon	Sti_e	Stv_e	Nti_e	Ntv_e	Codon	Sti_e	Stv_e	Nti_e	Ntv_e	Codon	Sti_e	Stv_e	Nti_e	Ntv_e	Codon	Sti_e	Stv_e	Nti_e	Ntv_e
UUU	1	0	2	6	UCU	1	2	2	4	UAU	1	0	2	4	UGU	1	0	2	5
UUC	1	0	2	6	UCC	1	2	2	4	UAC	1	0	2	4	UGC	1	0	2	5
UUA	2	0	1	4	UCA	1	2	2	2	UAA	X	X	X	X	UGA	X	X	X	X
UUG	2	0	1	5	UCG	1	2	2	3	UAG	X	X	X	X	UGG	0	0	1	6
CUU	1	2	2	4	CCU	1	2	2	4	CAU	1	0	2	6	CGU	1	2	2	4
CUC	1	2	2	4	CCC	1	2	2	4	CAC	1	0	2	6	CGC	1	2	2	4
CUA	2	2	1	4	CCA	1	2	2	4	CAA	1	0	1	6	CGA	1	3	1	3
CUG	2	2	1	4	CCG	1	2	2	4	CAG	1	0	1	6	CGG	1	3	2	3
AUU	1	1	2	5	ACU	1	2	2	4	AAU	1	0	2	6	AGU	1	0	2	6
AUC	1	1	2	5	ACC	1	2	2	4	AAC	1	0	2	6	AGC	1	0	2	6
AUA	0	2	3	4	ACA	1	2	2	4	AAA	1	0	2	5	AGA	1	1	2	4
AUG	0	0	3	6	ACG	1	2	2	4	AAG	1	0	2	5	AGG	1	1	2	5
GUU	1	2	2	4	GCU	1	2	2	4	GAU	1	0	2	6	GGU	1	2	2	4
GUC	1	2	2	4	GCC	1	2	2	4	GAC	1	0	2	6	GGC	1	2	2	4
GUA	1	2	2	4	GCA	1	2	2	4	GAA	1	0	2	5	GGA	1	2	2	3
GUG	1	2	2	4	GCG	1	2	2	4	GAG	1	0	2	5	GGG	1	2	2	4

Table 2. Total estimated synonymous *ti* (*tv*) and non-synonymous *ti* (*tv*) substitutions in the genetic

codon table. Asterisk (*) showed the total numbers of substitutions and ratio while excluding the

substitutions that lead to stop codons in PTC

ti & *tv*	Synonymous	Non-synonymous (*)	Total (*)
ti	62	121 (116)	183 (178)
tv	72	294 (276)	366 (348)
ti/tv	0.861	0.412 (0.420)	0.500 (0.511)

Table 3. Summary of the 2516 coding sequences selected for the present study. It represents the

minimum to maximum range of different parameters like GC%, TFD/FFD, PTC% and the mean and

standard deviation (S.D.) of the TFD%, FFD% and PTC%

Genes considered	Size bp (Min-Max)	GC% (Min-Max)	TFD/FFD (Min-Max)	PTC% (Min-Max)	TFD%	FFD%	PTC%	
					33.384	34.084	36.350	Mean
2516	93-4554	32.95-63.80	0.22-3.33	15.00-57.14	5.726	4.761	5.625	S. D