Synonymous and non-synonymous transitions/transversions vividly disclose purifying selection in *Escherichia coli* coding sequences

Pratyush Kumar Beura[1], Ruksana Aziz[1], Piyali Sen[2], Saurav Das[1], Nima Dondu Namsa[1,3], Edward J Feil[4], Siddharatha Sankar Satapathy*[2,3], Suvendra Kumar Ray*[1,3]

Department of [1]Molecular Biology and Biotechnology, [2]Computer Science and Engineering, [3]Centre for Multidisciplinary Research, Tezpur University, Tezpur, Assam, India, 784028, [4]The Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, United Kingdom

For correspondence*

Siddhartha Shankar Satapathy: ssankar@tezu.ernet.in

Suvendra Kumar Ray: suven@tezu.ernet.in

Full mailing addresses and zip codes:

Pratyush Kumar Beura, Ruksana Aziz, Piyali Sen, Saurav Das, Nima Dondu Namsa, Siddharatha Sankar Satapathy, Suvendra Kumar Ray: Tezpur University, Napaam, Tezpur, Assam, India 784028

Edward J Feil : Milner Centre for Evolution, University of Bath, Claverton Down**,** Bath, UK BA2 7AY

Running head: Substitutions and selection on coding sequences

Keywords: substitution mutation, nucleotide polymorphism, transition to transversion ratio, codon degeneracy, pretermination codon, purifying selection

Abstract

Transition (*ti*) and transversion (*tv*) are major events in genome variation. Therefore, it is crucial to estimate accurately these events for a better understanding of mutational and selection processes in organisms. Here, we developed a method (accessible at https://github.com/CBBILAB/CBBI.git) to estimate $\frac{ti}{tv}$ ratio by accounting degeneracy as well as pretermination nature of codons in coding sequences. Our findings revealed a distinct impact of codon degeneracy and pretermination codons on the $\frac{ti}{tv}$ ratio in the 2516 *Escherichia coli* coding sequences analyzed here. We observed the decreasing order of different base substitution frequencies such as *Sti* (synonymous transition) > *Stv* (synonymous transversion) > *Nti* (non-synonymous transition) > *Ntv* (non-synonymous transversion) in these coding sequences. The correlation between Sti and Stv values was strong (*r* value 0.795) whereas the correlation between *Sti* and *Nti* was weak (*r* value 0.192). Coding sequences with similar *Sti* values exhibited a wide range of *Nti* values. This indicated the varying strength of purifying selection acting on the coding sequences. In concordance with the assumption, the genes with high *Nti* values were observed to have lower CAI values than that of the genes with lower *Nti* values. Our approach is convenient to visualize the base substitution variation dynamics as well as selection in coding sequences. The proposed method is useful to estimate different $\frac{ti}{tv}$ ratios accurately in coding sequences and is insightful from an evolutionary perspective.

Article Summary

Genetic diversity is pivotal in evolution, with base substitution as a key driver. Transition (*ti*) surpasses transversion (*tv*) in genome frequency, making *ti*/*tv* ratios a valuable metric for studying mutation bias. Our improved estimator enhances *ti*/*tv* estimation in coding sequences, accounting in codon degeneracy and pretermination codons. Additionally, we unveil insights into substitution dynamics (*Sti*, *Stv*, *Nti*, *Ntv*) and demonstrate selection's impact on coding sequences by comparing *Sti* and *Nti* values.

Introduction

The genetic code table is a beautiful gift of nature to study molecular evolution (Crick *et al.* 1961; Khorana *et al.* 1966. The different degenerate codons influence the transition (*ti*) to transversion (*tv*) ratio ($\frac{ti}{tv}$) as follows: synonymous variation in a *two-fold degenerate* (TFD) codon occurs *via* only one transition whereas the same in a *four-fold degenerate* (FFD) codon occurs via one transition and two transversions (Supplementary Fig. 1a). In addition to the above, compositional variation of pretermination codons (PTC) (Supplementary Fig. 1b and 1c) (Modiano *et al.* 1981) among the coding sequences is also likely to influence the $\frac{ti}{tv}$ ratio because non-synonymous variations leading to termination codons purged out rapidly in a population due to strong purifying selection (Li *et al.* 1981; Eynden *et al.* 2016; Morales *et al.* 2021). Considering in general, a *ti* being more frequent than a *tv* (Seplyarskiy *et al.* 2012; Duchêne *et al.* 2015; Stoltzfus and Norris 2016; Lewis *et al.* 2016; Lyons and Lauring 2017; Schroeder *et al.* 2017; Sen *et al.* 2022) which is known as Kimura's two parameter model (Kimura 1980), and a stronger purifying selection on non-synonymous variation than a synonymous variation (Hurst and Pál 2001), the $\frac{ti}{tv}$ ratios of TFD codons and FFD codons are expected to be different.

Accordingly, it has been recently reported that the synonymous variation in TFD codons is

relatively higher than the FFD codons in *Escherichia coli* (Aziz *et al.* 2022). Therefore, in any

organism, the $\frac{ti}{tv}$ ratio is likely to vary across the coding sequences having differences in

degenerate codon composition (Beura *et al*. 2023). Hence, a comparative study of $\frac{ti}{tv}$ ratios

among coding sequences in a genome will be always insightful.

Considering the number of possibilities of the two types of base variations in genomic

DNA (Fersht and Knill-Jones 1981) (Supplementary Fig. 1d). The ratio of *ti* to *tv* is theoretically

expected to be 0.5. But the ratio is generally observed more than 0.5 in the genomes (Duchêne *et*

al. 2015; Stoltzfus and Norris 2016). Because of the following factors such as geometry of DNA

double helix (Topal and Fresco 1976), modification of DNA bases such as deamination of

cytosine and adenine (Kino and Sugiyama 2001; Rocha *et al.* 2006; Bhagwat *et al.* 2016),

secondary structure in the RNA (Sen *et al.* 2022) and codon degeneracy are known to elevate *ti*

phenomenon in a genome over *tv* (Muse and Gaut 1994; Aziz *et al.* 2022). In a standard genetic

code table out of 549 total single nucleotide variations (SNVs) involving 61 sense codons (9

SNVs per codon), only 134 SNVs are synonymous (24.4%). Among synonymous (*Sti* and *Stv*)

and non-synonymous (*Nti* and *Ntv*) variations, the theoretically estimated ($_e$) numbers of *Sti*, *Stv*,

Nti and *Ntv* in a genetic codon table are 62, 72, 121 and 294, respectively (Supplementary Table

1). Yet the observed synonymous variations are always higher than the expected proportion

across genes (Wolfe *et al.* 1989; Tamura 1992; Moriyama and Powell 1997; Ngandu *et al.* 2008)

due to stronger purifying selection on non-synonymous variations than the synonymous

variations (Hurst and Pál 2001). Hence, by accounting the codon degeneracy as well as PTC, the

Sti, *Stv*, *Nti* and *Ntv* in the genetic codon table can be estimated theoretically for each codon

(Supplementary Table 2).

The codon substitution model suggests that sites within codons evolve at different rates

and consequently they should not be equally treated (Muse and Gaut 1994; Shapiro *et al.* 2006;

Arenas 2015).   In the codon substitution model, using likelihood approach the $\frac{ti}{tv}$ ratio was

estimated in accordance with codon degeneracy class which was implemented in estimating

dN/dS in coding sequences (Muse and Gaut 1994; Goldman and Yang 1994). Recently, by

accounting $\frac{ti}{tv}$ substitution rate difference in codons as well as nonsense variations in PTC, the

dN/dS values have been improved (Aziz *et al.* 2022). This finding implies that factors such as

codon degeneracy and PTC composition should be considered while analyzing the evolution of

protein-coding genes. The ratio of *ti* to *tv* has been studied for reliable estimation of sequence

distance and phylogeny reconstruction (Yang and Yoder 1994). Using maximum likelihood

approach, the $\frac{ti}{tv}$ ratio (commonly known as kappa *κ*) has been used to mainly compare the $\frac{ti}{tv}$

values among different species (Yang and Yoder 1994). The parameter *κ* has not been

implemented to compare coding sequences within a genome in the context of their compositional

differences of different degenerate classes before. As the selection pressure on synonymous and

non-synonymous variations are heterogeneous (Bartolomé *et al.* 2005), it is important to analyze

the $\frac{ti}{tv}$ values among different genes within an organism by accounting synonymous sites and

non-synonymous sites regarding their compositional differences of different degenerate codons.

Given that there is a significant variation of substitution rates among protein coding

sequences within a genome (Wu and Li 1985), an appropriate methodology for the estimation of

ti to *tv* ratio is required. We are presenting an improved equation that builds upon the research

conducted by Yasuo Ina in 1998 (Ina 1998). Considering the differential selective constraint in

synonymous and non-synonymous variations it is crucial to separately analyze the synonymous

$\frac{ti}{tv}$ and non-synonymous $\frac{ti}{tv}$ to gain a comprehensive understanding of the overall $\frac{ti}{tv}$. The

improved equations used in this study are helpful to estimate the *ti* and *tv* rate differences in

coding sequences by accounting the compositional difference in terms of codon degeneracy and

PTC. Though it is known that TFD codons only undergo synonymous *ti*, the $\frac{Sti}{Stv}$ ratio among the

coding sequences in relation to their compositional differences (TFD, FFD and PTC) has not

been reported previously. Further, we observed a correlation between TFD: FFD and $\frac{Sti}{Stv}$ values

and the PTC% and $\frac{Nti}{Ntv}$ values. We tried to establish the biological significance of codon

composition and $\frac{ti}{tv}$ ratio across 2516 coding sequences of *E. coli*. Interestingly, coding sequences

with similar synonymous variation dynamics are found to have a profound variation regarding

their non-synonymous variation dynamics. The improved method developed in the present study

may be applied to study different $\frac{ti}{tv}$ $\left(\frac{Sti}{Stv} \text{ and } \frac{Nti}{Ntv}\right)$ across coding sequences.

Materials and Methods

Coding sequences of *E. coli* genome

In this study, we carried out a SNVs based computational analysis of 2516 coding sequences

across 157 strains of *E. coli* (Thrope *et al.* 2017). The information regarding the dataset selected

for the present study is provided in Supplementary Table 3. Although the public database has

~4500 coding sequences, we excluded coding sequences showing length variation and improper

alignment/annotation. Codons having variations in more than one position or variations leading

to ambiguous nucleotides were not considered in the analyses of the variations in this study. A

detailed procedure for estimating variation is represented in (Supplementary Table 4) using

example of a hypothetical sequence. This procedure is based on intra-species genome sequence

comparison to find out single nucleotide variations in bacterial genomes. A reference sequence

has been generated considering the most frequent nucleotide occurrence at a site in the gene

sequence. The logic is that the most abundant nucleotide in a species is the ancestral nucleotide

in the species (Sen et al. 2022; Aziz *et al.* 2022; Sen 2022, Beura *et al.* 2023).The observed

SNVs (SNV$_o$) at different positions of each codon were categorized into Sti_o, Stv_o, Nti_o and Ntv_o,

where S for synonymous, N for non-synonymous, ti_o for observed transition and tv_o for observed

transversion, and subsequently ti_o ($Sti_o + Nti_o$) and tv_o (Stv_{o+} Ntv_o) for each gene were calculated (

Supplementary Table 5). After finding observed variation values for a gene, we calculated the

theoretically estimated variation values Sti_e, Stv_e Nti_e, Ntv_e, ti_e and tv_e. We derived the reference

sequence for each gene based on the methodology explained (Supplementary Table 3). The

reference sequences were further used to find the codon counts of each gene. Then we multiplied

the count of each codon by the theoretically estimated variation values for each codon. For

example, if the codon frequency of GGG is observed to be 10 in a gene, then the total

theoretically estimated value for *Sti*, *Stv*, *Nti* and *Ntv* are 10, 20, 20 and 40 (as estimated *Sti*, *Stv*,

Nti and *Ntv* for each GGG is 1, 2, 2 and 4, respectively). We then summed up the individual

variation values for 61 sense codons to get the total theoretically estimated individual substitution

values (Sti_e, Stv_e, Nti_e and Ntv_e) for the gene (Supplementary Table 6). The values of Sti_o, Stv_o,

Nti_o, Ntv_o, ti_o, and tv_o and the values of Sti_e, Stv_e Nti_e, Ntv_e, ti_e, and tv_e of a coding sequence are

then applied in the modified equation to find out different $\frac{ti}{tv}$ ratio (Supplementary Table 7). A

detailed workflow is provided in Fig. 1.

Improved equations

We proposed an improved mathematical equation for the calculation of different $\frac{ti}{tv}$ ratio as

described below.

$$\frac{ti'}{tv'} = \frac{\left(ti_o/ti_e\right)}{\left(tv_o/tv_e\right)}$$ **Equation 1**

$$\frac{Sti'}{Stv'} = \frac{\left(Sti_o/Sti_e\right)}{\left(Stv_o/Stv_e\right)}$$ **Equation 2**

$$\frac{Nti'}{Ntv'} = \frac{\left(Nti_o/Nti_e\right)}{\left(Ntv_o/Ntv_e\right)}$$ **Equation 3**

Where, $\frac{ti'}{tv'}$ is improved $\frac{ti}{tv}$ ratio, $\frac{Sti'}{Stv'}$ is improved $\frac{Sti}{Stv}$ ratio, $\frac{Nti'}{Ntv'}$ is improved $\frac{Nti}{Ntv}$ ratio, ti_o is transition

observed, ti_e is the theoretically estimated transition, tv_o is transversion observed, tv_e is

transversion estimated, Sti_o is synonymous transition observed, Sti_e is the theoretically estimated

synonymous transition, Stv_o is synonymous transversion observed, Stv_e is the theoretically

estimated synonymous transversion, Nti_o is non-synonymous transition observed, Nti_e is the

theoretically estimated non-synonymous transition, Ntv_o is non-synonymous transversion

observed, Ntv_e is the theoretically estimated non-synonymous transversion. We calculated $\frac{ti}{tv}$ ratios

for all the genes using a program written in the Python language.

It is noteworthy that in a genome the number of potential replacements is greater for non-

synonymous alterations, whereas the frequency of actual variations is higher for synonymous

alterations among coding sequences. The improved estimators $\frac{ti'}{tv'}$, $\frac{Sti'}{Stv'}$, and $\frac{Nti'}{Ntv'}$ account for the

observed variations out of the total possible variations, on the contrary, the conventional method

only takes an account of the ti_o and tv_o. The possible sites for Sti and Stv are also different among

genes, which is dependent upon the codon composition belonging to different degeneracy classes.

The conventional $\frac{ti}{tv}$ ratio does not adequately account for the limitations of variable selection

against synonymous and non-synonymous variations. Hence, all these parameters make the

requirement of an improved estimator instead of the conventional estimator.

Comparative representation of different $\frac{ti}{tv}$ through conventional and improved method

We have done a comparative analysis of $\frac{ti}{tv}$ calculated using the existing and the improved method

to support the validation of method proposed in this study. Also, we have calculated the TFD: FFD

and the percentage of PTC (PTC%) of each gene and performed statistical analysis between

different parameters. A summary of the different parameters observed throughout the dataset is

represented in Table 1. We also calculated the percentage (%) change in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ values obtained

from conventional to improved method as follows.

(a) $\frac{Sti}{Stv}$ % change $= \dfrac{\frac{Sti}{Stv} - \frac{Sti'}{Stv'}}{\frac{Sti}{Stv}} \times 100$

(b) $\frac{Nti}{Ntv}$ % change $= \dfrac{\frac{Nti}{Ntv} - \frac{Nti'}{Ntv'}}{\frac{Nti}{Ntv}} \times 100$

We found out $\frac{ti}{tv}$ in all these 2516 genes using maximum likelihood (ML) approach

available in MEGA-X (Kumar *et al.* 2018). Using our improved approach, we computed the

normalized *Sti, Stv*, *Nti*, and *Ntv* values for the dataset. These normalized values were determined

by dividing the observed values by the corresponding estimated values in the dataset and further

analyzed in the variation dynamics study.

Statistical analyses

Comparison among the conventional, ML approach and the improved $\frac{ti}{tv}$ values were done using

different statistical methods to demonstrate the performance and accuracy of the improved

method. OriginPro, Version 2022, OriginLab Corporation, Northampton, MA, USA was used to

draw the Box-plot as well as to perform the Mann-Whitney test (Mann and Whitney 1947) to find

out the *p*-value. The correlation plot was drawn for R^2 value and Pearson's correlation coefficient

(Pearson 1896) was also calculated between the $\frac{ti}{tv}$ values obtained using the conventional and

improved method.

Results

Impact of codon degeneracy and PTC on $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ values in *E. coli*

We found out the total number of TFD, FFD, and PTC codons across 2516 coding sequences of *E.*

coli. The percentage of TFD, FFD, and PTC was calculated (Supplementary Table 3). The

percentage (%) compositional differences in the 2516 coding sequences represented as TFD%,

FFD%, and PTC% is shown in a box-plot (Fig. 2a). A variable range of TFD: FFD ratio in the

coding sequences of *E. coli* was observed. The minimum to maximum TFD:FFD ratio was observed

as 0.229 and 3.333 in *sugE* and *mntS*, respectively. It suggests that FFD codons are present

approximately four times more frequent than TFD codons in certain coding sequences. Similarly,

TFD codons are present approximately three times more frequent than FFD codons in the selected

coding sequences. Among 2516 coding sequences, only 35 coding sequences possessed the

TFD:FFD ratio of 1.000. Hence, this observation suggests that there is a compositional asymmetry

of TFD and FFD codons in the coding sequences. The minimum to maximum PTC% was observed

in the gene *atpE* as 15.00% and in the gene *chaB* as 57.14% (Fig. 2a). The compositional variation

(TFD, FFD and PTC) observed in coding sequences were considered to validate the accuracy of

different $\frac{ti}{tv}$ ratio using the conventional and improved method.

216     We estimated $\frac{ti}{tv}$ by accounting codon degeneracy as well as PTC in coding sequences and

compared with the conventional method of $\frac{ti}{tv}$ in coding sequences. Theoretically, a coding region

entirely composed of FFD codons will result into a twice $\frac{Sti'}{Stv'}$ value than $\frac{Sti}{Stv}$ value, similarly a coding

region composed of same proportion of FFD and TFD codons will result into equal values of $\frac{Sti'}{Stv'}$

and $\frac{Sti}{Stv}$. The increase in TFD codon proportion will directly impact the difference between $\frac{Sti}{Stv}$ values

and $\frac{Sti'}{Stv'}$ values. We have observed a similar pattern as well (Supplementary Fig. 2). The complete

set of observed and estimated values of different variations is enlisted (Supplementary Table 3).

These values were used to calculate the $\frac{ti}{tv}, \frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$. Further the improved equation was used to

calculate the $\frac{ti'}{tv'}, \frac{Sti'}{Stv'}$ and $\frac{Nti'}{Ntv'}$. We compared the values derived using the conventional method (

$\frac{ti}{tv}, \frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$) and improved method ($\frac{ti'}{tv'}, \frac{Sti'}{Stv'}$ and $\frac{Nti'}{Ntv'}$) (Supplementary Table 3). A box-plot

showing the comparison between the values of different $\frac{ti}{tv}$ ratio obtained through the conventional

and improved method is represented in Fig. 2b. The minimum % change in $\frac{Sti}{Stv}$ between conventional

and improved method was observed in the gene *garP* as -30.99% and maximum in the gene *yfhL* as

30.76%. Similarly, the minimum % change in $\frac{Nti}{Ntv}$ between conventional and improved method was

observed in the gene *thiS* as -194.675% and maximum in the gene *bhsA* as -129.600%. The detailed

% change in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ is represented (Supplementary Table 3). While it is evident that, TFD

codons and FFD codons have different probabilities of obtaining synonymous variations, the

composition of TFD: FFD was cross-checked against the % change in $\frac{Sti}{Stv}$ for each gene. Likewise,

the impact of PTC% in estimating the $\frac{Nti}{Ntv}$ values have been considered in this study.

The synonymous variation in the genes is likely to be different considering the differential

composition of TFD codons and FFD codons, because when FFD codon proportion is higher in a

gene, $\frac{ti}{tv}$ value is going to be lower. Similarly, when the FFD codon proportion is lower, $\frac{ti}{tv}$ value is

going to be higher. So, this observation is distinct, and the improved estimator is taking care of the

compositional differences (estimated numbers) of different degenerate codons. The Pearson

correlation coefficient value between TFD: FFD and % change in $\frac{Sti}{Stv}$ (conventional to improved)

was obtained to be 0.891, signifying a strong correlation. Similarly, the Pearson correlation

coefficient value between PTC% and % change in $\frac{Nti}{Ntv}$ was obtained to be -0.488, a moderate

negative correlation between the two variables. The results were found to be statistically significant

at $p<0.01$, for $\frac{ti}{tv}$ & $\frac{ti'}{tv'}$, $\frac{Sti}{Stv}$ & $\frac{Sti'}{Stv'}$ and $\frac{Nti}{Ntv}$ & $\frac{Nti'}{Ntv'}$. Further the correlation graph between % change in

$\frac{Sti}{Stv}$ and TFD: FFD and % change in $\frac{Nti}{Ntv}$ and PTC% also substantiate the role of codon composition

and $\frac{ti}{tv}$ ratio (Fig. 3a & 3b). In the trendline analyses the % change in $\frac{Sti}{Stv}$ was perceived to be directly

influenced by the TFD: FFD with a R^2 value of 0.795. A weak negative association was obtained

between % change in $\frac{Nti}{Ntv}$ and PTC% with a R^2 value of 0.216. Hence, it is pertinent to note that the

overall % change in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ were directly influenced by the TFD and PTC codon composition

across the coding sequences. We performed a simulation study to substantiate the impact of the

compositional difference of degeneracy class in estimating $\frac{Sti}{Stv}$ by using our improved method in

coding sequences. For detailed methodology and results (Supplementary Table 8, Supplementary

Fig. 3). The results of comparative analysis between ML based method and improved method are

presented in supplementary files for 2516 genes (Supplementary Table 9, Supplementary Fig. 4).

Sti, _Stv_, _Nti_ and _Ntv_ variation dynamics in coding sequences

The variation dynamics such as _Sti_, _Stv_, _Nti_, and _Ntv_ were analyzed in the 2516 coding sequences

where the mean values were 0.113, 0.034, 0.014 and 0.004, respectively (Supplementary Table 11,

Fig. 4a). The mean values suggested the decreasing order of the variation dynamics: _Sti_ > _Stv_ >_Nti_ >

Ntv. A comparison among the values suggested that _Sti_ was 3.3 times more frequent than _Stv_, while

Nti was 3.7 times more frequent than _Ntv_. In addition, _Sti_ is 8.2 times more frequent than _Nti_ while

Stv was 9.3 times more frequent than _Ntv_. This indicated that the relative purifying selection on _Nti_

and _Ntv_ in relation to their synonymous counter parts was similar. The Pearson correlation

coefficient _r_ values between _Sti_ and _Stv_ was 0.795 while the same between _Sti_ and _Nti_, was 0.192

(Table 2). This indicated that selection on non-synonymous variation is independent upon the

synonymous variation. In comparison with the conventional method, our improved method is more

appropriate in estimating the distinctive selection patterns between _Sti_ and _Nti_ (Table 2).

As part of our comparative study, we investigated the coding sequences of similar _Sti_ and

different _Nti_ values (Fig. 4b). In a set of 20 coding sequences with similar _Sti_ values such as 0.131,

the _Nti_ values were ranging from 0.000 to 0.030. This wide range of _Nti_ values, with a maximum 30-

fold variation, highlights the presence of a variable _Nti_ pattern within the respective coding

sequences. Similar observation could be found in other Sti values. We compared Sti and Nti values

in relation to gene expression represented with CAI values (Sharp and Li 1987; Sen et al. 2019). We

studied genes with size more than 500 bp. The top 100 genes with maximum Sti values (mean 0.255)

were compared with bottom 100 genes with minimum Sti values (mean 0.006). The mean CAI

values for the top 100 genes were 0.565 whereas the mean CAI values for the bottom 100 genes was

0.509. The two mean CAI values were found to be close. This indicates the Sti values might not be

influenced significantly due to gene expression. Similarly, the top 100 genes with maximum *Nti*

values (mean 0.036) were compared with bottom 100 genes with minimum *Nti* values (mean 0.001).

The mean CAI values for the top 100 genes were 0.491 whereas the mean CAI values for the bottom

100 genes was 0.638. The two mean CAI values of the coding sequences were found to be

significantly different ($p<0.01$). The selection due to gene expression is stronger on Nti than on Sti

among the strains of *E. coli*.

Discussion

Codon degeneracy influences the *ti* to *tv* ratio in coding sequences because a TFD codon

undergoes synonymous variation only by transition while a FFD codon undergoes synonymous

variation by transition as well as transversion (Aziz *et al.* 2022). In addition, there are only 18

pretermination codons in the genetic code table where nonsense substitution is rare to observe.

Coding sequences in a genome vary with composition of codons belonging to different

degeneracy as well as PTC. Therefore, it is inappropriate to compare coding sequences regarding

the $\frac{ti}{tv}$ ratio without accounting degeneracy and pretermination nature of codons. In this study we

have developed a methodology to estimate $\frac{ti}{tv}$ ratio by accounting degeneracy and pretermination

nature of codons, which has been implemented to analyze 2516 coding sequences in *E. coli*. Our

result suggested that coding sequences are different regarding their $\frac{ti}{tv}$ ratio, which is independent

of the composition of degenerate codons and PTC. In fact, the $\frac{Sti}{Stv}$ values are observed to be

variable among the coding sequences, which is intriguing. We did not observe any significant

correlation of $\frac{Sti}{Stv}$ values with the gene expression represented as CAI values. The role of context

dependent mutation (Sung *et al.* 2015; Zhu *et al.* 2017; Aikens *et al.* 2019; Ling *et al.* 2020),

which has been described recently, might be a contributing factor that needs to be investigated in

future.

Our investigation into *ti* and *tv* revealed that *Sti* is around three times more frequent than

Stv while it is eight times more frequent than *Nti*. The difference between *Sti* and *Stv* may be

attributed due to the structural changes in DNA happening due to mispairing between either

purine: purine or pyrimidine: pyrimidine but the difference between *Sti* and *Nti* may be

attributed to the purifying selection acting on the coding sequences because of the non-

synonymous changes. The magnitude of the purifying selection acting on non-synonymous

substitution in *E. coli* genome is vividly observed. The strength of purifying selection on *Nti* and

Ntv seems comparable considering the closeness of the $\frac{Sti}{Stv}$ and the $\frac{Nti}{Ntv}$ values. In a recent study

by Zou and Zhang (Zou and Zhang 2021) suggests that the purifying selection is more on *Ntv*

than on *Nti* which is unlike the observation reported in our study. Our different observation from

that of Zou and Zhang might be attributed to the intra-species study limited to *E. coli* whereas

Zou and Zhang worked on an inter-species approach covering 90 clades representing all domains

of life. Further, in our estimation of *Nti* and *Ntv* we have accounted codon degeneracy and PTC.

As transitions are more frequent events than transversion, transition can be considered

alone to understand the selection on coding sequences. Considering *Sti* representing the variation

dynamics in a coding sequence, the *Nti* representing the strength of purifying selection acting on

the coding sequence, a comparison between *Sti* and *Nti* in a coding sequence might represent the

strength of purifying selection acting on it. In several instances, we have observed that coding

sequences with equal *Sti* dynamics are highly different regarding their *Nti* dynamics, which

indicates disproportionate purifying selection on coding sequences. Additionally, when we

compared the mean CAI values of the coding sequences with higher *Sti* values against the mean

CAI values of the coding sequences with lower *Sti* values, we did not find any significant

difference. On the contrary, when we conducted the same analysis for the *Nti* values, we

observed a significant distinction between the CAI values of coding sequences with higher *Nti*

values and those with lower *Nti* values. This observation indicates that highly expressed genes

tend to have lower *Nti* values. Consequently, it suggests that purifying selection due to gene

expression is more pronounced for *Nti* values than for *Sti* values. Since our study on selection

relies on intra-species analysis, we anticipate that future research will unveil the underlying

mechanisms of selection governing variation dynamics in inter-species studies as well. A

probable discrepancy between the process of selection between intra-species and inter-species

can be drawn. This is a simple demonstration of observing purifying selection acting on coding

sequences. Though the difference between *ti* and *tv* are known since long, we could use it to

study selection on coding sequences because of the improved equation.

Various methods have been implemented in the estimation of the *ti* to *tv* ratio by

molecular evolutionary biologists to understand the mutation as well as selection bias. The

phylogeny-based method and Bayesian methods have been implemented by researchers in the

inter-species study of $\frac{ti}{tv}$ (Purvis and Bromham 1997; Huelsenbeck *et al.* 2001). Usually the

dN/dS approach has been used to visualize the selection in coding sequences (Aziz *et al.* 2022)

that considers both *ti* and *tv*. However, our method implemented in this study is simple, succinct,

easy to follow for researchers in the field with limitations in mathematics and provides an insight

to understand the evolution of protein coding sequences. We also did a comparison with the ML

method. There are several genes where we could observe the $\frac{ti}{tv}$ values estimated by the improved method is more in concordance with the observed *ti* and *tv* values than the ML method. It has been said that the likelihood methods are computer intensive and difficult to apply to large datasets and can fall into local traps (Golding and Felsenstein 1990). Furthermore, the extrapolation of our method in various inter-species studies, as well as phylogenetic tools can enhance the horizon of our understanding regarding mutational bias in coding sequences.

Data and software availability

A script written in Python-language is available at GitHub in the following link: (https://github.com/CBBILAB/CBBI.git). Further queries regarding the software may contact SSS (ssankar@tezu.ernet.in). The authors confirm that the data supporting the findings of this study are available within the article and Supporting information.

Acknowledgments

PKB is grateful to Tezpur University for its institutional fellowship. RA is thankful for the JRF fellowship from the DBT grant (BT/511/NE/TBP/2013) and (BT/403/NE/U-Excel/2013). PS is grateful to UGC, GoI New Delhi, for the JRF. SD is thankful to DBT for MSc. Scholarship. SSS and SKR are thankful to DBT, GoI for the twinning grant (BT/PR16361/NER/95/192/2015 date 18-10-2016) to them. NDN, SSS & SKR thankfully acknowledge the DBT funded Bioinformatics and Computational Biology Centre at Tezpur University. We are also thankful to Prof. Kini R Manjunatha, National University of Singapore for his kind suggestions during his visit to Tezpur University.

Funding

We received no external funding for the work.

Conflicts of interest

The author(s) declare no conflict of interest.

Literature cited

Aikens RC, Johnson KE, Voight BF. Signals of variation in human mutation rate at multiple levels of
sequence context. Mol Biol Evol. 2019;36(5): 955–965. doi:10.1093/molbev/msz023.

Arenas M. Trends in substitution models of molecular evolution. Front Genet. 2015; 6:319. doi:
10.3389/fgene.2015.00319

Aziz R. et al. Incorporation of transition to transversion ratio and nonsense mutations improves the
estimation of the number of synonymous and non-synonymous sites in codons. DNA Research.
2022;29: dsac023. doi:10.1093/dnares/dsac023

Bartolomé C, Maside X, Yi S, Grant AL, Charlesworth B. Patterns of selection on synonymous and
nonsynonymous variants in Drosophila miranda. Genetics. 2005; 169(3):1495-507.
doi: 10.1534/genetics.104.033068

Beura PK, Sen P, Aziz R, Satapathy SS, Ray S, Transcribed intergenic regions exhibit a lower
frequency of nucleotide polymorphism than the untranscribed intergenic regions in the genomes of
Escherichia coli and Salmonella enterica. J Genet. 2023; 102:22. doi:10.1007/s12041-023-01418-w

Beura PK et al. Difference in synonymous polymorphism related to codon degeneracy between co-
transcribed genes in the genome of *Escherichia coli*. Curr. Sci. 2023. Doi: 10.18520/cs/v125/i8/

Bhagwat AS et al. Strand-biased cytosine deamination at the replication fork causes cytosine to
thymine mutations in Escherichia coli. Proc Natl Acad Sci. 2016;113, 2176–2181.
doi:10.1073/pnas.1522325113.

Crick FHC, Barnett L, Brenner S, Watts-Tobin RJ, General nature of the genetic code for proteins.
Nature 1961;30; 192:1227-32. doi:10.1038/1921227a0

Duchêne S, Ho SYW, Holmes EC. Declining transition/transversion ratios through time reveal
limitations to the accuracy of nucleotide substitution models. BMC Evol Biol. 2015; 15:1–10.
doi:10.1186/s12862-015-0312-6

Fersht AR, Knill-Jones JW DNA polymerase accuracy and spontaneous mutation rates: frequencies of
purine. purine, purine. pyrimidine, and pyrimidine. pyrimidine mismatches during DNA
replication. Proc Natl Acad Sci. 1981;78(7):4251–4255. doi:10.1073/pnas.78.7.4251

Golding B, J. Felsenstein. A maximum likelihood approach to the detection of selection from a
phylogeny. J Mol Evol. 1990;31:511-523. doi:10.1007/BF02102078

Goldman N, Yang Z. A codon-based model of nucleotide substitution for protein-coding DNA
sequences. Mol Biol Evol. 1994;11(5), 725-736. doi:10.1093/oxfordjournals.molbev.a040153

Huelsenbeck J, Ronquist F. MrBayes: Bayesian inference of phylogenetic trees. Bioinform. 17, 754–
755 (2001). doi:10.1093/bioinformatics/17.8.754

Hurst LD, Pál C. Evidence for purifying selection acting on silent sites in BRCA1. TRENDS in
Genetics. 2001;17:62–65. doi:10.1016/s0168-9525(00)02173-9.

Ina Y. Estimation of the transition/transversion ratio. J Mol Evol. 1998; 46:521–533. doi:
10.1007/PL00006333

Khorana HG et al. Polynucleotide synthesis and the genetic code. Cold Spring Harbor Symp Quant
Biol. 1966; 31:39-49. doi:10.1101/sqb.1966.031.01.010

Kimura M. A simple method for estimating evolutionary rates of base substitutions through
comparative studies of nucleotide sequences. J Mol Evol. 1980; 16:111–120.
doi:10.1007/BF01731581

Kino K, Sugiyama H. Possible cause of G. C> C. G transversion mutation by guanine oxidation
product, imidazolone. Chem. Biol. 2001;8, 369–378. doi:10.1016/S1074-5521(01)00019-9

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGAX: Molecular evolutionary genetics analysis
across computing platforms. Mol Biol Evol. 2018;35(6):1547–1549. doi:10.1093/molbev/msy096.

Lewis Jr CA, Crayle J, Zhou S, Swanstrom R, Wolfenden R. Cytosine deamination and the
precipitous decline of spontaneous mutation during Earth's history. Proc Natl Acad Sci. 113:8194–
8199 (2016). doi:10.1073/pnas.1607580113.

Li W-H, Gojobori T, Nei M. Pseudogenes as a paradigm of neutral evolution. Nature 1981; 292:237–
239. doi:10.1038/292237a0

Ling G, Miller D, Nielsen R, Stern AA. Bayesian framework for inferring the influence of sequence
context on point mutations. Mol Biol Evol. 2020;37:893–903. doi:10.1093/molbev/msz248

Lyons DM, Lauring AS, Evidence for the selective basis of transition-to-transversion substitution bias
in two RNA viruses. Mol Biol Evol. 2017;34:3205–3215. doi:10.1093/molbev/msx251

Mann HB, Whitney DR. On a test of whether one of two random variables is stochastically larger than
the other. Ann Math Statist. 1947;18(1), 50-60. doi:10.1214/aoms/1177730491

Modiano G, Battistuzzi G, Motulsky AG. Nonrandom patterns of codon usage and of nucleotide

substitutions in human alpha-and beta-globin genes: an evolutionary strategy reducing the rate of

mutations with drastic effects? Proc Natl Acad Sci. 1981; 78:1110–1114.

doi:10.1073/pnas.78.2.1110

Morales AC et al. Causes and consequences of purifying selection on SARS-CoV-2. Genome Biol

Evol. 2021;13: evab196. doi:10.1093/gbe/evab196

Moriyama EN, Powell JR. Synonymous substitution rates in Drosophila: mitochondrial versus nuclear

genes. J Mol Evol. 1997; 45:378–391. doi:10.1007/pl00006243.

Muse SV, Gaut BS. A Likelihood Approach for Comparing Synonymous and Nonsynonymous

Nucleotide Substitution Rates, with Application to the Chloroplast Genome. Mol Bio Evol. 1994.;1

1(5):7 15-724. doi:10.1093/oxfordjournals.molbev.a040152

Ngandu NK et al. Extensive purifying selection acting on synonymous sites in HIV-1 Group M

sequences. Virol J. 2008; 5:1–11. doi:10.1186/1743-422X-5-160

Pearson K. Mathematical contributions to the theory of evolution. On a form of spurious correlation

which may arise when indices are used in the measurement of organs. Proc R Soc Lond.

1896;60(359-367), 489-498. doi:10.1098/rspl.1896.0076

Purvis A, Bromham L. Estimating the transition/transversion ratio from independent comparisons

with an assumed phylogeny. J Mol Evol. 1997;44, 112–119. doi:10.1007/pl00006117

Rocha EPC, M. Touchon, Feil EJ. Similar compositional biases are caused by very different

mutational effects. Genome Res. 16, 2006; 1537–1547. doi:10.1101/gr.5525106

Schroeder JW, Randall JR, Hirst WG, O'Donnel ME, Simmons LA. Cost of ribonucleotides in

bacterial DNA. Proc Natl Acad Sci. 2017; 114:11733–11738. doi:10.1073/pnas.1710995114

Sen P, Waris A, Ray SK, Satapathy SS. A Web Portal to Calculate Codon Adaptation Index (CAI)

with Organism Specific Reference Set of High Expression Genes for Diverse Bacteria Species.

ICICSCNT. 2019; doi:10.1007/978-981-15-0633-8_31.

Sen P. "Computational Analysis of Codon Usage Bias Single Nucleotide Polymorphism and RNA

Secondary Structures in Microbial Genome Sequences". Ph.D. Thesis, Tezpur University. 2022;

http://hdl.handle.net/10603/473607.

Sen P, Aziz R, Deka RC, Feil EJ, Ray SK, Satapathy SS. Stem region of tRNA genes favors transition

substitution towards keto bases in bacteria. J Mol Evol. 2022; 90:114–123. doi:10.1007/s00239-

021-10045-x.

Seplyarskiy VB, Kharchenko P, Kondrashov AS, Bazykin GA. Heterogeneity of the

transition/transversion ratio in Drosophila and Hominidae genomes. Mol Biol Evol. 2012;

29:1943–1955. doi:10.1093/molbev/mss071.

Shapiro B, Rambaut A, Drummond AJ, Choosing appropriate substitution models for the

phylogenetic analysis of protein-coding sequences. Mol Biol Evol. 2006;23, 7–9.

doi:10.1093/molbev/msj021

Sharp PM, Li W-H. The codon Adaptation Index--a measure of directional synonymous codon usage

bias, and its potential applications. Nucleic Acids Res. Feb 1987;11;15(3):1281-95.

doi:10.1093/nar/15.3.1281

Stoltzfus A, Norris RW, On the causes of evolutionary transition: transversion bias. Mol Biol Evol.

2016; 33:595–602. doi:10.1093/molbev/msv274.

Sung W. et al. Asymmetric context-dependent mutation patterns revealed through mutation-

accumulation experiments. Mol Biol Evol. 2015; 32:1672–1683. doi:10.1093/molbev/msv055.

Tamura K. The rate and pattern of nucleotide substitution in Drosophila mitochondrial DNA. Mol

Biol Evol. 1992; 9:814–825. doi:10.1093/oxfordjournals.molbev.a040763

Thorpe HA, Bayliss SC, Hurst LD, Feil EJ, Comparative analyses of selection operating on

nontranslated intergenic regions of diverse bacterial species. Genetics. 2017;206:363–376.

doi:10.1534/genetics.116.195784.

Topal MD, Fresco JR, Complementary base pairing and the origin of substitution mutations. Nature

1976;263(5575):285-289. doi:10.1038/263285a0.

van den Eynden J, Basu S, Larsson E. Somatic mutation patterns in hemizygous genomic regions

unveil purifying selection during tumor evolution. PLoS Genet. 12: e1006506 (2016).

doi:10.1371/journal.pgen.1006506

Yang Z, Yoder AD. Estimation of the Transition/Transversion Rate Bias and Species Sampling. J Mol

Evol. 1999;48:274–283. doi:10.1007/PL00006470

Wolfe KH, Sharp PM, Li W-H. Rates of synonymous substitution in plant nuclear genes. J Mol Evol.

1989;29:208–211. doi:10.1007/BF02100204

Wu C-I,   Li W-H. Evidence for higher rate of nucleotide substitution in rodents than in man. Proc

Natl Acad Sci. 1985;82(6):1741-1745. doi:10.1073/pnas.82.6.1741

Zhu Y, Neeman T, Yap VB, Huttley GA. Statistical methods for identifying sequence motifs affecting

point mutations. Genetics. 2017;205(2):843-856.   doi:10.1534/genetics.116.195677

Zou Z, J. Zhang, Are Nonsynonymous Transversions Generally More Deleterious than

Nonsynonymous   Transitions?   Mol   Biol   Evol.   38(1):181–191(2021).

doi:10.1093/molbev/msaa200.

494 **Figure_legend_Table**

Fig. 1. Schematic representation of the workflow showing the steps to execute our improved

method. By considering each codon as j, ti_e^j and tv_e^j can be calculated for each codon as presented

in Supplementary Table S2. The reference sequence needs to be calculated from the coding region

(gene/query). From the reference sequence the observed ($_o$) substitutions are calculated for the gene.

The other usage of the reference sequence is to find out the codon count of the gene. Then, the

respective codon count is multiplied by estimated transition and transversion for each codon. The

next step is to find out the sum of ti and sum of tv in all codons to represent the estimated ($_e$) ti and tv

for the gene. After getting the observed and estimated values for ti and tv, now the improved

equation is applied to find out the $\frac{ti'}{tv'}$ of the gene. This scenario is only showing the $\frac{ti'}{tv'}$ ratio, whereas

in similar way the $\frac{Sti'}{Stv'}$ & $\frac{Nti'}{Ntv'}$ can be calculated.

Fig. 2. (a) The box-plot presents the percentage (%) of TFD, FFD and PTC codons in the coding

sequences considered to validate the improved method proposed in this study. The minimum and

maximum TFD% in the dataset was observed as 10.38% and 53.25%, similarly the minimum and

maximum FFD% in the dataset was observed as 13.95% and 52.05%. The minimum and maximum

PTC% in the dataset was observed as 15.00% and 57.14%. (b) The box-plot presents the different $\frac{ti}{tv}$

ratio values. Here we have compared the different $\frac{ti}{tv}$ ratios of the entire set of genes through

conventional and improved method. $\frac{ti}{tv}$ shows the conventional method and $\frac{ti'}{tv'}$ shows the result of

improved method. Similarly, $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$ represent the conventional values and $\frac{Sti'}{Stv'}$ and $\frac{Nti'}{Ntv'}$ represent

the improved values. This illustration shows that the median values between $\frac{ti}{tv}$ & $\frac{ti'}{tv'}$, $\frac{Sti}{Stv}$ & $\frac{Sti'}{Stv'}$ and

$\frac{Nti}{Ntv}$ & $\frac{Nti'}{Ntv'}$ are statistically different (p-value < 0.01).

    (a)

    (b)

525 **Fig. 3.** Regression plot of coding sequence considered to validate the improved method. The x-axis

526 shows the codon composition (TFD: FFD and PTC%) and y-axis shows the % change in $\frac{Sti}{Stv}$ and $\frac{Nti}{Ntv}$.

527 (a) % change in $\frac{Sti}{Stv}$ is showing a strong correlation with the TFD: FFD with a R^2 value of 0.795. (b)

528 % change in $\frac{Nti}{Ntv}$ is showing a weak negative correlation with the PTC% with a a R^2 value of 0.216.

Fig. 4. (a) The figure illustrates the box-plot of the decreasing variation dynamics of *Sti*, *Stv*, *Nti*,

and *Ntv* values. The mean *Sti* and *Stv* were observed as 0.113 and 0.031 respectively, similarly the

mean *Nti* and *Ntv* values were observed as 0.014 and 0.004. *Sti* was observed to be highly frequent

than *Stv* and *Nti* was observed to be highly frequent than *Ntv*. There was a significant difference

between *Sti* and *Stv* values as well as between *Nti* and *Ntv* values($p<0.01$). (b) The figure illustrates

the Sti and Nti values of coding sequences used in the study through a double y-axis plot. The left y-

axis shows the Sti values of coding sequences, and the right y-axis shows the Nti values of the

coding sequences. The x-axis shows the numbers of coding sequences considered in this study. We

have arranged the Sti values in an increasing order. The black line represents the Sti values, and the

red dots represent the corresponding Nti values of a coding sequence. It is evident that many coding

sequences with similar Sti values have a wide range of Nti values. The Sti and Nti values may

represent the mutation and selection relation in a coding sequence.

Table 1. Summary of the 2516 coding sequences selected for the present study. It represents the
minimum to maximum range of different parameters like GC%, TFD/FFD, PTC% and the mean and
standard deviation (S.D.) of the TFD%, FFD% and PTC%

Genes considered	Size bp (Min-Max)	GC% (Min-Max)	TFD/FFD (Min-Max)	PTC% (Min-Max)	TFD%	FFD%	PTC%	
					33.384	34.084	36.350	Mean
2516	93-4554	32.95-63.80	0.22-3.33	15.00-57.14	5.726	4.761	5.625	S. D

556   **Table 2.** Comparative Pearson correlation coefficient study among Sti, Sti, Nti and Ntv through
557   conventional and improved estimator

Pearson correlation coefficient	Conventional method	Improved method
	Observed values	Normalized values (Obs/Est)
Sti & Stv	0.920	0.795
Nti & Ntv	0.789	0.612
Sti & Nti	0.527	0.192
Stv & Ntv	0.544	0.257