Phosphorylation, disorder, and phase separation govern the behavior of Frequency in the fungal circadian clock

Daniyal Tariq[a,*], Nicole Maurici[b,*], Bradley M. Bartholomai[c], Siddarth Chandrasekaran[a], Jay C. Dunlap[c], Alaji Bah[b,1] and Brian R. Crane[a,1].

[a]Department of Chemistry & Chemical Biology, Cornell University, Ithaca, NY,14853;

[b]Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, NY, 13210;

[c]Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth, Hanover, NH 03755

*Denotes equal contribution

[1]To whom correspondence may be addressed. BahA@upstate.edu; bc69@cornell.edu.

Abstract

Circadian clocks are composed from molecular oscillators that pace rhythms of gene expression to the diurnal cycle. Therein, transcriptional-translational negative feedback loops (TTFLs) generate oscillating levels of transcriptional repressor proteins that regulate their own gene expression. In the filamentous fungus *Neurospora crassa,* the proteins **F**requency (**F**RQ), the **F**RQ-interacting RNA helicase (FRH) and **C**asein-Kinase I (CK1) form the **FFC** complex that represses expression of clock-controlled genes activated by the White-Collar complex (WCC). A key question concerns how the FRQ protein orchestrates molecular interactions at the core of the clock despite containing little predicted tertiary structure. We present the reconstitution and characterization of FRQ and the FFC in unphosphorylated and highly phosphorylated states. An integrated biophysical approach incorporating site-specific spin labeling and pulse-dipolar ESR spectroscopy provides domain-specific structural details on the full-length, 989-residue intrinsically disordered FRQ and the FFC. Although extremely flexible, FRQ contains a compact core that associates and organizes FRH and CK1 to coordinate their roles in WCC repression. Phosphorylation of FRQ increases conformational flexibility and alters oligomeric state but the resulting changes in structure and dynamics are non-uniform. Full-length FRQ undergoes liquid-liquid phase separation (LLPS) in a phosphorylation dependent manner to sequester FRH and CK1 and influence CK1 enzymatic activity. Live imaging of *Neurospora* hyphae reveals FRQ foci near the nuclear periphery characteristic of condensates. Analog clock repressor proteins in

34 higher organisms share little position-specific sequence identity with FRQ; yet, they contain
35 amino-acid compositions that promote LLPS. Hence, condensate formation may be a conserved
36 feature of eukaryotic circadian clocks.
37
38 **Significance**
39
The central transcriptional repressor proteins of circadian clocks are often highly disordered,
undergo extensive post-translational modification and have poorly defined structural properties.
The fungal repressor protein Frequency (FRQ) nucleates a semi-ordered molecular assembly
that despite conformational heterogeneity arranges its components in close proximity and
facilitates their sequestration and enzymatic regulation through post-translational modification
and liquid-liquid phase separation. Analogous properties are predicted by the similar amino acid
compositions of clock proteins in insects and mammals.
47
48 **Introduction**
49
Eukaryotic circadian rhythms of biological phenomena arise from periodic gene expression
patterns that oscillate independent of external cues but can be entrained to light and
temperature (1). At the molecular level, circadian clocks function as transcription-translation
negative-feedback loops (TFFLs) that consist of core positive and negative elements (Fig. 1A
(2)). In the long-standing model clock of *Neurospora crassa*, the negative transcriptional
repressor complex is known as the FFC: named for its three principal components: Frequency
(FRQ, Fig. 1B), the FRQ-interacting RNA helicase (FRH) and Casein Kinase 1 (CK1) (1, 3).
FRQ is functionally analogous to Period (PER) in insects and mammals, whereas CK1 is an
ortholog of the mammalian CK1 and the Drosophila kinase Doubletime (DBT) (4). FRH is found
only in the core repressor of the fungal clock. Although FRH is homologous to the yeast
helicase MTR4, FRH helicase activity does not appear essential to clock function (5, 6). The
FFC represses the positive arm of the cycle, which involves two zinc-finger transcription factors,
White-Collar 1 (WC-1) and White-Collar 2 (WC-2) assembled into the White-Collar Complex
(WCC).
64
WCC binding to the clock-box (c-box) activates *frq* expression (Fig. 1A (7)). FRQ production is
also highly regulated at the post-transcriptional level through the use of rare codons to retard
translation, the expression of an anti-sense transcript and alternative splicing mechanisms (3, 4,

8). The latter leads to the production of either a long (L-FRQ) or short (S-FRQ) isoform, and the

ratio of these isoforms fine tunes period length in response to ambient temperatures (8-10).

Translated FRQ binds to FRH via its FFD domain and CK1 via its two FCD domains (1 and 2) to

form the repressive FFC (Fig. 1C). The FFC enters the nucleus where it directly inhibits the

WCC and thereby down-regulates *frq* expression (11). FRQ-mediated WCC phosphorylation by

CK1 (and associated kinases) reduces interactions between the WCC and the c-box to close

the feedback loop and restart the cycle (Fig. 1A (12)). FRH facilitates interactions between the

FFC and the WCC (6) and thus, repressor function likely depends upon the FRQ-mediated

association of CK1 and FRH. Whereas interaction regions on the FRQ protein sequence have

been identified (Fig. 1C), there is little information on the spatial arrangement of the FFC

components.

Despite low sequence conservation among the core negative arm proteins across species, they

all share a high degree of disorder (13). FRQ, the analogous Drosophila (d) Period (PER)

protein and the analogous mammalian (m) PER have > 65% disorder predicted in their

sequences with ~86% of the FRQ sequence assumed to have no defined structure (13). Both

FRQ and mPER2 have biochemical properties characteristic of intrinsically disordered proteins

(IDPs) (5, 11, 14).   These clock repressors also undergo extensive posttranslational

modification (PTM), particularly phosphorylation (13, 15). FRQ acquires over 100

phosphorylation sites over the course of the circadian day (similar to PER phosphorylation by

DBT), predominantly due to CK1 (16, 17). The phosphorylation events have two consequences:

1) those that weaken the interaction between FRQ and the WCC or CK1 (3, 18); and 2) those

that direct FRQ down the ubiquitin-proteasome pathway (19), although only the former are

required for circadian rhythmicity (19). Multi-site phosphorylation of FRQ influences its inhibitory

function and clock period length (18, 19). Although the consequences of phosphorylation-driven

conformational changes in FRQ have been probed by proteolytic sensitivity, the challenges of

producing FRQ have limited direct assessment of its conformational properties (13, 18, 20).

IDPs such as FRQ are known to participate in liquid-liquid phase separation (LLPS), which has

emerged as a key mechanism for organizing cellular components, coordinating metabolism and

directing signaling through, for example, the formation of membraneless organelles (MLOs) (21,

22). Emerging studies in plants (23) and flies (24) implicate LLPS in circadian clocks and in

Neurospora it has recently been shown that the Period-2 (PRD-2) RNA-binding protein

influences *frq* mRNA localization through a mechanism potentially mediated by LLPS (25).

Intriguing questions surround why clock repressor proteins are so disordered, undergo
extensive post-translational modification and whether LLPS has a role to play in their function.
103
Clock proteins and their complexes have been challenging to characterize due to their large size
(>100 kDa), low sequence complexity, extensive disorder and modification states. Herein, we
reconstitute FRQ and the FFC to assess the global and site-specific structural properties of
these moieties, as well as the dependence of their properties on multi-site phosphorylation.
FRQ undergoes non-uniform conformational change upon phosphorylation and arranges FRH
close to CK1 in the FFC. We also demonstrate that FRQ indeed participates in phosphorylation-
dependent LLPS, and as a condensate sequesters FRH and CK1 while altering CK1 activity.
Our results show how a massively disordered protein such as FRQ organizes the clock core
oscillator and reveal the effects of extensive phosphorylation on the conformational and phase-
separation properties of a large IDP.
114
115   **Results:**
116
<u>Sequence assessment and general properties of FRQ</u>
Analysis of an AlphaFold 2.0 (26) model of full-length FRQ indicates that FRQ lacks a defined
arrangement of large folded domains; instead, most of the molecule is unstructured (Fig. 1B), in
agreement with previous studies predicting that FRQ mostly comprises intrinsically disordered
regions (5). However, FRQ is also predicted to contain a small, clustered core, with several
secondary structural elements that collapse together to form a compact nucleus surrounded by
long flexible disordered regions (Fig. 1B). The AlphaFold pLDDT scores, which represent the
confidence in the structural prediction (26), are very low across the model, except within the
core (Fig. 1B, *SI Appendix*, Fig. S1), which contains the CK1 binding region. The FRH binding
region falls within a more flexible region with a lower pLDDT score than that of CK1. Helices
compose the majority of secondary structural elements predicted by AlphaFold (Fig. 1B). The
assignments of the helical regions were well supported by the Agadir program that predicts
helical propensity based on the helix/coil transition theory (27). Four regions distributed
throughout the length of FRQ (residues 194-212, 312-329, 709-719 and 801-817) have at least
10% helical propensity and these helical regions mostly agree with those of the AlphaFold
model (Fig. 1B). Together, these analyses primarily indicate that FRQ assumes a broad
conformational ensemble containing an ordered nucleus that interacts extensively with partially
ordered and flexible regions. However, much of the structural details of the model, including the

positioning and interactions between secondary structural elements, are predicted with low
confidence.

<u>Reconstitution of FRQ in unphosphorylated and highly phosphorylated states</u>

The size and intrinsic disorder of FRQ have made it challenging to recombinantly express and
purify, thereby limiting investigations of its biophysical properties. Previous purifications have
characterized FRQ as an IDP, but suffered from low yields and aggregated species (11, 28).
We also encountered extensive protein degradation and aggregation with conventional
purification methods and affinity tagging strategies; however, methods proven effective for IDPs
(29) yielded comparatively well-behaved protein. Briefly, we carried out Nickel affinity
chromatography and size exclusion chromatography (SEC) under denaturing conditions to
maximize yield and reduce aggregation of full-length L-FRQ (residues 1-989; Fig. 2A) (30).
Solubilization with high concentrations of guanidinium-HCl prevented proteolysis, aggregation
and limited co-purification of impurities that were otherwise difficult to remove. Once FRQ was
isolated, the protein remained soluble, monodisperse and could be concentrated after the
denaturant was removed by dialysis (Fig. 2A,B). To our knowledge, FRQ is one of the largest
native IDPs produced in amounts amenable to structural and biophysical characterization.

To generate phosphorylated FRQ, the protein was co-expressed with (untagged) CK1, which
forms a stable complex with FRQ and acts as its primary kinase (20). The same denaturing
purification procedure was followed as for FRQ expressed without CK1. Phosphorylation was
confirmed visually by SDS-PAGE (Fig. 2B) and mass spectrometry (*SI* Appendix, Fig. S2). The
latter revealed a network of > 80 phosphorylation sites (*SI* Appendix, Fig. S2), most of which
agree with those found on natively expressed FRQ (16, 17). Hereafter, we refer to this
preparation as phosphorylated FRQ (p-FRQ) whereas non-phosphorylated FRQ (np-FRQ)
refers to FRQ expressed without CK1. In general, we found that co-expression of FRQ with CK1
increased protein yield and stability, while limiting aggregation and degradation.

<u>Global properties of FRQ</u>
Size Exclusion Chromatography coupled to Multi Angle Light Scattering (SEC-MALS) of p-FRQ
gives a molecular weight measurement consistent with a primarily monomeric state (MW of
~143 kDa versus a predicted MW of ~110 kDa; Fig. 2B). In contrast, the np-FRQ measured by
MALS indicates a dimer and its measured mass distribution increases upon concentration (Fig.

2B) Hence, purified np-FRQ dimerizes, whereas phosphorylation destabilizes this association.

SEC-coupled small-angle x-ray scattering (SAXS) analyses revealed that both p-FRQ and np-

FRQ shared characteristics of highly flexible proteins, but that p-FRQ was more extended than

np-FRQ (*SI Appendix* Table S1). The dimensionless Kratky plots of p-FRQ and np-FRQ, which

report on compactness and conformational dispersity (31-34), revealed high degrees of

flexibility for both proteins (Fig. 2C). The peak values of the plot were shifted from those

expected for compact structures but were also not indicative of fully disordered proteins (33, 35,

36). Furthermore, p-FRQ appears more disordered than np-FRQ; a characteristic also reflected

in their pairwise distance distributions (*SI Appendix*, Fig. S3). These data then indicate that FRQ

generally adopts a relatively monodisperse conformational state that is partially compact but

highly flexible and then increases expanse and flexibility upon phosphorylation. This general

picture is consistent with the increased susceptibility to proteolysis that FRQ experiences with

phosphorylation (13, 18, 20).

<u>Local measures of FRQ structure and dynamics</u>

The properties of FRQ that include its intrinsic disorder and propensity to form heterogenous

ensembles are not well suited to commonly applied structural characterization methods. We

applied site-specific spin labeling (Fig. 3A) and electron spin resonance spectroscopy (ESR) to

monitor FRQ local structural properties. FRQ has four native cysteine residues that provide

reactive sites for nitroxide spin-labels (Fig. 3A,B (37)). The Cys residues distribute relatively

evenly throughout the protein sequence and fall within areas of varying structural order that

include the binding regions for the core clock proteins (FRH, CK1, WC1 and WC2; *SI Appendix*,

Table S2-3; Fig. 3B). Furthermore, we developed and applied enzymatic SortaseA & AEP1

peptide-coupling methods (38, 39) to spin-label FRQ at its N- and C- termini (Fig. 3A; *SI*

Appendix Fig. S4).

Line-shape analysis from continuous wave (cw)-ESR spectroscopy reports on local dynamics

(40, 41). For rapidly rotating nitroxide spin labels (SLs), the characteristic spectrum consists of

three narrow lines of equal width and intensity. As the rotational mobility decreases, the lines

broaden, especially the high-field line (42, 43). Cw-ESR reveals that FRQ has regions of varying

order that agree with computational predictions and that some of these sites undergo

phosphorylation-dependent conformational changes (Fig. 3C). Spin-labels attached to Cys490

and Cys658 (490SL and 658SL) are predicted to be within an α-helix and a β-strand

respectively. These sites correspondingly showed a broad low-field (left-most line) component
and a small high-field component (right-most line) indicative of slower tumbling compared to
sites predicted to be in flexible regions, such as 746SL (Fig. 3C). The 490SL site had a broader
profile compared to that of the 658SL site, although both are relatively well ordered. We
observed a similar trend with δ, which is the peak-to-peak width of the central line (44). 490SL
and 658SL had similar δ values that were higher than that of 746SL, indicating that they are less
flexible than 746SL (*SI Appendix* Table S4). Furthermore, the spectra also highlighted structural
differences between the phospho forms: p-FRQ and np-FRQ gave very different δ values at
490SL and the C-terminus, thereby demonstrating how phosphorylation can influence
conformational properties site-specifically. The C-terminus may have become more dynamic in
p-FRQ as compared to np-FRQ because the C-terminus contains a cluster of phospho-sites
(Fig. 3C). In contrast, the N-terminus and especially 490SL, which locates to the CK1-interaction
region of the FCD domain, are more ordered in p-FRQ. Such differences in mobility between p-
FRQ and np-FRQ in the FCD (490SL) may reflect changes in FRQ:CK1 interactions as FRQ
becomes progressively phosphorylated across the circadian period (18).
218
Double electron–electron resonance (DEER) spectroscopy provides distance distribution
between two nitroxide labels in the range of 15 Å to ~100 Å. We produced 6 cysteine pair
variants that each contained only two of the four native FRQ Cys residues to measure pair-wise
distances and one single Cys variant to test for oligomerization. Owing to association and
aggregation when concentrated, np-FRQ was not amenable to DEER measurements; however,
such issues were not prohibitive with p-FRQ. Consistent with the MALS data showing p-FRQ to
be monomeric, the single Cys spin-labeled variant 490SL produced very little dipolar signal that
would otherwise indicate dimerization (*SI Appendix Fig. S5*). Likewise, the double 370SL:746SL
p-FRQ variant also gave no dipolar signals and hence no dimerization as reported by those two
sites (Fig. 3D). The DEER experiments of the Cys pairs generally revealed slow decaying time
domain traces when at least one of the two cysteine residues resides outside of the predicted
core (Fig. 3D; *SI Appendix* Fig. S5-6). Such traces are indicative of long inter-spin distances and
conformational flexibility, which would be consistent with the large mobility and intrinsic disorder
of these peripheral regions (*SI Appendix*, Fig. S5-6). Only variant 490SL:658SL gave a fast-
decaying time-domain signal, indicative of short inter-spin distances and conformational rigidity
(*SI Appendix*, Fig. S5-6). Despite the overall uncertainty in the AlphaFold model (Fig. 1B) it is
notable that 490SL and 658SL both fall within the predicted ordered core.
236

<u>FFC reconstitution and component arrangement.</u>

The FFC was generated by mixing p-FRQ or np-FRQ with excess FRH and CK1 followed by
SEC purification to give p-FFC or np-FFC, respectively. FRHΔN (residues 100-1106), which
binds FRQ, was used to improve expression and enhance stability (6, 11). The resulting
symmetric SEC traces and SDS-PAGE gels revealed pure and homogenous ternary complexes
in each case (Fig. 4A). As expected, CK1 phosphorylates FRQ in np-FFC (Fig. 4A, *SI Appendix*
Fig. S7) and this activity does not require priming by other kinases (14). SEC-SAXS analysis of
the FFC showed that both FRQ phospho-forms produced complexes that were more globular
than FRQ alone, but still somewhat flexible, with the p-FFC being more flexible than the np-FFC
(Fig. 4B, *SI Appendix*, Fig. S3). MW estimates of the p-FFC and np-FFC from SAXS indicate a
1:1 stoichiometry of FRQ:FRH, with a likely single component of CK1 also contained within the
complex (*SI Appendix* Table S1).

DEER measurements from selectively spin-labeled components within the complex were used
to probe its overall architecture (Fig. 5; *SI Appendix* Fig. S8-S11). An ADP-βSL probe (45)
bound to both FRH and CK1 and revealed that the FRH and CK1 ATP-binding pockets are
closely positioned within the FFC complex (Fig. 5A; Fig. *SI Appendix* S8). Nonetheless, the
presence of broad and multiple peaks in the distance distributions indicated conformational
heterogeneity. Spin labels placed within the FCD2 and FFD domains of FRQ (490C and 746C),
which bind CK1 and FRH respectively, produced a distance distribution similar to that of the
labels that target the ATP binding sites of the two enzymes (Fig. 5A) (46). These independent
observations support the conclusion that FRQ organizes FRH and CK1 in close proximity within
the FFC.

Spin-label separations within p-FRQ changed substantially when FRH and CK1 bound (Fig. 5;
SI Appendix Fig. S5-6,S9-11). In general, the resulting distance distributions had smaller mean
distances and less breadth (Fig. 5B-H). However, the changes were not uniform; for example,
the 658SL-to-746SL distance increased in the complex compared to free FRQ (Fig. 5G). To
further assess how the distributions p-FRQ conformations changed when the p-FFC formed, the
distance data was fit to a model of one near and one far Gaussian functions (*SI Appendix*, Fig.
S11; *SI Appendix* Table S5). Shifts in population between the near and far components were
most prominent for the 370-658, 490-658 and 490-746 variants, which contain sites involved in
core clock interactions (*SI Appendix* Fig. S11). The proportion of the closer component

increased from ~30% to ~70% in the 370-658 variant (*SI Appendix* Fig. S11) and from ~10% to

~70% in the 490-746 variant (Fig. 5F) when bound to FRH and CK1. Similar trends were

observed when we compared the percentage change of the average separation observed in

FRQ alone versus when it is bound to FRH and CK1 (Fig. 5H). The 490, 658 and 746 sites

reside in regions important for WCC interactions (47) and FRH may organize such binding

determinants. Notably, the 370SL:746SL spin pair showed little dipolar coupling in free p-FRQ

but resolved into an observable mid-range separation in the presence of CK1 and FRH (Fig.

5C), perhaps because position 746 is within the FRH binding region. Thus, CK1 and FRH order

p-FRQ to an appreciable degree and the two enzymes are adjacently positioned within the

conformationally heterogeneous complex.

<u>FRQ exhibits behavior consistent with LLPS *in vitro* and *in vivo*</u>

The unstructured regions and multivalent interactions of IDPs can lead to liquid-liquid phase

separation (LLPS). Such de-mixing often manifests in the formation of biomolecular

condensates within cells (48). As has been previously noted (13), FRQ may be capable of LLPS

owing to its extensive disorder. We found that FRQ has a high propensity to undergo LLPS

based on a set of computational tools that evaluate charge distribution and clustering, long-

range aromatic and non-aromatic π:π contacts and nucleic acid binding propensies (49, 50).

The overall FRQ PScore, which grades LLPS propensity on these properties (50), was 6.39,

with three regions scoring above a value of 2, the threshold for favorable LLPS (*SI Appendix*,

Fig. S12). The region with the highest PScore also accounted for the majority of LLPS

propensity as determined by the catGRANULE prediction algorithm (49). Furthermore, we found

that these methods also predicted the functional analogs dPER and hPER1 to phase separate

(Fig. 6A, *SI Appendix*, Fig. S13). The FRQ functional analogs share physicochemical

characteristics with FRQ, particularly an enrichment of hydrophilic residues, increases in Gly

and Pro content and a depletion of hydrophobic residues, even though their positional sequence

conservation with FRQ is low (*SI Appendix*, Fig. S14 and S15). Similar amino acid composition

patterns among FRQ, dPER and hPER1, include the most common residues in each protein

and its UniRef50 cluster (sequences with at least 50% identity with the target protein) being

serine, proline and glycine (~30% of total sequence) accompanied by a high depletion in

aromatic residues (only ~7.5%) (*SI Appendix*, Fig. S15). In addition, all of the sequences have a

reduced proportion of hydrophobic residues such as A, I, L, and V and a slightly acidic

isoelectric point. Thus, FRQ, dPER and hPER1 all have low overall hydropathy (*SI Appendix*,

Fig. S14). However, in addition, FRQ and its homologs are unusual in their high Arg and Asp

content, which tend to separate in charge blocks throughout their sequences (as indicated by

high kappa values, *SI Appendix*, Fig. S15). Such charge clustering correlates with LLPS (51).

Overall, this analysis suggests that like intrinsic disorder, LLPS driven by the physiochemical

properties of the constituent residues, may be a conserved feature in the negative arm of the

circadian clock.

We tested whether both np-FRQ and p-FRQ can undergo LLPS *in vitro.* In each case, upon

exchanging the purified protein into their respective phase separation buffers, we observed the

appearance of a turbid solution that contained microscopic droplets (Fig. 6B). These droplets

scaled in size and number with increasing protein concentration, dissolved upon the addition of

1,6-hexanediol, and were shown to dock and fuse to one another, thereby confirming their

liquid-like properties (Fig. 6B, *SI Appendix*, Fig. S17). UV-Vis turbidity assays as a function of

temperature revealed that FRQ undergoes a low critical solution temperature (LCST) phase

transition that is concentration and phosphorylation status-dependent. (Fig. 6C, *SI Appendix*

Tables S6-7). As expected, phase separation increased as temperature and protein

concentration increased as indicated by growing amplitudes and earlier phase separation

initiation temperatures (Fig. 6C; *SI Appendix*, Table S6-7). Lower salt concentrations favored

the LLPS of p-FRQ, whereas higher salt disfavored LLPS, compared to higher salt

concentrations favoring LLPS for np-FRQ *(SI Appendix*, Fig. S17). Although the NaCl

concentration for np-FRQ LLPS conditions is non-physiological *per se*, the ionic strength

conditions and effective solute concentrations in cells are comparatively high (52). Furthermore,

the higher absorbance values of p-FRQ at lower temperatures relative to that of np-FRQ might

be indicative of some dual phase behavior for the phosphorylated isoform (Fig. 6C). Thus, FRQ

phosphorylation alters its behavior consistent with LLPS as evidenced by its non-zero,

concentration-dependent absorbance at lower temperatures, mostly likely by increasing the

relative negative charges, changing the charge patterns, and providing more sites for

multivalent interactions within the molecule (Fig. 6C, *SI Appendix*, Fig. S16-17)(51, 53).

Live cell imaging of *Neurospora crassa* expressing FRQ tagged with the fluorescent protein

mNeonGreen at its C-terminus (FRQ[mNeonGreen]) revealed heterogeneous patterning of FRQ in

nuclei (Fig. 6D,E). Bright and dynamic foci were observed near the nuclear periphery, which is

delineated by the cytoplasm facing nucleoporin Son-1 tagged with mApple at its C-terminus

(Fig. 6D,E *SI Appendix*, Movie S1). Such foci are characteristic of phase separated IDPs (25,

54-56) and share similar patterning to that seen for clock proteins in *Drosophila* nuclei (24).

Phase separated FRQ recruits FRH and CK1 and alters CK1 activity

Phase separated FRQ recruits FRH and CK1 and alters CK1 activity

The PScore and catGRANULE LLPS predictors gave low overall scores for the FRQ partners,

FRH and CK1 (*SI Appendix*, Fig. S13). Nonetheless, FRH and CK1 may still act as clients that

localize to the FRQ-formed LLPS-like droplets. Indeed, fluorescently-labeled FRH and CK1

were recruited into phase separated FRQ droplets, but a control protein labeled with the same

fluorophore, *E. coli* CheY, was not (Fig. 7A). The ability of FRQ to recruit its partner proteins into

phase separated droplets provides a means to sequester them and potentially control their

availabilities and activities.

LLPS has been suggested to modulate the activities of enzymes by creating a

microenvironment to alter substrate-sequestration and product-release (57). We tested the

effect of FRQ LLPS on its phosphorylation by CK1, a key reaction for regulating the period of

the clock (18). We compared the phosphorylation of FRQ by CK1 in phase separation buffer

under different temperatures, using the latter as a means to control the degree of LLPS. Both

the UV-Vis turbidity assays as well as the DIC micrographs of the protein at different

temperatures establish that FRQ LLPS increases substantially with temperature (Fig. 6C;

Fig.7C). CK1 autophosphorylation served as an internal control and its temperature

dependence was consistent with previous studies that show increased catalytic activity at higher

temperatures (14). As temperature increases drove more LLPS, the phosphorylation of FRQ

decreased, even as CK1 autophosphorylation increased (Fig. 7B; *SI Appendix* Fig. S19).

Quantitatively, we observed a reduction in FRQ phosphorylation by two-fold under LLPS

conditions relative to conditions under which it does not phase separate (Fig. 7B). Through

application of cw-ESR we probed FRQ dynamics within the condensates. We observed a

striking rigidification of spin labeled FRQ when LLPS was promoted by buffer conditions (Fig.

7D). The reduction of CK1-mediated FRQ phosphorylation, which likely requires substantial

dynamics given the distribution of phosphorylation sites throughout the FRQ sequence, may

reflect this change in molecular dynamics. Thus, in addition to affecting the localization of FRQ

and its partners, LLPS also impacts the enzymatic activity of the complex, by perhaps

modulating molecular encounters and motions essential for large scale phosphorylation.

Discussion

Discussion

Within the TTFLs that define eukaryotic circadian clocks, large repressor proteins like FRQ

compose cellular timing mechanisms by coordinating transient protein-protein interactions,

nuclear entry, targeted protein degradation, and transcriptional repression (13). They compose

dynamic macromolecular assemblies that achieve specificity in their functions despite assuming

highly variable structures (13). Our purification scheme enabled a multi-pronged biophysical

characterization of FRQ. The majority of previous such studies on IDPs, particularly those that

undergo de-mixing, have involved truncated proteins, excised domains or peptides of sizes less

than 200 residues (30, 42, 58-60). FRQ represents a full-length, nearly 1000-residue IDP that

can be investigated with respect to post-translational modification, interaction partners and

phase separation. As anticipated from its sequence (5, 13), this direct characterization of

structural properties shows that FRQ is indeed a highly dynamic protein, but even in isolation, it

is not completely disordered. The SAXS and ESR data indicate that FRQ has compact features

that localize to an ordered core interspersed with dynamic regions. Notably the SAXS and

DEER data suggest a degree of disorder that is substantially greater than that expected of a

molten globule (61), but far from that of a completely denatured protein (34, 62). Although FRQ

lacks the well-defined PAS domains of the analogous PER proteins (13), it also maintains a

structured nucleus to anchor its more dynamic elements. The regions that bind CK1 and FRQ

are the most ordered and they organize the two partners beside each other, in fact, they are in

remarkably close proximity, as judged by spin-labels in their ATP binding pockets or in their

binding domains. FRH mediates contacts to the WCC (6), whereas CK1 phosphorylates the

WCC to close the TTFL (3, 12, 18, 19); hence, CK1 targeting to the WCC likely depends on

FRQ associating FRH with CK1 (12).

Previous proteolytic sensitivity studies have suggested that increased phosphorylation across

the day transitions FRQ from a "closed" to a more "open" state (13, 18, 20). Our data paint a

more nuanced picture. The SAXS data reveal that p-FRQ and p-FFC are more expansive and

show greater flexibility than do np-FRQ and np-FFC, respectively. Furthermore, np-FRQ has a

greater tendency to self-associate. In the cell, completely unphosphorylated FRQ is likely short-

lived, if present at all (16). Phosphorylation may stabilize FRQ against aggregation, yet also

provides for enough conformational heterogeneity to poise FRQ near a degradation threshold

(19). This idea is supported by the cw-ESR data (Fig. 3), which shows that phosphorylation

causes non-uniform local structural changes with some regions increasing, while others

decreasing, in flexibility. The DEER data also demonstrates that binding to FRH and CK1

increases compactness of p-FRQ, particularly in the central-core that is responsible for binding

the partners. In contrast, the FRQ C-terminus gains considerably flexibility in the FFC. FRQ has

been suggested to assume a dimeric state because of a predicted coiled-coil region and

formation of mixed dimers formed in cellular pull-down assays with differentially tagged FRQ

subunits (63). Np-FRQ does indeed dimerize when isolated, whereas p-FRQ is primarily

monomeric alone. Furthermore, SAXS data indicates there is one copy of FRQ in both the

reconstituted np-FFC and p-FFC. Hence phosphorylation has the potential to regulate the FRQ

oligomeric state, which may in turn be linked to FFC assembly. Despite being a monomer in

isolation, the pull-down data (63) indicates that association of p-FRQ may be mediated by other

cellular factors, including LLPS.

FRQ has physicochemical features embedded within its sequence that give rise to LLPS and

the FRQ clock analogs exhibit similar characteristics (*SI Appendix*, Fig. S13-15) FRQ and

negative-arm proteins in general belong to the "flexible disorder" class of IDPs (57), wherein

amino acid patterns that generate intrinsic disorder are conserved, but amino acid sequence per

se, is not (64). Moreover, although these proteins all clearly contain regions of low-complexity

sequence (11, 13, 14) their conserved features include not only their intrinsic disorder, but also

their propensity for LLPS (Fig. 6A, *SI Appendix*, Fig. S13). These two properties, although

related, are not directly correlated (49, 50, 53). In particular, a preponderance of aromatic and

non-aromatic π-π planar contacts and charge-dense or charge-repeat regions drive LLPS (50,

53) and these features are consistently found in clock repressors, particularly FRQ. The similar

amino acid compositions of these proteins may then potentially give rise to their shared function.

Moreover, extensive phosphorylation may alter LLPS propensity because it increases extended

states, solvation of the peptide backbone, charge density and intermolecular hydrogen bonding

capacity (49, 50, 53). For FRQ, the sequence predictions are borne out experimentally, with p-

FRQ displaying a different phase behavior relative to np-FRQ. In addition to being more soluble,

dynamic and less aggregated than np-FRQ, p-FRQ also phase separates at lower temperatures

and under lower salt conditions relative to np-FRQ. Increased salt disfavors p-FRQ phase

separation, which when compared to np-FRQ, is perhaps consistent with more electrostatically

driven de-mixing due to the high degree of phosphorylation.

Live cell imaging of fluorescently-tagged FRQ proteins is consistent with FRQ phase separation

in *N. crassa* nuclei. The concentrated foci observed at specific positions in the nuclei indicate

condensate behavior and are highly dynamic in size and shape, similar to that observed for

other phase separating proteins (24, 25, 54-56). While ongoing experiments are exploring more

deeply the spatiotemporal dynamics of FRQ condensates in nuclei, the small size of fungal

nuclei as well as their rapid movement with cytoplasmic bulk flow through the hyphal syncytium

makes these experiments difficult. Of particular interest is drawing comparisons between FRQ

and the Drosophila Period protein, which has been observed in similar foci that change in size

and subnuclear localization throughout the circadian cycle (24).

Phosphorylation is known to modulate LLPS in other systems, but the effects are variable. For

example, phosphorylation of the transcriptional silencing factor HP1α also enhances LLPS,

whereas only the non-phosphorylated form of Xenopus CPEB4 involved in translation of mRNA

poly-A tails undergoes LLPS (30, 65). FRQ, especially p-FRQ, may achieve its inhibitory

function by sequestering the FFC and WCC into membraneless organelles. The large,

disordered domains and capacity for multivalency of the WC proteins (13) may also facilitate

their partitioning into such condensates. In the fly clock, PER and the WCC analog CLK,

produce liquid-like foci in the nuclei that are most prevalent during the repressive phase of the

circadian cycle (24). These large foci, which are few and localized to the nuclear periphery, may

serve to organize clock-controlled genes during repression (24). LLPS may also influence the

nuclear transport of FRQ by affecting interactions with the nuclear import machinery as has

been observed with proteins such as TD43 and FUS (66, 67). Conditions favoring LLPS reduce

the ability of FRQ to act as a substrate for CK1 (Fig. 7B), and this reactivity correlates with

reduced dynamics of FRQ in the phase separated state (Fig. 7D). Domains and disordered

regions from other spin-labeled proteins show similar reduced dynamics when induced to phase

separate and monitored by cw-ESR spectroscopy (30, 58, 59, 68), although the temperature

and salt dependencies of the ordering behavior can vary considerably, with some species

showing little change upon LLPS (60).

In the cell, the impact of LLPS on CK1 activity may modulate the timing of FRQ

phosphorylation, which is closely linked to clock period (14, 19). Moreover, and collectively, the

data from Fig. 6 and 7 show that as temperature increases, LLPS-like behavior increases, and

as LLPS increases, the ability of CK1 to phosphorylate FRQ decreases. The rate of FRQ

phosphorylation dictates period length (19). Therefore, this negative feedback of temperature

on period mediated by LLPS formation could provide a mechanism for circadian temperature

compensation of period length (*SI Appendix* Fig. S19). Sequestration of CK1 (and FRH) into

condensates may also render these factors unavailable to target other cellular substrates.

Phosphorylation alters FRQ interactions throughout the circadian cycle, not unlike the workings

of the cyanobacterial clock, which relies upon a timed phospho-code to regulate enzymatic

components (12, 69). We demonstrate that FRQ likewise responds to phosphorylation with non-

uniform conformational changes across its sequence and an increased propensity for LLPS. It is

the latter property that appears conserved among negative-arm repressor proteins in eukaryotic
clocks and may thus be a critical function of their intrinsic disorder and extensive post-
translational modifications.
478

Materials and Methods
FRQ was obtained by recombinant expression in *E. coli* and purification under denaturing
conditions followed by removal of denaturant. FRH and CK1 were produced using more
standard protocols. SEC-MALS was performed with Wyatt Technologies DAWN HELEOS-II
multi-angle light scattering detector and SEC-SAXS was carried out at the Cornell Synchrotron
G1 station. Protein labeling for ESR and microscopy experiments made use of enzymatic
labeling and reactive cysteine residue probes. Cw-ESR spectroscopy was carried out on a
Bruker E500 ESR spectrometer operating at X-band, whereas DEER spectroscopy utilized a
four-pulse protocol on a Bruker E580 spectrometer operating at Q band. Turbidity was as
measured by absorbance at 300 - 600 nm under different buffer conditions and a Peltier-driven
temperature ramp. Microscopy was carried out on a Zeiss Axio Imager Upright Trinocular
Fluorescence Microscope at 100x magnification. Phosphorylation assays involved reaction with
[γ-^{32}P] ATP, protein band resolution by SDS-PAGE and phosphorimage analysis. Mass-
spectrometry and phospho-peptide detection was carried out on tryptic peptides analyzed by
MS/MS with a CID ion-trap. *Neurospora crassa* strain generation targeted the endogenous *frq*
locus with DNA encoding *N. crassa* codon-optimized mNeonGreen appended to the C-terminus
of the *frq* ORF. For live-cell imaging a section near the edge of a growing mycelium was
removed and inverted on a chambered cover glass. Images were collected with 488 nm laser
excitation using a laser scanning confocal microscope equipped with a spinning disk scan head
and a CMOS camera. Detailed descriptions of the protocols can be found in the *SI Appendix*
Materials and Methods
500

501 Author contributions: D.T., N,M., B.M.B., J.C.D., A.B. and B.R.C. designed research; D.T., N.M.,
502 B.M.B., and S.C. performed research; D.T., N,M., B.M.B., S.C., J.C.D., A.B. and B.R.C.
503 analyzed data; D.T. and B.R.C. wrote the paper with contributions and editing from all authors.
504

505 The authors declare no competing interests.
506

Data Availability. The mass spectrometry FRQ phosphorylation data have been deposited to

the ProteomeXchange Consortium via the PRIDE partner repository with the dataset

identifier PXD037938. All other data are included in the article and/or supporting

Information.

Acknowledgements.

We thank the Cornell High Energy Synchrotron Source (CHESS) and the National Biomedical

Center for Advanced ESR Technologies (ACERT) for access to data collection facilities. This

work was supported by grants from the National Institutes of Health: R35GM122535 to BRC,

R35GM118021 to JCD; R35GM138097 to AB, 1F31GM143890 to NM, and T32-008704 to BMB

and from The Pew Charitable Trusts to AB. ACERT is supported by P41GM103521 and

R24GM146107. CHESS is supported by NSF award DMR-1332208 and NIH/NIGMS award

P30GM103485.

References

522
1.   J. M. Hurley, J. J. Loros, J. C. Dunlap, Circadian Oscillators: Around the Transcription-
Translation Feedback Loop and on to Output. *Trends in Biochemical Sciences* **41**, 834-
846 (2016).

2.   B. R. Crane, M. W. Young, Interactive Features of Proteins Composing Eukaryotic
Circadian Clocks. *Annual Review of Biochemistry* **83**, 191-219 (2014).

3.   J. Cha, M. Zhou, Y. Liu, Mechanism of the Neurospora Circadian Clock, a
FREQUENCY-centric View. *Biochemistry* **54**, 150-156 (2014).

4.   J. C. Dunlap, J. J. Loros, Making Time: Conservation of Biological Clocks from Fungi to
Animals. *Microbiology Spectrum* **5**, 3 (2017).

5.   J. M. Hurley, L. F. Larrondo, J. J. Loros, J. C. Dunlap, Conserved RNA Helicase FRH
Acts Nonenzymatically to Support the Intrinsically Disordered Neurospora Clock Protein
FRQ. *Molecular Cell* **52**, 832-843 (2013).

6.   K. S. Conrad *et al.*, Structure of the frequency-interacting RNA helicase: a protein
interaction hub for the circadian clock. *EMBO J.* **35**, 1707-1719 (2016).

7.   A. C. Froehlich, Y. Liu, J. J. Loros, J. C. Dunlap, White collar-1, a circadian blue light
photoreceptor, binding to the frequency promoter. *Science* **297**, 815-819 (2002).

8.   M. Zhou *et al.*, Non-optimal codon usage affects expression, structure and function of
clock protein FRQ. *Nature* **495**, 111-115 (2013).

9.   A. Diernfellner *et al.*, Long and short isoforms of Neurospora clock protein FRQ support
temperature-compensated circadian rhythms. *Febs Letters* **581**, 5759-5764 (2007).

10.   Y. Liu, N. Y. Garceau, J. J. Loros, J. C. Dunlap, Thermally regulated translational control
of FRQ mediates aspects of temperature responses in the Neurospora circadian clock.
Cell **89**, 477-486 (1997).

11.   J. M. Hurley, L. F. Larrondo, J. J. Loros, J. C. Dunlap, Conserved RNA helicase FRH
acts nonenzymatically to support the intrinsically disordered neurospora clock protein
FRQ. *Molecular Cell* **52**, 832-843 (2013).

12.   B. Wang, A. N. Kettenbach, X. Zhou, J. J. Loros, J. C. Dunlap, The Phospho-Code
Determining Circadian Feedback Loop Closure and Output in Neurospora. *Molecular*
Cell **74**, 771-784.e773 (2019).

13.   J. F. Pelham, J. C. Dunlap, J. M. Hurley, Intrinsic disorder is an essential characteristic
of components in the conserved circadian circuit. *Cell Communication and Signaling* **18**
(2020).

14.   D. Marzoll, F. E. Serrano, A. Shostak, C. Schunke, A. C. R. Diernfellner, Casein kinase 1
and disordered clock proteins form functionally equivalent , phospho-based circadian
modules in fungi and mammals. *Proc. Natl. Acad. Sci. USA* **119**, e2118286119 (2022).

15.   J. C. Dunlap, J. J. Loros, Just-So Stories and Origin Myths: Phosphorylation and
Structural Disorder in Circadian Clock Proteins. *Molecular Cell* **69**, 165-168 (2018).

16.   C. L. Baker, A. N. Kettenbach, J. J. Loros, S. A. Gerber, J. C. Dunlap, Quantitative
Proteomics Reveals a Dynamic Interactome and Phase-Specific Phosphorylation in the
Neurospora Circadian Clock. *Molecular Cell* **34**, 354-363 (2009).

17.   C. T. Tang *et al.*, Setting the pace of the Neurospora circadian clock by multiple
independent FRQ phosphorylation events. *Proc. Natl. Acad. Sci. USA* **106**, 10722-10727
(2009).

18.   X. Liu *et al.*, FRQ-CK1 interaction determines the period of circadian rhythms in
Neurospora. *Nature Communications* **10**, 4352 (2019).

19.   L. F. Larrondo, C. Olivares-Yañez, C. L. Baker, J. J. Loros, J. C. Dunlap, Decoupling
circadian clock protein turnover from circadian period determination. *Science* **347**,
1257277 (2015).

20.   C. Querfurth *et al.*, Circadian Conformational Change of the Neurospora Clock Protein
FREQUENCY Triggered by Clustered Hyperphosphorylation of a Basic Domain.
Molecular Cell **43**, 713-722 (2011).

21.   G. L. Dignon, R. B. Best, J. Mittal, Biomolecular Phase Separation: From Molecular
Driving Forces to Macroscopic Properties. *Annual Review of Physical Chemistry* **71**, 53-
75 (2020).

22.   P. E. Wright, H. J. Dyson, Intrinsically disordered proteins in cellular signalling and
regulation. *Nature Reviews Molecular Cell Biology* **16**, 18-29 (2014).

23.   J. H. Jung *et al.*, A prion-like domain in ELF3 functions as a thermosensor in
Arabidopsis. *Nature* **585**, 256-260 (2020).

24.   Y. B. Xiao, Y. Yuan, M. Jimenez, N. Soni, S. Yadlapalli, Clock proteins regulate
spatiotemporal organization of clock genes to control circadian rhythms. *Proc. Natl.*
Acad. Sci. USA **118**, e2019756118 (2021).

25.   B. M. Bartholomai, A. S. Gladfelter, Loros, J. J., J. C. Dunlap, PRD-2 mediates clock-
regulated perinuclear localization of clock gene RNAs within the circadian cycle of
Neurospora. *Proc. Natl. Acad. Sci. USA* **119**, e2203078119 (2022).

26.   J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature*
2021 596:7873 **596**, 583-589 (2021).

27.   V. Muñoz, L. Serrano, Elucidating the folding problem of helical peptides using empirical
parameters. *Nature Structural Biology* **1**, 399-409 (1994).

28.   L. Lauinger, A. Diernfellner, S. Falk, M. Brunner (2014) The RNA helicase FRH is an
ATP-dependent regulator of CK1a in the circadian clock of Neurospora crassa. in *Nature*
communications.

29.   C. P. Pedersen, P. Seiffert, I. Brakti, K. Bugge, Production of intrinsically disordered
proteins for biophysical studies: Tips and tricks. *Methods in Molecular Biology* **2141**,
195-209 (2020).

30.   M. Seal *et al.*, Evolution of CPEB4 Dynamics Across its Liquid–Liquid Phase Separation
Transition. *The Journal of Physical Chemistry B* **125**, 12947-12957 (2021).

31.   J. Trewhella *et al.*, 2017 publication guidelines for structural modelling of small-angle
scattering data from biomolecules in solution: An update. *Acta Crystallographica Section*
D: Structural Biology **73**, 710-728 (2017).

32.   D. Durand *et al.*, NADPH oxidase activator p67phox behaves in solution as a
multidomain protein with semi-flexible linkers. *Journal of Structural Biology* **169**, 45-53
(2010).

33.   A. Sagar, D. Svergun, P. Bernado, "Structural Analyses of Intrinsically Disordered
Proteins by Small-Angle X-Ray Scattering" in INTRINSICALLY DISORDERED
PROTEINS: Methods and Protocols*,* B. B. Kragelund, K. Skriver, Eds. (2020), vol. 2141,
pp. 249-269.

34.   E. W. Martin, J. B. Hopkins, T. Mittag, Small-angle X-ray scattering experiments of
monodisperse intrinsically disordered protein samples close to the solubility limit.
Methods in Enzymology **646**, 185-222 (2021).

35.   D. Durand *et al.*, NADPH oxidase activator P67(phox) behaves in solution as a
multidomain protein with semi-flexible linkers. *Journal of Structural Biology* **169**, 45-53
(2010).

36.   R. P. Rambo, J. A. Tainer, Characterizing Flexible and Intrinsically Unstructured
Biological Macromolecules by SAS Using the Porod-Debye Law. *Biopolymers* **95**, 559-
571 (2011).

37.   L. J. Berliner, J. Grunwald, H. O. Hankovszky, K. Hideg, A novel reversible thiol-specific
619   spin label - papain active-site labeling and inhibition. *Analytical Biochemistry* **119**, 450-
620   455 (1982).

621   38.   S. Chandrasekaran *et al.*, Tuning flavin environment to detect and control light-induced
622   conformational switching in Drosophila cryptochrome. *Communications Biology 2021 4:1*
623   **4**, 1-12 (2021).

624   39.   G. K. T. Nguyen, Y. Cao, W. Wang, C. F. Liu, J. P. Tam, Site-Specific N-Terminal
625   Labeling of Peptides and Proteins using Butelase 1 and Thiodepsipeptide. *Angewandte*
626   *Chemie-International Edition* **54**, 15694-15698 (2015).

627   40.   J. H. Freed (2005) ESR and Molecular Dynamics. in *Biomedical EPR, Part B:*
628   *Methodology, Instrumentation, and Dynamics* (Springer, Boston, MA), pp 239-268.

629   41.   J. M. Franck, S. Chandrasekaran, B. Dzikovski, C. R. Dunnam, J. H. Freed, Focus: Two-
630   dimensional electron-electron double resonance and molecular motions: The challenge
631   of higher frequencies. *Journal of Chemical Physics* **142**, 212302 (2015).

632   42.   L. Emmanouilidis *et al.*, NMR and EPR reveal a compaction of the RNA-binding protein
633   FUS upon droplet formation. *Nature Chemical Biology 2021 17:5* **17**, 608-614 (2021).

634   43.   A. Bonucci, M. G. Murrali, L. Banci, R. Pierattelli, A combined NMR and EPR
635   investigation on the effect of the disordered RGG regions in the structure and the activity
636   of the RRM domain of FUS. *Scientific Reports* **10**, 1-13 (2020).

637   44.   W. L. Hubbell, D. S. Cafiso, C. Altenbach, Identifying conformational changes with site-
638   directed spin labeling. *Nature Structural Biology* **7**, 735-739 (2000).

639   45.   A. R. Muok, T. K. Chua, H. Le, B. R. Crane, Nucleotide Spin Labeling for ESR
640   Spectroscopy of ATP-Binding Proteins. *Applied Magnetic Resonance* **49**, 1385-1395
641   (2018).

642   46.   P. Cheng, Q. He, Q. He, L. Wang, Y. Liu, Regulation of the Neurospora circadian clock
643   by an RNA helicase. *Genes and Development* **19**, 234-241 (2005).

644   47.   J. Guo, P. Cheng, Y. Liu, Functional significance of FRH in regulating the
645   phosphorylation and stability of Neurospora circadian clock protein FRQ. *J. Biol. Chem.*
646   **285**, 11508-11515 (2010).

647   48.   A. E. Posey, A. S. Holehouse, R. V. Pappu, Phase Separation of Intrinsically Disordered
648   Proteins. *Methods in enzymology* **611**, 1-30 (2018).

649   49.   B. Bolognesi *et al.*, A Concentration-Dependent Liquid Phase Separation Can Cause
650   Toxicity upon Increased Protein Expression. *CellReports* **16**, 1-10 (2016).

651   50.   R. M. Vernon *et al.*, Pi-Pi contacts are an overlooked protein feature relevant to phase
652   separation. *Elife* **7**, e31486 (2018).

51.   R. Somjee, D. M. Mitrea, R. W. Kriwacki (2020) Exploring Relationships between the
Density of Charged Tracts within Disordered Regions and Phase Separation. in *Pacific*
Symposium on Biocomputing (Hi), pp 207-218.

52.   B. Q. Liu, B. Poolman, A. J. Boersma, Ionic Strength Sensing in Living Cells. *Acs*
Chemical Biology **12**, 2510-2514 (2017).

53.   A. L. Szabó, A. Sánta, R. Pancsa, Z. Gáspári, Charged sequence motifs increase the
propensity towards liquid–liquid phase separation. *FEBS Letters* **596**, 1013-1028 (2022).

54.   E. Caragliano *et al.*, Human cytomegalovirus forms phase-separated compartments at
viral genomes to facilitate viral replication. *Cell Reports* **38** (2022).

55.   A. Gonzalez *et al.*, Mechanism of karyopherin-beta 2 binding and nuclear import of ALS
variants FUS(P525L) and FUS(R495X). *Scientific Reports* **11** (2021).

56.   R. Tatavosian *et al.*, Nuclear condensates of the Polycomb protein chromobox 2 (CBX2)
assemble through phase separation. *Journal of Biological Chemistry* **294**, 1451-1463
(2019).

57.   D. M. Shapiro, M. Ney, S. A. Eghtesadi, A. Chilkoti, Protein Phase Separation Arising
from Intrinsic Disorder: First-Principles to Bespoke Applications. *Journal of Physical*
Chemistry B **125**, 6740-6759 (2021).

58.   W. M. Babinchak *et al.*, The role of liquid-liquid phase separation in aggregation of the
TDP-43 low-complexity domain. *Journal of Biological Chemistry* **294**, 6306-6317 (2019).

59.   Y. X. Lin *et al.*, Liquid-Liquid Phase Separation of Tau Driven by Hydrophobic Interaction
Facilitates Fibrillization of Tau. *Journal of Molecular Biology* **433** (2021).

60.   Y. X. Lin *et al.*, Narrow equilibrium window for complex coacervation of tau and RNA
under cellular conditions. *Elife* **8** (2019).

61.   M. Kataoka, K. Kuwajima, F. Tokunaga, Y. Goto, Structural characterization of the
molten globule of alpha-lactalbumin by solution X-ray scattering. *Protein Science* **6**, 422-
430 (1997).

62.   A. G. Kikhney, D. I. Svergun, A practical guide to small angle X-ray scattering (SAXS) of
flexible and intrinsically disordered proteins. *Febs Letters* **589**, 2570-2577 (2015).

63.   P. Cheng, Y. Yang, C. Heintzen, Y. Liu, Coiled-coil domain-mediated FRQ–FRQ
interaction is essential for its circadian clock function in Neurospora. *EMBO J.* **20**, 101-
108 (2001).

64.   J. Bellay *et al.*, Bringing order to protein disorder through comparative genomics and
genetic interactions. *Genome Biology* **12**, R14 (2011).

65.   A. G. Larson *et al.*, Liquid droplet formation by HP1 alpha suggests a role for phase
separation in heterochromatin. *Nature* **547**, 236-241 (2017).

66. F. Gasset-Rosa *et al.*, Cytoplasmic TDP-43 De-mixing Independent of Stress Granules
Drives Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death. *Neuron*
102, 339-357 (2019).

67. A. Gonzalez *et al.*, Mechanism of karyopherin-β2 binding and nuclear import of ALS
variants FUS(P525L) and FUS(R495X). *Scientific Reports 2021 11:1* **11**, 1-15 (2021).

68. L. Emmanouilidis *et al.*, NMR and EPR reveal a compaction of the RNA-binding protein
FUS upon droplet formation. *Nature Chemical Biology* **17**, 608-614 (2021).

69. M. J. Rust, J. S. Markson, W. S. Lane, D. S. Fisher, E. K. O'Shea, Ordered
phosphorylation governs oscillation of a three-protein circadian clock. *Science* **318**, 809-
812 (2007).
698

Figure Legends

Figure 1: FRQ is a large, dynamic component of the clock oscillator (A) The TTFL in *Neurospora crassa*. In the core circadian oscillator, **F**RQ associates with **F**RH and **C**K1 to form a repressor complex (the **FFC**) This complex inhibits the positive-acting transcription-factor known as the white-collar complex WCC, composed of White Collar-1 (WC-1) and White Collar-2 (WC-2). The WCC activates *frq* expression by binding to the c-box region in the *frq* promoter. Stoichiometries of the components are not represented in the schematic. (B) AlphaFold derived model of LFRQ color coded by pLDDT score that highlights the confidence in each predicted region. (C) Functional domains of FRQ showing the start position for the LFRQ isoform (AUG$_{LFRQ}$.) and the four native cysteine residues. Characterized domains include: CC - coiled coil domain, FCD - FRQ:CK1 interacting domain, PEST - proline, glutamic acid, serine, and threonine-rich domain and FFD - FRQ:FRH interacting domain.

Figure 2: Purification of FRQ in non-phosphorylated and phosphorylated forms reveals IDP behavior (A) SDS-PAGE gel showing non-phosphorylated (np-FRQ) and phosphorylated FRQ (p-FRQ) alongside a molecular weight ladder. Both samples represent the purified full-length long isoform (L-FRQ, residues 1-989) in non-denaturing buffer. (B) Size exclusion chromatography–multiangle light scattering (SEC-MALS) of phosphorylated FRQ (p-FRQ) and non-phosphorylated (np-FRQ). The molecular weight (MW) of p-FRQ (~143 ± 1 kDa) is somewhat larger than that of an unphosphorylated monomer (~120 kDa), but substantially smaller than that of a dimer, whereas the high concentration MW value of np-FRQ (~261 ± 1 kDa) is slightly larger than that expected for a dimer (~240 kDa). Note that the p-FRQ MW is affected by >80 phospho sites. The shoulder to the right of the main peak had low light-scattering and a large MW error. The colored lines represent the MW of each sample. (C) Dimensionless Kratky plot of SAXS data for p-FRQ and np-FRQ, in both cases the peak positions of the curves are shifted from that expected for a globular protein ($\sqrt{3}$,1.104), but neither plateau at 2, which would be characteristic of a flexible, denatured polypeptide (32). See Table S1 for additional SAXS parameters. Total protein concentration was between 50-75 μM for the MALS and SAXS experiments.

Figure 3: ESR spectroscopy studies of spin-labeled FRQ. (A) Spin-labeling strategies for FRQ. MTSL was used for internal labeling of cysteine residues and AEP1 and SortaseA were used to label the N and C terminus, respectively. (B) AlphaFold generated model of FRQ with the positions of native spin-labeled cysteine residues highlighted in dark blue. (C) X-band (9.8 GHz) room temperature cw-ESR measurements for various spin-labeled FRQ variants, each containing a single Cys labeling site, as designated. Data represented as first-derivative spectra with the horizontal axes depicting the static field strength (typically 3330 to 3400 G), and the vertical axis depicting the change in magnetic susceptibility with respect to the static field, in arbitrary units. (D) Time-domain data of DEER measurements for double-cysteine labeled FRQ variants. The baseline-corrected time domain dipolar evolution trace fits are shown; details are given in Supplemental *SI Appendix*, Fig. S5-6. The 490-658 separation is noticeably closer and more ordered than the others. The total protein concentration was between 50-75 µM for all experiments.

Figure 4: Properties of the FFC. (A) SEC of the FFC; left inset: SDS-PAGE gel inset of the components; right inset: ^{32}P-autoradiogram of FRQ phosphorylation by CK1 and CK1 autophosphorylation at 10 and 20 µM FRQ all in the presence of FRH. Complete autoradiograms are shown in *SI Appendix*, Fig. S7. (B) Overlay of dimensionless Kratky plots of SAXS data from the FFC formed with either np- or p-FRQ (protein concentrations were between 50-75 µM). The SAXS-derived MW of np-FRQ FFC = 285 kDa, which compares to a predicted value of 273 kDa; MW of p-FFC = 340 kDa; *SI Appendix* Table S1).

Figure 5: PDS distance distributions of spin-labeled FRQ and the FFC. (A) DEER-derived distance distributions of the FFC wherein the spins were directed by ADP linkage to target the FRH and CK1 ATP-binding sites (blue) or in the form of MTSL labeling of cysteine residues within the FCD2 and FFD domains of FRQ (red). Time-domain data and distance distributions with error estimates are shown in *SI Appendix*, Fig. S5-S10. (B-G) DEER-derived distance distributions of cysteine residue pairs labeled with MTSL within p-FRQ, either alone (blue) or within the FFC (red). (H) Percentage change in the average separation of FRQ when it binds FRH and CK1. Errors are derived from the uncertainties in the distance distributions shown in *SI Appendix*, Fig. S9-10. In these experiments, only FRQ was labeled with MTSL at the positions noted. Note: Total protein concentrations were between 50-75 µM for all experiments.

Figure 6: FRQ undergoes LLPS in agreement with predictions from its sequence properties and those of its functional analogs. (A) LLPS propensity predictions using "Pi-Pi" and IUPred2A disorder predictions are shown for FRQ (purple), dPER1 (teal) and hPER1 (orange) as compared to values for proteins of similar length from their respective proteomes. The mean and standard deviations are represented as horizontal and vertical black bars, respectively. Note: **** denotes a p-value <0.0001 obtained from a Mann-Whitney U-test (B) DIC microscopy images of various concentrations of p-FRQ (in 150 mM NaCl, 25 mM Tris pH 8.0) and np-FRQ (in 500 mM NaCl, 25 mM Tris pH 8.0) at 25°C; images shown at 100x magnification. Scale bar = 10 microns (C) UV-Vis absorption spectrum of p-FRQ (in 150 mM NaCl, 25 mM Tris pH 8.0) and np-FRQ (in 500 mM NaCl, 25 mM Tris pH 8.0) as a function of temperature at increasing concentrations. Note: the shading around each curve represents the standard deviation from the mean absorbance from 2 technical replicates. 15 μM p-FRQ had the greatest standard deviation as seen by the largest shaded area. (D) Live cell fluorescent imaging of FRQ[mNeonGreen] in nuclei of *Neurospora* hyphae. Upper left panel shows the FRQ channel, represented as a multicolor heat map of fluorescence intensity; Upper right panel shows a surface plot derived from the raw FRQ image to emphasize regions of concentration; lower left shows nucleoporin SON-1[mApple], which localizes to the cytoplasmic face of the nuclear envelope; bottom right shows the FRQ:SON-1 merge image. The images were acquired on a Zeiss 880 laser scanning confocal microscope and were smoothed by bicubic interpolation during 10-fold enlargement from 42 X 42 pixels to 420 X 420 pixels. (E) FRQ[mNeonGreen] nuclear foci shown within an *N. crassa* syncytium mycelium (outlined by white lines), cropped from movie S1. Scale bar = 2μm.

Figure 7: Conditions favoring LLPS alter the structural and enzymatic properties of the FFC. (A) Differential Interference Contrast (DIC) and fluorescence microscopy images at 25 °C of phase separated np-FRQ or p-FRQ (5 µM each) droplets in 500 mM KCl, 20mM Na_2HPO_4, pH 7.4 with either equimolar Cy5-labeled FRH, CK1 or CheY (control) visualized using Cy5 fluorescence. Scale bar = 10 microns. (B) (Top) Autoradiography of FRQ (20 µM) (left) phosphorylated by CK1 in the presence of FRH at increasing temperatures. (Bottom) Schematic showing the nature and results of the autoradiography assay depicted above. The FRQ phosphorylation under LLPS conditions was reduced relative to non-LLPS conditions. The complete autoradiograms are shown in *SI Appendix*, Fig. S13. Quantification of FRQ phosphorylation by CK1 at RT under LLPS and non-LLPS conditions with phosphorylation levels under non-LLPS conditions normalized to 1. Note: *** denotes a p-value << 0.05 obtained from a student's t-test. (C) DIC microscopy images of 15 µM phase separated np-FRQ at the different temperatures studied in the phosphorylation assay shown in (B). Scale bar = 10 microns. (D) X- band (9.8GHz) RT cw-ESR spectra of FRQ labeled with MTSL at the 490 site in solubilizing buffer (black) and under conditions that promote LLPS (cyan). Data represented as first-derivative spectra with the horizontal axes depicting the static field strength (typically 3330 to 3400 G), and the vertical axis depicting the change in magnetic susceptibility with respect to the static field, in arbitrary units.

Supplementary Information

Phosphorylation, disorder, and phase separation govern the behavior of FRQ in the fungal circadian clock

Daniyal Tariq[a,*], Nicole Maurici[b,*], Bradley M. Bartholomai[c], Siddarth Chandrasekaran[a], Jay C. Dunlap[c], Alaji Bah[b] and Brian R. Crane[a].

[a]Department of Chemistry & Chemical Biology, Cornell University, Ithaca, NY,14853;
[b]Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, NY, 13210;
[c]Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth, Hanover, NH 03755.
*Denotes equal contribution

Contents

Materials and Methods:

Protein Expression and Purification

Full length L-FRQ (Uniprot ID:P19970), CK1 (CK1a; Uniprot ID:V5IR38) and FRHΔN (residues 100-1106) (Uniprot ID:Q1K502) were sub-cloned into a pET28a vector (Novagen) bearing an N-terminal His$_6$-tag and kanamycin resistance. Note: L-FRQ contained the S513A substitution to improve protein stability as noted in (1). The plasmids encoding the proteins were transformed into BL21 (DE3) *E. coli* cells, grown in LB media containing 50 µg ml^{-1} kanamycin to an optical density (OD$_{600}$) of ~0.6–0.8 at 37 °C and protein expression was induced with 1 mM IPTG at 17 °C or room temperature. Cells were harvested after 12–18 h by centrifugation at 5,000 rpm and 4 °C. FRH and CK1 were purified by standard Ni-NTA chromatography followed by ion-exchange and size-exclusion chromatography. Briefly, cells were resuspended in lysis buffer (50 mM Tris pH 8.0, 500 mM NaCl, 10% glycerol, 1 mM MgCl$_2$, 5 mM imidazole) and sonicated on ice. The lysate was clarified by centrifugation at 20,000 rpm for 1 hr. The clarified lysate was applied to a gravity Ni-NTA column and washed with 5 column volumes (CVs) of wash buffer (50 mM Tris pH 8.0, 500 mM NaCl, 10% glycerol, 1 mM MgCl$_2$, 20 mM imidazole.) The protein was eluted using the elution buffer (50 mM Tris pH 8.0, 500 mM NaCl, 10% glycerol, 1 mM MgCl$_2$, 500 mM imidazole). The protein was then diluted into IEX (ion-exchange) bufferA (50 mM HEPES pH 7.5, 10%

glycerol), applied to an S cation exchange column (GE Healthcare) and eluted over a 20 CV gradient against IEX bufferB (50 mM HEPES pH 7.5, 1 M NaCl, 10% glycerol). Fractions containing the protein were pooled, concentrated, and run over an S200 26/60 gel-filtration column pre-equilibrated in gel-filtration buffer (GFB; 50 mM HEPES, 200 mM NaCl, 10% glycerol.) L-FRQ was purified under denaturing conditions. Cells were resuspended in lysis buffer (50 mM Na_2PO_4 pH 7.42, 300 mM NaCl, 5 mM imidazole, 4 M guanidinium hydrochloride (GuHCl)) and sonicated on ice. The lysate was clarified by centrifugation at 20,000 rpm for 1 hr. The clarified lysate was applied to a gravity Ni-NTA column and washed with 3 CV of lysis buffer. The protein was eluted using the elution buffer (50 mM Na_2PO_4 pH 7.42, 300 mM NaCl, 500 mM imidazole, 4 M GuHCl). The protein was concentrated and applied to a S200 26/60 gel-filtration column pre-equilibrated in 50 mM Na_2PO_4 pH 7.42, 300 mM NaCl, 5mM Imidazole, 4 M GuHCl. Fractions containing the protein were pooled and concentrated. The concentrated protein was flash-frozen and stored at -80 °C. Prior to experiments, the purified L-FRQ was exchanged into a non-denaturing buffer (50 mM Na_2PO_4 pH 7.42, 300 mM NaCl, 200 mM arginine, 10% glycerol). For FFC preparations, GFB (50 mM HEPES pH 7.42, 200 mM NaCl, 10% glycerol) was used.

SEC-MALS

Protein samples between 50-75 µM were injected onto a Superose-6 (GE Lifesciences), equilibrated with 50 mM Na_2PO_4 pH 7.42, 300 mM NaCl, 200 mM arginine, 10% glycerol. The gel filtration column was coupled to a static 18-angle light scattering detector (DAWN HELEOS-II) and a refractive index detector (Optilab T-rEX) (Wyatt Technology, Goleta, CA). Data were collected every second at a flow rate of 0.5 mL/min and analysis was carried out using ASTRA VI, yielding the molar mass and mass distribution (polydispersity) of the sample. The monomeric fraction of BSA (Sigma, St. Louis, MO) was used to standardize the light scattering detectors and optimize data quality control.

SEC-SAXS

X-ray scattering experiments were performed at the Cornell High Energy Synchrotron Source (CHESS) G1 station. Size-exclusion chromatography-coupled SAXS (SEC–SAXS) experiments were performed using a Superdex 200 5/150 GL or Superose-6 5/150 GL (3 mL) column operated by a GE AKTA Purifier at 4 °C with the elution flowing directly into an in-vacuum X-ray sample cell. Samples were centrifuged at 14,000×g for 10 min at 4 °C before loading onto a column pre-equilibrated in a matched buffer (50 mM Na_2PO_4 pH 7.42, 300 mM NaCl, 200 mM arginine, 10% glycerol). Samples were eluted at flow rates of 0.05–0.1 mL min^{-1}. For each sample, 2-

s exposures were collected throughout elution until the elution profile had returned to buffer baseline, and scattering profiles of the elution buffer were averaged to produce a background-subtracted SEC–SAXS dataset. The scattering data was processed in RAW (2) and regions with homogenous R_G values were chosen for the Guinier and Kratky analyses.

Protein Labeling for ESR Spectroscopy and Microscopy Experiments

Sort/AEP1-tagged FRQ was cloned into a pET28a (Novagen) vector as described in (3). The production of the sortaseA and AEP1 enzymes and their associated peptides (GGGGC for Sortase A and CNGL for AEP1) is also described in (3) and (4), respectively. To enzymatically label FRQ with either SortaseA or AEP1, 1:1 protein and SortaseA/AEP1 as well as 2x excess peptide were incubated overnight to allow the reaction to reach completion. The following day, the reaction was run on a Superpose 6 10/300 analytical SEC column to remove unreacted peptide and enzyme, and the protein containing fractions were pooled and concentrated. To prepare the ADP-β-S-SL samples, ADP-β-S-SL was synthesized as previously reported in (5). 100 µL of buffer containing 100 µM of ADP-β-S-SL and 100 µM FRH or CK1 were incubated overnight and then washed with GFB (50 mM HEPES pH 7.42, 200 mM NaCl, 10% glycerol) through 5x dilution followed by re-concentration to remove any unbound ADP-β-S-SL. 30 µL of each sample were used for data collection. For MTSL labeling, the protein was incubated with 5x molar excess of MTSL overnight to allow the reaction to reach completion. The following day, the reaction was run on HiTrap Desalting column (GE Life Sciences) to remove any excess spin-label.

To prepare fluorophore labeled FRH and CK1, the purified proteins were incubated with 2x molar excess Cy5-maleimide dye (GE Life Sciences) overnight to label any accessible cysteines on the target proteins. The excess dye was removed by running the reaction over a HiTrap desalting column equilibrated with GFB (50 mM HEPES pH 7.42, 200 mM NaCl, 10% glycerol).

ESR Spectroscopy

The cw-ESR spectra were obtained on a Bruker E500 ESR spectrometer operating at X-band (~9.4 GHz). The ESR measurements were carried out at room temperature with a modulation amplitude of 0.5-2 G and modulation frequency of 100 kHz. All spectra were digitized to 1024 points with a sweep width of ~100 G and an average of 4–16 scans. The δ values were calculated using the display distance function on the Bruker Xepr 2.6b software.

For 4-pulse-DEER, the samples were exchanged into deuterated buffer containing 30% d8-glycerol, checked by cwESR and plunge frozen into liq. N_2. The pulse ESR measurements were carried out at Q-band (~34 GHz) on a Bruker E580 spectrometer equipped with a 10 W solid state amplifier (150W equivalent TWTA) and an arbitrary waveform generator (AWG). DEER was carried out using four pulses (π/2-τ1-π-τ1-πpump-τ2-π-τ2-echo) with 16-step phase cycling at 60 K. The pump and probe pulses were separated by 56 MHz (~20 G). The DEER data was background subtracted with an exponential background and the distance reconstruction was carried out using denoising and the SVD method (6, 7). Uncertainty in the distance distributions were calculated using the method described by Srivastava and Freed (8)

Turbidity Assays

Cysteine-less (Cys-less) FRQ was created in order to minimize disulfide cross-linking at the high protein concentrations needed for *in vitro* phase separation experiments. For this, all four of the native cysteine residues were substituted to serine using standard molecular biology techniques. For the phase separation experiments, this cys-less FRQ variant was purified and concentrated for dialysis into phase separation buffer. Cys-less FRQ was dialyzed overnight into 25 mM Tris pH 8.0, 150 mM NaCl or 25 mM Tris pH 8.0, 500 mM NaCl for p-FRQ and np-FRQ, respectively, at 4 °C. Phase separation was induced by dialyzing in either low or high salt and increasing temperature. The protein solution can be observed as a clear solution before dialysis and then turbid after dialysis indicating LLPS has occurred. Microscopy (See *DIC and Fluorescence Microscopy*) confirmed the turbid solution is due to LLPS and not aggregation. Turbidity as measured by absorbance at 300 - 600 nm was then recorded by an Agilent Cary 3500 UV-Vis Multicell Peltier using a temperature ramp rate of 5 °C/min increasing from 10 °C to 80 °C. A final protein concentration of 5-15 µM was used for all turbidity assays.

DIC and Fluorescence Microscopy

Phase separation was induced via overnight dialysis into 25 mM Tris pH 8.0, 150 mM NaCl or 25 mM Tris (pH 8.0), 500 mM NaCl for p-FRQ and np-FRQ, respectively. The final protein concentration used for microscopy were 5-15 µM. Images were taken on the Zeiss Axio Imager Z1 Upright Trinocular Fluorescence Microscope at 100x magnification. To test the interaction of FRQ with FRH and CK1, the phase separated FRQ was spiked with equimolar Cy5-labeled FRH or Cy5-labeled CK1 for final concentrations of 5 µM.

In vitro Phosphorylation Assay

22 µL samples containing 10-15 µM LFRQ were incubated in the presence or absence of 2 µM CK1 and 2 µM FRH in 50 mM Na_2HPO_4 pH 7.5, 300 mM NaCl, 200 mM arginine, 10 mM $MgCl_2$. For the LLPS assay, FRQ was dialyzed into a buffer containing 50 mM Na_2HPO_4 pH 7.5, 300 mM NaCl and 10 mM $MgCl_2$. 2 µL of 2.5 mM cold ATP and 0.1 µM of [γ-32P] ATP (3000 Ci/mmol, PerkinElmer) was added to produce a total volume of ~25 µL. After ~1h of incubation, the reaction was quenched with 25 µL of 4x SDS with 10 mM EDTA pH 8.0 and then subjected to gel electrophoresis on a 4–20% gradient Tris-glycine gel. The gel was dried with a GelAir dryer (Bio-Rad) and placed in an imaging cassette for at least 16 h and then imaged with a phosphorimager (GE, Typhoon FLA 7000). Band intensities we quantified using the Fiji software.

Mass Spectrometry and Phospho-Peptide Analysis

Phosphorylated FRQ was run on a 4-20% tris-glycine SDS-PAGE gel and the corresponding band was excised and trypsin digested. The digests were reconstituted in 0.5% formic acid (FA) for nanoLC-ESI-MS/MS analysis. The analysis was carried out using an Orbitrap Fusion™ Tribrid™ (Thermo-Fisher Scientific, San Jose, CA) mass spectrometer equipped with a nanospray Flex Ion Source, and coupled with a Dionex UltiMate 3000 RSLCnano system (Thermo, Sunnyvale, CA). The peptide samples (5 µL) were injected onto a PepMap C-18 RP nano trapping column (5 µm, 100 µm i.d x 20 mm) and then separated on a PepMap C-18 RP nano column (2 µm, 75 µm x 25 cm) at 35 °C. The Orbitrap Fusion was operated in positive ion mode with spray voltage set at 1.5 kV and source temperature at 275 °C. External calibration for FT, IT and quadrupole mass analyzers was performed. In data-dependent acquisition (DDA) analysis, the instrument was operated using FT mass analyzer in MS scan to select precursor ions followed by 3 second "Top Speed" data-dependent CID ion trap MS/MS scans at 1.6 m/z quadrupole isolation for precursor peptides with multiple charged ions above a threshold ion count of 10,000 and normalized collision energy of 30%. MS survey scans at a resolving power of 120,000 (fwhm at m/z 200), for the mass range of m/z 375-1600. Dynamic exclusion parameters were set at 35 s of exclusion duration with ±10 ppm exclusion mass width. All data were acquired under Xcalibur 4.4 operation software (Thermo-Fisher Scientific). The DDA raw files with MS and MS/MS were subjected to database searches using Proteome Discoverer (PD) 2.4 software (Thermo Fisher Scientific, Bremen, Germany) with the Sequest HT algorithm. The database search was conducted against a *Neurospora Crassa* database and *E. coli* database with adding LFRQ sequence into the database. The peptide precursor tolerance was set to 10 ppm and fragment ion tolerance was set to 0.6 Da with 2 miss-cleavage allowed for either trypsin or chymotrypsin. Variable modification

of methionine oxidation, deamidation of asparagines/glutamine, phosphorylation of serine/threonine/tyrosine, acetylation, M-loss and M-loss+acetylation on protein N-terminus and fixed modification of cysteine carbamidomethylation, were set for the database search. Only high confidence peptides defined by Sequest HT with a 1% FDR by Percolator were considered for confident peptide identification. All the mass spectra of phosphorylated peptides were double checked by manually inspection.

Integrated Bioinformatic Analyses and Computational Biology
All canonical protein and UniRef50 sequences for FRQ, FRH and CK1, and SWISS-Prot obtained proteome sequences of *Neurospora crassa*, *Drosophila melanogaster* and *Homo sapiens* were downloaded from the UniProt database (9) A combination of **(a)** in-house written Python scripts to allow us to analyze large datasets, **(b)** localCIDER (10), **(c)** IUPred2A (11), **(d)** Agadir (12), **(e)** AlphaFold 2.0 (13), **(f)** PScore (14) and (g) catGRANULE (15) were used to perform integrated bioinformatics analyses and computational biological studies of our proteins of interest. Unless mentioned, all programs were used by applying the default parameters. All scripts used to generate the data in this manuscript are provided as SI Datasets.

Neurospora crassa Strain Generation
Construction of the *frq[mNeonGreen]*, *son-1[mApple]* strain used for imaging was previously described (16). Briefly, a plasmid was constructed to target the endogenous *frq* locus with DNA encoding *N. crassa* codon-optimized mNeonGreen appended to the C-terminus of the *frq* ORF with a flexible linker. A gene encoding hygromycin B phosphotransferase (HPH) was introduced downstream from the 3' UTR of *frq* for selection. A plasmid was constructed containing the *N. crassa son-1* ORF with mApple appended to the C-terminus with a flexible linker for insertion at the *csr-1* locus. Transformation cassettes were amplified by PCR using NEB Q5® High-Fidelity 2X Master Mix (New England Bio Labs Cat. # M0492) and integrated into the genome by homologous recombination after electroporation of conidiophores. A homokaryotic strain containing the fluorescent fusion proteins was used for imaging.

Neurospora crassa Growth
N. crassa homokaryons containing *frq[mNeonGreen]* and *csr-1::son-1[mApple]*, were grown overnight on 2% agar gel pads made with Vogel's minimal medium (17) in constant light at 25 °C. Before imaging, a section near the edge of the growing mycelium was removed and inverted on a chambered cover glass (Thermo Scientific, Cat. # 155360).

Live Cell Image Acquisition and Processing

Images were acquired using a Zeiss AxioObserver laser scanning confocal microscope with a 100X/1.46 NA Oil immersion Plan-Apochromat objective. 561 nm and 488 nm lasers were used at 2% power with 2X averaging and a gain of 800 for detection of fluorescent signals by GaAsp PMTs. Images were processed for display using FIJI (18). The cropped images displaying a single nucleus were enlarged 10X using bicubic interpolation and contrasted to the minimum and maximum gray level values. A multi-color lookup table was applied to the FRQ[mNeonGreen] channel to better observe heterogeneity of the signal, and the SON-1[mApple] channel was pseudo-colored cyan to better contrast with the lookup table applied to the FRQ[mNeonGreen] channel. The surface plot of FRQ[mNeonGreen] was generated from the raw image. Movie S1, and the image in Figure 6E were acquired using a Nikon Ti-E based system, equipped with an Andor W1 spinning disk scan head, an Andor Zyla sCMOS camera, a 488 nm laser, and Chroma bandpass filters. Figure 6E is a still image from a single focal plane from a z-stack that was deconvolved using Nikon Elements and was acquired using a 100X/NA1.45 plan apo oil immersion objective. Figure 6E and Movie S1 were acquired using a 60X/NA1.45 plan apo oil immersion objective. The 1-minute movie was acquired with a 300 ms exposure per frame. Image processing for display was carried out using FIJI.

Results:

Table S1: SAXS parameters for p-FRQ, np-FRQ and their reconstitution into the FFC

	R_g [Guinier] (Å)	R_g [P(r)] (Å)	Dmax (Å)	MW [V_C] (kDa)
p-FRQ	123 ± 7	142 ± 5	530 ± 60	816
np-FRQ	100 ± 2	116 ± 4	446 ± 40	680
p-FFC	101 ± 2	136 ± 5	510 ± 20	314
np-FFC	80 ± 2	109 ± 7	480 ± 10	286

MW calculations for p-FRQ and np-FRQ are inaccurate because V_C (Volume of Correlation) plots (Integrated area of I(q)q vs q; q = $2\pi/d$) did not converge for these species. Expected MW for LFRQ is 110 kDa. Expected FFC MW for the 1:1:1 stoichiometry of FRQ:FRH:CK1 Is 273 kDa. Errors in Dmax were estimated from the range for which P(r) tailed to zero in the high distance range without constraint.

Table S2: FRQ Cysteine residue locations and context

Site	Domain	Binding Partner at this region	Reference
370C	N/A		
490C	FCD2	CK1	(19)
658C	N/A		
746C	FFD	FRH	(20)

Table S3: FRQ regions that bind clock proteins

Binding Partner	Region	Cysteine residue(s) present in this region	Reference
FRH	695-778	746C	(20)
CK1	319-326; 488-495	490C	(19)
WC2		490C,658C,746C	(21)

Table S4: δ values from the cw-ESR central peak for FRQ variants of Fig. 3C

Site	np-FRQ (G)	p-FRQ (G)
NT	1.75	1.87
490C	1.97	2.47
658C	1.97	2.01
746C	1.76	1.82
CT	2.30	1.70

Table S5: R-square values for the quality of the two component gaussian fits to the distance distributions.

	370	490	658	746
370		0.99, 0.99	0.96, 0.99	NA, 0.99
490			0.94, 0.99	0.99, 0.99
658				0.99, 0.95
746				

Note: There are two values reported in each cell – the first value corresponds to the R-square value for FRQ only distance while the second value is for the same variant in the FFC complex (i.e., when the FRQ is bound to CK1a and FRH).

Table S6: Calculated values from the UV-Vis turbidity assays of np-FRQ shown in Figure 6C.

	5 µM	10 µM	15 µM
Mean Initial Baseline Absorbance	0.02	0.08	0.13
Temperature of Initiation (°C)	56	23	21
Cloud Point (Inflection Point) (°C)	68	58	57
Absorbance at Inflection Point	0.07	0.21	0.33
Slope at Inflection Point	0.01	0.02	0.03
Mean Final Baseline Absorbance	0.14	0.47	0.64

Note: Initiation temperature was determined as baseline + 5X standard deviation of the baseline.

Table S7: Calculated values from the UV-Vis turbidity assays of p-FRQ shown in Figure 6C.

	5 µM	10 µM	15 µM
Mean Initial Baseline Absorbance	0.09	0.21	0.46
Temperature of Initiation (°C)	45	42	24
Cloud Point (Inflection Point) (°C)	-	-	-
Absorbance at Inflection Point	-	-	-
Slope	0.00	0.01	0.02
Mean Final Baseline Absorbance	-	-	-

Note: Initiation temperature was determined as baseline +5X standard deviation of the baseline. Also, because p-FRQ continues to undergo LLPS within the temperature range of the experiment, we cannot calculate a cloud point.

A   **B**

Figure S1: (A) AlphaFold model of LFRQ colored by pLDDT score. (B) Predicted error alignment plot that shows the uncertainty in the relative positions of two residues in the structure. (Image source: https://www.uniprot.org/uniprotkb/P19970/entry#structure)

Figure S2: Domain map of FRQ with phospho-sites (as identified by mass-spec) highlighted. The sites in red were also identified in (22, 23).

Figure S3: Pair-wise distance distributions calculated from SAXS data for p-FRQ (red), np-FRQ (dark blue), p-FFC (magenta), and np-FFC (light blue).

Figure S4: Enzymatic labeling of FRQ at its termini**.** X-band (9.8 GHz) RT cw-ESR measurements for the spin-labeled peptide by itself and when it has been grafted onto either the N or C-terminus of FRQ.

Figure S5: Time domain traces before (Raw) and after (data) background correction and trace fits (fit) from DEER experiments of different double-cysteine FRQ variants alone or within the FFC (i.e., when FRQ was bound to FRH and CK1a). The spins were present on the residue number of FRQ that is highlighted for example '370-490' refers to FRQ spin-labeled at the 370C and 490C site and the DEER experiment probed the dipolar coupling of the spins at these sites. The last graph shows the time domain data for the control single cysteine DEER experiment with p-FRQ 490C.

Figure S6: Baseline-corrected time domain fit traces from Q-band DEER experiments of different double-cysteine FRQ mutants alone (blue) and within the FFC (orange). Original data are shown in Supplementary Figures S5 and S9.

Figure S7: Full autoradiograph of np-FRQ phosphorylation by CK1 and γ-^{32}P-ATP from Figure 4A inset. Four technical replicates are shown for each np-FRQ concentration.

Figure S8: Background-corrected time domain trace and error analyses from DEER experiments targeting the ATP-binding pockets of FRH and CK1. The DEER experiment probed the dipolar coupling of the spins in the ADP-β-S-SL molecule that was present in ATP-binding pocket of FRH or CK1.

Figure S9: Error analysis of the distance distributions produced from DEER experiments of different double-cysteine FRQ variants. The spins were present on the amino acid number of FRQ that is highlighted for example '370-490' refers to FRQ spin-labeled at the 370C and 490C site and the DEER experiment probed the dipolar coupling of the spins at these sites.

Figure S10: Error analysis of the distance distributions produced from DEER experiments of different double-cysteine FRQ variants within the FFC (i.e., when FRQ was bound to FRH and CK1). The spins were present on the amino acid number of FRQ that is highlighted for example '370-490' refers to FRQ spin-labeled at the 370C and 490C site and the DEER experiment probed the dipolar coupling of the spins at these sites.

Figure S11: (Left)The two component Gaussian fitting of the distance distribution (P[r]) obtained from SVD for FFC 370-490 (i.e., spins were placed on resides 370 and 490 on FRQ, and this protein was bound to CK1 and FRH). No constraints were placed on the width, position, or amplitude of either component and the best fit values were obtained by minimizing the difference between the fits and the SVD distance distributions. Similar fits were carried out for all the different variants. The quality of the fit is reported by the R-square values and ranged between 0.94-0.99 for all the samples tested. They are reported in Table S5. (Right) The proportion of the closer distance peak in the distance distributions of FRQ (blue) or FRQ+FRH+CK1(FFC) (red) samples. The error bars reflect the uncertainty in estimating the population of each peak and were derived from uncertainties in the distance distributions shown in *SI Appendix*, Fig. S9 and S10.

Figure S12: FRQ, but not FRH or CK1, is highly disordered and predicted to undergo LLPS. Shown are the disorder and LLPS propensity scores of the members for the FFC complex, FRQ, FRH and CK1. Columns from left to right are disorder and phase separation predictions for FRQ, FRH and CK1, respectively. Disorder graphs (red) were computed using the IUPred2A program (11) where values above a 0.5 threshold (dashed line) are considered disordered. LLPS propensity predictions were computed using "Pi-Pi" (blue) (14) and catGRANULE (light blue) (15). Shown on the top right corner of each graph are the percent disorder or LLPS propensity scores of each protein.

Figure S13: FRQ and its functional homologs are predicted to undergo LLPS. Shown are the disorder and LLPS propensity scores for FRQ, dPER1 and hPER1. Data for the disorder graphs (red) were computed using the IUPred2A program where values above a 0.5 threshold (dashed line) are considered disordered. Pi-pi predictions (blue) were computed using (14), while catGRANULE predictions (light blue) were computed using catGRANULE server (15). Shown on the top-right corner of each graph are the percent disorder, PScore or catGRANULE score for each protein. FRQ, dPER1 and hPER1 are all predicted to be highly disordered (86.77%, 58.84% and 63.07%, respectively) and undergo LLPS.

Figure S14: FRQ and its homologs, dPER1 and hPER1, have similarly disparate physicochemical properties compared to their proteomes. (A) Shown are the distribution of fraction of charged residues (FCR), net charge per residue (NCPR), and hydropathy along the sequence of FRQ, dPER1 and hPER1. (B) Sequence characteristics of FRQ and its homologs, dPER1 and hPER1. Various protein sequence parameters are calculated using localCIDER (10) and displayed including fraction of charged residues (FCR), the degree of charge mixing (Kappa), net charge per residue (NCPR), and hydropathy (the hydrophobic and hydrophilic character of protein chain as defined by (24)) for the UniProtRef50 clusters of FRQ (purple), dPER1(blue) and hPER1 (orange). For comparison are shown protein sequences of similar length from the SWISS-Prot reviewed proteomes of *Neurospora crassa* (light purple), *Drosophila melanogaster* (light blue) and *Homo sapiens* (light orange). Hydropathy is consistently low for the clock proteins and FRQ orthologs have a high degree of charge separation, i.e., tracks of negative and positive residues (high kappa). High kappa values are common for proteins that phase separate (25). (C) Classification of the charge properties of the Ref50 clusters of FRQ and its functional homologs, dPER1 and hPER1. Das-Pappu phase plots showing the fraction of negatively vs positively charge residues for the UniProtRef50 clusters of FRQ (purple), *Neurospora crassa* proteome control (light purple), dPER1 (blue), *Drosophila melanogaster* proteome control (light blue), hPER1 (orange) and Homo sapiens proteome control (light orange). Shown are the fraction of negative residues versus fraction of positive residues for each cluster of homologs and their controls (similar size proteins from their respective proteomes). Despite the lack of amino acid conservation between the functional homologs of FRQ, dPER1 and hPER1, the ratio of fraction negative to fraction positive residues are similar among the clock proteins. dPER and hPER1 have a similarly low value of both fraction positive and fraction negative compared to their respective proteomes, whereas the fraction positive and fraction negative residues are both higher in FRQ. Note: Ref50 is a clustering method utilized by UniProt and is defined in (26).

Figure S15: FRQ and its functional homologs dPER1 and hPER1 have similar amino acid composition compared to their respective proteomes. FRQ and its functional homologs dPER1 and hPER1 have similar amino acid compositions, which differs substantially from that of their respective proteomes. The amino acid composition for FRQ, dPER1 and hPER1 and their respective homolog clusters and proteomes are shown. Generally, G, S, T and P residues are enriched whereas hydrophobic (A, I, L, V) residues are relatively depleted for FRQ, dPER1, hPER1 and their homologs compared to their proteomes. Unlike dPER1 and hPER1 and their homologs, FRQ and its homologs are also highly enriched in R and D.

Figure S16: Phosphorylation modulates the physicochemical properties of FRQ. Shown are the distribution of fraction of charged residues (FCR), net charge per residue (NCPR), and hydropathy along the sequence of np and p-FRQ (top panel). In the bottom panel, shown are the disorder and LLPS propensity scores for np and p-FRQ. Data for the disorder graphs (red) were computed using the IUPred2A program (11) where values above a 0.5 threshold (dashed line) are considered disordered. Pi-pi predictions (blue) were computed using, while catGRANULE predictions (light blue) were computed using catGRANULE server (15). Shown on the top-right corner of each graph are the percent disorder, PScore or catGRANULE score for each protein. Note: Glu(E) residues were employed as phospho-mimetics in all analysis.

Figure S17: Exploring the phase behavior of FRQ. (A) (Left) UV-Vis absorption spectrum of p-FRQ (in 150 mM NaCl, 25 mM Tris pH 8.0) and (Right) np-FRQ (in 500 mM NaCl, 25 mM Tris pH 8.0) as a function of temperature to check for reversibility of LLPS transition. (B) UV-Vis absorption spectrum of p-FRQ (Left) and np-FRQ (Right) as a function of temperature in either low salt (150 mM) or high salt (500 mM). (C) UV-Vis absorption spectrum of p-FRQ and np-FRQ in a non LLPS promoting buffer as a function of temperature. Note: p-FRQ appears to undergo a small amount of LLPS even in non-LLPS buffer as seen by its relatively high absorbance relative to np-FRQ. (D) DIC micrograph of FRQ showing droplets fusing and docking indicated within red circle.

Figure S18: Full radiograph of np-FRQ phosphorylation by CK1 and γ-^{32}P-ATP from Figure 7.

 Figure S19: Schematic diagram summarizing a potential mechanism for LLPS mediated circadian clock regulation. By reducing CK1 activity, especially at higher temperatures, LLPS provides a potential temperature compensation mechanism for the clock.

Movie S1: Visualization of FRQ [mNeonGreen] in living Neurospora, displaying heterogeneous, punctate intranuclear localization. Images of a single focal plane were acquired in a time series with 300 ms between exposures.

References:

1.   C. Querfurth *et al.*, Posttranslational regulation of Neurospora circadian clock by CK1a-dependent phosphorylation. *Cold Spring Harbor symposia on quantitative biology* **72**, 177-183 (2007).

2.   J. B. Hopkins, R. E. Gillilan, S. Skou, BioXTAS RAW: Improvements to a free open-source program for small-angle X-ray scattering data reduction and analysis. *Journal of Applied Crystallography* **50**, 1545-1553 (2017).

3.   S. Chandrasekaran *et al.*, Tuning flavin environment to detect and control light-induced conformational switching in Drosophila cryptochrome. *Communications Biology 2021 4:1* **4**, 1-12 (2021).

4.   G. K. T. Nguyen, Y. Cao, W. Wang, C. F. Liu, J. P. Tam, Site-Specific N-Terminal Labeling of Peptides and Proteins using Butelase 1 and Thiodepsipeptide. *Angewandte Chemie-International Edition* **54**, 15694-15698 (2015).

5.   A. R. Muok, T. K. Chua, H. Le, B. R. Crane, Nucleotide Spin Labeling for ESR Spectroscopy of ATP-Binding Proteins. *Applied Magnetic Resonance* **49**, 1385-1395 (2018).

6.   M. Srivastava, J. H. Freed, Singular Value Decomposition Method to Determine Distance Distributions in Pulsed Dipolar Electron Spin Resonance. *Journal of Physical Chemistry Letters* **8**, 5648-5655 (2017).

7.   M. Srivastava, E. R. Georgieva, J. H. Freed, A New Wavelet Denoising Method for Experimental Time-Domain Signals: Pulsed Dipolar Electron Spin Resonance. *Journal of Physical Chemistry A* **121**, 2452-2465 (2017).

8.   M. Srivastava, J. H. Freed, Singular Value Decomposition Method To Determine Distance Distributions in Pulsed Dipolar Electron Spin Resonance: II. Estimating Uncertainty. *Journal of Physical Chemistry A* **123**, 359-370 (2019).

9.   A. Bateman *et al.*, UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Research* **49**, D480-D489 (2021).

10.   A. S. Holehouse, R. K. Das, J. N. Ahad, M. O. G. Richardson, R. V. Pappu, CIDER: Resources to Analyze Sequence-Ensemble Relationships of Intrinsically Disordered Proteins. *Biophysical Journal* **112**, 16-21 (2017).

11.   B. Mészáros, G. Erdös, Z. Dosztányi, IUPred2A: context-dependent prediction of protein disorder as a function of redox state and protein binding. *Nucleic Acids Research* **46**, W329-W337 (2018).

12.   V. Muñoz, L. Serrano, Elucidating the folding problem of helical peptides using empirical parameters. *Nature Structural Biology* **1**, 399-409 (1994).

13.   J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature 2021 596:7873* **596**, 583-589 (2021).

14.   R. M. Vernon *et al.*, Pi-Pi contacts are an overlooked protein feature relevant to phase separation. *Elife* **7**, e31486 (2018).

15.   B. Bolognesi *et al.*, A Concentration-Dependent Liquid Phase Separation Can Cause Toxicity upon Increased Protein Expression. *CellReports* **16**, 1-10 (2016).

16.   B. M. Bartholomai (2022) Cell biological dissection of the Neurospora circadian oscillator: Tools, approaches and fundamental observations. in *Biochemistry and Genetics* (Dartmouth College, Hanover, NH).

17.   H. J. Vogel, A convenient growth medium for Neurospora (Medium N). *Microb. Genet. Bull.* **13**, 42-43 (1956).

18.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676-682 (2012).

19.   C. Querfurth *et al.*, Circadian Conformational Change of the Neurospora Clock Protein FREQUENCY Triggered by Clustered Hyperphosphorylation of a Basic Domain. *Molecular Cell* **43**, 713-722 (2011).

20.   P. Cheng, Q. He, Q. He, L. Wang, Y. Liu, Regulation of the Neurospora circadian clock by an RNA helicase. *Genes and Development* **19**, 234-241 (2005).

21.   J. Guo, P. Cheng, Y. Liu, Functional significance of FRH in regulating the phosphorylation and stability of Neurospora circadian clock protein FRQ. *J. Biol. Chem.* **285**, 11508-11515 (2010).

22.   C. L. Baker, A. N. Kettenbach, J. J. Loros, S. A. Gerber, J. C. Dunlap, Quantitative Proteomics Reveals a Dynamic Interactome and Phase-Specific Phosphorylation in the Neurospora Circadian Clock. *Molecular Cell* **34**, 354-363 (2009).

23.   C. T. Tang *et al.*, Setting the pace of the Neurospora circadian clock by multiple independent FRQ phosphorylation events. *Proc. Natl. Acad. Sci. USA* **106**, 10722-10727 (2009).

24.   J. Kyte, R. F. Doolittle, A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology* **157**, 105-132 (1982).

25.   R. Somjee, D. M. Mitrea, R. W. Kriwacki (2020) Exploring Relationships between the Density of Charged Tracts within Disordered Regions and Phase Separation. in *Pacific Symposium on Biocomputing* (Hi), pp 207-218.

26.   M. Steinegger, J. Söding, Clustering huge protein sequence sets in linear time. *Nature Communications* **9**, 2542 (2018).